

Phase transition in presence of surfactants: from discrete to continuum

ROBERTO ALICANDRO, MARCO CICALESE AND LAURA SIGALOTTI

Abstract

We study by Γ -convergence the atomistic-to-continuum limit of the Blume-Emery-Griffiths model describing the phase transition of a binary mixture in presence of a third surfactant phase. In the case of low surfactant concentration we study the dependence of the surface tension on the density of the surfactant and we show the microstructure of the ground states. We then consider more general (n -dimensional) energies modeling phase transitions in presence of different species of surfactants and, in the spirit of homogenization theory, we provide an integral representation result for their Γ -limit. As an application we study the ground states of the system for prescribed volume fractions of the phases.

Contents

1	Introduction	1
2	Notation and preliminaries	7
3	The Blume-Emery-Griffiths model	8
3.1	A brief description of the model: from bulk to surface scaling . . .	9
3.2	Low concentration of surfactants: discrete-to-continuum limit . . .	12
4	More general models	14
4.1	The main result	16
4.2	Prescribed volume-fractions	33
5	The Blume-Emery-Griffiths model: proof of Theorem 3.3	34

1 Introduction

In recent years a great effort has been made in order to study the atomistic-to-continuum limit of several discrete systems and in particular of Ising-type spin

systems (see for example [2], [3], [4], [7], [10], [11], [18], [27], [29] and [14] (Chapter 11) for a review on this subject). Despite a simple structure, the energy functionals of such systems (depending on the choice of the order parameter, the range of interactions and the scaling properties) may model a great variety of physical phenomena. An instructive example in this respect is provided by the functional

$$E_\varepsilon(u) = - \sum_{n.n.} \varepsilon^2 u(a) \cdot u(b), \quad (1.1)$$

which is the energy of a system of nearest-neighbors interacting spins defined on the points of a ε -lattice and taking values either in $\{\pm 1\}$ or in S^1 . The energy accounts for the interactions among nearest neighbors (*n.n.*); i.e., $a, b \in \varepsilon\mathbf{Z}^2 \cap \Omega$ such that $|a - b| = \varepsilon$ ($\Omega \subset \mathbf{R}^2$ being a bounded open set). The field u is defined on the lattice $\varepsilon\mathbf{Z}^2 \cap \Omega$ and takes values either in $\{\pm 1\}$ or in S^1 (in the latter case $u(a) \cdot u(b)$ denotes the scalar product in \mathbf{R}^2). On one hand, in the case $u \in \{\pm 1\}$ it has been proved in [2] that the discrete-to-continuum limit, as $\varepsilon \rightarrow 0$, of suitable power scaling of (1.1) resembles that of a Cahn-Hilliard type functional. Namely, it leads to a class of anisotropic surface tension energies of the form $\int_{\partial E} \varphi(\nu_E) d\mathcal{H}^1$, where ν_E stands for the inner normal to ∂E which, in turns, represents the interface between the $\{u = +1\}$ phase and the $\{u = -1\}$ phase. On the other hand, the vectorial case $u \in S^1$ leads to a completely different class of continuum limits. In [4] it has been proved that the asymptotic analysis, as $\varepsilon \rightarrow 0$, of a suitable logarithmic scaling of (1.1) resembles that of a complex Ginzburg-Landau functional. In particular, it leads to a limit energy model for the formation of vortex-type singularities described through the measure $\mu = \sum_{i=1}^N d_i \delta_{x_i}$ ($x_i \in \Omega$ being the location of the i -th singularity and $d_i \in \mathbf{Z}$ its topological degree) whose energy is of the type $|\mu|(\Omega)$.

In this paper we will prove that a different variant of the energies in (1.1) leads to a class of continuum functionals different from the two mentioned above. The model we consider can be described as follows. Given $\Omega \subset \mathbf{R}^2$ a bounded open set, we fix the two dimensional square lattice $\varepsilon\mathbf{Z}^2 \cap \Omega$ and we consider a ternary system driven by an energy defined on functions parameterized on the points of the lattice and taking only three different values (for simplicity, $-1, 0, 1$). In this framework, the values of u are usually thought to describe three admissible phases of the system. For a given configuration of particles, the free energy E of the system is given by

$$E(u) = - \sum_{n.n.} \varepsilon^2 (u(a)u(b) + k(u(a)u(b))^2). \quad (1.2)$$

Here $k > 0$ is the quotient between the so called bi-quadratic and quadratic exchange interaction strengths. The range of k , jointly with the scaling of the energy, will be chosen later as a result of a heuristic argument. The heuristic argument will justify the choice of the functional in (1.2) as the free energy of a system where the two phases $\{\pm 1\}$ coexist in presence of the phase $\{0\}$ representing a substance

called surfactant (a contraction for *surface-active-agent*) which, by being adsorbed onto the interfaces, may significantly reduce the surface tension of the system.

A variational description of the effects caused by the presence of surfactants in phase separation phenomena has been developed by modeling the physical system either as a continuum or as a discrete. Among the continuum theories, a first attempt to model phase transitions in presence of surfactants has been made by Laradji-Guo-Grant-Zuckermann in [24] and [25] (several generalizations have been further considered by Gompper and Schick in [23]) who proposed a variational model involving a two order parameters Ginzburg-Landau functional. Here one order parameter represents the local difference of density of the two phases (as in the standard Cahn-Hilliard model in the gradient theory of phase transitions) while the other one represents the local surfactant density. The two order parameters are energetically coupled to favor the segregation of the surfactant at the phase interface. The coarse-graining analysis of this model has been performed through Γ -convergence methods by Fonseca, Morini and Slastikov in [22] (the mathematical analysis of more general continuum models is the object of [1]). Concerning the discrete models, many of them use some variation of that in (1.2) which has been introduced by Blume, Emery and Griffiths (BEG) in [12] (see also [23] and the references therein). This model has been also object of interesting studies in the context of equilibrium statistical mechanics (see [21] for an exhaustive mean-field approach) and has been used to describe several ternary systems (e.g., solid-liquid-gas systems, semiconductor alloys, electronic conduction models).

In this paper we will perform a Γ -limit analysis of these energies and, as a result, we will be able to describe the behavior of the ground states of the BEG system as ε tends to 0. More in details, upon identifying arrays $\{u(a)\}$, $a \in \varepsilon\mathbf{Z}^2 \cap \Omega$ with their piecewise-constant interpolations, the energies E_ε can be interpreted as defined on (a subset of) $L^1(\Omega)$, and can therefore undergo a process of Γ -limit in that framework. As ε tends to 0, the Γ -limit E of E_ε is particularly simple, only giving the trivial constraint $|u| \leq 1$, and the constant (minimum) value $2|\Omega|(-1+k) \wedge 0$, corresponding to the uniform states $u = \pm 1$ and $u = 0$. By choosing $k < 1$ we set the uniform states $u = \pm 1$ to be the ground states. Note that, with this choice, the asymptotic analysis of the energy E_ε summarizes the fact that a sequence $(u_\varepsilon)_\varepsilon$ can arbitrarily mix the uniform states -1 and 1 at a mesoscopic scale with variation in energy from the value of the uniform states which is negligible as $\varepsilon \rightarrow 0$ (the asymptotic analysis of the bulk scaling of more general spin-type models has been performed in [6]). Thus, in order to have a better description of the ground states, in the spirit of *development by Γ -convergence* (see [17] for a general treatment of this topic and also [2], [5], [15], [16]), we select sequences that realize the minimum value with a sharper precision; *i.e.*, such that

$$E_\varepsilon(u_\varepsilon) = c_\varepsilon + O(\varepsilon),$$

where we have denoted by c_ε the absolute minimum of E_ε , that is $c_\varepsilon = \sum_{\text{n.n.}} \varepsilon^2(k-1)$. For such configurations the limit states u will take the values ± 1 only and the

limit energy will be an interfacial type energy which is interpreted as the surface tension of the system which undergoes a phase separation phenomenon between the phases $\{u = -1\}$ and $\{u = +1\}$. Thus, at this scaling, it is necessary to further specify the values of the parameter k in such a way that the phase 0 can be actually considered a surfactant phase (meaning that it contributes to lower the surface tension). In particular it can be easily showed (see Section 3) that, for $\frac{1}{3} < k < 1$, the energy for a transition from a -1 phase to a $+1$ phase is lowered if the surfactant particles are at the interface. With such a choice, the 0 phase can be rightly considered a surfactant phase and, for this reason, the previous scaling, in which in addition the measure of the phase 0 vanishes, is usually called the *low surfactant concentration regime*. We are interested in this scaling regime. Thus the scaled family of functionals we have to study is

$$E_\varepsilon^{(1)}(u) := \frac{E_\varepsilon(u) - c_\varepsilon}{\varepsilon} = \sum_{n.n.} \varepsilon(1 - u(a)u(b) - k(1 - (u(a)u(b))^2).$$

Observe that, within this scaling, the interaction energy for two particles of the same type -1 or $+1$ is zero, while the interaction of a surfactant particle 0 with respect to all the other particles is repulsive, its cost being the positive value $1 - k$. For this reason it is also said that the BEG functional describes a *repulsive surfactant model*. In Theorem 3.2 we show that $E_\varepsilon^{(1)}$ Γ -converges (in the $L^1(\Omega)$ -topology) to the interfacial-type energy functional

$$E^{(1)}(u) = \int_{S(u)} \psi(\nu_u) d\mathcal{H}^1,$$

where $u \in BV(\Omega; \{\pm 1\})$, $\psi(\nu) = (1 - k)(3|\nu_1| \vee |\nu_2| + |\nu_1| \wedge |\nu_2|)$ denotes the anisotropic surface tension of the model, $S(u)$ is the (essential) interface between the sets $\{u = 1\}$ and $\{u = -1\}$ and ν_u is the measure theoretic inner normal to $S(u)$.

As we have already observed, in the topology we have chosen, the limit order parameter u does not carry any information about the surfactant phase. Actually the role of the surfactant becomes clear when one looks at the minimizing microstructure leading to the computation of the limiting surface density ψ . In this direction, a natural further step in the analysis of the BEG model is the dependence of the surface tension of the continuum limit on the concentration of the surfactant. There is a big literature on this subject, both from the physical and the chemical point of view (see for example [23] and [26]). However no rigorous description of the microscopic geometry of the surfactant at the interface is present in literature and, as far as we know, all the attempts done to study this problem are based on merely numerical computations or on heuristic arguments. In order to rigorously address this problem we need to go beyond the standard formulation of the BEG model and we let the energy functional of the system depend explicitly on the distribution of the surfactant particles. To this end we set

$$I_0(u) = \{a \in \Omega_\varepsilon : u(a) = 0\},$$

and we introduce the following *surfactant measure*

$$\mu(u) = \sum_{a \in I_0(u)} \varepsilon \delta_a.$$

Then, with a slight abuse of notation, we can extend $E_\varepsilon^{(1)}$ to $L^1(\Omega) \times \mathcal{M}_+(\Omega) \rightarrow [0, +\infty]$ as

$$E_\varepsilon^{(1)}(u, \mu) = \begin{cases} E_\varepsilon^{(1)}(u) & \text{if } \mu = \mu(u) \\ +\infty & \text{otherwise.} \end{cases}$$

We observe that, in this discrete setting, the way we extend the functionals in order to track the energy of the surfactants is by decoupling the order parameter of the model instead of adding another variable as it has been done when dealing with continuum models (see [22] and [1]). We then endow the space $L^1(\Omega) \times \mathcal{M}_+(\Omega)$ with the topology $\tau_1 \times \tau_2$ where τ_1 denotes the strong topology in $L^1(\Omega)$ and τ_2 denotes the weak*-topology in the space of non-negative bounded Radon measures $\mathcal{M}_+(\Omega)$ and, by pursuing the Γ -limit with respect to this topology, in Theorem 3.3 we prove that $E_\varepsilon^{(1)}$ Γ -converges to the functional $E^{(1)} : L^1(\Omega) \times \mathcal{M}^+(\Omega) \rightarrow [0, +\infty]$ defined as

$$E^{(1)}(u, \mu) = \begin{cases} \int_{S(u)} \varphi\left(\frac{d\mu}{d\mathcal{H}^1|_{S(u)}}, \nu_u\right) d\mathcal{H}^1 + (2k-2)|\mu^s|(\Omega) & \text{if } u \in BV(\Omega; \{\pm 1\}) \\ +\infty & \text{otherwise,} \end{cases}$$

where the function $\varphi : \mathbf{R} \times S^1 \rightarrow [0, +\infty)$ is computed explicitly and its graphic is reported in Figure 1. As it is clear from the graphic of φ , an anisotropic threshold phenomenon occurs at the phase interface. Indeed, for a given $\nu \in S^1$ the surface tension $\varphi(z, \nu)$ decreases up to a certain value of the density z of the surfactant, namely $z = |\nu_1| \vee |\nu_2|$. Increasing further the density of the surfactant the surface tension may be constant, if the surfactants are not adsorbed onto the interface (in this case the singular part of the surfactant measure is increased), or otherwise it may increase. As an application of the previous Γ -convergence result, at the end of section 3 we address an optimization problem having as constraint the prescribed volume fractions of the different phases.

Inspired by the many different models of phase transitions in presence of surfactants studied in the physical/chemical literature, in section 4 we push forward the previous analysis to the case of a n -dimensional discrete systems driven by an energy accounting for quite general finite range pairwise interactions when different species of repulsive surfactant particles are present. For such a general system, in the spirit of homogenization theory, we obtain an integral representation result for the Γ -limit and we study some properties of its densities. More precisely, given $\Omega \subset \mathbf{R}^n$ and $u : \varepsilon\mathbf{Z}^n \cap \Omega \rightarrow K$ we define the functional F_ε as

$$F_\varepsilon(u) = \sum_{n.n.} \varepsilon^{n-1} f\left(\frac{b-a}{\varepsilon}, u(a), u(b)\right).$$




Figure 1: The local microstructure of a ground state of the BEG model at a fixed straight interface (the dashed line normal to ν) for three different values of the density of surfactants at the interface. Black, white and grey dots stand for the 0, +1 and -1 values of the spin field u , respectively.

where $R > 0$ is an interaction threshold, $K = \{m_1, m_2, s_1, s_2, \dots, s_M\} \subset \mathbf{R}$ describes the finite number of phases in the system and $f : \mathbf{Z}^n \times K^2 \rightarrow [0, +\infty)$ satisfies a sort of discrete isotropy condition (see Remark 4.1 and 4.5) and is such that $f(z, \cdot, \cdot)$ has $\{(m_1, m_1), (m_2, m_2)\}$ as absolute minima. As we have done in the case of the BEG model, to study the discrete-to-continuum limit of this discrete system we introduce the sets of points in $\varepsilon\mathbf{Z}^n \cap \Omega$ occupied by the different types of surfactant and suitable surfactant measures associated to them. For $l \in \{1, 2, \dots, M\}$ we set

$$I_l(u) := \{a \in \Omega_\varepsilon : u(a) = s_l\}, \quad I(u) := \bigcup_{l=1}^M I_l(u)$$

and we define

$$\mu_l(u) := \sum_{a \in I_l(u)} \varepsilon^{n-1} \delta_a, \quad \mu(u) = \{\mu_1(u), \mu_2(u), \dots, \mu_M(u)\}.$$

After extending F_ε to $L^1(\Omega) \times (\mathcal{M}_+(\Omega))^M \rightarrow [0, +\infty]$ as

$$F_\varepsilon(u, \mu) := \begin{cases} F_\varepsilon(u) & \text{if } \mu = \mu(u) \\ +\infty & \text{otherwise,} \end{cases}$$

in Theorem 4.4 we prove that F_ε Γ -converge with respect to the $\tau_1 \times \tau_2$ convergence (here τ_1 denotes the strong convergence in $L^1(\Omega)$ and τ_2 denotes the

weak*-convergence in $(\mathcal{M}_+(\Omega))^M$ to the functional

$$F(u, \mu) = \int_{S(u)} f_{hom} \left(\frac{d\mu}{d\mathcal{H}^{n-1} \llcorner S(u)}, \nu(u) \right) d\mathcal{H}^{n-1} + \int_{\Omega} g_{hom}(\mu^s) \quad (1.3)$$

if $u \in BV(\Omega; \{m_1, m_2\})$, $\mu = \frac{d\mu}{d\mathcal{H}^{n-1} \llcorner S(u)} \mathcal{H}^{n-1} \llcorner S(u) + \mu^s$, and $+\infty$ otherwise. The limit densities f_{hom} and g_{hom} are given by two asymptotic homogenization formulas in (4.19) and (4.20) respectively. About the derivation of the homogenization formulas, we want to stress the fact that, while for f_{hom} a standard argument in homogenization theory applies, the same does not hold true for g_{hom} . In this last case we are able to overcome the difficulties by combining some abstract arguments of measure theory with a reflection construction which exploits the discrete isotropy assumption on the interaction densities (see Remark 4.5).

As a final remark, we want to underline that in the models we consider here the surfactants are described as scalar particles without internal structure. More general models are known (see [19], [28] and [23]) where the surfactants are described as polar molecules whose heads and tails interact differently with the same phase. In this case it is also known that the presence of surfactants in a mixture may lead to self-assembling and that a number of different, and even topologically non trivial, microstructures may appear. We hope that the analysis performed in this paper may provide the basis to address the discrete-to-continuum limit for these systems.

Acknowledgements The work by Marco Cicalese was partially supported by the European Research Council under FP7, Advanced Grant n. 226234 ‘‘Analytic Techniques for Geometric and Functional Inequalities’’. This work is part of the third author’s Ph.D. thesis at the Department of Mathematics of Sapienza, Università di Roma.

2 Notation and preliminaries

In what follows, given $x, y \in \mathbf{R}^n$ we denote by (x, y) the usual scalar product in \mathbf{R}^n and we set $|x| = \sqrt{(x, x)}$. Moreover we denote by $\|\cdot\|_1$ the l_1 -norm in \mathbf{R}^n defined as $\|x\|_1 = |x_1| + \dots + |x_n|$. Given $t > 0$, we will denote by $[t]$ the integer part of t . For any measurable $A \subset \mathbf{R}^n$ we denote by $|A|$ the n -dimensional Lebesgue measure of A . Let Ω be a bounded open subset of \mathbf{R}^n with Lipschitz boundary. For fixed $\varepsilon > 0$ we consider the lattice $\varepsilon\mathbf{Z}^n \cap \Omega =: \Omega_\varepsilon$. Given $K \subset \mathbf{R}$ we denote by $\mathcal{A}_\varepsilon(\Omega; K)$ the set of functions

$$\mathcal{A}_\varepsilon(\Omega; K) := \{u : \Omega_\varepsilon \rightarrow K\}.$$

Remark 2.1 A function $u \in \mathcal{A}_\varepsilon(\Omega; K)$ will be identified with its piecewise-constant interpolation still denoted by u and given by $u(x) = u(z_x^\varepsilon)$, where $z_x^\varepsilon \in \mathbf{Z}^n$ is the closest point to x (which is uniquely defined up to a set of zero measure).

In this definition, we set $u(z) = 0$ if $z \in \varepsilon \mathbf{Z}^n \setminus \Omega$. In such a way $\mathcal{A}_\varepsilon(\Omega; K)$ will be regarded as a subset in $L^1(\Omega)$.

We denote by \mathcal{H}^{n-1} the $n-1$ -dimensional Hausdorff measure. Given $\nu = (\nu_1, \dots, \nu_n) \in S^{n-1}$ we set

$$Q_\nu := (-r_\nu, r_\nu)^n,$$

where $r_\nu > 0$ is such that

$$\mathcal{H}^{n-1}(Q_\nu \cap \Pi_\nu) = 1,$$

with $\Pi_\nu := \{x \in \mathbf{R}^n : (x, \nu) = 0\}$. We drop the dependence on ν whenever $\nu = e_i$ for $i \in \{1, 2, \dots, n\}$ and we set $Q := Q_{e_i} = (-\frac{1}{2}, \frac{1}{2})^n$.

For any $T > 0$ we set

$$\partial^\pm(TQ_\nu) := \{x \in \partial(TQ_\nu) : \pm(x, \nu) \geq 0\}$$

and then we introduce the discrete boundary of TQ_ν as

$$\partial_\varepsilon^\pm(TQ_\nu) := \{a \in \varepsilon \mathbf{Z}^n \cap TQ_\nu : (a + [-R\varepsilon, R\varepsilon]^n) \cap \partial^\pm(TQ_\nu) \neq \emptyset\}.$$

Next we recall some basic properties of BV functions with values in a finite set (see [8] for a general exposition of the subject). Let A be an open subset of \mathbf{R}^n and let J be a finite subset of \mathbf{R} . We denote by $BV(A; J)$ the set of measurable function $u : A \rightarrow J$ whose distributional derivative Du is a measure with bounded total variation. We denote by $S(u)$ the jump set of u and by $\nu_u(x)$ the measure theoretic inner normal to $S(u)$ at x , which is defined for \mathcal{H}^{n-1} a.e. $x \in S(u)$.

For the reader's convenience we recall the following compactness result (see [8]).

Theorem 2.2 *Let $u_k \in BV(A; J)$ such that*

$$\sup_n \mathcal{H}^{n-1}(S(u_k)) < +\infty.$$

Then there exists a subsequence (not relabelled) and $u \in BV(A; J)$ such that $u_k \rightarrow u$ in the L^1 convergence.

If Q is a cube we will denote by $BV^\#(Q; J)$ the set of Q -periodic functions belonging to $BV_{\text{loc}}(\mathbf{R}^n; J)$.

3 The Blume-Emery-Griffiths model

In this section we briefly introduce the Blume-Emery-Griffiths model for phase transitions in presence of surfactants.

3.1 A brief description of the model: from bulk to surface scaling

In its standard formulation the Blume-Emery-Griffiths model can be described as follows. Given a bounded open set $\Omega \subset \mathbf{R}^2$ with Lipschitz boundary, on the square lattice $\Omega_\varepsilon = \varepsilon\mathbf{Z}^2 \cap \Omega$ we consider the set $\mathcal{A}_\varepsilon(\Omega; \{\pm 1, 0\}) := \{u : \Omega_\varepsilon \rightarrow \{\pm 1, 0\}\}$ of those functions u whose values on the sites of the lattice correspond to a particle of water (or oil) (in this framework the scale is not precisely specified and the term particle means more in general a molecule or an aggregate of molecules) for $u = \pm 1$ or of surfactant for $u = 0$. We then introduce the family of energies $E_\varepsilon^{latt}(u) : \mathcal{A}_\varepsilon(\Omega; \{\pm 1, 0\}) \mapsto \mathbf{R}$

$$E_\varepsilon^{latt}(u) = \sum_{n.n.} \varepsilon^2 (-u(a)u(b) + k((u(a)u(b))^2)), \quad (3.4)$$

where *n.n.* means that the sum is performed over those $a, b \in \Omega_\varepsilon$ such that $|a - b| = \varepsilon$ and $k > 0$ is a parameter which measures the strength of the quadratic vs the bi-quadratic interactions. The asymptotic analysis of such a family of energies, as ε tends to 0, is particularly simple and can be obtained through a dual lattice approach as in [2]. Indeed, by identifying the functions $u \in \mathcal{A}_\varepsilon(\Omega; \{\pm 1, 0\})$ with their piecewise constant interpolations (see Remark 2.1), we first extend the energies E_ε^{latt} in (3.4) to a functional $E_\varepsilon : L^1(\Omega) \rightarrow [0, +\infty]$ as

$$E_\varepsilon(u) = \begin{cases} E_\varepsilon^{latt}(u) & \text{if } u \in \mathcal{A}_\varepsilon(\Omega; \{\pm 1, 0\}) \\ +\infty & \text{otherwise,} \end{cases}$$

and then compute the Γ -limit of (E_ε) with respect to the weak topology in $L^1(\Omega)$. As a result one obtains that the following Theorem holds true:

Theorem 3.1 *The family (E_ε) Γ -converges with respect to the $L^1(\Omega)$ -weak topology to the functional $E : L^1(\Omega) \rightarrow \mathbf{R} \cup \{+\infty\}$ defined as*

$$E(u) = \begin{cases} 2|\Omega|(k-1) \wedge 0 & \text{if } u \in L^1(\Omega; [-1, 1]) \\ +\infty & \text{otherwise.} \end{cases}$$

Let us comment the previous result in the interesting case when $k < 1$. In this regime the lattice energy has the two pure states $u = \pm 1$ as minimizers, and all the variations of the order parameter from these states count at order ε^2 in the discrete energy. This implies that, in the continuum limit, it is possible to obtain, with finite energy, any value of the order parameter u in $[-1, +1]$ by arbitrarily mixing the two ground states on a mesoscopic scale $\varepsilon \ll \delta \ll 1$. In particular this makes the energy of a phase separation to be negligible. More precisely the energy for a phase transition from a bulk -1 phase to a bulk $+1$ phase separated by an interface of finite length has an energy of order ε and suggests the correct scaling

to track the energetic behavior of a phase separation phenomenon. Observing that the absolute minimum value at scale ε is precisely given by

$$m_\varepsilon = \sum_{n.n.} \varepsilon^2(k-1),$$

in order to have more detailed description of the ground states we select those configurations described by functions u_ε that realize the minimum value with a sharper precision; *i.e.*, such that

$$E_\varepsilon(u_\varepsilon) = m_\varepsilon + O(\varepsilon).$$

In other words this amounts to study the family of discrete energies $E_\varepsilon^{(1)} : L^1(\Omega) \rightarrow [0, +\infty]$ defined as

$$E_\varepsilon^{(1)}(u) := \frac{E_\varepsilon(u) - m_\varepsilon}{\varepsilon},$$

that is

$$E_\varepsilon^{(1)}(u) = \begin{cases} \sum_{n.n.} \varepsilon(1 - u(a)u(b) - k(1 - (u(a)u(b))^2)) & \text{if } u \in \mathcal{A}_\varepsilon(\Omega; \{0, \pm 1\}), \\ +\infty & \text{otherwise.} \end{cases} \quad (3.5)$$

Within this scaling, the measure of the surfactant phase has to be negligible in the continuum limits. More precisely, it is easily seen that, since each interaction with a surfactant particle pays a positive energy $1 - k$, the following estimate

$$E_\varepsilon^{(1)}(u) \geq \#\{a \in \Omega_\varepsilon : u(a) = 0\} \varepsilon(1 - k) \geq \frac{C}{\varepsilon} |\{x \in \Omega : u(x) = 0\}|$$

implies that the measure of the surfactant phase scales as ε . As a result, the finite energy states u will only take the values ± 1 . It is now possible to further specify the values of the parameter k in such a way that the phase 0 can be actually considered a surfactant phase, meaning that it lowers the surface tension in the continuum limit. To obtain an estimate on the values of k we can proceed by computing the energy for a transition from -1 to $+1$ in the simplest case when the the interface is a straight line in one of the directions of the lattice (say e_1). Suppose for simplicity that $\Omega = Q$ and that the interface is the set $\{x \in Q : (x, e_2) = 0\}$. Our estimates is obtained by comparing the energy for such a macroscopic transition when the microscopic structure is either given by u_ε or by v_ε , where

$$u_\varepsilon(a) = \begin{cases} +1 & \text{if } (a, e_2) \geq 0 \\ -1 & \text{otherwise.} \end{cases}$$

and

$$v_\varepsilon(a) = \begin{cases} +1 & \text{if } (a, e_2) > 0 \\ 0 & \text{if } (a, e_2) = 0 \\ -1 & \text{otherwise.} \end{cases}$$

It holds that

$$\begin{aligned} E_\varepsilon^{(1)}(u_\varepsilon) &= 2 + o(1) \\ E_\varepsilon^{(1)}(v_\varepsilon) &= 3(1 - k) + o(1). \end{aligned}$$

Imposing now that the microstructure with the surfactant lowers the interface energy means to impose that $E_\varepsilon^{(1)}(v_\varepsilon) < E_\varepsilon^{(1)}(u_\varepsilon)$. This turns out to imply the condition $k > \frac{1}{3}$. Such an estimate is indeed proven to be sufficient to treat the case of a more general interface, as it is shown in Theorem 3.2.

Finally we remark that the previous heuristic derivation of the range of the parameter k leads us to call such a regime the *low surfactant concentration* regime. In this regime it can be also justified the absence, in the present model, of other terms originally present in the BEG energy, such as chemical potential of the surfactant phase, which in a general situation cannot be ignored (see [26]).

Theorem 3.2 *Let $\frac{1}{3} < k < 1$ and let $(E_\varepsilon^{(1)})_\varepsilon$ be the family of functionals defined as in (3.5). Then we have*

(i) *for any sequence $(u_\varepsilon) \subseteq L^1(\Omega)$ such that*

$$\sup_\varepsilon E_\varepsilon^{(1)}(u_\varepsilon) \leq C < +\infty$$

there exist $(u_{\varepsilon_k})_{k \in \mathbf{N}}$ and $u \in BV(\Omega; \{\pm 1\})$ such that

$$u_{\varepsilon_k} \rightarrow u \text{ for } k \rightarrow +\infty$$

with respect to the $L^1(\Omega)$ -topology;

(ii) *the family of functionals $(E_\varepsilon^{(1)})$ Γ -converges with respect to the $L^1(\Omega)$ -topology to the functional $E^{(1)} : L^1(\Omega) \rightarrow [0, +\infty]$ defined by*

$$E^{(1)}(u) = \begin{cases} \int_{S(u)} \psi(\nu_u) d\mathcal{H}^1 & \text{if } u \in BV(\Omega; \{\pm 1\}) \\ +\infty & \text{otherwise,} \end{cases} \quad (3.6)$$

where the function $\psi : S^1 \rightarrow [0, +\infty)$ is given by

$$\psi(\nu) = (1 - k)(3|\nu_1| \vee |\nu_2| + |\nu_1| \wedge |\nu_2|).$$

PROOF. Not to overburden the reader we prefer to derive the proof as a consequence of Theorem 3.3. The compactness result stated in (i) is a straightforward consequence of the analogous result stated in Theorem 3.3 (i). In order to prove the Γ -liminf inequality, let us first note that the function $\varphi(\cdot, \cdot)$ defined in (3.9) satisfies

$$\min\{\varphi(z, \nu) : z \in \mathbf{R}_+\} = \varphi(|\nu_1| \vee |\nu_2|, \nu) = \psi(\nu).$$

Hence the functionals $E^{(1)}(\cdot, \cdot)$ defined in (3.8) satisfies

$$E^{(1)}(u, \mu) \geq E^{(1)}(u), \quad \forall (u, \mu) \in BV(\Omega; \{\pm 1\}) \times \mathcal{M}^+(\Omega)$$

Let $u_\varepsilon \rightarrow u$ in $L^1(\Omega)$. By Theorem 3.3 (i), we may assume that $\mu(u_\varepsilon) \rightarrow \mu$ weakly in the sense of measures. Then

$$\liminf_\varepsilon E_\varepsilon^{(1)}(u_\varepsilon) \geq E^{(1)}(u, \mu) \geq E^{(1)}(u).$$

By a density argument it suffices to prove the Γ -lim sup inequality for a function u with a polyhedral jump set. Since the construction is local it is enough to consider $u = u_\nu$, where u_ν is defined in (5.74). For such a function the optimizing sequence is given by $v_{z,\nu}(\frac{\cdot}{\varepsilon})$, where $v_{z,\nu}$ is defined in (5.75), with $z = |\nu_1| \vee |\nu_2|$. \square

3.2 Low concentration of surfactants: discrete-to-continuum limit

As we have seen in the previous section, in the topology we have chosen, the limit order parameter u does not carry any information about the surfactant phase. Actually the role of the surfactant becomes clear when one looks at the minimizing microstructure leading to the computation of the limiting surface density ψ . A natural further step in the analysis of the BEG model is the dependence of the surface tension of the continuum limit on the concentration of the surfactant. To make it explicit, we need to go beyond the standard formulation of the BEG model and we let the energy functional of the system depend explicitly on the distribution of the surfactant particles. To this end, for all $u \in \mathcal{A}_\varepsilon(\Omega; \{0, \pm 1\})$ we set

$$I_0(u) = \{a \in \Omega_\varepsilon : u(a) = 0\},$$

and we introduce the following *surfactant measure*

$$\mu(u) = \sum_{a \in I_0(u)} \varepsilon \delta_a.$$

Then, with a slight abuse of notation, we can extend $E_\varepsilon^{(1)}$ to a functional $E_\varepsilon^{(1)} : L^1(\Omega) \times \mathcal{M}_+(\Omega) \rightarrow [0, +\infty]$ as

$$E_\varepsilon^{(1)}(u, \mu) = \begin{cases} E_\varepsilon^{(1)}(u) & \text{if } u \in \mathcal{A}_\varepsilon(\Omega; \{0, \pm 1\}), \mu = \mu(u) \\ +\infty & \text{otherwise.} \end{cases} \quad (3.7)$$

We endow the space $L^1(\Omega) \times \mathcal{M}_+(\Omega)$ with the topology $\tau_1 \times \tau_2$ where τ_1 denotes the strong topology in $L^1(\Omega)$ and τ_2 denotes the weak*-topology in the space of non-negative bounded Radon measures $\mathcal{M}_+(\Omega)$.

The following Theorem holds true.

Theorem 3.3 Let $E_\varepsilon^{(1)}$ be defined by (3.7). There holds:

(i) Let $\varepsilon_k \rightarrow 0$ and let $(u_k, \mu_k) \in L^1(\Omega) \times \mathcal{M}_+(\Omega)$ be such that

$$\sup_k E_{\varepsilon_k}^{(1)}(u_k, \mu_k) < +\infty.$$

Then there exist a subsequence (not relabeled) such that $(u_k, \mu_k) \rightarrow (u, \mu)$ with respect to the $\tau_1 \times \tau_2$ topology, for some $(u, \mu) \in L^1(\Omega) \times \mathcal{M}_+(\Omega)$.

(ii) the family $(E_\varepsilon^{(1)})$ Γ -converges with respect to the $\tau_1 \times \tau_2$ topology to the functional $E^{(1)} : L^1(\Omega) \times \mathcal{M}_+(\Omega) \rightarrow [0, +\infty]$ defined by

$$E^{(1)}(u, \mu) = \begin{cases} \int_{S(u)} \varphi\left(\frac{d\mu}{d\mathcal{H}^1 \llcorner S(u)}, \nu_u\right) d\mathcal{H}^1 + (2k-2)|\mu^s|(\Omega) & \text{if } u \in BV(\Omega; \{\pm 1\}) \\ +\infty & \text{otherwise,} \end{cases} \quad (3.8)$$

where, for $\mu \in \mathcal{M}_+(\Omega)$, we have set $\mu^s := \mu - \frac{d\mu}{d\mathcal{H}^1 \llcorner S(u)} \mathcal{H}^1 \llcorner S(u)$ and the function $\varphi : \mathbf{R}_+ \times S^1 \rightarrow [0, +\infty)$ is given by

$$\varphi(z, \nu) = \max\{\varphi_1(z, \nu), \varphi_2(z, \nu), \varphi_3(z, \nu)\}, \quad (3.9)$$

where

$$\begin{aligned} \varphi_1(z, \nu) &= -4kz + 2(|\nu_1| + |\nu_2|), \\ \varphi_2(z, \nu) &= (1-3k)z + 2(|\nu_1| \vee |\nu_2|) + (1-k)(|\nu_1| \wedge |\nu_2|), \\ \varphi_3(z, \nu) &= 2(1-k)z + (1-k)(|\nu_1| + |\nu_2|). \end{aligned}$$

We postpone the proof of the previous Theorem in Section 5, since it makes use of the integral representation result stated in Theorem 4.4.

Remark 3.4 As it is clear from the graphic of φ (see Figure 2), an anisotropic threshold phenomenon occurs at the phase interface. For a given $\nu \in S^1$ the surface tension $\varphi(z, \nu)$ of the system may decrease only up to a certain value of the density z of the surfactant, namely $z = |\nu_1| \vee |\nu_2|$. Increasing further the density of the surfactant in the system the surface tension may be constant if the surfactants are not adsorbed onto the interface, or otherwise it may increase. In the first case the singular part of the surfactant measure increases.

As an application of the previous result, one may study the asymptotic behavior, as $\varepsilon \rightarrow 0$, of the following constrained optimization problem:

$$m_\varepsilon := \left\{ E_\varepsilon^{(1)}(u), \varepsilon \# I_0(u) = \alpha_\varepsilon, \varepsilon^2 \# I_1(u) = \beta_\varepsilon \right\} \quad (3.10)$$

where

$$I_1(u) = \{a \in \Omega_\varepsilon : u(a) = 1\},$$




Figure 2: The graphic of the surface tension density $\varphi(z, \nu)$ as a function of the density z of surfactant at the phase interface.

$\lim_{\varepsilon} \alpha_{\varepsilon} = \alpha > 0$ and $\lim_{\varepsilon} \beta_{\varepsilon} = \beta > 0$. Since we are not interested in boundary layer effects, we consider the case when Ω is a torus, that we may identify with the semi-open cube $\mathcal{Q} := [0, 1)^2$, $\varepsilon = \frac{1}{k}$, $k \in \mathbf{N}$, and the admissible functions u in (3.10) are \mathcal{Q} -periodic. The solution to this problem is a particular case of the result stated in Corollary 4.12 (see Remark 4.13).

4 More general models

In this section we consider a class of energies that generalizes those involved in the BEG model and in which long range interactions and different types of surfactant are taken into account.

Let $\Omega \subset \mathbf{R}^n$ be a bounded open set with Lipschitz boundary. We consider the family of functionals $F_{\varepsilon} : \mathcal{A}_{\varepsilon}(\Omega; K) \rightarrow [0, +\infty)$ defined by

$$F_{\varepsilon}(u) = \sum_{a, b \in \Omega_{\varepsilon}, |a-b| \leq R\varepsilon} \varepsilon^{n-1} f\left(\frac{b-a}{\varepsilon}, u(a), u(b)\right) \quad (4.11)$$

where $R > 0$, $K = \{m_1, m_2, s_1, s_2, \dots, s_M\} \subset \mathbf{R}$, $M \in \mathbf{N}$, and $f : \mathbf{Z}^n \times K^2 \rightarrow [0, +\infty)$ satisfies the following hypotheses:

$$f^{-1}(0) = \mathbf{Z}^n \times \{(m_1, m_1), (m_2, m_2)\}, \quad (4.12)$$

$$f(R^i \xi, u, v) = f(\xi, u, v) \quad \forall i \in \{1, 2, \dots, n\}, \quad (4.13)$$

where $R^i(\xi_1, \xi_2, \dots, \xi_i, \dots, \xi_n) = (\xi_1, \xi_2, \dots, -\xi_i, \dots, \xi_n)$ is the reflection with respect to the i -th coordinate axis. We also define a localized energy for every $A \subset \Omega$ as

$$F_\varepsilon(u, A) = \sum_{a, b \in A \cap \varepsilon \mathbf{Z}^n, |a-b| \leq R\varepsilon} \varepsilon^{n-1} f\left(\frac{b-a}{\varepsilon}, u(a), u(b)\right) \quad (4.14)$$

Remark 4.1 We remark that (4.12) implies that the pure phases $u \equiv m_i$, $i = 1, 2$, are the ground states of the energy F_ε . Hypothesis (4.13) is a sort of discrete isotropy condition of the energy density and it is in particular satisfied if $f(\xi, u, v) = f(|\xi|, u, v)$.

Remark 4.2 We observe that the functional $E_\varepsilon^{(1)}$ defined in (3.5) is of the type (4.11) with

$$f(\xi, u, v) := \begin{cases} -uv - k(1-uv)^2 & \text{if } \xi = \pm e_i, i \in \{1, 2\} \\ 0 & \text{otherwise,} \end{cases} \quad (4.15)$$

which satisfies assumptions (4.12) and (4.13) with $K = \{\pm 1, 0\}$, being $m_1 = -1$ and $m_2 = 1$.

We now introduce the sets of points in Ω_ε occupied by the different types of surfactant and suitable measures associated to them. For $l \in \{1, 2, \dots, M\}$ we set, for all $A \subset \Omega$,

$$I_l(u, A) = \{a \in A \cap \Omega_\varepsilon : u(a) = s_l\},$$

$$I(u, A) = \bigcup_{l=1}^M I_l(u, A)$$

and, for the sake of simplicity, we set $I_l(u, \Omega) = I_l(u)$ and $I(u, \Omega) = I(u)$. Moreover we define

$$\mu_l(u) := \sum_{a \in I_l(u)} \varepsilon^{n-1} \delta_a$$

$$\mu(u) := \{\mu_1(u), \mu_2(u), \dots, \mu_M(u)\}. \quad (4.16)$$

With the identification given in Remark 2.1, and a slight abuse of notation, we can extend F_ε to a functional $F_\varepsilon : L^1(\Omega) \times (\mathcal{M}_+(\Omega))^M \rightarrow [0, +\infty]$ as

$$F_\varepsilon(u, \mu) := \begin{cases} F_\varepsilon(u) & \text{if } u \in \mathcal{A}_\varepsilon(\Omega; K), \mu = \mu(u) \\ +\infty & \text{otherwise.} \end{cases}$$

We endow the space $L^1(\Omega) \times (\mathcal{M}_+(\Omega))^M$ with the topology $\tau_1 \times \tau_2$ where τ_1 denotes the strong topology in $L^1(\Omega)$ and τ_2 denotes the weak*-topology in $(\mathcal{M}_+(\Omega))^M$. The choice of this topology is suggested by the following compactness result.

Proposition 4.3 *Let $\varepsilon_k \rightarrow 0$ and let (u_k, μ_k) be such that*

$$\sup_k F_{\varepsilon_k}(u_k, \mu_k) < +\infty.$$

Then there exist a subsequence (not relabeled) such that $(u_k, \mu_k) \rightarrow (u, \mu)$ w.r.t. the $\tau_1 \times \tau_2$ -topology, for some $(u, \mu) \in BV(\Omega; \{m_1, m_2\}) \times (\mathcal{M}_+(\Omega))^M$.

PROOF. First note that

$$\mathcal{H}^{n-1}(S(u_k)) + \mu_k(\Omega) \leq CF_{\varepsilon_k}(u_k, \mu_k). \quad (4.17)$$

By Theorem 2.2 and observing that

$$|\{x \in \Omega : u_k(x) \notin \{m_1, m_2\}\}| \leq C\varepsilon_k \mu_k(\Omega) \rightarrow 0,$$

one easily get the conclusion. \square

4.1 The main result

In this section we state and prove an integral representation result for the Γ -limit of the family F_ε . To this end we introduce, for any $\varepsilon > 0$ and $\nu \in S^{n-1}$, the class of discrete functions

$$\mathcal{B}_\varepsilon(TQ_\nu; K) := \{u \in \mathcal{A}_\varepsilon(TQ_\nu; K) : u(a) = m_1 \forall a \in \partial_\varepsilon^+(TQ_\nu), u(a) = m_2 \forall a \in \partial_\varepsilon^-(TQ_\nu)\}.$$

Theorem 4.4 *The family (F_ε) Γ -converges with respect to the $\tau_1 \times \tau_2$ -topology to the functional $F : L^1(\Omega) \times (\mathcal{M}_+(\Omega))^M \rightarrow [0, +\infty]$ defined by*

$$F(u, \mu) = \begin{cases} \int_{S(u)} f_{hom} \left(\frac{d\mu}{d\mathcal{H}^{n-1} \llcorner S(u)}, \nu(u) \right) d\mathcal{H}^{n-1} + \int_\Omega g_{hom}(\mu^s) & \text{if } u \in BV(\Omega; \{m_1, m_2\}) \\ +\infty & \text{otherwise,} \end{cases} \quad (4.18)$$

where, for $\mu \in (\mathcal{M}_+(\Omega))^M$, we have set $\mu^s := \mu - \frac{d\mu}{d\mathcal{H}^{n-1} \llcorner S(u)} \mathcal{H}^{n-1} \llcorner S(u)$. Here, $f_{hom} : (\mathbf{R}_+)^M \times S^{n-1} \rightarrow [0, +\infty)$ is defined as

$$f_{hom}(z, \nu) := \lim_{\delta \rightarrow 0^+} \lim_{T \rightarrow +\infty} \frac{1}{T^{n-1}} \inf \left\{ F_1(u, TQ_\nu) : u \in \mathcal{B}_1(TQ_\nu; K), \right. \\ \left. \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u, TQ_\nu)}{T^{n-1}} - z_l \right| < \delta \right\}, \quad (4.19)$$

while $g_{hom} : (\mathbf{R}_+)^M \rightarrow [0, +\infty)$ is 1-homogeneous and, for any $\zeta \in (\mathbf{R}_+)^M$ such that $\|\zeta\|_1 = 1$, is defined as

$$g_{hom}(\zeta) := \lim_{\delta \rightarrow 0^+} \liminf_{T \rightarrow +\infty} \inf \left\{ \frac{F_1(u, TQ)}{\#I(u, TQ)} : u \in \mathcal{A}_1(TQ; K), \right. \\ \left. \frac{F_1(u, TQ \setminus (T-R)Q)}{\#I(u, TQ)} < \delta, \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u, TQ)}{\#I(u, TQ)} - \zeta_l \right| < \delta \right\} \quad (4.20)$$

Remark 4.5 We observe that, while the formula for f_{hom} can be proved by using standard arguments in homogenization theory, the same does not hold for g_{hom} . In particular, as it will be clear in the proof of Theorem 4.4, optimizing sequences for $f_{hom}(z, \nu)$ can be constructed, as it is usual in this framework, by "periodically gluing" a solution of the minimum problem on TQ_ν given in (4.19). In such a construction, the energy due to the interactions which cross the boundary of the periodicity cell is asymptotically negligible thanks to the Dirichlet type condition we are allowed to impose by using a De Giorgi's cut-off construction (see Lemma 4.9). The same arguments do not apply to g_{hom} . In fact, for this term, we cannot be sure that, imposing the same type of boundary conditions, we do not modify too much the energy of minimal configurations in (4.20) since the distribution of the phases m_1 and m_2 for such configurations is not known. This fact rules out the standard "periodic gluing" construction. Instead, we first make use of an abstract argument from measure theory which allows us to prove that the minimal configurations do not concentrate energy at the boundary of the periodic cell and then, by exploiting hypothesis (4.13), we construct optimizing sequences for $g_{hom}(\zeta)$ by a reflection argument (see the proof of Proposition 4.10).

Before proving Theorem 4.4 we show some properties enjoyed by f_{hom} and g_{hom} . In the next Proposition we prove that the homogenization formula defining f_{hom} is well defined.

Proposition 4.6 *For any $z \in \mathbf{R}^M$, $\nu \in S^{n-1}$ and $\delta > 0$ there exists the limit*

$$\lim_{T \rightarrow +\infty} \frac{1}{T^{n-1}} \inf \left\{ F_1(u, TQ_\nu) : u \in \mathcal{B}_1(TQ_\nu; K), \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u, TQ_\nu)}{T^{n-1}} - z_l \right| < \delta \right\}.$$

PROOF. For simplicity of notation we develop the proof in the case $\nu = e_n$ but the argument obviously applies to the general case. We recall that we have set $Q := Q_{e_n} = (-\frac{1}{2}, \frac{1}{2})^n$. Let us define

$$\mathcal{I}_T(z, \delta) := \inf \left\{ F_1(u, TQ) : u \in \mathcal{B}_1(TQ; K), \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u, TQ)}{T^{n-1}} - z_l \right| < \delta \right\}, \quad (4.21)$$

and, given $\eta > 0$, let $u_T \in \mathcal{B}_1(TQ; K)$ be such that $\max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u_T, TQ)}{T^{n-1}} - z_l \right| < \delta$ and

$$F_1(u_T, TQ) \leq \mathcal{I}_T(z, \delta) + \eta.$$

Let u_T be extended on the stripe $((-\frac{T}{2}, \frac{T}{2})^{n-1} \times \mathbf{R}) \cap \mathbf{Z}^n$ by setting

$$u_T(a) = m_1 \text{ if } a_n \geq \frac{T}{2}, \quad u_T(a) = m_2 \text{ if } a_n \leq \frac{T}{2}.$$

Let us set $\hat{T} := 2 \lceil \frac{T}{2} \rceil$ and, for $S > T$, let $v_S \in \mathcal{B}_1(SQ; K)$ be defined as

$$v_S(a) = u_T(a) \quad \text{if } a \in \left\{ -\frac{\hat{T}}{2}, \dots, \frac{\hat{T}}{2} \right\}^{n-1} \times \mathbf{Z},$$

and then extended by periodicity in the e_1, \dots, e_{n-1} directions as

$$v_S(a + j\hat{T}e_i) = v_S(a) \quad \text{for all } j \in \mathbf{Z}, i \in \{1, \dots, n-1\}.$$

Note that, for S large enough, we have

$$\left| \frac{\#I_l(v_S, SQ)}{S^{n-1}} - z_l \right| < \delta.$$

Hence, splitting the energy into two terms, the first one accounting for the interactions inside each periodic cell of the type $\left[-\frac{\hat{T}}{2}, \frac{\hat{T}}{2}\right]^n + j\hat{T}e_i$, and the second one accounting for the interactions which cross the boundary of the same cells, we get

$$\begin{aligned} \frac{1}{S^{n-1}} \mathcal{I}_S(z, \delta) &\leq \frac{1}{S^{n-1}} F_1(v_S, SQ) \leq \frac{1}{S^{n-1}} \left[\frac{S}{\hat{T}} \right]^{n-1} (F_1(u_T, TQ) + C\hat{T}^{n-2}) \\ &\leq \frac{T^{n-1}}{S^{n-1}} \left[\frac{S}{\hat{T}} \right]^{n-1} \left(\frac{1}{T^{n-1}} \mathcal{I}_T(z, \delta) + \frac{C\hat{T}^{n-2} + \eta}{T^{n-1}} \right). \end{aligned}$$

By letting first S and then T go to $+\infty$, by the arbitrariness of η we finally get

$$\limsup_{S \rightarrow +\infty} \frac{1}{S^{n-1}} \mathcal{I}_S(z, \delta) \leq \liminf_{T \rightarrow +\infty} \frac{1}{T^{n-1}} \mathcal{I}_T(z, \delta).$$

□

In the next Proposition we prove growth and convexity properties of the functions f_{hom} and g_{hom} . We remark that, in the proof the Γ -convergence result, we only use the continuity of f_{hom} and g_{hom} and that their convexity would be a consequence of the lower semicontinuity of the Γ -limit. However the proof of the continuity would rely on the same argument we exploit here without providing any significant simplification.

Proposition 4.7 *Let f_{hom} and g_{hom} be defined as in (4.19) and (4.20). There holds*

(i) *the 1-homogeneous extension of f_{hom} in $(\mathbf{R}_+)^M \times \mathbf{R}^n$ is convex. Moreover there exists $C > 0$ such that*

$$f_{hom}(z, \nu) \leq C(|z| + 1), \quad \text{for all } (z, \nu) \in (\mathbf{R}_+)^M \times S^{n-1}. \quad (4.22)$$

(ii) *g_{hom} is convex.*

PROOF. **Proof of (i)**

By the 1-homogeneity of f_{hom} the proof of its convexity reduces to prove that, given $\nu, \nu^{(1)}, \nu^{(2)} \in S^{n-1}$ and $z, z^{(1)}, z^{(2)} \in (\mathbf{R}_+)^M$ such that $(z, \nu) = l_1(z^{(1)}, \nu^{(1)}) + l_2(z^{(2)}, \nu^{(2)})$, for $l_1, l_2 \in \mathbf{R}$, there holds

$$f_{hom}(z, \nu) \leq l_1 f_{hom}(z^{(1)}, \nu^{(1)}) + l_2 f_{hom}(z^{(2)}, \nu^{(2)}). \quad (4.23)$$

Without loss of generality, for simplicity of notation we prove (4.23) under the assumption that $l_1, l_2 > 0$, $\nu = e_n$, $\nu^{(1)}, \nu^{(2)} \in \{x \in \mathbf{R}^n : x_1 = \dots = x_{n-2} = 0\} =: \Pi$, $(\nu^{(1)}, \nu^{(2)}) \geq 0$ and the ordered base $\{\nu^{(1)}, \nu^{(2)}\}$ has the same orientation as $\{e_n, e_{n-1}\}$. Given $\eta > 0$, let $\delta > 0$, $T > \frac{1}{\eta}$, $u_1 \in \mathcal{B}_1(TQ_{\nu^{(1)}}; K)$, $u_2 \in \mathcal{B}_1(TQ_{\nu^{(2)}}; K)$ such that, for $i \in \{1, 2\}$ it holds

$$\frac{1}{T^{n-1}} F_1(u_i, TQ_{\nu^{(i)}}) \leq f_{hom}(z^{(i)}, \nu^{(i)}) + \eta$$

$$\max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u_i, TQ_{\nu^{(i)}})}{T^{n-1}} - z_l^{(i)} \right| < \delta.$$

Let us set, for $i \in \{1, 2\}$, $r_i := r_{\nu^{(i)}}$, so that $Q_{\nu^{(i)}} = [-\frac{r_i}{2}, \frac{r_i}{2}]^2$, and let us consider u_i identified with its piecewise-constant interpolation and extended to $\mathbf{R}^{n-2} \times T[-\frac{r_i}{2}, \frac{r_i}{2}]^2$ by periodicity in the $\{e_1, \dots, e_{n-2}\}$ directions. Moreover we set $L_i := \mathcal{H}^1(\{x \in \Pi : x = t(\nu^{(i)})^\perp, t \in \mathbf{R}\} \cap Q_{\nu^{(i)}})$ and $(\nu^{(i)})^\perp = (0, \dots, 0, \nu_n^{(i)}, -\nu_{n-1}^{(i)})$. We can then further extend u_i by periodicity in the direction $(\nu^{(i)})^\perp$ without renaming it, that is

$$u_i(x + L_i(\nu^{(i)})^\perp) = u_i(x), \quad x \in \mathbf{R}^{n-2} \times \bigcup_{j \in \mathbf{Z}} \left(T[-\frac{r_i}{2}, \frac{r_i}{2}]^2 + jL_i(\nu^{(i)})^\perp \right).$$

We now use a zig-zag construction to define a good test function $u_S \in \mathcal{B}_1(SQ; K)$, for $S \gg T$, in the minimum problem occurring in the definition of $f_{hom}(z, \nu)$ (see Figure 3). Let us set $V := \{x \in \mathbf{R}^n : 0 \leq x_{n-1} \leq 1\}$, $\Pi_\nu^\pm := \{x \in \mathbf{R}^n : \pm(x, \nu) \geq 0\}$, $\nu \in S^{n-1}$, and let $w : V \rightarrow \{m_1, m_2\}$ be defined as

$$w(x) = \begin{cases} m_1 & \text{if } x \in \Pi_{\nu^{(1)}}^+ \cup (\Pi_{\nu^{(2)}}^+ + l_1(\nu^{(1)})^\perp) \\ m_2 & \text{otherwise.} \end{cases}$$

Let $c_1, c_2 > 0$ be such that $c_1 \frac{TL_1}{l_1} = c_2 \frac{TL_2}{l_2} =: r$. Let $u : \mathbf{R}^n \rightarrow \mathbf{R}$ be the re_{n-1} -periodic function defined in rV as

$$u(x) = \begin{cases} u_1(x) & \text{if } 0 \leq x_{n-1} \leq [c_1]TL_1(\nu^{(1)}, e_n) \\ u_2(x - re_{n-1}) & \text{if } r - [c_2]TL_2(\nu^{(2)}, e_n) \leq x_{n-1} \leq r \\ w(\frac{x}{r}) & \text{otherwise in } rV. \end{cases}$$

and periodically extended in the direction e_{n-1} . Let $S \gg T$, we define $u_S \in \mathcal{B}_1(SQ_\nu; K)$ such that

$$u_S(a) = \begin{cases} u(a) & \text{if } a \in (S - R)Q_\nu \cap \mathbf{Z}^n \\ u_\nu(a) & \text{otherwise in } SQ_\nu \cap \mathbf{Z}^n. \end{cases}$$




Figure 3: The zig-zag construction leading to u_S in the proof of Proposition 4.7. On the right a zoom of one of the triangles on the left.

We can now estimate the energy of u_S :

$$\begin{aligned} \frac{1}{S^{n-1}} F_1(u_S, SQ_\nu) &\leq \frac{1}{S^{n-1}} \sum_{i=1}^2 \left[\frac{S}{r} \right] [c_i] \left[\frac{S}{Tr_i} \right]^{n-2} (F_1(u_i, TQ_{\nu^{(i)}}) + CT^{n-2}) \\ &\quad + \frac{1}{S^{n-1}} CS^{n-2}, \end{aligned} \quad (4.24)$$

where the term of the type CT^{n-2} is the energetic contribution due to the interactions near each set of the type $\partial(TQ_{\nu^{(i)}}) \cap \Pi_{\nu^{(i)}}$, while the term of the type CS^{n-2} is due to the interactions near $\partial(SQ_\nu) \cap \Pi_\nu$. By construction we have that, for $l \in \{1, \dots, M\}$,

$$\frac{\#I_l(u_S, SQ_\nu)}{S^{n-1}} = \frac{1}{S^{n-1}} \sum_{i=1}^2 \left[\frac{S}{r} \right] [c_i] \left[\frac{S}{Tr_i} \right]^{n-2} \#I_l(u_i, TQ_{\nu^{(i)}}) + o(1). \quad (4.25)$$

Taking into account the definition of c_i and the fact that by construction $r_i^{n-2}L_i = 1$ we have that, for T and S large enough,

$$\frac{1}{S^{n-1}} \left[\frac{S}{r} \right] [c_i] \left[\frac{S}{Tr_i} \right]^{n-2} \leq \frac{l_1 + C\eta}{T^{n-1}} \quad (4.26)$$

Then, by (4.25), for T and S large enough u_S is a good test function in the minimum problem defining $f_{hom}(z, \nu)$ and, by (4.24) we get

$$f_{hom}(z, \nu) \leq l_1 f_{hom}(z^{(1)}, \nu^{(1)}) + l_2 f_{hom}(z^{(2)}, \nu^{(2)}) + C\eta$$

which eventually gives (4.23) by the arbitrariness of η .

To obtain the estimate (4.22) we observe that we can rewrite the energy $F_1(u, TQ_\nu)$ as

$$F_1(u, TQ_\nu) = \sum_{(a,b) \in D(u)} f(b-a, u(a), u(b)) + F_1^{sur}(u, TQ_\nu) \quad (4.27)$$

where

$$D(u) := \{(a, b) \in (\mathbf{Z}^n \cap TQ_\nu)^2 : 0 < |b-a| \leq R, u(a) \neq u(b), \{u(a), u(b)\} = \{m_1 m_2\}\}.$$

Here F_1^{sur} is the energy accounting only for the contribution due to the interactions of surfactant type particles. Note that, since f is bounded, the first term in the right hand side of (4.27) is proportional to $\#D(u)$ and, since each particle has only an equi-bounded number of interactions, $F_1^{sur}(u, TQ_\nu)$ is proportional to $\#I(u, TQ_\nu)$. Then the estimate (4.22) is achieved by choosing, in the problem defining $f_{hom}(z, \nu)$ any test function u such that $\#D(u) \simeq CT^{n-1}$.

Proof of (ii). In order to prove (ii), by the 1-homogeneity of g_{hom} , it is enough to show that, given $\zeta^{(1)}, \zeta^{(2)} \in (\mathbf{R}_+)^M$ with $\|\zeta^{(1)}\|_1 = \|\zeta^{(2)}\|_1 = 1$ and $t \in (0, 1)$ it holds that

$$g_{hom}(t\zeta^{(1)} + (1-t)\zeta^{(2)}) \leq tg_{hom}(\zeta^{(1)}) + (1-t)g_{hom}(\zeta^{(2)}). \quad (4.28)$$

For any $\delta > 0$ and $\zeta \in (\mathbf{R}_+)^M$ with $\|\zeta\|_1 = 1$, we set

$$g(\delta, \zeta) := \liminf_{T \rightarrow +\infty} \inf \left\{ \frac{F_1(u, TQ)}{\#I(u, TQ)} : u \in \mathcal{A}_1(TQ; K), \right. \\ \left. \frac{F_1(u, TQ \setminus (T-R)Q)}{\#I(u, TQ)} < \delta, \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u, TQ)}{\#I(u, TQ)} - \zeta_l \right| < \delta \right\} \quad (4.29)$$

Given $\delta > 0$, for $i \in \{1, 2\}$ let $u_i \in \mathcal{A}(TQ; K)$ such that

$$\frac{F_1(u_i, TQ)}{\#I(u_i, TQ)} < g(\delta, \zeta^{(i)}) + \delta, \\ \frac{F_1(u_i, TQ \setminus (T-R)Q)}{\#I(u_i, TQ)} < \delta \\ \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u_i, TQ)}{\#I(u_i, TQ)} - \zeta_l^{(i)} \right| < \delta. \quad (4.30)$$

Set $\tilde{\zeta} = t\zeta^{(1)} + (1-t)\zeta^{(2)}$, we now construct a suitable test function in the minimum problem defining $g(\delta, \tilde{\zeta})$. Let $S = kT$ with $1 \ll k \in \mathbf{N}$. Let $h \in \mathbf{N}$, $h < k$ be such that, for k large enough, set

$$\lambda(h) := \frac{h\#I(u_1, TQ)}{h\#I(u_1, TQ) + (k-h)\#I(u_2, TQ)},$$

there holds

$$|t - \lambda(h)| < \delta. \quad (4.31)$$

Without loss of generality we choose T to be an even number. We extend u_i by reflection with respect to the coordinate axes. More precisely we set $u_{i,0} = u_i$ and, for all $j \in \{1, \dots, n\}$, we define $u_{i,j}$ recursively as follows: $u_{i,j}$ is the extension of $u_{i,j-1}$ on $\mathbf{R}^j \times [-\frac{T}{2}, \frac{T}{2}]^{n-j}$ satisfying the following property

$$u_{i,j}(a) = u_{i,j}(a + (T - 2a_j)e_j), \quad \forall a \in \mathbf{R}^{j-1} \times \left[-\frac{T}{2}, \frac{T}{2}\right]^{n-(j-1)} + mTe_j, \quad m \in \mathbf{Z}.$$

We have then obtained that the function $u_{i,n}$ extends u_i on all \mathbf{Z}^n . Let us observe that, by the symmetry hypotheses in (4.12), we have that

$$F_1(u_{i,n}, TQ + m) = F_1(u_i, TQ) \quad \forall m \in \mathbf{Z}^n. \quad (4.32)$$

Let $u : \mathbf{Z}^n \cap SQ \rightarrow K$ be defined as

$$u(a) = \begin{cases} u_{1,n}(a) & \text{if } -\frac{S}{2} \leq a_n \leq -\frac{S}{2} + h \\ u_{2,n}(a) & \text{otherwise.} \end{cases}$$

By (4.30) and (4.31) we have that

$$\begin{aligned} \left| \frac{\#I_l(u, SQ)}{\#I(u, SQ)} - \tilde{\zeta}_l \right| &= \left| \frac{k^{n-1}h\#I_l(u_1, TQ) + k^{n-1}(k-h)\#I_l(u_2, TQ)}{k^{n-1}h\#I(u_1, TQ) + k^{n-1}(k-h)\#I(u_2, TQ)} - \tilde{\zeta}_l \right| \\ &\leq \lambda(h) \left| \frac{\#I_l(u_1, TQ)}{\#I(u_1, TQ)} - \zeta_l^{(1)} \right| + (1 - \lambda(h)) \left| \frac{\#I_l(u_2, TQ)}{\#I(u_2, TQ)} - \zeta_l^{(2)} \right| \\ &\quad + (\|\zeta^{(1)}\|_1 \vee \|\zeta^{(2)}\|_1) \delta \\ &\leq (1 + \|\zeta^{(1)}\|_1 \vee \|\zeta^{(2)}\|_1) \delta. \end{aligned}$$

We now estimate the energy $\frac{F_1(u, SQ)}{\#I(u, SQ)}$:

$$\frac{F_1(u, SQ)}{\#I(u, SQ)} \leq \frac{k^{n-1}hF_1(u_1, TQ) + k^{n-1}(k-h)F_1(u_2, TQ)}{\#I(u, SQ)} + R_1 + R_2. \quad (4.33)$$

Here the first term in the right hand side is obtained by taking into account (4.32). Moreover R_1 is the energy due to the interactions which cross the set $SQ \cap \{x_n = -\frac{S}{2} + h\}$, where in the construction of u we pass from u_1 to u_2 , while R_2 accounts for all the other interactions which cross the boundary of the cubes of type $TQ + m$ with $m \in \mathbf{Z}$. An easy computation shows that

$$R_1 \leq C \frac{k^{n-1}T}{\#I(u, SQ)} \leq C \frac{T}{k}.$$

In addition, thanks to (4.30) we get

$$\begin{aligned}
R_2 &\leq \frac{k^{n-1}hF_1(u_1, TQ \setminus (T-R)Q) + k^{n-1}(k-h)F_1(u_2, TQ \setminus (T-R)Q)}{\#I(u, SQ)} \\
&= \frac{k^{n-1}hF_1(u_1, TQ \setminus (T-R)Q) + k^{n-1}(k-h)F_1(u_2, TQ \setminus (T-R)Q)}{k^{n-1}h\#I(u_1, TQ) + k^{n-1}(k-h)\#I(u_2, TQ)} \\
&\leq \lambda(h) \frac{F_1(u_1, TQ \setminus (T-R)Q)}{\#I(u_1, TQ)} + (1-\lambda(h)) \frac{F_1(u_2, TQ \setminus (T-R)Q)}{\#I(u_2, TQ)} \\
&\leq \delta.
\end{aligned}$$

Hence, using the same argument to estimate the first term in the right hand side of (4.33), by (4.30), (4.31) and the estimates of R_1 and R_2 , we have

$$\begin{aligned}
\frac{F_1(u, SQ)}{\#I(u, SQ)} &= \lambda(h) \frac{F_1(u_1, TQ)}{\#I(u_1, TQ)} + (1-\lambda(h)) \frac{F_1(u_2, TQ)}{\#I(u_2, TQ)} + R_1 + R_2 \\
&\leq tg(\delta, \zeta^{(1)}) + (1-t)g(\delta, \zeta^{(2)}) + C\left(\delta + \frac{T}{k}\right).
\end{aligned}$$

Letting k go to $+\infty$ we get

$$g(C\delta, \zeta) \leq tg(\delta, \zeta^{(1)}) + (1-t)g(\delta, \zeta^{(2)}) + C\delta$$

which eventually gives the conclusion letting δ go to 0. \square

The proof of Theorem 4.4 will follow by Propositions 4.8 and 4.10 in which we prove the Γ -lim inf and the Γ -lim sup inequality, respectively.

Proposition 4.8 (Γ -lim inf inequality) *We have*

$$\Gamma - \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u, \mu) \geq F(u, \mu) \quad (4.34)$$

PROOF. Let $\varepsilon_k \rightarrow 0$. Up to subsequences, it suffices to consider $(u_k, \mu_k) \rightarrow (u, \mu)$ w.r.t. the $\tau_1 \times \tau_2$ -topology such that

$$\liminf_k F_{\varepsilon_k}(u_k, \mu_k) = \lim_k F_{\varepsilon_k}(u_k, \mu_k) < +\infty. \quad (4.35)$$

By Proposition 4.3 we have that $u \in BV(\Omega; \{m_1, m_2\})$. We now consider the family of measures $(\lambda_k)_k \subset \mathcal{M}_+(\Omega)$ defined as

$$\lambda_k := \sum_{a \in \Omega_{\varepsilon_k}} \sum_{b \in \Omega_{\varepsilon_k}: |a-b| \leq R\varepsilon_k} \varepsilon_k^{n-1} f\left(\frac{b-a}{\varepsilon_k}, u(a), u(b)\right) \delta_a. \quad (4.36)$$

Note that $\lambda_k(\Omega) = F_{\varepsilon_k}(u_k, \mu_k)$. Then, by (4.35), we may suppose that, up to extracting a subsequence (not relabeled), there exist $\lambda \in \mathcal{M}_+(\Omega)$ such that $\lambda_k \rightharpoonup \lambda$. We now use a blow-up argument. By the Radon-Nikodym Theorem we may decompose μ into two mutually singular measures in $(\mathcal{M}_+(\Omega))^M$

$$\mu = z\mathcal{H}^{n-1} \llcorner S(u) + \mu^s = z\mathcal{H}^{n-1} \llcorner S(u) + \zeta \|\mu^s\|_1$$

and λ into three mutually singular non-negative measures

$$\lambda = \xi \mathcal{H}^{n-1} \llcorner S(u) + \eta \|\mu^s\|_1 + \lambda^s,$$

where $\|\mu^s\|_1 := \sum_{l=1}^M \mu_l^s$. Hence we complete the proof if we show that

$$\xi(x_0) \geq f_{hom}(z(x_0), \nu_u(x_0)), \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x_0 \in S(u), \quad (4.37)$$

and that

$$\eta(x_0) \geq g_{hom}(\zeta(x_0)), \quad \text{for } |\mu^s|\text{-a.e. } x_0 \in \Omega. \quad (4.38)$$

The proof of (4.37) and (4.38) will be performed in two steps.

Step 1. Proof of (4.37).

By the properties of BV functions (see [9]) we have that, for \mathcal{H}^{n-1} -a.e. $x_0 \in S(u)$ it holds

$$(i) \quad \lim_{\rho \rightarrow 0^+} \frac{1}{\rho^n} \int_{x_0 + \rho Q_{\nu_u^\pm(x_0)}} |u(x) - u^\pm(x_0)| dx = 0,$$

$$(ii) \quad \lim_{\rho \rightarrow 0^+} \frac{1}{\rho^{n-1}} \mathcal{H}^{n-1}(S(u) \cap \{x_0 + \rho Q_{\nu_u(x_0)}\}) = 1$$

$$(iii) \quad \xi(x_0) = \lim_{\rho \rightarrow 0^+} \frac{1}{\rho^{n-1}} \lambda(\{x_0 + \rho Q_{\nu_u(x_0)}\})$$

$$(iv) \quad z(x_0) = \lim_{\rho \rightarrow 0^+} \frac{1}{\rho^{n-1}} \mu(\{x_0 + \rho Q_{\nu_u(x_0)}\})$$

Fix such a $x_0 \in S(u)$ and let (ρ_m) be a sequence of positive numbers converging to zero such that

$$\lambda(\partial\{x_0 + \rho_m Q_{\nu_u(x_0)}\}) = 0, \quad |\mu|(\partial\{x_0 + \rho_m Q_{\nu_u(x_0)}\}) = 0$$

By (ii) and (iii) we get

$$\begin{aligned} \xi(x_0) &= \lim_m \frac{1}{\rho_m^{n-1}} \lambda(\{x_0 + \rho_m Q_{\nu_u(x_0)}\}) \\ &\geq \lim_m \lim_k \frac{1}{\rho_m^{n-1}} F_{\varepsilon_k}(u_k, \{x_0 + \rho_m Q_{\nu_u(x_0)}\}), \end{aligned}$$

and by (iv) we have

$$\lim_m \lim_k \frac{1}{\rho_m^{n-1}} \mu(u_k)(\{x_0 + \rho_m Q_{\nu_u(x_0)}\}) = z(x_0) \quad (4.39)$$

Observe that, for every m and k we can find $\rho_{m,k}$ with $\lim_k \rho_{m,k} = \rho_m$ and $x_0^k \in \varepsilon_k \mathbf{Z}^n$ with $\lim_k x_0^k = x_0$, such that

$$\varepsilon_k \mathbf{Z}^n \cap (\{x_0^k + \rho_{m,k} Q_{\nu_u(x_0)}\}) = \varepsilon_k \mathbf{Z}^n \cap \{x_0 + \rho_m Q_{\nu_u(x_0)}\}$$

which in turn implies

$$F_{\varepsilon_k}(u_k, \{x_0 + \rho_m Q_{\nu_u(x_0)}\}) = F_{\varepsilon_k}(u_k, \{x_0^k + \rho_{m,k} Q_{\nu_u(x_0)}\}).$$

Then

$$\xi(x_0) \geq \lim_m \lim_k \frac{1}{\rho_{m,k}^{n-1}} F_{\varepsilon_k}(u_k, \{x_0^k + \rho_{m,k} Q_{\nu_u(x_0)}\}) \quad (4.40)$$

Let us set

$$u_{m,k}(a) = u_k(x_0^k + \rho_{m,k} a), \quad a \in \frac{\varepsilon_k}{\rho_{m,k}} \mathbf{Z}^n \cap Q_{\nu_u(x_0)}$$

and

$$u_0(x) = \begin{cases} u^+(x_0) & \text{if } (x, \nu_u(x_0)) > 0 \\ u^-(x_0) & \text{if } (x, \nu_u(x_0)) \leq 0. \end{cases}$$

Since $u_k \rightarrow u$ in $L^1(\Omega)$, by (i) we get

$$\lim_m \lim_k \int_{Q_{\nu_u(x_0)}} |u_{m,k}(x) - u_0(x)| dx = 0 \quad (4.41)$$

Moreover, observe that, by (4.39), we get

$$\lim_m \lim_k \mu(u_{m,k})(Q_{\nu_u(x_0)}) = z(x_0) \quad (4.42)$$

and that inequality (4.40) can be written as

$$\xi(x_0) \geq \lim_m \lim_k F_{\frac{\varepsilon_k}{\rho_{m,k}}}(u_{m,k}, Q_{\nu_u(x_0)}) \quad (4.43)$$

Let us show now that the mass of $\mu(u_{m,k})$ does not concentrate near $\partial Q_{\nu_u(x_0)}$ for m and k large enough.

Given $\delta > 0$, by (iv) there exists $\rho(\delta)$ such that $\forall \rho < \rho(\delta)$ and $\forall l \in \{1, \dots, M\}$ it holds

$$\rho^{n-1}(z_l(x_0) - \frac{\delta}{4}) < \mu_l(x_0 + \rho Q_{\nu_u(x_0)}) < \rho^{n-1}(z_l(x_0) + \frac{\delta}{4}).$$

Let $m(\delta)$ be such that $\rho_m < \rho(\delta) \forall m > m(\delta)$. Then, for every $t \in (0, 1]$ there holds:

$$(t\rho_m)^{n-1}(z_l(x_0) - \frac{\delta}{4}) < \mu_l(x_0 + t\rho_m Q_{\nu_u(x_0)}) < (t\rho_m)^{n-1}(z_l(x_0) + \frac{\delta}{4}). \quad (4.44)$$

Thus, for all t such that $|\mu|(x_0 + \partial(t\rho_m Q_{\nu_u(x_0)})) = 0$, by (4.44) we get

$$\mu_l(x_0 + (\rho_m Q_{\nu_u(x_0)} \setminus t\rho_m Q_{\nu_u(x_0)})) = \mu_l(x_0 + \rho_m Q_{\nu_u(x_0)}) - \mu_l(x_0 + t\rho_m Q_{\nu_u(x_0)})$$

$$< \rho_m^{n-1} \left(z_l(x_0)(1 - t^{n-1}) + \frac{\delta}{2} \right) \quad (4.45)$$

Let $t(\delta) < 1$ be such that for every $l \in \{1, \dots, M\}$ there holds $z_l(0)(1 - (t(\delta))^{n-1}) < \frac{\delta}{2}$. Then, by (4.45), for every $t \in (t(\delta), 1)$ we get

$$\mu_l(x_0 + (\rho_m Q_{\nu_u(x_0)} \setminus t \rho_m Q_{\nu_u(x_0)})) < \delta \rho_m^{n-1} \quad (4.46)$$

Set $\bar{t}(\delta) := \frac{1+t(\delta)}{2}$ and let $t_m \in (t(\delta), \bar{t}(\delta))$ be such that $|\mu|(x_0 + \partial(t_m \rho_m Q_{\nu_u(x_0)})) = 0$. Thus

$$\lim_k \mu(u_k)(x_0 + (\rho_m Q_{\nu_u(x_0)} \setminus t_m \rho_m Q_{\nu_u(x_0)})) = \mu(x_0 + (\rho_m Q_{\nu_u(x_0)} \setminus t_m \rho_m Q_{\nu_u(x_0)}))$$

and, by (4.46), we may conclude that for any $m > m(\delta)$ there exists $k(m)$ such that for every $l \in \{1, \dots, M\}$ and $k > k(m)$ there holds

$$\mu_l(u_k)(x_0 + (\rho_m Q_{\nu_u(x_0)} \setminus \bar{t}(\delta) \rho_m Q_{\nu_u(x_0)})) < \delta \rho_m^{n-1} \quad (4.47)$$

Hence, by (4.47) we infer that for m and k large enough

$$\mu_l(u_{m,k})(Q_{\nu_u(x_0)} \setminus \bar{t}(\delta) Q_{\nu_u(x_0)}) < \delta. \quad (4.48)$$

Taking into account (4.41), (4.42), (4.43) and (4.48), by a standard diagonalization procedure we can then find a sequence of positive numbers $s_j \rightarrow 0$ and a sequence $w_j \in \mathcal{A}_{s_j}(Q_{\nu_u(x_0)}; K)$ such that $w_j \rightarrow u_0$ in $L^1(Q_{\nu_u(x_0)})$ and there holds

$$\lim_j \mu(w_j)(Q_{\nu_u(x_0)}) = z(x_0), \quad (4.49)$$

$$\mu_l(w_j)(Q_{\nu_u(x_0)} \setminus \bar{t}(\delta) Q_{\nu_u(x_0)}) < \delta \quad \forall l \in \{1, \dots, M\}, \quad (4.50)$$

$$\xi(x_0) \geq \lim_j F_{s_j}(w_j, Q_{\nu_u(x_0)}).$$

Then, by virtue of Lemma 4.9, we can find a sequence $(v_j) \subset B_{s_j}(Q_{\nu_u(x_0)}; K)$ such that (4.62) holds and

$$\xi(x_0) \geq \lim_j F_{s_j}(v_j, Q_{\nu_u(x_0)}). \quad (4.51)$$

Moreover, by (4.49), (4.50) and (4.62) we have that for j large enough

$$|\mu_l(v_j)(Q_{\nu_u(x_0)}) - z_l(x_0)| < \delta \quad \forall l \in \{1, \dots, M\}. \quad (4.52)$$

Set $T_j := \left[\frac{1}{s_j} \right]$ and let $\hat{v}_j \in B_1(T_j Q_{\nu_u(x_0)}; K)$ defined by

$$\hat{v}_j(a) := v(s_j a) \quad a \in \mathbf{Z}^n \cap T_j Q_{\nu_u(x_0)}.$$

Then (4.52) implies that for j large enough

$$\left| \frac{\#I_l(\hat{v}_j)}{T_j^{n-1}} - z_l \right| < \delta \quad \forall l \in \{1, \dots, M\}$$

and (4.51) reads

$$\xi(x_0) \geq \lim_j \frac{1}{T_j^{n-1}} F_1(\hat{v}_j, T_j Q_{\nu_u(x_0)}).$$

Hence (4.37) immediately follows by the definition of f_{hom} and by Proposition 4.6.

Step 2. Proof of (4.38).

For $|\mu^s|$ -a.e $x_0 \in \Omega$ we have

$$(v) \quad \eta(x_0) = \lim_{\rho \rightarrow 0} \frac{\lambda(x_0 + \rho Q)}{\|\mu\|_1(x_0 + \rho Q)};$$

$$(vi) \quad \zeta(x_0) = \lim_{\rho \rightarrow 0} \frac{\mu(x_0 + \rho Q)}{\|\mu\|_1(x_0 + \rho Q)}$$

Fix such a $x_0 \in \Omega$ and let (ρ_m) be a sequence of positive numbers converging to zero such that

$$\lambda(\partial\{x_0 + \rho_m Q_{\nu_u(x_0)}\}) = 0, \quad |\mu|(\partial\{x_0 + \rho_m Q_{\nu_u(x_0)}\}) = 0$$

By (v) and (vi) we get

$$\eta(x_0) = \lim_m \lim_k \frac{\lambda_k(x_0 + \rho_m Q)}{\|\mu(u_k)\|_1(x_0 + \rho_m Q)}, \quad (4.53)$$

$$\zeta(x_0) = \lim_m \lim_k \frac{\mu(u_k)(x_0 + \rho_m Q)}{\|\mu(u_k)\|_1(x_0 + \rho_m Q)}. \quad (4.54)$$

We now show that for a suitable sequence $k_m \in \mathbf{N}$ the mass of λ_{k_m} does not concentrate near $\partial(x_0 + \rho_m Q)$.

By the inner regularity of λ , given $\delta > 0$, for any $\rho > 0$ with $\lambda(\partial(x_0 + \rho Q)) = 0$ there exists $t(\rho)$ such that for all $t \in [t(\rho), 1]$ there holds

$$0 < \lambda(x_0 + \rho Q) - \lambda(x_0 + t\rho Q) < \delta \|\mu\|_1(x_0 + \rho Q).$$

Let $t_m \in [t(\rho_m), 1]$ be such that $\lambda(\partial(x_0 + t_m \rho_m Q)) = 0$. Then

$$\lambda(x_0 + (\rho_m Q \setminus t_m \rho_m Q)) = \lambda(x_0 + \rho_m Q) - \lambda(x_0 + t_m \rho_m Q) < \delta \|\mu\|_1(x_0 + \rho_m Q)$$

In particular, since $\lim_k \lambda_k(x_0 + (\rho_m Q \setminus t_m \rho_m Q)) = \lambda(x_0 + (\rho_m Q \setminus t_m \rho_m Q))$ and $\lim_k \|\mu(u_k)\|_1(x_0 + \rho_m Q) = \|\mu\|_1(x_0 + \rho_m Q)$, we have that for k large enough

$$\lambda_k(x_0 + (\rho_m Q \setminus t_m \rho_m Q)) < \delta \|\mu(u_k)\|_1(x_0 + \rho_m Q).$$

Hence, by the previous inequality and by (4.53) and (4.54), we can find a sequence k_m such that $\varepsilon_{k_m} \ll \rho_m$ and

$$\eta(x_0) = \lim_m \frac{\lambda_{k_m}(x_0 + \rho_m Q)}{\|\mu(u_{k_m})\|_1(x_0 + \rho_m Q)} \geq \lim_m \frac{F_{\varepsilon_{k_m}}(u_{k_m}, x_0 + \rho_m Q)}{\|\mu(u_{k_m})\|_1(x_0 + \rho_m Q)}, \quad (4.55)$$

$$\zeta(x_0) = \lim_m \frac{\mu(u_{k_m})(x_0 + \rho_m Q)}{\|\mu(u_{k_m})\|_1(x_0 + \rho_m Q)}, \quad (4.56)$$

$$\begin{aligned} \lambda_{k_m}(x_0 + (\rho_m Q \setminus t_m \rho_m Q)) &< \delta \|\mu(u_{k_m})\|_1(x_0 + \rho_m Q), \\ R\varepsilon_{k_m} &< \rho_m(1 - t_m). \end{aligned}$$

Note that the two last inequalities imply that

$$F_{\varepsilon_{k_m}}(u_{k_m}, x_0 + (\rho_m Q \setminus (\rho_m - R\varepsilon_{k_m})Q)) < \delta \|\mu(u_{k_m})\|_1(x_0 + \rho_m Q) \quad (4.57)$$

Observe that, for every m we can find $\tilde{\rho}_m$ with $\lim_m \frac{\tilde{\rho}_m}{\rho_m} = 1$ and $x_0^m \in \varepsilon_{k_m} \mathbf{Z}^n$ with $\lim_m x_0^m = x_0$, such that

$$\varepsilon_{k_m} \mathbf{Z}^n \cap (x_0^m + \tilde{\rho}_m Q) = \varepsilon_{k_m} \mathbf{Z}^n \cap (x_0 + \rho_m Q)$$

which in turn implies

$$F_{\varepsilon_{k_m}}(u_{k_m}, x_0 + \rho_m Q) = F_{\varepsilon_{k_m}}(u_{k_m}, x_0^m + \tilde{\rho}_m Q).$$

$$\mu(u_{k_m})(x_0 + \rho_m Q) = \mu(u_{k_m})(x_0^m + \tilde{\rho}_m Q).$$

Set $T_m := \frac{\tilde{\rho}_m}{\varepsilon_{k_m}}$ and let $\hat{u}_m \in \mathcal{A}_1(T_m Q; K)$ be defined by

$$\hat{u}_m(a) = u_{k_m}(x_0^m + \varepsilon_{k_m} a) \quad a \in \mathbf{Z}^n \cap T_m Q$$

Note that, by definition, $\|\mu(\hat{u}_m)\|_1(A) = \#I(\hat{u}_m, A)$ for any $A \subset \mathbf{R}^n$. Then (4.55) and (4.56) read

$$\eta(x_0) \geq \lim_m \frac{F_1(\hat{u}_m, T_m Q)}{\#I(\hat{u}_m, T_m Q)}. \quad (4.58)$$

$$\zeta(x_0) = \lim_m \frac{\mu(\tilde{u}_m)(Q)}{\#I(\hat{u}_m, T_m Q)}. \quad (4.59)$$

Moreover, by (4.57), we get

$$F_1(\hat{u}_m, T_m Q \setminus (T_m - R)Q) < \delta \#I(\hat{u}_m, T_m Q). \quad (4.60)$$

Hence, (4.38) immediately follows by the definition of g_{hom} , taking into account (4.58), (4.59) and (4.60).

□

Set, for $\nu \in S^{n-1}$,

$$u_\nu = \begin{cases} m_1 & \text{if } (x, \nu) > 0 \\ m_2 & \text{if } (x, \nu) \leq 0. \end{cases} \quad (4.61)$$

Moreover we recall that we have set $Q_\nu = (-r_\nu, r_\nu)^n$.

Lemma 4.9 *Let $s_j \rightarrow 0^+$, $\nu \in S^{n-1}$ and let $w_j \in \mathcal{A}_{s_j}(Q_\nu; K)$ be such that $w_j \rightarrow u_\nu$ in $L^1(Q_\nu)$. Then there exist $v_j \in B_{s_j}(Q_\nu; K)$ such that*

$$v_j \equiv u_\nu \text{ on } Q_\nu \setminus Q_j, \quad (4.62)$$

where $Q_j := (-r_j, r_j)^n \subset Q_\nu$, for some $r_j > 0$ such that $\lim_j r_j = r_\nu$, and

$$\liminf_j F_{s_j}(w_j, Q_\nu) \geq \liminf_j F_{s_j}(v_j, Q_\nu). \quad (4.63)$$

PROOF. Set

$$\delta_j := \int_{Q_\nu} |w_j - u_\nu| dx$$

Let $k_j \in \mathbf{N}$ be such that

$$\frac{\delta_j}{s_j} \ll k_j \ll \frac{1}{s_j} \quad (4.64)$$

and set, for $i \in \{0, \dots, k_j\}$,

$$r_j^i := \left\lceil \frac{r_\nu}{s_j} \right\rceil + (i - k_j)M$$

$$Q_j^i := (-r_j^i s_j, r_j^i s_j).$$

Then we get

$$\delta_j \geq \int_{Q_\nu \setminus Q_j^0} |w_j - u_\nu| dx \geq \sum_{i=0}^{k_j-1} \int_{Q_j^{i+1} \setminus Q_j^i} |w_j - u_\nu| dx.$$

Hence, there exists $i_j \in \{0, \dots, k_j - 1\}$ such that, set $S_j := Q_j^{i_j+1} \setminus Q_j^{i_j}$, we have

$$\delta_j \geq k_j \int_{S_j} |w_j - u_\nu| dx \geq C k_j s_j^n \#\{a \in s_j \mathbf{Z}^n \cap S_j : w_j(a) \neq u_\nu(a)\},$$

which in turn, by (4.64), implies that

$$s_j^{n-1} \#\{a \in \lambda_j \mathbf{Z}^n \cap S_j : w_j(a) \neq u_\nu(a)\} \rightarrow 0. \quad (4.65)$$

Let, then, $v_j \in B_{s_j}(Q_\nu; K)$ defined by

$$v_j(a) := \begin{cases} w_j(a) & \text{if } a \in s_j \mathbf{Z}^n \cap Q_j^{i_j} \\ u_\nu(a) & \text{otherwise.} \end{cases}$$

Thus, by (4.65), we get

$$\begin{aligned} F_{s_j}(v_j, Q_\nu) &\leq F_{s_j}(w_j, Q_j^{i_j}) + F_{s_j}(u_\nu, Q_\nu \setminus Q_j^{i_j}) \\ &\quad + C s_j^{n-1} \#\{a \in s_j \mathbf{Z}^n \cap S_j : w_j(a) \neq u_\nu(a)\} \leq F_j(w_j, Q) + o(1), \end{aligned}$$

from which we get the conclusion \square

Proposition 4.10 (Γ -lim sup inequality) *We have*

$$\Gamma\text{-lim sup}_{\varepsilon \rightarrow 0} F_\varepsilon(u, \mu) \leq F(u, \mu). \quad (4.66)$$

PROOF. We will use the notation $F'' := \Gamma\text{-lim sup}_{\varepsilon \rightarrow 0} F_\varepsilon$. We split the proof in several steps.

Step 1. Claim: (4.66) holds for every $(u, \mu) \in BV(\Omega; \{m_1, m_2\}) \times (\mathcal{M}_+(\Omega))^M$ such that $S(u)$ is a polyhedral set and μ is of the form $\mu = \varphi \mathcal{H}^{n-1} \llcorner S(u) + \sum_{j=1}^N w_j \delta_{x_j}$, where $\varphi : \Omega \rightarrow \mathbf{R}^M$ is a piecewise-constant function, $N \in \mathbf{N}$ and, for all $j \in \{1, 2, \dots, N\}$, $w_j \in (\mathbf{R}_+)^M$ and $x_j \in \Omega$.

Since the construction we provide is local, without loss of generality, we prove the claim in the particular case $u = u_\nu$, and $\mu = z \mathcal{H}^{n-1} \llcorner_{S(u)} + w \delta_0$ with $\nu \in S^{n-1}$, $z, w \in (\mathbf{R}_+)^M$. Here, without loss of generality, we also suppose $0 \in \Omega$. Note that

$$F(u, \mu) = f_{hom}(z, \nu) \mathcal{H}^{n-1}(S(u)) + g_{hom}(\zeta) \|w\|_1,$$

where $\zeta = \frac{w}{\|w\|_1}$. By the lower semicontinuity of F'' , in order to show (4.66), it suffices to prove that there exists $(\mu_j)_j \subset (\mathcal{M}_+(\Omega))^M$ weakly converging to μ as $j \rightarrow +\infty$ such that, for every $j \in \mathbf{N}$, there exists $u_\varepsilon \in \mathcal{A}_\varepsilon(\Omega; K)$ such that $(u_\varepsilon, \mu(u_\varepsilon)) \rightarrow (u, \mu_j)$ with respect to the $\tau_1 \times \tau_2$ -convergence and

$$\limsup_\varepsilon F_\varepsilon(u_\varepsilon) \leq F(u, \mu) + \frac{C}{j}. \quad (4.67)$$

For simplicity of notation we provide the construction of such u_ε in the case $\nu = e_n$, the same argument applying to the general case. Such a u_ε will be obtained by scaling the periodic extension of an optimal function for the problem defining f_{hom} in a neighborhood of $S(u)$ and a proper extension of an optimal function for the

problem defining g_{hom} in a suitable neighborhood of 0.

Let $0 < \delta_j < \frac{1}{j}$, $T_j > 0$, $u_j \in \mathcal{B}_1(T_j Q; K)$ and $v_j \in \mathcal{A}_1(T_j Q; K)$ be such that

$$\begin{aligned} \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(u_j, T_j Q)}{T_j^{n-1}} - z_l \right| &\leq \delta_j, & \max_{l \in \{1, \dots, M\}} \left| \frac{\#I_l(v_j, T_j Q)}{\#I(v_j, T_j Q)} - \zeta_l \right| &\leq \delta_j \\ \frac{1}{T_j^{n-1}} F_1(u_j, T_j Q) &\leq f_{hom}(z, e_n) + \frac{1}{j}, \\ \frac{F_1(v_j, T_j Q)}{\#I(v_j, T_j Q)} &\leq g_{hom}(\zeta) + \frac{1}{j}, \\ \frac{F_1(v_j, T_j Q \setminus (T_j - R)Q)}{\#I(v_j, T_j Q)} &\leq \delta_j. \end{aligned}$$

Without loss of generality we choose T_j to be an even number. With a little abuse of notation we consider u_j to be extended by periodicity to \mathbf{Z}^n . Moreover we extend v_j by reflection with respect to the coordinate axes. More precisely we set $v_{j,0} = v_j$ and, for all $k \in \{1, \dots, n\}$, we define $v_{j,k}$ recursively as follows: $v_{j,k}$ is the extension of $v_{j,k-1}$ on $\mathbf{R}^k \times \left[-\frac{T_j}{2}, \frac{T_j}{2}\right]^{n-k}$ satisfying the following property

$$v_{j,k}(a) = v_{j,k}(a + (T_j - 2a_k)e_k), \quad \forall a \in \mathbf{R}^{k-1} \times \left[-\frac{T_j}{2}, \frac{T_j}{2}\right]^{n-(k-1)} + hT_j e_k, \quad h \in \mathbf{Z}.$$

We have then obtained that the function $v_{j,n}$ extends v_j on all \mathbf{Z}^n . Let us observe that, by the symmetry hypotheses in (4.12), we have that

$$F_1(v_{j,n}, T_j Q + h) = F_1(v_j, T_j Q) \quad \forall h \in \mathbf{Z}^n. \quad (4.68)$$

Let $\tilde{u}_\varepsilon : \varepsilon \mathbf{Z}^n \rightarrow K$ be defined as

$$\tilde{u}_\varepsilon(a) = \begin{cases} m_1 & \text{if } a_n \geq \varepsilon \frac{T}{2} \\ u_j\left(\frac{a}{\varepsilon}\right) & \text{if } |a_n| \leq \varepsilon \frac{T}{2} \\ m_2 & \text{if } a_n \leq -\varepsilon \frac{T}{2} \end{cases}$$

and set

$$k_\varepsilon = \left[\left(\frac{\|w\|_1}{\varepsilon^{n-1} \#I(v_j, T_j Q)} \right)^{\frac{1}{n}} \right]. \quad (4.69)$$

Note that $\tilde{u}_\varepsilon \rightarrow u_\nu$ in $L^1(\Omega)$ and that $\varepsilon k_\varepsilon \rightarrow 0$. We now define $u_\varepsilon : \varepsilon \mathbf{Z}^n \rightarrow K$ as

$$u_\varepsilon(a) = \begin{cases} v_{j,n}\left(\frac{a}{\varepsilon}\right) & \text{if } a \in \varepsilon k_\varepsilon T_j Q \\ \tilde{u}_\varepsilon(a) & \text{otherwise.} \end{cases}$$

Then u_ε still converges to u_ν in $L^1(\Omega)$. Moreover, by construction, we have that $\mu(u_\varepsilon) \rightarrow \mu^j \in (\mathcal{M}_+(\Omega))^M$ where, for $l \in \{1, 2, \dots, M\}$,

$$\mu_l^j = \frac{\#I_l(u_j, T_j Q)}{T_j^{n-1}} \mathcal{H}^{n-1} \llcorner S(u) + \frac{\#I_l(v_j, T_j Q)}{\#I(v_j, T_j Q)} \|w\|_1 \delta_0.$$

We can now estimate the energy of u_ε . Taking into account the invariance of the energy under integer translations and (4.68), we get

$$F_\varepsilon(u_\varepsilon) \leq \left[\frac{\mathcal{H}^{n-1}(S(u))}{(T_j\varepsilon)^{n-1}} \right] \varepsilon^{n-1} F_1(u_j, T_j Q) + k_\varepsilon^n \varepsilon^{n-1} F_1(v_j, T_j Q) + C k_\varepsilon^n \varepsilon^{n-1} F_1(v_j, T_j Q \setminus (T_j - R)Q) + o(1),$$

where the third term in the right-hand-side is obtained by estimating the energy due to the interactions that cross the boundary of each cube of size εT_j contained in $\varepsilon K_\varepsilon T_j Q$. By (4.68) we eventually have

$$F_\varepsilon(u_\varepsilon) \leq \left[\frac{\mathcal{H}^{n-1}(S(u))}{(T_j\varepsilon)^{n-1}} \right] \varepsilon^{n-1} T_j^{n-1} \left(f_{hom}(z, e_n) + \frac{1}{j} \right) + k_\varepsilon^n \varepsilon^{n-1} \#I(v_j, T_j Q) \left(g_{hom}(\zeta) + C\delta_j + \frac{1}{j} \right) + o(1)$$

The conclusion follows passing to the limsup as ε tends to 0, taking into account (4.69).

Step 2. Claim: (4.66) holds for every (u, μ) as in Step 1 but with $\varphi \in C(\Omega; \mathbf{R}^M)$.

Let φ_k be a sequence of piecewise constant functions such that $\varphi_k \rightarrow \varphi$ with respect to the $L^1(S(u); \mathcal{H}^{n-1})$ and let $\mu_k = \varphi_k \mathcal{H}^{n-1} \lfloor S(u) + \sum_{j=1}^N w_j \delta_{x_j}$. Then $\mu_k \rightharpoonup \mu$, and by the convexity and growth properties of $f_{hom}(\cdot, \nu)$ stated in Proposition 4.7, $F(u, \mu_k) \rightarrow F(u, \mu)$. Eventually, by the lower semicontinuity of $F''(u, \mu)$ and by Step 1, we have

$$F''(u, \mu) \leq \liminf_k F''(u, \mu_k) \leq \liminf_k F(u, \mu_k) = F(u, \mu).$$

Step 3. Claim: (4.66) holds for every $(u, \mu) \in BV(\Omega; \{m_1, m_2\}) \times (\mathcal{M}_+(\Omega))^M$ such that $\mu = \varphi \mathcal{H}^{n-1} \lfloor S(u) + \sum_{j=1}^N w_j \delta_{x_j}$ with $\varphi \in C(\Omega; \mathbf{R}^M)$.

Let $u_k \in BV(\Omega; \{m_1, m_2\})$ be such that $u_k \rightarrow u$ in $L^1(\Omega; \{m_1, m_2\})$, $S(u_k)$ is a polyhedral set and $\mathcal{H}^{n-1}(S(u_k)) \rightarrow \mathcal{H}^{n-1}(S(u))$. Let $\mu_k = \varphi \mathcal{H}^{n-1} \lfloor S(u_k) + \sum_{j=1}^N w_j \delta_{x_j}$. Then we have that $\mu_k \rightharpoonup \mu$ and $|\varphi| \mathcal{H}^{n-1} \lfloor S(u_k)(\Omega) \rightarrow |\varphi| \mathcal{H}^{n-1} \lfloor S(u)(\Omega)$. Then, by the convexity of f_{hom} stated in Proposition 4.7 and by Reshetnyak's theorem we have that $F(u_k, \mu_k) \rightarrow F(u, \mu)$. Hence we conclude as in Step 2.

Step 4. Claim: (4.66) holds for every $(u, \mu) \in BV(\Omega; \{m_1, m_2\}) \times (\mathcal{M}_+(\Omega))^M$.

Let $\varphi_k \in C(\Omega; \mathbf{R}^M)$ be such that $\varphi_k \rightarrow \varphi$ in $L^1(S(u); \mathcal{H}^{n-1})$ and let $\mu_k^s = \sum_{j=1}^N w_j \delta_{x_j}$ be such that $\mu_k^s \rightharpoonup \mu^s$ and $|\mu_k^s|(\Omega) = |\mu^s|(\Omega)$. Let then $\mu_k = \varphi_k \mathcal{H}^{n-1} \lfloor S(u) + \mu_k^s$. We have that $\mu_k \rightharpoonup \mu$ and, by the convexity and growth properties of $f_{hom}(\cdot, \nu)$ and the convexity of g_{hom} stated in Proposition 4.7, applying Reshetnyak's theorem we get that $F(u_k, \mu_k) \rightarrow F(u, \mu)$. Hence we conclude as in Step 2. \square

4.2 Prescribed volume-fractions

In this Section we study a generalization of the constrained minimum problems introduce at the end of Section 3.2 in the case of the BEG model.

In what follows we set $\mathcal{Q} = [0, 1]^n$, $\varepsilon_k = \frac{1}{k}$, $k \in \mathbf{N}$ (for simplicity of notation we will drop the k and write ε instead of ε_k),

$$\mathcal{A}_\varepsilon^\#(\mathcal{Q}; K) := \{u : \varepsilon \mathbf{Z}^n \rightarrow K : u \text{ is } \mathcal{Q}\text{-periodic}\}$$

and define $F_\varepsilon^\# : \mathcal{A}_\varepsilon^\#(\mathcal{Q}; K) \rightarrow \mathbf{R}$ as

$$F_\varepsilon^\#(u) = \sum_{|\xi| \leq R} \sum_{a \in \mathcal{Q}_\varepsilon} \varepsilon^{n-1} f(\xi, u(a), u(a + \varepsilon \xi)).$$

Given $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_M) \in \mathbf{R}^M$ and $\beta \in \mathbf{R}$, let $\alpha_\varepsilon = (\alpha_{1,\varepsilon}, \alpha_{2,\varepsilon}, \dots, \alpha_{M,\varepsilon}) \rightarrow \alpha$ and $\beta_\varepsilon \rightarrow \beta$ as $\varepsilon \rightarrow 0$. We define the set of admissible functions $\mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)$ as

$$\mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K) := \left\{ u \in \mathcal{A}_\varepsilon^\#(\mathcal{Q}; K) : \begin{aligned} \varepsilon^{n-1} \# I_l(u, \mathcal{Q}) &= \alpha_{l,\varepsilon} \forall l \in \{1, \dots, M\}, \\ \varepsilon^n \# I_{m_1}(u, \mathcal{Q}) &= \beta_\varepsilon \end{aligned} \right\},$$

where we have set $I_{m_1}(u, \mathcal{Q}) := \{a \in \mathcal{Q}_\varepsilon : u(a) = m_1\}$, and consider the family of minimum problems

$$m_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon} := \min\{F_\varepsilon^\#(u) : u \in \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)\}.$$

Note that if $u \in \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)$ then $\mu(u_\varepsilon)(\mathcal{Q}) = \alpha_\varepsilon$. We are interested in studying the limit, as $\varepsilon \rightarrow 0$, of $m_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}$. To this end, we introduce the family of functionals $F_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon} : L_{loc}^1(\mathbf{R}^n; K) \times (\mathcal{M}_+^\#(\mathbf{R}^n))^M \rightarrow [0, +\infty]$ defined as

$$F_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(u, \mu) = \begin{cases} F_\varepsilon^\#(u) & \text{if } u \in \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K), \mu = \mu(u), \\ +\infty & \text{otherwise,} \end{cases}$$

where $u \in \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)$ is identified with its piecewise-constant interpolation on the cells of the lattice $\varepsilon \mathbf{Z}^n$, $\mu(u)$ is the surfactant measure defined in (4.16) and $\mathcal{M}_+^\#(\mathbf{R}^n)$ is the space of \mathcal{Q} -periodic non negative Radon measures. We endow the space $L_{loc}^1(\mathbf{R}^n; K) \times (\mathcal{M}_+^\#(\mathbf{R}^n))^M$ with the convergence $\tau_1 \times \tau_2$ where τ_1 denotes the strong convergence in $L_{loc}^1(\mathbf{R}^n)$ and τ_2 denotes the weak*-convergence in $(\mathcal{M}_+^\#(\mathbf{R}^n))^M$. Then the following Theorem holds true.

Theorem 4.11 *The family $(F_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon})$ Γ -converges with respect to the $\tau_1 \times \tau_2$ convergence to the functional $F^{\alpha, \beta} : L_{loc}^1(\mathbf{R}^n) \times (\mathcal{M}_+^\#(\mathbf{R}^n))^M \rightarrow [0, +\infty]$ defined by*

$$F^{\alpha, \beta}(u, \mu) = \begin{cases} \bar{F}(u, \mu) & \text{if } u \in BV^\#(\mathbf{R}^n; \{m_1, m_2\}), |\{x \in \mathcal{Q} : u(x) = m_1\}| = \beta, \\ & \text{and } \mu(\bar{\mathcal{Q}}) = \alpha, \\ +\infty & \text{otherwise,} \end{cases}$$

where

$$\bar{F}(u, \mu) = \int_{S(u) \cap \bar{\mathcal{Q}}} f_{hom} \left(\frac{d\mu}{d\mathcal{H}^{n-1} \llcorner S(u)}, \nu(u) \right) d\mathcal{H}^{n-1} + \int_{\bar{\mathcal{Q}}} g_{hom}(\mu^s),$$

$\mu^s = \mu - \frac{d\mu}{d\mathcal{H}^{n-1} \llcorner S(u)} \mathcal{H}^{n-1} \llcorner S(u)$, and the densities f_{hom} and g_{hom} are defined in (4.19) and in (4.20).

PROOF. It is easy to show that if $u_\varepsilon \in \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)$ and $(u_\varepsilon, \mu(u_\varepsilon)) \rightarrow (u, \mu)$ with respect to the $\tau_1 \times \tau_2$ convergence, then $(u, \mu) \in BV^\#(\mathbf{R}^n; \{m_1, m_2\}) \times (\mathcal{M}_+^\#(\mathbf{R}^n))^M$, $|\{x \in \mathcal{Q} : u(x) = m_1\}| = \beta$ and $\mu(\bar{\mathcal{Q}}) = \alpha$. The Γ -lim inf inequality follows by Theorem 4.4.

The proof of the opposite inequality can be obtained by following the lines of the proof of the Γ -lim sup inequality of Theorem 4.4, with some extra care to show that the recovery sequence u_ε for $(u, \mu) \in BV^\#(\mathbf{R}^n; \{m_1, m_2\}) \times (\mathcal{M}_+^\#(\mathbf{R}^n))^M$ such that $\mu(\bar{\mathcal{Q}}) = \alpha$ and $|\{x \in \mathcal{Q} : u(x) = m_1\}| = \beta$ can be slightly modified so that $u_\varepsilon \in \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)$. \square

As a consequence of the previous Theorem, by the standard properties of Γ -convergence (we refer the reader to [13] and [20]), we derive the following result about the convergence of the family of minimum problems defined above.

Corollary 4.12 *There holds*

$$\lim_\varepsilon m_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon} = \min\{\bar{F}(u, \mu), |\{x \in \mathcal{Q} : u(x) = m_1\}| = \beta, \mu(\bar{\mathcal{Q}}) = \alpha\}.$$

Moreover if $(u_\varepsilon) \subset \mathcal{A}_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(\mathcal{Q}; K)$ is such that

$$\lim_\varepsilon F_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(u_\varepsilon) = \lim_\varepsilon m_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon},$$

then any cluster point $(\bar{u}, \bar{\mu})$ of $(u_\varepsilon, \mu(u_\varepsilon))$ with respect to the $\tau_1 \times \tau_2$ convergence is a minimizer for $\min\{\bar{F}(u, \mu), |\{x \in \mathcal{Q} : u(x) = m_1\}| = \beta, \mu(\bar{\mathcal{Q}}) = \alpha\}$.

Remark 4.13 The previous Corollary applies to the case of the BEG model, $f(\xi, u, v)$ being defined in (4.15) and the limit energy densities f_{hom} and g_{hom} being given by

$$f_{hom}(z, \nu) = \varphi(z, \nu), \quad g_{hom}(1) = 2(1 - k),$$

with φ defined in (3.9).

5 The Blume-Emery-Griffiths model: proof of Theorem 3.3

By Remark 4.2, the functionals $E_\varepsilon^{(1)}$ satisfy all the hypotheses of Proposition 4.3 and Theorem 4.4. Hence the compactness result asserted in (i) follows by

Proposition 4.3. Moreover the integral representation result stated in Theorem 4.4 holds true for the Γ -limit of $E_\varepsilon^{(1)}$. Thus, in order to conclude, it is only left to prove that, for all $(z, \nu) \in \mathbf{R}^+ \times S^1$,

$$\begin{aligned} f_{hom}(z, \nu) &= \varphi(z, \nu), \\ g_{hom}(1) &= 2(1 - k). \end{aligned} \quad (5.70)$$

Without loss of generality we prove (5.70) for $\nu_1, \nu_2 > 0$.

Step 1 (lower bounds). In this Step we prove the following two inequalities:

$$\begin{aligned} f_{hom}(z, \nu) &\geq \varphi(z, \nu), \\ g_{hom}(1) &\geq 2(1 - k). \end{aligned} \quad (5.71)$$

Let us first prove the lower bound for f_{hom} . Without loss of generality we consider T to be an even number. Let $J_{1,T}, J_{2,T}$ be the following sets of integers:

$$\begin{aligned} J_{1,T} &= \left\{ -\left\lfloor \frac{T\nu_2}{2} \right\rfloor, -\left\lfloor \frac{T\nu_2}{2} \right\rfloor + 1, \dots, \left\lfloor \frac{T\nu_2}{2} \right\rfloor - 1, \left\lfloor \frac{T\nu_2}{2} \right\rfloor \right\} \\ J_{2,T} &= \left\{ -\left\lfloor \frac{T\nu_1}{2} \right\rfloor, -\left\lfloor \frac{T\nu_1}{2} \right\rfloor + 1, \dots, \left\lfloor \frac{T\nu_1}{2} \right\rfloor - 1, \left\lfloor \frac{T\nu_1}{2} \right\rfloor \right\}. \end{aligned}$$

Let u be an admissible test function in the problem defining $f_{hom}(z, \nu)$ in (4.19); that is $u \in \mathcal{B}_\varepsilon(TQ_\nu; \{\pm 1, 0\})$ and

$$\left| \frac{\#I_0(u, TQ_\nu)}{T} - z \right| < \delta. \quad (5.72)$$

We define

$$J_{1,T}^0(u) = \{i \in J_{1,T} : \exists j \in \mathbf{Z} \text{ such that } (i, j) \in I_0(u)\}$$

and

$$J_{2,T}^0(u) = \{j \in J_{2,T} : \exists i \in \mathbf{Z} \text{ such that } (i, j) \in I_0(u)\}.$$

Note that, by (5.72), for $i \in \{1, 2\}$

$$\#J_{i,T}^0(u) \leq \#I_0(u, TQ_\nu) \leq (z + \delta)T.$$

The proof will be the result of the following three estimates.

Estimate (i). By a slicing argument, splitting the energy into the contribution of the horizontal and the vertical interactions, we get

$$\begin{aligned} E_1^{(1)}(u) &\geq 2(1 - k)\#J_{1,T}^0(u) + 2(\#J_{1,T} - \#J_{1,T}^0(u)) \\ &\quad + 2(1 - k)\#J_{2,T}^0(u) + 2(\#J_{2,T} - \#J_{2,T}^0(u)) \\ &= -2k\#J_{1,T}^0(u) - 2k\#J_{2,T}^0(u) + 2(\#J_{1,T} + \#J_{2,T}) \\ &\geq -4k(z + \delta)T + 2\left(2\left\lfloor \frac{T\nu_2}{2} \right\rfloor + 2\left\lfloor \frac{T\nu_1}{2} \right\rfloor\right). \end{aligned}$$

There follows that:

$$\begin{aligned} f_{hom}(z, \nu) &\geq \lim_{\delta \rightarrow 0} \lim_{T \rightarrow +\infty} \frac{1}{T} \left(-4k(z + \delta)T + 2 \left(2 \left[\frac{T\nu_2}{2} \right] + 2 \left[\frac{T\nu_1}{2} \right] \right) \right) \\ &= -4kz + 2(\nu_1 + \nu_2). \end{aligned}$$

Estimate (ii). We observe that we may split the energy of the surfactants into two terms. The first term accounts for 2 vertical and 1 horizontal interaction for each column, while the second term accounts for the remaining interaction in each row. By counting the energy due to the non-surfactant particles as in estimate (i), we get

$$\begin{aligned} E_1^{(1)}(u) &\geq 3(1-k)\#J_{1,T}^0(u) + 2(\#J_{1,T} - \#J_{1,T}^0(u)) \\ &\quad + (1-k)\#J_{2,T}^0(u) + 2(\#J_{2,T} - \#J_{2,T}^0(u)) \\ &\geq (1-3k)\#J_{1,T}^0(u) + 2\#J_{1,T} \\ &\quad + (1-k)\#J_{2,T}^0(u) + (1-k)(\#J_{2,T} - \#J_{2,T}^0(u)) \\ &\geq (1-3k)(z + \delta)T + 2\#J_{1,T} + (1-k)\#J_{2,T}(u) \\ &\geq (1-3k)(z + \delta)T + 4 \left[\frac{T\nu_2}{2} \right] + 2(1-k) \left[\frac{T\nu_1}{2} \right] \end{aligned}$$

By exchanging the role of $J_{1,T}^0(u)$ with that of $J_{2,T}^0(u)$ in the previous estimate we have

$$E_1^{(1)}(u) \geq (1-3k)(z + \delta)T + 4 \left[\frac{T(\nu_1 \vee \nu_2)}{2} \right] + 2(1-k) \left[\frac{T(\nu_1 \wedge \nu_2)}{2} \right]$$

Therefore we get

$$\begin{aligned} f_{hom}(z, \nu) &\geq \lim_{\delta \rightarrow 0} \lim_{T \rightarrow +\infty} \frac{1}{T} (1-3k)(z + \delta)T + 4 \left[\frac{T(\nu_1 \vee \nu_2)}{2} \right] + 2(1-k) \left[\frac{T(\nu_1 \wedge \nu_2)}{2} \right] \\ &\geq (1-3k)z + 2(\nu_1 \vee \nu_2) + (1-k)(\nu_1 \wedge \nu_2). \end{aligned}$$

Estimate (iii). We observe that the we may split the energy of the surfactant into three terms. The first term takes into account two interactions for each surfactant particle. The other two terms take into account in each row and column containing a surfactant at least one interaction between a surfactant and a non-surfactant particle. Counting as in the previous estimates the remaining interactions, we have

$$\begin{aligned} E_1^{(1)}(u) &\geq 2(1-k)\#I_0(u, TQ\nu) + 2(\#J_{1,T} - \#J_{1,T}^0) + (1-k)\#J_{1,T}^0 \\ &\quad + 2(\#J_{2,T} - \#J_{2,T}^0) + (1-k)\#J_{2,T}^0 \\ &\geq 2(1-k)(z - \delta)T + (1-k)(\#J_{1,T} + \#J_{2,T}) \\ &\geq 2(1-k)(z - \delta)T + (1-k) \left(2 \left[\frac{T\nu_1}{2} \right] + 2 \left[\frac{T\nu_2}{2} \right] \right). \end{aligned}$$

Hence we have

$$\begin{aligned} f_{hom}(z, \nu) &\geq \lim_{\delta \rightarrow 0} \lim_{T \rightarrow +\infty} \frac{1}{T} \left(2(1-k)(z-\delta)T + (1-k) \left(2 \left[\frac{T\nu_1}{2} \right] + 2 \left[\frac{T\nu_2}{2} \right] \right) \right) \\ &\geq 2(1-k)z + (1-k)(\nu_1 + \nu_2). \end{aligned}$$

The lower bound in (5.71) for g_{hom} is straightforward. In fact, let us observe that, except for a negligible error due to the interactions at the boundary of TQ , the energy accounts for at least two interactions of each surfactant particle. Hence, for any test function u in the minimum problem defining $g_{hom}(1)$ we have

$$E_1^{(1)}(u, TQ) \geq 2(1-k)\#I_0(u, TQ) + o(1)$$

Then

$$g_{hom}(1) \geq \lim_{\delta} \liminf_T \frac{E_1^{(1)}(u, TQ)}{\#I_0(u, TQ)} \geq 2(1-k).$$

Step 2 (upper bounds). In this Step we conclude the proof by showing that the inequalities

$$\begin{aligned} f_{hom}(z, \nu) &\leq \varphi(z, \nu), \\ g_{hom}(1) &\leq 2(1-k). \end{aligned} \tag{5.73}$$

hold true. With a slight abuse of notation, for $\nu \in S^1$ we denote by $u_\nu : \mathbf{R}^2 \rightarrow \{\pm 1\}$ the function defined in (4.61), with -1 and 1 in place of m_1 and m_2 , that is

$$u_\nu(x) := \begin{cases} -1 & \text{if } (x, \nu) > 0 \\ +1 & \text{if } (x, \nu) \leq 0, \end{cases} \tag{5.74}$$

and, for $z \in \mathbf{R}^+$, we set

$$\mu_{z, \nu} := z\mathcal{H}^1 \llcorner S(u_\nu).$$

In order to prove the first inequality in (5.73), by Theorem 4.4 it is enough to construct $u_\varepsilon \in \mathcal{A}_\varepsilon(\Omega; \{0, \pm 1\})$ such that $(u_\varepsilon, \mu(u_\varepsilon)) \rightarrow (u_\nu, \mu_{z, \nu})$ w.r.t. the $\tau_1 \times \tau_2$ -convergence and

$$\limsup_{\varepsilon} E_\varepsilon^{(1)}(u_\varepsilon) \leq \varphi(z, \nu)\mathcal{H}^1(S(u_\nu) \cap \Omega).$$

To this end we find it useful to rewrite φ as

$$\varphi(z, \nu) = \begin{cases} \varphi_1(z, \nu) & \text{if } 0 \leq z < |\nu_1| \wedge |\nu_2| \\ \varphi_2(z, \nu) & \text{if } |\nu_1| \wedge |\nu_2| \leq z \leq |\nu_1| \vee |\nu_2| \\ \varphi_3(z, \nu) & \text{if } z > |\nu_1| \vee |\nu_2|. \end{cases}$$




Figure 4: $u_{z,\nu}$ in the periodicity cell $\{1, 2, \dots, q\} \times \{1, 2, \dots, p\}$ with $q = 5$, $p = 9$ and $z' = 3$.

The construction of u_ε differs in the three cases $0 < z < |\nu_1| \wedge |\nu_2|$, $|\nu_1| \wedge |\nu_2| \leq z \leq |\nu_1| \vee |\nu_2|$ or $z > |\nu_1| \vee |\nu_2|$. Without loss of generality, for simplicity of exposition, we may suppose that $-\nu_1 \geq \nu_2 > 0$. Moreover, by the continuity of $f_{hom}(z, \cdot)$ and $\varphi(z, \cdot)$, by a density argument we may assume that $\frac{-\nu_1}{\nu_2} \in \mathbf{Q}$. Let $p, q \in \mathbf{N}$ be such that $\frac{-\nu_1}{\nu_2} = \frac{p}{q}$. By the continuity of $f_{hom}(\cdot, \nu)$ and $\varphi(\cdot, \nu)$ we may further assume that $z' := z\sqrt{p^2 + q^2} \in \mathbf{Q}$. Hence, by possibly replacing (p, q) by (mp, mq) for some $m \in \mathbf{N}$, we may reduce to the case $z' \in \mathbf{N}$.

Let us set $\bar{\nu} = \frac{\varepsilon_1 + \varepsilon_2}{\sqrt{2}}$ and let $u^0 : \{1, 2, \dots, q\} \times \mathbf{Z} \rightarrow \{\pm 1, 0\}$ be defined as

$$u^0(a) = \begin{cases} 0 & \text{if } a_1 = a_2 \leq q \text{ or } a_1 = q, q < a_2 \leq p \\ u_{\bar{\nu}}(a) & \text{otherwise.} \end{cases}$$

where $a = (a_1, a_2)$.

Case 1: $0 < z < |\nu_1| \wedge |\nu_2|$. By the assumptions on ν_1, ν_2 this case corresponds to $z' < q$. Let $u_{z,\nu} : \mathbf{Z}^2 \rightarrow \{\pm 1, 0\}$ be such that

$$u_{z,\nu}(a + (p, q)) = u_{z,\nu}(a), \quad \forall a \in \mathbf{Z}^2$$

and, on $\{1, 2, \dots, q\} \times \mathbf{Z} \rightarrow \{\pm 1, 0\}$ is defined as (see Figure 4)

$$u_{z,\nu}(a) = \begin{cases} u^0(a) & \text{if } 0 < a_1 \leq z' \\ u_\nu & \text{otherwise.} \end{cases}$$

Let then $u_\varepsilon : \varepsilon\mathbf{Z}^2 \rightarrow \{\pm 1, 0\}$ be such that $u_\varepsilon(a) = u_{z,\nu}(\frac{a}{\varepsilon})$. It holds that $(u_\varepsilon, \mu(u_\varepsilon)) \rightarrow (u_\nu, \mu_{z,\nu})$ w.r.t. the $\tau_1 \times \tau_2$ -convergence. In order to estimate the




Figure 5: $u_{z,\nu}$ in the periodicity cell $\{1, 2, \dots, q\} \times \{1, 2, \dots, p\}$ with $q = 5$, $p = 9$ and $z' = 7$.

energy of u_ε we observe that it concentrates on each rectangle of the type $R_{\varepsilon,j} := (0, \varepsilon q] \times (0, \varepsilon p] + \varepsilon j(p, q)$, $j \in \mathbf{Z}$ where, by periodicity, it takes the constant value

$$E_\varepsilon^{(1)}(u_\varepsilon, R_{\varepsilon,j}) = \varepsilon (4(1-k)z' + 2(p-z') + 2(q-z')) = \varepsilon (-4kz' + 2(p+q)).$$

Then

$$\begin{aligned} E_\varepsilon^{(1)}(u_\varepsilon) &\leq E_\varepsilon^{(1)}(u_\varepsilon, R_{\varepsilon,0})(\#\{j \in \mathbf{Z} : R_{\varepsilon,j} \subset \Omega\} + 2) \\ &\leq \varepsilon (-4kz' + 2(p+q)) \left(\left[\frac{\mathcal{H}^1(S(u_\nu) \cap \Omega)}{\varepsilon \sqrt{p^2 + q^2}} \right] + 2 \right) \end{aligned}$$

Eventually, letting ε tend to 0 we obtain

$$\limsup_\varepsilon E_\varepsilon^{(1)}(u_\varepsilon) \leq \varphi_1(z, \nu) \mathcal{H}^1(S(u_\nu) \cap \Omega).$$

Case 2: $|\nu_1| \wedge |\nu_2| \leq z \leq |\nu_1| \vee |\nu_2|$. This case corresponds to $q \leq z' \leq p$. Let $v_{z,\nu} : \mathbf{Z}^2 \rightarrow \{\pm 1, 0\}$ be such that

$$v_{z,\nu}(a + (p, q)) = v_{z,\nu}(a), \quad \forall a \in \mathbf{Z}^2$$

and, on $\{1, 2, \dots, q\} \times \mathbf{Z} \rightarrow \{\pm 1, 0\}$ is defined as (see Figure 5)

$$v_{z,\nu}(a) = \begin{cases} -1 & \text{if } a_1 = q \text{ and } a_2 > z' \\ u^0 & \text{otherwise.} \end{cases} \quad (5.75)$$

Let then $v_\varepsilon : \varepsilon \mathbf{Z}^2 \rightarrow \{\pm 1, 0\}$ be such that $v_\varepsilon(a) = v_{z,\nu}(\frac{a}{\varepsilon})$. It holds that $(v_\varepsilon, \mu(v_\varepsilon)) \rightarrow (u_\nu, \mu_{z,\nu})$ w.r.t. the $\tau_1 \times \tau_2$ -convergence. By arguing as in Case 1, taking into account that

$$\begin{aligned} E_\varepsilon^{(1)}(v_\varepsilon, R_{\varepsilon,j}) &= \varepsilon (4(1-k)q + 3(1-k)(z' - q) + 2(p - z')) \\ &= \varepsilon ((1-3k)z' + (1-k)q + 2p), \end{aligned}$$

we get

$$E_\varepsilon^{(1)}(u_\varepsilon) \leq \varepsilon ((1-3k)z' + (1-k)q + 2p) \left(\left[\frac{\mathcal{H}^1(S(u_\nu) \cap \Omega)}{\varepsilon \sqrt{p^2 + q^2}} \right] + 2 \right).$$

Eventually, letting ε tend to 0 we obtain

$$\limsup_\varepsilon E_\varepsilon^{(1)}(v_\varepsilon) \leq \varphi_2(z, \nu) \mathcal{H}^1(S(u_\nu) \cap \Omega).$$

Case 3: $z > |\nu_1| \vee |\nu_2|$. This case corresponds to $z' > p$. Let us extend the function u^0 to \mathbf{Z}^2 in such a way that

$$u^0(a + (p, q)) = u^0(a), \quad \forall a \in \mathbf{Z}^2.$$

We now construct $w_{z,\nu} : \mathbf{Z}^2 \rightarrow \{\pm 1, 0\}$ by modifying the function u^0 suitably increasing the numbers of its zeros in order to match the density constraint on the surfactant phase. More precisely we set $z'' := z' - [\frac{z'}{p}]p$ and

$$\begin{aligned} \tilde{I}_0 &:= \bigcup_{m=0}^{[z'/p]} (I_0(u^0) + me_1) \cap \bigcup_{j \in \mathbf{Z}} \{a \in \mathbf{Z}^2 : jp + 1 \leq a_2 \leq jp + z''\} \\ \hat{I}_0 &:= \bigcup_{m=0}^{[z'/p]-1} (I_0(u^0) + me_1) \cap \bigcup_{j \in \mathbf{Z}} \{a \in \mathbf{Z}^2 : jp + z'' + 1 \leq a_2 \leq (j+1)p\} \end{aligned}$$

we define (see Figure 6)

$$w_{z,\nu}(a) = \begin{cases} 0 & \text{if } a \in \tilde{I}_0 \cup \hat{I}_0 \\ u^0 & \text{otherwise.} \end{cases}$$

Let then $w_\varepsilon : \varepsilon \mathbf{Z}^2 \rightarrow \{\pm 1, 0\}$ be such that $w_\varepsilon(a) = w_{z,\nu}(\frac{a}{\varepsilon})$. It holds that $(w_\varepsilon, \mu(w_\varepsilon)) \rightarrow (u_\nu, \mu_{z,\nu})$ w.r.t. the $\tau_1 \times \tau_2$ -convergence. An easy computation shows that, the energy of each stripe $S_{\varepsilon,j} := \mathbf{R} \times (jp, (j+1)p]$, $j \in \mathbf{Z}$, is

$$\begin{aligned} E_\varepsilon^{(1)}(w_\varepsilon, S_{\varepsilon,j}) &= \varepsilon ((1-3k)p + (1-k)q + 2p + 2(1-k)(z' - p)) \\ &= \varepsilon ((1-k)(p+q) + (1-k)z'). \end{aligned}$$




Figure 6: $u_{z,\nu}$ in the periodicity cell $\{1, 2, \dots, q\} \times \{1, 2, \dots, p\}$ with $q = 5$, $p = 9$ and $z' = 20$.

Then

$$\begin{aligned} E_\varepsilon^{(1)}(w_\varepsilon) &\leq E_\varepsilon^{(1)}(w_\varepsilon, S_{\varepsilon,0})(\#\{j \in \mathbf{Z} : S_{\varepsilon,j} \subset \Omega\} + 2) \\ &\leq \varepsilon((1-k)(p+q) + (1-k)z') \left(\left[\frac{\mathcal{H}^1(S(u_\nu) \cap \Omega)}{\varepsilon\sqrt{p^2+q^2}} \right] + 2 \right) \end{aligned}$$

Eventually, letting ε tend to 0 we obtain

$$\limsup_\varepsilon E_\varepsilon^{(1)}(w_\varepsilon) \leq \varphi_3(z, \nu) \mathcal{H}^1(S(u_\nu) \cap \Omega).$$

We now prove the second inequality in (5.73). To this end, by Theorem 4.4, given $x_0 \in \Omega$, it is enough to construct a sequence of functions $u_\varepsilon \in \mathcal{A}_\varepsilon(\Omega; \{0, \pm 1\})$ such that $(u_\varepsilon, \mu(u_\varepsilon)) \rightarrow (1, \delta_{x_0})$ w.r.t. the $\tau_1 \times \tau_2$ -convergence and that

$$\limsup_\varepsilon E_\varepsilon^{(1)}(u_\varepsilon) \leq 2(1-k).$$

We set

$$u_\varepsilon(a) = \begin{cases} 0 & \text{if } a \in \left(x_0 + \left(-\frac{\sqrt{\varepsilon}}{2}, \frac{\sqrt{\varepsilon}}{2} \right)^2 \right) \cap \varepsilon \mathbf{Z}^2 \\ 1 & \text{otherwise.} \end{cases}$$

Let us observe that $\varepsilon \# I_0(u_\varepsilon) = 1 + o(1)$ and that, each surfactant particle whose interactions do not cross the boundary of $x_0 + \left(-\frac{\sqrt{\varepsilon}}{2}, \frac{\sqrt{\varepsilon}}{2} \right)^2$ gives a contribution to the energy which is equal to $2(1-k)$. Moreover, since the number of the remaining surfactants scales as $\frac{1}{\sqrt{\varepsilon}}$, we have

$$E_\varepsilon^{(1)}(u_\varepsilon) = \varepsilon 2(1-k) \# I_0(u_\varepsilon) + o(1) = 2(1-k) + o(1).$$

Letting ε tend to 0 we get the conclusion. \square

References

- [1] E. Acerbi and G. Bouchitté, A general class of phase transition models with weighted interface energy, *Ann. I H. Poincaré-AN* **25** (2008), 1111–1143.
- [2] R. Alicandro, A. Braides and M. Cicalese, Phase and anti-phases boundaries in binary discrete systems: a variational viewpoint, *Netw. Heterog. Media* **1** (1) (2006), 85–107.
- [3] R. Alicandro, A. Braides and M. Cicalese, Continuum limits of discrete thin films with superlinear growth densities, *Calc. Var. Partial Differential Equations* **33** (2008), no. 3, 267–297.
- [4] R. Alicandro and M. Cicalese, Representation result for continuum limits of discrete energies with superlinear growth, *SIAM J. Math. Anal.* **36** (2004), 1–37.
- [5] R. Alicandro e M. Cicalese, Variational analysis of the asymptotics of the XY model, *Arch. Rat. Mech. Anal.* **192** (2009), no. 3, 501–536.
- [6] R. Alicandro, M. Cicalese and A. Gloria, Integral representation of the bulk limit of a general class of energies for bounded and unbounded spin systems, *Nonlinearity* **21** (2008), no. 8, 1881–1910.
- [7] R. Alicandro, M. Cicalese and M. Ponsiglione, Variational equivalence between Ginzburg-Landau, XY spin systems and screw dislocations energies, submitted (available online at www.cvgmt.sns.it).
- [8] L. Ambrosio and A. Braides, Functionals defined on partitions of sets of finite perimeter, I and II, *J. Math. Pures. Appl.*, **69** (1990), 285–305 and 307–333.
- [9] L. Ambrosio, N. Fusco and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford University Press, Oxford, 2000.
- [10] X. Blanc, C. Le Bris, P.L. Lions, From molecular models to continuum mechanics, *Arch. Rational Mech. Anal.* **164** (2002), no. 4, 341–381.
- [11] X. Blanc, C. Le Bris, P.L. Lions, Atomistic to continuum limits for computational materials science, *M2AN Math. Model. Numer. Anal.* **41** (2007), no. 2, 391–426.
- [12] M. Blume, V.J. Emery and R.b. Griffiths, Ising Model for the λ Transition and Phase Separation in $\text{He}^3\text{-He}^4$ Mixtures, *Physical Review A* **4** (3) (1971), 1071–1077.

- [13] A. Braides, *Γ -convergence for Beginners*, Oxford University Press, Oxford, 2002.
- [14] A. Braides, A handbook of Γ -convergence, in *Handbook of Differential Equations. Stationary Partial Differential Equations, Volume 3* (M. Chipot and P. Quittner, eds.) Elsevier 2006.
- [15] A. Braides e M. Cicalese, Surface energies in nonconvex discrete systems, *Math. Models Methods Appl. Sci.* **17**, no. 7 (2007), 985–1037.
- [16] A. Braides, A.J. Lew and M. Ortiz, Effective cohesive behavior of layers of interatomic planes, *Arch. Rational Mech. Anal.* **180** (2006), no. 2, 151–182.
- [17] A. Braides, L. Truskinowsky, Asymptotic expansions by Γ -convergence, *Cont. Mech. Therm.* **20** (2008), 21–62.
- [18] A. Braides and A. Piatnitski, Overall properties of a discrete membrane with randomly distributed defects, *Arch. Ration. Mech. Anal.* **189** (2008), no. 2, 301–323.
- [19] A. Ciach, J.S. Hoye and G. Stell, Microscopic model for microemulsions. I. Ground states properties, *J. Chem. Phys.* **90** (2) (1989), 1214–1221.
- [20] G. Dal Maso, An introduction to Γ -convergence, Progress in Nonlinear Differential Equations and their Applications, 8. Birkhuser Boston, Inc., Boston, MA, 1993.
- [21] Ellis, Richard S., Otto, Peter T. and Touchette, Hugo, Analysis of phase transitions in the mean-field Blume-Emery-Griffiths model, *Ann. Appl. Probab.* **15** (3) (2005) 2203–2254.
- [22] I. Fonseca, M. Morini and V. Slastikov, Surfactants in foam stability: a phase field model, *Arch. Rat. Mech. Anal.* **183** (3) (2007), 411–456.
- [23] G. Gompper and M. Schick, *Self-Assembling Amphiphilic Systems*, Academic Press, London, 1994.
- [24] M. Laradji, H. Guo, M. Grant and M.J. Zuckermann, Phase diagram of a lattice model for ternary mixtures of water, oil, and surfactants, *J. Phys. A* **24** L629 (1991).
- [25] M. Laradji, H. Guo, M. Grant and M.J. Zuckermann, The effect of surfactants on the dynamics of phase separation, *J. Phys. Condens. Matter* **4** 6715 (1992).
- [26] M. J. Rosen, *Surfactants and Interfacial Phenomena*, Wiley, New York, 1989.
- [27] M. Ponsiglione, Elastic energy stored in a crystal induced by screw dislocations: from discrete to continuous, *SIAM J. Math. Anal.* **39** (2007), no. 2, 449–469.

- [28] M. Schick and W. H. Shih, Simple microscopic model of a microemulsion
Phys. Rev. Letters, **59** (11), 1205–1208.
- [29] B. Schmidt, On the passage from atomic to continuum theory for thin films
Arch. Ration. Mech. Anal. **190** (2008), no. 1, 1–55.