

EXISTENCE OF WEAK SOLUTIONS FOR ELLIPTIC SYSTEMS WITH p, q -GROWTH

GIOVANNI CUPINI - FRANCESCO LEONETTI - ELVIRA MASCOLO

ABSTRACT. We consider a non-linear system of m equations in divergence form and a boundary condition:

$$\begin{cases} \sum_{i=1}^n \frac{\partial}{\partial x_i} (A_i^\alpha(x, Du(x))) = 0, & 1 \leq \alpha \leq m, & \text{in } \Omega \\ u = \tilde{u} & & \text{on } \partial\Omega. \end{cases}$$

The functions $A_i^\alpha(x, z)$ are Hölder continuous with respect to x and

$$|z|^p - c_1 \leq \sum_{\alpha=1}^m \sum_{i=1}^n A_i^\alpha(x, z) z_i^\alpha \leq c_2(1 + |z|)^q, \quad 2 \leq p \leq q.$$

We prove the existence of a weak solution u in $(\tilde{u} + W_0^{1,p}(\Omega; \mathbb{R}^m)) \cap W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m)$, provided p and q are close enough and under suitable summability assumptions on the boundary datum \tilde{u} .

1. INTRODUCTION

In this paper we are concerned with the existence and regularity of solutions to the Dirichlet problem associated to a non-linear system of m equations in divergence form

$$\begin{cases} \sum_{i=1}^n \frac{\partial}{\partial x_i} (A_i^\alpha(x, Du(x))) = 0, & 1 \leq \alpha \leq m, & \text{in } \Omega \\ u = \tilde{u} & & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Ω is a bounded open set in \mathbb{R}^n , $n \geq 2$, and \tilde{u} is a vector-valued function in a suitable Sobolev space.

We assume that $A_i^\alpha : \Omega \times \mathbb{R}^{nm} \rightarrow \mathbb{R}$, $1 \leq i \leq n$, $1 \leq \alpha \leq m$, are Carathéodory functions satisfying the following properties:

there exist $2 \leq p \leq q$, $0 < \gamma \leq 1$ and $\nu, M > 0$, such that for every $x, y \in \Omega$ and for every $z, \tilde{z} \in \mathbb{R}^{nm}$:

- (H1) $\nu (|z|^2 + |\tilde{z}|^2)^{\frac{p-2}{2}} |z - \tilde{z}|^2 \leq \sum_{\alpha=1}^m \sum_{i=1}^n [A_i^\alpha(x, z) - A_i^\alpha(x, \tilde{z})] [z_i^\alpha - \tilde{z}_i^\alpha]$,
- (H2) $|A_i^\alpha(x, z)| \leq M(1 + |z|)^{q-1}$,
- (H3) $\sum_{\alpha=1}^m \sum_{i=1}^n |A_i^\alpha(x, z) - A_i^\alpha(y, z)| \leq c|x - y|^\gamma (1 + |z|)^{q-1}$.

2000 *Mathematics Subject Classification*. Primary: 35J25; Secondary: 35J47, 49N60.

Key words and phrases. Existence, regularity, weak, solution, elliptic, system, growth.

Acknowledgement: We would like to thank Dominic Breit for raising a question that led us to the present study. The authors have been supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

Notice that we are in the framework of the p, q -growth, since (H1) and (H2) imply that there exist $c_1, c_2 > 0$ such that

$$\frac{\nu}{2}|z|^p - c_1 \leq \sum_{\alpha=1}^m \sum_{i=1}^n A_i^\alpha(x, z) z_i^\alpha \leq c_2(1 + |z|)^q. \quad (1.2)$$

For $\tilde{u} \in W^{1,q}(\Omega; \mathbb{R}^{nm})$, a *weak solution* u to problem (1.1) is a function u such that

$$u - \tilde{u} \in W_0^{1,p}(\Omega; \mathbb{R}^m) \cap W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m) \quad (1.3)$$

and

$$\int_{\Omega} \sum_{\alpha=1}^m \sum_{i=1}^n A_i^\alpha(x, Du) \varphi_{x_i}^\alpha(x) dx = 0 \quad \forall \varphi \in W^{1,q}(\Omega; \mathbb{R}^m), \text{ supp } \varphi \Subset \Omega. \quad (1.4)$$

If $p = q$, the existence of weak solutions to (1.1) can be established using the theory of coercive, monotone operators, see Leray-Lions [24], Browder [4] and Hartman-Stampacchia [20]. Moreover, by (H1) the solution is unique. Also the regularity issue has been extensively studied, see the monographs [18], [19] and the surveys [28] and [29].

If $p < q$ the above classical existence results cannot be applied due to the lack of coercivity in $W^{1,q}$. Moreover, the request $u \in W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m)$ in the definition of weak solution, needed to have a well defined integral in (1.4), is an additional difficulty. Notice that such a request is *a priori* assumed in some regularity results under the p, q -growth, see for example Leonetti [22], Bildhauer-Fuchs [1] and Cupini-Marcellini-Mascolo [10].

In this paper we prove the *existence* of a weak solution $u \in W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m)$ to (1.1). Moreover, under additional assumptions, we show that u is *locally bounded*.

The existence of weak solutions is proved assuming that the integrability of the boundary datum is high enough. Precisely, our result is the following.

Theorem 1.1. *Assume that (H1)-(H3) hold, with*

$$2 \leq p \leq q < p \frac{n + \gamma}{n}. \quad (1.5)$$

For every $\tilde{u} \in W^{1,p \frac{q-1}{p-1}}(\Omega; \mathbb{R}^m)$ there exists a weak solution u to the Dirichlet problem (1.1), that is u satisfies (1.3) and (1.4). Moreover, $u \in W_{\text{loc}}^{1,s}(\Omega; \mathbb{R}^m)$ for all $1 \leq s < p \frac{n}{n-\gamma}$.

As far as the regularity of solutions is concerned, the obstructions are essentially two: we are dealing with systems (if $m \geq 2$) and under non-standard growth ($p < q$). Indeed, in the vectorial case, even under the standard growth, the everywhere regularity of solutions for systems, or of minimizers of integrals, cannot be expected unless some structure conditions are assigned, and this holds also for the local boundedness, see e.g. the counterexamples by De Giorgi [13] and Sverák-Yan [30].

Since the pioneering paper by Marcellini [26], the theory of regularity in the framework of non-standard growth has been deeply investigated. The results and the contributions to regularity are so many, that it is a hard task to provide a comprehensive overview of the issue. For this, we refer to the survey of Mingione [28] for an accurate and interesting account on this subject. A common feature is that to get regularity results p and q must be not too far apart, as examples of irregular solutions by Giaquinta [17], Marcellini [25], [27] and Hong [21] show. Notice that the rate of Hölder continuity of $A(\cdot, z)$ interacts with the ratio q/p precisely as in (1.5), see Esposito-Leonetti-Mingione [15] and Colombo-Mingione [8] where the same bound appears. This is suggested by an example in [15], where the minimizer of a functional with $q/p > 1 + \gamma/n$ fails to be locally $W^{1,q}$ -regular; see also Fonseca-Malý-Mingione [16]. Moreover, the condition on the distance between the exponents p and q can usually be relaxed if the solutions/minimizers are assumed locally bounded, see e.g.

Choe [7], D'Ottavio [14], Bildhauer-Fuchs [2], Canale et al. [5], Carozza-Kristensen-Passarelli [6], Breit [3], Colombo-Mingione [9] and Cupini-Marcellini-Mascolo [12].

A particular case of systems such that its weak solutions in $W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m)$ are locally bounded is given by

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\sum_{j=1}^n a_{ij}(x, Du) u_{x_j}^\alpha \right) = 0, \quad \alpha = 1, \dots, m. \quad (1.6)$$

This result is a consequence of Theorem 4.1 in [10], see Theorem 5.1 below. See also Leonetti-Mascolo [23] and Cupini-Marcellini-Mascolo [11].

Collecting the existence result Theorem 1.1 and Theorem 5.1, we obtain the existence of a locally bounded weak solution to (1.1) under the structure condition (1.6), see Theorem 5.2. We emphasize that no boundedness condition is imposed on the boundary datum \tilde{u} .

Theorem 5.2 can be applied to the Euler equation of the functionals

$$\int_{\Omega} (|Du(x)|^2 + a(x)|Du(x)|^q) dx$$

and

$$\int_{\Omega} (1 + |Du(x)|^2)^{\frac{p(x)}{2}} dx,$$

as described at the end of paper.

The plan of the paper is briefly described. In Section 2 we give the strategy of the proof of the existence result Theorem 1.1, which relies on an approximation argument, and state useful preliminary results.

The a priori estimates needed to let the approximation work are proved in Section 3.

The conclusion of the proof of Theorem 1.1 is in section 4.

In the last section, we will apply the existence result to a particular class of non-linear systems, see (5.3), and we will prove that for this class there exists a locally bounded weak solution of the corresponding Dirichlet problem, see Theorem 5.2. Applications of our results are given.

2. PRELIMINARY RESULTS

We consider a bounded open set Ω in \mathbb{R}^n , $n \geq 2$, and Carathéodory functions $A_i^\alpha : \Omega \times \mathbb{R}^{nm} \rightarrow \mathbb{R}$, with $i = 1, \dots, n$ and $\alpha = 1, \dots, m$, satisfying (H1), (H2), (H3), with $2 \leq p \leq q$, $0 < \gamma \leq 1$ and $\nu, M > 0$.

Fixed $\tilde{u} \in W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m)$ we want to study the existence and the summability properties of the solutions to the elliptic system (1.1) in a weak sense; precisely, u is a weak solution to (1.1) if $u - \tilde{u} \in W_0^{1,p}(\Omega; \mathbb{R}^m) \cap W_{\text{loc}}^{1,q}(\Omega; \mathbb{R}^m)$ satisfies (1.4).

For the sake of clarity, we report the scheme of the proof of Theorem 1.1.

We approximate the system in (1.1) with the systems

$$\sum_{\alpha=1}^m \sum_{i=1}^n \frac{\partial}{\partial x_i} (A_{\epsilon,i}^\alpha(x, Du)) = 0, \quad 0 < \epsilon < 1, \quad (2.1)$$

where $A_{\epsilon,i}^\alpha : \Omega \times \mathbb{R}^{nm} \rightarrow \mathbb{R}$ is defined as

$$A_{\epsilon,i}^\alpha(z) := A_i^\alpha(z) + \epsilon|z|^{q-2}z_i^\alpha. \quad (2.2)$$

By (1.2), $A_{\epsilon,i}^\alpha(x, z)$ satisfies

$$\sum_{\alpha=1}^m \sum_{i=1}^n A_{\epsilon,i}^\alpha(x, z) z_i^\alpha \geq \epsilon|z|^q - \lambda.$$

Moreover, by (H1) and (H2), we have that

$$|A_{\epsilon,i}^\alpha(x, z)| \leq (M+1)(1+|z|)^{q-1} \quad (2.3)$$

and, since $z \mapsto |z|^{q-2}z$ is a monotone operator,

$$\nu (|z|^2 + |\tilde{z}|^2)^{\frac{p-2}{2}} |z - \tilde{z}|^2 \leq \sum_{\alpha=1}^m \sum_{i=1}^n [A_{\epsilon,i}^\alpha(x, z) - A_{\epsilon,i}^\alpha(x, \tilde{z})] [z_i^\alpha - \tilde{z}_i^\alpha]. \quad (2.4)$$

The classical theory of monotone operators, see e.g. [24], [4] and [20], permits to conclude that there exists (and, by (H1), unique) $u_\epsilon \in \tilde{u} + W_0^{1,q}(\Omega)$ such that

$$\int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) \varphi_{x_i}^\alpha dx = 0 \quad \forall \varphi \in W_0^{1,q}(\Omega; \mathbb{R}^m). \quad (2.5)$$

Then, we prove estimates in Lebesgue spaces for the gradient of the functions u_ϵ , uniform with respect to ϵ : the norms of Du_ϵ are bounded in $L^p(\Omega)$ and in $L^q(\Omega')$, $\Omega' \Subset \Omega$, see Proposition 3.1 and 3.2, respectively.

We conclude by proving that there exists $u_0 \in \tilde{u} + W_0^{1,p}(\Omega)$ such that $u_0 \in W_{\text{loc}}^{1,q}(\Omega)$ and, up to subsequences, $Du_\epsilon \rightarrow Du_0$ strongly in $L_{\text{loc}}^p(\Omega)$. This will imply that $A_{\epsilon,i}^\alpha(x, Du_\epsilon)$ converges, up to subsequences, to $A_i^\alpha(x, Du_0)$ in $L_{\text{loc}}^1(\Omega)$. Passing to the limit in the equation (2.5), with $\varphi \in W^{1,q}(\Omega; \mathbb{R}^m)$ with compact support in Ω , we conclude.

We here recall some known results.

To prove that the L^p norms of Du_ϵ are bounded with respect to ϵ , see Lemma 3.1, we need the following inequality, see also [27].

Lemma 2.1. *Assume that (H1) and (H2) hold. Let $A_{\epsilon,i}^\alpha$ be defined as in (2.2). Then there exists $c > 0$ such that for all $0 < \epsilon \leq 1$, for all $x \in \Omega$ and for all $\xi, \eta \in \mathbb{R}^{nm}$*

$$|\xi|^p \leq c \left\{ \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, \xi) (\xi_i^\alpha - \eta_i^\alpha) + (1 + |\eta|)^{\frac{p(q-1)}{p-1}} \right\}. \quad (2.6)$$

Proof. By (2.4),

$$\begin{aligned} |\xi|^p &\leq 2^{p-1} (|\xi - \eta|^{p-2} |\xi - \eta|^2 + |\eta|^p) \leq c \left((|\xi|^2 + |\eta|^2)^{\frac{p-2}{2}} |\xi - \eta|^2 + |\eta|^p \right) \\ &\leq \tilde{c} \left(\sum_{\alpha,i} (A_{\epsilon,i}^\alpha(x, \xi) - A_{\epsilon,i}^\alpha(x, \eta)) (\xi_i^\alpha - \eta_i^\alpha) + |\eta|^p \right), \quad \forall \xi, \eta \in \mathbb{R}^{nm}. \end{aligned}$$

By (H2) and the Young inequality,

$$\tilde{c} \left| \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, \eta) (\xi_i^\alpha - \eta_i^\alpha) \right| \leq c(M+1)(1+|\eta|)^{q-1} |\xi - \eta| \leq \frac{1}{2} |\xi|^p + c(1+|\eta|)^{\frac{p(q-1)}{p-1}}.$$

Thus, (2.6) follows. \square

To get that the L^q norms of Du_ϵ are locally bounded with respect to ϵ , see Proposition 3.2, we need technical results. For the reader's convenience we list them below.

The first one is based on an iteration argument, see e.g. [19].

Lemma 2.2. *Let $Z(t)$ be a bounded and non-negative function in the interval $[\rho, R]$. Assume that $\rho \leq r < d \leq R$*

$$Z(r) \leq \theta Z(d) + \frac{A}{(d-r)^\gamma} + B$$

with $A, B \geq 0$ and $0 \leq \theta < 1$. Then

$$Z(\rho) \leq c(\gamma) \left[\frac{A}{(R-\rho)^\gamma} + B \right].$$

The remaining results are related to fine properties of the difference quotient; they can be found e.g. in [15]. We recall that, given a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$, then $\tau_{s,h}f(x) = f(x + he_s) - f(x)$, where e_s is the s -th vector of the canonical basis in \mathbb{R}^n .

Lemma 2.3. *If $0 < \rho < R$, $|h| < R - \rho$, $1 \leq t < +\infty$, $s \in \{1, \dots, n\}$, and $G, D_s G \in L^t(B_R)$, then*

$$\int_{B_\rho} |\tau_{s,h}G(x)|^t dx \leq |h|^t \int_{B_R} |D_s G(x)|^t dx.$$

Lemma 2.4. *If $G : \mathbb{R}^n \rightarrow \mathbb{R}^k$, $G \in L^2(B_R)$, $0 < R \leq 1$ and for some $\rho \in (0, R)$, $d \in (0, 1)$, $M > 0$, $\eta : \mathbb{R}^n \rightarrow \mathbb{R}$ with $\eta \in C_c^1(B_{\frac{\rho+R}{2}})$, $0 \leq \eta \leq 1$ in \mathbb{R}^n , $|D\eta| \leq \frac{4}{R-\rho}$ in \mathbb{R}^n , $\eta = 1$ on B_ρ ,*

$$\sum_{s=1}^n \int_{B_R} |\tau_{s,h}G(x)|^2 \eta^2(x) dx \leq M^2 |h|^{2d}$$

for every h with $|h| \leq \frac{R-\rho}{4}$, then $G \in W^{b,2}(B_\rho; \mathbb{R}^k) \cap L^{\frac{2n}{n-2b}}(B_\rho; \mathbb{R}^k)$ for every $b \in (0, d)$ and

$$\|G\|_{L^{\frac{2n}{n-2b}}(B_\rho)} \leq \frac{c}{(R-\rho)^{2b+2d+2}} (M + \|G\|_{L^2(B_R)}),$$

where $c = c(n, k, b, d)$.

Lemma 2.5. *For every $p > 1$ and $G : B_R \rightarrow \mathbb{R}^k$, we have*

$$|\tau_{s,h}((\mu^2 + |G(x)|^2)^{\frac{p-2}{4}} G(x))|^2 \leq c(k, p)(\mu^2 + |G(x)|^2 + |G(x + he_s)|^2)^{\frac{p-2}{2}} |\tau_{s,h}G(x)|^2$$

for every $x \in B_\rho$, with $|h| \leq R - \rho$ and every $s = 1, \dots, n$, where $0 \leq \mu \leq 1$ and the constant $c = c(k, p)$ is independent of μ .

3. UNIFORM ESTIMATES

In this section we study the approximating systems (2.1).

The properties of $A_{\epsilon,i}^\alpha$, $i = 1, \dots, n$, $\alpha = 1, \dots, m$, ensure the existence of a unique function $u_\epsilon \in \tilde{u} + W_0^{1,q}(\Omega)$ such that

$$\int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) \varphi_{x_i}^\alpha dx = 0 \quad \forall \varphi \in W_0^{1,q}(\Omega; \mathbb{R}^m).$$

In the following, c is a positive constant that may vary from line to line.

First we claim that the L^p norms of Du_ϵ are bounded with respect to ϵ .

Proposition 3.1. *Assume that (H1) and (H2) hold and consider $\tilde{u} \in W^{1, \frac{p(q-1)}{p-1}}(\Omega)$.*

For all ϵ , let $u_\epsilon \in \tilde{u} + W_0^{1,q}(\Omega)$ satisfy (2.5). Then there exists a positive constant c_0 , independent of ϵ , such that

$$\int_{\Omega} |Du_\epsilon|^p dx \leq c_0 \int_{\Omega} (1 + |D\tilde{u}|)^{\frac{p(q-1)}{p-1}} dx. \quad (3.1)$$

Moreover, $\|u_\epsilon - \tilde{u}\|_{W_0^{1,p}(\Omega)} \leq c$ with c independent of ϵ .

Proof. By (2.5) with test function $\varphi = u_\epsilon - \tilde{u}$, and using Lemma 2.1, with $\xi = Du_\epsilon(x)$, $\eta = D\tilde{u}(x)$, we get (3.1). Moreover, since

$$\int_{\Omega} |Du_\epsilon - D\tilde{u}|^p dx \leq 2^{p-1} \int_{\Omega} (|Du_\epsilon|^p + |D\tilde{u}|^p) dx,$$

then (3.1) implies the boundedness of the $W_0^{1,p}$ -norm of $u_\epsilon - \tilde{u}$ with respect to ϵ . \square

In the following proposition we claim that the L^q norms of Du_ϵ are locally bounded with respect to ϵ . Precisely, we will prove the stronger result that the L^s -norms of Du_ϵ are locally bounded, with $q < s < p \frac{n}{n-\gamma}$.

Proposition 3.2. *Assume that (H1)–(H3) hold with $q < p \frac{n+\gamma}{n}$, and consider $\tilde{u} \in W^{1, \frac{p(q-1)}{p-1}}(\Omega)$.*

Let $u_\epsilon \in \tilde{u} + W_0^{1,q}(\Omega)$ satisfy (2.5), with $0 < \epsilon < 1$.

For all $s \in [q, p \frac{n}{n-\gamma})$ there exist σ, τ positive and independent of ϵ , such that for all $B_R(x_0) \Subset \Omega$, with $R \leq 1$, and for all $\rho < R$, we have that

$$\int_{B_\rho(x_0)} |Du_\epsilon(x)|^s dx \leq \frac{c_1}{(R-\rho)^\tau} \left(\int_{B_R(x_0)} (1 + |Du_\epsilon(x)|)^p dx \right)^\sigma + c_2 R^n, \quad (3.2)$$

with positive constants c_1, c_2 independent of ϵ .

Proof. From now on, we write u in place of u_ϵ .

Consider $B_R(x_0) \Subset \Omega$, $\rho \leq r < d \leq R \leq 1$, and define $\eta \in C_c^\infty(\Omega)$, $\text{supp } \eta \subseteq B_{\frac{d+r}{2}}(x_0)$, $\eta \equiv 1$ in B_r and $|D\eta| \leq \frac{4}{d-r}$.

Let $\varphi = \tau_{s,-h}(\eta^2 \tau_{s,h} u)$ with $|h| < \frac{d-r}{2}$. Then (2.5) implies

$$\sum_{\alpha,i} \int_{\Omega} \eta^2 \tau_{s,h} (A_{\epsilon,i}^\alpha(x, Du)) \tau_{s,h} u_{x_i}^\alpha dx = - \sum_{\alpha,i} \int_{\Omega} \tau_{s,h} (A_{\epsilon,i}^\alpha(x, Du)) 2\eta \eta_{x_i} \tau_{s,h} u^\alpha dx. \quad (3.3)$$

We now proceed by steps.

STEP 1.

We prove that

$$\int_{B_{\frac{d+r}{2}}} \eta^2 \left| \tau_{s,h} \left((|Du|^2)^{\frac{p-2}{4}} Du \right) \right|^2 dx \leq \frac{c}{d-r} \int_{B_d} (1 + |Du|)^q dx |h|^\gamma$$

where the constant c is independent of ϵ .

Indeed, since

$$\begin{aligned} & \sum_{\alpha,i} \int_{\Omega} \eta^2 \tau_{s,h} (A_{\epsilon,i}^\alpha(x, Du)) \tau_{s,h} u_{x_i}^\alpha dx \\ &= \sum_{\alpha,i} \int_{\Omega} \eta^2 (A_{\epsilon,i}^\alpha(x + he_s, Du(x + he_s)) - A_{\epsilon,i}^\alpha(x, Du(x + he_s))) \tau_{s,h} u_{x_i}^\alpha dx \\ &+ \sum_{\alpha,i} \int_{\Omega} \eta^2 (A_{\epsilon,i}^\alpha(x, Du(x + he_s)) - A_{\epsilon,i}^\alpha(x, Du(x))) \tau_{s,h} u_{x_i}^\alpha dx, \end{aligned}$$

then (3.3) implies

$$\sum_{\alpha,i} \int_{\Omega} \eta^2 (A_{\epsilon,i}^\alpha(x, Du(x + he_s)) - A_{\epsilon,i}^\alpha(x, Du(x))) \tau_{s,h} u_{x_i}^\alpha dx$$

$$\begin{aligned}
&= - \sum_{\alpha, i} \int_{\Omega} \tau_{s, h} (A_{\epsilon, i}^{\alpha}(x, Du)) 2\eta\eta_{x_i} \tau_{s, h} u^{\alpha} dx \\
&\quad - \sum_{\alpha, i} \int_{\Omega} \eta^2 (A_{\epsilon, i}^{\alpha}(x + he_s, Du(x + he_s)) - A_{\epsilon, i}^{\alpha}(x, Du(x + he_s))) \tau_{s, h} u_{x_i}^{\alpha} dx. \tag{3.4}
\end{aligned}$$

By Lemma 2.5,

$$\left| \tau_{s, h} \left((|Du|^2)^{\frac{p-2}{4}} Du \right) \right|^2 \leq c(n, m, p) (|Du(x)|^2 + |Du(x + he_s)|^2)^{\frac{p-2}{2}} |\tau_{s, h} Du(x)|^2.$$

Therefore, using (2.4), there exists a positive constant c , depending on n, m, p and ν , such that

$$\left| \tau_{s, h} \left((|Du|^2)^{\frac{p-2}{4}} Du \right) \right|^2 \leq c \sum_{\alpha, i} [A_{\epsilon, i}^{\alpha}(x, Du(x + he_s)) - A_{\epsilon, i}^{\alpha}(x, Du(x))] \tau_{s, h} u_{x_i}^{\alpha}.$$

This inequality, together with (3.4), implies

$$\begin{aligned}
&\frac{1}{c} \int_{\Omega} \eta^2 \left| \tau_{s, h} \left\{ (|Du|^2)^{\frac{p-2}{4}} Du \right\} \right|^2 dx \leq - \sum_{\alpha, i} \int_{\Omega} \tau_{s, h} (A_{\epsilon, i}^{\alpha}(x, Du)) 2\eta\eta_{x_i} \times \tau_{s, h} u^{\alpha} dx \\
&\quad - \sum_{\alpha, i} \int_{\Omega} \eta^2 (A_{\epsilon, i}^{\alpha}(x + he_s, Du(x + he_s)) - A_{\epsilon, i}^{\alpha}(x, Du(x + he_s))) \tau_{s, h} u_{x_i}^{\alpha} dx =: I_1 + I_2. \tag{3.5}
\end{aligned}$$

Let us estimate I_1 . By (2.3)

$$\begin{aligned}
|I_1| &\leq c \int_{\Omega} \eta |D\eta| (1 + |Du(x)| + |Du(x + he_s)|)^{q-1} |\tau_{s, h} u| dx \\
&\leq \frac{c}{d-r} \left\{ \int_{B_{\frac{r+d}{2}}} (1 + |Du(x)| + |Du(x + he_s)|)^q dx \right\}^{\frac{q-1}{q}} \left\{ \int_{B_{\frac{r+d}{2}}} |\tau_{s, h} u|^q dx \right\}^{\frac{1}{q}}.
\end{aligned}$$

Then, using Lemma 2.3 and taking into account that $|h| < 1$, we get

$$\begin{aligned}
|I_1| &\leq \frac{c}{d-r} \left\{ \int_{B_d} (1 + |Du(x)|)^q dx \right\}^{\frac{q-1}{q}} \left\{ \int_{B_d} |u_{x_s}|^q dx \right\}^{\frac{1}{q}} |h| \\
&\leq \frac{c}{d-r} \int_{B_d} (1 + |Du(x)|)^q dx |h|^{\gamma}. \tag{3.6}
\end{aligned}$$

Let us estimate I_2 . By Hölder continuity of $A(\cdot, z)$ in (H3),

$$\begin{aligned}
|I_2| &\leq c \int_{\Omega} \eta^2 (1 + |Du(x + he_s)|)^{q-1} |\tau_{s, h} Du| dx |h|^{\gamma} \\
&\leq c \int_{B_{\frac{r+d}{2}}} (1 + |Du(x)| + |Du(x + he_s)|)^q dx |h|^{\gamma} \leq c \int_{B_d} (1 + |Du|)^q dx |h|^{\gamma}. \tag{3.7}
\end{aligned}$$

Thus, collecting (3.5), (3.6) and (3.7), the claim follows.

STEP 2.

Since $\frac{q}{p} < \frac{n+\gamma}{n}$, then we can choose $0 < \theta < \gamma$ such that $\frac{q}{p} < \frac{n+\theta}{n}$. Such a choice is obtained when $0 \leq n \left(\frac{q}{p} - 1 \right) < \theta < \gamma$. We claim that $|Du| \in L_{\text{loc}}^{q\delta}(\Omega)$, where $\delta = \frac{p}{q} \frac{n}{n-\theta}$. Notice that $p < q\delta < p \frac{n}{n-\gamma}$.

Indeed, fixed a ball $B_R(x_0) \Subset \Omega$, with $\rho \leq r < d \leq R \leq 1$, by Step 1 and by Lemma 2.4, applied with $b = \frac{\theta}{2}$, we have $|Du|^{\frac{p}{2}} \in L^{\frac{2n}{n-\theta}}(B_r)$. Moreover,

$$\| |Du|^{\frac{p}{2}} \|_{L^{\frac{2n}{n-\theta}}(B_r)} \leq \frac{c}{(d-r)^{\theta+\gamma+2}} \left\{ \left(\frac{1}{d-r} \int_{B_d} (1 + |Du(x)|)^q dx \right)^{\frac{1}{2}} + \left(\int_{B_d} |Du(x)|^p dx \right)^{\frac{1}{2}} \right\}$$

with c independent of ϵ . Therefore, if we set $\alpha = (\theta + \gamma + \frac{5}{2}) \frac{2n}{n-\theta}$ the above inequality gives

$$\int_{B_r} |Du(x)|^{q\delta} dx \leq \frac{\bar{c}}{(d-r)^\alpha} \left(\int_{B_d} (1 + |Du(x)|)^q dx \right)^{\frac{q\delta}{p}}. \quad (3.8)$$

The arbitrariness of r , d and x_0 allows to conclude.

If $q = p$ then we go to Step 5. If $q > p$ we need Steps 3 and 4.

STEP 3.

Since $\frac{q}{p} < \frac{n+\theta}{n}$ then $\frac{\delta q}{p} < \frac{\delta q - p}{q-p}$. In this step we prove that there exist positive constants t, σ, c , all independent of ϵ , satisfying $\frac{\delta q}{p} < t \leq \frac{\delta q - p}{q-p}$, $0 < \sigma < 1$, such that

$$\int_{B_r} |Du(x)|^{q\delta} dx \leq \frac{c}{(d-r)^{\frac{\alpha}{1-\sigma}}} \left(\int_{B_d} (1 + |Du(x)|)^p dx \right)^{\frac{(t-1)\sigma}{1-\sigma}} + \frac{1}{2} \int_{B_d} |Du(x)|^{q\delta} dx + cR^n. \quad (3.9)$$

Indeed, $t \in \left(\frac{\delta q}{p}, \frac{\delta q - p}{q-p} \right]$ implies

$$1 < \frac{q\delta}{p} < t, \quad \frac{t-\delta}{t-1} \leq p. \quad (3.10)$$

By Hölder inequality

$$\begin{aligned} \left(\int_{B_d} (1 + |Du(x)|)^q dx \right)^{\frac{q\delta}{p}} &= \left(\int_{B_d} (1 + |Du(x)|)^{\frac{q\delta}{t}} (1 + |Du(x)|)^{q(1-\frac{\delta}{t})} dx \right)^{\frac{q\delta}{p}} \\ &\leq \left(\int_{B_d} (1 + |Du(x)|)^{q\delta} dx \right)^{\frac{q\delta}{pt}} \left(\int_{B_d} (1 + |Du(x)|)^{q\frac{t-\delta}{t-1}} dx \right)^{\frac{q\delta}{pt}(t-1)}. \end{aligned} \quad (3.11)$$

Thus, (3.8) and (3.11) imply

$$\int_{B_r} |Du(x)|^{q\delta} dx \leq \left(\int_{B_d} (1 + |Du(x)|)^{q\delta} dx \right)^\sigma \frac{\bar{c}}{(d-r)^\alpha} \left(\int_{B_d} (1 + |Du(x)|)^{qb} dx \right)^{(t-1)\sigma},$$

where \bar{c} is the constant in (3.8) and

$$\sigma := \frac{q\delta}{pt}, \quad b := \frac{t-\delta}{t-1}.$$

By Young inequality, we get

$$\int_{B_r} |Du(x)|^{q\delta} dx \leq \frac{c}{(d-r)^{\frac{\alpha}{1-\sigma}}} \left(\int_{B_d} (1 + |Du(x)|)^{qb} dx \right)^{\frac{(t-1)\sigma}{1-\sigma}} + \frac{1}{2^{q\delta}} \int_{B_d} (1 + |Du(x)|)^{q\delta} dx.$$

Notice that by Step 2 and (3.10) the right-hand side is finite. Since $qb \leq p$ and $(|x| + |y|)^{q\delta} \leq 2^{q\delta-1}(|x|^{q\delta} + |y|^{q\delta})$, we get (3.9).

STEP 4.

By using Lemma 2.2 with $Z(r) := \int_{B_r} |Du(x)|^{q\delta} dx$, the inequality (3.9) implies

$$\int_{B_\rho} |Du(x)|^{q\delta} dx \leq \frac{c}{(R-\rho)^{\frac{\alpha}{1-\sigma}}} \left(\int_{B_R} (1 + |Du(x)|)^p dx \right)^{\frac{(t-1)\sigma}{1-\sigma}} + cR^n.$$

STEP 5.

Notice that since θ is any number in $\left(n\left(\frac{q}{p} - 1\right), \gamma\right)$, then $s := q\delta$ can be any number in $\left(\frac{p^2}{2p-q}, p\frac{n}{n-\gamma}\right)$. Note that $q \leq \frac{p^2}{2p-q}$. The estimate (3.2) follows when s belongs to such an interval. For $s \in \left[p, \frac{p^2}{2p-q}\right]$ we use Hölder inequality and we get (3.2) also in this case. \square

4. PROOF OF THEOREM 1.1

We are now ready to prove the existence of a weak solution to (1.1).

Proof of Theorem 1.1. For all $0 < \epsilon < 1$, let us define the operators $A_{\epsilon,i}^\alpha$ as in (2.2) and let $u_\epsilon \in \tilde{u} + W_0^{1,q}(\Omega)$ satisfy

$$\int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) \varphi_{x_i}^\alpha dx = 0 \quad \forall \varphi \in W_0^{1,q}(\Omega), \quad (4.1)$$

as in Section 3.

STEP 1.

We claim that there exists $u_0 \in \tilde{u} + W_0^{1,p}(\Omega; \mathbb{R}^m)$ such that, up to subsequences, $Du_\epsilon \rightarrow Du_0$ strongly in $L_{loc}^p(\Omega)$.

By Proposition 3.1, the $W_0^{1,p}(\Omega)$ norm of $u_\epsilon - \tilde{u}$ is bounded with respect to ϵ . Moreover, the estimate (3.1) and Proposition 3.2 imply that, fixed $\Omega' \Subset \Omega$, Du_ϵ is bounded in $L^q(\Omega'; \mathbb{R}^{nm})$ uniformly with respect to ϵ . Thus, there exists $u_0 \in \tilde{u} + W_0^{1,p}(\Omega; \mathbb{R}^m)$ such that $Du_0 \in L_{loc}^q(\Omega; \mathbb{R}^{nm})$ and, up to subsequences,

$$u_\epsilon - \tilde{u} \rightarrow u_0 - \tilde{u} \text{ weakly in } W_0^{1,p}(\Omega; \mathbb{R}^m) \quad \text{and} \quad Du_\epsilon \rightarrow Du_0 \text{ weakly in } L_{loc}^q(\Omega; \mathbb{R}^{nm}).$$

Since $q < p\frac{n+\gamma}{n}$, by Rellich Theorem we get

$$u_\epsilon \rightarrow u_0 \text{ in } L_{loc}^q(\Omega; \mathbb{R}^m).$$

To conclude, consider $\eta \in C_c^\infty(\Omega)$ and define the test function $\varphi = (u_\epsilon - u_0)\eta$. Notice that $\varphi \in W^{1,q}(\Omega; \mathbb{R}^{nm})$, with $\text{supp } \varphi \Subset \Omega$. By (4.1)

$$\begin{aligned} \int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) (u_{\epsilon,x_i}^\alpha - u_{0,x_i}^\alpha) \eta dx &= - \int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) (u_\epsilon^\alpha - u_0^\alpha) \eta_{x_i}^\alpha dx \\ &\leq (M+1) \int_{\Omega} (1 + |Du_\epsilon|)^{q-1} |u_\epsilon - u_0| |D\eta| dx. \end{aligned} \quad (4.2)$$

Notice that (2.4) implies

$$\begin{aligned} |Du_\epsilon - Du_0|^p &= (|Du_\epsilon - Du_0|^2)^{\frac{p-2}{2}} |Du_\epsilon - Du_0|^2 \leq 2^{\frac{p-2}{2}} (|Du_\epsilon|^2 + |Du_0|^2)^{\frac{p-2}{2}} |Du_\epsilon - Du_0|^2 \\ &\leq \frac{2^{\frac{p-2}{2}}}{\nu} \sum_{\alpha,i} (A_{\epsilon,i}^\alpha(x, Du_\epsilon) - A_{\epsilon,i}^\alpha(x, Du_0)) (u_{\epsilon,x_i}^\alpha - u_{0,x_i}^\alpha). \end{aligned}$$

Thus, (4.2) and the definition of $A_{\epsilon,i}^\alpha$ imply

$$\begin{aligned} 2^{\frac{2-p}{2}} \nu \int_{\Omega} |Du_\epsilon - Du_0|^p \eta \, dx &\leq \int_{\Omega} \sum_{\alpha,i} (A_{\epsilon,i}^\alpha(x, Du_\epsilon) - A_{\epsilon,i}^\alpha(x, Du_0))(u_{\epsilon,x_i}^\alpha - u_{0,x_i}^\alpha) \eta \, dx \\ &\leq (M+1) \int_{\Omega} (1 + |Du_\epsilon|)^{q-1} |u_\epsilon - u_0| |D\eta| \, dx - \int_{\Omega} \sum_{\alpha,i} A_i^\alpha(x, Du_0)(u_{\epsilon,x_i}^\alpha - u_{0,x_i}^\alpha) \eta \, dx \\ &\quad - \epsilon \int_{\Omega} \sum_{\alpha,i} |u_{0,x_i}^\alpha|^{q-2} u_{0,x_i}^\alpha (u_{\epsilon,x_i}^\alpha - u_{0,x_i}^\alpha) \eta \, dx. \end{aligned}$$

Thus, passing to the limit, as ϵ goes to 0 we get that the right hand side goes to 0. We have so proved that

$$Du_\epsilon \rightarrow Du_0 \quad \text{in } L_{\text{loc}}^p(\Omega). \quad (4.3)$$

STEP 2.

We claim that, up to subsequences, $A_{\epsilon,i}^\alpha(x, Du_\epsilon)$ converges to $A_i^\alpha(x, Du_0)$ in $L_{\text{loc}}^1(\Omega)$. Indeed, for all $\Omega' \Subset \Omega$, adding and subtracting $A_i^\alpha(x, Du_\epsilon)$, we get

$$\int_{\Omega'} |A_{\epsilon,i}^\alpha(x, Du_\epsilon) - A_i^\alpha(x, Du_0)| \, dx \leq \epsilon \int_{\Omega'} |Du_\epsilon|^{q-1} \, dx + \int_{\Omega'} |A_i^\alpha(x, Du_\epsilon) - A_i^\alpha(x, Du_0)| \, dx.$$

By the uniform boundedness of the $L^q(\Omega')$ -norms of Du_ϵ the first term at the right hand side goes to zero. As far as the last term is concerned, define the functions $f_\epsilon(x) := |A_i^\alpha(x, Du_\epsilon) - A_i^\alpha(x, Du_0)|$, $g_\epsilon(x) := M \{(1 + |Du_\epsilon(x)|)^{q-1} + (1 + |Du_0(x)|)^{q-1}\}$ and $g_0(x) := 2M(1 + |Du_0(x)|)^{q-1}$.

By (H2) we get that $f_\epsilon(x) \leq g_\epsilon(x)$ for a.e. $x \in \Omega$. Moreover, by what previously proved, $g_\epsilon \rightarrow g_0$ almost everywhere. Let us now prove that $\int_{\Omega'} g_\epsilon \, dx \rightarrow \int_{\Omega'} g_0 \, dx$ as ϵ goes to 0. This easily follows by (4.3) and recalling that $q \leq p+1$. A generalization of the Lebesgue Convergence Theorem allows to conclude.

STEP 3.

By (4.1) we get that

$$\int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) \varphi_{x_i}^\alpha \, dx = 0 \quad \forall \varphi \in W^{1,q}(\Omega; \mathbb{R}^m), \text{ supp } \varphi \Subset \Omega.$$

By Step 2, up to subsequences,

$$\int_{\Omega} \sum_{\alpha,i} A_i^\alpha(x, Du) \varphi_{x_i}^\alpha \, dx = \lim_{\epsilon \rightarrow 0} \int_{\Omega} \sum_{\alpha,i} A_{\epsilon,i}^\alpha(x, Du_\epsilon) \varphi_{x_i}^\alpha \, dx = 0 \quad \forall \varphi \in W^{1,q}(\Omega), \text{ supp } \varphi \Subset \Omega.$$

Therefore, u is a weak solution to (1.1). \square

5. LOCAL BOUNDEDNESS OF THE WEAK SOLUTIONS

Let us assume that the functions A_i^α in (1.1) are such that

$$A_i^\alpha(x, z) := \sum_{j=1}^n a_{ij}(x, z) z_j^\alpha, \quad (5.1)$$

where $a_{ij} : \Omega \times \mathbb{R}^{nm} \rightarrow \mathbb{R}$ are Carathéodory functions such that

$$\sum_{i,j=1}^n a_{ij}(x, z) \lambda_i \lambda_j \geq 0 \quad \forall \lambda \in \mathbb{R}^n, \forall z \in \mathbb{R}^{nm} \text{ and a.e. } x \in \Omega. \quad (5.2)$$

Let us consider the corresponding system, that is

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\sum_{j=1}^n a_{ij}(x, Du(x)) u_{x_j}^\alpha(x) \right) = 0, \quad \alpha = 1, \dots, m. \quad (5.3)$$

The following boundedness result is consequence of Theorem 4.1 in [10].

Theorem 5.1. *Assume (5.1), (5.2) and*

$$\sum_{\alpha=1}^m \sum_{i=1}^n [A_i^\alpha(x, z) - A_i^\alpha(x, \tilde{z})] [z_i^\alpha - \tilde{z}_i^\alpha] \geq 0 \quad \forall z, \tilde{z} \in \mathbb{R}^{nm} \text{ and a.e. } x \in \Omega. \quad (5.4)$$

Assume also

$$\nu |z|^p - c_1 \leq \sum_{\alpha=1}^m \sum_{i=1}^n A_i^\alpha(x, z) z_i^\alpha \leq c_2 (1 + |z|)^q \quad \forall z, \tilde{z} \in \mathbb{R}^{nm} \text{ and a.e. } x \in \Omega,$$

for some $\nu, c_1, c_2 > 0$, with $1 < p \leq q$, and also $q < p \frac{n-1}{n-p}$ if $p < n$.

Let $u \in W_{\text{loc}}^{1,q}(\Omega)$ be a weak solution to (5.3), i.e.,

$$\int_{\Omega} \sum_{\alpha=1}^m \sum_{i=1}^n A_i^\alpha(x, Du) \varphi_{x_i}^\alpha(x) dx = 0, \quad \forall \varphi \in W^{1,q}(\Omega; \mathbb{R}^m), \quad \text{supp } \varphi \Subset \Omega.$$

Then $u \in L_{\text{loc}}^\infty(\Omega; \mathbb{R}^m)$.

Proof. It is easy to check that the proof of Theorem 4.1 in [10] works using (5.2) and (5.4) instead of the following condition assumed there:

$$\sum_{i,j=1}^n a_{ij}(x, z) \lambda_i \lambda_j \geq M_1 \sum_{i=1}^n \lambda_i^2 \left(\sum_{\alpha=1}^m |z_i^\alpha|^2 \right)^{\frac{p-2}{2}},$$

for all $z \in \mathbb{R}^{mn}$, $\lambda \in \mathbb{R}^n$ and with $M_1 > 0$. \square

By this result and Theorem 1.1, we obtain the following existence and regularity theorem.

Theorem 5.2. *Assume that (H1)-(H3) hold, with $2 \leq p \leq q < p \frac{n+\gamma}{n}$. Let A_i^α be as in (5.1), with (a_{ij}) satisfying (5.2).*

For every $\tilde{u} \in W^{1,p \frac{q-1}{p-1}}(\Omega; \mathbb{R}^m)$ there exists a weak solution u to the Dirichlet problem (1.1), such that u is locally bounded.

Proof. The existence follows by Theorem 1.1 and the local boundedness follows by Theorem 5.1. \square

We conclude with two applications of the above result.

Example 5.3. *Let us consider*

$$A_i^\alpha(x, z) := 2z_i^\alpha + a(x)q|z|^{q-2}z_i^\alpha, \quad (5.5)$$

where $2 \leq q$, $a \in C^{0,\gamma}(\bar{\Omega})$, $a(x) \geq 0$, $0 < \gamma \leq 1$. The assumptions of Theorems 1.1 and 5.2 are satisfied with $p = 2$. Therefore, if $q < 2 \frac{n+\gamma}{n}$ and $\tilde{u} \in W^{1,2(q-1)}(\Omega; \mathbb{R}^m)$ then there exists a locally bounded weak solution to the Dirichlet problem (1.1). Notice that (5.5) comes from the Euler equation of the functional

$$\int_{\Omega} (|Du(x)|^2 + a(x)|Du(x)|^q) dx.$$

Example 5.4. *Let us consider*

$$A_i^\alpha(x, z) := p(x)(1 + |z|^2)^{\frac{p(x)-2}{2}} z_i^\alpha. \quad (5.6)$$

with $p \in C^{0,\gamma}(\bar{\Omega})$, $0 < \gamma \leq 1$. Assume that $2 \leq p_1$ and $p_2 < p_1 \frac{n+\gamma}{n}$, where

$$p_1 := \min\{p(x)\} \quad \text{and} \quad p_2 := \max\{p(x)\}.$$

The assumptions of Theorems 1.1 and 5.2 are satisfied for $p = p_1$ and any q with $p_2 < q < p_1 \frac{n+\gamma}{n}$.

Therefore, if $\tilde{u} \in W^{1, \frac{p_1(q-1)}{p_1-1}}(\Omega; \mathbb{R}^m)$ then there exists a locally bounded weak solution to the Dirichlet problem (1.1). Notice that (5.6) comes from the Euler equation of the functional

$$\int_{\Omega} (1 + |Du(x)|^2)^{\frac{p(x)}{2}} dx.$$

REFERENCES

- [1] M. BILDHAUER, M. FUCHS: $C^{1,\alpha}$ -solutions to non-autonomous anisotropic variational problems, *Calc. Var. Partial Differential Equations* 24 (2005) 309340.
- [2] M. BILDHAUER, M. FUCHS: Higher integrability of the gradient for vectorial minimizers of decomposable variational integrals, *Manuscripta Math.* 123 (2007) 269-283.
- [3] D. BREIT: New regularity theorems for non-autonomous variational integrals with (p, q) -growth, *Calc. Var. Partial Differential Equations* 44 (2012) 101-129.
- [4] F. BROWDER: Nonlinear monotone operators and convex sets in Banach spaces, *Bull. Amer. Math. Soc.* 71 (1965) 780-785.
- [5] A. CANALE, A. D’OTTAVIO, F. LEONETTI, M. LONGOBARDI: Differentiability for bounded minimizers of some anisotropic integrals, *J. Math. Anal. Appl.* 253 (2001) 640-650.
- [6] M. CAROZZA, J. KRISTENSEN, A. PASSARELLI DI NAPOLI: Higher differentiability of minimizers of convex variational integrals, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 28 (2011) 395-411.
- [7] H.J. CHOE: Interior behaviour of minimizers for certain functionals with nonstandard growth, *Nonlinear Anal.* 19 (1992) 933-945.
- [8] M. COLOMBO, G. MINGIONE: Regularity for double phase variational problems, *Arch. Rat. Mech. Anal.* doi: 10.1007/s00205-014-0785-2.
- [9] M. COLOMBO, G. MINGIONE: Bounded minimisers of double phase variational integrals, *preprint* (2014).
- [10] G. CUPINI, P. MARCELLINI, E. MASCOLO: Local boundedness of solutions to quasilinear elliptic systems, *Manuscripta Math.* 137 (2012) 287-315.
- [11] G. CUPINI, P. MARCELLINI, E. MASCOLO: Local boundedness of solutions to some anisotropic elliptic systems, *Contemp. Math.* 595 (2013) 169-186.
- [12] G. CUPINI, P. MARCELLINI, E. MASCOLO: Existence and regularity for elliptic equations under p, q -growth, *Adv. Differential Equations* 19 (2014) 693-724.
- [13] E. DE GIORGI: Un esempio di estremali discontinue per un problema variazionale di tipo ellittico, *Boll. Un. Mat. Ital. (4)* 1 (1968) 135-137.
- [14] A. D’OTTAVIO: A remark on the paper: “Some remarks on the regularity of minimizers of integrals with anisotropic growth”, *Comment. Math. Univ. Carolin.* 36 (1995) 489-491.
- [15] L. ESPOSITO, F. LEONETTI, G. MINGIONE: Sharp regularity for functionals with (p, q) growth, *J. Differential Equations* 204 (2004) 5-55.
- [16] I. FONSECA, I. MALÝ, G. MINGIONE: Scalar minimizers with fractal singular sets, *Arch. Ration. Mech. Anal.* 172 (2004) 295-307.
- [17] M. GIAQUINTA: Growth conditions and regularity, a counterexample, *Manuscripta Math.* 59 (1987) 245-248.
- [18] M. GIAQUINTA: *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, Annals of Mathematics Studies, 105, Princeton University Press (1983).
- [19] E. GIUSTI: *Direct methods in the calculus of variations*, World Scientific Publishing Co., Inc., River Edge, NJ (2003).
- [20] P. HARTMAN, G. STAMPACCHIA: On some non-linear elliptic differential-functional equations, *Acta Math.* 115 (1966) 271-310.
- [21] HONG MIN-CHUN: Some remarks on the minimizers of variational integrals with nonstandard growth conditions, *Boll. Un. Mat. Ital. A* 6 (1992) 91-101.

- [22] F. LEONETTI: Weak differentiability for solutions to nonlinear elliptic systems with p, q -growth conditions, *Ann. Mat. Pura Appl.* 162 (1992) 349-366.
- [23] F. LEONETTI, E. MASCOLO: Local boundedness for vector valued minimizers of anisotropic functionals, *Z. Anal. Anwend.* 31 (2012) 357-378.
- [24] J. LERAY, J.L. LIONS: Quelques résultats de Višik sur les problèmes elliptiques nonlinéaires par les méthodes de Minty-Browder, *Bull. Soc. Math. France* 93 (1965) 97-107.
- [25] P. MARCELLINI: Un exemple de solution discontinue d'un problème variationnel dans le cas scalaire, *Preprint 11, Istituto Matematico "U.Dini"*, Università di Firenze, 1987.
- [26] P. MARCELLINI: Regularity of minimizers of integrals in the calculus of variations with nonstandard growth conditions, *Arch. Ration. Mech. Anal.* 105 (1989) 267-284.
- [27] P. MARCELLINI: Regularity and existence of solutions of elliptic equations with p, q -growth conditions, *J. Differential Equations* 90 (1991) 1-30.
- [28] G. MINGIONE: Regularity of minima: an invitation to the dark side of the calculus of variations, *Appl. Math.* 51 (2006) 355-426.
- [29] G. MINGIONE: Singularities of minima: a walk on the wild side of the calculus of variations, *J. Global Optim.* 40 (2008) 209-223.
- [30] V. SVERÁK, X. YAN: Non-Lipschitz minimizers of smooth uniformly convex functionals, *Proc. Natl. Acad. Sci. USA* 99 (2002), no. 24, 15269-15276.

GIOVANNI CUPINI: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA S.DONATO 5, 40126 - BOLOGNA, ITALY

E-mail address: giovanni.cupini@unibo.it

FRANCESCO LEONETTI: DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE, MATEMATICA, UNIVERSITÀ DI L'AQUILA, 67100 - L'AQUILA, ITALY

E-mail address: leonetti@univaq.it

ELVIRA MASCOLO: DIPARTIMENTO DI MATEMATICA E INFORMATICA "U. DINI", UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, 50134 - FIRENZE, ITALY

E-mail address: mascolo@math.unifi.it