

Evolution of spoon-shaped networks

Alessandra Pluda *

11th October 2015

Abstract

We consider a regular embedded network composed by two curves, one of them closed, in a convex and smooth domain Ω . The two curves meet only at one point, forming angles of 120 degrees. The non-closed curve has a fixed end point on $\partial\Omega$. We study the evolution by curvature of this network. We show that the maximal time of existence is finite and depends only on the area enclosed in the initial loop, if the length of the non-closed curve stays bounded from below during the evolution. Moreover, the closed curve shrinks to a point and the network is asymptotically approaching, after dilations and extraction of a subsequence, a Brakke spoon.

Mathematics Subject classification (2010). 53C44 (primary); 53A04, 35K55 (secondary).

1 Introduction

In this paper we study the evolution by curvature of two curves, one of them closed, inside a convex, open, smooth domain Ω . At the initial time the two curves meet only at one point, the 3-point O . At this point O the curves form angles of 120 degrees (Herring condition). The non-closed curve has one fixed end point on the boundary of Ω . We call such a network a *spoon-shaped network*.

The evolution is the formal L^2 gradient flow of the total length, which is the sum of the lengths of the two curves.

The first short time existence result of a problem similar to ours is due to Bronsard and Reitich; the existence of a unique solution for the problem of the evolution by curvature of three curves, meeting at only one point with prescribed angles, with Neumann boundary conditions of orthogonal intersection with $\partial\Omega$ was shown in [3]. If the planar network, with the Herring condition imposed at triple junctions, is initially close to the equilibrium configuration, global solutions of the flow were obtained by Kinderlehrer and Liu in [13].

In [17, 15, 12] it was investigated the long time behavior of the evolution by curvature of a triod, that is, a connected network composed by three curves meeting at a common point with angles of 120 degrees and with fixed end points on the boundary of a convex domain in \mathbb{R}^2 . In [17] the authors proved that at the first singular time, either the curvature blows-up or the length of at least one of the three curves goes to zero. In [15] it is shown that no singularity can arise during the evolution and this completes the analysis: either the inferior limit of the length of at least one of the three curves composing the initial triod goes to zero at some time T , or the evolution exists for all time and the evolving triod tends to the Steiner configuration connecting the three fixed points on the boundary. The authors mention the possibility to generalize several of the results to regular networks.

A precise definition of network is given in [16] but the case of network with loops is not taken into

*Dipartimento di Matematica, Università di Pisa, Largo Bruno Pontecorvo 5, 56127, Pisa, Italy

account.

In this paper, with the same technique as in [15], it is studied the simplest example of a regular network with loops. We call *Brakke spoon* a spoon-shaped network composed by a half-line which meets a closed curve shrinking in a self-similar way during the evolution.

The main result is the following:

Theorem 1.1. *Given a spoon-shaped network Γ_t evolving by curvature in a strictly convex and open set $\Omega \in \mathbb{R}^2$, there exists a finite time T such that either the inferior limit of the length of the non-closed curve goes to zero as $t \rightarrow T$, or there exists a point $x_0 \in \Omega$ such that for a subsequence of rescaled time t_j the associate rescaled network tends to a Brakke spoon as $j \rightarrow \infty$.*

This result can be seen as an analogue of Grayson's theorem for a closed curve, which says that an embedded evolving curve in \mathbb{R}^2 becomes eventually convex without developing singularities and then shrinks to a point. A new proof of this theorem is given in [14].

In Proposition 5.4, we obtain, up to subsequence, as possible limits, sets that satisfy the equation $k + \langle x | \nu \rangle = 0$ for all x , where k is the curvature and ν is the unit normal vector. Satisfying the previous equation, the sets shrink homothetically. The existence and uniqueness of self-similar shrinking solution of a problem similar to ours is proved in [4]: Chen and Guo consider an equivalent system that describes the motion by curvature, but they focus on the evolution of a curve symmetric about the x -axis, and consider the part of the curve in the upper half-plane, which forms fixed angles with the axis. To gain existence of a unique limit set, without self-intersections, with multiplicity one, with curvature not constantly zero, we will apply [4, Theorem 3], as our asymptotic solution is one of the Abresch-Langer curves, symmetric and convex.

Acknowledgments . *The author is warmly grateful to Matteo Novaga for the suggestion of the problem and for his constant support and encouragement. They author is very grateful to the anonymous referees for their valuable comments, which highly improved the paper.*

The author is partially supported by the PRIN 2010-2011 project "Calculus of Variations" and by GNAMPA of INDAM.

2 Definitions and preliminaries

For any C^2 curve $\sigma : [0, 1] \rightarrow \mathbb{R}^2$ we will denote with s its arclength parameter, defined by $s(x) = \int_0^x |\sigma_x(\xi)| d\xi$, with $\tau = \tau(x) = \frac{\sigma_x}{|\sigma_x|}$ its unit tangent vector and with $\nu = \nu(x) = R\tau(x) = R\frac{\sigma_x}{|\sigma_x|}$ its unit normal vector, where $R : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the counterclockwise rotation of $\pi/2$.

The curvature at the point $\sigma(x)$ will be $k = k(x) = \frac{\langle \sigma_{xx} | \nu \rangle}{|\sigma_x|^2} = \langle \partial_s \tau | \nu \rangle = -\langle \partial_s \nu | \tau \rangle$.

Moreover, we set $\underline{k} = k\nu$.

A curve $\sigma : [0, 1] \rightarrow \mathbb{R}^2$ is called regular if $\sigma_x \neq 0$ for all $x \in [0, 1]$.

Definition 2.1. *Let Ω be a smooth, open and convex subset of \mathbb{R}^2 ; a spoon-shaped network Γ is defined as: $\Gamma = \sigma^1([0, 1]) \cup \sigma^2([0, 1])$, where σ^1 and σ^2 are two regular, at least C^2 curves contained in Ω :*

$$\sigma^1 : [0, 1] \rightarrow \overline{\Omega}, \text{ with } \sigma^1(0) = \sigma^1(1),$$

$$\sigma^2 : [0, 1] \rightarrow \overline{\Omega}.$$

This network has only one point on the boundary of Ω , $\sigma^2(1) = P \in \partial\Omega$, which we call the end point of the network.

The two curves intersect each other only at one point $\sigma^1(0) = \sigma^1(1) = \sigma^2(0) = O$, the 3-point of the network, moreover $O \notin \partial\Omega$. At this junction point an angular condition is required, namely the unit

tangent vectors to the curves form angles of 120 degrees, that is, $\tau^1(0) - \tau^1(1) + \tau^2(0) = 0$.

We require that the network is embedded, that means

$$\begin{cases} \sigma^1(x) \neq \sigma^1(y) & \text{for } x, y \in (0, 1), x \neq y, \\ \sigma^2(x) \neq \sigma^2(y) & \text{for } x, y \in (0, 1), x \neq y, \\ \sigma^1(x) \neq \sigma^2(y) & \text{for } x, y \in (0, 1), x \neq y. \end{cases}$$

Remark 2.2. If Ω is not convex, it could happen that during the evolution the network intersects the boundary of Ω in a point different from the end point P . On the other hand, as a consequence of the strong maximum principle for parabolic equations, this cannot happen if Ω is convex.

The length of the curves is $L_i = \int_0^1 |\sigma_x^i(\xi)| d\xi = \int_{\sigma^i} ds$, the global length of the network is $L = L_1 + L_2$ and we call A the area enclosed in the loop.

The definitions given above of unit tangent vector, unit normal vector and curvature vector are the same for time depending curves $\gamma(x, t)$.

We define a spoon-shaped network $\Gamma_t = \gamma^1(x, t) \cup \gamma^2(x, t)$ as a union of two time depending curves $\gamma^1(x, t)$ and $\gamma^2(x, t)$ above, with the same properties of $\sigma^1(x)$ and $\sigma^2(x)$, for all the times at which they are defined.

For a given initial network Γ_0 , we define the following evolution problem:

Definition 2.3. We say that a family of spoon-shaped networks $\Gamma_t = \gamma^1(\cdot, t) \cup \gamma^2(\cdot, t)$ evolves by curvature in the time interval $[0, T]$ if the functions $\gamma^1 : [0, 1] \times [0, T] \rightarrow \overline{\Omega}$ and $\gamma^2 : [0, 1] \times [0, T] \rightarrow \overline{\Omega}$ are of class C^2 in space, C^1 in time and satisfy the following quasilinear parabolic system

$$\begin{cases} \gamma_x^i(x, t) \neq 0 & \text{regularity} \\ \gamma^1(x, t) = \gamma^1(y, t) & \text{iff } x = y \text{ or } x \in \{0, 1\}, y \in \{0, 1\} \\ \gamma^2(x, t) \neq \gamma^2(y, t) & \text{if } x \neq y & \text{simplicity} \\ \gamma^1(x, t) = \gamma^2(y, t) & \text{iff } x \in \{0, 1\}, y \in \{0\} & \text{intersection only at the 3-point} \\ \tau^1(0) - \tau^1(1) + \tau^2(0) = 0 & \text{angles of 120 degrees at the 3-point} \\ \gamma^2(1, t) = P & \text{fixed end point condition} \\ \gamma^i(x, 0) = \sigma^i(x) & \text{initial data} \\ \gamma_t^i(x, t) = \frac{\gamma_{xx}^i(x, t)}{|\gamma_x^i(x, t)|^2} & \text{motion by curvature} \end{cases} \quad (1)$$

for every $x \in [0, 1]$, $t \in [0, T]$ and $i \in \{1, 2\}$.

Remark 2.4. We can set an analogous Neumann problem requiring that the end point of the non-closed curve $\gamma^2(1, t)$ intersects orthogonally the boundary of Ω .

Often we will denote the flow with Γ_t and we will describe Γ_t as a map $F : \Gamma \rightarrow \overline{\Omega}$ from a fixed standard spoon-shaped network Γ in \mathbb{R}^2 . The evolution will be given by a map $F : \Gamma \times [0, T] \rightarrow \overline{\Omega}$, so $\Gamma_t = F(\Gamma, t)$.

Remark 2.5. In the evolution Problem (1) we define the velocity of the point $\gamma^i(x, t)$ as

$$\underline{v}^i(x, t) = \frac{\gamma_{xx}^i(x, t)}{|\gamma_x^i(x, t)|^2}.$$

Notice that $\langle \underline{v}^i, \nu^i \rangle$ is the curvature of the curve γ^i .

Lemma 2.6. *If γ satisfies the equation*

$$\gamma_t = \frac{\gamma_{xx}}{|\gamma_x|^2} = \lambda\tau + k\nu, \quad (2)$$

then the following commutation rule holds,

$$\partial_t \partial_s = \partial_s \partial_t + (k^2 - \lambda_s) \partial_s. \quad (3)$$

Proof. The thesis follows by direct computation, see [17], Lemma 2.6. \square

Lemma 2.7. *The evolution of the length of the curves is given by*

$$\begin{aligned} \frac{\partial L_i}{\partial t} &= - \int_{\gamma^i} (k^i)^2 - \lambda_s^i ds, \\ \frac{\partial L}{\partial t} &= \lambda^1(1, t) - \lambda^1(0, t) - \lambda^2(0, t) - \int_{\Gamma_t} k^2 ds \end{aligned}$$

and the evolution of the area enclosed in the loop is

$$\frac{\partial A}{\partial t} = -\frac{5\pi}{3}. \quad (4)$$

Proof. The evolution of the length of the curves and of the total length of the network can be seen as a particular case of a more general situation, studied in [16] or in [17]. The evolution of the area is an application of the Gauss-Bonnet theorem. \square

Lemma 2.8. *At the 3–point of a smooth spoon-shaped network Γ_t evolving as in Problem (1), there hold:*

$$\lambda^1(0, t) = \frac{k^1(1, t) + k^2(0, t)}{\sqrt{3}}, \quad \lambda^1(1, t) = \frac{k^2(0, t) - k^1(0, t)}{\sqrt{3}}, \quad \lambda^2(0, t) = \frac{-k^1(0, t) - k^1(1, t)}{\sqrt{3}}, \quad (5)$$

$$k^1(0, t) = \frac{-\lambda^1(1, t) - \lambda^2(0, t)}{\sqrt{3}}, \quad k^1(1, t) = \frac{\lambda^1(0, t) - \lambda^2(0, t)}{\sqrt{3}}, \quad k^2(0, t) = \frac{\lambda^1(0, t) + \lambda^1(1, t)}{\sqrt{3}}, \quad (6)$$

which imply

$$\begin{aligned} \lambda^1(0, t) - \lambda^1(1, t) + \lambda^2(0, t) &= 0, \\ k^1(0, t) - k^1(1, t) + k^2(0, t) &= 0. \end{aligned} \quad (7)$$

Hence,

$$\frac{dL}{dt} = \int_{\Gamma_t} k^2 ds. \quad (8)$$

Moreover

$$k_s^1(0, t) + \lambda^1(0, t)k^1(0, t) = k_s^1(1, t) + \lambda^1(1, t)k^1(1, t) = k_s^2(0, t) + \lambda^2(0, t)k^2(0, t). \quad (9)$$

Proof. In the system that describes the Problem (1), at the 3–point it is required the concurrency condition

$$\gamma^1(1, t) = \gamma^1(0, t) = \gamma^2(0, t). \quad (10)$$

If we differentiate in time (10), we obtain

$$\lambda^1(1, t)\tau^1(1, t) + k^1(1, t)\nu^1(1, t) = \lambda^1(0, t)\tau^1(0, t) + k^1(0, t)\nu^1(0, t), \quad (11)$$

$$\lambda^1(0, t)\tau^1(0, t) + k^1(0, t)\nu^1(0, t) = \lambda^2(0, t)\tau^2(0, t) + k^2(0, t)\nu^2(0, t), \quad (12)$$

$$\lambda^2(0, t)\tau^2(0, t) + k^2(0, t)\nu^2(0, t) = \lambda^1(1, t)\tau^1(1, t) + k^1(1, t)\nu^1(1, t). \quad (13)$$

Taking the scalar product of (10) by $\tau^2(0, t)$ and by $\lambda^2(0, t)$, of (11) by $\tau^1(1, t)$ and by $\lambda^1(1, t)$, of (12) by $\tau^1(0, t)$ and $\lambda^1(0, t)$ and using the angular condition $\tau^1(0, t) - \tau^1(1, t) + \tau^2(0, t) = 0$ we get

$$\begin{aligned} -\frac{1}{2}\lambda^1(0, t) - \frac{\sqrt{3}}{2}k^1(0, t) &= \frac{1}{2}\lambda^1(1, t) - \frac{\sqrt{3}}{2}k^1(1, t), \\ \frac{\sqrt{3}}{2}\lambda^1(0, t) - \frac{1}{2}k^1(0, t) &= \frac{\sqrt{3}}{2}\lambda^1(1, t) + \frac{1}{2}k^1(1, t), \\ \frac{1}{2}\lambda^1(1, t) + \frac{\sqrt{3}}{2}k^1(1, t) &= -\frac{1}{2}\lambda^2(0, t) + \frac{\sqrt{3}}{2}k^2(0, t), \\ -\frac{\sqrt{3}}{2}\lambda^1(1, t) + \frac{1}{2}k^1(1, t) &= -\frac{\sqrt{3}}{2}\lambda^2(0, t) - \frac{1}{2}k^2(0, t), \\ \frac{1}{2}\lambda^2(0, t) + \frac{\sqrt{3}}{2}k^2(0, t) &= \frac{1}{2}\lambda^1(0, t) - \frac{\sqrt{3}}{2}k^1(0, t), \\ -\frac{\sqrt{3}}{2}\lambda^2(0, t) + \frac{1}{2}k^2(0, t) &= -\frac{\sqrt{3}}{2}\lambda^1(0, t) + \frac{1}{2}k^1(0, t), \end{aligned}$$

from which we obtain (5) and (6). Summing up the previous conditions we have (7). Differentiating in time the angular condition at the 3–point $\tau^1(0, t) - \tau^1(1, t) + \tau^2(0, t) = 0$, using the commutation rule (3), which implies $\partial_t \tau = (k_s + k\lambda)\nu$, we have (9). \square

2.1 Short time existence

Definition 2.9. We say that for Problem (1) the compatibility conditions (of every order) are satisfied if at the end–point and at the 3–point there hold all the relations on the space derivatives of the functions σ^i obtained differentiating (any number of times) in time the boundary conditions and substitute them with the space derivatives, given by using the evolution equation (2).

Remark 2.10. Explicitly, for Problem (1), the compatibility conditions of order 2 are

$$\frac{\sigma_{xx}^2(1)}{|\sigma_x^2(1)|^2} = 0,$$

and

$$\frac{\sigma_{xx}^1(0)}{|\sigma_x^1(0)|^2} = \frac{\sigma_{xx}^1(1)}{|\sigma_x^1(1)|^2}, \quad \frac{\sigma_{xx}^1(0)}{|\sigma_x^1(0)|^2} = \frac{\sigma_{xx}^2(0)}{|\sigma_x^2(0)|^2}, \quad \frac{\sigma_{xx}^2(0)}{|\sigma_x^2(0)|^2} = \frac{\sigma_{xx}^1(1)}{|\sigma_x^1(1)|^2}.$$

Theorem 2.11. If Γ_0 is an initial $C^{2+2\alpha}$ network (with $0 < \alpha < 1/2$) satisfying the compatibility conditions of order 2, then, there exists a unique solution $\Gamma \in C^{2+2\alpha, 1+\alpha}([0, 1] \times [0, T))$ of Problem (1).

Proof. The proof is based on the results of [3]. In that paper they consider three curves lying in a convex and smooth domain $\Omega \subset \mathbb{R}^2$, intersecting at a point with prescribed angle and meeting the boundary of Ω orthogonally. Linearizing the problem about the initial data and using a fixed point argument, Bronsard and Reitich obtain a $C^{2+2\alpha, 1+\alpha}([0, 1] \times [0, T))$ solution. The proof can be easily adapted to our situation. \square

Theorem 2.12. *For any initial smooth, regular network Γ_0 in a smooth, convex, open set $\Omega \subset \mathbb{R}^2$ there exists a unique smooth solution of Problem (1) in a maximal time interval $[0, T)$.*

Proof. See [17], Theorem 3.1. □

2.2 Integral estimates

Following the line of proof of [17], we use integral estimates to describe the behavior of the evolution of the network. Thanks to the conditions (5), (6), (7) and (9) at the 3–point and to the Dirichlet condition at the end point P , the computation in [17], Section 3, pp.257-263, can be repeated, to obtain:

$$\int_{\Gamma_t} |\partial_s^j k|^2 ds \leq C \int_0^t \left(\int_{\Gamma_\xi} k^2 ds \right)^{2j+3} d\xi + C \left(\int_{\Gamma_t} k^2 ds \right)^{2j+1} + Ct + C,$$

where with $\partial_s^j k^2$ we denote the j -th derivative of k^2 along a curve with respect to the relative arclength parameter and C is a constant depending only on $j \in \mathbb{N}$ and the initial network Γ_0 . Passing from integral to L^∞ estimates by using Gagliardo-Nirenberg interpolation inequalities (see [18], equation (2.2), page 125), we have the following:

Proposition 2.13. *If the lengths of the curves are positively bounded from below and the L^2 norm of k is bounded, uniformly on $[0, T)$, then the curvature of Γ_t and all its space derivatives are uniformly bounded in the same time interval by some constants depending only on the L^2 integrals of the space derivatives of k on the initial network Γ_0 .*

Remark 2.14. If the length L_1 of the closed curve γ^1 is not bounded from below, then the L^2 norm of its curvature k is not bounded. In fact, from previous estimates and using Hölder inequality, we get $\frac{5\pi}{3} = |\int_{\gamma^1} k ds| \leq (L_1)^{\frac{1}{2}} \left(\int_{\gamma^1} k^2 ds \right)^{\frac{1}{2}}$, hence

$$\int_{\gamma^1} k^2 ds \geq \frac{25\pi^2}{9L_1}. \quad (14)$$

Moreover the maximal time T of existence of a smooth flow is finite and

$$T \leq \frac{3A_0}{5\pi}, \quad (15)$$

because, in order to avoid the presence of a singularity, the area $A(t)$ has to stay positive during the smooth evolution, otherwise the L^2 norm of the curvature is not bounded. From (4) we easily obtain (15).

The next theorem follows from Propositions 2.13 and 2.14 reasoning as in [17, Theorem 3.18].

Theorem 2.15. *If $[0, T)$, with $T < +\infty$, is the maximal time interval of existence of a smooth solution Γ_t of Problem (1), then at least one of the following possibilities holds:*

- $\liminf_{t \rightarrow T} L_2(t) = 0$;
- $\limsup_{t \rightarrow T} \int_{\Gamma_t} k^2 ds = +\infty$.

In the whole paper we will require that the length of the non-closed curve γ^2 is positively bounded from below during the flow, to avoid that the non-closed curve shrinks to a point faster than the closed one.

3 An isoperimetric estimate

Given the smooth flow $\Gamma_t = F(\Gamma, t)$ we take two points $p = F(x, t)$ and $q = F(y, t)$ belonging to Γ_t and we call $(\xi_{p,q})_i$ the geodesic curves contained in Γ_t connecting p and q .

Then, if p and q are both points of the closed curve γ^1 , we let A be the area of the region \mathcal{A} in \mathbb{R}^2 enclosed in the loop and $A_{i,p,q}$ the area of the open region $\mathcal{A}_{i,p,q}$ enclosed by the segment $[p, q]$ and one of the geodesic curve $(\xi_{p,q})_i$. (See Fig. 3.3).

When the $\mathcal{A}_{i,p,q}$ is not connected, we let $A_{i,p,q}$ to be the sum of the areas of its connected components. We define $\psi(A_{p,q})$ as

$$\psi(A_{p,q}) = \frac{A}{\pi} \sin\left(\frac{\pi}{A} A_i\right),$$

where A_i is the smallest of possible $A_{i,p,q}$, depending on the choice of $(\xi_{p,q})_i$.

If p and q are not both points of γ^1 , we let again $A_{i,p,q}$ be the area of the open region $\mathcal{A}_{i,p,q}$ enclosed by the segment $[p, q]$ and one of the geodesic curves $(\xi_{p,q})_i$. Like before, when the region $\mathcal{A}_{i,p,q}$ is not connected, we let $A_{i,p,q}$ be the sum of the areas of its connected components. Differently from the previous case, the possible $\mathcal{A}_{i,p,q}$ are not disjoint. We call $A_{p,q}$ the area of the smallest one (See Fig. 3.3).

We consider the function $\Phi_t : \Gamma \times \Gamma \rightarrow \mathbb{R} \cup \{+\infty\}$ defined as

$$\Phi_t(x, y) = \begin{cases} \frac{|p-q|^2}{\psi(A_{p,q})} & \text{if } x \neq y, x, y \text{ are points of } \gamma^1, \\ \frac{|p-q|^2}{A_{p,q}} & \text{if } x \neq y, x, y \text{ are not both points of } \gamma^1, \\ 4\sqrt{3} & \text{if } x \text{ and } y \text{ coincide with the 3-point } O \text{ of } \Gamma, \\ +\infty & \text{if } x = y \neq O, \end{cases}$$

where $p = F(x, t)$ and $q = F(y, t)$.

Since Γ_t is smooth and the 120 degrees condition holds, it is easy to check that Φ_t is a lower semicontinuous function. Hence, by the compactness of Γ , the following infimum is actually a minimum

$$E(t) = \inf_{x, y \in \Gamma} \Phi_t(x, y), \tag{16}$$

for every $t \in [0, T)$. We call $E(t)$ “embeddedness measure”. Similar geometric quantities have already been applied to similar problems in [9], [5] and [11].

The function Φ_t is locally Lipschitz because the numerator is the square of a distance function and until the flow exists the denominator never goes to zero.

Remark 3.1. Following the proof of Theorem 2.1 in [5], we can always find a minimizing pair (p, q) such that A_i is composed only by one connected component.

Remark 3.2. Following the line of proof by Huisken in [11], in the definition of the function Φ_t we distinguish two cases. We introduce the function $\psi(A_{p,q})$ when the two points belong to the closed curve γ^1 , because we want that Φ_t is a smooth function also when A_i is equal to $\frac{A}{2}$. As we can exclude the case that A_i has more than one connected component, as we said in the previous remark, it can happen that $A_1 = A_2$ only when both p and q belong to the closed curve γ^1 .

In the following computation we only consider the case in which p and q are both points of the closed curve γ^1 ; for the other situation, see Section 4 of [17].




Fig. 3.3. In the figure are shown two possible situations: in the first case the point $p \in \gamma^2$ and $q \in \gamma^1$, in the second case p and q are both point of γ^1 .

If the network Γ_t has no self-intersections we have $E(t) > 0$; the converse is clearly also true.

Moreover, $E(t) \leq \Phi_t(0, 0) = 4\sqrt{3}$ always holds, thus when $E(t) > 0$ the two points (p, q) of a minimizing pair (x, y) can coincide if and only if $p = q = O$.

Finally, since the evolution is smooth, it is easy to see that the function $E : [0, T] \rightarrow \mathbb{R}$ is continuous.

Since we are dealing with embedded networks, with a little abuse of notation, we consider the function Φ_t defined on $\Gamma_t \times \Gamma_t$ and we speak of a minimizing pair for the couple of points $(p, q) \in \Gamma_t \times \Gamma_t$ instead of $(x, y) \in \Gamma \times \Gamma$.

Proposition 3.4. *The function $E(t)$ is monotone increasing in every time interval where $0 < E(t) < \frac{5}{3}$.*

Proof. We assume that $0 < E(t) < \frac{5}{3}$. Since $E(t)$ is a locally Lipschitz function, to prove the statement it is then enough to show that $\frac{dE(t)}{dt} > 0$ for every time t at which this derivative exists (which happens almost everywhere in every time interval where $0 < E(t) < \frac{5}{3}$).

We can exclude the case in which p or q coincide with O , the proof goes like the one of Lemma 4.2, in [17].

Fixed a minimizing pair (p, q) at time t , we choose a value $\varepsilon > 0$ smaller than the intrinsic distances of p and q from the 3-point O of Γ_t and between them.

Possibly taking a smaller $\varepsilon > 0$, we fix an arclength coordinate $s \in (-\varepsilon, \varepsilon)$ and a local parametrization $p(s)$ of the curve containing in a neighborhood of $p = p(0)$, with the same orientation of the original one. Let $\eta(s) = |p(s) - q|$ and $A_i(s) = A_{i,p(s),q}$, since

$$E(t) = \min_{s \in (-\varepsilon, \varepsilon)} \frac{\eta^2(s)}{\psi(A_{p(s),q})} = \frac{\eta^2(0)}{\psi(A_{p(s),q}(0))},$$

if we differentiate in s we obtain

$$\frac{d\eta^2(0)}{ds} \psi(A_{p(s),q}(0)) = \frac{d\psi(A_{p(s),q}(0))}{ds} \eta^2(0). \quad (17)$$

As the intersection of the segment $[p, q]$ with the network is transversal, we have an angle $\alpha(p) \in (0, \pi)$ determined by the unit tangent $\tau(p)$ and the vector $q - p$.

We compute

$$\begin{aligned}
\frac{d\eta^2(0)}{ds} &= -2\langle \tau(p) | q - p \rangle = -2|p - q| \cos \alpha(p) \\
\frac{dA(0)}{ds} &= 0 \\
\frac{dA_i(0)}{ds} &= \frac{1}{2}|\tau(p) \wedge (q - p)| = \frac{1}{2}\langle \nu(p) | q - p \rangle = \frac{1}{2}|p - q| \sin \alpha(p) \\
\frac{d\psi(A_{p(s),q}(0))}{ds} &= \frac{dA_i(0)}{ds} \cos\left(\frac{\pi}{A}A_i(0)\right) \\
&= \frac{1}{2}|p - q| \sin \alpha(p) \cos\left(\frac{\pi}{A}A_i(0)\right).
\end{aligned}$$

Putting these derivatives in equation (17) and recalling that $\eta^2(0)/\psi(A_{p(s),q}(0)) = E(t)$, we get

$$\begin{aligned}
\cot \alpha(p) &= -\frac{|p - q|^2}{4\psi(A_{p(s),q}(0))} \cos\left(\frac{\pi}{A}A_i(0)\right) \\
&= -\frac{E(t)}{4} \cos\left(\frac{\pi}{A}A_i(0)\right).
\end{aligned} \tag{18}$$

Since $0 < E(t) < \frac{5}{3} < 4\sqrt{3}$ we get $-\sqrt{3} < \cot \alpha(p) < 0$ which implies

$$\frac{\pi}{2} < \alpha(p) < \frac{5}{6}\pi.$$

The same argument clearly holds for the point q , hence defining $\alpha(q) \in (0, \pi)$ to be the angle determined by the unit tangent $\tau(q)$ and the vector $p - q$, by equation (18) it follows that $\alpha(p) = \alpha(q)$ and we simply write α for both.

We consider now a different variation, moving at the same time the points p and q , in such a way that $\frac{dp(s)}{ds} = \tau(p(s))$ and $\frac{dq(s)}{ds} = \tau(q(s))$.

As above, letting $\eta(s) = |p(s) - q(s)|$ and $A(s) = A_i(p(s),q(s))$, by minimality we have

$$\frac{d\eta^2(0)}{ds}\psi(A_{p(s),q(s)}(0)) = \frac{d\psi(A_{p(s),q(s)}(0))}{ds}\eta^2(0) \quad \text{and} \quad \frac{d^2\eta^2(0)}{ds^2}\psi(A_{p(s),q(s)}(0)) \geq \frac{d^2\psi(A_{p(s),q(s)}(0))}{ds^2}\eta^2(0). \tag{19}$$

Computing as before,

$$\begin{aligned}
\frac{d\eta^2(0)}{ds} &= 2\langle p - q | \tau(p) - \tau(q) \rangle = -4|p - q| \cos \alpha \\
\frac{dA(0)}{ds} &= 0 \\
\frac{dA_i(0)}{ds} &= -\frac{1}{2}\langle p - q | \nu(p) + \nu(q) \rangle = +|p - q| \sin \alpha \\
\frac{d^2\eta^2(0)}{ds^2} &= 2\langle \tau(p) - \tau(q) | \tau(p) - \tau(q) \rangle + 2\langle p - q | k(p)\nu(p) - k(q)\nu(q) \rangle \\
&= 2|\tau(p) - \tau(q)|^2 + 2\langle p - q | k(p)\nu(p) - k(q)\nu(q) \rangle \\
&= 8\cos^2 \alpha + 2\langle p - q | k(p)\nu(p) - k(q)\nu(q) \rangle \\
\frac{d^2A_i(0)}{ds^2} &= -\frac{1}{2}\langle \tau(p) - \tau(q) | \nu(p) + \nu(q) \rangle + \frac{1}{2}\langle p - q | k(p)\tau(p) + k(q)\tau(q) \rangle \\
&= -\frac{1}{2}\langle \tau(p) | \nu(q) \rangle + \frac{1}{2}\langle \tau(q) | \nu(p) \rangle + \frac{1}{2}\langle p - q | k(p)\tau(p) + k(q)\tau(q) \rangle
\end{aligned}$$

$$\begin{aligned}
&= -2 \sin \alpha \cos \alpha - 1/2|p - q|(k(p) - k(q)) \cos \alpha \\
\frac{d^2\psi(A_{p(s),q(s)}(0))}{ds^2} &= \frac{d}{ds} \left\{ \frac{dA_i(s)}{ds} \cos \left(\frac{\pi}{A} A_i(s) \right) \right\} \Big|_{s=0} \\
&= \frac{d^2 A_i(0)}{ds^2} \cos \left(\frac{\pi}{A} A_i(0) \right) \\
&\quad - \frac{\pi}{A} \left(\frac{dA_i(0)}{ds} \right)^2 \sin \left(\frac{\pi}{A} A_i(0) \right) \\
&= \left(-2 \sin \alpha \cos \alpha - \frac{1}{2}|p - q|(k(p) - k(q)) \cos \alpha \right) \cos \left(\frac{\pi}{A} A_i(0) \right) \\
&\quad - \frac{\pi}{A} |p - q|^2 \sin^2 \alpha \sin \left(\frac{\pi}{A} A_i(0) \right).
\end{aligned}$$

Substituting the last two relations in the second inequality of (19), we get

$$\begin{aligned}
&(8 \cos^2 \alpha + 2(p - q | k(p)\nu(p) - k(q)\nu(q)))\psi(A_{p(s),q(s)}(0)) \\
&\geq |p - q|^2 \left\{ \left(-2 \sin \alpha \cos \alpha - \frac{1}{2}|p - q|(k(p) - k(q)) \cos \alpha \right) \cos \left(\frac{\pi}{A} A_i(0) \right) \right. \\
&\quad \left. - \frac{\pi}{A} |p - q|^2 \sin^2 \alpha \sin \left(\frac{\pi}{A} A_i(0) \right) \right\},
\end{aligned}$$

hence, keeping in mind that $\tan \alpha = \frac{-4}{E(t) \cos(\frac{\pi}{A} A_i)}$, we obtain

$$\begin{aligned}
&2\psi(A_{p(s),q(s)}(0))\langle p - q | k(p)\nu(p) - k(q)\nu(q) \rangle \\
&\quad + 1/2|p - q|^3(k(p) - k(q)) \cos \alpha \cos \left(\frac{\pi}{A} A_i(0) \right) \\
&\geq -2 \sin \alpha \cos \alpha |p - q|^2 \cos \left(\frac{\pi}{A} A_i(0) \right) \\
&\quad - 8\psi(A_{p(s),q(s)}(0)) \cos^2 \alpha + |p - q|^4 \sin^2 \alpha \left[-\frac{\pi}{A} \sin \left(\frac{\pi}{A} A_i(0) \right) \right] \\
&= -2\psi(A_{p(s),q(s)}(0)) \cos^2 \alpha \left(\tan \alpha \frac{|p - q|^2}{\psi(A_{p(s),q(s)})} \cos \left(\frac{\pi}{A} A_i(0) \right) + 4 \right) \\
&\quad + |p - q|^4 \sin^2 \alpha \left[-\frac{\pi}{A} \sin \left(\frac{\pi}{A} A_i(0) \right) \right] \\
&= + |p - q|^4 \sin^2 \alpha \left[-\frac{\pi}{A} \sin \left(\frac{\pi}{A} A_i(0) \right) \right].
\end{aligned} \tag{20}$$

We consider now a time t_0 such that the derivative $\frac{dE(t_0)}{dt}$ exists and we compute it with the Hamilton's trick (see [8]), that is,

$$\frac{dE(t_0)}{dt} = \frac{\partial}{\partial t} \Phi_t(p, q) \Big|_{t=t_0}$$

for *any* pair (p, q) such that $p, q \in \Gamma_{t_0}$ and $\frac{|p-q|^2}{\psi(A_{p,q})} = E(t_0)$.

Considering then a minimizing pair (p, q) for Φ_{t_0} with all the previous properties, by minimality, we are free to choose the “motion” of the points $p(t)$, $q(t)$ “inside” the networks Γ_t in computing such partial derivative.

Since locally the networks are moving by curvature and we know that neither p nor q coincides with the

3-point, we can find $\varepsilon > 0$ and two smooth curves $p(t), q(t) \in \Gamma_t$ for every $t \in (t_0 - \varepsilon, t_0 + \varepsilon)$ such that

$$\begin{aligned} p(t_0) = p & \quad \text{and} \quad \frac{dp(t)}{dt} = k(p(t), t) \nu(p(t), t), \\ q(t_0) = q & \quad \text{and} \quad \frac{dq(t)}{dt} = k(q(t), t) \nu(q(t), t). \end{aligned}$$

Then,

$$\frac{dE(t_0)}{dt} = \frac{\partial}{\partial t} \Phi_t(p, q) \Big|_{t=t_0} = \frac{1}{(\psi(A_{p,q}(t)))^2} \left(\psi(A_{p,q}(t)) \frac{d|p(t) - q(t)|^2}{dt} - |p - q|^2 \frac{d\psi(A_{p,q}(t))}{dt} \right) \Big|_{t=t_0}. \quad (21)$$

With a straightforward computation we get the following equalities,

$$\begin{aligned} \frac{d|p(t) - q(t)|^2}{dt} \Big|_{t=t_0} &= 2\langle p - q \mid k(p)\nu(p) - k(q)\nu(q) \rangle \\ \frac{d(A(t))}{dt} \Big|_{t=t_0} &= -\frac{5\pi}{3} \\ \frac{dA_i(t)}{dt} \Big|_{t=t_0} &= \int_{\Gamma_{p,q}} \langle \underline{k}(s) \mid \nu_{\xi_{p,q}} \rangle ds + \frac{1}{2} |p - q| \langle \nu_{[p,q]} \mid k(p)\nu(p) + k(q)\nu(q) \rangle \\ &= 2\alpha - \frac{5\pi}{3} - \frac{1}{2} |p - q| (k(p) - k(q)) \cos \alpha \\ \frac{d\psi(A_{p,q}(t))}{dt} \Big|_{t=t_0} &= \frac{d(A(t))}{dt} \Big|_{t=t_0} \left[\frac{1}{\pi} \sin \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \right] \\ &+ \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \left(\frac{d(A_i(t))}{dt} \Big|_{t=t_0} - \frac{A_i(t_0)}{A(t_0)} \frac{d(A(t))}{dt} \Big|_{t=t_0} \right) \\ &= \frac{d(A(t_0))}{dt} \left[\frac{1}{\pi} \sin \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) - \frac{A_i(t_0)}{A(t_0)} \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \right] \\ &+ \frac{d(A_i(t_0))}{dt} \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \\ &= -\frac{5\pi}{3} \left[\frac{1}{\pi} \sin \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) - \frac{A_i(t_0)}{A(t_0)} \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \right] \\ &+ \left(2\alpha - \frac{5\pi}{3} - \frac{1}{2} |p - q| (k(p) - k(q)) \cos \alpha \right) \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \end{aligned}$$

where we wrote $\nu_{\xi_{p,q}}$ and $\nu_{[p,q]}$ for the exterior unit normals to the region A_i , respectively at the points of the geodesic $\xi_{p,q}$ and of the segment $[p, q]$.

Substituting these derivatives in equation (21) we get

$$\begin{aligned} \frac{dE(t_0)}{dt} &= \frac{2\langle p - q \mid k(p)\nu(p) - k(q)\nu(q) \rangle}{\psi(A_{p,q}(t_0))} - \frac{|p - q|^2}{(\psi(A_{p,q}(t_0)))^2} \left\{ -\frac{5\pi}{3} \left[\frac{1}{\pi} \sin \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) - \frac{A_i(t_0)}{A(t_0)} \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \right] \right. \\ &\quad \left. + \left(2\alpha - \frac{5\pi}{3} - \frac{1}{2} |p - q| (k(p) - k(q)) \cos \alpha \right) \cos \left(\frac{\pi}{A(t_0)} A_i(t_0) \right) \right\} \end{aligned}$$

and, by equation (20),

$$\begin{aligned}
\frac{dE(t_0)}{dt} &\geq -\frac{|p-q|^2}{(\psi(A_{p,q}))^2} \left\{ -\frac{5}{3} \sin\left(\frac{\pi}{A}A_i\right) + \frac{5\pi}{3} \frac{A_i}{A} \cos\left(\frac{\pi}{A}A_i\right) \right. \\
&\quad \left. + \left(2\alpha - \frac{5\pi}{3}\right) \cos\left(\frac{\pi}{A}A_i\right) + \frac{\pi}{A}|p-q|^2 \sin^2(\alpha) \sin\left(\frac{\pi}{A}A_i\right) \right\} \\
&\geq \frac{|p-q|^2}{(\psi(A_{p,q}))^2} \left[\frac{5}{3} \sin\left(\frac{\pi}{A}A_i\right) - \frac{\pi}{A}|p-q|^2 \sin^2(\alpha) \sin\left(\frac{\pi}{A}A_i\right) \right] \\
&= \frac{\pi}{A} \frac{|p-q|^2}{(\psi(A_{p,q}))} \left[\frac{5}{3} - \frac{\pi}{A}|p-q| \sin^2(\alpha) \right] \\
&= \frac{\pi}{A} \frac{|p-q|^2}{(\psi(A_{p,q}))} \left[\frac{5}{3} - E(t) \sin\left(\frac{\pi}{A}A_i\right) \sin^2(\alpha) \right] \\
&\geq \frac{|p-q|^2}{(\psi(A_{p,q}))} \left[\frac{5}{3} - E(t) \right] \geq 0.
\end{aligned}$$

□

Now that we have proven the monotonicity of the quantity $E(t)$ also when the two points p and q belong to a closed curve, taking into account that the computation in [17, Section 4] is still valid for the other situation, the two subsequent theorems follow as in [17] and in [15]:

Proposition 3.5. *If Ω is bounded and strictly convex, there exists a constant $C > 0$ depending only on Γ_0 such that $E(t) > C > 0$ for every $t \in [0, T)$. Hence, the networks Γ_t remain embedded in all the maximal time interval of existence of the flow.*

Lemma 3.6. *If Ω is strictly convex, the function $E(\Gamma)$ defined on the class of C^1 networks without self-intersections (bounded or unbounded and with or without end points or 3-points), is upper semi continuous with respect to the C_{loc}^1 convergence. Moreover, E is dilation and translation invariant. Consequently, every C_{loc}^1 limit Γ_∞ of a sequence of rescaled networks $\tilde{\Gamma}_t$ has no self-intersections, has multiplicity one and satisfies $E(\Gamma_\infty) > C > 0$, where the constant C is given in Proposition 3.5*

4 Monotonicity formula

Now we introduce the Huisken's monotonicity formula, adapted to our situation. For every $x_0 \in \mathbb{R}^2$, we define the *backward heat kernel* relative to (x_0, T) as

$$\rho_{x_0}(x, t) = \frac{e^{-\frac{|x-x_0|^2}{4(T-t)}}}{\sqrt{4\pi(T-t)}},$$

and the *Gaussian density function* $\Theta : \Gamma \times [0, T) \rightarrow \mathbb{R}$ as

$$\Theta(x_0, t) = \int_{\Gamma_t} \rho_{x_0}(x, t) ds.$$

We call $\hat{\Theta}(x_0)$ the limit as $t \rightarrow T$ of $\Theta(x_0, t)$, that exists and is finite (see [15, Proposition 2.12]).

The following results can be obtained with minor modifications from [17, Proposition 6.4, Lemma 6.5].

Proposition 4.1 (Monotonicity Formula). *For every $x_0 \in \mathbb{R}^2$ and $t \in [0, T)$ the following identity holds*

$$\begin{aligned} \frac{d}{dt} \int_{\Gamma_t} \rho_{x_0}(x, t) ds &= - \int_{\Gamma_t} \left| \underline{k} + \frac{(x - x_0)^\perp}{2(T - t)} \right|^2 \rho_{x_0}(x, t) ds \\ &\quad + \left\langle \frac{P - x_0}{2(T - t)} \middle| \tau(1, t) \right\rangle \rho_{x_0}(P, t), \end{aligned} \quad (22)$$

where $\tau(1, t)$ is the unit tangent vector to $\gamma^2(x, t)$ in the point P .

Integrating between t_1 and t_2 with $0 \leq t_1 \leq t_2 < T$ we get

$$\begin{aligned} \int_{t_1}^{t_2} \int_{\Gamma_t} \left| \underline{k} + \frac{(x - x_0)^\perp}{2(T - t)} \right|^2 \rho_{x_0}(x, t) ds dt &= \int_{\Gamma_{t_1}} \rho_{x_0}(x, t_1) ds - \int_{\Gamma_{t_2}} \rho_{x_0}(x, t_2) ds \\ &\quad + \int_{t_1}^{t_2} \left\langle \frac{P - x_0}{2(T - t)} \middle| \tau(1, t) \right\rangle \rho_{x_0}(P, t) dt. \end{aligned} \quad (23)$$

Lemma 4.2. *Setting $|P - x_0| = d$, the following estimate holds*

$$\left| \int_t^T \left\langle \frac{P - x_0}{2(T - \xi)} \middle| \tau(1, \xi) \right\rangle \rho_{x_0}(P, \xi) d\xi \right| \leq \frac{1}{\sqrt{2\pi}} \int_{d/\sqrt{2(T-t)}}^{+\infty} e^{-y^2/2} dy \leq 1/2.$$

Then, for every $x_0 \in \mathbb{R}^2$,

$$\lim_{t \rightarrow T} \int_t^T \left\langle \frac{P - x_0}{2(T - \xi)} \middle| \tau(1, \xi) \right\rangle \rho_{x_0}(P, \xi) d\xi = 0.$$

We rescale now the flow in its maximal time interval $[0, T)$.

Fixed $x_0 \in \mathbb{R}^2$, let $\tilde{F}_{x_0} : \Gamma \times [-1/2 \log T, +\infty) \rightarrow \mathbb{R}^2$ be the map

$$\tilde{F}_{x_0}(p, t) = \frac{F(p, t) - x_0}{\sqrt{2(T - t)}} \quad \mathfrak{t}(t) = -\frac{1}{2} \log(T - t)$$

then, the rescaled networks are given by

$$\tilde{\Gamma}_{x_0, t} = \frac{\Gamma_t - x_0}{\sqrt{2(T - t)}}$$

and they evolve according to the equation

$$\frac{\partial}{\partial \mathfrak{t}} \tilde{F}_{x_0}(p, \mathfrak{t}) = \tilde{\nu}(p, \mathfrak{t}) + \tilde{F}_{x_0}(p, \mathfrak{t})$$

where

$$\tilde{\nu}(p, \mathfrak{t}) = \sqrt{2(T - t(\mathfrak{t}))} \cdot \underline{\nu}(p, t(\mathfrak{t})) = \tilde{\underline{k}} + \tilde{\underline{\lambda}} = \tilde{k}\tilde{\nu} + \tilde{\lambda}\tilde{\tau} \quad \text{and} \quad t(\mathfrak{t}) = T - e^{-2\mathfrak{t}}.$$

We obtained a smooth flow of spoon-shaped networks $\tilde{\Gamma}_t$ defined for $\mathfrak{t} \in [-\frac{1}{2} \log T, +\infty)$.

By a straightforward computation ([10], [17, Lemma 6.7]) we have the following rescaled version of the monotonicity formula.

Proposition 4.3 (Rescaled Monotonicity Formula). *Let $x_0 \in \mathbb{R}^2$ and set*

$$\tilde{\rho}(x) = e^{-\frac{|x|^2}{2}}.$$

For every $t \in [-1/2 \log T, +\infty)$ the following identity holds

$$\frac{d}{dt} \int_{\tilde{\Gamma}_{x_0, t}} \tilde{\rho}(x) d\sigma = - \int_{\tilde{\Gamma}_{x_0, t}} |\tilde{k} + x^\perp|^2 \tilde{\rho}(x) d\sigma + \left\langle \tilde{P}_{x_0, t} \mid \tau(1, t(t)) \right\rangle \tilde{\rho}(\tilde{P}_{x_0, t}) \quad (24)$$

where $\tilde{P}_{x_0, t} = \frac{P - x_0}{\sqrt{2(T - t(t))}}$.

Integrating between t_1 and t_2 with $-1/2 \log T \leq t_1 \leq t_2 < +\infty$ we get

$$\begin{aligned} \int_{t_1}^{t_2} \int_{\tilde{\Gamma}_{x_0, t}} |\tilde{k} + x^\perp|^2 \tilde{\rho}(x) d\sigma dt &= \int_{\tilde{\Gamma}_{x_0, t_1}} \tilde{\rho}(x) d\sigma - \int_{\tilde{\Gamma}_{x_0, t_2}} \tilde{\rho}(x) d\sigma \\ &\quad \int_{t_1}^{t_2} \left\langle \tilde{P}_{x_0, t} \mid \tau(1, t(t)) \right\rangle \tilde{\rho}(\tilde{P}_{x_0, t}) dt. \end{aligned} \quad (25)$$

Consequently, we have the analogue of Lemma 4.2, whose proof follows in the same way, substituting the rescaled quantities.

Lemma 4.4. *The following estimate holds*

$$\left| \int_t^{+\infty} \left\langle \tilde{P}_{x_0, \xi} \mid \tau(1, t(\xi)) \right\rangle \tilde{\rho}(\tilde{P}_{x_0, \xi}) d\xi \right| \leq \sqrt{\pi/2}.$$

Then, for every $x_0 \in \mathbb{R}^2$,

$$\lim_{t \rightarrow +\infty} \int_t^{+\infty} \left\langle \tilde{P}_{x_0, \xi} \mid \tau(1, t(\xi)) \right\rangle \tilde{\rho}(\tilde{P}_{x_0, \xi}) d\xi = 0.$$

5 Proof of Theorem 1.1

Definition 5.1. *We call infinite flat triod a connected planar set composed by three half-lines intersecting in the origin of \mathbb{R}^2 , forming angles of 120 degrees.*

Definition 5.2. *We call Brakke spoon a spoon-shaped network which shrinks in a self-similar way during the evolution.*




Fig. 5.3. A Brakke spoon.

A Brakke spoon is composed by a half-line which intersects a closed curve, forming angles of 120 degrees and satisfying the equation $k + \langle x | \nu \rangle = 0$ for all point x of the curve, where k is the curvature and ν the

unit normal vector in the point x . This configuration was mentioned in [2] as an example of evolving network with a loop shrinking down to a point, leaving a half-line that then vanishes instantaneously. If we require that this particular spoon-shaped network is embedded, with multiplicity 1, without self-intersections and with 120 degrees angles at the triple junction, then it is unique, up to translation, rotation and dilation.

Proposition 5.4. *Assume that the length of the non-closed curve $\gamma^2(x, t)$ is uniformly bounded away from zero for $t \in [0, T)$. Then, for every $x_0 \in \mathbb{R}^2$ and for every subset \mathcal{I} of $[-1/2 \log T, +\infty)$ with infinite Lebesgue measure, there exists a sequence of rescaled times $t_j \rightarrow +\infty$, with $t_j \in \mathcal{I}$, such that the sequence of rescaled networks $\tilde{\Gamma}_{x_0, t_j}$ converges in the C_{loc}^1 topology to a limit set Γ_∞ which, if not empty, is one of the following:*

- a half-line from the origin with multiplicity one (in this case $\widehat{\Theta}(x_0) = 1/2$).
- a straight line through the origin with multiplicity one (in this case $\widehat{\Theta}(x_0) = 1$);
- an infinite flat triod centered at the origin with multiplicity one (in this case $\widehat{\Theta}(x_0) = 3/2$);
- a Brakke spoon with multiplicity one (in this case $\widehat{\Theta}(x_0) > 3/2$).

Moreover, for every sequence of rescaled networks $\tilde{\Gamma}_{x_0, t_j}$ converging at least in the C_{loc}^1 topology to a limit Γ_∞ , as $t_j \rightarrow +\infty$, we have

$$\lim_{j \rightarrow \infty} \frac{1}{\sqrt{2\pi}} \int_{\tilde{\Gamma}_{x_0, t_j}} \tilde{\rho} d\sigma = \frac{1}{\sqrt{2\pi}} \int_{\Gamma_\infty} \tilde{\rho} d\sigma = \widehat{\Theta}(x_0). \quad (26)$$

Proof. We assume that we have a sequence of rescaled networks $\tilde{\Gamma}_{x_0, t_j}$ that converges to a limit Γ_∞ as $j \rightarrow \infty$ in the C_{loc}^1 topology. As in the proof of Proposition 2.19 in [15], we can pass to the limit in the following Gaussian integral: $\lim_{j \rightarrow \infty} \frac{1}{\sqrt{2\pi}} \int_{\tilde{\Gamma}_{x_0, t_j}} \tilde{\rho} d\sigma = \frac{1}{\sqrt{2\pi}} \int_{\Gamma_\infty} \tilde{\rho} d\sigma$. Consequently,

$$\frac{1}{\sqrt{2\pi}} \int_{\tilde{\Gamma}_{x_0, t_j}} \tilde{\rho} d\sigma = \int_{\Gamma_{t(t_j)}} \rho_{x_0}(x, t(t_j)) ds = \Theta(x_0, t(t_j)) \rightarrow \widehat{\Theta}(x_0),$$

as $j \rightarrow \infty$ and equality (26) follows.

If in the rescaled monotonicity formula (25) we set $t_1 = -\frac{1}{2} \log T$ and we let t_2 go to $+\infty$, considering the result in Lemma 4.4, we get $\int_{-1/2 \log T}^{+\infty} \int_{\tilde{\Gamma}_{x_0, t}} |\tilde{k} + \langle x | \nu \rangle|^2 \tilde{\rho} d\sigma dt < +\infty$, then $\int_{\mathcal{I}} \int_{\tilde{\Gamma}_{x_0, t}} |\tilde{k} + \langle x | \nu \rangle|^2 \tilde{\rho} d\sigma dt < +\infty$. Being the last integral finite and being the integrand a nonnegative function on a set of infinite Lebesgue measure, we can extract within \mathcal{I} a sequence of time $t_j \rightarrow +\infty$ such that

$$\lim_{j \rightarrow +\infty} \int_{\tilde{\Gamma}_{x_0, t_j}} |\tilde{k} + \langle x | \nu \rangle|^2 \tilde{\rho} d\sigma = 0$$

It follows that, for every ball of radius R in \mathbb{R}^2 , the networks $\tilde{\Gamma}_{x_0, t_j}$ have curvature uniformly bounded in $L^2(B_R)$.

Hence, we can extract a subsequence $\tilde{\Gamma}_{x_0, t_j}$ (not relabelled) which, after a possible reparametrization, converges to a limit (possibly empty) network Γ_∞ in the C_{loc}^1 topology.

We will call Γ_∞ a blow-up limit of Γ_t around the point x_0 (notice that the blow-up limit could depend on the choice of the subsequence).

If the limit set is not empty, it has multiplicity one thanks to Lemma 3.6. Also the embeddedness is granted by Proposition 3.5.

This limit set satisfies the equation

$$k_\infty + \langle x|\nu \rangle = 0 \text{ for all } x \in \Gamma_\infty, \quad (27)$$

where k_∞ is the curvature of Γ_∞ at x , because the integral $\int_{\tilde{\Gamma}} |\tilde{k} + \langle x|\nu \rangle|^2 \tilde{\rho} d\sigma$ is lower semicontinuous under the C_{loc}^1 convergence.

In principle, the limit set is composed of curves in $W_{loc}^{2,2}$, but from the relation $k_\infty + \langle x|\nu \rangle = 0$ it follows that k_∞ is continuous, since the curves are in C_{loc}^1 . By a bootstrap argument we gain the smoothness of Γ_∞ .

Now we classify all the possible limit sets that satisfy (27).

By the work of Abresch and Langer [1] the curves componing Γ_∞ can be only lines, half-lines, segments or pieces of curves of Abresch and Langer. If k is identically zero we can only have a straight line, a half-line or an infinite flat triod. If the point $x_0 \in \mathbb{R}^2$ is not the end point P , then Γ_∞ has no end points, otherwise if the point $x_0 \in \mathbb{R}^2$ is the end-point P , the set Γ_∞ is a half-line by the bound from below on the curve γ^2 .

Otherwise, if the curvature is not constantly zero, it is proved in [4, Theorem 3] that there exists a unique possible limit set without self-intersection and with multiplicity one and it is the Brakke spoon in Definition 5.2.

The value of gaussian density $\widehat{\Theta}(x_0)$ in the first three cases can be obtained by a direct computation (see [15]), for the last case see Lemma 8.4 in [12].

□

We define the set of reachable points of the flow by

$$R = \left\{ x \in \mathbb{R}^2 \mid \text{there exist } p_i \in \Gamma \text{ and } t_i \nearrow T \text{ such that } \lim_{i \rightarrow \infty} F(p_i, t_i) = x \right\}.$$

This set R is not empty and compact, if a point $x_0 \notin R$, it means that the flow is definitively far from x_0 , otherwise if $x_0 \in R$, for every $t \in [0, T)$ the closed ball of radius $\sqrt{2(T-t)}$ and center x_0 intersects Γ_t (for the proof see [14]). Hence we have two possibilities when we consider the blow-up around points of $\bar{\Omega}$:

- we are rescaling around a point in R and the limit of any sequence of rescaled network is not empty;
- the blow-up limit is empty.

Fixed $x_0 \in R$, repeating the previous arguments, there exists a not relabeled subsequence $t_j \rightarrow +\infty$ of rescaled times such that the rescaled networks $\tilde{\Gamma}_{x_0, t_j}$ converge in the C_{loc}^1 topology to a nonempty limit: a straight line, a half-line, an infinite flat triod or a Brakke spoon.

Proposition 5.5. *If the sequence of rescaled networks converges to a straight line or to a half-line, then the curvature of the evolving network is uniformly bounded for $t \in [0, T)$ in a ball around the point x_0 . If the sequence of rescaled networks converges to a infinite flat triod, then the full L^2 norm of the curvature of the evolving network is bounded.*

Proof. See [15], Propositions 3.2 and 3.3 and the proof of Theorem 2.4. □

Proof of Theorem 1.1. As we have imposed that the length of the non-closed curve γ^2 stays bounded away from zero during the evolution, at the time T of maximal existence of a smooth flow, only the second case of Theorem 2.15 can happen, that is $\limsup_{t \rightarrow T} \int_{\Gamma_t} k^2 ds = +\infty$. By Proposition 5.4 we know that, for every $x_0 \in \mathbb{R}^2$, there exists a sequence of rescaled times such that the sequence of

corresponding rescaled networks converges to a limit set Γ_∞ , which, if not empty, is a half-line from the origin, a straight line through the origin, an infinite flat triod centered at the origin or a Brakke spoon. If for every $x_0 \in R$ the limit set Γ_∞ is a half-line, a straight line or an infinite flat triod, then the curvature is uniformly bounded as $t \rightarrow T$ thanks to the previous proposition. This is in contradiction with Theorem 2.15. Hence, it exists a point $x_0 \in R$ such that the limit set Γ_∞ is a Brakke spoon and at x_0 the gaussian density is $\widehat{\Theta}(x_0) > 3/2$. This proves the main Theorem. \square

Corollary 5.6. *If the length of the non-closed curve γ^2 is positively bounded from below, at the maximal time of existence $T < +\infty$ of a smooth solution $\lim_{t \rightarrow T} A(t) = 0$, $\liminf_{t \rightarrow T} L_1(t) = 0$ and $\limsup_{t \rightarrow T} \int_{\gamma^1} k^2 ds = +\infty$. Moreover $T = \frac{3A_0}{5\pi}$ where A_0 is the initial area of the region contained in the closed curve γ^1 .*

Proof. From the equation (4) of the evolution of the area enclosed in the loop, we know that $A(t)$ is decreasing. If the length of the non-closed curve γ^2 stays positively bounded from below during all the smooth evolution of the flow and we know that at time $T < +\infty$ it appears a singularity, then from Theorem 1.1, the rescaled flow converges, up to subsequence, to a Brakke spoon. This implies that the area in the loop goes to zero and $\liminf_{t \rightarrow T} L_1(t) = 0$. Recalling the inequality (14) we also get that $\limsup_{t \rightarrow T} \int_{\gamma^1} k^2 ds = +\infty$. Moreover, thanks to the equation $\frac{\partial A}{\partial t} = \frac{-5\pi}{3}$, the time in which such singularity appears is given by

$$T = \frac{3A_0}{5\pi}. \quad (28)$$

\square

References

- [1] U. ABRESCH AND J. LANGER, *The normalized curve shortening flow and homothetic solutions*, J. Diff. Geom. **23** (1986), 175-196.
- [2] K .A. BRAKKE, *The Motion of a Surface by its Mean Curvature*, Princeton University Press, Princeton, NJ , 1978.
- [3] L. BRONSARD AND F. REITICH, *On three-phase boundary motion and the singular limit of a vector-valued Ginzburg-Landau equation*, Arch. Rational Mech. Anal. **124** (1993), 355-379.
- [4] X. CHEN AND J. GUO, *Motion by curvature of planar curves with end points moving freely on a line*, Math. Ann. **350** (2011), 277-311.
- [5] K. S. CHOU AND X. P. ZHU, *Shortening complete planar curves*, J. Diff. Geom. **50** (1998), 471-504.
- [6] M. E. GAGE AND R. S. HAMILTON, *The heat equation shrinking convex plane curves*, J. Diff. Geom. **23** (1986), 69-96.
- [7] M. A. GRAYSON, *The heat equation shrinks embedded plane curves to points*, J. Diff. Geom. **26** (1987), 285-314.
- [8] R. S. HAMILTON, *Four-manifolds with positive curvature operator*, J. Diff. Geom. **24** (1986), 153-179.
- [9] R. S. HAMILTON, *Isoperimetric estimates for the curve shrinking flow in the plane*, Modern Methods in Complex Analysis (Princeton, NJ, 1992), Princeton University Press, Princeton, NJ, 1995, 201-222.
- [10] G. HUISKEN, *Asymptotic behavior for singularities of the mean curvature flow*, J. Diff. Geom. **31** (1990), 285-299.

- [11] G. HUISKEN, *A distance comparison principle for evolving curves*, Asian J. Math. **2** (1998), 127-133.
- [12] T. ILMANEN, A. NEVES AND F. SCHULZE, *On short time existence for the planar network flow*, 2014, arXiv:1407.4756
- [13] D. KINDERLEHRER AND C. LIU, *Evolution of grain boundaries*, Math. Models Methods Appl. Sci. **11** (2001), no. 4,713-729.
- [14] A. MAGNI AND C. MANTEGAZZA, *A note on Grayson's theorem*, Rend. Semin. Mat. Univ. Padova **131** (2014), 263-279.
- [15] A. MAGNI, C. MANTEGAZZA AND M. NOVAGA, *Motion by curvature of planar networks II*, 2013, arXiv:1301.3352v1.
- [16] C. MANTEGAZZA, M. NOVAGA, A. PLUDA AND F. SCHULZE, *Evolution of network with multiple junctions*, Preprint 2015.
- [17] C. MANTEGAZZA, M. NOVAGA AND V. M. TORTORELLI, *Motion by curvature of planar networks*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **(5)3** (2004), 235-324.
- [18] L. NIRENBERG, *On elliptic partial differential equations*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **13** (1959), 116-162.
- [19] V.A. SOLONNIKOV, *Boundary Value Problems in Physics*, Proceeding of the Steklov Institute, Vol. LXXXIII, 1965.