

THE NEHARI MANIFOLD FOR FRACTIONAL SYSTEMS INVOLVING CRITICAL NONLINEARITIES

XIAOMING HE, MARCO SQUASSINA, AND WENMING ZOU

ABSTRACT. We study the combined effect of concave and convex nonlinearities on the number of positive solutions for a fractional system involving critical Sobolev exponents. With the help of the Nehari manifold, we prove that the system admits at least two positive solutions when the pair of parameters (λ, μ) belongs to a suitable subset of \mathbb{R}^2 .

CONTENTS

1.	Introduction	1
2.	Some preliminary facts	3
3.	The Palais-Smale condition	8
4.	The Nehari manifold	10
5.	Existence of Palais-Smale sequences	14
6.	Local minimization problems	17
7.	Proof of the main result concluded	22
	References	22

1. INTRODUCTION

This paper is concerned with the multiplicity of positive solutions for the following elliptic system involving the fractional Laplacian

$$(1.1) \quad \begin{cases} (-\Delta)^s u = \lambda |u|^{q-2} u + \frac{2\alpha}{\alpha+\beta} |u|^{\alpha-2} u |v|^\beta & \text{in } \Omega, \\ (-\Delta)^s v = \mu |v|^{q-2} v + \frac{2\beta}{\alpha+\beta} |u|^\alpha |v|^{\beta-2} v & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, $\lambda, \mu > 0$, $1 < q < 2$ and $\alpha > 1, \beta > 1$ satisfy $\alpha + \beta = 2_s^* = 2N/(N - 2s)$, $s \in (0, 1)$ and $N > 2s$. When $\alpha = \beta, \alpha + \beta = p \leq 2_s^*, \lambda = \mu$ and $u = v$, problem (1.1) reduces to the semilinear scalar fractional elliptic equation

$$(1.2) \quad \begin{cases} (-\Delta)^s u = \lambda |u|^{q-2} u + |u|^{p-2} u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Recently, a great attention has been focused on the study of nonlinear problems like (1.2) which involve the fractional Laplacian. This type of operators naturally arises in physical situations such as thin obstacle problems, optimization, population dynamics, geophysical fluid dynamics, mathematical finance, phases transitions, stratified materials, anomalous diffusion, crystal dislocation, soft thin films, semipermeable membranes, flames propagation, conservation laws, ultra-relativistic limits of quantum mechanics, quasi-geostrophic flows,

2000 *Mathematics Subject Classification.* 47G20, 35J50, 35B65.

Key words and phrases. Fractional elliptic system, concave-convex nonlinearities, Nehari manifold.

X. He is supported by NSFC (11371212, 10601063, 11271386) while W. Zou by NSFC (11371212, 11271386).

multiple scattering, minimal surfaces, materials science and water waves, see [21]. We refer to [12, 23, 25, 28, 29, 31, 34] for the subcritical case and to [3, 4, 11, 14, 15, 22, 24, 26, 30] for the critical case. In the remarkable paper [9], Caffarelli and Silvestre gave a new formulation of the fractional Laplacian through Dirichlet-Neumann maps. This is extensively used in the recent literature since it allows to transform nonlocal problems to local ones, which permits to use variational methods. For example, Barrios, Colorado, de Pablo and Sánchez [3] used the idea of the s -harmonic extension and studied the effect of lower order perturbations in the existence of positive solutions of (1.2). Brändle, Colorado and de Pablo [5] investigated the fractional elliptic equation (1.2) involving concave-convex nonlinearity, and obtained an analogue multiplicity result to the problem considered by Ambrosetti, Brézis and Cerami in [2]. In the case $q = 2$ and $p = 2_s^*$, Servadei and Valdinoci [26] studied (1.2) and extended the classical Brézis-Nirenberg result [6] to the nonlocal case. In [8], Cabré and Tan defined $(-\Delta)^{1/2}$ through the spectral decomposition of the Laplacian operator on Ω with zero Dirichlet boundary conditions. With classical local techniques, they established existence of positive solutions for problems with subcritical nonlinearities, regularity and L^∞ -estimates for weak solutions. In particular, Tan [30] considered

$$(1.3) \quad \begin{cases} (-\Delta)^{1/2}u = \lambda u + u^{\frac{N+1}{N-1}} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

investigating the solvability (see also [34] for a subcritical situation). Very recently, Colorado, de Pablo, and Sánchez [14] studied the following nonhomogeneous fractional equation involving critical Sobolev exponent

$$\begin{cases} (-\Delta)^s u = |u|^{2_s^*-2}u + f(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

and proved existence and multiplicity of solutions under appropriate conditions on the size of f . For the same problems, Shang, Zhang and Yang [22] obtained similar results.

The analogue problems to (1.1) for the Laplacian operator have been studied extensively in recent years, see [1, 13, 19, 20, 32, 33] and the references therein. In particular, Hu and Lin [20] studied the Laplacian system with critical growth and obtained the existence and multiplicity results of positive solutions by variational methods.

The purpose of this paper is to study system (1.1) in the critical case $\alpha + \beta = 2_s^*$. Using variational methods and a *Nehari manifold decomposition*, we prove that system (1.1) admits at least two positive solutions when the pair of parameters (λ, μ) belongs to a certain subset of \mathbb{R}^2 . To our best knowledge, there are just a few results in the literature on the fractional system (1.1) with both concave-convex nonlinearities and critical growth terms. We point out that we adopt in the paper the *spectral* (or *regional*) definition of the fractional laplacian in a bounded domain based upon a Caffarelli-Silvestre type extension (see [14]), and not the *integral definition*. We shall refer to [27] for a nice comparison between these two different notions. In [18], a problem like (1.1) with $q = 2$ is investigated, using the integral notion, from the point of view of existence, nonexistence and regularity.

To formulate the main result, we introduce

$$(1.4) \quad \Lambda_1 := \left(\frac{2_s^* - q}{2_s^* - 2} \mathcal{S}(s, N)^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} \right)^{-\frac{2}{2-q}} \left[\frac{2-q}{2(2_s^* - q)} (k_s \mathcal{S}_{\alpha, \beta})^{\frac{2_s^*}{2}} \right]^{\frac{2}{2_s^* - 2}},$$

where $|\Omega|$ is the Lebesgue measure of Ω , k_s is a normalization constant and $\mathcal{S}(s, N), \mathcal{S}_{\alpha, \beta}$ are best Sobolev constants that will be introduced later. For $\gamma > 0$, we also consider

$$\mathcal{C}_\gamma := \{(\lambda, \mu) \in \mathbb{R}_+^2 : 0 < \lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} < \gamma\}.$$

Then we have the following

Theorem 1.1. *The following facts holds*

- (i) *system (1.1) has at least one positive solution for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$.*
- (ii) *there is $\Lambda_2 < \Lambda_1$ such that (1.1) has at least two positive solutions for $(\lambda, \mu) \in \mathcal{C}_{\Lambda_2}$.*

Concerning regularity, one can get a priori estimates for the solutions to (1.1) and hence obtain, as in [3, Proposition 5.2], that $u, v \in C^\infty(\bar{\Omega})$ for $s = 1/2$, $u, v \in C^{0, 2s}(\bar{\Omega})$ if $0 < s < 1/2$ and $u, v \in C^{1, 2s-1}(\bar{\Omega})$ if $1/2 < s < 1$.

The paper is organized as follows. In Section 2 we introduce the variational setting of the problem and present some preliminary results. In Section 3 we show that the Palais-Smale condition holds for the energy functional associated with (1.1) at energy levels in a suitable range related to the best Sobolev constants. In Section 4 we give some properties about the Nehari manifold and fibering maps. In Section 5 we investigate the existence of Palais-Smale sequences. In Section 6 we obtain solutions to some related local minimization problems. Finally, the proof of Theorem 1.1 is given in Section 7.

2. SOME PRELIMINARY FACTS

In this section, we collect some preliminary facts in order to establish the functional setting. First of all, let us introduce the standard notations for future use in this paper. We denote the upper half-space in \mathbb{R}^{N+1} by

$$\mathbb{R}_+^{N+1} := \{z = (x, y) = (x_1, \dots, x_N, y) \in \mathbb{R}^{N+1} : y > 0\}.$$

Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain. Denote by

$$\mathcal{C}_\Omega := \Omega \times (0, \infty) \subset \mathbb{R}_+^{N+1},$$

the cylinder with base Ω and its lateral boundary by $\partial_L \mathcal{C}_\Omega := \partial\Omega \times (0, \infty)$. The powers $(-\Delta)^s$ of the positive Laplace operator $-\Delta$, in Ω , with zero Dirichlet boundary conditions are defined via its spectral decomposition, namely

$$(-\Delta)^s u(x) := \sum_{j=1}^{\infty} a_j \rho_j^s \varphi_j(x),$$

where (ρ_j, φ_j) is the sequence of eigenvalues and eigenfunctions of the operator $-\Delta$ in Ω under zero Dirichlet boundary data and a_j are the coefficients of u for the base $\{\varphi_j\}_{j=1}^{\infty}$ in $L^2(\Omega)$. In fact, the fractional Laplacian $(-\Delta)^s$ is well defined in the space of functions

$$H_0^s(\Omega) := \left\{ u = \sum_{j=1}^{\infty} a_j \varphi_j \in L^2(\Omega) : \|u\|_{H_0^s} = \left(\sum_{j=1}^{\infty} a_j^2 \rho_j^s \right)^{1/2} < \infty \right\},$$

and $\|u\|_{H_0^s} = \|(-\Delta)^{s/2} u\|_{L^2(\Omega)}$. The dual space $H^{-s}(\Omega)$ is defined in the standard way, as well as the inverse operator $(-\Delta)^{-s}$.

Definition 2.1. We say that $(u, v) \in H_0^s(\Omega) \times H_0^s(\Omega)$ is a solution of (1.1) if the identity

$$\begin{aligned} & \int_{\Omega} \left((-\Delta)^{\frac{s}{2}} u (-\Delta)^{\frac{s}{2}} \varphi_1 + (-\Delta)^{\frac{s}{2}} v (-\Delta)^{\frac{s}{2}} \varphi_2 \right) dx - \int_{\Omega} (\lambda |u|^{q-2} u \varphi_1 + \mu |v|^{q-2} v \varphi_2) dx \\ & - \frac{2\alpha}{\alpha + \beta} \int_{\Omega} |u|^{\alpha-2} u |v|^{\beta} \varphi_1 dx - \frac{2\beta}{\alpha + \beta} \int_{\Omega} |u|^{\alpha} |v|^{\beta-2} v \varphi_2 dx = 0, \end{aligned}$$

holds for all $(\varphi_1, \varphi_2) \in H_0^s(\Omega) \times H_0^s(\Omega)$.

Associated with problem (1.1), we consider the energy functional

$$\begin{aligned} \mathcal{J}_{\lambda, \mu}(u, v) &:= \frac{1}{2} \int_{\Omega} \left(|(-\Delta)^{\frac{s}{2}} u|^2 + |(-\Delta)^{\frac{s}{2}} v|^2 \right) dx - \frac{1}{q} \int_{\Omega} (\lambda |u|^q + \mu |v|^q) dx \\ & - \frac{2}{\alpha + \beta} \int_{\Omega} |u|^{\alpha} |v|^{\beta} dx. \end{aligned}$$

The functional is well defined in $H_0^s(\Omega) \times H_0^s(\Omega)$, and moreover, the critical points of the functional $\mathcal{J}_{\lambda, \mu}$ correspond to solutions of (1.1). We now conclude the main ingredients of a recently developed technique used in order to deal with fractional powers of the Laplacian. To treat the nonlocal problem (1.1), we shall study a corresponding extension problem, which allows us to investigate problem (1.1) by studying a local problem via classical variational methods. We first define the extension operator and fractional Laplacian for functions in $H_0^s(\Omega) \times H_0^s(\Omega)$. We refer the reader to [3–5, 10] and to the references therein.

Definition 2.2. For a function $u \in H_0^s(\Omega)$, we denote its s -harmonic extension $w = E_s(u)$ to the cylinder \mathcal{C}_{Ω} as the solution of the problem

$$\begin{cases} \operatorname{div}(y^{1-2s} \nabla w) = 0 & \text{in } \mathcal{C}_{\Omega} \\ w = 0 & \text{on } \partial_L \mathcal{C}_{\Omega} \\ w = u. & \text{on } \Omega \times \{0\}, \end{cases}$$

and

$$(-\Delta)^s u(x) = -k_s \lim_{y \rightarrow 0^+} y^{1-2s} \frac{\partial w}{\partial y}(x, y),$$

where $k_s = 2^{1-2s} \Gamma(1-s) / \Gamma(s)$ is a normalization constant.

The extension function $w(x, y)$ belongs to the space

$$X_0^s(\mathcal{C}_{\Omega}) := \overline{C_0^\infty(\Omega \times [0, \infty))}^{\|\cdot\|_{X_0^s(\mathcal{C}_{\Omega})}}$$

endowed with the norm

$$\|z\|_{X_0^s(\mathcal{C}_{\Omega})} := \left(k_s \int_{\mathcal{C}_{\Omega}} y^{1-2s} |\nabla z|^2 dx dy \right)^{1/2}.$$

The extension operator is an isometry between $H_0^s(\Omega)$ and $X_0^s(\mathcal{C}_{\Omega})$, namely

$$\|u\|_{H_0^s(\Omega)} = \|E_s(u)\|_{X_0^s(\mathcal{C}_{\Omega})}, \quad \text{for all } u \in H_0^s(\Omega).$$

With this extension we can reformulate (1.1) as the following local problem

$$(2.1) \quad \begin{cases} -\operatorname{div}(y^{1-2s}\nabla w_1) = 0, & -\operatorname{div}(y^{1-2s}\nabla w_2) = 0 & \text{in } \mathcal{C}_\Omega \\ w_1 = w_2 = 0 & & \text{on } \partial_L \mathcal{C}_\Omega \\ w_1 = u, \quad w_2 = v & & \text{on } \Omega \times \{0\} \\ \frac{\partial w_1}{\partial \nu^s} = \lambda |w_1|^{q-2} w_1 + \frac{2\alpha}{\alpha+\beta} |w_1|^{\alpha-2} w_1 |w_2|^\beta & & \text{on } \Omega \times \{0\} \\ \frac{\partial w_2}{\partial \nu^s} = \mu |w_2|^{q-2} w_2 + \frac{2\beta}{\alpha+\beta} |w_1|^\alpha |w_2|^{\beta-2} w_2 & & \text{on } \Omega \times \{0\}, \end{cases}$$

where

$$\frac{\partial w_i}{\partial \nu^s} := -k_s \lim_{y \rightarrow 0^+} y^{1-2s} \frac{\partial w_i}{\partial y}, \quad i = 1, 2$$

and $w_1, w_2 \in X_0^s(\mathcal{C}_\Omega)$ are the s -harmonic extension of $u, v \in H_0^s(\Omega)$, respectively. Let

$$E_0^s(\mathcal{C}_\Omega) := X_0^s(\mathcal{C}_\Omega) \times X_0^s(\mathcal{C}_\Omega).$$

An energy solution to this problem is a function $(w_1, w_2) \in E_0^\alpha(\mathcal{C}_\Omega)$ satisfying

$$\begin{aligned} & k_s \int_{\mathcal{C}_\Omega} y^{1-2s} \nabla w_1 \cdot \nabla \varphi_1 dx dy + k_s \int_{\mathcal{C}_\Omega} y^{1-2s} \nabla w_2 \cdot \nabla \varphi_2 dx dy \\ & = \lambda \int_{\Omega} |w_1|^{q-2} w_1 \varphi_1 dx + \frac{2\alpha}{\alpha+\beta} \int_{\Omega} |w_1|^{\alpha-2} w_1 |w_2|^\beta \varphi_1 dx \\ & \quad + \mu \int_{\Omega} |w_2|^{q-2} w_2 \varphi_2 dx + \frac{2\beta}{\alpha+\beta} \int_{\Omega} |w_1|^\alpha |w_2|^{\beta-2} w_2 \varphi_2 dx, \end{aligned}$$

for all $(\varphi_1, \varphi_2) \in E_0^s(\mathcal{C}_\Omega)$. If $(w_1, w_2) \in E_0^s(\mathcal{C}_\Omega)$ satisfies (2.1), then $(u, v) = (w_1(\cdot, 0), w_2(\cdot, 0))$, defined in the sense of traces, belongs to the space $H_0^s(\Omega) \times H_0^s(\Omega)$ and it is a solution of the original problem (1.1). The associated energy functional to the problem (2.1) is denoted by

$$\begin{aligned} \mathcal{I}_{\lambda, \mu}(w) & := \mathcal{I}_{\lambda, \mu}(w_1, w_2) = \frac{k_s}{2} \int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla w_1|^2 + |\nabla w_2|^2) dx dy \\ & \quad - \frac{1}{q} \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx - \frac{2}{\alpha+\beta} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx. \end{aligned}$$

Critical points of $\mathcal{I}_{\lambda, \mu}$ in $E_0^s(\mathcal{C}_\Omega)$ correspond to critical points of $\mathcal{J}_{\lambda, \mu} : H_0^s(\Omega) \times H_0^s(\Omega) \rightarrow \mathbb{R}$. In the following lemma we list some relevant inequalities from [5].

Lemma 2.3. *For any $1 \leq r \leq 2_s^*$ and any $z \in X_0^s(\mathcal{C}_\Omega)$, it holds*

$$(2.2) \quad \left(\int_{\Omega} |u(x)|^r dx \right)^{\frac{2}{r}} \leq C \int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla z(x, y)|^2 dx dy, \quad u := \operatorname{Tr}(z),$$

for some positive constant $C = C(r, s, N, \Omega)$. Furthermore, the space $X_0^s(\mathcal{C}_\Omega)$ is compactly embedded into $L^r(\Omega)$, for every $r < 2_s^*$.

Remark 2.4. When $r = 2_s^*$, the best constant in (2.2) is denoted by $\mathcal{S}(s, N)$, that is

$$(2.3) \quad \mathcal{S}(s, N) := \inf_{z \in X_0^s(\mathcal{C}_\Omega) \setminus \{0\}} \frac{\int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla z(x, y)|^2 dx dy}{\left(\int_{\Omega} |z(x, 0)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}}}.$$

It is not achieved in any bounded domain and, for all $z \in X^s(\mathbb{R}_+^{N+1})$,

$$(2.4) \quad \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla z(x, y)|^2 dx dy \geq \mathcal{S}(s, N) \left(\int_{\mathbb{R}^N} |z(x, 0)|^{\frac{2N}{N-2s}} dx \right)^{\frac{N-2s}{N}}.$$

$\mathcal{S}(s, N)$ is achieved for $\Omega = \mathbb{R}^N$ by functions w_ε which are the s -harmonic extensions of

$$(2.5) \quad u_\varepsilon(x) := \frac{\varepsilon^{(N-2s)/2}}{(\varepsilon^2 + |x|^2)^{(N-2s)/2}}, \quad \varepsilon > 0, \quad x \in \mathbb{R}^N.$$

Let $U(x) = (1 + |x|^2)^{\frac{2s-N}{2}}$ and let \mathcal{W} be the extension of U . Then

$$\mathcal{W}(x, y) = E_s(U) = c_{N,s} y^{2s} \int_{\mathbb{R}^N} \frac{U(z) dz}{(|x-z|^2 + y^2)^{\frac{N+2s}{2}}},$$

is the extreme function for the fractional Sobolev inequality (2.4). The constant $\mathcal{S}(s, N)$ given in (2.3) takes the exact value

$$\mathcal{S}(s, N) = \frac{2\pi^s \Gamma(\frac{N+2s}{2}) \Gamma(1-s) (\Gamma(\frac{N}{2}))^{\frac{2s}{N}}}{\Gamma(s) \Gamma(\frac{N-2s}{2}) (\Gamma(N))^s},$$

and it is achieved for $\Omega = \mathbb{R}^N$ by the functions w_ε .

Now, we consider the following minimization problem

$$(2.6) \quad \mathcal{S}_{s,\alpha,\beta} := \inf_{(w_1, w_2) \in E_0^s(\mathcal{C}_\Omega) \setminus \{0\}} \frac{\int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla w_1|^2 + |\nabla w_2|^2) dx dy}{\left(\int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \right)^{\frac{2}{2^*}}}.$$

Using ideas from [1], we establish a relationship between $\mathcal{S}(s, N)$ and $\mathcal{S}_{s,\alpha,\beta}$ (see also [18]).

Lemma 2.5. *For the constants $\mathcal{S}(s, N)$ and $\mathcal{S}_{s,\alpha,\beta}$ introduced in (2.3) and (2.6), it holds*

$$(2.7) \quad \mathcal{S}_{s,\alpha,\beta} = \left[\left(\frac{\alpha}{\beta} \right)^{\frac{\beta}{2^*}} + \left(\frac{\beta}{\alpha} \right)^{\frac{\alpha}{2^*}} \right] \mathcal{S}(s, N).$$

In particular, the constant $\mathcal{S}_{s,\alpha,\beta}$ is achieved for $\Omega = \mathbb{R}^N$.

Proof. Let $\{z_n\} \subset X_0^s(\mathcal{C}_\Omega)$ be a minimization sequence for $\mathcal{S}(s, N)$. Let $s, t > 0$ to be chosen later and consider the sequences $w_{1,n} := sz_n$ and $w_{2,n} := tz_n$ in $X_0^s(\mathcal{C}_\Omega)$. By (2.6), we have

$$\frac{s^2 + t^2}{(s^\alpha t^\beta)^{\frac{2}{2^*}}} \frac{\int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla z_n(x, y)|^2 dx dy}{\left(\int_{\Omega} |z_n|^{2^*} dx \right)^{\frac{2}{2^*}}} \geq \mathcal{S}_{s,\alpha,\beta}.$$

Defining $g: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ by setting $g(x) := x^{\frac{2\beta}{2^*}} + x^{\frac{-2\alpha}{2^*}}$, we have

$$\frac{s^2 + t^2}{(s^\alpha t^\beta)^{\frac{2}{2^*}}} = g\left(\frac{s}{t}\right), \quad \min_{\mathbb{R}^+} g = g(x_0) = g(\sqrt{\alpha/\beta}) = \left(\frac{\alpha}{\beta}\right)^{\frac{\beta}{2^*}} + \left(\frac{\beta}{\alpha}\right)^{\frac{\alpha}{2^*}}.$$

Choosing s, t in the previous inequality such that $s/t = \sqrt{\alpha/\beta}$ and letting $n \rightarrow \infty$ yields

$$\left[\left(\frac{\alpha}{\beta} \right)^{\frac{\beta}{2_s^*}} + \left(\frac{\beta}{\alpha} \right)^{\frac{\alpha}{2_s^*}} \right] \mathcal{S}(s, N) \geq \mathcal{S}_{s, \alpha, \beta}.$$

On the other hand, let $\{(w_{1,n}, w_{2,n})\} \subset E_0^s(\mathcal{C}_\Omega) \setminus \{0\}$ be a minimizing sequence for $\mathcal{S}_{s, \alpha, \beta}$. Set $h_n := s_n w_{2,n}$ for $s_n > 0$ with $\int_\Omega |w_{1,n}|^{2_s^*} dx = \int_\Omega |h_n|^{2_s^*} dx$. Then Young's inequality yields

$$\int_\Omega |w_{1,n}|^\alpha |h_n|^\beta dx \leq \frac{\alpha}{2_s^*} \int_\Omega |w_{1,n}|^{2_s^*} dx + \frac{\beta}{2_s^*} \int_\Omega |h_n|^{2_s^*} dx = \int_\Omega |h_n|^{2_s^*} dx = \int_\Omega |w_{1,n}|^{2_s^*} dx.$$

In turn, we can estimate

$$\begin{aligned} & \frac{\int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla w_{1,n}(x, y)|^2 + |\nabla w_{2,n}(x, y)|^2) dx dy}{\left(\int_\Omega |w_{1,n}|^\alpha |w_{2,n}|^\beta dx \right)^{\frac{2}{\alpha+\beta}}} \\ &= \frac{s_n^{\frac{2\beta}{2_s^*}} \int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla w_{1,n}(x, y)|^2 + |\nabla w_{2,n}(x, y)|^2) dx dy}{\left(\int_\Omega |w_{1,n}|^\alpha |h_n|^\beta dx \right)^{\frac{2}{\alpha+\beta}}} \\ &\geq \frac{s_n^{\frac{2\beta}{2_s^*}} \int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla w_{1,n}(x, y)|^2 dx dy}{\left(\int_\Omega |w_{1,n}|^{2_s^*} dx \right)^{2/2_s^*}} \\ &\quad + \frac{s_n^{\frac{2\beta}{2_s^*}} s_n^{-2} \int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla h_n(x, y)|^2 dx dy}{\left(\int_\Omega |h_n|^{2_s^*} dx \right)^{2/2_s^*}} \\ &\geq \mathcal{S}(s, N) g(s_n) \geq \mathcal{S}(s, N) g(\sqrt{\alpha/\beta}). \end{aligned}$$

Passing to the limit in the last inequality we obtain

$$\left[\left(\frac{\alpha}{\beta} \right)^{\frac{\beta}{2_s^*}} + \left(\frac{\beta}{\alpha} \right)^{\frac{\alpha}{2_s^*}} \right] \mathcal{S}(s, N) \leq \mathcal{S}_{s, \alpha, \beta}.$$

Whence, the conclusion follows by combining the previous inequalities. \square

In the end of this section, we fix some notations that will be used in the sequel.

Notations. In this paper we use the following notations:

- $L^p(\Omega)$, $1 \leq p \leq \infty$ denote Lebesgue spaces, with norm $\|\cdot\|_p$. $E = X_0^s(\mathcal{C}_\Omega) \times X_0^s(\mathcal{C}_\Omega)$ is equipped with the norm $\|z\|^2 = \|(w_1, w_2)\|^2 = \|w_1\|_{X_0^s(\mathcal{C}_\Omega)}^2 + \|w_2\|_{X_0^s(\mathcal{C}_\Omega)}^2$.
- The dual space of a Banach space E will be denoted by E^{-1} . We set $tz = t(w_1, w_2) = (tw_1, tw_2)$ for all $z \in E$ and $t \in \mathbb{R}$. $z = (w_1, w_2)$ is said to be non-negative in \mathcal{C}_Ω if $w_1(x, y) \geq 0, w_2(x, y) \geq 0$ in \mathcal{C}_Ω and to be positive if $w_1(x, y) > 0, w_2(x, y) > 0$ in \mathcal{C}_Ω .
- $|\Omega|$ is the Lebesgue measure of Ω . $B(0; r)$ is the ball at the origin with radius r .
- $\mathcal{O}(\varepsilon^t)$ denotes $|\mathcal{O}(\varepsilon^t)|/\varepsilon^t \leq C$ as $\varepsilon \rightarrow 0$ for $t \geq 0$. $o_n(1)$ denotes $o_n(1) \rightarrow 0$ as $n \rightarrow \infty$.
- C, C_i, c will denote various positive constants which may vary from line to line.

3. THE PALAIS-SMALE CONDITION

In this section we shall detect the range of values c for which the $(PS)_c$ -condition holds for the functional $\mathcal{I}_{\lambda,\mu}$. Let $c \in \mathbb{R}$ and set, for simplicity, $E := E_0^s(\mathcal{C}_\Omega)$. We say $\{z_n\} \subset E$ is a $(PS)_c$ -sequence in E for $\mathcal{I}_{\lambda,\mu}$ if $\mathcal{I}_{\lambda,\mu}(z_n) = c + o_n(1)$ and $\mathcal{I}'_{\lambda,\mu}(z_n) = o_n(1)$ strongly in E^{-1} , as $n \rightarrow \infty$. If any $(PS)_c$ -sequence $\{z_n\}$ in E for $\mathcal{I}_{\lambda,\mu}$ admits a convergent subsequence, we say that $\mathcal{I}_{\lambda,\mu}$ satisfies the $(PS)_c$ -condition. We shall need the following preliminary result.

Lemma 3.1. *Let $\{z_n\} \subset E$ be a $(PS)_c$ -sequence for $\mathcal{I}_{\lambda,\mu}$ with $z_n \rightharpoonup z$ in E . Then $\mathcal{I}'_{\lambda,\mu}(z) = 0$ and there exists a positive constant K_0 , depending only on $q, N, \mathcal{S}(s, N)$ and $|\Omega|$, such that*

$$\mathcal{I}_{\lambda,\mu}(z) \geq -K_0 \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right).$$

Proof. Consider $z_n = (w_{1,n}, w_{2,n}) \in E$ and $z = (w_1, w_2) \in E$. If $\{z_n\}$ is a $(PS)_c$ -sequence for $\mathcal{I}_{\lambda,\mu}$ with $z_n \rightharpoonup z$ in E , then $w_{1,n} \rightharpoonup w_1$ and $w_{2,n} \rightharpoonup w_2$ in $X_0^s(\mathcal{C}_\Omega)$, as $n \rightarrow \infty$. Then, by virtue of Sobolev embedding theorem (Lemma 2.3), we also have $w_{1,n}(\cdot, 0) \rightarrow w_1(\cdot, 0)$ and $w_{2,n}(\cdot, 0) \rightarrow w_2(\cdot, 0)$ strongly in $L^q(\Omega)$, as $n \rightarrow \infty$. Of course, up to a further subsequence, $w_{1,n}(\cdot, 0) \rightarrow w_1(\cdot, 0)$ and $w_{2,n}(\cdot, 0) \rightarrow w_2(\cdot, 0)$ a.e. in Ω . It is standard to check that $\mathcal{I}'_{\lambda,\mu}(z) = 0$. This implies that $\langle \mathcal{I}'_{\lambda,\mu}(z), z \rangle = 0$, namely

$$k_s \int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla w_1|^2 + |\nabla w_2|^2) dx dy = \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx + 2 \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx.$$

Consequently, we get

$$(3.1) \quad \begin{aligned} \mathcal{I}_{\lambda,\mu}(z) &= \left(\frac{1}{2} - \frac{1}{2_s^*} \right) k_s \int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla w_1|^2 + |\nabla w_2|^2) dx dy \\ &\quad - \left(\frac{1}{q} - \frac{1}{2_s^*} \right) \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx. \end{aligned}$$

By (3.1), Hölder and Young inequalities and the Sobolev embedding theorem, we obtain

$$\begin{aligned} \mathcal{I}_{\lambda,\mu}(z) &= \left(\frac{1}{2} - \frac{1}{2_s^*} \right) \|z\|^2 - \left(\frac{1}{q} - \frac{1}{2_s^*} \right) \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx \\ &\geq \frac{s}{N} \|z\|^2 - \frac{2_s^* - q}{q 2_s^*} |\Omega|^{(2_s^* - q)/2_s^*} (\lambda \|w_1\|_{2_s^*}^q + \mu \|w_2\|_{2_s^*}^q) \\ &\geq \frac{s}{N} \|z\|^2 - \frac{2_s^* - q}{q 2_s^*} |\Omega|^{(2_s^* - q)/2_s^*} \mathcal{S}(s, N)^{-\frac{q}{2}} (\lambda \|w_1\|^q + \mu \|w_2\|^q) \\ &\geq \frac{s}{N} \|z\|^2 - C \left(\frac{2-q}{2} \varepsilon \left[\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right] + \frac{q}{2} \varepsilon^{-\frac{2-q}{q}} [\|w_1\|^2 + \|w_2\|^2] \right) \\ &= \frac{s}{N} \|z\|^2 - \frac{s}{N} \|z\|^2 - K_0 \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right) \\ &= -K_0 \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right), \end{aligned}$$

which yields the assertion, where we have put

$$C := \frac{2_s^* - q}{q 2_s^*} |\Omega|^{(2_s^* - q)/2_s^*} \mathcal{S}(s, N)^{-\frac{q}{2}}, \quad \varepsilon := \left(\frac{NqC}{2s} \right)^{\frac{q}{2-q}}, \quad K_0 := \frac{\varepsilon(2-q)}{2} C,$$

the positive constants involving only $q, |\Omega|, \mathcal{S}(s, N)$ and N . \square

Lemma 3.2. *If $\{z_n\} \subset E$ is a $(PS)_c$ -sequence for $\mathcal{I}_{\lambda,\mu}$, then $\{z_n\}$ is bounded in E .*

Proof. Let $z_n = (w_{1,n}, w_{2,n}) \subset E$ be a $(PS)_c$ -sequence for $\mathcal{I}_{\lambda,\mu}$ and suppose, by contradiction, that $\|z_n\| \rightarrow \infty$, as $n \rightarrow \infty$. Put

$$\tilde{z}_n = (\tilde{w}_{1,n}, \tilde{w}_{2,n}) := \frac{z_n}{\|z_n\|} = \left(\frac{w_{1,n}}{\|z_n\|}, \frac{w_{2,n}}{\|z_n\|} \right).$$

We may assume that $\tilde{z}_n \rightharpoonup \tilde{z} = (\tilde{w}_1, \tilde{w}_2)$ in E . This implies that $\tilde{w}_{1,n}(\cdot, 0) \rightarrow \tilde{w}_1(\cdot, 0)$ and $\tilde{w}_{2,n}(\cdot, 0) \rightarrow \tilde{w}_2(\cdot, 0)$ strongly in $L^r(\Omega)$ for all $1 \leq r < 2_s^*$ and, thus,

$$\int_{\Omega} (\lambda |\tilde{w}_{1,n}|^q + \mu |\tilde{w}_{2,n}|^q) dx = \int_{\Omega} (\lambda |\tilde{w}_1|^q + \mu |\tilde{w}_2|^q) dx + o_n(1).$$

Since $\{z_n\}$ is a $(PS)_c$ sequence for $\mathcal{I}_{\lambda,\mu}$ and $\|z_n\| \rightarrow \infty$, we get

$$(3.2) \quad \begin{aligned} & \frac{k_s}{2} \int_{\mathcal{C}_{\Omega}} y^{1-2s} (|\nabla \tilde{w}_{1,n}|^2 + |\nabla \tilde{w}_{2,n}|^2) dx dy - \frac{\|z_n\|^{q-2}}{q} \int_{\Omega} (\lambda |\tilde{w}_{1,n}|^q + \mu |\tilde{w}_{2,n}|^q) dx \\ & - \frac{2\|z_n\|^{2_s^*-2}}{2_s^*} \int_{\Omega} |\tilde{w}_{1,n}|^{\alpha} |\tilde{w}_{2,n}|^{\beta} dx = o_n(1), \end{aligned}$$

and

$$(3.3) \quad \begin{aligned} & k_s \int_{\mathcal{C}_{\Omega}} y^{1-2s} (|\nabla \tilde{w}_{1,n}|^2 + |\nabla \tilde{w}_{2,n}|^2) dx dy - \|z_n\|^{q-2} \int_{\Omega} (\lambda |\tilde{w}_{1,n}|^q + \mu |\tilde{w}_{2,n}|^q) dx \\ & - 2\|z_n\|^{2_s^*-2} \int_{\Omega} |\tilde{w}_{1,n}|^{\alpha} |\tilde{w}_{2,n}|^{\beta} dx = o_n(1). \end{aligned}$$

Combining (3.2) and (3.3), as $n \rightarrow \infty$, we obtain

$$(3.4) \quad \begin{aligned} & k_s \int_{\mathcal{C}_{\Omega}} y^{1-2s} (|\nabla \tilde{w}_{1,n}|^2 + |\nabla \tilde{w}_{2,n}|^2) dx dy \\ & = \frac{2(2_s^* - q)}{q(2_s^* - 2)} \|z_n\|^{q-2} \int_{\Omega} (\lambda |\tilde{w}_{1,n}|^q + \mu |\tilde{w}_{2,n}|^q) dx + o_n(1). \end{aligned}$$

In view of $1 < q < 2$ and $\|z_n\| \rightarrow \infty$, (3.4) implies that

$$k_s \int_{\mathcal{C}_{\Omega}} y^{1-2s} (|\nabla \tilde{w}_{1,n}|^2 + |\nabla \tilde{w}_{2,n}|^2) dx dy \rightarrow 0,$$

as $n \rightarrow \infty$, which contradicts to the fact that $\|\tilde{z}_n\| = 1$ for any $n \geq 1$. \square

Lemma 3.3. $\mathcal{I}_{\lambda,\mu}$ satisfies the $(PS)_c$ condition with c satisfying

$$-\infty < c < c_{\infty} := \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s,\alpha,\beta}}{2} \right)^{\frac{N}{2s}} - K_0 \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right),$$

where K_0 is the positive constant introduced in Lemma 3.1

Proof. Let $\{z_n\} \subset E$ be a $(PS)_c$ -sequence for $\mathcal{I}_{\lambda,\mu}$ with $c \in (-\infty, c_{\infty})$. Write $z_n = (w_{1,n}, w_{2,n})$. By Lemma 3.2, we see that $\{z_n\}$ is bounded in E and $z_n \rightharpoonup z = (w_1, w_2)$ up to a subsequence and z is a critical point of $\mathcal{I}_{\lambda,\mu}$. Furthermore, $w_{1,n} \rightharpoonup w_1$ and $w_{2,n} \rightharpoonup w_2$ weakly in $X_0^s(\mathcal{C}_{\Omega})$, $w_{1,n}(\cdot, 0) \rightarrow w_1(\cdot, 0)$ and $w_{2,n}(\cdot, 0) \rightarrow w_2(\cdot, 0)$ strongly in $L^r(\Omega)$ for every

$1 \leq r < 2_s^*$ and $w_{1,n}(\cdot, 0) \rightarrow w_1(\cdot, 0)$, $w_{2,n}(\cdot, 0) \rightarrow w_2(\cdot, 0)$ a.e. in Ω , up to a subsequence. Hence, we have

$$(3.5) \quad \int_{\Omega} (\lambda |w_{1,n}|^q + \mu |w_{2,n}|^q) dx = \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx + o_n(1).$$

Let $\widehat{w}_{1,n} := w_{1,n} - w_1$, $\widehat{w}_{2,n} := w_{2,n} - w_2$ and $\widehat{z}_n := (\widehat{w}_{1,n}, \widehat{w}_{2,n})$. Then, we obtain

$$\|\widehat{z}_n\|^2 = \|z_n\|^2 - \|z\|^2 + o_n(1).$$

In light of [19, Lemma 2.1], we also get

$$(3.6) \quad \int_{\Omega} |\widehat{w}_{1,n}|^{\alpha} |\widehat{w}_{2,n}|^{\beta} dx = \int_{\Omega} |w_{1,n}|^{\alpha} |w_{2,n}|^{\beta} dx - \int_{\Omega} |w_1|^{\alpha} |w_2|^{\beta} dx + o_n(1).$$

Using $\mathcal{I}_{\lambda,\mu}(z_n) = c + o_n(1)$ and $\mathcal{I}'_{\lambda,\mu}(z_n) = o_n(1)$ and (3.5)-(3.6), we conclude

$$(3.7) \quad \begin{aligned} \frac{1}{2} \|\widehat{z}_n\|^2 - \frac{2}{2_s^*} \int_{\Omega} |\widehat{w}_{1,n}|^{\alpha} |\widehat{w}_{2,n}|^{\beta} dx &= c - \mathcal{I}_{\lambda,\mu}(z) + o_n(1), \\ \|\widehat{z}_n\|^2 - 2 \int_{\Omega} |\widehat{w}_{1,n}|^{\alpha} |\widehat{w}_{2,n}|^{\beta} dx &= \langle \mathcal{I}'_{\lambda,\mu}(z_n), z_n \rangle - \langle \mathcal{I}'_{\lambda,\mu}(z), z \rangle + o_n(1) = o_n(1). \end{aligned}$$

Hence, we may assume that

$$(3.8) \quad \|\widehat{z}_n\|^2 \rightarrow \ell, \quad 2 \int_{\Omega} |\widehat{w}_{1,n}|^{\alpha} |\widehat{w}_{2,n}|^{\beta} dx \rightarrow \ell.$$

If $\ell = 0$, the proof is complete. If, by contradiction, $\ell > 0$ then from (3.8) and the definition of $\mathcal{S}_{s,\alpha,\beta}$, we have

$$k_s \mathcal{S}_{s,\alpha,\beta} \left(\frac{\ell}{2} \right)^{\frac{2}{2_s^*}} = k_s \mathcal{S}_{s,\alpha,\beta} \lim_{n \rightarrow \infty} \left(\int_{\Omega} |\widehat{w}_{1,n}|^{\alpha} |\widehat{w}_{2,n}|^{\beta} dx \right)^{\frac{2}{2_s^*}} \leq \lim_{n \rightarrow \infty} \|\widehat{z}_n\|^2 = \ell,$$

which implies that $\ell \geq 2(k_s \mathcal{S}_{s,\alpha,\beta} / 2)^{\frac{N}{2_s}}$. On the other hand, from Lemma 3.1, formulas (3.7) and (3.8), we obtain

$$c = \left(\frac{1}{2} - \frac{1}{2_s^*} \right) \ell + \mathcal{I}_{\lambda,\mu}(z) \geq \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s,\alpha,\beta}}{2} \right)^{\frac{N}{2_s}} - K_0 \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right),$$

which contradicts $c < c_{\infty}$. \square

4. THE NEHARI MANIFOLD

Since the energy functional $\mathcal{I}_{\lambda,\mu}$ associated with (2.1) is not bounded on E , it is useful to consider the functional on the Nehari manifold

$$\mathcal{N}_{\lambda,\mu} := \{z \in E \setminus \{0\} : \langle \mathcal{I}'_{\lambda,\mu}(z), z \rangle = 0\}.$$

Thus, $z = (w_1, w_2) \in \mathcal{N}_{\lambda,\mu}$ if and only if

$$(4.1) \quad \langle \mathcal{I}'_{\lambda,\mu}(z), z \rangle = \|z\|^2 - Q_{\lambda,\mu}(z) - 2 \int_{\Omega} |w_1|^{\alpha} |w_2|^{\beta} dx = 0,$$

where

$$Q_{\lambda,\mu}(z) := \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx.$$

It is clear that all critical points of $\mathcal{I}_{\lambda,\mu}$ must lie on $\mathcal{N}_{\lambda,\mu}$ and, as we will see below, local minimizers on $\mathcal{N}_{\lambda,\mu}$ are actually critical points of $\mathcal{I}_{\lambda,\mu}$. We have the following results.

Lemma 4.1. *The energy functional $\mathcal{I}_{\lambda,\mu}$ is bounded below and coercive on $\mathcal{N}_{\lambda,\mu}$.*

Proof. Let $z = (w_1, w_2) \in \mathcal{N}_{\lambda,\mu}$, then by (4.1) and the Hölder inequality and the Sobolev embedding theorem

$$(4.2) \quad \begin{aligned} \mathcal{I}_{\lambda,\mu}(z) &= \frac{2_s^* - 2}{22_s^*} \|z\|^2 - \frac{2_s^* - q}{q2_s^*} Q_{\lambda,\mu}(z) \\ &\geq \frac{2_s^* - 2}{22_s^*} \|z\|^2 - \frac{2_s^* - q}{q2_s^*} S(s, N)^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} \left(\lambda^{\frac{2}{2_s^* - q}} + \mu^{\frac{2}{2_s^* - q}} \right)^{\frac{2-q}{2}} \|z\|^q. \end{aligned}$$

Since $1 < q < 2$, the functional $\mathcal{I}_{\lambda,\mu}$ is coercive and bounded below on $\mathcal{N}_{\lambda,\mu}$. \square

The Nehari manifold $\mathcal{N}_{\lambda,\mu}$ is closely linked to the fibering map $\Phi_z : t \rightarrow \mathcal{I}_{\lambda,\mu}(tz)$ given by

$$\Phi_z(t) := \mathcal{I}_{\lambda,\mu}(tz) = \frac{t^2}{2} \|z\|^2 - \frac{t^q}{q} \int_{\Omega} (\lambda|w_1|^q + \mu|w_2|^q) dx - \frac{2t^{2_s^*}}{\alpha + \beta} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx.$$

Such maps were introduced by Drabek and Pohozaev in [16] and later on used by Brown and Zhang [7]. Notice that we have

$$\begin{aligned} \Phi'_z(t) &= t\|z\|^2 - t^{q-1} \int_{\Omega} (\lambda|w_1|^q + \mu|w_2|^q) dx - 2t^{2_s^*-1} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx, \\ \Phi''_z(t) &= \|z\|^2 - (q-1)t^{q-2} \int_{\Omega} (\lambda|w_1|^q + \mu|w_2|^q) dx - 2(2_s^* - 1)t^{2_s^*-2} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx. \end{aligned}$$

It is clear that $\Phi'_z(t) = 0$ if and only if $tz \in \mathcal{N}_{\lambda,\mu}$. Hence, $z \in \mathcal{N}_{\lambda,\mu}$ if and only if $\Phi'_z(1) = 0$. Introduce now the functional

$$\mathcal{R}_{\lambda,\mu}(z) := \langle \mathcal{I}'_{\lambda,\mu}(z), z \rangle.$$

Then, for every $z \in \mathcal{N}_{\lambda,\mu}$, we have

$$(4.3) \quad \begin{aligned} \langle \mathcal{R}'_{\lambda,\mu}(z), z \rangle &= 2\|z\|^2 - qQ_{\lambda,\mu}(z) - 22_s^* \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \\ &= (2 - q)\|z\|^2 - 2(2_s^* - q) \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \\ &= (2_s^* - q)Q_{\lambda,\mu}(z) - (2_s^* - 2)\|z\|^2. \end{aligned}$$

Following the method used in [32], we split $\mathcal{N}_{\lambda,\mu}$ into three parts

$$\begin{aligned} \mathcal{N}_{\lambda,\mu}^+ &:= \{z \in \mathcal{N}_{\lambda,\mu} : \langle \mathcal{R}'_{\lambda,\mu}(z), z \rangle > 0\}, \\ \mathcal{N}_{\lambda,\mu}^0 &:= \{z \in \mathcal{N}_{\lambda,\mu} : \langle \mathcal{R}'_{\lambda,\mu}(z), z \rangle = 0\}, \\ \mathcal{N}_{\lambda,\mu}^- &:= \{z \in \mathcal{N}_{\lambda,\mu} : \langle \mathcal{R}'_{\lambda,\mu}(z), z \rangle < 0\}. \end{aligned}$$

Then, we have the following lemmas.

Lemma 4.2. *If z_0 is a local minimizer for $\mathcal{I}_{\lambda,\mu}$ on $\mathcal{N}_{\lambda,\mu}$ and $z_0 \notin \mathcal{N}_{\lambda,\mu}^0$, then $\mathcal{I}'_{\lambda,\mu}(z_0) = 0$.*

Proof. Let $z_0 = (w_{0,1}, w_{0,2}) \in \mathcal{N}_{\lambda,\mu}$ be a local minimizer for $\mathcal{I}_{\lambda,\mu}$ on $\mathcal{N}_{\lambda,\mu}$. Then z_0 solves

$$\text{Min } \{\mathcal{I}_{\lambda,\mu}(z) : \mathcal{R}_{\lambda,\mu}(z) = 0\}.$$

Hence, there exists a Lagrange multiplier $\gamma \in \mathbb{R}$ such that $\mathcal{I}'_{\lambda,\mu}(z_0) = \gamma \mathcal{R}'_{\lambda,\mu}(z_0)$. Thus,

$$\langle \mathcal{I}'_{\lambda,\mu}(z_0), z_0 \rangle = \gamma \langle \mathcal{R}'_{\lambda,\mu}(z_0), z_0 \rangle = 0.$$

Since $z_0 \notin \mathcal{N}_{\lambda,\mu}^0$, then $\langle \mathcal{R}'_{\lambda,\mu}(z_0), z_0 \rangle \neq 0$, yielding $\gamma = 0$. This completes the proof. \square

Let Λ_1 be the positive number defined in (1.4). Then we have the following result.

Lemma 4.3. *Assume that $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$. Then $\mathcal{N}_{\lambda, \mu}^0 = \emptyset$.*

Proof. Assume by contradiction that there exist $\lambda > 0$ and $\mu > 0$ with $0 < \lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} < \Lambda_1$ and such that $\mathcal{N}_{\lambda, \mu}^0 \neq \emptyset$. Let $z \in \mathcal{N}_{\lambda, \mu}^0$. Then, by virtue of (4.3), we get

$$\|z\|^2 = \frac{2(2_s^* - q)}{2 - q} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx, \quad \|z\|^2 = \frac{2_s^* - q}{2_s^* - 2} Q_{\lambda, \mu}(z).$$

By Hölder inequality and the Sobolev embedding theorem, we have

$$\begin{aligned} \|z\| &\geq \left[\frac{2 - q}{2(2_s^* - q)} (k_s \mathcal{S}_{s, \alpha, \beta})^{\frac{2_s^*}{2}} \right]^{\frac{1}{2_s^* - 2}}, \\ \|z\| &\leq \left(\frac{2_s^* - q}{2_s^* - 2} \mathcal{S}(s, N)^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} \right)^{\frac{1}{2 - q}} (\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}})^{\frac{1}{2}}, \end{aligned}$$

which leads to the inequality

$$\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \geq \left(\frac{2_s^* - q}{2_s^* - 2} \mathcal{S}(s, N)^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} \right)^{-\frac{2}{2-q}} \left[\frac{2 - q}{2(2_s^* - q)} (k_s \mathcal{S}_{s, \alpha, \beta})^{\frac{2_s^*}{2}} \right]^{\frac{2}{2_s^* - 2}} = \Lambda_1,$$

contradicting the assumption. \square

From Lemma 4.3, if $0 < \lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} < \Lambda_1$, we can write $\mathcal{N}_{\lambda, \mu} = \mathcal{N}_{\lambda, \mu}^+ \cup \mathcal{N}_{\lambda, \mu}^-$ and define

$$\alpha_{\lambda, \mu} := \inf_{z \in \mathcal{N}_{\lambda, \mu}} \mathcal{I}_{\lambda, \mu}(z), \quad \alpha_{\lambda, \mu}^+ := \inf_{z \in \mathcal{N}_{\lambda, \mu}^+} \mathcal{I}_{\lambda, \mu}(z), \quad \alpha_{\lambda, \mu}^- := \inf_{z \in \mathcal{N}_{\lambda, \mu}^-} \mathcal{I}_{\lambda, \mu}(z).$$

Moreover, we have the following properties about the Nehari manifold $\mathcal{N}_{\lambda, \mu}$.

Theorem 4.4. *The following facts holds*

- (i) *If $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$, then we have $\alpha_{\lambda, \mu} \leq \alpha_{\lambda, \mu}^+ < 0$;*
- (ii) *If $(\lambda, \mu) \in \mathcal{C}_{(q/2)^{2/(2-q)} \Lambda_1}$, then we have $\alpha_{\lambda, \mu}^- > c_0$ for some positive constant c_0 depending on λ, μ, N, s and $|\Omega|$.*

Proof. (i) Let $z = (w_1, w_2) \in \mathcal{N}_{\lambda, \mu}^+$. By formula (4.3), we have

$$\frac{2 - q}{2(2_s^* - q)} \|z\|^2 > \int_{\Omega} |w_1|^\alpha |w_2|^\beta$$

and so,

$$\begin{aligned} \mathcal{I}_{\lambda, \mu}(z) &= \left(\frac{1}{2} - \frac{1}{q} \right) \|z\|^2 + 2 \left(\frac{1}{q} - \frac{1}{2_s^*} \right) \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \\ &\leq \left[\left(\frac{1}{2} - \frac{1}{q} \right) + \left(\frac{1}{q} - \frac{1}{2_s^*} \right) \frac{2 - q}{2_s^* - q} \right] \|z\|^2 \\ &= -\frac{(2 - q)s}{Nq} \|z\|^2 < 0. \end{aligned}$$

Therefore, by the definition of $\alpha_{\lambda, \mu}, \alpha_{\lambda, \mu}^+$, we can deduce that $\alpha_{\lambda, \mu} \leq \alpha_{\lambda, \mu}^+ < 0$.

(ii) Let $z \in \mathcal{N}_{\lambda, \mu}^-$. By equation (4.3),

$$\frac{2 - q}{2(2_s^* - q)} \|z\|^2 < \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx.$$

By the Hölder inequality and the Sobolev embedding theorem, we have

$$\int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \leq (k_s \mathcal{S}_{s,\alpha,\beta})^{-\frac{2_s^*}{2}} \|z\|^{2_s^*}.$$

Hence, we obtain

$$\|z\| > \left(\frac{2-q}{2(2_s^* - q)} \right)^{\frac{1}{2_s^*-2}} (k_s \mathcal{S}_{s,\alpha,\beta})^{\frac{N}{4s}}, \quad \text{for all } z \in \mathcal{N}_{\lambda,\mu}^-.$$

From the last inequality we infer that

$$\begin{aligned} \mathcal{I}_{\lambda,\mu}(z) &= \frac{2_s^* - 2}{22_s^*} \|z\|^2 - \frac{2_s^* - q}{q2_s^*} Q_{\lambda,\mu}(z) \\ &\geq \|z\|^q \left[\frac{2_s^* - 2}{22_s^*} \|z\|^{2-q} - \frac{2_s^* - q}{q2_s^*} \mathcal{S}(s, N)^{-\frac{q}{2}} |\Omega|^{\frac{2_s^*-q}{2_s^*}} \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right)^{\frac{2-q}{2}} \right] \\ &> \left(\frac{2-q}{2(2_s^* - q)} \right)^{\frac{q}{2_s^*-2}} (k_s \mathcal{S}_{s,\alpha,\beta})^{\frac{qN}{4s}} \left[\frac{2_s^* - 2}{22_s^*} (k_s \mathcal{S}_{s,\alpha,\beta})^{\frac{(2-q)N}{4s}} \left(\frac{2-q}{2(2_s^* - q)} \right)^{\frac{2-q}{2_s^*-2}} \right. \\ &\quad \left. - \frac{2_s^* - q}{q2_s^*} \mathcal{S}(s, N)^{-\frac{q}{2}} |\Omega|^{\frac{2_s^*-q}{2_s^*}} \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right)^{\frac{2-q}{2}} \right]. \end{aligned}$$

Thus, if $\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} < (q/2)^{\frac{2}{2-q}} \Lambda_1$, then

$$\mathcal{I}_{\lambda,\mu}(z) > c_0, \quad \text{for all } z \in \mathcal{N}_{\lambda,\mu}^-,$$

for some positive constant $c_0 = c_0(\lambda, \mu, \mathcal{S}(s, N), |\Omega|)$. □

Theorem 4.5. *Let $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$. Then, for every $z = (w_1, w_2) \in E$ with*

$$\int_{\Omega} |w_1|^\alpha |w_2|^\beta dx > 0,$$

then there exist (unique) $t^- = t^-(z) > 0$ and $t^+ = t^+(z) > 0$ such that

$$t^+ z \in \mathcal{N}_{\lambda,\mu}^+, \quad t^- z \in \mathcal{N}_{\lambda,\mu}^-.$$

In particular, we have

$$t^+ < t_{\max} < t^-, \quad t_{\max} := \left(\frac{(2-q)\|z\|^2}{2(2_s^* - q) \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx} \right)^{\frac{1}{2_s^*-2}}$$

as well as

$$\mathcal{I}_{\lambda,\mu}(t^+ z) = \min_{0 \leq t \leq t_{\max}} \mathcal{I}_{\lambda,\mu}(tz), \quad \mathcal{I}_{\lambda,\mu}(t^- z) = \max_{t \geq 0} \mathcal{I}_{\lambda,\mu}(tz).$$

Proof. Fix $z = (w_1, w_2) \in E$ with $\int_{\Omega} |w_1|^\alpha |w_2|^\beta dx > 0$. Put

$$(4.4) \quad \mathbf{m}(t) := t^{2-q} \|z\|^2 - 2t^{2_s^*-q} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \quad \text{for } t \geq 0.$$

Clearly, $\mathbf{m}(0) = 0$ and $\mathbf{m}(t) \rightarrow -\infty$ as $t \rightarrow \infty$. Notice that

$$\mathbf{m}'(t) = (2-q)t^{1-q} \|z\|^2 - 2(2_s^* - q)t^{2_s^*-q-1} \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx.$$

Clearly, $\mathbf{m}'(t) = 0$ at $t = t_{\max}$, $\mathbf{m}'(t) > 0$ for $t \in (0, t_{\max})$ and $\mathbf{m}'(t) < 0$ for $t \in (t_{\max}, \infty)$. Moreover, by a direct calculation we obtain

$$\begin{aligned} \mathbf{m}(t_{\max}) &= \|z\|^q \left[\left(\frac{2-q}{2(2_s^* - q)} \right)^{\frac{2-q}{2_s^* - 2}} - 2 \left(\frac{2-q}{2(2_s^* - q)} \right)^{\frac{2_s^* - q}{2_s^* - 2}} \right] \left(\frac{\|z\|^{2_s^*}}{\int_{\Omega} |w_1|^{\alpha} |w_2|^{\beta} dx} \right)^{\frac{2-q}{2_s^* - 2}} \\ &\geq \|z\|^q \left(\frac{2_s^* - 2}{2_s^* - q} \right) \left(\frac{2-q}{2(2_s^* - q)} \right) (k_s \mathcal{S}_{\alpha, \beta})^{\frac{2_s^*}{2}} \frac{2-q}{2_s^* - 2}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \mathbf{m}(0) &= 0 \\ &< Q_{\lambda, \mu}(w_1, w_2) \\ (4.5) \quad &\leq |\Omega|^{\frac{2_s^* - q}{2_s^*}} \mathcal{S}(s, N)^{-\frac{q}{2}} (\lambda^{2-q} + \mu^{2-q})^{\frac{2-q}{2}} \|z\|^q \\ &< \|z\|^q \left(\frac{2_s^* - 2}{2_s^* - q} \right) \left(\frac{2-q}{2(2_s^* - q)} \right) (k_s \mathcal{S}_{\alpha, \beta})^{\frac{2_s^*}{2}} \frac{2-q}{2_s^* - 2} \\ &\leq \mathbf{m}(t_{\max}) \end{aligned}$$

for $0 < \lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} < \Lambda_1$. By the equation for $\Phi'_z(t)$ and (4.4) we get

$$(4.6) \quad \Phi'_z(t) = t^{q-1} \left[\mathbf{m}(t) - \int_{\Omega} (\lambda w_1^q + \mu w_2^q) dx \right].$$

Furthermore, if $tz \in \mathcal{N}_{\lambda, \mu}$, then

$$(4.7) \quad t^{q-1} \mathbf{m}'(t) = \Phi''_z(t).$$

By (4.5) and (4.6), there exist unique t^+ and t^- such that $0 < t^+ < t_{\max} < t^-$ such that

$$\mathbf{m}(t^+) = Q_{\lambda, \mu}(w_1, w_2) = \mathbf{m}(t^-), \quad \mathbf{m}'(t^+) > 0 > \mathbf{m}'(t^-),$$

and (4.6) implies $\Phi'_z(t^+) = \Phi'_z(t^-) = 0$. By (4.7), we have $\Phi''_z(t^+) > 0, \Phi''_z(t^-) < 0$. This shows that $\Phi_z(t)$ has a local minimum at t^+ and local maximum at t^- with $t^+z \in \mathcal{N}_{\lambda, \mu}^+, t^-z \in \mathcal{N}_{\lambda, \mu}^-$. Moreover, $\mathcal{I}_{\lambda, \mu}(t^-z) \geq \mathcal{I}_{\lambda, \mu}(tz) \geq \mathcal{I}_{\lambda, \mu}(t^+z)$ for all $t \in [t^+, t^-]$ and $\mathcal{I}_{\lambda, \mu}(t^+z) \leq \mathcal{I}_{\lambda, \mu}(tz)$ for all $t \in [0, t^+]$. Hence, $\mathcal{I}_{\lambda, \mu}(t^+z) = \min_{0 \leq t \leq t_{\max}} \mathcal{I}_{\lambda, \mu}(tz)$ and $\mathcal{I}_{\lambda, \mu}(t^-z) = \max_{t \geq 0} \mathcal{I}_{\lambda, \mu}(tz)$, concluding the proof. \square

5. EXISTENCE OF PALAIS-SMALE SEQUENCES

Lemma 5.1. *Let $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$. Then, for any $z \in \mathcal{N}_{\lambda, \mu}$, there exists $r > 0$ and a differentiable map $\xi : B(0; r) \subset E \rightarrow \mathbb{R}^+$ such that $\xi(0) = 1$ and $\xi(h)(z - h) \in \mathcal{N}_{\lambda, \mu}$ for every $h \in B(0; r)$. Let us set*

$$\begin{aligned} \mathcal{T}_1 &:= 2k_s \int_{\mathcal{C}_{\Omega}} y^{1-2s} (\nabla w_1 \cdot \nabla h_1 + \nabla w_2 \cdot \nabla h_2) dx dy, \\ \mathcal{T}_2 &:= q \int_{\Omega} (\lambda |w_1|^{q-2} w_1 h_1 + \mu |w_2|^{q-2} w_2 h_2) dx, \\ \mathcal{T}_3 &:= 2 \int_{\Omega} \left(\alpha |w_1|^{\alpha-2} w_1 h_1 |w_2|^{\beta} + \beta |w_1|^{\alpha} |w_2|^{\beta-2} w_2 h_2 \right) dx, \end{aligned}$$

for all $(h_1, h_2) \in E$ and $(w_1, w_2) \in E$. Then

$$(5.1) \quad \langle \xi'(0), h \rangle = \frac{\mathcal{T}_3 + \mathcal{T}_2 - \mathcal{T}_1}{(2-q)\|z\|^2 - 2(2_s^* - q) \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx},$$

for all $(h_1, h_2) \in E$.

Proof. For $z = (w_1, w_2) \in \mathcal{N}_{\lambda, \mu}$, define a function $\mathcal{H}_z : \mathbb{R} \times E \rightarrow \mathbb{R}$ by

$$\begin{aligned} \mathcal{H}_z(\xi, p) &:= \langle \mathcal{I}'_{\lambda, \mu}(\xi(z-p)), \xi(z-p) \rangle \\ &= \xi^2 k_s \int_{\mathcal{C}_\Omega} y^{1-2s} (|\nabla(w_1 - p_1)|^2 + |\nabla(w_2 - p_2)|^2) dx dy \\ &\quad - \xi^q \int_{\Omega} (\lambda |w_1 - p_1|^q + \mu |w_2 - p_2|^q) dx - 2\xi^{2_s^*} \int_{\Omega} |w_1 - p_1|^\alpha |w_2 - p_2|^\beta dx. \end{aligned}$$

Then $\mathcal{H}_z(1, 0) = \langle \mathcal{I}'_{\lambda, \mu}(z), z \rangle = 0$ and, by Lemma 4.3, we have

$$\begin{aligned} \frac{d\mathcal{H}_z(1, (0, 0))}{d\xi} &= 2\|z\|^2 - q \int_{\Omega} (\lambda |w_1|^q + \mu |w_2|^q) dx - 22_s^* \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \\ &= (2-q)\|z\|^2 - 2(2_s^* - q) \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx \neq 0. \end{aligned}$$

In turn, by virtue of the Implicit Function Theorem, there exists $r > 0$ and a function $\xi : B(0; r) \subset E \rightarrow \mathbb{R}$ of class C^1 such that $\xi(0) = 1$ and formula (5.1) holds, via direct computation. Moreover, $\mathcal{H}_z(\xi(h), h) = 0$, for all $h \in B(0; r)$, is equivalent to

$$\langle \mathcal{I}'_{\lambda, \mu}(\xi(h)(z-h)), \xi(h)(z-h) \rangle = 0, \quad \text{for all } h \in B(0; r),$$

namely $\xi(h)(z-h) \in \mathcal{N}_{\lambda, \mu}$. \square

Lemma 5.2. *Let $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$. Then, for each $z \in \mathcal{N}_{\lambda, \mu}^-$, there exists $r > 0$ and a differentiable function $\xi^- : B(0; r) \subset E \rightarrow \mathbb{R}^+$ such that $\xi^-(0) = 1$, $\xi^-(h)(z-h) \in \mathcal{N}_{\lambda, \mu}^-$ for every $h \in B(0; r)$ and formula (5.1) holds.*

Proof. Arguing as for the proof of Lemma 5.1, there exists $r > 0$ and a differentiable function $\xi^- : B(0; r) \subset E \rightarrow \mathbb{R}^+$ such that $\xi^-(0) = 1$, $\xi^-(h)(z-h) \in \mathcal{N}_{\lambda, \mu}$ for all $h \in B(0; r)$ and formula (5.1) holds. Since

$$\langle \mathcal{R}'_{\lambda, \mu}(z), z \rangle = (2-q)\|z\|^2 - 2(2_s^* - q) \int_{\Omega} |w_1|^\alpha |w_2|^\beta dx < 0,$$

by the continuity of the functions $\mathcal{R}'_{\lambda, \mu}$ and ξ^- , up to reducing the size of $r > 0$, we get

$$\begin{aligned} \langle \mathcal{R}'_{\lambda, \mu}(\xi^-(h)(z-h)), \xi^-(h)(z-h) \rangle &= (2-q)\|\xi^-(h)(z-h)\|^2 \\ &\quad - 2(2_s^* - q) \int_{\Omega} |(\xi^-(h)(z-h))_1|^\alpha |(\xi^-(h)(z-h))_2|^\beta dx < 0, \end{aligned}$$

where $(\xi^-(h)(z-h))_i \in X_0^s(\mathcal{C}_\Omega)$ denote the components of $\xi^-(h)(z-h)$. This implies that the functions $\xi^-(h)(z-h)$ belong to $\mathcal{N}_{\lambda, \mu}^-$. \square

Proposition 5.3. *The following facts hold.*

- (i) *Let $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$. Then there is a $(PS)_{\alpha_{\lambda, \mu}}$ -sequence $\{z_n\} \subset \mathcal{N}_{\lambda, \mu}$ for $\mathcal{I}_{\lambda, \mu}$.*
- (ii) *Let $(\lambda, \mu) \in \mathcal{C}_{(q/2)^{2/(2-q)}\Lambda_1}$. Then there is a $(PS)_{\alpha_{\lambda, \mu}^-}$ -sequence $\{z_n\} \subset \mathcal{N}_{\lambda, \mu}^-$ for $\mathcal{I}_{\lambda, \mu}$.*

Proof. (i) By Lemma 4.1 and Ekeland Variational Principle [17], there exists a minimizing sequence $\{z_n\} \subset \mathcal{N}_{\lambda,\mu}$ such that

$$(5.2) \quad \begin{aligned} \mathcal{I}_{\lambda,\mu}(z_n) &< \alpha_{\lambda,\mu} + \frac{1}{n}, \\ \mathcal{I}_{\lambda,\mu}(z_n) &< \mathcal{I}_{\lambda,\mu}(w) + \frac{1}{n}\|w - z_n\|, \quad \text{for each } w \in \mathcal{N}_{\lambda,\mu}. \end{aligned}$$

Taking n large and using $\alpha_{\lambda,\mu} < 0$, we have

$$(5.3) \quad \begin{aligned} \mathcal{I}_{\lambda,\mu}(z_n) &= \left(\frac{1}{2} - \frac{1}{2_s^*}\right)\|z_n\|^2 - \left(\frac{1}{q} - \frac{1}{2_s^*}\right) \int_{\Omega} (\lambda|w_{1,n}|^q + \mu|w_{2,n}|^q) dx \\ &< \alpha_{\lambda,\mu} + \frac{1}{n} < \frac{\alpha_{\lambda,\mu}}{2}. \end{aligned}$$

This yields that

$$(5.4) \quad \begin{aligned} -\frac{q2_s^*}{2(2_s^* - q)}\alpha_{\lambda,\mu} &< \int_{\Omega} (\lambda|w_{1,n}|^q + \mu|w_{2,n}|^q) dx \\ &\leq |\Omega|^{\frac{2_s^* - q}{2_s^*}} \mathcal{S}(s, N)^{-\frac{q}{2}} (\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}})^{\frac{2-q}{2}} \|z_n\|^q. \end{aligned}$$

Consequently, $z_n \neq 0$ and combining with (5.3) and (5.4) and using Hölder inequality

$$\|z_n\| > \left[-\frac{q2_s^*}{2(2_s^* - q)}\alpha_{\lambda,\mu} |\Omega|^{\frac{q-2_s^*}{2_s^*}} \mathcal{S}(s, N)^{\frac{q}{2}} (\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}})^{\frac{q-2}{2}} \right]^{\frac{1}{q}},$$

and

$$(5.5) \quad \|z_n\| < \left[\frac{2(2_s^* - q)}{q(2_s^* - 2)} |\Omega|^{\frac{2_s^* - q}{2_s^*}} \mathcal{S}(s, N)^{-\frac{q}{2}} (\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}})^{\frac{2-q}{2}} \right]^{\frac{1}{2-q}}.$$

Now we prove that

$$\|\mathcal{I}'_{\lambda,\mu}(z_n)\|_{E^{-1}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Fix $n \in \mathbb{N}$. By applying Lemma 5.1 to z_n , we obtain the function $\xi_n : B(0; r_n) \rightarrow \mathbb{R}^+$ for some $r_n > 0$, such that $\xi_n(h)(z_n - h) \in \mathcal{N}_{\lambda,\mu}$. Take $0 < \rho < r_n$. Let $w \in E$ with $w \neq 0$ and put $h^* = \frac{\rho w}{\|w\|}$. We set $h_\rho = \xi_n(h^*)(z_n - h^*)$, then $h_\rho \in \mathcal{N}_{\lambda,\mu}$, and we have from (5.2) that

$$\mathcal{I}_{\lambda,\mu}(h_\rho) - \mathcal{I}_{\lambda,\mu}(z_n) \geq -\frac{1}{n}\|h_\rho - z_n\|.$$

By the Mean Value Theorem, we get

$$\langle \mathcal{I}'_{\lambda,\mu}(z_n), h_\rho - z_n \rangle + o(\|h_\rho - z_n\|) \geq -\frac{1}{n}\|h_\rho - z_n\|.$$

Thus, we have

$$\langle \mathcal{I}'_{\lambda,\mu}(z_n), -h^* \rangle + (\xi_n(h^*) - 1) \langle \mathcal{I}'_{\lambda,\mu}(z_n), z_n - h^* \rangle \geq -\frac{1}{n}\|h_\rho - z_n\| + o(\|h_\rho - z_n\|).$$

Whence, from $\xi_n(h^*)(z_n - h^*) \in \mathcal{N}_{\lambda,\mu}$, it follows that

$$\begin{aligned} -\rho \left\langle \mathcal{I}'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \right\rangle + (\xi_n(h^*) - 1) \langle \mathcal{I}'_{\lambda,\mu}(z_n) - \mathcal{I}'_{\lambda,\mu}(h_\rho), z_n - h^* \rangle \\ \geq -\frac{1}{n}\|h_\rho - z_n\| + o(\|h_\rho - z_n\|). \end{aligned}$$

So, we get

$$(5.6) \quad \begin{aligned} \left\langle \mathcal{I}'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \right\rangle &\leq \frac{1}{n\rho} \|h_\rho - z_n\| + \frac{o(\|h_\rho - z_n\|)}{\rho} \\ &\quad + \frac{(\xi_n(h^*) - 1)}{\rho} \langle \mathcal{I}'_{\lambda,\mu}(z_n) - \mathcal{I}'_{\lambda,\mu}(h_\rho), z_n - h^* \rangle. \end{aligned}$$

Since $\|h_\rho - z_n\| \leq \rho|\xi_n(h^*)| + |\xi_n(h^*) - 1|\|z_n\|$ and

$$\lim_{\rho \rightarrow 0} \frac{|\xi_n(h^*) - 1|}{\rho} \leq \|\xi'_n(0)\|.$$

For fixed $n \in \mathbb{N}$, if we let $\rho \rightarrow 0$ in (5.6), then by virtue of (5.5) we can choose a constant $C > 0$ independent of ρ such that

$$\left\langle \mathcal{I}'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \right\rangle \leq \frac{C}{n} (1 + \|\xi'_n(0)\|).$$

Thus, we are done once we prove that $\|\xi'_n(0)\|$ remains uniformly bounded. By (5.1), (5.5) and Hölder inequality, we have

$$|\langle \xi'_n(0), h \rangle| \leq \frac{C_1 \|h\|}{\left| (2-q)\|z_n\|^2 - 2(2_s^* - q) \int_{\Omega} |w_{1,n}|^\alpha |w_{2,n}|^\beta dx \right|}$$

for some $C_1 > 0$. We only need to prove that

$$\left| (2-q)\|z_n\|^2 - 2(2_s^* - q) \int_{\Omega} |w_{1,n}|^\alpha |w_{2,n}|^\beta dx \right| \geq C_2,$$

for some $C_2 > 0$ and n large enough. We argue by contradiction. Suppose that there exists a subsequence $\{z_n\}$ such that

$$(5.7) \quad (2-q)\|z_n\|^2 - 2(2_s^* - q) \int_{\Omega} |w_{1,n}|^\alpha |w_{2,n}|^\beta dx = o_n(1).$$

By virtue of (5.7) and the fact that $z_n \in \mathcal{N}_{\lambda,\mu}$, we have

$$\|z_n\|^2 = \frac{2(2_s^* - q)}{2 - q} \int_{\Omega} |w_{1,n}|^\alpha |w_{2,n}|^\beta dx + o_n(1), \quad \|z_n\|^2 = \frac{2_s^* - q}{2_s^* - 2} Q_{\lambda,\mu}(z_n) + o_n(1).$$

Taking into account that $\mathcal{I}_{\lambda,\mu}(z_n) \rightarrow \alpha_{\lambda,\mu} < 0$ as $n \rightarrow \infty$, we have $\|z_n\| \not\rightarrow 0$ as $n \rightarrow \infty$. Then, arguing as in the proof of Lemma 4.3 yields $(\lambda, \mu) \notin \mathcal{C}_{\Lambda_1}$, a contradiction. Then,

$$\left\langle \mathcal{I}'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \right\rangle \leq \frac{C}{n}.$$

This proves (i). By Lemma 5.2, one can prove (ii), but we shall omit the details here. \square

6. LOCAL MINIMIZATION PROBLEMS

Now, we establish the existence of a local minimizer for $\mathcal{I}_{\lambda,\mu}$ in $\mathcal{N}_{\lambda,\mu}^+$.

Proposition 6.1. *Let $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$. Then $\mathcal{I}_{\lambda,\mu}$ has a local minimizer z^+ in $\mathcal{N}_{\lambda,\mu}^+$ satisfying the following conditions:*

- (i) $\mathcal{I}_{\lambda,\mu}(z^+) = \alpha_{\lambda,\mu} = \alpha_{\lambda,\mu}^+ < 0$;
- (ii) z^+ is a positive solution of (2.1).

Proof. By (i) of Proposition 5.3, there exists a minimizing sequence $\{z_n\} = \{(w_{1,n}, w_{2,n})\}$ for $\mathcal{I}_{\lambda,\mu}$ in $\mathcal{N}_{\lambda,\mu}$ such that, as $n \rightarrow \infty$,

$$(6.1) \quad \mathcal{I}_{\lambda,\mu}(z_n) = \alpha_{\lambda,\mu} + o_n(1) \quad \text{and} \quad \mathcal{I}'_{\lambda,\mu}(z_n) = o_n(1) \quad \text{in } E^{-1}.$$

By Lemma 4.1, we see that $\mathcal{I}_{\lambda,\mu}$ is coercive on $\mathcal{N}_{\lambda,\mu}$, and $\{z_n\}$ is bounded in E . Then there exists a subsequence, still denoted by $\{z_n\}$ and $z^+ = (w_1^+, w_2^+) \in E$ such that, as $n \rightarrow \infty$,

$$\begin{cases} w_{1,n} \rightharpoonup w_1^+, & w_{2,n} \rightharpoonup w_2^+, & \text{weakly in } X_0^s(\Omega), \\ w_{1,n} \rightarrow w_1^+, & w_{2,n} \rightarrow w_2^+, & \text{strongly in } L^r(\Omega) \text{ for all } 1 \leq r < 2_s^*, \\ w_{1,n} \rightarrow w_1^+, & w_{2,n} \rightarrow w_2^+, & \text{a.e. in } \Omega, \end{cases}$$

up to subsequences. This implies that, as $n \rightarrow \infty$,

$$(6.2) \quad Q_{\lambda,\mu}(z_n) = Q_{\lambda,\mu}(z^+) + o_n(1).$$

We claim that z^+ is a nontrivial solution of (2.1). From (6.1)-(6.2), it is easy to verify that z^+ is a weak solution of (2.1). From $z_n \in \mathcal{N}_{\lambda,\mu}$ and (4.2) we deduce that

$$(6.3) \quad Q_{\lambda,\mu}(z_n) = \frac{q(2_s^* - 2)}{2(2_s^* - q)} \|z_n\|^2 - \frac{q2_s^*}{2_s^* - q} \mathcal{I}_{\lambda,\mu}(z_n).$$

Let $n \rightarrow \infty$ in (6.3), by (6.1), (6.2) and $\alpha_{\lambda,\mu} < 0$, we have

$$Q_{\lambda,\mu}(z^+) \geq -\frac{q2_s^*}{2_s^* - q} \alpha_{\lambda,\mu} > 0.$$

Therefore, $z^+ \in \mathcal{N}_{\lambda,\mu}$ is a nontrivial solution of (2.1). Now we show that $z_n \rightarrow z^+$ strongly in E and $\mathcal{I}_{\lambda,\mu}(z^+) = \alpha_{\lambda,\mu}$. Since $z^+ \in \mathcal{N}_{\lambda,\mu}$, then by (6.3), we obtain

$$\begin{aligned} \alpha_{\lambda,\mu} &\leq \mathcal{I}_{\lambda,\mu}(z^+) \\ &= \frac{s}{N} \|z^+\|^2 - \frac{2_s^* - q}{q2_s^*} Q_{\lambda,\mu}(z^+) \\ &\leq \lim_{n \rightarrow \infty} \left(\frac{s}{N} \|z_n\|^2 - \frac{2_s^* - q}{q2_s^*} Q_{\lambda,\mu}(z_n) \right) \\ &= \lim_{n \rightarrow \infty} \mathcal{I}_{\lambda,\mu}(z_n) = \alpha_{\lambda,\mu}. \end{aligned}$$

This implies that $\mathcal{I}_{\lambda,\mu}(z^+) = \alpha_{\lambda,\mu}$ and $\lim_{n \rightarrow \infty} \|z_n\|^2 = \|z^+\|^2$. Set $\widehat{z}_n = z_n - z^+$. Then, that

$$\|\widehat{z}_n\|^2 = \|z_n\|^2 - \|z^+\|^2 + o_n(1).$$

Hence, $z_n \rightarrow z^+$ in E . We claim that $z^+ \in \mathcal{N}_{\lambda,\mu}^+$. Assume by contradiction that $z^+ \in \mathcal{N}_{\lambda,\mu}^-$. Then, by Theorem 4.5, there exist (unique) t_1^+ and t_1^- with $t_1^+ z^+ \in \mathcal{N}_{\lambda,\mu}^+$ and $t_1^- z^+ \in \mathcal{N}_{\lambda,\mu}^-$. In particular, we have $t_1^+ < t_1^- = 1$. Since

$$\frac{d}{dt} \mathcal{I}_{\lambda,\mu}(tz^+) \Big|_{t=t_1^+} = 0, \quad \text{and} \quad \frac{d^2}{dt^2} \mathcal{I}_{\lambda,\mu}(tz^+) \Big|_{t=t_1^+} > 0,$$

there exists $t_1^+ < t^* \leq t_1^-$ such that $\mathcal{I}_{\lambda,\mu}(t_1^+ z^+) < \mathcal{I}_{\lambda,\mu}(t^* z^+)$. By Theorem 4.5, we have

$$\mathcal{I}_{\lambda,\mu}(t_1^+ z^+) < \mathcal{I}_{\lambda,\mu}(t^* z^+) \leq \mathcal{I}_{\lambda,\mu}(t_1^- z^+) = \mathcal{I}_{\lambda,\mu}(z^+),$$

a contradiction. Since $\mathcal{I}_{\lambda,\mu}(z^+) = \mathcal{I}_{\lambda,\mu}(|w_1^+|, |w_2^+|)$ and $(|w_1^+|, |w_2^+|) \in \mathcal{N}_{\lambda,\mu}$, by Lemma 4.2 we may assume that z^+ is a nontrivial nonnegative solution of (2.1). Then by the Strong Maximum Principle [10, Lemma 2.4], we have $w_1^+, w_2^+ > 0$ in \mathcal{C}_Ω , hence, z^+ is a positive solution for (2.1). \square

Next we will use $w_\varepsilon = E_s(u_\varepsilon)$, the family of minimizers for the trace inequality (2.4), where u_ε is given in (2.5). Without loss of generality, we may assume that $0 \in \Omega$. We then define the cut-off function $\phi \in C_0^\infty(\mathcal{C}_\Omega)$, $0 \leq \phi \leq 1$ and for small fixed $\rho > 0$,

$$\phi(x, y) = \begin{cases} 1, & (x, y) \in B_\rho, \\ 0, & (x, y) \notin \overline{B_{2\rho}}, \end{cases}$$

where $B_\rho = \{(x, y) : |x|^2 + y^2 < \rho^2, y > 0\}$. We take ρ so small that $\overline{B_{2\rho}} \subset \overline{\mathcal{C}_\Omega}$. Recall \mathcal{W} is the extension of U introduced in Section 2, we have (cf. [3]) $|\nabla \mathcal{W}(x, y)| \leq Cy^{-1}\mathcal{W}(x, y)$. Let

$$U_\varepsilon(x) = \frac{1}{(\varepsilon^2 + |x|^2)^{\frac{N-2s}{2}}}, \quad \varepsilon > 0.$$

Then the extension of $U_\varepsilon(x)$ has the form

$$\mathcal{W}_\varepsilon(x, y) = c_{N,s} y^{2s} \int_{\mathbb{R}^N} \frac{U_\varepsilon(z) dz}{(|x-z|^2 + y^2)^{\frac{N+2s}{2}}} = \varepsilon^{2s-N} \mathcal{W}\left(\frac{x}{\varepsilon}, \frac{y}{\varepsilon}\right).$$

Notice that $\phi \mathcal{W}_\varepsilon \in X_0^s(\mathcal{C}_\Omega)$, for $\varepsilon > 0$ small enough.

Lemma 6.2. *There is $z \in E \setminus \{0\}$ nonnegative and $\Lambda^* > 0$ such that for $(\lambda, \mu) \in \mathcal{C}_{\Lambda^*}$*

$$\sup_{t \geq 0} \mathcal{I}_{\lambda, \mu}(tz) < c_\infty,$$

where c_∞ is given in Lemma 3.3. In particular, $\alpha_{\lambda, \mu}^- < c_\infty$ for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda^*}$.

Proof. By an argument similar to that of the proof of [3, formula (3.26)], we get

$$(6.4) \quad \begin{aligned} \|\phi \mathcal{W}_\varepsilon\|_{X_0^s}^2 &= k_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \mathcal{W}_\varepsilon|^2 dx dy + \mathcal{O}(1) \\ &= \varepsilon^{2s-N} k_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \mathcal{W}(x, y)|^2 dx dy + \mathcal{O}(1). \end{aligned}$$

We notice that

$$\begin{aligned} \|\phi U_\varepsilon\|_{2_s^*}^{2_s^*} &= \int_{\Omega} |\phi U_\varepsilon|^{2_s^*} dx = \int_{\Omega} \frac{\phi(x)^{2_s^*}}{(\varepsilon^2 + |x|^2)^N} dx, \\ \|U_\varepsilon\|_{2_s^*}^{2_s^*} &= \int_{\mathbb{R}^N} \frac{1}{(\varepsilon^2 + |x|^2)^N} dx = \varepsilon^{-N} \|U\|_{2_s^*}^{2_s^*}. \end{aligned}$$

Then, one has that

$$\|\phi U_\varepsilon\|_{2_s^*}^{2_s^*} - \varepsilon^{-N} \|U\|_{2_s^*}^{2_s^*} = \int_{\Omega} \frac{\phi^{2_s^*}(x) - 1}{(\varepsilon^2 + |x|^2)^N} dx - \int_{\mathbb{R}^N \setminus \Omega} \frac{dx}{(\varepsilon^2 + |x|^2)^N},$$

which yields

$$\begin{aligned} \left| \|\phi U_\varepsilon\|_{2_s^*}^{2_s^*} - \varepsilon^{-N} \|U\|_{2_s^*}^{2_s^*} \right| &\leq \int_{\Omega \setminus B(0; \rho)} \frac{1}{(\varepsilon^2 + |x|^2)^N} dx + \int_{\mathbb{R}^N \setminus \Omega} \frac{dx}{(\varepsilon^2 + |x|^2)^N} \\ &= \int_{\mathbb{R}^N \setminus B(0; \rho)} \frac{dx}{(\varepsilon^2 + |x|^2)^N} \leq \int_{\mathbb{R}^N \setminus B(0; \rho)} \frac{dx}{|x|^{2N}} = C_3. \end{aligned}$$

This implies that

$$1 - C_3 \varepsilon^N \|U\|_{2_s^*}^{-2_s^*} \leq \varepsilon^N \|\phi U_\varepsilon\|_{2_s^*}^{2_s^*} \|U\|_{2_s^*}^{-2_s^*} \leq 1 + C_3 \varepsilon^N \|U\|_{2_s^*}^{-2_s^*}.$$

Taking ε so small that $C_2\varepsilon^N\|U\|_{2_s^*}^{-2_s^*} < 1$, since $2/2_s^* = (N-2s)/N < 1$, we obtain

$$\begin{aligned} 1 - \varepsilon^N C_3 \|U\|_{2_s^*}^{-2_s^*} &\leq (1 - \varepsilon^N C_3 \|U\|_{2_s^*}^{-2_s^*})^{2/2_s^*} \leq \varepsilon^{N-2s} \|\phi U_\varepsilon\|_{2_s^*}^2 \|U\|_{2_s^*}^{-2} \\ &\leq (1 + \varepsilon^N C_3 \|U\|_{2_s^*}^{-2_s^*})^{2/2_s^*} \leq 1 + \varepsilon^N C_3 \|U\|_{2_s^*}^{-2_s^*}. \end{aligned}$$

Hence $\|\phi U_\varepsilon\|_{2_s^*}^2 = \varepsilon^{2s-N} \|U\|_{2_s^*}^2 + \mathcal{O}(\varepsilon^{2s})$. Since $\mathcal{W} = E_s(U)$ optimizes (2.4), by (6.4) we have

$$\begin{aligned} (6.5) \quad \frac{\|\phi \mathcal{W}_\varepsilon\|_{X_0^s}^2}{\|\phi U_\varepsilon\|_{2_s^*}^2} &= \frac{\varepsilon^{2s-N} k_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \mathcal{W}(x, y)|^2 dx dy + \mathcal{O}(1)}{\varepsilon^{2s-N} \|U\|_{2_s^*}^2 + \mathcal{O}(\varepsilon^{2s})} \\ &= \frac{k_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \mathcal{W}(x, y)|^2 dx dy}{\|U\|_{2_s^*}^2} \left(1 + \mathcal{O}(\varepsilon^{N-2s})\right) \\ &= k_s \mathcal{S}(s, N) + \mathcal{O}(\varepsilon^{N-2s}). \end{aligned}$$

Now we consider the function $J : E \rightarrow \mathbb{R}$ defined by $J(z) := 1/2 \|z\|^2 - 2/2_s^* \int_\Omega |w_1|^\alpha |w_2|^\beta dx$. Set $w_{0,1} := \sqrt{\alpha} \phi \mathcal{W}_\varepsilon$, $w_{0,2} := \sqrt{\beta} \phi \mathcal{W}_\varepsilon$ and $z_0 := (w_{0,1}, w_{0,2}) \in E$. Notice that $J(0) = 0$, $J(tz_0) > 0$ for $t > 0$ small and $J(tz_0) < 0$ for $t > 0$ large. The map $t \mapsto J(tz_0)$ maximizes at

$$(6.6) \quad t_0 := \left(\frac{\|z_0\|^2}{2 \int_\Omega |w_{0,1}|^\alpha |w_{0,2}|^\beta dx} \right)^{\frac{1}{2_s^*-2}}.$$

Then from (2.7), (6.5) and (6.6), we conclude that

$$\begin{aligned} (6.7) \quad \sup_{t \geq 0} J(tz_0) &= J(t_0 z_0) = \left(\frac{1}{2} - \frac{1}{2_s^*} \right) \frac{\|z_0\|_{2_s^*}^{\frac{22_s^*}{2_s^*-2}}}{\left(2 \int_\Omega |w_{0,1}|^\alpha |w_{0,2}|^\beta dx \right)^{\frac{2}{2_s^*-2}}} \\ &= \frac{s}{N} \left[\frac{(\alpha + \beta) k_s \int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla(\phi \mathcal{W}_\varepsilon)|^2 dx dy}{\left(\alpha^{\frac{\alpha}{2}} \beta^{\frac{\beta}{2}} \int_\Omega |\phi U_\varepsilon|_{2_s^*}^2 dx \right)^{\frac{2}{2_s^*}}} \right]^{\frac{2_s^*}{2_s^*-2}} \cdot \frac{1}{2^{\frac{N-2s}{2s}}} \\ &= \frac{s}{N 2^{\frac{N-2s}{2s}}} \left[\left(\frac{\alpha}{\beta} \right)^{\frac{\beta}{\alpha+\beta}} + \left(\frac{\beta}{\alpha} \right)^{\frac{\alpha}{\alpha+\beta}} \right]^{\frac{N}{2s}} \left[\frac{k_s \int_{\mathcal{C}_\Omega} y^{1-2s} |\nabla(\phi \mathcal{W}_\varepsilon)|^2}{\left(\int_\Omega |\phi U_\varepsilon|_{2_s^*}^2 dx \right)^{\frac{2}{2_s^*}}} \right]^{\frac{N}{2s}} \\ &= \frac{s}{N 2^{\frac{N-2s}{2s}}} \left[\left(\frac{\alpha}{\beta} \right)^{\frac{\beta}{\alpha+\beta}} + \left(\frac{\beta}{\alpha} \right)^{\frac{\alpha}{\alpha+\beta}} \right]^{\frac{N}{2s}} \left[k_s \mathcal{S}(s, N) + \mathcal{O}(\varepsilon^{N-2s}) \right]^{\frac{N}{2s}} \\ &= \frac{s}{N 2^{\frac{N-2s}{2s}}} \left[k_s \mathcal{S}_{s, \alpha, \beta} + \mathcal{O}(\varepsilon^{N-2s}) \right]^{\frac{N}{2s}} \\ &= \frac{s}{N} \frac{1}{2^{\frac{N-2s}{2s}}} (k_s \mathcal{S}_{s, \alpha, \beta})^{\frac{N}{2s}} + \mathcal{O}(\varepsilon^{N-2s}) = \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s, \alpha, \beta}}{2} \right)^{\frac{N}{2s}} + \mathcal{O}(\varepsilon^{N-2s}). \end{aligned}$$

We now choose $\delta_1 > 0$ so small that, for all $(\lambda, \mu) \in \mathcal{C}_{\delta_1}$, we get

$$c_\infty = \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s,\alpha,\beta}}{2} \right)^{\frac{N}{2s}} - K_0 (\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}}) > 0.$$

By the definition of $\mathcal{I}_{\lambda,\mu}$ and z_0 , we have

$$\mathcal{I}_{\lambda,\mu}(tz_0) \leq \frac{t^2}{2} \|z_0\|^2, \quad \text{for all } t \geq 0 \text{ and } \lambda, \mu > 0,$$

which implies that there exists $t_0 \in (0, 1)$ satisfying

$$\sup_{t \in [0, t_0]} \mathcal{I}_{\lambda,\mu}(tz_0) < c_\infty, \quad \text{for all } (\lambda, \mu) \in \mathcal{C}_{\delta_1}.$$

Hence, from (6.7) and $\alpha, \beta > 1$ we see that

$$\begin{aligned} (6.8) \quad & \sup_{t \geq t_0} \mathcal{I}_{\lambda,\mu}(tz_0) \\ &= \sup_{t \geq t_0} \left(J(tz_0) - \frac{t^q}{q} Q_{\lambda,\mu}(tz_0) \right) \\ &\leq \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s,\alpha,\beta}}{2} \right)^{\frac{N}{2s}} + \mathcal{O}(\varepsilon^{N-2s}) - \frac{t_0^q}{q} \left(\lambda \alpha^{\frac{q}{2}} + \mu \beta^{\frac{q}{2}} \right) \int_{B(0;\rho)} |U_\varepsilon|^q dx \\ &\leq \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s,\alpha,\beta}}{2} \right)^{\frac{N}{2s}} + \mathcal{O}(\varepsilon^{N-2s}) - \frac{t_0^q}{q} (\lambda + \mu) \int_{B(0;\rho)} |U_\varepsilon|^q dx. \end{aligned}$$

Letting $0 < \varepsilon \leq \rho$, we have

$$\int_{B(0;\rho)} |U_\varepsilon|^q dx = \int_{B(0;\rho)} \frac{1}{(\varepsilon^2 + |x|^2)^{\frac{q(N-2s)}{2}}} dx \geq \int_{B(0;\rho)} \frac{1}{(2\rho^2)^{\frac{q(N-2s)}{2}}} dx = C_4,$$

for some $C_4 = C_4(N, s, \rho)$. Combining this with (6.8), for $\varepsilon = (\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}})^{\frac{1}{N-2s}} < \rho$,

$$\sup_{t \geq t_0} \mathcal{I}_{\lambda,\mu}(tz_0) \leq \frac{2s}{N} \left(\frac{k_s \mathcal{S}_{s,\alpha,\beta}}{2} \right)^{\frac{N}{2s}} + \mathcal{O} \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right) - \frac{t_0^q}{q} (\lambda + \mu) C_4.$$

Choosing $\delta_2 > 0$ small enough, for all $(\lambda, \mu) \in \mathcal{C}_{\delta_2}$, we have

$$\mathcal{O} \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right) - \frac{t_0^q}{q} (\lambda + \mu) C_4 < -K_0 \left(\lambda^{\frac{2}{2-q}} + \mu^{\frac{2}{2-q}} \right).$$

If we set $\Lambda^* = \min\{\delta_1, \rho^{N-2s}, \delta_2\} > 0$, then for $(\lambda, \mu) \in \mathcal{C}_{\Lambda^*}$,

$$(6.9) \quad \sup_{t \geq 0} \mathcal{I}_{\lambda,\mu}(tz_0) < c_\infty.$$

Finally, we prove that $\alpha_{\lambda,\mu}^- < c_\infty$ for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda^*}$. Recall that

$$z_0 = (w_{0,1}, w_{0,2}) = (\sqrt{\alpha} \phi \mathcal{W}_\varepsilon, \sqrt{\beta} \phi \mathcal{W}_\varepsilon).$$

Since $\int_\Omega |w_{0,1}|^\alpha |w_{0,2}|^\beta dx > 0$, by Theorem 4.5 there exists $t_0 > 0$ such that $t_0 z_0 \in \mathcal{N}_{\lambda,\mu}^-$. By the definition of $\alpha_{\lambda,\mu}^-$ and (6.9), we conclude that

$$\alpha_{\lambda,\mu}^- \leq \mathcal{I}_{\lambda,\mu}(t_0 z_0) \leq \sup_{t \geq 0} \mathcal{I}_{\lambda,\mu}(tz_0) < c_\infty$$

for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda^*}$. □

Let Λ^* be as in Lemma 6.2. We prove the existence a local minimizer for $\mathcal{I}_{\lambda,\mu}$ on $\mathcal{N}_{\lambda,\mu}^-$.

Proposition 6.3. *Let $\Lambda^* > 0$ be as in Lemma 6.2 and set*

$$\Lambda_2 := \min\{\Lambda^*, (q/2)^{\frac{2}{2-q}} \Lambda_1\}.$$

For $(\lambda, \mu) \in \mathcal{C}_{\Lambda_2}$, $\mathcal{I}_{\lambda,\mu}$ has a minimizer z^- in $\mathcal{N}_{\lambda,\mu}^-$ with $\mathcal{I}_{\lambda,\mu}(z^-) = \alpha_{\lambda,\mu}^-$. Furthermore, z^- is a positive solution of (2.1).

Proof. By (ii) of Proposition 5.3, there is a $(PS)_{\alpha_{\lambda,\mu}^-}$ sequence $\{z_n\} \subset \mathcal{N}_{\lambda,\mu}^-$ for $\mathcal{I}_{\lambda,\mu}$ for all

$$(\lambda, \mu) \in \mathcal{C}_{(q/2)^{2/(2-q)} \Lambda_1}.$$

By Lemmas 3.3 and 6.2 and (ii) of Theorem 4.4, for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda^*}$, $\mathcal{I}_{\lambda,\mu}$ satisfies the $(PS)_{\alpha_{\lambda,\mu}^-}$ condition and $\alpha_{\lambda,\mu}^- > 0$. Then, there exists a subsequence still denoted by $\{z_n\}$ and $z^- = (w_1^-(x, y), w_2^-(x, y)) \in \mathcal{N}_{\lambda,\mu}^-$ such that $z_n \rightarrow z^-$ strongly in E and $\mathcal{I}_{\lambda,\mu}(z^-) = \alpha_{\lambda,\mu}^- > 0$, for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda_2}$. Arguing as in the proof of Proposition 6.1, for $(\lambda, \mu) \in \mathcal{C}_{\Lambda_2}$, we obtain that z^- is a positive solution of (2.1). \square

7. PROOF OF THE MAIN RESULT CONCLUDED

By Proposition 6.1, for $(\lambda, \mu) \in \mathcal{C}_{\Lambda_1}$, system (2.1) admits a positive solution $z^+ \in \mathcal{N}_{\lambda,\mu}^+$. By Proposition 6.3, a positive solution $z^- \in \mathcal{N}_{\lambda,\mu}^-$ exists for all $(\lambda, \mu) \in \mathcal{C}_{\Lambda_2}$. Furthermore, since $\mathcal{N}_{\lambda,\mu}^+ \cap \mathcal{N}_{\lambda,\mu}^- = \emptyset$, then z^+, z^- are distinct positive solutions of system (2.1). In turn, $(u^\pm(x), v^\pm(x)) = (w_1^\pm(x, 0), w_2^\pm(x, 0))$ are distinct positive solutions of (1.1). \square

REFERENCES

- [1] C.O. Alves, D.C. de Morais Filho, M.A.S. Souto, On systems of elliptic equations involving subcritical or critical Sobolev exponents, *Nonlinear Anal.* **42** (2000), 771–787. [2](#), [6](#)
- [2] A. Ambrosetti, H. Brezis, G. Cerami, Combined effects of concave-convex nonlinearities in some elliptic problems, *J. Funct. Anal.* **122** (1994), 519–543. [2](#)
- [3] B. Barrios, E. Colorado, A. de Pablo, U. Sánchez, On some critical problems for the fractional Laplacian, *J. Differential Equations* **252** (2012), 6133–6162. [2](#), [3](#), [4](#), [19](#)
- [4] B. Barrios, E. Colorado, R. Servadei, F. Soria, A critical fractional equation with concave-convex power nonlinearities, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **32** (2015), 875–900. [2](#), [4](#)
- [5] C. Brändle, E. Colorado, A. de Pablo, A concave-convex elliptic problem involving the fractional Laplacian, *Proc. Roy. Soc. Edinburgh Sect. A. Math.* **142** (2013), 39–71. [2](#), [4](#), [5](#)
- [6] H. Brézis, L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.* **36** (1983), 437–477. [2](#)
- [7] K.J. Brown, Y. Zhang, The Nehari manifold for a semilinear elliptic equation with a sign-changing weight function, *J. Differential Equations* **193** (2003), 481–499. [11](#)
- [8] X. Cabré, J. Tan, Positive solutions of nonlinear problems involving the square root of the Laplacian, *Adv. Math.* **224** (2010), 2052–2093. [2](#)
- [9] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian, *Comm. Partial Differential Equations* **32** (2007), 1245–1260. [2](#)
- [10] A. Capella, J. Dávila, L. Dupaigne, Y. Sire, Regularity of radial extremal solutions for some non-local semilinear equations, *Comm. Partial Differential Equations* **36** (2011), 1353–1384. [4](#), [18](#)
- [11] W. Chen, S. Deng, The Nehari manifold for a nonlinear elliptic operators involving concave-convex nonlinearities, *Z. Angew. Math. Phys.* **66**, (2015), 1387–1400. [2](#)
- [12] W. Chen, C. Li, B. Ou, Classification of solutions for an integral equation, *Comm. Pure Appl. Math.* **59** (2006), 330–343. [2](#)

- [13] X. Cheng, S. Ma, Existence of three nontrivial solutions for elliptic systems with critical exponents and weights, *Nonlinear Anal.* **69** (2008), 3537–3548. [2](#)
- [14] E. Colorado, A. de Pablo, U. Sánchez, Perturbation of a critical fractional equations, *Pacific J. Math.* **271** (2014), 65–85. [2](#)
- [15] A. Cotsoioli, N. Tavoularis, Best constant for Sobolev inequalities for higher order fractional derivatives, *J. Math. Anal. Appl.* **295** (2004), 225–236. [2](#)
- [16] P. Drabek, S.I. Pohozaev, Positive solutions for the p -Laplacian: application of the fibering methods, *Proc. Roy. Soc. Edinburgh Sect. A* **127** (1997), 703–726. [11](#)
- [17] I. Ekeland, On the variational principle, *J. Math. Anal. Appl.* **47** (1974), 324–353. [16](#)
- [18] L.F.O. Faria, O.H. Miyagaki, F.R. Pereira, M. Squassina, C. Zhang, The Brezis-Nirenberg problem for nonlocal systems, *Adv. Nonlinear Anal.*, to appear. [2](#), [6](#)
- [19] P. Han, The effect of the domain topology of the number of positive solutions of elliptic systems involving critical Sobolev exponents, *Houston J. Math.* **32** (2006), 1241–1257. [2](#), [10](#)
- [20] T. Hsu, H. Lin, Multiple positive solutions for a critical elliptic system with concave-convex nonlinearities, *Proc. Roy. Soc. Edinburgh Sect. A* **139** (2009), 1163–1177. [2](#)
- [21] E. Di Nezza, G. Palatucci, E. Valdinoci, Hithiker’s guide to the fractional Sobolev spaces, *Bull. Sci. Math.* **136** (2012), 521–573. [2](#)
- [22] X. Shang, J. Zhang, Y. Yang, Positive solutions of nonhomogeneous fractional Laplacian problem with critical exponent, *Comm. Pure Appl. Anal.* **13** (2014), 567–584. [2](#)
- [23] R. Servadei, E. Valdinoci, Mountain pass solutions for nonlinear elliptic operators, *J. Math. Anal. Appl.* **389** (2012), 887–898. [2](#)
- [24] R. Servadei, E. Valdinoci, A Brezis-Nirenberg result for non-local critical equations in lower dimension, *Comm. Pure Appl. Anal.* **12** (2013), 2445–2464. [2](#)
- [25] R. Servadei, E. Valdinoci, Lewy-Stampacchia type estimates for variational inequalities driven by (non) local operators, *Rev. Mat. Iberoamericana* **29** (2013), 1091–1126. [2](#)
- [26] R. Servadei, E. Valdinoci, The Brezis-Nirenberg result for the fractional Laplacian, *Trans. Amer. Math. Soc.* **367** (2015), 67–102. [2](#)
- [27] R. Servadei, E. Valdinoci, On the spectrum of two different fractional operators, *Proc. Roy. Soc. Edinburgh Sect. A* **144** (2014), 831–855. [2](#)
- [28] J. Serra, X. Ros-Oton, The Dirichlet problem for the fractional Laplacian: regularity up to the boundary, *J. Math. Pures Appl.* **101** (2014), 275–302. [2](#)
- [29] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, *Comm. Pure Appl. Math.* **60** (2007), 67–112. [2](#)
- [30] J. Tan, The Brézis-Nirenberg type problem involving the square root of the Laplacian, *Calc. Var. Partial Differential Equations* **36** (2011), 21–41. [2](#)
- [31] Y. Wei, X. Su, Multiplicity of solutions for non-local elliptic equations driven by the fractional Laplacian, *Calc. Var. Partial Differential Equations* **52** (2015), 95–124. [2](#)
- [32] T.F. Wu, On semilinear elliptic equations involving concave-convex nonlinearities and sign-changing weight function, *J. Math. Anal. Appl.* **318** (2006), 253–270. [2](#), [11](#)
- [33] T.F. Wu, The Nehari manifold for a semilinear elliptic system involving sign-changing weight functions, *Nonlinear Anal.* **68** (2008), 1733–1745. [2](#)
- [34] X. Yu, The Nehari manifold for elliptic equation involving the square root of the laplacian, *J. Differential Equations* **252** (2012), 1283–1308. [2](#)

(X. He) COLLEGE OF SCIENCE, MINZU UNIVERSITY OF CHINA
 BEIJING 100081, CHINA
E-mail address: xmhe923@muc.edu.cn

(M. Squassina) DIPARTIMENTO DI INFORMATICA, UNIVERSITÀ DEGLI STUDI DI VERONA
 STRADA LE GRAZIE 15, I-37134 VERONA, ITALY
E-mail address: marco.squassina@univr.it

(W. Zou) DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY
 BEIJING 100084, CHINA
E-mail address: wzou@math.tsinghua.edu.cn