

# DIRICHLET PROBLEMS FOR SINGULAR ELLIPTIC EQUATIONS WITH GENERAL NONLINEARITIES

VIRGINIA DE CICCO, DANIELA GIACHETTI, FRANCESCANTONIO OLIVA, AND FRANCESCO PETITTA

ABSTRACT. In this paper, under very general assumptions, we prove existence and regularity of distributional solutions to homogeneous Dirichlet problems of the form

$$\begin{cases} -\Delta_1 u = h(u)f & \text{in } \Omega, \\ u \geq 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where,  $\Delta_1$  is the 1-laplace operator,  $\Omega$  is a bounded open subset of  $\mathbb{R}^N$  with Lipschitz boundary,  $h(s)$  is a continuous function which may become singular at  $s = 0^+$ , and  $f$  is a nonnegative datum in  $L^{N,\infty}(\Omega)$  with suitable small norm. Uniqueness of solutions is also shown provided  $h$  is decreasing and  $f > 0$ . As a by-product of our method a general theory for the same problem involving the  $p$ -laplacian as principal part, which is missed in the literature, is established. Most of the assumptions are further discussed in order to show their optimality.

## CONTENTS

|   |    |
|---|----|
| 1. Introduction   | 1  |
| 2. Notations and Preliminaries  | 4  |
| 2.1. Notations  | 6  |
| 3. Main assumptions and core results  | 6  |
| 4. Existence of a solution for $p > 1$ and for a nonnegative $f \in L^{(p^*)'}(\Omega)$ | 8  |
| 4.1. A priori estimates   | 9  |
| 4.2. Proof of Theorem 4.3   | 10 |
| 4.3. More regular data $f \in L^m(\Omega)$ with $(p^*)' \leq m < N$                     | 12 |
| 4.4. Uniform estimates in the case $f \in L^N(\Omega)$                                  | 14 |
| 5. The limit as $p \rightarrow 1^+$ for a positive $f$                                  | 17 |
| 6. Nonnegative data $f$   | 23 |
| 7. The case $f$ in $L^{N,\infty}(\Omega)$   | 26 |
| 8. Further extensions, remarks, and examples  | 27 |
| 8.1. Some global $BV$ solutions   | 27 |
| 8.2. More general growths   | 29 |
| References  | 31 |

## 1. INTRODUCTION

The main goal of this paper is to deal with existence, regularity (and uniqueness, if attainable) of distributional solutions to problems involving 1-laplacian as a principal part and lower order terms which can become singular on the set where the solution  $u$  vanishes. Formally the problem looks

---

2010 *Mathematics Subject Classification.* 35J60, 35J75, 34B16, 35R99, 35A02.

*Key words and phrases.* 1-Laplacian, p-Laplacian, Nonlinear elliptic equations, Singular elliptic equations.

like

$$\begin{cases} -\Delta_1 u := -\operatorname{div}\left(\frac{Du}{|Du|}\right) = h(u)f & \text{in } \Omega, \\ u \geq 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where  $\Omega$  is a bounded open subset of  $\mathbb{R}^N$  with Lipschitz boundary,  $0 \leq f \in L^{N,\infty}(\Omega)$ ,  $h(s)$  is a non-negative continuous function defined in  $[0, \infty)$ , bounded at infinity, possibly singular at  $s = 0$  (i.e.  $h(0) = \infty$ ) without any monotonicity property.

The natural space for this kind of problems is  $BV$  (or its local version  $BV_{\text{loc}}$ ), the space of functions of bounded variation, i.e. the space of  $L^1$  functions whose gradient is a Radon measure with finite (or locally finite) total variation. In (1.1)  $\frac{Du}{|Du|}$  is the Radon-Nikodym derivative of the measure  $Du$  with respect to its total variation  $|Du|$ .

Problems involving the 1-laplace operator, which is known to be closely related to the mean curvature operator ([37]), enter in a variety of both practical and theoretical issues as for instance in image restoration and in torsion problems ([26, 27, 39, 30, 9]). We refer the interested reader to the monograph [6] for a more complete review on applications.

From the mathematical point of view, in order to give sense to the left hand side term of the equation in (1.1), Andreu, Ballester, Caselles and Mazón ([4]) proposed to use the Anzellotti theory of pairings  $(z, Du)$  of  $L^\infty$ -divergence-measure vector fields  $z$  and the gradient of a BV-function  $u$ . In their definition the vector field  $z$  belongs to  $\mathcal{DM}^\infty(\Omega)$  (the space of  $L^\infty$  vector fields whose distributional divergence is a Radon measure with bounded total variation), and is such that both  $\|z\|_\infty \leq 1$  and  $(z, Du) = |Du|$ ; in this way the vector  $z$  is intended to play the role of the ratio  $\frac{Du}{|Du|}$ .

Problem (1.1) has been studied by several authors when  $h \equiv 1$  (see [16, 28, 31] and references therein) under suitable smallness assumption on the datum  $f$ . The non-autonomous case has also been treated; the eigenvalue problem is discussed in [28], the absorption case is handled in [34], while the (sub-)critical exponent problem has been also addressed (see for instance [22, 33]). Finally, some early results can be found in [21] concerning the mild singular model case  $h(s) = s^{-\gamma}$ ,  $0 < \gamma \leq 1$ . As one of our main point also concerns the singular case of a nonlinearity  $h(s)$  which is unbounded near the origin  $s = 0$ , it is expected that, if  $f$  vanishes in a portion of  $\Omega$  of positive Lebesgue measure, then also the region where  $u$  degenerates at zero plays a nontrivial role; in this case, in fact, the characteristic function  $\chi_{\{u>0\}}$  may appear in the definition of solution of problem (1.1) (see Definition 6.1 in Section 6 below).

To be more transparent and to fix the ideas, we assume, at first, that  $f$  is a strictly positive function in  $L^N(\Omega)$ ; most of the main issues are already present in this less general case in which, although, some further properties for solutions to (1.1) can be proven and these will ease the presentation. In this case, for instance,  $h(u)f \in L^1(\Omega)$  and uniqueness holds, in a suitable class of functions, if  $h$  is decreasing.

Let us explain the meaning of solution in this particular non-degenerate case. If  $0 < f \in L^N(\Omega)$ , a function  $u \in BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$  will be a distributional solution to problem (1.1) if there exists  $z \in \mathcal{DM}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  such that  $-\operatorname{div} z = h(u)f$  in  $\mathcal{D}'(\Omega)$ ,  $(z, Du) = |Du|$  as measures in  $\Omega$ , and one of the following conditions holds:

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \cap B(x, \epsilon)} u(y) dy = 0 \quad \text{or} \quad [z, \nu](x) = -1 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega. \quad (1.2)$$

Condition (1.2) is a (very weak) way to give sense to the boundary condition  $u = 0$  at  $\partial\Omega$  in this case where  $u$  is only in  $BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$ , and where  $[z, \nu]$  is the weak normal trace of  $z$  defined in [7] (see Section 2 below).

In order to describe the core of our results, we will also assume that  $h : [0, \infty) \rightarrow [0, \infty]$  is a continuous function, such that  $h(0) \neq 0$ ,

$$\exists c_1, \gamma, k_0 > 0 \quad \text{such that} \quad h(s) \leq \frac{c_1}{s^\gamma} \quad \text{if } s < k_0,$$

and

$$\lim_{s \rightarrow \infty} h(s) := h(\infty) < \infty.$$

More general right hand sides (for (1.1))  $F(x, u)$  are considered in Section 8.2 (see Theorem 8.6) in order to include more general growth as, for instance,

$$h(s) \sim \exp\left(\exp\left(\frac{1}{s}\right)\right),$$

as well as the general non-autonomous case.

As far as the right hand side is then concerned in (1.1), in order to get existence, we require a smallness condition on the datum that  $0 < f \in L^N(\Omega)$  depending on the behavior of the function  $h$  at infinity, i.e.

$$\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}, \quad (1.3)$$

where  $\mathcal{S}_1$  is the best constant in the Sobolev inequality for functions in  $W_0^{1,1}(\Omega)$ . In particular, (1.3) implies that no condition is assumed if  $h(\infty) = 0$ . This allows us to conclude that the behavior of  $h$  at infinity can induce a first regularizing effect on the problem if compared to the nonsingular case (e.g.  $h \equiv 1$ ). In fact, if  $h(\infty)$  is a suitable positive constant then the smallness assumption (1.3) (or its general version for data in  $L^{N,\infty}(\Omega)$ , see Section 7) is necessary and sharp as it can be deduced by comparing with the results in [16] and [31]. A second regularizing effect appears if  $h(0) = \infty$ ; the presence of a singular term will imply that  $u > 0$  in  $\Omega$  that is again in contrast with the nonsingular case; we shall come back on this fact in a while.

We also remark that the equality  $(z, Du) = |Du|$  in the definition of solutions of (1.1) does not depend on  $h$ , which is in some sense natural if we think to the homogeneity of the principal part. However, this requires a formal proof (in particular, to handle the jump part of the pairing) which is based on a result in [17]. Similarly, we are able to prove the weak boundary condition (1.2) independently of the function  $h$ .

As we already mentioned, this case of a strictly positive datum  $f$  enjoys particular features most of them summarized in Theorem 3.4 below. In particular  $h(u)f \in L^1(\Omega)$  and  $u^\sigma \in BV(\Omega)$  with  $\sigma = \max(1, \gamma)$ . These two conditions will allow us to obtain uniqueness of solutions, if we assume that  $h$  is decreasing (see Theorem 3.5). Observe that uniqueness does not hold, in general, for merely nonnegative data (see [21]).

Let us briefly describe the technique we exploit in order to get existence. As noticed in [27], one can naturally handle with (1.1) via approximation with problems having  $p$ -laplacians principal part (with  $p > 1$ ) which in our case can look like

$$\begin{cases} -\Delta_p u_p = h(u_p)f & \text{in } \Omega, \\ u_p \geq 0 & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.4)$$

Note that, at the best of our knowledge, even the existence of solutions  $u_p$  for (1.4) is still mostly unknown under this generality and it requires a preliminary study that we shall present in Section 4 in the more general case of a nonnegative  $f \in L^{(p^*)'}(\Omega)$ . In fact, existence of solutions  $u_p$  of (1.4) has been proven in the model case, i.e.  $h(s) = s^{-\gamma}$ ,  $\gamma > 0$ , ([12, 18, 21]), where, if  $\gamma \leq 1$ ,  $u_p$  is shown to have global finite energy in the sense that  $u_p \in W_0^{1,p}(\Omega)$ , as well as in the case  $p = 2$  (see [11, 24, 36]). While, if  $\gamma > 1$ , solutions  $u_p$  have in general only locally finite energy, that is  $u_p \in W_{loc}^{1,p}(\Omega)$ , so that the boundary datum needs to be assigned through a suitable weaker condition than the usual trace sense (see [11, 18, 19, 25, 35, 36] for further remarks on this fact) eventually producing (1.2) in the limit as  $p \rightarrow 1^+$ . Only recently, existence of solutions for [20] has been shown in case of a general function  $h$  and a measure datum  $f$ .

Note that this approximation approach with respect to the parameter  $p$  requires a priori estimates on the solutions  $u_p$  of (1.4) that are independent of  $p$ , and that will be proven (see Section 4.4) provided (1.3) holds. This will allow us to pass to the limit obtaining a solution to (1.1). As we

already mentioned, if  $h(0) = \infty$  then  $u_p \rightarrow u$  a.e. on  $\Omega$  as  $p \rightarrow 1^+$ , and  $u > 0$ ; this should be compared with the nonsingular case in which, under assumption (1.3),  $u_p \rightarrow 0$  instead (see for instance [16]).

In Section 6 we will show how our results extend to the case of a general nonnegative datum  $f$ . Indeed, if  $f$  can vanish on a subset of  $\Omega$  of positive measure, the situation becomes much more delicate and, as we said, it will involve the region  $\{u > 0\}$  in a substantial way. Among other technical points that will be discussed later, as in [21], we will be forced to ask, in the definition of solutions to (1.4), that  $\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega)$  and that the equation

$$-(\operatorname{div} z)\chi_{\{u>0\}}^* = h(u)f \quad (1.5)$$

is satisfied in the sense of distributions, where  $\chi_{\{u>0\}}$  is the characteristic function of the region  $\{u > 0\}$  and  $\chi_{\{u>0\}}^*$  is its precise representative (in the sense of the  $BV$ -function). This fact will lead to some technical complications in the proof of existence of a solution. Observe that the notion of solution we use here will essentially coincide with the previous if  $f > 0$  (see Remark 6.3 below).

We specify some further peculiarities of this case. First notice that requiring  $\chi_{\{u>0\}}$  to be a locally  $BV$ -function is equivalent to the fact that the region  $\{u > 0\}$  is a set of locally finite perimeter. Also, one may observe that equation (1.5) can also be written as follows (see Remark 6.2 below)

$$-\operatorname{div}(\mathbf{z}\chi_{\{u>0\}}) + |D\chi_{\{u>0\}}| = h(u)f,$$

where the left hand side is a sum of an operator in divergence form and an additional term  $|D\chi_{\{u>0\}}|$ , which is a measure concentrated on the reduced boundary  $\partial^*\{u > 0\}$ . Moreover, in the case where  $f$  is merely nonnegative, we are just able to prove that  $h(u)f \in L^1_{\text{loc}}(\Omega)$  (instead of  $h(u)f \in L^1(\Omega)$  which holds true in the case of positive  $f$ ) and, as we mentioned, no uniqueness of solutions holds (see [21]).

In Section 7 we handle with  $L^{N,\infty}$ -data and here one needs to use the machinery of Lorentz spaces that will be briefly summarized for the convenience of the reader. The extension given in this section is not only technical as it can be shown to be optimal (see Remark 7.2 below).

Finally, in Section 8.1, we will also discuss the possibility to have finite energy solutions, i.e.  $u \in BV(\Omega)$ . As we said, global energy estimates were only known in the mild model case (i.e.  $\gamma \leq 1$ ). Although we get rid of this fact by working with suitable compositions of the solutions, in the general framework of strong singularity the *global*  $BV$  regularity of the solution  $u$  is still an open question; Section 8.1 will be devoted to give some partial answers and insights on this issue. We conclude by investigating the case where a more general right-hand side of the form  $F(x, u)$  is considered (Section 8.2).

## 2. NOTATIONS AND PRELIMINARIES

In the entire paper  $\mathcal{H}^{N-1}(E)$  denotes the  $(N-1)$ -dimensional Hausdorff measure of a set  $E$  while, for simplicity,  $|E|$  will stand for the classical  $N$ -dimensional Lebesgue measure. Here  $\Omega$  is an open bounded subset of  $\mathbb{R}^N$  ( $N \geq 1$ ) with Lipschitz boundary and, in case  $N = 1$ , we will mean  $\frac{N}{N-1} := \infty$ . We will denote by  $\mathcal{M}(\Omega)$  the space of Radon measures with finite total variation over  $\Omega$ .

We denote by

$$\mathcal{D}\mathcal{M}^\infty(\Omega) := \{z \in L^\infty(\Omega; \mathbb{R}^N) : \operatorname{div} z \in \mathcal{M}(\Omega)\},$$

and by  $\mathcal{D}\mathcal{M}^\infty_{\text{loc}}(\Omega)$  the vector fields  $z \in L^\infty(\Omega; \mathbb{R}^N)$  such that  $\operatorname{div} z \in \mathcal{M}(\omega)$ ,  $\forall \omega \subset\subset \Omega$ .

As usual

$$BV(\Omega) := \{u \in L^1(\Omega) : Du \in \mathcal{M}(\Omega, \mathbb{R}^N)\},$$

and its local counterpart, which is the space of functions  $u \in BV(\omega)$  for all  $\omega \subset\subset \Omega$ , is denoted by  $BV_{\text{loc}}(\Omega)$ . We recall that for  $BV(\Omega)$  a norm is given by

$$\|u\|_{BV(\Omega)} = \int_{\Omega} |u| + \int_{\Omega} |Du|,$$

or by

$$\|u\|_{BV(\Omega)} = \int_{\partial\Omega} |u| d\mathcal{H}^{N-1} + \int_{\Omega} |Du|.$$

By  $S_u$  we mean the set of  $x \in \Omega$  such that  $x$  is not a Lebesgue point of  $u$ , by  $J_u$  the jump set and by  $u^*$  the precise representative of  $u$ . For more properties regarding  $BV$  spaces we refer to [3], from which we mainly derive our notations. We also refer to [23, 43].

The theory of  $L^\infty$ -divergence-measure vector fields is due to Anzellotti [7] and to Chen and Frid [15]. We define the following distribution  $(z, Dv) : C_c^1(\Omega) \rightarrow \mathbb{R}$  as

$$\langle (z, Dv), \varphi \rangle := - \int_{\Omega} v^* \varphi \operatorname{div} z - \int_{\Omega} vz \cdot \nabla \varphi \quad \varphi \in C_c^1(\Omega). \quad (2.1)$$

In Anzellotti's theory we need some compatibility conditions, such as  $\operatorname{div} z \in L^1(\Omega)$  and  $v \in BV(\Omega) \cap L^\infty(\Omega)$ , or  $\operatorname{div} z$  a Radon measure with finite total variation and  $v \in BV(\Omega) \cap L^\infty(\Omega) \cap C(\Omega)$ . Anzellotti's definition of  $(z, Dv)$  can be extended to the case in which  $\operatorname{div} z$  is a Radon measure with finite total variation and  $v \in BV(\Omega) \cap L^\infty(\Omega)$  (see [32, Appendix A] and [13, Section 5]). Moreover ([21]), if  $v \in BV_{\text{loc}}(\Omega) \cap L^1(\Omega, \operatorname{div} z)$  and  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$ , the distribution defined in (2.1) is a Radon measure having locally finite total variation satisfying

$$|\langle (z, Dv), \varphi \rangle| \leq \|\varphi\|_{L^\infty(\omega)} \|z\|_{\infty, \omega} \int_{\omega} |Dv|,$$

for all open set  $\omega \subset\subset \Omega$  and for all  $\varphi \in C_c^1(\omega)$ . Here, and throughout the paper, we use the simplified notation  $\|\cdot\|_{q, \omega}$  to indicate the norm of the vector space  $L^q(\omega, \mathbb{R}^N)$ ; also  $\|\cdot\|_q := \|\cdot\|_{q, \Omega}$ .

Moreover the measure  $|(z, Dv)|$  is absolutely continuous with respect to the measure  $|Dv|$  and that, if  $v \in BV(\Omega)$ , then  $(z, Dv)$  has finite total variation.

We have the following proposition proved in [21].

**Proposition 2.1.** *Let  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  and let  $v \in BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$ . Then  $zv \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$ . Moreover the following formula holds in the sense of measures*

$$\operatorname{div}(zv) = (\operatorname{div} z)v^* + (z, Dv).$$

We observe that, since for every  $v \in BV_{\text{loc}}(\Omega)$  the measure  $(z, Dv)$  is absolutely continuous with respect to  $|Dv|$ , it holds

$$(z, Dv) = \theta(z, Dv, x) |Dv|,$$

where  $\theta(z, Dv, \cdot)$  denotes the Radon-Nikodým derivative of  $(z, Dv)$  with respect to  $|Dv|$ . By Proposition 4.5 (iii) of [17], for every  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  and  $u \in BV_{\text{loc}}(\Omega)$ , we have

$$\theta(z, D\Lambda(u), x) = \theta(z, Du, x), \quad \text{for } |D\Lambda(u)|\text{-a.e. } x \in \Omega,$$

where  $\Lambda : \mathbb{R} \rightarrow \mathbb{R}$  is a locally Lipschitz function. We remark that, if  $\Lambda$  is an increasing function, then

$$\theta(z, D\Lambda(u), x) = \theta(z, Du, x), \quad \text{for } |Du|\text{-a.e. } x \in \Omega. \quad (2.2)$$

In fact, by the Chain rule formula (see [3, Theorem 3.99]), we have that

$$D^d \Lambda(u) = \Lambda'(\tilde{u}) D^d u, \quad D^j \Lambda(u) = (\Lambda(u^+) - \Lambda(u^-)) \nu \mathcal{H}^{N-1} \llcorner J_u.$$

and  $(\Lambda(u))^\pm = \Lambda(u^\pm)$ .

The outward normal unit vector  $\nu(x)$  is defined for  $\mathcal{H}^{N-1}$ -almost every  $x \in \partial\Omega$ . It follows from Anzellotti's theory that every  $z \in \mathcal{D}\mathcal{M}^\infty(\Omega)$  has a weak trace on  $\partial\Omega$  of the normal component of  $z$  which is denoted by  $[z, \nu]$ . Moreover, it satisfies

$$\|[z, \nu]\|_{L^\infty(\partial\Omega)} \leq \|z\|_\infty. \quad (2.3)$$

We explicitly point out that if  $z \in \mathcal{D}\mathcal{M}^\infty(\Omega)$  and  $v \in BV(\Omega) \cap L^\infty(\Omega)$ , then

$$v[z, \nu] = [vz, \nu] \quad (2.4)$$

holds (see [13, Lemma 5.6] or [5, Proposition 2]).

The following generalized Green type formula for merely local vector fields  $z$  also holds true ([21]).

**Proposition 2.2.** *Let  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  and set  $\mu = \text{div } z$ . Let  $v \in BV(\Omega) \cap L^\infty(\Omega)$  be such that  $v^* \in L^1(\Omega, \mu)$ . Then  $vz \in \mathcal{D}\mathcal{M}^\infty(\Omega)$  and the following holds:*

$$\int_{\Omega} v^* d\mu + \int_{\Omega} (z, Dv) = \int_{\partial\Omega} [vz, \nu] d\mathcal{H}^{N-1}. \quad (2.5)$$

Analogously to (2.3), it can be proved that, for  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  such that the product  $vz \in \mathcal{D}\mathcal{M}^\infty(\Omega)$  for some  $v \in BV(\Omega) \cap L^\infty(\Omega)$ ,

$$|[vz, \nu]| \leq |v|_{\perp \partial\Omega} \|z\|_{\infty} \quad \mathcal{H}^{N-1}\text{-a.e. on } \partial\Omega.$$

We will finally use the symbol  $\mathcal{S}_p$  to denote the best constant in the Sobolev inequality ( $1 \leq p < N$ ), that is

$$\|v\|_{L^{p^*}(\Omega)} \leq \mathcal{S}_p \|v\|_{W_0^{1,p}(\Omega)}, \quad \forall v \in W_0^{1,p}(\Omega),$$

where  $p^* = \frac{Np}{N-p}$ . Recall that

$$\lim_{p \rightarrow 1^+} \mathcal{S}_p = \mathcal{S}_1 = (N\omega_N^{\frac{1}{N}})^{-1},$$

where  $\omega_N$  is the volume of the unit sphere of  $\mathbb{R}^N$  (see for instance [42]). We conclude by recalling a well known consequence of the Egorov theorem.

**Lemma 2.3.** *Let  $f_n$  be a sequence converging to  $f$  weakly in  $L^1(\Omega)$  and let  $g_n$  be a sequence converging to  $g$  almost everywhere in  $\Omega$  and  $*$ -weakly in  $L^\infty(\Omega)$ . Then*

$$\lim_{n \rightarrow \infty} \int_{\Omega} f_n g_n = \int_{\Omega} f g.$$

**2.1. Notations.** In our arguments we will use several truncating functions. In particular, for a fixed  $k > 0$ , we introduce  $T_k$  and  $G_k$  as the function defined by

$$T_k(s) = \max(-k, \min(s, k)),$$

and

$$G_k(s) = (|s| - k)^+ \text{sign}(s).$$

Note in particular that  $T_k(s) + G_k(s) = s$ , for any  $s \in \mathbb{R}$ .

If no otherwise specified, we denote by  $C$  several constants whose value may change from line to line and, sometimes, on the same line. These values only depend on the data but they do not depend on the indexes of the sequences. We underline the use of the standard convention to do not relabel an extracted compact subsequence.

### 3. MAIN ASSUMPTIONS AND CORE RESULTS

One of the main aims of this paper is to deal with the following problem

$$\begin{cases} -\Delta_1 u = h(u)f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (3.1)$$

where  $f \in L^N(\Omega)$  is positive and  $\Delta_1 u = \text{div} \left( \frac{Du}{|Du|} \right)$  is the 1-Laplace operator. As we already mentioned, the general case of a nonnegative  $f \in L^{N,\infty}(\Omega)$  will be studied in Sections 6 and 7.

On the nonlinearity  $h : [0, \infty) \rightarrow [0, \infty]$  we assume that it is continuous and finite outside the origin,  $h(0) \neq 0$ ,

$$\exists c_1, \gamma, k_0 > 0 \text{ such that } h(s) \leq \frac{c_1}{s^\gamma} \text{ if } s \leq k_0, \quad (h1)$$

and

$$\lim_{s \rightarrow \infty} h(s) := h(\infty) < \infty. \quad (h2)$$

Let us note that the function  $h$  is not necessarily blowing up at the origin so that a bounded continuous function is an admissible choice.

We start providing the definition of solution to problem (3.1).

**Definition 3.1.** Let  $0 < f \in L^N(\Omega)$  then a function  $u \in BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$  is a solution to problem (3.1) if there exists  $z \in \mathcal{D}\mathcal{M}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  such that

$$h(u)f \in L^1_{\text{loc}}(\Omega), \quad (3.2)$$

$$-\operatorname{div} z = h(u)f \quad \text{in } \mathcal{D}'(\Omega), \quad (3.3)$$

$$(z, Du) = |Du| \quad \text{as measures in } \Omega, \quad (3.4)$$

and one of the following conditions holds:

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \cap B(x, \epsilon)} u(y) dy = 0 \quad \text{or} \quad [z, \nu](x) = -1 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega. \quad (3.5)$$

**Remark 3.2.** Notice that Definition 3.1 does not depend explicitly on the parameter  $\gamma$  in (h1), this fact suggesting that an extension to more general nonlinearities is allowed (see Section 8.2 below). Furthermore, condition (3.5) is a way to give meaning to the homogeneous Dirichlet boundary datum. It is well known that  $BV$  solutions to problems involving the 1-Laplace operator do not necessarily assume the boundary datum pointwise. A standard weaker request in this framework is that a solution  $u \in BV(\Omega)$  satisfies

$$u(1 + [z, \nu])(x) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega, \quad (3.6)$$

i.e., for  $\mathcal{H}^{N-1}$ -a.e.  $x \in \partial\Omega$ , it holds that  $u(x) = 0$  or  $[z, \nu](x) = -1$  (see for instance [31, 21]). It is not clear whether problem (3.1) admits, in general, finite energy solutions, that is solutions that are  $BV$  up to the boundary of  $\Omega$ . This fact leads to impose (3.5) which is a weaker assumption than (3.6) for nonnegative functions (see for instance [3, Theorem 3.87]). A similar argument was already exploited in [36] when dealing with infinite energy solutions to similar problems involving laplacian type operators. We refer to Section 8.1 for further instances of solutions belonging to the natural energy space  $BV(\Omega)$  and how this fact can be related to the smoothness of the domain  $\Omega$ .

Here is our main existence result in the case of a positive datum  $f$ :

**Theorem 3.3.** Let  $0 < f \in L^N(\Omega)$  such that  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$ , where  $h$  satisfies (h1) and (h2). Then there exists a solution  $u$  to problem (3.1) in the sense of Definition 3.1.

In the following result we collect some further qualitative properties enjoyed by the solution we found in Theorem 3.3. Here and below, in order to unify the presentation, for  $\gamma > 0$  we set

$$\sigma := \max(1, \gamma). \quad (3.7)$$

**Theorem 3.4.** Let  $0 < f \in L^N(\Omega)$  such that  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$ , where  $h$  satisfies (h1) and (h2). Then the solution  $u$  to problem (3.1) found in Theorem 3.3 is such that

$$u^\sigma \in BV(\Omega). \quad (3.8)$$

Moreover  $\operatorname{div} z \in L^1(\Omega)$  and it holds

$$-\int_{\Omega} v \operatorname{div} z = \int_{\Omega} h(u)fv, \quad \forall v \in BV(\Omega) \cap L^\infty(\Omega). \quad (3.9)$$

Finally if  $h(0) = \infty$  then  $u > 0$  a.e. in  $\Omega$  while, if  $h \in L^\infty([0, \infty))$  and  $\|f\|_{L^N(\Omega)} < (\mathcal{S}_1 \|h\|_{L^\infty([0, \infty))})^{-1}$ , then  $u \equiv 0$  a.e. in  $\Omega$ .

If  $h$  is decreasing, then the solutions obtained in the previous theorems are unique in the sense specified by the following theorem.

**Theorem 3.5.** Let  $h$  be a decreasing function. Then, under the assumptions of Theorem 3.3, there is only one solution  $u \in BV_{\text{loc}}(\Omega)$  to problem (3.1) in the sense of Definition 3.1 such that  $u^\sigma \in BV(\Omega)$ .

**Remark 3.6.** We stress that Theorem 3.3 is new also in the case of a bounded nonlinearity  $h$ . A remark is in order concerning (3.8). First of all, if  $\gamma \leq 1$  then (3.8) says that solutions always belong to the natural space of functions with finite  $BV$  norm, accordingly with the case  $p > 1$  where solutions

with finite  $W_0^{1,p}$  energy are always achieved when the datum  $f$  belongs to  $L^{(p^*)'}(\Omega)$  ([11, 12, 18, 24]). If  $\gamma > 1$  then (3.8) is a sort of counterpart, as  $p$  tends to  $1^+$ , of the regularity for the solutions to the Dirichlet problem associated to  $-\Delta_p v = f v^{-\gamma}$ . In [11, 18, 25] it is shown that, although  $v$  has no finite energy, in general  $v^{\frac{\gamma-1+p}{p}}$  does, and this also gives a (weak) sense to the boundary datum. Again, we refer to Section 8.1 for more comments on this and on how the belonging of  $u$  to  $BV(\Omega)$  can depend on both the degeneracy of the datum  $f$  and the geometry of  $\partial\Omega$ .

We finally emphasize the regularizing effect given by the, possibly singular, nonlinear term  $h$ . If  $h(0) = \infty$  then  $u > 0$  a.e. in  $\Omega$ ; this is a striking difference with the bounded case; as a consequence of a result in [16], in fact, if  $h(s) \equiv 1$  and  $\|f\|_{L^N(\Omega)} < \mathcal{S}_1^{-1}$ , then  $u = 0$ . The extension of this property to the case of a general bounded nonlinearity is given by the last assertion of Theorem 3.4.

As expected, also the behavior at infinity of the nonlinearity  $h$  plays a role; in fact, if  $h(\infty) = 0$  then no smallness assumptions need to be imposed on the data in contrast with the *linear* right hand side case ([16, 26, 28, 31]).

#### 4. EXISTENCE OF A SOLUTION FOR $p > 1$ AND FOR A NONNEGATIVE $f \in L^{(p^*)'}(\Omega)$

Here we set the theory in the case of a  $p$ -laplace principal part for a fixed  $1 < p < N$ . This case represents the basis of the approximation scheme of Theorem 3.3. Existence of solutions (and uniqueness when expected) for this kind of problems has its own interest and we will present it in full generality. Let us consider

$$\begin{cases} -\Delta_p u_p = h(u_p)f & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.10)$$

where  $f \in L^{(p^*)'}(\Omega)$  is a nonnegative function and  $h$  satisfies both (h1) and (h2). We precise the notion of solution to problem (4.10) we adopt.

**Definition 4.1.** A nonnegative function  $u_p \in W_{\text{loc}}^{1,p}(\Omega)$  is a *distributional solution* to problem (4.10) if

$$h(u_p)f \in L_{\text{loc}}^1(\Omega), \quad (4.11)$$

$$G_k(u_p) \in W_0^{1,p}(\Omega) \quad \text{for all } k > 0, \quad (4.12)$$

and

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} h(u_p)f \varphi, \quad (4.13)$$

for every  $\varphi \in C_c^1(\Omega)$ .

**Remark 4.2.** Notice that condition (4.12) is the same used in [12] for  $p > 1$  in the model case  $h(s) = s^{-\gamma}$  in order to give sense to the boundary datum and to ensure uniqueness under suitable assumptions on both the datum and/or the domain. As a matter of fact, if  $h$  is nonincreasing, then a straightforward re-adaptation of the proof of Theorem 1.5 in [12] shows that the same uniqueness property holds for problem (4.10). In particular, if  $\Omega$  star-shaped, one can show that uniqueness of distributional solutions in the sense of Definition 4.1 holds if  $f \in L^1(\Omega)$ , while some regularity on  $f$  is needed if  $\gamma > 1$  and the domain is more general.

As we already mentioned, an alternative way one can weakly intend the boundary datum in problem (4.10) is by requesting that

$$u_p^{\frac{\sigma-1+p}{p}} \in W_0^{1,p}(\Omega). \quad (4.14)$$

Property (4.14) is the same given in [18], it is consistent with the case  $p = 2$  ([11]), and, as we said, it has its counterpart as  $p \rightarrow 1^+$  (i.e.  $u^\sigma \in BV(\Omega)$ ). Let us observe that, if  $f \in L^N(\Omega)$ , then the solutions we will construct in Theorem 4.3 enjoy (4.14) (see (4.37) and Section 4.3 below) that, by a direct computation, can be shown to imply (4.12).

We finally want to stress some striking differences with the model case  $h(s) = s^{-\gamma}$ ,  $\gamma > 0$ . In this



case, in fact, the behavior of  $h$  at infinity also plays a role and (4.14) can be shown to hold for any nonnegative  $f \in L^1(\Omega)$  if  $\gamma \geq 1$ . If  $\gamma < 1$  then solutions in  $W_0^{1,p}(\Omega)$  exist for a datum  $f \in L^{(\frac{p^*}{1-\gamma})'}(\Omega)$  (see [18]). In this sense, as we do not assume any behavior for  $h$  at infinity, our summability assumption on the datum  $f$  can be considered to be optimal. See also Theorem 4.7 below for further regularity results depending on the summability of the datum  $f$ .

The following existence result holds.

**Theorem 4.3.** *Let  $0 \leq f \in L^{(p^*)'}(\Omega)$  and let  $h$  satisfy (h1) and (h2). Then there exists a distributional solution  $u_p$  to problem (4.10).*

**4.1. A priori estimates.** In order to prove Theorem 4.3 we need to establish some general a priori estimates that will be the content of this section. We look for a priori estimates for a weak solution to the following problem

$$\begin{cases} -\Delta_p v = \bar{h}(v)f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.15)$$

where  $1 < p < N$ ,  $\bar{h} : [0, \infty) \rightarrow [0, \infty)$  is bounded continuous function satisfying (h2), and  $f$  is a nonnegative function in  $L^{(p^*)'}(\Omega)$ . For a *weak solution* to problem (4.15) we mean a function  $v \in W_0^{1,p}(\Omega)$  such that

$$\int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla \varphi = \int_{\Omega} \bar{h}(v) f \varphi \quad \forall \varphi \in W_0^{1,p}(\Omega). \quad (4.16)$$

We recall that  $\sigma$  is defined by (3.7), and, for any  $k > 0$ , we set for simplicity

$$\epsilon_k := \sup_{s \in [k, \infty)} \bar{h}(s) - \bar{h}(\infty).$$

Observe that  $\epsilon_k \geq 0$  and

$$\lim_{k \rightarrow \infty} \epsilon_k = 0.$$

We have the following

**Lemma 4.4.** *Let  $0 \leq f \in L^{(p^*)'}(\Omega)$  and let  $\bar{h}$  be a bounded continuous function satisfying (h2). Then every weak solution  $v$  to problem (4.15) satisfies*

$$\|G_k(v)\|_{W_0^{1,p}(\Omega)} \leq C(p, \mathcal{S}_p, \epsilon_k, \|f\|_{L^{(p^*)'}(\Omega)}) \quad \text{for all } k > 0. \quad (4.17)$$

$$\|v\|_{W^{1,p}(\omega)} \leq C(p, \mathcal{S}_p, \bar{h}(\infty), \|f\|_{L^{(p^*)'}(\Omega)}, \omega) \quad \text{for all } \omega \subset\subset \Omega. \quad (4.18)$$

*Proof.* We first look for (4.17). For a fixed  $k > 0$ , one takes  $G_k(v)$  as test function in (4.16), and using the Hölder and the Sobolev inequality, one readily gets

$$\begin{aligned} \int_{\Omega} |\nabla G_k(v)|^p &= \int_{\Omega} \bar{h}(v) f G_k(v) \leq (\bar{h}(\infty) + \epsilon_k) \|f\|_{L^{(p^*)'}(\Omega)} \left( \int_{\Omega} G_k^{p^*}(v) \right)^{\frac{1}{p^*}} \\ &\leq (\bar{h}(\infty) + \epsilon_k) \|f\|_{L^{(p^*)'}(\Omega)} \mathcal{S}_p \left( \int_{\Omega} |\nabla G_k(v)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

that gives

$$\|G_k(v)\|_{W_0^{1,p}(\Omega)} \leq [(\bar{h}(\infty) + \epsilon_k) \|f\|_{L^{(p^*)'}(\Omega)} \mathcal{S}_p]^{\frac{1}{p-1}}. \quad (4.19)$$

Now, in order to obtain (4.18), we look for a local estimate on  $T_k(v)$ . Consider  $\omega \subset\subset \Omega$  and  $\phi \in C_c^1(\Omega)$  as a cut-off function for  $\omega$ , i. e.  $0 \leq \phi \leq 1$ ,  $\phi = 1$  on  $\omega$  and  $|\nabla \phi| \leq c_\omega$ , where  $c_\omega$  is a constant that only depends on  $\text{dist}(\omega, \partial\Omega)$ . We take  $T_k(v)\phi^p$  as test function in (4.16), we have

$$\int_{\Omega} |\nabla T_k(v)|^p \phi^p + p \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla \phi \phi^{p-1} T_k(v) = \int_{\Omega} \bar{h}(v) f T_k(v) \phi^p,$$

and so

$$\begin{aligned} \int_{\Omega} |\nabla T_k(v)|^p \phi^p &\leq pk \left| \int_{\Omega} |\nabla T_k(v)|^{p-2} \nabla T_k(v) \cdot \nabla \phi \phi^{p-1} \right| \\ &\quad + pk \left| \int_{\Omega} |\nabla G_k(v)|^{p-2} \nabla G_k(v) \cdot \nabla \phi \phi^{p-1} \right| + \int_{\Omega} \bar{h}(v) f T_k(v) \phi^p. \end{aligned} \quad (4.20)$$

By (4.16), choosing  $\varphi = \phi^p$ , we have

$$\begin{aligned} \int_{\Omega} \bar{h}(v) f T_k(v) \phi^p &\leq k \int_{\Omega} \bar{h}(v) f \phi^p = pk \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla \phi \phi^{p-1} \\ &\leq pk \left| \int_{\Omega} |\nabla T_k(v)|^{p-2} \nabla T_k(v) \cdot \nabla \phi \phi^{p-1} \right| \\ &\quad + pk \left| \int_{\Omega} |\nabla G_k(v)|^{p-2} \nabla G_k(v) \cdot \nabla \phi \phi^{p-1} \right|, \end{aligned} \quad (4.21)$$

and collecting (4.21) and (4.20) one deduces

$$\begin{aligned} \int_{\Omega} |\nabla T_k(v)|^p \phi^p &\leq 2pk \left| \int_{\Omega} |\nabla T_k(v)|^{p-2} \nabla T_k(v) \cdot \nabla \phi \phi^{p-1} \right| \\ &\quad + 2pk \left| \int_{\Omega} |\nabla G_k(v)|^{p-2} \nabla G_k(v) \cdot \nabla \phi \phi^{p-1} \right|. \end{aligned}$$

By the Young inequality and by (4.19) the previous implies, for a positive  $\eta$  to be fixed later, that

$$\begin{aligned} \int_{\Omega} |\nabla T_k(v)|^p \phi^p &\leq (p-1)\eta^{p'} \int_{\Omega} |\nabla T_k(v)|^p \phi^p + \frac{(2k)^p}{\eta^p} \int_{\Omega} |\nabla \phi|^p \\ &\quad + (p-1) \int_{\Omega} |\nabla G_k(v)|^p \phi^p + (2k)^p \int_{\Omega} |\nabla \phi|^p \\ &\stackrel{(4.19)}{\leq} (p-1)\eta^{p'} \int_{\Omega} |\nabla T_k(v)|^p \phi^p + \frac{(2k)^p}{\eta^p} \int_{\Omega} |\nabla \phi|^p \\ &\quad + (p-1)[(\bar{h}(\infty) + \epsilon_k) \|f\|_{L^{(p^*)'}(\Omega)} \mathcal{S}_p]^{\frac{p}{p-1}} + (2k)^p \int_{\Omega} |\nabla \phi|^p. \end{aligned} \quad (4.22)$$

We fix  $\eta$  such that  $1 - (p-1)\eta^{p'} = \frac{1}{2}$  that is  $\eta = \left(\frac{1}{2(p-1)}\right)^{\frac{1}{p'}}$ , then (4.22) implies

$$\begin{aligned} \int_{\omega} |\nabla T_k(v)|^p &\leq 2^{p+1} k^p [(2(p-1))^{p-1} + 1] \int_{\Omega} |\nabla \phi|^p \\ &\quad + 2(p-1)[(\bar{h}(\infty) + \epsilon_k) \|f\|_{L^{(p^*)'}(\Omega)} \mathcal{S}_p]^{\frac{p}{p-1}} \\ &\leq 2^{p+1} k^p [(2(p-1))^{p-1} + 1] c_{\omega}^p |\Omega| \\ &\quad + 2(p-1)[(\bar{h}(\infty) + \epsilon_k) \|f\|_{L^{(p^*)'}(\Omega)} \mathcal{S}_p]^{\frac{p}{p-1}}. \end{aligned} \quad (4.23)$$

Now we fix  $k$  large enough in order to have  $\epsilon_k \leq 1$  and we collect (4.19) and (4.23) yielding (4.18).  $\square$

#### 4.2. Proof of Theorem 4.3.

*Proof of Theorem 4.3.* If  $h(0) < \infty$  then the existence of a solution belonging to  $W_0^{1,p}(\Omega)$  follows by standard application of a fixed point argument, so that, without losing generality, we assume  $h(0) = \infty$ . Let us introduce the following scheme of approximation

$$\begin{cases} -\Delta_p u_n = h_n(u_n) f & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.24)$$

where  $h_n(s) = T_n(h(s))$ , for  $s \in [0, \infty)$ , and  $n > h(\infty)$ . The existence of such  $u_n \in W_0^{1,p}(\Omega)$  follows again by standard Schauder fixed point theorem. Moreover,  $u_n$  is easily seen to be nonnegative. We apply Lemma 4.4 to  $u_n$ , deducing

$$\|u_n\|_{W^{1,p}(\omega)} \leq C_{\omega}, \quad \forall \omega \subset\subset \Omega \quad (4.25)$$

which implies that, up to subsequences,  $u_n$  weakly converges in  $W^{1,p}(\omega)$  and a.e. in  $\Omega$  to a function  $u_p$ . Moreover, by weak lower semicontinuity in (4.17) (applied to  $u_n$ ) one also gets the boundary condition (4.12). Now we prove (4.13). First of all, using (4.24) and (4.25) we observe that  $h_n(u_n)f$  is locally bounded in  $L^1(\Omega)$ ; in fact, for any  $\varphi \in C_c^1(\Omega)$  one obtains

$$\int_{\Omega} h_n(u_n)f\varphi = \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla \varphi \leq \frac{1}{p'} \int_{\text{supp} \varphi} |\nabla u_n|^p + \frac{1}{p} \int_{\Omega} |\nabla \varphi|^p \leq C,$$

and  $\varphi$  can be chosen to be a cut-off function for any compact subset  $\omega$  of  $\Omega$ . We can then apply Theorem 2.1 of [10] in order to deduce that  $\nabla u_n$  converges a.e. in  $\Omega$  to  $\nabla u_p$ . In particular,  $|\nabla u_n|^{p-2} \nabla u_n$  locally strongly converges to  $|\nabla u_p|^{p-2} \nabla u_p$  in  $L^q(\Omega, \mathbb{R}^N)$ , for any  $q < p'$ . Moreover, by the Fatou lemma, it follows

$$\int_{\Omega} h(u_p)f\varphi \leq C, \quad (4.26)$$

for any nonnegative  $\varphi \in C_c^1(\Omega)$ , which implies (recall we are assuming  $h(0) = \infty$ )

$$\{u_p = 0\} \subset \{f = 0\}, \quad (4.27)$$

up to a set of zero Lebesgue measure. Now we consider  $V_{\delta}(u_n)\varphi$ , as a test for the weak formulation of (4.24) where

$$V_{\delta}(s) := \begin{cases} 1 & s \leq \delta, \\ \frac{2\delta - s}{\delta} & \delta < s < 2\delta, \\ 0 & s \geq 2\delta, \end{cases} \quad (4.28)$$

and  $0 \leq \varphi \in C_c^1(\Omega)$ . Recalling that  $V'_{\delta}(s) \leq 0$  one deduces that

$$\int_{\Omega} h_n(u_n)fV_{\delta}(u_n)\varphi \leq \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla \varphi V_{\delta}(u_n).$$

We first pass to the limit with respect to  $n$ . Using the strong convergence of  $|\nabla u_n|^{p-2} \nabla u_n$  and the a.e. and  $*$ -weak convergence of  $V_{\delta}(u_n)$  in  $L^{\infty}(\Omega)$ , we can pass to the limit on the right hand side. By Fatou's lemma on the left hand side we then deduce

$$\int_{\{u_p \leq \delta\}} h(u_p)f\varphi \leq \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi V_{\delta}(u_p).$$

Now we take in the previous  $\delta \rightarrow 0^+$  obtaining, by  $*$ -weak convergence

$$\lim_{\delta \rightarrow 0^+} \int_{\{u_p \leq \delta\}} h(u_p)f\varphi \leq \int_{\{u_p = 0\}} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = 0.$$

Therefore,

$$\int_{\Omega} h_n(u_n)f\varphi = \int_{\{u_n > \delta\}} h_n(u_n)f\varphi + \epsilon(n, \delta),$$

where  $\epsilon(n, \delta)$  is a quantity that vanishes as first  $n$  goes to  $\infty$  and then  $\delta$  goes to zero. We observe that, without loss of generality, we can always assume that  $\delta \notin \{\eta : |\{u_p = \eta\}| > 0\}$  which is at most a countable set; this will imply, in particular, that  $\chi_{\{u_n > \delta\}}$  converges a.e. in  $\Omega$  to  $\chi_{\{u_p > \delta\}}$  as  $n \rightarrow \infty$ . Moreover, we both have

$$h_n(u_n)f\chi_{\{u_n > \delta\}}\varphi \leq \sup_{s \in [\delta, \infty)} h(s)f\varphi \in L^1(\Omega),$$

and

$$h(u_p)f\chi_{\{u_p > \delta\}}\varphi \leq h(u_p)f\varphi \stackrel{(4.26)}{\in} L^1(\Omega).$$

We can then apply the Lebesgue theorem in order to deduce that

$$\lim_{\delta \rightarrow 0^+} \lim_{n \rightarrow \infty} \int_{\Omega} h_n(u_n)f\varphi = \int_{\{u_p > 0\}} h(u_p)f\varphi \stackrel{(4.27)}{=} \int_{\Omega} h(u_p)f\varphi.$$

On the left hand side of the weak formulation of (4.24) we pass to the limit using the weak convergence and a.e. convergence of  $\nabla u_n$  to  $\nabla u_p$ , finally obtaining

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} h(u_p) f \varphi.$$

for every nonnegative  $\varphi \in C_c^1(\Omega)$  from which easily (4.13) follows.  $\square$

**Remark 4.5.** An important remark is that, a careful re-adaptation of the proof of Theorem 4.3, can allow to slightly improve the set of admissible test function in (4.13). Precisely, following the same steps, one can actually realize that (4.13) holds true for  $\varphi \in W^{1,p}(\Omega)$  having compact support in  $\Omega$ . We will use this property later once we will pass to the limit as  $p \rightarrow 1^+$ .

**Remark 4.6.** We also highlight the fact that if  $h(\tilde{k}) = 0$  for some  $\tilde{k} > 0$  then we can retrieve some more informations on  $u_p$ . Indeed Theorem 4.3 guarantees the existence of a distributional solution to problem

$$\begin{cases} -\Delta_p u_p = \underline{h}(u_p) f & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega, \end{cases}$$

where

$$\underline{h}(s) := \begin{cases} h(s) & \text{if } s \leq \tilde{k}, \\ 0 & \text{if } s > \tilde{k}. \end{cases}$$

As far as it has been obtained,  $u_p$  is the almost everywhere limit of the  $u_n$  solutions of (4.24) with  $\underline{h}_n$  in place of  $h_n$ . By considering  $G_{\tilde{k}}(u_n)$  as a test function in (4.24) one thus obtain

$$\int_{\Omega} |\nabla G_{\tilde{k}}(u_n)|^p \leq \int_{\Omega} \underline{h}_n(u_n) f G_{\tilde{k}}(u_n) = 0,$$

that implies

$$\|u_n\|_{L^\infty(\Omega)} \leq \tilde{k},$$

and, so

$$\|u_p\|_{L^\infty(\Omega)} \leq \tilde{k}. \quad (4.29)$$

Since  $u_p$  satisfies

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} \underline{h}(u_p) f \varphi,$$

for every  $\varphi \in C_c^1(\Omega)$ , using (4.29), one also has

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} h(u_p) f \varphi,$$

namely  $u_p$  solves

$$\begin{cases} -\Delta_p u_p = h(u_p) f & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega, \end{cases}$$

as it also satisfies (4.12). A trivial observation is that, in this case,  $h$  needs not to be bounded at infinity.

**4.3. More regular data**  $f \in L^m(\Omega)$  **with**  $(p^*)' \leq m < N$ . As we already observed, a natural question concerning solutions  $u_p$  to problem (4.10) is whether they enjoy property (4.14), as this is the case, for instance, in the model  $h(s) = s^{-\gamma}$  (even for merely integrable data). As far as our nonlinear term is concerned the behavior at infinity of  $h$  plays a crucial role so that (4.14) is not expected in general for such a large class of data. What is true in general, for  $f \in L^{(p^*)'}(\Omega)$ , is that the solutions  $u_p$  of (4.10) found in Theorem 4.3 satisfy

$$T_k^{\frac{\sigma-1+p}{p}}(u_p) \in W_0^{1,p}(\Omega) \quad \text{and} \quad G_k(u_p) \in W_0^{1,p}(\Omega) \quad \text{for all } k > 0. \quad (4.30)$$

The first in (4.30) can be easily obtained by taking  $T_k^\sigma(u_n)$  as a test function in (4.24) while the second one has been already shown to follow from (4.17).

However, if the datum  $f$  is more regular something more can be said

**Theorem 4.7.** *Let  $0 \leq f \in L^m(\Omega)$  with  $m$  such that  $(p^*)' \leq m < \frac{N}{p}$  and let  $k > 0$ . Then there exists  $C > 0$  such that the solution  $u_p$  to (4.10) found in Theorem 4.3 satisfies*

$$\|G_k^{\frac{q-1+p}{p}}(u_p)\|_{W_0^{1,p}(\Omega)} \leq C, \quad (4.31)$$

for any  $q$  such that  $1 \leq q \leq \frac{p^*(p-1)}{m'p-p^*}$ . Moreover, if  $f \in L^m(\Omega)$  with  $m = \frac{pN-N+N\sigma}{pN-N+p\sigma}$ , then

$$\|u_p^{\frac{\sigma-1+p}{p}}\|_{W_0^{1,p}(\Omega)} \leq C. \quad (4.32)$$

Finally, if  $f \in L^m(\Omega)$  with  $m > \frac{N}{p}$  then  $u_p \in L^\infty(\Omega)$ .

*Proof.* For a fixed  $k > 0$  let us define the following auxiliary function

$$\Phi(t) := \begin{cases} t^\sigma & \text{if } t \leq k, \\ (t-k)^q + k^\sigma & \text{if } t > k, \end{cases}$$

and take  $\Phi(u_n)$  as test in (4.24) obtaining, using the Hölder inequality and the assumption on  $q$ , that

$$\begin{aligned} & \left(\frac{p}{\sigma-1+p}\right)^p \sigma \int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_n)|^p + \left(\frac{p}{q-1+p}\right)^p q \int_{\Omega} |\nabla G_k^{\frac{q-1+p}{p}}(u_n)|^p \\ & \leq \max_{s \in [0,k]} h(s) s^\sigma \int_{\{u_n \leq k\}} f + \sup_{s \in [k,\infty)} h(s) \int_{\{u_n > k\}} f G_k^q(u_n) + \sup_{s \in [k,\infty)} h(s) k^\sigma \int_{\Omega} f \\ & \leq C + \sup_{s \in [k,\infty)} h(s) \|f\|_{L^m(\Omega)} |\Omega|^{\frac{1}{m'} - \frac{qp}{(q-1+p)p^*}} \left( \int_{\Omega} G_k^{\frac{(q-1+p)p^*}{p}}(u_n) \right)^{\frac{qp}{(q-1+p)p^*}}. \end{aligned} \quad (4.33)$$

From (4.33), using the Sobolev inequality we deduce

$$\begin{aligned} & \left(\frac{p}{q-1+p}\right)^p q \mathcal{S}_p^{-p} \left( \int_{\Omega} G_k^{\frac{(q-1+p)p^*}{p}}(u_n) \right)^{\frac{p}{p^*}} \leq \left(\frac{p}{q-1+p}\right)^p q \int_{\Omega} |\nabla G_k^{\frac{q-1+p}{p}}(u_n)|^p \\ & \leq C + \sup_{s \in [k,\infty)} h(s) \|f\|_{L^m(\Omega)} |\Omega|^{\frac{1}{m'} - \frac{qp}{(q-1+p)p^*}} \left( \int_{\Omega} G_k^{\frac{(q-1+p)p^*}{p}}(u_n) \right)^{\frac{qp}{(q-1+p)p^*}} \\ & \leq C + \varepsilon \sup_{s \in [k,\infty)} h(s) \|f\|_{L^m(\Omega)} |\Omega|^{\frac{1}{m'} - \frac{qp}{(q-1+p)p^*}} \left( \int_{\Omega} G_k^{\frac{(q-1+p)p^*}{p}}(u_n) \right)^{\frac{p}{p^*}} \\ & \quad + C_\varepsilon \sup_{s \in [k,\infty)} h(s) \|f\|_{L^m(\Omega)} |\Omega|^{\frac{1}{m'} - \frac{qp}{(q-1+p)p^*}}, \end{aligned}$$

where in the last step we also used Young's inequality. Up to suitably choose  $\varepsilon$ , this gives that  $G_k(u_n)$  is bounded in  $L^{\frac{(q-1+p)p^*}{p}}(\Omega)$ . Hence by (4.33) one gets

$$\int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_n)|^p + \int_{\Omega} |\nabla G_k^{\frac{q-1+p}{p}}(u_n)|^p \leq C. \quad (4.34)$$

that implies (4.31) by weak lower semicontinuity.

If  $f \in L^m(\Omega)$  with  $m = \frac{pN-N+N\sigma}{pN-N+p\sigma}$ , then  $q = \sigma$  and one obtains (4.32) by weak lower semicontinuity in (4.34) and considering, for instance,  $k = 1$  (recall  $T_1(s) + G_1(s) = s$ ).

Now, let  $f \in L^m(\Omega)$  with  $m > \frac{N}{p}$ . It suffices to observe that in (4.24) it is not restrictive to choose  $n > \sup_{s \in [k_0,\infty)} h(s)$ , that is one can possibly truncate only near the singularity. Hence, if we multiply (4.24) by  $G_k(u_n)$  one readily has

$$\int_{\Omega} |\nabla G_k(u_n)|^p \leq \sup_{s \in [k_0,\infty)} h(s) \int_{\Omega} f G_k(u_n),$$

and one can apply standard Stampacchia's method in order to get the boundedness of  $u_p$ .  $\square$

**Remark 4.8.** First of all observe that, if  $\gamma \leq 1$  (i.e.  $\sigma = 1$ ), then (4.32) holds for  $m = (p^*)'$  and one has that the solutions are globally  $W_0^{1,p}(\Omega)$  no matter of the behavior of  $h$  at infinity.

What actually holds for any  $\gamma > 0$ , is that  $q = \frac{p^*(p-1)}{m'p-p^*} = 1$  if  $m = (p^*)'$  and so (4.31) is in continuity with the result of Theorem 4.3 (i.e.  $G_k(u_p) \in W_0^{1,p}(\Omega)$ ).

Also notice that  $q \rightarrow \infty$  as  $m \rightarrow \frac{N}{p}$ , formally implying that every power of  $G_k(u_p)$  stands in  $W_0^{1,p}(\Omega)$ . Observe that  $\frac{pN-N+N\sigma}{pN-N+p\sigma} < \frac{N}{p} < N$  for any  $p > 1$ . In the following section we will consider data in  $L^N(\Omega)$ ; hence, we shall be backed to look for (bounded, in fact) solutions satisfying  $u_p^{\frac{\sigma-1+p}{p}} \in W_0^{1,p}(\Omega)$  (see (4.37) below).

**4.4. Uniform estimates in the case  $f \in L^N(\Omega)$ .** In this section we are going to prepare the proofs of Theorems 3.3, 3.4, and 3.5. As one would like to let  $p \rightarrow 1^+$  we will need some *uniform estimates* with respect to  $p$ . In order to do that we introduce a family of test functions that behaves differently as  $u_p \sim 0$  and  $u_p \sim \infty$ . For fixed  $p > 1$ , we define the following auxiliary function

$$\psi_p(s) := \begin{cases} s^\sigma k_0^{-\frac{(\sigma-1)(p-1)}{p}} & \text{if } s \leq k_0, \\ s^{\frac{\sigma-1+p}{p}} & \text{if } s > k_0. \end{cases} \quad (4.35)$$

Observe that

$$\psi_p(s) \rightarrow s^\sigma, \text{ as } p \rightarrow 1^+.$$

We also define  $\Gamma_p(s) := \int_0^s \psi_p^{\frac{1}{p}}(t) dt$ . We have the following

**Lemma 4.9.** *Let  $h$  satisfy (h1) and (h2),  $0 \leq f \in L^N(\Omega)$  such that  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$ , and let  $u_p$  be the solution to (4.10) found in Theorem 4.3. Then there exists  $p_0 > 1$  such that for any  $p \in (1, p_0)$  one has*

$$\|u_p\|_{W^{1,p}(\omega)} \leq C(\mathcal{S}_1, h(\infty), \|f\|_{L^N(\Omega)}, \omega) \text{ for all } \omega \subset\subset \Omega, \quad (4.36)$$

$$\|u_p^{\frac{\sigma-1+p}{p}}\|_{W_0^{1,p}(\Omega)} \leq C(\mathcal{S}_1, \sup_{s \in [k_0, \infty)} h(s), \|f\|_{L^N(\Omega)}, c_1, |\Omega|). \quad (4.37)$$

Moreover, there exists  $\bar{k} > 0$  such that, for any  $k \geq \bar{k}$

$$\|G_k(u_p)\|_{L^{\frac{N}{N-1}}(\Omega)} \leq \frac{p-1}{p} |\Omega| C(\mathcal{S}_1, h(\infty), \|f\|_{L^N(\Omega)}, \bar{k}). \quad (4.38)$$

Finally,

$$\int_{\Omega} |\nabla \Gamma_p(u_p)|^p \leq \int_{\Omega} h(u_p) f \psi_p(u_p). \quad (4.39)$$

*Proof.* We divide the proof in few steps.

*Proof of (4.36).* We consider the solutions  $u_n$  to (4.24). We apply (4.19) to  $u_n$  and we apply the Hölder inequality to obtain

$$\int_{\Omega} |\nabla G_k(u_n)|^p \leq ((h(\infty) + \epsilon_k) \|f\|_{L^N(\Omega)} \mathcal{S}_p)^{\frac{p}{p-1}} |\Omega|.$$

Recalling that  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$ , we fix a constant  $c$  such that  $h(\infty) \|f\|_{L^N(\Omega)} \mathcal{S}_1 < c < 1$ . By continuity, there exist  $p_0$  sufficiently near to  $1^+$  and  $\bar{k}$  large enough such that,

$$(h(\infty) + \epsilon_k) \|f\|_{L^N(\Omega)} \mathcal{S}_p < c < 1,$$

for any  $p \in (1, p_0)$  and  $k \geq \bar{k}$ . In particular,

$$\int_{\Omega} |\nabla G_{\bar{k}}(u_n)|^p \leq c^{\frac{p-1}{p}} |\Omega|. \quad (4.40)$$

To prove (4.36) we reason as in the proof of (4.23) on  $T_{\bar{k}}(u_n)$ , and, again by Hölder's inequality, we obtain

$$\begin{aligned} \int_{\omega} |\nabla T_{\bar{k}}(u_n)|^p &\leq 4\bar{k}^p c_{\omega} [(2(p-1))^{p-1} + 1] |\Omega| \\ &\quad + 4(p-1) [(h(\infty) + \epsilon_{\bar{k}}) \|f\|_{L^N(\Omega), \mathcal{S}_p}]^{\frac{p}{p-1}} |\Omega|. \end{aligned}$$

Now observe that, thanks to the choice of  $p_0$  and  $\bar{k}$  all terms at the right hand side of the previous expression are bounded uniformly with respect to  $p \in (1, p_0)$ . This fact together with (4.40), and using weak lower semicontinuity, shows that (4.36) holds.

*Global estimate (4.37)*. In order to show (4.37) we take  $u_n^{\sigma}$  as a test function in (4.16) obtaining (for  $k_1 > k_0$ )

$$\begin{aligned} \left(\frac{p}{\sigma-1+p}\right)^p \sigma \int_{\Omega} |\nabla u_n^{\frac{\sigma-1+p}{p}}|^p &= \int_{\Omega} h_n(u_n) f u_n^{\sigma} \leq c_1 k_0^{\sigma-\gamma} \int_{\{u_n \leq k_0\}} f \\ &\quad + \left(\max_{s \in [k_0, k_1]} h(s)\right) k_1^{\sigma} \int_{\{k_0 < u_n < k_1\}} f \\ &\quad + (h(\infty) + \epsilon_{k_1}) \int_{\{u_n \geq k_1\}} f u_n^{\sigma}. \end{aligned} \quad (4.41)$$

We estimate the last term in the right hand side of (4.41). Observing that  $\frac{\sigma N}{N-1} < \frac{(\sigma-1+p)p^*}{p}$ , one can apply Hölder's inequality and then Young's inequality to get

$$\begin{aligned} (h(\infty) + \epsilon_{k_1}) \int_{\{u_n \geq k_1\}} f u_n^{\sigma} &\leq (h(\infty) + \epsilon_{k_1}) \|f\|_{L^N(\Omega)} \left( \int_{\{u_n \geq k_1\}} u_n^{\frac{\sigma N}{N-1}} \right)^{\frac{N-1}{N}} \\ &\leq (h(\infty) + \epsilon_{k_1}) \|f\|_{L^N(\Omega)} \left( \int_{\Omega} u_n^{\frac{(\sigma-1+p)p^*}{p}} \right)^{\frac{p\sigma}{(\sigma-1+p)p^*}} |\Omega|^{\frac{(p-1)(\sigma-1+N)}{N(\sigma-1+p)}} \\ &\leq (h(\infty) + \epsilon_{k_1}) \|f\|_{L^N(\Omega)} \left( \int_{\Omega} u_n^{\frac{(\sigma-1+p)p^*}{p}} \right)^{\frac{p}{p^*}} \\ &\quad + (h(\infty) + \epsilon_{k_1}) \|f\|_{L^N(\Omega)} |\Omega|^{\frac{\sigma-1+N}{N}}. \end{aligned}$$

Concerning the left hand side of (4.41) we apply the Sobolev inequality and we have

$$\left(\frac{p}{\sigma-1+p}\right)^p \sigma \int_{\Omega} |\nabla u_n^{\frac{\sigma-1+p}{p}}|^p \geq \left(\frac{p}{\sigma-1+p}\right)^p \frac{\sigma}{\mathcal{S}_p^p} \left( \int_{\Omega} u_n^{\frac{(\sigma-1+p)p^*}{p}} \right)^{\frac{p}{p^*}}.$$

Collecting the previous two inequalities gathered with (4.41) we deduce

$$\left( \left(\frac{p}{\sigma-1+p}\right)^p \frac{\sigma}{\mathcal{S}_p^p} - (h(\infty) + \epsilon_{k_1}) \|f\|_{L^N(\Omega)} \right) \left( \int_{\Omega} u_n^{\frac{(\sigma-1+p)p^*}{p}} \right)^{\frac{p}{p^*}} \leq C. \quad (4.42)$$

Now, since  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$ , for  $p$  sufficiently near to 1 and  $k_1$  sufficiently large, one has that

$$\left( \left(\frac{p}{\sigma-1+p}\right)^p \frac{\sigma}{\mathcal{S}_p^p} - (h(\infty) + \epsilon_{k_1}) \|f\|_{L^N(\Omega)} \right) > c > 0,$$

for some constant  $c$  not depending on both  $p < p_0$  and  $k_1$ . Using estimate (4.42) in (4.41) then we deduce (4.37) by weak lower semicontinuity.

*Proof of estimate (4.38)*. Now in order to show (4.38) we take  $G_k(u_n)$  as a test function in (4.24) obtaining

$$\int_{\Omega} |\nabla G_k(u_n)|^p \leq (h(\infty) + \epsilon_k) \int_{\Omega} f G_k(u_n).$$

Moreover, by the Sobolev, the Young and the Hölder inequalities, we have (recall  $p > 1$ )

$$\begin{aligned} \frac{1}{\mathcal{S}_1} \left( \int_{\Omega} G_k^{\frac{N}{N-1}}(u_n) \right)^{\frac{N-1}{N}} &\leq \int_{\Omega} |\nabla G_k(u_n)| \leq \frac{1}{p} \int_{\Omega} |\nabla G_k(u_n)|^p + \frac{p-1}{p} |\Omega| \\ &\leq (h(\infty) + \epsilon_k) \int_{\Omega} f G_k(u_n) + \frac{p-1}{p} |\Omega| \\ &\leq (h(\infty) + \epsilon_k) \|f\|_{L^N(\Omega)} \left( \int_{\Omega} G_k^{\frac{N}{N-1}}(u_n) \right)^{\frac{N-1}{N}} + \frac{p-1}{p} |\Omega|, \end{aligned}$$

that implies

$$\left( \frac{1}{\mathcal{S}_1} - (h(\infty) + \epsilon_k) \|f\|_{L^N(\Omega)} \right) \|G_k(u_n)\|_{L^{\frac{N}{N-1}}(\Omega)} \leq \frac{p-1}{p} |\Omega|.$$

As before, recalling  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$ , it is possible to pick  $\bar{k}$  large enough so that  $\frac{1}{\mathcal{S}_1} - (h(\infty) + \epsilon_k) \|f\|_{L^N(\Omega)} > c > 0$ , for any  $k \geq \bar{k}$ , where  $c$  only depends on  $\mathcal{S}_1, h(\infty)$ , and  $\|f\|_{L^N(\Omega)}$ . Then (4.38) follows by Fatou's lemma.

*Proof of (4.39).* Observe that by (4.37) one readily gets, by compact embeddings, that  $u_n^{\frac{\sigma-1+p}{p}}$  strongly converges to  $u_p^{\frac{\sigma-1+p}{p}}$  in  $L^{\frac{N}{N-1}}(\Omega)$ . As already done, by possibly decreasing its value we assume, without loss of generality, that  $k_0 \notin \{\eta : |\{u_p = \eta\}| > 0\}$ . Recalling (4.35), we consider  $\psi_p(u_n)$  as a test function for (4.24) getting

$$\int_{\Omega} |\nabla \Gamma_p(u_n)|^p = \int_{\Omega} h_n(u_n) f \psi_p(u_n) \leq c_1 k_0^{\frac{\sigma+p-1-\gamma p}{p}} \int_{\{u_n \leq k_0\}} f + \sup_{s \in [k_0, \infty)} h(s) \int_{\{u_n > k_0\}} f u_n^{\frac{\sigma-1+p}{p}} \leq C.$$

In particular  $\Gamma_p(u_n)$  is bounded in  $W_0^{1,p}(\Omega)$  with respect to  $n$  and we can use weak lower semicontinuity of the norm in order to get

$$\int_{\Omega} |\nabla \Gamma_p(u_p)|^p \leq \liminf_{n \rightarrow \infty} \int_{\Omega} h_n(u_n) f \psi_p(u_n)$$

What is left is to identify the limit, as  $n$  goes to infinity, of the right hand side of the previous expression.

We write

$$h_n(u_n) f \psi_p(u_n) = h_n(u_n) f \psi_p(u_n) \chi_{\{u_n \leq k_0\}} + h_n(u_n) f u_n^{\frac{\sigma-1+p}{p}} \chi_{\{u_n > k_0\}};$$

as

$$h_n(u_n) f \psi_p(u_n) \chi_{\{u_n \leq k_0\}} \leq c_1 k_0^{\frac{\sigma+p-1-\gamma p}{p}} f,$$

we can pass to the limit in the first term by dominated convergence. On the other hand, as  $f u_n^{\frac{\sigma-1+p}{p}}$  strongly converges in  $L^1(\Omega)$  to  $f u_p^{\frac{\sigma-1+p}{p}}$  and  $h_n(u_n) \chi_{\{u_n > k_0\}}$  converges to  $h(u_p) \chi_{\{u_p > k_0\}}$  both a. e. and \*-weak in  $L^\infty(\Omega)$  we can use the Egorov theorem in order to finally get

$$\lim_{n \rightarrow \infty} \int_{\Omega} h_n(u_n) f \psi_p(u_n) = \int_{\Omega} h(u_p) f \psi_p(u_p),$$

and so (4.39). □

**Remark 4.10.** Let us note that Theorem 4.3 holds true even for a problem like

$$\begin{cases} -\operatorname{div}(a(x, \nabla u_p)) = h(u_p) f & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega, \end{cases}$$



where  $a(x, \xi) : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}^N$  is a classical Leray-Lions operator satisfying the following structure conditions

$$\begin{aligned} a(x, \xi) \cdot \xi &\geq \alpha |\xi|^p, \quad \alpha > 0, \\ |a(x, \xi)| &\leq \beta |\xi|^{p-1}, \quad \beta > 0, \\ (a(x, \xi) - a(x, \xi')) \cdot (\xi - \xi') &> 0, \end{aligned}$$

for every  $\xi \neq \xi'$  in  $\mathbb{R}^N$  and for almost every  $x$  in  $\Omega$ .

### 5. THE LIMIT AS $p \rightarrow 1^+$ FOR A POSITIVE $f$

In this section we prove our main results concerning the case  $p = 1$ , namely Theorems 3.3, 3.4 and 3.5. As before, throughout this section,  $h$  satisfies (h1) and (h2) and  $f \in L^N(\Omega)$  is positive. The proofs of Theorems 3.3 and 3.4 will be split into those of various lemmata.

Preliminarily, let us recall that the solutions  $u_p$  found in Theorem 4.3 satisfy (4.36). This implies that  $u_p$  is locally uniformly bounded in  $BV(\Omega)$  with respect to  $p$ . Indeed for every  $\omega \subset\subset \Omega$

$$\int_{\omega} |\nabla u_p| \leq \frac{1}{p} \int_{\omega} |\nabla u_p|^p + \frac{p-1}{p} |\omega| \leq C.$$

Hence by compactness in  $BV$ , and a standard diagonal argument, we deduce the existence of a function  $u \in BV_{\text{loc}}(\Omega)$  such that (up to not relabeled subsequences and recalling (4.38))

$$u_p \rightarrow u \text{ in } L^q(\Omega) \text{ with } q < \frac{N}{N-1} \text{ and a.e. in } \Omega, \quad (5.1)$$

$$\nabla u_p \rightarrow Du \text{ locally } * \text{-weakly as measures,}$$

as  $p \rightarrow 1^+$ . First we have the following

**Lemma 5.1.** *Let  $0 < f \in L^N(\Omega)$  such that  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1 h(\infty)}$  with  $h$  satisfying both (h1) and (h2). Then  $u$ , defined by (5.1), belongs to  $L^\infty(\Omega)$ . Moreover  $u^\sigma \in BV(\Omega)$ .*

*Proof.* Using the Fatou lemma in (4.38) we have that there exists  $\bar{k}$  such that for every  $k \geq \bar{k}$

$$\|G_k(u)\|_{L^{\frac{N}{N-1}}(\Omega)} = 0,$$

that is  $0 \leq u \leq k$  a.e. in  $\Omega$ . Moreover it follows from the Young inequality and from (4.37) that

$$\int_{\Omega} |\nabla u_p^{\frac{\sigma-1+p}{p}}| \leq \frac{1}{p} \int_{\Omega} |\nabla u_p^{\frac{\sigma-1+p}{p}}|^p + \frac{1}{p} |\Omega| \leq C,$$

for some constant  $C$  not depending on  $p$ . This implies that  $u_p^{\frac{\sigma-1+p}{p}}$  is bounded in  $BV(\Omega)$ . Then there exists  $w \in BV(\Omega)$  such that  $u_p^{\frac{\sigma-1+p}{p}}$  converges to  $w$  in  $L^q(\Omega)$  for  $q < \frac{N}{N-1}$  and a.e. in  $\Omega$ , and  $\nabla u_p^{\frac{\sigma-1+p}{p}}$  converges  $Dw$   $*$ -weakly as measures. As  $u_p$  converges a.e. to  $u$  this implies that  $w = u^\sigma$  which concludes the proof. We observe that we have just shown that

$$u_p^{\frac{\sigma-1+p}{p}} \rightarrow u^\sigma \text{ in } L^q(\Omega), \text{ for every } q < \frac{N}{N-1}. \quad (5.2)$$

□

The following Lemma shows the existence (and the identification) of the vector field  $z$ .

**Lemma 5.2.** *Under the same assumptions of Lemma 5.1, there exists  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  such that*

$$h(u)f \in L_{\text{loc}}^1(\Omega), \quad (5.3)$$

$$-\operatorname{div} z = h(u)f \text{ in } \mathcal{D}'(\Omega). \quad (5.4)$$

Moreover

$$(z, Du) = |Du| \text{ as measures in } \Omega. \quad (5.5)$$

*Proof.* We divide the proof into few steps.

*Existence of the field  $z$ .* Recalling (4.36) we have, for  $1 \leq q < p'$  and for any  $\omega \subset\subset \Omega$

$$\int_{\omega} |\nabla u_p|^{p-2} \nabla u_p|^q = \int_{\omega} |\nabla u_p|^{q(p-1)} \leq \left( \int_{\omega} |\nabla u_p|^p \right)^{\frac{q}{p'}} |\omega|^{1-\frac{q}{p'}} \leq C_{\omega}^{\frac{q}{p'}} |\Omega|^{1-\frac{q}{p'}}$$

and thus,

$$\| |\nabla u_p|^{p-2} \nabla u_p \|_{q,\omega} \leq C_{\omega}^{\frac{1}{p'}} |\Omega|^{\frac{1}{q} - \frac{1}{p'}}. \quad (5.6)$$

The previous implies that  $|\nabla u_p|^{p-2} \nabla u_p$  is bounded in  $L^q(\omega, \mathbb{R}^N)$  with respect to  $p$ . Then there exists  $z_q \in L^q(\omega, \mathbb{R}^N)$  such that

$$|\nabla u_p|^{p-2} \nabla u_p \rightharpoonup z_q, \quad \text{weakly in } L^q(\omega, \mathbb{R}^N).$$

A standard diagonal argument shows that there exists a unique vector field  $z$  which is defined on  $\Omega$  independently of  $q$ , such that

$$|\nabla u_p|^{p-2} \nabla u_p \rightharpoonup z, \quad \text{weakly in } L^q(\Omega, \mathbb{R}^N), \quad \forall q < \infty. \quad (5.7)$$

Moreover, it follows from the lower semicontinuity in (5.6) with respect to  $p$  that

$$\|z\|_{q,\omega} \leq |\Omega|^{\frac{1}{q}}, \quad \forall q < \infty$$

and thus if  $q \rightarrow \infty$  then  $z \in L^{\infty}(\omega, \mathbb{R}^N)$  and  $\|z\|_{\infty,\omega} \leq 1$ . Since this estimate is independent of  $\omega$ , then  $\|z\|_{\infty} \leq 1$ .

*Distributional formulation.* We prove that (5.4) holds. Note that by (5.7) one can pass to limit with respect to  $p$  in the left hand side of the distributional formulation of (4.10). Concerning the right hand side, we first notice that, if  $h(0) < \infty$  then one passes to the limit using the a.e. convergence of  $u_p$  and the Lebesgue theorem. We then assume that  $h(0) = \infty$ .

Let  $0 \leq \varphi \in C_c^1(\Omega)$ , then applying the Young inequality in the distributional formulation of (4.10) it yields

$$\int_{\Omega} h(u_p) f \varphi \leq \frac{1}{p'} \int_{supp \varphi} |\nabla u_p|^p + \frac{1}{p} \int_{\Omega} |\nabla \varphi|^p \leq C,$$

and then the Fatou lemma gives

$$\int_{\Omega} h(u) f \varphi \leq \liminf_{p \rightarrow 1^+} \int_{\Omega} h(u_p) f \varphi \leq C, \quad (5.8)$$

which implies (5.3) and that  $u > 0$  a.e. in  $\Omega$  since  $f > 0$  a.e. in  $\Omega$ . Now, recalling Remark 4.5, we are able to take  $V_{\delta}(u_p)\varphi$  as a test function in (4.10) where  $0 \leq \varphi \in C_c^1(\Omega)$  and  $V_{\delta}$  is defined as in (4.28), we have

$$\int_{\Omega} |\nabla u_p|^p V_{\delta}'(u_p) \varphi + \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi V_{\delta}(u_p) = \int_{\Omega} h(u_p) f V_{\delta}(u_p) \varphi,$$

which, since  $V_{\delta}'(s) \leq 0$ , takes to

$$\int_{\{u_p \leq \delta\}} h(u_p) f \varphi \leq \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi V_{\delta}(u_p).$$

Hence the Egorov theorem (recall (5.7)) implies

$$\limsup_{p \rightarrow 1^+} \int_{\{u_p \leq \delta\}} h(u_p) f \varphi \leq \int_{\Omega} z \cdot \nabla \varphi V_{\delta}(u).$$

Then the Lebesgue theorem implies that

$$\lim_{\delta \rightarrow 0^+} \limsup_{p \rightarrow 1^+} \int_{\{u_p \leq \delta\}} h(u_p) f \varphi \leq \int_{\{u=0\}} z \cdot \nabla \varphi = 0, \quad (5.9)$$

since  $u > 0$  a.e. in  $\Omega$ . Observe now that, by a standard density argument, (5.9) can be shown to hold for any  $\varphi \in C_c^1(\Omega)$ .

In order to pass to the limit (with respect to  $p$ ) in

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} h(u_p) f \varphi,$$

we then let

$$\int_{\Omega} h(u_p) f \varphi = \int_{\{u_p \leq \delta\}} h(u_p) f \varphi + \int_{\{u_p > \delta\}} h(u_p) f \varphi,$$

and, using the same agreement on  $\delta$  used in the proof of Theorem 4.3 (namely  $\delta \notin \{\eta : |\{u = \eta\}| > 0\}$ ), and recalling (5.8) we have, again by Lebesgue theorem that

$$\lim_{\delta \rightarrow 0^+} \lim_{p \rightarrow 1^+} \int_{\{u_p > \delta\}} h(u_p) f \varphi = \int_{\Omega} h(u) f \varphi,$$

for any  $\varphi \in C_c^1(\Omega)$ . This fact, together with (5.9) implies (5.4). Observe that this also implies that  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$ .

*A variational identity.* In order to show (5.5), the first step consists in proving the following

$$-(u^\sigma)^* \operatorname{div} z = h(u) f u^\sigma \text{ in } \mathcal{D}'(\Omega). \quad (5.10)$$

To do that we re-adapt the idea in [21]; we test (5.4) with  $(\rho_\epsilon * u^\sigma)\varphi$ , where  $\rho_\epsilon$  is a standard mollifier and  $\varphi \in C_c^1(\Omega)$ . One has

$$-\int_{\Omega} (\rho_\epsilon * u^\sigma)\varphi \operatorname{div} z = \int_{\Omega} h(u) f (\rho_\epsilon * u^\sigma)\varphi. \quad (5.11)$$

Since  $u \in L^\infty(\Omega)$  then we have  $(u^\sigma)^* \leq \|u^\sigma\|_{L^\infty(\Omega)} \mathcal{H}^{N-1}$ -a.e. and so  $\operatorname{div} z$ -a.e. (recall that  $\operatorname{div} z \ll \mathcal{H}^{N-1}$ ). It is then standard (see for instance Propositions 3.64 (b) and 3.69 (b) of [3]) that  $\rho_\epsilon * u^\sigma \rightarrow (u^\sigma)^* \mathcal{H}^{N-1}$ -a.e. and then  $\operatorname{div} z$ -a.e. Therefore, we can pass to the limit in both sides of (5.11) by dominated convergence theorem also using (5.8) and the fact that  $|\rho_\epsilon * u^\sigma| \leq \|u^\sigma\|_{L^\infty(\Omega)}$ . Then (5.10) holds.

*A first identification result.* A second ingredient for (5.5) is the following identification identity involving  $u^\sigma$ :

$$|Du^\sigma| = (z, Du^\sigma) \text{ as measures.}$$

We use  $u_p^{\frac{\sigma-1+p}{p}}$  as a test function in (4.10) where  $0 \leq \varphi \in C_c^1(\Omega)$ , and we get

$$\left(\frac{\sigma-1+p}{p}\right) \left(\frac{p^2}{\sigma-1+p^2}\right)^p \int_{\Omega} \varphi |\nabla u_p^{\frac{\sigma-1+p^2}{p^2}}|^p + \int_{\Omega} u_p^{\frac{\sigma-1+p}{p}} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi.$$

Thus, by Young's inequality, we deduce

$$\begin{aligned} \left(\frac{\sigma-1+p}{p}\right)^{\frac{1}{p}} \left(\frac{p^2}{\sigma-1+p^2}\right) \int_{\Omega} \varphi |\nabla u_p^{\frac{\sigma-1+p^2}{p^2}}| + \int_{\Omega} u_p^{\frac{\sigma-1+p}{p}} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi \\ \leq \int_{\Omega} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi + \frac{p-1}{p} \int_{\Omega} \varphi. \end{aligned} \quad (5.12)$$

By (5.2) we observe that  $u_p^{\frac{\sigma-1+p}{p}}$  converges to  $u^\sigma$  in  $L^r(\Omega)$  for  $r < \frac{N}{N-1}$  and a.e. in  $\Omega$  and (see (5.7)) that  $|\nabla u_p|^{p-2} \nabla u_p$  converges weakly to  $z$  in  $L^q(\Omega, \mathbb{R}^N)$  for any  $q < \infty$ ; this is sufficient to pass to the limit in the second term on the left hand side of the previous.

Concerning the right hand side of (5.12) we have, for  $\delta > 0$ ,

$$\int_{\Omega} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi = \int_{\{u_p \leq \delta\}} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi + \int_{\{u_p > \delta\}} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi.$$

On one hand, we treat the first term on the right hand side of the previous as follows

$$\lim_{\delta \rightarrow 0^+} \limsup_{p \rightarrow 1^+} \int_{\{u_p \leq \delta\}} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi \leq \lim_{\delta \rightarrow 0^+} \limsup_{p \rightarrow 1^+} \delta^{\frac{\sigma-1+p}{p}} \int_{\{u_p \leq \delta\}} h(u_p) f \varphi = 0.$$

reasoning as for (5.9). For the second term, using the usual convention for the choice of  $\delta$ , the a.e. convergence and the weak convergence of  $h(u_p)u_p^{\frac{\sigma-1+p}{p}} \chi_{\{u_p>\delta\}}$  to  $h(u)u^\sigma \chi_{\{u>\delta\}}$  in  $L^{\frac{N}{N-1}}(\Omega)$ , with respect to  $p$ , recalling that  $f \in L^N(\Omega)$ , implies that

$$\lim_{p \rightarrow 1^+} \int_{\{u_p>\delta\}} h(u_p) f u_p^{\frac{\sigma-1+p}{p}} \varphi = \int_{\{u>\delta\}} h(u) f u^\sigma \varphi.$$

Then by the Lebesgue theorem, we can pass to the limit also as  $\delta \rightarrow 0^+$  since  $h(u) f u^\sigma \varphi \in L^1(\Omega)$  (recall  $u$  is bounded). In particular observe that, for fixed  $0 \leq \varphi \in C_c^1(\Omega)$ , all but the first term in (5.12) are uniformly bounded with respect to  $p$ .

Therefore  $u_p^{\frac{\sigma-1+p^2}{p^2}}$  is bounded in  $BV_{\text{loc}}(\Omega)$  and it locally converges, up to subsequences, to  $u^\sigma$  a.e. in  $\Omega$ , in  $L^r(\Omega)$  with  $r < \frac{N}{N-1}$  and  $\nabla u_p^{\frac{\sigma-1+p^2}{p^2}}$  converges \*-weakly locally as measures to  $Du^\sigma$ . Then by weak lower semicontinuity in the first term we obtain, recalling (5.10), that

$$\int_{\Omega} \varphi |Du^\sigma| + \int_{\Omega} u^\sigma z \cdot \nabla \varphi \leq \int_{\Omega} h(u) f u^\sigma \varphi = - \int_{\Omega} (u^\sigma)^* \varphi \operatorname{div} z, \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0.$$

Moreover it follows from Proposition 2.1 that

$$\int_{\Omega} \varphi |Du^\sigma| \leq - \int_{\Omega} u^\sigma z \cdot \nabla \varphi - \int_{\Omega} (u^\sigma)^* \varphi \operatorname{div} z = \int_{\Omega} \varphi(z, Du^\sigma), \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0,$$

then

$$\int_{\Omega} \varphi |Du^\sigma| = \int_{\Omega} \varphi(z, Du^\sigma), \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0, \quad (5.13)$$

the reverse inequality being trivial since  $\|z\|_\infty \leq 1$ .

*Proof completed.* Here we show that (5.13) implies (5.5). Let us choose in (2.2)  $\Lambda : [0, \infty) \rightarrow [0, \infty)$  defined by  $\Lambda(s) = s^\sigma$ , which is a Lipschitz increasing function, since  $\sigma \geq 1$ . Then

$$\frac{(z, Du)}{|Du|} = \theta(z, Du, x) = \theta(z, Du^\sigma, x) = \frac{(z, Du^\sigma)}{|Du^\sigma|} \quad \text{for } |Du|\text{-a.e. } x \in \Omega,$$

namely (5.5). □

In order to prove Theorem 3.3 we need to show that  $z \in \mathcal{D}\mathcal{M}^\infty(\Omega)$ . In fact, we have the following stronger general fact:

**Lemma 5.3.** *Let  $0 \leq g \in L^1_{\text{loc}}(\Omega)$  and let  $z \in \mathcal{D}\mathcal{M}^\infty_{\text{loc}}(\Omega)$  with  $\|z\|_\infty \leq 1$  such that*

$$- \operatorname{div} z = g \text{ in } \mathcal{D}'(\Omega), \quad (5.14)$$

then

$$g \in L^1(\Omega). \quad (5.15)$$

In particular,  $\operatorname{div} z \in L^1(\Omega)$  and the following holds

$$- \int_{\Omega} v \operatorname{div} z = \int_{\Omega} g v, \quad \forall v \in BV(\Omega) \cap L^\infty(\Omega). \quad (5.16)$$

*Proof.* Let  $0 \leq v \in W_0^{1,1}(\Omega)$  and let  $\varphi_n \in C_c^1(\Omega)$  be a sequence of nonnegative functions converging in  $W_0^{1,1}(\Omega)$  to  $v$ . Let us take  $\rho_\eta * (v \wedge \varphi_n)$  as test function in (5.14) where  $\rho_\eta$  ( $\eta > 0$ ) is a standard mollifier. We obtain

$$\int_{\Omega} z \cdot \nabla(\rho_\eta * (v \wedge \varphi_n)) = \int_{\Omega} g \rho_\eta * (v \wedge \varphi_n), \quad (5.17)$$

and we are able to pass to the limit in the left hand side of the previous as  $\eta \rightarrow 0$  since  $\rho_\eta * (v \wedge \varphi_n) \rightarrow v \wedge \varphi_n$  strongly in  $W_0^{1,1}(\Omega)$ . For the right hand side we observe that, for  $\eta > 0$  small enough,  $\operatorname{supp}(\rho_\eta * (v \wedge \varphi_n)) \subseteq \omega_n$ , where  $\omega_n \subset\subset \Omega$ . Moreover  $\|\rho_\eta * (v \wedge \varphi_n)\|_{L^\infty(\Omega)} \leq \|v \wedge \varphi_n\|_{L^\infty(\Omega)}$  and, as  $\eta \rightarrow 0$ ,

$\rho_\eta * (v \wedge \varphi_n)$  converges a.e. in  $\Omega$  to  $v \wedge \varphi_n$ . Thus  $\rho_\eta * (v \wedge \varphi_n)$  converges  $*$ -weak in  $L^\infty(\Omega)$  to  $v \wedge \varphi_n$ . Then, since  $g \in L^1(\omega_n)$ , we have that

$$\lim_{\eta \rightarrow 0} \int_{\Omega} g(\rho_\eta * (v \wedge \varphi_n)) = \int_{\Omega} g(v \wedge \varphi_n). \quad (5.18)$$

By (5.17) and (5.18) we deduce

$$\int_{\Omega} z \cdot \nabla(v \wedge \varphi_n) = \int_{\Omega} g(v \wedge \varphi_n). \quad (5.19)$$

Now, we need to pass to the limit the previous as  $n \rightarrow \infty$ . Since  $v \wedge \varphi_n$  converges to  $v$  in  $W_0^{1,1}(\Omega)$  then we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} z \cdot \nabla(v \wedge \varphi_n) = \int_{\Omega} z \cdot \nabla v.$$

For the right hand side of (5.19) we observe that  $g(v \wedge \varphi_n)$  converges a.e. in  $\Omega$  to  $gv$  and that  $0 \leq g(v \wedge \varphi_n) \leq gv$ . Then, in order to apply the Lebesgue theorem, it is sufficient to show that  $gv \in L^1(\Omega)$ . Indeed we have

$$\int_{\Omega} g\varphi_n = \int_{\Omega} z \cdot \nabla\varphi_n \leq \|z\|_{L^\infty(\Omega)} \int_{\Omega} |\nabla\varphi_n| \leq C,$$

where the last inequality follows from the fact that  $\varphi_n$  converges to  $v$  in  $W_0^{1,1}(\Omega)$ . Then an application of the Fatou lemma implies  $gv \in L^1(\Omega)$ . Hence we have proved that

$$\int_{\Omega} z \cdot \nabla v = \int_{\Omega} gv, \quad \forall v \in W_0^{1,1}(\Omega), \quad v \geq 0. \quad (5.20)$$

Now we take  $\tilde{v} \in W^{1,1}(\Omega)$  and then it follows from [7, Lemma 5.5] the existence of  $w_n \in W^{1,1}(\Omega) \cap C(\Omega)$  having  $w_n|_{\partial\Omega} = \tilde{v}|_{\partial\Omega}$ ,  $\int_{\Omega} |\nabla w_n| dx \leq \int_{\partial\Omega} \tilde{v} d\mathcal{H}^{N-1} + \frac{1}{n}$ , and such that  $w_n$  tends to 0 in  $\Omega$ . Clearly, we can take  $|v - w_n| \in W_0^{1,1}(\Omega)$  as a test function in (5.20), obtaining

$$\begin{aligned} \int_{\Omega} g|v - w_n| &= \int_{\Omega} z \cdot \nabla|v - w_n| \leq \|z\|_{\infty} \int_{\Omega} |\nabla\tilde{v}| + \|z\|_{\infty} \int_{\Omega} |\nabla w_n| \\ &\leq \int_{\Omega} |\nabla\tilde{v}| + \int_{\partial\Omega} \tilde{v} d\mathcal{H}^{N-1} + \frac{1}{n}. \end{aligned}$$

Once again an application of the Fatou lemma implies

$$\int_{\Omega} g\tilde{v} \leq \int_{\Omega} |\nabla\tilde{v}| + \int_{\partial\Omega} \tilde{v} d\mathcal{H}^{N-1},$$

where, taking  $\tilde{v} \equiv 1$ , one deduces that  $g$  belongs to  $L^1(\Omega)$ . Since  $\operatorname{div} z \in L^1(\Omega)$ , one can apply Anzellotti's theory in order to prove (5.16).  $\square$

The following lemma shows that the boundary datum is attained in the sense of Definition 3.1.

**Lemma 5.4.** *Under the same assumptions of Lemma 5.1, the vector field  $z$  found in Lemma 5.2 is such that one of the following holds:*

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \cap B(x, \epsilon)} u(y) dy = 0 \quad \text{or} \quad [z, \nu](x) = -1 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega. \quad (5.21)$$

*Proof.* In order to prove (5.21) we observe that (4.39) together with the Young inequality and the fact that  $u_p^{\frac{\sigma-1+p}{p}}$  has zero trace in  $W_0^{1,p}(\Omega)$ , gives

$$\int_{\Omega} |\nabla \Gamma_p(u_p)| + \int_{\partial\Omega} u_p^{\frac{\sigma-1+p}{p}} d\mathcal{H}^{N-1} \leq \int_{\Omega} h(u_p) f \psi_p(u_p) + \frac{p-1}{p} |\Omega|.$$

Now, we use weak lower semicontinuity on the left hand side, while, reasoning as in the proof of Lemma 5.2 (splitting the integral the function in the two zones  $\{u_p \leq k_0\}$  and  $\{u_p > k_0\}$  and using the definition of  $\psi_p$ ) it is not difficult to use Lebesgue theorem in order to get

$$\int_{\Omega} |Du^{\sigma}| + \int_{\partial\Omega} u^{\sigma} d\mathcal{H}^{N-1} \leq \int_{\Omega} h(u)f u^{\sigma} = - \int_{\Omega} (u^{\sigma})^* \operatorname{div} z,$$

where in the last equality we used (5.10). By the Gauss–Green formula (2.5) we have

$$\int_{\Omega} |Du^{\sigma}| + \int_{\partial\Omega} u^{\sigma} d\mathcal{H}^{N-1} \leq \int_{\Omega} (z, Du^{\sigma}) - \int_{\partial\Omega} [u^{\sigma} z, \nu] d\mathcal{H}^{N-1}.$$

Now we can apply Lemma 5.3 with  $g = h(u)f$  in order to deduce that  $z \in \mathcal{D}\mathcal{M}^{\infty}(\Omega)$  and then, since by (5.13)  $|Du^{\sigma}| = (z, Du^{\sigma})$ , the previous implies

$$u^{\sigma}(1 + [z, \nu]) = 0 \quad \mathcal{H}^{N-1}\text{-a.e. on } \partial\Omega.$$

Therefore, either  $[z, \nu](x) = -1$  or  $u^{\sigma}(x) = 0$  for  $\mathcal{H}^{N-1}$ -a.e.  $x \in \partial\Omega$ . In particular (see Theorem 3.87, [3]), if  $u^{\sigma}(x) = 0$ , then

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \cap B(x, \epsilon)} u^{\sigma}(y) dy = 0.$$

If  $\sigma > 1$ , using Hölder inequality one gets,

$$\int_{\Omega \cap B(x, \epsilon)} u(y) dy \leq \left( \int_{\Omega \cap B(x, \epsilon)} u^{\sigma}(y) dy \right)^{\frac{1}{\sigma}} \frac{|\Omega \cap B(x, \epsilon)|^{\frac{1}{\sigma'}}}{\epsilon^{\frac{N}{\sigma'}}} \leq C \left( \int_{\Omega \cap B(x, \epsilon)} u^{\sigma}(y) dy \right)^{\frac{1}{\sigma}} \xrightarrow{\epsilon \rightarrow 0} 0,$$

that is (5.21) holds.  $\square$

*Proof of Theorem 3.3.* The proof follows by gathering together Lemmata 5.1 – 5.3.  $\square$

*Proof of Theorem 3.4.* By Lemma 5.1 we have that  $u^{\sigma} \in BV(\Omega)$ . Moreover, if  $h(0) = \infty$ , it was already observed that, as (5.8) is in force, then  $u > 0$  a.e. in  $\Omega$ . Moreover we can apply Lemma 5.3 with  $g = h(u)f$  in order to deduce that  $\operatorname{div} z \in L^1(\Omega)$  and that (3.9) holds.

Therefore let  $h(0) < \infty$ , we want to show that  $u \equiv 0$ . We consider the solution  $0 \leq w_p \in W_0^{1,p}(\Omega) \cap L^{\infty}(\Omega)$  to

$$\begin{cases} -\Delta_p w_p = \|h\|_{L^{\infty}([0, \infty))} f & \text{in } \Omega, \\ w_p = 0 & \text{on } \partial\Omega. \end{cases} \quad (5.22)$$

If  $\|h\|_{L^{\infty}([0, \infty))} \|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{S}_1}$  then using [16, Theorem 4.1] we deduce that  $w_p$  goes to zero a.e. in  $\Omega$  as  $p \rightarrow 1^+$ .

On the other hand, we recall that  $u$  is the a.e. limit in  $\Omega$  of the solutions to

$$\begin{cases} -\Delta_p u_p = h(u_p) f & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega, \end{cases} \quad (5.23)$$

where  $0 \leq u_p \in W_0^{1,p}(\Omega) \cap L^{\infty}(\Omega)$ . We take  $(w_p - u_p)^-$  as a test function in the difference between weak formulations (5.22) and (5.23)

$$- \int_{\{w_p < u_p\}} (|\nabla w_p|^{p-2} \nabla w_p - |\nabla u_p|^{p-2} \nabla u_p) \cdot \nabla (w_p - u_p) = \int_{\Omega} (\|h\|_{L^{\infty}([0, \infty))} - h(u_p)) f (w_p - u_p)^- \geq 0,$$

which, by monotonicity, implies

$$\int_{\{w_p < u_p\}} (|\nabla w_p|^{p-2} \nabla w_p - |\nabla u_p|^{p-2} \nabla u_p) \cdot \nabla (w_p - u_p) = 0,$$

that is  $w_p \geq u_p \geq 0$  a.e. in  $\Omega$ . Therefore, taking  $p \rightarrow 1^+$ , one obtains  $u \equiv 0$ .  $\square$

We conclude this section by proving our uniqueness result.

*Proof of Theorem 3.5.* Since  $h(u)f \in L^1_{\text{loc}}(\Omega)$  (see (3.2)) we can apply Lemma 5.3 deducing

$$-\int_{\Omega} v \operatorname{div} z = \int_{\Omega} h(u)fv, \quad \forall v \in BV(\Omega) \cap L^\infty(\Omega).$$

Moreover we apply Proposition 2.2 and, recalling (2.4), we deduce

$$\int_{\Omega} (z, Dv) - \int_{\partial\Omega} v[z, \nu] d\mathcal{H}^{N-1} = \int_{\Omega} h(u)fv, \quad \forall v \in BV(\Omega) \cap L^\infty(\Omega). \quad (5.24)$$

Let  $u_1$  and  $u_2$  be solutions to problem (3.1) satisfying (3.8) and we denote by, respectively,  $z_1$  and  $z_2$  the vector fields appearing in Definition 3.1. Now we take  $v = u_1^\sigma - u_2^\sigma$  in the difference of weak formulations (5.24) solved by  $u_1, u_2$ . Thus

$$\begin{aligned} \int_{\Omega} (z_1, Du_1^\sigma) - \int_{\Omega} (z_2, Du_1^\sigma) + \int_{\Omega} (z_2, Du_2^\sigma) - \int_{\Omega} (z_1, Du_2^\sigma) - \int_{\partial\Omega} (u_1^\sigma - u_2^\sigma)[z_1, \nu] d\mathcal{H}^{N-1} \\ + \int_{\partial\Omega} (u_1^\sigma - u_2^\sigma)[z_2, \nu] d\mathcal{H}^{N-1} = \int_{\Omega} (h(u_1) - h(u_2))f(u_1^\sigma - u_2^\sigma). \end{aligned}$$

Then we can reason as in the last step of the proof Lemma 5.2 in order to deduce

$$\int_{\Omega} (z_i, Du_i^\sigma) = \int_{\Omega} |Du_i^\sigma| \quad \text{for } i = 1, 2.$$

Moreover observe that for a nonnegative function  $u \in L^\infty(\Omega)$  such that  $u^\sigma$  in  $BV(\Omega)$  one has that

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \cap B(x, \epsilon)} u(y) dy = 0$$

implies

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \cap B(x, \epsilon)} u^\sigma(y) dy = 0.$$

In particular

$$u_i^\sigma(1 + [z_i, \nu]) = 0 \quad \mathcal{H}^{N-1} \text{ - a.e. on } \partial\Omega \quad \text{for } i = 1, 2$$

Then, it follows

$$\begin{aligned} \int_{\Omega} |Du_1^\sigma| - \int_{\Omega} (z_2, Du_1^\sigma) + \int_{\Omega} |Du_2^\sigma| - \int_{\Omega} (z_1, Du_2^\sigma) + \int_{\partial\Omega} (u_1^\sigma + u_1^\sigma[z_2, \nu]) d\mathcal{H}^{N-1} \\ + \int_{\partial\Omega} (u_2^\sigma[z_1, \nu] + u_2^\sigma) d\mathcal{H}^{N-1} = \int_{\Omega} (h(u_1) - h(u_2))f(u_1^\sigma - u_2^\sigma). \end{aligned}$$

Hence recalling that  $\|z_i\|_\infty \leq 1$  and that  $[z_i, \nu] \in [-1, 1]$  for  $i = 1, 2$  then the left hand side of the previous is nonnegative. This gives that

$$\int_{\Omega} (h(u_1) - h(u_2))f(u_1^\sigma - u_2^\sigma) \geq 0,$$

which implies  $u_1 = u_2$  a.e. in  $\Omega$  since  $f > 0$  a.e. in  $\Omega$ .  $\square$

## 6. NONNEGATIVE DATA $f$

Here we extend existence Theorem 3.3 to the case of a nonnegative  $f$  in  $L^N(\Omega)$  in (3.1). Here we focus on the purely singular case  $h(0) = \infty$ ; if this is not the case (i.e.  $h(0) < \infty$ ), one can easily re-adapt (with many simplifications) the argument of the previous section in order to obtain a solution to problem (3.1) which satisfies (3.2)-(3.5).

As suggested in [21], when  $h$  actually blows up at the origin then the notion of solution should be suitably modified. In fact, roughly speaking, the approximating solutions  $u_p$  could converge to a limit function  $u$  that may have a nontrivial set  $\{u = 0\}$ . This fact, in the  $BV$  context amounts to the fact that an additional term (namely a measure) appears in the limit equation (see Remark 6.2); this additional term can be absorbed in the principal part of the equation by formally multiplying it by  $\chi_{\{u > 0\}}^*$ . Moreover, in this case the vector field  $z$  will actually belong to  $\mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  and this leads to a different formulation for the boundary datum that involves the power  $\sigma = \max(1, \gamma)$  of  $u$ . As a

matter of fact, in the case  $f > 0$ , the two definitions do essentially coincide (see Remark 6.3). We set the following

**Definition 6.1.** Let  $0 \leq f \in L^N(\Omega)$  then a function  $u \in BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$  having  $\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega)$  and  $u^\sigma \in BV(\Omega)$  is a solution to problem (3.1) if there exists  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  such that

$$h(u)f \in L^1_{\text{loc}}(\Omega), \quad (6.1)$$

$$-(\operatorname{div} z)\chi_{\{u>0\}}^* = h(u)f \quad \text{in } \mathcal{D}'(\Omega), \quad (6.2)$$

$$(z, Du) = |Du| \quad \text{as measures in } \Omega, \quad (6.3)$$

$$u^\sigma(x) + [u^\sigma z, \nu](x) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega. \quad (6.4)$$

**Remark 6.2.** It is worth noting that, reasoning as in [21], one can prove that  $(z, D\chi_{\{u>0\}}) = |D\chi_{\{u>0\}}|$ . This means that, by the Anzellotti theory, one has

$$-(\operatorname{div} z)\chi_{\{u>0\}}^* = -\operatorname{div}(z\chi_{\{u>0\}}) + (z, D\chi_{\{u>0\}}),$$

and then the equation (6.2) reads as

$$-\operatorname{div}(z\chi_{\{u>0\}}) + |D\chi_{\{u>0\}}| = h(u)f,$$

that is, it reduces to a sum of an operator in divergence form and an additional term  $|D\chi_{\{u>0\}}|$ , which is a measure concentrated on the, nontrivial in this case, reduced boundary  $\partial^*\{u > 0\}$ , or equivalently on the reduced boundary  $\partial^*\{u = 0\}$ .

**Remark 6.3.** Let us stress that a solution in the sense of Definition 6.1 is also a solution in the sense of Definition 3.1 in case of  $f > 0$  a.e. in  $\Omega$ . Indeed, since  $h(u)f$  is locally integrable then (recall we are assuming  $h(0) = \infty$ )  $u > 0$  a.e. in  $\Omega$  and (6.2) reads as

$$-\operatorname{div} z = h(u)f \quad \text{in } \mathcal{D}'(\Omega).$$

Then we can apply Lemma 5.3 in order to deduce that  $z \in \mathcal{D}\mathcal{M}^\infty(\Omega)$ . Finally we only need to show that (3.5) holds. We observe that having  $u^\sigma \in BV(\Omega) \cap L^\infty(\Omega)$  and  $z \in \mathcal{D}\mathcal{M}^\infty(\Omega)$  we can use (2.4) in order to deduce from (6.4) that

$$u^\sigma(x)(1 + [z, \nu](x)) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega,$$

that implies (3.5) reasoning as in the proof of Lemma 5.4.

On the other hand it is easy to see that a solution in the sense of Definition 3.1 also satisfies Definition 6.1 provided  $u^\sigma \in BV(\Omega)$ .

Let us also finally remark that uniqueness of solution in the sense of Definition 6.1 is not expected in general as some 1-D examples in the model case with  $\gamma \leq 1$  show (see [21]).

We have the following counterpart of Theorem 3.3 for general nonnegative  $f$

**Theorem 6.4.** Let  $0 \leq f \in L^N(\Omega)$  such that  $\|f\|_{L^N(\Omega)} < \frac{1}{\mathcal{F}_1 h(\infty)}$  and let  $h$  satisfy (h1) and (h2). Then there exists a solution  $u$  to problem (3.1) in the sense of Definition 6.1.

*Proof.* The proof strictly follows the lines of the one of Theorem 3.3 so we only sketch it by highlighting the main differences. One reasons by approximation with the distributional solutions  $u_p$  to (4.10) and use the estimates given in Lemma 4.9. The existence of both an a.e. limit function  $u \in BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$  and a vector field  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  (\*-weak limit in  $L^\infty(\Omega, \mathbb{R}^N)$ ) of  $|\nabla u_p|^{p-2} \nabla u_p$ ) then follows as before. Moreover  $u^\sigma \in BV(\Omega)$  and (6.3) is in force. What are left are the proofs that  $\chi_{\{u>0\}}$  belongs to  $BV_{\text{loc}}(\Omega)$ , that  $h(u)f \in L^1_{\text{loc}}(\Omega)$ , and that (6.2) holds. It follows by the Fatou lemma applied to (4.13) that

$$\int_\Omega h(u)f\varphi \leq \int_\Omega z \cdot \nabla\varphi = - \int_\Omega \varphi \operatorname{div} z \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0; \quad (6.5)$$



in particular  $h(u)f \in L^1_{\text{loc}}(\Omega)$ . Now we test (4.10) with  $S_\delta(u_p)\varphi$  (see Remark 4.5), where  $S_\delta(s) := 1 - V_\delta(s)$ ,  $V_\delta$  is defined in (4.28), and  $0 \leq \varphi \in C^1_c(\Omega)$ , obtaining

$$\int_{\Omega} |\nabla u_p|^p S'_\delta(u_p)\varphi + \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi S_\delta(u_p) = \int_{\Omega} h(u_p) f S_\delta(u_p) \varphi. \quad (6.6)$$

Thus from (6.6), using Young's inequality (recall  $p > 1$ ), we have

$$\begin{aligned} & \int_{\Omega} |\nabla S_\delta(u_p)|\varphi + \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi S_\delta(u_p) \\ & \leq \frac{1}{p} \int_{\Omega} |\nabla u_p|^p S'_\delta(u_p)\varphi + \frac{p-1}{p} \int_{\Omega} S'_\delta(u_p)\varphi + \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi S_\delta(u_p) \\ & \leq \frac{p-1}{p} \int_{\Omega} S'_\delta(u_p)\varphi + \int_{\Omega} h(u_p) f S_\delta(u_p)\varphi. \end{aligned} \quad (6.7)$$

We want to pass to the limit as  $p \rightarrow 1^+$  first, and then we will let  $\delta \rightarrow 0^+$ . First of all, using that  $|\nabla S_\delta(u_p)| = |S'_\delta(u_p)\nabla u_p| \leq \frac{1}{\delta} |\nabla u_p|$  then it follows from (4.36) the uniform local boundedness of  $S_\delta(u_p)$  in  $BV_{\text{loc}}(\Omega)$  with respect to  $p$  and we can pass to the limit in (6.7) by weak lower semicontinuity in the first term on the left hand side. Also the second term easily passes to the limit by the Egorov theorem. On the right hand side, the first term vanishes (as  $S'_\delta$  is bounded) while for the second term we have

$$h(u_p) f S_\delta(u_p)\varphi \leq h(u_p) f \varphi \chi_{\{u_p > \delta\}} \leq \sup_{s \in [\delta, \infty)} h(s) f \varphi$$

so that, by dominated convergence theorem, we can pass to the limit in this term as well finally get

$$\int_{\Omega} |DS_\delta(u)|\varphi + \int_{\Omega} z \cdot \nabla \varphi S_\delta(u) \leq \int_{\Omega} h(u) f S_\delta(u)\varphi.$$

Thanks to the fact that  $z \in \mathcal{DM}^\infty_{\text{loc}}(\Omega)$  and to (6.5), we have that all but the first term in the previous are uniformly bounded with respect to  $\delta$ . Hence  $S_\delta(u)$  is bounded in  $BV_{\text{loc}}(\Omega)$  and we are allowed to pass to the limit in  $\delta$  (using once again weak lower semicontinuity in the first term) and Lebesgue's theorem for the remaining terms, getting

$$\int_{\Omega} |D\chi_{\{u>0\}}|\varphi + \int_{\Omega} z \cdot \nabla \varphi \chi_{\{u>0\}} \leq \int_{\Omega} h(u) f \chi_{\{u>0\}}\varphi.$$

Observe, in particular, that

$$\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega).$$

Therefore, recalling Proposition 2.1,

$$-(\text{div } z)\chi_{\{u>0\}}^* = -\text{div}(z\chi_{\{u>0\}}) + (z, D\chi_{\{u>0\}}),$$

and so

$$-\int_{\Omega} \varphi \chi_{\{u>0\}}^* \text{div } z \leq \int_{\Omega} h(u) f \chi_{\{u>0\}}\varphi \leq \int_{\Omega} h(u) f \varphi. \quad (6.8)$$

We prove the reverse inequality. In (6.5) we take  $\varphi = (\chi_{\{u>0\}} * \rho_\epsilon)\phi$  where  $0 \leq \phi \in C^1_c(\Omega)$  and  $\rho_\epsilon$  is a mollifier. Passing to the limit in  $\epsilon$  (Lebesgue's theorem on the left hand side and Fatou's lemma on the right hand side) we obtain

$$-\int_{\Omega} \phi \chi_{\{u>0\}}^* \text{div } z \geq \int_{\Omega} h(u) f \chi_{\{u>0\}}\phi \stackrel{\{u=0\} \subseteq \{f=0\}}{=} \int_{\Omega} h(u) f \phi \quad \forall \phi \in C^1_c(\Omega), \phi \geq 0, \quad (6.9)$$

then (6.8) and (6.9) imply that (6.2) holds.  $\square$

7. THE CASE  $f$  IN  $L^{N,\infty}(\Omega)$ 

The main results proven in the previous section can be extended to the case of a slightly more general nonnegative datum in the Lorentz space  $f \in L^{N,\infty}(\Omega)$ , also called Marcinkiewicz space, this extension being optimal in some sense (see Remark 7.2 below). We refer, for instance, to the monograph [38] for a smooth introduction to the subject of Lorentz spaces and their main properties. We only recall that an Hölder's inequality is available in this case and that the conjugate space associated to  $L^{p,q}(\Omega)$  for  $p > 1$  and  $q \in [1, \infty]$  is  $L^{p',q'}(\Omega)$ . Also, a Sobolev embedding inequality for  $W_0^{1,p}(\Omega)$  holds, that is

$$\|v\|_{L^{p^*,p}(\Omega)} \leq \tilde{\mathcal{S}}_p \|\nabla v\|_{W_0^{1,p}(\Omega)}, \quad \forall v \in W_0^{1,p}(\Omega). \quad (7.10)$$

The involved constants are explicit and one has

$$\tilde{\mathcal{S}}_p = \frac{p\Gamma(1 + \frac{N}{2})^{\frac{1}{N}}}{\sqrt{\pi}(N-p)} \xrightarrow{p \rightarrow 1^+} \tilde{\mathcal{S}}_1 = [(N-1)\omega_N^{\frac{1}{N}}]^{-1},$$

where  $\Gamma$  is the usual Gamma function (see [2, 14]).

We consider problem

$$\begin{cases} -\Delta_1 u = h(u)f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (7.11)$$

where  $f \in L^{N,\infty}(\Omega)$  is nonnegative and  $h$ , as before, is a continuous function satisfying (h1) and (h2). Definition 6.1 can be straightforwardly re-adapted to this case with many simplifications if  $f > 0$  (as in Definition 3.1). We summarize the results one can obtain in the following

**Theorem 7.1.** *Let  $0 \leq f \in L^{N,\infty}(\Omega)$  such that  $\|f\|_{L^{N,\infty}(\Omega)} < \frac{1}{\tilde{\mathcal{S}}_1 h(\infty)}$ , where  $h$  satisfies (h1) and (h2). Then there exists a (unique, if  $h$  is decreasing and  $f > 0$  a.e. in  $\Omega$ ) solution  $u$  to problem (7.11) in the sense of Definition 6.1.*

*Moreover, if  $h(0) = \infty$  then  $u > 0$  a.e. in  $\Omega$ , and, if  $h \in L^\infty([0, \infty))$  and*

$$\|f\|_{L^{N,\infty}(\Omega)} < \frac{1}{\tilde{\mathcal{S}}_1 \|h\|_{L^\infty([0, \infty))}}, \quad (7.12)$$

*then  $u \equiv 0$  a.e. in  $\Omega$ .*

**Remark 7.2.** Observe that condition (7.12) is optimal in the sense that, if  $h \equiv 1$ , one can construct a datum  $\bar{f}$  with  $\|\bar{f}\|_{L^{N,\infty}(\Omega)} = \tilde{\mathcal{S}}_1^{-1}$  such that problem (7.11) relative to  $\bar{f}$  admits a nontrivial solution (ie.  $u \neq 0$ ) (see [16, Theorem 3.4, Remark 3.2]).

As far as the proofs of our existence, uniqueness and regularity results in the case  $f \in L^N(\Omega)$  are concerned, the proof of Theorem 7.1 is a standard re-adaptation once the analogous of Lemma 4.9 is established. That is, if we consider

$$\begin{cases} -\Delta_p u_n = h_n(u_n)f_n & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega, \end{cases} \quad (7.13)$$

where  $f \in L^{N,\infty}(\Omega)$  is nonnegative and  $f_n = T_n(f)$ ,  $h_n(s) = T_n(h(s))$ . Hence Theorem 7.1 is a consequence of the following

**Lemma 7.3.** *Let  $u_n$  be a solution to (7.13). If  $\|f\|_{L^{N,\infty}(\Omega)} < \frac{1}{\tilde{\mathcal{S}}_1 h(\infty)}$  then there exists  $\bar{k} > 0$  such that  $u_n$  satisfies:*

$$\|G_k(u_n)\|_{L^{\frac{N}{N-1}}(\Omega)} \leq \frac{p-1}{p} |\Omega| C(\tilde{\mathcal{S}}_1, h(\infty), \|f\|_{L^{N,\infty}(\Omega)}, \bar{k}), \quad \text{for all } k \geq \bar{k}, \quad (7.14)$$

$$\|u_n\|_{W^{1,p}(\omega)} \leq C(\tilde{\mathcal{S}}_1, h(\infty), \|f\|_{L^{N,\infty}(\Omega)}, \omega, |\Omega|), \quad \text{for all } \omega \subset\subset \Omega, \quad (7.15)$$

$$\|u_n\|_{W_0^{1,p}(\Omega)}^{\frac{\sigma-1+p}{p}} \leq C(\tilde{\mathcal{S}}_1, \sup_{s \in [k_0, \infty)} h(s), \|f\|_{L^{N,\infty}(\Omega)}, c_1, |\Omega|), \quad (7.16)$$

*Proof.* The proofs of both (7.14) and (7.15) strictly follow the ones who led to (4.36) and (4.38) where, systematically, the Hölder inequality is replaced by the generalized Hölder inequality in Lorentz spaces and (7.10) substitutes the usual Sobolev's embedding inequality in Lebesgue's spaces. The only estimate that needs some further efforts is (7.16) and we focus on it. As in (4.41) one fixes  $k_1 > k_0$  and then multiplies (7.13) by  $u_n^\sigma$ , obtaining

$$\begin{aligned} \left(\frac{p}{\sigma-1+p}\right)^p \sigma \int_{\Omega} |\nabla u_n^{\frac{\sigma-1+p}{p}}|^p &= \int_{\Omega} h_n(u_n) f u_n^\sigma \leq c_1 k_0^{\sigma-\gamma} \int_{\{u_n \leq k_0\}} f \\ &+ \max_{s \in [k_0, k_1]} h(s) k_1^\sigma \int_{\{k_0 < u_n < k_1\}} f + (h(\infty) + \epsilon_{k_1}) \int_{\{u_n \geq k_1\}} f u_n^\sigma \\ &\leq C(k_0, \max_{s \in [k_0, k_1]} h(s), \|f\|_{L^1(\Omega)}, c_1) \\ &+ (h(\infty) + \epsilon_{k_1}) \|f\|_{L^{N,\infty}(\Omega)} \|u_n^\sigma\|_{L^{\frac{N}{N-1},1}(\Omega)}. \end{aligned} \quad (7.17)$$

Now, let  $u^*(t)$  be the nonincreasing rearrangement of  $u$  for  $t \in (0, |\Omega|)$ , and observe that  $(u^q)^* = (u^*)^q$  for  $q > 0$ . Using the Hölder inequality with exponent  $\frac{\sigma-1+p}{\sigma}$  we have

$$\begin{aligned} \|u_n^\sigma\|_{L^{\frac{N}{N-1},1}(\Omega)} &= \int_0^{|\Omega|} t^{-\frac{1}{N}} u_n^*(t)^\sigma dt \leq \left( \int_0^{|\Omega|} t^{-\frac{\sigma-1+p}{N\sigma}} u_n^*(t)^{\sigma-1+p} dt \right)^{\frac{\sigma}{\sigma-1+p}} |\Omega|^{\frac{p-1}{\sigma-1+p}} \\ &= \left( \int_0^{|\Omega|} \left( t^{-\frac{\sigma(1-N)-1+p}{Np\sigma}} u_n^*(t)^{\frac{\sigma-1+p}{p}} \right)^p \frac{dt}{t} \right)^{\frac{\sigma}{\sigma-1+p}} |\Omega|^{\frac{p-1}{\sigma-1+p}} \\ &= \left( \int_0^{|\Omega|} \left( t^{\frac{1}{p^*}} u_n^*(t)^{\frac{\sigma-1+p}{p}} \right)^p t^{\frac{(\sigma-1)(p-1)}{N\sigma}} \frac{dt}{t} \right)^{\frac{\sigma}{\sigma-1+p}} |\Omega|^{\frac{p-1}{\sigma-1+p}} \\ &\leq \left( \int_0^{|\Omega|} \left( t^{\frac{1}{p^*}} u_n^*(t)^{\frac{\sigma-1+p}{p}} \right)^p \frac{dt}{t} \right)^{\frac{\sigma}{\sigma-1+p}} |\Omega|^{\frac{(p-1)(\sigma-1+N)}{N(\sigma-1+p)}} \\ &= \|u_n^{\frac{\sigma-1+p}{p}}\|_{L^{p^*,p}(\Omega)}^{\frac{p\sigma}{\sigma-1+p}} |\Omega|^{\frac{(p-1)(\sigma-1+N)}{N(\sigma-1+p)}} \\ &\leq \frac{\sigma}{\sigma-1+p} \|u_n^{\frac{\sigma-1+p}{p}}\|_{L^{p^*,p}(\Omega)}^p + \frac{p-1}{\sigma-1+p} |\Omega|^{\frac{\sigma-1+N}{N}}, \end{aligned}$$

where in the last step we used Young's inequality. Concerning the left hand side of (7.17) we have

$$\left(\frac{p}{\sigma-1+p}\right)^p \sigma \int_{\Omega} |\nabla u_n^{\frac{\sigma-1+p}{p}}|^p \geq \left(\frac{p}{\sigma-1+p}\right)^p \frac{\sigma}{\mathcal{F}_p^p} \|u_n^{\frac{\sigma-1+p}{p}}\|_{L^{p^*,p}(\Omega)}^p$$

and gathering the previous two inequalities with (7.17) we deduce

$$\left( \left(\frac{p}{\sigma-1+p}\right)^p \frac{\sigma}{\mathcal{F}_p^p} - (h(\infty) + \epsilon_{k_1}) \|f\|_{L^{N,\infty}(\Omega)} \frac{\sigma}{\sigma-1+p} \right) \|u_n^{\frac{\sigma-1+p}{p}}\|_{L^{p^*,p}(\Omega)}^p \leq C, \quad (7.18)$$

where, since  $\|f\|_{L^{N,\infty}(\Omega)} < \frac{1}{\mathcal{F}_1 h(\infty)}$ , one can pick  $p$  near to 1 and  $k_1$  large enough such that

$$\left( \left(\frac{p}{\sigma-1+p}\right)^p \frac{\sigma}{\mathcal{F}_p^p} - (h(\infty) + \epsilon_{k_1}) \|f\|_{L^{N,\infty}(\Omega)} \frac{\sigma}{\sigma-1+p} \right) > c > 0$$

for a constant  $c$  that does not depend on both  $p$  and  $k_1$ . Using estimate (7.18) in (7.17) one finally deduces (7.16).  $\square$

## 8. FURTHER EXTENSIONS, REMARKS, AND EXAMPLES

**8.1. Some global BV solutions.** As we have seen the fact  $u^\sigma \in BV(\Omega)$  is crucial in order to prove the existence and uniqueness of a solution to (3.1). Due to the possible degeneracy of the datum  $f$

and to the weak requests one assumes on the nonlinearity  $h$  this step seems to be needed, in general, in order to conclude.

Although, a natural question is whether the solution to (3.1) enjoys itself the further property to have global finite energy in the natural space  $BV(\Omega)$ . To fix the ideas, if  $\Omega$  is a smooth domain, consider the problem

$$\begin{cases} -\Delta_p u = \frac{f}{u^\gamma} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (8.1)$$

where  $\gamma > 0$  and  $f$  is an Hölder continuous function that is bounded away from zero on  $\Omega$ . If  $p = 2$  and  $\gamma \geq 1$  it can be proven that solutions belong to the natural space  $H_0^1(\Omega)$  if and only if  $\gamma < 3$  ([29], see also [41, 36] for further refinements). Extensions to the case  $p > 1$  are also available ([40]) and the threshold becomes  $\gamma < \frac{2p-1}{p-1}$ , suggesting that, as  $p \rightarrow 1^+$ , one should recover the global  $BV$  regularity of the solutions for any  $\gamma > 0$  at least for both a nondegenerate datum and a smooth domain. Recall that, as for the case  $p > 1$  with sufficiently integrable data, if  $\gamma \leq 1$  (see [21]) then solutions to problem (8.1) always have finite energy if  $p = 1$  (compare with Theorem 3.4 above).

In order to better understand this phenomenon one can look at the proof of the result in [29]. One immediately realizes that, at regular boundary points, the slope of the solutions to (8.1) become larger and larger as  $\gamma$  grows eventually leading the solution to loose its  $C^1$  regularity (beyond  $\gamma = 1$ ) and its  $H^1$  regularity (at  $\gamma = 3$ ). Hence, due to the fact that the boundary datum needs not to be attained in the classical sense, and to the particular nature of  $BV$  (that allows jumps), it seems reasonable that, for any fixed  $\gamma > 0$  solutions to (8.1) may become globally  $BV$  as  $p$  reaches 1 (at least if the datum  $f$  does not degenerate at zero).

In this section we want to present some further evidences of this fact. Though a more general right hand side can be considered, in order to simplify the exposition we consider the simplest model

$$\begin{cases} -\Delta_1 u = u^{-\gamma} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (8.2)$$

We will construct an example showing that, for a rich enough class of domains, solutions to (8.2) belongs to  $BV(\Omega)$ , for any  $\gamma > 0$ . We first need the following

**Definition 8.1.** We say that a bounded convex set  $E$  of class  $C^{1,1}$  is *calibrable* if there exists a vector field  $\xi \in L^\infty(\mathbb{R}^N, \mathbb{R}^N)$  such that  $\|\xi\|_\infty \leq 1$ ,  $(\xi, D\chi_E) = |D\chi_E|$  as measures, and

$$-\operatorname{div} \xi = \lambda_E \chi_E \quad \text{in } \mathcal{D}'(\mathbb{R}^N)$$

for some constant  $\lambda_E$ . In this case  $\lambda_E = \frac{\operatorname{Per}(E)}{|E|}$  and  $[\xi, \nu^E] = -1$ ,  $\mathcal{H}^{N-1}$ -a.e in  $\partial E$  (see [1, Section 2.3] and [34]).

As a consequence of [1, Theorem 9] a bounded and convex set  $E$  is calibrable if and only if the following condition holds:

$$(N-1)\|\mathbf{H}_E\|_{L^\infty(\partial E)} \leq \lambda_E = \frac{\operatorname{Per}(E)}{|E|},$$

where  $\mathbf{H}_E$  denotes the ( $\mathcal{H}^{N-1}$ -a.e. defined) mean curvature of  $\partial E$ . In particular, if  $E = B_R(0)$ , for some  $R > 0$ , then  $E$  is calibrable.

**Example 1.** If  $\Omega$  is a calibrable set, let us prove that  $u = \left(\frac{|\Omega|}{\operatorname{Per}(\Omega)}\right)^{\frac{1}{\gamma}}$  is the unique solution to (8.2) in the sense of Definition 3.1. It suffices to take the restriction to  $\Omega$  of the vector field in the definition of calibrability; i.e.:  $z := \xi|_{\Omega}$ . In fact, due to the properties of  $\xi$  one has

$$-\operatorname{div} z = \frac{\operatorname{Per}(\Omega)}{|\Omega|} = u^{-\gamma} \quad \text{and} \quad [\xi, \nu^\Omega] = -1. \quad (8.3)$$

Moreover, using both (2.5) and (8.3), one finally gets

$$\begin{aligned} (z, Du)(\Omega) &= \int_{\Omega} \left( \frac{|\Omega|}{Per(\Omega)} \right)^{\frac{1}{\gamma}} \frac{Per(\Omega)}{|\Omega|} dx \\ &+ \int_{\partial\Omega} [\xi, \nu^{\Omega}] \left( \frac{|\Omega|}{Per(\Omega)} \right)^{\frac{1}{\gamma}} d\mathcal{H}^{N-1} = 0 = |Du|(\Omega). \end{aligned}$$

**Remark 8.2.** Observe that the solutions of (8.2) given by Theorem 3.3 belong to  $BV(\Omega)$  once they are bounded away from zero on  $\Omega$ ; in fact, let  $u \geq a > 0$ , then, due to property (3.8) one has  $u = S(u^\gamma) \in BV(\Omega)$ , where  $S$  is the Lipschitz continuous function defined by

$$S(s) = \max(a, s^{\frac{1}{\gamma}}).$$

Let us show a situation where, at least formally,  $u \geq a > 0$  holds. Let  $\Omega$  be a convex open set. In [34] it is shown that  $-H_{\Omega}(x)$  is a (so called) large solution to  $\Delta_1 v = v$ , i. e.

$$\begin{cases} \Delta_1 v = v & \text{in } \Omega, \\ v = \infty & \text{on } \partial\Omega, \end{cases} \quad (8.4)$$

where  $H_{\Omega}(x)$  is the variational mean curvature of  $\Omega$  (see [8] for details). Without entering into technicalities, only recall that  $\|H_{\Omega}\|_{L^{\infty}(\mathbb{R}^N)} < \infty$  if and only if  $\Omega$  is of class  $C^{1,1}$ ; in particular, these solutions only assume the (large) datum  $\infty$  at nonregular points of  $\Omega$  (e.g. at corners).

As through the change of variable  $u = v^{-\frac{1}{\gamma}}$  problem (8.4) formally transforms into (8.2) (using also the homogeneity of the operator) then one can expect that solutions to problem (8.2) always belong to  $BV(\Omega)$  if  $\Omega$  is a convex bounded  $C^{1,1}$  domain and that, in general, the Dirichlet homogeneous boundary datum is only assumed pointwise at nonsmooth points of  $\Omega$ .

**8.2. More general growths.** Here we show how the assumption on the control of  $h$  near zero can be removed allowing more general growths not satisfying (h1). Consider the general problem

$$\begin{cases} -\Delta_1 u = F(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (8.5)$$

where  $F(x, s)$  is a nonnegative Carathéodory function satisfying

$$F(x, s) \leq h(s)f(x), \quad \forall (x, s) \in \Omega \times [0, \infty), \quad (8.6)$$

with  $0 \leq f \in L^{N, \infty}(\Omega)$ , and  $h$  is a continuous function in  $[0, \infty)$  satisfying (h2).

First of all, without loss of generality, we can assume that  $h$  such that

$$h \text{ is decreasing, } h \in C^1((0, \infty)), h^{-1} \in C^1([0, \infty)), h^{-1}(0) = 0, \quad (8.7)$$

where  $h^{-1}$  stands for the reciprocal of  $h$ . Indeed, for any given  $h$  satisfying our assumptions one can construct (see for instance [25, Remark 2.1, vii]) a function  $\bar{h}$  such that (8.7) holds and

$$h(s) \leq \bar{h}(s), \quad \forall s \geq 0.$$

As for the previous sections we look for a solution of (8.5) through an approximation argument, letting  $p \rightarrow 1^+$  in the solutions to

$$\begin{cases} -\Delta_p u_p = F(x, u_p) & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega. \end{cases} \quad (8.8)$$

The notion of solution to (8.8) for  $p > 1$  is the following one.

**Definition 8.3.** A nonnegative function  $u_p \in W_{\text{loc}}^{1,p}(\Omega)$  is a *distributional solution* to problem (8.8) if

$$F(x, u_p) \in L_{\text{loc}}^1(\Omega), \quad (8.9)$$

$$G_k(u_p) \in W_0^{1,p}(\Omega), \quad \text{for all } k > 0, \quad (8.10)$$

and

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} F(x, u) \varphi, \quad (8.11)$$

for every  $\varphi \in C_c^1(\Omega)$ .

We have the following result whose proof, using (8.6), easily follows line by line the proof of Theorem 4.3. Only observe that, in this case, the approximating problems read as

$$\begin{cases} -\Delta_p u_n = F_n(x, u_n) & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega, \end{cases} \quad (8.12)$$

where  $F_n(x, s) = F(x, T_n(s))$ . Moreover we set

$$\beta_p(s) = \int_0^s ((h^{-1}(t))')^{\frac{1}{p}} dt. \quad (8.13)$$

**Theorem 8.4.** *Let  $F$  satisfy (8.6) where  $0 \leq f \in L^{(p^*)'}(\Omega)$  and let  $h$  satisfy (8.7) and (h2). Then there exists a solution  $u_p$  to problem (8.8) in the sense of Definition 8.3 such that  $\beta_p(u_p) \in W_0^{1,p}(\Omega)$ .*

Moreover the following counterpart of (4.39) can be proven:

$$\int_{\Omega} |\nabla \Gamma_{-p}(u_p)|^p \leq \int_{\Omega} F(x, u_p) \underline{\psi}_{-p}(u_p), \quad (8.14)$$

where, for  $\delta > 0$ ,

$$\underline{\psi}_{-p}(s) = \begin{cases} h^{-1}(s) \frac{\beta_p(\delta)}{h^{-1}(\delta)} & \text{if } s < \delta, \\ \beta_p(s) & \text{if } s \geq \delta, \end{cases}$$

and  $\Gamma_{-p}$  is the primitive of  $\underline{\psi}_{-p}^{\frac{1}{p}}(s)$  (such that  $\Gamma_{-p} = 0$ ).

Here is how the definition of solution to problem (8.5) can be suitably modified:

**Definition 8.5.** A function  $u \in BV_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$  having  $\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega)$  and  $\beta_1(u) \in BV(\Omega)$  ( $\beta_1$  is defined in (8.13)) is a solution to problem (8.5) if there exists  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  such that

$$F(x, u) \in L_{\text{loc}}^1(\Omega), \quad (8.15)$$

$$-(\text{div } z) \chi_{\{u>0\}}^* = F(x, u) \quad \text{in } \mathcal{D}'(\Omega), \quad (8.16)$$

$$(z, Du) = |Du| \quad \text{as measures in } \Omega, \quad (8.17)$$

$$\beta_1(u(x)) + [\beta_1(u)z, \nu](x) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega. \quad (8.18)$$

One finally has the following

**Theorem 8.6.** *Let  $0 \leq f \in L^{N,\infty}(\Omega)$  such that  $\|f\|_{L^{N,\infty}(\Omega)} < \frac{1}{\mathcal{F}_1 h(\infty)}$  and let  $h$  satisfy (h2). Then there exists a solution  $u$  to problem (8.5) in the sense of Definition 8.5. Moreover if  $F(x, 0) = \infty$  then  $u > 0$  a.e. in  $\Omega$ . Otherwise if  $F(x, 0) < \infty$  and if  $\|f\|_{L^{N,\infty}(\Omega)} < (\mathcal{F}_1 \|h\|_{L^\infty(\{0,\infty\})})^{-1}$  then  $u \equiv 0$  a.e. in  $\Omega$ .*

*Proof.* The proof is a suitable modification of the one of Theorems 3.3, 6.4, and 7.1; we only highlight the main differences. One starts with the solutions  $u_n$  of (8.12). As  $f \in L^{N,\infty}(\Omega)$  with  $\|f\|_{L^{N,\infty}(\Omega)} < \frac{1}{\mathcal{F}_1 h(\infty)}$ , then, uniform estimates hold. In fact, using (8.6), both (7.14) and (7.15) continue to hold, and, recalling (8.7), one can show that  $p_0$  exists such that for any  $p \in (1, p_0)$

$$\|\beta_p(u_n)\|_{W_0^{1,p}(\Omega)} \leq C(\mathcal{F}_1, \sup_{s \in [1,\infty)} h(s), \|f\|_{L^{N,\infty}(\Omega)}). \quad (8.19)$$

This is done by considering the solutions to (8.12) and taking  $h^{-1}(u_n)$  as test. By weak lower semi-continuity the same holds for  $u_p$ . One then deduces, reasoning as in the proof of Lemma 5.2, the

existence of a vector field  $z \in \mathcal{D}\mathcal{M}_{\text{loc}}^\infty(\Omega)$  with  $\|z\|_\infty \leq 1$  which is the  $*$ -weak limit of  $|\nabla u_p|^{p-2} \nabla u_p$ . The proof of (8.16) can be derived as for (6.2). Now recalling (8.19), we have

$$\int_{\Omega} |\nabla \beta_p(u_p)| \leq \frac{1}{p} \int_{\Omega} |\nabla \beta_p(u_p)|^p + \frac{1}{p'} |\Omega| \leq C,$$

which implies that  $\beta_p(u_p)$  is bounded in  $BV(\Omega)$  and then it strongly converges in  $L^q(\Omega)$  to its a.e. limit  $\beta_1(u) \in BV(\Omega)$ . Moreover  $\nabla \beta_p(u_p)$  converges  $*$ -weakly in the sense of measures to  $D\beta_1(u)$ . Hence reasoning exactly as in the proof of (5.10) it yields

$$-\beta_1(u)^* \operatorname{div} z = F(x, u) \beta_1(u) \text{ in } \mathcal{D}'(\Omega). \quad (8.20)$$

In order to prove (8.17), one lets  $0 \leq \varphi \in C_c^1(\Omega)$  and  $\beta_p(u_p)\varphi$  to test (8.8) obtaining

$$\int_{\Omega} \varphi |\nabla u_p|^p \beta_p'(u_p) + \int_{\Omega} \beta_p(u_p) |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi = \int_{\Omega} F(x, u_p) \beta_p(u_p) \varphi.$$

Thus, by Young's inequality, one deduces

$$\int_{\Omega} \varphi |\nabla u_p \beta_p'(u_p)^{\frac{1}{p}}| + \int_{\Omega} \beta_p(u_p) |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi \leq \int_{\Omega} F(x, u_p) \beta_p(u_p) \varphi + \frac{p-1}{p} \int_{\Omega} \varphi.$$

Reasoning as in the proof of Lemma 5.2 all but the first term are shown to pass to the limit with respect to  $p$ . Then by lower semicontinuity we obtain, recalling (8.20),

$$\int_{\Omega} \varphi |D\beta_1(u)| + \int_{\Omega} \beta_1(u) z \cdot \nabla \varphi \leq \int_{\Omega} F(x, u) \beta_1(u) \varphi = - \int_{\Omega} (\beta_1(u))^* \varphi \operatorname{div} z, \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0,$$

and we can apply Proposition 2.1 to deduce

$$\int_{\Omega} \varphi |D\beta_1(u)| \leq - \int_{\Omega} \beta_1(u) z \cdot \nabla \varphi - \int_{\Omega} \beta_1(u)^* \varphi \operatorname{div} z = \int_{\Omega} \varphi (z, D\beta_1(u)), \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0,$$

then

$$\int_{\Omega} \varphi |D\beta_1(u)| \leq \int_{\Omega} \varphi (z, D\beta_1(u)), \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0,$$

that implies  $|D\beta_1(u)| = (z, D\beta_1(u))$  as  $\|z\|_\infty \leq 1$ . Now since  $\beta_1(s)$  is locally Lipschitz one obtains (8.17) by the same argument as in the last step of the proof of Lemma 5.2.

It is left to prove (8.18). It follows by using (8.14) and Young's inequality that

$$\int_{\Omega} |\nabla \Gamma_p(u_p)| + \int_{\partial\Omega} \beta_p(u_p) d\mathcal{H}^{N-1} \leq \int_{\Omega} F(x, u_p) \psi_p^-(u_p) + \frac{p-1}{p} |\Omega|,$$

then by weak lower semicontinuity one can use Gauss-Green formula (2.5) and (8.20) to have

$$\int_{\Omega} |D\beta_1(u)| + \int_{\partial\Omega} \beta_1(u) d\mathcal{H}^{N-1} \leq \int_{\Omega} (z, D\beta_1(u)) - \int_{\partial\Omega} [\beta_1(u) z, \nu] d\mathcal{H}^{N-1},$$

that, as  $|D\beta_1(u)| = (z, D\beta_1(u))$ , gives (8.18).  $\square$

## REFERENCES

- [1] F. Alter, V. Caselles and A. Chambolle: A characterization of convex calibrable sets in  $\mathbb{R}^N$ . *Math. Ann.* **332** 329–366 (2005)
- [2] A. Alvino: Sulla disuguaglianza di Sobolev in spazi di Lorentz. *Boll. Un. Mat. Ital. A (5)* **14**(1) 148–156 (1977)
- [3] L. Ambrosio, N. Fusco and D. Pallara: *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford Mathematical Monographs, 2000
- [4] F. Andreu, C. Ballester, V. Caselles and J. M. Mazón: The Dirichlet problem for the total variation flow. *J. Funct. Anal.* **180** (2), 347–403 (2001)
- [5] F. Andreu, A. Dall'Aglio and S. Segura de León: Bounded solutions to the 1-Laplacian equation with a critical gradient term. *Asymptot. Anal.* **80** (1-2), 21–43 (2012)
- [6] F. Andreu, V. Caselles and J. M. Mazón: *Parabolic quasilinear equations minimizing linear growth functionals*. Progress in Mathematics, 223, Birkhäuser Verlag, Basel, 2004
- [7] G. Anzellotti: Pairings between measures and bounded functions and compensated compactness. *Ann. Mat. Pura Appl.* **135** (4), 293–318 (1983)
- [8] E. Barozzi, E. Gonzalez, and I. Tamanini: The mean curvature of a set of finite perimeter. *Proc. Amer. Math. Soc.* **99**, 313–316 (1987)

- [9] M. Bertalmio, V. Caselles, B. Rougé, A. Solé: TV based image restoration with local constraints. Special issue in honor of the sixtieth birthday of Stanley Osher, *J. Sci. Comput.* **19** (1-3), 95–122 (2003)
- [10] L. Boccardo and F. Murat: Almost Everywhere Convergence of the Gradients of Solutions to Elliptic and Parabolic Equations. *Nonlinear Anal.* **19**, 581–597 (1992)
- [11] L. Boccardo and L. Orsina, Semilinear elliptic equations with singular nonlinearities. *Calc. Var. Partial Differential Equations* **37** (2010), 363–380.
- [12] A. Canino, B. Sciunzi and A. Trombetta: Existence and uniqueness for  $p$ -Laplace equations involving singular nonlinearities. *NoDEA Nonlinear Differential Equations Appl.* **23**, 8–18 (2016)
- [13] V. Caselles: On the entropy conditions for some flux limited diffusion equations. *J. Differential Equations* **250**, 3311–3348 (2011)
- [14] D. Cassani, B. Ruf, C. Tarsi: Optimal Sobolev type inequalities in Lorentz spaces. *Potential Anal.* **39** (3), 265–285 (2013)
- [15] G.-Q. Chen and H. Frid: Divergence–measure fields and hyperbolic conservation laws. *Arch. Ration. Mech. Anal.* **147** (2), 89–118 (1999)
- [16] M. Cicalese and C. Trombetti: Asymptotic behaviour of solutions to  $p$ -Laplacian equation. *Asymptotic Anal.* **35**, 27–40 (2003)
- [17] G. Crasta and V. De Cicco: Anzellotti’s pairing theory and the Gauss–Green theorem. Preprint: arXiv:1708.00792
- [18] L. De Cave: Nonlinear elliptic equations with singular nonlinearities. *Asymptot. Anal.* **84** (3-4), 181–195 (2013)
- [19] L. M. De Cave and F. Oliva: Elliptic equations with general singular lower order term and measure data. *Nonlinear Anal.* **128**, 391–411 (2015)
- [20] L. M. De Cave, R. Durastanti and F. Oliva: Existence and uniqueness results for possibly singular nonlinear elliptic equations with measure data. Preprint: arXiv:1709.06042
- [21] V. De Cicco, D. Giachetti and S. Segura de León: Elliptic problems involving the 1-laplacian and a singular lower order term. Preprint
- [22] F. Demengel: On some nonlinear partial differential equations involving the "1"-Laplacian and critical Sobolev exponent. *ESAIM Control Optim. Calc. Var.* **4**, 667–686 (1999)
- [23] L. C. Evans and R.F. Gariepy: *Measure Theory and Fine Properties of Functions*. Studies in Advanced Mathematics, CRC Press, 1992
- [24] D. Giachetti, P.J. Martínez-Aparicio and F. Murat: A semilinear elliptic equation with a mild singularity at  $u = 0$ : Existence and homogenization. *J. Math. Pures Appl.* **107**, 41–77 (2017)
- [25] D. Giachetti, P. J. Martínez-Aparicio and F. Murat: Definition, existence, stability and uniqueness of the solution to a semilinear elliptic problem with a strong singularity at  $u = 0$ . *Ann. Scuola Normale Pisa*, in press
- [26] B. Kawohl: On a family of torsional creep problems. *J. Reine Angew. Math.* **410**, 1–22 (1990)
- [27] B. Kawohl: From  $p$ -Laplace to mean curvature operator and related questions. In : *Progress in Partial Differential Equations: the Metz Surveys*, Pitman Res. Notes Math. Ser., Vol. 249, Longman Sci. Tech., Harlow, pp. 40–56, 1991
- [28] B. Kawohl, F. Schuricht: Dirichlet problems for the 1-Laplace operator, including the eigenvalue problem. *Commun. Contemp. Math.* **9** (4), 515–543 (2007)
- [29] A. C. Lazer and P. J. McKenna: On a singular nonlinear elliptic boundary-value problem. *Proc. Amer. Math. Soc.* **111**, 721–730 (1991)
- [30] Y. Meyer: *Oscillating Patterns in Image Processing and Nonlinear Evolution Equations: The Fifteenth Dean Jacqueline B. Lewis Memorial Lectures*. Providence, RI: American Mathematical Society, 2001
- [31] A. Mercaldo, S. Segura de León and C. Trombetti: On the behaviour of the solutions to  $p$ -Laplacian equations as  $p$  goes to 1. *Publ. Mat.* **52**, 377–411 (2008)
- [32] A. Mercaldo, S. Segura de León and C. Trombetti: On the solutions to 1-Laplacian equation with  $L^1$  data. *J. Funct. Anal.* **256** (8), 2387–2416 (2009)
- [33] A. Molino, S. Segura de León: Elliptic equations involving the 1-Laplacian and a subcritical source term. Preprint: arXiv:1707.01934
- [34] S. Moll and F. Petitta: Large solutions for the elliptic 1-laplacian with absorption. *J. Anal. Math.* **125** (1), 113–138 (2015)
- [35] F. Oliva and F. Petitta: On singular elliptic equations with measure sources. *ESAIM Control Optim. Calc. Var.* **22** (1), 289–308 (2016)
- [36] F. Oliva and F. Petitta: Finite and infinite energy solutions of singular elliptic problems: Existence and uniqueness. *J. Differential Equations* **264** (1), 311–340. (2018)
- [37] S. Osher, J. Sethian: Fronts propagating with curvature-dependent speed: algorithms based on Hamilton-Jacobi formulations. *Journal of Computational Physics* **79** (1), 12–49 (1988)
- [38] L. Pick, A. Kufner, J. Oldrich, S. Fucik: *Function Spaces*, 1. Berlin, Boston: De Gruyter, 2012
- [39] G. Sapiro: *Geometric partial differential equations and image analysis*. Cambridge University Press, 2001.
- [40] G. Singh: Regularity of weak solutions for singular elliptic problems driven by  $m$ -Laplace operator. arXiv:1511.03219v1
- [41] Y. Sun, D. Zhang: The role of the power 3 for elliptic equations with negative exponents. *Calc. Var. Partial Differential Equations* **49** (3-4) 909–922 (2014)
- [42] G. Talenti: Best constant in Sobolev inequality. *Ann. Mat. Pura Appl.* **110** 353–372 (1976)
- [43] W. P. Ziemer: *Weakly differentiable functions. Sobolev spaces and functions of bounded variation*. Graduate Texts in Mathematics, 120, Springer–Verlag, New York, 1989



DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L' INGEGNERIA, "SAPIENZA" UNIVERSITÀ DI ROMA, VIA SCARPA 16, 00161 ROMA, ITALIA:

(Virginia De Cicco) VIRGINIA.DECICCO@SBAI.UNIROMA1.IT

(Daniela Giachetti) DANIELA.GIACHETTI@SBAI.UNIROMA1.IT

(Francescantonio Oliva) FRANCESCO.OLIVA@SBAI.UNIROMA1.IT

(Francesco Petitta) FRANCESCO.PETITTA@SBAI.UNIROMA1.IT