

On the notion of parallel transport on RCD spaces

Nicola Gigli* Enrico Pasqualetto†

February 28, 2018

Abstract

We propose a general notion of parallel transport on RCD spaces, prove an unconditioned uniqueness result and existence under suitable assumptions on the space.

MSC2010: 30Lxx, 51Fxx

Keywords: Parallel transport, RCD spaces.

Contents

1	Introduction	2
2	Some basic notions	3
2.1	Curves in Banach spaces	3
2.2	Pullback of an L^0 -normed module	7
2.3	Some properties of test plans	9
3	Introduction of appropriate functional spaces	11
3.1	Test vector fields along π	12
3.2	The space $\mathcal{L}^2(\pi)$	14
3.3	The space $\mathcal{C}(\pi)$	17
3.4	The spaces $\mathcal{W}^{1,2}(\pi)$ and $\mathcal{H}^{1,2}(\pi)$	18
4	Parallel transport on RCD spaces	25
4.1	Definition and basic properties of parallel transport	25
4.1.1	Definition and uniqueness	25
4.1.2	Some consequences of existence of parallel transport	25
4.2	Existence of the parallel transport in a special case	28
A	Sobolev base of the tangent module	32

*SISSA, ngigli@sissa.it

†SISSA, epasqual@sissa.it

1 Introduction

More than ten years ago Sturm [21, 22] and Lott-Villani [18] introduced the concept of lower Ricci curvature bounds for metric measure spaces. Their approach has been refined in [3] and [10] with the introduction of the class of metric measure spaces with Riemannian Ricci curvature bounded from below, RCD spaces in short, which is currently a very active research area. We refer to the surveys [23, 24] for an overview of the topic and detailed references.

Among the various recent contributions, of particular relevance for the current manuscript is the paper [8] by the first author, where a second order calculus has been built. In particular, on $\text{RCD}(K, \infty)$ spaces the covariant derivative of vector fields is well defined. Let us mention that in [8] ‘vector field’ is intended in the sense of abstract L^2 -normed L^∞ -modules and that in our previous paper [12] we showed that on $\text{RCD}(K, N)$ spaces this abstract notion has a canonically more concrete counterpart described in terms of pointed-measured-Gromov-Hausdorff limits of rescaled spaces (this uses the rectifiability results obtained in [19] and in [17], [11]).

In the classical smooth Riemannian framework, covariant derivative and parallel transport are two closely related concepts, thus given the existence of covariant derivative on RCD spaces it is natural to ask: is there a notion of parallel transport in the same setting? In this paper we address this question, our main results being:

- i) We provide a precise framework and give a rigorous meaning to the ‘PDE’ defining the parallel transport (see Definitions 3.19, 3.22 and 4.1).
- ii) By the nature of our definition, norm-preservation and linearity of the parallel transport are easy to derive, and these in turn will give uniqueness (see Corollary 4.3).
- iii) On RCD spaces satisfying a certain regularity property, we are able to show existence of the parallel transport (see Section 4.2). The regularity condition that we use is closely related to the existence of Sobolev vector fields with bounded covariant derivative (see Definition 4.9 for the precise assumption).

We believe that in fact the parallel transport exists on any RCD space, but we are currently unable to get the full proof. An insight on why this should not be too easy to prove is the following: on a space where the parallel transport exists, the dimension of the tangent module must be constant (see Theorem 4.8) and thanks to the aforementioned paper [12] this would in turn imply that the dimension of the pmGH-limits of rescaled spaces is constant. This very same result has been extremely elusive in the context of Ricci-limit spaces and has been obtained only relatively recently by Colding-Naber in [4]. Therefore it would be perhaps too optimistic to hope that the language proposed in [8] and here allows for an easy generalization of such ‘constant dimension’ result to the RCD setting. In this direction we remark that the assumptions that we put in order to obtain existence of the parallel transport are rather ad hoc and not really interesting from the geometric perspective: the intent with our existence result is just to show that the approach we propose is non-void. In our forthcoming paper [13] we shall study the same problem for more interesting geometric objects like Alexandrov spaces.

Let us also mention that in the appendix (see Theorem A.2) we prove that

- iv) Any $\text{RCD}(K, \infty)$ space admits a base of the tangent module made of vectors in $W_C^{1,2}(TX)$.

That is: if we relax the condition of ‘bounded covariant derivative’ present in (iii) above into ‘covariant derivative in L^2 ’, then every RCD space meets the requirement.

Let us now briefly describe our approach. The crucial idea is that we don’t study the problem of parallel transport along a single Lipschitz curve, but rather we study the problem along π -a.e. curve, where π is a Borel probability measure on the space $\Gamma(X)$ of continuous curves such that:

$$\begin{aligned} \pi &\text{ is concentrated on equi-Lipschitz curves,} \\ \text{for some } C > 0 &\text{ we have } (e_t)_*\pi \leq C\mathfrak{m} \text{ for every } t \in [0, 1], \end{aligned}$$

where \mathfrak{m} is the given reference measure on our RCD space X and $e_t : \Gamma(X) \rightarrow X$ is the evaluation map defined by $e_t(\gamma) := \gamma_t$. Measures π of this sort are a special case of so-called *test plans* introduced in [2]; these can be used to define Sobolev functions on metric measure spaces by a kind of duality argument. The advantage of working with these plans rather than with single curves is that they are naturally linked to Sobolev calculus and thus also to all the functional-analytic machinery built in [8].

Let us now pretend, for the sake of this introduction, that our space X is in fact a smooth Riemannian manifold. In this case a time dependent vector field (v_t) along π is, roughly said, given by a choice, for π -a.e. γ , of time dependent vector fields (v_t^γ) on X . Then we say that (v_t) is a parallel transport along π provided for π -a.e. γ the vector field $t \mapsto v_t^\gamma(\gamma_t)$ is a parallel transport along γ . This happens if and only if

$$\text{for } \pi\text{-a.e. } \gamma \text{ we have } \quad \partial_t v_t^\gamma + \nabla_{\gamma_t'} v_t^\gamma = 0 \quad \text{a.e. } t.$$

A relevant part of our paper (the whole Chapter 3) is devoted to showing that the above PDE can be stated even in the non-smooth setting, the key point being that it is possible to define a closed operator acting on L^2 vector fields along π which plays the role of $(\partial_t + \nabla_{\gamma_t'})$, see Definitions 3.17, 3.19 and Proposition 3.20.

We conclude this introduction recalling that Petrunin proved in [20] that a certain notion of parallel transport exists along geodesic on Alexandrov spaces. Uniqueness for his construction is still an open problem. The question of comparing our notion and his one is certainly interesting, but outside the scope of this manuscript.

Acknowledgments This research has been supported by the MIUR SIR-grant ‘Nonsmooth Differential Geometry’ (RBSI147UG4).

2 Some basic notions

To keep the presentation short, we shall assume the reader familiar with the language proposed in [8] (see also [6]).

2.1 Curves in Banach spaces

We recall here some basic results about measurability and integration of Banach-valued maps of a single variable $t \in [0, 1]$. A detailed discussion about this topic can be found e.g. in [5].

Let \mathbb{B} be a fixed Banach space. We will denote by \mathbb{B}' its dual space. A *simple function* is any map $y : [0, 1] \rightarrow \mathbb{B}$ that can be written in the form

$$y = \sum_{i=1}^k \chi_{E_i} v_i, \quad \text{for some } E_1, \dots, E_k \in \mathcal{B}([0, 1]) \text{ and } v_1, \dots, v_k \in \mathbb{B},$$

where for any topological space X we denote by $\mathcal{B}(X)$ the set of Borel subsets of X . A map $y : [0, 1] \rightarrow \mathbb{B}$ is said to be *strongly measurable* provided there exists a sequence $(y_n)_n$ of simple functions $y_n : [0, 1] \rightarrow \mathbb{B}$ such that $\lim_n \|y_n(t) - y(t)\|_{\mathbb{B}} = 0$ for \mathcal{L}^1 -a.e. $t \in [0, 1]$, while it is said to be *weakly measurable* provided $[0, 1] \ni t \mapsto \omega(y(t)) \in \mathbb{R}$ is a Borel map for every $\omega \in \mathbb{B}'$. It directly follows from the very definition that linear combinations of strongly (resp. weakly) measurable functions are strongly (resp. weakly) measurable. Moreover, if a map $y : [0, 1] \rightarrow \mathbb{B}$ is strongly measurable, then the function $[0, 1] \ni t \mapsto \|y(t)\|_{\mathbb{B}} \in \mathbb{R}$ is Borel.

The relation between the strongly measurable functions and the weakly measurable ones is fully described by a theorem of Pettis, which states that a function $y : [0, 1] \rightarrow \mathbb{B}$ is strongly measurable if and only if it is weakly measurable and there exists a Borel set $N \subseteq [0, 1]$ of null \mathcal{L}^1 -measure such that $y([0, 1] \setminus N)$ is a separable subset of \mathbb{B} .

We now describe how to define \mathbb{B} -valued integrals, the so-called *Bochner integrals*. First of all, given a simple function $y : [0, 1] \rightarrow \mathbb{B}$, written as $y = \sum_{i=1}^k \chi_{E_i} v_i$, we define

$$\int_0^1 y(t) dt := \sum_{i=1}^k \mathcal{L}^1(E_i) v_i \in \mathbb{B}.$$

It can be readily checked that this definition does not depend on the particular way of expressing the function y . Further, we say that any strongly measurable function $y : [0, 1] \rightarrow \mathbb{B}$ is *Bochner integrable* provided there exists a sequence $(y_n)_n$ of simple functions $y_n : [0, 1] \rightarrow \mathbb{B}$ such that $\lim_n \int_0^1 \|y_n(t) - y(t)\|_{\mathbb{B}} dt = 0$. In particular, the sequence $(\int_0^1 y_n(t) dt)_n \subseteq \mathbb{B}$ is Cauchy, so that it makes sense to define

$$\int_0^1 y(t) dt := \lim_{n \rightarrow \infty} \int_0^1 y_n(t) dt \in \mathbb{B}.$$

It turns out that the value of $\int_0^1 y(t) dt$ just defined is independent of the approximating simple functions $(y_n)_n$ and that it satisfies the fundamental inequality

$$\left\| \int_0^1 y(t) dt \right\|_{\mathbb{B}} \leq \int_0^1 \|y(t)\|_{\mathbb{B}} dt. \quad (2.1)$$

An alternative characterisation of the \mathbb{B} -valued integrable maps is given by the following theorem, which is due to Bochner: a strongly measurable function $y : [0, 1] \rightarrow \mathbb{B}$ is Bochner integrable if and only if it satisfies $\int_0^1 \|y(t)\|_{\mathbb{B}} dt < +\infty$.

The previous result naturally leads to the notion of \mathbb{B} -valued L^p space: given $p \in [1, \infty]$, we define $L^p([0, 1], \mathbb{B})$ as the space of all (equivalence classes of) those strongly measurable maps $y : [0, 1] \rightarrow \mathbb{B}$ for which the quantity $\|y\|_{L^p([0, 1], \mathbb{B})}$ is finite, where

$$\|y\|_{L^p([0, 1], \mathbb{B})} := \begin{cases} \left(\int_0^1 \|y(t)\|_{\mathbb{B}}^p dt \right)^{1/p} & \text{if } p < \infty, \\ \text{ess sup}_{t \in [0, 1]} \|y(t)\|_{\mathbb{B}} & \text{if } p = \infty. \end{cases}$$

Hence $L^p([0, 1], \mathbb{B})$ itself is a Banach space, for any $p \in [1, \infty]$.

Definition 2.1 (Vector-valued Sobolev/absolutely continuous maps) Let $p \in [1, \infty]$. The space $W^{1,p}([0, 1], \mathbb{B})$ consists of those curves $y \in L^p([0, 1], \mathbb{B})$ for which there is $y' \in L^p([0, 1], \mathbb{B})$ such that

$$\int_0^1 \varphi'(t)y(t) dt = - \int_0^1 \varphi(t)y'(t) dt \quad \forall \varphi \in C_c^\infty(0, 1).$$

It is equipped with the norm

$$\|y\|_{W^{1,p}([0,1],\mathbb{B})} := \left(\|y\|_{L^p([0,1],\mathbb{B})}^p + \|y'\|_{L^p([0,1],\mathbb{B})}^p \right)^{1/p}.$$

The space $AC^p([0, 1], \mathbb{B})$ consists of those curves $y : [0, 1] \rightarrow \mathbb{B}$ for which there is $f \in L^p(0, 1)$ such that

$$\|y(s) - y(t)\|_{\mathbb{B}} \leq \int_t^s f(r) dr, \quad \forall t, s \in [0, 1], t \leq s.$$

Proposition 2.2 (Absolutely continuous representative) Let $y \in W^{1,p}([0, 1], \mathbb{B})$. Then there is $\tilde{y} \in AC^p([0, 1], \mathbb{B})$ such that $y(t) = \tilde{y}(t)$ for a.e. t . Moreover, such \tilde{y} satisfies

$$\tilde{y}(s) - \tilde{y}(t) = \int_t^s y'(r) dr \quad \forall t, s \in [0, 1], t < s.$$

Proof. The curve $t \mapsto z(t) := \int_0^t y'(s) ds$ belongs to $AC^p \cap W^{1,p}([0, 1], \mathbb{B})$ and thus in particular the curve $t \mapsto y(t) - z(t)$ belongs to $W^{1,p}([0, 1], \mathbb{B})$ and, obviously, has derivative a.e. equal to 0. Hence to conclude it is sufficient to show that any such curve is a.e. constant. This follows noticing that for any $\ell \in \mathbb{B}'$ the map $t \mapsto \ell(y(t) - z(t))$ is in $W^{1,p}([0, 1])$ (by direct verification) and has derivative a.e. equal to 0. \square

Proposition 2.3 (Characterization of curves in $W^{1,p}([0, 1], \mathbb{B})$) Let $y, z \in L^p([0, 1], \mathbb{B})$. Then $y \in W^{1,p}([0, 1], \mathbb{B})$ and $z = y'$ if and only if for some dense set $D \subset \mathbb{B}'$ we have that $\ell \circ y \in W^{1,1}([0, 1])$ with $(\ell \circ y)' = \ell \circ z$ a.e. for every $\ell \in D$.

In particular, if $\mathbb{B} = L^{\tilde{p}}(\mu)$ for some Radon measure μ , then $y \in W^{1,p}([0, 1], L^{\tilde{p}}(\mu))$ and $z = y'$ if and only if for every Borel set E we have that $t \mapsto \int_E y(t) d\mu$ is in $W^{1,1}(0, 1)$ with derivative given by $t \mapsto \int_E z(t) d\mu$.

Proof. By assumption and using the fact that the Bochner integral commutes with the application of ℓ we have that

$$\ell \left(\int_0^1 \varphi'(t)y(t) dt \right) = \ell \left(- \int_0^1 \varphi(t)z(t) dt \right) \quad \forall \varphi \in C_c^\infty(0, 1),$$

for every $\ell \in D$. The conclusion follows by the density of D in \mathbb{B}' .

For the second claim just observe that the linear span of the set of characteristic functions of Borel sets is dense in $L^{\tilde{q}}(\mu) \sim (L^{\tilde{p}}(\mu))'$. \square

It is important to underline that in general absolute continuity does not imply a.e. differentiability: this has to do with the so-called Radon-Nikodym property of the target Banach space. A sufficient condition for this implication to hold is given in the next theorem:

Theorem 2.4 *Let \mathbb{B} be a reflexive Banach space, $p \in [1, \infty]$ and $y \in AC^p([0, 1], \mathbb{B})$. Then for a.e. $t \in [0, 1]$ the limit of $\frac{y(t+h)-y(t)}{h}$ as $h \rightarrow 0$ exists in \mathbb{B} .*

In particular, $AC^p([0, 1], \mathbb{B}) \sim W^{1,p}([0, 1], \mathbb{B})$, i.e. every absolutely continuous curve is the (only) continuous representative of a curve in $W^{1,p}([0, 1], \mathbb{B})$.

Given any Banach space \mathbb{B} , we shall denote by $\text{End}(\mathbb{B})$ the space of all linear and continuous maps of \mathbb{B} to itself, which is a Banach space if endowed with the operator norm.

The space $\Gamma(\mathbb{B}) := C([0, 1], \mathbb{B})$ is a Banach space with respect to the norm $\|\cdot\|_{\Gamma(\mathbb{B})}$, given by $\|y\|_{\Gamma(\mathbb{B})} := \max \{\|y_t\|_{\mathbb{B}} : t \in [0, 1]\}$ for every $y \in \Gamma(\mathbb{B})$.

Theorem 2.5 (Integral solutions to vector-valued linear ODEs) *Let \mathbb{B} be a Banach space. Let $z \in \Gamma(\mathbb{B})$. Let $\lambda : [0, 1] \rightarrow \text{End}(\mathbb{B})$ be a bounded function, i.e. there exists $c > 0$ such that $\|\lambda(t)\|_{\text{End}(\mathbb{B})} \leq c$ for every $t \in [0, 1]$. Assume that $[0, 1] \ni t \mapsto \lambda(t)v \in \mathbb{B}$ is strongly measurable for every $v \in \mathbb{B}$. Then there exists a unique curve $y \in \Gamma(\mathbb{B})$ such that*

$$y(t) = z(t) + \int_0^t \lambda(s)y(s) ds \quad \text{for every } t \in [0, 1]. \quad (2.2)$$

Moreover, the solution y satisfies $\|y\|_{\Gamma(\mathbb{B})} \leq e^c \|z\|_{\Gamma(\mathbb{B})}$.

Proof. Given any simple function $t \mapsto y_t = \sum_{i=1}^k \chi_{A_i}(t) v_i$, with $A_1, \dots, A_k \in \mathcal{B}([0, 1])$ and $v_1, \dots, v_k \in \mathbb{B}$, we have that $t \mapsto \lambda(t)y_t = \sum_{i=1}^k \chi_{A_i}(t) \lambda(t)v_i$ is strongly measurable by hypothesis on λ . Now fix $y \in \Gamma(\mathbb{B})$. In particular, $y : [0, 1] \rightarrow \mathbb{B}$ is strongly measurable, hence there exists a sequence $(y^k)_k$ of simple functions $y^k : [0, 1] \rightarrow \mathbb{B}$ such that $\lim_k \|y_t^k - y_t\|_{\mathbb{B}} = 0$ holds for \mathcal{L}^1 -a.e. $t \in [0, 1]$. This grants that $\|\lambda(t)y_t^k - \lambda(t)y_t\|_{\mathbb{B}} \leq c \|y_t^k - y_t\|_{\mathbb{B}} \xrightarrow{k} 0$ is satisfied for \mathcal{L}^1 -a.e. $t \in [0, 1]$, thus accordingly the map $t \mapsto \lambda(t)y_t$ is strongly measurable as pointwise limit of strongly measurable functions. Moreover, since $\|\lambda(t)y_t\|_{\mathbb{B}} \leq c \|y\|_{\Gamma(\mathbb{B})}$ for all $t \in [0, 1]$, one has that $t \mapsto \lambda(t)y_t$ actually belongs to $L^\infty([0, 1], \mathbb{B})$. Therefore it makes sense to define the function $\Lambda y : [0, 1] \rightarrow \mathbb{B}$ as $(\Lambda y)(t) := \int_0^t \lambda(s)y_s ds$ for every $t \in [0, 1]$. Note that

$$\|\Lambda y(t_1) - \Lambda y(t_0)\|_{\mathbb{B}} \leq c \|y\|_{\Gamma(\mathbb{B})} (t_1 - t_0) \quad \text{for every } t_0, t_1 \in [0, 1] \text{ with } t_0 < t_1. \quad (2.3)$$

Then Λy is Lipschitz with $\text{Lip}(\Lambda y) \leq c \|y\|_{\Gamma(\mathbb{B})}$, so in particular $\Lambda y \in \Gamma(\mathbb{B})$. By plugging $t_1 = t$ and $t_0 = 0$ into (2.3), we deduce that $\|\Lambda y(t)\|_{\mathbb{B}} \leq c \|y\|_{\Gamma(\mathbb{B})} t$ for all $t \in [0, 1]$ and accordingly that $\|\Lambda y\|_{\Gamma(\mathbb{B})} \leq c \|y\|_{\Gamma(\mathbb{B})}$. This guarantees that the mapping $\Lambda : \Gamma(\mathbb{B}) \rightarrow \Gamma(\mathbb{B})$ is linear and continuous, with $\|\Lambda\|_{\text{End}(\Gamma(\mathbb{B}))} \leq c$. Now observe that

$$y \in \Gamma(\mathbb{B}) \text{ satisfies (2.2)} \iff (\text{Id}_{\Gamma(\mathbb{B})} - \Lambda)(y) = z. \quad (2.4)$$

For any $n \in \mathbb{N}^+$, the iterated operator $\Lambda^n = \Lambda \circ \dots \circ \Lambda$ satisfies

$$\begin{aligned}
\|\Lambda^n y(t)\|_{\mathbb{B}} &\leq c \int_0^t \|\Lambda^{n-1} y(t_n)\|_{\mathbb{B}} dt_n \\
&\leq c^2 \int_0^t \int_0^{t_n} \|\Lambda^{n-2} y(t_{n-1})\|_{\mathbb{B}} dt_{n-1} dt_n \\
&\leq \dots \dots \\
&\leq c^n \int_0^t \int_0^{t_n} \dots \int_0^{t_2} \|y(t_1)\|_{\mathbb{B}} dt_1 \dots dt_{n-1} dt_n \\
&\leq c^n \|y\|_{\Gamma(\mathbb{B})} \int_0^t \int_0^{t_n} \dots \int_0^{t_2} dt_1 \dots dt_{n-1} dt_n \\
&= c^n \|y\|_{\Gamma(\mathbb{B})} \frac{t^n}{n!}
\end{aligned}$$

for every $y \in \Gamma(\mathbb{B})$ and $t \in [0, 1]$, whence $\|\Lambda^n\|_{\text{End}(\Gamma(\mathbb{B}))} \leq c^n/n!$. Hence $\text{Id}_{\Gamma(\mathbb{B})} - \Lambda$ is invertible and the operator norm of its inverse $(\text{Id}_{\Gamma(\mathbb{B})} - \Lambda)^{-1} = \sum_{n=0}^{\infty} \Lambda^n$ is bounded above by e^c . In light of (2.4), we finally conclude that there exists a unique curve $y \in \Gamma(\mathbb{B})$ fulfilling (2.2), namely $y := (\text{Id}_{\Gamma(\mathbb{B})} - \Lambda)^{-1}(z)$, which also satisfies $\|y\|_{\Gamma(\mathbb{B})} \leq e^c \|z\|_{\Gamma(\mathbb{B})}$. \square

We will actually make use of is the following consequence of Theorem 2.5.

Corollary 2.6 (Differential solutions to vector-valued linear ODEs) *Fix a reflexive Banach space \mathbb{B} . Let $\bar{y} \in \mathbb{B}$. Let $\lambda : [0, 1] \rightarrow \text{End}(\mathbb{B})$ be a bounded function. Suppose that the map $[0, 1] \ni t \mapsto \lambda(t)v \in \mathbb{B}$ is strongly measurable for every $v \in \mathbb{B}$. Then there exists a unique curve $y \in \text{LIP}([0, 1], \mathbb{B})$ such that*

$$\begin{cases} y'(t) = \lambda(t)y(t) & \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1], \\ y(0) = \bar{y}. \end{cases} \quad (2.5)$$

Moreover, the solution y satisfies $\|y\|_{\Gamma(\mathbb{B})} \leq e^c \|\bar{y}\|_{\mathbb{B}}$, where $c := \max_{t \in [0, 1]} \|\lambda(t)\|_{\text{End}(\mathbb{B})}$.

Proof. Define $z(t) := \bar{y}$ for every $t \in [0, 1]$ and consider the curve $y \in \Gamma(\mathbb{B})$, whose existence is granted by Theorem 2.5. For every $t, s \in [0, 1]$, with $s < t$, we have that

$$\|y(t) - y(s)\|_{\mathbb{B}} = \left\| \int_s^t \lambda(r)y(r) dr \right\|_{\mathbb{B}} \stackrel{(2.1)}{\leq} \int_s^t \|\lambda(r)y(r)\|_{\mathbb{B}} dr \leq c \int_s^t \|y(r)\|_{\mathbb{B}} dr.$$

Since $\|y(\cdot)\|_{\mathbb{B}} \in L^\infty(0, 1)$, we deduce that the function y is Lipschitz, so that by Theorem 2.4 y is a.e. differentiable. Then (2.5) trivially follows from (2.2).

Conversely, let $y \in \text{LIP}([0, 1], \mathbb{B})$ be any curve such that (2.5) holds true. By integration we conclude that y satisfies also property (2.2), proving uniqueness. \square

2.2 Pullback of an L^0 -normed module

The aim of this subsection is to introduce the concept of pullback of an L^0 -normed module and to study its main properties. The following definitions and results mimic the analogous ones for L^p -normed modules, which are treated in [8, Subsection 1.6]; a digression similar to the one below has been done in [15].

Theorem 2.7 Let $(X, \mathcal{A}_X, \mathfrak{m}_X)$, $(Y, \mathcal{A}_Y, \mathfrak{m}_Y)$ be σ -finite measured spaces. Let $\varphi : X \rightarrow Y$ be a map of bounded compression (i.e. $\varphi_* \mathfrak{m}_X \leq C \mathfrak{m}_Y$ for some $C > 0$). Let \mathcal{M}^0 be an $L^0(\mathfrak{m}_Y)$ -normed module. Then there exists (up to unique isomorphism) a unique couple $(\mathcal{N}^0, \mathbb{T})$, where \mathcal{N}^0 is an $L^0(\mathfrak{m}_X)$ -normed module and $\mathbb{T} : \mathcal{M}^0 \rightarrow \mathcal{N}^0$ is a linear map, such that

- (i) $|\mathbb{T}v| = |v| \circ \varphi$ holds \mathfrak{m}_X -a.e. in X , for every $v \in \mathcal{M}^0$,
- (ii) the set of all the elements of the form $\sum_{i=1}^n \chi_{A_i} \mathbb{T}v_i$, with $(A_i)_{i=1}^n \subseteq \mathcal{A}_X$ partition of X and $v_1, \dots, v_n \in \mathcal{M}^0$, is dense in \mathcal{N}^0 .

Namely, if two couples $(\mathcal{N}_1^0, \mathbb{T}_1)$ and $(\mathcal{N}_2^0, \mathbb{T}_2)$ as above fulfill both (i) and (ii), then there exists a unique module isomorphism $\Phi : \mathcal{N}_1^0 \rightarrow \mathcal{N}_2^0$ such that the diagram

$$\begin{array}{ccc} \mathcal{M}^0 & \xrightarrow{\mathbb{T}_1} & \mathcal{N}_1^0 \\ & \searrow \mathbb{T}_2 & \downarrow \Phi \\ & & \mathcal{N}_2^0 \end{array}$$

is commutative.

Proof. Existence. Fix $p \in [1, \infty)$ and let $\mathcal{M} := \{v \in \mathcal{M}^0 : |v| \in L^p(\mathfrak{m}_Y)\}$. Hence \mathcal{M} is an $L^p(\mathfrak{m}_Y)$ -normed module and \mathcal{M}^0 is the L^0 -completion of \mathcal{M} . Define $\mathcal{N}^0 := (\varphi^* \mathcal{M})^0$. We construct the map \mathbb{T} in the following way: since the space \mathcal{M} is dense in \mathcal{M}^0 and $\varphi^* \mathcal{M}$ is continuously embedded into \mathcal{N}^0 , the map $\varphi^* : \mathcal{M} \rightarrow \varphi^* \mathcal{M}$ can be uniquely extended to a linear continuous function $\mathbb{T} : \mathcal{M}^0 \rightarrow \mathcal{N}^0$. Such function \mathbb{T} also satisfies (i). Moreover, it is clear that the diagram

$$\begin{array}{ccc} \mathcal{M} & \hookrightarrow & \mathcal{M}^0 \\ \varphi^* \downarrow & & \downarrow \mathbb{T} \\ \varphi^* \mathcal{M} & \hookrightarrow & \mathcal{N}^0 \end{array}$$

commutes. Therefore, since the set $\{\varphi^* v : v \in \mathcal{M}\}$ generates \mathcal{N}^0 as $L^0(\mathfrak{m}_X)$ -normed module, we have in particular that the set $\{\mathbb{T}v : v \in \mathcal{M}^0\}$ generates \mathcal{N}^0 as $L^0(\mathfrak{m}_X)$ -normed module, proving (ii).

Uniqueness. Let us choose $(\mathcal{N}_1^0, \mathbb{T}_1)$, $(\mathcal{N}_2^0, \mathbb{T}_2)$ satisfying (i) and (ii). For $j = 1, 2$, denote by V_j the set of $\sum_{i=1}^n \chi_{A_i} \mathbb{T}_j v_i$ as in (ii). Then the unique $L^0(\mathfrak{m}_X)$ -linear map $\Psi : V_1 \rightarrow V_2$, which satisfies $\Psi \circ \mathbb{T}_1 = \mathbb{T}_2$, is necessarily given by $\Psi(\sum_{i=1}^n \chi_{A_i} \mathbb{T}_1 v_i) = \sum_{i=1}^n \chi_{A_i} \mathbb{T}_2 v_i$. By requiring the condition (i), we force the \mathfrak{m}_X -a.e. equality

$$\left| \sum_{i=1}^n \chi_{A_i} \mathbb{T}_2 v_i \right| = \sum_{i=1}^n \chi_{A_i} |v_i| \circ \varphi = \left| \sum_{i=1}^n \chi_{A_i} \mathbb{T}_1 v_i \right|,$$

which shows that the map Ψ is actually well-defined and continuous. There exists a unique module morphism $\Phi : \mathcal{N}_1^0 \rightarrow \mathcal{N}_2^0$ that extends Ψ , by density of V_1 in \mathcal{N}_1^0 . Such map Φ clearly satisfies $\Phi \circ \mathbb{T}_1 = \mathbb{T}_2$. Finally, by interchanging the roles of \mathcal{N}_1^0 and \mathcal{N}_2^0 , one can easily conclude that Φ is an isomorphism, getting the thesis. \square

Definition 2.8 (Pullback module) Any couple $(\mathcal{N}^0, \mathbb{T})$ that satisfies Theorem 2.7 will be unambiguously denoted by $(\varphi^* \mathcal{M}^0, \varphi^*)$. Moreover, we shall call $\varphi^* \mathcal{M}^0$ the pullback module of \mathcal{M}^0 and φ^* the pullback map.

Proposition 2.9 (Universal property of the pullback) *Let $(X, \mathcal{A}_X, \mathbf{m}_X)$, $(Y, \mathcal{A}_Y, \mathbf{m}_Y)$ be σ -finite measured spaces. Let $\varphi : X \rightarrow Y$ be a map of bounded compression. Let \mathcal{M}^0 be an $L^0(\mathbf{m}_Y)$ -normed module and let \mathcal{N}^0 be an $L^0(\mathbf{m}_X)$ -normed module. Consider a linear operator $\mathbb{T} : \mathcal{M}^0 \rightarrow \mathcal{N}^0$ such that*

$$|\mathbb{T}v| \leq \ell |v| \circ \varphi \quad \mathbf{m}_X\text{-a.e. in } X, \quad \text{for every } v \in \mathcal{M}^0, \quad (2.6)$$

for a suitable map $\ell \in L^0(\mathbf{m}_X)$. Then there exists a unique $L^0(\mathbf{m}_X)$ -linear and continuous operator $\widehat{\mathbb{T}} : \varphi^* \mathcal{M}^0 \rightarrow \mathcal{N}^0$ such that $\widehat{\mathbb{T}} \circ \varphi^* = \mathbb{T}$ and

$$|\widehat{\mathbb{T}}w| \leq \ell |w| \quad \mathbf{m}_X\text{-a.e. in } X, \quad \text{for every } w \in \varphi^* \mathcal{M}^0. \quad (2.7)$$

Proof. Denote by V the set of all elements of the form $\sum_{i=1}^n \chi_{A_i} \varphi^* v_i$, with $A_1, \dots, A_n \in \mathcal{A}_X$ partition of X and $v_1, \dots, v_n \in \mathcal{M}^0$, so that V is a dense linear subspace of $\varphi^* \mathcal{M}^0$ by property (ii) of Theorem 2.7. Any $L^0(\mathbf{m}_X)$ -linear map $\widehat{\mathbb{T}} : \varphi^* \mathcal{M}^0 \rightarrow \mathcal{N}^0$ with $\widehat{\mathbb{T}} \circ \varphi^* = \mathbb{T}$ must satisfy

$$\widehat{\mathbb{T}}w = \sum_{i=1}^n \chi_{A_i} \widehat{\mathbb{T}}(\varphi^* v_i) = \sum_{i=1}^n \chi_{A_i} \mathbb{T}v_i \quad \text{for } w = \sum_{i=1}^n \chi_{A_i} \varphi^* v_i \in V. \quad (2.8)$$

Consider $\widehat{\mathbb{T}} : V \rightarrow \mathcal{N}^0$ defined as in (2.8), then (2.6) grants that

$$|\widehat{\mathbb{T}}w| = \sum_{i=1}^n \chi_{A_i} |\mathbb{T}v_i| \leq \ell \sum_{i=1}^n \chi_{A_i} |v_i| \circ \varphi = \ell \sum_{i=1}^n \chi_{A_i} |\varphi^* v_i| = \ell |w| \quad \text{holds } \mathbf{m}_X\text{-a.e.}, \quad (2.9)$$

which shows that $\widehat{\mathbb{T}} : V \rightarrow \mathcal{N}^0$ is well-defined (in the sense that $\widehat{\mathbb{T}}w$ depends only on w and not on the way of representing it) and continuous. Therefore $\widehat{\mathbb{T}}$ can be uniquely extended to a linear continuous operator $\widetilde{\mathbb{T}} : \varphi^* \mathcal{M}^0 \rightarrow \mathcal{N}^0$. We readily deduce from the definition (2.8) that the equality $f \widehat{\mathbb{T}}w = \widehat{\mathbb{T}}(fw)$ holds for $f : X \rightarrow \mathbb{R}$ simple function, so that $\widehat{\mathbb{T}}$ can be shown to be $L^0(\mathbf{m}_X)$ -linear by an approximation argument. Finally, it follows from (2.9) that the inequality (2.7) is satisfied for $w \in V$, whence (2.7) holds by density of V in $\varphi^* \mathcal{M}^0$. \square

2.3 Some properties of test plans

For the sake of brevity, hereafter we shall use the notation \mathcal{L}_1 to indicate the 1-dimensional Lebesgue measure restricted to $[0, 1]$, namely

$$\mathcal{L}_1 := \mathcal{L}^1|_{[0,1]}.$$

Let $\boldsymbol{\pi} \in \mathcal{P}(\Gamma(X))$ be any fixed test plan on X , whence $(\Gamma(X), d_{\Gamma(X)}, \boldsymbol{\pi})$ is a metric measure space. Given that the map e_t is of bounded compression, it makes sense to consider the pullback module $e_t^* L^2(TX)$. Observe that $e_t^* L^2(TX)$ is a Hilbert module as soon as (X, d, \mathbf{m}) is infinitesimally Hilbertian.

Remark 2.10 Let us define the map $e : \Gamma(X) \times [0, 1] \rightarrow X$ as $e(\gamma, t) := \gamma_t$ for every $\gamma \in \Gamma(X)$ and $t \in [0, 1]$. It can be easily proved that the map e is continuous. This grants that, given any Borel map $f : X \rightarrow \mathbb{R}$, the function $f \circ e$ is Borel. Moreover, observe that

$$(\boldsymbol{\pi} \times \mathcal{L}_1)(e^{-1}(A)) = \int_0^1 \boldsymbol{\pi}(e_t^{-1}(A)) dt \leq C(\boldsymbol{\pi}) \mathbf{m}(A) \quad \text{for every } A \in \mathcal{B}(X)$$

by Fubini theorem, in other words it holds that $e_*(\boldsymbol{\pi} \times \mathcal{L}_1) \leq C(\boldsymbol{\pi}) \mathbf{m}$. Therefore one has that the composition $f \circ e \in L^0(\boldsymbol{\pi} \times \mathcal{L}_1)$ is well-defined for any $f \in L^0(\mathbf{m})$. \blacksquare

Theorem 2.11 *Let (X, d, \mathbf{m}) be a metric measure space. Let π be a test plan on X . Then*

$$\text{for every } f \in L^1(\mathbf{m}) \text{ the map } [0, 1] \ni t \mapsto f \circ e_t \in L^1(\pi) \text{ is continuous.} \quad (2.10)$$

In particular, the map $[0, 1] \ni t \mapsto \int f \circ e_t d\pi$ is continuous for every $f \in L^1(\mathbf{m})$.

Proof. First of all, we claim that

$$\lim_{s \rightarrow t} \int |f \circ e_s - f \circ e_t| d\pi = 0 \quad \text{if } f \in C_b(X) \cap L^1(\mathbf{m}) \text{ and } t \in [0, 1]. \quad (2.11)$$

To prove it, note that $|f \circ e_s - f \circ e_t|(\gamma) \leq 2\|f\|_{L^\infty(\mathbf{m})}$ for every $\gamma \in \Gamma(X)$ and $t, s \in [0, 1]$. Moreover, $|f(\gamma_s) - f(\gamma_t)| \rightarrow 0$ as $s \rightarrow t$ by continuity, for every $\gamma \in \Gamma(X)$ and $t \in [0, 1]$. Hence we obtain (2.11) as a consequence of the dominated convergence theorem. Observe also that

$$L^1(\mathbf{m}) \ni f \mapsto f \circ e_t \in L^1(\pi) \text{ is a linear bounded map, for every } t \in [0, 1]. \quad (2.12)$$

Indeed, $\int |f \circ e_t| d\pi \leq C(\pi) \int |f| d\mathbf{m}$ is satisfied for every $f \in L^1(\mathbf{m})$. Now fix $f \in L^1(\mathbf{m})$. Choose a sequence $(f_n)_n \subseteq C_b(X) \cap L^1(\mathbf{m})$ that converges to f with respect to the L^1 -norm. Given $t \in [0, 1]$ and $n \in \mathbb{N}$, we have that (2.11) and (2.12) yield

$$\begin{aligned} \overline{\lim}_{s \rightarrow t} \int |f \circ e_s - f \circ e_t| d\pi &\leq 2C(\pi) \|f - f_n\|_{L^1(\mathbf{m})} + \overline{\lim}_{s \rightarrow t} \int |f_n \circ e_s - f_n \circ e_t| d\pi \\ &= 2C(\pi) \|f - f_n\|_{L^1(\mathbf{m})}. \end{aligned} \quad (2.13)$$

By letting $n \rightarrow \infty$ in (2.13), we finally conclude that $\int |f \circ e_s - f \circ e_t| d\pi \rightarrow 0$ as $s \rightarrow t$, which proves (2.10). The last statement is obvious. \square

Under further assumptions on (X, d, \mathbf{m}) , we have at disposal also a notion of ‘speed’ π'_t of the test plan π at time t , as described in the following result. For the proof of such fact, we refer to [8, Theorem 2.3.18] or [6, Theorem/Definition 1.32].

Theorem 2.12 (Speed of a test plan) *Let (X, d, \mathbf{m}) be a metric measure space such that $L^2(TX)$ is separable. Let π be a test plan on X . Then there exists a unique (up to \mathcal{L}_1 -a.e. equality) family $\pi'_t \in e_t^* L^2(TX)$ such that*

$$\exists L^1(\pi)\text{-}\lim_{h \rightarrow 0} \frac{f \circ e_{t+h} - f \circ e_t}{h} = (e_t^* df)(\pi'_t) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1], \quad (2.14)$$

for every $f \in W^{1,2}(X)$. Moreover, the function $(\gamma, t) \mapsto |\pi'_t|(\gamma)$ is (the equivalence class of) a Borel map such that for \mathcal{L}^1 -a.e. $t \in [0, 1]$ it holds that

$$|\pi'_t|(\gamma) = |\dot{\gamma}_t| \quad \text{for } \pi\text{-a.e. } \gamma \in AC([0, 1], X). \quad (2.15)$$

Proposition 2.13 *Let (X, d, \mathbf{m}) be a metric measure space such that $L^2(TX)$ is separable, let π be a test plan on X and $f \in W^{1,2}(X)$. Then the a.e. defined map $[0, 1] \ni t \mapsto e_t^* df(\pi'_t) \in L^1(\pi)$ is a.e. equal to a Borel map.*

Proof. For every $h \in (0, 1)$ the map $[0, 1-h] \ni t \mapsto (f \circ e_{t+h} - f \circ e_t)/h \in L^1(\pi)$ is continuous. Thus by classical arguments the set of t 's for which the limit as $h \rightarrow 0$ exists is Borel and the limit function, set, say, to 0 when the limit does not exist, is Borel. \square

In this paper we shall work only on RCD spaces, which in particular are so that $W^{1,2}(X)$ is reflexive. In turn this implies, by the arguments in [1], that the tangent module is separable, so that the assumptions of Theorem 2.12 and Proposition 2.13 above are fulfilled.

In the sequel, we will mainly focus our attention on those test plans π that are concentrated on an equiLipschitz family of curves. As illustrated in the next definition, we will refer to them as ‘Lipschitz test plans’.

Definition 2.14 (Lipschitz test plan) *Let (X, d, \mathbf{m}) be a metric measure space such that $L^2(TX)$ is separable. Then a test plan π on X is said to be a Lipschitz test plan provided there exists a constant $L \geq 0$ such that*

$$|\pi'_t| \leq L \text{ holds } \pi\text{-a.e. in } \Gamma(X), \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1], \quad (2.16)$$

or, equivalently, such that π is concentrated on the family of all the L -Lipschitz curves in X . The smallest constant $L \geq 0$ for which (2.16) is satisfied will be denoted by $L(\pi)$.

Whenever the test plan π is Lipschitz, we have $(e_t^* df)(\pi'_t) \in L^2(\pi)$ for every $f \in W^{1,2}(X)$. One is then led to wonder whether in this case the limit in (2.14) takes place not only in $L^1(\pi)$, but also in $L^2(\pi)$. The answer is affirmative, as shown in the following simple result:

Proposition 2.15 *Let (X, d, \mathbf{m}) be a metric measure space such that $L^2(TX)$ is separable. Let π be a Lipschitz test plan on X . Let $f \in W^{1,2}(X)$. Then the mapping $t \mapsto f \circ e_t \in L^2(\pi)$ is Lipschitz and*

$$L^2(\pi)\text{-}\frac{d}{dt}(f \circ e_t) = (e_t^* df)(\pi'_t) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \quad (2.17)$$

Proof. Given any $t, s \in [0, 1]$ with $s < t$, one has that

$$\begin{aligned} \|f \circ e_t - f \circ e_s\|_{L^2(\pi)}^2 &= \int |f(\gamma_t) - f(\gamma_s)|^2 d\pi(\gamma) \\ \text{(by definition of Sobolev functions)} \quad &\leq \int \left(\int_s^t |Df|(\gamma_r) |\dot{\gamma}_r| dr \right)^2 d\pi(\gamma) \\ \text{(by Hölder inequality)} \quad &\leq (t-s) L(\pi)^2 \iint_s^t |Df|^2(\gamma_r) dr d\pi(\gamma) \\ &\leq C(\pi) L(\pi)^2 \|f\|_{W^{1,2}(X)}^2 (t-s)^2, \end{aligned}$$

which shows that $t \mapsto f \circ e_t \in L^2(\pi)$ is a Lipschitz map. In particular, it is differentiable at almost every $t \in [0, 1]$ by Theorem 2.4, so that (2.17) follows from (2.14). \square

3 Introduction of appropriate functional spaces

Throughout all this chapter, (X, d, \mathbf{m}) is a given $\text{RCD}(K, \infty)$ space and π a test plan on it.

Recall that the space $\text{TestF}(X)$ of *test functions on X* is defined as

$$\text{TestF}(X) := \left\{ f \in L^\infty \cap \text{LIP} \cap W^{1,2}(X) \cap D(\Delta) : \Delta f \in W^{1,2}(X) \right\}$$

and that the space of *test vector fields on X* is defined as

$$\text{TestV}(X) := \left\{ \sum_{i=1}^n f_i \nabla g_i : n \in \mathbb{N}^+, f_i, g_i \in \text{TestF}(X) \text{ for every } i \right\} \subset L^2(TX).$$

3.1 Test vector fields along π

We define the space of *vector fields along π* as:

$$\text{VF}(\boldsymbol{\pi}) := \prod_{t \in [0,1]} e_t^* L^2(TX).$$

Thus $\text{VF}(\boldsymbol{\pi})$ is the collection of maps assigning to each $t \in [0, 1]$ an element of $e_t^* L^2(TX)$; it is a vector space w.r.t. pointwise operation.

To each $V \in \text{VF}(\boldsymbol{\pi})$ we associate the function $\llbracket V \rrbracket : [0, 1] \rightarrow [0, +\infty)$, defined by

$$\llbracket V \rrbracket_t := \|V_t\|_{e_t^* L^2(TX)} \quad \text{for every } t \in [0, 1].$$

The subspace $\text{TestVF}(\boldsymbol{\pi}) \subset \text{VF}(\boldsymbol{\pi})$ of *test vector fields along π* is defined as:

$$\text{TestVF}(\boldsymbol{\pi}) := \left\{ t \mapsto \sum_{i=1}^n \varphi_i(t) \chi_{A_i} e_t^* v_i \mid \begin{array}{l} n \in \mathbb{N}^+, A_i \in \mathcal{B}(\Gamma(X)), \varphi_i \in \text{LIP}([0, 1]), \\ \text{and } v_i \in \text{TestV}(X) \text{ for every } i = 1, \dots, n \end{array} \right\}.$$

Since $\text{TestV}(X) \subseteq L^\infty(TX)$ we see that for any $V \in \text{TestVF}(\boldsymbol{\pi})$ the function $(\gamma, t) \mapsto |V_t|(\gamma)$ belongs to $L^\infty(\mathcal{L}_1 \times \boldsymbol{\pi})$.

Proposition 3.1 (Continuity of the test vector fields along π) *For any $V, W \in \text{TestVF}(\boldsymbol{\pi})$ we have that*

$$[0, 1] \ni t \rightarrow \langle V_t, W_t \rangle \in L^1(\boldsymbol{\pi}) \quad \text{is continuous.} \quad (3.1)$$

In particular, the function $\llbracket V \rrbracket : [0, 1] \rightarrow [0, +\infty)$ is continuous for every $V \in \text{TestVF}(\boldsymbol{\pi})$.

Proof. By linearity, it is clear that it is sufficient to prove the claim for V, W of the form $V = \chi_A e_t^* v$, $W = \chi_B e_t^* w$ for $v, w \in \text{TestV}(X)$. In this case the claim (3.1) is a direct consequence of

$$\langle V_t, W_t \rangle = \chi_{A \cap B} \langle v, w \rangle \circ e_t$$

and Theorem 2.11. The last statement follows by choosing $W = V$. \square

We now define two norms on $\text{TestVF}(\boldsymbol{\pi})$:

$$\begin{aligned} \|V\|_{\mathcal{L}^2(\boldsymbol{\pi})}^2 &:= \int_0^1 \llbracket V \rrbracket_t^2 dt \\ \|V\|_{\mathcal{C}(\boldsymbol{\pi})} &:= \max_{t \in [0,1]} \llbracket V \rrbracket_t. \end{aligned}$$

Notice that Proposition 3.1 ensures that $t \mapsto \llbracket V \rrbracket_t$ is Borel, hence $\|\cdot\|_{\mathcal{L}^2(\boldsymbol{\pi})}$ is well defined; also, routine computations show that $\|\cdot\|_{\mathcal{L}^2(\boldsymbol{\pi})}, \|\cdot\|_{\mathcal{C}(\boldsymbol{\pi})}$ are norms on $\text{TestVF}(\boldsymbol{\pi})$ with $\|\cdot\|_{\mathcal{L}^2(\boldsymbol{\pi})} \leq \|\cdot\|_{\mathcal{C}(\boldsymbol{\pi})}$.

We now want to show that $(\text{TestVF}(\boldsymbol{\pi}), \|\cdot\|_{\mathcal{C}(\boldsymbol{\pi})})$ is separable by exhibiting a countable dense subset. To this aim, we first choose three countable families $\mathcal{F}_1 \subseteq \{\text{open sets of } \Gamma(X)\}$, $\mathcal{F}_2 \subseteq \text{LIP}([0, 1])$ and $\mathcal{F}_3 \subseteq \text{TestV}(X)$ such that

- given $A \subseteq \Gamma(X)$ Borel and $\varepsilon > 0$, there exists $U \in \mathcal{F}_1$ with $\pi(A \Delta U) < \varepsilon$,
- \mathcal{F}_2 is dense in $C([0, 1])$ and stable by product and \mathbb{Q} -linear combinations,
- \mathcal{F}_3 is a \mathbb{Q} -vector space of functions in $W^{1,2}(X)$ whose gradients generate $L^2(TX)$.

We proceed in the following way:

\mathcal{F}_1 : Since $\Gamma(X)$ is separable, there exists a countable family $\tilde{\mathcal{F}}_1$ of open subsets of $\Gamma(X)$ that is a neighbourhood basis for each point $\gamma \in \Gamma(X)$. Let us denote by \mathcal{F}_1 the set of finite unions of elements of $\tilde{\mathcal{F}}_1$, so that \mathcal{F}_1 is countable. Fix $A \in \mathcal{B}(\Gamma(X))$ and $\varepsilon > 0$. The measure π is regular, since $(\Gamma(X), \mathbf{d}_{\Gamma(X)})$ is complete and separable. By inner regularity of π , there exists a compact subset $K \subseteq A$ such that $\pi(A \setminus K) < \varepsilon/2$. By outer regularity of π , there exists $V \subseteq \Gamma(X)$ open such that $K \subseteq V$ and $\pi(V \setminus K) < \varepsilon/2$. We can then associate to any $\gamma \in K$ a set $U_\gamma \in \tilde{\mathcal{F}}_1$ such that $\gamma \in U_\gamma \subseteq V$. By compactness of K , one has $K \subseteq U_{\gamma_1} \cup \dots \cup U_{\gamma_n} \subseteq V$ for some finite choice $\gamma_1, \dots, \gamma_n \in K$. Let us call $U := U_{\gamma_1} \cup \dots \cup U_{\gamma_n} \in \mathcal{F}_1$. We thus have that

$$\pi(A \Delta U) = \pi(A \setminus U) + \pi(U \setminus A) \leq \pi(A \setminus K) + \pi(V \setminus K) < \varepsilon.$$

\mathcal{F}_2 : By the separability of $C([0, 1])$ such \mathcal{F}_2 exists.

\mathcal{F}_3 : Since the space $(X, \mathbf{d}, \mathbf{m})$ is infinitesimally Hilbertian, we have that $W^{1,2}(X)$ is reflexive and therefore, by [1], separable. Thus let \mathcal{F}_3 be any countable dense \mathbb{Q} -vector subspace of $W^{1,2}(X)$ and notice that since gradients of functions in $W^{1,2}(X)$ generate $L^2(TX)$, the same holds for functions in \mathcal{F}_3 .

We now define the class of test vector fields $\text{TestVF}_{\mathbb{N}}(\pi)$ as:

$$\text{TestVF}_{\mathbb{N}}(\pi) := \left\{ t \mapsto \sum_{i=1}^n \psi_i(t) \chi_{U_i} e_t^* \nabla f_i \mid \begin{array}{l} n \in \mathbb{N}^+ \text{ and } U_i \in \mathcal{F}_1, \psi_i \in \mathcal{F}_2, \\ f_i \in \mathcal{F}_3 \text{ for every } i = 1, \dots, n \end{array} \right\}.$$

Clearly $\text{TestVF}_{\mathbb{N}}(\pi)$ is a countable subset of $\text{TestVF}(\pi)$. Also, notice that the inequalities

$$\begin{aligned} \|\chi_{Ae_t^*} v - \chi_{Ae_t^*} w\|_t &\leq \sqrt{\mathbf{C}(\pi)} \|v - w\|_{L^2(TX)}, \\ \|\chi_{Ae_t^*} v - \chi_{Ue_t^*} v\|_t &\leq \sqrt{\pi(A \Delta U)} \|v\|_{L^\infty(TX)}, \end{aligned}$$

valid for any $t \in [0, 1]$, $A, U \subset \Gamma(X)$ Borel and $v, w \in L^2(TX)$ and the very definition of pullback module, show that

$$\text{for any } t \in [0, 1] \text{ the set } \{W_t : W \in \text{TestVF}_{\mathbb{N}}(\pi)\} \text{ is dense in } e_t^* L^2(TX). \quad (3.2)$$

Lemma 3.2 (Separability of $\text{TestVF}(\pi)$) *The family $\text{TestVF}_{\mathbb{N}}(\pi)$ is dense in $\text{TestVF}(\pi)$ with respect to the norm $\|\cdot\|_{\mathcal{G}(\pi)}$ (and thus also w.r.t. $\|\cdot\|_{\mathcal{L}^2(\pi)}$).*

Proof. Let $V \in \text{TestVF}(\pi)$ be arbitrary, $\varepsilon > 0$ and $t_0 \in [0, 1]$. By (3.2) there is $W \in \text{TestVF}_{\mathbb{N}}(\pi)$ such that $\|V - W\|_{t_0} < \varepsilon$. Since $t \mapsto \|V - W\|_t^2 = \|V\|_t^2 + \|W\|_t^2 - 2 \int \langle V_t, W_t \rangle d\pi$ is continuous, we see that $\|V - W\|_t < \varepsilon$ for every t in a neighbourhood of t_0 . By compactness of $[0, 1]$ we can then find a finite number of open intervals I_1, \dots, I_n covering $[0, 1]$ and elements $W_1, \dots, W_n \in \text{TestVF}_{\mathbb{N}}(\pi)$ such that

$$\|V - W_i\|_t < \varepsilon \quad \forall t \in I_i \cap [0, 1], \quad i = 1, \dots, n. \quad (3.3)$$

By multiplying W_i by an appropriate function in \mathcal{F}_2 we can also assume that

$$\|W_i\|_t < \|V\|_{\mathcal{G}(\pi)} + 2\varepsilon \quad \forall t \in [0, 1]. \quad (3.4)$$

Now let (ϕ_i) be a Lipschitz partition of the unity subordinate to the cover made with the I_i 's and for any i let $\psi_i \in \mathcal{F}_2$ be such that $|\phi_i(t) - \psi_i(t)| < \varepsilon$ for every $t \in [0, 1]$. Then we have $W_t := \sum_i \psi_i(t)W_{i,t} \in \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ and

$$\llbracket V - W \rrbracket_t \leq \llbracket V - \sum_i \phi_i(t)W_i \rrbracket_t + \llbracket \sum_i (\psi_i(t) - \phi_i(t))W_i \rrbracket_t \stackrel{(3.3),(3.4)}{\leq} \varepsilon + \varepsilon(\|V\|_{\mathcal{E}(\boldsymbol{\pi})} + 2\varepsilon)$$

for any $t \in [0, 1]$. The conclusion follows by the arbitrariness of $\varepsilon > 0$. \square

3.2 The space $\mathcal{L}^2(\boldsymbol{\pi})$

We start defining the class of Borel vector fields along $\boldsymbol{\pi}$:

Definition 3.3 (Borel vector fields along $\boldsymbol{\pi}$) *We say that $V \in \text{VF}(\boldsymbol{\pi})$ is Borel provided*

$$[0, 1] \ni t \longmapsto \int \langle V_t, W_t \rangle d\boldsymbol{\pi} \text{ is a Borel function,} \quad (3.5)$$

for every $W \in \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$.

Notice that thanks to Lemma 3.2 this notion would be unaltered if we require (3.5) to hold for any $W \in \text{TestVF}(\boldsymbol{\pi})$. Also, Proposition 3.1 ensures that test vector fields are Borel. We have the following basic result:

Proposition 3.4 *Let $V \in \text{VF}(\boldsymbol{\pi})$ be Borel. Then the map $\llbracket V \rrbracket : [0, 1] \rightarrow [0, \infty)$ is Borel.*

Proof. From (3.2) we deduce that

$$\llbracket V \rrbracket_t^2 = \sup_{W \in \text{Test}\Gamma_{\mathbb{N}}(X)} \left(2 \int \langle V_t, W_t \rangle d\boldsymbol{\pi} - \llbracket W \rrbracket_t^2 \right) \quad \text{for every } t \in [0, 1].$$

and the thesis follows. \square

We can now define the space $\mathcal{L}^2(\boldsymbol{\pi})$:

Definition 3.5 (The space $\mathcal{L}^2(\boldsymbol{\pi})$) *The space $\mathcal{L}^2(\boldsymbol{\pi})$ is the space of all Borel vector fields $V \in \text{VF}(\boldsymbol{\pi})$ such that*

$$\|V\|_{\mathcal{L}^2(\boldsymbol{\pi})}^2 := \int_0^1 \llbracket V \rrbracket_t^2 dt = \int_0^1 \int |V_t|^2 d\boldsymbol{\pi} dt < +\infty,$$

where we identify $V, \tilde{V} \in \text{VF}(\boldsymbol{\pi})$ if $V_t = \tilde{V}_t$ for a.e. $t \in [0, 1]$.

Clearly $(\mathcal{L}^2(\boldsymbol{\pi}), \|\cdot\|_{\mathcal{L}^2(\boldsymbol{\pi})})$ is a normed space, wherein $\text{TestVF}(\boldsymbol{\pi})$ is embedded. Adapting the classical arguments concerning the standard L^2 spaces we have the following:

Proposition 3.6 (Basic properties of $\mathcal{L}^2(\boldsymbol{\pi})$) *The space $\mathcal{L}^2(\boldsymbol{\pi})$ is a Hilbert space and if $V_n \rightarrow V$ in $\mathcal{L}^2(\boldsymbol{\pi})$ then there is a subsequence such that $V_{n,t} \rightarrow V_t$ in $\mathfrak{e}_t^*L^2(TX)$ for a.e. $t \in [0, 1]$.*

Proof. It is clear that the $\mathcal{L}^2(\boldsymbol{\pi})$ norm comes from the scalar product

$$\langle V, W \rangle_{\mathcal{L}^2(\boldsymbol{\pi})} := \int_0^1 \int \langle V_t, W_t \rangle d\boldsymbol{\pi} dt.$$

To conclude the proof we shall show that if (V_n) is a sequence of Borel vector fields in $\mathcal{L}^2(\boldsymbol{\pi})$ such that $\sum_n \|V_{n+1} - V_n\|_{\mathcal{L}^2(\boldsymbol{\pi})} < \infty$, then such sequence has a limit $V \in \mathcal{L}^2(\boldsymbol{\pi})$ and for a.e. t it holds $V_{n,t} \rightarrow V_t$ in $e_t^* L^2(TX)$.

Define the Borel function $g : [0, 1] \rightarrow [0, +\infty]$ as $g := \sum_n \llbracket V_{n+1} - V_n \rrbracket$ and notice that since

$$\left\| \sum_{n=1}^N \llbracket V_{n+1} - V_n \rrbracket \right\|_{L^2(0,1)} \leq \sum_{n=1}^N \|V_{n+1} - V_n\|_{\mathcal{L}^2(\boldsymbol{\pi})} \leq \sum_{n=1}^{\infty} \|V_{n+1} - V_n\|_{\mathcal{L}^2(\boldsymbol{\pi})} < \infty \quad \forall N \in \mathbb{N},$$

we have that $g \in L^2(0, 1)$. Let $N := \{t : g(t) = +\infty\}$ and notice that for every $t \in [0, 1] \setminus N$ we have

$$\sum_{n=1}^{\infty} \|V_{n+1,t} - V_{n,t}\|_{e_t^* L^2(TX)} = \sum_{n=1}^{\infty} \llbracket V_{n+1} - V_n \rrbracket_t = g(t) < \infty, \quad (3.6)$$

proving that $(V_{n,t})$ is a Cauchy sequence in $e_t^* L^2(TX)$. Then define

$$V_t := \begin{cases} \lim_n V_{n,t} \in e_t^* L^2(TX) & \text{if } t \in [0, 1] \setminus N. \\ 0 \in e_t^* L^2(TX) & \text{if } t \in N. \end{cases}$$

Notice that for every $W \in \text{TestVF}(\boldsymbol{\pi})$ we have $\int \langle V_t, W_t \rangle d\boldsymbol{\pi} = \lim_n \int \langle V_{n,t}, W_t \rangle d\boldsymbol{\pi}$ for all $t \in [0, 1] \setminus N$, hence the map $[0, 1] \ni t \mapsto \int \langle V_t, W_t \rangle d\boldsymbol{\pi}$ is Borel and, by arbitrariness of W , this shows that V is Borel. Since trivially we have $\llbracket V \rrbracket_t \leq \llbracket V_1 \rrbracket_t + \sum_{n=1}^{\infty} \llbracket V_{n+1} - V_n \rrbracket_t$, by (3.6) we see that $V \in \mathcal{L}^2(\boldsymbol{\pi})$. Now to check that $V_n \rightarrow V$ in $\mathcal{L}^2(\boldsymbol{\pi})$ notice that, again by (3.6), the sequence $\llbracket V - V_n \rrbracket_t$ is dominated in $L^2(0, 1)$ and that for every $t \in [0, 1] \setminus N$ we have

$$\lim_{n \rightarrow \infty} \llbracket V - V_n \rrbracket_t \leq \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \sum_{i=n}^m \llbracket V_{i+1} - V_i \rrbracket_t \stackrel{(3.6)}{=} 0,$$

so that the conclusion follows by the dominated convergence theorem. \square

Proposition 3.7 (Density of $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ in $\mathcal{L}^2(\boldsymbol{\pi})$) *The space $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ is dense in $\mathcal{L}^2(\boldsymbol{\pi})$. In particular, $\mathcal{L}^2(\boldsymbol{\pi})$ is separable.*

Proof. Let (Z_k) be an enumeration of the elements in $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$, pick a Borel vector field $V \in \mathcal{L}^2(\boldsymbol{\pi})$ and choose $\varepsilon > 0$. Then for every $k \in \mathbb{N}$ let $\tilde{G}_k := \{t \in [0, 1] : \llbracket V - Z_k \rrbracket_t < \varepsilon\}$ and put $G_1 := \tilde{G}_1$ and $G_k := \tilde{G}_k \setminus (\tilde{G}_1 \cup \dots \cup \tilde{G}_{k-1})$ for $k > 1$. Then (3.2) grants that $(G_k)_{k \geq 1}$ is a Borel partition of $[0, 1]$.

Now for $m \in \mathbb{N} \cup \{\infty\}$ define $W_m \in \mathcal{L}^2(\boldsymbol{\pi})$ as $W_{m,t} := \sum_{k=1}^m \chi_{G_k}(t) Z_{k,t}$. Observe that $\|W_{\infty} - V\|_{\mathcal{L}^2(\boldsymbol{\pi})} < \varepsilon$ by definition of G_k . Moreover, for each $m \geq 1$ one has that

$$\|W_m - W_{\infty}\|_{\mathcal{L}^2(\boldsymbol{\pi})}^2 = \sum_{k=m+1}^{\infty} \int_{G_k} \llbracket Z_k \rrbracket_t^2 dt \leq \int_{\bigcup_{k>m} G_k} 2(\llbracket V \rrbracket_t^2 + \varepsilon^2) dt,$$

so that accordingly $\lim_{m \rightarrow \infty} \|W_m - W_{\infty}\|_{\mathcal{L}^2(\boldsymbol{\pi})} = 0$ by the dominated convergence theorem.

Hence to conclude it is sufficient to show that each W_m belongs to the $\mathcal{L}^2(\boldsymbol{\pi})$ -closure of $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ and in turn this will follow if we prove that for $Z \in \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ and $G \subset [0, 1]$ Borel the vector field $\chi_G Z$ belongs to the $\mathcal{L}^2(\boldsymbol{\pi})$ -closure of $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$. To see this, simply let $(\varphi_n) \subset \text{LIP}([0, 1])$ be uniformly bounded and a.e. converging to χ_G , notice that $\varphi_n Z \in \text{TestVF}(\boldsymbol{\pi})$ and that an application of the dominate convergence theorem shows that $\varphi_n Z \rightarrow \chi_G Z$ in $\mathcal{L}^2(\boldsymbol{\pi})$. \square

Now consider the speed $\boldsymbol{\pi}'_t$, associated to any test plan $\boldsymbol{\pi}$ by Theorem 2.12.

Proposition 3.8 *The (equivalence class up to a.e. equality of the) map $t \mapsto \boldsymbol{\pi}'_t$ is an element of the space $\mathcal{L}^2(\boldsymbol{\pi})$.*

Proof. We have $\boldsymbol{\pi}'_t \in e_t^* L^2(TX)$ for a.e. $t \in [0, 1]$ and

$$\int_0^1 \int |\boldsymbol{\pi}'_t|^2 d\boldsymbol{\pi} dt \stackrel{(2.15)}{=} \int_0^1 \int |\dot{\gamma}_t|^2 d\boldsymbol{\pi}(\gamma) dt < +\infty$$

by the very definition of test plan. Hence we need only to show that $t \mapsto \boldsymbol{\pi}'_t$ has a Borel representative in the sense of Definition 3.3.

Notice that for any $f \in W^{1,2}(X)$ by Proposition 2.13 we have that the map $t \mapsto (e_t^* df)(\boldsymbol{\pi}'_t)$ has a Borel representative. Hence the same holds for $t \mapsto \psi(t) \chi_U \langle e_t^* \nabla f, \boldsymbol{\pi}'_t \rangle$ for every $\psi \in \text{LIP}([0, 1])$ and $U \subset \Gamma(X)$ Borel. Therefore there exists a Borel negligible set $N \subset [0, 1]$ such that for every $V \in \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ the map $t \mapsto \int \langle V_t, \boldsymbol{\pi}'_t \rangle d\boldsymbol{\pi}$, set to 0 on N , is Borel. This is sufficient to conclude. \square

We conclude the section by pointing out that $\mathcal{L}^2(\boldsymbol{\pi})$ can also be seen as the pullback of $L^2(TX)$ via the evaluation map $e : \Gamma(X) \times [0, 1] \rightarrow X$ defined as $e(\gamma, t) := \gamma_t$. To this aim, let us start by defining the following operations:

(i) Given $f \in L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)$ and $V \in \mathcal{L}^2(\boldsymbol{\pi})$, we define $fV \in \mathcal{L}^2(\boldsymbol{\pi})$ as

$$(fV)_t := f(\cdot, t) V_t \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \quad (3.7)$$

(ii) To each $V \in \mathcal{L}^2(\boldsymbol{\pi})$ we associate the map $|V| \in L^2(\boldsymbol{\pi} \times \mathcal{L}_1)$, defined by

$$|V|(\gamma, t) := |V_t|(\gamma) \quad \text{for } (\boldsymbol{\pi} \times \mathcal{L}_1)\text{-a.e. } (\gamma, t) \in \Gamma(X) \times [0, 1].$$

It is clear that these operations give $\mathcal{L}^2(\boldsymbol{\pi})$ the structure of an $L^2(\boldsymbol{\pi} \times \mathcal{L}_1)$ -normed module.

We then define the linear continuous operator $\Phi : L^2(TX) \rightarrow \mathcal{L}^2(\boldsymbol{\pi})$ by putting

$$\Phi(v)_t := e_t^* v, \quad \text{for } \mathcal{L}_1\text{-a.e. } t \in [0, 1].$$

We then have:

Proposition 3.9 ($\mathcal{L}^2(\boldsymbol{\pi})$ as pullback) *We have $(\mathcal{L}^2(\boldsymbol{\pi}), \Phi) \sim (e^* L^2(TX), e^*)$, i.e.:*

$$\begin{aligned} |\Phi(v)| &= |v| \circ e \quad \text{holds } (\boldsymbol{\pi} \times \mathcal{L}_1)\text{-a.e., for any } v \in L^2(TX), \\ \{\Phi(v) : v \in L^2(TX)\} &\quad \text{generates } \mathcal{L}^2(\boldsymbol{\pi}) \text{ as a module.} \end{aligned} \quad (3.8)$$

Proof. The first in (3.8) follows by noticing that $|\Phi(v)|(\gamma, t) = |e_t^* v|(\gamma) = (|v| \circ e)(\gamma, t)$ holds for $(\boldsymbol{\pi} \times \mathcal{L}_1)$ -a.e. (γ, t) , the second one stems from the density of $\text{TestVF}(\boldsymbol{\pi})$ in $\mathcal{L}^2(\boldsymbol{\pi})$. \square

Notice that the notion of pullback module $(e^*L^2(TX), e^*)$ makes no (explicit) reference to the concept of ‘test vector field’ as defined it in Section 3.1. Thus this last proposition is also telling that the choice of using these test object to check Borel regularity, which a priori might seem arbitrary, leads in fact to a canonical interpretation of $\mathcal{L}^2(\boldsymbol{\pi})$.

Remark 3.10 ($\mathcal{L}^2(\boldsymbol{\pi})$ as direct integral) The construction of $\mathcal{L}^2(\boldsymbol{\pi})$ can be summarized by saying that such space is the direct integral of the $e_t^*L^2(TX)$, the space of Borel vector fields being the so-called ‘measurable sections’ and the set $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ being the one used to check measurability. \blacksquare

3.3 The space $\mathcal{C}(\boldsymbol{\pi})$

Here we introduce and briefly study those vector fields in $\text{VF}(\boldsymbol{\pi})$ which are ‘continuous in time’. We start with the following definition:

Definition 3.11 (The space $\mathcal{C}(\boldsymbol{\pi})$) *Let $V \in \text{VF}(\boldsymbol{\pi})$. Then we say that V is a continuous vector field provided*

$$[0, 1] \ni t \mapsto \int \langle V_t, W_t \rangle d\boldsymbol{\pi} \quad \text{is continuous} \quad (3.9)$$

for every $W \in \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ and

$$[0, 1] \ni t \mapsto \llbracket V \rrbracket_t \quad \text{is continuous.} \quad (3.10)$$

We denote the family of all continuous vector fields by $\mathcal{C}(\boldsymbol{\pi})$ and, for every $V \in \mathcal{C}(\boldsymbol{\pi})$, we put

$$\|V\|_{\mathcal{C}(\boldsymbol{\pi})} := \max_{t \in [0, 1]} \llbracket V \rrbracket_t.$$

Lemma 3.2 ensures that this definition would be unaltered if we require (3.9) to hold for any $W \in \text{TestVF}(\boldsymbol{\pi})$. Also, Proposition 3.1 gives that $\text{TestVF}(\boldsymbol{\pi}) \subset \mathcal{C}(\boldsymbol{\pi})$.

It is not obvious that $\mathcal{C}(\boldsymbol{\pi})$ is a vector space, the problem being in checking that (3.10) holds for linear combinations. This will be a consequence of the density of $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ in $\mathcal{C}(\boldsymbol{\pi})$, which is part of the content of the next result:

Proposition 3.12 ($\mathcal{C}(\boldsymbol{\pi}), \|\cdot\|_{\mathcal{C}(\boldsymbol{\pi})}$) *is a separable Banach space, with $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ being dense.*

Proof. Let $V_1, V_2 \in \mathcal{C}(\boldsymbol{\pi})$ and notice that using (3.9), (3.10) and arguing exactly as in the proof of Lemma 3.2 we can find $(W_{1,n}), (W_{2,n}) \subset \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ such that the functions $t \mapsto \llbracket V_i - W_{i,n} \rrbracket_t$ uniformly converge to 0 as $n \rightarrow \infty$, $i = 1, 2$.

Now observe that since $W_{1,n} + W_{2,n} \in \text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ the map $t \mapsto \llbracket W_{1,n} + W_{2,n} \rrbracket_t$ is continuous and that for every $t \in [0, 1]$ we have

$$\left| \llbracket V_1 + V_2 \rrbracket_t - \llbracket W_{1,n} + W_{2,n} \rrbracket_t \right| \leq \llbracket V_1 - W_{1,n} + V_2 - W_{2,n} \rrbracket_t \leq \llbracket V_1 - W_{1,n} \rrbracket_t + \llbracket V_2 - W_{2,n} \rrbracket_t.$$

Hence $t \mapsto \llbracket V_1 + V_2 \rrbracket_t$ is the uniform limit of continuous functions and thus continuous itself. Since trivially $\mathcal{C}(\boldsymbol{\pi})$ is closed by multiplication by scalars we proved that it is a vector space. That $\|\cdot\|_{\mathcal{C}(\boldsymbol{\pi})}$ is a complete norm on it is trivial and the density of $\text{TestVF}_{\mathbb{N}}(\boldsymbol{\pi})$ has already been shown, hence the proof is finished. \square

A useful consequence of the density of test vector fields is the following strengthening of the continuity property:

Corollary 3.13 *Let $V \in \mathcal{C}(\boldsymbol{\pi})$. Then the map $t \mapsto |V_t|^2 \in L^1(\boldsymbol{\pi})$ is continuous.*

Proof. For $V \in \text{TestVF}(\boldsymbol{\pi})$ the claim has been proved in Proposition 3.1. Now notice that for $V, W \in \mathcal{C}(\boldsymbol{\pi})$ we have

$$\int \| |V_t|^2 - |W_t|^2 \| \, d\boldsymbol{\pi} \leq \int |V_t + W_t| |V_t - W_t| \, d\boldsymbol{\pi} \leq (\|V\|_{\mathcal{C}(\boldsymbol{\pi})} + \|W\|_{\mathcal{C}(\boldsymbol{\pi})}) \llbracket V - W \rrbracket_t$$

thus showing that if $V_n \rightarrow V$ in $\mathcal{C}(\boldsymbol{\pi})$ then $t \mapsto |V_{n,t}|^2 \in L^1(\boldsymbol{\pi})$ uniformly converge to $t \mapsto |V_t|^2 \in L^1(\boldsymbol{\pi})$. The conclusion then follows from the density of $\text{TestVF}(\boldsymbol{\pi})$ in $\mathcal{C}(\boldsymbol{\pi})$. \square

3.4 The spaces $\mathcal{W}^{1,2}(\boldsymbol{\pi})$ and $\mathcal{H}^{1,2}(\boldsymbol{\pi})$

Throughout all this section we shall further assume that the test plan $\boldsymbol{\pi}$ is Lipschitz in the sense of Definition 2.14.

Let $v \in W_C^{1,2}(TX)$ and notice that the map from $L^0(TX)$ to $e_t^* L^0(TX)$ defined by

$$w \mapsto e_t^*(\nabla_w v)$$

satisfies

$$|e_t^*(\nabla_w v)| \leq |\nabla v|_{\text{HS}} \circ e_t |w| \circ e_t \quad \boldsymbol{\pi} - a.e..$$

Hence by the universal property of the pullback given in Proposition 2.9 we know that there exists a unique $L^0(\boldsymbol{\pi})$ -linear continuous operator, which we shall call $\text{Cov}(v, \cdot)$ from $e_t^* L^0(TX)$ to $e_t^* L^0(TX)$ such that

$$\text{Cov}(v, e_t^* w) = e_t^*(\nabla_w v) \quad \forall w \in L^0(TX)$$

and such operator satisfies the bound

$$|\text{Cov}(v, W)| \leq |\nabla v|_{\text{HS}} \circ e_t |W| \quad \boldsymbol{\pi} - a.e.. \quad (3.11)$$

We shall be interested in such covariant differentiation along the speed of our test plan: for every $t \in [0, 1]$ such that $\boldsymbol{\pi}'_t$ exists we define the map $\text{Cov}_t : W_C^{1,2}(TX) \rightarrow e_t^* L^0(TX)$ as

$$\text{Cov}_t(v) := \text{Cov}(v, \boldsymbol{\pi}'_t).$$

Notice the following simple proposition:

Proposition 3.14 *For every $t \in [0, 1]$ such that $\boldsymbol{\pi}'_t$ exists, the map Cov_t is linear and continuous from $W_C^{1,2}(TX)$ to $e_t^* L^2(TX)$.*

Moreover, for every $v \in W_C^{1,2}(TX)$ the (equivalence class up to a.e. equality of the) a.e. defined map $t \mapsto \text{Cov}_t(v) \in e_t^ L^2(TX)$ is an element of $\mathcal{L}^2(\boldsymbol{\pi})$.*

Proof. The continuity of Cov_t as map from $W_C^{1,2}(TX)$ to $e_t^*L^2(TX)$ is a direct consequence of the bounds (3.11) and our assumption (2.16):

$$\|\text{Cov}_t(v)\|_{e_t^*L^2(TX)}^2 = \int |\text{Cov}_t(v)|^2 d\pi \stackrel{(3.11)}{\leq} \int |\nabla v|_{\text{HS}}^2 \circ e_t |\pi'_t|^2 d\pi \leq C(\pi) L(\pi)^2 \|v\|_{W_C^{1,2}(TX)}^2.$$

Thanks to this bound, to conclude it is sufficient to show that for any $v \in W_C^{1,2}(TX)$ the map $t \mapsto \text{Cov}_t(v) = \text{Cov}(v, \pi'_t)$ is a.e. equal to a Borel element of $\text{VF}(\pi)$. Taking into account that $t \mapsto \pi'_t \in \mathcal{L}^2(\pi)$ by Proposition 3.8, that $\text{TestVF}(\pi)$ is dense in $\mathcal{L}^2(\pi)$, the second claim in Proposition 3.6 and the bound (3.11), we see that to conclude it is sufficient to show that $t \mapsto \text{Cov}(v, V_t)$ is a Borel vector field in $\text{VF}(\pi)$ for any $V \in \text{TestVF}(\pi)$.

Thus fix such V , say $V_t = \sum_i \phi_i(t) \chi_{A_i} e_t^* v_i$, and let $W_t = \sum_j \psi_j(t) \chi_{B_j} e_t^* w_j \in \text{TestVF}(\pi)$ be arbitrary. Notice that since $|v_i|, |w_j| \in L^2 \cap L^\infty(X)$, we have that $\langle v_i, \nabla_{w_j} v \rangle \in L^1(X)$ and thus by Theorem 2.11 we deduce that $t \mapsto \langle v_i, \nabla_{w_j} v \rangle \circ e_t \in L^1(\pi)$ is continuous for every i, j . Therefore

$$t \mapsto \int \langle V_t, \text{Cov}_t(v, W_t) \rangle d\pi = \sum_{i,j} \varphi_i(t) \psi_j(t) \int \chi_{A_i \cap B_j} \langle v_i, \nabla_{w_j} v \rangle \circ e_t d\pi$$

is continuous, thus establishing, by the arbitrariness of W , the Borel regularity of $t \mapsto \text{Cov}(v, V_t)$. \square

The ‘compatibility with the metric’ of the covariant derivative yields the following simple but crucial lemma:

Lemma 3.15 *Let $v, w \in \text{TestV}(X)$. Then the map $t \mapsto \langle v, w \rangle \circ e_t \in L^2(\pi)$, which is Lipschitz by Proposition 2.15, satisfies*

$$L^2(\pi) \text{-} \frac{d}{dt} \langle v, w \rangle \circ e_t = \langle \text{Cov}_\pi(v)_t, e_t^* w \rangle + \langle e_t^* v, \text{Cov}_\pi(w)_t \rangle \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \quad (3.12)$$

Proof. Recall from [8] that it holds

$$d(\langle v, w \rangle)(z) = \langle \nabla_z v, w \rangle + \langle v, \nabla_z w \rangle \quad \mathfrak{m} \text{- a.e.} \quad \forall z \in L^0(TX)$$

and notice that from the defining property of pointwise norm in the pullback and by polarization we obtain that $\langle e_t^* v_1, e_t^* v_2 \rangle = \langle v_1, v_2 \rangle \circ e_t$ for every $v_1, v_2 \in L^0(TX)$. Thus we have that the identity

$$(e_t^* d\langle v, w \rangle)(Z) = \langle \text{Cov}_t(v, Z), e_t^* w \rangle + \langle e_t^* v, \text{Cov}_t(w, Z) \rangle \quad (3.13)$$

holds for every $Z \in e_t^*L^2(TX)$ of the form $Z_t = e_t^*z$ for some $z \in L^2(TX)$. Since both sides of this identity are $L^\infty(\pi)$ -linear and continuous in Z , we see that (3.13) holds for generic $Z \in e_t^*L^2(TX)$. The conclusion comes picking $Z = \pi'_t$ and recalling Proposition 2.15. \square

We now want to introduce a new differential operator, initially defined only on $\text{TestVF}(\pi)$ and then extended to more general vector fields. To this aim the following lemma will be useful.

Lemma 3.16 *Let $(\varphi_i), (\psi_j) \subset \text{LIP}([0, 1])$, $(A_i), (B_j)$ Borel partitions of $\Gamma(X)$ and $(v_i), (w_j) \subset \text{TestV}(X)$, where $i = 1, \dots, n$ and $j = 1, \dots, m$. Assume that*

$$\sum_i \chi_{A_i} \varphi_i(t) e_t^* v_i = \sum_j \chi_{B_j} \psi_j(t) e_t^* w_j \quad \text{for every } t \in [0, 1]. \quad (3.14)$$

Then for a.e. t it holds:

$$\begin{aligned} \sum_i \chi_{A_i} \varphi'_i(t) e_t^* v_i &= \sum_j \chi_{B_j} \psi'_j(t) e_t^* w_j, \\ \sum_i \chi_{A_i} \varphi_i(t) \text{Cov}_t(v_i) &= \sum_j \chi_{B_j} \psi_j(t) \text{Cov}_t(w_j). \end{aligned} \quad (3.15)$$

Proof. For the first in (3.15) we notice that our assumption (3.14) and Proposition 3.9 yield that $\sum_i \chi_{A_i \times [0,1]}(\cdot, t) \varphi_i(t) e_t^* v_i = \sum_j \chi_{B_j \times [0,1]}(\cdot, t) \psi_j(t) e_t^* w_j$ as elements in $e^* L^2(TX) \sim \mathcal{L}^2(\boldsymbol{\pi})$, thus we can differentiate in time and conclude using again Proposition 3.9.

For the second in (3.15) start noticing that our assumption (3.14) and the very definition of pullback imply that for any i, j and every $t \in [0, 1]$ it holds $\chi_C \varphi_i(t) v_i = \chi_C \psi_j(t) w_j$, where $C := \left\{ \frac{d(e_t)_*(\chi_{A_i \cap B_j} \boldsymbol{\pi})}{d(e_t)_* \boldsymbol{\pi}} > 0 \right\}$. This identity and the locality of the covariant derivative give that $\chi_C \varphi_i(t) \nabla_z v_i = \chi_C \psi_j(t) \nabla_z w_j$ for every $z \in L^2(TX)$. Applying the pullback map on both sides and noticing that $\chi_C \circ e_t \geq \chi_{A_i \cap B_j}$ we deduce that

$$\chi_{A_i \cap B_j} \varphi_i(t) \text{Cov}(v_i, Z) = \chi_{A_i \cap B_j} \psi_j(t) \text{Cov}(w_j, Z)$$

for every Z of the form $Z_t = e_t^* z$. From the $L^\infty(\boldsymbol{\pi})$ -linearity in Z of both sides and the arbitrariness of i, j the conclusion follows. \square

We can now define the convective derivative of test vector fields:

Definition 3.17 (Convective derivative along a test plan) *We define the convective derivative operator $\tilde{D}_\boldsymbol{\pi} : \text{TestVF}(\boldsymbol{\pi}) \rightarrow \mathcal{L}^2(\boldsymbol{\pi})$ as follows: to the element $V \in \text{TestVF}(\boldsymbol{\pi})$, of the form $V_t = \sum_{i=1}^n \varphi_i(t) \chi_{A_i} e_t^* v_i$, we associate the vector field $\tilde{D}_\boldsymbol{\pi} V \in \mathcal{L}^2(\boldsymbol{\pi})$ given by*

$$(\tilde{D}_\boldsymbol{\pi} V)_t := \sum_{i=1}^n \chi_{A_i} \left(\varphi'_i(t) e_t^* v_i + \varphi_i(t) \text{Cov}_t(v_i) \right) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \quad (3.16)$$

For the sake of simplicity, we will briefly write $\tilde{D}_\boldsymbol{\pi} V_t$ instead of $(\tilde{D}_\boldsymbol{\pi} V)_t$.

Notice that Lemma 3.16 ensures that the right hand side of (3.16) depends only on V and not on the way we write it as $V_t = \sum_{i=1}^n \varphi_i(t) \chi_{A_i} e_t^* v_i$. The fact that the right hand side of (3.16) defines a Borel vector field in $\text{VF}(\boldsymbol{\pi})$ follows directly from Proposition 3.14; to see that it belongs to $\mathcal{L}^2(\boldsymbol{\pi})$ notice that $(t \mapsto e_t^* v_i), \text{Cov}_\boldsymbol{\pi}(v_i) \in \mathcal{L}^2(\boldsymbol{\pi})$ for every i and that the φ_i 's are Lipschitz.

Therefore the definition is well posed and is then clear that $\tilde{D}_\boldsymbol{\pi}$ is a linear operator.

The convective derivative has the following simple and crucial property, which is a direct consequence of Lemma 3.15.

Proposition 3.18 *Let $V, W \in \text{TestVF}(\boldsymbol{\pi})$. Then the map $t \mapsto \langle V_t, W_t \rangle \in L^2(\boldsymbol{\pi})$ is Lipschitz and satisfies*

$$L^2(\boldsymbol{\pi})\text{-}\frac{d}{dt} \langle V_t, W_t \rangle = \langle \tilde{D}_\boldsymbol{\pi} V_t, W_t \rangle + \langle V_t, \tilde{D}_\boldsymbol{\pi} W_t \rangle \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \quad (3.17)$$

Proof. By bilinearity, to prove (3.17) is sufficient to consider the case $V_t = \varphi(t) \chi_A e_t^* v$ and $W_t = \psi(t) \chi_B e_t^* w$ for $v, w \in \text{TestV}(X)$. Lemma 3.15 ensures that $t \mapsto \langle e_t^* v, e_t^* w \rangle = \langle v, w \rangle \circ e_t \in L^2(\boldsymbol{\pi})$ is Lipschitz and it is then clear that $t \mapsto \langle V_t, W_t \rangle = \chi_{A \cap B} \varphi(t) \psi(t) \langle e_t^* v, e_t^* w \rangle$ is also Lipschitz. The identity (3.17) now follows from (3.12) and the Leibniz rule. \square

This last proposition allows to ‘integrate by parts’ and extend the definition of convective derivative to ‘Sobolev vector fields along $\boldsymbol{\pi}$ ’.

Let us define the *support* $\text{spt}(V)$ of a test vector field $V \in \text{TestVF}(\boldsymbol{\pi})$ as the closure of the set of t ’s for which $V_t \neq 0$ and let us introduce the space of sections with compact support in $(0, 1)$:

$$\text{TestVF}_c(\boldsymbol{\pi}) := \{V \in \text{TestVF}(\boldsymbol{\pi}) : \text{spt}(V) \subseteq (0, 1)\}.$$

A simple cut-off argument shows that $\text{TestVF}_c(\boldsymbol{\pi})$ is $\mathcal{L}^2(\boldsymbol{\pi})$ -dense in $\text{TestVF}(\boldsymbol{\pi})$ and hence in $\mathcal{L}^2(\boldsymbol{\pi})$.

Definition 3.19 (The space $\mathscr{W}^{1,2}(\boldsymbol{\pi})$) *The Sobolev space $\mathscr{W}^{1,2}(\boldsymbol{\pi})$ is the vector subspace of $\mathcal{L}^2(\boldsymbol{\pi})$ consisting of all those $V \in \mathcal{L}^2(\boldsymbol{\pi})$ such that there exists $Z \in \mathcal{L}^2(\boldsymbol{\pi})$ satisfying*

$$\int_0^1 \int \langle V_t, \tilde{D}_\pi W_t \rangle d\boldsymbol{\pi} dt = - \int_0^1 \int \langle Z_t, W_t \rangle d\boldsymbol{\pi} dt \quad \text{for every } W \in \text{TestVF}_c(\boldsymbol{\pi}). \quad (3.18)$$

In this case the section Z , whose uniqueness is granted by density of $\text{TestVF}_c(\boldsymbol{\pi})$ in $\mathcal{L}^2(\boldsymbol{\pi})$, can be unambiguously denoted by $D_\pi V$ and called convective derivative of V . We endow $\mathscr{W}^{1,2}(\boldsymbol{\pi})$ with the norm $\|\cdot\|_{\mathscr{W}^{1,2}(\boldsymbol{\pi})}$, defined by

$$\|V\|_{\mathscr{W}^{1,2}(\boldsymbol{\pi})} := \sqrt{\|V\|_{\mathcal{L}^2(\boldsymbol{\pi})}^2 + \|D_\pi V\|_{\mathcal{L}^2(\boldsymbol{\pi})}^2} \quad \text{for every } V \in \mathscr{W}^{1,2}(\boldsymbol{\pi}).$$

This choice of terminology is consistent with that of Definition 3.17:

Proposition 3.20 *Let $V \in \text{TestVF}(\boldsymbol{\pi})$. Then $V \in \mathscr{W}^{1,2}(\boldsymbol{\pi})$ and $D_\pi V = \tilde{D}_\pi V$.*

Proof. Fix $W \in \text{TestVF}_c(\boldsymbol{\pi})$. We know from Proposition 3.18 that $[0, 1] \ni t \mapsto \int \langle V_t, W_t \rangle d\boldsymbol{\pi}$ is an absolutely continuous function, so that (3.17) gives, after integration, that

$$0 = \int \langle V_1, W_1 \rangle d\boldsymbol{\pi} - \int \langle V_0, W_0 \rangle d\boldsymbol{\pi} = \int_0^1 \int \langle \tilde{D}_\pi V_t, W_t \rangle d\boldsymbol{\pi} dt + \int_0^1 \int \langle V_t, \tilde{D}_\pi W_t \rangle d\boldsymbol{\pi} dt.$$

This proves that V satisfies (3.18) with $Z = \tilde{D}_\pi V$. \square

Proposition 3.21 (Basic properties of $\mathscr{W}^{1,2}(\boldsymbol{\pi})$) *The following hold:*

- i) D_π is a closed operator from $\mathcal{L}^2(\boldsymbol{\pi})$ into itself, i.e. its graph is closed in the product space $\mathcal{L}^2(\boldsymbol{\pi}) \times \mathcal{L}^2(\boldsymbol{\pi})$.
- ii) $\mathscr{W}^{1,2}(\boldsymbol{\pi})$ is a separable Hilbert space.
- iii) Let $V, Z \in \mathcal{L}^2(\boldsymbol{\pi})$. Then $V \in \mathscr{W}^{1,2}(\boldsymbol{\pi})$ and $Z = D_\pi V$ if and only if for every $W \in \text{TestVF}(\boldsymbol{\pi})$ the map $t \mapsto \langle V_t, W_t \rangle$ belongs to $W^{1,1}([0, 1], L^1(\boldsymbol{\pi}))$ with derivative given by

$$\frac{d}{dt} \langle V_t, W_t \rangle = \langle V_t, D_\pi W_t \rangle + \langle Z_t, W_t \rangle \quad \text{a.e. } t. \quad (3.19)$$

Proof.

(i) Let $(V_n) \subseteq \mathscr{W}^{1,2}(\boldsymbol{\pi})$ be a sequence such that $V_n \rightarrow V$ and $D_\pi V_n \rightarrow Z$ in $\mathcal{L}^2(\boldsymbol{\pi})$ for some $V, Z \in \mathcal{L}^2(\boldsymbol{\pi})$. Then for arbitrary $W \in \text{TestVF}_c(\boldsymbol{\pi})$ we have

$$\begin{aligned} \int_0^1 \int \langle V_t, D_\pi W_t \rangle d\boldsymbol{\pi} dt &= \lim_{n \rightarrow \infty} \int_0^1 \int \langle V_t^n, D_\pi W_t \rangle d\boldsymbol{\pi} dt \\ &= - \lim_{n \rightarrow \infty} \int_0^1 \int \langle D_\pi V_t^n, W_t \rangle d\boldsymbol{\pi} dt = - \int_0^1 \int \langle Z_t, W_t \rangle d\boldsymbol{\pi} dt, \end{aligned}$$

proving that $V \in \mathscr{W}^{1,2}(\boldsymbol{\pi})$ with $D_{\boldsymbol{\pi}}V = Z$, which was the claim.

(ii) Consequence of what just proved and the fact that the map

$$\mathscr{W}^{1,2}(\boldsymbol{\pi}) \ni V \mapsto (V, D_{\boldsymbol{\pi}}V) \in \mathscr{L}^2(\boldsymbol{\pi}) \times \mathscr{L}^2(\boldsymbol{\pi})$$

is an isometry, provided we endow $\mathscr{L}^2(\boldsymbol{\pi}) \times \mathscr{L}^2(\boldsymbol{\pi})$, with the (separable, by Proposition 3.7) norm $\|(V, Z)\|^2 := \|V\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 + \|Z\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2$.

(iii) The ‘if’ trivially follows from (3.19) by integration. For the ‘only if’, fix $W \in \text{TestVF}(\boldsymbol{\pi})$ and let $\varphi \in C_c^1(0, 1)$ and $\Gamma \subset \Gamma(X)$ Borel. Then $t \mapsto \varphi(t)\chi_{\Gamma}W_t$ is in $\text{TestVF}_c(\boldsymbol{\pi})$ and a direct computation shows that $D_{\boldsymbol{\pi}}(\varphi\chi_{\Gamma}W)_t = \varphi'(t)\chi_{\Gamma}W_t + \varphi(t)\chi_{\Gamma}D_{\boldsymbol{\pi}}W_t$. Hence writing the defining property (3.18) with $\varphi\chi_{\Gamma}W$ in place of W we obtain, after rearrangement, that

$$\int_0^1 \varphi'(t) \int_{\Gamma} \langle V_t, W_t \rangle d\boldsymbol{\pi} dt = - \int_0^1 \varphi(t) \int_{\Gamma} \langle V_t, D_{\boldsymbol{\pi}}W_t \rangle + \langle Z_t, W_t \rangle d\boldsymbol{\pi} dt.$$

The arbitrariness of φ, Γ and Proposition 2.3 yield the claim. \square

We just proved that $\text{TestVF}(\boldsymbol{\pi})$ is contained in $\mathscr{W}^{1,2}(\boldsymbol{\pi})$, but we don’t know if it is dense. Hence the following definition is meaningful:

Definition 3.22 (The space $\mathscr{H}^{1,2}(\boldsymbol{\pi})$) $\mathscr{H}^{1,2}(\boldsymbol{\pi})$ is the $\mathscr{W}^{1,2}(\boldsymbol{\pi})$ -closure of $\text{TestVF}(\boldsymbol{\pi})$.

Clearly, $\mathscr{H}^{1,2}(\boldsymbol{\pi})$ is a separable Hilbert space. A key feature of elements of $\mathscr{H}^{1,2}(\boldsymbol{\pi})$ is that they admit a continuous representative (much like Sobolev functions on the interval):

Theorem 3.23 The inclusion $\text{TestVF}(\boldsymbol{\pi}) \hookrightarrow \mathscr{C}(\boldsymbol{\pi})$ uniquely extends to a linear continuous and injective operator $\iota : \mathscr{H}^{1,2}(\boldsymbol{\pi}) \rightarrow \mathscr{C}(\boldsymbol{\pi})$.

Proof. We claim that

$$\|V\|_{\mathscr{C}(\boldsymbol{\pi})} \leq \sqrt{2} \|V\|_{\mathscr{W}^{1,2}(\boldsymbol{\pi})} \quad \forall V \in \text{TestVF}(\boldsymbol{\pi}). \quad (3.20)$$

By the density of $\text{TestVF}(\boldsymbol{\pi})$ in $\mathscr{H}^{1,2}(\boldsymbol{\pi})$ this will be enough to obtain the existence of ι . Thus let $V \in \text{TestVF}(\boldsymbol{\pi})$, pick $W = V$ in (3.17) and integrate in $[t_1, t_2] \subset [0, 1]$ and w.r.t. $\boldsymbol{\pi}$ to obtain

$$\begin{aligned} |[\![V]\!]_{t_2}^2 - [\![V]\!]_{t_1}^2| &= 2 \left| \int_{t_1}^{t_2} \int \langle V_t, D_{\boldsymbol{\pi}}V_t \rangle d\boldsymbol{\pi} dt \right| \\ &\leq 2 \int_{t_1}^{t_2} \int |V_t| |D_{\boldsymbol{\pi}}V_t| d\boldsymbol{\pi} dt \leq \|V\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 + \|D_{\boldsymbol{\pi}}V\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2. \end{aligned}$$

Hence for any $t \in [0, 1]$ one has

$$[\![V]\!]_t^2 = \int_0^1 [\![V]\!]_t^2 ds \leq \int_0^1 |[\![V]\!]_t^2 - [\![V]\!]_s^2| ds + \|V\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 \leq 2 \|V\|_{\mathscr{W}^{1,2}(\boldsymbol{\pi})}^2,$$

which is our claim (3.20).

To prove injectivity, let $V \in \mathscr{H}^{1,2}(\boldsymbol{\pi})$ be such that $\iota(V) = 0$. Choose a sequence $(V_n) \subseteq \text{TestVF}(\boldsymbol{\pi})$ which is $\mathscr{W}^{1,2}(\boldsymbol{\pi})$ -converging to V and notice that, up to pass to a subsequence and using Proposition 3.6, we can assume that $V_t^n \rightarrow V_t$ for \mathscr{L}^1 -a.e. $t \in [0, 1]$. By continuity of the operator ι , one also has $\|V^n\|_{\mathscr{C}(\boldsymbol{\pi})} = \|\iota(V^n) - \iota(V)\|_{\mathscr{C}(\boldsymbol{\pi})} \rightarrow 0$ and thus in particular $V_t^n \rightarrow 0$ for all $t \in [0, 1]$. Therefore $V_t = 0$ for \mathscr{L}^1 -a.e. $t \in [0, 1]$, yielding the required injectivity of ι . \square

Whenever we will consider an element V of $\mathcal{H}^{1,2}(\boldsymbol{\pi})$, we will always implicitly refer to its unique continuous representative $\iota(V) \in \mathcal{C}(\boldsymbol{\pi})$.

Among the several properties of the test sections that can be carried over to the elements of $\mathcal{H}^{1,2}(\boldsymbol{\pi})$, the most important one is the Leibniz formula for the convective derivatives:

Proposition 3.24 (Leibniz formula for D_π) *Let $V \in \mathcal{W}^{1,2}(\boldsymbol{\pi})$ and $W \in \mathcal{H}^{1,2}(\boldsymbol{\pi})$. Then the function $t \mapsto \langle V_t, W_t \rangle$ is in $W^{1,1}([0, 1], L^1(\boldsymbol{\pi}))$ and its derivative is given by*

$$\frac{d}{dt} \langle V_t, W_t \rangle = \langle D_\pi V_t, W_t \rangle + \langle V_t, D_\pi W_t \rangle \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1].$$

Proof. For $W \in \text{TestVF}(\boldsymbol{\pi})$ the claim is a direct consequence of point (iii) in Proposition 3.21. The general case can be achieved by approximation noticing that the simple inequalities

$$\begin{aligned} \|\langle V_t, W_t \rangle\|_{L^1(\boldsymbol{\pi} \times \mathcal{L}_1)} &\leq \|V\|_{\mathcal{L}^2(\boldsymbol{\pi})} \|W\|_{\mathcal{L}^2(\boldsymbol{\pi})}, \\ \|\langle D_\pi V_t, W_t \rangle + \langle V_t, D_\pi W_t \rangle\|_{L^1(\boldsymbol{\pi} \times \mathcal{L}_1)} &\leq 2\|V\|_{\mathcal{W}^{1,2}(\boldsymbol{\pi})} \|W\|_{\mathcal{W}^{1,2}(\boldsymbol{\pi})}, \end{aligned}$$

allow to pass to the limit in the distributional formulation of $\frac{d}{dt} \langle V_t, W_t \rangle$ as W varies in $\mathcal{H}^{1,2}(\boldsymbol{\pi})$. \square

In the next proposition we collect some examples of elements of $\mathcal{H}^{1,2}(\boldsymbol{\pi})$:

Proposition 3.25 *Let $\boldsymbol{\pi}$ be a Lipschitz test plan. Then:*

i) *For every $w \in H_C^{1,2}(TX)$ the vector field $t \mapsto W_t := e_t^* w$ belongs to $\mathcal{H}^{1,2}(\boldsymbol{\pi})$ and*

$$D_\pi W_t = \text{Cov}_t(w) \quad \text{a.e. } t. \quad (3.21)$$

ii) *Let $W \in \mathcal{H}^{1,2}(\boldsymbol{\pi})$ be such that $|W|, |D_\pi W| \in L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)$ and $a \in W^{1,2}([0, 1], L^2(\boldsymbol{\pi}))$. Then $aW \in \mathcal{H}^{1,2}(\boldsymbol{\pi})$ with*

$$D_\pi(aW)_t = a'_t W_t + a_t D_\pi W_t \quad \text{a.e. } t. \quad (3.22)$$

Moreover, if $W \in \mathcal{C}(\boldsymbol{\pi})$ and $a \in AC^2([0, 1], L^2(\boldsymbol{\pi}))$, then $aW \in \mathcal{C}(\boldsymbol{\pi})$.

Proof.

(i) If $w \in \text{TestV}(X)$ we have that $W \in \text{TestVF}(\boldsymbol{\pi})$ by definition and in this case formula (3.21) holds by the definition (3.16) and Proposition 3.20. The general case can then be obtained by approximating w with vector fields in $\text{TestV}(X)$ w.r.t. the $W_C^{1,2}(TX)$ topology, using the bounds

$$\begin{aligned} \int_0^1 \int |e_t^*(v)|^2 d\boldsymbol{\pi} dt &= \int_0^1 \int |v|^2 \circ e_t d\boldsymbol{\pi} dt \leq C(\boldsymbol{\pi}) \|v\|_{W_C^{1,2}(TX)}^2, \\ \int_0^1 \int |\text{Cov}_\pi(v)_t|^2 d\boldsymbol{\pi} dt &\stackrel{(3.11)}{\leq} C(\boldsymbol{\pi}) L(\boldsymbol{\pi})^2 \|v\|_{W_C^{1,2}(TX)}^2 \end{aligned}$$

and recalling the closure of D_π .

(ii) The claim about continuity is obvious, so we concentrate on the other one. Assume at first that a belongs to the space \mathcal{A} defined as

$$\mathcal{A} := \left\{ \sum_{i=1}^n \varphi_i \chi_{E_i} : n \in \mathbb{N}, \varphi_i \in \text{LIP}([0, 1]), (E_i) \text{ Borel partition of } \Gamma(X) \right\}$$

and that $W \in \text{TestVF}(\boldsymbol{\pi})$. In this case aW belongs to $\text{TestVF}(\boldsymbol{\pi})$ as well and formula (3.22) is a direct consequence of the definitions. Then using the trivial bounds

$$\begin{aligned} \|aW\|_{\mathcal{L}^2(\boldsymbol{\pi})} &\leq \|a\|_{L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)} \|W\|_{\mathcal{L}^2(\boldsymbol{\pi})}, \\ \|a'W + aD_{\boldsymbol{\pi}}W\|_{\mathcal{L}^2(\boldsymbol{\pi})} &\leq (\|a\|_{L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)} + \|a'\|_{L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)}) \|W\|_{\mathcal{W}^{1,2}(\boldsymbol{\pi})}, \end{aligned}$$

the $\mathcal{W}^{1,2}(\boldsymbol{\pi})$ -density of $\text{TestVF}(\boldsymbol{\pi})$ in $\mathcal{H}^{1,2}(\boldsymbol{\pi})$ and the closure of $D_{\boldsymbol{\pi}}$, we conclude that $aW \in \mathcal{H}^{1,2}(\boldsymbol{\pi})$ for every $a \in \mathcal{A}$ and $W \in \mathcal{H}^{1,2}(\boldsymbol{\pi})$ and that (3.22) holds in this case.

Now let W be as in the assumptions and notice that we also have the bounds

$$\begin{aligned} \|aW\|_{\mathcal{L}^2(\boldsymbol{\pi})} &\leq \|a\|_{L^2([0,1], L^2(\boldsymbol{\pi}))} \|W\|_{L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)}, \\ \|a'W + aD_{\boldsymbol{\pi}}W\|_{\mathcal{L}^2(\boldsymbol{\pi})} &\leq \|a\|_{W^{1,2}([0,1], L^2(\boldsymbol{\pi}))} (\|W\|_{L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)} + \|D_{\boldsymbol{\pi}}W\|_{L^\infty(\boldsymbol{\pi} \times \mathcal{L}_1)}), \end{aligned}$$

therefore using again the closure of $D_{\boldsymbol{\pi}}$, to conclude it is sufficient to prove that \mathcal{A} is dense in $W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$. To this aim we argue as follows: for every $n \in \mathbb{N}$ let $(E_i^n)_{i \in \mathbb{N}}$ be a Borel partition of $\text{supp}(\boldsymbol{\pi}) \subset \Gamma(\mathbf{X})$ made of sets with positive $\boldsymbol{\pi}$ -measure and diameter $\leq \frac{1}{n}$. Then for every $n, N \in \mathbb{N}$ let $P_n^N : L^2(\boldsymbol{\pi}) \rightarrow L^2(\boldsymbol{\pi})$ be defined by

$$P_n^N(f) := \sum_{i=1}^N \chi_{E_i^n} \frac{1}{\boldsymbol{\pi}(E_i^n)} \int_{E_i^n} f \, d\boldsymbol{\pi}.$$

It is clear that P_n^N has operator norm ≤ 1 for every $n, N \in \mathbb{N}$ and an application of the dominated convergence theorem shows that

$$\lim_n \lim_N P_n^N(f) = f \tag{3.23}$$

for every $f \in C_b(\Gamma(\mathbf{X}))$, the limits being intended in $L^2(\boldsymbol{\pi})$. Therefore (3.23) also holds for every $f \in L^2(\boldsymbol{\pi})$. The linearity and continuity of P_n^N also grants that if $t \mapsto a_t$ belongs to $W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$, then also $t \mapsto P_n^N(a)_t := P_n^N(a_t)$ belongs to $W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$ with

$$(P_n^N(a))'_t = P_n^N(a'_t) \quad a.e. \, t. \tag{3.24}$$

All these considerations imply that

$$\lim_n \lim_N P_n^N(a) = a \quad \text{in } W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$$

for every $a \in W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$ and thus to conclude it is sufficient to prove that $P_n^N(a)$ belongs to the $W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$ -closure of \mathcal{A} for every $n, N \in \mathbb{N}$ and $a \in W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$.

It is clear by construction and (3.24) that $P_n^N(a) = \sum_{i=1}^N g_i \chi_{E_i^n}$ for some $g_i \in W^{1,2}([0,1])$. Now for every $i = 1, \dots, N$ find $(g_{i,j}) \subset \text{LIP}([0,1])$ which $W^{1,2}([0,1])$ -converges to g_i and notice that

$$\left\| \sum_{i=1}^N (g_{i,j} - g_i) \chi_{E_i^n} \right\|_{W^{1,2}([0,1], L^2(\boldsymbol{\pi}))}^2 = \sum_{i=1}^N \boldsymbol{\pi}(E_i^n) \|g_{i,j} - g_i\|_{W^{1,2}([0,1])}^2 \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Since $\sum_{i=1}^N g_{i,j} \chi_{E_i^n} \in \mathcal{A}$ for every j , the proof is finished. \square

4 Parallel transport on RCD spaces

4.1 Definition and basic properties of parallel transport

4.1.1 Definition and uniqueness

We shall frequently use the fact that, since $\mathcal{H}^{1,2}(\pi)$ is continuously embedded into $\mathcal{C}(\pi)$ by Theorem 3.23, any vector field $V \in \mathcal{H}^{1,2}(\pi)$ has pointwise values $V_t \in e_t^* L^2(TX)$ defined at every time $t \in [0, 1]$.

Definition 4.1 (Parallel transport) *Let $K \in \mathbb{R}$, (X, d, \mathfrak{m}) an RCD(K, ∞) space and π be a Lipschitz test plan on X . A parallel transport along π is an element $V \in \mathcal{H}^{1,2}(\pi)$ such that $D_\pi V = 0$.*

The linearity of the requirement $D_\pi V = 0$ ensures that the set of parallel transports along π is a vector space. From Proposition 3.24 we deduce the following simple but crucial result:

Proposition 4.2 (Norm preservation) *Let V be a parallel transport along the Lipschitz test plan π . Then $t \mapsto |V_t|^2 \in L^1(\pi)$ is constant.*

Proof. We know from Corollary 3.13 that $t \mapsto |V_t|^2 \in L^1(\pi)$ is continuous. Hence the choice $W = V$ in Proposition 3.24 tells that such map is absolutely continuous with derivative given by

$$\frac{d}{dt} |V_t|^2 = 2 \langle D_\pi V_t, V_t \rangle = 0, \quad a.e. \ t.$$

This is sufficient to conclude. □

Linearity and norm preservation imply uniqueness:

Corollary 4.3 (Uniqueness of parallel transport) *Let π be a Lipschitz test plan and V_1, V_2 two parallel transports along it such that for some $t_0 \in [0, 1]$ it holds $V_{1,t_0} = V_{2,t_0}$. Then $V_1 = V_2$.*

Proof. Since $D_\pi(V_1 - V_2) = D_\pi V_1 - D_\pi V_2 = 0$, we have that $V_1 - V_2$ is a parallel transport and by assumption we know that $|V_{1,t_0} - V_{2,t_0}| = 0$ π -a.e.. Thus Proposition 4.2 above grants that for every $t \in [0, 1]$ it holds $|V_{1,t} - V_{2,t}| = 0$ π -a.e., i.e. that $V_{1,t} = V_{2,t}$. □

Remark 4.4 We emphasize that the norm preservation property is a consequence of the Leibniz formula in Proposition 3.24. We don't know if such formula holds for $V, W \in \mathcal{W}^{1,2}(\pi)$ and this is why we defined the parallel transport as an element of $\mathcal{H}^{1,2}(\pi)$ with null convective derivative, as opposed to an element of $\mathcal{W}^{1,2}(\pi)$ with the same property. ■

4.1.2 Some consequences of existence of parallel transport

In this section we assume existence of parallel transport along some/all Lipschitz test plans and see what can be derived from such assumption.

It will be convenient to recall the concept of *base* of a module, referring to [8] for a more detailed discussion. Let μ be a Borel measure on a Polish space Y , \mathcal{M} a $L^2(\mu)$ -normed module, $v_1, \dots, v_n \in \mathcal{M}$ and $E \subset Y$ a Borel set. Then the v_i 's are said to be *independent* on E provided for any $f_i \in L^\infty(\mu)$ we have

$$\sum_i f_i v_i = 0 \quad \Rightarrow \quad f_i = 0 \quad \mu - a.e. \text{ on } E$$

and generators of \mathcal{M} on E provided $L^\infty(\mu)$ -linear combinations of the v_i 's are dense in $\{\chi_{Ev} : v \in \mathcal{M}\}$. If v_1, \dots, v_n are both independent and generators of \mathcal{M} on E we say that they are a base of \mathcal{M} on E and in this case we say that the dimension of \mathcal{M} on E is n .

Recall that there always exists a (unique up to μ -negligible sets) Borel partition $(E_i)_{i \in \mathbb{N} \cup \{\infty\}}$ of Y , called *dimensional decomposition of \mathcal{M}* , such that the dimension of \mathcal{M} on E_i is i for every $i \in \mathbb{N}$ and for no Borel subset F of E_∞ with positive measure the dimension of \mathcal{M} on F is finite.

For a separable Hilbert module \mathcal{H} on a space with finite measure μ one can always find an *orthonormal base* $(v_n)_{n \in \mathbb{N}}$ i.e. a sequence whose L^∞ -linear combinations are dense in \mathcal{H} and such that for some Borel partition $(E_n)_{n \in \mathbb{N} \cup \{\infty\}}$ the following hold:

$$\begin{aligned} \forall n \in \mathbb{N} \cup \{\infty\} \text{ we have} & \quad \langle v_i, v_j \rangle = \delta_{ij} & \quad \mu - a.e. \text{ on } E_n \quad \forall i, j \in \mathbb{N}, \quad i, j < n \\ \forall n \in \mathbb{N} \text{ we have} & \quad |v_i| = 0 & \quad \mu - a.e. \text{ on } E_n \quad \forall i \in \mathbb{N}, \quad i \geq n \end{aligned}$$

and it is easily verified that if these hold, then necessarily the E_n 's form the dimensional decomposition of \mathcal{H} (the role of the assumption about finiteness of μ is to ensure that the v_i 's are elements of \mathcal{H} : their pointwise norm is in $L^\infty(\mu)$ and thus in general it may be not in $L^2(\mu)$).

Given such a base and $v \in \mathcal{H}$ there are functions $a_n \in L^2(\mu)$ such that

$$v = \sum_{n \in \mathbb{N}} a_n v_n \tag{4.1}$$

meaning that the sequence converges absolutely in \mathcal{H} . A choice for the a_n is $a_n := \langle v, v_n \rangle$ and for any two sequences $(a_n), (\tilde{a}_n)$ for which (4.1) holds we have

$$a_n = \tilde{a}_n \quad \mu - a.e. \text{ on } E_m, \quad \forall m \in \mathbb{N} \cup \{\infty\}, \quad m > n.$$

With this said, it is easy to see that parallel transport sends bases into bases:

Proposition 4.5 *Let π be a Lipschitz test plan such that for every $t \in [0, 1]$ and $\bar{V}_t \in e_t^* L^2(TX)$ there exists a (necessarily unique) parallel transport V along π such that $V_t = \bar{V}_t$.*

Also, let $(E_n)_{n \in \mathbb{N} \cup \{\infty\}}$ be the dimensional decomposition of $e_0^ L^2(TX)$ and $(\bar{V}_n) \subset e_0^* L^2(TX)$, $n \in \mathbb{N}$ an orthonormal base of $e_0^* L^2(TX)$ and denote by $t \mapsto V_{n,t}$ the parallel transport of \bar{V}_n along π .*

Then for every $t \in [0, 1]$ the partition $(E_n)_{n \in \mathbb{N} \cup \{\infty\}}$ is also the dimensional decomposition of $e_t^ L^2(TX)$ and the set $(V_{n,t})$ is an orthonormal base of $e_t^* L^2(TX)$.*

Proof. For every $t, s \in [0, 1]$ consider the map sending $\bar{V} \in e_t^* L^2(TX)$ to $V_s \in e_s^* L^2(TX)$ where $V \in \mathcal{H}^{1,2}(\pi)$ is the parallel transport such that $V_t = \bar{V}$. Proposition 4.2 ensures that this map preserves the pointwise norm. Since it is clearly linear, it is easily verified that it must be an isomorphism of $e_t^* L^2(TX)$ and $e_s^* L^2(TX)$.

The conclusions follow. \square

We shall apply this result to show that, under the same assumptions, we have $\mathcal{W}^{1,2}(\pi) = \mathcal{H}^{1,2}(\pi)$:

Proposition 4.6 ($\mathcal{H} = \mathcal{W}$) *Let π be a Lipschitz test plan such that for every $t \in [0, 1]$ and $\bar{V}_t \in e_t^* L^2(TX)$ there exists a (necessarily unique) parallel transport V along π such that $V_t = \bar{V}_t$.*

Then $\mathcal{H}^{1,2}(\pi) = \mathcal{W}^{1,2}(\pi)$.

Proof. Let $V \in \mathscr{W}^{1,2}(\boldsymbol{\pi})$, $(\bar{V}_i) \subset e_0^*L^2(T\mathbf{X})$, $i \in \mathbb{N}$, an orthonormal base of $e_0^*L^2(T\mathbf{X})$ and $t \mapsto V_{i,t}$ the parallel transport of \bar{V}_i along $\boldsymbol{\pi}$. Then by Proposition 4.5 above we see that

$$V_t = \sum_{i \in \mathbb{N}} a_{i,t} V_{i,t} \quad \text{where} \quad a_{i,t} := \langle V_t, V_{i,t} \rangle \quad \text{a.e. } t, \quad (4.2)$$

being intended that the series converges absolutely in $e_t^*L^2(T\mathbf{X})$ for a.e. t . By Proposition 3.24 we see that $t \mapsto a_{i,t}$ is in $W^{1,1}([0,1], L^1(\boldsymbol{\pi}))$ with derivative given by

$$a'_{i,t} = \langle D_{\boldsymbol{\pi}} V_t, V_{i,t} \rangle. \quad (4.3)$$

In particular, since $|V_{i,t}| \leq 1$ we see that $a_{i,t}, a'_{i,t} \in L^2([0,1], L^2(\boldsymbol{\pi}))$ and in turn this implies - by Proposition 2.3 - that $t \mapsto a_{i,t}$ is in $W^{1,2}([0,1], L^2(\boldsymbol{\pi}))$. This fact and point (ii) in Proposition 3.25 give that $(t \mapsto a_{i,t} V_{i,t}) \in \mathscr{H}^{1,2}(\boldsymbol{\pi})$ for every $i \in \mathbb{N}$ and therefore $(t \mapsto \sum_{i=0}^n a_{i,t} V_{i,t}) \in \mathscr{H}^{1,2}(\boldsymbol{\pi})$ for every $n \in \mathbb{N}$.

Hence to conclude it is sufficient to show that these partial sums are a $\mathscr{W}^{1,2}(\boldsymbol{\pi})$ -Cauchy sequence, as then it is clear from (4.2) that the limit coincides with V . From (4.2) and (4.3) we have that

$$\sum_{i \in \mathbb{N}} \iint_0^1 |a_{i,t}|^2 + |a'_{i,t}|^2 dt d\boldsymbol{\pi} = \|V\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 + \|D_{\boldsymbol{\pi}} V\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 < \infty,$$

hence the conclusion follows from the identity

$$\left\| \sum_{i=n}^m a_i V_i \right\|_{\mathscr{W}^{1,2}(\boldsymbol{\pi})}^2 = \left\| \sum_{i=n}^m a_i V_i \right\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 + \left\| \sum_{i=n}^m a'_i V_i \right\|_{\mathscr{L}^2(\boldsymbol{\pi})}^2 = \sum_{i=n}^m \iint_0^1 |a_{i,t}|^2 + |a'_{i,t}|^2 dt d\boldsymbol{\pi}.$$

□

We shall now prove that if the parallel transport exists along all Lipschitz test plans, then the dimension of \mathbf{X} , intended here as the dimension of the tangent module, must be constant. We shall use the following simple lemma (for simplicity we state it for Hilbert modules, but in fact the same holds for general ones):

Lemma 4.7 *Let $(\mathbf{X}, d_{\mathbf{X}}, \mathfrak{m}_{\mathbf{X}})$ and $(\mathbf{Y}, d_{\mathbf{Y}}, \mathfrak{m}_{\mathbf{Y}})$ be two metric measure spaces, $\varphi : \mathbf{Y} \rightarrow \mathbf{X}$ of bounded compression and \mathscr{H} an Hilbert module on \mathbf{X} . Then for every $E \subset \mathbf{X}$ Borel we have that the dimension of \mathscr{H} on E is n if and only if the dimension of $\varphi^* \mathscr{H}$ on $\varphi^{-1}(E)$ is n .*

Proof. From the well-posedness of the definition of dimension we see that it is sufficient to prove the *only if*. Thus let v_0, \dots, v_{n-1} be an orthonormal base of \mathscr{H} on E , so in particular $\langle v_i, v_j \rangle = \delta_{ij}$ $\mathfrak{m}_{\mathbf{X}}$ -a.e. on E .

From the fact that v_0, \dots, v_{n-1} generate \mathscr{H} on E and the very definition of pullback we deduce that $\varphi^* v_0, \dots, \varphi^* v_{n-1}$ generate $\varphi^* \mathscr{H}$ on $\varphi^{-1}(E)$.

To see that they are independent, let $f_0, \dots, f_{n-1} \in L^\infty(\mathbf{Y})$ be such that $\sum_i f_i \varphi^* v_i = 0$ on $\varphi^{-1}(E)$. Then it holds

$$0 = \left| \sum_i f_i \varphi^* v_i \right|^2 = \sum_{i,j} f_i f_j \langle \varphi^* v_i, \varphi^* v_j \rangle = \sum_{i,j} f_i f_j \langle v_i, v_j \rangle \circ \varphi = \sum_i f_i^2 \quad \mathfrak{m}_{\mathbf{Y}}\text{-a.e. on } \varphi^{-1}(E),$$

and therefore $f_i = 0$ $\mathfrak{m}_{\mathbf{Y}}$ -a.e. on $\varphi^{-1}(E)$ for every $i = 0, \dots, n-1$. □

We now prove that the dimension is constant:

Theorem 4.8 (From parallel transport to constant dimension) *Let (X, d, \mathbf{m}) be a $\text{RCD}(K, \infty)$ space such that for any Lipschitz test plan π , any $t \in [0, 1]$ and any $\bar{V} \in e_t^* L^2(TX)$ there exists the parallel transport V along π such that $V_t = \bar{V}$.*

Then the tangent module has constant dimension, i.e. in its dimensional decomposition $(E_i)_{i \in \mathbb{N} \cup \{\infty\}}$ one of the E_i 's has full measure and the others are negligible.

Proof. We argue by contradiction, thus we shall assume that for some $i, j \in \mathbb{N} \cup \{\infty\}$, $i \neq j$, we have $\mathbf{m}(E_i), \mathbf{m}(E_j) > 0$. Let $F_0 \subset E_i$ and $F_1 \subset E_j$ be bounded, with positive and finite measure, consider

$$\mu_0 := \mathbf{m}(F_0)^{-1} \mathbf{m}|_{F_0} \quad \mu_1 := \mathbf{m}(F_1)^{-1} \mathbf{m}|_{F_1},$$

let π be the unique optimal geodesic plan connecting them and recall that it is a test plan (see [14]). Since $\pi(e_0^{-1}(F_0)) = \mu_0(F_0) = 1$ and the dimension of $L^2(TX)$ on F_0 is i , by Lemma 4.7 above we see that for the dimensional decomposition $(\tilde{E}_n^0)_{n \in \mathbb{N} \cup \{\infty\}}$ of $e_0^* L^2(TX)$ we have $\pi(\tilde{E}_i^0) = 1$ and $\pi(\tilde{E}_k^0) = 0$ for every $k \neq i$. Similarly, for the the dimensional decomposition $(\tilde{E}_n^1)_{n \in \mathbb{N} \cup \{\infty\}}$ of $e_1^* L^2(TX)$ we have $\pi(\tilde{E}_j^1) = 1$ and $\pi(\tilde{E}_k^1) = 0$ for every $k \neq j$. In particular we have

$$\pi(\tilde{E}_i^0 \Delta \tilde{E}_i^1) = \pi(\tilde{E}_i^0) = 1 > 0. \quad (4.4)$$

Now notice that from basic considerations about optimal transport we have that π is concentrated on geodesics starting from F_0 and ending in F_1 . The constant speed of any such geodesic is bounded from above by $\sup_{x \in F_0, y \in F_1} d(x, y) < \infty$ so that π is a Lipschitz test plan. Therefore from Proposition 4.5 we know that the dimensional decomposition of $e_t^* L^2(TX)$ does not depend on t . This however contradicts (4.4), hence the proof is completed. \square

4.2 Existence of the parallel transport in a special case

It is unclear to us whether on general RCD spaces the parallel transport exists or not. Aim of this section it to show at least that the theory we propose is not empty, i.e. that under suitable assumptions on the space, the parallel transport exists. We won't insist in trying to make such assumptions as general as possible (for instance, the 'good base' defined below could consist in different vector fields on different open sets covering our space) as our main concern is only to show that in some circumstances our notion of parallel transport can be shown to exist.

We shall work with spaces admitting the following sort of base for the tangent module:

Definition 4.9 (Good base) *Let (X, d, \mathbf{m}) be a given $\text{RCD}(K, N)$ space, for some $K \in \mathbb{R}$ and $N \in (1, \infty)$. Let us denote by $(A_k)_{k=1}^n$ the dimensional decomposition of X . Then a family $\mathcal{W} = \{w_1, \dots, w_n\} \subseteq H_C^{1,2}(TX)$ of Sobolev vector fields on X is said to be a good basis for $L^2(TX)$ provided there exists $M > 0$ such that the following properties are satisfied:*

(i) *For any $k = 1, \dots, n$, we have that w_1, \dots, w_k constitute a basis for $L^2(TX)$ on A_k and*

$$\begin{cases} |w_i| \in (M^{-1}, M), \\ |\langle w_i, w_j \rangle| < \frac{1}{M^{2k}} \end{cases} \quad \mathbf{m}\text{-a.e. in } A_k, \quad \text{for every } i, j = 1, \dots, k \text{ with } i \neq j, \quad (4.5)$$

(ii) *We have*

$$|\nabla w_i|_{\text{HS}} \leq M \quad \mathbf{m}\text{-a.e. in } X, \quad \forall i = 1, \dots, n. \quad (4.6)$$

Let us notice that the ‘hard’ assumption here is given by point (ii) (perhaps coupled with the lower bound in (i)), which imposes an L^∞ bound on covariant derivative, when in our setting L^2 -ones are more natural (compare with Theorem A.2). Let us mention in particular that for spaces admitting a good base it is not hard to prove, regardless of parallel transport, that the dimension is constant (see Proposition 4.11 below).

Let us start the technical work with the following simple lemma:

Lemma 4.10 *Let \mathcal{H} be a Hilbert module on X , $A \subset \mathsf{X}$ be a Borel set and $M > 1$. Also, for $k \in \mathbb{N}$ let $w_1, \dots, w_k \in \mathcal{H}$ be such that*

$$\left\{ \begin{array}{l} |w_i| \in (M^{-1}, M) \\ |\langle w_i, w_j \rangle| \leq \frac{1}{M^2 k} \end{array} \right. \text{ hold } \mathbf{m}\text{-a.e. in } A, \quad \text{for every } i, j = 1, \dots, k \text{ with } i \neq j.$$

For $h_1, \dots, h_k \in L^0(\mathfrak{m})|_A$ put $w := \sum_{i=1}^k h_i w_i \in \mathcal{H}^0$ (the L^0 -completion of \mathcal{H}). Then it holds

$$\frac{1}{M^2 k} \sum_{i=1}^k |h_i|^2 \leq |w|^2 \leq M^2 k \sum_{i=1}^k |h_i|^2 \quad \mathbf{m} - \text{a.e. on } A \quad (4.7)$$

and in particular $w \in \mathcal{H}|_A$ if and only if $h_i \in L^2(\mathfrak{m})|_A$ for every $i = 1, \dots, k$.

Proof. For the second in (4.7) we notice that $\langle w_i, w_j \rangle \leq M^2$ \mathbf{m} -a.e. on A for every i, j , thus

$$|w|^2 = \left| \sum_{i=1}^k h_i w_i \right|^2 = \sum_{i,j=1}^k h_i h_j \langle w_i, w_j \rangle \leq M^2 \sum_{i,j=1}^k \frac{1}{2} |h_i|^2 + \frac{1}{2} |h_j|^2 = M^2 k \sum_{i=1}^k |h_i|^2.$$

For the first inequality we recall that $|w_i| > M^{-1}$ and $\langle w_i, w_j \rangle \geq -\frac{1}{M^2 k}$ for $i \neq j$ \mathbf{m} -a.e. on A to deduce

$$\begin{aligned} |w|^2 &= \left| \sum_{i=1}^k h_i w_i \right|^2 = \sum_{i=1}^k |h_i|^2 |w_i|^2 + \sum_{i \neq j} h_i h_j \langle w_i, w_j \rangle \geq \frac{1}{M^2} \sum_{i=1}^k |h_i|^2 - \frac{1}{M^2 k} \sum_{i \neq j} |h_i h_j| \\ &\geq \frac{1}{M^2} \sum_{i=1}^k |h_i|^2 - \frac{1}{M^2 k} \sum_{i \neq j} \frac{1}{2} |h_i|^2 + \frac{1}{2} |h_j|^2 = \frac{1}{M^2 k} \sum_{i=1}^k |h_i|^2. \end{aligned}$$

□

The constant dimension now easily follows from the lemma and the fact that if a good base exists, then there is another one for which the functions $\langle w_i, w_j \rangle$ are Lipschitz, as shown in the proof of the following proposition.

Proposition 4.11 *Let $(\mathsf{X}, \mathbf{d}, \mathfrak{m})$ be a $\text{RCD}(K, N)$ space admitting a good base for $L^2(\text{TX})$. Then the tangent module has constant dimension, i.e. in its dimensional decomposition $(E_k)_{k=1}^n$ one of the E_k ’s has full measure and the others are negligible.*

Proof. Let $k \in \mathbb{N}$ be the maximal index such that $\mathfrak{m}(E_k) > 0$ (its existence follows from the finiteness results in [16] and [12]). To conclude it is enough to show that on a neighbourhood of E_k the tangent module has dimension $\geq k$. Let $(w_i)_{i=1}^k$ be a good base, $f \in C_c^\infty(\mathbb{R})$ be such that $f(z) = z$ for every $z \in [0, M]$ and consider the vector fields $\tilde{w}_i := f(|w_i|^2) w_i$. Notice that by [8] we know that $f(|w_i|^2) \in W^{1,2}(\mathsf{X})$ with $\nabla f(|w_i|^2) = 2f'(|w_i|^2) \nabla w_i(\cdot, w_i)$, hence by

(4.6) and the choice of f we see that $f(|w_i|^2)$ is bounded with bounded gradient. It follows (see [8]) that $\tilde{w}_i \in H_C^{1,2}(TX)$ with

$$\nabla \tilde{w}_i = \nabla f(|w_i|^2) \otimes w_i + f(|w_i|^2) \nabla w_i,$$

so that from the expression of $\nabla f(|w_i|^2)$ we see that \tilde{w}_i is bounded with bounded covariant derivative. Hence $g_{i,j} := \langle \tilde{w}_i, \tilde{w}_j \rangle$ belongs to $W^{1,2}(X)$ (see [8]) and is bounded with bounded gradient as well. By the Sobolev-to-Lipschitz property (see [3], [7], [9]) we deduce that $g_{i,j}$ has a Lipschitz, in particular continuous, representative. By construction, the bounds (4.5) hold on E_k for the \tilde{w}_i 's, hence the continuity of $g_{i,j}$ grants that they hold also on some neighbourhood of E_k , and by Lemma 4.10 above this is sufficient to conclude that the vector fields \tilde{w}_i are independent - by the first in (4.7) - on such neighbourhood, thus concluding the proof. \square

We now prove existence of the parallel transport for the class of those RCD spaces that admit a good base for their tangent module. In the proof we shall use, for simplicity, the fact just proved that the dimension must be constant, but in fact the same argument works even without knowing a priori this fact (albeit since constant dimension follows so directly from the existence of a good base, this remark is perhaps irrelevant).

Theorem 4.12 (Existence of the parallel transport) *Let (X, d, m) be any RCD(K, N) space, $K \in \mathbb{R}$ and $N \in (1, \infty)$, that admits a good base. Let π be a Lipschitz test plan on X and fix $\bar{V} \in e_0^* L^2(TX)$.*

Then there exists the parallel transport $V \in \mathcal{H}^{1,2}(\pi)$ along π such that $V_0 = \bar{V}$.

Proof. We know by Proposition 4.11 that the dimension of the tangent module must be constant. From this fact and Definition 4.9 we know that for some n we have $w_1, \dots, w_n \in H_C^{1,2}(TX)$ for which (4.5) and (4.6) hold on X . Put $W_{i,t} := e_t^* w_i$. By point (i) in Proposition 3.25 we have that $W_i \in \mathcal{H}^{1,2}(\pi)$ with

$$D_\pi W_{i,t} = \text{Cov}_t(w_i)$$

and therefore from (3.11), (4.6) and the assumption that π is Lipschitz we get

$$|D_\pi W_{i,t}| \leq ML(\pi). \quad (4.8)$$

Also, by the defining property of the pullback map and from (4.5) we have that for every $t \in [0, 1]$ it holds

$$\begin{cases} |W_{i,t}| \in (M^{-1}, M), \\ |\langle W_{i,t}, W_{j,t} \rangle| \leq (M^2 k)^{-1} \end{cases} \quad \pi\text{-a.e.}, \quad \text{for every } i, j = 1, \dots, n \text{ with } i \neq j,$$

thus Lemma 4.10 grants that there are functions $\bar{g}_1, \dots, \bar{g}_n \in L^2(\pi)$ such that $\bar{V} = \sum_{i=1}^n \bar{g}_i e_0^* w_i$. A similar argument applied to the pullback of the map e (recall Proposition 3.9 and the definition (3.7)) and based on the bound (4.8) shows that there are functions $H_{i,j} \in L^\infty(\pi \times \mathcal{L}_1)$ such that

$$D_\pi W_{i,t} = \sum_j H_{i,j,t} W_{j,t} \quad a.e. \ t. \quad (4.9)$$

It will be technically convenient to fix once and for all Borel representatives, still denoted by $H_{i,j}$, of these functions such that

$$\sup_{\gamma,t} |H_{i,j,t}(\gamma)| = \|H_{i,j}\|_{L^\infty(\pi \times \mathcal{L}_1)} \quad \forall i = 1, \dots, n. \quad (4.10)$$

We shall look for a parallel transport of the form $V := \sum_i g_i W_i$ with $g_i \in AC^2([0, 1], L^2(\pi))$. Notice that Lemma 4.10 grants that any such V belongs to $\mathcal{H}^{1,2}(\pi)$ with

$$\begin{aligned} D_\pi V_t &\stackrel{(3.22)}{=} \sum_{i=1}^n g'_{i,t} W_{i,t} + \sum_{i=1}^n g_{i,t} D_\pi W_{i,t} \stackrel{(4.9)}{=} \sum_{i=1}^n g'_{i,t} W_{i,t} + \sum_{i,j=1}^n g_{i,t} H_{i,j,t} W_{j,t} \\ &= \sum_{i=1}^n \left(g'_{i,t} + \sum_{j=1}^n H_{j,i,t} g_{j,t} \right) W_{i,t} \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \end{aligned}$$

Hence our V is the desired parallel transport if and only if the functions g_1, \dots, g_n solve the system

$$\begin{cases} g_{i,0} = \bar{g}_i & \forall i = 1, \dots, n, \\ g'_{i,t} + \sum_{j=1}^n H_{j,i,t} g_{j,t} = 0 & \forall i = 1, \dots, n \text{ and for } \mathcal{L}^1\text{-a.e. } t \in [0, 1]. \end{cases} \quad (4.11)$$

To solve this system we shall apply Theorem 2.6 to the Banach (in fact Hilbert) space $\mathbb{B} := [L^2(\pi)]^n$ equipped with the norm

$$\|f\|_{\mathbb{B}}^2 := \sum_{i=1}^n \int |f_i|^2 d\pi \quad \forall f = (f_1, \dots, f_n) \in \mathbb{B}.$$

For every $t \in [0, 1]$ define $\lambda_t \in \text{End}(\mathbb{B})$ as

$$(\lambda_t f)_i := - \sum_{j=1}^n H_{j,i,t} f_j \quad \forall i = 1, \dots, n \text{ and } f = (f_1, \dots, f_n) \in \mathbb{B},$$

so that the system (4.11) can be rewritten as

$$\begin{cases} g_0 = \bar{g}, \\ g'_t = \lambda_t g_t \quad \text{for a.e. } t \in [0, 1], \end{cases}$$

where $\bar{g} := (\bar{g}_1, \dots, \bar{g}_n)$. Theorem 2.6 grants that a solution in $\text{LIP}([0, 1], \mathbb{B}) \subset AC^2([0, 1], \mathbb{B})$ exists provided the λ_t 's are equibounded and $t \mapsto \lambda_t f$ is strongly measurable for every $f \in \mathbb{B}$. The former follows from

$$\begin{aligned} \|\lambda_t f\|_{\mathbb{B}}^2 &= \sum_{i=1}^n \left\| \sum_{j=1}^n H_{j,i,t} f_j \right\|_{L^2(\pi)}^2 \leq n \sum_{i,j=1}^n \|H_{j,i,t} f_j\|_{L^2(\pi)}^2 \\ &\leq n \max_{i,j,t} \|H_{j,i,t}\|_{L^\infty(\pi)}^2 \sum_{i,j=1}^n \|f_j\|_{L^2(\pi)}^2 \stackrel{(4.10)}{\leq} n^2 \max_{i,j} \|H_{i,j}\|_{L^\infty(\pi \times \mathcal{L}_1)}^2 \|f\|_{\mathbb{B}}^2. \end{aligned}$$

For the latter, notice that since \mathbb{B} is separable it is sufficient to prove that for any $f \in \mathbb{B}$ the map $t \mapsto \lambda_t f \in \mathbb{B}$ is weakly measurable. Since \mathbb{B} is also Hilbert we need to show that for every $f, g \in \mathbb{B}$ the map $t \mapsto \langle \lambda_t f, g \rangle_{\mathbb{B}} \in \mathbb{R}$ is measurable. Since we have

$$\langle \lambda_t f, g \rangle_{\mathbb{B}} = - \sum_{i,j=1}^n \int H_{j,i,t} f_j g_i d\pi,$$

the conclusion follows from Fubini's theorem. \square

A Sobolev base of the tangent module

In this appendix we show that one can always build a base of the tangent module of a RCD space which has at least Sobolev regularity, as opposed to just L^2 regularity. The basic idea used in the construction is based on the observation that ‘being a base’ is a non-linear requirement. Technically speaking, the crucial argument is contained in the following lemma:

Lemma A.1 *Let (X, d, \mathfrak{m}) be a $\text{RCD}(K, \infty)$ space and $w \in L^2(TX)$.*

Then there exists $v \in H_C^{1,2}(TX)$ such that $\langle v, w \rangle \neq 0$ holds \mathfrak{m} -a.e. on $\{|w| \neq 0\}$.

Proof. We can assume $w \neq 0$ or otherwise there is nothing to prove; then replacing if necessary w with $(\chi_{\{|w| \leq 1\}} + \chi_{\{|w| > 1\}}|w|^{-1})w$ we can also assume that $|w| \leq 1$ \mathfrak{m} -a.e.. Let $(w_n) \subset \text{TestV}(X)$ be $L^2(TX)$ -converging to w and $\tilde{\mathfrak{m}}$ a Borel probability measure on X such that $\mathfrak{m} \ll \tilde{\mathfrak{m}} \leq \mathfrak{m}$. Then $\langle w_n, w \rangle \rightarrow |w|^2$ in $L^2(X, \tilde{\mathfrak{m}})$ and thus

$$m_n := \tilde{\mathfrak{m}}(\{|\langle w_n, w \rangle| > 0\}) \quad \rightarrow \quad m_\infty := \tilde{\mathfrak{m}}(\{|w| > 0\}). \quad (\text{A.1})$$

We now observe that

$$\begin{aligned} &\text{for every } v, \tilde{w} \in L^2(TX) \text{ and } a > 0 \text{ there is } b \in (0, a) \text{ such that} \\ &\tilde{\mathfrak{m}}\left(\{|\langle \tilde{w}, w \rangle| > 0\} \cap \{v + b\tilde{w}, w = 0\}\right) = 0. \end{aligned} \quad (\text{A.2})$$

Indeed, putting for brevity $E_b := \{|\langle \tilde{w}, w \rangle| > 0\} \cap \{v + b\tilde{w}, w = 0\}$ we have

$$\tilde{\mathfrak{m}}(E_b \cap E_{b'}) \leq \tilde{\mathfrak{m}}\left(\{|\langle \tilde{w}, w \rangle| > 0\} \cap \{(b - b')\langle \tilde{w}, w \rangle = 0\}\right) = 0 \quad \forall b \neq b',$$

so that the claim follows from the finiteness of $\tilde{\mathfrak{m}}$ and the fact that the interval $(0, a)$ is uncountable.

Now put $\alpha_n := \| |w_n| \|_{L^\infty(X)} + \|w_n\|_{W_C^{1,2}(TX)}$ and recursively define decreasing sequences $(\beta_n), (\gamma_n) \subset (0, \infty)$ such that $\beta_1 = 1$ and for every $n \in \mathbb{N}$ we have

$$3\beta_{n+1} \leq \gamma_{n+1} \leq \beta_n \quad \text{and for } E_n := \left\{ \left| \left\langle \sum_{i=1}^n \frac{\beta_i}{\alpha_i} w_i, w \right\rangle \right| \geq \gamma_{n+1} \right\} \quad \text{it holds } \tilde{\mathfrak{m}}(E_n) \geq \frac{m_n}{1 + \frac{1}{n}}.$$

To see that this is possible, let $\beta_1 = 1$, and notice that trivially $\{|\langle \frac{\beta_1}{\alpha_1} w_1, w \rangle| > 0\} = \{|\langle w_1, w \rangle| > 0\}$ so that for $\gamma_2 \in (0, \beta_1)$ sufficiently small the above holds. Now assume that β_{n-1} and γ_n have been found, use (A.2) for $v := \sum_{i=1}^{n-1} \frac{\beta_i}{\alpha_i} w_i$, $\tilde{w} := w_n$ and $a := \gamma_n/3$ to find $\beta_n := b < \gamma_n/3$ such that $\tilde{\mathfrak{m}}(\{|\langle \sum_{i=1}^n \frac{\beta_i}{\alpha_i} w_i, w \rangle| > 0\}) \geq \tilde{\mathfrak{m}}(\{|\langle w_n, w \rangle| > 0\}) = m_n$. Hence for $\gamma_{n+1} \in (0, \beta_n)$ sufficiently small the claim holds.

We claim that the vector $v := \sum_{i \geq 1} \frac{\beta_i}{\alpha_i} w_i$ satisfies the conclusion of the statement and start observing that $\beta_i \leq 3^{-i}$ and thus $\| \frac{\beta_i}{\alpha_i} w_i \|_{W_C^{1,2}(TX)} \leq 3^{-i} \| \alpha_i^{-1} w_i \|_{W_C^{1,2}(TX)} \leq 3^{-i}$ by definition of α_i . Hence the series converges in $W_C^{1,2}(TX)$, so that v is well defined and belongs to $H_C^{1,2}(TX)$. Now notice that by construction and (A.1) we have $\tilde{\mathfrak{m}}(E_n) \rightarrow m_\infty$ and $\tilde{\mathfrak{m}}(E_n \setminus \{|w| > 0\}) = 0$, so that $\tilde{\mathfrak{m}}(\{|w| > 0\} \setminus \cup_n E_n) = 0$. Hence to conclude it is sufficient to show that for every $n \geq 1$ it holds $\langle v, w \rangle \neq 0$ $\tilde{\mathfrak{m}}$ -a.e. on E_n . Fix such n , let $m > n$ and observe that by definition of the α_i 's and β_i 's we have

$$\left| \left\langle \frac{\beta_m}{\alpha_m} w_m, w \right\rangle \right| \leq 3^{n-m+1} \beta_{n+1} \left| \langle \alpha_m^{-1} w_m, w \rangle \right| \leq 3^{n-m+1} \beta_{n+1}, \quad \tilde{\mathfrak{m}} - a.e.,$$

so that $|\sum_{m>n} \langle \frac{\beta_m}{\alpha_m} w_m, w \rangle| \leq \frac{3}{2} \beta_{n+1} \leq \frac{1}{2} \gamma_{n+1}$. On the other hand by construction we have that $|\sum_{i=1}^n \langle \frac{\beta_i}{\alpha_i} w_i, w \rangle| \geq \gamma_{n+1}$ holds $\tilde{\mathbf{m}}$ -a.e. on E_n , granting that $|\langle v, w \rangle| \geq \frac{1}{2} \gamma_{n+1}$ $\tilde{\mathbf{m}}$ -a.e. on E_n . \square

By repeatedly applying Lemma A.1, we can find a family of $H_C^{1,2}(TX)$ -Sobolev generators of the tangent module on any $\text{RCD}(K, \infty)$ space X , as follows:

Theorem A.2 (Sobolev base of the tangent module) *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space, for some constant $K \in \mathbb{R}$. Suppose that the dimensional decomposition of X is given by $(A_n)_{n \in \mathbb{N}}$. Then there exists a sequence of vector fields $(v_n)_{n \geq 1} \subseteq H_C^{1,2}(TX)$ such that*

$$v_1, \dots, v_n \text{ is a basis for } L^2(TX) \text{ on } A_n, \quad \text{for every } n \in \mathbb{N}^+.$$

Proof. The thesis can be equivalently rewritten in the following way:

$$v_1, \dots, v_n \text{ are independent on } \bigcup_{k \geq n} A_k, \quad \text{for every } n \in \mathbb{N}^+. \quad (\text{A.3})$$

We build the sequence $(v_n)_n$ by means of a recursive argument. First of all, choose a vector field $w \in L^2(TX)$ such that $0 < |w| \leq 1$ \mathbf{m} -a.e. in $\bigcup_{k \geq 1} A_k$, then pick $v_1 \in H_C^{1,2}(TX)$ such that $\langle v_1, w \rangle \neq 0$ \mathbf{m} -a.e. in $\bigcup_{k \geq 1} A_k$, whose existence is granted by Lemma A.1. Thus in particular we have $|v_1| > 0$ \mathbf{m} -a.e. in $\bigcup_{k \geq 1} A_k$, proving (A.3) for $n = 1$. Now suppose to have already found v_1, \dots, v_n satisfying the required property. It can be easily seen that there exists $w \in L^2(TX)$ such that $\langle v_1, w \rangle = \dots = \langle v_n, w \rangle = 0$ and $0 < |w| \leq 1$ hold \mathbf{m} -a.e. in the set $\bigcup_{k > n} A_k$. Hence Lemma A.1 ensures the existence of a vector field $v_{n+1} \in H_C^{1,2}(TX)$ such that $\langle v_{n+1}, w \rangle \neq 0$ \mathbf{m} -a.e. in $\bigcup_{k > n} A_k$.

Now take any $f_1, \dots, f_{n+1} \in L^\infty(\mathbf{m})$ such that $\sum_{i=1}^{n+1} f_i v_i = 0$ \mathbf{m} -a.e. in $\bigcup_{k > n} A_k$, thus one has $f_{n+1} \langle v_{n+1}, w \rangle = \sum_{i=1}^{n+1} f_i \langle v_i, w \rangle = 0$ \mathbf{m} -a.e. in $\bigcup_{k > n} A_k$, from which we can deduce that $f_{n+1} = 0$ holds \mathbf{m} -a.e. in $\bigcup_{k > n} A_k$. Therefore $\sum_{i=1}^n f_i v_i = 0$ \mathbf{m} -a.e. in $\bigcup_{k > n} A_k$ and accordingly also $f_1 = \dots = f_n = 0$ \mathbf{m} -a.e. in $\bigcup_{k > n} A_k$, as a consequence of the independence of v_1, \dots, v_n . This grants that the vector fields v_1, \dots, v_{n+1} are independent on $\bigcup_{k > n} A_k$, proving (A.3) for $n + 1$. The thesis is then achieved. \square

References

- [1] L. AMBROSIO, M. COLOMBO, AND S. DI MARINO, *Sobolev spaces in metric measure spaces: reflexivity and lower semicontinuity of slope*. Accepted at Adv. St. in Pure Math., arXiv:1212.3779, 2014.
- [2] L. AMBROSIO, N. GIGLI, AND G. SAVARÉ, *Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below*, Invent. Math., 195 (2014), pp. 289–391.
- [3] —, *Metric measure spaces with Riemannian Ricci curvature bounded from below*, Duke Math. J., 163 (2014), pp. 1405–1490.
- [4] T. H. COLDING AND A. NABER, *Sharp Hölder continuity of tangent cones for spaces with a lower Ricci curvature bound and applications*, Ann. of Math. (2), 176 (2012), pp. 1173–1229.

- [5] J. DIESTEL AND J. J. UHL, JR., *Vector measures*, American Mathematical Society, Providence, R.I., 1977. With a foreword by B. J. Pettis, Mathematical Surveys, No. 15.
- [6] N. GIGLI, *Lecture notes on differential calculus on RCD spaces*. Preprint, arXiv:1703.06829.
- [7] —, *The splitting theorem in non-smooth context*. Preprint, arXiv:1302.5555, 2013.
- [8] —, *Nonsmooth differential geometry - an approach tailored for spaces with Ricci curvature bounded from below*. Accepted at Mem. Amer. Math. Soc., arXiv:1407.0809, 2014.
- [9] —, *An overview of the proof of the splitting theorem in spaces with non-negative Ricci curvature*, Analysis and Geometry in Metric Spaces, 2 (2014), pp. 169–213.
- [10] —, *On the differential structure of metric measure spaces and applications*, Mem. Amer. Math. Soc., 236 (2015), pp. vi+91.
- [11] N. GIGLI AND E. PASQUALETTO, *Behaviour of the reference measure on RCD spaces under charts*. Accepted at Comm. Anal. Geom., arxiv: 1607.05188, 2016.
- [12] —, *Equivalence of two different notions of tangent bundle on rectifiable metric measure spaces*. Preprint, arXiv:1611.09645, 2016.
- [13] —, *work in progress*.
- [14] N. GIGLI, T. RAJALA, AND K.-T. STURM, *Optimal Maps and Exponentiation on Finite-Dimensional Spaces with Ricci Curvature Bounded from Below*, J. Geom. Anal., 26 (2016), pp. 2914–2929.
- [15] N. GIGLI AND C. RIGONI, *Recognizing the flat torus among $\text{RCD}^*(0, N)$ spaces via the study of the first cohomology group*. Work in progress.
- [16] B. HAN, *Ricci tensor on $\text{RCD}^*(K, N)$ spaces*. Preprint, arXiv: 1412.0441.
- [17] M. KELL AND A. MONDINO, *On the volume measure of non-smooth spaces with Ricci curvature bounded below*. Preprint, arXiv:1607.02036.
- [18] J. LOTT AND C. VILLANI, *Ricci curvature for metric-measure spaces via optimal transport*, Ann. of Math. (2), 169 (2009), pp. 903–991.
- [19] A. MONDINO AND A. NABER, *Structure Theory of Metric-Measure Spaces with Lower Ricci Curvature Bounds*. Accepted at J. Eur. Math. Soc., arXiv:1405.2222, 2014.
- [20] A. PETRUNIN, *Parallel transportation for Alexandrov space with curvature bounded below*, Geom. Funct. Anal., 8 (1998), pp. 123–148.
- [21] K.-T. STURM, *On the geometry of metric measure spaces. I*, Acta Math., 196 (2006), pp. 65–131.
- [22] —, *On the geometry of metric measure spaces. II*, Acta Math., 196 (2006), pp. 133–177.
- [23] C. VILLANI, *Inégalités isopérimétriques dans les espaces métriques mesurés [d’après F. Cavalletti & A. Mondino]*. Séminaire Bourbaki, available at: <http://www.bourbaki.ens.fr/TEXTES/1127.pdf>.

- [24] —, *Synthetic theory of ricci curvature bounds*, Japanese Journal of Mathematics, 11 (2016), pp. 219–263.