

HOMOGENIZATION OF HIGH-CONTRAST MUMFORD-SHAH ENERGIES

XAVIER PELLET, LUCIA SCARDIA, AND CATERINA IDA ZEPPIERI

ABSTRACT. We prove a homogenization result for Mumford-Shah-type energies associated to a brittle composite material with weak inclusions distributed periodically at a scale $\varepsilon > 0$. The matrix and the inclusions in the material have the same elastic moduli but very different toughness moduli, with the ratio of the toughness modulus in the matrix and in the inclusions being $1/\beta_\varepsilon$, with $\beta_\varepsilon > 0$ small. We show that the high-contrast behaviour of the composite leads to the emergence of interesting effects in the limit: The volume and surface energy densities interact by Γ -convergence, and the limit volume energy is not a quadratic form in the critical scaling $\beta_\varepsilon = \varepsilon$, unlike the ε -energies, and unlike the extremal limit cases.

Keywords: Homogenization, Γ -convergence, free-discontinuity problems, high-contrast materials, brittle fracture.

MSC 2010: 49J45, 49Q20, 74Q05.

1. INTRODUCTION

We study the homogenization of a family of Mumford-Shah-type free-discontinuity functionals representing the (linearly) elastic energy of a *high-contrast* composite material constituted by a brittle matrix with weak inclusions distributed periodically. Our analysis is restricted to the case of an anti-plane shear, namely to scalar displacements $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$, where $\Omega \subset \mathbb{R}^n$ is a bounded and open set with Lipschitz boundary representing the cross-section of the reference configuration $\Omega \times \mathbb{R}$. The energy we consider is

$$F_\varepsilon(u) = \int_{\Omega} |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap \varepsilon P) + \beta_\varepsilon \mathcal{H}^{n-1}(S_u \cap (\Omega \setminus \varepsilon P)), \quad (1.1)$$

where $\varepsilon > 0$ is the ratio between the size of the microstructure and the observable length scale, and εP is the ε -scaled copy of a connected, Q -periodic, open set $P \subset \mathbb{R}^n$, with $Q = (-1/2, 1/2)^n$, which models the matrix of the composite. The displacement u is in the class $SBV^2(\Omega)$ of special functions of bounded variation, ∇u denotes its approximate gradient and S_u its discontinuity set. The volume term in F_ε represents the linearly elastic energy of the body, and the surface integral describes the energy needed to open a crack in the material. Note that the matrix and the inclusions have the same elastic moduli (normalised to 1), but very different toughness moduli: the toughness modulus is 1 in the matrix and $\beta_\varepsilon > 0$ in the inclusions, with $\beta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. This is why we call the brittle composite high-contrast.

The literature on high-contrast materials and on the derivation of their effective behaviour by homogenization is vast. In the classical Sobolev case, it is well known that interesting effects appear in the limit when the volume energy density does not satisfy uniform lower bounds (see, e.g., [2, 14, 15], and [8] for the case of discrete energies). It is then natural to try to extend this analysis to the case of free-discontinuity functionals, and there has been a recent effort in this direction. Note that for free-discontinuity functionals the high contrast can be in the volume part of the energy [3, 5], in the surface part [23, 24], or in both [12, 4, 22, 11].

For the functional (1.1) the high-contrast behaviour is in the surface term. We show that depending on how small β_ε is, with respect to the ‘critical’ value ε , the effective behaviour of the high-contrast material is different. To see this we introduce the parameter

$$\ell := \lim_{\varepsilon \rightarrow 0} \frac{\beta_\varepsilon}{\varepsilon} \in [0, +\infty],$$




FIGURE 1. Schematic of a periodic brittle material with weak inclusions

and characterise the Γ -limit of F_ε in the case $\ell = 0$ (subcritical regime), $\ell \in (0, +\infty)$ (critical regime), and $\ell = +\infty$ (supercritical regime).

1.1. Abstract Γ -convergence result and the choice of the convergence. As a first step we prove that there exists an infinitesimal sequence (ε_k) along which F_{ε_k} Γ -converges to a functional $F_{\text{hom}}^{\ell, (\varepsilon_k)}$ (depending on the sequence (ε_k)) which can be represented in an integral form as

$$F_{\text{hom}}^{\ell, (\varepsilon_k)}(u) = \begin{cases} \int_{\Omega} f_{\text{hom}}^{\ell, (\varepsilon_k)}(\nabla u) dx + \int_{S_u} g_{\text{hom}}^{\ell, (\varepsilon_k)}([u], \nu_u) d\mathcal{H}^{n-1} & \text{if } u \in GSBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega), \end{cases}$$

where $f_{\text{hom}}^{\ell, (\varepsilon_k)}$ and $g_{\text{hom}}^{\ell, (\varepsilon_k)}$ depend on the mutual vanishing rate of β_ε and ε , that is on ℓ , and $GSBV^2(\Omega)$ is the space of generalised special functions with bounded variation.

In this step the choice of the convergence plays a crucial role and introduces some difficulties. The functional F_ε in (1.1) is non-coercive with respect to the $L^1(\Omega)$ topology, due to the infinitesimal prefactor β_ε for the measure of the jump set in the inclusions $\Omega \setminus \varepsilon P$. In [12], where the authors considered the Mumford-Shah functional on a perforated domain, namely

$$\widehat{F}_\varepsilon(u) := \begin{cases} \int_{\Omega \cap \varepsilon P} |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap \varepsilon P) & \text{if } u|_{\Omega \cap \varepsilon P} \in GSBV^2(\Omega \cap \varepsilon P), \\ +\infty & \text{otherwise in } L^1(\Omega), \end{cases} \quad (1.2)$$

the (even more prominent) lack of coerciveness was solved by means of an extension result. Namely, each sequence (u_ε) with equibounded energy \widehat{F}_ε was replaced by a new, improved sequence \tilde{u}_ε which coincides with u_ε outside the perforations and which is precompact in $L^1(\Omega)$. Since $\widehat{F}_\varepsilon(u_\varepsilon) = \widehat{F}_\varepsilon(\tilde{u}_\varepsilon)$, this substitution does not affect the energy, and hence it is natural to study the Γ -convergence of \widehat{F}_ε with respect to the strong convergence in $L^1(\Omega)$ (see also [4, 22, 11]).

The situation in the case of the functional F_ε in (1.1) is quite different, since $F_\varepsilon(u_\varepsilon) \neq F_\varepsilon(\tilde{u}_\varepsilon)$. It is therefore necessary to formulate a compactness result for the original sequence (u_ε) . The convergence we introduce only looks at the behaviour of (u_ε) on the matrix $\Omega \cap \varepsilon P$: We say that (u_ε) converges to u in $\Omega \cap \varepsilon P$ if there exists a sequence $(\tilde{u}_\varepsilon) \subset L^1(\Omega)$ such that $\tilde{u}_\varepsilon = u_\varepsilon$ in $\Omega \cap \varepsilon P$, and \tilde{u}_ε converges to u strongly in $L^1(\Omega)$ (see Definition 3.1).

This new convergence is natural since it guarantees the convergence of minimisers and minimum values of F_ε , up to the addition of a forcing term (see Proposition 3.8). It introduces, however, several difficulties. First of all, we can only work with a *sequential* notion of Γ -convergence for F_ε . As a result, the so-called localisation method of Γ -convergence that is usually employed to prove the existence of a Γ -limit of integral type does not apply directly. In particular, the proof of the

fundamental estimate, which is a crucial step to guarantee that the Γ -limit is a measure, requires some care (see Lemma 4.4). Moreover, dealing with a weaker convergence (than the usual $L^1(\Omega)$) implies that there are more converging sequences, and therefore proving an optimal lower bound for $f^{\ell,(\varepsilon_k)}$ and $g^{\ell,(\varepsilon_k)}$ is more subtle.

1.2. The surface energy. We show in Theorem 5.2 that the homogenized surface integrand $g_{\text{hom}}^{\ell,(\varepsilon_k)}$ does not depend on ℓ or on the subsequence (ε_k) and

$$g_{\text{hom}}^{\ell}(\nu) = \hat{g}(\nu) \quad (1.3)$$

for every $\ell \in [0, +\infty]$ and every $\nu \in \mathbb{S}^{n-1}$, where \hat{g} denotes the limit surface density of the Mumford-Shah functional in a periodically perforated domain, $\widehat{F}_{\varepsilon}$, in (1.2). This result is surprising since the functionals F_{ε} in (1.1) exhibit a high-contrast, β_{ε} -dependent behaviour in the surface term, while the limit surface energy density is independent of ℓ and coincides with the case of $\beta_{\varepsilon} = 0$. In other words, the effective cost of introducing a crack in the material is the lowest possible, and coincides with the case where the inclusions are replaced by perforations.

Note that, since $F_{\varepsilon} \geq \widehat{F}_{\varepsilon}$, the bound $g_{\text{hom}}^{\ell} \geq \hat{g}$ is immediate. However, proving the opposite inequality is nontrivial and it requires extending partitions of a perforated domain inside the perforations, without essentially increasing the perimeter of the partition.

1.3. The volume energy. The homogenized volume integrand $f_{\text{hom}}^{\ell,(\varepsilon_k)}$, instead, shows a nontrivial dependence on ℓ . The dependence on the subsequence (ε_k) , however, is only present in the critical regime $\ell \in (0, +\infty)$ and not in the extremal cases $\ell = 0$ and $\ell = +\infty$.

Specifically, in the subcritical regime $\ell = 0$, for every $\xi \in \mathbb{R}^n$

$$f_{\text{hom}}^0(\xi) = \hat{f}(\xi), \quad (1.4)$$

where \hat{f} denotes the limit volume density of $\widehat{F}_{\varepsilon}$ (Theorem 6.2). So, in the subcritical regime $\beta_{\varepsilon} \ll \varepsilon$, F_{ε} and $\widehat{F}_{\varepsilon}$ are asymptotically equivalent, since ‘cutting out’ all the inclusions has an infinitesimal energy-cost of order $\beta_{\varepsilon}/\varepsilon$ (given by the perimeter of the inclusion, which is proportional to $\beta_{\varepsilon}\varepsilon^{n-1}$, for each of the $1/\varepsilon^n$ ε -cells in Ω). In other words, having very weak inclusions is equivalent to having a perforated material.

In the other extremal regime; *i.e.*, $\ell = +\infty$, the homogenized volume integrand is the highest possible, namely $f_{\text{hom}}^{\infty}(\xi) = |\xi|^2$ (Theorem 6.4). In this case it is the upper bound $f_{\text{hom}}^{\infty}(\xi) \leq |\xi|^2$ that is immediate; the difficulty in proving the opposite inequality is due to having to prove that every sequence u_{ε} converging to $u_{\xi} = \xi \cdot x$ in $\Omega \cap \varepsilon P$ satisfies the lower bound

$$\liminf_{\varepsilon \rightarrow 0} F_{\varepsilon}(u_{\varepsilon}) \geq \mathcal{L}^n(\Omega)|\xi|^2. \quad (1.5)$$

Note that (1.5) means that the microscopic cracks at scale ε do not lower the elastic moduli of the material. We prove the lower bound (1.5) by classifying each ε -cell in Ω . Either the measure of the jump set of u_{ε} in the cell is large, in which case we ‘cut out’ the inclusion from the cell as in the subcritical case. Or, alternatively, the measure of the jump set of u_{ε} in the cell is small, in which case the function is essentially smooth, thanks to an ‘Elimination Property’ for the jump set due to De Giorgi, Carriero and Leaci [21] (see also [19]). Due to the relatively high cost of creating a fracture in the supercritical case, we show that it is energetically convenient to have no fracture at all in the majority of the cubes, and this gives (1.5).

In the critical case $\ell \in (0, +\infty)$, we cannot exclude that the volume energy density $f_{\text{hom}}^{\ell,(\varepsilon_k)}$ depends on the subsequence (ε_k) along which we prove Γ -convergence. Each (subsequence dependent) $f_{\text{hom}}^{\ell,(\varepsilon_k)}$, however, satisfies some (subsequence independent) properties. First of all, $f_{\text{hom}}^{\ell,(\varepsilon_k)} = |\xi|^2$ for small $|\xi|$; namely, for small ξ the effective elastic behaviour of the material is the same as in the supercritical case. Moreover, unlike the extremal cases $\ell \in \{0, +\infty\}$, the volume energy density *is not 2-homogeneous* even if the volume density of F_{ε} is. This shows emergence of non-standard constitutive laws in the homogenized limit of high-contrast brittle materials.

1.4. Comparison with previous work. This work has interesting similarities and differences with previous results on the homogenization of free-discontinuity energies.

1.4.1. *Interaction of the volume and surface term in the Γ -limit.* Our result shows that the volume and surface terms of F_ε interact in the limit. This case is different from the analysis in [13], where the authors devise a list of assumptions ensuring that the volume and surface energies do not interact in the homogenization of free-discontinuity functionals. The functionals F_ε in (1.1) are not covered by the analysis in [13] due to their degenerate growth conditions. Such a degeneracy, and the consequent lack of coerciveness, however, is not sufficient to cause the interaction of the terms of the energy, since this was not the case for the functionals \widehat{F}_ε .

1.4.2. *Other high-contrast Mumford-Shah energies.* The general form of high-contrast Mumford-Shah energies is

$$H_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(u) = \int_{\Omega \cap \varepsilon P} |\nabla u|^2 dx + \alpha_\varepsilon \int_{\Omega \cap (\Omega \setminus \varepsilon P)} |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap \varepsilon P) + \beta_\varepsilon \mathcal{H}^{n-1}(S_u \cap (\Omega \setminus \varepsilon P)),$$

where $\alpha_\varepsilon, \beta_\varepsilon \geq 0$, with either α_ε or β_ε being infinitesimal for $\varepsilon \rightarrow 0$. Note that $\widehat{F}_\varepsilon = H_\varepsilon^{0,0}$, $F_\varepsilon = H_\varepsilon^{1, \beta_\varepsilon}$, while the case studied in [5] corresponds to $H_\varepsilon^{\alpha_\varepsilon, 1}$. In [5] the authors proved that the volume energy density of the Γ -limit of $H_\varepsilon^{\alpha_\varepsilon, 1}$ is \hat{f} , regardless of the smallness of α_ε relative to ε . Combining this result with the identification of the Γ -limit of $F_\varepsilon = H_\varepsilon^{1, \beta_\varepsilon}$ proved in the present paper, and of the Γ -limit of $\widehat{F}_\varepsilon = H_\varepsilon^{0,0}$ in [12, 4, 22], we can deduce the expression of the Γ -limit of $H_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}$, when both $\alpha_\varepsilon \rightarrow 0$ and $\beta_\varepsilon \rightarrow 0$. This follows by letting $\varepsilon \rightarrow 0$ in the estimate

$$H_\varepsilon^{0,0}(u) \leq H_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}(u) \leq \min\{H_\varepsilon^{1, \beta_\varepsilon}(u), H_\varepsilon^{\alpha_\varepsilon, 1}(u)\}. \quad (1.6)$$

Indeed, the limit volume energy density in the left- and right-hand sides of the inequality (1.6) is \hat{f} , and the limit surface energy density in both sides of the inequality is \hat{g} , so that the Γ -limit of $H_\varepsilon^{\alpha_\varepsilon, \beta_\varepsilon}$ is the same as that of $H_\varepsilon^{0,0}$; *i.e.*, the same as the Γ -limit of the perforated Mumford-Shah functional \widehat{F}_ε (see also [11]).

1.5. **Structure of the paper.** In Section 2 we introduce the notation used in the paper and we recall some previous results on the homogenization of Mumford-Shah-type energies. Section 3 is devoted to the statement of our main Γ -convergence result, Theorem 3.7, and of its consequences. The proof of Theorem 3.7 is split into the remaining three sections: The abstract Γ -convergence and integral representation result is proved in Section 4, while Sections 5 and 6 are devoted to the characterisation of the surface and volume energy densities, respectively.

2. NOTATION AND PRELIMINARIES

In this section we fix the notation and recall some definitions and results that we are going to use throughout the paper.

2.1. **Notation.** Let $n \geq 2$ and let $\Omega \subset \mathbb{R}^n$ be open bounded and with Lipschitz boundary. We denote with $\mathcal{A}(\Omega)$ the class of all open subsets of Ω , and with $\mathcal{B}(\Omega)$ the σ -algebra of Borel sets in Ω . The n -dimensional Lebesgue measure is denoted by \mathcal{L}^n , and the $(n-1)$ -dimensional Hausdorff measure by \mathcal{H}^{n-1} . For $U, V \in \mathcal{B}(\Omega)$ we set $\mathcal{H}^{n-1} \llcorner U(V) := \mathcal{H}^{n-1}(U \cap V)$. For every $x \in \mathbb{R}^n$ and $r > 0$, $B_r(x)$ will be the open ball with centre x and radius r , with $B_r := B_r(0)$; for $0 < s < r$ we also set $B_{s,r} := B_r \setminus \overline{B}_s$. The boundary of the ball B_1 will be denoted with \mathbb{S}^{n-1} .

For $r > 0$, Q_r denotes the open cube centred at the origin, with side-length r . We write $Q = Q_1$. For $0 < s < r$ we also set $Q_{s,r} := Q_r \setminus \overline{Q}_s$.

We also define the periodic set

$$P := \mathbb{R}^n \setminus \left(\bigcup_{i \in \mathbb{Z}^n} (i + Q_{1/2}) \right). \quad (2.1)$$

The functional setting for our analysis is that of *special functions of bounded variation* in Ω ; *i.e.*,

$$SBV(\Omega) := \{u \in BV(\Omega) : Du = \nabla u \mathcal{L}^n + (u^+ - u^-) \nu_u d\mathcal{H}^{n-1} \llcorner S_u\}.$$

Here S_u denotes the discontinuity set of u , ν_u is the generalised normal to S_u , u^+ and u^- are the traces of u on both sides of S_u . More precisely, we work with the following vector subspace of $SBV(\Omega)$:

$$SBV^2(\Omega) := \{u \in SBV(\Omega) : \nabla u \in L^2(\Omega) \text{ and } \mathcal{H}^{n-1}(S_u) < +\infty\}.$$

We consider also the larger space of *generalised special functions of bounded variations* in Ω ,

$$GSBV(\Omega) := \{u \in L^1(\Omega) : (u \wedge m) \vee (-m) \in SBV(\Omega) \text{ for all } m \in \mathbb{N}\}.$$

By analogy with the case of SBV functions, we write

$$GSBV^2(\Omega) := \{u \in GSBV(\Omega) : \nabla u \in L^2(\Omega) \text{ and } \mathcal{H}^{n-1}(S_u) < +\infty\}.$$

We also consider the space

$$SBV^{\text{pc}}(\Omega) := \{u \in SBV(\Omega) : \nabla u = 0 \text{ } \mathcal{L}^n\text{-a.e.}, \mathcal{H}^{n-1}(S_u) < +\infty\};$$

it is known (see [1, Theorem 4.23]) that every u in $SBV^{\text{pc}}(\Omega) \cap L^\infty(\Omega)$ is piecewise constant in the sense of [1, Definition 4.21], namely there exists a Caccioppoli partition (E_i) of Ω such that u is constant \mathcal{L}^n -a.e. in each set E_i .

Moreover, we set

$$\mathcal{P}(\Omega) := \{u \in SBV^{\text{pc}}(\Omega) : u(x) \in \{0, 1\} \text{ } \mathcal{L}^n\text{-a.e. in } \Omega\}.$$

2.2. Mumford-Shah-type energies. For $\varepsilon > 0$ let $0 \leq \alpha_\varepsilon, \beta_\varepsilon \leq 1$; we define the functional $H_\varepsilon : L^1(\Omega) \rightarrow [0, +\infty]$ as follows

$$H_\varepsilon(u) := \begin{cases} \int_\Omega a_\varepsilon\left(\frac{x}{\varepsilon}\right) |\nabla u|^2 dx + \int_{S_u \cap \Omega} b_\varepsilon\left(\frac{x}{\varepsilon}\right) d\mathcal{H}^{n-1}(x) & \text{if } u \in GSBV^2(\Omega) \\ +\infty & \text{otherwise,} \end{cases} \quad (2.2)$$

where $a_\varepsilon, b_\varepsilon : \mathbb{R}^n \rightarrow \mathbb{R}$ are the Q -periodic functions defined as

$$a_\varepsilon(y) := \begin{cases} 1 & \text{if } y \in Q_{\frac{1}{2}, 1} \\ \alpha_\varepsilon & \text{if } y \in Q_{\frac{1}{2}} \end{cases}, \quad b_\varepsilon(y) := \begin{cases} 1 & \text{if } y \in Q_{\frac{1}{2}, 1} \\ \beta_\varepsilon & \text{if } y \in Q_{\frac{1}{2}} \end{cases},$$

on the periodicity cell Q . The particular choice of the geometry of the inclusions (or of the set P) is not relevant for the subsequent analysis: Instead of $Q_{1/2}$ we could consider any Lipschitz open subset U of Q , with $U \subset \subset Q$. In what follows however we only consider P as in (2.1) for the sake of the exposition.

2.2.1. The extreme cases $\alpha_\varepsilon = \beta_\varepsilon = 1$ and $\alpha_\varepsilon = \beta_\varepsilon = 0$. The case $\alpha_\varepsilon = \beta_\varepsilon = 1$ corresponds to the (ε -independent) Mumford-Shah functional, which we denote with

$$MS(u) := \begin{cases} \int_\Omega |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap \Omega) & \text{if } u \in GSBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega). \end{cases} \quad (2.3)$$

The other extreme case $\alpha_\varepsilon = \beta_\varepsilon = 0$ corresponds to the Mumford-Shah functional on a periodically perforated domain, which we denote with

$$\widehat{F}_\varepsilon(u) := \begin{cases} \int_{\Omega \cap \varepsilon P} |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap \varepsilon P) & \text{if } u|_{\Omega \cap \varepsilon P} \in GSBV^2(\Omega \cap \varepsilon P), \\ +\infty & \text{otherwise in } L^1(\Omega). \end{cases} \quad (2.4)$$

We recall that the Γ -limit of \widehat{F}_ε with respect to the $L^1(\Omega)$ -topology, which has been studied in [22, 12, 4, 11], is the anisotropic free-discontinuity functional $\widehat{F} : L^1(\Omega) \rightarrow [0, +\infty]$ defined as

$$\widehat{F}(u) := \begin{cases} \int_\Omega \widehat{f}(\nabla u) dx + \int_{S_u \cap \Omega} \widehat{g}(\nu_u) d\mathcal{H}^{n-1} & \text{if } u \in GSBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega), \end{cases} \quad (2.5)$$

(see, e.g., [4, Theorem 4]). In (2.5), the volume density $\hat{f}: \mathbb{R}^n \rightarrow [0, +\infty)$ is the quadratic form given by

$$\hat{f}(\xi) := \inf \left\{ \int_{Q \cap P} |\xi + Dw|^2 dx : w \in H_{\text{per}}^1(Q \cap P) \right\}, \quad (2.6)$$

moreover, there exists a constant $c_1 := c_1(n, P) > 0$ such that

$$c_1 |\xi|^2 \leq \hat{f}(\xi) \leq |\xi|^2 \quad \text{for every } \xi \in \mathbb{R}^n. \quad (2.7)$$

The surface density $\hat{g}: \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ is defined as

$$\hat{g}(\nu) := \liminf_{\varepsilon \rightarrow 0^+} \left\{ \mathcal{H}^{n-1}(S_w \cap (Q^\nu \cap \varepsilon P)) : w \in \mathcal{P}(Q^\nu \cap \varepsilon P), \right. \\ \left. w = u_{0,1}^\nu \text{ in a neighbourhood of } \partial Q^\nu \right\}, \quad (2.8)$$

where Q^ν stands for the unit cube centred in 0 with one face orthogonal to ν and

$$u_{0,1}^\nu(x) = \begin{cases} 1 & \text{if } \langle x, \nu \rangle \geq 0 \\ 0 & \text{if } \langle x, \nu \rangle < 0. \end{cases}$$

It can be seen that the function \hat{g} in (2.8) is continuous on \mathbb{S}^{n-1} and satisfies

$$c_2 \leq \hat{g}(\nu) \leq 1 \quad \text{for every } \nu \in \mathbb{S}^{n-1}, \quad (2.9)$$

for some constant $c_2 := c_2(n, P) > 0$. Therefore, gathering (2.7) and (2.9) we obtain the following lower bound for \hat{F}

$$\min\{c_1, c_2\} MS \leq \hat{F}. \quad (2.10)$$

Remark 2.1. By changing variables, we can equivalently write

$$\hat{g}(\nu) = \lim_{t \rightarrow +\infty} \frac{1}{t^{n-1}} \min \{ \mathcal{H}^{n-1}(S_w \cap P \cap tQ^\nu) : u \in \mathcal{P}(tQ^\nu \cap P), \\ w = u_{0,1}^\nu \text{ in a neighbourhood of } \partial(tQ^\nu) \}.$$

Remark 2.2. In [11] Braides and Solci proved, among other things, that \hat{F}_ε Γ -converges to \hat{F} also with respect to the following convergence, that is weaker than convergence in $L^1(\Omega)$: Given $u_\varepsilon, u \in L^1(\Omega)$ we say that u_ε converges to u if

$$u_\varepsilon \chi_{\Omega \cap \varepsilon P} \rightharpoonup C_P u \quad \text{weakly in } L^1(\Omega),$$

where $C_P := \mathcal{L}^n(Q \cap P) = \mathcal{L}^n(Q_{\frac{1}{2}, 1}) = 1 - \left(\frac{1}{2}\right)^n$.

2.2.2. High-contrast Mumford-Shah functionals. This is the case of the functional H_ε in (2.2) where either $\alpha_\varepsilon \rightarrow 0$ or $\beta_\varepsilon \rightarrow 0$, as $\varepsilon \rightarrow 0$. For this choice of α_ε and β_ε , H_ε represents the energy associated to a high-contrast composite material, the latter being characterised by two constituents (the matrix and the inclusions) with significantly different mechanical properties.

The case where both $\alpha_\varepsilon \rightarrow 0$ and $\beta_\varepsilon \rightarrow 0$, as $\varepsilon \rightarrow 0$ was considered by Braides and Solci in [11]. In this case, independently of the vanishing rate of α_ε and β_ε , the functionals H_ε Γ -converge to \hat{F} with respect to the convergence introduced in Remark 2.2. This means that, no matter how small the weights α_ε and β_ε are, as long as they are *both* infinitesimal, the effective behaviour of the functional H_ε is the same as for \hat{F}_ε in (2.4), namely it is the same as for $\alpha_\varepsilon = \beta_\varepsilon = 0$.

The case $\alpha_\varepsilon \rightarrow 0$ and $\beta_\varepsilon = 1$ was treated in [5], where the authors focus on the critical case $\alpha_\varepsilon = \varepsilon$ and study the Γ -convergence of H_ε with respect to the strong convergence in $L^1(\Omega)$. They prove that the Γ -limit of H_ε , which exists up to a subsequence, is of free-discontinuity type. More precisely, the limit volume density is the function \hat{f} in (2.6) corresponding to the case $\alpha_\varepsilon = 0$. The surface energy density, instead, depends non trivially on the jump opening $[u]$, even if the surface energy density in H_ε is identically equal to 1. In particular, the case considered in [5] is an example where volume and surface energies interact in the Γ -limit thus giving rise to a homogenized effective energy of completely different nature with respect to the microscopic ones.

Here we consider the complementary case to [5], namely in H_ε we choose $\alpha_\varepsilon = 1$ and let $\beta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. This choice corresponds to having inclusions which are much more brittle than the matrix, but whose elastic behaviour is the same as that of the matrix.

3. SETTING OF THE PROBLEM AND STATEMENT OF THE MAIN RESULT

In this section we state our main result, that is a Γ -convergence theorem for the functionals $F_\varepsilon: L^1(\Omega) \rightarrow [0, +\infty]$ defined as

$$F_\varepsilon(u) := \begin{cases} \int_{\Omega} |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap (\Omega \cap \varepsilon P)) + \beta_\varepsilon \mathcal{H}^{n-1}(S_u \cap (\Omega \setminus \varepsilon P)) & \text{if } u \in GSBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega), \end{cases} \quad (3.1)$$

where $\beta_\varepsilon \searrow 0$ as ε tends to zero. We analyse the asymptotic behaviour of F_ε in three possible scaling-regimes; *i.e.*, $\beta_\varepsilon \ll \varepsilon$ (subcritical regime), $\beta_\varepsilon \sim \varepsilon$ (critical regime), and $\beta_\varepsilon \gg \varepsilon$ (supercritical regime). We note that trivially

$$\widehat{F}_\varepsilon \leq F_\varepsilon \leq MS, \quad (3.2)$$

where \widehat{F}_ε and MS are defined in (2.4) and (2.3), respectively.

For the Γ -convergence result to be meaningful we first need to identify a notion of convergence on the space $L^1(\Omega)$ for which the equicoerciveness of the functionals F_ε is guaranteed.

3.1. Choice of the convergence. Let $(u_\varepsilon) \subset L^1(\Omega)$ be a sequence satisfying

$$\sup_\varepsilon \|u_\varepsilon\|_{L^\infty(\Omega \cap \varepsilon P)} < +\infty \quad \text{and} \quad \sup_\varepsilon F_\varepsilon(u_\varepsilon) < +\infty. \quad (3.3)$$

Since β_ε is a vanishing sequence, the uniform bound on $F_\varepsilon(u_\varepsilon)$ gives no control on the surface term $\mathcal{H}^{n-1}(S_{u_\varepsilon} \cap (\Omega \setminus \varepsilon P))$ so that, in particular, the sequence (u_ε) is not uniformly bounded in $BV(\Omega)$. However, by the extension result [12, Theorem 1.1] we can find a new sequence $\tilde{u}_\varepsilon \subset L^1(\Omega)$ satisfying

$$\begin{cases} \tilde{u}_\varepsilon = u_\varepsilon & \text{in } \Omega \cap \varepsilon P \\ \sup_\varepsilon \|\tilde{u}_\varepsilon\|_{L^\infty(\Omega)} < +\infty \\ MS(\tilde{u}_\varepsilon) \leq CF_\varepsilon(u_\varepsilon), \end{cases} \quad (3.4)$$

for some $C > 0$. Therefore, by combining (3.3)-(3.4) with the Ambrosio Compactness Theorem (see *e.g.*, [1, Theorem 4.8]) we deduce that there exist $u \in SBV^2(\Omega)$ and a subsequence of (\tilde{u}_ε) , (not relabelled) such that $\tilde{u}_\varepsilon \rightarrow u$ in $L^1(\Omega)$. As a consequence we get

$$\int_{\Omega \cap \varepsilon P} |u_\varepsilon - u| dx = \int_{\Omega \cap \varepsilon P} |\tilde{u}_\varepsilon - u| dx \leq \int_{\Omega} |\tilde{u}_\varepsilon - u| dx \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

The above observation motivates the following definition.

Definition 3.1 (Convergence). Let (u_ε) be a sequence in $L^1(\Omega)$. We say that (u_ε) converges in $\Omega \cap \varepsilon P$ to a function $u \in L^1(\Omega)$, and we write $u_\varepsilon \rightarrow u$, if there exists a sequence $(\tilde{u}_\varepsilon) \subset L^1(\Omega)$ such that $\tilde{u}_\varepsilon = u_\varepsilon$ in $\Omega \cap \varepsilon P$, and \tilde{u}_ε converges to u strongly in $L^1(\Omega)$.

Remark 3.2. We observe that Definition 3.1 is well-posed. Indeed let $(\tilde{u}_{1,\varepsilon}), (\tilde{u}_{2,\varepsilon}) \subset L^1(\Omega)$ be such that $\tilde{u}_{1,\varepsilon} = \tilde{u}_{2,\varepsilon} = u_\varepsilon$ in $\Omega \cap \varepsilon P$ and $\tilde{u}_{1,\varepsilon} \rightarrow u_1$ and $\tilde{u}_{2,\varepsilon} \rightarrow u_2$ strongly in $L^1(\Omega)$. Then

$$0 = \lim_{\varepsilon \rightarrow 0} \int_{\Omega \cap \varepsilon P} |\tilde{u}_{1,\varepsilon} - \tilde{u}_{2,\varepsilon}| dx = \lim_{\varepsilon \rightarrow 0} \int_{\Omega} |\tilde{u}_{1,\varepsilon} - \tilde{u}_{2,\varepsilon}| \chi_{\Omega \cap \varepsilon P} dx = C_P \int_{\Omega} |u_1 - u_2| dx,$$

hence, since $C_P = \mathcal{L}^n(Q_{\frac{1}{2},1}) > 0$ we necessarily have $u_1 = u_2$.

Remark 3.3. Let $(u_\varepsilon) \subset L^1(\Omega)$ be such that $u_\varepsilon \rightarrow u$ in the sense of Definition 3.1, for some $u \in L^1(\Omega)$. It is immediate to see that

- i. $\lim_\varepsilon \|u_\varepsilon - u\|_{L^1(\Omega \cap \varepsilon P)} = 0$;
- ii. the sequence $(u_\varepsilon \chi_{\Omega \cap \varepsilon P}) \subset L^1(\Omega)$ converges to $C_P u$ weakly in $L^1(\Omega)$;

- iii. if (u_ε^m) denotes the sequence of truncated functions of u_ε at level $m \in \mathbb{N}$, then $u_\varepsilon^m \rightarrow u^m$ in the sense of Definition 3.1, where u^m denotes the truncated function of u at level m .

In view of the above considerations, in what follows we study the Γ -convergence of the functionals F_ε with respect to the convergence as in Definition 3.1. To this end we give the following definition.

Definition 3.4 (Sequential Γ -convergence). Let $F_\varepsilon, F: L^1(\Omega) \rightarrow [0, +\infty]$; we say that the functionals F_ε Γ -converge to F with respect to the convergence as in Definition 3.1 if for every $u \in L^1(\Omega)$ the two following conditions are satisfied:

- (i) (Ansatz-free lower bound) For every $(u_\varepsilon) \subset L^1(\Omega)$ with $u_\varepsilon \rightarrow u$ in the sense of Definition 3.1 we have

$$F(u) \leq \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon);$$

- (ii) (Existence of a recovery sequence) There exists $(\bar{u}_\varepsilon) \subset L^1(\Omega)$ with $\bar{u}_\varepsilon \rightarrow u$ in the sense of Definition 3.1 such that

$$F(u) \geq \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(\bar{u}_\varepsilon).$$

Remark 3.5. It is easy to check that F is lower semicontinuous with respect to the convergence as in Definition 3.1 and thus, in particular, with respect to the strong $L^1(\Omega)$ convergence.

For every $u \in L^1(\Omega)$ we set

$$\Gamma\text{-}\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u) := \inf \left\{ \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) : u_\varepsilon \rightarrow u \right\} \quad (3.5)$$

and

$$\Gamma\text{-}\limsup_{\varepsilon \rightarrow 0} F_\varepsilon(u) := \inf \left\{ \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) : u_\varepsilon \rightarrow u \right\}, \quad (3.6)$$

where the convergence $u_\varepsilon \rightarrow u$ is understood in the sense of Definition 3.1. We also introduce the more compact notation $F'(u) := \Gamma\text{-}\liminf_{\varepsilon} F_\varepsilon(u)$ and $F''(u) := \Gamma\text{-}\limsup_{\varepsilon} F_\varepsilon(u)$. Then, it is immediate to see that Definition 3.4 is equivalent to $F' = F'' = F$ in $L^1(\Omega)$. Further, it can be easily shown that the infima in (3.5) and (3.6) are actually minima.

3.2. Compactness and domain of the Γ -limit. If not otherwise specified, in all that follows the Γ -convergence will be always understood in the sense of Definition 3.4.

The following proposition shows that the domain of the Γ -limit of F_ε (if it exists) is the space $GSBV^2(\Omega)$.

Proposition 3.6 (Domain of the Γ -limit). *Let $(u_\varepsilon) \subset L^1(\Omega)$ be such that $\sup_\varepsilon F_\varepsilon(u_\varepsilon) < +\infty$. Assume moreover that u_ε converges to u in the sense of Definition 3.1, then $u \in GSBV^2(\Omega)$.*

Proof. By Definition 3.1 there exists $(\tilde{u}_\varepsilon) \subset L^1(\Omega)$ such that $\tilde{u}_\varepsilon = u_\varepsilon$ in $\Omega \cap \varepsilon P$, and \tilde{u}_ε converges to u strongly in $L^1(\Omega)$. Let $m \in \mathbb{N}$ and let u_ε^m be the truncated function of u_ε at level m . Appealing to [12, Theorem 1.1] we deduce the existence of a sequence $(\tilde{u}_\varepsilon^m)$ such that $\tilde{u}_\varepsilon^m = u_\varepsilon^m$ in $\Omega \cap \varepsilon P$ and

$$MS(\tilde{u}_\varepsilon^m) \leq CF_\varepsilon(u_\varepsilon^m) \leq CF_\varepsilon(u_\varepsilon),$$

for some $C > 0$, where to establish the last inequality we have used the fact that F_ε decreases by truncations. Therefore, in view of the bound on the energy, invoking [1, Theorem 4.8] yields the existence of a function $v \in SBV^2(\Omega)$ and a subsequence of $(\tilde{u}_\varepsilon^m)$ (not relabelled) such that $\tilde{u}_\varepsilon^m \rightarrow v$ strongly in $L^1(\Omega)$. For every fixed $m \in \mathbb{N}$ we have

$$0 = \lim_{\varepsilon \rightarrow 0} \int_{\Omega \cap \varepsilon P} |\tilde{u}_\varepsilon^m - u_\varepsilon^m| dx = \lim_{\varepsilon \rightarrow 0} \int_{\Omega} |\tilde{u}_\varepsilon^m - (\tilde{u}_\varepsilon)^m| \chi_{\Omega \cap \varepsilon P} dx = C_P \int_{\Omega} |v - u^m| dx,$$

therefore $v = u^m$ and $u^m \in SBV^2(\Omega)$ for every $m \in \mathbb{N}$. This implies that $u \in GSBV^2(\Omega)$. \square

3.3. Γ -convergence. Our main result is the following theorem.

Theorem 3.7. *Let $F_\varepsilon : L^1(\Omega) \rightarrow (0, +\infty]$ be the functionals defined in (3.1) and let $\ell := \lim_{\varepsilon \rightarrow 0} \frac{\beta_\varepsilon}{\varepsilon}$. Then for every sequence of positive numbers decreasing to zero, there exists a subsequence (ε_k) such that the functionals F_{ε_k} Γ -converge to $F_{\text{hom}}^\ell : L^1(\Omega) \rightarrow [0, +\infty)$ defined by*

$$F_{\text{hom}}^\ell(u) := \begin{cases} \int_{\Omega} f_{\text{hom}}^\ell(\nabla u) dx + \int_{S_u \cap \Omega} g_{\text{hom}}^\ell(\nu_u) d\mathcal{H}^{n-1} & \text{if } u \in GSBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega). \end{cases}$$

Moreover, the function g_{hom}^ℓ is independent of ℓ and it coincides with \hat{g} as in (2.8), while the function f_{hom}^ℓ satisfies, for every $\xi \in \mathbb{R}^n$ and every $\ell \in [0, +\infty]$, the bounds

$$\hat{f}(\xi) \leq f_{\text{hom}}^\ell(\xi) \leq |\xi|^2,$$

where \hat{f} is the quadratic form in (2.6). Furthermore, in the extreme regimes $\ell = 0$ and $\ell = +\infty$ we have, respectively, $f_{\text{hom}}^0(\xi) = \hat{f}(\xi)$ and $f_{\text{hom}}^\infty(\xi) = |\xi|^2$ for every $\xi \in \mathbb{R}^n$. Therefore for $\ell = 0$ and $\ell = +\infty$ the whole sequence (F_ε) Γ -converges.

We divide the proof of Theorem 3.7 into two main steps carried out, respectively, in Section 4 and in Sections 5-6. Specifically, the first step deals with the existence of a subsequence of F_ε which Γ -converges to a homogeneous free-discontinuity functional of the form

$$\int_{\Omega} f^\ell(\nabla u) dx + \int_{S_u} g^\ell(u^+ - u^-, \nu_u) d\mathcal{H}^{n-1}.$$

Then in the second step we identify the integrands g^ℓ and f^ℓ in the three different scaling regimes.

3.4. Convergence of minimisation problems. Thanks to the Γ -convergence result Theorem 3.7, we are able to establish a convergence result for the minimisation problems associated with a suitable perturbation of the functionals F_ε .

Let $g \in L^\infty(\Omega)$, and let $G_\varepsilon : L^1(\Omega) \rightarrow [0, +\infty)$ be the functionals defined as

$$G_\varepsilon(u) := \int_{\Omega \cap \varepsilon P} |u(x) - g(x)| dx.$$

Let moreover $G : L^1(\Omega) \rightarrow [0, +\infty)$ be given by

$$G(u) := C_P \int_{\Omega} |u(x) - g(x)| dx,$$

where $C_P := \mathcal{L}^n(Q_{\frac{1}{2}, 1})$.

Let $\varepsilon > 0$ be fixed; we start by observing that $F_\varepsilon + G_\varepsilon$ is lower-semicontinuous with respect to the strong $L^1(\Omega)$ -convergence. Moreover, since F_ε decreases by truncations, we can readily deduce that a minimising sequence (u_j) for $F_\varepsilon + G_\varepsilon$ satisfies the uniform bound $\|u_j\|_{L^\infty(\Omega)} \leq \|g\|_{L^\infty(\Omega)}$. The latter allows us to invoke the compactness result [1, Theorem 4.7] and thus to use the direct methods to deduce the existence of a minimiser $u_\varepsilon \in SBV^2(\Omega)$ for $F_\varepsilon + G_\varepsilon$.

The following proposition establishes a convergence result for minimisers and minimum values of $F_\varepsilon + G_\varepsilon$.

Proposition 3.8. *Let (ε_k) and F_{hom}^ℓ be, respectively, the subsequence and the functional whose existence is established in Theorem 3.7. Let $k \in \mathbb{N}$ be fixed and let $u_k \in L^1(\Omega)$ be a solution to*

$$m_k := \min \{F_{\varepsilon_k}(u) + G_{\varepsilon_k}(u) : u \in L^1(\Omega)\}.$$

Then, up to subsequences (not relabelled), u_k converges in the sense of Definition 3.1 to a function $\bar{u} \in SBV^2(\Omega)$ which solves

$$m := \min \{F_{\text{hom}}^\ell(u) + G(u) : u \in L^1(\Omega)\}.$$

Moreover we have $m_k \rightarrow m$, as $k \rightarrow +\infty$.

Proof. Let $(u_k) \subset L^1(\Omega)$ be a sequence of minimisers for $F_{\varepsilon_k} + G_{\varepsilon_k}$; i.e.,

$$m_k = \min \{F_{\varepsilon_k}(u_k) + G_{\varepsilon_k}(u_k) : u \in L^1(\Omega)\},$$

hence in particular $\sup_{k \in \mathbb{N}} F_{\varepsilon_k}(u_k) < +\infty$. A truncation argument also yields $\|u_k\|_{L^\infty(\Omega)} \leq \|g\|_{L^\infty(\Omega)}$, so that we can readily deduce the existence of a subsequence of (u_k) (not relabelled) and a function $\bar{u} \in SBV^2(\Omega)$ such that $u_k \rightarrow \bar{u}$ in the sense of Definition 3.1. Thus, in particular

$$\lim_{k \rightarrow +\infty} \int_{\Omega \cap \varepsilon_k P} |u_k - g| dx = C_P \int_{\Omega} |\bar{u} - g| dx = G(\bar{u}),$$

therefore in view of Theorem 3.7 we get

$$F_{\text{hom}}^\ell(\bar{u}) + G(\bar{u}) \leq \liminf_{k \rightarrow +\infty} (F_{\varepsilon_k}(u_k) + G_{\varepsilon_k}(u_k)) = \liminf_{k \rightarrow +\infty} m_k. \quad (3.7)$$

Now let $w \in SBV^2(\Omega)$ be arbitrary; again appealing to Theorem 3.7 we can find $(w_k) \subset L^1(\Omega)$ such that $w_k \rightarrow w$ in the sense of Definition 3.1 and

$$\lim_{k \rightarrow +\infty} F_{\varepsilon_k}(w_k) = F_{\text{hom}}^\ell(w).$$

Since

$$\lim_{k \rightarrow +\infty} \int_{\Omega \cap \varepsilon P} |w_k - g| dx = G(w),$$

we immediately deduce

$$\limsup_{k \rightarrow +\infty} m_k \leq \limsup_{k \rightarrow +\infty} (F_{\varepsilon_k}(w_k) + G_{\varepsilon_k}(w_k)) = F_{\text{hom}}^\ell(w) + G(w). \quad (3.8)$$

Thus, putting together (3.7) and (3.8) we obtain

$$\begin{aligned} F_{\text{hom}}^\ell(\bar{u}) + G(\bar{u}) &\leq \liminf_{k \rightarrow +\infty} m_k \leq \limsup_{k \rightarrow +\infty} m_k \\ &\leq F_{\text{hom}}^\ell(w) + G(w), \end{aligned}$$

hence by the arbitrariness of w we deduce that \bar{u} is a minimiser for $F_{\text{hom}}^\ell + G$. Finally, taking $w = \bar{u}$ also implies $m_k \rightarrow m$. Since moreover this limit does not depend on the subsequence, the convergence holds true for the whole (m_k) . \square

4. Γ -CONVERGENCE AND INTEGRAL REPRESENTATION

In this section we prove the existence of a Γ -convergent subsequence of F_ε to a homogeneous free-discontinuity functional of the form

$$\int_{\Omega} f^\ell(\nabla u) dx + \int_{S_u} g^\ell(u^+ - u^-, \nu_u) d\mathcal{H}^{n-1}.$$

To do so, we follow the so-called localisation method of Γ -convergence [18], with the caveat that, since we prove a sequential Γ -convergence result, we cannot apply the abstract general theory directly.

As a first step, we localise the functionals, and define $F_\varepsilon : L^1(\Omega) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ as

$$F_\varepsilon(u, U) := \begin{cases} \int_U |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap (U \cap \varepsilon P)) + \beta_\varepsilon \mathcal{H}^{n-1}(S_u \cap (U \setminus \varepsilon P)) & \text{if } u \in GSBV^2(U), \\ +\infty & \text{otherwise in } L^1(\Omega). \end{cases} \quad (4.1)$$

Remark 4.1. For every fixed $\varepsilon > 0$ the functional $F_\varepsilon : L^1(\Omega) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ satisfies the following properties, for every $u \in L^1(\Omega)$, and $U \in \mathcal{A}(\Omega)$:

- (1) $F_\varepsilon(u, \cdot)$ is increasing: $F_\varepsilon(u, V) \leq F_\varepsilon(u, U)$ for every $V \in \mathcal{A}(\Omega)$, $V \subset U$;
- (2) $F_\varepsilon(u, \cdot)$ is super-additive: for $U, V \in \mathcal{A}(\Omega)$, $U \cap V = \emptyset$, then

$$F_\varepsilon(u, U \cup V) \geq F_\varepsilon(u, U) + F_\varepsilon(u, V);$$

- (3) F_ε is local: $F_\varepsilon(u, U) = F_\varepsilon(v, U)$ for every $v \in L^1(\Omega)$ such that $u = v$ \mathcal{L}^n -a.e. in U ;
- (4) F_ε is invariant by translations in u : $F_\varepsilon(u + s, U) = F_\varepsilon(u, U)$ for every $s \in \mathbb{R}$;

- (5) $F_\varepsilon(\cdot, U)$ is decreasing by truncations: $F_\varepsilon(u^m, U) \leq F_\varepsilon(u, U)$ for every $m > 0$, where $u^m := (u \wedge m) \vee (-m)$.

Let (ε_k) be a positive sequence of real numbers decreasing to 0. We define the localised versions $F', F'' : L^1(\Omega) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ of (3.5) and (3.6) as

$$F'(\cdot, U) := \Gamma\text{-}\liminf_{k \rightarrow +\infty} F_{\varepsilon_k}(\cdot, U), \quad F''(\cdot, U) := \Gamma\text{-}\limsup_{k \rightarrow +\infty} F_{\varepsilon_k}(\cdot, U), \quad (4.2)$$

for every $U \in \mathcal{A}(\Omega)$. We notice that in our case the functionals F' and F'' will depend on ℓ , however to simplify the notation at this stage we prefer to omit this dependence.

It is easy to prove that F' and F'' are both lower semicontinuous with respect to the convergence in Definition 3.1. In view of Remark 4.1, they are easily seen to be increasing and local; further, F' is superadditive. Moreover, by combining (5) of Remark 4.1 and iii. of Remark 3.3 it can be immediately checked that both F' and F'' decrease by truncation.

In general $F'(u, \cdot)$ and $F''(u, \cdot)$ are not inner regular. Therefore we also consider their inner regular envelopes, that is the functionals $F'_-, F''_- : L^1(\Omega) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ defined as

$$F'_-(u, U) := \sup \{F'(u, V) : V \subset\subset U, V \in \mathcal{A}(\Omega)\}.$$

and

$$F''_-(u, U) := \sup \{F''(u, V) : V \subset\subset U, V \in \mathcal{A}(\Omega)\}.$$

Remark 4.2. The functionals F'_- and F''_- are both increasing, lower semicontinuous [18, Remark 15.10], and local [18, Remark 15.25], and decrease by truncations. Moreover, F'_- is superadditive [18, Remark 15.10].

The next proposition is the analogue of the compactness result [18, Theorem 16.9], when the sequential notion of Γ -convergence in Definition 3.4 is taken into account.

Proposition 4.3. *Let $F_\varepsilon : L^1(\Omega) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ be the functionals defined in (4.1). Then for every sequence of positive numbers decreasing to zero, there exists a subsequence (ε_k) such that the corresponding functionals F' and F'' in (4.2) satisfy $F'_- = F''_-$.*

Proof. Let $\mathcal{R}(\Omega)$ be the class of all finite unions of open rectangles of Ω with rational vertices. Let $R \in \mathcal{R}(\Omega)$ and $u \in L^1(R)$ be fixed. Since F' in (4.2) is actually attained along a sequence we can find $(u_\varepsilon) \subset L^1(R)$ converging to u in the sense of Definition 3.1 such that

$$\begin{aligned} F''(u, R) \geq F'(u, R) &= \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon, R) = \lim_{k \rightarrow +\infty} F_{\varepsilon_k}(u_{\varepsilon_k}, R) \\ &= \limsup_{k \rightarrow +\infty} F_{\varepsilon_k}(u_{\varepsilon_k}, R) \geq F''(u, R), \end{aligned}$$

where the subsequence ε_k depends on R . Since $\mathcal{R}(\Omega)$ is countable, by a diagonal argument, we can find a subsequence $\tilde{\varepsilon}_k$ such that $F'(u, R) = F''(u, R)$ for every $u \in L^1(\Omega)$ and for every $R \in \mathcal{R}(\Omega)$. Since $\mathcal{R}(\Omega)$ is dense in $\mathcal{A}(\Omega)$, we have

$$\begin{aligned} F'_-(u, U) &= \sup \{F'(u, V) : V \in \mathcal{A}(\Omega), V \subset\subset U\} = \sup \{F'(u, R) : R \in \mathcal{R}(\Omega), R \subset\subset U\} \\ &= \sup \{F''(u, R) : R \in \mathcal{R}(\Omega), R \subset\subset U\} = \sup \{F'(u, V) : V \in \mathcal{A}(\Omega), V \subset\subset U\} \\ &= F''_-(u, U), \end{aligned}$$

which concludes the proof. \square

We now reintroduce the ℓ -dependence in our notation and define the functional

$$F_{\text{hom}}^\ell := F'_- = F''_-. \quad (4.3)$$

We observe that by monotonicity we always have

$$F''_- = F'_- \leq F' \leq F''. \quad (4.4)$$

Therefore, if we show that F'' is inner regular; *i.e.*, $F'' = F''_-$, by combining (4.3) and (4.4) we get $F' = F'' = F_{\text{hom}}^\ell$, and therefore the Γ -convergence of the subsequence (F_{ε_k}) to the functional F_{hom}^ℓ .

A first step towards the proof of the inner-regularity of F'' consists in proving that the functionals F_ε satisfy a fundamental estimate, uniformly in ε . The fundamental estimate we prove is non-standard. Indeed, we need an error term that is infinitesimal for the convergence in Definition 3.1, and hence that only weights the functions outside the weak inclusion. We note that an analogous estimate was also established by Braides and Garroni in [10, Proposition 3.3].

Lemma 4.4. *(Fundamental Estimate in perforated domains). For every $\eta > 0$, and for every $U', U'', V \in \mathcal{A}(\Omega)$, with $U' \subset\subset U''$, there exists a constant $M(\eta) > 0$ satisfying the following property: for every $\varepsilon > 0$, for every $u \in L^1(\Omega)$ with $u \in SBV^2(U'')$, and for every $v \in L^1(\Omega)$ with $v \in SBV^2(V)$, there exists a function $\varphi \in C_0^\infty(\Omega)$ with $\varphi = 1$ in a neighbourhood of U' , $\text{supp } \varphi \subset U''$ and $0 \leq \varphi \leq 1$ such that*

$$F_\varepsilon(\varphi u + (1 - \varphi)v, U' \cup V) \leq (1 + \eta) (F_\varepsilon(u, U'') + F_\varepsilon(v, V)) + M(\eta) \|u - v\|_{L^2(S \cap \varepsilon P)} \quad (4.5)$$

with $S := (U'' \setminus U') \cap V$.

Proof. Let $0 < \eta < 1$, U', U'' , and $V \in \mathcal{A}(\Omega)$ be as in the statement. Let $\delta > 0$ be small enough so that $Q_{\frac{1}{2}+\delta} \subset\subset Q$, and let $\psi \in C_0^\infty(Q)$ be a cut-off function between $Q_{\frac{1}{2}}$ and $Q_{\frac{1}{2}+\delta}$, namely a function such that $0 \leq \psi \leq 1$, $\psi \equiv 1$ on $Q_{\frac{1}{2}}$, and $\text{supp } \psi \subset Q_{\frac{1}{2}+\delta}$.

For $\varepsilon > 0$ and $i \in \mathbb{Z}^n$, we define the operator $R_i^\varepsilon : W_{\text{loc}}^{1,\infty}(\mathbb{R}^n) \rightarrow W_{\text{loc}}^{1,\infty}(\mathbb{R}^n)$ as

$$R_i^\varepsilon(\phi)(x) := \left(1 - \psi\left(\frac{x}{\varepsilon} - i\right)\right) \phi(x) + \psi\left(\frac{x}{\varepsilon} - i\right) \int_{\varepsilon i + \varepsilon Q_{\frac{1}{2}+\delta}} \phi(y) \, dy.$$

It is easy to see that $R_i^\varepsilon(\phi)(x) = \phi(x)$ if $x \notin \varepsilon i + \varepsilon Q_{\frac{1}{2}+\delta}$, and $R_i^\varepsilon(w)$ is instead constant in $\varepsilon i + \varepsilon Q_{\frac{1}{2}}$. Finally, we define the operator $\mathcal{R}^\varepsilon : W_{\text{loc}}^{1,\infty}(\mathbb{R}^n) \rightarrow W_{\text{loc}}^{1,\infty}(\mathbb{R}^n)$ as

$$\mathcal{R}^\varepsilon(\phi)(x) := \begin{cases} R_i^\varepsilon(\phi)(x) & \text{if } x \in \varepsilon i + \varepsilon Q_{\frac{1}{2}+\delta}, \, i \in \mathbb{Z}^n, \\ \phi(x) & \text{otherwise.} \end{cases}$$

Let $W', W'' \in \mathcal{A}(\Omega)$ be such that $U' \subset\subset W' \subset\subset W'' \subset\subset U''$. Let ϕ be a cut-off function between the sets W' and W'' and set $\varphi := \mathcal{R}^\varepsilon(\phi)$; then φ is a cut-off between the sets U' and U'' , provided δ is small enough. Moreover, by construction, $D\varphi = 0$ on $\Omega \setminus \varepsilon P$ and $\|D\varphi\|_{L^\infty(\Omega)} \leq C \|D\phi\|_{L^\infty(\Omega)}$, with a uniform constant independent of ε (see e.g., [10, Remark 2.7]).

Let u and v be as in the statement and let $w := \varphi u + (1 - \varphi)v$; clearly, $w \in SBV^2(U' \cup V)$. Then

$$F_\varepsilon(w, U' \cup V) = F_\varepsilon(u, U') + F_\varepsilon^*(v, V \setminus U'') + F_\varepsilon^*(w, S), \quad (4.6)$$

where, for fixed $w \in L^1(\Omega)$, $F_\varepsilon^*(w, \cdot)$ denotes the measure that extends $F_\varepsilon(w, \cdot)$ to the σ -algebra $\mathcal{B}(\Omega)$ of Borel subsets of Ω , and is defined as

$$F_\varepsilon^*(w, B) := \inf\{F_\varepsilon(w, U) : U \in \mathcal{A}(\Omega), B \subset U\},$$

for every $B \in \mathcal{B}(\Omega)$. Now we estimate the last term in the right-hand side of (4.6). For any fixed $\eta \in (0, 1)$ by convexity and by the definition of φ we have

$$\begin{aligned}
F_\varepsilon^*(w, S) &\leq \int_{S \cap \varepsilon P} \left| (1-\eta) \frac{\varphi \nabla u + (1-\varphi) \nabla v}{1-\eta} + \eta \frac{D\varphi(u-v)}{\eta} \right|^2 dx \\
&\quad + \int_{S \setminus \varepsilon P} \left| (1-\eta) \frac{\varphi \nabla u + (1-\varphi) \nabla v}{1-\eta} \right|^2 dx \\
&\quad + \mathcal{H}^{n-1}(S_u \cap (S \cap \varepsilon P)) + \mathcal{H}^{n-1}(S_v \cap (S \cap \varepsilon P)) \\
&\quad + \beta_\varepsilon \mathcal{H}^{n-1}(S_u \cap (S \setminus \varepsilon P)) + \beta_\varepsilon \mathcal{H}^{n-1}(S_v \cap (S \setminus \varepsilon P)) \\
&\leq \frac{1}{1-\eta} \left(\int_S |\nabla u|^2 dx + \int_S |\nabla v|^2 dx \right) + \frac{1}{\eta} \int_{S \cap \varepsilon P} |D\varphi|^2 |u-v|^2 dx \\
&\quad + \mathcal{H}^{n-1}(S_u \cap (S \cap \varepsilon P)) + \mathcal{H}^{n-1}(S_v \cap (S \cap \varepsilon P)) \\
&\quad + \beta_\varepsilon \mathcal{H}^{n-1}(S_u \cap (S \setminus \varepsilon P)) + \beta_\varepsilon \mathcal{H}^{n-1}(S_v \cap (S \setminus \varepsilon P)) \\
&\leq \frac{1}{1-\eta} (F_\varepsilon^*(u, S) + (F_\varepsilon^*(v, S))) + \frac{1}{\eta} \|D\varphi\|_{L^\infty(\Omega)}^2 \int_{S \cap \varepsilon P} |u-v|^2 dx \\
&\leq \frac{1}{1-\eta} (F_\varepsilon^*(u, S) + (F_\varepsilon^*(v, S))) + \frac{C}{\eta} \int_{S \cap \varepsilon P} |u-v|^2 dx.
\end{aligned}$$

This, together with (4.6), concludes the proof. \square

Theorem 4.5 (Γ -convergence and properties of the Γ -limit). *Let (ε_k) be the sequence for which $F'_- = F''_-$. Then F_{hom}^ℓ defined in (4.3) satisfies the following properties:*

- i) (locality and lower semicontinuity) *for every $U \in \mathcal{A}(\Omega)$, the functional $F_{\text{hom}}^\ell(\cdot, U)$ is local and lower semicontinuous with respect to the strong $L^1(\Omega)$ -topology;*
- ii) (measure property) *for every $u \in GSBV^2(\Omega)$, the set function $F_{\text{hom}}^\ell(u, \cdot)$ is the restriction to $\mathcal{A}(\Omega)$ of a Radon measure on Ω ;*
- iii) (Γ -convergence) *for every $U \in \mathcal{A}(\Omega)$*

$$F_{\text{hom}}^\ell(\cdot, U) = F'(\cdot, U) = F''(\cdot, U) \quad \text{on } GSBV^2(\Omega);$$

- iv) (translational invariance in u) *for every $u \in L^1(\Omega)$ and $U \in \mathcal{A}(\Omega)$*

$$F_{\text{hom}}^\ell(u + s, U) = F_{\text{hom}}^\ell(u, U) \quad \text{for every } s \in \mathbb{R};$$

- v) (translational invariance in x) *for every $u \in L^1(\Omega)$ and $U \in \mathcal{A}(\Omega)$*

$$F_{\text{hom}}^\ell(u(\cdot - y), U + y) = F_{\text{hom}}^\ell(u, U) \quad \text{for every } y \in \mathbb{R}^n \text{ such that } U + y \subset\subset \Omega.$$

Proof. Property i) is a straightforward consequence of Remark 4.2, since lower semicontinuity with respect to the convergence in Definition 3.1 implies lower semicontinuity in $L^1(\Omega)$. Property ii) follows by the measure-property criterion of the De Giorgi and Letta (see [18, Theorem 14.23]) once we show that for every $u \in GSBV^2(\Omega)$ the set function $F_{\text{hom}}^\ell(u, \cdot)$ is subadditive. The proof of the subadditivity of $F_{\text{hom}}^\ell(u, \cdot)$ follows from the fundamental estimate Lemma 4.4. In our case the main difference with respect to a standard situation is that the remainder in (4.5) is given in terms of $\|u - v\|_{L^2(S \cap \varepsilon P)}$, while we are studying the Γ -convergence of F_ε with respect to the convergence in Definition 3.1 which only ensures that $\|u - v\|_{L^1(S \cap \varepsilon P)}$ tends to zero. Therefore we provide a detailed proof of the subadditivity of $F_{\text{hom}}^\ell(u, \cdot)$.

We start observing that on $GSBV^2(\Omega) \cap L^\infty(\Omega)$ the equality $F'_- = F''_-$ in particular implies that the following limsup-type inequality is satisfied: for every $u \in GSBV^2(\Omega) \cap L^\infty(\Omega)$ and for every $U, U' \in \mathcal{A}(\Omega)$ with $U' \subset\subset U$, there exists a sequence $(u_k) \subset GSBV^2(U') \cap L^1(\Omega)$ with $u_k \rightarrow u$ in the sense of Definition 3.1 such that

$$\limsup_{k \rightarrow +\infty} F_{\varepsilon_k}(u_k, U') \leq F_{\text{hom}}^\ell(u, U)$$

(see e.g. [18, Proposition 16.4 and Remark 16.5], since both the infima in the definition of F' and F'' are actually minima).

Now let $U, V \in \mathcal{A}(\Omega)$ and let $u \in GSBV^2(\Omega) \cap L^\infty(\Omega)$. Fix any $U' \subset\subset U$, $V' \subset\subset V$, $U', V' \in \mathcal{A}(\Omega)$. Choose an open set U'' such that $U' \subset\subset U'' \subset\subset U$ and two sequences $(u_k) \subset GSBV^2(U'') \cap L^1(\Omega)$ and $(v_k) \subset GSBV^2(V') \cap L^1(\Omega)$, with $u_k \rightarrow u$ and $v_k \rightarrow u$ in the sense of Definition 3.1, and such that

$$\limsup_{k \rightarrow +\infty} F_{\varepsilon_k}(u_k, U'') \leq F_{\text{hom}}^\ell(u, U), \quad \limsup_{k \rightarrow +\infty} F_{\varepsilon_k}(v_k, V') \leq F_{\text{hom}}^\ell(u, V).$$

Moreover, since the functionals F_{ε_k} decrease by truncation, we can additionally assume that $\|\tilde{u}_k\|_{L^\infty(\Omega)} \leq \|u\|_{L^\infty(\Omega)}$, $\|\tilde{v}_k\|_{L^\infty(\Omega)} \leq \|u\|_{L^\infty(\Omega)}$, where \tilde{u}_k and \tilde{v}_k are the extended sequences in Definition 3.1. In particular $\tilde{u}_k \rightarrow u$ in $L^2(\Omega)$ and $\tilde{v}_k \rightarrow u$ in $L^2(\Omega)$.

Let $\eta > 0$ be arbitrary but fixed. Then, the fundamental estimate Lemma 4.4 provided us with a constant $M(\eta) > 0$ and a sequence (φ_k) of cut-off functions between U' and U'' such that

$$\begin{aligned} & F_{\varepsilon_k}(\varphi_k u_k + (1 - \varphi_k)v_k, U' \cup V') \\ & \leq (1 + \eta) \left(F_{\varepsilon_k}(u_k, U'') + F_{\varepsilon_k}(v_k, V') \right) + M(\eta) \|u_k - v_k\|_{L^2(\Omega \cap \varepsilon P)}^2. \end{aligned}$$

Hence, taking the limit as $k \rightarrow +\infty$, and noticing that $\varphi_k u_k + (1 - \varphi_k)v_k \rightarrow u$ in the sense of Definition 3.1, we get

$$F_{\text{hom}}^\ell(u, U' \cup V') \leq (1 + \eta) \left(F_{\text{hom}}^\ell(u, U) + F_{\text{hom}}^\ell(u, V) \right).$$

Now letting $\eta \rightarrow 0$, and then $U' \nearrow U$, $V' \nearrow V$ in view of the inner-regularity of F_{hom}^ℓ we get

$$F_{\text{hom}}^\ell(u, U \cup V) \leq F_{\text{hom}}^\ell(u, U) + F_{\text{hom}}^\ell(u, V),$$

hence the subadditivity of $F_{\text{hom}}^\ell(u, \cdot)$ for $u \in GSBV^2(\Omega) \cap L^\infty(\Omega)$.

Now let $u \in GSBV^2(\Omega)$ and, for every $m \in \mathbb{N}$, set $u^m := (u \wedge m) \vee (-m)$. Since $u^m \in L^\infty(\Omega)$ and F_{hom}^ℓ decreases by truncation (see Remark 4.2), we have

$$F_{\text{hom}}^\ell(u^m, U \cup V) \leq F_{\text{hom}}^\ell(u^m, U) + F_{\text{hom}}^\ell(u^m, V) \leq F_{\text{hom}}^\ell(u, U) + F_{\text{hom}}^\ell(u, V).$$

On the other hand, since $u^m \rightarrow u$ in $L^1(\Omega)$, the lower semicontinuity of F_{hom}^ℓ yields

$$F_{\text{hom}}^\ell(u, U \cup V) \leq \liminf_{m \rightarrow +\infty} F_{\text{hom}}^\ell(u^m, U \cup V) \leq F_{\text{hom}}^\ell(u, U) + F_{\text{hom}}^\ell(u, V),$$

thus the subadditivity of $F_{\text{hom}}^\ell(u, \cdot)$ for every $u \in GSBV^2(\Omega)$.

We now turn to the proof of iii). This will be achieved by showing that F'' is inner-regular. As a consequence we will have $F'' = F''_- = F'_- = F_{\text{hom}}^\ell \leq F'$ and therefore the Γ -convergence of F_{ε_k} to F_{hom}^ℓ .

The inner regularity of F'' is again a consequence of the fundamental estimate Lemma 4.4. In fact, let MS be as in (2.3) and fix $W \in \mathcal{A}(\Omega)$. Let $u \in GSBV^2(\Omega)$; since $MS(u, \cdot)$ is (the restriction of) a Radon measure on $\mathcal{A}(\Omega)$, for every $\eta > 0$ there exists a compact set $K \subset W$ such that $MS(u, W \setminus K) < \eta$.

Now choose $U, U' \in \mathcal{A}(\Omega)$ satisfying $K \subset U' \subset\subset U \subset\subset W$ and set $V := W \setminus K$. Recalling that $F''(u, \cdot)$ is increasing, Lemma 4.4 easily yields

$$F''(u, W) \leq F''(u, U' \cup V) \leq F''(u, U) + F''(u, V) = F''(u, U) + F''(u, W \setminus K).$$

Moreover, by the definition of F''_- and by the trivial estimate $F_\varepsilon \leq MS$ we have

$$F''(u, W) \leq F''_-(u, W) + MS(u, W \setminus K) \leq F''_-(u, W) + \eta.$$

Hence by the arbitrariness of η we get

$$F''(u, W) \leq F''_-(u, W) \quad \text{for every } W \in \mathcal{A}(\Omega), u \in GSBV^2(\Omega).$$

Therefore, as the opposite inequality is trivial, we deduce that $F''(u, \cdot)$ is inner regular.

Finally, the proof of iv) and v) is standard and follows as in e.g. [9, Lemma 3.7]. \square

4.1. Integral representation of the Γ -limit. In this subsection we show that the Γ -limit F_{hom}^ℓ can be represented in an integral form.

Theorem 4.6 (Integral representation). *Let F_{hom}^ℓ be the functional whose existence is established in Theorem 4.5. Then, for every $u \in GSBV^2(\Omega)$ and every $U \in \mathcal{A}(\Omega)$ we have*

$$F_{\text{hom}}^\ell(u, U) = \int_U f_{\text{hom}}^\ell(\nabla u) dx + \int_{S_u \cap U} g_{\text{hom}}^\ell([u], \nu_u) d\mathcal{H}^{n-1} \quad (4.7)$$

for a convex function $f_{\text{hom}}^\ell: \mathbb{R}^n \rightarrow [0, +\infty)$ satisfying for every $\ell \in [0, +\infty]$ and every $\xi \in \mathbb{R}^n$ the following bounds

$$\hat{f}(\xi) \leq f_{\text{hom}}^\ell(\xi) \leq |\xi|^2 \quad (4.8)$$

and a Borel function $g_{\text{hom}}^\ell: \mathbb{R} \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ satisfying:

- i) (monotonicity in t and symmetry) for any fixed $\nu \in \mathbb{S}^{n-1}$, $g_{\text{hom}}^\ell(\cdot, \nu)$ is nondecreasing on $(0, +\infty)$ and satisfies the symmetry condition $g_{\text{hom}}^\ell(-t, -\nu) = g_{\text{hom}}^\ell(t, \nu)$ for $t \in \mathbb{R}$;
- ii) (subadditivity in t) for any $\nu \in \mathbb{S}^{n-1}$

$$g_{\text{hom}}^\ell(t_1 + t_2, \nu) \leq g_{\text{hom}}^\ell(t_1, \nu) + g_{\text{hom}}^\ell(t_2, \nu),$$

for every $t_1, t_2 \in \mathbb{R}$;

- iii) (convexity in ν) for any $t \in \mathbb{R}$, the 1-homogeneous extension of $g_{\text{hom}}^\ell(t, \cdot): \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ to \mathbb{R}^n is convex. This condition can be also equivalently expressed in terms of the function g_{hom}^ℓ as

$$g_{\text{hom}}^\ell(t, \nu) \leq \lambda_1 g_{\text{hom}}^\ell(t, \nu_1) + \lambda_2 g_{\text{hom}}^\ell(t, \nu_2),$$

for every $\nu, \nu_1, \nu_2 \in \mathbb{S}^{n-1}$, $\lambda_1, \lambda_2 \geq 0$ such that $\lambda_1 \nu_1 + \lambda_2 \nu_2 = \nu$.

Proof. We recall that F_ε satisfies the bound (3.2), namely $\hat{F}_\varepsilon \leq F_\varepsilon \leq MS$ in $L^1(\Omega)$. Therefore by Γ -convergence and in view of Remark 2.2 and Remark 3.3 ii. we get

$$\hat{F} \leq F_{\text{hom}}^\ell \leq MS \quad \text{in } L^1(\Omega), \quad (4.9)$$

thus invoking (2.10) we deduce

$$\min\{c_1, c_2\} MS \leq F_{\text{hom}}^\ell \leq MS \quad \text{in } L^1(\Omega), \quad (4.10)$$

where c_1 and c_2 are as in (2.7) and (2.9), respectively. Hence, Theorem 4.5, (4.10), [6, Theorem 1], and a standard perturbation argument (see e.g. [20, Theorem 2.2]) yield the integral representation (4.7) on the space $SBV^2(\Omega)$. Then a standard truncation and continuity argument allows to extend this integral representation to the whole space $GSBV^2(\Omega)$ and thus to get exactly (4.7).

The convexity of f_{hom}^ℓ , the subadditivity of g_{hom}^ℓ in t and the convexity in ν of its 1-homogeneous extension to \mathbb{R}^n are straightforward consequences of the $L^1(\Omega)$ -lower semicontinuity of F_{hom}^ℓ . Since moreover $f_{\text{hom}}^\ell(\xi) = F_{\text{hom}}^\ell(u_\xi, Q)$ where $u_\xi(x) := \xi \cdot x$, the bounds in (4.8) are an immediate consequence of (4.9). Finally, the monotonicity in t and the symmetry of g_{hom}^ℓ follow from [6, Theorem 1]. \square

5. IDENTIFICATION OF THE HOMOGENIZED SURFACE INTEGRAND

In this section we identify the limit surface integrand g_{hom}^ℓ . To do so we make use of the following technical lemma (see [16, Lemma 4.5], and see also [17, Lemma 2.5] for a more general version of the result).

Lemma 5.1 (“Fracture Lemma”). *Let $n \geq 2$ and $\eta \in (0, 1]$ be fixed. There exists a constant $\gamma = \gamma(n, \eta) > 0$ such that if $0 < s \leq r$, and $u \in \mathcal{P}(B_{r, r+s})$ verify the following hypotheses:*

- (H1) $\mathcal{H}^{n-1}(S_u \cap B_{r, r+s}) \leq \mathcal{H}^{n-1}(S_v \cap B_{r, r+s})$ for every competitor $v \in \mathcal{P}(B_{r, r+s})$ satisfying $\text{supp}(u - v) \subset B_{r, r+s}$;
- (H2) $\mathcal{H}^{n-1}(S_u \cap B_{r, r+s}) \leq \gamma s^{n-1}$;

then for every r_0 and s_0 such that $r \leq r_0 < r_0 + s_0 \leq r + s$ and $s_0 \geq \eta s$, there exists a radius $\bar{r} \in (r + s_0/3, r_0 + 2s_0/3)$ with the property that

$$S_u \cap \partial B_{\bar{r}} = \emptyset.$$

The following theorem is the main result of this section.

Theorem 5.2 (Identification of the homogenized surface integrand). *Let g_{hom}^ℓ be the function as in Theorem 4.6 and let \hat{g} be as in (2.8). Then, for every $(t, \nu) \in (\mathbb{R} \setminus \{0\}) \times \mathbb{S}^{n-1}$ and every $\ell \in [0, +\infty]$ we have $g_{\text{hom}}^\ell(t, \nu) = \hat{g}(\nu)$.*

Proof. For $\varepsilon, \delta > 0$ we define the comparison functional $G_\varepsilon^\delta : L^1(\Omega) \rightarrow [0, +\infty]$ as

$$G_\varepsilon^\delta(u) := \begin{cases} \int_\Omega |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u \cap \varepsilon P) + \delta \int_{S_u} (1 + |[u]|) d\mathcal{H}^{n-1} & \text{if } u \in SBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega). \end{cases}$$

Let $\delta > 0$ be fixed; by [9, Theorem 2.3] we deduce that, as ε tends to zero, the functionals G_ε^δ Γ -converge, with respect to the $L^1(\Omega)$ -convergence, to the functional $G^\delta : L^1(\Omega) \rightarrow [0, +\infty]$ given by

$$G^\delta(u) := \begin{cases} \int_\Omega |\nabla u|^2 dx + \int_{S_u} g^\delta([u], \nu_u) d\mathcal{H}^{n-1}, & \text{if } u \in SBV^2(\Omega), \\ +\infty, & \text{otherwise in } L^1(\Omega), \end{cases}$$

where $g^\delta : \mathbb{R} \times \mathbb{S}^{n-1} \rightarrow [0, +\infty]$ is defined as

$$g^\delta(z, \nu) := \lim_{t \rightarrow +\infty} \frac{1}{t^{n-1}} \inf \left\{ \int_{S_u \cap tQ^\nu} (\chi_P + \delta(1 + |[u]|)) d\mathcal{H}^{n-1} : u \in SBV^{\text{pc}}(tQ^\nu), \right. \\ \left. u = u_{0,z}^\nu \text{ in a neighbourhood of } \partial(tQ^\nu) \right\}. \quad (5.1)$$

Let $\varepsilon > 0$ be small enough to have that $\beta_\varepsilon < \delta$; then we immediately deduce the bounds

$$\widehat{F}_\varepsilon(u) \leq F_\varepsilon(u) \leq G_\varepsilon^\delta(u), \quad \text{for every } u \in L^1(\Omega). \quad (5.2)$$

From (5.2) it follows that for every $z \in \mathbb{R} \setminus \{0\}$, $\nu \in \mathbb{S}^{n-1}$, and $\ell \in [0, +\infty]$

$$\hat{g}(\nu) \leq g_{\text{hom}}^\ell(z, \nu) \leq g^\delta(z, \nu). \quad (5.3)$$

Indeed let $(u_\varepsilon) \subset L^1(\Omega)$ be a recovery sequence for G_ε^δ , with $u_\varepsilon \rightarrow u_{0,z}^\nu$ strongly in $L^1(\Omega)$. Then, since $u_\varepsilon \rightarrow u_{0,z}^\nu$ in the sense of Definition 3.1, we have

$$g^\delta(z, \nu) \mathcal{H}^{n-1}(\Omega \cap \Pi_0^\nu) = G^\delta(u_{0,z}^\nu) = \lim_{\varepsilon \rightarrow 0} G_\varepsilon^\delta(u_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \\ \geq F^\ell(u_{0,z}^\nu) = g_{\text{hom}}^\ell(z, \nu) \mathcal{H}^{n-1}(\Omega \cap \Pi_0^\nu).$$

Similarly, let $(u_\varepsilon) \subset L^1(\Omega)$ be a recovery sequence for F_ε , with $u_\varepsilon \rightarrow u_{0,z}^\nu$ in the sense of Definition 3.1. Then there exists a sequence \tilde{u}_ε with $\tilde{u}_\varepsilon = u_\varepsilon$ in $\Omega \cap \varepsilon P$ and such that \tilde{u}_ε converges to $u_{0,z}^\nu$ strongly in $L^1(\Omega)$; hence

$$g_{\text{hom}}^\ell(z, \nu) \mathcal{H}^{n-1}(\Omega \cap \Pi_0^\nu) = F^\ell(u_{0,z}^\nu) = \lim_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} \widehat{F}_\varepsilon(u_\varepsilon) = \liminf_{\varepsilon \rightarrow 0} \widehat{F}_\varepsilon(\tilde{u}_\varepsilon) \\ \geq \widehat{F}(u_{0,z}^\nu) = \hat{g}(z, \nu) \mathcal{H}^{n-1}(\Omega \cap \Pi_0^\nu),$$

where $\Pi_0^\nu := \{y \in \mathbb{R}^n : y \cdot \nu = 0\}$.

We now claim that for every $z \in \mathbb{R} \setminus \{0\}$, $\nu \in \mathbb{S}^{n-1}$

$$g^\delta(z, \nu) \leq \hat{g}(\nu) + o(1), \quad (5.4)$$

as $\delta \rightarrow 0$; hence (5.4) together with (5.3) will imply the thesis.

To prove (5.4) let $\nu \in \mathbb{S}^{n-1}$ and $t > 0$ be fixed, and let $\bar{u} \in \mathcal{P}(tQ^\nu \cap P)$ be such that $\bar{u} = u_{0,1}^\nu$ in a neighbourhood of $\partial(tQ^\nu)$ and

$$\mathcal{H}^{n-1}(S_{\bar{u}} \cap tQ^\nu \cap P) = \min \left\{ \mathcal{H}^{n-1}(S_u \cap tQ^\nu \cap P) : u \in \mathcal{P}(tQ^\nu \cap P), u = u_{0,1}^\nu \text{ on } \partial(tQ^\nu) \right\}. \quad (5.5)$$

Note that the minimiser in (5.5) exists: indeed any minimising sequence is weakly convergent in $BV(tQ^\nu \cap P)$, and the Hausdorff measure of the jump set in $tQ^\nu \cap P$ is lower semicontinuous.

We now modify \bar{u} in order to get a competitor for the minimisation problem in the definition of g^δ . More precisely, we construct from \bar{u} a Caccioppoli partition w defined on the whole tQ^ν and such that

$$\mathcal{H}^{n-1}(S_w \cap tQ^\nu \cap P) + \delta \int_{S_w \cap tQ^\nu} (1 + |[w]|) d\mathcal{H}^{n-1} \leq (1 + \delta c(z)) \mathcal{H}^{n-1}(S_{\bar{u}} \cap tQ^\nu \cap P) + \delta c(z) t^{n-1} \quad (5.6)$$

holds true for some constant $c(z) > 0$ independent of t and δ .

It is convenient to write

$$tQ^\nu \cap P = \bigcup_{k \in \mathcal{I}^1} (Q^k \cap P) \cup \bigcup_{k \in \mathcal{I}^2} (Q^k \cap tQ^\nu \cap P),$$

where $Q^k := Q + k$, $\mathcal{I}^1 := \{k \in \mathbb{Z}^n : Q^k \subset tQ^\nu\}$, and $\mathcal{I}^2 := \{k \in \mathbb{Z}^n : Q^k \cap \partial(tQ^\nu) \neq \emptyset\}$. We now illustrate in detail the multi-step construction of the desired function w .

Step 1: Modification of \bar{u} in the “internal” cubes Q^k , for $k \in \mathcal{I}^1$. Let $k \in \mathcal{I}^1$, and set $u^k := \bar{u}|_{Q^k \cap P}$. We recall that for our choice of P we have $Q^k \cap P = Q_{1/2,1}^k$, where we defined $Q_{a,b}^k := Q_{a,b} + k$ for any $0 < a < b \leq 1$. We also define $B_{a,b}^k := B_{a,b} + k$, for any $0 < a < b \leq 1$.

Let $\sqrt{2}/4 \leq r < r+s < 1$ be fixed. We extend the function u^k differently depending on whether the function u^k admits a *small* or a *large* jump set in $B_{r,r+s}^k$ (note that $B_{r,r+s}^k \subseteq Q_{1/2,1}^k$). More precisely, we say that u^k has a small jump set in $B_{r,r+s}^k$ if

$$\mathcal{H}^{n-1}(S_{u^k} \cap B_{r,r+s}^k) \leq \gamma s^{n-1}, \quad (5.7)$$

where $\gamma = \gamma(n, \eta)$ is as in Lemma 5.1, corresponding to $\eta = 1$.

Note that for the cubes with small jump, the assumptions (H1) and (H2) of Lemma 5.1 are satisfied by u^k . Indeed, (H1) follows by the minimality of \bar{u} (defined in (5.5)) in $tQ^\nu \cap P$, which implies its local minimality in every subset, and hence in particular the local minimality of u^k in $B_{r,r+s}^k$, for every $k \in \mathcal{I}^1$. Finally, (H2) is exactly (5.7).

By Lemma 5.1 there exists $\bar{r} \in (r + s/3, r + 2s/3)$ such that $S_{u^k} \cap \partial B_{\bar{r}}^k = \emptyset$, namely the trace of u^k on $\partial B_{\bar{r}}^k$ is constant. We denote this constant value by m^k . Then we define the function v^k on the whole Q^k as follows

$$v^k := \begin{cases} u^k & \text{in } Q^k \setminus \bar{B}_{\bar{r}}^k, \\ m^k & \text{in } \bar{B}_{\bar{r}}^k. \end{cases} \quad (5.8)$$

Clearly $v^k \in \mathcal{P}(Q^k)$, and

$$\mathcal{H}^{n-1}(S_{v^k} \cap Q^k) = \mathcal{H}^{n-1}(S_{u^k} \cap (Q^k \setminus \bar{B}_{\bar{r}}^k)) \leq \mathcal{H}^{n-1}(S_{\bar{u}} \cap Q^k \cap P).$$

Hence in a cube with small jump we have replaced \bar{u} with a function $v^k \in \mathcal{P}(Q^k)$ whose Mumford-Shah energy in Q^k is controlled by the energy of \bar{u} in $Q^k \cap P$.

If now u^k has a large jump in $B_{r,r+s}^k$; *i.e.*, if (5.7) is not satisfied, then we extend u^k to Q^k by simply setting

$$v^k := \begin{cases} u^k & \text{in } Q_{1/2,1}^k, \\ 0 & \text{in } Q_{1/2}^k. \end{cases} \quad (5.9)$$

Clearly $v^k \in \mathcal{P}(Q^k)$, and

$$\begin{aligned} \mathcal{H}^{n-1}(S_{v^k} \cap Q^k) &\leq 2n(1/2)^{n-1} + \mathcal{H}^{n-1}(S_{\bar{u}} \cap Q^k \cap P) \\ &< \frac{2n(1/2)^{n-1}}{\gamma s^{n-1}} \mathcal{H}^{n-1}(S_{u^k} \cap B_{r,r+s}^k) + \mathcal{H}^{n-1}(S_{\bar{u}} \cap Q^k \cap P) \\ &\leq C \mathcal{H}^{n-1}(S_{\bar{u}} \cap Q^k \cap P), \end{aligned}$$

where ω_{n-1} is the surface of the unit sphere \mathbb{S}^{n-1} . Thus finally, for every $k \in \mathcal{I}^1$ we have replaced \bar{u} with a function $v^k \in \mathcal{P}(Q^k) \subset SBV^{\text{pc}}(Q^k)$ whose energy in Q^k is controlled by the energy of \bar{u} in $Q^k \cap P$; moreover, $\|\bar{u}\|_{L^\infty(Q^k \cap P)} = \|v^k\|_{L^\infty(Q^k)}$.

Step 2: Modification of \bar{u} in the “boundary” cubes Q^k , $k \in \mathcal{I}^2$. In this step we consider only the cubes Q^k such that $Q^k \cap \partial(tQ^\nu) \neq \emptyset$. In order to preserve the boundary condition we need to distinguish between two cases. If $Q^k \cap \partial(tQ^\nu \cap \{x \cdot \nu > 0\}) \neq \emptyset$, in Q^k we set

$$v^k := \begin{cases} \bar{u} & \text{in } Q_{1/2,1}^k, \\ 1 & \text{in } Q_{1/2}^k, \end{cases} \quad (5.10)$$

while in those cubes Q^k such that $Q^k \cap \partial(tQ^\nu \cap \{x \cdot \nu < 0\}) \neq \emptyset$, we set

$$v^k := \begin{cases} \bar{u} & \text{in } Q_{1/2,1}^k, \\ 0 & \text{in } Q_{1/2}^k. \end{cases} \quad (5.11)$$

The additional energy contribution of the boundary cubes is proportional to the perimeter of tQ^ν ; *i.e.*, of order Ct^{n-1} for some $C > 0$ independent of t .

Step 3: Adding up all the cubes. We now denote with $v \in \mathcal{P}(tQ^\nu)$ the function defined as $v = v^k$ in Q^k for every k , where v^k is as in (5.8) or (5.9) if $k \in \mathcal{I}^1$, as in (5.10) if $k \in \mathcal{I}^2$ and Q^k intersects $\partial(tQ^\nu \cap \{x \cdot \nu > 0\})$, or (5.11) if $k \in \mathcal{I}^2$ and Q^k intersects $\partial(tQ^\nu \cap \{x \cdot \nu < 0\})$.

By construction the function v satisfies the following properties:

- i. $v \in \mathcal{P}(tQ^\nu)$,
- ii. $v = u_{0,1}^\nu$ in a neighbourhood of $\partial(tQ^\nu)$,
- iii. $\mathcal{H}^{n-1}(S_v \cap (tQ^\nu \cap P)) \leq \mathcal{H}^{n-1}(S_{\bar{u}} \cap (tQ^\nu \cap P))$,
- iv. $\mathcal{H}^{n-1}(S_v \cap tQ^\nu) \leq C(\mathcal{H}^{n-1}(S_{\bar{u}} \cap (tQ^\nu \cap P)) + t^{n-1})$,
- iv. $\|v\|_{L^\infty(tQ^\nu)} = \|\bar{u}\|_{L^\infty(tQ^\nu \cap P)} = 1$.

We finally set $w := zv$ so that $w = u_{0,z}^\nu$ in a neighbourhood of $\partial(tQ^\nu)$ and therefore it can be used as a competitor in the minimisation problem defining $g^\delta(z, \nu)$.

We are now able to compare $g^\delta(z, \nu)$ and $\hat{g}(\nu)$. By definition of w have

$$\begin{aligned} & \mathcal{H}^{n-1}(S_w \cap tQ^\nu \cap P) + \delta \int_{S_w \cap tQ^\nu} (1 + |[w]|) d\mathcal{H}^{n-1} \\ & \leq \mathcal{H}^{n-1}(S_{\bar{u}} \cap (tQ^\nu \cap P)) + \delta(1 + |z|)\mathcal{H}^{n-1}(S_v \cap tQ^\nu) \\ & \leq \mathcal{H}^{n-1}(S_{\bar{u}} \cap (tQ^\nu \cap P)) + \delta C(1 + |z|)(\mathcal{H}^{n-1}(S_{\bar{u}} \cap (tQ^\nu \cap P)) + t^{n-1}), \end{aligned}$$

which is exactly the claim (5.6). Then in view of (5.1) by dividing the above expression by t^{n-1} and letting $t \rightarrow +\infty$, we get

$$g^\delta(z, \nu) \leq \lim_{t \rightarrow +\infty} \frac{1}{t^{n-1}} \left(\mathcal{H}^{n-1}(S_{\bar{u}} \cap (tQ^\nu \cap P))(1 + \delta C(1 + |z|)) + \delta C(1 + |z|) \right).$$

By virtue of (5.5) the previous estimate yields

$$g^\delta(z, \nu) \leq \hat{g}(\nu) (1 + \delta C(1 + |z|)) + \delta C(1 + |z|),$$

and hence the claim. \square


6. IDENTIFICATION OF THE HOMOGENIZED VOLUME INTEGRAND

In this section we identify the limit volume integrand f_{hom}^ℓ . We start with a preliminary result.

Lemma 6.1. *Let f_{hom}^ℓ be as in Theorem 4.6 and let \hat{f} be as in (2.6). For every $\xi \in \mathbb{R}^n$ we have*

$$\hat{f}(\xi) \leq f_{\text{hom}}^\ell(\xi) \leq \min\{|\xi|^2, \hat{f}(\xi) + C\ell\}, \quad (6.1)$$

where $C > 0$ is a constant depending only on P and on Ω .

FIGURE 2. Bounds on f_{hom}^ℓ

Proof. Let $\xi \in \mathbb{R}^n \setminus \{0\}$. By the classical homogenization result for the Dirichlet functional on a perforated domain [7, Theorem 5.1], there exists a sequence $(v_\varepsilon) \subset H^1(\Omega)$ which converges strongly in $L^2(\Omega)$, as $\varepsilon \rightarrow 0$, to the linear function $u_\xi(x) := \xi \cdot x$, such that $\|Dv_\varepsilon\|_{L^2(\Omega)} \leq C$, and

$$\limsup_{\varepsilon \rightarrow 0} \int_{\Omega \cap \varepsilon P} |Dv_\varepsilon|^2 dx = \mathcal{L}^n(\Omega) \hat{f}(\xi), \quad (6.2)$$

where \hat{f} is the quadratic form defined in (2.6). We define the new sequence $u_\varepsilon : \Omega \rightarrow \mathbb{R}$ as

$$u_\varepsilon := \begin{cases} v_\varepsilon & \text{in } Q_{\frac{1}{2}, 1}^{k, \varepsilon} \cap \Omega \\ m_\varepsilon^k & \text{in } Q_{\frac{1}{2}}^{k, \varepsilon} \cap \Omega \end{cases} \quad k \in \mathbb{Z}^n \quad (6.3)$$

where for $0 < s < r$ we set $Q_{s, r}^{k, \varepsilon} := \varepsilon(Q_{s, r} + k)$ for $k \in \mathbb{Z}^n$, and

$$m_\varepsilon^k := \int_{Q_{\frac{1}{2}}^{k, \varepsilon} \cap \Omega} v_\varepsilon(x) dx.$$

By (6.3) we immediately get

$$\|v_\varepsilon - u_\varepsilon\|_{L^2(\Omega)}^2 = \sum_{k \in \mathbb{Z}^n} \int_{Q_{\frac{1}{2}}^{k, \varepsilon} \cap \Omega} |v_\varepsilon(x) - m_\varepsilon^k|^2 dx,$$

moreover by the Poincaré-Wirtinger inequality, for every $k \in \mathbb{Z}^n$ we have

$$\int_{Q_{\frac{1}{2}}^{k, \varepsilon} \cap \Omega} |v_\varepsilon(x) - m_\varepsilon^k|^2 dx \leq C\varepsilon^2 \int_{Q_{\frac{1}{2}}^{k, \varepsilon} \cap \Omega} |Dv_\varepsilon(x)|^2 dx,$$

with a constant $C > 0$ independent of ε and of k . From the bound $\|Dv_\varepsilon\|_{L^2(\Omega)} \leq C$ we eventually deduce

$$\|v_\varepsilon - u_\varepsilon\|_{L^2(\Omega)} \leq C\varepsilon,$$

therefore, since $v_\varepsilon \rightarrow u_\xi$ strongly in $L^2(\Omega)$, also (u_ε) converges to u_ξ strongly in $L^2(\Omega)$. We now estimate

$$\begin{aligned} F_\varepsilon(u_\varepsilon) &= \int_{\Omega} |\nabla u_\varepsilon|^2 dx + \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap (\Omega \cap \varepsilon P)) + \beta_\varepsilon \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap (\Omega \setminus \varepsilon P)) \\ &\leq \int_{\Omega \cap \varepsilon P} |Dv_\varepsilon|^2 dx + \beta_\varepsilon \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap \partial(\Omega \setminus \varepsilon P)). \end{aligned} \quad (6.4)$$

For the second term in (6.4) we have

$$\beta_\varepsilon \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap \partial(\Omega \setminus \varepsilon P)) \leq \beta_\varepsilon N(\varepsilon) \mathcal{H}^{n-1}(\partial Q_{\varepsilon/2}) + \beta_\varepsilon \mathcal{H}^{n-1}(\partial\Omega),$$

where $N(\varepsilon)$ is the cardinality of the set $\{k \in \mathbb{Z}^n : \overline{Q^{k,\varepsilon}} \cap \Omega \neq \emptyset\}$. Since Ω is a bounded set with Lipschitz boundary, we have that $N(\varepsilon) \leq c/\varepsilon^n$ and $\mathcal{H}^{n-1}(\partial\Omega) < c$, which gives

$$F_\varepsilon(u_\varepsilon) \leq \int_{\Omega \cap \varepsilon P} |Dv_\varepsilon|^2 dx + C \frac{\beta_\varepsilon}{\varepsilon}.$$

Now, as in particular $u_\varepsilon \rightarrow u_\xi$ in the sense of Definition 3.1, by Theorem 4.5 and Theorem 4.6, we have

$$\mathcal{L}^n(\Omega) f_{\text{hom}}^\ell(\xi) = F_{\text{hom}}^\ell(u_\xi) \leq \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \leq \limsup_{\varepsilon \rightarrow 0} \int_{\Omega \cap \varepsilon P} |Dv_\varepsilon|^2 dx + C\ell,$$

which, together with (6.2), gives

$$\mathcal{L}^n(\Omega) f_{\text{hom}}^\ell(\xi) \leq \limsup_{\varepsilon \rightarrow 0} \int_{\Omega \cap \varepsilon P} |Dv_\varepsilon|^2 dx + C\ell \leq \mathcal{L}^n(\Omega) \hat{f}(\xi) + C\ell.$$

Dividing by $\mathcal{L}^n(\Omega)$ and using (4.8) concludes the proof. \square

6.1. Subcritical case: $\ell = 0$.

Theorem 6.2 (Identification of the homogenized volume integrand for $\ell = 0$). *Let f_{hom}^0 be as in Theorem 4.6 and corresponding to the choice $\ell = 0$, and let \hat{f} be as in (2.6). Then for every $\xi \in \mathbb{R}^n$ we have $f_{\text{hom}}^0(\xi) = \hat{f}(\xi)$.*

Proof. The thesis readily follows from (6.1) by using the assumption that $\ell = 0$, and the upper bound in (2.7). \square

Remark 6.3. Note that, although in Theorem 4.5 the Γ -convergence of F_ε has been established only up to subsequences (and the Γ -limit might be in principle different along different subsequences), in the subcritical case the situation is different. Indeed, thanks to Theorem 5.2 and Theorem 6.2, we deduce that the Γ -limit is the same for every subsequence, and is given by the functional \widehat{F} defined in (2.5).

6.2. Supercritical case: $\ell = +\infty$.

Theorem 6.4 (Identification of the homogenized volume integrand for $\ell = +\infty$). *Let f_{hom}^∞ be as in Theorem 4.6 and corresponding to the choice $\ell = +\infty$. Then, for every $\xi \in \mathbb{R}^n$, we have that $f_{\text{hom}}^\infty(\xi) = |\xi|^2$.*

Remark 6.5. Theorem 5.2 and Theorem 6.4 imply that, for $\ell = +\infty$, the Γ -limit of (F_ε) is

$$F_{\text{hom}}^\infty(u) := \begin{cases} \int_{\Omega} |\nabla u|^2 dx + \int_{S_u} \hat{g}(\nu_u) d\mathcal{H}^{n-1} & \text{if } u \in GSBV^2(\Omega), \\ +\infty & \text{otherwise in } L^1(\Omega). \end{cases}$$

In particular, the whole sequence (F_ε) Γ -converges to F_{hom}^∞ .

Before proving Theorem 6.4 above we need to recall the Elimination Property proved in [19, Lemma 0.7] (see also [21, Theorem 3.6]). For the definition of local minimiser of the Mumford-Shah functional we refer to [1, Definition 6.6].

Theorem 6.6 (Elimination property). *Let $\Omega \subset \mathbb{R}^n$ be open. There exists a strictly positive dimensional constant $\theta = \theta(n)$ independent of Ω such that, if $u \in SBV^2(\Omega)$ is a local minimiser of the Mumford-Shah functional and $B_\rho(x_0) \subset \Omega$ is any ball with centre x_0 and with*

$$\mathcal{H}^{n-1}(S_u \cap B_\rho(x_0)) < \theta \rho^{n-1},$$

then $S_u \cap B_{\frac{\rho}{2}}(x_0) = \emptyset$.

We now introduce some auxiliary functionals which will be used in the proof of Theorem 6.4.

Let $\frac{1}{2} < r < 1$ and let $\varphi \in H^{1/2}(\partial Q_r)$. For $h \in \mathbb{N}$ and $t > 1$ we define the functionals $I_\varphi, I_\varphi^{h,t} : L^1(Q_r) \rightarrow [0, +\infty]$ as follows:

$$I_\varphi(u) := \begin{cases} \int_{Q_r} |Du|^2 dx & \text{if } u \in H^1(Q_r), u = \varphi \text{ on } \partial Q_r, \\ +\infty & \text{otherwise in } L^1(Q_r), \end{cases} \quad (6.5)$$

and

$$I_\varphi^{h,t}(u) := \begin{cases} \int_{Q_r} |\nabla u|^2 dx + t\mathcal{H}^{n-1}(S_u \cap Q_{\frac{1}{2},r}) + \mathcal{H}^{n-1}(S_u \cap Q_{\frac{1}{2}}) \\ \text{if } u \in SBV^2(Q_r), \mathcal{H}^{n-1}(S_u) \leq \frac{1}{h}, w = \varphi \text{ on } \partial Q_r, \\ +\infty & \text{otherwise in } L^1(Q_r). \end{cases} \quad (6.6)$$

The next result is a straightforward adaption of [23, Lemma 4.3].

Lemma 6.7. *Let $\frac{1}{2} < r < 1$; let $\varphi \in H^{1/2}(\partial Q_r)$, and let $(\varphi_h) \subset H^{1/2}(\partial Q_r)$ be a sequence with*

$$\varphi_h \rightarrow \varphi \text{ in } H^{1/2}(\partial Q_r), \text{ as } h \rightarrow +\infty.$$

Then the functionals I_{φ_h} and $I_{\varphi_h}^{h,t}$ defined, respectively, as in (6.5) and (6.6), with φ replaced by φ_h , Γ -converge with respect to the strong $L^1(\Omega)$ -topology, as $h \rightarrow +\infty$, to the Dirichlet functional $I_\varphi : L^1(Q_r) \rightarrow [0, +\infty]$ defined in (6.5).

We now state and prove a technical result which is the heart of the proof of Theorem 6.4.

Theorem 6.8 (Lower bound for F_ε). *Let $\xi \neq 0$ and let $(u_\varepsilon) \subset L^1(\Omega)$ be a sequence such that $\sup_\varepsilon F_\varepsilon(u_\varepsilon) < +\infty$ and $u_\varepsilon \rightarrow u_\xi$ in the sense of Definition 3.1. Then*

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \geq \mathcal{L}^n(\Omega) |\xi|^2. \quad (6.7)$$

Proof. Assume that $F_\varepsilon(u_\varepsilon) \leq c$. The proof strategy consists of replacing the sequence u_ε with an improved sequence w_ε (in a sense that will be clarified below) which converges to u_ξ strongly in $L^1(\Omega)$, and whose energy is asymptotically smaller than $F_\varepsilon(u_\varepsilon)$.

Since the energy F_ε decreases by truncations, we can truncate the sequence (u_ε) at level $\|u_\xi\|_{L^\infty(\Omega)}$ and preserve both the uniform bound on $F_\varepsilon(u_\varepsilon)$ and the convergence of (u_ε) to u_ξ . Hence in what follows we assume that $\|u_\varepsilon\|_{L^\infty(\Omega)} \leq \|u_\xi\|_{L^\infty(\Omega)}$.

As an initial step we rewrite Ω as

$$\Omega = \left(\bigcup_{k \in \mathcal{I}_\varepsilon^1} Q^{k,\varepsilon} \right) \cup \left(\bigcup_{k \in \mathcal{I}_\varepsilon^2} Q^{k,\varepsilon} \cap \Omega \right),$$

where $Q^{k,\varepsilon} := \varepsilon(Q + k)$, $\mathcal{I}_\varepsilon^1 := \{z \in \mathbb{Z}^n : Q^{k,\varepsilon} \subset \Omega\}$, and $\mathcal{I}_\varepsilon^2 := \{z \in \mathbb{Z}^n : Q^{k,\varepsilon} \cap \partial\Omega \neq \emptyset\}$. Clearly

$$F_\varepsilon(u_\varepsilon) = \sum_{k \in \mathcal{I}_\varepsilon^1} F_\varepsilon(u_\varepsilon, Q^{k,\varepsilon}) + \sum_{k \in \mathcal{I}_\varepsilon^2} F_\varepsilon(u_\varepsilon, Q^{k,\varepsilon} \cap \Omega).$$

Step 1: Classification of the interior cubes. We estimate the energy in $Q^{k,\varepsilon}$, for $k \in \mathcal{I}_\varepsilon^1$, namely

$$F_\varepsilon(u_\varepsilon, Q^{k,\varepsilon}) = \int_{Q^{k,\varepsilon}} |\nabla u_\varepsilon|^2 dx + \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap Q_{\frac{1}{2},1}^{k,\varepsilon}) + \beta_\varepsilon \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap Q_{\frac{1}{2}}^{k,\varepsilon}).$$

For $y \in Q^k$ set $v_\varepsilon(y) := (\sqrt{\beta_\varepsilon \varepsilon})^{-1} u_\varepsilon(\varepsilon y)$; by changing variables we have

$$\begin{aligned} F_\varepsilon(u_\varepsilon, Q^{k,\varepsilon}) &= \beta_\varepsilon \varepsilon^{n-1} \left(\int_{Q^k} |\nabla v_\varepsilon|^2 dy + \frac{1}{\beta_\varepsilon} \mathcal{H}^{n-1}(S_{v_\varepsilon} \cap Q_{\frac{1}{2},1}^k) + \mathcal{H}^{n-1}(S_{v_\varepsilon} \cap Q_{\frac{1}{2}}^k) \right) \\ &=: \beta_\varepsilon \varepsilon^{n-1} \mathcal{F}_\varepsilon(v_\varepsilon, Q^k). \end{aligned}$$

Let $\vartheta > 0$ be a fixed constant, and let $\varepsilon > 0$ be fixed. We call Q^k a *good* cube if

$$\mathcal{F}_\varepsilon(v_\varepsilon, Q^k) \leq C \quad \text{for some } C > 0 \quad \text{and} \quad \mathcal{H}^{n-1}(S_{v_\varepsilon} \cap Q^k) \leq \vartheta, \quad (6.8)$$

namely if both the energy of v_ε and the total measure of the jump of v_ε are bounded in the cube. Otherwise, we say that Q^k is a *bad* cube. Let $\mathcal{I}_\varepsilon^{1,g}, \mathcal{I}_\varepsilon^{1,b} \subset \mathcal{I}_\varepsilon^1$ denote the set of internal good and bad cubes, respectively; we denote with N_ε^g and N_ε^b their cardinalities.

We can easily estimate the number of bad cubes, by using the fact that either one of the conditions (6.8) is not satisfied. Namely, if the first condition in (6.8) is not satisfied, then

$$c \geq F_\varepsilon(u_\varepsilon) \geq \beta_\varepsilon \varepsilon^{n-1} \sum_{k \in \mathcal{I}_\varepsilon^{1,b}} \mathcal{F}_\varepsilon(v_\varepsilon, Q^k) \geq C \beta_\varepsilon \varepsilon^{n-1} N_\varepsilon^b.$$

Similarly, if the second condition in (6.8) is not satisfied, then

$$c \geq \beta_\varepsilon \varepsilon^{n-1} N_\varepsilon^b \vartheta.$$

Hence, we have the bound

$$N_\varepsilon^b \leq \frac{C(\vartheta)}{\beta_\varepsilon \varepsilon^{n-1}}. \quad (6.9)$$

Step 2: Energy estimate on the good cubes. This is the most delicate part of the proof, and is split into a number of sub-steps.

Step 2.1: Elimination property. Let Q^k , for $k \in \mathcal{I}_\varepsilon^{1,g}$, be an arbitrary good cube in the sense of (6.8). We omit now the superscript k for the sake of notation. Let $\frac{1}{2} < \varsigma < 1$; for fixed $\varepsilon > 0$ consider the following (local) minimisation problem:

$$(\text{LMS})_\varepsilon \text{ loc min} \left\{ \mathcal{F}_\varepsilon(v, Q_\varsigma) : v \in SBV^2(Q_\varsigma), \mathcal{F}_\varepsilon(v, Q_\varsigma) \leq C \text{ for some } C > 0, \mathcal{H}^{n-1}(S_v \cap Q_\varsigma) \leq \vartheta \right\}.$$

Let $\mathcal{M}_\vartheta^\varepsilon$ denote the class of solutions of $(\text{LMS})_\varepsilon$, and let $\hat{v}_\varepsilon \in \mathcal{M}_\vartheta^\varepsilon$. We recall that, following [1, Definition 6.6], for every open set $A \subset\subset Q_\varsigma$ we have that $\mathcal{F}_\varepsilon(\hat{v}_\varepsilon, A) \leq \mathcal{F}_\varepsilon(v, A)$, whenever $\{\hat{v}_\varepsilon \neq v\} \subset\subset A$.

With no loss of generality we can assume that \hat{v}_ε is bounded in $L^\infty(Q_\varsigma)$ (with a possibly ε -dependent constant) as v_ε , since the energy \mathcal{F}_ε decreases by truncations. We observe that \hat{v}_ε is also a local minimiser of $\mathcal{F}_\varepsilon(\cdot, Q_{\frac{1}{2}, \varsigma})$.

By setting $\hat{w}_\varepsilon(y) := \sqrt{\beta_\varepsilon} \hat{v}_\varepsilon(y)$, we have

$$\mathcal{F}_\varepsilon(\hat{v}_\varepsilon, Q_{\frac{1}{2}, \varsigma}) = \frac{1}{\beta_\varepsilon} \left(\int_{Q_{\frac{1}{2}, \varsigma}} |\nabla \hat{w}_\varepsilon|^2 dy + \mathcal{H}^{n-1}(S_{\hat{w}_\varepsilon} \cap Q_{\frac{1}{2}, \varsigma}) \right),$$

and

$$\mathcal{H}^{n-1}(S_{\hat{v}_\varepsilon} \cap Q_{\frac{1}{2}, \varsigma}) = \mathcal{H}^{n-1}(S_{\hat{w}_\varepsilon} \cap Q_{\frac{1}{2}, \varsigma}) \leq C \beta_\varepsilon.$$

Hence the function \hat{w}_ε is a local minimiser of the Mumford-Shah functional in $Q_{\frac{1}{2}, \varsigma}$, and its jump set has small Hausdorff measure. We then apply Theorem 6.6 to \hat{w}_ε ; namely, we fix $x_0 \in Q_{\frac{1}{2}, \varsigma}$ and consider a ball $B_{\rho(\varepsilon)}(x_0) \subset\subset Q_{\frac{1}{2}, \varsigma}$, where $\rho(\varepsilon)$ is such that

$$\mathcal{H}^{n-1}(S_{\hat{w}_\varepsilon} \cap B_{\rho(\varepsilon)}(x_0)) \leq C \beta_\varepsilon \leq \theta \rho(\varepsilon)^{n-1} \quad (6.10)$$

is satisfied for $\varepsilon > 0$ fixed, where θ is the elimination constant in Theorem 6.6 (note that $\rho(\varepsilon)$ can be very small, of the order of $(\beta_\varepsilon)^{\frac{1}{n-1}}$). Then Theorem 6.6 guarantees that

$$S_{\hat{w}_\varepsilon} \cap B_{\rho(\varepsilon)}(x_0) = \emptyset.$$

This same argument can be repeated for every $x \in Q_{\frac{1}{2}, \varsigma}$ with any radius $\rho(\varepsilon) > 0$ satisfying (6.10). In this way we conclude that the jump of \hat{w}_ε has to be contained in a neighbourhood of $\partial Q_{\frac{1}{2}, \varsigma}$ of order $(\beta_\varepsilon)^{\frac{1}{n-1}}$. In particular, there exist $\frac{1}{2} < \rho_1 < \rho_2 < \varsigma$ such that $Q_{\rho_1, \rho_2} \subset\subset Q_{\frac{1}{2}, \varsigma}$ and $S_{\hat{w}_\varepsilon} \cap \overline{Q}_{\rho_1, \rho_2} = \emptyset$.

From the definition of \hat{w}_ε we deduce that, for any $\hat{v}_\varepsilon \in \mathcal{M}_\vartheta^\varepsilon$ (and for sufficiently small ε)

$$\mathcal{H}^{n-1}(S_{\hat{v}_\varepsilon} \cap \overline{Q}_{\rho_1, \rho_2}) = 0,$$

hence $\hat{v}_\varepsilon \in H^1(Q_{\rho_1, \rho_2})$, and $S_{\hat{v}_\varepsilon} \subseteq Q_{\rho_1} \cup Q_{\rho_2, \varepsilon}$. Moreover, since the Mumford-Shah functional is invariant under translations, we can assume with no loss of generality that any local minimiser $\hat{v}_\varepsilon \in \mathcal{M}_{\hat{\vartheta}}^\varepsilon$ satisfies

$$\int_{Q_{\rho_1, \rho_2}} \hat{v}_\varepsilon dx = 0. \quad (6.11)$$

Step 2.2: Comparison between \hat{v}_ε and its harmonic extension. For a given $\hat{v}_\varepsilon \in \mathcal{M}_{\hat{\vartheta}}^\varepsilon$ we define the function $\tilde{v}_\varepsilon \in H^1(Q_{\rho_2})$ as the solution of the following Dirichlet problem for the Laplace equation

$$(\text{Dir}) \begin{cases} \Delta w = 0 & \text{in } Q_{\rho_1}, \\ w = \hat{v}_\varepsilon & \text{in } \bar{Q}_{\rho_1, \rho_2}. \end{cases}$$

Throughout this sub-step we simply write \hat{v} and \tilde{v} instead of \hat{v}_ε and \tilde{v}_ε .

We now claim that for every $\eta > 0$ there exists $\vartheta^* = \vartheta^*(\eta) > 0$ such that for every $\hat{v} \in \mathcal{M}_{\hat{\vartheta}^*}^\varepsilon$ and every corresponding \tilde{v} as in (Dir) we have

$$\int_{Q_{\rho_2}} |D\tilde{v}|^2 dx \leq (1 + \eta) \int_{Q_{\rho_2}} |\nabla \hat{v}|^2 dx. \quad (6.12)$$

We note that the claim is true for constant \tilde{v} . Therefore we only need to prove (6.12) when \tilde{v} is not constant. Arguing by contradiction we assume that there exists $\eta > 0$ such that for every $h \in \mathbb{N}$ there are $\hat{v}_h \in \mathcal{M}_{1/h}^\varepsilon$ and \tilde{v}_h defined as in (Dir) satisfying

$$\int_{Q_{\rho_2}} |D\tilde{v}_h|^2 dx > (1 + \eta) \int_{Q_{\rho_2}} |\nabla \hat{v}_h|^2 dx.$$

Using that $\hat{v}_h = \tilde{v}_h$ in Q_{ρ_1, ρ_2} , and that $\hat{v}_h \in H^1(Q_{\rho_1, \rho_2})$ by Step 2.1, the previous estimate gives

$$\int_{Q_{\rho_1}} |D\tilde{v}_h|^2 dx > (1 + \eta) \int_{Q_{\rho_1}} |\nabla \hat{v}_h|^2 dx + \eta \int_{Q_{\rho_1, \rho_2}} |D\hat{v}_h|^2 dx \quad (6.13)$$

From the normalisation condition in (6.11) and the energy bound in $(\text{LMS})_\varepsilon$ satisfied by \hat{v}_ε , we can apply the Poincaré-Wirtinger inequality to deduce that there exists a constant $C > 0$ (independent of h) such that $\|\hat{v}_h\|_{H^1(Q_{\rho_1, \rho_2})} \leq C$. Therefore \hat{v}_h converges weakly in $H^1(Q_{\rho_1, \rho_2})$, hence in particular

$$\varphi_h := (\hat{v}_h)|_{\partial Q_{\rho_1}} \rightarrow \varphi \text{ strongly in } H^{1/2}(\partial Q_{\rho_1}), \quad (6.14)$$

for some $\varphi \in H^{1/2}(\partial Q_{\rho_1})$. Moreover, $(\tilde{v}_h)|_{\partial Q_{\rho_1}} = \varphi_h$. Then, since

$$\|\tilde{v}_h\|_{H^1(Q_{\rho_1})} \leq C \|\varphi_h\|_{H^{1/2}(\partial Q_{\rho_1})},$$

we immediately deduce that \tilde{v}_h is uniformly bounded in $H^1(Q_{\rho_1})$.

We now apply Lemma 6.7 with $r = \rho_1$, $t = \frac{1}{\beta_\varepsilon}$ and the functions φ_h and φ defined in (6.14). By the fundamental theorem of Γ -convergence, the sequence \tilde{v}_h , which is a compact sequence of minimisers for the functionals I_{φ_h} , converges weakly in $H^1(Q_{\rho_1})$ to the unique minimiser of I_φ , which we denote with \tilde{v} . Furthermore we have convergence of the corresponding minimum values:

$$\begin{aligned} \lim_{h \rightarrow +\infty} \int_{Q_{\rho_1}} |D\tilde{v}_h|^2 dx &= \lim_{h \rightarrow +\infty} I_{\varphi_h}(\tilde{v}_h) = \lim_{h \rightarrow +\infty} \inf_{v \in H^1(Q_{\rho_1})} I_{\varphi_h}(v) \\ &= \min_{v \in H^1(Q_{\rho_1})} I_\varphi(v) = \int_{Q_{\rho_1}} |D\tilde{v}|^2 dx. \end{aligned}$$

Similarly, the sequence $\hat{v}_h \in \mathcal{M}_{1/h}^\varepsilon$, which is a compact sequence of minimisers of the functionals $I_{\varphi_h}^{h,t}$, converges in $L^1(Q_{\rho_1})$ to the unique minimiser of I_φ ; *i.e.*, to \tilde{v} . Furthermore we have convergence of the minimum values:

$$\lim_{h \rightarrow +\infty} I_{\varphi_h}^{h,t}(\hat{v}_h) = \lim_{h \rightarrow +\infty} \inf_{v \in \mathcal{M}_{1/h}^\varepsilon} I_{\varphi_h}^{h,t}(v) = \min_{v \in H^1(Q_{\rho_1})} I_\varphi(v) = \int_{Q_{\rho_1}} |D\tilde{v}|^2 dx.$$

On the other hand, we clearly have, by the definition of good cubes (6.8), with $\vartheta = \frac{1}{h}$, that

$$\lim_{h \rightarrow +\infty} I_{\varphi_h}^{h,t}(\hat{v}_h) = \lim_{h \rightarrow +\infty} \int_{Q_{\rho_1}} |\nabla \hat{v}_h|^2 dx,$$

and hence

$$\lim_{h \rightarrow +\infty} \int_{Q_{\rho_1}} |\nabla \hat{v}_h|^2 dx = \int_{Q_{\rho_1}} |D\tilde{v}|^2 dx.$$

By passing to the limit in (6.13) we then have in particular that

$$(1 + \eta) \int_{Q_{\rho_1}} |D\tilde{v}|^2 dx \leq \int_{Q_{\rho_1}} |D\tilde{v}|^2 dx,$$

which gives a contradiction since $D\tilde{u} \not\equiv 0$ and $\eta > 0$, and thus proves (6.12).

In view of (6.12), in what follows we will choose $\vartheta = \vartheta^*$ in the definition of good cubes (6.8).

Step 2.3: Energy bound on the good cubes. Let ϑ^* be as in Step 2.2; we consider the following minimisation problems

$$(MS)_\varepsilon \min \left\{ \mathcal{F}_\varepsilon(v, Q_{\rho_2}) : v \in SBV^2(Q_{\rho_2}), S_v \subset Q_{\rho_1}, \mathcal{H}^{n-1}(S_v \cap Q_{\rho_2}) \leq \vartheta^*, \right. \\ \left. \mathcal{F}_\varepsilon(v, Q_{\rho_2}) \leq C \text{ for some } C > 0, v = v_\varepsilon \text{ on } \partial Q_{\rho_2} \right\},$$

where $\frac{1}{2} < \rho_1 < \rho_2 < 1$. For a minimiser \hat{v}_ε of $(MS)_\varepsilon$, let \tilde{v}_ε be the corresponding function as in (Dir). Since \hat{v}_ε is also a local minimiser of the same functional, from (6.12) we have

$$\int_{Q_{\rho_2}} |D\tilde{v}_\varepsilon|^2 dx \leq (1 + \eta) \int_{Q_{\rho_2}} |\nabla \hat{v}_\varepsilon|^2 dx.$$

Then for the sequence (v_ε) defined at the beginning of Step 1 we have

$$\begin{aligned} \mathcal{F}(v_\varepsilon, Q_{\rho_2}) &= \int_{Q_{\rho_2}} |\nabla v_\varepsilon|^2 dy + \frac{1}{\beta_\varepsilon} \mathcal{H}^{n-1}(S_{v_\varepsilon} \cap Q_{\frac{1}{2}, \rho_2}) + \mathcal{H}^{n-1}(S_{v_\varepsilon} \cap Q_{\frac{1}{2}}) \\ &\geq \int_{Q_{\rho_2}} |\nabla \hat{v}_\varepsilon|^2 dy + \frac{1}{\beta_\varepsilon} \mathcal{H}^{n-1}(S_{\hat{v}_\varepsilon} \cap Q_{\frac{1}{2}, \rho_2}) + \mathcal{H}^{n-1}(S_{\hat{v}_\varepsilon} \cap Q_{\frac{1}{2}}) \\ &\geq \left(1 - \frac{\eta}{1 + \eta}\right) \int_{Q_{\rho_2}} |D\tilde{v}_\varepsilon|^2 dy. \end{aligned}$$

By the change of variables $\tilde{u}_\varepsilon(\varepsilon y) := \sqrt{\beta_\varepsilon} \varepsilon \tilde{v}_\varepsilon(y)$, we get

$$\begin{aligned} F(u_\varepsilon, Q_{\rho_2}^\varepsilon) &= \int_{Q_{\rho_2}^\varepsilon} |\nabla u_\varepsilon|^2 dx + \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap Q_{\frac{1}{2}, \rho_2}^\varepsilon) + \beta_\varepsilon \mathcal{H}^{n-1}(S_{u_\varepsilon} \cap Q_{\frac{1}{2}}^\varepsilon) \\ &\geq \left(1 - \frac{\eta}{1 + \eta}\right) \int_{Q_{\rho_2}^\varepsilon} |D\tilde{u}_\varepsilon|^2 dx. \end{aligned}$$

Hence, for every $k \in \mathcal{I}_\varepsilon^{1,g}$,

$$F_\varepsilon(u_\varepsilon, Q_{\rho_2}^{k,\varepsilon}) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \int_{Q_{\rho_2}^{k,\varepsilon}} |D\tilde{u}_{k,\varepsilon}|^2 dx, \quad (6.15)$$

where the subscript k has now been added to highlight the dependence of the construction of $\tilde{u}_{k,\varepsilon}$ on the cube $Q^{k,\varepsilon}$.

Step 3: Energy estimate on the bad cubes and on the boundary cubes. Let now $k \in \mathcal{I}_\varepsilon^{1,b} \cup \mathcal{I}_\varepsilon^2$. We bound the energy of u_ε on $Q^{k,\varepsilon} \cap \Omega$ as

$$F_\varepsilon(u_\varepsilon, Q^{k,\varepsilon} \cap \Omega) \geq \widehat{F}_\varepsilon(\hat{u}_{k,\varepsilon}, Q^{k,\varepsilon} \cap \Omega), \quad (6.16)$$

where \widehat{F}_ε is defined as in (2.4) and $\hat{u}_{k,\varepsilon} : Q^{k,\varepsilon} \cap \Omega \rightarrow \mathbb{R}$ is defined as

$$\hat{u}_{k,\varepsilon} := T_\varepsilon((u_\varepsilon)|_{(Q^{k,\varepsilon} \cap \Omega) \cap P}), \quad (6.17)$$

where T_ε denotes the extension operator provided by [12, Theorem 1.1].

Step 4: Construction of an improved sequence converging to u_ξ . We define the sequence $w_\varepsilon : \Omega \rightarrow \mathbb{R}$ as

$$w_\varepsilon := \begin{cases} \tilde{u}_{k,\varepsilon} & \text{in } Q_{\rho_2}^{k,\varepsilon}, k \in \mathcal{I}_\varepsilon^{1,g}, \\ u_\varepsilon & \text{in } Q_{\rho_2,1}^{k,\varepsilon}, k \in \mathcal{I}_\varepsilon^{1,g}, \\ \hat{u}_{k,\varepsilon} & \text{in } Q^{k,\varepsilon} \cap \Omega, k \in \mathcal{I}_\varepsilon^{1,b} \cup \mathcal{I}_\varepsilon^2. \end{cases}$$

Clearly $w_\varepsilon \in SBV^2(\Omega)$; we now show that w_ε converges to u_ξ strongly in $L^1(\Omega)$.

As a first step we show that $\sup_\varepsilon MS(w_\varepsilon) < +\infty$.

From (6.15) applied to the good cubes, we have that for every $\varepsilon > 0$

$$MS\left(w_\varepsilon, \bigcup_{k \in \mathcal{I}_\varepsilon^{1,g}} Q^{k,\varepsilon}\right) = \sum_{k \in \mathcal{I}_\varepsilon^{1,g}} \left(\int_{Q_{\rho_2}^{k,\varepsilon}} |D\tilde{u}_{k,\varepsilon}|^2 dx + MS(u_\varepsilon, Q_{\rho_2,1}^{k,\varepsilon}) \right) \leq (2 + \eta)F_\varepsilon(u_\varepsilon).$$

For the bad cubes, by (6.17) and (6.9) we have, for $\varepsilon > 0$,

$$MS\left(w_\varepsilon, \bigcup_{k \in \mathcal{I}_\varepsilon^{1,b}} Q^{k,\varepsilon}\right) = \sum_{k \in \mathcal{I}_\varepsilon^{1,b}} MS(\hat{u}_{k,\varepsilon}, Q^{k,\varepsilon}) \leq C \sum_{k \in \mathcal{I}_\varepsilon^{1,b}} MS(u_\varepsilon, Q_{\frac{1}{2},1}^{k,\varepsilon}) \leq F_\varepsilon(u_\varepsilon),$$

where C_n is the perimeter of $Q \setminus P$ in Q . Similarly, for the boundary cubes we have

$$MS\left(w_\varepsilon, \bigcup_{k \in \mathcal{I}_\varepsilon^2} (Q^{k,\varepsilon} \cap \Omega)\right) \leq \sum_{k \in \mathcal{I}_\varepsilon^2} MS(u_\varepsilon, Q_{\frac{1}{2},1}^{k,\varepsilon} \cap \Omega) + \mathcal{H}^{n-1}(\partial\Omega) \leq F_\varepsilon(u_\varepsilon) + C.$$

Eventually, since $F_\varepsilon(u_\varepsilon) \leq C$ for every $\varepsilon > 0$, we get the desired uniform bound on $MS(w_\varepsilon)$.

Moreover, since $\|u_\varepsilon\|_{L^\infty(\Omega)} \leq \|u_\xi\|_{L^\infty(\Omega)}$ the function w_ε above can be constructed in a way such that $\|w_\varepsilon\|_{L^\infty(\Omega)} \leq \|u_\xi\|_{L^\infty(\Omega)}$. Hence, we can apply the compactness result [1, Theorem 4.8] to the sequence (w_ε) to deduce the existence of $w \in SBV^2(\Omega)$ such that (up to a subsequence not relabelled) w_ε converges to w weakly* in $BV(\Omega)$ and hence strongly in $L^1(\Omega)$.

It remains to show that $w = u_\xi$. We observe that

$$w_\varepsilon = u_\varepsilon \text{ in } A_{\rho_2}^\varepsilon := \bigcup_{k \in \mathcal{I}_\varepsilon^{1,b} \cup \mathcal{I}_\varepsilon^2} Q_{\rho_2,1}^{k,\varepsilon} \cap \Omega,$$

and $\chi_{A_{\rho_2}^\varepsilon} \rightarrow C(\rho_2)$ weakly* in $L^\infty(\Omega)$, for some constant $0 < C(\rho_2) < 1$. Moreover by assumption $u_\varepsilon \rightarrow u_\xi$ in the sense of Definition 3.1 so that there exists a sequence $(\tilde{u}_\varepsilon) \subset L^1(\Omega)$ such that $\tilde{u}_\varepsilon = u_\varepsilon$ in $\Omega \cap \varepsilon P$, and \tilde{u}_ε converges to u_ξ strongly in $L^1(\Omega)$. Hence, since $A_{\rho_2}^\varepsilon \subset \Omega \cap \varepsilon P$,

$$0 = \int_{A_{\rho_2}^\varepsilon} |\tilde{u}_\varepsilon - w_\varepsilon| dx = \int_\Omega |\tilde{u}_\varepsilon - w_\varepsilon| \chi_{A_{\rho_2}^\varepsilon} dx \rightarrow C(\rho_2) \int_\Omega |u_\xi - w| dx,$$

as $\varepsilon \rightarrow 0$. Since $C(\rho_2) > 0$ then necessarily $w = u_\xi$.

Step 5: Energy estimate for w_ε . First of all, from (6.15) and (6.16) we have

$$F_\varepsilon(u_\varepsilon) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \int_\Omega \phi_\varepsilon(x) |\nabla w_\varepsilon|^2 dx, \quad (6.18)$$

where

$$\phi_\varepsilon(x) := \begin{cases} 0 & \text{if } x \in Q_{\frac{1}{2}}^{k,\varepsilon} \cap \Omega, k \in \mathcal{I}_\varepsilon^{1,b} \cup \mathcal{I}_\varepsilon^2, \\ 1 & \text{otherwise in } \Omega. \end{cases}$$

Note that $\phi_\varepsilon \rightarrow 1$ in measure as $\varepsilon \rightarrow 0$, by (6.9) and since the cardinality of $\mathcal{I}_\varepsilon^2$ is of order ε^{1-n} .

By the previous step and by [1, Theorem 4.7] we have

$$\liminf_{\varepsilon \rightarrow 0} \mathcal{H}^{n-1}(S_{w_\varepsilon} \cap \Omega) \geq \mathcal{H}^{n-1}(S_{u_\xi} \cap \Omega) = 0,$$

and since ∇w_ε converges to ξ weakly in $L^2(\Omega)$, and $\phi_\varepsilon \rightarrow 1$ in measure, we conclude that

$$\liminf_{\varepsilon \rightarrow 0} \int_{\Omega} \phi_\varepsilon(x) |\nabla w_\varepsilon|^2 dx \geq \int_{\Omega} |Du_\xi|^2 dx = \mathcal{L}^n(\Omega) |\xi|^2. \quad (6.19)$$

Eventually passing to the liminf in (6.18) and appealing to (6.19) gives

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \mathcal{L}^n(\Omega) |\xi|^2.$$

Finally, by letting $\eta \rightarrow 0^+$ we deduce that

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \geq \mathcal{L}^n(\Omega) |\xi|^2,$$

which concludes the proof. \square

We are now ready to prove Theorem 6.4.

Proof of Theorem 6.4. Lemma 6.1 gives $f_{\text{hom}}^\infty(\xi) \leq |\xi|^2$ for every $\xi \in \mathbb{R}^n$, hence it only remains to prove the opposite inequality. By Γ -convergence we have that there exists a sequence (u_ε) converging to u_ξ in the sense of Definition 3.1 such that

$$\mathcal{L}^n(\Omega) f_{\text{hom}}^\infty(\xi) = F_{\text{hom}}^\infty(u_\xi) = \lim_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon),$$

hence the desired inequality immediately follows from (6.7). \square

6.3. Critical case: $\ell \in (0, +\infty)$. We start by proving a simple result, which is essentially a corollary of Lemma 6.1. Then the main result of this section is Corollary 6.11 which asserts that the homogenized volume integrand f_{hom}^ℓ is *not* 2-homogeneous, unlike the extreme cases $\ell = 0$ and $\ell = +\infty$, and unlike the volume integrand of the functionals F_ε .

Lemma 6.9. *Let $\ell \in (0, +\infty)$, and let f_{hom}^ℓ be as in Theorem 4.6. Then f_{hom}^ℓ is 2-homogeneous if and only if $f_{\text{hom}}^\ell(\xi) = \hat{f}(\xi)$ for every $\xi \in \mathbb{R}^n$.*

Proof. Assume that f_{hom}^ℓ is 2-homogeneous. Replacing ξ by $\lambda\xi$ in (6.1), with $\lambda \neq 0$, gives

$$\lambda^2 \hat{f}(\xi) \leq \lambda^2 f_{\text{hom}}^\ell(\xi) \leq \min \left\{ \lambda^2 |\xi|^2, \lambda^2 \hat{f}(\xi) + C\ell \right\},$$

which can be rewritten as

$$\hat{f}(\xi) \leq f_{\text{hom}}^\ell(\xi) \leq \min \left\{ |\xi|^2, \hat{f}(\xi) + \frac{C\ell}{\lambda^2} \right\}.$$

By letting $|\lambda| \rightarrow +\infty$, we have

$$f_{\text{hom}}^\ell(\xi) = \hat{f}(\xi),$$

where we have used the obvious bound $\hat{f}(\xi) \leq |\xi|^2$. \square

In the following result we assume that $\beta_\varepsilon = \varepsilon$ for convenience.

Proposition 6.10. *For every $\xi \in \mathbb{R}^n \setminus \{0\}$, and every $\ell \in (0, +\infty)$, we have*

$$f_{\text{hom}}^\ell(\xi) \neq \hat{f}(\xi).$$

Proof. Clearly the statement reduces to proving that $\hat{f}(\xi) < f_{\text{hom}}^\ell(\xi)$ for every $\xi \in \mathbb{R}^n \setminus \{0\}$.

We first note that from the definition of \hat{f} in (2.6) we have that, if $\xi \neq 0$,

$$\hat{f}(\xi) \leq \int_{Q_{\frac{1}{2}, 1}} |\xi|^2 dx < \int_Q |\xi|^2 dx = |\xi|^2.$$

Hence

$$\hat{f}(\xi) < |\xi|^2 \quad \text{for every } \xi \in \mathbb{R}^n \setminus \{0\}. \quad (6.20)$$

To prove the claim, it is enough to show that for every $\xi \neq 0$ and for every admissible sequence u_ε which converges to u_ξ in the sense of Definition 3.1 we have

$$\hat{f}(\xi) < \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon, Q). \quad (6.21)$$

Indeed, if the statement (6.21) is proven, then we can choose u_ε to be the recovery sequence of F_ε for u_ξ and deduce, from the Γ -convergence of F_ε to F_{hom}^ℓ for $0 < \ell < +\infty$, that for

$$\hat{f}(\xi) < \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon, Q) = f_{\text{hom}}^\ell(\xi).$$

We can assume $\sup_\varepsilon F_\varepsilon(u_\varepsilon, Q) < +\infty$, otherwise there is nothing to prove. We can also assume that $\|u_\varepsilon\|_{L^\infty(Q)} \leq \|u_\xi\|_{L^\infty(Q)}$.

Proceeding as in the proof of Theorem 6.8, we fix $\vartheta > 0$ independent of ε and introduce a classification of the cubes of the form $Q^{k,\varepsilon} := \varepsilon(Q + k)$, with $k \in \mathbb{Z}^n$, for the cubes well contained in Q , as follows. We call a cube $Q^{k,\varepsilon}$ undamaged if it satisfies

$$\mathcal{H}^{n-1}(S_{u_\varepsilon} \cap Q^{k,\varepsilon}) \leq \vartheta \varepsilon^{n-1},$$

and damaged otherwise, namely if

$$\mathcal{H}^{n-1}(S_{u_\varepsilon} \cap Q^{k,\varepsilon}) > \vartheta \varepsilon^{n-1}.$$

Let $\mathcal{I}_\varepsilon^d$ be the set of damaged or boundary cubes, and let $N_d(\varepsilon)$ denote its cardinality.

Again, as in the proof of Theorem 6.8, we can construct an improved sequence $(w^\varepsilon) \subset L^1(Q)$ such that w_ε converges to u_ξ weakly in $BV(Q)$, and

$$F_\varepsilon(u_\varepsilon) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \int_Q \phi_\varepsilon(x) |\nabla w_\varepsilon|^2 dx + \varepsilon N_d(\varepsilon) \vartheta \varepsilon^{n-1}, \quad (6.22)$$

where

$$\phi_\varepsilon(x) := \begin{cases} 0 & \text{if } x \in Q^{\frac{k,\varepsilon}{2}} \cap \Omega, \ k \in \mathcal{I}_\varepsilon^d, \\ 1 & \text{otherwise in } \Omega. \end{cases} \quad (6.23)$$

Note that $\|\phi_\varepsilon - 1\|_{L^1(Q)} \leq c N_d(\varepsilon) \varepsilon^n$.

We consider two different cases.

If the number of damaged cubes is small; *i.e.*,

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon^n N_d(\varepsilon) = 0,$$

since $\phi_\varepsilon \rightarrow 1$ in measure and $\nabla w_\varepsilon \rightharpoonup u_\xi$ weakly in $L^2(Q)$, by taking the liminf in (6.22) we obtain directly that

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \geq \left(1 - \frac{\eta}{1 + \eta}\right) |\xi|^2,$$

which by (6.20) and by the arbitrariness of $\eta > 0$ implies (6.21).

On the other hand, if the number of damaged cubes is large; *i.e.*,

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon^n N_d(\varepsilon) = C > 0,$$

then there exists an infinitesimal subsequence (ε_k) such that

$$\lim_{k \rightarrow +\infty} \varepsilon_k^n N_d(\varepsilon_k) = \limsup_{\varepsilon \rightarrow 0} \varepsilon^n N_d(\varepsilon) = C.$$

From (6.22) we deduce that

$$F_{\varepsilon_k}(u_{\varepsilon_k}) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \widehat{F}_{\varepsilon_k}(w_{\varepsilon_k}) + \varepsilon_k N_d(\varepsilon_k) \vartheta \varepsilon_k^{n-1},$$

which, by the Γ -convergence of \widehat{F}_ε to \widehat{F} , implies that

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) &\geq \limsup_{k \rightarrow \infty} F_{\varepsilon_k}(u_{\varepsilon_k}) \geq \liminf_{k \rightarrow \infty} F_{\varepsilon_k}(u_{\varepsilon_k}) \\ &\geq \left(1 - \frac{\eta}{1 + \eta}\right) \liminf_{k \rightarrow \infty} \widehat{F}_{\varepsilon_k}(w_{\varepsilon_k}) + \vartheta C \geq \left(1 - \frac{\eta}{1 + \eta}\right) \widehat{f}(\xi) + \vartheta C \end{aligned}$$

and hence the claim (6.21), by the arbitrariness of $\eta > 0$. This concludes the proof. \square

A direct consequence of the previous results is that f_{hom}^ℓ is not 2-homogeneous.

Corollary 6.11 (f_{hom}^ℓ is not 2-homogeneous). *Let $\ell \in (0, +\infty)$; then the function f_{hom}^ℓ is not 2-homogeneous.*

Proof. The conclusion is a straightforward consequence of Lemma 6.9 and Proposition 6.10. \square

We can also say something about the behaviour of f_{hom}^ℓ close to zero.

Corollary 6.12 ($f_{\text{hom}}^\ell = |\xi|^2$ close to zero). *Let $\ell \in (0, \infty)$; then there exists a constant $\gamma_0 > 0$ such that $f_{\text{hom}}^\ell(\xi) = |\xi|^2$ for every $|\xi| \leq \gamma_0$.*

Proof. Let $\xi \in \mathbb{R}$, let u_ε be a recovery sequence for F_ε , converging to u_ξ in the sense of Definition 3.1, and let w_ε be the improved sequence satisfying

$$F_\varepsilon(u_\varepsilon) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \int_Q \phi_\varepsilon(x) |\nabla w_\varepsilon|^2 dx + \varepsilon N_d(\varepsilon) \vartheta \varepsilon^{n-1},$$

with ϕ_ε as in (6.23). Clearly

$$|\xi|^2 \geq f_{\text{hom}}^\ell(\xi) \geq \left(1 - \frac{\eta}{1 + \eta}\right) \liminf_{\varepsilon \rightarrow 0} \int_Q \phi_\varepsilon(x) |\nabla w_\varepsilon|^2 dx + \liminf_{\varepsilon \rightarrow 0} \varepsilon N_d(\varepsilon) \vartheta \varepsilon^{n-1}. \quad (6.24)$$

Note that, if ξ is small, then the number of damaged cubes for w_ε has to be small. If not, then from (6.24) and by the arbitrariness of $\eta > 0$ we would deduce that

$$|\xi|^2 \geq f_{\text{hom}}^\ell(\xi) \geq \widehat{f}(\xi) + C\vartheta > |\xi|^2,$$

where the last inequality follows by the smallness of ξ , since $C > 0$, and this would lead to a contradiction. Hence there exists $\gamma_0 > 0$ such that whenever $|\xi| < \gamma_0$, the improved sequence w_ε has only a negligible number of damaged cubes, in which case, from (6.24), we would have that

$$|\xi|^2 \geq f_{\text{hom}}^\ell(\xi) \geq \left(1 - \frac{\eta}{1 + \eta}\right) |\xi|^2,$$

and so the claim. \square

Remark 6.13 (Asymptotic behaviour of F_{hom}^ℓ at infinity). From Lemma 6.1 we know that for every $\xi \in \mathbb{R}^n$ and every $\ell \in (0, +\infty)$

$$\widehat{f}(\xi) \leq f_{\text{hom}}^\ell(\xi) \leq \min\{|\xi|^2, \widehat{f}(\xi) + C\ell\}, \quad (6.25)$$

where $C > 0$ is a constant depending only on P and on Ω . Note that for $|\xi|$ sufficiently large (6.25) simplifies to

$$\widehat{f}(\xi) \leq f_{\text{hom}}^\ell(\xi) \leq \widehat{f}(\xi) + C\ell,$$

which implies that, for $|\xi|$ large, $|f_{\text{hom}}^\ell(\xi) - \widehat{f}(\xi)| \leq C$, uniformly in ξ . In particular, if we divide by $\widehat{f}(\xi)$ the previous estimate and let $|\xi| \rightarrow +\infty$ we deduce the limit behaviour

$$\lim_{|\xi| \rightarrow +\infty} \frac{f_{\text{hom}}^\ell(\xi)}{\widehat{f}(\xi)} = 1.$$

ACKNOWLEDGMENTS

The authors wish to thank Gianni Dal Maso and Massimiliano Morini for stimulating discussions. L. Scardia acknowledges support by the EPSRC under the Grant EP/N035631/1 ‘‘Dislocation patterns beyond optimality’’.

REFERENCES

- [1] Ambrosio L., Fusco N., and Pallara D.: Functions of bounded variations and free discontinuity problems. Clarendon Press, Oxford, 2000.
- [2] Babych N.O., Kamotski I.V., and Smyshlyaev V.P.: Homogenization in periodic media with doubly high contrasts. *Netw. Heterog. Media*, **3** (3) (2008), 413–436.
- [3] Barchiesi M.: Toughening by Crack Deflection in the Homogenization of Brittle Composites with Soft Inclusions. *Arch. Ration. Mech. Anal.* **227** (2018), 749–766.
- [4] Barchiesi M., and Focardi M.: Homogenization of the Neumann problem in perforated domains: an alternative approach. *Calc. Var. Partial Differential Equations*, **42** (2011), 257–288.
- [5] Barchiesi M., Lazzaroni G., and Zeppieri C.I. A bridging mechanism in the homogenization of brittle composites with soft inclusions, *SIAM J. Math. Anal.* **48** (2016), no. 2, 1178–1209.
- [6] Bouchitté G., Fonseca I., Leoni G., and Mascarenhas L.: A global method for relaxation in $W^{1,p}$ and in $SBVP$. *Arch. Ration. Mech. Anal.* **165** (2002), 187–242.
- [7] Braides A.: A handbook of Γ -convergence. In *Handbook of Differential Equations. Stationary Partial Differential Equations*, Volume 3 (M. Chipot and P. Quittner, eds.), Elsevier, 2006.
- [8] Braides A., Chiadò Piat V., and Piatnitski A.: Homogenization of discrete high-contrast energies. *SIAM J. Math. Anal.* **47** (2015), no. 4, 3064–3091.
- [9] Braides A., Defranceschi A., and Vitali E.: Homogenization of free discontinuity problems. *Arch. Ration. Mech. Anal.* **135** (1996), 297–356.
- [10] Braides A., and Garroni A.: Homogenization of periodic nonlinear media with stiff and soft inclusions. *Math. Models Methods Appl. Sci.* **05** (1995), 543–564.
- [11] Braides A., and Solci M.: Multi-scale free-discontinuity problems with soft inclusions. *Boll. Unione Mat. Ital.* (9) **6** (2013), no. 1, 29–51.
- [12] Cagnetti F., and Scardia L.: An extension theorem in SBV and an application to the homogenization of the Mumford-Shah functional in perforated domains. *J. Math. Pures Appl.*, **95** (2011), 349–381.
- [13] Cagnetti F., Dal Maso G., Scardia L., and Zeppieri C.I.: Γ -convergence of free-discontinuity problems, *Arxiv preprint arXiv:1712.07093*.
- [14] Cherednichenko K. B., and Cherdantsev M.: Two-scale Γ -convergence of integral functionals and its application to homogenization of nonlinear high-contrast periodic composites. *Arch. Ration. Mech. Anal.* **204** (2012), 445–478.
- [15] Cherednichenko K. B., and Cooper S.: Homogenization of the system of high-contrast Maxwell equations. *Mathematika*, **61** (2015), 475–500.
- [16] Congedo G., and Tamanini I.: Density theorems for local minimizers of area-type functionals. *Rend. Sem. Mat. Univ. Padova*, **85** (1991), 217–248.
- [17] Congedo G., and Tamanini I.: On the existence of solutions to a problem in multidimensional segmentation, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **8** (1991), no. 2, 175–195.
- [18] Dal Maso G.: An introduction to Γ -convergence. Birkhäuser, Boston, 1993.
- [19] Dal Maso G., Morel J.M., and Solimini S.: A variational method in image segmentation: Existence and approximation results, *Acta Mathematica*, **168** (1992), 89–151.
- [20] Dal Maso G., and Zeppieri C.I.: Homogenization of fiber reinforced brittle materials: the intermediate case. *Adv. Calc. Var.* **3** (2010), no. 4, 345–370.
- [21] De Giorgi E., Carriero M., and Leaci A.: Existence theorem for a minimum problem with free discontinuity set, *Arch. Ration. Mech. Anal.* **108** (1989), 195–218.
- [22] Focardi M., Gelli M.S., and Ponsiglione M.: Fracture mechanics in perforated domains: a variational model for brittle porous media. *Math. Models Methods Appl. Sci.* **19** (2009), 2065–2100.
- [23] Scardia L.: Damage as Γ -limit of microfractures in anti-plane linearized elasticity. *Math. Models Methods Appl. Sci.*, **18/10** (2008), 1703–1740.
- [24] Scardia L.: Damage as the Γ -limit of microfractures in linearized elasticity under the non-interpenetration constraint. *Adv. Calc. Var.*, **3** (2010), 423–458.

(X. Pellet and L. Scardia) DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH, UNITED KINGDOM

E-mail address: X.P.J.Pellet@bath.ac.uk

E-mail address: L.Scardia@bath.ac.uk

(C. I. Zeppieri) ANGEWANDTE MATHEMATIK, WWU MÜNSTER, GERMANY

E-mail address: caterina.zeppieri@uni-muenster.de