

STABLE HYPERSURFACES IN THE COMPLEX PROJECTIVE SPACE

ERIKA BATTAGLIA, ROBERTO MONTI, AND ALBERTO RIGHINI

ABSTRACT. We characterize the sphere with radius $\tan^2 r = 2n + 1$ in the complex projective space \mathbf{CP}^n as the unique stable hypersurface subject to certain bounds on the curvatures.

1. INTRODUCTION

In [BdCE88] Barbosa, Do Carmo and Eschenburg proved among other results that the geodesic sphere with radius $r \in (0, \pi/2)$ in the complex projective space \mathbf{CP}^n , $n \geq 2$, is stable for the area functional with fixed enclosed volume if and only if $\tan^2 r \leq 2n + 1$. They also computed the stability-intervals for the radius of a geodesic tube around $\mathbf{CP}^k \subset \mathbf{CP}^n$, with $1 \leq k < n$.

The classification of complete oriented stable hypersurfaces in \mathbf{CP}^n could be an important step towards the classification of isoperimetric sets in \mathbf{CP}^n . Indeed, the boundary of an isoperimetric set, if smooth, is a hypersurface with constant mean curvature that is stable for variations fixing the volume. Since [BdCE88], there was apparently no progress on the problem of the classification of stable hypersurfaces in \mathbf{CP}^n .

In this paper, we characterize the geodesic sphere with radius $\tan^2 r = 2n + 1$ as the unique stable connected and complete hypersurface subject to a bound either on the characteristic curvature or on the restriction of the second fundamental form to the complex tangent space. See Definition 6.1 for the precise definition of stability.

The characteristic curvature κ of a hypersurface $\Sigma \subset \mathbf{CP}^n$ is the curvature in direction JN , where N is the normal to Σ and J is the complex structure of \mathbf{CP}^n , i.e., $\kappa = h(JN, JN)$ where h is the second fundamental form of Σ .

For fixed $H \in \mathbb{R}$ and $n \in \mathbb{N}$, let $p(\cdot; H, n)$ be the quadratic polynomial of the real variable $t \in \mathbb{R}$

$$p(t; H, n) = (2n + 1)t^2 - 2Ht - H^2 - 4(n^2 - 1). \quad (1.1)$$

Our first result is the following theorem.

2010 *Mathematics Subject Classification.* 49Q10, 53C42.

Key words and phrases. Stable hypersurfaces in complex projective space.

Theorem 1.1. *Let $\Sigma \subset \mathbf{CP}^n$, $n \geq 2$, be a complete connected stable hypersurface with constant $H = \text{tr}(h)$. If the characteristic curvature κ of Σ satisfies $p(\kappa; H, n) \geq 0$ then Σ is a geodesic sphere of radius $r > 0$ with $\tan^2 r = 2n + 1$.*

Let $\mathbf{CT}\Sigma$ be the complex tangent space of a hypersurface Σ , let T_1 and T_2 be nonzero subbundles of $\mathbf{CT}\Sigma$ such that

$$\mathbf{CT}\Sigma = T_1 \oplus T_2, \quad (1.2)$$

and denote their dimensions by $\alpha = \dim_{\mathbb{R}}(T_1)$ and $\beta = \dim_{\mathbb{R}}(T_2)$. Then we have $\alpha + \beta = \dim_{\mathbb{R}}(\mathbf{CT}\Sigma) = 2(n - 1)$ and $1 \leq \alpha < 2(n - 1)$. Let h_1 and h_2 be the restrictions of the second fundamental form h of Σ to T_1 and T_2 , and denote by $H_1 = \text{tr}(h_1)$ and $H_2 = \text{tr}(h_2)$ their respective traces.

For fixed $H \in \mathbb{R}$, $n \in \mathbb{N}$ with $n \geq 2$, and $1 \leq \alpha < 2(n - 1)$ let $p(\cdot; H, n, \alpha)$ be the quadratic convex polynomial of the variables $(s, t) \in \mathbb{R}^2$

$$p(s, t; H, n, \alpha) = \frac{s^2}{\alpha} + \frac{t^2}{\beta} + (s + t - H)^2 + \frac{(s + t - 2H)^2}{2(n^2 - 1)} - \frac{H^2}{n - 1} - 2n. \quad (1.3)$$

Our second result is a refined version of Theorem 1.1.

Theorem 1.2. *Let $\Sigma \subset \mathbf{CP}^n$, $n \geq 3$, be a complete connected stable hypersurface with constant $H = \text{tr}(h)$. If for the decomposition (1.2), with $1 \leq \alpha = \dim_{\mathbb{R}}(T_1) < 2(n - 1)$, we have $p(H_1, H_2; H, n, \alpha) \geq 0$ then Σ is a geodesic sphere of radius $r > 0$ with $\tan^2 r = 2n + 1$.*

Both Theorems 1.1 and 1.2 are a consequence of the following geometric inequality that is implied by stability. Let $\nabla_{JN}^{\Sigma} N \in T\Sigma$ be the covariant derivative of the normal N to Σ in the direction JN . We denote by $h_N \in \mathbf{CT}\Sigma$ the projection of $\nabla_{JN}^{\Sigma} N$ onto $\mathbf{CT}\Sigma$. By $|h|^2$ we denote the squared norm of h .

Theorem 1.3. *Let $\Sigma \subset \mathbf{CP}^n$, $n \geq 2$, be a complete stable hypersurface with constant $H = \text{tr}(h)$. Then we have*

$$\int_{\Sigma} \left\{ |h|^2 + \frac{(H + \kappa)^2 + |h_N|^2}{2(n^2 - 1)} - \frac{H^2}{n + 1} - 2n \right\} d\mu \leq 0, \quad (1.4)$$

where μ is the Riemannian hypersurface measure.

The method for obtaining formula (1.4) starts from an idea contained in the proof of [BdC84] that geodesic spheres are the unique stable complete hypersurfaces in the standard sphere. Our first step is the isometric embedding of \mathbf{CP}^n into H^{n+1} , the space of $(n + 1) \times (n + 1)$ Hermitian matrices, see [Ros83] and [Tai68]. Once the hypersurface Σ is embedded in H^{n+1} , we can consider the *position matrix* $A \in \Sigma$ and compute its tangential Laplacian, see Theorem 4.1,

$$\Delta A = \text{tr}(\sigma) - \sigma(N, N) - \text{tr}(h)N, \quad (1.5)$$

where Δ is the Laplace-Beltrami operator of Σ and $\sigma : T_A \mathbf{CP}^n \times T_A \mathbf{CP}^n \rightarrow T_A^\perp \mathbf{CP}^n$ is the second fundamental form of the immersion $\mathbf{CP}^n \subset H^{n+1}$. We shall make systematic use of the geometric formulas concerning σ proved by Ros in [Ros83] and [Ros84]. They are reviewed in Section 3.

For a smooth function $u : \Sigma \rightarrow \mathbb{R}$, the second variation of the area functional in the normal direction uN is given by the formula

$$\mathcal{A}''(u) = - \int_{\Sigma} u \mathcal{L}u \, d\mu,$$

where $\mathcal{L}u = \Delta u + (|h|^2 + \text{Ric}(N))u$ is the Jacobi operator. When u has zero mean, the deformation of Σ encloses a region with the same volume as Σ , at the infinitesimal level.

For any fixed $V \in H^{n+1}$, the function $u_V = \langle \Delta A, V \rangle$ has zero mean. In Section 6, we compute the trace of the quadratic form Q_{Σ} on H^{n+1} defined by $Q_{\Sigma}(V) = \mathcal{A}''(u_V)$. If Σ is stable, this trace is nonnegative and this fact is precisely inequality (1.4).

From (1.5), it is clear that in the computation for $\mathcal{L}u_V$ we need geometric formulas for $\Delta \text{tr}(\sigma)$, $\Delta \sigma(N, N)$, and ΔN . The computation for ΔN is done in Theorem 4.2. The formula for $\Delta \text{tr}(\sigma)$ follows easily from the formula for the trace of σ , see (3.18). The difficult task is to compute the tangential Laplacian of $\sigma(N, N)$. This is done in Section 5 and the resulting formula is in (5.4). In Section 6, we collect all these preliminary computations and we finish the proof of Theorem 1.3.

The proof of Theorem 1.1 now follows from Theorem 1.3 using Takagi's and Cecil-Ryan's characterization of the sphere in \mathbf{CP}^n as the unique hypersurface having precisely two (constant) different curvatures. The proof of Theorem 1.2 uses Takagi's classification of hypersurfaces in \mathbf{CP}^n , $n \geq 3$, having precisely three different constant curvatures: they are either geodesics tubes around \mathbf{CP}^k for some $k = 1, \dots, n-1$ or geodesics tubes around the real projective space \mathbf{RP}^n . The details are in the final Section 7, while the preliminary results on spheres and tubes are reviewed in Section 2.

In this paper, by “hypersurface” we always mean “embedded hypersurface” in \mathbf{CP}^n .

2. GEOMETRY OF SPHERES AND TUBES

The n -dimensional complex projective space is the quotient of the unit sphere $\mathbf{S}^{2n+1} = \{z \in \mathbf{C}^{n+1} : |z| = 1\}$ by the Hopf-action of \mathbf{S}^1 , $(e^{i\vartheta}, z) \mapsto e^{i\vartheta}z$. We denote by $[z]$ the equivalence class of $z \in \mathbf{S}^{2n+1}$. The tangent space of \mathbf{CP}^n at the point $[z]$ is

$$T_{[z]} \mathbf{CP}^n = \{w \in \mathbf{C}^{n+1} : z \cdot \bar{w} = 0\},$$

where $z \cdot \bar{w} = z_1 \bar{w}_1 + \dots + z_{n+1} \bar{w}_{n+1}$ is the standard Hermitian product of \mathbf{C}^{n+1} . The complex structure on $T_{[z]} \mathbf{CP}^n$ is given by $Jw = iw$, the standard multiplication by i of $w \in T_{[z]} \mathbf{CP}^n \subset \mathbf{C}^{n+1}$.

The metric $\langle \zeta, w \rangle_{FS} = \operatorname{Re}(\zeta \cdot \bar{w})$, with $\zeta, w \in T_{[z]} \mathbf{CP}^n$, is the *Fubini-Study* metric of \mathbf{CP}^n , that makes the complex projective space a Riemannian manifold. The induced distance function $d : \mathbf{CP}^n \times \mathbf{CP}^n \rightarrow [0, \pi/2]$ is $d([z], [w]) = \arccos |z \cdot \bar{w}|$.

Let $\Sigma \subset \mathbf{CP}^n$ by a C^∞ -smooth hypersurface oriented by the unit normal N . We define the second fundamental form h of Σ with the following sign convention

$$h(X, Y) = \langle \nabla_X^\Sigma N, Y \rangle, \quad X, Y \in T_{[z]} \Sigma,$$

where ∇^Σ is the Levi-Civita connection of Σ induced by the one of \mathbf{CP}^n . The *characteristic curvature* of Σ at the point $[z] \in \Sigma$ is $\kappa = h(JN, JN)$.

For any fixed $[w] \in \mathbf{CP}^n$ and $0 < r < \pi/2$, the geodesic sphere centered at $[w]$ with radius r is

$$\Sigma_r = \{[z] \in \mathbf{CP}^n : |z \cdot \bar{w}| = \cos r\}.$$

We omit reference to the center. The curvatures of Σ_r are well-known, see e.g. [CR82, Example 1 page 493]. Letting $t = \tan r$, they are

$$\begin{aligned} \lambda &= \cot r = \frac{1}{t}, \quad \text{with multiplicity } 2(n-1), \\ \kappa &= 2 \cot(2r) = \frac{1}{t} - t, \quad \text{the characteristic curvature.} \end{aligned} \tag{2.1}$$

These two curvatures are constant and distinct for each value of $t > 0$. In [Tak75a], Takagi proved that this property characterizes the sphere.

Proposition 2.1. *If $\Sigma \subset \mathbf{CP}^n$, $n \geq 2$, is a connected hyperface with precisely two distinct constant curvatures, then Σ is a subset of a sphere Σ_r .*

In fact, the constancy assumption on the curvatures can be dropped, see [CR82].

We now discuss tubes around \mathbf{CP}^k . For $k = 1, \dots, n-1$, the natural inclusion $\mathbf{S}^{2k+1} = \{z \in \mathbf{S}^{2n+1} : z_{k+2} = \dots = z_{n+1} = 0\} \subset \mathbf{S}^{2n+1}$ induces the inclusion $\mathbf{CP}^k \subset \mathbf{CP}^n$. For $0 < r < \pi/2$, we define the tube

$$\begin{aligned} T_r^k &= \{[z] \in \mathbf{CP}^n : \operatorname{dist}([z], \mathbf{CP}^k) = r\} \\ &= \{[z] = [(z', z'')] \in \mathbf{CP}^n : |z| = 1, z' \in \mathbf{C}^{k+1}, |z'| = \cos r\}. \end{aligned}$$

The curvatures of T_r^k are computed in [CR82]. Letting $t = \tan r$, they are

$$\begin{aligned} \lambda_1 &= \cot \left(r - \frac{\pi}{2} \right) = -t, \quad \text{with multiplicity } 2k, \\ \lambda_2 &= \cot r = \frac{1}{t}, \quad \text{with multiplicity } 2\ell = 2(n-1-k), \\ \kappa &= 2 \cot(2r) = \frac{1}{t} - t, \quad \text{the characteristic curvature.} \end{aligned} \tag{2.2}$$

These three curvatures are constant and distinct for each value of $t > 0$. In particular, T_r^k has constant mean curvature. For $r + s = \pi/2$ and $k + \ell = n-1$ the hypersurfaces T_r^k and T_s^ℓ are congruent.

The tubes T_r^k share with the sphere Σ_r the property of being stable for some value of r . The following theorem is proved in [BdCE88, Theorem 1.3] using the method of G -stability.

Theorem 2.2. *Let $n \geq 2$ and $k = 1, \dots, n-1$. Then:*

- 1) *The sphere Σ_r is stable if and only if $\tan^2 r \leq 2n+1$.*
- 2) *The tube T_r^k is stable if and only if*

$$\frac{2n-2k-1}{2k+3} \leq \tan^2 r \leq \frac{2n-2k+1}{2k+1}.$$

Finally, we consider geodesics tubes around $\mathbf{R}P^n$. We start from the following embedding of the sphere \mathbf{S}^n into \mathbf{S}^{2n+1} :

$$\mathbf{S}^n = \{z \in \mathbf{C}^{n+1} : |z| = 1, z = \bar{z}\} \subset \mathbf{S}^{2n+1}.$$

Passing to the quotient, the inclusion $\mathbf{S}^n \subset \mathbf{S}^{2n+1}$ gives an embedding of $\mathbf{R}P^n$ into $\mathbf{C}P^n$. The distance of $z \in \mathbf{S}^{2n+1}$ from \mathbf{S}^n is

$$\text{dist}(z, \mathbf{S}^n) = \frac{1}{2} \arccos \left(\left| \sum_{j=1}^{n+1} z_j^2 \right| \right),$$

and it does not depend on the equivalence class of z . The level-sets of this distance form the isoparametric family of hypersurfaces in \mathbf{S}^{2n+1} studied in [Nom73, Theorem 1]. Hence, the geodesic tube in $\mathbf{C}P^n$ with radius r around $\mathbf{R}P^n$ is

$$V_r = \left\{ [z] \in \mathbf{C}P^n : \left| \sum_{j=1}^{n+1} z_j^2 \right| = \cos 2r \right\}.$$

The curvatures of V_r are computed in [CR82] starting from the formulas in [Nom73] for the preimage of V_r in \mathbf{S}^{2n+1} . Letting $t = \tan r$, they are

$$\begin{aligned} \lambda_1 &= -\cot r = -\frac{1}{t}, \quad \text{with multiplicity } n-1, \\ \lambda_2 &= -\cot \left(\frac{\pi}{2} - r \right) = t, \quad \text{with multiplicity } n-1, \\ \kappa &= 2 \cot \left(\frac{\pi}{2} - 2r \right) = \frac{4t}{1-t^2}, \quad \text{the characteristic curvature.} \end{aligned} \tag{2.3}$$

These three curvatures are constant and distinct for each value of $t \in (0, 1)$. We have a third example of a complete constant mean curvatures hypersurface in $\mathbf{C}P^n$. We will see in Lemma 6.5 that V_r is not stable for any $0 < r < \pi/4$.

In [Tak75b] Takagi proved that the tubes V_r and the tubes T_r^k are characterized by the fact of having precisely three *distinct and constant* curvatures.

Proposition 2.3. *Let $n \geq 3$. If $\Sigma \subset \mathbf{C}P^n$ is a connected hypersurface with three distinct constant curvatures, then Σ is a subset of some tube T_r^k or V_r .*

We shall use Proposition 2.3 and Theorem 2.2 in the proof of Theorem 1.2 in Section 7.

3. GEOMETRY OF THE ISOMETRIC IMMERSION OF \mathbf{CP}^n INTO H^{n+1}

Let $H^{n+1} = \{A \in \text{gl}(n+1, \mathbf{C}) : \overline{A} = A^t\}$ be the set of $(n+1)$ -dimensional Hermitian matrices. This is a $(n+1)^2$ -dimensional real subspace of $\text{gl}(n+1, \mathbf{C})$. The standard scalar product on H^{n+1} is

$$\langle A, B \rangle := \frac{1}{2} \text{tr}(AB), \quad A, B \in H^{n+1}. \quad (3.1)$$

Let $\Phi : \mathbf{CP}^n \rightarrow H^{n+1}$ be the mapping that takes the equivalence class $[z] \in \mathbf{CP}^n$ to the Hermitian matrix $A = \Phi([z]) \in H^{n+1}$ of the projection of \mathbf{C}^{n+1} onto the complex line $[z]$. The matrix A satisfies $A^2 = A$ because it is a projection and $\text{tr}(A) = 1$, because it projects onto a complex line. It can be checked that Φ is an isometry from \mathbf{CP}^n with the Fubini-Study metric into H^{n+1} with the metric (3.1). Hence, from now on we identify the complex projective space with

$$\mathbf{CP}^n = \{A \in H^{n+1} : A^2 = A, \text{tr}(A) = 1\}.$$

For details on this identification and for the proof of the following lemmas, we refer the reader to [Ros83]. Our normalization in (3.1) of the scalar product is different from the one by Ros. Namely, the relation between the metric g used by Ross and the metric in (3.1) is $g(A, B) = 4 \langle A, B \rangle$. The isometric embedding Φ was introduced in [Tai68].

For any $A \in \mathbf{CP}^n$, we denote by $T_A \mathbf{CP}^n$ and $T_A^\perp \mathbf{CP}^n$ the tangent space and the normal space of \mathbf{CP}^n at the point $A \in H^{n+1}$, respectively.

Lemma 3.1. *For any $A \in \mathbf{CP}^n$, we have:*

$$T_A \mathbf{CP}^n = \{X \in H^{n+1} : XA + AX = X\}, \quad (3.2)$$

$$T_A^\perp \mathbf{CP}^n = \{Z \in H^{n+1} : AZ = ZA\}. \quad (3.3)$$

For the proof see [Ros83]. We easily see that $A, I \in T_A^\perp \mathbf{CP}^n$, where I is the identity matrix. We call the matrix $A_0 \in \mathbf{CP}^n$

$$A_0 := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

the *origin* of \mathbf{CP}^n . This is the projection onto the complex line of the versor $e_0 = (1, 0, \dots, 0) \in \mathbf{C}^{n+1}$. At the point A_0 , an orthonormal basis for the tangent space of \mathbf{CP}^n is given by the matrices $X_1, \dots, X_n, \widehat{X}_1, \dots, \widehat{X}_n$, where for $j = 1, \dots, n$

$$X_j = \begin{pmatrix} 0 & e_j \\ e_j^t & 0 \end{pmatrix} \quad \text{and} \quad \widehat{X}_j = \begin{pmatrix} 0 & -ie_j \\ ie_j^t & 0 \end{pmatrix}. \quad (3.4)$$

We will see that $\widehat{X}_j = JX_j$.

For any point $A \in \mathbf{CP}^n$ there exists a (non-unique) unitary transformation $Q \in U(n+1)$ such that $A = QA_0Q^{-1}$. The conjugation $T_Q : H^{n+1} \rightarrow H^{n+1}$, $T_Q X = QXQ^{-1}$, preserves the metric of H^{n+1} :

$$\langle T_Q X, T_Q Y \rangle = \frac{1}{2} \text{tr}(QXYQ^{-1}) = \frac{1}{2} \text{tr}(XY) = \langle X, Y \rangle, \quad X, Y \in H^{n+1}.$$

In particular, T_Q maps isometrically the tangent space $T_{A_0}\mathbf{CP}^n$ onto $T_A\mathbf{CP}^n$. We will use isometries to reduce computations of isometric-invariant quantities to the origin A_0 .

By elementary computations based on the projection equation $A^2 = A$ and on the equation $X = AX + XA$ for tangent vectors, it is possible to check the following algebraic identities.

Lemma 3.2. *For any point $A \in \mathbf{CP}^n$ and for any vector fields $X, Y \in T_A\mathbf{CP}^n$, we have:*

$$AXY = XYA, \quad (3.5)$$

$$AXA = 0, \quad (3.6)$$

$$X(I - 2A) = -(I - 2A)X, \quad (3.7)$$

$$(I - 2A)^2 = I, \quad (3.8)$$

$$(I - 2A)XY = XY(I - 2A). \quad (3.9)$$

Using the isometric identification Φ , the natural complex structure on $T_{[z]}\mathbf{CP}^n \subset \mathbf{C}^{n+1}$ can be taken to $T_A\mathbf{CP}^n \subset H^{n+1}$. The resulting mapping is described in the following proposition.

Proposition 3.3. *For any $A \in \mathbf{CP}^n$, the mapping $J_A : T_A\mathbf{CP}^n \rightarrow T_A\mathbf{CP}^n$ defined by the formula*

$$J_A X = i(I - 2A)X, \quad (3.10)$$

satisfies the following properties:

- i) *it is an isometry;*
- ii) *it satisfies $J_A^2 = -\text{Id}$, where Id is the identity mapping;*
- iii) *it commutes with the isometries T_Q , i.e., for any $A, B \in \mathbf{CP}^n$ with $A = T_Q B$ for some $Q \in U(n+1)$, and for any $X \in T_B\mathbf{CP}^n$, we have*

$$J_A T_Q X = T_Q J_B X. \quad (3.11)$$

We compute the mean curvature of the immersion of \mathbf{CP}^n into H^{n+1} . For any $A \in \mathbf{CP}^n$ we define the orthogonal projections $\pi_A^\top : H^{n+1} \rightarrow T_A\mathbf{CP}^n$ and $\pi_A^\perp : H^{n+1} \rightarrow$

$T_A^\perp \mathbf{CP}^n$. Explicit formulas for π^\top and π^\perp can be expressed using the symmetric product $\pi : H^{n+1} \times H^{n+1} \rightarrow H^{n+1}$

$$\pi(X, Y) = XY + YX.$$

Notice that by (3.3) and (3.5), we have $\pi : T_A \mathbf{CP}^n \times T_A \mathbf{CP}^n \rightarrow T_A^\perp \mathbf{CP}^n$.

Lemma 3.4. *For any $A \in \mathbf{CP}^n$ and $X \in H^{n+1}$, we have*

$$\pi_A^\top(X) = \pi(A, X) - 2AXA, \quad (3.12)$$

$$\pi_A^\perp(X) = X - \pi(A, X) + 2AXA. \quad (3.13)$$

Proof. The mapping defined by formula (3.12) is linear and is the identity on $T_A \mathbf{CP}^n$. Indeed, for any $X \in T_A \mathbf{CP}^n$, by (3.6) and (3.2) we have $\pi_A^\top(X) = AX + XA = X$. We claim that $\pi_A^\top(X) \in T_A \mathbf{CP}^n$ for any $X \in H^{n+1}$. Indeed, we have

$$\begin{aligned} \pi_A^\top(X)A + A\pi_A^\top(X) &= AXA + XA - 2AXA + AX + AXA - 2AXA \\ &= \pi(A, X) - 2AXA = \pi_A^\top(X), \end{aligned}$$

where we used $A^2 = A$.

Then formula (3.12) defines the projection onto $T_A \mathbf{CP}^n$. Formula (3.13) follows from (3.12). \square

We split the standard connection ∇ of H^{n+1} into the part that is tangent to \mathbf{CP}^n and the part that is normal. Namely, for $X \in \Gamma(T\mathbf{CP}^n)$, $Y \in \Gamma(TH^{n+1})$, and $A \in \mathbf{CP}^n$ we let $\nabla_X^\top Y(A) = \pi_A^\top(\nabla_X Y)$, and $\nabla_X^\perp Y(A) = \pi_A^\perp(\nabla_X Y)$. By (3.12) and (3.13), we have the formulas

$$\nabla_X^\top Y = \pi(A, \nabla_X Y) - 2A(\nabla_X Y)A, \quad (3.14)$$

$$\nabla_X^\perp Y = \nabla_X Y - \pi(A, \nabla_X Y) + 2A(\nabla_X Y)A. \quad (3.15)$$

The second fundamental form of the immersion of \mathbf{CP}^n into H^{n+1} is the mapping $\sigma_A : T_A \mathbf{CP}^n \times T_A \mathbf{CP}^n \rightarrow T_A^\perp \mathbf{CP}^n$, $A \in \mathbf{CP}^n$, defined by $\sigma_A(X, Y) = \nabla_X^\perp Y(A)$. When no confusion arises, we drop the subscript A and write $\sigma = \sigma_A$.

The non-normalized mean curvature vector of the immersion is the trace of σ , i.e.,

$$\mathcal{H} = \text{tr}(\sigma) = \sum_{i=1}^{2n} \sigma(X_i, X_i), \quad (3.16)$$

where X_1, \dots, X_{2n} is any orthonormal frame of $T_A \mathbf{CP}^n$.

Proposition 3.5. *For any $A \in \mathbf{CP}^n$ and $X, Y \in T_A \mathbf{CP}^n$ we have*

$$\sigma(X, Y) = \pi(X, Y)(I - 2A), \quad (3.17)$$

and the trace of σ is

$$\mathcal{H} = 4(I - (n+1)A), \quad A \in \mathbf{CP}^n. \quad (3.18)$$

Proof. For a proof of (3.17) see [Ros83, Proposition 1.4]. Here, we prove (3.18). It is enough to verify the identity at A_0 . We use the orthonormal basis (3.4). We denote by $E_{jj} \in H^{n+1}$ the matrix with 1 at the entry jj and 0 elsewhere, so that $E_{00} = A_0$. By (3.17), for any $j = 1, \dots, n$ we get

$$\sigma(X_j, X_j) = 2X_j X_j (I - 2A_0) = 2(E_{jj} - A_0), \quad (3.19)$$

and, moreover, $\sigma(\widehat{X}_j, \widehat{X}_j) = \sigma(X_j, X_j)$. So we obtain

$$\mathcal{H} = 4 \sum_{j=1}^n (E_{jj} - A_0) = 4((I - A_0) - nA_0) = 4(I - (n+1)A_0).$$

□

Remark 3.6. Using (3.17) and (3.10), it is possible to check the following identity for any $X, Y, V, W \in T_A \mathbf{CP}^n$:

$$\begin{aligned} \langle \sigma(X, Y), \sigma(V, W) \rangle = & 2 \langle X, Y \rangle \langle V, W \rangle + \langle X, W \rangle \langle Y, V \rangle + \langle X, V \rangle \langle Y, W \rangle + \\ & + \langle X, JW \rangle \langle Y, JV \rangle + \langle X, JV \rangle \langle Y, JW \rangle. \end{aligned} \quad (3.20)$$

The *Weingarten endomorphism* of the immersion is the mapping $\Lambda : T_A \mathbf{CP}^n \times T_A^\perp \mathbf{CP}^n \rightarrow T_A \mathbf{CP}^n$, $A \in \mathbf{CP}^n$, defined by the formula $\Lambda(X, Z) = \Lambda_Z(X) = -\nabla_X^\top Z$.

Proposition 3.7. *For any $A \in \mathbf{CP}^n$, $X \in T_A \mathbf{CP}^n$, and $Z \in T_A^\perp \mathbf{CP}^n$ we have*

$$\Lambda_Z(X) = (XZ - ZX)(I - 2A). \quad (3.21)$$

For the proof see [Ros83, Proposition 1.4]. We establish some identities linking Λ and σ . Let X_1, \dots, X_{2n} be an orthonormal frame for $T\mathbf{CP}^n$ and we use the notation $N = X_{2n}$. In the sequel, we also let

$$\begin{aligned} \pi_{ij} &= \pi(X_i, X_j) \quad \text{and} \quad \pi_{i,N} = \pi(X_i, N), \\ \sigma_{ij} &= \sigma(X_i, X_j) \quad \text{and} \quad \sigma_{i,N} = \sigma(X_i, N). \end{aligned}$$

Lemma 3.8. *Let X_1, \dots, X_{2n-1}, N be an orthonormal frame of \mathbf{CP}^n . Then for any $i, j = 1, \dots, 2n-1$ we have*

$$\Lambda_{\sigma_{j,N}}(X_i) = \pi(\pi_{j,N}, X_i) = 2\pi_{j,N}X_i - \sigma(\sigma_{ij}, N) - \sigma(X_j, \sigma_{i,N}). \quad (3.22)$$

Proof. We prove the identity on the left of (3.22). By (3.17), (3.21), (3.7), and (3.8) we get

$$\begin{aligned} \Lambda_{\sigma_{j,N}}(X_i) &= (X_i \sigma_{j,N} - \sigma_{j,N} X_i)(I - 2A) \\ &= (X_i \pi_{j,N}(I - 2A) - \pi_{j,N}(I - 2A)X_i)(I - 2A) \\ &= X_i \pi_{j,N} + \pi_{j,N} X_i = \pi(\pi_{j,N}, X_i). \end{aligned} \quad (3.23)$$

Now we check the identity on the right. Using (3.7) and (3.8) we have:

$$\begin{aligned}
& 2\pi_{j,N}X_i - [\pi(\sigma_{ij}, N) + \pi(X_j, \sigma_{i,N})](I - 2A) \\
&= -(\pi_{ij}(I - 2A)N + N\pi_{ij}(I - 2A) + X_j\pi_{i,N}(I - 2A) + \\
&\quad + \pi_{i,N}(I - 2A)X_j)(I - 2A) + 2\pi_{j,N}X_i \\
&= \pi_{ij}N - N\pi_{ij} - X_j\pi_{i,N} + \pi_{i,N}X_j + 2\pi_{j,N}X_i \\
&= X_iX_jN + NX_jX_i + X_jNX_i + X_iNX_j \\
&= X_i\pi_{j,N} + \pi_{j,N}X_i \\
&= \pi(X_i, \pi_{j,N}).
\end{aligned}$$

□

Lemma 3.9. *For any orthonormal frame X_1, \dots, X_{2n-1}, N of \mathbf{CP}^n we have*

$$\sum_{i=1}^{2n-1} \Lambda_{\sigma_{i,N}}(X_i) = 2(n-1)N. \quad (3.24)$$

Proof. It is enough to verify (3.24) at the point $A_0 \in \mathbf{CP}^n$, with the frame (3.4) where $N = \widehat{X}_n$. Using formula (3.21) for Λ and the identities (3.7) and (3.8), for any $i = 1, \dots, n$ and $j = 1, \dots, n-1$ we find

$$\begin{aligned}
\Lambda_{\sigma(X_i, N)}(X_i) &= \delta_{in} (N - 2\widehat{X}_i) + N, \\
\Lambda_{\sigma(\widehat{X}_j, N)}(\widehat{X}_j) &= \delta_{jn} (2\widehat{X}_j + N) + N = N.
\end{aligned}$$

Summing up, we obtain (3.24).

□

4. LAPLACIAN OF POSITION AND NORMAL

Let $\Sigma \subset \mathbf{CP}^n$ be a hypersurface oriented by the unit normal N . In the following we adopt the short notation $\sigma_N = \sigma(N, N)$ and $\pi_N = \pi(N, N)$. In this section, we compute ΔA and ΔN , where Δ is the Laplace-Beltrami operator of Σ .

We denote by ∇^Σ the restriction of ∇^\top to Σ . The second fundamental form of Σ is the mapping $h : T_A\Sigma \times T_A\Sigma \rightarrow \mathbb{R}$, $h(X, Y) = \langle \nabla_X^\top N, Y \rangle$, and we denote its trace by $H = \text{tr}(h)$.

Notation. From now on, we will omit the symbol of sum over repeated indices. The repeated index always runs from 1 to $2n-1$. In the other cases, we will write the sum.

Theorem 4.1. *Let $\Sigma \subset \mathbf{CP}^n$ be an oriented hypersurface with normal N . The position matrix A satisfies the equation*

$$\Delta A = \text{tr}(\sigma) - \sigma_N - \text{tr}(h)N, \quad A \in \Sigma, \quad (4.1)$$

where σ is the second fundamental form of the immersion $\mathbf{CP}^n \subset H^{n+1}$.

Proof. Without loss of generality we can assume that $A_0 \in \Sigma$ and we check formula (4.1) at the point A_0 . Let X_1, \dots, X_{2n-1} be a frame of vector fields tangent to Σ given by normal coordinates at A_0 . Namely, for all $i, j = 1, \dots, 2n-1$, we have

$$\nabla_{X_i}^\Sigma X_j(A_0) = 0. \quad (4.2)$$

In the next line and in the rest of the paper, we shall use the identity

$$\nabla_X A = X.$$

With sum over repeated indices for $j = 1, \dots, 2n-1$, we have

$$\begin{aligned} \Delta A|_{A=A_0} &= \nabla_{X_j} \nabla_{X_j} A|_{A=A_0} = \nabla_{X_j} X_j(A_0) = \\ &= \nabla_{X_j}^\top X_j(A_0) + \sigma_{A_0}(X_j, X_j). \end{aligned}$$

In the last equality, we used the definition of the second fundamental form σ . Again in A_0 , by (4.2) we obtain

$$\nabla_{X_j}^\top X_j = \langle \nabla_{X_j}^\top X_j, N \rangle N = -\langle X_j, \nabla_{X_j}^\top N \rangle N = -\text{tr}(h)N.$$

Since X_1, \dots, X_{2n-1}, N is an orthonormal frame of \mathbf{CP}^n , from the definition (3.16) of \mathcal{H} we have $\sigma(X_j, X_j) = \mathcal{H} - \sigma_N$, and this ends the proof. \square

In the next theorem, we compute a formula for ΔN . The second fundamental form h of Σ can be identified with a linear operator on $T_A \Sigma$. The restriction of σ to $T_A \Sigma$ can be identified with a linear operator from $T_A \Sigma$ to $\text{End}(T_A \Sigma, T_A^\perp \mathbf{CP}^n)$. Hence, the composition $\sigma h = \sigma \circ h$ is a linear operator from $T_A \Sigma$ to $\text{End}(T_A \Sigma, T_A^\perp \mathbf{CP}^n)$. Namely, for any $X, Y \in T_A \Sigma$ we have $\sigma h(X)[Y] = \sigma(h(X), Y)$. We denote its trace by

$$\text{tr}(\sigma h) = \sigma h(X_i)[X_i] = \sigma(h_{ij}X_j, X_i) = h_{ij}\sigma_{ij} \in T_A^\perp \mathbf{CP}^n,$$

where $\sigma_{ij} = \sigma(X_i, X_j)$ and $h_{ij} = h(X_i, X_j)$ for any orthonormal frame X_1, \dots, X_{2n-1} of $T_A \Sigma$.

Theorem 4.2. *Let $\Sigma \subset \mathbf{CP}^n$ be an oriented hypersurface with constant mean curvature. The normal N to the hypersurface satisfies the equation*

$$\Delta N = 2\text{tr}(\sigma h) - (|h|^2 + 2(n-1))N - \text{tr}(h)\sigma_N. \quad (4.3)$$

The proof is preceded by a numbers of lemmas. We are using a frame of vector fields satisfying (4.2).

Lemma 4.3. *Let $\Sigma \subset \mathbf{CP}^n$ be an orientable hypersurface with constant mean curvature. At the center $A_0 \in \Sigma$ of normal coordinates, the entries of the second fundamental form $h_{ij} = h(X_i, X_j)$ satisfy for each $j = 1, \dots, 2n-1$ the equations*

$$X_i h_{ij} = 0, \quad (4.4)$$

with sum over the repeated index.

Proof. By the Codazzi's equations, we have the identities for $i, j, \ell = 1, \dots, 2n-1$

$$\begin{aligned} X_j h_{i\ell} - X_i h_{j\ell} &= \langle R(X_i, X_j)N, X_\ell \rangle + \langle \nabla_{X_j}^\top N, \nabla_{X_i}^\top X_\ell \rangle - \langle \nabla_{X_i}^\top N, \nabla_{X_j}^\top X_\ell \rangle + \\ &\quad - \langle N, \nabla_{[X_i, X_j]}^\top X_\ell \rangle, \end{aligned}$$

where N is the normal to Σ and R is the Riemann curvature tensor of \mathbf{CP}^n . Since ∇^\top has vanishing torsion, at the point A_0 we have by (4.2)

$$[X_i, X_j] = \nabla_{X_i}^\top X_j - \nabla_{X_j}^\top X_i = 0.$$

Thus, at the point A_0 the previous identity reduces to

$$X_j h_{i\ell} - X_i h_{j\ell} = \langle R(X_i, X_j)N, X_\ell \rangle$$

Letting $i = \ell$, summing up in $i = 1, \dots, 2n-1$, and using the fact that Σ has constant mean curvature, we obtain

$$X_i h_{ij} = -\langle R(X_i, X_j)N, X_i \rangle = -\text{Ric}(X_j, N).$$

The last equality follows from the standard symmetries of the curvature operator. In fact, we have $\langle R(N, X_j)N, N \rangle = \langle R(X_j, N)N, N \rangle = -\langle R(N, X_j)N, N \rangle$ and thus $\langle R(N, X_j)N, N \rangle = 0$. The complex projective space is a Kähler manifold, and thus it is an Einstein manifold. From the orthogonality of X_j and N it follows that $\text{Ric}(X_j, N) = 0$. \square

Lemma 4.4. *Let $\Sigma \subset \mathbf{CP}^n$ be an orientable hypersurface with normal N and $H = \text{tr}(h)$. At the center of normal coordinates, we have the identity*

$$\nabla_{X_i} \sigma(X_i, N) = \text{tr}(\sigma h) - 2(n-1)N - H\sigma_N. \quad (4.5)$$

Proof. By (3.17), (3.21) and the definition of second fundamental form σ we have

$$\begin{aligned} \nabla_{X_i} \sigma(X_i, N) &= \nabla_{X_i} \pi_{i,N}(I - 2A) + \pi_{i,N} \nabla_{X_i}(I - 2A) \\ &= \sigma(\nabla_{X_i} X_i, N) + \sigma(X_i, \nabla_{X_i} N) - 2\pi_{i,N} X_i \\ &= -h_{ii} \sigma_N + \sigma(\sigma_{ii}, N) + h_{ij} \sigma_{ij} + \sigma(X_i, \sigma_{i,N}) - 2\pi_{i,N} X_i. \\ &= -h_{ii} \sigma_N + h_{ij} \sigma_{ij} + (*), \end{aligned}$$

where $(*) = \sigma(\sigma_{ii}, N) + \sigma(X_i, \sigma_{i,N}) - 2\pi_{i,N} X_i$. Using (3.22) in the particular case when $i = j$, we deduce that $(*) = -\Lambda_{\sigma_{i,N}}(X_i)$.

Hence, we proved that

$$\nabla_{X_i} \sigma(X_i, N) = -h_{ii} \sigma_N + h_{ij} \sigma_{ij} - \Lambda_{\sigma_{i,N}}(X_i), \quad (4.6)$$

and our claim (4.5) follows from (3.24). \square

Remark 4.5. Starting from

$$\nabla_{X_i} \sigma(X_i, N) = \nabla_{X_i}^\top \sigma(X_i, N) + \nabla_{X_i}^\perp \sigma(X_i, N) = -\Lambda_{\sigma_{i,N}}(X_i) + \nabla_{X_i}^\perp \sigma_{i,N},$$

and using (4.6) we obtain

$$\nabla_{X_i}^\perp \sigma(X_i, N) = -H\sigma_N + h_{ij}\sigma_{ij}. \quad (4.7)$$

Proof of Theorem 4.2. We check formula (4.3) using normal coordinates at the point $A_0 \in \Sigma$. Using $|N|^2 = 1$ and $\nabla_{X_i}^\perp N = \sigma(X_i, N)$ we obtain

$$\Delta N = \nabla_{X_i} \nabla_{X_i} N = \nabla_{X_i} (\nabla_{X_i}^\perp N + \nabla_{X_i}^\top N) = \nabla_{X_i} \left\{ \sigma(X_i, N) + h_{ij} X_j \right\}. \quad (4.8)$$

From (4.4) and (4.2), we deduce that, at the point A_0 ,

$$\nabla_{X_i} (h_{ij} X_j) = h_{ij} \nabla_{X_i} X_j = h_{ij} \{ \sigma_{ij} - h_{ij} N \} = -|h|^2 N + h_{ij} \sigma_{ij}.$$

By (4.5) we have

$$\nabla_{X_i} \sigma(X_i, N) = -H\sigma_N + h_{ij}\sigma_{ij} - 2(n-1)N,$$

and then $\Delta N = -(|h|^2 + 2(n-1))N + 2h_{ij}\sigma_{ij} - H\sigma_N$. \square

5. LAPLACIAN OF σ_N

Let A be the position matrix. In the second variation of the area of a hypersurface Σ we shall use normal deformations associated with the coordinate-functions of the matrix $u = \Delta A$. In our formula (4.1) for ΔA there is the term σ_N and thus to know Δu we need $\Delta \sigma_N$. In this section we compute this crucial quantity.

Let X_1, \dots, X_{2n-1} be an orthonormal frame of $T\Sigma$ such that $X_{n+j} = JX_j$ for all $j = 1, \dots, n-1$. Then we have $X_n = -JN$, where N is the normal to Σ and J is the complex structure. The tangent vector

$$h_N = \sum_{j=1, j \neq n}^{2n-1} h(JN, X_j) X_j \in \mathbf{CT}\Sigma$$

does not depend on the given frame. The vector h_N is the projection onto $\mathbf{CT}\Sigma$ of $\nabla_{JN}^\Sigma N$. So, we have $h_N = 0$ if and only if $\nabla_{JN}^\Sigma N = \kappa JN$ with $\kappa = h(JN, JN)$, i.e., JN is a principal direction with characteristic curvature κ .

Lemma 5.1. *Let $\Sigma \subset \mathbf{CP}^n$ be an oriented hypersurface with normal N , characteristic curvature κ , and $H = \text{tr}(h)$. Then we have*

$$h_{ij} \Lambda_{\sigma(X_j, N)}(X_i) = (H - \kappa)N + Jh_N. \quad (5.1)$$

Proof. The proof is a computation based on the following relation between the second fundamental form σ and the *Weingarten* endomorphism, that is a consequence of (3.20):

$$\Lambda_{\sigma(X,Y)}(V) = 2 \langle X, Y \rangle V + \langle Y, V \rangle X + \langle X, V \rangle Y + \langle JY, V \rangle JX + \langle JX, V \rangle JY, \quad (5.2)$$

for every $X, Y, V \in T_A \mathbf{CP}^n$. For details see [Ros84, Section 1] and recall the normalization $g(A, B) = 4 \langle A, B \rangle$.

Indeed, by (5.2) we get:

$$\begin{aligned} \Lambda_{\sigma(X_j, N)}(X_i) &= \delta_{ij} N - \delta_{in} JX_j, & j &= 1, \dots, n; \\ \Lambda_{\sigma(JX_j, N)}(X_i) &= -\delta_{ij} JN + \delta_{in} X_j, & j &= 1, \dots, n-1, \end{aligned}$$

for any $i = 1, \dots, n$, and

$$\begin{aligned} \Lambda_{\sigma(X_j, N)}(JX_i) &= \delta_{ij} JN, & j &= 1, \dots, n; \\ \Lambda_{\sigma(JX_j, N)}(JX_i) &= \delta_{ij} N, & j &= 1, \dots, n-1, \end{aligned}$$

for every $i = 1, \dots, n-1$. So, adding up appropriately we prove the thesis. \square

The following lemma is a technical computation.

Lemma 5.2. *Let $\Sigma \subset \mathbf{CP}^n$ be an oriented hypersurface with normal N . Letting, for any orthonormal frame of $T\Sigma$,*

$$S_1 = \sigma(\sigma_{i,N}, \sigma_{i,N}) - 4\pi(\sigma_{i,N}, N)X_i \quad \text{and} \quad S_2 = -\pi(N, N)\Delta A,$$

we have the identity

$$S_1 + S_2 = 2\text{tr}(h)N - \text{tr}(\sigma) + 2(n-1)\sigma_N. \quad (5.3)$$

The proof is postponed at the end of the section. In the previous section we introduced the linear operator $\sigma h : T_A \Sigma \rightarrow \text{End}(T_A \Sigma, T_A^\perp \mathbf{CP}^n)$. In the same way, we define the linear operator $\sigma h^2(X)[Y] = \sigma(h^2(X), Y)$, for $X, Y \in T_A \Sigma$. Its trace is

$$\text{tr}(\sigma h^2) = \sigma h^2(X_j)[X_j] = \sigma(h_{ij}h(X_i), X_j) = h_{ij}h_{ik}\sigma_{jk} \in T_A^\perp \mathbf{CP}^n.$$

Now we are ready to prove the main formula of this section.

Theorem 5.3. *Let $\Sigma \subset \mathbf{CP}^n$ be an orientable hypersurface with constant mean curvature and normal N . Then we have*

$$\Delta \sigma_N = 4\kappa N + 2\text{tr}(\sigma h^2 - \sigma) - 2|h|^2 \sigma_N - 4Jh_N, \quad (5.4)$$

where κ is the characteristic curvature of Σ .

Proof. We check formula (5.4) using normal coordinates at the point $A_0 \in \Sigma$. Using (4.3), (3.17) and the short notation $\pi_N = \pi(N, N)$ we have

$$\begin{aligned}
\Delta\sigma_N &= \nabla_{X_i} \nabla_{X_i} \sigma_N = \nabla_{X_i} (2\pi(\nabla_{X_i} N, N)(I - 2A) - 2\pi_N X_i) \\
&= 2\{\sigma(\nabla_{X_i} \nabla_{X_i} N, N) + \sigma(\nabla_{X_i} N, \nabla_{X_i} N) - 4\pi(\nabla_{X_i} N, N)X_i - \pi_N \nabla_{X_i} X_i\} \\
&= 2\{\sigma(\Delta N, N) + 2\sigma(h_{ij}X_j + \sigma_{i,N}, h_{ik}X_k + \sigma_{i,N}) + \\
&\quad - 4\pi(h_{ij}X_j + \sigma_{i,N}, N)X_i - \pi_N(-h_{ii}N + \sigma_{ii})\} \\
&= -2(|h|^2 + 2(n-1))\sigma_N + 2h_{ij}h_{ik}\sigma_{jk} + \\
&\quad + 4h_{ij}\{\sigma(\sigma_{ij}, N) + \sigma(X_j, \sigma_{i,N}) - 2\pi_{j,N}X_i\} \\
&\quad + 2(\sigma(\sigma_{i,N}, \sigma_{i,N}) - 4\pi(\sigma_{i,N}, N)X_i) - 2\pi_N \Delta A.
\end{aligned}$$

We used the identity $\sigma(\sigma_N, N) = 0$. By (3.22) and (5.3), we have

$$\begin{aligned}
\Delta\sigma_N &= -2(|h|^2 + 2(n-1))\sigma_N + 2h_{ij}h_{ik}\sigma_{jk} - 4h_{ij}\Lambda_{\sigma(X_j, N)}(X_i) \\
&\quad + 4(n-1)\sigma_N - 2\text{tr}(\sigma) + 4\text{tr}(h)N \\
&= 4\text{tr}(h)N - 2\text{tr}(\sigma) - 2|h|^2\sigma_N + 2h_{ij}h_{ik}\sigma_{jk} - 4h_{ij}\Lambda_{\sigma(X_j, N)}(X_i).
\end{aligned}$$

By (5.1), this ends the proof. □

Proof of Lemma 5.2. We check the formula at the point $A_0 \in \Sigma$. Using the formulas (3.8), (3.9), and (3.7) we obtain

$$\begin{aligned}
S_1 &= 2\pi(X_i, N)(I - 2A)\pi(X_i, N)(I - 2A)^2 - 4\pi(\sigma(X_i, N), N)X_i \\
&= 2[\pi_{i,N}^2(I - 2A) - 2(\pi_{i,N}(I - 2A)N + N\pi_{i,N}(I - 2A))X_i] \\
&= 2\{\pi_{i,N}^2 - 2(\pi_{i,N}N - N\pi_{i,N})X_i\}(I - 2A).
\end{aligned}$$

A simple computation gives

$$\begin{aligned}
\pi(X_i, N)^2(I - 2A_0) &= -\delta_{in}(E_{in} + E_{ni}) + E_{nn} + E_{ii}, \\
\pi(\widehat{X}_j, N)^2(I - 2A_0) &= E_{nn} + E_{jj},
\end{aligned}$$

and also

$$\begin{aligned}
(\pi(N, X_i)NX_i - N\pi(X_i, N)X_i)(I - 2A_0) &= -\delta_{in}E_{ni} + E_{ii} + (1 - \delta_{in})A_0, \\
(\pi(N, \widehat{X}_j)N\widehat{X}_j - N\pi(\widehat{X}_j, N)\widehat{X}_j)(I - 2A_0) &= E_{jj} + A_0.
\end{aligned}$$

Therefore at A_0 we have

$$\begin{aligned}
S_1 &= 2 \sum_{i=1}^n [-\delta_{in}(E_{in} + E_{ni}) + E_{nn} + E_{ii} - 2(-\delta_{in}E_{ni} + E_{ii} + (1 - \delta_{in})A_0)] + \\
&\quad + 2 \sum_{i=1}^{n-1} [E_{nn} + E_{ii} - 2(E_{ii} + A_0)] \\
&= 2[2nE_{nn} - 2(I - A_0) + (4 - 4n)A_0] \\
&= 4nE_{nn} - 4I + (12 - 8n)A_0.
\end{aligned}$$

Moreover, using (3.18) and $\sigma_N = 2(E_{nn} - A_0)$, we have $4I = \text{tr}(\sigma) + 4(n+1)A_0$ and $2E_{nn} = \sigma_N + 2A_0$, and hence we get

$$S_1 = 2n\sigma_N - \text{tr}(\sigma) + 8(1 - n)A_0. \quad (5.5)$$

Now, we compute S_2 at A_0 . Using $\pi(N, N)N = 2N$, $\pi(N, N)\text{tr}(\sigma) = -8nA_0 + 8E_{nn}$, and $\pi(N, N)\sigma_N = -4A_0 + 4E_{nn}$, we get

$$\begin{aligned}
S_2 &= -\pi(N, N)(-\text{tr}(h)N + \text{tr}(\sigma) - \sigma_N) \\
&= -(-2\text{tr}(h)N - 8nA_0 + 8E_{nn} + 4A_0 - 4E_{nn}) \\
&= 2\text{tr}(h)N - 8(1 - n)A_0 - 2\sigma_N.
\end{aligned}$$

Adding S_1 and S_2 we get the claim. \square

6. TRACE OF THE SECOND VARIATION OF THE AREA

Let Σ be a C^∞ hypersurface with normal N and without boundary and let $u \in C^\infty(\Sigma)$ be a function with zero mean:

$$\int_{\Sigma} u d\mu = 0, \quad (6.1)$$

where μ is the Riemannian hypersurface measure in \mathbf{CP}^n . For $t \in \mathbb{R}$ and $p \in \Sigma$ let $t \mapsto \gamma_p(t)$ be the curve (geodesic) in \mathbf{CP}^n solving $\nabla_{\dot{\gamma}}^\top \dot{\gamma} = 0$ with $\gamma(0) = p$ and $\dot{\gamma}(0) = u(p)N(p)$. For small t , the hypersurface $\Sigma(t; u) = \{\gamma_p(t) \in \mathbf{CP}^n : p \in \Sigma\}$ is well defined and we denote its area by $\mathcal{A}(t; u) = \mu(\Sigma(t; u))$. If we have

$$\left. \frac{d\mathcal{A}(t; u)}{dt} \right|_{t=0} = 0$$

for any $u \in C^\infty(\Sigma)$ satisfying (6.1) then Σ has constant mean curvature. If $\Sigma(t; u)$ is the boundary of a region with volume $V(t)$, then condition (6.1) implies that $V'(0) = 0$.

The second variation of the area functional is given by the formula

$$\mathcal{A}''(u) = \left. \frac{d^2 \mathcal{A}(t; u)}{dt^2} \right|_{t=0} = - \int_{\Sigma} u \mathcal{L} u d\mu, \quad (6.2)$$

where

$$\mathcal{L}u = \Delta u + (|h|^2 + \text{Ric}(N))u$$

is the Jacobi operator, see e.g. [BdCE88]. The quantity $\text{Ric}(N)$ is the Ricci curvature of \mathbf{CP}^n in the direction N .

Definition 6.1 (Stability). We say that an oriented hypersurfaces Σ without boundary and with constant mean curvature is *stable* if $\mathcal{A}''(u) \geq 0$ for any $u \in C^\infty(\Sigma)$ satisfying (6.1).

It is well known that the quantity $\text{Ric}(N)$ in the formula for the second variation is a geometric constant in \mathbf{CP}^n . We need the precise value of this constant. The sectional curvature of \mathbf{CP}^n is

$$K(X \wedge Y) = 1 + 3\langle JX, Y \rangle^2, \quad (6.3)$$

where $X, Y \in T_A \mathbf{CP}^n$ is an orthonormal basis of the 2-plane $X \wedge Y$ and J is the complex structure of $T_A \mathbf{CP}^n$. Let $X_1, \dots, X_n, \hat{X}_1, \dots, \hat{X}_{n-1}, N$ be an orthonormal basis of $T_A \mathbf{CP}^n$. Then the Ricci curvature $\text{Ric}(N)$ is

$$\text{Ric}(N) = \sum_{j=1}^n K(N, X_j) + \sum_{j=1}^{n-1} K(N, \hat{X}_j). \quad (6.4)$$

By the fact that $N = JX_n$, one has:

$$K(N, X_j) = 1 + 3\langle JN, X_j \rangle^2 = 1 + 3\langle X_n, X_j \rangle^2 = \begin{cases} 1, & \text{for } j \neq n \\ 4, & \text{for } j = n \end{cases}$$

$$K(N, \hat{X}_j) = 1 + 3\langle JN, \hat{X}_j \rangle^2 = 1 + 3\langle X_n, \hat{X}_j \rangle^2 = 1,$$

because $j \leq n-1$. Thus formula (6.4) yields $\text{Ric}(N) = 2n + 2$.

On Σ we consider the matrix valued function $u = \Delta A$, where A is the position matrix. For any $V \in H^{n+1}$, we define the scalarization $u_V = \langle u, V \rangle$. By the divergence theorem, the function u_V satisfies the zero-mean condition (6.1) because Σ has no boundary.

The mapping $Q_\Sigma : H^{n+1} \rightarrow \mathbb{R}$ defined by $Q_\Sigma(V) = \mathcal{A}''(u_V)$ is a quadratic form. If the surface Σ is stable then Q_Σ is positive semidefinite, i.e., $Q_\Sigma(V) \geq 0$ for any $V \in H^{n+1}$. It follows that $\text{tr}(Q_\Sigma) \geq 0$. In the next theorem we compute an explicit expression for this trace.

Theorem 6.2. *Let Σ be an oriented complete hypersurface with constant mean curvature. The trace of the quadratic form Q_Σ is*

$$\text{tr}(Q_\Sigma) = 4 \int_\Sigma \left\{ 2(n+1)H^2 + 2(n^2-1)(2n-|h|^2) - (H+\kappa)^2 - |h_N|^2 \right\} d\mu. \quad (6.5)$$

Proof. For any orthonormal basis \mathcal{V} of H^{n+1} , we have

$$\begin{aligned} \operatorname{tr}(Q_\Sigma) &= \sum_{V \in \mathcal{V}} Q_\Sigma(V) = - \int_\Sigma \sum_{V \in \mathcal{V}} u_V \mathcal{L} u_V d\mu \\ &= - \int_\Sigma \{ \langle u, \Delta u \rangle + (|h|^2 + 2(n+1))|u|^2 \} d\mu. \end{aligned}$$

We compute first the norm of u . By formula (4.1) we have:

$$\begin{aligned} |u|^2 &= |\Delta A|^2 = \langle \mathcal{H} - \sigma_N - \operatorname{tr}(h)N, \mathcal{H} - \sigma_N - \operatorname{tr}(h)N \rangle \\ &= H^2 + |\mathcal{H}|^2 - 2 \langle \mathcal{H}, \sigma_N \rangle + |\sigma_N|^2, \end{aligned}$$

because the matrices N and $\mathcal{H} - \sigma_N$ are orthogonal. Using the identity $\sigma(X, Y) = \sigma(JX, JY)$ for any $X, Y \in T_A \mathbf{CP}^n$ (see [Ros83, Proposition 1.6]), we obtain

$$|\mathcal{H}|^2 = \sum_{i,j=1}^{2n} \langle \sigma_{ii}, \sigma_{jj} \rangle = 4 \sum_{i,j=1}^n \langle \sigma_{ii}, \sigma_{jj} \rangle.$$

Now by (3.20) we have for any $i, j = 1, \dots, n$

$$\langle \sigma_{ii}, \sigma_{jj} \rangle = 2(1 + \delta_{ij}), \quad (6.6)$$

and hence

$$\begin{aligned} |\mathcal{H}|^2 &= 4 \sum_{i,j=1}^n 2(1 + \delta_{ij}) = 8 \left(n^2 + \sum_{i,j=1}^n \delta_{ij} \right) \\ &= 8(n^2 + n) = 8n(n+1). \end{aligned} \quad (6.7)$$

In the same way, we have

$$\langle \mathcal{H}, \sigma_N \rangle = \sum_{i=1}^{2n} \langle \sigma_{ii}, \sigma_N \rangle = 2 \sum_{i=1}^n \langle \sigma_{ii}, \sigma_N \rangle = 4 \sum_{i=1}^n (1 + \delta_{in}) = 4(n+1), \quad (6.8)$$

where we used (6.6) with $j = n$. Finally, by (6.6) with $i = j = n$, we have

$$|\sigma_N|^2 = 4. \quad (6.9)$$

Now, by (6.7), (6.8) and (6.9) we get

$$\begin{aligned} |u|^2 &= H^2 + 8n(n+1) - 8(n+1) + 4 = H^2 + 8(n+1)(n-1) + 4 \\ &= H^2 + 8n^2 - 8 + 4 = H^2 + 4(2n^2 - 1). \end{aligned} \quad (6.10)$$

By formula (3.18) we have $\Delta \mathcal{H} = -4(n+1)\Delta A = -4(n+1)u$. Hence, from formula (4.1) we find

$$\begin{aligned} \langle u, \Delta u \rangle &= \langle u, -H\Delta N - 4(n+1)u - \Delta \sigma_N \rangle \\ &= -4(n+1)|u|^2 - H\langle u, \Delta N \rangle - \langle u, \Delta \sigma_N \rangle. \end{aligned}$$

Since N is orthogonal to \mathcal{H} , σ_N , and σ_{ij} we have

$$\begin{aligned}\langle u, \Delta N \rangle &= \langle -\text{tr}(h)N + \mathcal{H} - \sigma_N, -(|h|^2 + 2(n-1))N + 2h_{ij}\sigma_{ij} - \text{tr}(h)\sigma_N \rangle \\ &= H(|h|^2 + 2(n-1)) + 2\langle \mathcal{H}, \text{tr}(\sigma h) \rangle - H\langle \mathcal{H}, \sigma_N \rangle \\ &\quad - 2\langle \sigma_N, \text{tr}(\sigma h) \rangle + H|\sigma_N|^2.\end{aligned}$$

By (3.20) we have

$$\langle \sigma(X_i, X_i), \sigma(X_j, X_k) \rangle = 2(\delta_{jk} + \delta_{ik}\delta_{ij}), \quad (6.11)$$

$$\langle \sigma(X_i, X_i), \sigma(X_j, JX_k) \rangle = 0, \quad (6.12)$$

for every $i, j, k = 1, \dots, n$. Hence, by (6.11) and (6.12) and also using the notation $h_{\hat{j}\hat{k}} := h(JX_j, JX_k)$, we get

$$\begin{aligned}\langle \mathcal{H}, \text{tr}(\sigma h) \rangle &= 2 \sum_{i=1}^n \sum_{j,k=1}^{2n-1} h_{jk} \langle \sigma_{ii}, \sigma_{jk} \rangle = \\ &= 4 \sum_{i,j,k=1}^n h_{jk} (\delta_{jk} + \delta_{ik}\delta_{ij}) + 4 \sum_{i=1}^n \sum_{j,k=1}^{n-1} h_{\hat{j}\hat{k}} (\delta_{jk} + \delta_{ik}\delta_{ij}) \\ &= 4nh_{jj} + 4h_{ii} = 4(n+1)H.\end{aligned} \quad (6.13)$$

Again by (6.11) and (6.12),

$$\langle \sigma_N, \text{tr}(\sigma h) \rangle = h_{ij} \langle \sigma_N, \sigma_{ij} \rangle = 2h_{ij}(\delta_{ij} + \delta_{jn}\delta_{in}) = 2H + 2\kappa. \quad (6.14)$$

Finally, using (6.8), (6.9), (6.13) and (6.14) we get

$$\begin{aligned}\langle u, \Delta N \rangle &= H(|h|^2 + 2(n-1)) + 8(n+1)H - 4(n+1)H - 4(H + \kappa) + 4H \\ &= (|h|^2 + 6n + 2)H - 4\kappa,\end{aligned}$$

and so we obtain the formula

$$\langle u, \Delta N \rangle = H(|h|^2 + 6n + 2) - 4\kappa. \quad (6.15)$$

We are left with the computation of $\langle u, \Delta \sigma_N \rangle$. By formula (5.4), also using $\langle u, h_N \rangle = 0$, we obtain

$$\begin{aligned}\langle u, \Delta \sigma_N \rangle &= \langle -\text{tr}(h)N + \mathcal{H} - \sigma_N, 4\kappa N - 2\mathcal{H} - 2|h|^2\sigma_N + 2\text{tr}(\sigma h^2) \rangle \\ &= -4\kappa H - 2|\mathcal{H}|^2 - 2(|h|^2 - 1)\langle \mathcal{H}, \sigma_N \rangle + 2\langle \mathcal{H}, \text{tr}(\sigma h^2) \rangle + 2|h|^2|\sigma_N|^2 \\ &\quad - 2\langle \sigma_N, \text{tr}(\sigma h^2) \rangle.\end{aligned}$$

Again by (6.11) and (6.12),

$$\begin{aligned}
\langle \mathcal{H}, \text{tr}(\sigma h^2) \rangle &= 2 \sum_{i=1}^n \sum_{\ell, j, k=1}^{2n-1} h_{\ell j} h_{\ell k} \langle \sigma_{ii}, \sigma_{jk} \rangle = \\
&= 4 \sum_{i=1}^n \sum_{\ell=1}^{2n-1} \left\{ \sum_{j, k=1}^n h_{\ell j} h_{\ell k} (\delta_{jk} + \delta_{ik} \delta_{ij}) + \sum_{j, k=1}^{n-1} h_{\ell j} h_{\ell k} (\delta_{jk} + \delta_{ik} \delta_{ij}) \right\} \\
&= 4n \sum_{\ell, j=1}^{2n-1} h_{\ell j}^2 + 4 \sum_{\ell, i=1}^{2n-1} h_{\ell i}^2 = 4(n+1)|h|^2.
\end{aligned}$$

So, we get

$$\langle \mathcal{H}, \text{tr}(\sigma h^2) \rangle = 4(n+1)|h|^2. \quad (6.16)$$

Moreover, we have

$$\begin{aligned}
\langle \sigma_N, \text{tr}(\sigma h^2) \rangle &= h_{ij} h_{ik} \langle \sigma_N, \sigma_{jk} \rangle = 2h_{ij} h_{ik} (\delta_{jk} + \delta_{kn} \delta_{jn}) \\
&= 2h_{ij}^2 + 2h_{in}^2 = 2|h|^2 + 2(|h_N|^2 + \kappa^2).
\end{aligned} \quad (6.17)$$

Adding (6.7), (6.8), (6.9), (6.16) and (6.17), we get the identity

$$\langle u, \Delta \sigma_N \rangle = 4|h|^2 - 8(2n-1)(n+1) - 4H\kappa - 4\kappa^2 - 4|h_N|^2. \quad (6.18)$$

Now, in order to get (6.5), we just have to use the formulas (6.10), (6.15) and (6.18) and sum them up.

□

In the next lemmas, we test the trace formula (6.5) on geodesic spheres and on the tubes introduced in Section 2.

Lemma 6.3. *For the sphere $\Sigma_r \subset \mathbf{CP}^n$ we have $\text{tr}(Q_{\Sigma_r}) \geq 0$ if and only if $\tan^2 r \leq 2n+1$. The trace is zero if and only if $\tan^2 r = 2n+1$.*

Proof. Letting $t = \tan r$, by the formulas (2.1) we have:

$$\begin{aligned}
|h|^2 &= (2n-1) \frac{1}{t^2} + t^2 - 2, \\
H^2 &= (2n-1)^2 \frac{1}{t^2} + t^2 - 2(2n-1), \\
(H + \kappa)^2 &= 4(n-1)^2 \frac{1}{t^2} + 4t^2 - 8(n-1).
\end{aligned}$$

Inserting these values into the trace formula (6.5) we find

$$\text{tr}(Q_{\Sigma_r}) = -\frac{8n(n-1)}{t^2} \mu(\Sigma_r) \{t^4 - 2nt^2 - (2n+1)\}.$$

Then we have $\text{tr}(Q_{\Sigma_r}) \geq 0$ if and only if $t^4 - 2nt^2 - (2n+1) \leq 0$, which holds if and only if $t^2 \leq 2n+1$. In particular, the trace is zero precisely when $t^2 = 2n+1$.

□

Lemma 6.3 shows that the formula (6.5) detects the sharp stability interval for the radius of a geodesic sphere, see Theorem 2.2. In the next lemma we consider the tubes T_r^k .

Lemma 6.4. *Let $k = 1, \dots, n-1$. For the tubes $T_r^k \subset \mathbf{CP}^n$ we have:*

- 1) *If $n = 2k + 1$ then $\text{tr}(Q_{T_r^k}) > 0$ for any $0 < r < \pi/2$.*
- 2) *If $n \neq 2k + 1$ then $\text{tr}(Q_{T_r^k}) \geq 0$ if and only if $\tan^2 r \leq d(n, k)$ for a certain positive number $d(n, k)$ that satisfies*

$$d(n, k) > \frac{2n - 2k + 1}{2k + 1}. \quad (6.19)$$

Moreover, $\text{tr}(Q_{T_r^k}) = 0$ precisely when $\tan^2 r = d(n, k)$.

Proof. We use the short notation $t = \tan r$ and $m = 2k + 1$. By the formulas (2.2) we have:

$$|h|^2 = mt^2 + (2n - m)\frac{1}{t^2} - 2,$$

$$H^2 = m^2 t^2 + (2n - m)^2 \frac{1}{t^2} - 2m(2n - m),$$

$$(H + \kappa)^2 = (m + 1)^2 t^2 + (2n - m + 1)^2 \frac{1}{t^2} - 2(m + 1)(2n - m + 1).$$

Inserting these values into the formula (6.5), we get the following expression for the trace of $Q_{T_r^k}$

$$\text{tr}(Q_{T_r^k}) = \frac{8\mu(T_r^k)}{t^2}(at^4 + bt^2 + c),$$

where a, b, c are coefficients depending on n and k , and namely

$$a = m(n + 1)(m - n + 1) - \frac{1}{2}(m + 1)^2,$$

$$b/2 = (n + 1)[(n^2 - 1) - m(2n - m)] + \frac{1}{2}(m + 1)(2n - m + 1),$$

$$c = (n + 1)(2n - m)(n - m + 1) - \frac{1}{2}(2n - m + 1)^2.$$

It is easy to check that $b = a + c$, which means that $t^2 = -1$ is a root of $at^4 + bt^2 + c = 0$. So we have the decomposition

$$\text{tr}(Q_{T_r^k}) = \frac{8\mu(T_r^k)}{t^2}(t^2 + 1)(at^2 + c).$$

Now there are two cases: $n = m$ and $n \neq m$. When $n = m = 2k + 1$ then n must be odd and in this case it is $a = c = \frac{1}{2}(n^2 - 1) > 0$. It follows that $\text{tr}(Q_{T_r^k}) > 0$ for any $t > 0$.

In the case $n \neq m$ we have $a < 0$ and $c > 0$. We conclude that $\text{tr}(Q_{T_r^k}) \geq 0$ if and only if $t^2 \leq d(n, k) := -c/a$. After some computations, inequality (6.19) is equivalent to $m > 1$, that is $k > 0$. \square

Lemma 6.4 shows that Theorem 6.5 is not sharp in the case of the tubes T_r^k . In fact, when $\tan^2 r = d(n, k)$ then the trace of $Q_{T_r^k}$ is zero but the tube T_r^k is unstable by Theorem 2.2 part 2.

We finish this section proving the non-stability of the tubes V_r around $\mathbf{R}P^n$.

Lemma 6.5. *The tube $V_r \subset \mathbf{C}P^n$ is unstable for any $r > 0$.*

Proof. Using the formulas (2.3) we compute:

$$\begin{aligned} |h|^2 &= (n-1)\left(t^2 + \frac{1}{t^2}\right) + \frac{16t^2}{(1-t^2)^2}, \\ H^2 &= (n-1)^2 \frac{(1-t^2)^2}{t^2} + \frac{16t^2}{(1-t^2)^2} - 8(n-1), \\ (H+\kappa)^2 &= (n-1)^2 \frac{(1-t^2)^2}{t^2} + \frac{64t^2}{(1-t^2)^2} - 16(n-1). \end{aligned}$$

Inserting these values into the formula (6.5), we obtain the value for the trace of Q_{V_r}

$$\mathrm{tr}(Q_{V_r}) = -4(n-1)\mu(V_r) \left\{ (n-1)\left(t^2 + \frac{1}{t^2}\right) + \frac{32nt^2}{(1-t^2)^2} + 2(5n-1) \right\},$$

and we see that $\mathrm{tr}(Q_{V_r}) < 0$ for any $t = \tan r \in (0, 1)$ and $n \geq 2$. We deduce that the surfaces V_r are not stable. \square

7. PROOF OF THEOREMS 1.1 AND 1.2

In this section, we prove Theorems 1.1 and 1.2. Let $\Sigma \subset \mathbf{C}P^n$ be a complete stable oriented hypersurface that is connected.

Proof of Theorem 1.1. We denote by \hat{h} the restriction of the second fundamental form h of Σ to the complex tangent space $\mathbf{C}T\Sigma$ and by \hat{H} the trace of \hat{h} . At any point of Σ , we have the identities

$$H = \hat{H} + \kappa \quad \text{and} \quad |h|^2 = |\hat{h}|^2 + 2|h_N|^2 + \kappa^2,$$

and the inequalities

$$|h|^2 \geq |\hat{h}|^2 + \kappa^2 \quad \text{and} \quad |\hat{h}|^2 \geq \frac{\hat{H}^2}{2(n-1)} = \frac{(H-\kappa)^2}{2(n-1)}. \quad (7.1)$$

Inserting these inequalities and $|h_N| \geq 0$ into (6.5) we obtain

$$\begin{aligned} \mathrm{tr}(Q_\Sigma) &\leq 4 \int_\Sigma \left\{ 2(n+1)H^2 + 2(n^2-1)\left(2n-\kappa^2 - \frac{(H-\kappa)^2}{2(n-1)}\right) - (H+\kappa)^2 \right\} d\mu \\ &= -4n \int_\Sigma p(\kappa; H, n) d\mu, \end{aligned} \quad (7.2)$$

where $p(\cdot; H, n)$ is the polynomial in (1.1). By our assumption $p(\kappa; H, n) \geq 0$ on Σ , we deduce that $\mathrm{tr}(Q_\Sigma) \leq 0$. On the other hand, the stability of Σ implies that

$\text{tr}(Q_\Sigma) \geq 0$. We deduce that $\text{tr}(Q_\Sigma) = 0$ and that we have equality in (7.2). In turn, the equality in (7.2) implies that $p(\kappa; H, n) = 0$, that

$$|h|^2 = |\hat{h}|^2 + \kappa^2 \quad \text{and} \quad |\hat{h}|^2 = \frac{\hat{H}^2}{2(n-1)}, \quad (7.3)$$

and also that $h_N = 0$ on Σ .

The equation $h_N = 0$ means that JN is an eigenvector of h . By Maeda's theorem [Mae76], this implies that the characteristic curvature κ is constant. This also simply follows from the fact that κ is one of the roots of $p(\kappa; H, n) = 0$. Here we use the fact that Σ is connected.

The identity in the right-hand side of (7.3) implies that Σ is umbilical in $\mathbf{CT}\Sigma$, i.e., each unit vector in $\mathbf{CT}\Sigma$ is an eigenvector of h with eigenvalue $\lambda = \hat{H}/2(n-1)$. Moreover, λ is constant on Σ , because $\hat{H} = H - \kappa$ is constant.

The two constants κ and λ are different, because in \mathbf{CP}^n there are no totally umbilical hypersurfaces. By Takagi's theorem, Proposition 2.1, Σ is a geodesic sphere: up to a suitable choice of the center of the sphere, we have $\Sigma = \Sigma_r$ for some $r \in (0, \pi/2)$. By Lemma 6.3 the equation $\text{tr}(Q_{\Sigma_r}) = 0$ implies that $\tan^2 r = 2n + 1$. \square

Finally, we prove Theorem 1.2. We shall use Takagi's characterization of tubes in Proposition 2.3 and the computations of Lemmas 6.4 and 6.5.

Proof of Theorem 1.2. We have the decomposition $\mathbf{CT}\Sigma = T_1 \oplus T_2$ where, at each point of Σ , T_1 and T_2 are subspaces of real dimension α and $\beta = 2(n-1) - \alpha$. We denote by h_1 and h_2 the restrictions of h to T_1 and T_2 , respectively, and we let $H_1 = \text{tr}(h_1)$ and $H_2 = \text{tr}(h_2)$.

We have the identity $H = H_1 + H_2 + \kappa$ and the inequalities

$$|h|^2 \geq |h_1|^2 + |h_2|^2 + \kappa^2, \quad |h_1|^2 \geq \frac{H_1^2}{\alpha} \quad \text{and} \quad |h_2|^2 \geq \frac{H_2^2}{\beta}. \quad (7.4)$$

Inserting these inequalities and $|h_N| \geq 0$ into (6.5), we obtain

$$\text{tr}(Q_\Sigma) \leq -8(n^2 - 1) \int_\Sigma p(H_1, H_2; H, n, \alpha) d\mu, \quad (7.5)$$

where $p(\cdot; H, n, \alpha)$ is the polynomial in (1.3). By our assumption $p(H_1, H_2; H, n, \alpha) \geq 0$ on Σ , we deduce that $\text{tr}(Q_\Sigma) \leq 0$. On the other hand, the stability of Σ implies that $\text{tr}(Q_\Sigma) \geq 0$. We deduce that $\text{tr}(Q_\Sigma) = 0$ and that we have equality in (7.5). In turn, the equality in (7.5) implies that $p(H_1, H_2; H, n, \alpha) = 0$, that

$$|h|^2 = |h_1|^2 + |h_2|^2 + \kappa^2 \quad \text{and} \quad |h_1|^2 = \frac{H_1^2}{\alpha}, \quad |h_2|^2 = \frac{H_2^2}{\beta} \quad (7.6)$$

and also that $h_N = 0$ on Σ .

The equation $h_N = 0$ means that JN is an eigenvector of h . By Maeda's theorem [Mae76], the characteristic curvature κ is constant.

The identities in (7.6) imply that T_1 is an eigenspace of h for a curvature λ_1 , and T_2 is an eigenspace of h for a curvature λ_2 . We clearly have $H_1 = \alpha\lambda_1$ and $H_2 = \beta\lambda_2$. From $p(H_1, H_2; H, n, \alpha) = 0$ and $H_1 + H_2 = \text{constant}$, we deduce that λ_1 and λ_2 are constant.

Now we have three cases:

- 1) $\kappa = \lambda_1 = \lambda_2$. This case is empty, because in \mathbf{CP}^n there are no totally umbilical hypersurfaces.
- 2) Precisely two of the numbers κ, λ_1 , and λ_2 are equal. By Proposition 2.1, Σ is a geodesic sphere. Hence, it must be $\lambda_1 = \lambda_2$ and the radius of the sphere is $\tan^2 r = 2n + 1$, as explained at the end of the proof of Theorem 1.1.
- 3) The three numbers κ, λ_1 , and λ_2 are different. By Proposition 2.3, the surface Σ is either a tube around \mathbf{CP}^k , $\Sigma = T_r^k$ with $\alpha = 2k$ even, or a tube around \mathbf{RP}^n , $\Sigma = V_r$ with $\alpha = n - 1$. The latter case $\Sigma = V_r$ is excluded because V_r is unstable for any $r > 0$, by Proposition 6.5. We are left with the case $\Sigma = T_r^k$ with $\alpha = 2k$ and for some $r > 0$. The radius is determined by the equation $\text{tr}(Q_{T_r^k}) = 0$. However, this equation either has no solution (this happens in the case $\alpha = \beta$), or its unique solution $r > 0$ has the property that the tube T_k^r is unstable, as shown in Proposition 6.4.

The only possible case is that Σ is a geodesic sphere with radius $\tan^2 r = 2n + 1$. □

REFERENCES

- [BdC84] J. Lucas Barbosa and Manfredo do Carmo, *Stability of hypersurfaces with constant mean curvature*, Math. Z. **185** (1984), no. 3, 339–353. MR 731682
- [BdCE88] J. Lucas Barbosa, Manfredo do Carmo, and Jost Eschenburg, *Stability of hypersurfaces of constant mean curvature in Riemannian manifolds*, Math. Z. **197** (1988), no. 1, 123–138. MR 917854
- [CR82] Thomas E. Cecil and Patrick J. Ryan, *Focal sets and real hypersurfaces in complex projective space*, Trans. Amer. Math. Soc. **269** (1982), no. 2, 481–499. MR 637703
- [Mae76] Yoshiaki Maeda, *On real hypersurfaces of a complex projective space*, J. Math. Soc. Japan **28** (1976), no. 3, 529–540. MR 0407772
- [Nom73] Katsumi Nomizu, *Some results in E. Cartan's theory of isoparametric families of hypersurfaces*, Bull. Amer. Math. Soc. **79** (1973), 1184–1188. MR 0326625
- [Ros83] Antonio Ros, *Spectral geometry of CR-minimal submanifolds in the complex projective space*, Kodai Math. J. **6** (1983), no. 1, 88–99. MR 698330
- [Ros84] ———, *On spectral geometry of Kähler submanifolds*, J. Math. Soc. Japan **36** (1984), no. 3, 433–448. MR 746704
- [Tai68] Shin-Sheng Tai, *Minimum imbeddings of compact symmetric spaces of rank one*, J. Differential Geometry **2** (1968), 55–66. MR 0231395

- [Tak75a] Ryoichi Takagi, *Real hypersurfaces in a complex projective space with constant principal curvatures*, J. Math. Soc. Japan **27** (1975), 43–53. MR 0355906
- [Tak75b] ———, *Real hypersurfaces in a complex projective space with constant principal curvatures. II*, J. Math. Soc. Japan **27** (1975), no. 4, 507–516. MR 0400120

DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”. UNIVERSITÀ DI PADOVA (ITALY)
E-mail address: `erika.battaglia@unipd.it`

DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”. UNIVERSITÀ DI PADOVA (ITALY)
E-mail address: `monti@math.unipd.it`

DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”. UNIVERSITÀ DI PADOVA (ITALY)
E-mail address: `righini@math.unipd.it`