

SINGULAR BEHAVIOR AND GENERIC REGULARITY OF MIN-MAX MINIMAL HYPERSURFACES

OTIS CHODOSH, YEVGENY LIOKUMOVICH, AND LUCA SPOLAOR

ABSTRACT. We show that for a generic 8-dimensional Riemannian manifold with positive Ricci curvature, there exists a smooth minimal hypersurface. Without the curvature condition, we show that for a dense set of 8-dimensional Riemannian metrics there exists a minimal hypersurface with at most one singular point. This extends previous work on generic regularity that only dealt with area-minimizing hypersurfaces.

These results are a consequence of a more general estimate for a one-parameter min-max minimal hypersurface $\Sigma \subset (M, g)$ (valid in any dimension):

$$\mathcal{H}^0(\mathcal{S}_{\text{nm}}(\Sigma)) + \text{Index}(\Sigma) \leq 1$$

where $\mathcal{S}_{\text{nm}}(\Sigma)$ denotes the set of singular points of Σ with a unique tangent cone non-area minimizing on either side.

INTRODUCTION

It is well known that 7-dimensional area minimizing hypersurfaces can have isolated singularities. Using work of Hardt–Simon [HS85], Smale proved in [Sma93] that in an 8-dimensional manifold M with $H_7(M; \mathbb{Z}) \neq 0$, there exists a smooth embedded area minimizing hypersurface for a generic choice of metric. In other words, he showed that isolated singularities of an area-minimizing 7-dimensional hypersurface can generically be perturbed away.

One may thus seek to find a smooth embedded minimal hypersurface in all 8-manifolds M equipped with a generic metric g , even when $H_7(M; \mathbb{Z}) = 0$. Here, we find such a hypersurface in the case of positive Ricci curvature, and give a partial answer in general. We let $\text{Met}^{2,\alpha}(M)$ denote the space of Riemannian metrics of regularity $C^{2,\alpha}$ on M and $\text{Met}_{\text{Ric}>0}^{2,\alpha}(M) \subset \text{Met}^{2,\alpha}(M)$ denote the open subset of Ricci positive metrics.

Theorem 1 (Generic regularity with positive Ricci in dimension 8). *Let M^8 be a compact smooth 8-manifold. There is an open and dense set $\mathcal{G} \subset \text{Met}_{\text{Ric}>0}^{2,\alpha}(M)$ so that for $g \in \mathcal{G}$, there exists a smooth embedded minimal hypersurface $\Sigma \subset M$.*

Without the curvature condition, we have the following partial result.

Theorem 2 (Generic almost regularity in dimension 8). *Let M^8 be a compact smooth 8-manifold. There exists a dense set $\mathcal{G} \subset \text{Met}^{2,\alpha}(M)$ so that for $g \in \mathcal{G}$, there exists a smooth embedded minimal hypersurface $\Sigma \subset M$ with at most one singular point.*

We actually prove more general results valid in all dimensions, see Theorem 3 below.

As mentioned above, the principal motivation for such results is to study generic regularity of non-minimizing, high-dimensional minimal submanifolds. This contrasts with previous works on generic regularity:

- Hardt–Simon [HS85] (resp. Smale [Sma93]), cf. [Liu19], show that regular singularities of (one-sided) minimizing hypersurfaces can be perturbed away by perturbing the boundary (resp. metric).
- White [Whi85, Whi19] shows that minimizing integral 2-cycles are smoothly embedded surfaces for a generic metric.
- Moore [Moo06, Moo07] shows that parametrized minimal (2-dimensional) surfaces are free of branch points for a generic ambient metric.

In fact, our work proves that generically there exists a minimal hypersurface of optimal regularity avoiding *certain* singularities in ambient dimensions beyond the singular dimension. Indeed, Theorem 2 is a consequence of a more general result stated below.

Theorem 3 (Generic removability of isolated singularities). *Consider a compact smooth $(n + 1)$ -manifold, for $n \geq 7$. There is a dense set $\mathcal{G} \subset \text{Met}^{2,\alpha}(M)$ with the following properties:*

- *If $g \in \mathcal{G}$ then there exists a minimal hypersurface Σ , smooth away from a closed singular set of Hausdorff dimension at most $n - 7$, so that for $\mathcal{S}_0 \subset \text{Sing}(\Sigma)$ the set of singular points with regular tangent cones, we have $\mathcal{H}^0(\mathcal{S}_0) \leq 1$.*
- *If $g \in \mathcal{G} \cap \text{Met}_{\text{Ric} > 0}^{2,\alpha}(M)$ then the same statement holds, except we can conclude that $\mathcal{H}^0(\mathcal{S}_0) = 0$.*

In order to remove the topological condition $H_7(M; \mathbb{Z}) \neq 0$ of Smale, we will use the Almgren–Pitts min-max construction [Pit81], which guarantees the existence of a minimal hypersurface Σ^n in a closed Riemannian manifold (M^{n+1}, g) . As in the area-minimizing case, when the dimension n satisfies $2 \leq n \leq 6$, the Almgren–Pitts minimal hypersurface is smooth, but for larger values of n there may be at most $(n - 7)$ -dimensional singular set (this follows from work of Schoen–Simon [SS81]). However tangent cones to min-max hypersurfaces are *a priori* only stable, while only area-minimizing cones have complements that are foliated by smooth minimal hypersurfaces (cf. [BDGG69, Law91]) and it seems that such a foliation is needed (at least on one side) to perturb the singularity away by adjusting the metric [HS85].

The key technical result of this paper is that (for one-parameter min-max) at all points—except possibly one—of the singular set with a regular tangent cone, the tangent cone is area minimizing on at least one side. Put another way, we show that tangent cones that are not area minimizing on either side “contribute to the Morse index” from the point of view of min-max (and these are precisely the cones that we are unable to perturb away using Hardt–Simon [HS85]).

0.1. Detailed description of results. Let (M^{n+1}, g) be a closed Riemannian manifold. By a *sweepout* of M we will mean a family of (possibly singular) hypersurfaces $\{\Phi(x) = \partial\Omega(x)\}_{x \in [0,1]}$, where each hypersurface $\Phi(x)$ is the boundary of an open set

$\Omega(x)$ with $\Omega(0) = \emptyset$ and $\Omega(1) = M$, and we denote the family of such sweepouts by \mathcal{S} (see Section 2 for the precise definition). The width, $W(M)$, is then defined by

$$W(M) = \inf_{\Phi \in \mathcal{S}} \left\{ \sup_x \mathbf{M}(\Phi(x)) \right\}.$$

Given a stationary integral varifold V , with $\text{supp } V$ regular outside of a set of $n - 7$ Hausdorff dimension, we define

$$\mathfrak{h}_{\text{nm}}(V) := \left\{ p \in \text{supp}(V) : \begin{array}{l} \text{for all } r > 0 \text{ small, } \text{supp } V \cap B_r(p) \text{ is not one-sided} \\ \text{homotopy area minimizing on either side (in } B_r(p)\text{).} \end{array} \right\}$$

In other words, $p \in \mathfrak{h}_{\text{nm}}(V)$ implies that in any small ball there are one-sided homotopies on both sides of $\text{supp } V$ that strictly decrease area without ever increasing area. Let \mathcal{R} denote the set of integral varifolds, whose support is a complete embedded minimal hypersurface regular away from a closed singular set of Hausdorff dimension $n - 7$. Finally, we let $\text{Index}(V)$ denote the Morse index of the regular part of the support of V , that is

$$\text{Index}(V) = \text{Index}(\text{supp}(\text{Reg}(V))).$$

Then the main technical estimate of this paper is the following result.

Theorem 4 (Index plus non-area minimizing singularities bound). *For $n \geq 7$, let (M^{n+1}, g) be a closed Riemannian manifold of class C^2 . There exists a stationary integral varifold $V \in \mathcal{R}$ such that $|V|(M) = W$, which satisfies*

$$(1) \quad \mathcal{H}^0(\mathfrak{h}_{\text{nm}}(V)) + \text{Index}(V) \leq 1.$$

If equality holds in (1), then for any point $p \in \text{supp } V \setminus \mathfrak{h}_{\text{nm}}(V)$ there is $\varepsilon > 0$ so that $\text{supp } V$ is area-minimizing to one side in $B_\varepsilon(p)$. Finally, if we write $V = \sum_i \kappa_i |\Sigma_i|$, then $\kappa_i \leq 2$ for every i ; if Σ_i is one-sided then $\kappa_i = 2$ and if $\kappa_j = 2$ for some j then each Σ_i is stable.

The above bound is valid in all dimensions and can be seen as a generalization of the work of Calabi–Cao concerning min-max on surfaces [CC92]. Indeed if we define $\mathcal{S}_{\text{nm}}(V)$ by¹

$$\mathcal{S}_{\text{nm}}(V) := \left\{ p \in \text{supp}(V) : \begin{array}{l} V \text{ is locally a } C^{1,\omega} \text{ graph over its unique tangent cone } \mathcal{C} \\ \text{at } p \text{ and both sides of } \mathcal{C} \text{ are not one-sided minimizing} \end{array} \right\}$$

then we will see that $\mathcal{S}_{\text{nm}}(V) \subset \mathfrak{h}_{\text{nm}}(V)$ in Lemma 26. In particular, (1) implies that

$$\mathcal{H}^0(\mathcal{S}_{\text{nm}}(V)) + \text{Index}(V) \leq 1.$$

Thus, if we are guaranteed to have $\text{Index}(V) = 1$ (e.g., in positive curvature) we see that $\mathcal{S}_{\text{nm}}(V) = \emptyset$. This is precisely the higher dimensional analogue of the result of Calabi–Cao (cf. Figure 1 and the remark below).

See also the more recent work of Mantoulidis [Man17] which makes a more explicit connection with Morse index, using the Allen–Cahn approach (as developed by Guaraco

¹Here ω is a modulus of continuity, and we could take it to be logarithmic, as suggested by the work of [Sim83a]. Notice in fact that at all isolated singularities \mathcal{S}_0 , minimal surfaces have unique tangent cone and are locally $C^{1,\log}$ deformation of the cone itself.

and Gaspar [Gua18, GG19]) rather than Almgren–Pitts; it would be interesting to elucidate the relationship between Mantoulidis’s Allen–Cahn techniques and our proof of Theorem 4.




FIGURE 1. The figure eight geodesic c is an example of a min-max closed geodesic that is stable and has one singularity with non-area minimizing tangent cone.

Remark. By the index bound in Theorem 4, any tangent cone to V has stable regular part. Moreover, we note that the Simons cones [Sim68] in \mathbb{R}^8 (formed from products of two spheres) are all stable and area minimizing on (at least) one side (cf. [Law91]). We particularly emphasize that the Simons cone

$$\mathbf{C}^{1,5} := \{(x, y) \in \mathbb{R}^2 \times \mathbb{R}^6 : 5|x|^2 = |y|^2\}$$

is one-sided minimizing (and stable), but is not minimizing on the other side. It seems to be an open question whether or not there exists an n -dimensional stable cone that does not minimize area on either side, for $n \geq 7$.

Even assuming the existence of a stable minimal cone which is not area minimizing on either sides, it is hard to decide if the above bound is optimal. In dimension $n = 1$, such an example is provided by the classical starfish example (cf. Figure 1), whose tangent cone at the singular point (the union of two lines through the origin) is indeed stable non-area minimizing on either sides (and the starfish fails to be one-sided homotopy minimizing on either side).

We conjecture that if there is a regular stable minimal cone that is not area-minimizing on either side, then it can arise as the tangent cone to a min-max minimal hypersurface (possibly in a manifold geometrically similar to the starfish); note that were this to occur, Theorem 4 would imply that the resulting hypersurface would necessarily be stable.

Theorem 4 generalizes the index upper bound of Marques and Neves [MN16] for Riemannian manifolds M^{n+1} , $3 \leq n + 1 \leq 7$ (see also [Zho15]). In recent years there has been tremendous progress in the understanding of the geometry of minimal hypersurfaces constructed using min-max methods in these dimensions (see [DLT13a], [MN19], [CM20], [Zho19], [Son18] and references therein).

For manifolds of dimension $n + 1 \geq 8$ much less is known. When Ricci curvature is positive Zhou obtained index and multiplicity bounds for one-parameter min-max minimal hypersurface [Zho17] (see also the work of Ramírez-Luna [RL19] and Bellettini [Bel20]). Upper Morse index bounds are known to hold in arbitrary manifolds of any dimensions for hypersurfaces constructed by Allen–Cahn, as proven by Hiesmayr and

Gaspar [Hie18, Gas20] (see also the recent work of Dey showing that the Almgren–Pitts and Allen–Cahn approaches are equivalent [Dey20]). Li proved [Li19] existence of infinitely many distinct minimal hypersurface constructed via min-max methods for a generic set of metrics, using the Weyl law of Liokumovich–Marques–Neves [LMN18].

0.2. Overview of the proof. The construction of a minimal hypersurface in Almgren–Pitts min-max theory proceeds by considering a sequence of sweepouts $\{\Phi_i(x)\}$ with the supremum of the mass $\sup_x \mathbf{M}(\Phi_i(x)) \rightarrow W(M)$ as $i \rightarrow \infty$. It is then proved that we can find a subsequence $\{i_k\}$ and $\{\Phi_{i_k}(x_k)\}$ with mass tending to W , so that $|\Phi_{i_k}|(x_k)$ converges to some $V \in \mathcal{R}$.

We outline the proof of Theorem 4. For the sake of simplicity, let’s focus on the non-cancellation case, i.e., when all multiplicities of V are one (in the case of cancellation we must argue slightly differently but the main strategy is the same). The main geometric idea is to show that there cannot be two disjoint open sets U_1, U_2 so that $\Sigma = \text{supp } V$ fails to be one-sided homotopy minimizing on the same side in both U_1 and U_2 . This property is reminiscent of (but different from) almost minimizing property introduced by Pitts to prove regularity of min-max minimal hypersurfaces.

Granted this fact, it is easy to deduce the bound (1). For example, if $\text{Index}(\Sigma) = 1$ and $\mathfrak{h}_{\text{nm}}(\Sigma) = \{p\}$, then we can localize the index in some U disjoint from p . Because Σ is unstable in U , we can find area decreasing homotopies to both sides there, and we can also find $B_r(p)$ disjoint from U with area decreasing homotopies (by definition). This contradicts the above fact.

As such, we want to show the one-sided homotopy minimizing property in pairs by using the fact that V is a min-max minimal hypersurface. However, this leads us to a major difficulty. Indeed, the approximating currents $\Phi_{i_k}(x_k)$ might cross Σ many times, making it difficult to glue in one-sided homotopies to push down the mass.

At a technical level, the main tool used in this paper is that it is possible to simplify the one-parameter case of min-max theory by constructing a nested optimal sweepout $\Phi(x)$ with $\sup \mathbf{M}(\Phi(x)) = W$. This allows us to work with one sweepout $\Phi(x)$ instead of a sequence of sweepouts. The nested property allows us to directly “glue in” the one-sided homotopies to push down the mass.

The existence of a nested optimal sweepout follows from a monotone technique from [CL20]. There Chambers and Liokumovich proved that each sweepout $\Phi_i(x)$ can be replaced by a nested sweepout $\Psi_i(x)$ with $\sup \mathbf{M}(\Psi_i(x)) \leq \sup \mathbf{M}(\Phi_i(x)) + \frac{1}{i}$. “Nested” here means that $\Psi_i(x) = \partial\Omega(x)$ for a family of open sets with $\Omega(x) \subset \Omega(y)$ if $x < y$. The proof used ideas of Chambers and Rotman [CR18] on existence of monotone homotopies of closed curves on surfaces.

After we reparametrize $\Psi_i(x)$ by the volume swept out we obtain a sequence of families that is uniformly Lipschitz in flat topology. By Arzelà–Ascoli a subsequence will converge to an optimal sweepout.

In the Almgren–Pitts theory, a “pull-tight” procedure is used to find a varifold achieving the width with good properties. We can apply this procedure to our sweepout to deduce that one of the critical varifolds in the sweepout is a smooth (up to small singular set) minimal hypersurface. We would then like to prove that this hypersurface satisfies (1). However, this poses another issue, namely that it could a priori be possible




FIGURE 2. A diagram of how to use two disjoint regions that are not one-sided homotopy minimizing to either side to push down the mass. The “bumps” have smaller mass than the original surface and are homotopic along one-sided homotopies that never increase mass. By doing one homotopy fully and then the other, we can see that the mass is always strictly decreased in this range. This can then be glued into a nested sweepout without increasing mass of the adjacent leaves.

to push the mass of the sweepout near this critical value down, while not decreasing the global width. This could lead to an infinite sequence of “pushing down” operations that could create extra critical points not dealt with previously.

As such, our second main technical tool is that we construct an optimal sweepout $\Phi(x)$ with the following special property. For every point x_0 with $\sup_{x \in U} \mathbf{M}(\Phi(x)) = W$ for every neighbourhood U of x_0 , there *does not* exist an open interval $I \ni x_0$ and a family $\{\Phi'(x)\}$ that coincides with Φ for $x \notin I$ and satisfying $\sup_{x \in I'} \mathbf{M}(\Phi'(x)) < W$ for every closed interval $I' \subset I$. (In fact, we will prove a somewhat stronger property that holds for open, half-open and closed intervals I). In other words, we can not make a small “dip” in the graph of $\mathbf{M}(x)$, pushing it below W in the neighbourhood of x_0 .

This property of Φ allows us to easily prove the “homotopy minimizing to one-side” property of V discussed above (and thus Theorem 4). A diagram of the procedure to prove this can be found in Figure 2.

It thus remains to explain how we perturb remaining singularities. Hardt-Simon (cf. [HS85]) proved that one sided perturbations of area minimizing cones with isolated singularities are smooth, and Liu extended this result to one sided stationary minimizers (cf. [Liu19]). The proofs of the bounds in Theorems 1, 2, and 3 are then obtained by combining a weak generalization of White [Whi94], where “stability” is replaced by the stronger hypothesis that the surface is homotopic minimizing to one side, with a simple surgery procedure in an annulus around the singularity, to show that singular points with regular tangent cones in $\text{Sing}(V) \setminus \mathfrak{h}_{\text{nm}}(V)$ are not generic. As such, the main results follow from (1) when combined with a result of Simon on uniqueness of the blow up at certain singularities [Sim83a]. The rest of the theorem follows from the index lower bound in manifolds with positive Ricci curvature.

0.3. Organization of the paper. The paper is divided into four sections. The first section contains the basic definitions and some useful geometric tools. The second

section introduces the notion of non-excessive optimal sweepouts, which is a key idea in the present work. The third section proves Theorem 4, while the last section is dedicated to the proofs of Theorems 2 and 3.

0.4. Acknowledgements. We are grateful to Zhihan Wang for pointing out an error in the original version of this paper and for pointing out an interesting alternative approach to prove Theorem 1.

O.C. was partially supported by a Terman Fellowship, a Sloan Fellowship, and an NSF grant DMS-1811059/2016403. Y. L. was partially supported by NSERC Discovery grant. L. S. acknowledges the support of the NSF grant DMS-1951070.

1. GEOMETRIC PRELIMINARIES

In this section we introduce the main notations, we prove the existence of optimal nested sweepouts and we recall some useful geometric tools.

1.1. Notations. Let (M^{n+1}, g) be a closed Riemannian manifold. By scaling, it suffices to consider $\text{Vol}(M, g) = 1$, which we will always assume below. We use $\mathcal{Z}_n(M; \mathbb{Z}_2)$ to denote the space of mod 2 flat cycles in M . The topology on the space $\mathcal{Z}_n(M; \mathbb{Z}_2)$ is induced by the usual flat norm \mathcal{F} .

We will make extensive use of the notion of *Caccioppoli set*. A measurable set $E \subset M$ is Caccioppoli if

$$\text{Per}(E) := \sup \left\{ \int_M \chi_E \text{div} \omega : \omega \in X(M), \|\omega\|_{C^0} \leq 1 \right\} < \infty,$$

where χ_E denotes the indicator function of E . By De Giorgi's structure theorem we have that the distributional derivative $D\chi_E$ (which is a Radon measure) of a set of finite perimeter E is given by $D\chi_E = \nu_E \mathcal{H}^n \llcorner \partial^* E$, where $\partial^* E$ is the reduced boundary of E , which is a n -rectifiable set, and ν_E is the normal direction to $\partial^* E$ pointing outside E defined \mathcal{H}^n -a.e.. This allows us to identify $D\chi_E$ with an element of $\mathcal{Z}_n(M; \mathbb{Z}_2)$, which we will abuse notation and denote by

$$\partial E := \nu_E \mathcal{H}^n \llcorner \partial^* E.$$

In particular, with this identification we have

$$\mathbf{M}(\partial E) = \text{Per}(E) \quad \text{and} \quad \mathcal{F}(\partial E, \partial F) = \|\chi_E - \chi_F\|_{L^1} = \text{Vol}(E \Delta F),$$

where $E \Delta F$ denotes the symmetric difference between two sets. As usual, the perimeter of E in an open set U , denoted by $\text{Per}(E | U)$, is the total variation of $D\chi_E$ in the set U .

We let \mathcal{V} denote the set of varifolds of (M^{n+1}, g) . Given a cycle $\Gamma \in \mathcal{Z}_n(M; \mathbb{Z}_2)$, we will write $|\Gamma|$ for the associated integral varifold. In particular if $\Gamma = \partial\Omega$, then

$$|\Gamma| = \mathcal{H}^n \llcorner \partial^* \Omega \otimes \delta_{T\partial^* \Omega} \quad \text{and} \quad \mu_\Gamma := \mathcal{H}^n \llcorner \partial^* \Omega$$

is the total variation measure of the measure $D\chi_\Omega$. Finally, given a set Σ regular outside a set of dimension $n - 7$, we will denote with $|\Sigma|$ the associated integral varifold.

1.2. Optimal nested sweepouts. We start by recalling the notion of sweepouts that we will use in this paper.

Definition 5 (Sweepout). A sweepout of M is a map $\Phi : [0, 1] \rightarrow \mathcal{Z}_n(M; \mathbb{Z}_2)$ continuous in \mathcal{F} -topology such that $\Phi(x) = \partial\Omega(x)$, where $\{\Omega(x) : x \in [0, 1]\}$ is a family of Caccioppoli sets with $\Omega(0)$ the 0-cycle and $\Omega(1) = M$. We will denote with \mathcal{S} the collection of all such sweepouts. Moreover, we define the width W to be

$$W = \inf_{\Phi \in \mathcal{S}} \sup_{x \in [0, 1]} \mathbf{M}(\Phi(x)).$$

It is a consequence of the isoperimetric inequality that $W > 0$.

We will switch freely between the equivalent notation $\mathbf{M}(\Phi(x))$ and $\text{Per}(\Omega(x))$. We now introduce the notion of optimal nested sweepouts and prove their existence.

Definition 6 (Optimal nested volume parametrized (ONVP) sweepout). A sweepout $\{\Phi(x) = \partial\Omega(x) : x \in [0, 1]\}$ is called

- *optimal* if $\sup_{x \in [0, 1]} \mathbf{M}(\Phi(x)) = W$;
- *nested* if $\Omega(x_1) \subset \Omega(x_2)$, for all $0 \leq x_1 \leq x_2 \leq 1$;
- *volume parametrized* if $\text{Vol}(\Omega(x)) = x$, for every $x \in [0, 1]$ (recall that we have assumed $\text{Vol}(M, g) = 1$).

Nested volume parametrized sweepouts enjoy nice compactness properties.

Lemma 7 (Compactness for nested volume parametrized sweepouts). *Let $(\Phi_i)_i$ be a sequence of nested volume-parametrized sweepouts with mass uniformly bounded, that is*

$$(2) \quad \sup_{i \in \mathbb{N}} \sup_{x \in [0, 1]} \mathbf{M}(\Phi_i(x)) \leq M < \infty.$$

Then there exists a subsequence $(\Phi_{i_k})_k$ converging uniformly to a nested volume parametrized sweepout Ψ such that

$$(3) \quad \sup_x \mathbf{M}(\Psi(x)) \leq \liminf_k \left(\sup_x \mathbf{M}(\Phi_{i_k}(x)) \right).$$

Proof. The sequence of continuous functions $\Phi_i : [0, 1] \rightarrow \mathcal{Z}_{n-1}(M; \mathbb{Z}_2)$ is uniformly Lipschitz continuous, since for every $0 \leq x < y \leq 1$ we have

$$\mathcal{F}(\Phi_i(x), \Phi_i(y)) \leq \text{Vol}(\Omega_i(y) \setminus \Omega_i(x)) = \text{Vol}(\Omega_i(y)) - \text{Vol}(\Omega_i(x)) = y - x$$

and $\Phi_i(0) = \emptyset$ for every i , so by Arzelà–Ascoli Theorem there exists a subsequence Φ_{i_k} and a nested volume parametrized sweepout $\Psi : [0, 1] \rightarrow \mathcal{Z}_n(M; \mathbb{Z}_2)$ such that Φ_{i_k} converges uniformly to Ψ . Then (3) follows from (2) and the lower semi-continuity of \mathbf{M} with respect to the flat topology. \square

Optimal nested volume parametrized sweepouts exist.

Theorem 8 (Existence of (ONVP) sweepouts). *For any closed Riemannian manifold (M, g) there exists an optimal nested volume-parametrized sweepout.*

Proof. Let $\{\Psi_i\}_i$ be a min-max sequence of sweepouts with $\lim_{i \rightarrow \infty} \sup_x \mathbf{M}(\Psi_i(x)) = W$. By [CL20, Theorem 1.4] we can replace $\{\Psi_i(x)\}_i$ by a sequence of nested sweepouts $\{\Phi_i\}_i$, such that $\Phi_i(x) = f_i^{-1}(x)$ for some Morse function $f_i : M \rightarrow [0, 1]$ and $\lim_{i \rightarrow \infty} \sup_x \mathbf{M}(\Phi_i(x)) = W$. Let $\phi_i(x) = \text{Vol}(f_i^{-1}([0, x]))$. Note that $\phi_i : [0, 1] \rightarrow [0, 1]$ is a continuous strictly increasing function. Then $(\Phi_i \circ \phi_i^{-1})_i$ is a sequence of nested volume-parametrized sweepouts. By Lemma 7, a subsequence of $\{\Phi_i \circ \phi_i^{-1}\}_i$ converges to a (ONVP) sweepout. \square

Finally we recall the definition of critical set for a sweepout.

Definition 9 (Critical set). Given a sweepout Φ , we define

$$M(x) = \limsup_{r \rightarrow 0} \{\mathbf{M}(\Phi(y)) : |y - x| < r\}.$$

If Φ is an optimal sweepout, we define the *critical domain* of Φ to be the set

$$\mathbf{m}(\Phi) = \{x \in [0, 1] : M(x) = W\}.$$

We will say that a sequence $x_i \rightarrow x \in \mathbf{m}(\Phi)$ is a *min-max sequence* if $|\Phi|(x_i)$ converges in the varifold sense to a varifold V of mass W , i.e. $|V|(M) = W$. We denote the set of such varifolds by $\mathbf{C}(\Psi)$.

In fact, it is convenient to refine this definition somewhat.

Definition 10 (Left and right critical set). Given a sweepout Φ , we say that $x \in \mathbf{m}_L(\Phi)$ if there are $x_i \nearrow x$ with

$$\mathbf{M}(\Psi(x_i)) \rightarrow W$$

and similarly, $x \in \mathbf{m}_R(\Phi)$ if there are $x_i \searrow x$ with

$$\mathbf{M}(\Psi(x_i)) \rightarrow W.$$

Note that $\mathbf{m}(\Phi) = \mathbf{m}_L(\Phi) \cup \mathbf{m}_R(\Phi)$ (and $\mathbf{m}_L(\Phi) \cap \mathbf{m}_R(\Phi)$ need not necessarily be empty).

Definition 11 (Varifolds of optimal regularity). For an open set $U \subset M$ we say that a varifold V is in $\mathcal{R}(U)$ if

$$V \llcorner U = \sum_{k=1}^K \kappa_k |\Sigma_k|,$$

for $\Sigma_i \subset U$ embedded minimal hypersurfaces that are regular away from a closed singular set of Hausdorff dimension $n - 7$, and $\kappa_i \in \mathbb{N}$ integer multiplicities. We set $\mathcal{R} = \mathcal{R}(M)$.

Abusing notation, we will say that a (singular) hypersurface Σ is in $\mathcal{R}(U)$ if the associated varifold satisfies $|\Sigma| \in \mathcal{R}(U)$.

As we will discuss later, Almgren–Pitts theory [Pit81] implies that given an optimal sweepout Ψ , there is $V \in \mathbf{C}(\Psi)$ with $V \in \mathcal{R}$.

1.3. (Homotopic) one sided minimizers. The notions of one sided minimizers and homotopic minimizers will be useful when dealing with our deformations theorems, as we will see in Section 3.

Let $U \subset M^{n+1}$ be an open set. Given a Caccioppoli set $E \subset \Omega$ with $E\Delta\Omega \subset U$ and $\varepsilon \geq 0$, we denote the inner families of deformations between E and Ω in U which do not increase the volume more than ε by

$$\mathcal{I}_\varepsilon(\Omega, E | U) := \left\{ \begin{array}{l} \Omega(0) = E, \Omega(1) = \Omega, (\Omega(t)\Delta\Omega) \setminus U = \emptyset, \\ \{\Omega(t)\}_{t \in [0,1]} : \quad \Omega(t_1) \subset \Omega(t_2) \text{ for } t_1 < t_2, \\ \text{Per}(\Omega_t) + \varepsilon \leq \text{Per}(\Omega) \end{array} \right\},$$

and analogously for $\Omega \subset E$ with $E\Delta\Omega \subset U$, we define the outer families of deformations between Ω and E in U which do not increase the volume more than ε by

$$\mathcal{O}_\varepsilon(\Omega, E | U) := \left\{ \begin{array}{l} \Omega(0) = \Omega, E = \Omega(1), (\Omega(t)\Delta\Omega) \setminus U = \emptyset, \\ \{\Omega(t)\}_{t \in [0,1]} : \quad \Omega(t_1) \subset \Omega(t_2) \text{ for } t_1 < t_2 \\ \text{Per}(\Omega_t) \leq \text{Per}(\Omega) + \varepsilon \end{array} \right\}.$$

Moreover, given Ω and $\varepsilon \geq 0$, we denote the collections of inner and outer Caccioppoli sets that can be reached by an inner or outer family of deformations by

$$\begin{aligned} \mathcal{I}_\varepsilon(\Omega | U) &= \{E \subset \Omega : E\Delta\Omega \subset U, \mathcal{I}_\varepsilon(\Omega, E | U) \neq \emptyset\}, \\ \mathcal{O}_\varepsilon(\Omega | U) &= \{E \supset \Omega : E\Delta\Omega \subset U, \mathcal{O}_\varepsilon(\Omega, E | U) \neq \emptyset\}. \end{aligned}$$

In both definitions, if we do not include an ε -subscript, it should be understood that we are taking $\varepsilon = 0$; this will happen most of the time below, but we will crucially rely on the definition with $\varepsilon > 0$ to obtain *regular* homotopic minimizers in certain situations.

Definition 12 (Homotopic inner and outer minimizers). Given a Caccioppoli set Ω we say that a Caccioppoli set $L(\Omega | U) \in \mathcal{I}(\Omega | U)$ is a *homotopic inner minimizer* for Ω in U , if

- (1) $\text{Per}(L(\Omega | U) | U) \leq \text{Per}(\Omega' | U)$, for every $\Omega' \in \mathcal{I}(\Omega | U)$ and
- (2) if $E \in \mathcal{I}(\Omega | U)$ satisfies (1) and $E\Delta L(\Omega | U) \subset M \setminus L(\Omega | U)$ then $E = L(\Omega | U)$.

Similarly, define $R(\Omega | U) \in \mathcal{O}(\Omega | U)$ to be a *homotopic outer minimizer* for Ω in U , if

- (1) $\text{Per}(R(\Omega | U) | U) \leq \text{Per}(\Omega' | U)$, for every $\Omega' \in \mathcal{I}(\Omega | U)$;
- (2) if $E \in \mathcal{I}(\Omega | U)$ satisfies (1) and $E\Delta R(\Omega | U) \subset R(\Omega | U) \setminus E$ then $E = R(\Omega | U)$.

We say that a Caccioppoli set Ω is an *inner (resp. outer) homotopic minimizer* in U if Ω is a homotopic inner (resp. outer) minimizer relative to itself.

It is easy to see that inner and outer homotopic minimizers for a fixed set Ω always exist.

Lemma 13 (Existence of homotopic minimizers). *For any Caccioppoli set Ω and open set U we can find a homotopic inner (resp. outer) minimizer $L(\Omega | U)$ (resp. $R(\Omega | U)$) for Ω in U . Moreover, if $E \in \mathcal{I}(\Omega | U)$ (resp. $E \in \mathcal{O}(\Omega | U)$) then*

- (4) $\text{Per}(L(\Omega | U) | U) \leq \text{Per}(E | U)$ (resp. $\text{Per}(R(\Omega | U) | U) \leq \text{Per}(E | U)$).

In particular, if

$$\text{Per}(E | U) < \text{Per}(\Omega | U)$$

then $L(\Omega | U)$ (resp. $R(\Omega | U)$) does not coincide with Ω .

Proof. We consider only the case of inner minimizers as the outer minimizers are handled identically.

This is once again an application of Arzelà–Ascoli theorem. Indeed, notice that $\mathcal{I}(\Omega | U) \neq \emptyset$, since $\Omega \in \mathcal{I}(\Omega | U)$, so we can consider a minimizing sequence $(E_j)_j$, that is

$$\lim_j \text{Per}(E_j | U) = \inf \{ \text{Per}(E | U) : E \in \mathcal{I}(\Omega | U) \}$$

and let $\{E_j(x) : x \in [0, 1]\} \in \mathcal{I}(\Omega, E_j; U)$ be the corresponding inner volume non increasing sweepout between E_j and Ω . We can assume that it is volume parametrized (being nested). Moreover $\text{Per}(E_j(x) | U)$ is uniformly bounded by $\text{Per}(\Omega | U)$, so by Arzelà–Ascoli there is a subsequence converging to $\{E_\infty(x)\} \in \mathcal{I}(\Omega, E_\infty | U)$, with E_∞ satisfying the desired minimality property by lower semi-continuity of the perimeter.

Finally, again by Arzelà–Ascoli, we can find $L(\Omega | U) \subset \Omega$ in the set of minimizers, which infimizes the flat distance to $\partial\Omega$, and so satisfies condition (2) (otherwise there would be a competitor closer to Ω in flat norm). \square

We recall the definition of one-sided minimizers, which will be useful in the sequel when we perform cut and paste arguments.

Definition 14 (One sided minimizers). Let E be a Caccioppoli set. We say that E is *locally one-sided inner* (resp. *outer*) *area-minimizing* in U if for every $A \Subset U$ and V with $V \Delta E \subset A$, we have

$$\text{Per}(E | A) \leq \text{Per}(V | A)$$

whenever $V \subset E$ (resp. $E \subset V$). We say that E is *strictly locally one-sided inner* (resp. *outer*) *area-minimizing* if the inequality holds strictly except when $E = V$ as Caccioppoli sets.

We show that homotopic minimizers are in fact strict one sided minimizers into the region they sweep out.

Lemma 15 (Homotopic minimizers are one sided minimizers in the swept out region). *Suppose $L(\Omega | U)$ is an homotopic inner (resp. outer) minimizer for Ω in U . Then $L(\Omega | U)$ (resp. $R(\Omega | U)$) is strict locally outer (resp. inner) one-sided minimizing in $U \cap \Omega$ (resp. $U \setminus \Omega$).*

Proof. We consider homotopic inner minimizers; the case of outer minimizers is similar.

If $L(\Omega | U)$ is not a strict outer minimizer in $U \cap \Omega$ then there is V' with $L(\Omega | U) \subset V'$ and $L(\Omega | U) \Delta V' \subset A \Subset U$ and

$$\text{Per}(V' | A) \leq \text{Per}(L(\Omega | U) | A).$$

We can minimize perimeter in A among all such V' to find V . Namely,

$$(5) \quad \text{Per}(V | A) \leq \text{Per}(W | A)$$

for all W with $W\Delta V \subset A \setminus L(\Omega|U)$. Since $L(\Omega|U) \in \mathcal{I}(\Omega|U)$, there is $\{U(x) : x \in [0, 1]\} \in \mathcal{I}(\Omega, L(\Omega|U)|U)$. Set $\Omega(x) = U(x) \cup V$. Since V satisfies (5), we have that

$$\text{Per}(\Omega_t|A) \leq \text{Per}(U_t|A).$$

This implies that $\Omega(1) = V$ satisfies (1) of Definition 12 and $V\Delta L(\Omega|U) \subset A \setminus L(\Omega|U)$, therefore by (2) of Definition 12, it follows that $V = L(\Omega|U)$. This completes the proof. \square

We have the following lemma that will allow us to find bounded mass homotopies in certain situations.

Lemma 16 (Interpolation lemma). *Fix $L > 0$. For every $\varepsilon > 0$ there exists $\delta > 0$, such that the following holds. If Ω_0, Ω_1 are two sets of finite perimeter, such that $\Omega_0 \subset \Omega_1$, $\text{Per}(\Omega_i) \leq L$ and $\text{Vol}(\Omega_1 \setminus \Omega_0) \leq \delta$, then there exists a nested \mathcal{F} -continuous family $\{\partial\Omega_t\}_{t \in [0,1]}$ with*

$$\text{Per}(\Omega_t) \leq \max\{\text{Per}(\Omega_1), \text{Per}(\Omega_2)\} + \varepsilon$$

for all $t \in [0, 1]$

Proof. Let Ω be a Caccioppoli set that minimizes perimeter among sets Ω' with $\Omega_0 \subset \Omega' \subset \Omega_1$.

Fix $r > 0$ such that for every $x \in M$ the ball $B(x, 2r)$ is 2-bi-Lipschitz diffeomorphic to the Euclidean ball of radius $2r$. Let $\{B(x_i, r)\}_{i=1}^N$ be a collection of balls covering M . By coarea inequality we can find a radius $r_i \in [r, 2r]$, so that $\mathbf{M}(\partial B(x_i, r_i) \cap \Omega \setminus \Omega_0) \leq \frac{\delta}{r}$.

Let $U_1 = B(x_1, r_1) \cap \Omega \setminus \Omega_0$. By a result of Falconer (see [Fal80], [Gut07, Appendix 6]) there exists a family of hypersurfaces sweeping out U_1 of area bounded by $c(n)\delta^{\frac{n}{n+1}}$. It follows (see [CL20, Lemma 5.3]) that there exists a nested family $\{\Xi^1(t)\}$ of Caccioppoli sets with $\Xi^1(0) = \Omega_0$ and $\Xi^1(1) = \Omega_0 \cup U_1$ and satisfying

$$\text{Per}(\Xi^1(t)) \leq \text{Per}(\Omega_0) + 2c(n)\delta^{\frac{n}{n+1}}$$

Let $\Omega^1 = \Omega_0 \cup U_1$. Observe, that the minimality of Ω implies that

$$\text{Per}(\Omega^1) \leq \text{Per}(\Omega_0) + \frac{2\delta}{r}$$

Inductively, we define $\Omega^k = \Omega^{k-1} \cup U_k$ and $U_k = B(x_k, r_k) \cap \Omega \setminus \Omega^{k-1}$. As above we can construct a nested homotopy of Caccioppoli sets $\Xi^k(t)$ from Ω^{k-1} to Ω^k , satisfying

$$\text{Per}(\Xi^k(t)) \leq \text{Per}(\Omega_0) + 2c(n)\delta^{\frac{n}{n+1}} + \frac{2N\delta}{r}$$

We choose $\delta > 0$ so small that $\text{Per}(\Xi^k(t)) < \text{Per}(\Omega_0) + \varepsilon$. It follows then that we have obtained a homotopy from Ω_0 to Ω satisfying the desired perimeter bound. Similarly, we construct a homotopy from Ω to Ω_1 . \square

Finally, we have the following result. Recall that White [Whi94] proves that strictly stable *smooth* minimal hypersurfaces are locally area-minimizing. A generalization of such a result to the case of hypersurfaces with singularities (i.e., elements of \mathcal{R}) would be very interesting. The following (weaker) result will suffice for our needs; it can be

seen as a result along these lines, except “stability” is replaced by a stronger hypothesis: the surface is homotopic minimizing to one side.²

Proposition 17 (Comparing the notions of minimizing vs. homotopic minimizing for minimal surfaces). *Suppose that Ω is a Caccioppoli set and for some strictly convex open set $U \subset M$ with smooth boundary, the associated varifold $V = |\partial\Omega|$ satisfies $V \in \mathcal{R}(U)$. Assume that $\text{supp } V \cap U$ is connected.*

Suppose that Ω is inner (resp. outer) homotopy minimizing in U . Then, at least one of the following two situations holds:

- (1) *for all $p \in \text{supp } V \cap U$, there is $\rho_0 > 0$ so that for $\rho < \rho_0$, $B_\rho(p) \subset U$ and Ω is inner (resp. outer) minimizing in $B_\rho(p)$, or*
- (2) *there exists a sequence of Caccioppoli sets $E_i \neq \Omega$ with $|\partial E_i| \in \mathcal{R}(U)$ so that $E_i \Delta \Omega \subset \Omega \cap U$ (resp. $\Omega^c \cap U$), $|\partial E_i|$ has stable regular part, and $\partial E_i \rightarrow \partial\Omega$ in the flat norm.*

Remark. It is interesting to ask if the second possibility occurs; it seems possible that one could rule this out in the case where V has regular tangent cones that are all strictly minimizing in the sense of Hardt–Simon [HS85, §3].

Proof of Proposition 17. We consider the “inner” case, as the “outer” case is similar. Let $E^\varepsilon \in \mathcal{I}_\varepsilon(\Omega|U)$ minimize perimeter among all sets in $\mathcal{I}_\varepsilon(\Omega|U)$ (as usual, the existence of E^ε follows from Arzelà–Ascoli). We claim that E^ε is area-minimizing to the inside of Ω in sufficiently small balls.

More precisely, for $r > 0$ sufficiently small, suppose there was a Caccioppoli set E' so that $E' \Delta E^\varepsilon \subset B_r(p) \cap U \cap \Omega$ and $\text{Per}(E'|U) < \text{Per}(E^\varepsilon|U)$. As long as r was chosen sufficiently small, Lemma 16 guarantees that $E' \in \mathcal{I}_\varepsilon(\Omega|U)$. This is a contradiction.

Now, consider $p \in \text{Reg } V \cap U$. We note that E^ε is almost minimizing (with no constraint coming from Ω) in the sense of [Tam84], and thus has $C^{1,\alpha}$ boundary in $B_r(p) \cap U$, thanks to standard results on the obstacle problem; see [Tam84, §1.9, §1.14(iv)]. As such, away from $\text{Sing } V$ (which has Hausdorff dimension at most $n - 7$) we can thus conclude that $\partial^* E^\varepsilon$ is regular, stationary and stable.³ A capacity argument then implies that $|\partial E^\varepsilon| \in \mathcal{R}(U)$ and $\partial^* E^\varepsilon$ is stable. Therefore, the maximum principle for (possibly singular) hypersurfaces [Ilm96] implies that either $E^\varepsilon = \Omega$ or $\partial^* E^\varepsilon \cap \text{supp } V = \emptyset$. In the first case, we can conclude that Ω is inner minimizing in small balls (since E^ε is).

We can thus assume that the latter possibility holds for all $\varepsilon > 0$ sufficiently small. Taking $\varepsilon_j \rightarrow 0$, there is $E \in \mathcal{I}_0(\Omega|U)$ so that $E^{\varepsilon_j} \rightarrow E$ with respect to the flat norm. If $E = \Omega$, then the second possibility in the conclusion of the proposition holds for $E_j = E^{\varepsilon_j}$.

The final case to consider is $E \neq \Omega$. By curvature estimates for stable minimal hypersurfaces [SS81], $|\partial E| \in \mathcal{R}(\Omega)$ and thus $\partial^* E \cap \text{supp } V = \emptyset$ again by the maximum

²Note that one certainly needs a condition on the singularities rather than just a condition on the regular part like strict stability, since as we show in Proposition 24, the existence of (regular) non-minimizing tangent cones implies that the hypersurface is not homotopic minimizing irrespective of any stability condition that might hold on the regular part.

³Cf. the proof of [Liu19, Proposition 2.1] for the proof of stability.

principle. By assumption that Ω is inner perimeter minimizing, we have that

$$\text{Per}(E | U) = \text{Per}(\Omega | U).$$

Using Lemma 15, we thus see that Ω is minimizing in $\Omega \cap E^c \cap U$, which implies that it is inner minimizing in small balls, as asserted. \square

2. NON-EXCESSIVE SWEEPOUTS

In this section we introduce the concept of excessive intervals and excessive points for a sweepout and prove that there is a sweepout, such that every point in the critical domain is not excessive.

Definition 18 (Excessive points and intervals). Suppose $\{\Phi(x) = \partial\Omega(x)\}$ is a sweepout. Given a connected interval I (we allow I to be open, closed, or half-open) we will say that $\{\Phi^I(x) = \partial\Omega^I(x)\}_{x \in \bar{I}}$ is an I -replacement family for Φ if $\Omega^I(a) = \Omega(a)$, $\Omega^I(b) = \Omega(b)$ and for all $x \in I$,

$$\limsup_{I \ni y \rightarrow x} \mathbf{M}(\Phi^I(y)) < W.$$

We say that a connected interval I is an *excessive interval* for Φ if there is an I -replacement family for Φ . We say that a point x is *left (resp. right) excessive* for Φ if there is an excessive interval I for Φ so that $(x - \varepsilon, x] \subset I$ (resp. $[x, x + \varepsilon) \subset I$) for some $\varepsilon > 0$.

The goal of this section is to prove the following result.

Theorem 19 (Existence of non-excessive min-max hypersurface). *There exists a (ONVP) sweepout Ψ such that every $x \in \mathbf{m}_L(\Psi)$ is not left excessive and every $x \in \mathbf{m}_R(\Psi)$ is not right excessive.*

2.1. Preliminary results. We establish several results that will be used in the proof Theorem 19.

Lemma 20 (Extension lemma I). *If I, J are excessive for Φ and $I \cap J \neq \emptyset$, then $I \cup J$ is excessive for Φ .*

Proof. Let $\{\partial\Omega^I(x)\}_{x \in I}$ and $\{\partial\Omega^J(x)\}_{x \in J}$ be I and J replacement families for Φ .

Let $a_1 = \inf\{x \in I\}$, $a_2 = \inf\{x \in J\}$ and $b_1 = \sup\{x \in I\}$, $b_2 = \sup\{x \in J\}$. Assume without any loss of generality that $a_1 \leq a_2$ and $b_1 \leq b_2$ and at least one of the two inequalities is strict.

Let $K = I \cap J$; let a, b denote, respectively, left and right boundary points of K and $c = \frac{a+b}{2} \in K$. Let $\tilde{\Omega}$ be a Caccioppoli set minimizing perimeter among all Ω' with $\Omega(a) \subset \Omega' \subset \Omega(b)$. Define $\phi_1 : [a_1, c] \rightarrow [a_1, b_1]$ and $\phi_2 : [c, b_2] \rightarrow [a_2, b_2]$ given by $\phi_1(x) = a_1 + \frac{b_1 - a_1}{c - a_1}(x - a_1)$ and $\phi_2(x) = a_2 + \frac{b_2 - a_2}{b_2 - c}(x - c)$. We define an $I \cup J$ replacement family for Φ by setting

$$\Phi^{I \cup J}(x) = \begin{cases} \partial(\Omega^I(\phi_1(x)) \cap \tilde{\Omega}) & x \in [a_1, c] \\ \partial(\Omega^J(\phi_2(x)) \cup \tilde{\Omega}) & x \in [c, b_2] \end{cases}$$

Observe that $\Phi^{I \cup J}$ is continuous since $\Phi^{I \cup J}(c) = \partial \tilde{\Omega}$. It follows from our choice of $\tilde{\Omega}$ that $\mathbf{M}(\Phi^{I \cup J}(x)) \leq \mathbf{M}(\Phi^I(\phi_1^{-1}(x))) < W$ for $x \in I \cap (-\infty, c]$ and $\mathbf{M}(\Phi^{I \cup J}(x)) \leq \mathbf{M}(\Phi^J(\phi_2^{-1}(x))) < W$ for $x \in J \cap [c, \infty)$. \square

Lemma 21 (Extension lemma II). *If I is excessive for Φ and J has $J \cap I \neq \emptyset$ and is excessive for*

$$\Psi(x) := \begin{cases} \Phi^I(x) & x \in I \\ \Phi(x) & x \notin I \end{cases}$$

then $J \cup I$ is excessive for Φ .

Proof. Define an $I \cup J$ -replacement family $\Phi^{I \cup J}$ for Φ by

$$\Phi^{I \cup J}(x) = \begin{cases} \Phi(x) & x \in [0, 1] \setminus (I \cup J) \\ \Phi^I(x) & x \in I \setminus J \\ \Psi^J(x) & x \in J \end{cases}$$

where Ψ^J is a J -replacement family for Ψ . \square

The following is the technical core of the proof of Theorem 19.

Proposition 22 (Existence of maximal excessive intervals). *Given an (ONVP) sweep-out Φ , if \hat{J} is excessive for $\{\Phi(x) = \partial \Omega(x)\}$, then there exists an excessive interval $J \supset \hat{J}$ so that J is maximal in the sense that if \tilde{J} is excessive with $\tilde{J} \cap J \neq \emptyset$, then $\tilde{J} \subset J$.*

Proof. Let

$$\alpha := \sup\{|\tilde{J}| : \tilde{J} \text{ excessive, } \hat{J} \cap \tilde{J} \neq \emptyset\}.$$

Choose excessive intervals \tilde{J}_n with $\tilde{J}_n \cap \hat{J} \neq \emptyset$ and $|\tilde{J}_n| \rightarrow \alpha$. By Lemma 20, we can replace \tilde{J}_n by $\tilde{J}_n \cup \hat{J}$, and thus assume that $\hat{J} \subset \tilde{J}_n$. In particular $\tilde{J}_n \cap \tilde{J}_m \neq \emptyset$ for all m, n . Using Lemma 20 again, we can replace \tilde{J}_n by

$$\bigcup_{m=1}^n \tilde{J}_m$$

so that the \tilde{J}_n form an increasing sequence of excessive intervals (still with $|\tilde{J}_n| \rightarrow \alpha$). Note that the interior of an excessive interval is still excessive, so we can consider $J_n := (\tilde{J}_n)^\circ$. Note that $|J_n| \rightarrow \alpha$ and the J_n are increasing.

We will show below that

$$J' := \bigcup_n J_n$$

is excessive. Write $J' = (a, b)$. Granted the fact that J' is excessive, we claim that one of the intervals (a, b) , $(a, b]$, $[a, b)$, or $[a, b]$ is the desired maximal excessive interval. Note that by Lemma 20, if $[a, b)$ and $(a, b]$ are excessive, then so is $[a, b]$, so we can choose the largest excessive interval out of these four choices and call it J . Suppose that \tilde{J} is excessive with $\tilde{J} \cap J \neq \emptyset$. Then, $J \cup \tilde{J}$ is excessive by Lemma 20 and $\hat{J} \subset J \cup \tilde{J}$. Thus,

$$|J \cup \tilde{J}| \leq \alpha,$$

so $\tilde{J} \subset \bar{J}$ (where \bar{J} is the closure of J). Now, $J \cup \tilde{J}$ is excessive, but strictly larger than \tilde{J} (by assumption). This contradicts the choice of J as the largest excessive interval out of (a, b) , $(a, b]$, $[a, b)$, and $[a, b]$. This shows that J is maximal, as desired.

It thus remains to prove that $J' = \cup_n J_n$ is excessive for a nested sequence of open excessive intervals J_n . Write $J_n = (a_n, b_n)$ and set $a'_n = a_n + \frac{1}{n}$, $b'_n = b_n - \frac{1}{n}$.

Fix $i = 0, 1, \dots$ and assume we have real numbers $0 < A_1, \dots, A_i < W$ and integers $n_i \geq i$ (with $n_1 < n_2 < \dots < n_i$) so that for $n \geq n_i$, there is a J_n -replacement $\{\Phi_i^n(x) = \partial\Omega_i^n(x)\}$ so that

$$\text{Per}(\Omega_i^n(x)) \leq A_j$$

for $x \in [a'_j, b'_j]$ and $1 \leq j \leq i$. (Note that for $i = 0$, we can find such objects because the J_n are excessive.)

We will choose $0 < A_{i+1} < W$, and $n_{i+1} > \max\{n_i, i + 1\}$ so that we can construct J_n -replacements $\{\Phi_{i+1}^n(x) = \partial\Omega_{i+1}^n(x)\}$ for $n \geq n_{i+1}$ with

$$\text{Per}(\Omega_{i+1}^n(x)) \leq A_j$$

for $x \in [a'_j, b'_j]$ and $1 \leq j \leq i + 1$. Granted this, we can easily (inductively) complete the proof by passing Φ_{i+1}^n to a subsequential limit (using Arzelà–Ascoli).

It is useful to introduce the following notation, used in the construction of Φ_{i+1}^n . Given two nested sets of finite perimeter $V \subset W$, we let

- $\mathcal{M}_{V,W}$ an outermost Caccioppoli set minimizing perimeter among all the Caccioppoli sets Ω with $V \subset \Omega \subset W$;
- $\{\mathcal{V}_{(V,W)}(x)\}_x$ the optimal nested homotopy from V to W .

For $n \geq n_i$, we set

$$L_n := \mathcal{M}_{\Omega(a_n), \Omega_i^{n_i}(a'_{i+1})}, \quad U_n := \mathcal{M}_{\Omega_i^{n_i}(b'_{i+1}), \Omega(b_n)}$$

Note that for $n \leq m$, $L_m \subset L_n$ and $U_n \subset U_m$. Hence, L_n and U_n have \mathcal{F} -limits as $n \rightarrow \infty$. For $\varepsilon > 0$ fixed so that

$$\max\{\text{Per}(\Omega_i^{n_i}(a'_{i+1})), \text{Per}(\Omega_i^{n_i}(b'_{i+1}))\} + \varepsilon < W,$$

Lemma 16 thus guarantees that there is $n_{i+1} \geq i + 1$ sufficiently large so that for $n \geq n_{i+1}$,

$$\sup_t \text{Per}(\mathcal{V}_{(L_n, L_{n_1})}(t)) < W, \quad \sup_t \text{Per}(\mathcal{V}_{(U_{n_1}, U_n)}(t)) < W.$$

For $n \geq n_{i+1}$, we define

$$\tilde{\Phi}_{i+1}^n(x) = \begin{cases} \partial(\Omega_i^n(x+1) \cap L_n) & x \in [a_n - 1, b_n - 1] \\ \partial\tilde{\mathcal{V}}_{L_n, L_{n_i}}(x) & x \in [b_n - 1, a_{n_i}] \\ \partial(\Omega_i^{n_i}(x) \cup L_{n_i} \cap U_{n_i}) & x \in [a_{n_i}, b_{n_i}] \\ \partial\tilde{\mathcal{V}}_{U_{n_i}, U_n}(x) & x \in [b_{n_i}, a_n + 1] \\ \partial(\Omega_i^n(x-1) \cup U_n) & x \in [a_n + 1, b_n + 1]. \end{cases}$$

Here, the $\tilde{\mathcal{V}}$ are the homotopies \mathcal{V} reparametrized to be defined on the given intervals (the exact parametrization is immaterial). It is easy to check that $\tilde{\Phi}_{i+1}^n$ is continuous.

Let Φ_{i+1}^n denote the reparametrization of $\tilde{\Phi}_{i+1}^n$ by volume. We have arranged that Φ_{i+1}^n is a J_n -replacement. Moreover, for $x \in [a'_{i+1}, b'_{i+1}]$, we have that $\Phi_{i+1}^n(x) = \Phi_i^{n_i}(x)$, so

$$\mathbf{M}(\Phi_{i+1}^n(x)) \leq A_j$$

for $x \in [a'_j, b'_j]$ and $1 \leq j \leq i$. Finally, we can set

$$A_{i+1} := \sup_{x \in [a'_{i+1}, b'_{i+1}]} \mathbf{M}(\Phi_{i+1}^{n_i}(x)) < W$$

(which is independent of n). This completes the proof. □

2.2. Proof Theorem 19.

We are now able to complete proof of Theorem 19. Let Φ be a nested optimal sweepout. Consider the collection \mathcal{A} of the maximal (with respect to inclusion) excessive intervals for Φ , that is $I \in \mathcal{A}$ if for every excessive interval I' such that $I' \cap I \neq \emptyset$, we have $I \supset I'$. The existence of maximal intervals follows from Proposition 22 proven above.

Notice that by definition $I \neq J \in \mathcal{A}$ implies that $I \cap J = \emptyset$, so we can define a new sweepout Ψ in the following way

$$\Psi(x) = \begin{cases} \Phi^I(x) & \text{if } x \in I \in \mathcal{A} \\ \Phi(x) & \text{otherwise.} \end{cases}$$

Note that Ψ is a nested optimal sweepout, so up to reparametrization we can assume it is (ONVP), and moreover by construction $\mathbf{m}(\Psi) \subset \mathbf{m}(\Phi)$. Suppose that $x \in \mathbf{m}_L(\Psi)$ is left excessive. Then, there is a Ψ -excessive interval J with $(x - \varepsilon, x] \subset J$. We claim that there is $I \in \mathcal{A}$ with $J \subset I$. Indeed, if $J \cap I = \emptyset$ for all $I \in \mathcal{A}$, then J is a Φ -excessive interval, contradicting the definition of \mathcal{A} . On the other hand, if there is $I \in \mathcal{A}$ with $J \cap I \neq \emptyset$, then $J \cup I$ is excessive by Lemma 21. Thus, $J \subset I$ by definition of \mathcal{A} again. Thus, for $y \in (x - \varepsilon, x] \subset I$, $\Psi(y) = \Phi^I(y)$. By the definition of replacement family, we know that if $x_i \in (x - \varepsilon, x]$ has $x_i \rightarrow x$, then

$$\limsup_{i \rightarrow \infty} \mathbf{M}(\Psi^I(x_i)) < W.$$

However, this contradicts the assumption that $x \in \mathbf{m}_L(\Psi)$. The same proof works to prove that $x \in \mathbf{m}_R(\Psi)$ is not right excessive. This finishes the proof. □

3. DEFORMATION THEOREMS AND PROOF OF THEOREM 4

In this section we conclude the proof of Theorem 4. By Theorem 19, there exists an (ONVP) sweepout Φ so that every $x \in \mathbf{m}_L(\Phi)$ is not left excessive and every $x \in \mathbf{m}_R(\Phi)$ is not right excessive. By Almgren–Pitts pull-tight and regularity theory [Pit81], we find that for some $x_0 \in \mathbf{m}(\Phi)$, there is a min-max sequence $x_i \rightarrow x_0$ so that $|\Phi(x_i)|$ converges to some $V \in \mathcal{R}$. Indeed, we can pull-tight Φ to find a sweepout (in the sense of Almgren–Pitts, not in the (ONVP) sense considered in this paper) $\tilde{\Phi}$; we have that $\mathbf{C}(\tilde{\Phi}) \subset \mathbf{C}(\Phi)$ and some $V \in \mathbf{C}(\tilde{\Phi})$ is in \mathcal{R} . By replacing $\Phi(x)$ by $\Phi(1 - x)$ if necessary, we can then assume for the rest of this section that:

- (6) there is a (ONVP) sweepout $\{\Phi(x) = \partial\Omega(x)\}$ and $x_i \nearrow x_0 \in \mathbf{m}_L(\Phi)$, so that
- $$|\Phi(x_i)| \rightarrow V \in \mathcal{R} \text{ and } \Phi \text{ is not left excessive at } x_0$$

We then consider two cases: $\mathbf{M}(\Phi(x_0)) = W$ (no cancellation) and $\mathbf{M}(\Phi(x_0)) < W$ (cancellation). We analyze the geometric properties of V in both cases separately, proving deformation theorems reminiscent of those in [MN16].

3.1. No cancellation. Throughout this subsection we will assume the no cancellation condition

$$\mathbf{M}(\Phi(x_0)) = W .$$

In this case we have that $|\Phi(x_i)| \rightarrow |\partial\Omega|$, see for instance [DLT13b, Proposition A.1], so we can rephrase our assumption (6) as

$$(7) \quad \begin{aligned} & \text{there is a (ONVP) sweepout } \{\Phi(x) = \partial\Omega(x)\} \text{ and } x_i \nearrow x_0 \in \mathbf{m}_L(\Phi), \text{ so that} \\ & |\Phi(x_i)| \rightarrow |\Sigma| := |\partial\Omega| \in \mathcal{R} \text{ and } \Phi \text{ is not left excessive at } x_0. \end{aligned}$$

In particular, in this case the multiplicity bound of Theorem 4 follows immediately.

Proposition 23. *Let Σ be as in (7). Suppose Σ is not homotopic minimizing to either side in some open set U . Then the following holds:*

- (1) *for every $x \notin \bar{U}$ there exists $r > 0$, such that Σ is minimizing to one side in $B_r(x)$;*
- (2) *for every open set U' disjoint from U , we have that Σ is homotopic minimizing to one side in U' .*

Proof. We prove statement (1). There is $\delta > 0$ and Caccioppoli sets $E_1^- \in \mathcal{I}(\Omega | U)$ and $E_1^+ \in \mathcal{O}(\Omega | U)$ with

$$(8) \quad \text{Per}(E_1^\pm | U) \leq \text{Per}(\Omega | U) - \delta ,$$

and nested families $\{\Omega_1^-(x) : x \in [0, 1]\} \in \mathcal{I}(\Omega, E_1^- | U)$ and $\{\Omega_1^+(x) : x \in [0, 1]\} \in \mathcal{O}(\Omega, E_1^+ | U)$. Furthermore, by Lemma 13, we can assume that E_1^+ are inner and E_1^- are outer homotopic minimizers in U .

Let $x \in \Sigma \setminus \bar{U}$ and assume, for contradiction, that Σ is not area minimizing on both sides in every ball $B_r(x)$, $r < \text{dist}(x, U)$. Let $E_2^- \subset \Omega$, with $\Omega \setminus E_2^- \subset B_r(x)$, denote a Caccioppoli set that is a strict outer minimizer in $\Omega \cap B_r(x)$. Similarly, let $\Omega \subset E_2^+$, with $E_2^+ \setminus \Omega \subset B_r(x)$, denote a Caccioppoli set that is a strict inner minimizer in $\Omega \cap B_r(x)$. We have

$$\text{Per}(\Omega) > \max\{\text{Per}(E_2^\pm)\}$$

If we choose $r > 0$ sufficiently small, then, by Lemma 16, there exist nested families $\{\Omega_2^-(x) : x \in [0, 1]\}$ and $\{\Omega_2^+(x) : x \in [0, 1]\}$ that interpolate between E_2^- and Ω and between Ω and E_2^+ and satisfying

$$(9) \quad \text{Per}(\Omega_2^\pm(x)) \leq \text{Per}(\Omega) + \frac{\delta}{2}$$

Let $(x_l, x_r) \neq \emptyset$ be the interval (since Φ is nested) such that

$$\Phi(x) \cap (\cup_i E_i^+ \setminus \cup_i E_i^-) \neq \emptyset$$

Then we define a family $\bar{\Psi}: [x_l - 2, x_r + 2] \rightarrow \mathcal{Z}_n(M; \mathbb{Z}_2)$ by setting

$$\bar{\Psi}(x) := \begin{cases} \partial (\Omega(x+2) \cap E_1^- \cap E_2^-) & \text{if } x \in (x_l - 2, x_0 - 2] \\ \partial (\Omega_1^-(x - x_0 + 2) \cap E_2^-) & \text{if } x \in [x_0 - 2, x_0 - 1] \\ \partial (\Omega_1^+(x - x_0 + 1) \cap E_2^-) & \text{if } x \in [x_0 - 1, x_0] \\ \partial (\Omega_2^-(x - x_0) \cup E_1^+) & \text{if } x \in [x_0, x_0 + 1] \\ \partial (\Omega_2^+(x - x_0 - 1) \cup E_1^+) & \text{if } x \in [x_0 + 1, x_0 + 2] \\ \partial (\Omega(x-2) \cup E_1^+ \cup E_2^+) & \text{if } x \in [x_0 + 2, x_r + 2] \end{cases}$$

It is easy to see that $\bar{\Psi}$ is continuous, and moreover notice that, since by Lemma 15 E_1^+ is a strict inner minimizer in U and E_1^- strict outer minimizers in U , we have that

$$\limsup_{y \rightarrow x} \mathbf{M}(\bar{\Psi}(y)) < \limsup_{y \rightarrow x} \mathbf{M}(\bar{\Phi}(y)) \leq W$$

for $x \in (x_l - 2, x_0 - 2] \cup [x_0 - 2, x_r - 2)$. Since the families $\Omega_1^\pm(x)$ do not increase the volume of Σ in U_i and using (8) and (9), we also have

$$\mathbf{M}(\bar{\Psi}(x)) \leq W - \frac{\delta}{2} \quad \forall x \in [x_0 - 2, x_0 + 2].$$

We let Ψ be the volume reparametrization of the nested sweepout $\bar{\Psi}$, then Ψ is a (x_l, x_r) -replacement for Φ , thus giving a contradiction with the fact that $x_0 \in (x_l, x_r)$ and $x_0 \in \mathbf{m}_L(\Phi)$.

The proof of statement (2) is completely analogous. \square

Proposition 24. *Let Σ be as in (7), then the following holds*

- (1) $\text{Index}(\Sigma) \leq 1$;
- (2) *If $\text{Index}(\Sigma) = 1$, then for every point $x \in \Sigma$ there exists $r > 0$, such that Σ is minimizing to one side in $B_r(x)$;*
- (3) *If $\mathfrak{h}_{\text{nm}}(\Sigma)$ is non-empty, then Σ is stable, $\mathcal{H}^0(\mathfrak{h}_{\text{nm}}(\Sigma)) = 1$ and for every point $x \in \Sigma \setminus \mathfrak{h}_{\text{nm}}(\Sigma)$ there exists $r > 0$, such that Σ is minimizing to one side in $B_r(x)$.*

In particular, Theorem 4 holds in the case of no cancellations.

Proof. Note that if $U \cap \Sigma$ is smooth and unstable, it is easy to see that Σ is not homotopic minimizing to either side in U (just consider the normal flow generated by a compactly supported unstable variation of fixed sign). Statements (2) and (3) of the Proposition now immediately follow from Proposition 23. The upper bound on the index (1) follows from (2) of Proposition 23 and Lemma 25 below. \square

Lemma 25 (Localizing the index). *Suppose that $\Sigma \in \mathcal{R}$ is two-sided and has $\text{Index}(\Sigma) \geq 2$. Then, there is $\Sigma_1^*, \Sigma_2^* \subset \Sigma$ smooth hypersurfaces with boundary so that the Σ_i^* are both unstable (for variations fixing the boundary).*

Proof. A standard capacity argument implies that there is a subset $\Sigma' \subset \Sigma$ where Σ' is a smooth minimal surface with smooth boundary and $\text{Index}(\Sigma') \geq 2$ (with Dirichlet boundary conditions). Let u denote the second (Dirichlet) eigenfunction (with eigenvalue $\lambda < 0$) for the stability operator for Σ . Because u must change sign, there are

at least two nodal domains $\Sigma_1, \Sigma_2 \subset \Sigma$. One can find subsets with smooth boundary $\Sigma_i^* \subset \Sigma_i$ so that Σ_i^* are unstable. This follows from the argument in [Cha84, p. 21] (namely, by considering $(u|_{\Sigma_i} - \varepsilon)_+$ in the stability operator for $\varepsilon \rightarrow 0$ chosen so that $\{u|_{\Sigma_i} > \varepsilon\}$ has smooth boundary). \square

Lemma 26. $\mathcal{S}_{\text{nm}}(\Sigma) \subset \mathfrak{h}_{\text{nm}}(\Sigma)$.

Proof. Suppose that $p \in \mathcal{S}_{\text{nm}}(V)$, we claim that Σ is not homotopic minimizing to either side in $B_\varepsilon(p)$ for any $\varepsilon > 0$ sufficiently small. Indeed, by assumption, the unique tangent cone $\mathbf{C} = \partial\Omega_{\mathbf{C}}$ to Σ at p is not minimizing to either side. This implies that there are Caccioppoli sets $E_{\mathbf{C}}^- \subset \Omega_{\mathbf{C}} \subset E_{\mathbf{C}}^+$ so that $E_{\mathbf{C}}^\pm \Delta \Omega_{\mathbf{C}} \subset B_1 \subset \mathbb{R}^{n+1}$ and so that

$$\text{Per}_{\mathbb{R}^{n+1}}(E_{\mathbf{C}}^\pm | B_1) \leq \text{Per}_{\mathbb{R}^{n+1}}(\Omega_{\mathbf{C}} | B_1) - \delta.$$

Choose $C^{1,\omega}$ coordinates on M around p so that $\Omega = \Omega_{\mathbf{C}}$ in $B_\varepsilon(p)$ and so that $g_{ij}(p) = \delta_{ij}$, which we can do since $g \in C^2$ and Σ is a $C^{1,\omega}$ deformation of \mathbf{C} near p by assumption. Then, set

$$E(x) := \begin{cases} (\Omega \setminus B_\varepsilon) \cup (|x|E_{\mathbf{C}}^- \cap B_\varepsilon) & x < 0 \\ \Omega & x = 0 \\ (\Omega \setminus B_\varepsilon) \cup (|x|E_{\mathbf{C}}^+ \cap B_\varepsilon) & x > 0 \end{cases}$$

We have that

$$\text{Per}_g(E(x)) - \text{Per}_g(\Omega) = -|x|^n \delta(1 + o(1))$$

as $x \rightarrow 0$ (since the metric g_{ij} converges to the flat metric δ_{ij} after rescaling $|x| \rightarrow 1$, by the $C^{1,\omega}$ regularity of the chart). This shows that Σ is not homotopic minimizing to either side in $B_\varepsilon(p)$, so $p \in \mathfrak{h}_{\text{nm}}(\Sigma)$ as claimed. \square

3.2. Cancellation. We will assume the cancellation condition

$$\mathbf{M}(\Phi(x_0)) < W$$

throughout this subsection. In particular, we can find $q \in \text{Reg } V$ so that for all $\varepsilon > 0$ sufficiently small,

$$\text{Per}(\Omega | B_\varepsilon(q)) < |V|(B_\varepsilon(q))$$

where $\partial\Omega = \Phi(x_0)$. Like in the previous section we set $\Sigma := \text{supp } V$.

Furthermore we set $V = \sum_i \kappa_i |\Sigma_i|$, where each Σ_i is a minimal hypersurface with optimal regularity and $\kappa_i \in \mathbb{N}$ are constant multiplicities, by the constancy theorem [Sim83b, Theorem 41.1]. So (6) becomes

$$(10) \quad \begin{aligned} & \text{there is a (ONVP) sweepout } \{\Phi(x) = \partial\Omega(x)\} \text{ and } x_i \nearrow x_0 \in \mathbf{m}_L(\Phi), \text{ so that} \\ & |\Phi(x_i)| \rightarrow V = \sum_i \kappa_i |\Sigma_i| \in \mathcal{R}, \Phi \text{ is not left excessive at } x_0 \text{ and} \end{aligned}$$

$$\text{there is } q \in \Sigma \text{ such that } \text{Per}(\Omega | B_\varepsilon(q)) \leq |V|(B_\varepsilon(q)) - \delta(\varepsilon) \text{ for all } \varepsilon > 0.$$

We write $\Omega = \Omega(x_0)$ and observe that $\Sigma \subset \bar{\Omega}$. We would like to claim that Σ is homotopically minimizing, but this condition might not make sense if Σ is one-sided. However, thanks to the cancellation we can actually prove that Σ is area-minimizing in its neighborhood in Ω away from a small ball around q .

Definition 27. We will call a set Ω' a $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor if

$$(\Omega \setminus B_\tau(\Sigma)) \cup (B_\varepsilon(q) \setminus \Sigma) \subset \Omega' \subsetneq \Omega \setminus \Sigma$$

An $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor Ω' will be called a minimizing competitor if its perimeter is strictly less than perimeter of any $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor Ω'' with $\Omega' \subset \Omega''$. (Note that we do not require $\text{Per}(\Omega')$ to be less than the perimeter of all competitors, but only those that contain Ω').

Proposition 28. *Suppose (10) holds, then for every $\varepsilon > 0$ there is $\tau > 0$, such that minimizing $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor does not exist.*

Proof. For contradiction suppose there exists a minimizing $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor U . Observe that for every $\delta > 0$ we can find $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitors Ω' with $\text{Per}(\Omega') \leq \text{Per}(\Omega) + \delta$. It follows that

$$\text{Per}(U) \leq \text{Per}(\Omega) < W$$

by the cancellation assumption. If we choose $\tau > 0$ sufficiently small, then by Lemma 16 there exists a nested family $\{E(x) : x \in [0, 1]\}$ with $E(0) = U$, $E(1) = \Omega$ and

$$\text{Per}(E(x)) < W$$

Let $(x_l, x_0]$ be the connected interval such that $\Omega(x) \setminus U \neq \emptyset$, where $\{\Phi(x) = \partial\Omega(x)\}$, and define family $\Psi : (x_l, x_0 + 1] \rightarrow \mathcal{Z}_n(M, \mathbb{Z}_2)$ by

$$\Psi(x) := \begin{cases} \partial(\Omega(x) \cap U) & \text{if } x \in (x_l, x_0] \\ \partial E(x - x_0) & \text{if } x \in [x_0, x_0 + 1] \end{cases}$$

Clearly Ψ is continuous, since $\Omega = \Omega(x_0)$ and moreover we have that

$$\limsup_{y \rightarrow x} \mathbf{M}(\Psi(y)) < \limsup_{y \rightarrow x} \mathbf{M}(\Phi(x)) \leq W$$

for every $x \in (x_l, x_0)$ by strict minimality condition in Definition 27. For every $x \in [x_0, x_0 + 1]$ we also have $\mathbf{M}(\Psi(x)) = \mathbf{M}(\partial E(x)) < W$. This implies that x_0 is left excessive for Φ which is a contradiction. \square

Proposition 29. *Suppose $V = \sum_i \kappa_i |\Sigma_i|$ is as in (10), then each Σ_i has stable regular part and $\mathfrak{h}_{\text{nm}}(V) = \emptyset$. Moreover, for every point $x \in \text{support}(V)$ there exists $r > 0$, such that the support of V is minimizing to one side in $B_r(x)$*

Proof. First we observe that we can find two points q_1 and q_2 in $\text{Reg } V$, such that for all $\varepsilon > 0$ sufficiently small,

$$\text{Per}(\Omega | B_\varepsilon(q_j)) < |V|(B_\varepsilon(q))$$

By Proposition 28 we have non-existence of minimizing $(q_j, \varepsilon, \tau, \Sigma, \Omega)$ -competitors for $j = 1, 2$. This implies that Σ_i is area minimizing to one side in a small ball around every point of V . In particular, we have $\mathcal{H}^0(\mathfrak{h}_{\text{nm}}(V)) = 0$.

The stability of the regular part of each Σ_i also follows from the non-existence of minimizing $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitors. Indeed, if a component Σ_i has index ≥ 1 , then for $\varepsilon > 0$ sufficiently small, the minimal hypersurface $\Sigma_i \setminus B_\varepsilon(q)$ with fixed boundary will be unstable by a standard capacity argument. If Σ_i is two-sided, then by considering a

minimization problem to one side of Σ_i in $B_\tau(\Sigma_i) \setminus B_\varepsilon(q)$ we can find open set $U \subset \Omega$, such that $\Omega \setminus U$ is a minimizing $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor.

Suppose Σ_i is one-sided. Since $\Sigma_i \subset \bar{\Omega}$ we have that $B_\tau(\Sigma_i) \setminus \Sigma_i \subset \Omega$ for all sufficiently small $\tau > 0$. In particular, for small $\tau < \varepsilon$ we can minimize in the class of hypersurfaces $\{S \subset B_\tau(\Sigma_i) : S \cap B_\varepsilon(q) = \Sigma_i \cap B_\varepsilon(q)\}$ to obtain a minimizer Σ'_i in the same homology class and open set $U \subset \Omega$ with $\partial U = \Sigma_i \cup \Sigma'_i$. Then $\Omega \setminus U$ is a minimizing $(q, \varepsilon, \tau, \Sigma, \Omega)$ -competitor. \square

3.3. Multiplicity 2 bound. In this subsection we show that if $\kappa_i > 2$ for some i , then x_0 is excessive, by using simple comparisons with disks. Notice that if any multiplicity satisfies $\kappa_i \geq 2$ then we must be in the cancellation case considered above.

Lemma 30 (Multiplicity 2 bound). *Let $V = \sum_i \kappa_i |\Sigma_i|$ be as in (6). Then $\kappa_i \leq 2$ for every i .*

Proof. Suppose by contradiction $\kappa_i \geq 3$ for some i . Then let $p \in \text{Reg}(\Sigma_i)$, $p \neq q$ (where q is the cancellation point considered above). Consider a ball $B_r(p)$, $r < \frac{1}{2} \text{dist}(p, q)$, sufficiently small so that $\Sigma_i \cap B_r(p)$ is two-sided. Let $\tau(r) > 0$ be a small constant to be chosen later and set $U = B_r(p) \cap B_\tau(\Sigma_i)$.

Consider sequence $x_j \nearrow x_0$ with $|\partial\Omega(x_j)| \rightarrow V$. We can assume that the radius r was chosen sufficiently small, so that

$$(11) \quad \mathbf{M}(\partial\Omega(x_j) \cap U) \geq \left(\kappa_i - \frac{1}{10} \right) \omega_n r^n,$$

for all j large enough, where ω_n denotes the measure of the n -dimensional ball of radius one.

Let $\Omega'_j \subset \Omega(x_j)$, $\Omega'_j \setminus U = \Omega(x_j) \setminus U$, be a strict one-sided outer area minimizer in $\Omega(x_j) \cap U$. Observe that if Ω'_j does not converge to $\Omega(x_0)$, then $\lim \Omega'_j$ is a $(q, \frac{1}{2} \text{dist}(p, q), \tau, \Sigma, \Omega(x_0))$ -competitor, which contradicts Proposition 28.

We conclude that $\lim \Omega'_j = \Omega(x_0)$. On the other hand, by comparing $\Omega(x_j) \setminus U$ to Ω'_j and assuming that $\tau(r)$ was chosen sufficiently small, we have that one-sided area minimizing property of Ω'_j implies

$$\mathbf{M}(\partial\Omega'_j \cap U) \leq \text{Per}(U) \leq \left(2 + \frac{1}{10} \right) \omega_n r^n,$$

For $\tau(r)$ sufficiently small and j large we can apply Lemma 16 to find a nested family $E(x)$ interpolating between Ω'_j and Ω , such that

$$\begin{aligned} \text{Per}(E(x)) &\leq \max\{\mathbf{M}(\partial\Omega'_j \setminus U), \mathbf{M}(\partial\Omega(x_0) \setminus U)\} + \left(2 + \frac{2}{10} \right) \omega_n r^n \\ &\leq W - \left(1 - \frac{3}{10} \right) \omega_n r^n. \end{aligned}$$

By combining families $\Omega(x) \cap \Omega'_j$ and $E(x)$ we obtain that x_0 is left-excessive. \square

3.4. Proof of Theorem 4. The result follow immediately by combining Corollary 19 with Propositions 23, 24, 29 and Lemma 30. \square

4. PROOF OF THEOREMS 1, 2, AND 3

In this section we prove Theorem 3 (Theorems 1 and 2 follow immediately from Theorem 3 when combined with the facts that when $n = 8$ all singularities are regular and that the set of bumpy metrics is open and dense [Whi91, Whi17]). Theorem 3 will follow from Theorem 4 and Proposition 17, together with a simple surgery procedure.

4.1. Surgery procedure. We show here how to regularize minimal hypersurfaces with regular singularities under the assumption that the hypersurface minimizes area in a small ball around each singularity.

Proposition 31 (Perturbing away regular singularities of locally area minimizing surfaces). *For (M^{n+1}, g) a compact $C^{2,\alpha}$ -Riemannian metric and $\Sigma \in \mathcal{R}$ a minimal hypersurface, recall that $\mathcal{S}_0(\Sigma) \subset \text{Sing } \Sigma$ is defined to be the set of singular points with a regular tangent cone. There is $\tilde{g} \in \text{Met}^{2,\alpha}(M)$ arbitrarily close to g and $\tilde{\Sigma}$ arbitrarily close in the Hausdorff sense to Σ so that $\tilde{\Sigma}$ is minimal with respect to \tilde{g} and $\mathcal{S}_0(\tilde{\Sigma}) \subset \mathfrak{h}_{\text{nm}}(\tilde{\Sigma}) = \mathfrak{h}_{\text{nm}}(\Sigma)$.*

Proof. For every $p \in \mathcal{S}_0(\Sigma) \setminus \mathfrak{h}_{\text{nm}}(\Sigma)$, and $\varepsilon_0 = \varepsilon_0(p)$ so that $\Sigma \cap (B_{\varepsilon_0}(p) \setminus p)$ is regular, we will show how to perturb g and Σ so that p becomes regular. We will do this by making an arbitrarily small change to g , Σ supported in $B_{\varepsilon_0}(p)$. Because \mathcal{S}_0 is discrete (but not necessarily closed when $n \geq 9$) it is easy to enumerate the elements of $\mathcal{S}_0(\Sigma) \setminus \mathfrak{h}_{\text{nm}}(\Sigma)$ and make a summably small change around each point. As such, it suffices to consider just the perturbation near p .

By definition, taking $\varepsilon < \varepsilon_0$ sufficiently small, $\Sigma \cap B_\varepsilon(p)$ is one-sided homotopy area-minimizing. For concreteness write $\Sigma \cap B_\varepsilon(p) = \partial\Omega$ in $B_\varepsilon(p)$ and assume that Ω is inner homotopy minimizing. By Lemma 26, the tangent cone at p is area-minimizing (to the same side).

We claim that (after taking $\varepsilon > 0$ smaller if necessary) there is a sequence of $\Sigma_i \in \mathcal{R}(B_\varepsilon(p))$ with stable regular part, with $\Sigma_i \subset \Omega$, Σ_i disjoint from Σ , and $\Sigma_i \rightarrow \Sigma$. Indeed, we can apply Proposition 17 to conclude that either (after shrinking $\varepsilon > 0$), Ω is area-minimizing to the inside, or there are Σ_i as asserted.

In the case that Ω is area-minimizing to the inside, we can still construct the Σ_i by shrinking $\varepsilon > 0$ even further so that Ω is strictly area-minimizing to the inside and then minimizing area with respect to a boundary $\Sigma \cap \partial B_\varepsilon(p) + \delta_i$, for a sequence $\delta_i \rightarrow 0$; i.e., the boundary of $\Sigma \cap B_\varepsilon(p)$ pushed slightly into Ω . By the unique minimizing property, the minimizers will converge back to Σ in $B_\varepsilon(p)$.

For i sufficiently large we can write the intersection of Σ_i with the annulus $A(p, \varepsilon/5, \varepsilon)$ as a graph of function u_i over Σ .

Reasoning as in Hardt–Simon [HS85, Theorem 5.6] (cf. [Liu19, Theorem 3.1]), for i sufficiently large, Σ_i will be regular in $B_{\varepsilon/2}(p)$. We now set

$$\tilde{\Sigma}_i = (\Sigma_i \cap B_{\varepsilon/5}) \cup (\Sigma \setminus B_\varepsilon(p)) \cup ((\Sigma + \chi u_i) \cap A(p, \varepsilon/5, \varepsilon))$$

where χ is a smooth cutoff function with $\chi \equiv 1$ on $B_{\varepsilon/5}$ and $\chi \equiv 0$ on $B_{3\varepsilon/5}$. Note that

$$H_g(\tilde{\Sigma}_i) \quad \text{is supported in } B_{4\varepsilon/5}(p) \setminus B_{\varepsilon/5}(p)$$

and $\|H_g(\tilde{\Sigma}_i)\|_{C^{2,\alpha}} = o(1)$ as $i \rightarrow \infty$.

Now, define $\tilde{g} = e^f g$, in this new metric, since $\tilde{\Sigma}$ is smooth, we have the transformation

$$H_{\tilde{g}}(\tilde{\Sigma}) = e^{-f} \left(H_g(\tilde{\Sigma}) + \frac{\partial f}{\partial \nu} \right),$$

where ν is the normal direction to Σ . Setting $H_{\tilde{g}}(\tilde{\Sigma})$, this reduces to the equation

$$H_g(\tilde{\Sigma}) + \frac{\partial f}{\partial \nu} = 0$$

which implies that $f = -H_g(\tilde{\Sigma})\zeta(\nu)$, for a function $\zeta(t)$ such that $\zeta'(0) = 1$ and $\zeta \equiv 0$ for $|t| \geq \varepsilon/100$ is a solution. Since, as observed, $H_g(\tilde{\Sigma})$ is supported in $A(p, \varepsilon/5, 4\varepsilon/5)$, so is the metric change, and since $\|u\|_{C^{4,\alpha}} \leq o(1)$ and χ is smooth, we have

$$\|g - \tilde{g}\|_{2,\alpha} = \|e^f - 1\|_{C^{2,\alpha}} \|g\|_{2,\alpha} \leq C \|u\|_{C^{4,\alpha}} \|g\|_{2,\alpha} = o(1)$$

as $i \rightarrow \infty$. This completes the proof. \square

4.2. Proof of Theorem 3. For $g \in \text{Met}^{2,\alpha}(M)$, apply Theorem 4 to find $V \in \mathcal{R}$ with

$$\mathcal{H}^0(\mathfrak{h}_{\text{nm}}(V)) + \text{Index}(V) \leq 1.$$

We can apply Proposition 31 to $\Sigma = \text{supp } V$ to find a metric \tilde{g} that is arbitrarily $C^{2,\alpha}$ -close to g and a \tilde{g} minimal hypersurface $\tilde{\Sigma} \in \mathcal{R}$ so that $\mathcal{S}_0(\Sigma) \subset \mathfrak{h}_{\text{nm}}(\Sigma)$. (Note that if $\text{Index}(V) = 1$, then $\mathfrak{h}_{\text{nm}}(\Sigma) = \emptyset$, so $\mathcal{S}_0(\Sigma) = \emptyset$.) This completes the first part of the proof.

We now consider $g \in \text{Met}_{\text{Ric} > 0}^{2,\alpha}(M)$.⁴ If Σ is two-sided, then $\text{Index}(\Sigma) \geq 1$, so we can argue as above. On the other hand if Σ is one-sided, then $[\Sigma] \neq 0 \in H_n(M, \mathbb{Z}_2)$. We can then find $\hat{\Sigma} \in [\Sigma]$ by minimizing area in the homology class. The surface $\hat{\Sigma}$ may have singularities, but they are all locally area minimizing. Thus, we can apply Proposition 31 to Σ yielding $\tilde{\Sigma}$ and \tilde{g} with $\mathcal{S}_0(\tilde{\Sigma}) = \emptyset$. \square

REFERENCES

- [BDGG69] Enrico Bombieri, Ennio De Giorgi, and Enrico Giusti. Minimal cones and the Bernstein problem. *Inventiones mathematicae*, 7:243–268, 1969.
- [Bel20] Costante Bellettini. Multiplicity-1 minmax minimal hypersurfaces in manifolds with positive Ricci curvature. <https://arxiv.org/abs/2004.10112>, 2020.
- [CC92] Eugenio Calabi and Jian Guo Cao. Simple closed geodesics on convex surfaces. *J. Differential Geom.*, 36(3):517–549, 1992.
- [Cha84] Isaac Chavel. *Eigenvalues in Riemannian geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1984. Including a chapter by Burton Randol, With an appendix by Jozef Dodziuk.
- [CL20] Gregory R. Chambers and Yevgeny Liokumovich. Existence of minimal hypersurfaces in complete manifolds of finite volume. *Invent. Math.*, 219(1):179–217, 2020.
- [CM20] Otis Chodosh and Christos Mantoulidis. Minimal surfaces and the allen–cahn equation on 3-manifolds: index, multiplicity, and curvature estimates. *Annals of Mathematics*, 191(1):213–328, 2020.

⁴The idea is that positive Ricci curvature rules out stable hypersurfaces but this requires the hypersurface to be two-sided. As such, we must consider two cases, depending on whether Σ is one or two-sided.

- [CR18] Gregory R. Chambers and Regina Rotman. Monotone homotopies and contracting discs on riemannian surfaces. *Journal of Topology and Analysis*, 10(02):323–354, 2018.
- [Dey20] Akashdeep Dey. A comparison of the Almgren-Pitts and the Allen-Cahn min-max theory. <https://arxiv.org/abs/2004.05120>, 2020.
- [DLT13a] Camillo De Lellis and Dominik Tasnady. The existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 95(3):355–388, 11 2013.
- [DLT13b] Camillo De Lellis and Dominik Tasnady. The existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 95(3):355–388, 2013.
- [Fal80] K. J. Falconer. Continuity properties of k -plane integrals and Besicovitch sets. *Math. Proc. Cambridge Philos. Soc.*, 87(2):221–226, 1980.
- [Gas20] Pedro Gaspar. The second inner variation of energy and the Morse index of limit interfaces. *J. Geom. Anal.*, 30(1):69–85, 2020.
- [GG19] Pedro Gaspar and Marco A. M. Guaraco. The Weyl law for the phase transition spectrum and density of limit interfaces. *Geom. Funct. Anal.*, 29(2):382–410, 2019.
- [Gua18] Marco A. M. Guaraco. Min-max for phase transitions and the existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 108(1):91–133, 2018.
- [Gut07] Larry Guth. The width-volume inequality. *Geom. Funct. Anal.*, 17(4):1139–1179, 2007.
- [Hie18] Fritz Hiesmayr. Spectrum and index of two-sided Allen-Cahn minimal hypersurfaces. *Comm. Partial Differential Equations*, 43(11):1541–1565, 2018.
- [HS85] Robert Hardt and Leon Simon. Area minimizing hypersurfaces with isolated singularities. *J. Reine Angew. Math.*, 362:102–129, 1985.
- [Ilm96] T. Ilmanen. A strong maximum principle for singular minimal hypersurfaces. *Calc. Var. Partial Differential Equations*, 4(5):443–467, 1996.
- [Law91] Gary R. Lawlor. A sufficient criterion for a cone to be area-minimizing. *Mem. Amer. Math. Soc.*, 91(446):vi+111, 1991.
- [Li19] Yangyang Li. Existence of infinitely many minimal hypersurfaces in higher-dimensional closed manifolds with generic metrics. <https://arxiv.org/abs/1901.08440>, 2019.
- [Liu19] Zhenhua Liu. Stationary one-sided area-minimizing hypersurfaces with isolated singularities. <https://arxiv.org/abs/1904.02289>, 2019.
- [LMN18] Yevgeny Liokumovich, Fernando C. Marques, and Andre Neves. Weyl law for the volume spectrum. *Annals of Mathematics*, 187(3):933–961, 2018.
- [Man17] Christos Mantoulidis. Allen-Cahn min-max on surfaces. <https://arxiv.org/abs/1706.05946>, to appear in *Jour. Diff. Geom.*, 2017.
- [MN16] Fernando C. Marques and André Neves. Morse index and multiplicity of min-max minimal hypersurfaces. *Camb. J. Math.*, 4(4):463–511, 2016.
- [MN19] Fernando Codá Marques and Andr’e Neves. Morse index of multiplicity one min-max minimal hypersurfaces. *arXiv: Differential Geometry*, 2019.
- [Moo06] John Douglas Moore. Bumpy metrics and closed parametrized minimal surfaces in Riemannian manifolds. *Trans. Amer. Math. Soc.*, 358(12):5193–5256, 2006.
- [Moo07] John Douglas Moore. Correction for: “Bumpy metrics and closed parametrized minimal surfaces in Riemannian manifolds” [Trans. Amer. Math. Soc. **358** (2006), no. 12, 5193–5256 (electronic); mr2238914]. *Trans. Amer. Math. Soc.*, 359(10):5117–5123, 2007.
- [Pit81] Jon T. Pitts. *Existence and regularity of minimal surfaces on Riemannian manifolds*, volume 27 of *Mathematical Notes*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1981.
- [RL19] Alejandra Ramírez-Luna. Orientability of min-max hypersurfaces in manifolds of positive Ricci curvature. <https://arxiv.org/abs/1907.12519>, 2019.
- [Sim68] James Simons. Minimal varieties in riemannian manifolds. *Ann. of Math. (2)*, 88:62–105, 1968.
- [Sim83a] Leon Simon. Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems. *Ann. of Math. (2)*, 118(3):525–571, 1983.

- [Sim83b] Leon Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis, Australian National University*. Australian National University Centre for Mathematical Analysis, Canberra, 1983.
- [Sma93] Nathan Smale. Generic regularity of homologically area minimizing hypersurfaces in eight-dimensional manifolds. *Comm. Anal. Geom.*, 1(2):217–228, 1993.
- [Son18] Antoine Song. Existence of infinitely many minimal hypersurfaces in closed manifolds. <https://arxiv.org/abs/1806.08816>, 2018.
- [SS81] Richard Schoen and Leon Simon. Regularity of stable minimal hypersurfaces. *Comm. Pure Appl. Math.*, 34(6):741–797, 1981.
- [Tam84] Italo Tamanini. Regularity results for almost minimal oriented hypersurfaces in \mathbb{R}^n . *Quaderni del Dipartimento di Matematica dell' Università di Lecce*, 1:1–92, 1984.
- [Whi85] Brian White. Generic regularity of unoriented two-dimensional area minimizing surfaces. *Ann. of Math. (2)*, 121(3):595–603, 1985.
- [Whi91] Brian White. The space of minimal submanifolds for varying Riemannian metrics. *Indiana Univ. Math. J.*, 40(1):161–200, 1991.
- [Whi94] Brian White. A strong minimax property of nondegenerate minimal submanifolds. *J. Reine Angew. Math.*, 457:203–218, 1994.
- [Whi17] Brian White. On the bumpy metrics theorem for minimal submanifolds. *Amer. J. Math.*, 139(4):1149–1155, 2017.
- [Whi19] Brian White. Generic transversality of minimal submanifolds and generic regularity of two-dimensional area-minimizing integral currents. <https://arxiv.org/abs/1901.05148>, 2019.
- [Zho15] Xin Zhou. Min-max minimal hypersurface in (M^{n+1}, g) with $Ric > 0$ and $2 \leq n \leq 6$. *J. Differential Geom.*, 100(1):129–160, 2015.
- [Zho17] Xin Zhou. Min-max hypersurface in manifold of positive Ricci curvature. *J. Differential Geom.*, 105(2):291–343, 02 2017.
- [Zho19] Xin Zhou. On the multiplicity one conjecture in min-max theory. <https://arxiv.org/abs/1901.01173>, 2019.

OC: DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, BUILDING 380, STANFORD, CALIFORNIA 94305, USA.

Email address: ochodosh@stanford.edu

YL: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST GEORGE STREET, TORONTO, ON M5S 2E4, CANADA.

Email address: ylio@math.toronto.edu

LS: DEPARTMENT OF MATHEMATICS, UC SAN DIEGO, AP&M, LA JOLLA, CALIFORNIA, 92093, USA.

Email address: lspolaor@ucsd.edu