

# An approximation for the Mumford-Shah functional

Luca Lussardi

Dipartimento di Matematica  
Facoltà di Ingegneria, Università degli Studi di Brescia  
via Valotti 9, 25133 Brescia, Italy  
email: luca.lussardi@ing.unibs.it

## Abstract

We approximate, in the sense of  $\Gamma$ -convergence, the Mumford-Shah functional by means of a sequence of non-local integral functionals depending on the average of the absolute value of the gradient.

**Mathematics Subject Classification:** 49Q20

**Keywords:**  $\Gamma$ -convergence, Mumford-Shah functional

## 1 Introduction

In the variational approach to many problems in computer vision (image segmentation, signal processing and so on) an important rôle has been played by the Mumford-Shah functional, which is the most famous example of a free discontinuity functional (terminology introduced by DeGiorgi in [11]). The Mumford-Shah functional is given by

$$MS(u) = \int_{\Omega} |\nabla u|^2 dx + c\mathcal{H}^{n-1}(S_u)$$

where  $u \in SBV(\Omega)$ , the space of special functions of bounded variation;  $S_u$  is the approximate discontinuity set of  $u$  and  $\mathcal{H}^{n-1}$  is the  $(n-1)$ -dimensional Hausdorff measure. Several approximation methods are known for the Mumford-Shah functional and, more in general, for free discontinuity functionals: the Ambrosio & Tortorelli approximation (see [1] and [3]) via elliptic functionals, the Gobbino's approximation by finite difference methods (see [12]) and many others (see [6], [7], [9], [10]).

In [5] Braides & Dal Maso approximate the Mumford-Shah functional by means of a sequence of non-local integral functionals given by

$$F_{\varepsilon}(u) = \frac{1}{\varepsilon} \int_{\Omega} f \left( \varepsilon \int_{B_{\varepsilon}(x) \cap \Omega} |\nabla u|^2 dy \right) dx \quad (1)$$

with  $u \in H^1(\Omega)$  and, for instance,  $f(t) = t \wedge 1/2$ . A variant of this method is investigated in [14], [15] and [13] where the problem of the convergence of

$$F_\varepsilon(u) = \frac{1}{\varepsilon} \int_{\Omega} f_\varepsilon \left( \varepsilon \int_{B_\varepsilon(x) \cap \Omega} |\nabla u| dy \right) dx \quad (2)$$

is considered; here  $f_\varepsilon$  is a convex-concave function with  $f_\varepsilon(\varepsilon t)/\varepsilon \rightarrow \phi(t)$ , as  $(\varepsilon, t) \rightarrow (0, 0)$ , where  $\phi$  has linear growth at infinity and plays the rôle of the bulk energy density in the limit of  $F_\varepsilon$ . Then, under the assumption on  $f_\varepsilon$  in [13], the Mumford-Shah functional cannot be recovered by the  $\Gamma$ -convergence of  $F_\varepsilon$ , since the bulk term in  $MS$  is given by

$$\int_{\Omega} |\nabla u|^2 dx$$

and it has superlinear growth at infinity. A question arise: is it possible to recover the Mumford-Shah functional from (2) instead of (1)? The aim of this paper is to prove an approximation results for the Mumford-Shah functional, obtained adapting the results contained in [14], [15] and [13], by means of a sequence of functionals of type (2). The core of the proof is Theorem 4.1 where the lower bound for the  $\Gamma$ -limit is optimized by a sup of measures argument, while the upper bound descends from standard density results and general properties of Minkowsky content.

## 2 Preliminary Notes

*Functions of bounded variation.* For a thorough treatment of  $BV$  functions we refer to [2]. Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ; the space  $BV(\Omega)$  of real *functions of bounded variation* is the space of the functions  $u \in L^1(\Omega)$  whose distributional derivative is representable by a measure  $\mathbb{R}^n$ -valued measure  $Du$  on  $\Omega$ . We denote by  $S_u$  the *approximate discontinuity set* of  $u$  and by  $J_u$  the set of approximate jump points of  $u$ .

For a function  $u \in BV(\Omega)$  let  $Du = D^a u + D^s u$  be the (Lebesgue) decomposition of  $Du$  into absolutely continuous and singular part. We denote by  $\nabla u$  the density of  $D^a u$ ; the measures  $D^j u := D^s u \llcorner J_u$ ,  $D^c u := D^s u \llcorner (\Omega \setminus S_u)$  are called the *jump part* and the *Cantor part* of the derivative, respectively.

We say that a function  $u \in BV(\Omega)$  is a *special function of bounded variation* ( $u \in SBV(\Omega)$ ) if  $|D^c u|(\Omega) = 0$ ; moreover we say that a function  $u \in L^1(\Omega)$  is a *generalized special function of bounded variation* ( $u \in GSBV(\Omega)$ ) if  $u^T := (-T) \vee u \wedge T$  belongs to  $SBV(\Omega)$  for every  $T \geq 0$ . If  $u \in GSBV(\Omega)$ , the function  $\nabla u$  given by  $\nabla u = \nabla u^T$  for  $\mathcal{L}^n$ -a.e. on  $\{|u| \leq T\}$  turns out to be well-defined. Moreover, the set function  $T \mapsto S_{u^T}$  is monotone increasing; therefore, we set  $S_u = \bigcup_{T>0} S_{u^T}$ .

*Supremum of measures.* We recall the following useful result from measure theory, which can be found in [4].

**Lemma 2.1 (supremum of measures)** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$  and denote by  $\mathcal{A}(\Omega)$  the family of its open subsets. Let  $\lambda$  be a positive Borel measure on  $\Omega$ , and  $\mu: \mathcal{A}(\Omega) \rightarrow [0, +\infty)$  a set function which is superadditive on open sets with disjoint compact closures (i.e. if  $A, B \subset\subset \Omega$  and  $\overline{A} \cap \overline{B} = \emptyset$ , then  $\mu(A \cup B) \geq \mu(A) + \mu(B)$ ). Let  $(\psi_i)_{i \in I}$  be a family of positive Borel functions. Suppose that*

$$\mu(A) \geq \int_A \psi_i d\lambda \quad \text{for every } A \in \mathcal{A}(\Omega) \text{ and } i \in I;$$

then

$$\mu(A) \geq \int_A \sup_i \psi_i d\lambda \quad \text{for every } A \in \mathcal{A}(\Omega).$$

### 3 Main Results

Let  $\Omega \subseteq \mathbb{R}^n$  be a bounded open set with Lipschitz boundary, and consider the family  $(F_\varepsilon)_{\varepsilon > 0}$  of non-local functionals  $L^1(\Omega) \rightarrow [0, +\infty]$  given by

$$F_\varepsilon(u) = \begin{cases} \frac{1}{\varepsilon} \int_\Omega f_\varepsilon \left( \varepsilon \int_{B_\varepsilon(x) \cap \Omega} |\nabla u| dy \right) dx & \text{if } u \in W^{1,1}(\Omega) \\ +\infty & \text{otherwise,} \end{cases} \quad (3)$$

where  $f_\varepsilon: [0, +\infty) \rightarrow [0, +\infty)$  is requested to satisfy the following conditions:

**(A1)** for every  $\varepsilon > 0$ ,  $f_\varepsilon$  is a non-decreasing continuous function with  $f_\varepsilon(0) = 0$ ; moreover, there exists  $a_\varepsilon > 0$  such that  $a_\varepsilon \rightarrow 0$  as  $\varepsilon \rightarrow 0$  and  $f_\varepsilon$  is concave in  $(a_\varepsilon, +\infty)$ .

**(A2)** 
$$\lim_{(\varepsilon, t) \rightarrow (0, 0)} \frac{\varepsilon f_\varepsilon(t)}{t^2} = 1.$$

**(A3)**  $f_\varepsilon \nearrow f$  uniformly on the compact subsets of  $(0, +\infty)$ , where  $f(t) = f_\infty > 0$  is a constant function.

A possible choice for  $f_\varepsilon$  is  $f_\varepsilon(t) = (t^2/\varepsilon) \wedge f_\infty$ .

**Remark 3.1** *Let  $\delta \in (0, 1)$ ; by **(A2)** there exists  $t_\delta > 0$  and  $\varepsilon_\delta > 0$  such that  $f_\varepsilon(t) \leq (1 + \delta)t^2/\varepsilon$  for any  $0 \leq t \leq t_\delta$  and  $0 \leq \varepsilon \leq \varepsilon_\delta$ . Since  $\phi(t) = t^2$  is convex and  $f_\varepsilon$  is concave in  $(a_\varepsilon, +\infty)$ , with  $a_\varepsilon \rightarrow 0$ , we get  $f_\varepsilon(t) \leq (1 + \delta)t^2/\varepsilon$  for any  $t \geq 0$  and  $\varepsilon$  sufficiently small. Then  $f_\varepsilon(\varepsilon t)/\varepsilon \leq (1 + \delta)t^2$  for any  $t \geq 0$ .*

The main result is the following convergence result.

**Theorem 3.2** *Let  $(F_\varepsilon)_{\varepsilon>0}$  be as in (3), with  $f_\varepsilon$  satisfying conditions (A1)-(A2)-(A3). Then  $(F_\varepsilon)$   $\Gamma$ -converges, w.r.t. the strong  $L^1$ -topology, as  $\varepsilon \rightarrow 0$ , to  $\mathcal{F}: L^1(\Omega) \rightarrow [0, +\infty]$  given by*

$$\mathcal{F}(u) = \begin{cases} \int_{\Omega} |\nabla u|^2 dx + 2f_\infty \mathcal{H}^{n-1}(S_u) & \text{if } u \in GSBV(\Omega) \\ +\infty & \text{otherwise.} \end{cases}$$

Moreover we have a compactness property:

**Theorem 3.3 (compactness)** *Let  $(\varepsilon_j)$  be a positive infinitesimal sequence and let  $(u_j)$  be a sequence in  $L^1(\Omega)$  such that  $\|u_j\|_\infty \leq M$ , and  $F_{\varepsilon_j}(u_j) \leq M$  for a suitable constant  $M$  independent of  $j$ ; then there exists a subsequence  $(u_{j_k})$  converging in  $L^1(\Omega)$  to a function  $u \in SBV(\Omega)$ .*

For the sequel we will need a ‘‘localization’’ of  $F_\varepsilon$ : for every open subset  $A$  of  $\Omega$ , we set

$$F_\varepsilon(u, A) = \begin{cases} \frac{1}{\varepsilon} \int_A f_\varepsilon \left( \varepsilon \int_{B_\varepsilon(x) \cap \Omega} |\nabla u| dy \right) dx & \text{if } u \in W^{1,1}(\Omega) \\ +\infty & \text{otherwise.} \end{cases}$$

Clearly,  $F_\varepsilon(\cdot, \Omega)$  coincides with the functional  $F_\varepsilon$  defined in (3). The lower and upper  $\Gamma$ -limits of  $(F_\varepsilon(\cdot, A))$  will be denoted by  $F'(\cdot, A)$  and  $F''(\cdot, A)$ , respectively.

## 4 Lower bound and compactness

**Theorem 4.1** *For any  $u \in GSBV(\Omega)$  and for any open subset  $A$  of  $\Omega$*

$$F'(u, A) \geq \int_A |\nabla u|^2 dx + 2f_\infty \mathcal{H}^{n-1}(S_u \cap A).$$

*Proof.* *Step 1.* First we show that

$$F'(u, A) \geq \int_{\Omega} |\nabla u|^2 dx + 2f_\infty \mathcal{H}^{n-1}(S_u \cap A), \quad \forall u \in SBV(\Omega).$$

Fix  $\delta \in (0, 1)$ ,  $T > 0$  and  $\eta > 0$  small; consider the family  $(g_\varepsilon)_{\varepsilon>0}$  given by

$$g_\varepsilon(t) = (1 - \delta)\varepsilon\phi^T\left(\frac{t}{\varepsilon}\right)$$

if  $0 \leq t < \sqrt{\varepsilon}$  and

$$g_\varepsilon(t) = \left\{ (1 - \delta) \left[ \varepsilon \phi^T \left( \frac{\sqrt{\varepsilon}}{\varepsilon} \right) + (\phi^T)' \left( \frac{\sqrt{\varepsilon}}{\varepsilon} \right) (t - \sqrt{\varepsilon}) \right] \right\} \wedge (f_\infty - \eta)$$

if  $t \geq \sqrt{\varepsilon}$ , with  $\phi^T(t) = t^2$  if  $0 \leq t < T$  and  $\phi^T(t) = 2Tt - T^2$  if  $t \geq T$ . The function  $g_\varepsilon$  depends on  $\varepsilon, \delta, T$  and  $\eta$ , but, for simplicity, we drop the dependence by  $\delta, T$  and  $\eta$ . By **(A2)** there exists  $t_\delta > 0$  such that, for  $\varepsilon$  sufficiently small,  $f_\varepsilon(t) \geq (1 - \delta)\varepsilon\phi^T(t/\varepsilon)$  whenever  $0 \leq t \leq t_\delta$ ; from convexity of  $\phi^T$  and from uniform convergence of  $f_\varepsilon$  on compact subsets of  $(0, +\infty)$  we get  $f_\varepsilon \geq g_\varepsilon$ , for  $\varepsilon$  sufficiently small. Thus:

- (1) for every  $\varepsilon > 0$ ,  $g_\varepsilon$  is a non-decreasing continuous function with  $g_\varepsilon(0) = 0$ ; moreover, there exists  $a_\varepsilon > 0$  ( $a_\varepsilon = \sqrt{\varepsilon}$ ) such that  $a_\varepsilon \rightarrow 0$  as  $\varepsilon \rightarrow 0$  and  $g_\varepsilon$  is concave in  $(a_\varepsilon, +\infty)$ .

(2) 
$$\lim_{(\varepsilon, t) \rightarrow (0, 0)} \frac{g_\varepsilon(t)}{(1 - \delta)\varepsilon\phi^T(t/\varepsilon)} = 1.$$

Moreover it turns out that, denoting by  $g(t) = 2(1 - \delta)Tt \wedge (f_\infty - \eta)$ ,

- (3)  $g_\varepsilon \rightarrow g$  uniformly on the compact subsets of  $[0, +\infty)$ .

- (4) There exists  $L > 0$  such that

$$|g_\varepsilon(s) - g_\varepsilon(t)| \leq L|s - t|, \quad \forall s, t > 0.$$

Then, since

$$F_\varepsilon(u, A) \geq \frac{1}{\varepsilon} \int_A g_\varepsilon \left( \varepsilon \int_{B_\varepsilon(x) \cap \Omega} |\nabla u| dy \right) dx, \quad u \in W^{1,1}(\Omega), \quad (4)$$

we get, by Theorem 3.1 in [13],

$$F'(u, A) \geq (1 - \delta) \int_A \phi^T(|\nabla u|) dx + 2 \int_{S_u \cap A} \int_0^1 \vartheta(x, t) dt d\mathcal{H}^{n-1}(x) + 2(1 - \delta)T|D^c u|(\Omega)$$

for all  $u \in BV(\Omega)$ , where

$$\vartheta(x, t) = g \left( \frac{\omega_{n-1}}{\omega_n} |u^+(x) - u^-(x)| (\sqrt{1 - t^2})^{n-1} \right).$$

By arbitrariness of  $\delta \in (0, 1)$  we have

$$F'(u, A) \geq \int_A \phi^T(|\nabla u|) dx + 2 \int_{(S_u \cap A) \times [0, 1]} \vartheta(x, t) dt d\mathcal{H}^{n-1}(x) + 2T|D^c u|(\Omega). \quad (5)$$

As  $\sup_T \phi^T(t) = t^2$  and  $\sup_{T,\eta} [Tt(f_\infty - \eta)] = f_\infty$ , for  $t > 0$ , by Lemma 2.1 we obtain

$$\begin{aligned} F'(u, A) &\geq \int_A |\nabla u|^2 dx + 2 \int_{(S_u \cap A) \times [0,1]} f_\infty dt d\mathcal{H}^{n-1}(x) \\ &= \int_A |\nabla u|^2 dx + 2f_\infty \mathcal{H}^{n-1}(S_u \cap A). \end{aligned}$$

*Step 2.* Let  $u \in GSBV(\Omega)$ , and  $T > 0$ . By definition,  $u^T \in SBV(\Omega)$  and  $|\nabla u| \geq |\nabla u^T|$ . Thus for every sequence  $u_j \rightarrow u$  in  $L^1(\Omega)$  we get  $F'(u, A) \geq \liminf_{j \rightarrow +\infty} F_{\varepsilon_j}(u_j^T)$ . By *Step 1*, as  $u_j^T \rightarrow u^T$  in  $L^1(\Omega)$ , we obtain

$$F'(u, A) \geq \int_A |\nabla u^T|^2 dx + 2f_\infty \mathcal{H}^{n-1}(S_{u^T} \cap A).$$

By taking the limit as  $T \rightarrow +\infty$  and recalling the definition of  $\nabla u$  and  $S_u$  we conclude. ■

*Proof of Theorem 3.3.* Let  $(\varepsilon_j)$  be a positive infinitesimal sequence and let  $(u_j)$  be a sequence in  $L^1(\Omega)$  such that  $\|u_j\|_\infty \leq M$ , and  $F_{\varepsilon_j}(u_j) \leq M$  for a suitable constant  $M$  independent of  $j$ . Then by (4) and by compactness Theorem 3.2 in [13], there exists a subsequence  $(u_{j_k})$  converging to  $u \in BV(\Omega)$ . Suppose  $|D^c u|(\Omega) \neq 0$ ; then, by taking the limit as  $T \rightarrow +\infty$  in (5),  $F'(u)$  would be  $+\infty$ , which contradicts  $F_{\varepsilon_j}(u_j) \leq M$ . Thus  $|D^c u|(\Omega) = 0$  and then  $u \in SBV(\Omega)$ . ■

## 5 Upper bound

In this last section we conclude the proof of Theorem 3.2. As usual, first we will take into account a suitable dense subset of  $SBV(\Omega)$ : let  $\mathcal{W}(\Omega)$  be the space of all functions  $w \in SBV(\Omega)$  satisfying the following properties:

- i)  $\mathcal{H}^{n-1}(\overline{S}_w \setminus S_w) = 0$ ;
- ii)  $\overline{S}_w$  is the intersection of  $\Omega$  with the union of a finite member of  $(n-1)$ -dimensional simplexes;
- iii)  $w \in W^{k,\infty}(\Omega \setminus \overline{S}_w)$  for every  $k \in \mathbb{N}$

where  $SBV^2(\Omega) = \{u \in SBV(\Omega) : |\nabla u| \in L^2(\Omega), \mathcal{H}^{n-1}(S_u) < +\infty\}$ . In [8] the density property of  $\mathcal{W}(\Omega)$  in  $SBV(\Omega)$  is proved. More precisely:

**Theorem 5.1** *Assume that  $\partial\Omega$  is Lipschitz. Let  $u \in SBV^2(\Omega) \cap L^\infty(\Omega)$ . Then there exists a sequence  $(w_j)$  in  $\mathcal{W}(\Omega)$  such that  $w_j \rightarrow u$  strongly in  $L^1(\Omega)$ ,  $\nabla w_j \rightarrow \nabla u$  strongly in  $L^2(\Omega, \mathbb{R}^n)$ ,  $\limsup_h \|w_j\|_\infty \leq \|u\|_\infty$  and*

$$\limsup_{j \rightarrow +\infty} \int_{S_{w_j}} \phi(w_j^+, w_j^-, \nu_{w_j}) d\mathcal{H}^{n-1} \leq \int_{S_u} \phi(u^+, u^-, \nu_u) d\mathcal{H}^{n-1}$$

for every upper semicontinuous function  $\phi$  such that  $\phi(a, b, \nu) = \phi(b, a, -\nu)$  whenever  $a, b \in \mathbb{R}$  and  $\nu \in S^{n-1}$ .

**Theorem 5.2** *Let  $u \in GSBV(\Omega)$ ; then*

$$F''(u) \leq \int_{\Omega} |\nabla u|^2 dx + 2f_{\infty} \mathcal{H}^{n-1}(S_u).$$

*Proof.* Since the upper  $\Gamma$ -limit of  $F_{\varepsilon}$  coincides with the upper  $\Gamma$ -limit of the relaxed functional  $\overline{F}_{\varepsilon}$ , we get  $F''(u) \leq \limsup_{\varepsilon \rightarrow 0} \overline{F}_{\varepsilon}(u)$ . It can be easily seen (see [15], Proposition 3.6) that, for  $\varepsilon > 0$  fixed, we have

$$\overline{F}_{\varepsilon}(u) = \frac{1}{\varepsilon} \int_{\Omega} f_{\varepsilon} \left( \frac{\varepsilon}{|B_{\varepsilon}(x) \cap \Omega|} |Du|(B_{\varepsilon}(x) \cap \Omega) \right) dx. \quad (6)$$

*Step 1.* First we consider the case  $u \in \mathcal{W}(\Omega)$ . Let  $S_{\varepsilon} = \{x \in \Omega : d(x, S_u) < \varepsilon\}$ ; then we can split  $\overline{F}_{\varepsilon}$  as follows:

$$\begin{aligned} \overline{F}_{\varepsilon}(u) &= \frac{1}{\varepsilon} \int_{\Omega \setminus S_{\varepsilon}} f_{\varepsilon} \left( \frac{\varepsilon}{|B_{\varepsilon}(x) \cap \Omega|} |Du|(B_{\varepsilon}(x) \cap \Omega) \right) dx \\ &\quad + \frac{1}{\varepsilon} \int_{S_{\varepsilon}} f_{\varepsilon} \left( \frac{\varepsilon}{|B_{\varepsilon}(x) \cap \Omega|} |Du|(B_{\varepsilon}(x) \cap \Omega) \right) dx. \end{aligned}$$

Since  $u \in W^{1,1}(\Omega \setminus S_{\varepsilon})$ , the first integral becomes

$$\frac{1}{\varepsilon} \int_{\Omega \setminus S_{\varepsilon}} f_{\varepsilon} \left( \varepsilon \int_{B_{\varepsilon}(x) \cap \Omega} |\nabla u| dy \right) dx \leq \frac{1}{\varepsilon} \int_{\Omega} f_{\varepsilon} \left( \varepsilon \int_{B_{\varepsilon}(x) \cap \Omega} |\nabla u| dy \right) dx.$$

Moreover since

$$\int_{B_{\varepsilon}(x) \cap \Omega} |\nabla u| dy \rightarrow |\nabla u(x)|$$

a.e.  $x \in \Omega$ , by **(A2)** and from the dominated convergence Theorem (see Remark 3.1) we get

$$\frac{1}{\varepsilon} \int_{\Omega} f_{\varepsilon} \left( \varepsilon \int_{B_{\varepsilon}(x) \cap \Omega} |\nabla u| dy \right) dx \rightarrow \int_{\Omega} |\nabla u|^2 dx.$$

We estimate now the second integral

$$\frac{1}{\varepsilon} \int_{S_{\varepsilon}} f_{\varepsilon} \left( \frac{\varepsilon}{|B_{\varepsilon}(x) \cap \Omega|} |Du|(B_{\varepsilon}(x) \cap \Omega) \right) dx.$$

Let  $\eta > 0$  small; by uniform convergence of  $f_{\varepsilon}$  on compact subsets of  $(0, +\infty)$  and by monotonicity property of  $f_{\varepsilon}$ , for  $\varepsilon$  sufficiently small it holds  $f_{\varepsilon}(t) \leq f_{\infty} + \eta$ , for any  $t \geq 0$ . Thus

$$\frac{1}{\varepsilon} \int_{S_{\varepsilon}} f_{\varepsilon} \left( \frac{\varepsilon}{|B_{\varepsilon}(x) \cap \Omega|} |Du|(B_{\varepsilon}(x) \cap \Omega) \right) dx \leq \frac{|S_{\varepsilon}|}{\varepsilon} (f_{\infty} + \eta).$$

Since  $S_u$  is the union of  $(n - 1)$ -dimensional simplexes, by standard results on Minkowsky content we have  $|S_\varepsilon|/2\varepsilon \rightarrow \mathcal{H}^{n-1}(S_u)$ , and then

$$\frac{|S_\varepsilon|}{\varepsilon}(f_\infty + \eta) \rightarrow 2(f_\infty + \eta)\mathcal{H}^{n-1}(S_u).$$

We conclude by arbitrariness of  $\eta$ .

*Step 2.* In the case  $u \in SBV^2(\Omega) \cap L^\infty(\Omega)$  the thesis descends from Theorem 5.1 and from lower semicontinuity of  $F''$ . Finally it is easy to conclude by truncation arguments and again by lower semicontinuity of  $F''$ . ■

## References

- [1] L. Ambrosio and V.M. Tortorelli, Approximation of functionals depending on jumps by elliptic functionals via  $\Gamma$ -convergence, *Comm. Pure Appl. Math.*, **XLIII** (1990), 999–1036.
- [2] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford University Press, 2000.
- [3] L. Ambrosio and V.M. Tortorelli, On the approximation of free discontinuity problems, *Boll. Un. Mat. Ital. B (7)*, **VI(1)** (1992), 105–123.
- [4] A. Braides, Approximation of free-discontinuity problems, volume 1694 of *Lecture Notes in Mathematics*, Springer Verlag, Berlin, 1998.
- [5] A. Braides and G. Dal Maso, Non-local approximation of the Mumford-Shah functional, *Calc. Var.*, **5(1)** (1997), 293–322.
- [6] A. Braides and A. Garroni, On the non-local approximation of free-discontinuity problems, *Comm. Partial Differential Equations*, **23(5-6)** (1998), 817–829.
- [7] G. Cortesani, Sequence of non-local functionals which approximate free-discontinuity problems, *Arch. Rational Mech. Anal.*, **144** (1998), 357–402.
- [8] G. Cortesani and R. Toader, A density result in SBV with respect to non-isotropic energies, *Nonlinear Anal.*, **38(5)** (1999), 585–604.
- [9] G. Cortesani and R. Toader, Finite element approximation of non-isotropic free-discontinuity problems, *Numer. Funct. Anal. Optim.*, **18(9-10)** (1997), 921–940.

- [10] G. Cortesani and R. Toader, Nonlocal approximation of nonisotropic free-discontinuity problems, *SIAM J. Appl. Math.*, **59(4)** (1999), 1507–1519.
- [11] E. De Giorgi, Free discontinuity problems in calculus of variations, *Frontiers in pure and applied mathematics. A collection of papers dedicated to Jacques-Louis Lions on the occasion of his sixtieth birthday, 55 - 62*, North-Holland Publishing Co, Amsterdam, 1991.
- [12] M. Gobbino, Finite difference approximation of the Mumford-Shah functional, *Comm. Pure Appl. Math.*, **51** (1998), 197–228.
- [13] L. Lussardi, On the variational approximation of free-discontinuity functionals, *submitted paper*, (2007), 1–11.
- [14] L. Lussardi and E. Vitali, Non-local approximation of free-discontinuity functionals with linear growth: the one-dimensional case, *Ann. Mat. Pura e Appl.*, **186(4)** (2007), 722–744.
- [15] L. Lussardi and E. Vitali, Non-local approximation of free-discontinuity problems with linear growth, *ESAIM: Control. Optim. and Calc. of Var.*, **13(1)** (2007), 135–162.