

THE LOCATION OF THE HOT SPOT IN A GROUNDED CONVEX CONDUCTOR

LORENZO BRASCO, ROLANDO MAGNANINI, AND PAOLO SALANI

ABSTRACT. We investigate the location of the (unique) hot spot in a convex heat conductor with uniform initial temperature and with boundary grounded at zero temperature. We present two methods to locate the hot spot: the first is based on ideas related to the Alexandrov-Bakelmann-Pucci maximum principle and Monge-Ampère equations; the second relies on Alexandrov's reflection principle. We then show how such a problem can be simplified in case the conductor is a polyhedron. Finally, we present some numerical computations.

1. INTRODUCTION

Consider a heat conductor Ω having (positive) constant initial temperature while its boundary is constantly kept at zero temperature. This physical situation can be described by the following initial-boundary value problem for heat equation:

$$(1.1) \quad \begin{aligned} u_t = \Delta u & \quad \text{in} \quad \Omega \times (0, \infty), \\ u = 1 & \quad \text{on} \quad \Omega \times \{0\}, \\ u = 0 & \quad \text{on} \quad \partial\Omega \times (0, \infty). \end{aligned}$$

Here Ω — the *heat conductor* — is a bounded domain in the Euclidean space \mathbb{R}^N , $N \geq 2$, with Lipschitz boundary and $u = u(x, t)$ denotes the normalized temperature of the conductor at a point $x \in \Omega$ and time $t > 0$.

A *hot spot* $x(t)$ is a point at which the temperature u attains its maximum at each given time t , that is such that

$$u(x(t), t) = \max_{y \in \bar{\Omega}} u(y, t).$$

If Ω is convex (in this case $\bar{\Omega}$ is said a *convex body*), it is well-known by a result of Brascamp and Lieb [1] that $\log u(x, t)$ is concave in x for every $t > 0$ and this, together with the analyticity of u in x , implies that for every $t > 0$ there is a unique point $x(t) \in \Omega$ at which the gradient ∇u of u vanishes (see also [13]).

The aim of this paper is to give quantitative information on the hot spot's location in a convex body.

A description of the evolution with time of the hot spot can be found in [16]; we summarize it here for the reader's convenience. A classical result of Varadhan's [20] tells us where $x(t)$ is located for small times: since

$$-4t \log\{1 - u(x, t)\} \rightarrow \text{dist}(x, \partial\Omega)^2 \quad \text{uniformly for } x \in \bar{\Omega} \text{ as } t \rightarrow 0^+$$

(here $\text{dist}(x, \partial\Omega)$ is the distance of x from $\partial\Omega$), we have that

$$\text{dist}(x(t), \mathcal{M}) \rightarrow 0 \quad \text{as } t \rightarrow 0^+,$$

where

$$(1.2) \quad \mathcal{M} = \{x \in \Omega : \text{dist}(x, \partial\Omega) = r_\Omega\}$$

2010 *Mathematics Subject Classification.* 35K05; 35B38; 35B50.

Key words and phrases. Heat equation, hot spot, eigenfunctions, Santalò point.

and

$$r_\Omega = \max\{\text{dist}(y, \partial\Omega) : y \in \overline{\Omega}\}$$

is the *inradius* of Ω . In particular, we have that

$$(1.3) \quad \text{dist}(x(t), \partial\Omega) \rightarrow r_\Omega \quad \text{as } t \rightarrow 0^+,$$

For large times instead, we know that $x(t)$ must be close to the maximum point x_∞ of the first Dirichlet eigenfunction ϕ_1 of $-\Delta$. Indeed, denoting with $\lambda_1 = \lambda_1(\Omega)$ the eigenvalue corresponding to ϕ_1 , we have that $e^{\lambda_1 t} u(\cdot, t)$ converges to ϕ_1 locally in C^2 as t goes to ∞ ; therefore (see [16])

$$(1.4) \quad x(t) \rightarrow x_\infty \quad \text{as } t \rightarrow \infty.$$

While it is relatively easy to locate the set \mathcal{M} by geometrical means, (1.4) does not give much information: locating either $x(t)$ or x_∞ has more or less the same difficulty. In this paper, we shall develop geometrical means to estimate the location of $x(t)$ (or x_∞), based on two kinds of arguments.

The first is somehow reminiscent of the proof of the maximum principle of Alexandrov, Bakelmann and Pucci and of some ideas contained in [19], concerning properties of solutions of the Monge-Ampère equation. The estimates obtained in this way are applicable to any open bounded set, not necessarily convex.

Let Ω be a bounded open set and define

$$(1.5) \quad \mathcal{K} = \text{closure of the convex hull of } \Omega;$$

we shall prove the following inequality (see Theorem 2.7):

$$(1.6) \quad \text{dist}(x_\infty, \partial\mathcal{K}) \geq \alpha_N \frac{\text{diam}(\Omega)}{(|\Omega|^{1/N} \text{diam}(\Omega) \lambda_1(\Omega))^N}.$$

Here, $\text{diam}(\Omega)$ is the *diameter* of Ω and α_N is a constant, depending only on the dimension N , for which we will give the precise expression; observe that the quantity $|\Omega|^{1/N} \text{diam}(\Omega) \lambda_1(\Omega)$ is scale invariant.

When Ω is convex, more refined bounds can be derived; for instance, the following one relates the distance of x_∞ from $\partial\Omega$ to the *inradius* r_Ω and the ratio $2r_\Omega/\text{diam}(\Omega)$, that we call the *eccentricity* of Ω and denote by $\mathcal{E}(\Omega)$:

$$(1.7) \quad \text{dist}(x_\infty, \partial\Omega) \geq \beta_N r_\Omega \mathcal{E}(\Omega)^{N^2-1},$$

where again β_N is an explicit constant depending only on N (see Theorem 2.8 for its expression). We point out that the so called *Santalò point* of Ω always satisfies (1.7), hence this can also be used to locate such a point (see Section 2 and Remark 2.11).

It is worth noticing at this point that the technique employed to derive (1.6) and (1.7) applies to more general (nonlinear) equations, yielding similar estimates; these give an answer to a problem raised by Gidas, Ni and Nirenberg in [5] (see Remark 2.5, for details).

The second argument relies instead on the following idea from [3, 5]. Let \mathbb{S}^{N-1} be the unit sphere in \mathbb{R}^N . For $\omega \in \mathbb{S}^{N-1}$ and $\lambda \in \mathbb{R}$ define the hyperplane

$$(1.8) \quad \pi(\lambda, \omega) = \{x \in \mathbb{R}^N : x \cdot \omega = \lambda\},$$

and the two half-spaces

$$(1.9) \quad \pi^+(\lambda, \omega) = \{x \in \mathbb{R}^N : x \cdot \omega > \lambda\} \quad \pi^-(\lambda, \omega) = \{x \in \mathbb{R}^N : x \cdot \omega \leq \lambda\}$$

(here the symbol \cdot denotes the usual scalar product in \mathbb{R}^N). Suppose $\pi(\lambda, \omega)$ has non-empty intersection with the interior of the conductor Ω and set

$$\Omega_{\lambda, \omega}^+ = \Omega \cap \pi^+(\lambda, \omega).$$

Then if the reflection $\mathcal{T}_{\lambda,\omega}(\Omega_{\lambda,\omega}^+)$ of $\Omega_{\lambda,\omega}$ with respect to the hyperplane $\pi(\lambda,\omega)$ lies in Ω , then $\pi(\lambda,\omega)$ cannot contain any critical point of u . This is a simple consequence of *Alexandrov's reflection principle* based on Hopf's boundary point lemma (see Section 3 for details).

Based on this remark, for a convex body \mathcal{K} we can define a (convex) set $\heartsuit(\mathcal{K})$ — the *heart* of \mathcal{K} — such that $x(t) \in \heartsuit(\mathcal{K})$ for every $t > 0$ (in fact, we will prove that $\mathcal{K} \setminus \heartsuit(\mathcal{K})$ cannot contain the hot spot $x(t)$ for any $t > 0$). The heart $\heartsuit(\mathcal{K})$ of \mathcal{K} is easily obtained as the set

$$\heartsuit(\mathcal{K}) = \bigcap \{ \pi^-(\lambda,\omega) : \mathcal{T}_{\lambda,\omega}(\mathcal{K}_{\lambda,\omega}^+) \subset \mathcal{K} \}.$$

As we shall see, the two methods have their advantages and drawbacks, but they are, in a sense, complementary. On the one hand, while inequalities (1.6) and (1.7) are quite rough in the case in which Ω has some symmetry (e.g. they do not allow to precisely locate x_∞ even when Ω is a ball), by the second argument, the problem of locating $x(t)$ is quite trivial; on the other hand, while in some cases (e.g. when Ω has no symmetries or $\partial\Omega$ contains some flat parts, as Example 4.3 explains), we cannot exclude that the heart of \mathcal{K} extends up to the boundary $\partial\mathcal{K}$ of \mathcal{K} , estimates (1.6) and (1.7) turn out to be useful to quantitatively bound $x(t)$ away from $\partial\mathcal{K}$. Thus, we believe that a joint use of both of them provides a very useful method to locate $x(t)$ or x_∞ .

Studies on the problem of locating x_∞ can also be found in [6]: there, by arguments different from ours and for the two-dimensional case, the location of x_∞ is estimated within a distance comparable to the inradius, uniformly for arbitrarily large diameter.

In Section 4 we shall relate $\heartsuit(\mathcal{K})$ to a function $\mathcal{R}_\mathcal{K}$ of the direction ω — the *maximal folding function* — and we will construct ways to characterize it. We will also connect $\mathcal{R}_\mathcal{K}$ to the Fourier transform of the characteristic function of \mathcal{K} : this should have some interest from a numerical point of view. Finally, in Section 5, we will present an algorithm to compute $\mathcal{R}_\mathcal{K}$ when \mathcal{K} is a polyhedron: based on this algorithm, we shall present some numerical computations.

2. HOT SPOTS AND POLAR SETS

In this section, if not otherwise specified, Ω is a bounded open set and we denote by \mathcal{K} the set defined by (1.5), i.e. the closure of the convex hull of Ω . Notice that \mathcal{K} is a convex body, that is a compact convex set, with non empty interior. In what follows, $|E|$ denotes the N -dimensional Lebesgue measure of a set $E \subset \mathbb{R}^N$ and $|\partial E|$ the $(N-1)$ -dimensional Hausdorff measure of its boundary; also, ω_k will be the volume of the unit ball in \mathbb{R}^k .

2.1. Preliminaries. We recall here some notations from [18]. The *gauge function* j_p of \mathcal{K} centered at a point $p \in \mathcal{K}$ is the function defined by

$$j_p(x) = \min\{\lambda \geq 0 : x - p \in \lambda(-p + \mathcal{K})\}, \quad x \in \mathbb{R}^N.$$

Observe that we have $j_p(t(x-p) + p) = t j_p(x)$ for every $t > 0$; in particular, if $0 \in \mathcal{K}$ then j_0 is 1-homogeneous. We set

$$(2.1) \quad g_p(x) = \begin{cases} j_p(x) - 1, & \text{if } x \in \mathbb{R}^N \setminus \{p\}, \\ -1, & \text{if } x = p. \end{cases}$$

so that g_p is the convex function whose graph is the cone projecting $\partial\mathcal{K}$ from the point $(p, -1) \in \mathbb{R}^{N+1}$.

It is also useful to recall the definition of the *support function* $h_\mathcal{K}$ of \mathcal{K} , that is

$$(2.2) \quad h_\mathcal{K}(\xi) = \max\{x \cdot \xi : x \in \mathcal{K}\}, \quad \xi \in \mathbb{R}^N.$$

As it is easily seen, $h_{\mathcal{K}}$ is a 1-homogeneous convex function; viceversa, to any convex 1-homogeneous function h , it corresponds exactly one convex body whose support function is h (refer to [18], for instance).

The *polar set of \mathcal{K} with respect to p* is the convex set \mathcal{K}_p^* coinciding with the unit ball of the “norm” $\|\cdot\|_* = h_{\mathcal{K}}(\cdot)$ centered at p , that is

$$\mathcal{K}_p^* = \{y \in \mathbb{R}^N : (x-p) \cdot (y-p) \leq 1 \text{ for every } x \in \mathcal{K}\}.$$

Observe that properly speaking $\|\cdot\|_*$ is not a norm, since in general we have $\| -x \|_* \neq \|x\|_*$. It is easy to see that \mathcal{K}_p^* can be equivalently defined as

$$\mathcal{K}_p^* = \{y \in \mathbb{R}^N : (x-p) \cdot (y-p) \leq j_p(x) \text{ for every } x \in \mathbb{R}^N\},$$

and that if p is in the interior of \mathcal{K} , then \mathcal{K}_p^* is compact.

We also recall that for every convex body \mathcal{K} the function $\psi : \mathcal{K} \rightarrow [0, \infty]$ defined by $\psi(x) = |\mathcal{K}_x^*|$ attains a positive minimum at some point $s_{\mathcal{K}} \in \mathcal{K}$, which is called the *Santalò point* of \mathcal{K} (see [18]). When referring to the polar of \mathcal{K} with respect to its Santalò point, we simply write \mathcal{K}^* , instead of $\mathcal{K}_{s_{\mathcal{K}}}^*$: we will see that the method developed in the next subsection for estimating the hot spot, applies to the Santalò point as well.

It is not difficult to see that \mathcal{K}_p^* coincides (up to a translation) with the *subdifferential* ∂g_p of g_p at the point p , i.e.

$$(2.3) \quad \partial g_p(p) = \mathcal{K}_p^* - p,$$

where for every x_0

$$\partial g_p(x_0) = \{\xi \in \mathbb{R}^N : g_p(x) \geq g_p(x_0) + \xi \cdot (x - x_0), x \in \mathbb{R}^N\}.$$

Finally, we will need the following monotonicity property of the subdifferential of a function: the proof can be found in [8].

Lemma 2.1. *Let u_1 and u_2 be continuous convex functions on \mathcal{K} such that $u_1 = u_2$ on $\partial\mathcal{K}$. Define:*

$$\partial u_i(\mathcal{K}) = \bigcup_{x \in \mathcal{K}} \partial u_i(x), \quad i = 1, 2.$$

If $u_1 \leq u_2$ in \mathcal{K} , then $\partial u_2(\mathcal{K}) \subseteq \partial u_1(\mathcal{K})$.

2.2. The polar set with respect to the hot spot. The following result holds for a general domain Ω and is the cornerstone of our estimates: we recall that the set \mathcal{K} is given by (1.5).

Theorem 2.2. *(i) Let u be the solution of the initial-boundary value problem (1.1) and, for every fixed time $t \in (0, +\infty)$, let $x(t) \in \Omega$ be a hot spot at time t , that is a point where the value*

$$M(t) = \max_{\overline{\Omega}} u(\cdot, t)$$

is attained.

Then

$$(2.4) \quad |\mathcal{K}_{x(t)}^*| \leq [N M(t)]^{-N} \int_{\mathcal{C}(t)} |u_t(x, t)|^N dx,$$

where $\mathcal{C}(t)$ is the contact set at time t , i.e. the subset of Ω where $-u(\cdot, t)$ coincides with its convex envelope.

(ii) Let $\lambda_1(\Omega)$ and ϕ_1 be respectively the first Dirichlet eigenvalue and eigenfunction of $-\Delta$ in Ω . Let x_{∞} be a maximum point of ϕ_1 in Ω and set $M_{\infty} = \phi_1(x_{\infty})$.

Then

$$(2.5) \quad |\mathcal{K}_{x_{\infty}}^*| \leq \left[\frac{\lambda_1(\Omega)}{N M_{\infty}} \right]^N \int_{\mathcal{C}} \phi_1(x)^N dx,$$

where \mathcal{C} is the contact set of ϕ_1 , i.e. the subset of Ω where $-\phi_1$ coincides with its convex envelope.

Proof. (i) For $t \in (0, \infty)$, let $U^{(t)} : \mathcal{K} \rightarrow \mathbb{R}$ denote the convex envelope of $-u(\cdot, t)$; then, $U^{(t)}$ is a continuous convex function in \mathcal{K} , such that $U^{(t)} = 0$ on $\partial\mathcal{K}$.

The function G defined by $G(x) = M(t)g_{x(t)}(x)$, for $x \in \mathcal{K}$, is such that

$$G \geq U^{(t)} \quad \text{in } \mathcal{K} \quad \text{and} \quad G = U^{(t)} \quad \text{on } \partial\mathcal{K},$$

and hence $\partial G(\mathcal{K}) \subseteq \partial U^{(t)}(\mathcal{K})$ by Lemma 2.1. By the rescaling properties of the subdifferential and (2.3), we know that

$$\partial G(\mathcal{K}) = M(t) (\mathcal{K}_{x(t)}^* - x(t)),$$

thus,

$$|\mathcal{K}_{x(t)}^*| \leq M(t)^{-N} |\partial U^{(t)}(\mathcal{K})|.$$

On the other hand, by Sard's Lemma and the formula for change of variables (see for instance [8, Section 1.4.2]), we obtain

$$|\partial U^{(t)}(\mathcal{K})| \leq \int_{\mathcal{C}(t)} |\det D^2 u(x, t)| \, dx,$$

with $\mathcal{C}(t) = \{x \in \Omega : U^{(t)}(x) = -u(x, t)\}$. Observe that the contact set is not empty, thanks to the fact that $x(t) \in \mathcal{C}(t)$ and moreover we have $|\mathcal{C}(t)| > 0$. Now, by the arithmetic-geometric mean inequality, we have in $\mathcal{C}(t)$ that

$$|\det D^2 u(x, t)|^{1/N} \leq \frac{|\Delta u(x, t)|}{N},$$

which yields

$$\int_{\mathcal{C}(t)} |\det D^2 u(x, t)| \, dx \leq N^{-N} \int_{\mathcal{C}(t)} |\Delta u(x, t)|^N \, dx.$$

Therefore, we finally obtain that

$$|\mathcal{K}_{x(t)}^*| \leq [N M(t)]^{-N} \int_{\mathcal{C}(t)} |\Delta u(x, t)|^N \, dx.$$

and we conclude the proof by simply using the equation $\Delta u = u_t$.

(ii) The proof runs similarly to case (i). \square

Estimates (2.4) and (2.5) are generally difficult to handle. The following weaker forms of (2.5) may be more useful.

Corollary 2.3. *Under the same assumptions as in Theorem 2.2, we have:*

$$(2.6) \quad |\mathcal{K}_{x_\infty}^*| \leq \left[\frac{\lambda_1(\Omega)}{N} \right]^N |\Omega|,$$

and

$$(2.7) \quad |\mathcal{C}| \geq \left[\frac{N}{\lambda_1(\Omega)} \right]^N |\mathcal{K}^*|;$$

we recall that \mathcal{K}^* denotes the polar set of \mathcal{K} with respect to the Santalò point.

Remark 2.4. As is well-known, $|\mathcal{K}^*|$ can be estimated from below by $m_N/|\mathcal{K}|$, where m_N is a positive constant (see [18]). Thus, (2.7) becomes

$$\frac{|\mathcal{C}|}{|\mathcal{K}|} \geq m_N \left[\frac{N}{\lambda_1(\Omega)|\mathcal{K}|^{2/N}} \right]^N.$$

Remark 2.5. In [5, p. 223] the following problem is posed:

[...] Suppose $u > 0$ is a solution of

$$(2.8) \quad -\Delta u = f(u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega$$

in a bounded domain Ω in \mathbb{R}^N , say $u \in C^2(\bar{\Omega})$. Is there some $\varepsilon > 0$ depending only on Ω (i.e., independent of f and u) such that u has no stationary point in an ε -neighbourhood of $\partial\Omega$?

This is true for $N = 2$ in case $f(u) \geq 0$ for $u \geq 0$, but for $N > 2$ the problem is open. [...]

Here, we point out that by the same arguments used for Theorem 2.2, we can easily prove the following estimate:

$$(2.9) \quad |\mathcal{K}_{x_0}^*| \leq \left(\frac{|f(0)| + LM_0}{NM_0} \right)^N |\Omega|,$$

where L is the Lipschitz constant for f and x_0 is a point where u achieves its maximum M_0 . When $f(0) = 0$, we obtain the inequality

$$(2.10) \quad |\mathcal{K}_{x_0}^*| \leq \left(\frac{L}{N} \right)^N |\Omega|$$

— an estimate, similar to (2.6), that can be used to bound $\text{dist}(x_0, \partial\mathcal{K})$ from below in a way similar to that of Theorem 2.7 below.

An interesting instance of (2.9) occurs when $f \equiv 1$ — in this case u is the *torsional creep* of an infinite bar with cross-section Ω ; we thus obtain:

$$|\mathcal{K}_{x_0}^*| \leq \frac{|\Omega|}{(NM_0)^N}.$$

This inequality can also be viewed as an estimate for the maximum M_0 in the spirit of the Alexandrov-Bakelman-Pucci principle.

Using the definition of the polar set, it is easy to see that $|\mathcal{K}_x^*|$ goes to ∞ as the point x approaches the boundary. The following lemma gives a quantitative version of this fact and helps us to provide explicit estimates of the position of x_∞ .

Lemma 2.6. *Let p be any point belonging to the interior of \mathcal{K} and define $R(p) = \max\{|p - y| : y \in \partial\mathcal{K}\}$. Then*

$$(2.11) \quad |\mathcal{K}_p^*| \geq \frac{\omega_{N-1}/N}{R(p)^{N-1} \text{dist}(p, \partial\mathcal{K})}.$$

Proof. Set $d = \text{dist}(p, \partial\mathcal{K})$ and $R = R(p)$. Obviously \mathcal{K} is contained in the ball $B(p, R)$ centered at p with radius R and in the halfspace

$$H^+ = \{y \in \mathbb{R}^N : (y - \bar{p}) \cdot (\bar{p} - p) \leq d^2\}$$

supporting \mathcal{K} at any point \bar{p} such that $|p - \bar{p}| = d$. Set $E = B(p, R) \cap H^+$; then $\mathcal{K} \subseteq E$ whence

$$\mathcal{K}_p^* \supseteq E_p^*.$$

Now notice that E_p^* is the convex envelope of the union of the ball $B(0, R^{-1})$ and the point $q = p + d^{-2}(\bar{p} - p)$; its volume is explicitly computed:

$$|E_p^*| = \frac{\omega_{N-1}}{N R^{N-1} d} \left\{ (1 - \sigma^2)^{\frac{N+1}{2}} + N \sigma \int_{-1}^{\sigma} (1 - \tau^2)^{\frac{N-1}{2}} d\tau \right\},$$

with $\sigma = d/R \in [0, 1]$. Thus, (2.11) is readily obtained by observing that the function of σ into the braces is increasing and hence bounded below by 1. \square

2.3. Estimates of critical points. We are now ready to prove the first quantitative estimate on the location of x_∞ : this will result from a combination of the previous lemma and (2.6).

Theorem 2.7. *Let Ω be an open bounded set, then estimate (1.6) holds true with constant α_N given by*

$$\alpha_N = N^{N-1} \omega_{N-1}.$$

Proof. Applying Lemma 2.6 with $p = x_\infty$ and Corollary 2.3 gives:

$$\left(\frac{\lambda_1(\Omega)}{N}\right)^N |\Omega| \geq \frac{\omega_{N-1}/N}{R(x_\infty)^{N-1} \text{dist}(x_\infty, \partial\mathcal{K})}.$$

Thus, (1.6) easily follows by observing that $\text{diam}(\mathcal{K}) = \text{diam}(\Omega) \geq R(x_\infty)$. \square

Estimate (1.6) involves the first eigenvalue $\lambda_1(\Omega)$, which in general is not easy to compute explicitly; when Ω is convex, we can estimate $\lambda_1(\Omega)$ from above by means of basic geometric quantities, thus providing an easily computable lower bound on $\text{dist}(x_\infty, \partial\Omega)$.

This is the content of the following Theorem, which represents the main contribution of this section.

Theorem 2.8. *If Ω is convex, then*

$$(2.12) \quad \text{dist}(x_\infty, \partial\Omega) \geq C_N r_\Omega \frac{\mathcal{E}(\Omega)^{N-1}}{|\partial\Omega|^N |\Omega|^{1-N}},$$

where r_Ω is the inradius of Ω , $\lambda_1(B_1)$ denotes the first Dirichlet eigenvalue of $-\Delta$ in the unit ball, $\mathcal{E}(\Omega)$ is the eccentricity of Ω (see Section 1) and the constant C_N is given by

$$C_N = \left(\frac{N}{2}\right)^{N-1} N^N \lambda_1(B_1)^{-N} \omega_{N-1}.$$

In particular, estimate (1.7) holds true with constant β_N given by

$$\beta_N = \left(\frac{N}{2}\right)^{N-1} \lambda_1(B_1)^{-N} \frac{\omega_{N-1}}{\omega_N}.$$

Proof. The proof of (2.12) readily follows by combining (1.6) to the following upper bound on $\lambda_1(\Omega)$

$$(2.13) \quad \lambda_1(\Omega) \leq \frac{\lambda_1(B_1)}{N} \frac{|\partial\Omega|}{r_\Omega |\Omega|},$$

proved in [4, Theorem 2]. Using (2.12) and the two inequalities

$$|\Omega| \geq \omega_N r_\Omega^N \quad \text{and} \quad |\partial\Omega| \leq N \omega_N \left[\frac{\text{diam}(\Omega)}{2}\right]^{N-1},$$

we obtain

$$|\partial\Omega| |\Omega|^{(1-N)/N} \leq N \omega_N^{1/N} \mathcal{E}(\Omega)^{1-N}.$$

Inserting the latter into (2.12), we end up with (1.7). \square

Inequality (2.13) is in fact a corollary of a sharper inequality holding for star-shaped sets. We can then give a refinement of (2.12) which holds in this larger class: to this end, we borrow some notations from [4].

A set Ω is said to be *strictly starshaped* with respect to a point $x_0 \in \Omega$ if it is starshaped with respect to x_0 and if its support function centered at x_0 , i.e.

$$h_{\Omega, x_0}(x) = \max_{y \in \Omega} (y - x_0) \cdot x,$$

is uniformly positive, that is $\inf_{x \in \partial\Omega} h_{\Omega, x_0}(x) > 0$. Let Ω be a strictly starshaped set with locally Lipschitz boundary, as in [4] we define

$$W(\Omega) = \inf \left\{ \int_{\partial\Omega} \frac{1}{h_{\Omega, x_0}(x)} d\sigma(x) : x_0 \in \Omega \right\},$$

where $d\sigma$ denotes surface measure on $\partial\Omega$. According to this notation, [4, Theorem 3] states:

$$(2.14) \quad \lambda_1(\Omega) \leq \frac{\lambda_1(B_1)}{N} \frac{W(\Omega)}{|\Omega|}.$$

Arguing as in Theorem 2.8 and using (2.14) in place of (2.13) gives the following estimate.

Theorem 2.9. *Let Ω be a strictly starshaped set with locally Lipschitz boundary and denote by \mathcal{K} the closure of the convex hull of Ω . Then*

$$(2.15) \quad \text{dist}(x_\infty, \partial\mathcal{K}) \geq \frac{c_N}{W(\Omega)} \left(\frac{|\Omega|}{\text{diam}(\Omega) W(\Omega)} \right)^{1-N}.$$

with constant c_N given by

$$c_N = \frac{N^{2N-1} \omega_{N-1}}{\lambda_1(B_1)^N}.$$

Remark 2.10. We remark that (2.15) is sharper and more general than (2.12), and it is at the same time more explicit than (1.6), in the sense that, differently from $\lambda_1(\Omega)$, the number $W(\Omega)$ can be computed directly from the support function (which exactly determines a convex set).

Remark 2.11. It is worth noticing that the Santalò point $s_{\mathcal{K}}$ of \mathcal{K} always satisfies (2.6) (as well as (2.4) for every $t > 0$), then it satisfies all the estimates we proved for x_∞ in this section. In particular, Theorem 2.8 (or Theorem 2.9) can be used as well to estimate the location of the Santalò point of a convex set.

3. ALEXANDROV'S REFLECTION PRINCIPLE

In this section, for the reader's convenience, we recall some relevant facts about *Aleksandrov's symmetry principle*, which has been extensively used in many situations and with various generalizations (see [3] for a good reference).

For $\omega \in \mathbb{S}^{N-1}$, let $\pi(\lambda, \omega)$, $\pi^+(\lambda, \omega)$, and $\pi^-(\lambda, \omega)$ be the sets defined in (1.8) and (1.9). Also, define a linear transformation $A_\omega : \mathbb{R}^N \rightarrow \mathbb{R}^N$ by the matrix:

$$A_\omega = (\delta_{ij} - 2\omega_i \omega_j)_{i,j=1,\dots,N}$$

where δ_{ij} is the Kronecker symbol and the ω_i are the components of ω . Then the application $\mathcal{T}_{\lambda, \omega} : \mathbb{R}^N \rightarrow \mathbb{R}^N$ defined by

$$\mathcal{T}_{\lambda, \omega}(x) = A_\omega x + 2\lambda\omega, \quad x \in \mathbb{R}^N,$$

represents the reflection with respect to $\pi(\lambda, \omega)$. As already mentioned, if Ω is a subset of \mathbb{R}^N , we set $\Omega_{\lambda, \omega}^+ = \Omega \cap \pi^+(\lambda, \omega)$.

Proposition 3.1. *Let Ω be a bounded domain in \mathbb{R}^N with Lipschitz continuous boundary $\partial\Omega$ and suppose the hyperplane $\pi(\lambda, \omega)$ defined by (1.8) has non-empty intersection with Ω . Assume that $\mathcal{T}_{\lambda, \omega}(\Omega_{\lambda, \omega}^+) \subset \Omega$.*

If Ω is not symmetric with respect to $\pi(\lambda, \omega)$, then $\pi(\lambda, \omega)$ does not contain any (spatial) critical point of the solution u of (1.1).

Proof. For $x \in \Omega_{\lambda,\omega}^+$ and $t > 0$ the function

$$v(x, t) = u(\mathcal{T}_{\lambda,\omega} x, t) - u(x, t)$$

is well-defined and is such that

$$(3.1) \quad \begin{aligned} v_t &= \Delta v & \text{in } & \Omega_{\lambda,\omega}^+ \times (0, \infty), \\ v &= 0 & \text{on } & \Omega_{\lambda,\omega}^+ \times \{0\}, \\ v &\geq 0 & \text{on } & \partial\Omega_{\lambda,\omega}^+ \times (0, \infty). \end{aligned}$$

Hence $v > 0$ in $\Omega_{\lambda,\omega}^+ \times (0, \infty)$, by the strong maximum principle for parabolic operators (see [15]). Since $v = 0$ on $(\partial\Omega_{\lambda,\omega}^+ \cap \pi(\lambda, \omega)) \times (0, \infty)$, we obtain that $\frac{\partial v}{\partial \omega} > 0$ on it (ω is in fact the interior normal unit vector), by Hopf's boundary lemma for parabolic operators. We conclude by noticing that $\frac{\partial v}{\partial \omega} = -2 \frac{\partial u}{\partial \omega}$ on $(\partial\Omega_{\lambda,\omega}^+ \cap \pi(\lambda, \omega)) \times (0, \infty)$. \square

With the same arguments and a little more work, one can extend this result to more general situations, involving nonlinearities both for elliptic and parabolic operators. As an example, here we present the following result.

Proposition 3.2. *Let Ω and $\Omega_{\lambda,\omega}^+$ satisfy the same assumptions as those of Proposition 3.1; in particular suppose that $\mathcal{T}_{\lambda,\omega}(\Omega_{\lambda,\omega}^+) \subset \Omega$.*

Let $u = u(x)$ be a solution of class $C^1(\bar{\Omega}) \cap C^2(\Omega)$ of the system:

$$(3.2) \quad \begin{aligned} \Delta u + f(u) &= 0 & \text{and } u > 0 & \text{ in } \Omega, \\ u &= 0 & \text{on } \partial\Omega, \end{aligned}$$

where f is a locally Lipschitz continuous function.

If Ω is not symmetric with respect to $\pi(\lambda, \omega)$, then $\pi(\lambda, \omega)$ does not contain any critical point of u .

Proof. The proof runs similarly to that of Proposition 3.1; the relevant changes follow. The function

$$v(x) = u(\mathcal{T}_{\lambda,\omega} x) - u(x),$$

defined for $x \in \Omega_{\lambda,\omega}$, satisfies the conditions:

$$\begin{aligned} \Delta v + c(x)v &= 0 & \text{in } & \Omega_{\lambda,\omega}^+, \\ v &\geq 0 & \text{on } & \partial\Omega_{\lambda,\omega}^+, \end{aligned}$$

where the function $c(x)$, defined by

$$c(x) = \begin{cases} \frac{f(u(\mathcal{T}_{\lambda,\omega} x)) - f(u(x))}{u(\mathcal{T}_{\lambda,\omega} x) - u(x)} & \text{for } u(\mathcal{T}_{\lambda,\omega} x) \neq u(x), \\ 0 & \text{for } u(\mathcal{T}_{\lambda,\omega} x) = u(x), \end{cases}$$

is bounded by the Lipschitz constant of f in the interval $[0, \max_{\bar{\Omega}} u]$. Hence $v \geq 0$ in $\Omega_{\lambda,\omega}^+$, by the arguments used in [3]. Let $c^-(x) = \max(-c(x), 0)$; then

$$\Delta v - c^-(x)v \leq 0 \quad \text{and} \quad v \geq 0 \quad \text{in } \Omega_{\lambda,\omega}^+$$

and the strong maximum principle can be applied to obtain that $v > 0$ in $\Omega_{\lambda,\omega}^+$. The conclusion then follows as before by Hopf's boundary lemma. \square

An immediate consequence of this theorem is the following result.

Corollary 3.3. *Let Ω and $\Omega_{\lambda,\omega}^+$ satisfy the same assumptions as those of Proposition 3.1. Let u_1 be the first (positive) eigenfunction of $-\Delta$ with homogeneous Dirichlet boundary conditions.*

If Ω is not symmetric with respect to $\pi(\lambda, \omega)$, then $\pi(\lambda, \omega)$ does not contain any critical point of u_1 .

4. THE HEART OF A CONVEX BODY

In what follows, we shall assume that $\mathcal{K} \subset \mathbb{R}^N$ is a convex body, that is a compact convex set with non-empty interior. Occasionally, we will suppose that $\mathcal{K} \subset \mathbb{R}^N$ is of class C^1 , i.e. a set whose boundary $\partial\mathcal{K}$ is an $(N - 1)$ -dimensional submanifold of \mathbb{R}^N of class C^1 .

4.1. The maximal folding function. We are interested in determining the function given by

$$(4.1) \quad \mathcal{R}_{\mathcal{K}}(\omega) := \min\{\lambda \in \mathbb{R} : \mathcal{T}_{\lambda, \omega}(\mathcal{K}_{\lambda, \omega}^+) \subseteq \mathcal{K}\}, \quad \omega \in \mathbb{S}^{N-1},$$

which will be called the *maximal folding function* of \mathcal{K} ; $\mathcal{R}_{\mathcal{K}}$ defines in turn a subset of \mathcal{K} – the *heart* of \mathcal{K} – as

$$\heartsuit(\mathcal{K}) = \{x \in \mathcal{K} : x \cdot \omega \leq \mathcal{R}_{\mathcal{K}}(\omega), \text{ for every } \omega \in \mathbb{S}^{N-1}\}.$$

Of course, $\heartsuit(\mathcal{K})$ is a closed convex subset of \mathcal{K} . Observe that $\mathcal{R}_{\mathcal{K}}$ can be bounded below and above by means of the support functions of $\heartsuit(\mathcal{K})$ and \mathcal{K} :

$$(4.2) \quad h_{\heartsuit(\mathcal{K})}(\omega) \leq \mathcal{R}_{\mathcal{K}}(\omega) \leq h_{\mathcal{K}}(\omega), \quad \omega \in \mathbb{S}^{N-1}.$$

The following results motivate our interest on $\heartsuit(\mathcal{K})$ and $\mathcal{R}_{\mathcal{K}}$.

Proposition 4.1. *Let \mathcal{K} be a convex body.*

- (i) *The hot spot $x(t)$ of \mathcal{K} , the point x_{∞} and any limit point of $x(t)$ as $t \rightarrow 0^+$ always belong to $\heartsuit(\mathcal{K})$; moreover, $x(t)$ and x_{∞} must fall in the interior of $\heartsuit(\mathcal{K})$, whenever this is non-empty.*
- (ii) *The center of mass of \mathcal{K} ,*

$$\bar{x}_{\mathcal{K}} = \frac{1}{|\mathcal{K}|} \int_{\mathcal{K}} y \, dy,$$

always belongs to the heart $\heartsuit(\mathcal{K})$ of \mathcal{K} .

- (iii) *If \mathcal{K} is strictly convex, the incenter $x_{\mathcal{K}}^+$ of \mathcal{K} belongs to $\heartsuit(\mathcal{K})$.*
- (iv) *Let $\bar{x}_{\mathcal{K}} = 0$. If there exist ℓ ($1 \leq \ell \leq N$) independent directions $\omega_1, \dots, \omega_{\ell}$ such that $\mathcal{R}_{\mathcal{K}}(\omega_j) = 0$, $j = 1, \dots, \ell$, then*

$$\heartsuit(\mathcal{K}) \subset \mathcal{K} \cap \bigcap_{j=1}^{\ell} \pi(0, \omega_j).$$

In particular, if $\ell = N$, then $\heartsuit(\mathcal{K})$ reduces to $\bar{x}_{\mathcal{K}}$ and the hot spot of \mathcal{K} is stationary.

- (v) *Let*

$$(4.3) \quad \mathfrak{r}_{\mathcal{K}} = \max_{\theta \in \mathbb{S}^{N-1}} \left\{ \min_{\omega \cdot \theta > 0} \frac{\mathcal{R}_{\mathcal{K}}(\omega) - \bar{x}_{\mathcal{K}} \cdot \omega}{\theta \cdot \omega} \right\},$$

then

$$\heartsuit(\mathcal{K}) \subseteq B(\bar{x}_{\mathcal{K}}, \mathfrak{r}_{\mathcal{K}}).$$

Proof. Items (i) and (iv) follow by observing that, for $\lambda = \mathcal{R}_{\mathcal{K}}(\omega)$, the set $\mathcal{K}_{\lambda, \omega}^+ \cup \mathcal{T}_{\lambda, \omega}(\mathcal{K}_{\lambda, \omega}^+)$ is contained in \mathcal{K} and is symmetric with respect to $\pi(\lambda, \omega)$. Hence,

$$\lambda - \bar{x}_{\mathcal{K}} \cdot \omega = \frac{1}{|\mathcal{K}|} \int_{\mathcal{K} \setminus (\mathcal{K}_{\lambda, \omega}^+ \cup \mathcal{T}_{\lambda, \omega}(\mathcal{K}_{\lambda, \omega}^+))} [\lambda - y \cdot \omega] \, dy$$

and the last term is non-negative, vanishing if and only if \mathcal{K} is ω -symmetric.

Items (ii) and (iii) are easy consequences of Proposition 3.1 and Corollary 3.3.

For a fixed $\theta \in \mathbb{S}^{N-1}$, let us define

$$\alpha(\theta) = \max\{t : \bar{x}_{\mathcal{K}} + t\theta \in \heartsuit(\mathcal{K})\},$$

which is non negative, thanks to (ii). Then $x = \bar{x}_{\mathcal{K}} + \alpha(\theta)\theta \in \heartsuit(\mathcal{K})$ and

$$\bar{x}_{\mathcal{K}} \cdot \omega + \alpha \theta \cdot \omega \leq \mathcal{R}_{\mathcal{K}}(\omega),$$

for every $\omega \in \mathbb{S}^{N-1}$ such that $\omega \cdot \theta > 0$. Hence

$$\alpha \leq \min_{\omega \in \mathbb{S}^{N-1}} \frac{\mathcal{R}_{\mathcal{K}}(\omega) - \bar{x}_{\mathcal{K}} \cdot \omega}{\theta \cdot \omega},$$

thus taking the maximum as θ varies on \mathbb{S}^{N-1} we obtain (4.3). \square

Informations on convex heat conductors with a stationary hot spot can be found in [2, 7, 10, 11, 13, 14].

Remark 4.2. Formula (4.3) deserves some comments: observe that for every fixed $\theta \in \mathbb{S}^{N-1}$, the minimum problem inside the braces amounts to finding a direction ω close to θ , so to maximize $\theta \cdot \omega$, and such that at the same time we can fold \mathcal{K} as much as possible, so to minimize the difference $\mathcal{R}_{\mathcal{K}}(\omega) - \bar{x}_{\mathcal{K}} \cdot \omega$.

We conclude this subsection by an example that shows how the simultaneous application of Proposition 4.1 and the results of Section 2 substantially benefits the problem of locating x_{∞} .

Example 4.3. Let us consider a spherical cap

$$B_{\mu}^{+} = \left\{ (x_1, \dots, x_N) \in \mathbb{R}^N : \sum_{i=1}^N x_i^2 = R^2, x_N \geq \mu \right\},$$

with $0 \leq \mu < R$. Thanks to the symmetry of B_{μ}^{+} , it is easily seen that its heart is given by

$$\heartsuit(B_{\mu}^{+}) = \{(x_1, \dots, x_N) \in \mathbb{R}^N : x_1 = \dots = x_{N-1} = 0, \mu \leq x_N \leq (R + \mu)/2\},$$

which is a vertical segment touching the boundary ∂B_{μ}^{+} at the point $(0, \dots, 0, \mu)$. In particular, by this method we can not exclude that the hotspot $x(t)$ (or the point x_{∞}) is on the boundary. However, we can now use the results of Section 2, to further sharpen this estimate on the location of x_{∞} : indeed, applying Theorem 2.8, we get

$$\text{dist}(x_{\infty}, \partial B_{\mu}^{+}) \geq (R - \mu) \left[\frac{2^{1-N-N^2} N^{N-1}}{\lambda_1(B_1)^N} \frac{\omega_{N-1}}{\omega_N} \left(\frac{R - \mu}{R + \mu} \right)^{(N^2-1)/2} \right],$$

where we used that $\text{diam}(B_{\mu}^{+}) = 2\sqrt{R^2 - \mu^2}$ and $r_{B_{\mu}^{+}} = (R - \mu)/2$.

4.2. Computing $\mathcal{R}_{\mathcal{K}}$. The following theorem whose proof can be found in [3, Theorem 5.7] guarantees that, for a regular set (not necessarily convex), the maximal folding function is never trivial.

Theorem 4.4. *Let Ω be a bounded open (not necessarily convex) subset of \mathbb{R}^N , with C^1 boundary $\partial\Omega$, and denote by \mathcal{K} the convex hull of Ω .*

For every $\omega \in \mathbb{S}^{N-1}$, there exists $\varepsilon > 0$ such that, for every λ in the interval $(h_{\mathcal{K}}(\omega) - \varepsilon, h_{\mathcal{K}}(\omega))$, we have:

- (i) $\mathcal{T}_{\lambda, \omega}(\Omega_{\lambda, \omega}^{+}) \subset \Omega$;
- (ii) $\nu(x) \cdot \omega > 0$, for every $x \in \partial\Omega \cap \pi^{+}(\lambda, \omega)$.

Unfortunately, the previous result is just qualitative and does not give any quantitative information about the maximal folding function. Moreover, notice that the C^1 assumption on $\partial\Omega$ cannot be dropped, even in the case of a convex domain: think of the spherical cap in Example 4.3, for which we have $\mathcal{R}_{B_{\mu}^{+}}(-e_N) = h_{B_{\mu}^{+}}(-e_N)$.

In order to compute $\mathcal{R}_{\mathcal{K}}$, we need some more definitions. We set

$$\omega^\perp = \pi(0, \omega)$$

and for every $y \in \omega^\perp$ we define the segment

$$\sigma_\omega(y) = \{x \in \mathcal{K} : x = y + t\omega, t \in \mathbb{R}\}.$$

Then, we denote by $\mathcal{P}_\omega : \mathbb{R}^N \rightarrow \omega^\perp$ the projection operator on ω^\perp , that is the application defined by

$$\mathcal{P}_\omega(x) = x - (x \cdot \omega)\omega, \quad x \in \mathbb{R}^N,$$

and, for y in the set

$$\mathcal{S}_\omega(\mathcal{K}) = \omega^\perp \cap \mathcal{P}_\omega(\mathcal{K})$$

– the *shadow of \mathcal{K} in the direction ω* – we define:

$$a_\omega(y) = \min\{t \in \mathbb{R} : y + t\omega \in \mathcal{K}\} \quad \text{and} \quad b_\omega(y) = \max\{t \in \mathbb{R} : y + t\omega \in \mathcal{K}\}.$$

We say that a convex body \mathcal{K} is *ω -strictly convex* if $\partial\mathcal{K}$ does not contain any segment parallel to ω . If \mathcal{K} is ω -strictly convex, then for every $x = y + t\omega \in \partial\mathcal{K}$ (equivalently $y \in \partial\mathcal{S}_\omega(\mathcal{K})$) such that the normal $\nu(x)$ to $\partial\mathcal{K}$ at x is orthogonal to ω , the set $\sigma_\omega(x)$ degenerates to the singleton $\{x\}$.

Remark 4.5. We point out that a_ω is a convex function on $\mathcal{S}_\omega(\mathcal{K})$, while b_ω is concave; moreover, if we set

$$\begin{aligned} \text{graph}^+(a_\omega) &= \{(y, t\omega) : y \in \mathcal{S}_\omega(\mathcal{K}), t \geq a_\omega(y)\}, \\ \text{graph}^-(b_\omega) &= \{(y, t\omega) : y \in \mathcal{S}_\omega(\mathcal{K}), t \leq b_\omega(y)\}, \end{aligned}$$

we have that

$$\text{graph}^+(a_\omega) \cap \text{graph}^-(b_\omega) = \mathcal{K}$$

and, as soon as \mathcal{K} is ω -strictly convex,

$$\text{graph}(a_\omega) \cup \text{graph}(b_\omega) = \partial\mathcal{K},$$

where obviously $\text{graph}(\cdot)$ denotes the graph of the relevant functions.

Theorem 4.6. *Let $\mathcal{K} \subset \mathbb{R}^N$ be a convex body. For $\omega \in \mathbb{S}^{N-1}$ consider the function $f : \mathcal{S}_\omega(\mathcal{K}) \rightarrow \mathbb{R}$ given by*

$$(4.4) \quad f_\omega(y) = \frac{a_\omega(y) + b_\omega(y)}{2}, \quad y \in \mathcal{S}_\omega(\mathcal{K}).$$

Then

$$(4.5) \quad \mathcal{R}_{\mathcal{K}}(\omega) = \max_{y \in \mathcal{S}_\omega(\mathcal{K})} f_\omega(y).$$

Proof. Observe that

$$\mathcal{K}_{\lambda, \omega}^+ = \{y + t\omega : y \in \mathcal{S}_\omega(\mathcal{K}), \lambda < t < b_\omega(y)\}.$$

Let $\bar{\lambda} = \mathcal{R}_{\mathcal{K}}(\omega)$; since $\mathcal{T}_{\bar{\lambda}, \omega}^+(\mathcal{K}_{\bar{\lambda}, \omega}^+) \subset \mathcal{K}$, then, for every point $y + t\omega$ with $y \in \mathcal{S}_\omega(\mathcal{K})$ and $\lambda < t < b_\omega(y)$, we have that $\mathcal{T}_{\bar{\lambda}, \omega}^+(y + t\omega) \in \mathcal{K}$; in particular, for $t = b_\omega(y)$, we obtain that $2\bar{\lambda} - b_\omega(y) \geq a_\omega(y)$ and hence $\bar{\lambda} \geq f_\omega(y)$. Thus,

$$\mathcal{R}_{\mathcal{K}}(\omega) \geq \max_{y \in \mathcal{S}_\omega(\mathcal{K})} f_\omega(y).$$

If $y_0 \in \mathcal{S}_\omega(\mathcal{K})$ maximizes f_ω , by taking $\lambda = f_\omega(y_0)$, we see that $\mathcal{T}_{\bar{\lambda}, \omega}^+(y + t\omega) \in \mathcal{K}$ for every $y \in \mathcal{S}_\omega(\mathcal{K})$ and $\lambda < t < b_\omega(y)$. Therefore, $\mathcal{T}_{\bar{\lambda}, \omega}^+(\mathcal{K}_{\bar{\lambda}, \omega}^+) \subset \mathcal{K}$ and hence $\mathcal{R}_{\mathcal{K}}(\omega) \leq f_\omega(y_0)$. \square

If we now remember that, for a convex domain \mathcal{K} , the quantity

$$w_{\mathcal{K}}(\omega) = h_{\mathcal{K}}(\omega) + h_{\mathcal{K}}(-\omega),$$

is the *width of \mathcal{K} in the direction ω* , we immediately get a nice consequence of the previous Theorem.

Corollary 4.7. *Let $\mathcal{K} \subset \mathbb{R}^N$ be a convex body. Then we have the following estimate for the width of $\heartsuit(\mathcal{K})$ in the direction ω :*

$$(4.6) \quad w_{\heartsuit(\mathcal{K})}(\omega) \leq \operatorname{osc}_{y \in \mathcal{S}_{\omega}(\mathcal{K})} f_{\omega}(y).$$

Proof. We first observe that $\mathcal{S}_{-\omega}(\mathcal{K}) = \mathcal{S}_{\omega}(\mathcal{K})$, so that

$$f_{-\omega}(y) = -f_{\omega}(y), \quad y \in \mathcal{S}_{\omega}(\mathcal{K}),$$

and (4.1) yields

$$(4.7) \quad \mathcal{R}_{\mathcal{K}}(-\omega) = - \min_{y \in \mathcal{S}_{\omega}(\mathcal{K})} f_{\omega}(y).$$

Then, from the definition of width, using (4.2), (4.1) and (4.7), we get

$$\begin{aligned} w_{\heartsuit(\mathcal{K})}(\omega) &= h_{\heartsuit(\mathcal{K})}(\omega) + h_{\heartsuit(\mathcal{K})}(-\omega) \\ &\leq \mathcal{R}_{\mathcal{K}}(\omega) + \mathcal{R}_{\mathcal{K}}(-\omega) \\ &= \max_{y \in \mathcal{S}_{\omega}(\mathcal{K})} f_{\omega}(y) - \min_{y \in \mathcal{S}_{\omega}(\mathcal{K})} f_{\omega}(y) = \operatorname{osc}_{y \in \mathcal{S}_{\omega}(\mathcal{K})} f_{\omega}(y), \end{aligned}$$

thus concluding the proof. \square

Example 4.8. In general, inequality (4.6) is strict. For example, in \mathbb{R}^2 consider the ellipse given by

$$\mathcal{K} = \left\{ (t, s) \in \mathbb{R}^2 : \frac{t^2}{a^2} + \frac{s^2}{b^2} = 1 \right\},$$

with $0 < b \leq a$. The function $\mathcal{R}_{\mathcal{K}}$ can be easily computed in this case: for every $\omega = (\omega_1, \omega_2) \in \mathbb{S}^1$ we get

$$\mathcal{R}_{\mathcal{K}}(\omega) = \frac{a^2 - b^2}{\sqrt{b^2\omega_1^2 + a^2\omega_2^2}} |\omega_1\omega_2|$$

— the set $\{\mathcal{R}_{\mathcal{K}}(\omega) \omega : \omega \in \mathbb{S}^1\}$ is the image of a *quadrifolium* (a *rhodonea* with 4 petals) by the mapping $(x, y) \mapsto (x/a, y/b)$.

Thus, for example, by choosing the direction $\omega = (1/\sqrt{2}, 1/\sqrt{2})$, the right-hand side of (4.6) equals

$$\frac{a^2 - b^2}{2\sqrt{2}} \sqrt{\frac{1}{a^2 + b^2}},$$

while clearly the left-hand side is zero since $\heartsuit(\mathcal{K}) = \{(0, 0)\}$, due to the symmetries of \mathcal{K} .

This example also highlights the interest of the quantity $\operatorname{osc}_{\mathcal{S}_{\omega}(\mathcal{K})} f_{\omega} - w_{\heartsuit(\mathcal{K})}$, which can be seen as a measure of the lack of symmetry of \mathcal{K} in the direction of ω .

The function f_{ω} in (4.4) can be explicitly computed by the use of the Fourier transform: this is the content of the next result.

Theorem 4.9. *Let \mathcal{K} be a convex body and for $\omega \in \mathbb{S}^{N-1}$, let f_{ω} be the function defined in (4.4).*

$$(4.8) \quad f_{\omega}(y) = \frac{i \int_{\omega^{\perp}} \partial_{\omega} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) e^{iy \cdot \eta} d\eta}{\int_{\omega^{\perp}} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) e^{iy \cdot \eta} d\eta}, \quad y \in \mathcal{S}_{\omega}(\mathcal{K}),$$

where $\widehat{\mathcal{X}}_{\mathcal{K}}$ denotes the Fourier transform of the characteristic function of \mathcal{K} and ∂_{ω} differentiation in the direction ω .

Proof. For $x \in \mathcal{K}$ and $\xi \in \mathbb{R}^N$ we write $x = y + t\omega$ and $\xi = \eta + \tau\omega$, with $y \in \mathcal{S}_{\omega}(\mathcal{K})$, $\eta \in \omega^{\perp}$ and $t, \tau \in \mathbb{R}$. By Fubini's theorem we compute

$$\begin{aligned} \widehat{\mathcal{X}}_{\mathcal{K}}(\xi) &= \int_{\mathcal{K}} e^{-ix \cdot \xi} dx \\ &= \int_{\mathcal{S}_{\omega}(\mathcal{K})} \left(\int_{-\infty}^{\infty} \mathcal{X}_{\mathcal{K}}(y + t\omega) e^{-it\tau} dt \right) e^{-iy \cdot \eta} dy \\ (4.9) \quad &= \int_{\mathcal{S}_{\omega}(\mathcal{K})} \left(\int_{a_{\omega}(y)}^{b_{\omega}(y)} e^{-it\tau} dt \right) e^{-iy \cdot \eta} dy. \end{aligned}$$

For $\tau = 0$ we then obtain:

$$(4.10) \quad \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) = \int_{\mathcal{S}_{\omega}(\mathcal{K})} [b_{\omega}(y) - a_{\omega}(y)] e^{-iy \cdot \eta} dy.$$

Therefore, by the inversion formula for the Fourier transform, we have:

$$(4.11) \quad \frac{1}{(2\pi)^{N-1}} \int_{\omega^{\perp}} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) e^{iy \cdot \eta} d\eta = \begin{cases} b_{\omega}(y) - a_{\omega}(y), & y \in \mathcal{S}_{\omega}(\mathcal{K}), \\ 0, & y \in \omega^{\perp} \setminus \mathcal{S}_{\omega}(\mathcal{K}). \end{cases}$$

By (4.9), we also obtain that

$$\begin{aligned} \partial_{\omega} \widehat{\mathcal{X}}_{\mathcal{K}}(\xi) &= \frac{d}{d\tau} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta + \tau\omega) = -i \int_{\mathcal{S}_{\omega}(\mathcal{K})} \left(\int_{a_{\omega}(y)}^{b_{\omega}(y)} t e^{-it\tau} dt \right) e^{-iy \cdot \eta} dy, \\ \partial_{\omega} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) &= -i \int_{\mathcal{S}_{\omega}(\mathcal{K})} \frac{b_{\omega}(y)^2 - a_{\omega}(y)^2}{2} e^{-iy \cdot \eta} dy, \quad \eta \in \omega^{\perp}, \end{aligned}$$

and hence

$$(4.12) \quad \frac{i}{(2\pi)^{N-1}} \int_{\omega^{\perp}} \partial_{\omega} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) e^{iy \cdot \eta} d\eta = \begin{cases} \frac{b_{\omega}(y)^2 - a_{\omega}(y)^2}{2}, & y \in \mathcal{S}_{\omega}(\mathcal{K}), \\ 0, & y \in \omega^{\perp} \setminus \mathcal{S}_{\omega}(\mathcal{K}). \end{cases}$$

Formula (4.8) follows from (4.12) and (4.11) at once. \square

Remark 4.10. If \mathcal{K} is a polygon, $\widehat{\mathcal{X}}_{\mathcal{K}}$ can be explicitly computed in terms of the vertices of \mathcal{K} . Let $\mathcal{K} \subset \mathbb{R}^2$ be a (convex) polygon with vertices p_1, \dots, p_n ; we assume that p_1, \dots, p_n are ordered counterclockwise and we set $p_{n+1} = p_1$.

Rewriting $\widehat{\mathcal{X}}_{\mathcal{K}}$ as a boundary integral (see [9]) by means of the divergence theorem, we have that

$$\widehat{\mathcal{X}}_{\mathcal{K}}(\xi) = -\frac{1}{|\xi|^2} \sum_{j=1}^n |p_{j+1} - p_j| (\nu_j \cdot \xi) \frac{e^{-ip_{j+1} \cdot \xi} - e^{-ip_j \cdot \xi}}{(p_{j+1} - p_j) \cdot \xi},$$

where

$$\nu_j = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \frac{p_{j+1} - p_j}{|p_{j+1} - p_j|}, \quad j = 1, \dots, n,$$

is the exterior normal to the j -th side of \mathcal{K} . Also, $\partial_{\omega} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta)$ is easily computed from the previous expression:

$$\begin{aligned} \partial_{\omega} \widehat{\mathcal{X}}_{\mathcal{K}}(\eta) &= \frac{1}{|\eta|} \sum_{j=1}^n |p_{j+1} - p_j|^2 \frac{e^{-ip_{j+1} \cdot \eta} - e^{-ip_j \cdot \eta}}{[(p_{j+1} - p_j) \cdot \eta]^2} \\ &\quad + \frac{i}{|\eta|^2} \sum_{j=1}^n |p_{j+1} - p_j| (\nu_j \cdot \eta) \frac{(p_{j+1} \cdot \omega) e^{-ip_{j+1} \cdot \eta} - (p_j \cdot \omega) e^{-ip_j \cdot \eta}}{(p_{j+1} - p_j) \cdot \eta}. \end{aligned}$$

4.3. Necessary optimality conditions. We conclude this section by presenting some necessary conditions for the optimality of f_ω . To this aim, we first state and prove an easy technical result for the subdifferential of a function.

Lemma 4.11. *Let $\Omega \subset \mathbb{R}^k$ be a convex open set. Let φ and ψ be a convex and, respectively, a concave function from Ω to \mathbb{R} . If $\varphi + \psi$ attains its maximum at a point $y_0 \in \Omega$, then*

$$(4.13) \quad \partial\varphi(y_0) \subset \partial(-\psi)(y_0).$$

Proof. It is clear that both $\partial\varphi(y_0)$ and $\partial(-\psi)(y_0)$ are non-empty. Since y_0 is a maximum point, we get

$$\varphi(y_0) + \psi(y_0) \geq \varphi(y) + \psi(y) \text{ for every } y \in \Omega,$$

and hence

$$\varphi(y) - \varphi(y_0) + \xi \cdot (y - y_0) \leq -\psi(y) + \psi(y_0) + \xi \cdot (y - y_0)$$

for every $\xi \in \mathbb{R}^N$, and $y \in \Omega$. If $\xi \in \partial\varphi(y_0)$, then we have $\xi \in \partial(-\psi)(y_0)$. \square

As a consequence of the definitions of a_ω and b_ω , we have that $\partial a_\omega(y_0) \cup \partial(-b)_\omega(y_0) = \emptyset$ implies that y_0 belongs to the boundary of $\mathcal{S}_\omega(\mathcal{K})$.

We are now in a position to state a necessary optimality condition.

Theorem 4.12. *Let $\mathcal{K} \subset \mathbb{R}^N$ be a convex body and $\omega \in \mathbb{S}^{N-1}$. Suppose that f_ω attains its maximum at a point $y_0 \in \mathcal{S}_\omega(\mathcal{K})$, that is*

$$\mathcal{R}_\mathcal{K}(\omega) = f_\omega(y_0).$$

Set $\lambda = \mathcal{R}_\mathcal{K}(\omega)$ and for every $x_0 \in \mathcal{P}_\omega^{-1}(y_0) \cap \partial\mathcal{K}$ denote by x_0^λ its reflection with respect to the optimal hyperplane, that is $x_0^\lambda = \mathcal{T}_{\lambda, \omega} x_0$.

Then:

(i) *if $x_0 \neq x_0^\lambda$, we have*

$$(4.14) \quad \mathcal{A}_\omega(N_\mathcal{K}(x_0^\lambda)) \subseteq N_\mathcal{K}(x_0),$$

where

$$N_\mathcal{K}(x) = \{\xi \in \mathbb{R}^N \setminus \{0\} : x \cdot \xi = h_\mathcal{K}(\xi)\} \cup \{0\}.$$

denotes the normal cone of \mathcal{K} at a point $x \in \partial\mathcal{K}$;

(ii) *if $x_0 = x_0^\lambda$, there holds*

$$(4.15) \quad \mathcal{A}_\omega(N_\mathcal{K}^-(x_0)) \subseteq N_\mathcal{K}^+(x_0),$$

where

$$N_\mathcal{K}^-(x) = \{\xi \in N_\mathcal{K}(x) : \xi \cdot \omega \leq 0\} \quad \text{and} \quad N_\mathcal{K}^+(x) = \{\xi \in N_\mathcal{K}(x) : \xi \cdot \omega \geq 0\}.$$

Proof. We can suppose for simplicity that $\omega = e_N = (0, \dots, 0, 1)$. We first suppose that y_0 is an interior point. Since by its very definition f_ω is the sum of a convex function and a concave one, by Lemma 4.11

$$(4.16) \quad \partial a_\omega(y_0) \subset \partial(-b_\omega)(y_0).$$

Let us now set $x_0 = y_0 + b_\omega(y_0)\omega$; the reflection of x_0 in the optimal hyperplane is $x_0^\lambda = y_0 + a_\omega(y_0)\omega$ and

$$N_\mathcal{K}(x_0) = \{(\eta, 1) : \eta \in \partial(-b_\omega)(y_0)\} \quad N_\mathcal{K}(x_0^\lambda) = \{(\eta, -1) : \eta \in \partial a_\omega(y_0)\},$$

for $\text{graph}^+(a_\omega) \cap \text{graph}^-(b_\omega) = \mathcal{K}$. Then, since $\mathcal{A}_\omega(\eta, 1) = (\eta, -1)$, (4.16) implies (4.14).

If y_0 is on the boundary of $\mathcal{S}_\omega(\mathcal{K})$, then $\partial(-b_\omega)(y_0)$ or $\partial a_\omega(y_0)$ may be empty: this is clearly the case if $\text{graph}(a_\omega)$ or $\text{graph}(b_\omega)$ have some vertical parts. Observe that actually we have the following possibilities:

- (1) $\partial(-b_\omega)(y_0) = \partial a_\omega(y_0) = \emptyset$;
- (2) $\partial(-b_\omega)(y_0) \neq \emptyset$.

If (1) holds, then at every $x_0 \in \mathcal{P}^{-1}(y_0) \cap \partial\mathcal{K}$, the convex body \mathcal{K} has only supporting hyperplanes parallel to e_N : these are invariant with respect to the action of \mathcal{A}_ω , so that their reflections are supporting hyperplanes for \mathcal{K} at x_0^λ and formula (4.14) or (4.15) easily follows.

If (2) holds, we have $x_0 = x_0^\lambda$ and let us call $\Omega = \mathcal{K}_{\lambda,\omega}^+ \cup \mathcal{T}_{\lambda,\omega}(\mathcal{K}_{\lambda,\omega}^+)$. Then

$$N_{\mathcal{K}}(x_0) \subset N_\Omega(x_0)$$

and we have $N_\Omega^+(x_0) = N_{\mathcal{K}}^+(x_0)$, so that

$$N_\Omega^-(x_0) = N_\Omega(x_0) \setminus N_\Omega^+(x_0) \supset N_{\mathcal{K}}(x_0) \setminus N_{\mathcal{K}}^+(x_0) = N_{\mathcal{K}}^-(x_0).$$

By observing that $N_\Omega^-(x_0) = \mathcal{A}_\omega(N_\Omega^+(x_0)) = \mathcal{A}_\omega(N_{\mathcal{K}}^+(x_0))$, (4.15) follows. \square

Corollary 4.13. *Under the same notations of Theorem 4.12, if $\partial\mathcal{K}$ admits a (unique) unit normal ν at the point x_0 , then it admits a unit normal at the point x_0^λ too and*

$$(4.17) \quad \mathcal{A}_\omega \nu(x_0^\lambda) = \nu(x_0).$$

In particular, if $x_0 = x_0^\lambda$ we have $\nu(x_0) \in \omega^\perp$.

Proof. It is sufficient to observe that in this case

$$N_{\mathcal{K}}(x_0) = \{\xi : \xi = t\nu(x_0), t > 0\} \cup \{0\},$$

and hence (4.17) is a consequence of (4.14) or (4.15). \square

Using Theorem 4.12, we obtain an interesting upper bound on the maximal folding function for a strictly convex domain, in terms of its support function.

Proposition 4.14. *If \mathcal{K} is strictly convex, then for every $\omega \in \mathbb{S}^{N-1}$*

$$(4.18) \quad \mathcal{R}_{\mathcal{K}}(\omega) \leq \max \left\{ \left(\frac{\nabla h_{\mathcal{K}}(\xi) + \nabla h_{\mathcal{K}}(\mathcal{A}_\omega \xi)}{2} \right) \cdot \omega : \xi \in \mathbb{R}^N \setminus \{0\} \right\}.$$

Proof. First observe that thanks to the 1-homogeneity of the support function, the maximization problem in (4.18) can be equivalently settled in \mathbb{S}^{N-1} .

The strict convexity of \mathcal{K} implies that $h_{\mathcal{K}} \in C^1(\mathbb{R}^N \setminus \{0\})$ (see [18]). Moreover, $\nabla h_{\mathcal{K}}(\theta) = x$ for every $\theta \in N_{\mathcal{K}}(x)$, with $x \in \partial\mathcal{K}$. Thus, with the same notations as in Theorem 4.12, $x_0^\lambda = \nabla h_{\mathcal{K}}(\theta)$ for every $\theta \in N_{\mathcal{K}}(x_0^\lambda)$ and using the condition (4.14), we also get $x_0 = \nabla h_{\mathcal{K}}(\mathcal{A}_\omega(\theta))$; on the other hand, $a_\omega(y_0) = x_0^\lambda \cdot \omega$ and $b_\omega(y_0) = x_0 \cdot \omega$, which implies the following

$$f_\omega(y_0) = \frac{\nabla h_{\mathcal{K}}(\theta) + \nabla h_{\mathcal{K}}(\mathcal{A}_\omega \theta)}{2} \cdot \omega.$$

Hence we can conclude by simply applying Theorem 4.6. \square

Remark 4.15. If \mathcal{K} is symmetric with respect to a hyperplane orthogonal to ω , then equality holds in (4.18) and both quantities equal $\bar{x}_{\mathcal{K}} \cdot \omega$. Otherwise, in general inequality (4.18) is strict as Figure 1 informs us.

Notice that, following an argument similar to that of the proof of Proposition 4.18, we can in fact give a precise characterization of the maximal folding function $\mathcal{R}_{\mathcal{K}}$ in terms of the support function $h_{\mathcal{K}}$. Precisely the following holds

$$(4.19) \quad \mathcal{R}_{\mathcal{K}}(\omega) = \max_{\xi \in \Sigma(\omega)} \frac{\nabla h_{\mathcal{K}}(\xi) + \nabla h_{\mathcal{K}}(\mathcal{A}_\omega \xi)}{2},$$

where

$$\Sigma(\omega) = \{\xi \in \mathbb{R}^N \setminus \{0\} : \nabla h_{\mathcal{K}}(\xi) = \nabla h_{\mathcal{K}}(\mathcal{A}_\omega \xi) + \mu \omega \text{ for some } \mu \in \mathbb{R}\}.$$




FIGURE 1. Here $\omega = (1, 0, \dots, 0)$, $x = \nabla h_{\mathcal{K}}(\xi)$, $y = \nabla h_{\mathcal{K}}(\mathcal{A}_{\omega}(\xi))$; the intersection of the two straight lines corresponds to $\mathcal{R}_{\mathcal{K}}(\omega)$, the dark dot corresponds to $\frac{1}{2}(x + y)$.

If \mathcal{K} is not strictly convex (and then $h_{\mathcal{K}}$ is not C^1) the above formula still remains valid, up to suitably interpreting the gradient of $h_{\mathcal{K}}$ as the subdifferential $\partial h_{\mathcal{K}}$.

5. NUMERICAL EXAMPLES

5.1. The case of convex polyhedrons. If \mathcal{K} is a convex polyhedron, then the conclusions of Theorem 4.6 can be improved: roughly speaking, we can discretize the optimization problem (4.1), by only visiting the projections of the vertices of \mathcal{K} on ω^{\perp} . We begin with the following general result.

Lemma 5.1. *Let A and B be convex sets such that $A \subseteq B$ and let $x \in \partial A \cap \partial B$.*

If ∂A contains a segment ℓ and x belongs to the relative interior of ℓ , then ℓ is also contained in ∂B .

Proof. In other words, if A “touches” B from the interior at x and x is contained in the interior of some segment on the boundary of A , then the boundary of B must contain all the segment at the same.

Indeed, let π be a support hyperplane to B at x and denote by π^+ the half-space delimited by π and containing B ; then π is also a support hyperplane to A at x and $A \subseteq \pi^+$. Thus, $\ell \subset \pi^+$ while $x \in \ell \cap \pi \neq \emptyset$; this implies $\ell \subset \pi$, since x is not an endpoint of ℓ , and hence $\ell \subset \partial B$. \square

Corollary 5.2. *Under the same assumptions and notations of Theorem 4.12, if x_0 belongs to the relative interior of a segment ℓ contained in $\partial \mathcal{K}$, then $\mathcal{T}_{\lambda, \omega}(\ell) \subset \partial \mathcal{K}$ and*

$$(5.1) \quad f_{\omega}(y) = \mathcal{R}_{\mathcal{K}}(\omega) \quad \text{for every } y \in \mathcal{P}_{\omega}(\ell).$$

Proof. The proof follows from the previous lemma by setting $B = \mathcal{K}$ and $A = \mathcal{K} \cap \mathcal{T}_{\lambda, \omega}(\mathcal{K})$ and using the definition of f_{ω} . \square

Theorem 5.3. *Let $x_1, \dots, x_s \in \mathbb{R}^N$ be the vertices of an N -dimensional convex polyhedron $\mathcal{K} \subset \mathbb{R}^N$, so that*

$$\mathcal{K} = \left\{ x \in \mathbb{R}^N : x = \sum_{i=1}^s \lambda_i x_i, \text{ with } \sum_{i=1}^s \lambda_i = 1, \lambda_i \in [0, 1] \right\},$$




FIGURE 2. The two cases (i) and (ii).

For a fixed $\omega \in \mathbb{S}^{N-1}$, let f_ω be the function defined by (4.4).

Then $\mathcal{R}_\mathcal{K}$ is the solution of the following discrete optimization problem

$$(5.2) \quad \mathcal{R}_\mathcal{K}(\omega) = \max \{f_\omega(y_j) : j = 1, \dots, s\},$$

where $y_j = \mathcal{P}_\omega(x_j)$ is the projection of x_j on $\mathcal{S}_\omega(\mathcal{K})$, for every $j = 1, \dots, s$.

Proof. By definition (4.1) and Theorem 4.6, we know that the value $\lambda = \mathcal{R}_\mathcal{K}(\omega)$ can possibly be achieved when the boundary of the reflected cap $\mathcal{T}_{\lambda, \omega}(\mathcal{K}_{\lambda, \omega}^+)$ is tangent to that of \mathcal{K} either

- (i) at a point $p \notin \pi(\lambda, \omega)$, or
- (ii) at a point $q \in \pi(\lambda, \omega)$

(see Figure 2). Thus, the maximum of f_ω is attained at the projection of either p or q on $\mathcal{S}_\omega(\mathcal{K})$.

Now, let \mathcal{K} be a convex polyhedron. If p is not a vertex of \mathcal{K} , then p belongs to the relative interior of some m -dimensional facet of $\partial\mathcal{K}$, with $1 \leq m \leq N - 1$, and hence it belongs to the relative interior of a segment ℓ with (at least) one end at some vertex v of $\partial\mathcal{K}$.




FIGURE 3. The sets \mathcal{K} and $\heartsuit(\mathcal{K})$ in two examples. In the first picture observe that, by means of (1.7), we also know that x_∞ is at a positive (and computable) distance from the boundary of \mathcal{K} .

By Corollary 5.2, we then have:

$$\mathcal{R}_{\mathcal{K}}(\omega) = f_{\omega}(\mathcal{P}_{\omega}(p)) = f_{\omega}(\mathcal{P}_{\omega}(v)).$$

In case (ii), the m -dimensional facet F of $\partial\mathcal{K}$ containing q must be orthogonal to the hyperplane $\pi(\lambda, \omega)$; however, the same argument used for case (i) can easily be worked out in F . \square

The pictures in Figure 3 show two convex polygons with their relative hearts. The hearts have been drawn, by using MATLAB, by an algorithm based on Theorem 5.3.

Acknowledgements. The first author was partially supported by the European Research Council under FP7, Advanced Grant n. 226234 “Analytic Techniques for Geometric and Functional Inequalities”, while the second and third authors were partially supported by the PRIN-MIUR grant “Aspetti geometrici delle equazioni alle derivate parziali e questioni connesse”.

REFERENCES

- [1] H. L. Brascamp, E. H. Lieb, *On extensions of Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log-concave functions, and with an application to the diffusion equation*, J. Funct. Anal. 22 (1976), 366–389.
- [2] M. Chamberland, D. Siegel, *Convex domains with stationary hot spots*, Math. Meth. Appl. Sci. 20 (1997) 1163–1169.
- [3] L. E. Fraenkel, *Introduction to maximum principles and symmetry in elliptic problems*, Cambridge University Press 2000.
- [4] P. Freitas, D. Krejčířík, *A sharp upper bound for the first Dirichlet eigenvalue and the growth of the Isoperimetric Constant of convex domains*, Proc. American Math. Soc. 136 (2008), 2997–3006.
- [5] B. Gidas, W. M. Ni, L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. 68 (1979), no. 3, 209–243.
- [6] D. Grieser, D. Jerison, *The size of the first eigenfunction on a convex planar domain*, J. Amer. Math. Soc. 11 (1998), 41–72.
- [7] R. Gulliver, N.B. Willms, *A conjectured heat flow problem*, In Solutions, SIAM Review 37, (1995) 100–104.
- [8] C. E. Gutierrez, *The Monge-Ampère equation*, Progress in Nonlinear Differential Equations and their Applications, 44. Birkhäuser Boston, Inc., Boston, MA, 2001.
- [9] C. S. Herz, *Fourier transforms related to convex sets*, Ann. Math. 75 (1962) 81–92.
- [10] B. Kawohl, *A conjectured heat flow problem*, In Solutions, SIAM Review 37 (1995) 104–105.
- [11] M. S. Klamkin, *A conjectured heat flow problem*, In Problems, SIAM Review 36 (1994) 107.
- [12] N. Korevaar, *Convex solutions to nonlinear elliptic and parabolic boundary value problems*, Indiana Univ. Math. J. 32 (1983), 603–614.
- [13] R. Magnanini, S. Sakaguchi, *On heat conductors with a stationary hot spot*, Ann. Mat. Pura Appl. 183 (2004), 1–23.
- [14] R. Magnanini, S. Sakaguchi, *Polygonal Heat Conductors with a Stationary Hot Spot*, J. Anal. Math. 105 (2008), 1–18.
- [15] M. H. Protter, H. F. Weinberger, *Maximum Principles in Differential Equations*, Prentice-Hall, Englewood Cliffs, N. J., 1967.
- [16] S. Sakaguchi, *Behavior of spatial critical points and zeros of solutions of diffusion equations*, Selected papers on differential equations and analysis, 15–31, Amer. Math. Soc. Transl. Ser. 2, 215, Amer. Math. Soc., Providence, RI, 2005.
- [17] J. Serrin, *A symmetry problem in potential theory*, Arch. Rational Mech. Anal. 43 (1971), 304–318.
- [18] R. Schneider, *Convex bodies: the Brunn-Minkowski theory*, Cambridge University Press 1993.
- [19] G. Talenti, *Some estimates of solutions to Monge-Ampère type equations in dimension two*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 8 (1981), 183–230.
- [20] S. R. S. Varadhan, *On the behaviour of the fundamental solution of the heat equation with variable coefficients*, Comm. Pure Appl. Math. 20, (1967), 431–455.

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI "R. CACCIOPPOLI", UNIVERSITÀ DI NAPOLI
"FEDERICO II", COMPLESSO DI MONTE S. ANGELO, VIA CINTIA, 80126 NAPOLI, ITALY
E-mail address: `lorenzo.brasco@unina.it`

DIPARTIMENTO DI MATEMATICA "U. DINI", UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A,
50134 FIRENZE, ITALY
E-mail address: `magnanin@math.unifi.it`, `paolo.salani@math.unifi.it`