

## On a class of differential equations with self-reference

MICHELE MIRANDA Jr. – EDUARDO PASCALI

ABSTRACT: *We give a result of local existence and uniqueness for equations of the type*

$$\frac{\partial^2}{\partial t^2}u(x, t) = k_1u \left( \frac{\partial^2}{\partial t^2}u(x, t) + k_2u(x, t), t \right)$$

where  $k_1$  and  $k_2$  are given real numbers or real functions  $k_i = k_i(x, t)$ .

### 1 – Introduction

In previous notes we have started the study of a new type of differential equations that we have called self-referred and hereditary (see [4], [5]). The abstract scheme that we can think in order to embed such equations can be stated as follows. Given two functionals  $A : X \rightarrow R$  and  $B : X \rightarrow R$ , where  $X$  is a space of functions  $u = u(x, t)$ , we look for a solution of the equation:

$$(Au)(x, t) = u(Bu(x, t), t)$$

under the condition that  $u(x, 0)$  is a given function.

When  $Bu$  is an "hereditary" operator as for instance

$$Bu(x, t) = \int_0^t u(x, \tau) d\tau$$

and similar, the previous equation is said of hereditary and self-referred tupe. General equations with "hereditary" operator has been studied since the beginnig of XX-th century (see [2], [3] and [8]).

---

KEY WORDS AND PHRASES: **MISSING**  
A.M.S. CLASSIFICATION: 47J35 – 45G10

At the previous general scheme can be associated the class of equations that we will study in this note; in fact, the equations that we will consider are of the type:

$$(1.1) \quad \frac{\partial^2}{\partial t^2} u(x, t) = k_1 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_2 u(x, t), t \right),$$

where  $k_1, k_2$  are given real numbers or also suitable real functions. For such equation, arguing as in [5] [4], we can establish a result of local existence and uniqueness, under suitable conditions on  $k_1, k_2$  and on the initial datum  $u(x, 0)$ . We can also establish similar results for more general equations than the previous one, for instance for equations of the type:

$$(1.2) \quad \frac{\partial^2}{\partial t^2} u(x, t) = k_1 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_2 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_3 u(x, t), t \right), t \right),$$

The given general scheme is also related to recent papers as [1], [6] and [7].

## 2 – The local existence and uniqueness result

If we consider the equation

$$(2.3) \quad \frac{\partial^2}{\partial t^2} u(x, t) = k_1 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_2 u(x, t), t \right)$$

we will assume, for the sake of simplicity, that  $|k_1| = |k_2| = 1$ . Let  $\alpha, \beta \in L^\infty(\mathbb{R}, \mathbb{R}) \cap Lip(\mathbb{R}, \mathbb{R})$  be given functions, and let us denote by  $L_\alpha, L_\beta$  the Lipchitz contants; we will also assume that  $L_\alpha < \frac{1}{2}$ . We define:

$$u_0(x, t) = \alpha(x) + t\beta(x)$$

and we consider

$$u_1(x, t) = \alpha(x) + t\beta(x) + k_1 \int_0^t \int_0^\tau u_0(k_2 u_0(x, s)) ds d\tau$$

and then, for all  $n \in \mathbb{N}$ :

$$u_{n+1}(x, t) = \alpha(x) + t\beta(x) + k_1 \int_0^t \int_0^\tau u_n \left( \frac{\partial^2}{\partial s^2} u_n(x, s) + k_2 u_n(x, s), s \right) ds d\tau.$$

We want to prove that the sequence  $u_n$  is a Cauchy sequence and that the limit function  $u_\infty$  is a solution of equation (2.3). It is easy to prove that

$$|u_n(x, t)| \leq (||\alpha||_\infty + ||\beta||_\infty) e^t \quad \forall x \in \mathbb{R}, t \geq 0, n \in \mathbb{N}.$$

We notice that:

$$|u_1(x, t) - u_0(x, t)| \leq \left( \|\alpha\|_\infty \frac{t^2}{2!} + \|\beta\|_\infty \frac{t^3}{3!} \right) = A_1(t).$$

Moreover, for the second derivative there holds the following estimate:

$$\left| \frac{\partial^2}{\partial t^2} u_1(x, t) - \underbrace{\frac{\partial^2}{\partial t^2} u_0(x, t)}_{=0} \right| \leq (\|\alpha\|_\infty + t\|\beta\|_\infty) = B_1(t)$$

and there holds:

$$A_1(t) = \int_0^t \int_0^\tau B_1(s) ds d\tau.$$

Arguing in the same way, for the second step of the sequence, there holds:

$$\begin{aligned} |u_2(x, t) - u_1(x, t)| &\leq \int_0^t \int_0^\tau \left| u_1 \left( \frac{\partial^2}{\partial s^2} u_1(x, s) + k_2 u_1(x, s), s \right) + \right. \\ &\quad - u_0 \left( \frac{\partial^2}{\partial s^2} u_1(x, s) + k_2 u_1(x, s) \right) + \\ &\quad + u_0 \left( \frac{\partial^2}{\partial s^2} u_1(x, s) + k_2 u_1(x, s) \right) + \\ &\quad \left. - u_0 \left( \frac{\partial^2}{\partial s^2} u_0(x, s) + k_2 u_0(x, s) \right) \right| ds d\tau \leq \\ &\leq \int_0^t \int_0^\tau \left( A_1(s) + \right. \\ &\quad \left. + L_0(s) \left( \left| \frac{\partial^2}{\partial s^2} u_1(x, s) - \frac{\partial^2}{\partial s^2} u_0(x, s) \right| + \right. \right. \\ &\quad \left. \left. + A_1(s) \right) \right) ds d\tau, \end{aligned}$$

where we have defined  $L_0(t) = L_\alpha + tL_\beta$ . Then, by induction, if we define;

$$(2.4) \quad A_{n+1}(t) = \int_0^t \int_0^\tau \left( A_n(s)(1 + L_{n-1}(s)) + L_{n-1}(s)B_n(s) \right) ds,$$

it is now easy to notice that:

$$|u_{n+1}(x, t) - u_n(x, t)| \leq A_{n+1}(t).$$

In addition, if we define;

$$(2.5) \quad B_{n+1}(t) = A_n(t)(1 + L_{n-1}(t)) + L_{n-1}(t)B_n(t)$$

there holds:

$$A_{n+1}(t) = \int_0^t \int_0^\tau B_{n+1}(s) ds d\tau,$$

and we obtain that:

$$\left| \frac{\partial}{\partial t} u_{n+1}(x, t) - \frac{\partial}{\partial t} u_n(x, t) \right| \leq B_{n+1}(t).$$

We want now prove that the  $(u_n)_n$  is an equi-Lipschitz sequence. Let us then consider;

$$\begin{aligned} |u_1(x, t) - u_1(y, t)| &\leq (L_\alpha + tL_\beta)|x - y| + \\ &+ \left| \int_0^t \int_0^\tau u_0 \left( \frac{\partial^2}{\partial s^2} u_0(x, s) + k_2 u_0(x, s) \right) + \right. \\ &\left. - u_0 \left( \frac{\partial^2}{\partial s^2} u_0(y, s) + k_2 u_0(y, s) \right) \right| \leq \\ &\leq \left( L_0(t) + \int_0^t \int_0^\tau L_0(s)^2 ds d\tau \right) |x - y| \end{aligned}$$

Then, if we define, by recurrence:

$$\begin{cases} C_0(t) \equiv 0 \\ C_{n+1}(t) = L_n(t)(L_n(t) + C_n(t)), \quad \forall n \in \mathbb{N} \end{cases}$$

and

$$L_{n+1}(t) = L_0(t) + \int_0^t \int_0^\tau L_n(s)(L_n(s) + C_n(s)) ds d\tau,$$

we obtain that, for every  $n \in \mathbb{N}$ :

$$|u_{n+1}(x, t) - u_n(x, t)| \leq L_{n+1}(t)|x - y|;$$

Moreover:

$$\begin{aligned} \left| \frac{\partial^2}{\partial t^2} u_1(x, t) - \frac{\partial^2}{\partial t^2} u_1(y, t) \right| &\leq \left| u_0 \left( \frac{\partial^2}{\partial t^2} u_0(x, t) + k_2 u_0(x, t) \right) + \right. \\ &\left. - u_0 \left( \frac{\partial^2}{\partial t^2} u_0(y, t) + k_2 u_0(y, t) \right) \right|. \end{aligned}$$

Since  $0 < L_\alpha < 1/2$ , there follows that there exists  $M > 0$  such that:

$$L_\alpha < M \leq \frac{1}{2}.$$

Let us fix  $T_0 > 0$  in such a way that:

$$0 \leq t \leq T_0 \Rightarrow \begin{cases} L_\alpha + tL_\beta + M\frac{t^2}{2} \leq M \\ M + (1+M)\frac{t^2}{2} \leq h < 1 \end{cases}$$

We observe what follows:

$$L_1(t) = L_\alpha + tL_\beta + L_\alpha^2 \frac{t^2}{2} \leq L_\alpha + tL_\beta + M\frac{t^2}{2} \leq M.$$

Since

$$C_1(t) = L_0^2(t) = (L_\alpha + tL_\beta)^2 \leq L_\alpha + tL_\beta \leq M,$$

there follows

$$C_2(t) = L_1^2(t) + L_1(t)C_1(t) \leq M^2 + M \cdot M = 2M^2 \leq M.$$

Moreover

$$\begin{aligned} L_2(t) &= L_0(t) + \int_0^t \int_0^\tau (L_1^2(s) + L_1(s)C_1(s)) ds d\tau = \\ &= L_\alpha + tL_\beta + \int_0^t \int_0^\tau (M^2 + M^2) ds d\tau = \\ &\leq L_\alpha + tL_\beta + M\frac{t^2}{2} \leq M; \end{aligned}$$

and then

$$C_3(t) = L_2^2(t) + L_2(t)C_2(t) \leq M^2 + M^2 \leq M.$$

In conclusion, it is possible to prove, by induction:

$$0 \leq L_n(t) \leq M, \quad \forall n \in \mathbb{N}, \forall t \in [0, T_0].$$

Using formulae (2.4) and (2.5), we obtain, for all  $t \in [0, T_0]$  and  $n \in \mathbb{N}$ :

$$\begin{aligned} 0 \leq B_{n+1}(t) &\leq A_n(t)(1+M) + MB_n(t) \leq \\ &\leq \|B_n\|_\infty \left( \frac{t^2}{2}(1+M) + M \right). \end{aligned}$$

Hence, there follows that for all  $n \in \mathbb{N}$ :

$$\frac{\|B_{n+1}\|_\infty}{\|B_n\|_\infty} \leq h < 1.$$

Then the series

$$\sum B_{n+1}(t)$$

is totally convergent and then there exists  $v_\infty$  such that:

$$\frac{\partial^2}{\partial t^2} u_n(x, t) \rightrightarrows v_\infty(x, t), \quad x \in \mathbb{R}, t \in [0, T_0],$$

and, since

$$\|A_{n+1}\|_\infty \leq \|B_{n+1}\|_\infty \frac{T_0^2}{2}$$

we can deduce that

$$\sum A_{n+1}(t)$$

is totally convergent. This implies that there exists  $u_\infty$  such that:

$$u_n(x, t) \rightrightarrows u_\infty(x, t), \quad x \in \mathbb{R}, t \in [0, T_0].$$

Since

$$\begin{aligned} & \left| u_n \left( \frac{\partial^2}{\partial t^2} u_n(x, t) + k_2 u_n(x, t), t \right) - u_\infty(v_\infty(x, t) + k_2 u_\infty(x, t), t) \right| \\ & \leq \|u_n - u_\infty\|_\infty + L_\infty(t) \left( \|u_n - u_\infty\|_\infty + \left\| \frac{\partial^2}{\partial t^2} u_n - v_\infty \right\|_\infty \right) \rightarrow 0, \end{aligned}$$

we can deduce that:

$$u_\infty(x, t) = \alpha(x) + t\beta(x) + \int_0^t k_1 \int_0^\tau u_\infty(v_\infty(x, s) + k_2 u_\infty(x, s), s) ds.$$

Then the conclusion follows if we prove that  $v_\infty = \partial^2 u / \partial t^2$ ; but we since

$$\begin{aligned} \left| v_\infty(x, t) - \frac{\partial^2}{\partial t^2} u_\infty(x, t) \right| & \leq \left\| v_\infty - \frac{\partial^2}{\partial t^2} u_n \right\|_\infty + \\ & \quad + \left| \frac{\partial^2}{\partial t^2} u_n(x, t) + \frac{\partial^2}{\partial t^2} u_\infty(x, t) \right| + \\ & \leq \left\| v_\infty - \frac{\partial^2}{\partial t^2} u_n \right\|_\infty + \\ & \quad + \left| u_{n-1} \left( \frac{\partial^2}{\partial t^2} u_{n-1}(x, t) + k_2 u_{n-1}(x, t), t \right) + \right. \\ & \quad \left. + u_\infty(v_\infty(x, t) + k_2 u_\infty(x, t), t) \right| \leq \\ & \leq \left\| v_\infty - \frac{\partial^2}{\partial t^2} u_{n-1} \right\|_\infty + \|u_{n-1} - u_\infty\|_\infty + \\ & \quad + L_\infty(t) \left( \|u_{n-1} - u_\infty\|_\infty + \left\| \frac{\partial^2}{\partial t^2} u_{n-1} - v_\infty \right\|_\infty \right) \\ & \rightarrow 0, \end{aligned}$$

we can deduce that:

$$u_\infty(x, t) = \alpha(x) + t\beta(x) + \int_0^t \int_0^\tau k_1 u_\infty \left( \frac{\partial^2}{\partial s^2} u_\infty(x, s) + k_2 u_\infty(x, s), s \right) ds d\tau,$$

with  $x \in \mathbb{R}$ ,  $t \in [0, T_0]$ , and then  $u_\infty$  is a solution of (2.3). We prove now that (2.3) has a unique solution; let us then assume that there exists an other solution  $v = v(x, t)$ ;

$$\begin{aligned} |v(x, t) - u_\infty(x, t)| &\leq \int_0^t \int_0^\tau \left| v \left( \frac{\partial^2}{\partial s^2} v(x, s) + k_2 v(x, s), s \right) + \right. \\ &\quad \left. + u_\infty \left( \frac{\partial^2}{\partial s^2} v(x, s) + k_2 v(x, s), s \right) + \right. \\ &\quad \left. + u_\infty \left( \frac{\partial^2}{\partial s^2} v(x, s) + k_2 v(x, s), s \right) + \right. \\ &\quad \left. - u_\infty \left( \frac{\partial^2}{\partial s^2} u_\infty(x, s) + k_2 u_\infty(x, s), s \right) \right| ds d\tau. \end{aligned}$$

Hence:

$$(2.6) \quad \begin{aligned} |v(x, t) - u_\infty(x, t)| &\leq \|v - u_\infty\|_\infty \int_0^t \int_0^\tau (1 + L_\infty(s)) ds d\tau + \\ &\quad + \int_0^t \int_0^\tau L_\infty(s) \left| \frac{\partial^2}{\partial s^2} u_\infty(x, s) - \frac{\partial^2}{\partial s^2} v(x, s) \right| ds d\tau. \end{aligned}$$

But:

$$\begin{aligned} \left| \frac{\partial^2}{\partial t^2} v(x, t) - \frac{\partial^2}{\partial t^2} u_\infty(x, t) \right| &\leq \left| v \left( \frac{\partial^2}{\partial t^2} v(x, t) + k_2 v(x, t), t \right) + \right. \\ &\quad \left. - u_\infty \left( \frac{\partial^2}{\partial t^2} v(x, t) + k_2 v(x, t), t \right) + \right. \\ &\quad \left. + u_\infty \left( \frac{\partial^2}{\partial t^2} v(x, t) + k_2 v(x, t), t \right) + \right. \\ &\quad \left. - u_\infty \left( \frac{\partial^2}{\partial t^2} u_\infty(x, t) + k_2 u_\infty(x, t), t \right) \right| \leq \\ &\leq \|v - u_\infty\|_\infty + L_\infty \left( \|v - u_\infty\|_\infty + \right. \\ &\quad \left. + \left\| \frac{\partial^2}{\partial t^2} v - \frac{\partial^2}{\partial t^2} u_\infty \right\|_\infty \right) \\ &\leq (M + 1) \|v - u_\infty\|_\infty + \\ &\quad + M \left\| \frac{\partial^2}{\partial s^2} v - \frac{\partial^2}{\partial s^2} u_\infty \right\|_\infty. \end{aligned}$$

This implies that:

$$\left\| \frac{\partial^2}{\partial t^2} v - \frac{\partial^2}{\partial t^2} u_\infty \right\|_\infty (1 - M) \leq (1 + M) \|v - u_\infty\|_\infty.$$

Then

$$\left\| \frac{\partial^2}{\partial t^2} v - \frac{\partial^2}{\partial t^2} u_\infty \right\|_\infty \leq \frac{1 + M}{1 - M} \|v - u_\infty\|_\infty.$$

From (2.6) we obtain that, for every  $t \in [0, T_0]$ :

$$\begin{aligned} |v(x, t) - u_\infty(x, t)| &\leq \|v - u_\infty\|_\infty \left( (1 + M) \frac{t^2}{2} \right) + \frac{1 + M}{1 - M} \frac{t^2}{2} M \|v - u_\infty\|_\infty = \\ &= \|v - u_\infty\|_\infty \left( \frac{t^2}{2} \left( 1 + M + \frac{1 + M}{1 - M} M \right) \right) = \\ &= \|v - u_\infty\|_\infty \frac{t^2}{2} (1 + M) \frac{1 - M + M}{1 - M} = \\ &= \|v - u_\infty\|_\infty \frac{t^2}{2} \frac{1 + M}{1 - M} < \|v - u_\infty\|_\infty, \end{aligned}$$

whence the uniqueness.

In this way we have proved the following local existence and uniqueness Theorem.

**THEOREM 1.** *Let  $\alpha, \beta \in L^\infty(\mathbb{R}, \mathbb{R}) \cap Lip(\mathbb{R}, \mathbb{R})$  be given functions with Lipschitz constant  $L_\alpha$  and  $L_\beta$  respectively and such that  $L_\alpha < 1/2$ ; then there exists  $T_0 > 0$  and there exists a unique  $u : \mathbb{R} \times [0, T_0] \rightarrow \mathbb{R}$  bounded, Lipschitz in the  $x$  variable, uniformly with respect to  $t$  (and Lipschitz in  $t$  uniformly with respect to  $x$  and continuous in both variables) with second derivative in  $t$  continuous and such that*

$$u(x, t) = \alpha(x) + t\beta(x) + \int_0^t \int_0^\tau k_1 u \left( \frac{\partial^2}{\partial s^2} u(x, s) + k_2 u(x, s), s \right) ds d\tau$$

for all  $x \in \mathbb{R}$ ,  $t \in [0, T_0]$  and then

$$\begin{cases} \frac{\partial^2}{\partial t^2} u(x, t) = k_1 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_2 u(x, t), t \right) \\ u(x, 0) = \alpha(x) \\ \frac{\partial}{\partial t} u(x, 0) = \beta(x). \end{cases}$$



## REMARK 2

1. From the previous proof it is clear that the condition  $|k_1| = |k_2| = 1$  is not a restrictively hypothesis; in fact we can also assume that  $k_i = k_i(x, t)$ , being careful to insure that  $|k_i(x, t)| \leq 1 \quad \forall x \in R, t \geq 0$ .
2. It is easy to notice that similar theorems of existence and uniqueness for small time can be established also for equations of the type:

$$\frac{\partial^2}{\partial t^2} u(x, t) = k_1 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_2 u \left( \frac{\partial^2}{\partial t^2} u(x, t) + k_3 u(x, t), t \right), t \right),$$

that we can denote by  $E_{(k_1, k_2, k_3)}$ . For these equations the existence time for the solution established using the proof of the Theorem is sensibly reduced.

In view of the previous remarks and from the boundedness of the approximating sequences used in the proof of the Theorem, we can state the following problems.

- A) Find some condition assuring that the solution of  $E_{(k_1, k_2, \dots, k_i)}$  exists for all time.
- B) An interesting problem is to give existence result for the equation when  $\alpha$  and  $\beta$  are continuous but not Lipschitz functions.
- C) Given a sequence  $(k_i)$  of real numbers with  $|k_i| \leq 1 \quad \forall i$  and assuming that each equation  $E_{(k_1, k_2, \dots, k_n)}$  admits a unique solution  $v_n = v_n(x, t)$ , defined for every time and satisfying the initial condition  $v_n(x, 0) = \alpha(x)$ ,  $\frac{\partial}{\partial t} v_n(x, 0) = \beta(x)$ , being  $\alpha$  and  $\beta$  given, study the behaviour of the sequence  $(v_n)$ .

## REFERENCES

- [1] E. EDER.: *The functional-differential equation  $x'(t) = x(x(t))$* , J. Differential Equations, **54**(3) (1984), 390-400.
- [2] G. FICHERA: *Having a long memory creates serious problems*, Arch. Rational Mech. Anal., **70**(2) (1979), 101-112.
- [3] G. FICHERA: *Analytic problems of materials with memory*, In: "Theoretical and applied mechanics" (Proc. 15th Internat. Congr., Univ. Toronto, Toronto, Ont., 1980), 1980, pp. 223-230, North-Holland, Amsterdam.
- [4] M. JR. MIRANDA – E. PASCALI: *On a type of evolution of self-referred and hereditary phenomena*, Submitted to.
- [5] M. JR. MIRANDA – E. PASCALI: *Other classes of self-referred equations*, Preprint n.15/2004, Dipartimento di Matematica, Università di Lecce.
- [6] JIAN-GUO SI – SUI SUN CHENG: *Analytic solutions of a functional-differential equation with state dependent argument*, Taiwanese J. Math., **1** (4) (1997), 471-480.

- [7] JIAN-GUO SI– XIN-PING WANG – SUI SUN CHENG: *Analytic solutions of a functional-differential equation with a state derivative dependent delay*, *Aequationes Math.*, **57**(1)(1999), 75-86.
- [8] V. VOLTERRA: *Opere matematiche: Memorie e note*, Vol. V 1926-1940. Pubblicate a cura dell'Accademia Nazionale dei Lincei col concorso del Consiglio Nazionale delle Ricerche. Accademia Nazionale dei Lincei, Rome, 1962.

*Lavoro pervenuto alla redazione il 17 settembre 2004  
ed accettato per la pubblicazione il 17 gennaio 2005.  
Bozze licenziate il ???*

INDIRIZZO DELL'AUTORE:

Michele Miranda Jr.– Eduardo Pascali – Dipartimento di Matematica “Ennio De Giorgi” –  
Università di Lecce – C.P. 193, 73100 Lecce – Italy