

# SPHERICALIZATION AND FLATTENING

ZOLTÁN M. BALOGH AND STEPHEN M. BUCKLEY

ABSTRACT. The conformal deformations of flattening and sphericalization of length metric spaces are considered. These deformations are dual to each other if the space satisfies a simple quantitative connectivity property. Moreover, the quasihyperbolic metrics corresponding to the flat and the spherical metrics are bilipschitz equivalent.

## 0. INTRODUCTION

Recent progress in geometric function theory: [AG], [AK], [Se1], [Se2], led to the study of conformal deformations of domains in Euclidean space. Elaborate results on quite general deformations defined on the unit ball have been developed in several recent papers, notably [BKR], [BK], [BHR]. The extension of the theory of quasiconformal mappings from Euclidean to general metric spaces [HK],[BKo], [Ty1], [Ty2] motivated the study of conformal deformations in general metric spaces [BHK], [BB], [He1], [He2].

One of the most important conformal deformations is the so-called *quasihyperbolic metric*  $k$  which, on a domain  $\Omega \subsetneq \mathbb{R}^n$ , is defined by the equation

$$k_{\Omega}(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_{\gamma} \frac{dl(z)}{d(x, \partial\Omega)}.$$

where  $\Gamma(x, y)$  is the class of rectifiable paths  $\lambda : [0, T] \rightarrow X$  for which  $\lambda(0) = x$  and  $\lambda(T) = y$ ,  $dl$  is the Euclidean length element, and  $d(x, \partial\Omega)$  is Euclidean distance to the boundary. This metric was extensively studied in [BHK], [He2], [BB]. Notice however that if we instead use the spherical length element and spherical boundary distance, we get an alternate quasihyperbolic metric  $k'_{\Omega}$ . Both  $k_{\Omega}$  and  $k'_{\Omega}$  are used frequently in the literature, but how do they compare? This question arises naturally from the results of [BHK] and [BB] which were proven for the spherical quasihyperbolic metric  $k'_{\Omega}$ ; but in many instances one would like to know what happens for the case of the usual quasihyperbolic metric  $k_{\Omega}$ .

Notice first that  $k_{\Omega}(x, y)$  can be much smaller than  $k'_{\Omega}(x, y)$ . For instance, if  $\Omega$  is a ball of Euclidean radius  $r > 2$  centered at the origin,  $x$  is the origin, and  $y$  is a point on the Euclidean unit circle then, as  $r \rightarrow \infty$ ,  $k_{\Omega}(x, y)$  tends to 0 but  $k'_{\Omega}(x, y)$  is bounded away from 0. Despite this example, we shall prove the following result which is essentially a special case of one of the main results of this paper.

---

*Date:* 14.10.2004

This research was partially supported by the Swiss Nationalfond and Enterprise Ireland. It was partly conducted during a visit by the second author to the University of Bern; the hospitality of the Mathematics Department was much appreciated.

**Theorem 0.1.** *Given  $x_0 \in \mathbb{R}^n$  and  $r_0 > 0$ , let  $\mathcal{C}$  be the class of all domains  $\Omega \subsetneq \mathbb{R}^n$  containing  $x_0$  for which  $\text{dist}(x_0, \partial\Omega) \leq r_0$ . For  $\Omega \in \mathcal{C}$  the quasihyperbolic metrics  $k_\Omega, k'_\Omega$  defined above are bilipschitz equivalent to each other, with a bilipschitz constant dependent only on  $|x_0|$  and  $r_0$ .*

In order to obtain this statement we have to consider carefully the relation between the spherical and the Euclidean metric. In fact we examine the relation between two classes of conformal deformations in a general metric space setting. The first class of deformations, which we call *sphericalization*, generalizes the process of obtaining the Riemann sphere from Euclidean space. If we sphericalize an unbounded length space, we get an incomplete bounded length space. The second class of deformations, which we call *flattening*, generalizes the process of obtaining Euclidean space from the Riemann sphere. If we flatten an incomplete bounded length space, we get an unbounded length space. In the process of flattening, a selected part of the spherical boundary is pushed out to infinity, while sphericalization brings back the boundary at infinity to a bounded distance. These two procedures are dual to each other in a similar way as the procedures of quasihyperbolization and dampening from [BHK]. We show in a general metric setting that a necessary and sufficient condition for the quasihyperbolic metric associated with the original metric to be bilipschitz equivalent to the quasihyperbolic metric associated with the sphericalized metric (or flattened metric) is a surprisingly simple geometric property of the original space which we call an *escape property* (or *reverse escape property*). Since all Euclidean domains have such properties, Theorem 0.1 will follow almost immediately.

As already indicated, the main motivation for this investigation lies in the results of [BHK] and [BB]. Those papers investigate the question of when the quasihyperbolic metric associated with a general length metric is Gromov hyperbolic. The assumption of boundedness is essential for some of the key results in those papers. Since one is often interested in quasihyperbolic metrics on unbounded Euclidean domains and other unbounded metric spaces, this analysis at first seems to be of limited use in that context. However on a large class of unbounded spaces (including all Euclidean domains), the quasihyperbolic metric on the original (unbounded) space is Gromov hyperbolic if and only if the quasihyperbolic metric on the sphericalized (bounded) space is Gromov hyperbolic, allowing us in theory to apply the results of those previous papers to unbounded spaces. Since sphericalization is a fairly simple conformal distortion, it is often easy to get a rather explicit description of the sphericalized space in terms of the original space, so this method of appealing to the results of [BHK] and [BB] is in practice quite feasible.

After defining our main notation in Section 1, we introduce the class of conformal deformations called sphericalization and flattening and discuss the procedures of sphericalization in Section 2, and flattening in Section 3. We obtain in conclusion the comparability of the spherical and the flat quasihyperbolic metrics. In Section 4, we examine to what extent sphericalization and flattening are inverse processes. We introduce the notion of dual deformations in the context of sphericalization-flattening. We show that if we flatten a sphericalized space (or sphericalize a flattened space) using “dual” sphericalizing and flattening functions, then we get a space bilipschitz equivalent to the original one if and only if the original space satisfies a stronger variant of the escape property (or reverse escape property). This makes it easy to give examples of incomplete bounded spaces that are not produced by sphericalization, and unbounded spaces that are not produced by flattening.

1. NOTATION AND TERMINOLOGY

We write  $A \lesssim B$  or  $B \gtrsim A$  whenever  $A \leq CB$  for some positive constant  $C$  that depends only on explicitly allowed parameters, and we write  $A \approx B$  if  $A \lesssim B \lesssim A$ . We denote by  $a \vee b$  and  $a \wedge b$  the maximum and minimum, respectively, of any  $a, b \in [-\infty, \infty]$ .

Suppose that  $(X, d)$  is a metric space. Given  $x \in X$  and  $r \in \mathbb{R}$ , we define  $B_d(x, r) = \{y \in X : d(y, x) < r\}$ . If  $\gamma$  is a path in  $(X, d)$ , then  $\gamma(T-; d)$  is short-hand for  $\lim_{t \rightarrow T-} \gamma(t)$ , assuming this  $d$ -limit exists in  $\overline{X}_d$ , and  $\text{len}_d(\gamma)$  is the  $d$ -length of  $\gamma$ . We do not distinguish notationally between paths and their images.

We denote by  $\overline{X}_d$  the metric completion of  $X$  and  $\partial X_d = \overline{X}_d \setminus X$ . We say that  $(X, d)$  is *locally complete* if  $d(x, \partial X_d) > 0$  for all  $x \in X$ . Given  $x, y \in \overline{X}_d$ ,  $\Gamma(x, y)$  denotes the class of rectifiable paths  $\gamma : [0, T] \rightarrow \overline{X}_d$  for which  $\gamma|_{(0, T)}$  is a rectifiable path in  $X$ ,  $\gamma(0) = x$ , and  $\gamma(T) = y$ . We also define  $\Gamma_d(x, y)$  to be the subset of  $\Gamma(x, y)$  consisting of paths that are parametrized by  $d$ -arclength.

Suppose  $(X, d, o)$  is a rectifiably connected pointed metric space which is locally complete, and either incomplete or unbounded. Writing  $\delta_{d, C}(x) = d(x, \partial X_d) \wedge (C + d(x, o))$  for  $0 < C \leq \infty$ , we define the  $C$ -*quasihyperbolic metric* on  $X$  by the equation

$$k_{d, C}(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_{\gamma} \frac{dl(z)}{\delta_{d, C}(z)}. \tag{1.1}$$

Whenever we omit the subscript  $C$  in this notation, it is to be assumed that  $C = \infty$ . Thus  $\delta_d$  and  $k_d$  are the boundary distance and quasihyperbolic metric, respectively, as they are usually defined for incomplete spaces but, taking  $C < \infty$ , we get variants of these concepts that are also defined on complete spaces. Note that if  $X$  is incomplete then  $\delta_d = \delta_{d, C}$  and  $k_d = k_{d, C}$ , for all  $C \geq d(o, \partial X_d)$ . We shall see that the resulting metrics  $k_{d, C}$  behave in a similar manner whether  $C$  is finite or infinite. In this paper, we compare two such quasihyperbolic metrics,  $k_{l, C}$  and  $k_{\sigma}$ , where  $l$  is an unbounded metric,  $\sigma$  is typically a bounded incomplete metric, and  $l, \sigma$  are related via sphericalization or flattening. Note that local completeness is essential to ensure that the right-hand side of (1.1) is finite.

2. SPHERICALIZATION

In this section we “sphericalize” a fairly general unbounded metric space  $(X, l)$  to get a related length space  $(X, \sigma)$ ; we use the term “sphericalization” because this process generalizes that of obtaining the Riemann sphere from Euclidean space. The main result, Theorem 2.28, compares the corresponding quasihyperbolic metrics  $k_{l, C}$  and  $k_{\sigma}$ .

A couple of results require that  $l$  is a length metric but, since we want our main result to apply to Euclidean domains with the Euclidean metric attached (in order to deduce Theorem 0.1), we mostly require  $l$  to be merely a local length metric, a concept we now define.

Suppose  $(X, l)$  is a metric space and let  $l'(x, y) = \inf_{\gamma \in \Gamma_l(x, y)} \text{len}_l(\gamma)$ ,  $x, y \in \overline{X}_l$ . If  $l'|_{X \times X}$  is always finite, we call it the *inner metric* associated with  $l$ . A rectifiably connected metric space  $(X, l)$  is a *local length space*, and  $l$  is a *local length metric*, if  $l(x, y) = l'(x, y)$  whenever  $x \in X$ ,  $y \in \overline{X}_l$ , and  $l(x, y) \leq l(x, \partial X_l)$ . Recall that  $(X, l)$  is a *length space* if  $l(x, y) = l'(x, y)$

for all  $x, y \in X$ ; such an equality clearly extends to points  $x, y \in \overline{X}_l$ , so a length space is always a local length space.

Every Euclidean domain is a local length space when equipped with the Euclidean metric, but the slit disk is not a length space. To make a Euclidean domain into a length space, we can equip it with either the *inner Euclidean* or *inner spherical metric*, i.e. the inner metrics associated with either the Euclidean metric or the intrinsic metric of the Riemann sphere.

Throughout the rest of this section,  $(X, l, o)$  is an unbounded pointed local length space, and we write  $|x| = l(x, o)$ ,  $x \in \overline{X}_l$ . Given  $g : [0, \infty) \rightarrow (0, \infty)$ , we define a length metric  $\mathfrak{S}(l, o, g)$  on  $X$  by the equation

$$\mathfrak{S}(l, o, g)(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_{\gamma} g(|z|) dl(z), \quad x, y \in X.$$

We normally write  $\sigma$  in place of  $\mathfrak{S}(l, o, g)$ , and we take  $g$  to be any *sphericalizing function*, a concept we now define.

### 2.1. Sphericalizing functions.

A continuous function  $g : [0, \infty) \rightarrow (0, \infty)$  is a  $C$ -*sphericalizing function*,  $C > 2$ , if it has the following properties:

- (S1)  $g(r) \leq Cg(s)$  whenever  $r, s > 0$ ,  $r \leq 2s + 1$ , and  $s \leq 2r + 1$ ;
- (S2)  $\int_r^\infty g(t) dt \leq Crg(r)$ ,  $r \geq 1$ .

We shall see that this pair of properties is quantitatively equivalent to the combination of (S1) and the following condition, where  $0 < \epsilon \leq 1 \leq C'$  are other constants:

- (S3)  $\frac{g(s)}{g(r)} \leq C' \left(\frac{r}{s}\right)^{1+\epsilon}$ , for all  $1 \leq r \leq s$ .

Notice that (S1) merely says that all values of  $g$  on any one of the intervals  $I_j$  are mutually comparable, where  $I_0 = [0, 1]$  and  $I_j = [2^{j-1}, 2^j]$ ,  $j \in \mathbb{N}$ .

Typical sphericalizing functions include the “standard sphericalizing function”  $g(t) = 2/(1+t^2)$  and, more generally,  $g(t) = c \log^q(2+t)/(1+t^p)$ , where  $p > 1$ ,  $q \in \mathbb{R}$ , and  $c > 0$ . Note that (S1) and (S3) imply that  $g$  is quasidecreasing in the sense that

- (S4)  $g(s) \leq C''g(r)$ ,  $0 \leq r \leq s$ ,

where  $C'' = CC'$ . Moreover  $h(r) := \sup_{r \geq s} g(s)$  defines a monotonically decreasing  $C_1$ -sphericalizing function comparable to  $g$ , where  $C_1 = C^2C'$ . Thus we may assume without loss of generality that a sphericalizing function is decreasing whenever switching to a comparable sphericalizing function and making a quantitatively controlled change in  $C$  is harmless.

We now state and prove two simple lemmas.

**Lemma 2.2.** *A continuous function  $g : [0, \infty) \rightarrow (0, \infty)$  is a sphericalizing function if and only if it satisfies (S1) and (S3), with quantitative control of parameters.*

*Proof.* By integration, (S3) immediately implies (S2), with  $C = C'/\epsilon$ . Conversely suppose (S1) and (S2) hold. Since  $g(t) \geq g(r)/C$  for  $r \leq t \leq 2r$ , we get the useful estimate

$$\frac{rg(r)}{C} \leq \int_r^\infty g(t) dt \leq Crg(r), \quad r \geq 1. \quad (2.3)$$

Writing  $u(r) := r^{1/C} \int_r^\infty g(t) dt$ , we see that for all  $r > 1$ ,

$$u'(r) = \frac{r^{-1+1/C}}{C} \left( \int_r^\infty g(t) dt - Crg(r) \right) \leq 0,$$

Thus  $u$  is a decreasing function and

$$\int_s^\infty g(t) dt \leq \left( \frac{r}{s} \right)^{1/C} \int_r^\infty g(t) dt, \quad 1 \leq r \leq s.$$

This inequality plus (2.3) gives (S3) with  $C' = C^2$  and  $\epsilon = 1/C$ .  $\square$

**Lemma 2.4.** *Suppose  $g$  is a  $C$ -sphericalizing function,  $C > 2$ . Then for every  $K > 0$  there exists  $\alpha = \alpha(C, K) \in (0, 1)$  such that*

$$\int_{\alpha r}^r g(t) dt \geq Krg(r), \quad r \geq 1/\alpha.$$

*Proof.* Replacing  $g(t)$  by the lower bound  $g(r)(r/t)^{1+\epsilon}/C'$  provided by Lemma 2.2, and integrating, we get

$$\int_{\alpha r}^r g(t) dt \geq rg(r) \cdot \left( \frac{\alpha^{-\epsilon} - 1}{\epsilon C'} \right).$$

Taking  $\alpha > 0$  small enough, the parenthesized expression can be made arbitrarily large.  $\square$

We record the following change of variables lemma, which is essentially a special case of [V, Theorem 5.7].

**Lemma 2.5.** *Let  $I \subset [0, \infty)$  be an interval. If  $\alpha : [a, b] \rightarrow I$  satisfies the 1-Lipschitz condition  $|\alpha(s) - \alpha(t)| \leq |s - t|$ ,  $s, t \in [a, b]$ , and  $\rho : I \rightarrow (0, \infty)$  is a Borel function, then*

$$\int_a^b \rho(\alpha(t)) dt \geq \left| \int_{\alpha(a)}^{\alpha(b)} \rho(s) ds \right|.$$

We now come to the first of the three results in this section that require  $l$  to be a length metric, and not just a local length metric.

**Proposition 2.6.** *If  $(X, l, o)$  is an unbounded pointed length space, and  $\sigma = \mathfrak{S}(l, o, g)$  for some  $C$ -sphericalizing function  $g$ , then  $\text{dia}_\sigma(X) \leq 4Cg(1)$ .*

*Proof.* Let  $\gamma_\epsilon \in \Gamma_l(o, y)$  be a path of  $l$ -length  $L \leq |y| + \epsilon$  for some  $\epsilon \in (0, 1]$ , and so

$$\text{len}_\sigma(\gamma) = \int_0^L g(|\gamma_\epsilon(t)|) dt.$$

By the triangle inequality,  $t - \epsilon \leq |\gamma(t)| \leq t$ . Letting  $\epsilon$  tend to zero, and using the uniform continuity of  $g$  on  $[0, |y| + 1]$ , we deduce that  $\text{len}_\sigma(\gamma_\epsilon)$  tends to  $\int_0^{|y|} g(t) dt$ . Using (S1) and (S2), the last integral is bounded by  $2Cg(1)$ .  $\square$

*Remark 2.7.* The above lemma would fail if we merely required  $l$  to be a local length metric. For instance, if  $l$  is the Euclidean metric on a domain  $\Omega \subset \mathbb{R}^n$  then, using the standard sphericalizing function,  $\sigma$  is the usual inner spherical metric, and this can be unbounded if  $\Omega$  tightly spirals in the vicinity of some point  $z \in \partial\Omega$ . If we instead sphericalized  $\Omega$  with respect to the inner Euclidean metric, we would always get a bounded space, although this space is typically not a subdomain of the Riemann sphere.

*Remark 2.8.* If  $\gamma$  is a path in  $X$  containing points  $x, y$  with  $|x| \leq |y|$ , then

$$\text{len}_\sigma(\gamma) \geq \int_{|x|}^{|y|} g(t) dt. \quad (2.9)$$

This follows from Lemma 2.5 with  $\rho = g$  and  $\alpha(t) = |\nu(t)|$ , where  $\nu$  is the reparametrization of  $\gamma$  by arclength. Since we already know from the proof of Proposition 2.6 that  $\sigma(o, y) \leq \int_0^{|y|} g(t) dt$ , it follows that  $\sigma(o, y) = \int_0^{|y|} g(t) dt$ , assuming  $l$  is a length metric.

## 2.10. Escape properties.

**Definition 2.11.** Suppose  $(X, l, o)$  is a pointed local length space,  $0 < c \leq 1$ , and  $\rho \geq 0$ . We say that  $\gamma$  is a  $c$ -escape path for  $x \in X$  if  $\gamma : [0, T) \rightarrow X$  is parametrized by  $l$ -arclength,  $\gamma(0) = x$ ,  $|\gamma(t)| \geq c(t + |x|)$ , and either  $T = \infty$ , or  $T < \infty$  with  $\gamma(T-; l) = y \in \partial X_l$ ; recall that  $\gamma(T-; l)$  indicates the  $l$ -limit  $\lim_{t \rightarrow T-} \gamma(t)$ . We call  $\gamma$  a *strong  $c$ -escape path* if  $T = \infty$ . We say that  $(X, l, o)$  has the (strong)  $(c, \rho)$ -escape property if every  $x \in X$  satisfying  $|x| \geq \rho$  has a (strong)  $c$ -escape path.

Trivially, the strong  $(c, \rho)$ -escape property implies the  $(c, \rho)$ -escape property. By the triangle inequality, a  $c$ -escape path for  $x$  must satisfy  $|\gamma(t)|/(t + |x|) \in [c, 1]$ . This implies the following fact.

**Fact 2.12.** *If  $g$  is a  $C_0$ -sphericalization function, and  $\gamma : [0, T) \rightarrow X$  is a  $c$ -escape path for a point  $x \in X$ , then there exists  $C = C(C_0, c) \geq 1$  such that  $g(|\gamma(t)|)/g(t + |x|) \in [1/C, C]$ ,  $0 \leq t < T$ .*

Modulo a change in the values of  $c$  and  $\rho$ , (strong) escape properties are independent of the basepoint  $o$ . Moreover the next lemma shows that if any point in a length space has a strong escape path, so does  $o$ .

**Lemma 2.13.** *Let  $(X, l, o)$  be an unbounded pointed length space, and suppose that some point  $x \in X$  has a strong  $c$ -escape path for some  $0 < c < 1$ . Then  $o$  also has a strong  $c$ -escape path.*

*Proof.* We claim that there is a path  $\nu$  from  $o$  to  $x_0$ , parametrized by  $l$ -arclength and satisfying  $|\nu(t)| \geq ct$ . Once we prove this claim, the lemma follows by concatenating  $\nu$  with the strong escape path for  $x_0$ . To prove the claim, we construct  $\nu$  inductively. Letting  $\delta = (1 - c)/4$ ,

we first pick  $\lambda_1$  from  $x_0$  to  $o$  of length less than  $(1 + \delta)|x_0|$ . Let  $\nu_1$  be the initial segment of  $\lambda_1$  as far as  $x_1 = \lambda_1(t_1)$ , the first point on  $\lambda_1$  whose distance from  $o$  is  $|x_0|/2$ . Inductively, given  $x_j$ , we find a path  $\lambda_{j+1}$  from  $x_j$  to  $o$  of length less than  $(1 + \delta)|x_j|$  and let  $\nu_{j+1}$  be the initial segment of  $\lambda_{j+1}$  as far as  $x_{j+1} = \lambda_{j+1}(t_{j+1})$ , the first point on  $\lambda_{j+1}$  whose distance from  $o$  is  $|x_j|/2$ . Concatenating the paths  $\nu_j$ ,  $j \in \mathbb{N}$ , and reparametrizing the resulting path by  $l$ -arclength in the direction from  $o$  to  $x_0$ , it is clear that the resulting path  $\nu$  has the desired property.  $\square$

**Example 2.14.** Suppose that  $\Omega \subset \mathbb{R}^n$  is a Euclidean domain containing the origin. Let  $o$  be the origin and attach the inner Euclidean metric  $l$ . Then  $\Omega$  has the  $(1, 0)$ -escape property, since if  $x \in \Omega$ ,  $x \neq 0$ , we can take the path  $\gamma : [0, T) \rightarrow \Omega$ ,  $\gamma(t) = x + t \cdot x$ , with  $T$  being the least value of  $t > 0$  for which  $x + t \cdot x \in \partial_l \Omega$  (or  $T = \infty$  if all there is no such number  $t$ ). If  $x = 0$ , take any ray beginning at the origin, and let  $\gamma$  be the initial part which ends (if at all) when this ray first reaches  $\partial_l \Omega$ . It is, however, easy to give an example of such a domain where  $o$  (and hence every point) fails to have any strong escape paths, for instance the complement of an Archimidean spiral  $S \not\ni o$ .

**Example 2.15.** There are also domains that in which there are strong escape paths from all points but which fail to have a strong escape property because the escape parameter  $C$  depends on the point. Consider, for instance, the domain whose complement consists of the two curves given by  $t \mapsto (t, 1 + t^2 \sin t)$ ,  $t \geq 0$ , and  $t \mapsto (t, 3 + t^2 \sin t)$ ,  $t \geq 0$ . There is a strong escape path from every point  $x \in \Omega$ , but the strong escape parameter  $c$  depends on  $x$ . For instance, if we write

$$\Omega' = \{(x_1, x_2) : x_1 \geq 0, |x_2 - 2 - x_1^2 \sin x_1| < 1\} \subset \Omega,$$

then for points in  $(x_1, x_2) \in \Omega'$ , the parameter  $c$  must be chosen to be very small if  $x_1$  is very large. Thus  $\Omega$  fails to have any strong escape property. By contrast, there is a strong 1-escape path from all points  $(x_1, x_2)$  with  $x_1 < 0$ .

**Example 2.16.** Consider  $X \subset \mathbb{R}^2$ , where  $X$  consists of the horizontal half-line  $(x_1, 0)$ ,  $x_1 \geq 0$ , together with the attached vertical line segments  $(a_n, x_2)$ ,  $0 \leq x_2 \leq b_n$ ,  $n \in \mathbb{N}$ , where  $(a_n)$  is a strictly increasing sequence of positive numbers tending to infinity, and  $(b_n)$  is a non-negative sequence of numbers. Let  $o = (0, 0)$  and equip  $X$  with its natural Euclidean length metric  $l$ . Then it is readily verified that  $(X, l, o)$  has the  $(c, \rho)$ -escape property for a particular  $0 < c \leq 1$  if and only if  $b_n \leq (1 - c)a_n/2c$  whenever  $a_n + b_n \geq \rho$ . In particular,  $(X, l, o)$  has some escape property if and only if  $(b_n/a_n)_{n=1}^\infty$  is bounded. The same condition also characterizes when  $(X, l, o)$  has a strong escape property.

**Example 2.17.** Suppose  $(X, l, o)$  is as in the previous example, with  $b_n > 0$  for all  $n \in \mathbb{N}$ . Given  $T \subset \mathbb{N}$ , we define  $X_T$  to be the subset of  $X$  consisting of all points except  $(a_n, b_n)$ ,  $n \in T$ , and let us also denote by  $l$  the subspace metric inherited by  $X_T$ . Then  $(X_T, l, o)$  is always a length space, and it has an escape property if and only if the set  $\{b_n/a_n : n \in \mathbb{N} \setminus T\}$  is bounded. In the extreme case  $T = \mathbb{N}$ ,  $(X_{\mathbb{N}}, l, o)$  always satisfies the  $(1, 0)$ -escape property. Notice however that strong escape properties for spaces of the form  $(X_T, l, o)$  are more restrictive than escape properties whenever  $T$  is nonempty: although there always exists a strong escape path for  $o$  and all other points  $(x_1, 0) \in X_T$ , nevertheless  $(X_T, l, o)$  fails to satisfy a strong escape property unless the full sequence  $(b_n/a_n)_{n=1}^\infty$  is bounded.

**Example 2.18.** Let  $X$  consist of a bouquet of half-open line segments  $I_n$ , where  $I_n$  has length  $n$ . Here, we identify the included endpoints of all the intervals and call this point  $o$ . There is a natural length metric  $l$  on  $X$  which restricts to the length metric on each interval  $I_n$ . Then  $(X, l, o)$  satisfies a  $(1, 0)$ -escape property, but there is no strong escape path from any point of  $X$ .

### 2.19. The spherical boundary.

Given a local length space  $(X, l)$ , let  $\partial_0 X_l$  denote the set of all points  $y \in \partial X_l$  for which  $\Gamma(x, y)$  is nonempty for some (and hence all)  $x \in X$ . Equivalently  $\partial_0 X_l$  is the set of all  $y \in \overline{X}_l$  whose inner distance from some (and hence all)  $x \in X$  is finite. If  $l$  is a length metric, then  $\partial_0 X_l = \partial X_l$ , but equality may fail if  $l$  is merely a local length space. For instance, if  $l$  is the Euclidean metric on a domain  $\Omega \subset \mathbb{R}^n$  which spirals sufficiently tightly near some point  $y \in \partial\Omega$ , then  $\Gamma(x, y)$  is empty for all  $x \in \Omega$ . However,  $\Gamma(x, y)$  is always nonempty if there exists  $z \in X$  such that  $d(z, y) = \text{dist}_d(z, \partial\Omega)$ .

Suppose  $\sigma = \mathfrak{S}(l, o, g)$ , where  $g$  is a sphericalizing function. Given  $x \in X$ ,  $y \in \partial_0 X_l$ , and  $\gamma \in \Gamma_l(x, y)$ , it is clear that  $\gamma$  is also of finite  $\sigma$ -length. Thus  $\partial_0 X_l$  can be viewed in a natural way as a subset of  $\partial X_\sigma$ . We define the *spherical boundary of  $X$*  to be  $\partial_S X := \partial X_\sigma \setminus \partial_0 X_l$ . It is clear that if  $x \in X$ ,  $y \in \partial_S X$ , and  $\gamma \in \Gamma_\sigma(x, y)$ , then  $\gamma$  must exit from all balls  $B_l(o, r)$ . Thus  $\partial_S X$  is a type of “boundary at infinity” of the space  $(X, l)$  which generalizes the role of the point at infinity on the Riemann sphere.

**Proposition 2.20.** *Suppose  $(X, l, o)$  is an unbounded pointed local length space and  $\sigma = \mathfrak{S}(l, o, g)$  for some  $C$ -sphericalizing function  $g$ . If some point  $x \in X$  has a strong escape path, then  $\partial_S X$  is nonempty. In any case,  $\partial_S X$  is always a closed subset of  $\overline{X}_\sigma$ .*

*Proof.* Let  $\gamma : [0, \infty) \rightarrow X$  be a strong escape path, so that  $\gamma$  is parametrized by  $l$ -arclength,  $\gamma(0) = x$ , and  $|\gamma(t)| \geq c(t + |x|)$ . Using Fact 2.12, (S1) and (S2), we get

$$\text{len}_\sigma(\gamma) = \int_0^\infty g(|\gamma(t)|) dt \approx \int_0^\infty g(t + |x|) dt = \int_{|x|}^\infty g(s) ds \lesssim (1 + |x|)g(|x|). \quad (2.21)$$

Since  $\gamma$  has finite  $\sigma$ -length,  $\gamma(t)$  has a  $\sigma$ -limit  $w \in \overline{X}_\sigma$ . Now  $w$  cannot be in  $\overline{X}_l$ , since then  $\gamma$  would have to  $l$ -converge to  $w$ . Thus  $w \in \partial_S X$ , and  $\partial_S X$  is nonempty.

Since paths converging to  $\partial_S X$  must leave all balls  $B_l(o, r)$ , it follows from (2.9) that

$$\text{dist}_\sigma(y, \partial_S X) \geq \int_{|y|}^\infty g(t) dt, \quad y \in \overline{X}_l. \quad (2.22)$$

So a sequence in  $\partial_S X$  cannot converge to a point in  $\overline{X}_l$ , and  $\partial_S X$  is  $\sigma$ -closed.  $\square$

**Example 2.23.** In the absence of the strong escape path assumption,  $\partial_S X$  might be empty. This is indeed the case if  $X = \bigcup_{j=1}^\infty S_j$ , where  $S_j \subset \mathbb{R}^2$  is the circle centered at  $(j, 0)$  of radius  $j$ ; as usual,  $l$  is the inner Euclidean metric and  $o$  is the origin. Then  $\partial X_l$  and  $\partial_S X$  are both empty. A similar example is the space in Example 2.18; there  $\partial_S X$  is empty but  $\partial X_l$  is nonempty.



## 2.24. Comparability of quasihyperbolic metrics.

Let us first introduce some notation and estimates that will be used throughout the remainder of this section. Suppose  $g$  is a  $C$ -sphericalizing function. Given  $x \in X$ , let  $B_x = B_l(x, r_x)$ , where  $r_x := 1 \vee (|x|/2)$ . Let  $A_j = B_l(o, 2^j) \setminus B_l(o, 2^{j-1})$ ,  $j \in \mathbb{N}$ ,  $A_0 = B_l(o, 1)$ , and let  $A_{-1}$  be the empty set. We also define fatter annuli  $A_{i,j} = \bigcup_{m=i}^j A_m$  for all  $-1 \leq i \leq j \leq \infty$ . We denote the  $l$ -closure of any of these sets by means of a bar:  $\overline{B}_x, \overline{A}_j$ , etc. Note that  $B_x \subset A_{j-1, j+1}$  for all  $x \in A_j$ .

It follows from (S1) that  $g(|y|)/g(|x|) \in [1/C, C]$  for all  $y \in \overline{B}_x$ , and that  $g(|y|)/g(|z|) \in [1/C, C]$  for all  $y, z \in \overline{A}_j$ . Thus if  $\gamma$  is a path in  $X$  then

$$C^{-1} \leq \frac{\text{len}_\sigma(\gamma)}{\text{len}_l(\gamma)g(|x|)} \leq C, \quad \gamma \subset \overline{B}_x. \quad (2.25)$$

$$C^{i-j-1} \leq \frac{\text{len}_\sigma(\gamma)}{\text{len}_l(\gamma)g(|x|)} \leq C^{j+1-i}, \quad \gamma \subset \overline{A}_{i,j}, \quad x \in \overline{A}_{i,j}, \quad 0 \leq i \leq j. \quad (2.26)$$

If  $\gamma$  is a path that passes through  $x$  of length at least  $r_x$ , then it has a piece of length at least  $r_x$  inside  $B_x$ , and so (2.25) implies that

$$\text{len}_\sigma(\gamma) \geq C^{-1}r_x g(|x|). \quad (2.27)$$

In particular, this estimate holds for any path through  $x$  that is not contained in  $B_x$ , and for any path through  $x \in \overline{A}_i$  that is not contained in  $A_{i-1, i+1}$ .

We now state and prove the main result of this section which concerns the relationship between quasihyperbolic metrics associated with  $l$  and  $\sigma$ . We do not want to restrict  $(X, l)$  to be incomplete, so we use  $k_{l,C}$  for some  $C < \infty$  as the quasihyperbolic metric associated with  $l$ . By contrast, only incomplete spaces  $(X, \sigma)$  are of interest, so we use  $k_\sigma$  as the quasihyperbolic metric associated with  $\sigma$ . Note that

$$k_{l,C}(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma \frac{dl(z)}{\delta_{l,C}(z)},$$

$$k_\sigma(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma \frac{d\sigma(z)}{\delta_\sigma(z)} = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma \frac{g(|z|) dl(z)}{\delta_\sigma(z)},$$

where  $dl$  and  $d\sigma$  are the length elements with respect to  $l$  and  $\sigma$ , respectively. Note also that if  $(X, l)$  is locally incomplete, then so is  $(X, \sigma)$ .

**Theorem 2.28.** *Let  $(X, l, o)$  be a locally complete unbounded pointed local length space, let  $\sigma := \mathfrak{S}(l, o, g)$  for some  $C_g$ -sphericalizing function  $g$ , and let  $1 \leq C_l < \infty$ .*

- (a) *If  $\partial X_\sigma$  is nonempty, then  $k_\sigma \leq C' k_{l, C_l}$ , where  $C' = C'(C_g, C_l)$ .*
- (b) *If  $(X, l, o)$  has some  $(c, \rho)$ -escape property, then  $\partial X_\sigma$  is nonempty and  $k_\sigma \geq c' k_{l, C_l}$  for some  $c' = c'(c, \rho, C_g) > 0$ .*
- (c) *Conversely if  $k_\sigma \geq c' k_{l, C_l}$  for some  $c' > 0$ , then  $(X, l, o)$  has a  $(c, \rho)$ -escape property, for some  $c, \rho$  dependent only on  $c', C_g$ , and  $C_l$ .*

*Proof.* Since  $k_{l,C_l}$  is the conformal deformation of  $l$  with density function  $1/\delta_{l,C_l}(x)$ , and  $k_\sigma$  is the conformal deformation of  $l$  with density function  $g(|x|)/\delta_\sigma(x)$ , inequalities between  $k_{l,C_l}$  and  $k_\sigma$  are equivalent to the corresponding reverse inequalities between  $\delta_\sigma(x)$  and  $g(|x|)\delta_{l,C_l}(x)$ .

To prove (a), we need to prove that  $\delta_\sigma(x) \gtrsim g(|x|)\delta_{l,C_l}(x)$ . Thus we need to find  $C' = C'(C_g, C_l)$  such that for all  $y \in \partial X_\sigma$ ,  $\gamma \in \Gamma_\sigma(x, y)$ ,

$$\text{len}_\sigma(\gamma) \geq g(|x|)\delta_{l,C_l}(x)/C'. \quad (2.29)$$

Let  $j \geq 0$  be such that  $x \in A_j$ , and let  $\gamma^-$  be the half-open curve  $\gamma \setminus \{y\}$ . If  $\gamma^- \subset A_{j-1, j+1}$ , then necessarily  $y \in \partial X_l$  and so (2.26) implies that  $\text{len}_\sigma(\gamma) \geq g(|x|)\text{len}_l(\gamma)/C_g^3 \geq g(|x|)\delta_{l,C_l}(x)/C_g^3$ . On the other hand if  $\gamma^- \not\subset A_{j-1, j+1}$ , then by (2.27),  $\text{len}_\sigma(\gamma \cap A_{j-1, j+1}) \gtrsim g(|x|)2^j$ . But  $\delta_{l,C_l}(x) \leq C_l + |x| \lesssim 2^j$ , so again  $\text{len}_\sigma(\gamma) \gtrsim g(|x|)\delta_{l,C_l}(x)$ .

We now turn to the proof of (b). Suppose that  $(X, l, o)$  has the  $(c, \rho)$ -escape property; we assume without loss of generality that  $\rho \geq 2$ . We must prove that  $\delta_\sigma(x) \lesssim g(|x|)\delta_{l,C_l}(x)$ . If  $\delta_{l,C_l}(x) < r_x$ , then  $\delta_{l,C_l}(x) = \delta_l(x)$ , so the local length property ensures that we can choose  $y \in \partial X_l$  and  $\gamma \in \Gamma_l(x, y)$  with  $L := \text{len}_l(\gamma) < r_x \wedge (2\delta_l(x))$ . Now (2.25) tells us that

$$\delta_\sigma(x) \leq \text{len}_\sigma(\gamma) \leq C_g g(|x|)L \leq 2C_g g(|x|)\delta_{l,C_l}(x).$$

We may therefore assume that  $\delta_{l,C_l}(x) \geq r_x \geq 1$ .

Suppose  $|x| \geq \rho$ . Let  $\gamma : [0, T) \rightarrow X$  be parametrized by  $l$ -arclength such that  $\gamma(0) = x$ ,  $|\gamma(t)| \geq c(t + |x|)$ , and either  $T = \infty$ , or  $T < \infty$  with  $\gamma(T-; l) = y \in \partial X_l$ . Now  $|x| \geq \rho \geq 2$  and  $r_x = |x|/2 \leq \delta_{l,C_l}(x)$ , so if  $T < \infty$ , then  $y \in \partial_0 X_l \subset \partial X_\sigma$ , so  $\partial X_\sigma$  is nonempty. Furthermore, using Fact 2.12 and (S2), we get

$$\delta_\sigma(x) \leq \int_0^T g(|\gamma(t)|) dt \approx \int_0^T g(t + |x|) dt \leq C_g g(|x|)|x| \leq 2C_g g(|x|)\delta_{l,C_l}(x),$$

as desired. If instead  $T = \infty$ , it follows similarly that  $\text{len}_\sigma(\gamma) \lesssim g(|x|)|x|$ , and so  $\gamma(t)$  is  $\sigma$ -convergent to some point  $w \in \overline{X}_\sigma$ . But  $w$  cannot be in  $X \cup \partial_0 X_l$ , since  $\gamma$  would then have to  $l$ -converge to  $w$ . Thus  $w \in \partial_S X$ , so  $\partial X_\sigma$  is again nonempty and  $\delta_\sigma(x) \leq \text{len}_\sigma(\gamma) \lesssim g(|x|)\delta_{l,C_l}(x)$ .

Suppose instead that  $|x| < \rho$ . Since  $X$  is an unbounded length space, there exists  $x_0 \in X$  with  $|x_0| = \rho$ . By the previous case with  $C_l = 1$ , we see that  $\delta_\sigma(x_0) \lesssim g(\rho)(1 + \rho) \lesssim g(|x|)$ . As in the proof of Proposition 2.6, we see that  $\sigma(x, x_0) \leq 4C_g(1) \lesssim g(|x|)$ , and so

$$\delta_\sigma(x) \leq \sigma(x, x_0) + \delta_\sigma(x_0) \lesssim g(|x|) \approx g(|x|)\delta_{l,C_l}(x).$$

This finishes the proof of (b).

Finally we prove (c). Assume that  $\delta_\sigma(z) \leq g(|z|)\delta_{l,C_l}(z)/c'$ ,  $z \in X$ . We also assume without loss of generality that  $0 < c' < 1$ . Let  $K = 2(C_l + 1)/c'$  and  $\rho = \alpha^{-1} > 1$ , where  $\alpha = \alpha(C_g, K)$  is as in Lemma 2.4. We prove that  $(X, l, o)$  has a  $(c, \rho)$ -escape property for some  $0 < c < 1$  by constructing an escape path  $\gamma$  for an arbitrary point  $x$ ,  $|x| \geq \rho$ . Let  $\lambda_0 : [0, T_0) \rightarrow X$  be a path parametrized by  $\sigma$ -arclength such that  $\lambda_0(0) = x$ ,  $\lambda_0(T_0-; \sigma) = w_0 \in \partial X_\sigma$ , and  $T_0 < 2g(|x|)\delta_{l,C_l}(x)/c'$ . Let  $\nu_0 : [0, S_0) \rightarrow X$  be the reparametrization of  $\lambda_0$  by  $l$ -arclength. Note that  $S_0$  may be finite or infinite.

We first show that  $\lambda_0$  stays outside the ball  $B_l(o, |x|/\rho)$ . If this were not the case, then according to Lemma 2.4, we would have

$$T_0 = \text{len}_\sigma(\lambda_0) > \int_{|x|/\rho}^{|x|} g(t) dt > K|x|g(|x|).$$

Since  $\delta_{l, C_l}(x) \leq (C_l + 1)|x|$ , this contradicts the fact that  $T_0 < 2g(|x|)\delta_{l, C_l}(x)/c' \leq K|x|g(|x|)$ .

Let  $s_1$  be the least number  $s$  for which  $|\nu_0(s)| = 2|x|$ , or let  $s_1 = S_0$  if  $|\nu_0(s)| < 2|x|$  for all  $0 \leq s \leq S_0$ . We define  $x_1 = \nu_0(s_1)$  and  $\gamma_1 = \nu_0|_{[0, s_1]}$ . Since the values  $g(t)$ ,  $t \in [|x|/\rho, 2|x|]$ , are all mutually comparable, the bound  $T_0 < K|x|g(|x|)$  implies that  $s_1 \lesssim |x|$ . If  $s_1 = S_0$ , we have reached a stopping point and  $\gamma_1$  is the desired escape path; notice that  $|\gamma_1(t)| \geq |x|/\rho$  for  $t \in [0, s_1]$ . Otherwise we iterate the argument as follows.

We choose a path  $\lambda_1 : [0, T_1) \rightarrow X$ , parametrized by  $\sigma$ -arclength, such that  $\lambda_1(0) = x_1$ ,  $\lambda_1(T_1-; \sigma) = w_1 \in \partial X_\sigma$ , and  $T_1 < 2g(|x_1|)\delta_{l, C_l}(x_1)/c'$ . Let  $\nu_1 : [0, S_1) \rightarrow X$  be the reparametrization of  $\lambda_1$  by  $l$ -arclength. Replacing the input data  $(x, w_0, \lambda_0, \nu_0)$  in the above argument by  $(x_1, w_1, \lambda_1, \nu_1)$ , we get as output  $(s_2, x_2, \gamma_2)$  where  $s_2 \lesssim |x_1|$ ,  $x_2 = \nu_1(s_1)$ , and  $\gamma_2 = \nu_1|_{[0, s_2]}$ . Furthermore, either  $s_2 = S_1$ , in which case we have reached a stopping point, or  $s_2 < S_1$  and  $|x_2| = 2|x_1|$ , in which case we continue to iterate the process.

We continue this iterative procedure until we reach a stopping point, or forever if this never occurs. All the paths  $\gamma_j$  obtained by this process can be joined to form a single path  $\gamma$  parametrized by  $l$ -arclength. Since each  $\gamma_j$  has  $l$ -arclength  $s_j \lesssim 2^j$ , and each point  $y \in \gamma_j$  has the property that  $|y| \approx 2^j|x|$ , it readily follows that  $\gamma$  is an escape path for  $x$ .  $\square$

Note that if  $(X, l)$  is incomplete, then  $\delta_{l, C} = \delta_l$  and  $k_{l, C} = k_l$  for all  $C \geq l(o, \partial X)$ . Thus Theorem 2.28 says that, in the presence of an escape property, the usual quasihyperbolic metrics  $k_l$  and  $k_\sigma$  are comparable with a constant dependent only on the escape parameters,  $C_g$ , and  $l(o, \partial X)$ .

Since Euclidean domains are locally complete and satisfy a  $(1, 0)$ -escape property, we can as a special case apply this last statement with  $(X, l)$  being the Euclidean domain  $\Omega \subsetneq \mathbb{R}^n$ ,  $g(t) = 2/(1 + t^2)$ , and  $o = x_0 \in \Omega$ . The resulting sphericalized metric  $\sigma$  is a bilipschitz distortion of the usual inner spherical metric  $\sigma'$ , with a bilipschitz constant dependent only on  $|x_0|$ ; note that  $\sigma = \sigma'$  only if  $x_0 = 0$ ; note that the case  $n = 1$  is a little different, but the result is still valid in this case. Thus we have proven Theorem 0.1.

### 2.30. Reverse escape properties.

In §3.4, we shall discuss the concept of (strong) reverse escape properties in more detail but for now we wish to show only that the sphericalization process always produces data with such a property. Given  $\sigma := \mathfrak{S}(l, o, g)$  as usual, and  $0 < c \leq 1$ , we say that  $(X, \sigma, o)$  has the *reverse  $c$ -escape property* if, for every  $x \in X$ , there exists  $y \in \{o\} \cup \partial_0 X$  and  $\gamma \in \Gamma_\sigma(x, y)$  such that  $\delta_S(\gamma(t)) \geq c(t + \delta_S(x))$ ,  $0 \leq t \leq T = \text{len}_\sigma(\gamma)$ , where  $\delta_S(y) = \text{dist}(x, \partial_S X)$ . If we can always take  $y = o$  in this condition, we say that  $(X, \sigma, o)$  has the *strong reverse  $c$ -escape property*.

**Theorem 2.31.** *Let  $(X, l, o)$  be a pointed local length space, and let  $\sigma = \mathfrak{S}(l, o, g)$ , for some  $C_g$ -sphericalizing function  $g$ . Then*

- (a)  $(X, \sigma, o)$  has a reverse  $c$ -escape property, and
- (b) if  $l$  is a length metric, then  $(X, \sigma, o)$  has a strong reverse  $c$ -escape property.

Moreover,  $c > 0$  can be taken to depend only on  $C_g$ .

*Proof.* Let us prove (b). We need to find a strong reverse escape path  $\gamma$  for a general point  $x \in X$ . If  $|x| \leq 1$ , we can take  $\gamma$  to be any path of  $l$ -length at most 2: such a path must remain in the ball  $B_l(o, 2)$ , and so has  $\sigma$ -length comparable to  $g(1)$  which dominates  $\delta_S(y)$  for every  $y \in B_l(o, 2)$  by (2.9) and (S2).

We may therefore assume that  $|x| > 1$ . For  $z \in X$ , define  $j(z)$  to be the non-negative integer satisfying  $z \in A_{j(z)}$ . Let  $\gamma \in \Gamma_\sigma(o, x)$  have  $l$ -length  $L$  and  $\sigma$ -length  $M$ , where  $L \leq |x| + 1$ . We cut  $\gamma$  into segments  $\gamma_i = \gamma|_{[s_{i-1}, s_i]}$ , where  $s_0 = 0$ ,  $s_{j(x)} = M$  and, for  $0 < i < j(x)$ ,  $\gamma(s_i)$  is the last point on  $\gamma$  contained in  $\overline{A_i}$ . By construction,  $\gamma_i \subset A_{i-1}^{i+1}$  and  $\text{len}_l(\gamma_i) \lesssim 2^i + 1$ . Thus  $\text{len}_\sigma(\gamma_i) \lesssim 2^i g(2^i)$ . But by (S3), the series  $\sum_{i=j(w)}^\infty 2^i g(2^i)$  is dominated by a convergent geometric series, and so  $M - t \lesssim 2^{j(w)} g(j(w))$  whenever  $w = \gamma(t)$ .

If we prove the estimate  $\delta_S(w) \gtrsim 2^{j(w)} g(j(w))$ , it therefore follows that the reverse parametrization  $-\gamma \in \Gamma_\sigma(x, o)$  is a reverse escape path. But this last estimate follows rather easily. Indeed if  $\lambda$  is a path from  $w$  to  $\partial_S X$ , then  $\lambda$  exits all  $l$ -balls and so (2.9) and the properties of a sphericalization function tell us that

$$\text{len}_\sigma(\lambda) \geq \int_{|w|}^{2^{|w|+1}} g(t) dt \gtrsim (|w| + 1)g(|w|).$$

Part (a) follows from (b) if  $(X, l)$  is complete, since then  $l$  is a length metric. If  $(X, l)$  is incomplete, the proof of (a) is similar to that of (b) except that we use a path  $\gamma \in \Gamma_\sigma(y, x)$  of length at most  $l(x, y) + 1$ , where  $y \in \partial_l X$  and  $l(x, y) = \delta_l(x)$ .  $\square$

We cannot conclude in Theorem 2.31(a) that  $(X, \sigma, o)$  has a strong reverse  $c$ -escape property, as the following example indicates.

**Example 2.32.** Consider the domain  $X \subset \mathbb{R}^2$  given by the curve complement

$$\{(s, t) \mid s, t \in \mathbb{R}, s \geq 1, t \neq s^2 \sin^2 s\} \subset \mathbb{R}^2.$$

Taking  $l$  to be the Euclidean metric,  $o = (0, 0)$ , and  $g(t) = 2/(1+t^2)$ , the metric  $\sigma = \mathfrak{S}(l, o, g)$  is the inner spherical metric. The space  $(X, \sigma, o)$  fails to have any strong reverse  $c$ -escape property, as we see by taking  $x = (j\pi, 1) \in X$  for sufficiently large  $j = j_c \in \mathbb{N}$ .

### 3. FLATTENING

In this section we proceed in the direction opposite to that of the last section: we “flatten” a bounded incomplete local length space  $(X, \sigma)$  to get an unbounded length space  $(X, l)$ . We use the term “flattening” because this process generalizes that of obtaining Euclidean

space from the Riemann sphere. The main result, Theorem 3.11, compares the corresponding quasihyperbolic metrics  $k_\sigma$  and  $k_{l,C}$ .

Throughout this section, we assume that the data  $(X, \sigma, \partial_S X)$  are  $d_0$ -spherical data for some  $0 < d_0 < \infty$ , which we define to mean that the data have the following pair of properties:

- (1)  $(X, \sigma)$  is an incomplete local length space of diameter at most  $d_0$ ;
- (2)  $\partial_S X$  is a nonempty closed subset of  $\overline{X}_\sigma$ .

We shall denote by  $\partial_0 X$  the set of all points in  $\partial X_\sigma \setminus \partial_S X$  that are accessible by  $\sigma$ -rectifiable paths from points in  $X$ . Note that  $\partial_0 X$  equals  $\partial X_\sigma \setminus \partial_S X$  if  $\sigma$  is a length metric. We write  $\delta_S(x) = \sigma(x, \partial_S X)$ , for all  $x \in X \cup \partial_0 X$ .

By the results of Section 2, we know that if we sphericalize a length space in which  $o$  has a strong escape path, we always get spherical data  $(X, \sigma, \partial_S X)$ , with  $\sigma$  being a length metric. However, we shall see in the next section that there are such spherical data that do not arise in this way, even modulo a bilipschitz distortion.

Fixing a function  $f : (0, d_0] \rightarrow (0, \infty)$ , we define a new metric  $\mathfrak{L}(\sigma, \partial_S X, f)$  on  $X$  by the equation

$$\mathfrak{L}(\sigma, \partial_S X, f)(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma f(\delta_S(z)) d\sigma(z), \quad x, y \in X.$$

We normally write  $l$  in place of  $\mathfrak{L}(\sigma, \partial_S X, f)$ , and we take  $f$  to be any *flattening function*, a concept we now define.

### 3.1. Flattening functions.

A continuous function  $f : (0, d_0] \rightarrow (0, \infty)$  is a  $(C, d_0)$ -flattening function,  $C > 2$ ,  $d_0 > 0$ , if it has the following properties:

- (F1)  $f(r) \leq C f(s)$  whenever  $r, s \in (0, d_0]$ ,  $1/2 \leq r/s \leq 2$ ;
- (F2)  $\int_r^{d_0} f(t) dt \leq C r f(r)$ , whenever  $0 < r \leq d_0$ .

Typical flattening functions include  $f(t) = \log^q(2 + 1/t)t^{-p}$ , for  $p > 1$ ,  $q \in \mathbb{R}$ . The usual flattening of the Riemann sphere into the Euclidean metric uses  $f(t) = \sec^2((\pi - t)/2)/2$ ,  $0 < t \leq d_0 := \pi$ ; note that  $f$  is comparable with  $t \mapsto t^{-2}$ .

It follows from (F2) that if  $f$  is a flattening function, then  $f(r) \geq c/r$  for all  $0 < r < d_0/2$ , where  $c = C^{-1} \int_{d_0/2}^{d_0} f(t) dt$ . Consequently,  $\int_0^{d_0} f(t) dt = \infty$ ; the divergence of this integral pushes  $\partial_S X$  “infinitely far” from every point in  $X$  with respect to the metric  $l = \mathfrak{L}(\sigma, \partial_S X, f)$ . Consequently, we see that we can naturally identify  $\partial X_l$  with  $\partial_0 X$ . In particular if  $\sigma$  is a length metric, we can identify  $\partial X_\sigma$  with the disjoint union of  $\partial_S X$  and  $\partial X_l$ . Note also that if  $y \in \partial X_\sigma$  is such that  $\sigma(x, y) = \text{dist}_\sigma(x, \partial X_\sigma)$  for some  $x \in \Omega$ , then there are paths  $\gamma \in \Gamma_\sigma(x, y)$  of  $\sigma$ -length arbitrarily close to  $\sigma(x, y)$ .

We now state two lemmas. The first gives a useful alternative description of flattening functions, and it readily implies the second lemma. We omit both proofs, as they are very similar to those of Lemmas 2.2 and 2.4.

**Lemma 3.2.** *A Borel function  $f : (0, d_0] \rightarrow (0, \infty)$  is a flattening function if and only if it satisfies (F1) and a condition of the following type for some constants  $C' \geq 1$ ,  $\epsilon > 0$ :*

$$(F3) \quad \frac{f(r)}{f(s)} \leq C' \left(\frac{s}{r}\right)^{1+\epsilon}, \text{ for all } 0 < s \leq r \leq d_0.$$

*Furthermore, this equivalence holds with quantitative control of parameters.*

**Lemma 3.3.** *Given  $C > 2$  and  $d_0, K > 0$ , there exists  $\alpha = \alpha(C, K)$  such that, whenever  $f$  is a  $C$ -flattening function and  $0 < r \leq d_0$ ,*

$$\int_{\alpha r}^r f(t) dt \geq Kr f(r).$$

We shall see that the quasihyperbolic metrics associated with  $(X, \sigma)$  and  $(X, l)$  are bilipschitz equivalent if and only if  $(X, \sigma, o)$  satisfies a reverse escape property similar to the one at the end of Section 2.

### 3.4. Reverse escape properties revisited.

Suppose  $(X, \sigma, \partial_S X)$  are spherical data. Given  $0 < c \leq 1$ , we say that  $(X, \sigma, o)$  has the *reverse  $c$ -escape property* if, for every  $x \in X$ , there exists  $y \in \{o\} \cup \partial_0 X$  and  $\gamma \in \Gamma_\sigma(x, y)$  such that  $\delta_S(\gamma(t)) \geq c(t + \delta_S(x))$ , for all  $0 \leq t < T = \text{len}_\sigma(\gamma)$ . If we can always take  $y = o$  in this condition, we say that  $(X, \sigma, o)$  has the *strong reverse  $c$ -escape property*. Since not all spherical data arise via sphericalization, this generalizes the previous definition of a strong reverse escape property.

**Example 3.5.** Suppose that  $\Omega$  is an unbounded domain in  $\mathbb{R}^n$  that includes the origin  $o$ , let  $\sigma$  be the spherical metric (or the inner spherical metric if  $\Omega$  is of bounded inner spherical diameter), and let  $\partial_S \Omega = \{\infty\}$ . Given  $x \in \Omega$ , the Euclidean line segment from  $x$  to  $o$  provides us with a path  $\gamma_1$ , parametrized by spherical arclength, which satisfies the estimate  $\delta_S(\gamma_1(t)) \geq t + \delta_S(x)$ . We pick  $\gamma$  to be an initial segment of  $\gamma_1$  that ends the first time that  $\gamma_1$  hits  $\partial\Omega \cup \{o\}$ . Thus  $(\Omega, \sigma, o)$  always has a reverse 1-escape property. However,  $(\Omega, \sigma, o)$  may fail to have a strong reverse escape property; one example of this type, with bounded inner spherical diameter, is the domain in Example 2.32.

**Example 3.6.** Consider  $X \subset \mathbb{R}^2$ , where  $X$  consists of the horizontal line segment  $(x_1, 0)$ ,  $0 < x_1 \leq 1$ , together with vertical line segments of the form  $(a_n, x_2)$ ,  $0 \leq x_2 \leq b_n$ ,  $n \in \mathbb{N}$ , where  $a_1 < 1$ ,  $(a_n)$  is a strictly decreasing sequence of positive numbers tending to zero, and  $(b_n)$  is a bounded sequence of positive numbers. We equip  $X$  with the inner Euclidean metric, which we denote by  $\sigma$ . Then  $\partial X_\sigma$  contains only the single point  $(0, 0)$  and we define  $\partial_S X = \partial X_\sigma$ . We also write  $o = (1, 0) \in X$ . It is clear that  $(X, \sigma, o)$  has a (strong) reverse escape property if and only if  $(b_n/a_n)$  is bounded. In particular, if we choose  $a_n = 1/(n+1)!$  and  $b_n = 1/n!$ , then  $(X, \sigma, o)$  fails to have a reverse escape property despite the fact that it is in other ways a rather nice metric space, being for instance a compact Ahlfors regular tree.

**Example 3.7.** Let the data  $(X, \sigma, \partial_S X)$  be as in the previous example. Given  $T \subset \mathbb{N}$ , define  $A_T := \{(a_n, b_n) \mid n \in T\} \subset X$ ,  $X_T = X \setminus A_T$ ,  $\partial_0 X_T = \partial_0 X \cup A_T$ , and denote by  $\sigma$  the subspace metric inherited by  $X_T$ . Then  $(X_T, \sigma, \partial_T X)$  are spherical data, and  $(X_T, \sigma, o)$  has a reverse escape property if and only if  $b_n/a_n$  is uniformly bounded for  $n \in \mathbb{N} \setminus T$ . In the extreme case  $T = \mathbb{N}$ ,  $(X_{\mathbb{N}}, \sigma, o)$  always has a reverse 1-escape property. A strong reverse escape property is more restrictive: it always requires the full sequence  $(b_n/a_n)$  to be bounded.

We pause to introduce some notation and estimates that will be used throughout the remainder of this section. Suppose  $f$  is a  $(C, d_0)$ -flattening function. Given  $x \in X$ , let  $B'_x = B_\sigma(x, r'_x)$ , where  $r'_x := \delta_S(x)/2$ . Let

$$L_j = \{x \in X \mid 2^{-j}d_0 < \delta_S(x) \leq 2^{-j+1}d_0\}, \quad j \in \mathbb{N},$$

and let  $L_{-1}$  be the empty set. We also define fatter layers  $L_{i,j} = \bigcup_{m=i}^j L_m$  for all  $-1 \leq i \leq j \leq \infty$ . We denote the  $\sigma$ -closure of any of these sets by means of a bar:  $\overline{B'_x}$ ,  $\overline{L}_j$ , etc. Note that  $B'_x \subset L_{j-1, j+1}$  for all  $x \in L_j$ .

It follows from (F1) that  $f(\delta_S(y))/f(\delta_S(x)) \in [1/C, C]$  for all  $y \in \overline{B'_x}$ , and that  $f(\delta_S(y))/f(\delta_S(z)) \in [1/C, C]$  for all  $y, z \in \overline{L}_j$ . Thus if  $\gamma$  is a path in  $X$  then

$$C^{-1} \leq \frac{\text{len}_l(\gamma)}{\text{len}_\sigma(\gamma)f(\delta_S(x))} \leq C, \quad \gamma \subset \overline{B'_x}. \quad (3.8)$$

$$C^{i-j-1} \leq \frac{\text{len}_l(\gamma)}{\text{len}_\sigma(\gamma)f(\delta_S(x))} \leq C^{j+1-i}, \quad \gamma \subset \overline{L}_{i,j}, \quad x \in \overline{L}_{i,j}, \quad 0 \leq i \leq j. \quad (3.9)$$

If  $\gamma$  is a path that passes through  $x$  of  $\sigma$ -length at least  $r'_x$ , then it has a piece of  $\sigma$ -length at least  $r'_x$  inside  $B'_x$ , and so (3.8) implies that

$$\text{len}_l(\gamma) \geq \delta_S(x)f(\delta_S(x))/2C, \quad (3.10)$$

In particular, this estimate holds for any path through  $x$  that is not contained in  $B'_x$ , and for any path through  $x \in \overline{L}_i$  that is not contained in  $L_{i-1, i+1}$ .

We now state and prove the main result of this section which concerns the relationship between quasihyperbolic metrics associated with  $\sigma$  and  $l$ . We do not want to restrict  $(X, l)$  to be incomplete, so we use  $k_{l,C}$  for some  $C < \infty$  as the quasihyperbolic metric associated with  $l$ . By contrast, only incomplete spaces  $(X, \sigma)$  are of interest, so we use  $k_\sigma$  as the quasihyperbolic metric associated with  $\sigma$ . Note that

$$k_\sigma(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma \frac{d\sigma(z)}{\delta_\sigma(z)},$$

$$k_{l,C}(x, y) = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma \frac{dl(z)}{\delta_{l,C}(z)} = \inf_{\gamma \in \Gamma(x, y)} \int_\gamma \frac{f(\delta_S(z)) d\sigma(z)}{\delta_{l,C}(z)},$$

where  $dl$  and  $d\sigma$  are the length elements with respect to  $l$  and  $\sigma$ , respectively. As before,  $\delta_{l,C}(z)$  is defined relative to some fixed basepoint  $o \in X$ , and we write  $|x| = l(x, o)$ .

**Theorem 3.11.** *Let  $(X, \sigma, \partial_S X)$  be  $d_0$ -spherical data, where  $(X, \sigma)$  is locally complete, let  $l := \mathfrak{L}(\sigma, \partial_S X, f)$  for some  $(C_f, d_0)$ -flattening function  $f$ , let  $0 < C_l < \infty$ , and define  $h(t) := tf(t)/C_l$ .*

- (a)  $k_{l, C_l} \leq C' k_\sigma$ , where  $C'$  depends only on  $C_f$  and  $h(\delta_S(o))$ .
- (b) If  $(X, \sigma, o)$  has a reverse  $c$ -escape property, then  $k_{l, C_l} \geq c' k_\sigma$  for some  $c' > 0$  dependent only on  $c$ ,  $C_f$ , and  $h(d_0)$ .
- (c) Conversely if  $k_{l, C_l} \geq c' k_\sigma$  for some  $c' > 0$ , then  $(X, \sigma, o)$  has a reverse  $c$ -escape property for some  $c = c(c', C_f)$ .

*Proof.* Since  $k_\sigma$  is the conformal deformation of  $\sigma$  with density function  $1/\delta_\sigma(x)$ , and  $k_{l, C_l}$  is the conformal deformation of  $\sigma$  with density function  $f(\delta_S(x))/\delta_{l, C_l}(x)$ , inequalities between  $k_{l, C_l}$  and  $k_\sigma$  are equivalent with the corresponding reverse inequalities between  $\delta_{l, C_l}(x)$  and  $f(\delta_S(x))\delta_\sigma(x)$ . Note also that if we replace  $f$  by  $sf$  and  $C_l$  by  $sC_l$  for any number  $s > 0$ , then both the hypotheses and conclusions remain unchanged. This allows us to assume without loss of generality a normalization  $t_0 f(t_0) = 1$ , where  $t_0 \in (0, d_0]$  is any value that suits our purposes; dependence on  $h(t_0)$  is then equivalent to dependence on  $C_l$ .

Part (a) is equivalent to the inequality  $f(\delta_S(x))\delta_\sigma(x) \lesssim \delta_{l, C_l}(x)$ . We assume without loss of generality that  $\delta_S(o)f(\delta_S(o)) = 1$ . We first wish to prove that  $f(\delta_S(x))\delta_\sigma(x) \lesssim C_l + |x|$ . If  $\delta_S(x) \leq \delta_S(o)/2$ , then using (F1) we see that

$$f(\delta_S(x))\delta_\sigma(x) \leq f(\delta_S(x))\delta_S(x) \leq C_f \int_{\delta_S(x)}^{2\delta_S(x)} f(t) dt \leq C_f |x|$$

Note that the last inequality follows by similar reasoning to that in Remark 2.8. If instead  $\delta_S(x) > \delta_S(o)/2$ , then (F1), (F3), and our normalization ensure that  $f(\delta_S(x))\delta_\sigma(x) \lesssim 1 \lesssim C_l$ .

It remains to prove that  $f(\delta_S(x))\delta_\sigma(x) \lesssim \delta_l(x)$ ; in particular we may assume that  $X_l$  is incomplete. By (3.8), a path  $\gamma \in \Gamma_\sigma(x, y)$ ,  $y \in \partial_0 X = \partial X_l$ , has  $l$ -length comparable to  $\text{len}_\sigma(\gamma)f(\delta_S(x))$  if  $\gamma \subset \overline{B_x} \cup \{y\}$ ; this gives the desired inequality if such a path  $\gamma$  has  $l$ -length less than  $2\delta_l(x)$ . Alternatively if no such path exists, then we may apply (3.10) to every path  $\gamma$  from  $x$  to  $\partial X_l$  to get that  $\text{len}_l(\gamma) \gtrsim f(\delta_S(x)) \text{len}_\sigma(\gamma)$ . Taking a minimum over all such connecting paths  $\gamma$ , we deduce (a).

The conclusion of (b) is equivalent to the inequality  $\delta_{l, C_l}(x) \lesssim f(\delta_S(x))\delta_\sigma(x)$ . We assume without loss of generality that  $d_0 f(d_0) = 1$ . The reverse escape property provides us with a path  $\gamma \in \Gamma_\sigma(x, y)$  such that  $y \in \{o\} \cup \partial_0 X$  and  $\delta_S(\gamma(t)) \geq c_0(t + \delta_S(x))$ ,  $0 \leq t < T = \text{len}_\sigma(\gamma) \leq d_0$ . By this last inequality, a change of variables, (F3), and (F2), we see that

$$\begin{aligned} \text{len}_l(\gamma) &= \int_0^T f(\delta_S(\gamma(t))) dt \lesssim \int_0^T f(c_0(t + \delta_S(x))) dt \\ &\leq \int_{c_0 \delta_S(x)}^{d_0} f(s) ds \\ &\lesssim f(\delta_S(x))\delta_S(x). \end{aligned} \tag{3.12}$$



Assuming that  $\delta_\sigma(x) \geq \delta_S(x)/2$ , it follows that  $|x| \wedge \delta_l(x) \lesssim f(\delta_S(x))\delta_\sigma(x)$ . The desired inequality now follows since also

$$C_l \lesssim 1 = d_0 f(d_0) \lesssim t f(t), \quad 0 < t \leq d_0.$$

Suppose instead that  $\delta_\sigma(x) < \delta_S(x)/2$  and so we can find  $\gamma \in \Gamma_\sigma(x, y)$ ,  $y \in \partial_0 X$ , such that  $\text{len}_\sigma(\gamma) < (2\delta_\sigma(x)) \wedge (\delta_S(x)/2)$ . Then  $\gamma \subset B'_x \cup \{y\}$  and by (3.8), we have  $\delta_l(x) \leq \text{len}_l(\gamma) \lesssim f(\delta_S(x))\delta_\sigma(x)$ . This concludes the proof of (b).

For (c), we need to show that the inequality  $\delta_{l, C_l}(x) \leq f(\delta_S(x))\delta_\sigma(x)/c'$ ,  $x \in X$ , implies a reverse escape property. Let  $x \in X$  be fixed but arbitrary. There exists a point  $w_0 \in \{o\} \cup \partial_0 X$ , and a path  $\lambda_0 : [0, T_0) \rightarrow X$ , parametrized by  $l$ -arclength, such that  $\lambda_0(0) = x$ ,  $\lambda_0(T_0-; l) = w_0$ , and  $T_0 \leq 2f(\delta_S(x))\delta_\sigma(x)/c'$ . Let  $\nu_0 : [0, S_0) \rightarrow X$  be the reparametrization of  $\lambda_0$  by  $\sigma$ -arclength.

Using Lemma 3.3 and the fact that  $\delta_S(\cdot) \geq \delta_\sigma(\cdot)$ , we see that  $\delta_S(\lambda_0(t)) \gtrsim \delta_S(x)$ . Next, using the bound  $T_0 \leq 2f(\delta_S(x))\delta_\sigma(x)/c'$  and the comparability of all values of  $f(s)$  for  $s \approx \delta_S(x)$ , we see that there must exist a positive number  $s_1 \lesssim \delta_S(x)$  such that either  $s_1 = S_0$ , or  $s_1 < S_0$  and  $\delta_S(z_1) > 2\delta_S(x)$ , where  $z_1 = \nu(s_1)$ . In the first case, we have reached a stopping point, while in the second case we iterate. As in the proof of Theorem 2.28, we get the desired path by joining together the segments  $\gamma_j$  obtained by this process; of course this time the process must stop after finitely many steps because  $\delta_S(\cdot)$  is bounded.  $\square$

### 3.13. Flattened spaces have a strong escape property.

We saw in Theorem 2.31 that if we sphericalize a length metric  $l$ , we get a metric  $\sigma$  satisfying a strong reverse escape property. We now prove the dual of this result: if we flatten a length metric  $\sigma$ , we get a metric  $l$  satisfying a strong escape property.

**Theorem 3.14.** *Let  $(X, \sigma, \partial_S X)$  be  $d_0$ -spherical data, where  $\sigma$  is a local length metric, and let  $l := \mathfrak{L}(\sigma, \partial_S X, f)$  for some  $(C, d_0)$ -flattening function  $f$ . Given  $o \in X$ , there exist  $0 < c < 1$ , and  $0 < \rho < \infty$  such that*

- (a)  $(X, l, o)$  has a  $(c, \rho)$ -escape property, and
- (b) if  $\sigma$  is a length metric, then  $(X, l, o)$  has a strong  $(c, \rho)$ -escape property.

Moreover, we can take  $c$  to depend only on  $C$ , and  $\rho$  to depend only on  $C$  and  $\delta_S(o)f(\delta_S(o))$ .

*Proof.* We prove only (b), since the proof of (a) is very similar. Let us fix  $o \in X$ . Given  $x \in X$ , with  $|x|$  sufficiently large, we shall construct a strong escape path  $\gamma$  by concatenating a sequence of path segments. First, write  $x_0 = x$ ,  $S = \delta_S(x)$ , and choose a path  $\lambda_0 : [0, S_0) \rightarrow X$ , parametrized by  $\sigma$ -arclength, with  $\lambda_0(0) = x_0$ ,  $\lambda_0(S_0-; \sigma) = y_0 \in \partial_S X$ , and  $S_0 < 2S$ . Let  $x_1 = \lambda_0(t_1)$ , where  $t_1 \in (0, S_0)$  is such that  $\delta_S(x_1) = S/2$  but  $\delta_S(\lambda_0(t)) > S/2$  for all  $0 < t < t_1$ . Let  $\gamma_1 = \lambda_0|_{[0, t_1]}$ . Inductively, given  $x_j \in X$  with  $\delta_S(x_j) = 2^{-j}S$ , we let  $\lambda_j : [0, S_j) \rightarrow X$  be parametrized by  $\sigma$ -arclength with  $\lambda_j(0) = x_j$ ,  $\lambda_j(S_j-; \sigma) = y_j \in \partial_S X$ , and  $S_j < 2\delta_S(x_j)$ . As for  $j = 0$ , we find  $x_{j+1} = \lambda_j(t_{j+1})$  such that  $\delta_S(x_{j+1}) = \delta_S(x_j)/2$  but  $\delta_S(\lambda_j(t)) > \delta_S(x_j)/2$  for all  $0 < t < t_{j+1}$ , and we let  $\gamma_{j+1} = \lambda_j|_{[0, t_{j+1}]}$ . Note that  $\text{len}_\sigma(\gamma_j) < 2^{2-j}S$ . Joining together the paths  $\gamma_j$ ,  $j \in \mathbb{N}$ , and parametrizing the resulting curve by  $l$ -arclength, we get a path  $\gamma : [0, \infty) \rightarrow X$  with  $\gamma(0) = x$  and of infinite  $l$ -length (since it connects  $x$  to  $\partial_S X$ ). It is convenient to denote by  $\gamma^j$  the initial segment of  $\gamma$  obtained by gluing together  $\gamma_i$ ,  $i \leq j$ .

By construction,  $\delta_S(z)/2^{1-j}S \in [1/2, 2]$  for  $z \in \gamma_j$ , and so  $f(\delta_S(z))/f(2^{1-j}S) \in [1/C, C]$ . Since the  $\sigma$ -length of  $\gamma_j$  is at most  $2^{2-j}S$ , we get that

$$\text{len}_l(\gamma_j) \leq C2^{2-j}Sf(2^{1-j}S) \leq 4C^2 \int_{2^{-j}S}^{2^{1-j}S} f(s) ds,$$

where the second inequality follows from (F1). If  $z = \gamma(t) \in \gamma_j$ , it follows from the last inequality and (F2) that

$$t \leq \text{len}_l(\gamma^j) \leq 4C^2 \int_{2^{-j}S}^S f(s) ds \leq C^3 2^{2-j}Sf(2^{-j}S). \quad (3.15)$$

Our goal is to obtain a lower estimate for  $|z| = \gamma(t)$  in terms of  $t$ . Suppose first that  $z \in \gamma_j$ , where  $2^{3-j}S \leq \delta_S(o)$ ; in particular  $\delta_S(z) < \delta_S(o)$ . Given  $\nu \in \Gamma_\sigma(z, o)$ , we have

$$\text{len}_l(\nu) = \int_0^{\text{len}_\sigma(\nu)} f(\delta_S(\nu(t))) dt.$$

Applying Lemma 2.5 with  $\alpha = \delta_S \circ \nu$  to this last equation and taking an infimum over all such  $\nu$ , we get

$$|z| \geq \int_{\delta_S(z)}^{\delta_S(o)} f(t) dt. \quad (3.16)$$

Since  $z \in \gamma_j$  and  $\delta_S(o) \geq 2^{3-j}S$ , it follows from (F1) and (F2) that

$$|z| \geq \int_{2^{2-j}S}^{\delta_S(o)} f(t) dt \geq \int_{2^{2-j}S}^{2^{3-j}S} f(t) dt \geq C^{-3} 2^{2-j}Sf(2^{-j}S). \quad (3.17)$$

Putting together (3.17) and (3.15), we see that  $|\gamma(t)| \geq t/C^6$ . This last inequality readily implies that  $|\gamma(t)| \geq c(t + |x|)$ , where  $c = 1/3C^6$ ; simply consider separately the possibilities  $t \geq |x|/2$  and  $t < |x|/2$ , and use the triangle inequality in the latter case. Thus we have proved the defining inequality for a strong escape path on the final segment of  $\gamma$  consisting of  $\gamma_j$  for all  $j \in \mathbb{N}$  such that  $2^{-j}S \leq \delta_S(o)/8$ . In particular the full path  $\gamma$  is a strong escape path for  $x$  if  $\delta_S(x) < \delta_S(o)/4$ .

Suppose instead that  $z \in \gamma_j$ , where  $2^{3-j}S > \delta_S(o)$ . According to (3.15),  $\text{len}_l(\gamma^j) \leq C^3 2^{-j+2}Sf(2^{-j}S)$ . Since  $2^{-j}S \gtrsim \delta_S(o)$ , it follows from this last inequality, (F1), and (F3) that  $\text{len}_l(\gamma^j) \leq C'$  for some  $C' \lesssim 1$ . It follows that if  $|x| \geq \rho := 2C'$ , then  $|z| \geq (t + |x|)/3$ .  $\square$

We cannot conclude in Theorem 3.14(a) that  $(X, l, o)$  has a strong  $c$ -escape property. For instance, if  $X \subset \mathbb{R}^2$  is given by the curve complement

$$\{(s, t) \mid s, t \in \mathbb{R}, s \geq 1, t \neq s^2 \sin^2 s\} \subset \mathbb{R}^2.$$

with the spherical metric attached,  $\partial_S X = \{\infty\}$ , and we use the standard flattening function  $f$ , then  $l = \mathcal{L}(\sigma, \partial_S X, f)$  is the inner Euclidean metric on  $X$ , and  $(X, l, o)$  fails to have any strong  $c$ -escape property, as we can see by taking  $x = ((2j+1)\pi/2, j^2)$  for sufficiently large  $j = j_c \in \mathbb{N}$ .

4. ARE SPHERICALIZATION AND FLATTENING INVERSE PROCESSES?

Throughout this section, we write  $\mathcal{A} = \mathcal{U} \cup \mathcal{B}$ , where  $\mathcal{U}$  is the class of all unbounded incomplete pointed length spaces  $(X, l, o)$  for which  $o$  has a strong escape property, and  $\mathcal{B}$  is the class of all bounded pointed length spaces  $(X, \sigma, o)$  with associated spherical data. Fixing a sphericalization function  $g$  and a flattening function  $f$ , we have seen that sphericalization associates an element of  $\mathcal{B}$  with each element of  $\mathcal{U}$ , and that flattening associates an element of  $\mathcal{U}$  with each element of  $\mathcal{B}$ . In each case let us call the associated space the *sf-dual* of the original space. More formally, we should talk of original and dual data since flattening requires spherical data to be specified, but we prefer to refer to the spherical data and the functions  $f, g$  only when needed. We also call a space a *bsf-dual* if it is bilipschitz equivalent to the sf-dual of some space.

We say that a space in  $\mathcal{A}$  is *sf-reflexive* if it is bilipschitz equivalent to its double sf-dual. For instance Euclidean space and the Riemann sphere (with north pole omitted) are sf-reflexive with respect to the standard sphericalization and flattening functions. It is clear that sf-reflexivity requires that the functions  $f, g$  satisfy some sort of duality relation; the desired condition, which we call *quasiduality* will be defined in §4.2. A necessary condition for a space to be sf-reflexive is that it is a bsf-dual. We now state the main theorem in this last section which says that being a bsf-dual is also a sufficient condition for a space to be sf-reflexive.

**Theorem 4.1.** *Suppose  $X$  is a space in the class  $\mathcal{A}$  and sf-duality is defined via quasidual functions  $f, g$ . Then the following are equivalent:*

- (a)  $X$  is sf-reflexive.
- (b)  $X$  is a bsf-dual.
- (c)  $(X, l, o)$  has a strong escape property (if  $(X, l, o)$  is in  $\mathcal{U}$ ) or  $(X, \sigma, o)$  has a strong reverse escape property (if  $(X, \sigma, o)$  is in  $\mathcal{B}$ ).

This result says in particular that the sf-dual of every space in  $\mathcal{A}$  is sf-reflexive. However not all spaces in  $\mathcal{A}$  are sf-reflexive. Consider for example the spaces  $(X_T, l, o)$  in Example 2.17 and the spaces  $(X_T, \sigma, o)$  in Example 3.7. According to Theorem 4.1, these spaces are sf-reflexive if and only if the full sequence  $(b_n/a_n)_{n=1}^\infty$  is bounded. For counterexamples in  $\overline{\mathbb{R}^n}$ , see Examples 2.14 and 2.32.

**4.2. Quasiduality.**

For a space to be sf-reflexive, with respect to a particular choice of sphericalization and flattening functions  $g$  and  $f$ , requires something of the geometry of the space as indicated in Theorem 4.1, but it also requires that  $g$  and  $f$  are in some sense well matched. For instance, we shall see that the “standard” sphericalization and flattening functions  $g(t) = 2/(1 + t^2)$ ,  $0 \leq t < \infty$ , and  $f(t) = \sec^2((\pi - t)/2)/2$ ,  $0 < t \leq d_0 := \pi$ , are quasidual. Since  $f(t)$  is comparable with  $t^{-2}$ , this is essentially a special case of the fact that for  $p, q > 1$ ,  $d_0 > 0$ ,  $g_p(t) := 1/(1 + t^p)$  is comparable with  $f_q(t) := t^{-q}$ ,  $0 < t \leq d_0$ , if and only if  $q = p/(p - 1)$ . Before we formally define the notion of quasiduality, let us consider Euclidean space and the Riemann sphere as motivational examples.

Let  $X = \mathbb{R}^n$ ,  $n > 1$ , with the Euclidean metric  $l$  attached, and let  $o$  be the usual origin. If  $\sigma = \mathfrak{S}(l, o, g)$  for some sphericalization function  $g$ , then the resulting space  $(X, \sigma)$  is topologically a sphere minus a single point at infinity  $w$  (the only element of the spherical boundary). Moreover rays from  $o$  are geodesic both for  $l$  and  $\sigma$ , and  $\delta_S(x) = \sigma(x, w) = \int_{|x|}^{\infty} g(s) ds$ . If we now define  $l' = \mathfrak{L}(\sigma, \{w\}, f)$  for some flattening function  $f$  whose  $d_0$  parameter is at least as large as  $\int_0^{\infty} g(s) ds$ , then

$$dl'(x) = f(\delta_S(x)) d\sigma(x) = f\left(\int_{|x|}^{\infty} g(s) ds\right) \cdot g(|x|) dl(x).$$

Thus  $l'$  is bilipschitz equivalent to  $l$  if and only if there is some constant  $C \geq 1$  such that

$$\frac{1}{C} \leq f\left(\int_t^{\infty} g(s) ds\right) \cdot g(t) \leq C, \quad 0 \leq t < \infty. \quad (4.3)$$

In the opposite direction, let us begin with  $X = \overline{\mathbb{R}^n} \setminus \{w\}$ ,  $n > 1$ , the Riemann sphere with the north pole removed, and let  $d_0 = \pi$ . Flattening and then sphericalizing using some flattening and sphericalization functions  $f : [0, d_0] \rightarrow (0, \infty)$  and  $g$ , respectively, we get a new metric  $\sigma'$  which is comparable to  $\sigma$  if and only if there is some constant  $C \geq 1$  such that

$$\frac{1}{C} \leq g\left(\int_t^{d_0} f(s) ds\right) \cdot f(t) \leq C, \quad 0 < t \leq d_0. \quad (4.4)$$

We leave the straightforward verification of the above statements to the reader.

We could use (4.3) and (4.4) as our definition of quasiduality, but we would like to replace them by quantitatively equivalent conditions that are simpler to verify and to use. Note that (S1) and (S2) tell us that  $\int_t^{\infty} g(s) ds$  is comparable with  $tg(t)$  when  $t \geq 1$ , so using also (F1), we see that (4.3) can be recast (at least for  $t \geq 1$ ) as

$$f(tg(t)) tg(t) \approx t, \quad 1 \leq t < \infty, \quad (4.5)$$

as long as  $f(tg(t))$  is always defined. Similarly (4.4) can be recast as

$$g(tf(t)) tf(t) \approx t, \quad 0 < t \leq d_0/2. \quad (4.6)$$

Our official definition of quasiduality, which we now give, is similar to this last pair of conditions, but modified to ensure that all function values are defined.

**Definition 4.7.** Suppose that  $C \geq 1$ ,  $d_0 > 0$ , and that  $f : (0, d_0] \rightarrow (0, \infty)$  and  $g : [0, \infty) \rightarrow (0, \infty)$  are continuous. Let  $F, G$  be defined by the equations

$$F(t) = \begin{cases} tf(t), & 0 < t < d_0, \\ d_0^2 f(d_0)/t, & d_0 \leq t. \end{cases} \quad (4.8)$$

$$G(t) = \begin{cases} g(1)/t, & 0 < t < 1, \\ tg(t), & 1 \leq t. \end{cases} \quad (4.9)$$

We say that  $f$  is a left  $C$ -quasidual of  $g$ , or that  $g$  is a right  $C$ -quasidual of  $f$  if  $F(G(t)) \in [t/C, Ct]$  for all  $0 < t < \infty$ . We say that  $f$  is a right  $C$ -quasidual of  $g$ , or that  $g$  is a left  $C$ -quasidual of  $f$ , if  $G(F(t)) \in [t/C, Ct]$  for all  $0 < t < \infty$ . We say that  $f$  and  $g$  are  $C$ -quasidual if  $f$  is a left and a right  $C$ -quasidual of  $g$ .

It is easy to see that quasiduality remains true (with a quantitatively controlled change in the parameter  $C$ ) if the sphericalization and flattening functions  $g$  and  $f$  are replaced by comparable functions, and  $d_0$  is replaced by a comparable positive number, for instance if the standard choices  $g(t) = 2/(1 + t^2)$ ,  $0 \leq t < \infty$ , and  $f(t) = \sec^2((\pi - t)/2)/2$ ,  $0 < t \leq \pi$ , are replaced by  $\tilde{g}(t) = 1 \wedge t^{-2}$ ,  $0 \leq t < \infty$ , and  $\tilde{f}(t) = t^{-2}$ ,  $0 < t \leq 1$ . Verifying quasiduality for  $\tilde{f}$  and  $\tilde{g}$  is trivial. More generally, if  $p, q > 1$ ,  $d_0 > 0$ , then  $g(t) = 1/(1 + t^p)$  is comparable with  $t^{-p}$  for  $1 \leq t < \infty$ , and so it is clear that  $g$  is quasidual to  $f(t) = t^{-q}$ ,  $0 < t \leq d_0$ , if and only if  $q = p/(p - 1)$ .

The statement of Theorem 4.1 leads naturally to the following question: do all sphericalizing and flattening functions have quasidual flattening and sphericalizing functions, respectively? We shall see that the answer is affirmative. In fact, as we shall see later, it is rather easy to prove that sphericalizing and flattening functions have one-sided quasiduals but this then leads to the following pair of questions. Are one-sided quasiduals of sphericalizing and flattening functions automatically two-sided quasiduals? Are quasiduals of sphericalizing and flattening functions automatically flattening and sphericalizing functions, respectively? We next state four lemmas which essentially give affirmative answers to these two questions.

**Lemma 4.10.** *Suppose that  $f : (0, d_0] \rightarrow (0, \infty)$  is a  $C_f$ -flattening function, and that  $g : [0, \infty) \rightarrow (0, \infty)$  is a left or right  $C_g$ -quasidual of  $f$ . Then  $g$  is a  $C$ -quasidual of  $g$  for some  $C = C(C_f, C_g)$ .*

**Lemma 4.11.** *Suppose that  $g$  is a  $C_g$ -sphericalizing function and that  $f : (0, d_0] \rightarrow (0, \infty)$  is a left or right  $C_f$ -quasidual of  $g$ . Then  $f$  is a  $C$ -quasidual of  $g$ , for some  $C = C(C_f, C_g)$ .*

**Lemma 4.12.** *Suppose that  $f : (0, d_0] \rightarrow (0, \infty)$  is a  $C_f$ -quasidual of a  $C_g$ -sphericalizing function  $g$ . Then  $f$  is a  $(C, d_0)$ -flattening function for some  $C = C(C_f, C_g)$ .*

**Lemma 4.13.** *Suppose that  $g$  is a  $C_g$ -quasidual of a  $(C_f, d_0)$ -flattening function  $f$ , and that  $g(r)/g(s) \in [1/C_g, C_g]$  for all  $0 \leq r, s \leq 1$ . Then  $g$  is a  $C$ -sphericalizing function for some  $C = C(C_f, C_g)$ .*

Note that comparability of the values of  $g$  on  $[0, 1]$  needs to be assumed in this last lemma since quasiduality tells us nothing about these values. The proof of Lemma 4.11 is very similar to that of Lemma 4.10, and the proof of Lemma 4.13 is very similar to that of Lemma 4.12, so we omit both of them.

Before proving Lemmas 4.10 and 4.12, it is useful to rewrite the basic assumptions (F1), (F3), (G1), and (G3) as decay estimates for the functions  $F$  and  $G$  in (4.8) and (4.9). Suppose

$f$  is a  $(C_f, d_0)$ -flattening function and  $g$  is a  $C_g$ -sphericalizing function. Then there are positive constants  $K_f, K_g, C'_f, C'_g, \epsilon_f, \epsilon_g$  such that

$$\frac{1}{C_f} \left(\frac{s}{r}\right)^{K_f} \leq \frac{F(r)}{F(s)} \leq C'_f \left(\frac{s}{r}\right)^{\epsilon_f}, \quad 0 < s \leq r < \infty \quad (4.14)$$

$$\frac{1}{C_g} \left(\frac{s}{r}\right)^{K_g} \leq \frac{G(r)}{G(s)} \leq C'_g \left(\frac{s}{r}\right)^{\epsilon_g}, \quad 0 < s \leq r < \infty \quad (4.15)$$

Indeed, we can pick  $K_f = \log_2 C_f$ ,  $K_g = \log_2 C_g$ , let  $C'_f$  and  $\epsilon_f$  be the constants  $C'$  and  $\epsilon$  in (F3), and let  $C'_g$  and  $\epsilon_g$  be the constants  $C'$  and  $\epsilon$  in (G3). We leave it to the reader to verify these estimates.

*Proof of Lemma 4.10.* Let  $F, G$  be defined by (4.8) and (4.9). Suppose that  $g$  is a right quasidual of  $f$ , and so  $F(G(t)) \approx t$ ,  $t > 0$ . Taking  $t = F(s)$ , we get that  $F(G(F(s))) \approx F(s)$  for all  $s > 0$ . Using (4.14) we see that if  $F(r) \approx F(r')$ , then  $r \approx r'$ , and so we deduce that  $G(F(s)) \approx s$  for all  $s > 0$ , as required.

Suppose instead that  $g$  is a left quasidual of  $f$ , and so  $G(F(t)) \approx t$ ,  $t > 0$ . Again by (4.14), we see that if  $t \approx t'$ , then  $F(t) \approx F(t')$ , and so  $F(G(F(t))) \approx F(t)$ . But it is clear from the definition of  $F$  and (F3) that  $F$  has range  $(0, \infty)$ , and so  $F(G(s)) \approx s$  for all  $s > 0$ .  $\square$

*Proof of Lemma 4.12.* Throughout this proof, we assume that  $0 < s \leq s' \leq d_0$ ,  $t, t' > 0$ ,  $G(t) = s$ , and  $G(t') = s'$ . Since  $G$  has range  $(0, \infty)$ , such numbers  $t, t' > 0$  exist regardless of the values of  $r, r'$ . Quasiduality implies the useful estimate

$$\frac{t'F(s)}{tF(s')} = \frac{t'F(G(t))}{tF(G(t'))} \approx 1. \quad (4.16)$$

To prove (F1), it suffices to show that if  $s, s'$  satisfy the additional constraint  $s' \leq 2s$ , then  $f(s) \approx f(s')$ , or equivalently  $F(s) \approx F(s')$ . Using (4.15), it follows that  $t \approx t'$  whenever  $s'/s \in [1, 2]$ . Thus using (4.16), we see that  $F(s) \approx F(s')$ , as required.

It remains to prove (F3), or equivalently  $F(s)/F(s') \gtrsim (s'/s)^\delta$  for some  $\delta > 0$ . This already follows from (F1) if  $s \approx s'$ , so we may assume that  $s'/s$  is so large that the second inequality in (4.15) implies that  $t' \leq t$ . But now by the first part of (4.15), we get  $t/t' \gtrsim (s'/s)^{1/K_g}$  and, combining this inequality with (4.16), we deduce that  $F(s)/F(s') \gtrsim (s'/s)^{1/K_g}$ .  $\square$

By *data dilation* (by a factor  $a > 0$ ), we shall mean replacing  $f$  by  $t \mapsto af(at)$ ,  $g$  by  $t \mapsto g(t)/a$ , and  $d_0$  by  $d_0/a$ . Data dilation by a factor  $a$  leads to the associated functions  $F$  and  $G$  being replaced by  $t \mapsto F(at)$  and  $t \mapsto a^{-1}G(t)$ , respectively. It follows that  $C$ -quasiduality is invariant under data dilations. Note also that the assumptions that  $g$  is a  $C$ -sphericalizing function and that  $f$  is a  $(C, d_0)$ -flattening function for some  $d_0 > 0$  are invariant under data dilations. Data dilation is used later to assume without loss of generality that  $d_0 = 1$ , or to normalize some value of  $f$  or  $g$ .

With the aid of the above lemmas, it is now a straightforward task to prove the following result which answers the remaining one of the three questions posed above.

**Proposition 4.17.** *Suppose  $d_0 > 0$ . Given a  $C_1$ -sphericalizing function  $g$ , there exists a  $(C_2, d_0)$ -flattening function  $f$  which is  $C_3$ -quasidual to  $g$ . If instead we are given a  $(C_1, d_0)$ -flattening function  $f$ , there exists a  $C_2$ -sphericalizing function  $g$  which is  $C_3$ -quasidual to  $f$ . For both statements  $C_2$  and  $C_3$  depend only on  $C_1$ .*

*Proof.* By data dilation, we may assume that  $g(1) = 1$ . Suppose that  $g$  is a  $C_1$ -sphericalizing function and that  $G$  is defined by (4.9). Since  $G$  is continuous with range  $(0, \infty)$  there exists, for any given  $t > 0$ , a least non-negative number  $s =: H(t)$  for which  $G(s) = t/d_0$ . Note that  $H(t) = d_0/t$  for all  $t \geq d_0$ , and  $H(t) > 1$  for all  $t < d_0$ . Now  $H$  is strictly decreasing but it might be discontinuous on  $(0, d_0]$ . To fix this, let  $F(t) = H(t)$  for all  $t \geq d_0$  and for  $t = 2^{-j}d_0$ ,  $j \in \mathbb{N}$ , and define  $F(t)$  by linear interpolation for all other  $t \in (0, d_0)$ . If we define  $f(t) = F(t)/t$  for  $0 < t \leq d_0$ , then  $f$  and  $F$  are related as in (4.8). It follows from (4.15) that the values of  $H$ , and hence those of  $f$ , are mutually comparable on each of the subintervals  $I_j = [2^{-j}d_0, 2^{-j+1}d_0]$ ,  $j \in \mathbb{N}$ . Using (S1), it now follows that  $f$  is a right quasidual of  $g$ . Using Lemmas 4.11 and 4.12, we deduce that  $f$  is both a quasidual of  $g$  and a flattening function.

We omit the analogous proof of the second part of this result.  $\square$

#### 4.18. Proof of Theorem 4.1.

We first prove that a space  $(X, l, o)$  in  $\mathcal{U}$  is sf-reflexive with respect to a quasidual pair  $f, g$  if and only if  $(X, l, o)$  has a strong escape property.

**Theorem 4.19.** *Let  $(X, l, o)$  be an incomplete pointed length space, and let  $\sigma := \mathfrak{S}(l, o, g)$ , where  $g$  is a  $C$ -sphericalizing function with a  $C$ -quasidual  $f : (0, d_0] \rightarrow (0, \infty)$ ,  $d_0 = \text{dia}_\sigma(X)$ , and  $\partial_S X$  is nonempty. Also let  $\partial_S X = \partial X_\sigma \setminus \partial X_l$ , and  $l' = \mathfrak{L}(\sigma, \partial_S X, f)$ . Then:*

- (a)  $l' \leq C'l$ , where  $C' = C'(C)$ .
- (b) If  $(X, l, o)$  has the strong  $(c_0, r_0)$ -escape property, then  $l' \geq cl$ , for some  $c = c(C, c_0, r_0) > 0$ .
- (c) If  $l' \geq cl$  for some  $c > 0$ , then  $(X, l, o)$  has the strong  $(c_0, r_0)$ -escape property, for some  $c_0, r_0$  dependent only on  $c$  and  $C$ .

*Proof.* Let  $F, G$  be as defined in (4.8) and (4.9). Propositions 2.6 and 2.20 ensure that  $d_0 \leq 4Cg(1)$  and that  $(X, \sigma, \partial_S X)$  are  $d_0$ -spherical data. Thus it makes sense to talk of  $l'$ . Note that Lemma 4.12 implies that  $f$  is a  $C''$ -flattening function for some  $C'' \lesssim 1$ . Since  $l'$  is the conformal deformation of  $\sigma$  with density function  $f(\delta_S(x))$ , and  $\sigma$  is the conformal deformation of  $l$  with density function  $g(|x|)$ , the conclusions in (a) and (b) are equivalent to the statements that  $f(\delta_S(x))g(|x|) \lesssim 1$  and  $f(\delta_S(x))g(|x|) \gtrsim 1$ , respectively. Since the hypotheses and conclusions are invariant under data dilation, we may assume that  $g(1) = 1$ . By Proposition 2.6 and (2.9), we see that  $d_0 \approx 1$ , and by quasiduality,  $f(d_0) \approx 1$ .

By (2.22) and (S2),  $\delta_S(x) \gtrsim G(|x|)$  for  $|x| \geq 1$  and  $\delta_S(x) \gtrsim g(1) = 1$  for  $|x| < 1$ . Using quasiduality, we see that  $f(\delta_S(x))g(|x|) \lesssim F(G(|x|))/|x| \approx 1$  for  $|x| \geq 1$ , and that  $f(\delta_S(x))g(|x|) \lesssim F(G(1)) \approx 1$  for  $|x| < 1$ . Thus we have proven (a).

Similarly, the conclusion of (b) follows once we prove that  $\delta_S(x) \lesssim 1$  for  $|x| \leq r_0$  and  $\delta_S(x) \lesssim G(|x|)$  for  $|x| \geq r_0$ . The first estimate is trivial since  $\delta_S(x) \leq d_0 \approx 1 \approx r_0$ . If instead  $|x| \geq r_0$ , then we choose a strong  $c_0$ -escape path for  $x$ , and (2.21) tells us that

$\text{len}_\sigma(\gamma) \lesssim G(|x|)$ . Since  $\gamma$  has finite  $\sigma$ -length but infinite  $l$ -length, it connects  $x$  to  $\partial_S X$ , and so  $\delta_S(x) \leq \text{len}_\sigma(\gamma) \lesssim G(|x|)$ .

Finally we prove (c). Suppose  $l' \gtrsim l$  and so  $f(\delta_S(x))g(|x|) \gtrsim 1$ . By quasiduality of  $f$  and  $g$ ,  $f(G(|x|))g(|x|) \approx 1$ ,  $|x| \geq 1$ , and (F3) ensures that there exists  $1 \leq r_0 \lesssim 1$  such that  $\delta_S(x) \lesssim G(|x|)$  for all  $|x| \geq r_0$ . Thus if  $|x| \geq r_0$ , we can find a path  $\lambda_0 : [0, T_0) \rightarrow X$ , parametrized by  $\sigma$ -arclength, such that  $\lambda_0(0) = x$ ,  $\lambda_0(T_0-; \sigma) = w_0 \in \partial_S X$ , and  $T_0 \lesssim g(|x|)|x|$ . Arguing as in Theorem 2.28, we get a strong escape path. We leave the details to the reader; note that the stopping condition never occurs since all paths  $\lambda_i$  are of infinite  $l$ -length.  $\square$

We next tackle the reverse direction: flattening followed by sphericalization.

**Theorem 4.20.** *Suppose that  $(X, \sigma, \partial_S X)$  are  $d_0$ -spherical data and that  $l := \mathfrak{L}(\sigma, \partial_S X, f)$ , where  $f$  is a  $(C, d_0)$ -flattening function with a  $C$ -quasidual  $g$ . Fix  $o \in X$  and define  $\sigma' := \mathfrak{S}(l, o, g)$ . Then*

- (a)  $\sigma' \leq C'\sigma$ , for some  $C'$  dependent only on  $C$ ,  $d_0 f(d_0)$ , and  $\delta_S(o)/d_0$ .
- (b) If  $(X, \sigma, o)$  satisfies a strong reverse  $c_0$ -escape property, then  $\sigma' \geq c\sigma$ , for some  $c = c(C, c_0, d_0 f(d_0))$ .
- (c) If  $\sigma' \geq c\sigma$ ,  $c > 0$ , then  $(X, \sigma, o)$  satisfies a strong reverse  $c_0$ -escape property, for some  $c_0 = c_0(C, c, d_0 f(d_0))$ .

*Proof.* According to Lemma 4.13,  $g$  is a  $C''$ -sphericalizing function for some  $C'' = C''(C)$ . We write  $|x| = l(x, o)$  as usual, and also write  $tB_y = B_\sigma(y, t\delta_S(y)/4)$ , for all  $y \in X$ ,  $t > 0$ . The combination of data dilation by a factor  $a$  and dilation of  $\sigma$ -distances by a factor  $a^{-1}$  leaves both the assumptions and the conclusions invariant, so we may assume that  $d_0 = 1$ . Thus  $F(1) = f(1)$  and, because of the dependence of all conclusions on  $d_0 f(d_0)$ , we may assume that  $F(1) \approx 1$  throughout.

By quasiduality, it follows that

$$g(F(\delta_S(x)))f(\delta_S(x)) \approx 1. \quad (4.21)$$

In fact this is immediate when  $F(\delta_S(x)) \geq 1$ , and it holds in general because  $F(\delta_S(x)) \gtrsim F(1) \approx 1$ . But, as in the proof of the previous theorem, the conclusions in (a) and (b) are equivalent to the statements that  $g(|x|)f(\delta_S(x)) \lesssim 1$  and  $g(|x|)f(\delta_S(x)) \gtrsim 1$ , respectively. In view of (4.21), (S1), and (S3), the conclusions in (a) and (b) are thus equivalent to the statements that  $F(\delta_S(x)) \lesssim |x| \vee 1$  and  $F(\delta_S(x)) \gtrsim |x| \vee 1$ , respectively.

Since  $f$  is flattening, the values  $f(\delta_S(z))$ ,  $z \in 2B_x$ , are all mutually comparable. Therefore the  $l$ -arclength of a path in any such ball is comparable to  $f(\delta_S(x))$  times its  $\sigma$ -arclength. Thus if we define  $r_x$  to be the largest radius with the property that  $B_x \supset B_l(x, r_x)$ , and  $R_x$  to be the smallest radius with the property that  $2B_x \subset B_l(x, R_x)$ , then  $r_x \approx R_x \approx F(\delta_S(x))$ .

Let us now prove (a). If  $x \in X$ ,  $|x| \geq R_o$ , then  $x \notin 2B_o$ . Thus  $\delta_S(x) \leq \sigma(x, o) + \delta_S(o) \leq 3\sigma(x, o)$ , and so  $o \notin B_x$ . Thus  $|x| \geq r_x \approx F(\delta_S(x))$ . It remains to show that  $F(\delta_S(x)) \lesssim |x| \vee 1$  when  $|x| < R_o$ . Since  $R_o \lesssim F(\delta_S(o)) \approx 1$ , this follows from the case  $|x| = R_o$  (which we already know) and (F1), once we show that  $\delta_S(x)$  is approximately constant on  $\overline{B}_l(o, R_o)$ .



Certainly  $\delta_S(x) \leq d_0 = 1 \approx \delta_S(o)$ . If  $\delta_S(x) < t$  for some  $t$  much smaller than  $\delta_S(x)$ , then we can apply Lemmas 2.5 and 3.3 to deduce that

$$|x| \geq \int_t^{\delta_S(o)} f(s) ds \geq K\delta_S(o)f(\delta_S(o)) \approx K,$$

where  $K \rightarrow \infty$  as  $t/\delta_S(o) \rightarrow 0$ . But  $|x| \leq R_o$ , so the resulting upper bound on  $K$  translates into a lower bound on  $t$ , as required.

We now turn to (b). Let  $\gamma$  be the path from  $x$  to  $o$  provided by the strong reverse escape property. Since  $S := \text{len}_\sigma(\gamma) \leq \delta_S(o)/c_0 - \delta_S(x) \leq 1/c_0 - \delta_S(x)$ , we see by a change of variables, (F1), (F3), and (F2) that

$$\begin{aligned} \text{len}_l(\gamma) &= \int_0^S f(\delta_S(\gamma(t))) dt \lesssim \int_0^S f(c_0(t + \delta_S(x))) dt \\ &= \int_{c_0\delta_S(x)}^1 f(s) ds \\ &\lesssim F(\delta_S(x)). \end{aligned} \tag{4.22}$$

Thus  $|x| \lesssim F(\delta_S(x))$  and (b) follows.

Finally, we prove (c). The hypothesis  $\sigma' \gtrsim \sigma$  is, as we have seen, equivalent to the statement that  $F(\delta_S(x)) \gtrsim |x| \vee 1$ . Thus we can find a path  $\lambda_0 : [0, T_0] \rightarrow X$ , parametrized by  $l$ -arclength, such that  $\lambda_0(0) = x$ ,  $\lambda_0(T_0) = o$ , and  $T_0 \lesssim F(\delta_S(x))$ . Arguing as in Theorem 3.11, we get a strong reverse escape property for  $(X, \sigma, o)$ ; the details are left to the reader.  $\square$

It is now straightforward to complete the proof of Theorem 4.1. The fact that (a) implies (b) is trivial, and (b) implies (c) according to Theorems 2.31 and 3.14. Finally, (c) implies (a) according to Theorems 4.19 and 4.20.

## REFERENCES

- [AG] K. Astala and F. W. Gehring, *Quasiconformal analogues of theorems of Koebe and Hardy-Littlewood*, Mich. Math. J. **32** (1985), 99–107.
- [AK] K. Astala and P. Koskela, *Quasiconformal mappings and global integrability of the derivative*, J. Anal. Math. **57** (1991), 99–107.
- [BB] Z. M. Balogh and S.M. Buckley, *Geometric characterizations of Gromov hyperbolicity*, Invent. Math **153** (2003), 261–301.
- [BK] Z. M. Balogh and P. Koskela, *Quasiconformality, quasisymmetry and removability in Loewner spaces*, Duke Math. J. **101** (2000), 554–577.
- [BHK] M. Bonk, J. Heinonen, and P. Koskela, *Uniformizing Gromov hyperbolic spaces*, Astérisque **270** (2001).
- [BHR] M. Bonk, J. Heinonen, and S. Rohde, *Doubling conformal densities*, J. reine angew. Math. **541** (2001), 117–141.
- [BKo] M. Bonk and P. Koskela, *Conformal metrics and the size of the boundary*, Amer. J. Math. **124** (2002), 1247–1287.
- [BKR] M. Bonk, P. Koskela, and S. Rohde, *Conformal metrics on the unit ball in Euclidean space*, Proc. London Math. Soc. **77** (1998), 635–664.

- [HK] J. Heinonen and P. Koskela, *Quasiconformal maps in metric spaces with controlled geometry*, Acta Math. **181** (1998), 1–61.
- [He1] D. Herron, *Quasiconformal deformations and volume growth*, preprint.
- [He2] D. Herron, *Conformal deformations of uniform Loewner spaces*, Math. Proc. Camb. Phil. Soc. **136** (2004), 325–360.
- [Se1] S. Semmes, *Bilipschitz mappings and strong  $A_\infty$  weights*, Ann. Acad. Sci. Fenn. Ser. A. I. Math. **18** (1993), 211–248.
- [Se2] S. Semmes, *On the nonexistence of bilipschitz parametrization and geometric problems about  $A_\infty$  weights*, Rev. Math. Iberoamericana **12** (1996), 345–360.
- [Ty1] J. T. Tyson, *Quasisymmetry and quasiconformality in metric measure spaces*, Ann. Acad. Sci. Fenn. Math. **23** (1998), 525–548.
- [Ty2] J. T. Tyson, *Metric and geometric quasiconformality in Ahlfors regular Loewner spaces*, Conform. Geom. Dyn. **5** (2001), 21–73.
- [V] J. Väisälä, *Lectures on  $n$ -dimensional quasiconformal mappings*, Lecture Notes in Mathematics 229, Springer-Verlag, Berlin, 1971.

DEPARTMENT MATHEMATIK, UNIVERSITÄT BERN, SIDLERSTRASSE 5, 3012, BERN, SCHWEIZ.

*E-mail address:* `zoltan@math-stat.unibe.ch`

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF IRELAND, MAYNOOTH, CO. KILDARE, IRELAND.

*E-mail address:* `sbuckley@maths.may.ie`