

The p -Laplace eigenvalue problem as $p \rightarrow 1$ and Cheeger sets in a Finsler metric*

B. Kawohl M. Novaga

Dedicated to the memory of Thomas Lachand-Robert

Abstract

We consider the p -Laplacian operator on a domain equipped with a Finsler metric. After deriving and recalling relevant properties of its first eigenfunction for $p > 1$, we investigate the limit problem as $p \rightarrow 1$.

Keywords: p -Laplace, eigenfunction, Finsler metric, Cheeger set, anisotropic isoperimetric inequality

Math. Sci. Classification: 35J20, 35J70, 49A52, 49G05, 52A05, 52A40

1 Introduction

Imagine a nonlinear elastic membrane, fixed on a boundary $\partial\Omega$ of a plane domain Ω . If $u(x)$ denotes its vertical displacement, and if its deformation energy is given by $\int_{\Omega} |\nabla u|^p dx$, then a minimizer of the Rayleigh quotient

$$\frac{\int_{\Omega} |\nabla u|^p dx}{\int_{\Omega} |u|^p dx}$$

on $W_0^{1,p}(\Omega)$ satisfies the Euler-Lagrange equation

$$-\Delta_p u = \lambda_p |u|^{p-2} u \quad \text{in } \Omega, \tag{1.1}$$

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is the well-known p -Laplace operator. This eigenvalue problem has been extensively studied in the literature. As $p \rightarrow 1$, formally the limit equation reads

$$\begin{aligned} -\operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right) &= \lambda_1(\Omega) && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega. \end{aligned} \tag{1.2}$$

*SHORT TITLE: p -Laplace eigenvalue problem in a Finsler metric

For a precise interpretation of (1.2) see [21] or [30]. Naturally, here $\lambda_1(\Omega) := \lim_{p \rightarrow 1+} \lambda_p(\Omega)$. A somewhat surprising recent result is that the family of eigenfunctions $\{u_p\}$ converges in L^1 cum grano salis to (a multiple of) the characteristic function χ_{C_Ω} of a subset C_Ω of Ω , a so called Cheeger-set, see [19]. A Cheeger set of Ω is characterized as a domain that minimizes

$$h(\Omega) := \inf_D \frac{|\partial D|}{|D|}$$

with D varying over all smooth subdomains of Ω whose boundary ∂D does not touch $\partial\Omega$, and with $|\partial D|$ and $|D|$ denoting $(n-1)$ - and n -dimensional Lebesgue measure of ∂D and D . The existence, uniqueness, regularity and construction of such sets is discussed in [19] and [20] and its continuous dependence on Ω in [16]. The paper [23] contains a numerical method for the calculation of n -dimensional Cheeger sets and some three-dimensional examples. Cheeger sets are of significant importance in the modelling of landslides, see [17], [18], or in fracture mechanics, see [22]. Notice that a set $D \subseteq \Omega$ is a Cheeger set if and only if it is a minimizer of

$$|\partial E| - h(\Omega)|E| \quad \text{for } E \subseteq \Omega. \quad (1.3)$$

Now suppose that the membrane is not isotropic. It is for instance woven out of elastic strings like a piece of material. Then the deformation energy can be anisotropic, see [5]. Another way to describe this effect is by stating that the Euclidean distance in Ω is somehow distorted. It is the purpose of the present paper to generalize the above result on eigenfunctions and their convergence as $p \rightarrow 1$ to the situation, where $\Omega \subset \mathbb{R}^n$ is no longer equipped with the Euclidean norm, but instead with a general norm ϕ . In that case a Lipschitz continuous function $u : \Omega \mapsto \mathbb{R}$ (in a convex domain Ω) has Lipschitz constant $L = \sup_{z \in \Omega} \phi^*(\nabla u(z))$, where ϕ^* denotes the dual norm to ϕ . Therefore the Rayleigh quotient studied in this paper is given by

$$R_p(u) := \frac{\int_{\Omega} (\phi^*(\nabla u))^p dx}{\int_{\Omega} |u|^p dx} \quad (1.4)$$

on $W_0^{1,p}(\Omega)$ and the Cheeger constant by

$$h(\Omega) := \inf_{D \subset \Omega} \frac{P_\phi(D)}{|D|}, \quad (1.5)$$

with P_ϕ denoting anisotropic perimeter in \mathbb{R}^n (see (2.10) below). The minimizer u_p of R_p satisfies the Euler-Lagrange equation

$$-Q_p u := -\operatorname{div} \left((\phi^*(\nabla u))^{p-2} J(\nabla u) \right) \ni \lambda_p |u|^{p-2} u \quad \text{in } \Omega \quad (1.6)$$

in the weak sense [8], i.e.

$$\int_{\Omega} (\phi^*(\nabla u_p))^{p-2} \langle \eta, \nabla v \rangle dx = \lambda_p \int_{\Omega} |u_p|^{p-2} u_p \cdot v dx \quad (1.7)$$

for any $v \in W_0^{1,p}(\Omega)$ and for a measurable selection $\eta \in J(\nabla u_p)$, where the function $J : \mathbb{R}^n \rightarrow \mathcal{P}(\mathbb{R}^n)$ is defined as the subdifferential

$$J(\xi) := \partial \left(\frac{\phi^*(\xi)^2}{2} \right). \quad (1.8)$$

Note that the function J is single-valued iff the norm ϕ is strictly convex, i.e. if its unit sphere $\{x : \phi(x) = 1\}$ contains no nontrivial line segments [34, pag. 400]. Note further that $J(0) = 0$ and that for the Euclidean norm the duality map reduces to the identity $J(\nabla u) = \nabla u$.

The paper is organized as follows. In Section 2 we fix some notation. In Section 3 we recall and derive the existence, uniqueness, regularity and log-concavity of solutions for $p > 1$. In Section 4 we derive the limit equation for $p \rightarrow 1$. In Section 5, we discuss in detail the two-dimensional case, proving uniqueness of Cheeger sets in the convex case. In Section 6 we provide some instructive examples.

2 Notation

We say that the norm ϕ is *regular* if $\phi^2, (\phi^*)^2 \in C^2(\mathbb{R}^n)$. This includes for instance $\phi(x) = \|x\|_q$ with $q \in (1, \infty)$ but excludes the crystalline cases $q = 1$ or $q = \infty$, see Section 6.

Given $E \subset \mathbb{R}^n$ and $x \in \mathbb{R}^n$, we set

$$\text{dist}_{\phi}(x, E) := \inf_{y \in E} \phi(x - y), \quad d_{\phi}^E(x) := \text{dist}_{\phi}(x, E) - \text{dist}_{\phi}(\mathbb{R}^n \setminus E, x).$$

Notice that, at each point where d_{ϕ}^E is differentiable, there holds

$$\phi^*(\nabla d_{\phi}^E) = 1. \quad (2.9)$$

Let us define the (anisotropic) perimeter of E as

$$P_{\phi}(E) := \sup \left\{ \int_E \text{div} \eta dx \mid \eta \in C_c^1(\mathbb{R}^n), \phi(\eta) \leq 1 \right\} = \int_{\partial^* E} \phi^*(\nu^E) d\mathcal{H}^{n-1}, \quad (2.10)$$

where $\partial^* E$ and ν^E denote the reduced boundary of E and the (Euclidean) unit normal to $\partial^* E$.

Given an open set $\Omega \subseteq \mathbb{R}^n$ we define the BV -seminorm of $v \in BV(\Omega)$ as

$$\int_{\Omega} \phi^*(Dv) := \sup \left\{ \int_{\Omega} v \operatorname{div} \eta \, dx \mid \eta \in C_c^1(\mathbb{R}^n), \phi(\eta) \leq 1 \right\}.$$

Given $\delta > 0$, we define

$$\begin{aligned} E_+^\delta &:= \left\{ x \in \mathbb{R}^n \mid d_\phi^E < \delta \right\} = E + \delta W_\phi, \\ E_-^\delta &:= \left\{ x \in \mathbb{R}^n \mid d_\phi^E > -\delta \right\}, \\ E_\pm^\delta &:= \left(E_\pm^\delta \right)_+^\delta \subseteq E, \end{aligned}$$

where $W_\phi := \{x \mid \phi(x) < 1\}$, also called *Wulff shape*, denotes the unit ball with respect to the norm ϕ .

Given a compact set $E \subset \mathbb{R}^n$ with Lipschitz boundary, we denote by $n_\phi : \partial E \rightarrow \mathbb{R}^n$ any Lipschitz vector field satisfying $n_\phi \in J(\nabla d_\phi^E)$ a.e. on ∂E . Moreover, we set

$$\|\kappa_\phi\|_{L^\infty(\partial E)} := \inf_{n_\phi \in J(\nabla d_\phi^E)} \|\operatorname{div}_\tau n_\phi\|_{L^\infty(\partial E)},$$

which represents the L^∞ -norm of the ϕ -mean curvature of ∂E . We make the convention that $\|\kappa_\phi\|_{L^\infty(\partial E)} = +\infty$ if the set E does not admit any Lipschitz vector field $n_\phi \in J(\nabla d_\phi^E)$. We say that E is ϕ -regular if $\|\kappa_\phi\|_{L^\infty(\partial E)} < +\infty$.

Notice that in the Euclidean case E is ϕ -regular iff ∂E is of class $C^{1,1}$. Moreover, the unit ball W_ϕ is always ϕ -regular and $\|\kappa_\phi\|_{L^\infty(\partial W_\phi)} = n - 1$. To see this, it is enough to consider the vector field $n_\phi(x) = x/\phi(x)$.

3 Existence, uniqueness, regularity and log-concavity of solutions

Let $\Omega \subset \mathbb{R}^n$ be a bounded open set. If we minimize the functional

$$I_p(v) = \int_{\Omega} \phi^*(\nabla v)^p \, dx \quad \text{on} \quad K := \{ v \in W_0^{1,p}(\Omega) \mid \|v\|_{L^p(\Omega)} = 1 \}, \quad (3.1)$$

then via standard arguments (see [6]) a minimizer u_p exists for every $p > 1$ and it is a weak solution to the equation (1.6), with $\lambda_p = I_p(u_p)$. Note that $\Lambda_p := I_p(u_p)^{1/p}$ is the minimum of the Rayleigh quotient

$$R_p(v) := \frac{\left(\int_{\Omega} (\phi^*(\nabla v))^p \, dx \right)^{1/p}}{\|v\|_p} \quad (3.2)$$

on $W_0^{1,p}(\Omega) \setminus \{0\}$. Without loss of generality we may assume that u_p is non-negative. Otherwise we can replace it by its modulus.

Moreover, as shown in [6] any nonnegative weak solution of (1.6) is necessarily bounded and positive in Ω . If $p > n$, then u_p is also uniformly Hölder continuous because of the Sobolev-embedding theorem and the equivalence of the usual Sobolev norm with

$$\|u\|_{1,p} := \left(\int_{\Omega} |u|^p dx \right)^{1/p} + \left(\int_{\Omega} (\phi^*(\nabla u))^p dx \right)^{1/p}. \quad (3.3)$$

If the norm ϕ is regular and $p > 1$, one can even show that $u_p \in C^{1,\alpha}(\Omega)$. Indeed, the function u_p minimizes

$$J_p(v) := \int_{\Omega} (\phi^*(\nabla v))^p - \lambda_p(\Omega)|u|^p dx,$$

and the theory for quasiminima in [15] implies that minimizers are bounded (Thm. 7.5), Hölder continuous (Thm. 7.16) and satisfy a strong maximum principle (Thm. 7.12), because one can easily check that u_p satisfies (7.71) in [15]. Therefore u_p is positive. Once positivity is known, the uniqueness follows from a simple convexity argument, see [4] or [6]. Moreover, from the result in [11] one can conclude that $u_p \in C^{0,\beta}(\Omega)$ for any $\beta \in (0, 1)$. Finally, if ϕ is regular, then $u_p \in C^{1,\alpha}(\Omega)$ according to [7], [25], [32], [33] or [12]. Let us summarize these statements.

Theorem 3.1. *For every $p \in (1, \infty)$ the nonnegative minimizer u_p of (3.1) is positive, unique, belongs to $C^{0,\beta}(\Omega)$ for any $\beta \in (0, 1)$ and it solves (1.6) in the weak sense. Moreover, if the norm ϕ is regular then u_p is of class $C^{1,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$. Finally, if Ω is convex, then u_p is log-concave and the level sets set $\{u_p > t\} \subseteq \Omega$ are convex for all $t > 0$.*

To prove the last statement, we follow Sakaguchi's approach from [28], first for strictly convex Ω and for a smooth norm ϕ . The general case follows then from approximation arguments for Ω and ϕ . Log-concavity of a sequence $u_{p,n}$ is preserved under pointwise limits as $n \rightarrow \infty$, because the inequality

$$\log u_{p,n} \left(\frac{x_1 + x_2}{2} \right) \geq \frac{1}{2} \log u_{p,n}(x_1) + \frac{1}{2} \log u_{p,n}(x_2) \quad \text{in } \Omega \times \Omega$$

is stable under such limits. If u_p solves (1.6), then $v_p := \log u_p$ solves

$$-\operatorname{div} \left((\phi^*(\nabla v))^{p-2} J(\nabla v) \right) = (p-1)\phi^*(\nabla v)^p + \lambda_p \quad \text{in } \Omega \quad (3.4)$$

and this degenerate elliptic equation can be approximated by a nondegenerate one

$$-\operatorname{div} \left((\varepsilon + (\phi^*(\nabla v))^2)^{\frac{p-2}{2}} J(\nabla v) \right)$$

$$= (p - 1 - \varepsilon)(\phi^*(\nabla v))^2(\varepsilon + (\phi^*(\nabla v))^2)^{\frac{p-2}{2}} + \lambda_p. \quad (3.5)$$

Modulo yet another approximation by a right hand side which is strictly monotone in v , equation (3.5) is now amenable to Korevaar's concavity maximum principle which states that the concavity function

$$C(x_1, x_2) := v\left(\frac{x_1 + x_2}{2}\right) - \frac{1}{2}v(x_1) - \frac{1}{2}v(x_2) \quad \in \Omega \times \Omega$$

can attain a negative minimum only on the boundary of $\Omega \times \Omega$. The latter is ruled out, however, because of the boundary condition.

Remark 3.2. *We should point out that without uniqueness of u_p the approximation arguments would only yield log-concavity of a solution and not the solution u_p .*

4 The limit eigenvalue and eigenfunction for $p \rightarrow 1$

The following estimate for λ_p is optimal (as $p \rightarrow 1$) for any shape of Ω (see [6]).

Theorem 4.1. (Cheeger type inequality) *For every $p \in (1, \infty)$ the eigenvalue $\lambda_p(\Omega)$ can be estimated from below as follows:*

$$\lambda_p(\Omega) \geq \left(\frac{h(\Omega)}{p}\right)^p. \quad (4.1)$$

Here $h(\Omega)$ is the Cheeger constant of Ω as defined in (1.5). Moreover, as $p \rightarrow 1$, the eigenvalue $\lambda_p(\Omega)$ converges to $h(\Omega)$.

In the Euclidean case this is Cheeger's original estimate [10] when $p = 2$, and for general p it can be found in [24], [2], [26] and [31]. For a more general ϕ one can easily modify their proofs by using the generalized coarea formula from [13] or [14]. To prove the limiting behaviour of $\lambda_p(\Omega)$ as $p \rightarrow 1$ we proceed as in [19] and observe that (4.1) implies $\liminf_{p \rightarrow 1} \lambda_p(\Omega) \geq h(\Omega)$. Therefore it suffices to find a suitable upper bound. Let $\{D_k\}_{k=1,2,\dots}$ be a sequence of regular domains for which $P_\phi(D_k)/|D_k|$ converges to $h(\Omega)$. We approximate the characteristic function of each D_k by a function w_k with the following properties: $w \equiv 1$ on $\overline{D_k}$, $w \equiv 0$ outside an ε -neighborhood of D_k and $\phi^*(\nabla w_k) = 1/\varepsilon$ in an ε -layer outside D_k . For small ε the function w_k is in $W_0^{1,\infty}(\Omega)$ and provides the upper bound

$$\lambda_p(\Omega) \leq \frac{P_\phi(D_k)}{|D_k|} (\alpha\varepsilon)^{1-p}. \quad (4.2)$$

Now one sends first $p \rightarrow 1$, then $k \rightarrow \infty$ to complete the proof of Theorem 4.1.

Let us normalize the eigenfunctions u_p to have L^1 -norm equal to 1 and study the sequence (λ_p, u_p) of eigenvalues and normalized eigenfunctions as $p \rightarrow 1$. For every $p > 1$ the function u_p minimizes

$$J_p(v) := \int_{\Omega} (\phi^*(\nabla v))^p - \lambda_p(\Omega)|u|^p dx$$

on $X_p = \{v \in W_0^{1,p}(\Omega); v \geq 0 \text{ in } \Omega; \|v\|_1 = 1\}$. If one extends J_p to $BV(\Omega)$ by setting it ∞ on $BV \setminus X_p$, the family J_p can be shown to Γ -converge to

$$J_1(v) := \int_{\Omega} \phi^*(Dv) - \lambda_1(\Omega) \int_{\Omega} |v| dx$$

on $X_1 := \{v \in BV(\Omega); v \geq 0 \text{ in } \Omega; \|v\|_1 = 1\}$. Notice that $J_1 \geq 0$ on X_1 . Moreover u_p forms a minimizing sequence for J_1 since

$$\begin{aligned} J_1(u_p) &= \int_{\Omega} \phi^*(\nabla u_p) - \lambda_1 \int_{\Omega} |u_p| dx \\ &\leq \left[\int_{\Omega} (\phi^*(\nabla u_p))^p \right]^{\frac{1}{p}} |\Omega|^{\frac{p-1}{p}} - \lambda_1 \int_{\Omega} |u_p| dx \\ &\leq \frac{1}{p} \int_{\Omega} (\phi^*(\nabla u_p))^p dx + \frac{p-1}{p} |\Omega| \\ &\quad + (\lambda_p - \lambda_1) \int_{\Omega} |u_p| dx - \lambda_p \int_{\Omega} |u_p|^p dx \\ &\leq J_p(u_p) + \frac{p-1}{p} |\Omega| + (\lambda_p - \lambda_1) \int_{\Omega} |u_p| dx \\ &\leq p-1 + \lambda_p - \lambda_1. \end{aligned} \tag{4.3}$$

Here we have used the fact that $|u_p| \leq 1$. Hence $J_1(u_p) \rightarrow 0$ as $p \rightarrow 1$.

As a consequence, the family $\{u_p\}_{p>1}$ is bounded in $BV(\Omega)$ and, after possibly passing to a subsequence, it converges strongly in L^1 to a limit function $u_1 \in X_1$ such that $J_1(u_1) = 0$. Using the coarea formula, one can see that for all $t \in [0, \max_{\Omega} u_1)$ the level set $\Omega_t := \{u_1 > t\}$ is a Cheeger set. Thus we have shown

Theorem 4.2. (Convergence of eigenfunctions) *As $p \rightarrow 1$, a subsequence converges to a limit function u_1 in X_1 , and almost all level sets $\Omega_t := \{u_1 > t\}$ of u_1 are Cheeger sets.*

Remark 4.3. *As a consequence of Theorem 4.2 and the logconcavity of u_p , for convex Ω there exists a convex Cheeger set. Moreover, it follows from the results of [9] that there exists a convex Cheeger set $D \subseteq \Omega$ which is maximal, in the sense that any other Cheeger set of Ω must be contained in D . The uniqueness of Cheeger sets is in general not true for nonconvex domains [20], and an open problem for convex domains in dimension $n > 2$.*




Figure 1: The Cheeger sets D , \bar{D} of Theorem 5.1.

5 The planar case

In this section we derive further properties of the function u_1 , under the additional assumption $n = 2$. Let us begin with the following theorem, which extends the analogous result in the Euclidean case [20, Th. 1].

Theorem 5.1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded open convex set. Then, there exists a unique Cheeger set $D \subseteq \Omega$. Moreover, D is convex and we have*

$$h(\Omega) = \frac{1}{t^*}, \quad D = \Omega_{\pm}^{t^*}, \quad (5.1)$$

where $t^* > 0$ is the (unique) value t such that $|\Omega_-^t| = t^2 |W_\phi|$.

Proof. Let D be a Cheeger set of Ω . Notice first that D is a convex set, since otherwise we could replace it by its convex hull and reduce (1.3) (see [3, Th. 7.1]). Moreover, from the first variation of (1.3) it follows that the anisotropic curvature of ∂D is bounded by $h(\Omega)$, and each connected component of $\partial D \cap \Omega$ is contained up to translation in $\frac{1}{h(\Omega)} \partial W_\phi$ (see [27, Theorem 4.5]). Let \bar{D} be the open maximal Cheeger set of Ω (recall Remark 4.3), and let $\Gamma \subset \frac{1}{h(\Omega)} \partial W_\phi$ be a connected component of $\partial D \cap \bar{D}$. We denote by $x, y \in \Gamma \cap \partial \bar{D}$ the extremal points of Γ , and we let Γ' be the arc of $\partial \bar{D}$ with extrema x, y and lying in the same halfplane of Γ with respect to the straight line r passing through x, y (see Figure 1). Reasoning as in [3, Lemma 7.3], it is easy to show that both Γ and Γ' can be written as graphs on r along some directions. More precisely, there exists a vector $v \in \mathbb{R}^2$, with $|v| = 1$, and two functions $f_1, f_2 : r \rightarrow \mathbb{R}$ such that $0 \leq f_1 \leq f_2$ on $[x, y]$, that $\min\{f_2(x), f_2(y)\} = 0$,

and that $\Gamma = F_1([x, y])$ and $\Gamma' = F_2([x, y])$, with $F_i(x) := f_i(x)v$, for $i = 1, 2$. Without loss of generality, we shall assume that $v \perp r$. Since D and \overline{D} are both minimizers of (1.3), it follows that both f_1 and f_2 are minimizers of

$$G(f) := \int_{[x, y]} \phi^*(-f'(s), 1) - h(\Omega)f(s) \, ds. \quad (5.2)$$

If ϕ is a regular norm, then the functional G is strictly convex, which implies $f_1 = f_2$, i.e. $D = \overline{D}$. For a general norm, one has to be more careful, since the functional G is not strictly convex, but only convex. However, reasoning as in [3, Lemma 8.2], the inclusion $\Gamma \subset \frac{1}{h(\Omega)}\partial W_\phi$ and the inequality $f_1 \leq f_2$ imply $\|\kappa_\phi\|_{L^\infty(\Gamma')} \geq h(\Omega)$, with equality iff $\Gamma = \Gamma'$, which proves the uniqueness of the Cheeger set D .

Let us now prove (5.1), reasoning as in [20, Th. 1]. It has been proved in [3] that the convex set $D = \Omega_{\pm}^{1/h(\Omega)}$ is a Cheeger set of Ω , hence it is the unique Cheeger set of Ω . Therefore, it remains to prove that $t^* = 1/h(\Omega)$, i.e.

$$|\Omega_{-}^{\frac{1}{h(\Omega)}}| = \frac{|W_\phi|}{h(\Omega)^2}.$$

Let us recall from [1, Section 2.7],[29] the following Steiner-type formulae

$$\begin{aligned} |C^\delta| &= |C| + \delta P_\phi(C) + \delta^2 |W_\phi|, \\ P_\phi(C^\delta) &= P_\phi(C) + \delta P_\phi(W_\phi). \end{aligned} \quad (5.3)$$

Incidentally, the second equation follows from the first one and, as in the Euclidean case, $P_\phi(W_\phi) = 2|W_\phi|$. This follows from integrating $\operatorname{div} x$ on W_ϕ . Applying (5.3) to $C = D_{-}^{1/h(\Omega)}$ and recalling that $h(\Omega) = P_\phi(D)/|D|$, we get

$$|D_{-}^{1/h(\Omega)}| = \frac{|W_\phi|}{h(\Omega)^2}.$$

The claim now follows if we observe that

$$\Omega_{-}^{\frac{1}{h(\Omega)}} = D_{-}^{\frac{1}{h(\Omega)}}.$$

□

Corollary 5.2. *If $n = 2$ and Ω is a bounded convex set, then the sequence of functions u_p converges to a multiple of the characteristic function of D . Moreover, $D = \Omega$ if and only if*

$$\|\kappa_\phi\|_{L^\infty(\partial\Omega)} \leq h(\Omega). \quad (5.4)$$

In particular, (5.4) always holds in the case $\Omega = W_\phi$.

6 Example and concluding remarks

If the norm under consideration for $x \in \Omega$ is the usual ℓ_q - norm, i.e. for $\phi_q(x) = (\sum_{i=1}^n |x_i|^q)^{1/q}$, $q \geq 1$. When $q > 1$, the dual norm of ϕ_q is given by $\phi_q^* = \phi_{q'}$, with $q' = q/(q-1)$, and the duality map according to (1.8) is

$$J_i(y) = (|y|_{q'})^{2-q'} |y_i|^{q'-2} y_i.$$

Then the p -Laplace operator in this metric is given by (see [6])

$$Q_{p,q}u = \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\phi_{q'}(\nabla u)^{p-q'} \left| \frac{\partial u}{\partial x_i} \right|^{q'-2} \frac{\partial u}{\partial x_i} \right),$$

and for $q = 2 = q'$ the norm $\phi_{q'}$ is just the Euclidean norm and $Q_{p,q}$ reduces to the well-known p -Laplace Operator

$$Q_{p,q}u = \Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u).$$

For general q and $p \rightarrow 1$ the operator $Q_{1,q}$ is formally given by

$$Q_{1,q}u = \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\left[\frac{|u_{x_i}|}{\phi_{q'}(\nabla u)} \right]^{q'-2} \frac{u_{x_i}}{\phi_{q'}(\nabla u)} \right).$$

Again for $q = 2 = q'$ this expression shrinks down to the customary

$$Q_{1,2}u = \Delta_1 u = \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right).$$

We complete this section with the construction of a particular Cheeger set for a nonregular anisotropy. Let us fix $n = 2$ and consider the norm $\phi = \phi_1$. Notice that in this case the Wulff Shape W_ϕ has the shape of a rhombus. To be precise, it is square of sidelength $\sqrt{2}$, centered in the origin and rotated by $\pi/2$ with respect to the coordinate axes. Moreover, the dual norm ϕ^* is given by $\phi^*(y) = \max\{|y_1|, |y_2|\}$. To better illustrate the results of Section 5, let us compute the Cheeger set (and Cheeger constant) of a square Q of sidelength 1 (see Figure 2).

Since in this case $|W_\phi| = 2$ and Q_-^t is a square of sidelength $1 - 2t$, from Theorem 5.1 we get $t^* = 1 - \sqrt{2}/2$ and $h(Q) = 2 + \sqrt{2}$. It is interesting to note that the Cheeger set of Q is a regular octahedron.




Figure 2: The Cheeger set of a square with respect to the norm ϕ_1 .

References

- [1] B.Andrews, Volume-preserving anisotropic mean curvature flow, *Indiana Univ. Math. J.*, **50** (2001) pp. 783–827.
- [2] A.Avinyo, Isoperimetric constants and some lower bounds for the eigenvalues of the p -Laplacian, *Nonlin. Anal., Theory, Methods and Appl.* **30** (1997) pp. 177–180.
- [3] G.Bellettini, M.Novaga & M.Paolini, Characterization of facet-breaking for nonsmooth mean curvature flow in the convex case, *Interfaces and Free Boundaries*, **3** (2001) pp. 415–446.
- [4] M.Belloni & B.Kawohl, A direct uniqueness proof for equations involving the p -Laplace operator, *manuscripta math.*, **109** (2002) pp. 229–231.
- [5] M.Belloni & B.Kawohl, The pseudo- p -Laplace eigenvalue problem and viscosity solutions as $p \rightarrow \infty$, *ESAIM COCV*, **10** (2004) pp. 28–52.
- [6] M.Belloni, V.Ferone & B.Kawohl, Isoperimetric inequalities, Wulff shape and related questions for strongly nonlinear elliptic operators. *J. Appl. Math. Phys. (ZAMP)*, **54** (2003) pp. 771–783.
- [7] M.Belloni, B.Kawohl & P.Juutinen: The p -Laplace eigenvalue problem as $p \rightarrow \infty$ in a Finsler metric, *J. Europ. Math. Soc.* **8** (2006) pp. 123–138.
- [8] H.Brezis, *Operateurs Maximaux Monotones*, North-Holland, Amsterdam (1973).

- [9] V.Caselles, A.Chambolle, S.Moll & M.Novaga, A characterization of convex calibrable sets in \mathbb{R}^N with respect to anisotropic norms, Preprint No. 2.560.1613, Univ. Pisa (2005).
- [10] J.Cheeger, A lower bound for the smallest eigenvalue of the Laplacian, in: *Problems in Analysis, A Symposium in Honor of Salomon Bochner, Ed.: R.C.Gunning*, Princeton Univ. Press (1970) pp. 195–199.
- [11] G.Cupini, N.Fusco, R.Petti, Hölder continuity of local minimizers. *J. Math. Anal. Appl.* **235** (1999) pp. 578–597.
- [12] E.DiBenedetto, $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. *Nonlin. Anal. TMA* **7** (1983) pp. 827–850.
- [13] L. Esposito & C. Trombetti, Convex symmetrization and Pólya-Szegő inequality, *Nonlinear Anal.* **56** (2004) pp. 43–62.
- [14] A. Ferone & R.Volpicelli, Convex symmetrization: the equality case in Pólya-Szegő inequality, *Calc. Var. Partial Differential Equations* **21** (2004) pp. 259–272.
- [15] E.Giusti, *Metodi diretti nel calcolo delle variazioni*. Unione Matematica Italiana, Bologna (1994)
- [16] E.Hebey & N.Saintier, Stability and perturbation of the domain for the first eigenvalue of the 1-Laplacian, *Archiv der Math.*, to appear
- [17] P.Hild, I.R. Ionescu, Th. Lachand-Robert & I.Rosca, The blocking of an inhomogeneous Bingham fluid. Applications to landslides. *Math. Modelling & Num. Anal.* **36**(2002) pp. 1013–1026.
- [18] I.R.Ionescu & Th.Lachand-Robert, Generalized Cheeger sets related to landslides, *Calc. Var.* **23** (2005) pp. 227–249.
- [19] B. Kawohl & V.Fridman, Isoperimetric estimates for the first eigenvalue of the p -Laplace operator and the Cheeger constant, *Comment. Math. Univ. Carolinae*, **44** (2003) pp. 659-667.
- [20] B.Kawohl & T.Lachand-Robert, Characterization of Cheeger sets for convex subsets of the plane, *Pacific J. Math.*, to appear
- [21] B.Kawohl & F.Schuricht, Dirichlet problems for the 1-laplace operator, including the eigenvalue problem, submitted 2005
- [22] J.B. Keller, Plate failure under pressure, *SIAM Review* **22** (1980) pp. 227–228.
- [23] Th.Lachand-Robert & É.Oudet, Minimizing within convex bodies using a convex hull method. *SIAM J. Optim.* **16** (2006) pp. 368–379.
- [24] L.Lefton & D.Wei, Numerical approximation of the first eigenpair of the p -Laplacian using finite elements and the penalty method, *Numer. Funct. Anal. Optim.* **18** (1997) 389–399.

- [25] G.M.Lieberman, Boundary regularity for solutions of degenerate elliptic equations, *Nonlin. Anal. TMA* **12** (1988) pp. 1203–1219.
- [26] A.M.Matei, First eigenvalue for the p -Laplace operator, *Nonlinear Anal. TMA* **39** (2000) 1051–1061.
- [27] M.Novaga & E.Paolini, Regularity results for boundaries in \mathbb{R}^2 with prescribed anisotropic curvature. *Ann. Mat. Pura Appl.* **184** (2005) pp. 239–261.
- [28] S.Sakaguchi, Concavity properties of solutions to some degenerate quasilinear elliptic Dirichlet problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **14** (1987) pp. 403–421.
- [29] R.Schneider, *Convex bodies: The Brunn-Minkowski theory*, Cambridge University Press (1993).
- [30] F.Schuricht, An alternative derivation of the eigenvalue equation for the 1-Laplace operator, *Arch. d. Math.*, to appear
- [31] H.Takeuchi, On the first eigenvalue of the p -Laplacian in a Riemannian manifold. *Tokyo J. Math.* **21**(1998) 135–140.
- [32] P.Tolksdorf, On the Dirichlet problem for quasilinear equations in domains with conical boundary points. *Comm. Part. Differ. Eq.* **8** (1983) pp. 773–817.
- [33] P.Tolksdorf, Regularity for a more general class of quasilinear elliptic equations. *J. Differential Equations* **51** (1984) pp. 126–150.
- [34] E.Zeidler, *Nonlinear Functional Analysis and Applications III, Variational Methods and Optimization*, Springer Verlag, Heidelberg (1984)

Author’s addresses:

B. Kawohl, Mathematisches Institut, Universität zu Köln, D-50923 Köln, Germany; kawohl@math.uni-koeln.de

M. Novaga, Dipartimento di Matematica, Università di Pisa, Largo B. Pontecorvo 5, 56127 Pisa, Italy; novaga@dm.unipi.it