

EXISTENCE OF MINIMIZERS FOR SPECTRAL PROBLEMS

DARIO MAZZOLENI AND ALDO PRATELLI

ABSTRACT. In this paper we show that any increasing functional of the first k eigenvalues of the Dirichlet Laplacian admits a (quasi-)open minimizer among the subsets of \mathbb{R}^N of unit measure. In particular, there exists such a minimizer which is bounded, where the bound depends on k and N , but not on the functional.

Dans ce papier, on montre que toute fonctionnelle croissante des premières k valeurs propres du Laplacien-Dirichlet admet un minimiseur (quasi-)ouvert, parmi les sous-ensembles de \mathbb{R}^N ayant mesure unitaire. En particulier, il existe un tel minimiseur qui est borné, la limitation en dépendant de k et de N , mais pas de la fonctionnelle.

CONTENTS

1. Introduction	1
1.1. Notation and preliminaries	3
2. Proof of Theorem A	5
2.1. Boundedness of the tails	6
2.2. Boundedness of the interior	15
2.3. Proof of Proposition 2.1	21
3. Proof of Theorem B	22
References	25

1. INTRODUCTION

This paper deals with the existence of minimizers for an increasing functional of the first k eigenvalues of the Dirichlet Laplacian, among (quasi-)open sets in \mathbb{R}^N with unit measure.

This kind of spectral minimization problems is well-studied and natural in many situations: for instance, one may be interested in minimizing a certain eigenvalue, or a linear combination of eigenvalues, or a product of eigenvalues, and so on. However, despite the large interest in the question, very little is known in the general situation when the problem is stated in the whole \mathbb{R}^N . The reason is basically the lack of compactness for generic sequences of open sets; in fact, given a sequence of sets of unit measure, it is not easy to tell whether or not it converges to a limit set in a suitable sense. The first existence result ([7, Theorem 2.5], see also [10,

Key words and phrases. Dirichlet Laplacian, Eigenvalue problems, Shape Optimization, Minimization of Spectral Functionals.

Théorème 4.7.6]), which is now classical, was found by Buttazzo and Dal Maso in the context of bounded sets.

Theorem 1.1 (Buttazzo–Dal Maso, 1993). *Let $D \subseteq \mathbb{R}^N$ be a bounded open set, and let $\{A_n\} \subseteq D$ be a sequence of (quasi-)open sets of unit measure. Then, there exists a subsequence $\{A_{n(m)}\}$ and a (quasi-)open set $A \subseteq D$ of unit measure such that*

$$\lambda_i(A) \leq \liminf_{m \rightarrow \infty} \lambda_i(A_{n(m)})$$

for every $i \in \mathbb{N}$. As an immediate consequence, for every l.s.c. functional $\mathcal{F} : \mathbb{R}^k \rightarrow \mathbb{R}$ increasing in each variable, there exists A minimizing the value of $\mathcal{F}(\lambda_1(A), \lambda_2(A), \dots, \lambda_k(A))$ among the (quasi-)open sets of unit volume contained in D .

Thanks to a concentration-compactness argument (see for instance [9, Theorem 5.3.1]), it is now well-known that the boundedness assumption ($A_n \subseteq D$ for every n) in the theorem above can be replaced by the following slightly weaker assumption: for every $\varepsilon > 0$ there exists $R = R(\varepsilon) > 0$ such that every set A_n is contained in a cube of side R up to a volume at most ε .

The results described above make it simple to study spectral minimization problems inside an ambient space which is essentially bounded. On the other hand, very little is known concerning general problems in \mathbb{R}^N , because minimizing sequences, in principle, could have a significant portion of volume moving to infinity. More precisely, the existence of a minimizer for λ_1 and λ_2 is clear, since it is well-known that a ball of unit volume minimizes λ_1 , while two disjoint balls of half volume minimize λ_2 : these two classical results are usually referred to as the Faber–Krahn inequality and the Krahn–Szegő inequality respectively. On the other hand, the existence of a minimizer for λ_3 has been proved only in 2000 by Bucur and Henrot (see [6]), and it is still presently not known which set is the minimizer (and this is a major open problem). For other open problems and partial results see [8, 3].

In this paper, we show that every increasing functional of the first k eigenvalues admits a minimizer, which is in turn a bounded set. In other words, we prove that the Buttazzo–Dal Maso Theorem 1.1 is true in the whole \mathbb{R}^N , with no need of the *a priori* boundedness assumption.

Theorem A (Existence of bounded minimizers). *Let $k \in \mathbb{N}$, and let $\mathcal{F} : \mathbb{R}^k \rightarrow \mathbb{R}$ be a l.s.c. functional, increasing in each variable. Then there exists a bounded minimizer for the problem*

$$\inf \left\{ \mathcal{F}(\lambda_1(A), \lambda_2(A), \dots, \lambda_k(A)) : A \subseteq \mathbb{R}^N, |A| = 1 \right\} \quad (1.1)$$

among the (quasi-)open sets. More precisely, a minimizer A is contained in a cube of side R , where R depends on k and on N , but not on the particular functional \mathcal{F} .

The strategy of the proof consists in taking a generic open set $\Omega \subseteq \mathbb{R}^N$ of unit volume, and showing that there exists a modified open set $\widehat{\Omega}$ which is uniformly bounded and which has all the first k eigenvalues lower than those of Ω . Roughly speaking, the basic idea why this works is that, if a set of unit volume has huge diameter, then there must be some very thin sections. This works against the smallness of the Rayleigh quotients of the eigenfunctions, since by definition they vanish on the boundary. As an immediate consequence of the above boundedness claim,

there will clearly exist a uniformly bounded minimizing sequence, and therefore Theorem A will follow by Theorem 1.1.

During our construction, we will need the following result.

Theorem B (Boundedness of the ratio λ_k/λ_1). *There exists a constant $M = M(k, N)$ such that for every quasi-open set A one has $\lambda_k(A) \leq M\lambda_1(A)$.*

In the particular case when $k = 2$, Theorem B is a consequence of the stronger result by Ashbaugh and Benguria [2], which states that not only the ratio λ_2/λ_1 is bounded, but also that balls maximize this ratio. Many stronger results were proved also for generic k (see [1] for an overview), but in order to keep the present paper self contained, we give a simple proof of Theorem B in Section 3. Notice that one cannot apply Theorem A to this problem, since the functional λ_k/λ_1 is not increasing in the first variable. Notice also that Theorem B is stated for sets of any volume; in fact, since by trivial rescaling one finds that for every $\alpha > 0$ and every $i \in \mathbb{N}$ one has

$$\lambda_i(\alpha A) = \alpha^{-2}\lambda_i(A), \quad (1.2)$$

the ratio λ_k/λ_1 does not change by rescaling.

The plan of the paper is the following. In Section 1.1 below we give a brief list of the notation that will be used through the paper, and some preliminaries. Then, in Section 2 we show Theorem A, while in Section 3 we show Theorem B.

NOTE: *The preprint version of this paper appeared contemporarily to the preprint version of the paper [4] by Dorin Bucur, in which he gives the proof –by means of completely independent ideas– of the existence of a minimizer for the k -th eigenvalue of the Dirichlet Laplacian in \mathbb{R}^N (which corresponds, in the language of our Theorem A, to the particular case when \mathcal{F} is projection on the last variable). In addition, Bucur is able to show that the quasi-open minimizers of λ_k are sets of finite perimeter.*

1.1. Notation and preliminaries. Throughout this paper, the ambient space will be \mathbb{R}^N for some given $N \geq 2$. A generic point of \mathbb{R}^N will be denoted by $z \equiv (x, y) \in \mathbb{R} \times \mathbb{R}^{N-1}$, or sometimes as $z \equiv (z_1, z_2, \dots, z_N)$, while a generic open set will be $\Omega \subseteq \mathbb{R}^N$. The eigenvalues of the Dirichlet Laplacian on Ω will be denoted by λ_i , $i \in \mathbb{N}$, while $\{u_i\} \subseteq W_0^{1,2}(\Omega)$ will be a corresponding sequence of orthogonal eigenfunctions, always normalized to have unit L^2 norm. For any function $v : \Omega \rightarrow \mathbb{R}$ and any set $D \subseteq \Omega$, we will consider the Rayleigh quotient

$$\mathcal{R}(v, D) := \frac{\int_D |Dv|^2}{\int_D v^2};$$

hence, in particular $\mathcal{R}(u_i, \Omega) = \lambda_i(\Omega)$ for every $i \in \mathbb{N}$. The following is a very useful characterization of the eigenvalues, which we will need later, and whose proof can be found for instance in [9]. It is often referred to as *min-max principle*.

Theorem 1.2. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set. Then for every $j \in \mathbb{N}$ one has*

$$\lambda_j(\Omega) = \min \left\{ \max \left\{ \mathcal{R}(w, \Omega), w \in K_j \setminus \{0\} \right\} \right\},$$

where the minimum is taken among all the j -dimensional subspaces K_j of $W_0^{1,2}(\Omega)$.

In our result, we need to use quasi-open sets. However, the construction of this paper only deals with open sets, and actually no knowledge of quasi-open sets is needed; therefore, we only recall here the very basic definition, addressing the interested reader for instance to [5, 10] for a complete treatment.

Definition 1.3. *Let Ω be an open set, and $V \subset\subset \Omega$ a compactly supported subset. The capacity of V in Ω is defined as*

$$\text{cap}_\Omega(V) := \inf \left\{ \int_\Omega |D\varphi|^2 : \varphi \in C_0^\infty(\Omega), \varphi \geq 1 \text{ on } V \right\}.$$

Let then $A \subseteq \mathbb{R}^N$ be a bounded set, and let Ω be an open set such that $A \subset\subset \Omega$. The set A is said quasi-open if for every $\varepsilon > 0$ there exists an open set $A \subseteq \Omega_\varepsilon \subset\subset \Omega$ such that $\text{cap}_\Omega(\Omega_\varepsilon \setminus A) < \varepsilon$. This definition does not depend on the choice of Ω . Finally, a generic set $A \subseteq \mathbb{R}^N$ is said quasi-open if the intersection of A with any ball of \mathbb{R}^N is a quasi-open bounded set.

It is well-known that the notion of eigenvalues can be extended to the realm of quasi-open sets. An important feature of the quasi-open sets is given by the Theorem 1.1 by Buttazzo and Dal Maso; namely, that any bounded sequence of open (or quasi-open) sets converges –up to a subsequence– to a quasi-open set, in the sense that all the eigenvalues converge.




FIGURE 1. A set Ω and the corresponding sets Ω_t^l , Ω_t^r and Ω_t .

Through the paper, $k \in \mathbb{N}$ will be a fixed integer, and $\mathcal{F} : \mathbb{R}^k \rightarrow \mathbb{R}$ will be a l.s.c. function, increasing in each variable. For the sake of brevity, we will often write $\mathcal{F}(\Omega)$ in place of $\mathcal{F}(\lambda_1(\Omega), \lambda_2(\Omega), \dots, \lambda_k(\Omega))$; hence, the goal of this paper can be rephrased by saying that we seek for a quasi-open set A minimizing $\mathcal{F}(A)$. By density, it is clear that the infimum of \mathcal{F} among the quasi-open sets equals the infimum among the open (or even smooth) sets.

The letter C will be always used to denote a big geometric constant, possibly increasing from line to line; the constant C will always depend *only* on N and on k (sometimes, possibly also on some constant K , which in turn will eventually be chosen only depending on N and k), thus *not* on the particular choice of \mathcal{F} , and *not* on the set Ω . Sometimes, we will label the constants in our results as $C_1, C_2, C_3 \dots$ for successive reference.

For any $t \in \mathbb{R}$, we will define

$$\Omega_t^l := \{(x, y) \in \Omega : x < t\}, \quad \Omega_t := \{y \in \mathbb{R}^{N-1} : (t, y) \in \Omega\}, \quad \Omega_t^r := \{(x, y) \in \Omega : x > t\};$$

notice that Ω_t^l and Ω_t^r are subsets of \mathbb{R}^N , while Ω_t is a subset of \mathbb{R}^{N-1} . Figure 1 shows an example of a generic set Ω with Ω_t^l , Ω_t^r and Ω_t . On the other hand, given $0 \leq m \leq |\Omega|$ and $0 \leq m_1 \leq m_2 \leq |\Omega|$, we define the level $\tau(\Omega, m) \in \overline{\mathbb{R}}$ and the width $W(\Omega, m_1, m_2)$ as

$$\tau(\Omega, m) := \inf \left\{ t \in \mathbb{R} : |\Omega_t^l| \geq m \right\}, \quad W(\Omega, m_1, m_2) := \tau(\Omega, m_2) - \tau(\Omega, m_1).$$

Observe that one surely has $-\infty < \tau(\Omega, m) < +\infty$ whenever $0 < m < |\Omega|$, as well as $W(\Omega, m_1, m_2) < +\infty$ if $0 < m_1 \leq m_2 < |\Omega|$.

Finally, given any set $\Omega \subseteq \mathbb{R}^N$, we define its 1-dimensional projections for $1 \leq p \leq N$ as

$$\pi_p(\Omega) := \left\{ t \in \mathbb{R} : \exists (z_1, z_2, \dots, z_N) \in \Omega, z_p = t \right\}.$$

2. PROOF OF THEOREM A

In this section we present the proof of Theorem A. As already anticipated in the introduction, our strategy basically consists in showing that to minimize \mathcal{F} it is enough to concentrate on uniformly bounded sets. More precisely, we will show the following result.

Proposition 2.1. *For every $K > 0$ there exists a constant $R = R(k, K, N)$, such that the following holds. If $\Omega \subseteq \mathbb{R}^N$ is an open set of unit volume and with $\lambda_k(\Omega) \leq K$, there exists another open set $\widehat{\Omega}$, still of unit volume but contained in a cube of side R , and with $\lambda_i(\widehat{\Omega}) \leq \lambda_i(\Omega)$ for every $1 \leq i \leq k$.*

Let us immediately see how Theorem A follows from this proposition; then the rest of the section will be devoted to showing the proposition.

Proof of Theorem A. Let us take a minimizing sequence of open sets $\{\Omega_n\}$ for problem (1.1). Fix a generic $n \in \mathbb{N}$, and assume for a moment that $\lambda_k(\Omega_n) \geq M\lambda_k(B_N)$, being B_N the ball of unit volume and M the constant of Theorem B. If it is so, then by Theorem B one has $\lambda_1(\Omega_n) \geq \lambda_k(B_N)$, thus for every $1 \leq i \leq k$ it is $\lambda_i(\Omega_n) \geq \lambda_i(B_N)$, hence $\mathcal{F}(\Omega_n) \geq \mathcal{F}(B_N)$, being \mathcal{F} increasing in each variable. Thanks to this observation, it is admissible to assume that $\lambda_k(\Omega_n) \leq K := M\lambda_k(B_N)$ for every n . By Proposition 2.1, then, there exists another sequence $\{\widehat{\Omega}_n\}$, made by open sets of unit volume contained in a cube of side R , with $\lambda_i(\widehat{\Omega}_n) \leq \lambda_i(\Omega_n)$ for every $1 \leq i \leq k$ and every $n \in \mathbb{N}$. Again by the assumption that \mathcal{F} is increasing in each variable, we derive that also $\{\widehat{\Omega}_n\}$ is a minimizing sequence for (1.1).

We can then apply Theorem 1.1 to find a quasi-open set A , still contained in the cube of side R , and such that, up to extract a subsequence of $\{\widehat{\Omega}_n\}$, one has $\lambda_i(A) \leq \liminf_n \lambda_i(\widehat{\Omega}_n)$ for every $1 \leq i \leq k$. By the lower semi-continuity of \mathcal{F} , we derive that A is a minimizer for \mathcal{F} , thus the proof is concluded. \square

The rest of this section is devoted to show Proposition 2.1. For the ease of presentation, we divide the construction in three subsections. In the first one we obtain the boundedness of the

“tails” (Lemma 2.2), while in the second one we consider the internal part (Lemma 2.8). Then, in the last subsection we put everything together to give the proof of Proposition 2.1.

2.1. Boundedness of the tails. This subsection is devoted to show that, under the assumptions of Proposition 2.1, we can reduce to the case when the “tails” of Ω are bounded. More precisely, we fix once for all a small positive number $\widehat{m} = \widehat{m}(K, N) \in (0, 1/4)$ in such a way that

$$\frac{(4\widehat{m})^{\frac{2}{N}}}{\lambda_1(B_N)} K \leq \frac{1}{2}, \quad (2.1)$$

being B_N the ball of unit volume in \mathbb{R}^N . We aim to show the following result.

Lemma 2.2. *For every $K > 0$ there exist $R_1 = R_1(k, K, N)$ and $\Gamma_1 = \Gamma_1(k, K, N)$ such that, for any open set $\Omega \subseteq \mathbb{R}^N$ of unit volume and with $\lambda_k(\Omega) \leq K$, there exists another open set $\widehat{\Omega} \subseteq \mathbb{R}^N$, still of unit volume, such that $\lambda_i(\widehat{\Omega}) \leq \lambda_i(\Omega)$ for every $1 \leq i \leq k$, and that for every $2 \leq p \leq N$*

$$W(\widehat{\Omega}, 0, \widehat{m}) \leq R_1, \quad (2.2)$$

$$W(\widehat{\Omega}, \widehat{m}, 1) \leq \Gamma_1 \left(W(\Omega, \widehat{m}, 1) \right), \quad \text{diam}(\pi_p(\widehat{\Omega})) \leq \Gamma_1 \text{diam}(\pi_p(\Omega)). \quad (2.3)$$

The claim of the lemma, roughly speaking, says that it is always possible to assume that the “tail” of Ω , i.e., the set $\Omega_{\tau(\Omega, \widehat{m})}^l$ of volume \widehat{m} , has horizontal projection of length at most R_1 . More precisely, condition (2.2) says that one can modify Ω in such a way that the tail is uniformly horizontally bounded, while condition (2.3) says that this modification does not excessively worsen the remaining part of the set Ω , nor its extension in the $N - 1$ non-horizontal directions.

To prove the lemma, we start setting for brevity $\bar{t} = \tau(\Omega, 2\widehat{m})$, and for every $t \leq \bar{t}$ we define

$$\Omega^+(t) := \Omega_t^r, \quad \Omega^-(t) := \Omega_t^l, \quad \varepsilon(t) := \mathcal{H}^{N-1}(\Omega_t). \quad (2.4)$$

Observe that

$$m(t) := |\Omega^-(t)| = \int_{-\infty}^t \varepsilon(s) ds \leq 2\widehat{m}. \quad (2.5)$$

Moreover, we let as usual $\{u_1, u_2, \dots, u_k\}$ be an orthonormal set of eigenfunctions with unit L^2 norm and corresponding to the first k eigenvalues of Ω . We define then also, for every $1 \leq i \leq k$ and every $t \leq \bar{t}$,

$$\delta_i(t) := \int_{\Omega_t} |Du_i(t, y)|^2 d\mathcal{H}^{N-1}(y), \quad \mu_i(t) := \int_{\Omega_t} u_i(t, y)^2 d\mathcal{H}^{N-1}(y), \quad (2.6)$$

which makes sense since every u_i is smooth. It is convenient to give the further notation

$$\delta(t) := \sum_{i=1}^k \delta_i(t) = \sum_{i=1}^k \int_{\Omega_t} |Du_i(t, y)|^2 d\mathcal{H}^{N-1}(y),$$

and in analogy with (2.5) we also set

$$\phi(t) := \sum_{i=1}^k \int_{\Omega^-(t)} |Du_i|^2 = \int_{-\infty}^t \delta(s) ds. \quad (2.7)$$

Applying the Faber–Krahn inequality in \mathbb{R}^{N-1} to the set Ω_t , and using (1.2) on \mathbb{R}^{N-1} , we know that

$$\varepsilon(t)^{\frac{2}{N-1}} \lambda_1(\Omega_t) = \mathcal{H}^{N-1}(\Omega_t)^{\frac{2}{N-1}} \lambda_1(\Omega_t) \geq \lambda_1(B_{N-1}),$$

calling B_{N-1} the unit ball in \mathbb{R}^{N-1} . As a trivial consequence, we can estimate μ_i in terms of ε and δ_i : in fact, noticing that $u_i(t, \cdot) \in W_0^{1,2}(\Omega_t)$ and writing $Du_i = (D_1u_i, D_yu_i)$, we have

$$\mu_i(t) = \int_{\Omega_t} u_i(t, \cdot)^2 d\mathcal{H}^{N-1} \leq \frac{1}{\lambda_1(\Omega_t)} \int_{\Omega_t} |D_yu_i(t, \cdot)|^2 d\mathcal{H}^{N-1} \leq C\varepsilon(t)^{\frac{2}{N-1}} \delta_i(t). \quad (2.8)$$

We can now present two estimates which assure that u_i and Du_i can not be too big in $\Omega^-(t)$.

Lemma 2.3. *Under the assumptions of Lemma 2.2, for every $1 \leq i \leq k$ and $t \leq \bar{t}$ the following inequalities hold:*

$$\int_{\Omega^-(t)} u_i^2 \leq C_1 \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t), \quad \int_{\Omega^-(t)} |Du_i|^2 \leq C_1 \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t), \quad (2.9)$$

for some $C_1 = C_1(k, K, N)$.

Proof. Let us fix $t \leq \bar{t}$. Consider the set Ω_S^- obtained by the union of $\Omega^-(t)$ and its reflection with respect to the plane $\{x = t\}$, and call $u_S \in W_0^{1,2}(\Omega_S)$ the function obtained by reflecting u_i . Calling B_N the unit ball in \mathbb{R}^N , we find then

$$\frac{\lambda_1(B_N)}{(2m(t))^{\frac{2}{N}}} = \frac{\lambda_1(B_N)}{|\Omega_S^-|^{\frac{2}{N}}} \leq \lambda_1(\Omega_S^-) \leq \mathcal{R}(u_S, \Omega_S^-) = \mathcal{R}(u_i, \Omega^-(t)) = \frac{\int_{\Omega^-(t)} |Du_i|^2}{\int_{\Omega^-(t)} u_i^2},$$

by the symmetry of Ω_S^- , and using again (1.2). This estimate gives

$$\int_{\Omega^-(t)} u_i^2 \leq \frac{(2m(t))^{\frac{2}{N}}}{\lambda_1(B_N)} \int_{\Omega^-(t)} |Du_i|^2 \quad (2.10)$$

which in particular, being $m(t) \leq 2\hat{m}$ and recalling (2.1), implies

$$\int_{\Omega^-(t)} u_i^2 \leq \frac{1}{2}. \quad (2.11)$$

On the other hand, recalling that $-\Delta u_i = \lambda_i u_i$, by Schwarz inequality and using (2.8) we have

$$\begin{aligned} \int_{\Omega^-(t)} |Du_i|^2 &= \int_{\Omega^-(t)} \lambda_i u_i^2 + \int_{\Omega_t} u_i \frac{\partial u_i}{\partial \nu} \leq K \int_{\Omega^-(t)} u_i^2 + \sqrt{\int_{\Omega_t} u_i^2 \int_{\Omega_t} |Du_i|^2} \\ &\leq K \int_{\Omega^-(t)} u_i^2 + C\varepsilon(t)^{\frac{1}{N-1}} \delta_i(t). \end{aligned} \quad (2.12)$$

It is now easy to obtain (2.9) combining (2.10) and (2.12). In fact, by inserting the latter into the first, we find

$$\int_{\Omega^-(t)} u_i^2 \leq \frac{(2m(t))^{\frac{2}{N}}}{\lambda_1(B_N)} \left(K \int_{\Omega^-(t)} u_i^2 + C\varepsilon(t)^{\frac{1}{N-1}} \delta_i(t) \right),$$

which by (2.1) again yields

$$\frac{1}{2} \int_{\Omega^-(t)} u_i^2 \leq \frac{(2m(t))^{\frac{2}{N}}}{\lambda_1(B_N)} C\varepsilon(t)^{\frac{1}{N-1}} \delta_i(t) \leq C\varepsilon(t)^{\frac{1}{N-1}} \delta_i(t). \quad (2.13)$$

The left estimate in (2.9) is then obtained. To get the right one, one has then just to insert (2.13) into (2.12). \square

Let us go further into our construction, giving some definitions. For any $t \leq \bar{t}$ and $\sigma(t) > 0$, we define the cylinder $Q(t)$, shown in Figure 2, as

$$Q(t) := \left\{ (x, y) \in \mathbb{R}^N : t - \sigma < x < t, (t, y) \in \Omega \right\} = (t - \sigma, t) \times \Omega_t, \quad (2.14)$$

where for any $t \leq \bar{t}$ we set

$$\sigma(t) = \varepsilon(t)^{\frac{1}{N-1}}. \quad (2.15)$$

We let also $\tilde{\Omega}(t) = \Omega^+(t) \cup Q(t)$, and we introduce $\tilde{u}_i \in W_0^{1,2}(\tilde{\Omega}(t))$ as

$$\tilde{u}_i(x, y) := \begin{cases} u_i(x, y) & \text{if } (x, y) \in \Omega^+(t), \\ \frac{x - t + \sigma}{\sigma} u_i(t, y) & \text{if } (x, y) \in Q(t). \end{cases} \quad (2.16)$$

The fact that \tilde{u}_i vanishes on $\partial\tilde{\Omega}(t)$ is obvious; moreover, $Du_i = D\tilde{u}_i$ on $\Omega^+(t)$, while on $Q(t)$ one has

$$D\tilde{u}_i(x, y) = \left(\frac{u_i(t, y)}{\sigma}, \frac{x - t + \sigma}{\sigma} D_y u_i(t, y) \right). \quad (2.17)$$

A simple calculation allows us to estimate the integrals of \tilde{u}_i and $D\tilde{u}_i$ on $Q(t)$.




FIGURE 2. A set Ω with the cylinder $Q(t)$ (shaded).

Lemma 2.4. *For every $t \leq \bar{t}$ and $1 \leq i \leq k$, one has*

$$\int_{Q(t)} |D\tilde{u}_i|^2 \leq C_2 \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t), \quad \int_{Q(t)} \tilde{u}_i^2 \leq C_2 \varepsilon(t)^{\frac{3}{N-1}} \delta_i(t), \quad (2.18)$$

for a suitable constant $C_2 = C_2(k, K, N)$.

Proof. Thanks to (2.17), and using also (2.8) and (2.15), one obtains the first estimate in (2.18) since

$$\begin{aligned} \int_{Q(t)} |D\tilde{u}_i(x, y)|^2 dx dy &= \int_{t-\sigma}^t \int_{\Omega_t} \frac{u_i^2(t, y)}{\sigma^2} + \frac{(x-t+\sigma)^2}{\sigma^2} |D_y u_i(t, y)|^2 dy dx = \frac{\mu_i}{\sigma} + \frac{\delta_i \sigma}{3} \\ &\leq C \frac{\varepsilon(t)^{\frac{2}{N-1}} \delta_i(t)}{\sigma} + \frac{\delta_i \sigma}{3} = C \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t). \end{aligned}$$

On the other hand, the second estimate in (2.18) follows, also again using (2.8) and (2.15), by

$$\int_{Q(t)} \tilde{u}_i(x, y)^2 dx dy = \int_{t-\sigma}^t \int_{\Omega_t} \tilde{u}_i^2(t, y) dy dx = \frac{\sigma \mu_i}{3} \leq C \sigma \varepsilon(t)^{\frac{2}{N-1}} \delta_i(t) = C \varepsilon(t)^{\frac{3}{N-1}} \delta_i(t).$$

□

Another simple but useful estimate concerns the Rayleigh quotients of the functions \tilde{u}_i on the sets $\tilde{\Omega}(t)$ and the integral of the products $\tilde{u}_i \tilde{u}_j$.

Lemma 2.5. *For every $t \leq \bar{t}$ and $1 \leq i \leq k$, one has*

$$\mathcal{R}(\tilde{u}_i, \tilde{\Omega}(t)) \leq \lambda_i(\Omega) + C \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t). \quad (2.19)$$

Moreover, for every $i \neq j \in \{1, 2, \dots, k\}$, one has

$$\left| \int_{\tilde{\Omega}(t)} \tilde{u}_i \tilde{u}_j + D\tilde{u}_i \cdot D\tilde{u}_j \right| \leq C \left(\varepsilon(t)^{\frac{3}{N-1}} + \varepsilon(t)^{\frac{1}{N-1}} \right) \sqrt{\delta_i(t) \delta_j(t)}. \quad (2.20)$$

Proof. Recalling that $-\Delta u_i = \lambda_i(\Omega) u_i$, making use of (2.11) and (2.18) and arguing as in (2.12), we obtain

$$\begin{aligned} \mathcal{R}(\tilde{u}_i, \tilde{\Omega}(t)) &= \frac{\int_{\Omega^+(t)} |D\tilde{u}_i|^2 + \int_{Q(t)} |D\tilde{u}_i|^2}{\int_{\Omega^+(t)} \tilde{u}_i^2 + \int_{Q(t)} \tilde{u}_i^2} \leq \frac{\int_{\Omega^+(t)} |Du_i|^2 + \int_{Q(t)} |D\tilde{u}_i|^2}{\int_{\Omega^+(t)} u_i^2} \\ &= \frac{\lambda_i(\Omega) \int_{\Omega^+(t)} u_i^2 + \int_{\Omega_t} u_i \frac{\partial u_i}{\partial \nu} + \int_{Q(t)} |D\tilde{u}_i|^2}{\int_{\Omega^+(t)} u_i^2} \leq \lambda_i(\Omega) + C \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t), \end{aligned}$$

hence (2.19) is proved.

On the other hand, recall that u_i and u_j are orthogonal on Ω both in L^2 and in $W_0^{1,2}$ sense by definition, hence by using (2.9) and (2.18) we find

$$\begin{aligned} \left| \int_{\tilde{\Omega}(t)} \tilde{u}_i \tilde{u}_j \right| &\leq \left| \int_{\Omega^+(t)} u_i u_j \right| + \left| \int_{Q(t)} \tilde{u}_i \tilde{u}_j \right| = \left| \int_{\Omega^-(t)} u_i u_j \right| + \left| \int_{Q(t)} \tilde{u}_i \tilde{u}_j \right| \\ &\leq C \left(\varepsilon(t)^{\frac{1}{N-1}} \sqrt{\delta_i(t) \delta_j(t)} + \varepsilon(t)^{\frac{3}{N-1}} \sqrt{\delta_i(t) \delta_j(t)} \right). \end{aligned}$$

In the very same way, concerning $D\tilde{u}_i$ and $D\tilde{u}_j$, we have

$$\begin{aligned} \left| \int_{\tilde{\Omega}(t)} D\tilde{u}_i \cdot D\tilde{u}_j \right| &\leq \left| \int_{\Omega^+(t)} Du_i \cdot Du_j \right| + \left| \int_{Q(t)} D\tilde{u}_i \cdot D\tilde{u}_j \right| \\ &= \left| \int_{\Omega^-(t)} Du_i \cdot Du_j \right| + \left| \int_{Q(t)} D\tilde{u}_i \cdot D\tilde{u}_j \right| \leq C \varepsilon(t)^{\frac{1}{N-1}} \sqrt{\delta_i(t) \delta_j(t)}. \end{aligned}$$

Adding up the last two estimates yields (2.20). \square

In order to prove Lemma 2.2, we need to compare the eigenvalues of Ω and those of $\tilde{\Omega}(t)$; this can be done by means of the min-max principle, Theorem 1.2, which relates the eigenvalues with the Rayleigh quotients of $W_0^{1,2}$ functions.

Lemma 2.6. *There exist a small constant $\nu = \nu(k, K, N) < 1$ and a constant $C_3 = C_3(k, K, N)$ such that, if $\varepsilon(t), \delta_i(t) \leq \nu$ for every $1 \leq i \leq k$, then*

$$\lambda_j(\tilde{\Omega}(t)) \leq \lambda_j(\Omega) + C_3 \varepsilon(t)^{\frac{1}{N-1}} \delta(t) \quad \forall 1 \leq j \leq k. \quad (2.21)$$

Proof. We aim to use the characterization given by Theorem 1.2. To do so, for every $1 \leq j \leq k$ we define K_j as the linear subspace of $W_0^{1,2}(\tilde{\Omega}(t))$ spanned by the functions \tilde{u}_i with $1 \leq i \leq j$. First of all, we give the

Claim A. For every $1 \leq j \leq k$, the space K_j has dimension j .

Suppose that the claim is not true. Then, there should exist some $1 \leq \ell \leq k$ and some coefficients β_i for $i \neq \ell$ with all $|\beta_i| \leq 1$ and

$$\tilde{u}_\ell = \sum_{1 \leq i \leq k, i \neq \ell} \beta_i \tilde{u}_i.$$

Notice now that by (2.11) we know that $\int_{\Omega^+(t)} u_\ell^2 \geq 1/2$, hence also by (2.20) we deduce

$$\frac{1}{2} \leq \int_{\tilde{\Omega}(t)} \tilde{u}_\ell^2 = \int_{\tilde{\Omega}(t)} \sum_{1 \leq i \leq k, i \neq \ell} \beta_i \tilde{u}_i \tilde{u}_\ell \leq \sum_{1 \leq i \leq k, i \neq \ell} \left| \int_{\tilde{\Omega}(t)} \tilde{u}_i \tilde{u}_\ell \right| \leq k C \left(\nu^{\frac{3}{N-1}} + \nu^{\frac{1}{N-1}} \right) \nu < \frac{1}{2},$$

where the last inequality is true provided that $\nu = \nu(k, K, N)$ is chosen small enough. The absurd shows the validity of Claim A.

We can now show (2.21): to do so, pick a generic function $w \in K_j$, which can be written (up to a rescaling) as $w = \sum_{i=1}^j \beta_i \tilde{u}_i$, where $\max\{|\beta_i|, 1 \leq i \leq j\} = 1$. We need to evaluate $\mathcal{R}(w, \tilde{\Omega}(t))$: we start by noticing that

$$\begin{aligned} \mathcal{R}(w, \tilde{\Omega}(t)) &= \frac{\int_{\tilde{\Omega}(t)} |Dw|^2}{\int_{\tilde{\Omega}(t)} w^2} = \frac{\sum_{i=1}^j \beta_i^2 \int_{\tilde{\Omega}(t)} |D\tilde{u}_i|^2 + \sum_{i \neq j} \beta_i \beta_j \int_{\tilde{\Omega}(t)} D\tilde{u}_i \cdot D\tilde{u}_j}{\sum_{i=1}^j \beta_i^2 \int_{\tilde{\Omega}(t)} \tilde{u}_i^2 + \sum_{i \neq j} \beta_i \beta_j \int_{\tilde{\Omega}(t)} \tilde{u}_i \tilde{u}_j} \\ &\leq \frac{\sum_{i=1}^j \beta_i^2 \int_{\tilde{\Omega}(t)} |D\tilde{u}_i|^2 + Ck^2 \varepsilon(t)^{\frac{1}{N-1}} \delta(t)}{\sum_{i=1}^j \beta_i^2 \int_{\tilde{\Omega}(t)} \tilde{u}_i^2 - Ck^2 \varepsilon(t)^{\frac{1}{N-1}} \delta(t)}, \end{aligned} \quad (2.22)$$

where the last inequality comes by (2.20). If $\nu(k, K, N)$ is small enough, then

$$Ck^2 \varepsilon(t)^{\frac{1}{N-1}} \delta(t) \leq Ck^2 \nu^{\frac{N}{N-1}} \leq \frac{1}{4},$$

hence by the choice of β_i and by (2.11) the denominator in the last fraction of (2.22) is bigger than $1/4$ (in particular, it is strictly positive). As a consequence, recalling also that by (2.19) one has for every $1 \leq i \leq j$

$$\frac{\int_{\tilde{\Omega}(t)} |D\tilde{u}_i|^2}{\int_{\tilde{\Omega}(t)} \tilde{u}_i^2} \leq \lambda_i(\Omega) + C \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t) \leq \lambda_j(\Omega) + C \varepsilon(t)^{\frac{1}{N-1}} \delta(t),$$

from (2.22) we deduce

$$\begin{aligned} \mathcal{R}(w, \tilde{\Omega}(t)) &\leq \frac{\left(\lambda_j(\Omega) + C\varepsilon(t)^{\frac{1}{N-1}}\delta(t)\right) \left(\sum_{i=1}^j \beta_i^2 \int_{\tilde{\Omega}(t)} \tilde{u}_i^2\right) + C\varepsilon(t)^{\frac{1}{N-1}}\delta(t)}{\sum_{i=1}^j \beta_i^2 \int_{\tilde{\Omega}(t)} \tilde{u}_i^2 - C\varepsilon(t)^{\frac{1}{N-1}}\delta(t)} \\ &\leq \frac{\left(\lambda_j(\Omega) + C\varepsilon(t)^{\frac{1}{N-1}}\delta(t)\right) + 2C\varepsilon(t)^{\frac{1}{N-1}}\delta(t)}{1 - 2C\varepsilon(t)^{\frac{1}{N-1}}\delta(t)} \leq \lambda_j(\Omega) + C\varepsilon(t)^{\frac{1}{N-1}}\delta(t) \end{aligned}$$

(keep in mind that the constant $C = C(k, K, N)$ may increase from line to line). The validity of (2.21) is then now an immediate consequence of Theorem 1.2 and of Claim A. \square

We can now enter in the central part of our construction. Basically, we aim to show that either Ω already satisfies the requirements of Lemma 2.2, or some $\tilde{\Omega}(t)$ does it, up to a rescaling. To do so, we need another definition, namely, for every $t \leq \bar{t}$ we define the rescaled set

$$\widehat{\Omega}(t) := |\tilde{\Omega}(t)|^{-\frac{1}{N}} \tilde{\Omega}(t),$$

so that $|\widehat{\Omega}(t)| = 1$. We can now show the following result.

Lemma 2.7. *Let Ω be as in the assumptions of Lemma 2.2, and let $t \leq \bar{t}$. There exists $C_4 = C_4(k, K, N)$ such that exactly one of the three following conditions hold:*

- (1) $\max\{\varepsilon(t), \delta(t)\} > \nu$;
- (2) (1) does not hold and $m(t) \leq C_4(\varepsilon(t) + \delta(t))\varepsilon(t)^{\frac{1}{N-1}}$;
- (3) (1) and (2) do not hold and for every $1 \leq i \leq k$, one has $\lambda_i(\widehat{\Omega}(t)) < \lambda_i(\Omega)$.

In particular, if condition (3) holds for t and $m(t) \geq \widehat{m}$, then for every $1 \leq i \leq k$ one has $\lambda_i(\widehat{\Omega}(t)) < \lambda_i(\Omega) - \eta$, being $\eta = \eta(k, K, N) > 0$.

Proof. If (1) holds true, there is of course nothing to prove. Otherwise, it is possible to apply Lemma 2.6, hence we have

$$\lambda_i(\tilde{\Omega}(t)) \leq \lambda_i(\Omega) + C_3\varepsilon(t)^{\frac{1}{N-1}}\delta(t) \quad (2.23)$$

for every $1 \leq i \leq k$. By (1.2) and the fact that $|\widehat{\Omega}(t)| = 1$, we know that

$$\lambda_i(\widehat{\Omega}(t)) = |\tilde{\Omega}(t)|^{\frac{2}{N}} \lambda_i(\tilde{\Omega}(t)).$$

By construction,

$$|\tilde{\Omega}(t)| = |\Omega^+(t)| + |Q(t)| = 1 - m(t) + \varepsilon(t)^{\frac{N}{N-1}},$$

hence the above estimates and (2.23) lead to

$$\begin{aligned} \lambda_i(\widehat{\Omega}(t)) &= \left(1 - m(t) + \varepsilon(t)^{\frac{N}{N-1}}\right)^{\frac{2}{N}} \lambda_i(\tilde{\Omega}(t)) \\ &\leq \left(1 - \frac{2}{N}m(t) + \frac{2}{N}\varepsilon(t)^{\frac{N}{N-1}}\right) \left(\lambda_i(\Omega) + C_3\varepsilon(t)^{\frac{1}{N-1}}\delta(t)\right) \\ &\leq \lambda_i(\Omega) - \frac{2\lambda_1(B_N)}{N}m(t) + \frac{2K}{N}\varepsilon(t)^{\frac{N}{N-1}} + \left(C_3 + \frac{2}{N}\right)\varepsilon(t)^{\frac{1}{N-1}}\delta(t). \end{aligned} \quad (2.24)$$

This allows us to conclude. In fact, defining $C_4 := \frac{2(K+1)}{N} + C_3$, if $m(t) \leq C_4(\varepsilon(t) + \delta(t))\varepsilon(t)^{\frac{1}{N-1}}$, then condition (2) holds true. Otherwise, (2.24) directly implies that $\lambda_i(\widehat{\Omega}(t)) < \lambda_i(\Omega)$.

Finally, assume that condition (3) holds and $m(t) \geq \widehat{m}$: in this case, (2.24) directly implies

$$\lambda_i(\widehat{\Omega}(t)) - \lambda_i(\Omega) \leq -\frac{2\lambda_1(B_N)}{N} \widehat{m} + C_4 \nu^{\frac{N}{N-1}} \leq -\eta,$$

where $\eta = \lambda_1(B_N)\widehat{m}/N$ and the last inequality is true up to decrease ν (notice that *decreasing* the value of the constant ν of Lemma 2.6 does not change the value of C_3). \square

We are finally in position to give the proof of Lemma 2.2.

Proof of Lemma 2.2. Let us start defining

$$\hat{t} := \sup \left\{ t \leq \bar{t} : \text{condition (3) of Lemma 2.7 holds for } t \right\}, \quad (2.25)$$

with the usual convention that, if condition (3) is false for every $t \leq \bar{t}$, then $\hat{t} = -\infty$. We introduce now the following subsets of (\hat{t}, \bar{t}) ,

$$\begin{aligned} A &:= \left\{ t \in (\hat{t}, \bar{t}) : \text{condition (1) of Lemma 2.7 holds for } t \right\}, \\ B &:= \left\{ t \in (\hat{t}, \bar{t}) : \text{condition (2) of Lemma 2.7 holds for } t \text{ and } m(t) > 0 \right\}, \end{aligned}$$

and we further subdivide them as

$$\begin{aligned} A_1 &:= \left\{ t \in A : \varepsilon(t) \geq \delta(t) \right\}, & A_2 &:= \left\{ t \in A : \varepsilon(t) < \delta(t) \right\}, \\ B_1 &:= \left\{ t \in B : \varepsilon(t) \geq \delta(t) \right\}, & B_2 &:= \left\{ t \in B : \varepsilon(t) < \delta(t) \right\}. \end{aligned}$$

We aim to show that both A and B are uniformly bounded. Concerning A_1 , observe that

$$\nu |A_1| \leq \int_{A_1} \varepsilon(t) dt = \left| \left\{ (x, y) \in \Omega : x \in A_1 \right\} \right| \leq |\Omega| = 1,$$

so that $|A_1| \leq 1/\nu$. Concerning A_2 , in the same way and also recalling that $\lambda_i(\Omega) \leq K$ for every $i \leq k$, we have

$$\nu |A_2| \leq \int_{A_2} \delta(t) dt = \sum_{i=1}^k \int_{A_2} \int_{\Omega_t} |Du_i(t, y)|^2 d\mathcal{H}^{N-1}(y) dt \leq \sum_{i=1}^k \int_{\Omega} |Du_i|^2 \leq kK,$$

so that $|A_2| \leq kK/\nu$. Summarizing, we have proved that

$$|A| \leq \frac{1 + kK}{\nu}. \quad (2.26)$$

Let us then pass to the set B_1 . To deal with it, we need a further subdivision, namely, we write $B_1 = \cup_{n \in \mathbb{N}} B_1^n$, where

$$B_1^n := \left\{ t \in B_1 : \frac{\widehat{m}}{2^n} < m(t) \leq \frac{\widehat{m}}{2^{n-1}} \right\}. \quad (2.27)$$

Keeping in mind (2.5), we know that $t \mapsto m(t)$ is an increasing function, and that for a.e. $t \in \mathbb{R}$ one has $m'(t) = \varepsilon(t)$. Moreover, for every $t \in B_1$ one has by construction that

$$m(t) \leq C_4(\varepsilon(t) + \delta(t))\varepsilon(t)^{\frac{1}{N-1}} \leq 2C_4 \varepsilon(t)^{\frac{N}{N-1}}.$$

As a consequence, for every $t \in B_1^n$ one has

$$m'(t) = \varepsilon(t) \geq \frac{1}{C} m(t)^{\frac{N-1}{N}} \geq \frac{1}{C} \widehat{m}^{\frac{N-1}{N}} \frac{1}{(2^{\frac{N-1}{N}})^n}.$$

This readily implies

$$\frac{1}{C} \widehat{m}^{\frac{N-1}{N}} \frac{1}{(2^{\frac{N-1}{N}})^n} |B_1^n| \leq \int_{B_1^n} m'(t) \leq \frac{\widehat{m}}{2^n},$$

which in turn gives

$$|B_1^n| \leq C \widehat{m}^{\frac{1}{N}} (2^{-\frac{1}{N}})^n.$$

Finally, we deduce

$$|B_1| = \sum_{n \in \mathbb{N}} |B_1^n| \leq C \widehat{m}^{\frac{1}{N}} \sum_{n \in \mathbb{N}} (2^{-\frac{1}{N}})^n = C \widehat{m}^{\frac{1}{N}} \frac{2^{\frac{1}{N}}}{2^{\frac{1}{N}} - 1}. \quad (2.28)$$

Notice that basically our argument consisted in using the fact that in B_1 one has

$$m(t) \leq C \varepsilon(t)^{\frac{N}{N-1}}, \quad \text{with } \varepsilon(t) = m'(t). \quad (2.29)$$

Concerning B_2 , we can almost repeat the same argument: in fact, thanks to (2.9), for every $t \in B_2$ we have

$$\phi(t) = \sum_{i=1}^k \int_{\Omega^-(t)} |Du_i|^2 \leq C_1 \varepsilon(t)^{\frac{1}{N-1}} \delta(t) \leq C_1 \delta(t)^{\frac{N}{N-1}}, \quad \text{with } \delta(t) = \phi'(t).$$

which is the perfect analogous of (2.29) with δ and ϕ in place of ε and m respectively. Since as already observed $\phi(\bar{t}) \leq \sum \int_{\Omega} |Du_i|^2 \leq kK$, in analogy with (2.27) we can define

$$B_2^n := \left\{ t \in B_2 : \frac{kK}{2^{n+1}} < \phi(t) \leq \frac{kK}{2^n} \right\},$$

thus the very same argument which lead to (2.28) now gives

$$|B_2| = \sum_{n \in \mathbb{N}} |B_2^n| \leq C (kK)^{\frac{1}{N}} \sum_{n \in \mathbb{N}} (2^{-\frac{1}{N}})^n = C (kK)^{\frac{1}{N}} \frac{2^{\frac{1}{N}}}{2^{\frac{1}{N}} - 1}. \quad (2.30)$$

Putting (2.26), (2.28) and (2.30) together, we find

$$|A| + |B| \leq C_5 = C_5(k, K, N). \quad (2.31)$$

We will prove the validity of the lemma with the following choice of R_1 and Γ_1 ,

$$R_1 = 2C_5 + 4, \quad \Gamma_1 = 2^{\lceil K/\eta \rceil + 1},$$

where $C_5 = C_5(k, K, N)$ and $\eta = \eta(k, K, N)$ have been introduced in (2.31) and in Lemma 2.7 respectively. To obtain our proof, we will distinguish the possible cases for Ω .

Case I. One has $\hat{t} = -\infty$.

If this case happens, then condition (3) of Lemma 2.7 never holds true, i.e., for every $t \leq \bar{t}$ either condition (1) or (2) holds. Recalling the definition of A and B and (2.31), we deduce that $W(\Omega, 0, \widehat{m}) \leq C_5$. Therefore, the claim of Lemma 2.2 is immediately obtained simply taking $\widehat{\Omega} = \Omega$, since $R_1 \geq C_5$ and $\Gamma_1 \geq 1$.

Case II. One has $\hat{t} > -\infty$.

In this case, let us notice that it must be $m(\hat{t}) > 0$, hence $(\hat{t}, \bar{t}) \subseteq A \cup B$ and thus by (2.31) $\hat{t} \geq \bar{t} - C_5$. Let us now pick some $t^* \in [\hat{t} - 1, \hat{t}]$ for which condition (3) holds, and define $U_1 := \widehat{\Omega}(t^*)$. By definition, U_1 has unit volume, and $\lambda_i(U_1) < \lambda_i(\Omega)$ for every $1 \leq i \leq k$, being condition (3) true for t^* .

Observe now that by definition for every $2 \leq p \leq N$ one has $\pi_p(\widetilde{\Omega}(t^*)) = \pi_p(\Omega^+(t^*))$, hence

$$\begin{aligned} \text{diam}(\pi_p(U_1)) &= \text{diam}(\pi_p(\widehat{\Omega}(t^*))) = \text{diam}\left(\pi_p\left(|\widetilde{\Omega}(t^*)|^{-\frac{1}{N}}\widetilde{\Omega}(t^*)\right)\right) \leq 2 \text{diam}\left(\pi_p(\widetilde{\Omega}(t^*))\right) \\ &= 2 \text{diam}\left(\pi_p(\Omega^+(t^*))\right) \leq 2 \text{diam}(\pi_p(\Omega)), \end{aligned}$$

where we have used that $|\widetilde{\Omega}(t^*)| \geq 1/2$. Concerning the widths of U_1 and Ω , we can start observing that

$$W(U_1, \widehat{m}, 1) = |\widetilde{\Omega}(t^*)|^{-\frac{1}{N}} \left(W\left(\widetilde{\Omega}(t^*), \widehat{m}|\widetilde{\Omega}(t^*)|, |\widetilde{\Omega}(t^*)|\right) \right).$$

Moreover, since it is admissible to assume $\nu^{\frac{N}{N-1}} < \frac{\widehat{m}}{2}$ and then

$$|\widetilde{\Omega}(t^*)|_{t^*}^l = |\varepsilon(t^*)^{\frac{N}{N-1}}| < |\widetilde{\Omega}(t^*)|\widehat{m},$$

we have

$$\tau\left(\widetilde{\Omega}(t^*), \widehat{m}|\widetilde{\Omega}(t^*)|\right) = \tau\left(\Omega, \widehat{m}|\widetilde{\Omega}(t^*)| + 1 - |\widetilde{\Omega}(t^*)|\right);$$

as a consequence, we evaluate

$$\begin{aligned} W\left(\widetilde{\Omega}(t^*), \widehat{m}|\widetilde{\Omega}(t^*)|, |\widetilde{\Omega}(t^*)|\right) &= \tau\left(\widetilde{\Omega}(t^*), |\widetilde{\Omega}(t^*)|\right) - \tau\left(\widetilde{\Omega}(t^*), \widehat{m}|\widetilde{\Omega}(t^*)|\right) \\ &= \tau(\Omega, 1) - \tau\left(\Omega, \widehat{m}|\widetilde{\Omega}(t^*)| + 1 - |\widetilde{\Omega}(t^*)|\right) \leq \tau(\Omega, 1) - \tau(\Omega, \widehat{m}) \\ &= W(\Omega, \widehat{m}, 1), \end{aligned}$$

thus we deduce, again recalling $|\widetilde{\Omega}(t^*)| \geq 1/2$, that

$$W(U_1, \widehat{m}, 1) \leq 2W(\Omega, \widehat{m}, 1).$$

Summarizing, we have found that

$$\lambda_i(U_1) < \lambda_i(\Omega), \quad \text{diam}(\pi_p(U_1)) \leq 2 \text{diam}(\pi_p(\Omega)), \quad W(U_1, \widehat{m}, 1) \leq 2W(\Omega, \widehat{m}, 1). \quad (2.32)$$

As a consequence, the choice $\widehat{\Omega} = U_1$ satisfies all the requirements of Lemma 2.2, except possibly condition (2.2). To deal with this last condition, we need to further subdivide this case.

Case IIa. One has $\hat{t} > -\infty$ and $m(t^*) < \widehat{m}$.

In this case, we can show that the choice $\widehat{\Omega} = U_1$ actually works. As noticed above, we have only to prove the validity of (2.2). To do so, we assume for simplicity that $\bar{t} = 0$, which is clearly admissible by translation. Hence, $t^* \geq \hat{t} - 1 \geq \bar{t} - C_5 - 1 = -C_5 - 1$, and thus

$$\widetilde{\Omega}(t^*) = \Omega^+(t^*) \cup Q(t^*) \subseteq \{(x, y) : x > t^* - 1\} \subseteq \{(x, y) : x > -C_5 - 2\}.$$

Recalling that $|\widetilde{\Omega}(t^*)| \geq |\Omega^+(t^*)| \geq |\Omega^+(\bar{t})| = 1 - 2\widehat{m} \geq 1/2$, we deduce that

$$\widehat{\Omega} = \widehat{\Omega}(t^*) = |\widetilde{\Omega}(t^*)|^{-\frac{1}{N}}\widetilde{\Omega}(t^*) \subseteq 2\widetilde{\Omega}(t^*) \subseteq \{(x, y) : x > -2C_5 - 4\}. \quad (2.33)$$

Moreover, since $m(\hat{t}) < \hat{m}$,

$$\begin{aligned} |\widehat{\Omega}'_0| &= |\widehat{\Omega}(t^*)'_0| \geq |\widetilde{\Omega}(t^*)'_0| = \left| \left(\Omega^+(t^*) \cup Q(t^*) \right)'_0 \right| \geq \left| \left(\Omega^+(t^*) \right)'_0 \right| \\ &= \left| \{(x, y) \in \Omega : t^* < x < 0\} \right| = m(0) - m(t^*) \geq m(0) - m(\hat{t}) \geq \hat{m}, \end{aligned}$$

and this implies that $\tau(\widehat{\Omega}, \hat{m}) \leq 0$. The inclusion (2.33) ensures then that $\tau(\widehat{\Omega}, 0) \geq -2C_5 - 4$, and then (2.2) holds true since $R_1 = 2C_5 + 4$.

Case IIb. One has $\hat{t} > -\infty$ and $m(t^*) \geq \hat{m}$.

We have now to face the last possible case, namely, when \hat{t} is finite but $m(t^*) \geq \hat{m}$. In this case, thanks to Lemma 2.7 the estimates (2.32) can be strengthened as

$$\lambda_i(U_1) < \lambda_i(\Omega) - \eta, \quad \text{diam}(\pi_p(U_1)) \leq 2 \text{diam}(\pi_p(\Omega)), \quad W(U_1, \hat{m}, 1) \leq 2W(\Omega, \hat{m}, 1). \quad (2.34)$$

Concerning the validity of (2.2), it does not follow by (2.33) because the assumption $m(\hat{t}) \geq \hat{m}$ does not imply that $\tau(\widehat{\Omega}, \hat{m}) \leq 0$. However, we can argue as follow: if (2.2) holds true for U_1 , then of course we are done by setting $\widehat{\Omega} = U_1$. Otherwise, we apply the above construction to the set U_1 in place of Ω : since U_1 does not satisfy (2.2), then Case I is impossible, thus we are in Case II and then by (2.32) we find an open set U_2 of unit measure such that

$$\begin{cases} \lambda_i(U_2) < \lambda_i(U_1) < \lambda_i(\Omega) - \eta \quad \forall 1 \leq i \leq k, \\ \text{diam}(\pi_p(U_2)) \leq 2 \text{diam}(\pi_p(U_1)) \leq 4 \text{diam}(\pi_p(\Omega)), \\ W(U_2, \hat{m}, 1) \leq 2W(U_1, \hat{m}, 1) \leq 4W(\Omega, \hat{m}, 1). \end{cases}$$

If U_1 is in Case IIa, then as before we are done with the choice of $\widehat{\Omega} = U_2$. Otherwise, if we are in Case IIb, then (2.34) becomes

$$\lambda_i(U_2) < \lambda_i(\Omega) - 2\eta, \quad \text{diam}(\pi_p(U_2)) \leq 4 \text{diam}(\pi_p(\Omega)), \quad W(U_2, \hat{m}, 1) \leq 4W(\Omega, \hat{m}, 1).$$

Going on with the obvious iteration, if the proof has not been concluded after $\ell \in \mathbb{N}$ steps then we have found an open set U_ℓ satisfying

$$\lambda_i(U_\ell) < \lambda_i(\Omega) - \ell\eta, \quad \text{diam}(\pi_p(U_\ell)) \leq 2^\ell \text{diam}(\pi_p(\Omega)), \quad W(U_\ell, \hat{m}, 1) \leq 2^\ell W(\Omega, \hat{m}, 1).$$

This is of course impossible if $\ell\eta \geq K$, being $\lambda_k(\Omega) \leq K$: as a consequence, our iteration must stop after less than K/η steps, thus our thesis is concluded with our choice of Γ_1 . \square

2.2. Boundedness of the interior. The goal of this subsection is to obtain a uniform bound also for the interior part of a set Ω , in the sense of Lemma 2.2. Most of the arguments of this case will be identical to those that we made for the tails in Section 2.1, but some modifications are essential. In particular we give new definitions for ε , δ_i , μ_i , $\widetilde{\Omega}(t)$ and $\widehat{\Omega}(t)$ in order to maintain the analogy with what was done in Section 2.1. The result that we are going to prove is the following.

Lemma 2.8. *For every $K > 0$ there exist $R_2 = R_2(k, K, N)$ and $\Gamma_2 = \Gamma_2(k, K, N)$ such that, for any open set $\Omega \subseteq \mathbb{R}^N$ of unit volume and with $\lambda_k(\Omega) \leq K$, and for any choice of $\bar{m} \in (\hat{m}, 1 - \frac{\hat{m}}{2})$,*

there exists another open set $\widehat{\Omega} \subseteq \mathbb{R}^N$, still of unit volume, such that $\lambda_i(\widehat{\Omega}) \leq \lambda_i(\Omega)$ for every $1 \leq i \leq k$, and such that for every $2 \leq p \leq N$

$$W(\widehat{\Omega}, 0, \bar{m}) \leq R_2 + \Gamma_2 W(\Omega, 0, \bar{m} - \widehat{m}), \quad \text{diam}(\pi_p(\widehat{\Omega})) \leq \Gamma_2 \text{diam}(\pi_p(\Omega)). \quad (2.35)$$

To start with, we give the analogous of the definitions (2.4), (2.5) and (2.6) of Section 2.1 that we need now; Figure 3 helps to visualize the new situation. More precisely, we set for brevity

$$t_0 := \frac{\tau(\Omega, \bar{m} + \frac{\widehat{m}}{2}) + \tau(\Omega, \bar{m} - \widehat{m})}{2}, \quad \bar{t} := \frac{\tau(\Omega, \bar{m} + \frac{\widehat{m}}{2}) - \tau(\Omega, \bar{m} - \widehat{m})}{2};$$

keep in mind that, since $\bar{m} \in (\widehat{m}, 1 - \frac{\widehat{m}}{2})$, then $-\infty < \tau(\Omega, \bar{m} - \widehat{m}) < \tau(\Omega, \bar{m} + \frac{\widehat{m}}{2}) < +\infty$. For any $0 \leq t \leq \bar{t}$, we define

$$\begin{aligned} \Omega^+(t) &:= \Omega_{t_0-t}^l \cup \Omega_{t_0+t}^r, & \Omega^-(t) &:= \Omega_{t_0-t}^r \cap \Omega_{t_0+t}^l = \Omega \setminus \Omega^+(t), \\ \varepsilon(t) &:= \mathcal{H}^{N-1}(\Omega_{t_0-t}) + \mathcal{H}^{N-1}(\Omega_{t_0+t}), & m(t) &:= |\Omega^-(t)| = \int_0^t \varepsilon(s) ds \leq \frac{3}{2} \widehat{m}. \end{aligned}$$

Moreover, having fixed an orthonormal set $\{u_1, u_2, \dots, u_k\}$ of eigenfunctions with unit L^2 norm corresponding to the first k eigenvalues of Ω , for every $1 \leq i \leq k$ and $0 \leq t \leq \bar{t}$ we define

$$\delta_i(t) := \int_{\Omega_{t_0-t}} |Du_i|^2 + \int_{\Omega_{t_0+t}} |Du_i|^2, \quad \mu_i(t) := \int_{\Omega_{t_0-t}} u_i^2 + \int_{\Omega_{t_0+t}} u_i^2.$$

In analogy with (2.7), we define again $\delta(t) = \sum_{i=1}^k \delta_i(t)$, and we set again

$$\phi(t) := \sum_{i=1}^k \int_{\Omega^-(t)} |Du_i|^2 = \int_0^t \delta(s) ds.$$

Our strategy to prove Lemma 2.8 is very similar to what we did to show Lemma 2.2; in fact,




FIGURE 3. A set Ω and the corresponding quantities t_0 , \bar{t} and sets $\Omega^+(t)$ (white) and $\Omega^-(t)$ (shaded).

basically the only difference is that to show the analogous of Lemma 2.3 we cannot rely on the symmetrization of $\Omega^-(t)$. Let us see how to overcome this difficulty.

Lemma 2.9. *There exists a small constant $\nu = \nu(k, K, N) < 1$ such that, if Ω and \bar{m} are as in the assumptions of Lemma 2.8, and $0 \leq t \leq \bar{t}$ is such that $\varepsilon(t), \delta(t) \leq \nu$, then for every $1 \leq i \leq k$ one has*

$$\int_{\Omega^-(t)} u_i^2 \leq C\varepsilon(t)^{\frac{1}{N-1}} \delta_i(t), \quad \int_{\Omega^-(t)} |Du_i|^2 \leq C\varepsilon(t)^{\frac{1}{N-1}} \delta_i(t). \quad (2.36)$$

Proof. Consider the “external cylinders”

$$Q_1 := (t_0 - t - \sigma_1, t_0 - t) \times \Omega_{t_0-t}, \quad Q_2 := (t_0 + t, t_0 + t + \sigma_2) \times \Omega_{t_0+t},$$

where

$$\sigma_1 = \mathcal{H}^{N-1}(\Omega_{t_0-t})^{\frac{1}{N-1}}, \quad \sigma_2 = \mathcal{H}^{N-1}(\Omega_{t_0+t})^{\frac{1}{N-1}},$$

in perfect analogy with (2.14) and (2.15). Calling $U = \Omega^-(t) \cup Q_1 \cup Q_2$, we can extend (2.16) to obtain the following definition of $\tilde{u}_i \in W_0^{1,2}(U)$,

$$\tilde{u}_i(x, y) := \begin{cases} u_i(x, y) & \text{if } (x, y) \in \Omega^-(t), \\ \frac{x - (t_0 - t - \sigma_1)}{\sigma_1} u_i(t_0 - t, y) & \text{if } (x, y) \in Q_1, \\ \frac{(t_0 + t + \sigma_2) - x}{\sigma_2} u_i(t_0 + t, y) & \text{if } (x, y) \in Q_2. \end{cases}$$

Applying Lemma 2.4 to the two cylinders Q_1 and Q_2 , and comparing the present definitions of ε and δ_i with those that we used in Section 2.1, (2.18) gives us

$$\int_{Q_1 \cup Q_2} |D\tilde{u}_i|^2 \leq C_2 \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t).$$

We can then obtain an estimate between $\int_{\Omega^-(t)} u_i^2$ and $\int_{\Omega^-(t)} |Du_i|^2$ similar to (2.10), first noticing that

$$\frac{\lambda_1(B_N)}{(m(t) + \varepsilon(t)^{\frac{N}{N-1}})^{\frac{2}{N}}} \leq \frac{\lambda_1(B_N)}{|U|^{\frac{2}{N}}} \leq \lambda_1(U) \leq \mathcal{R}(\tilde{u}_i, U) \leq \frac{\int_{\Omega^-(t)} |Du_i|^2 + \int_{Q_1 \cup Q_2} |D\tilde{u}_i|^2}{\int_{\Omega^-(t)} u_i^2},$$

and then deducing, recalling (2.1) and choosing ν small enough,

$$\begin{aligned} \int_{\Omega^-(t)} u_i^2 &\leq \frac{(m(t) + \varepsilon(t)^{\frac{N}{N-1}})^{\frac{2}{N}}}{\lambda_1(B_N)} \left(\int_{\Omega^-(t)} |Du_i|^2 + C_2 \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t) \right) \\ &\leq \frac{1}{2^{1+\frac{1}{N}} K} \int_{\Omega^-(t)} |Du_i|^2 + \frac{C_2}{2^{1+\frac{1}{N}} K} \varepsilon(t)^{\frac{1}{N-1}} \delta_i(t). \end{aligned} \quad (2.37)$$

Observe that $\nu = \nu(k, K, N)$ can be chosen so small that the last estimate implies

$$\int_{\Omega^-(t)} u_i^2 \leq \frac{1}{2}.$$

We can also generalize (2.12); in fact, since the very same argument used in (2.8) again ensures that $\mu_i(t) \leq C\varepsilon(t)^{\frac{2}{N-1}}\delta_i(t)$, we can obtain

$$\begin{aligned} \int_{\Omega^-(t)} |Du_i|^2 &= \int_{\Omega^-(t)} \lambda_i u_i^2 + \int_{\Omega_{t_0-t} \cup \Omega_{t_0+t}} u_i \frac{\partial u_i}{\partial \nu} \leq K \int_{\Omega^-(t)} u_i^2 + \sqrt{\mu_i(t)\delta_i(t)} \\ &\leq K \int_{\Omega^-(t)} u_i^2 + C\varepsilon(t)^{\frac{1}{N-1}}\delta_i(t). \end{aligned} \quad (2.38)$$

Putting together (2.37) and (2.38) gives (2.36). \square

We need now to extend the result of Lemma 2.6 to our new setting. To do so, going on in analogy with Section 2.1, we give the following definition.

Definition 2.10. *Let Ω , \bar{m} and $1 \leq t \leq \bar{t}$ be as in the assumptions of Lemma 2.9. Consider the “internal cylinders”*

$$Q_1 := (t_0 - t, t_0 - t + \sigma_1) \times \Omega_{t_0-t}, \quad Q_2 := (t_0 + t - \sigma_2, t_0 + t) \times \Omega_{t_0+t},$$

where

$$\sigma_1 = \mathcal{H}^{N-1}(\Omega_{t_0-t})^{\frac{1}{N-1}}, \quad \sigma_2 = \mathcal{H}^{N-1}(\Omega_{t_0+t})^{\frac{1}{N-1}},$$

and notice that by the assumption on $\varepsilon(t)$ and the fact that $t \geq 1$ one has $Q_1 \cap Q_2 = \emptyset$. The set $\tilde{\Omega}(t)$ is defined as

$$\begin{aligned} \tilde{\Omega}(t) := \left\{ (x, y) \in \mathbb{R}^N : \text{either } x \leq t_0, (x - t + \sigma_1, y) \in \Omega^+(t) \cup Q_1, \right. \\ \left. \text{or } x \geq t_0, (x + t - \sigma_2, y) \in \Omega^+(t) \cup Q_2 \right\}, \end{aligned}$$

see Figure 4. Notice that

$$\begin{aligned} |\tilde{\Omega}(t)| &= |\Omega^+(t)| + |Q_1| + |Q_2| = 1 - m(t) + \mathcal{H}^{N-1}(\Omega_{t_0-t})^{\frac{N}{N-1}} + \mathcal{H}^{N-1}(\Omega_{t_0+t})^{\frac{N}{N-1}} \\ &\leq 1 - m(t) + \varepsilon(t)^{\frac{N}{N-1}}. \end{aligned}$$

Moreover, define again the rescaled set

$$\hat{\Omega}(t) := |\tilde{\Omega}(t)|^{-\frac{1}{N}} \tilde{\Omega}(t).$$




FIGURE 4. A set Ω and the corresponding set $\tilde{\Omega}(t)$.

With this definition of the sets $\tilde{\Omega}(t)$ and with the obvious extension of (2.16) in order to define $\tilde{u}_i \in W_0^{1,2}(\tilde{\Omega}(t))$, we can now literally repeat the proofs of Lemmas 2.4, 2.5, 2.6 and 2.7, the

unique difference being the substitution of $Q(t)$ with $Q_1 \cup Q_2$, and of Ω_t with $\Omega_{t_0+t} \cup \Omega_{t_0-t}$. We obtain then the following result, which holds up to possibly decrease the constant $\nu = \nu(k, K, N)$ of Lemma 2.9.

Lemma 2.11. *Let Ω be as in the assumptions of Lemma 2.8, and let $1 \leq t \leq \bar{t}$. There exists $C_6 = C_6(k, K, N)$ such that exactly one of the three following conditions hold:*

- (1) $\max\{\varepsilon(t), \delta(t)\} > \nu$;
- (2) (1) does not hold and $m(t) \leq C_6(\varepsilon(t) + \delta(t))\varepsilon(t)^{\frac{1}{N-1}}$;
- (3) (1) and (2) do not hold and for every $1 \leq i \leq k$, one has $\lambda_i(\widehat{\Omega}(t)) < \lambda_i(\Omega)$.

In particular, if condition (3) holds for t and $m(t) \geq \widehat{m}/2$, then for every $1 \leq i \leq k$ one has $\lambda_i(\widehat{\Omega}(t)) < \lambda_i(\Omega) - \eta$, being $\eta = \eta(k, K, N) > 0$.

We can now conclude this section by presenting the proof of Lemma 2.8, which will be a minor modification of the proof of Lemma 2.2.

Proof of Lemma 2.8. First of all, we want to show that it is admissible to assume

$$m(t) > 0 \quad \forall t > 0. \quad (2.39)$$

In fact, suppose that it is not so, and let $\tau = \max\{0 \leq t \leq \bar{t} : m(t) = 0\} > 0$. Then, Ω is the disjoint union of $\Omega \cap \{x > t_0 + \tau\}$ and $\Omega \cap \{x < t_0 - \tau\}$, and it does not intersect the whole strip $\{t_0 - \tau < x < t_0 + \tau\}$. Therefore, replacing Ω with $\{(x + \tau, y) : x < t_0, (x, y) \in \Omega\} \cup \{(x - \tau, y) : x > t_0, (x, y) \in \Omega\}$, that is, moving closer the two disjoint parts of Ω , does not change any of the eigenvalues of Ω and is clearly admissible for the proof of the lemma; moreover, the property (2.39) of course holds true for this new set. Hence, from now on we directly assume that (2.39) holds true for Ω .

Define now \hat{t} analogously to (2.25) by setting

$$\hat{t} := \sup \left\{ 1 \leq t \leq \bar{t} : \text{condition (3) of Lemma 2.11 holds for } t \right\},$$

with the convention that, if condition (3) is false for every $1 \leq t \leq \bar{t}$, then $\hat{t} = 1$. Again we define A and B as

$$\begin{aligned} A &:= \left\{ t \in (\hat{t}, \bar{t}) : \text{condition (1) of Lemma 2.11 holds for } t \right\}, \\ B &:= \left\{ t \in (\hat{t}, \bar{t}) : \text{condition (2) of Lemma 2.11 holds for } t \text{ and } m(t) > 0 \right\}. \end{aligned}$$

The same argument of the proof of Lemma 2.2 gives then

$$|A| + |B| \leq C_7 = C_7(k, K, N), \quad (2.40)$$

and we are going to show the thesis with the choice

$$R_2 = 4C_7 + 8, \quad \Gamma_2 = 2^{\lfloor K/\eta \rfloor + 1},$$

being η the constant of Lemma 2.11. We can again subdivide the possible cases for Ω .

Case I. One has $\hat{t} = 1$.

In this case, by (2.39) one has that every $1 < t \leq \bar{t}$ belongs either to A or to B , thus by (2.40)

$\bar{t} \leq C_7 + 1$; as a consequence, the choice $\widehat{\Omega} = \Omega$ satisfies the requirements of the lemma. Indeed, while the right condition of (2.35) is obviously true, the left one follows just noticing that

$$\begin{aligned} W(\widehat{\Omega}, 0, \bar{m}) &= W(\Omega, 0, \bar{m}) = \tau(\Omega, \bar{m}) - \tau(\Omega, \bar{m} - \widehat{m}) + W(\Omega, 0, \bar{m} - \widehat{m}) \\ &\leq \tau\left(\Omega, \bar{m} + \frac{\widehat{m}}{2}\right) - \tau(\Omega, \bar{m} - \widehat{m}) + W(\Omega, 0, \bar{m} - \widehat{m}) = 2\bar{t} + W(\Omega, 0, \bar{m} - \widehat{m}) \\ &\leq 2(C_7 + 1) + W(\Omega, 0, \bar{m} - \widehat{m}). \end{aligned}$$

Case II. One has $\hat{t} > 1$.

In this case, again by (2.39) we know that $A \cup B$ contains the whole segment (\hat{t}, \bar{t}) , thus $\bar{t} \leq \hat{t} + C_7$ by (2.40). If we choose $t^* \in (\hat{t} - 1, \hat{t})$ for which condition (3) holds and define $U_1 := \widehat{\Omega}(t^*)$, we know by construction that $|U_1| = 1$ and $\lambda_i(U_1) < \lambda_i(\Omega)$ for every $1 \leq i \leq k$. As in the proof of Lemma 2.2, the fact that $|\widetilde{\Omega}(t^*)| \geq 1 - \frac{3}{2}\widehat{m} \geq 1/2$ ensures that $\text{diam}(\pi_p(U_1)) \leq 2 \text{diam}(\pi_p(\Omega))$ for each $2 \leq p \leq N$. On the other hand, concerning the width of U_1 , recalling the definition of $\widetilde{\Omega}(t^*)$ and observing that $(\bar{m} - \widehat{m})|\widetilde{\Omega}(t^*)| < \bar{m} - \widehat{m}$ one finds

$$\begin{aligned} W(U_1, 0, \bar{m} - \widehat{m}) &= W(\widehat{\Omega}(t^*), 0, \bar{m} - \widehat{m}) = |\widetilde{\Omega}(t^*)|^{-\frac{1}{N}} W(\widetilde{\Omega}(t^*), 0, (\bar{m} - \widehat{m})|\widetilde{\Omega}(t^*)|) \\ &\leq 2\left(\tau\left(\widetilde{\Omega}(t^*), (\bar{m} - \widehat{m})|\widetilde{\Omega}(t^*)|\right) - \tau(\widetilde{\Omega}(t^*), 0)\right) \\ &= 2\left(\tau\left(\Omega, (\bar{m} - \widehat{m})|\widetilde{\Omega}(t^*)|\right) - \tau(\Omega, 0)\right) = 2W(\Omega, 0, (\bar{m} - \widehat{m})|\widetilde{\Omega}(t^*)|) \\ &\leq 2W(\Omega, 0, \bar{m} - \widehat{m}). \end{aligned}$$

Summarizing, we have found that

$$\lambda_i(U_1) < \lambda_i(\Omega), \quad \frac{\text{diam}(\pi_p(U_1))}{\text{diam}(\pi_p(\Omega))} \leq 2, \quad \frac{W(U_1, 0, \bar{m} - \widehat{m})}{W(\Omega, 0, \bar{m} - \widehat{m})} \leq 2. \quad (2.41)$$

To conclude the inspection of the validity of (2.35), we will again need to consider separately two subcases.

Case IIa. One has $\hat{t} > 1$ and $m(t^*) \leq \widehat{m}/2$.

In this case, we can quickly observe that

$$\begin{aligned} W(U_1, 0, \bar{m}) &= |\widetilde{\Omega}(t^*)|^{-\frac{1}{N}} W(\widetilde{\Omega}(t^*), 0, \bar{m}|\widetilde{\Omega}(t^*)|) \leq 2W(\widetilde{\Omega}(t^*), 0, \bar{m} + \frac{\widehat{m}}{2} - |\Omega^-(t^*)|) \\ &\leq 2W(\Omega, 0, \bar{m} - \widehat{m}) + 4(\bar{t} - t^* + 1) \leq 2W(\Omega, 0, \bar{m} - \widehat{m}) + 4(C_7 + 2), \end{aligned} \quad (2.42)$$

and, together with (2.41), this concludes the proof of (2.35) and of the lemma with the choice $\widehat{\Omega} = U_1$.

Case IIb. One has $\hat{t} > 1$ and $m(t^*) > \widehat{m}/2$.

Let us conclude with this last case. By Lemma 2.11, in place of (2.41) we have then

$$\lambda_i(U_1) < \lambda_i(\Omega) - \eta, \quad \frac{\text{diam}(\pi_p(U_1))}{\text{diam}(\pi_p(\Omega))} \leq 2, \quad \frac{W(U_1, 0, \bar{m} - \widehat{m})}{W(\Omega, 0, \bar{m} - \widehat{m})} \leq 2,$$

which still does not guarantee the validity of (2.35). However, we can apply the construction above to U_1 : if U_1 is in Case I, then the choice $\widehat{\Omega} = U_1$ concludes the proof; otherwise, there

exists an open set U_2 of unit measure satisfying

$$\lambda_i(U_2) < \lambda_i(\Omega) - \eta, \quad \frac{\text{diam}(\pi_p(U_2))}{\text{diam}(\pi_p(\Omega))} \leq 4, \quad \frac{W(U_2, 0, \bar{m} - \hat{m})}{W(\Omega, 0, \bar{m} - \hat{m})} \leq 4. \quad (2.43)$$

If U_1 is in Case IIa then (2.42) gives

$$W(U_2, 0, \bar{m}) \leq 2W(U_1, 0, \bar{m} - \hat{m}) + 4(C_7 + 2) \leq 4W(\Omega, 0, \bar{m} - \hat{m}) + 4(C_7 + 2),$$

so the choice $\hat{\Omega} = U_2$ concludes the proof. Instead, if U_1 is in Case IIb then the first inequality of (2.43) becomes $\lambda_i(U_2) < \lambda_i(\Omega) - 2\eta$ for every $1 \leq i \leq k$. The obvious iteration ensures us that, if the proof has not been obtained after ℓ steps, then there must be some open set U_ℓ satisfying

$$\lambda_i(U_\ell) < \lambda_i(\Omega) - \ell\eta, \quad \frac{\text{diam}(\pi_p(U_\ell))}{\text{diam}(\pi_p(\Omega))} \leq 2^\ell, \quad \frac{W(U_\ell, 0, \bar{m} - \hat{m})}{W(\Omega, 0, \bar{m} - \hat{m})} \leq 2^\ell.$$

Since this is not possible for $\ell > K/\eta$, the iteration must stop at some $\ell \leq [K/\eta]$, and thus we conclude the proof thanks to the choice of Γ_2 . \square

2.3. Proof of Proposition 2.1. We are finally in position to give the proof of Proposition 2.1, which is now a simple consequence of Lemma 2.2 and Lemma 2.8.

Proof of Proposition 2.1. Let us pick a generic open set Ω with $\lambda_k(\Omega) \leq K$. Applying Lemma 2.2 to Ω , we find a set E_1 with

$$\lambda_i(E_1) \leq \lambda_i(\Omega), \quad W(E_1, 0, \hat{m}) \leq R_1, \quad \frac{\text{diam}(\pi_p(E_1))}{\text{diam}(\pi_p(\Omega))} \leq \Gamma_1,$$

for every $2 \leq p \leq N$. Then, we apply Lemma 2.8 to E_1 with $\bar{m} = 2\hat{m}$ finding E_2 which satisfies

$$\lambda_i(E_2) \leq \lambda_i(\Omega), \quad W(E_2, 0, 2\hat{m}) \leq R_2 + \Gamma_2 R_1, \quad \frac{\text{diam}(\pi_p(E_2))}{\text{diam}(\pi_p(\Omega))} \leq \Gamma_2 \Gamma_1.$$

Iterating, for any $\ell \geq 3$ such that $\ell\hat{m} \leq 1 - \frac{\hat{m}}{2}$ we apply Lemma 2.8 to $E_{\ell-1}$ with $\bar{m} = \ell\hat{m}$ finding E_ℓ such that

$$\lambda_i(E_\ell) \leq \lambda_i(\Omega), \quad W(E_\ell, 0, \ell\hat{m}) \leq R_2 \frac{\Gamma_2^{\ell-1} - 1}{\Gamma_2 - 1} + \Gamma_2^{\ell-1} R_1, \quad \frac{\text{diam}(\pi_p(E_\ell))}{\text{diam}(\pi_p(\Omega))} \leq \Gamma_2^{\ell-1} \Gamma_1.$$

Possibly applying a last time Lemma 2.8 with $\bar{m} = 1 - \hat{m}$, we have then found an open set E satisfying

$$\lambda_i(E) \leq \lambda_i(\Omega), \quad W(E, 0, 1 - \hat{m}) \leq R_2 \frac{\Gamma_2^{[1/\hat{m}] - 1} - 1}{\Gamma_2 - 1} + \Gamma_2^{[1/\hat{m}] - 1} R_1, \quad \frac{\text{diam}(\pi_p(E))}{\text{diam}(\pi_p(\Omega))} \leq \Gamma_2^{[1/\hat{m}] - 1} \Gamma_1.$$

Calling E' the set obtained by reflecting E with respect to the plane $\{x = 0\}$, the above estimates become

$$\lambda_i(E') \leq \lambda_i(\Omega), \quad W(E', \hat{m}, 1) \leq R_2 \frac{\Gamma_2^{[1/\hat{m}] - 1} - 1}{\Gamma_2 - 1} + \Gamma_2^{[1/\hat{m}] - 1} R_1, \quad \frac{\text{diam}(\pi_p(E'))}{\text{diam}(\pi_p(\Omega))} \leq \Gamma_2^{[1/\hat{m}] - 1} \Gamma_1,$$

so that applying once again Lemma 2.2 to E' we find a set F_1 satisfying

$$\lambda_i(F_1) \leq \lambda_i(\Omega), \quad \text{diam}(\pi_1(F_1)) = W(F_1, 0, 1) \leq R_3, \quad \frac{\text{diam}(\pi_p(F_1))}{\text{diam}(\pi_p(\Omega))} \leq \Gamma_3,$$

having set

$$R_3 := R_1 + \Gamma_1 R_2 \frac{\Gamma_2^{[1/\widehat{m}]-1} - 1}{\Gamma_2 - 1} + \Gamma_1 \Gamma_2^{[1/\widehat{m}]-1} R_1, \quad \Gamma_3 := \Gamma_2^{[1/\widehat{m}]-1} \Gamma_1^2.$$

We can now repeat the whole construction, using as starting set F_1 in place of Ω , and using the second coordinate in place of the first one. We will end up with a set F_2 with

$$\lambda_i(F_2) \leq \lambda_i(F_1) \leq \lambda_i(\Omega) \quad \forall 1 \leq i \leq k, \quad \text{diam}(\pi_2(F_2)) \leq R_3,$$

and such that for every $p \neq 2$ it is $\text{diam}(\pi_p(F_2)) \leq \Gamma_3 \text{diam}(\pi_p(F_1))$. In particular, choosing $p = 1$ we discover that $\text{diam}(\pi_1(F_2)) \leq \Gamma_3 R_3$. We have now to iterate also this argument: for any $3 \leq j \leq N$ we repeat the above construction starting from F_{j-1} and using the j -th coordinate in the whole procedure, obtaining a set F_j which satisfies

$$\lambda_i(F_j) \leq \lambda_i(\Omega) \quad \forall 1 \leq i \leq k, \quad \text{diam}(\pi_p(F_j)) \leq \Gamma_3^{j-p} R_3 \quad \forall 1 \leq p \leq j.$$

The thesis is then finally obtained by defining $\widehat{\Omega} := F_N$, being $R := \Gamma_3^{N-1} R_3$. \square

3. PROOF OF THEOREM B

This last section is devoted to the proof of Theorem B. For the ease of presentation, we will begin with a couple of technical lemmas, then pass to the proof of the theorem. The first simple step of our construction states that functions with bounded Rayleigh quotients cannot concentrate too much on small regions.

Lemma 3.1. *For every $m \in (0, 1]$ and $K > 0$ there exists $\rho = \rho(m, K, N) > 0$ such that the following holds. Let $u \in W^{1,2}(\mathbb{R}^N)$ with*

$$\int_{\mathbb{R}^N} u^2 = 1, \quad \int_{\mathbb{R}^N} |Du|^2 \leq K.$$

Then for every cube $Q \subseteq \mathbb{R}^N$ with half-side ρ one has

$$\int_Q u^2 \leq m.$$

Proof. Suppose that the claim is not true. Then there exists a sequence $\{u_n\} \subseteq W^{1,2}(\mathbb{R}^N)$ satisfying

$$\int_{\mathbb{R}^N} u_n^2 = 1, \quad \int_{\mathbb{R}^N} |Du_n|^2 \leq K, \quad \int_{Q_{1/n}} u_n^2 \geq m, \quad (3.1)$$

being $Q_r = [-r, r]^N$ the cube of half-side r centered at the origin. By definition, this sequence is bounded in $W^{1,2}(\mathbb{R}^N)$, hence up to a subsequence we have that u_n weakly converges to some function $u \in W^{1,2}(\mathbb{R}^N)$. In particular, for any $\varepsilon > 0$, u_n strongly converges to u in $L^2(Q_\varepsilon)$, so that, thanks to (3.1), one has $\int_{Q_\varepsilon} u^2 \geq m$. Since this is absurd, the claim follows. \square

The second lemma, which is the core of our proof of Theorem B, ensures that every set with bounded first eigenvalue can be split into two subregions, each of them having first eigenvalue not too large.

Lemma 3.2. *For every $K > 0$ there exists $K' = K'(K, N)$ such that, if Ω is an open subset of \mathbb{R}^N with $\lambda_1(\Omega) \leq K$, then there are two disjoint open subsets Ω_1, Ω_2 of Ω with $\lambda_1(\Omega_i) \leq K'$ for $i = 1, 2$.*

Proof. We start applying Lemma 3.1 with K and with $m = 1/2$, thus getting a positive number ρ . Let then $\Omega \subseteq \mathbb{R}^N$ be an open set with $\lambda_1(\Omega) \leq K$, and let $u \in W_0^{1,2}(\Omega)$ be a first eigenfunction of Ω with unit L^2 norm. Extending u by 0 outside Ω , we have then by definition

$$\int_{\mathbb{R}^N} u^2 = \int_{\Omega} u^2 = 1, \quad \int_{\mathbb{R}^N} |Du|^2 = \int_{\Omega} |Du|^2 \leq K. \quad (3.2)$$

Let now $t^- < t^+$ be identified by

$$\int_{\Omega_{t^-}^l} u^2 = \int_{\Omega_{t^+}^r} u^2 = \frac{1}{4N}. \quad (3.3)$$

We claim that it is possible to assume

$$t^+ - t^- \geq 2\rho. \quad (3.4)$$

In fact, if it is not so, this means that there is a vertical stripe of width 2ρ out of which the squared L^2 norm of u is less than $1/(2N)$ (by ‘‘vertical’’ we mean orthogonal to e_1). If this happens for every direction e_1, e_2, \dots, e_N , the intersection of the corresponding stripes is a square of half-side ρ out of which the squared L^2 norm of u is less than $1/2$. Since this is in contradiction with Lemma 3.1, we obtain the validity of (3.4), up to a rotation.

Let us now call $t = (t^+ + t^-)/2$, define $\Omega_1 = \Omega_t^l$ and $\Omega_2 = \Omega_t^r$, and let $\tilde{u} \in W^{1,2}(\Omega_1)$ be defined as

$$\tilde{u}(x, y) := \begin{cases} u(x, y) & \text{for } x \leq t - \rho, \\ \frac{t-x}{\rho} u(x, y) & \text{for } t - \rho \leq x \leq t. \end{cases}$$

Since $u \in W_0^{1,2}(\Omega)$, it is clear that $\tilde{u} \in W_0^{1,2}(\Omega_1)$. Moreover, writing $Du = (D_1u, D_yu)$, one has

$$D\tilde{u}(x, y) = \left(\frac{t-x}{\rho} D_1u(x, y) - \frac{1}{\rho} u(x, y), \frac{t-x}{\rho} D_yu(x, y) \right)$$

for every $(x, y) \in \Omega_1$ with $x \geq t - \rho$. As a consequence, minding (3.2) one gets

$$\int_{\Omega_1} |D\tilde{u}|^2 \leq 2 \int_{\Omega_1} |Du|^2 + \frac{2}{\rho^2} \int_{\Omega_1} u^2 \leq 2K + \frac{2}{\rho^2}. \quad (3.5)$$

On the other hand, recalling (3.4) and (3.3) it is

$$\int_{\Omega_1} \tilde{u}^2 \geq \int_{\Omega_{t^-}^l} u^2 = \frac{1}{4N}. \quad (3.6)$$

Putting together (3.5) and (3.6) one immediately obtains

$$\lambda_1(\Omega_1) \leq \mathcal{R}(\tilde{u}, \Omega_1) = \frac{\int_{\Omega_1} |D\tilde{u}|^2}{\int_{\Omega_1} \tilde{u}^2} \leq 8N \left(K + \frac{1}{\rho^2} \right).$$

Finally, we can set $K' = 8N(K + 1/\rho^2)$: since we have shown that $\lambda_1(\Omega_1) \leq K'$, and since by symmetry it is also $\lambda_1(\Omega_2) \leq K'$, the thesis follows. \square

We are now in position to prove a first boundedness result of λ_k in terms of λ_1 , from which Theorem B will then readily follow.

Lemma 3.3. *For every $K > 0$ there exists $M' = M'(k, K, N) > 0$ such that, for all open sets $\Omega \subseteq \mathbb{R}^N$, if $\lambda_1(\Omega) \leq K$ then $\lambda_k(\Omega) \leq M'$.*

Proof. Let us start by setting $K_1 = K$, and then, applying Lemma 3.2, we let recursively $K_{l+1} = K'(K_l, N)$ for every $l \geq 1$. Finally, we define $M' = K_{j+1}$, where j is the smallest natural number such that $2^j \geq k$. We will show the claim of the theorem with such constant M' .

To do so, we pick any open set Ω with $\lambda_1(\Omega) \leq K = K_1$. Applying Lemma 3.2 to Ω with constant K_1 , we find two disjoint open sets $\Omega_1, \Omega_2 \subseteq \Omega$ with $\lambda_1(\Omega_i) \leq K'(K_1, N) = K_2$ for $i = 1, 2$. Then, we can apply Lemma 3.2 to Ω_1 and Ω_2 with constant K_2 , finding four disjoint subsets $\Omega_{11}, \Omega_{12}, \Omega_{21}, \Omega_{22}$ of Ω , each of them with first eigenvalue smaller than K_3 . Continuing with the obvious induction, we end up with 2^j disjoint open subsets of Ω , say Ω^i for $1 \leq i \leq 2^j$, having $\lambda_1(\Omega^i) \leq K_{j+1} = M'$ for each i .

To conclude the thesis, it is thus enough to show that

$$\lambda_k(\Omega) \leq \lambda_{2^j}(\Omega) \leq \max \left\{ \lambda_1(\Omega^i) : 1 \leq i \leq 2^j \right\} \leq M', \quad (3.7)$$

and in fact only the second inequality is to be shown, being the first and the last true by construction.

To get (3.7), for every $1 \leq i \leq 2^j$ let u_i be a first eigenfunction of Ω^i , again extended by 0 on $\Omega \setminus \Omega^i$, so that

$$\int_{\Omega} u_i^2 = \int_{\Omega^i} u_i^2 = 1, \quad \int_{\Omega} |Du_i|^2 = \int_{\Omega^i} |Du_i|^2 = \lambda_1(\Omega^i) \leq M',$$

and then $\mathcal{R}(u_i, \Omega) \leq M'$. Observe that the functions u_i are mutually orthogonal (both in the L^2 and in the $W^{1,2}$ sense) by construction, since they are supported on disjoint sets. Hence, the linear subspace K_{2^j} of $W_0^{1,2}(\Omega)$ spanned by the functions u_i for $1 \leq i \leq 2^j$ is 2^j -dimensional. Thanks to Theorem 1.2, to prove (3.7) it is enough to show that $\mathcal{R}(w) \leq \max \{ \lambda_1(\Omega^i) : 1 \leq i \leq 2^j \}$ for every $w \in K_{2^j}$. And in fact, writing the generic function $w \in K_{2^j}$ as $w = \sum \beta_i u_i$, by the

orthogonality of the different u_i one has clearly

$$\begin{aligned} \mathcal{R}(w, \Omega) &= \frac{\int_{\Omega} \left| \sum \beta_i D u_i \right|^2}{\int_{\Omega} \left(\sum \beta_i u_i \right)^2} = \frac{\sum \beta_i^2 \int_{\Omega} |D u_i|^2}{\sum \beta_i^2 \int_{\Omega} u_i^2} = \frac{\sum \beta_i^2 \mathcal{R}(u_i, \Omega) \int_{\Omega} u_i^2}{\sum \beta_i^2 \int_{\Omega} u_i^2} \\ &= \frac{\sum \beta_i^2 \lambda_1(\Omega^i) \int_{\Omega} u_i^2}{\sum \beta_i^2 \int_{\Omega} u_i^2} \leq \max \left\{ \lambda_1(\Omega^i) : 1 \leq i \leq 2^j \right\}. \end{aligned}$$

As noticed before, this gives the validity of (3.7), hence the proof is concluded. \square

To obtain Theorem B, we now only need a trivial rescaling argument.

Proof of Theorem B. First of all notice that, by density, it is admissible to consider only the case of the open sets. We apply Lemma 3.3 with $K = 1$, so defining $M := M'(k, 1, N)$. We will prove Theorem B with such M . Let $\Omega \subseteq \mathbb{R}^N$ be an open set, and apply the rescaling formula (1.2) choosing $\alpha = \lambda_1(\Omega)^{\frac{1}{2}}$, thus getting $\lambda_1(\alpha \Omega) = 1$. By Lemma 3.3, we derive $\lambda_k(\alpha \Omega) \leq M$, and then by (1.2) again we find $\lambda_k(\Omega) = \alpha^2 \lambda_k(\alpha \Omega) \leq M \lambda_1(\Omega)$, thus the proof is concluded. \square

Acknowledgments. We warmly thank Michiel van den Berg and Frank Morgan for their suggestions on the paper. This work has been supported by the ERC Starting Grant n. 258685 “AnOptSetCon”, and the ERC Advanced Grant n. 226234 “AnTeGeFI”. The work of this paper was done while both the authors were at the University of Pavia.

REFERENCES

- [1] M.S. Ashbaugh, The universal eigenvalue bounds of Payne-Pölya-Weinberger, Hile-Protter, and H C Yang, Proc. Indian Acad. Sci. (Math. Sci.) Vol. 112, No. 1, February 2002, pp. 330.
- [2] M.S. Ashbaugh & R. Benguria, Proof of the Payne-Pölya-Weinberger conjecture, Bull. Amer. Math. Soc., **25** no. 1 (1991), 19–29.
- [3] M. van den Berg & M. Iversen, On the minimization of Dirichlet eigenvalues of the Laplace operator, J. Geom. Anal., to appear.
- [4] D. Bucur, Minimization of the k -th eigenvalue of the Dirichlet Laplacian, Arch. Ration. Mech. Anal., to appear.
- [5] D. Bucur & G. Buttazzo, Variational methods in shape optimization problems, Progress in nonlinear differential equations and their applications, Birkhäuser Verlag, Boston (2005).
- [6] D. Bucur & A. Henrot, Minimization of the third eigenvalue of the Dirichlet Laplacian, Proc. Roy. Soc. London, **456** (2000), 985–996.
- [7] G. Buttazzo & G. Dal Maso, An existence result for a class of shape optimization problems, Arch. Ration. Mech. Anal. **122** (1993), 183–195.
- [8] A. Henrot, Minimization problems for eigenvalues of the Laplacian, Journal of Evolution Equations **3** (2003), 443–461.
- [9] A. Henrot, Extremum Problems for Eigenvalues of Elliptic Operators, Frontiers in Mathematics, Birkhäuser Verlag, Basel (2006).
- [10] A. Henrot & M. Pierre, Variation et optimisation de formes, Mathématiques et Applications **48**, Springer (2005).

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI PAVIA, VIA FERRATA,1, 27100 PAVIA (ITALY)

DEPARTMENT MATHEMATIK, FRIEDERICH-ALEXANDER UNIVERSITÄT ERLANGEN-NÜRNBERG, CAUERSTRASSE,11,
91058 ERLANGEN (GERMANY)

E-mail address: `dario.mazzoleni@unipv.it`, `mazzoleni@math.fau.de`

DEPARTMENT MATHEMATIK, FRIEDERICH-ALEXANDER UNIVERSITÄT ERLANGEN-NÜRNBERG, CAUERSTRASSE,11,
91058 ERLANGEN (GERMANY)

E-mail address: `pratelli@math.fau.de`