

**STABILITY THEOREMS FOR GAGLIARDO-NIRENBERG-SOBOLEV
INEQUALITIES:
A REDUCTION PRINCIPLE TO THE RADIAL CASE**

B. RUFFINI

ABSTRACT. In this paper we investigate the quantitative stability for Gagliardo-Nirenberg-Sobolev inequalities. The main result is a reduction theorem, which states that, to solve the problem of the stability for Gagliardo-Nirenberg-Sobolev inequalities, one can consider only the class of radial decreasing functions.

1. INTRODUCTION

1.1. **Background.** The sharp Gagliardo-Nirenberg-Sobolev inequality (briefly: GNS) in \mathbb{R}^n , with $n \geq 2$, takes the form, for a suitable constant $G = G(n, p, s, q) > 0$,

$$G\|u\|_q \leq \|\nabla u\|_p^\theta \|u\|_s^{1-\theta} \quad (1.1)$$

where the parameters s, q, p satisfy

$$\begin{aligned} 1 < p < n, \\ 1 \leq s < q < p^*, \quad p^* = \frac{np}{n-p} \quad \text{and} \\ \frac{\theta}{p^*} + \frac{1-\theta}{s} = \frac{1}{q}. \end{aligned} \quad (1.2)$$

In (1.1), and in what follows in this paper, the symbol $\|u\|_r$ denotes the L^r -norm of a function u on \mathbb{R}^n

$$\|u\|_r^r = \int_{\mathbb{R}^n} |u|^r = \int |u|^r$$

while if the set of integration is a proper subset E of \mathbb{R}^n , we use the notation $\|\cdot\|_{L^r(E)}$. Analogously, to lighten the notations, we shall omit the set of integration when an integral is defined on the whole \mathbb{R}^n . The function u is taken in $D^{p,s}(\mathbb{R}^n)$, that is the closure of $C_c^\infty(\mathbb{R}^n)$ under the norm $\|u\|_{D^{p,s}} = \|\nabla u\|_p + \|u\|_s$, anyhow we can always suppose, in this paper, to deal with regular functions, so that the gradient ∇u of a function u is well defined.

Inequality (1.1) belongs to the class of the celebrated *Sobolev inequalities*, which are a fundamental tool in several applications in the Calculus of Variation, and it can be derived by combining the interpolation inequality between the L^s, L^q and L^{p^*} norms on \mathbb{R}^n with the classical Sobolev inequality

$$S(n, p)\|u\|_{p^*} \leq \|\nabla u\|_p. \quad (1.3)$$

Here $S(n, p)$ is the optimal Sobolev constant, namely

$$S(n, p) = \inf \left\{ \frac{\|\nabla u\|_p}{\|u\|_{p^*}} : u \in W^{1,p}(\mathbb{R}^n) \setminus \{0\} \right\}, \quad (1.4)$$

2010 *Mathematics Subject Classification.* 26D10; 46E35; 49J40.

Key words and phrases. Sobolev inequality; Gagliardo-Nirenberg inequalities; Quantitative stability inequalities .

being $W^{1,p}(\mathbb{R}^n)$ the Sobolev space of functions which are, together with their weak gradient, in $L^p(\mathbb{R}^n)$ (see [10]). Explicit formulas for $S(n, p)$ and minimizers in (1.4) are known since the work of Aubin [1] and Talenti [24]. We stress that the same result *is not available* for the optimal constants and functions in (1.1) with the exception of the one-parameter family of exponents

$$p(s-1) = q(p-1) \text{ if } q, s > p \quad p(q-1) = s(p-1) \text{ if } q, s < p, \quad (1.5)$$

see [9, 6]. What can be said in full generality, is that optimal functions in (1.1) exist and, according to the Pólya-Szegő inequality (see for instance [2]), are non-negative, radially symmetric functions with decreasing profile (see [2, 23]). Furthermore, they are unique up to translations, rescaling and multiplication by (non-zero) constants. In this paper we deal with the problem of getting a quantitative stability version of inequality (1.1). To state rigorously this problem, we introduce the (GNS) *deficit* $\delta(u)$ of a function $u \in D^{p,s}(\mathbb{R}^n)$ as

$$\delta(u) = \frac{\|\nabla u\|_p^\theta \|u\|_s^{1-\theta}}{G\|u\|_q} - 1, \quad (1.6)$$

and notice that inequality (1.1) reads, in terms of $\delta(u)$, as

$$\delta(u) \geq 0. \quad (1.7)$$

Then, for a quantitative version of the GNS inequality, we mean an improvement of inequality (1.7) of the form

$$\delta(u) \geq \kappa_0 \text{dist}(u, M)^{\alpha_0}, \quad (1.8)$$

where κ_0, α_0 are positive constants not depending on u and $\text{dist}(\cdot, M)$ indicates an appropriate distance from M , the set of the optimizers for (1.1). The concept of distance we shall adopt is the following

$$\lambda(u) = \inf \left\{ \frac{\|u-v\|_q^q}{\|u\|_q^q} : v \text{ is optimal for (1.1), } \|v\|_q = \|u\|_q \right\} \quad (1.9)$$

and we will call it *asymmetry of u* . Results in this direction have been recently obtained with some ad-hoc techniques valid for special classes of parameters among those in (1.2), see [7] and [4]. In particular, the parameters considered in [7] are contained in those introduced in (1.5) (although the authors focus, because of their later applications, just on the particular case $p=2, q=6, s=4$), and the knowledge of minimizers is exploited in a crucial way. In [4], the authors address a class of parameters, $p=s=2, q>2$, for which the minimizers are not explicitly known and they follow a strategy developed by Bianchi and Egnell in [5], which heavily relies on the Hilbertian structure corresponding to $p=2$ and seems complicate to generalize. Thus the above techniques seem adaptable to prove the stability for the GNS inequalities only for a particular class of parameters. In this paper, thanks to a general symmetrization technique introduced by Cianchi, Fusco, Maggi and Pratelli in [8], we are able to prove a reduction principle which is valid for the *whole class* of parameters (1.2). Namely we reduce the problem to that of showing the stability just for radial symmetric functions, reducing the complexity of the task in its generality from a n -dimensional to a 1-dimensional problem. Although this does not solve completely the problem, it offers a more simple way to attack it.

1.2. Main result and plan of the paper. Consider the class of parameters (1.2). The main result we shall prove is the following

Main Theorem. Consider the functionals $\delta(\cdot)$ and $\lambda(\cdot)$ defined in (1.6) and (1.9) respectively. Suppose that there exist two positive constants k_0 and α_0 such that the stability inequality

$$\delta(u) \geq \kappa_0 \lambda(u)^{\alpha_0} \quad (1.10)$$

holds for any radial non-increasing function $u \in D^{p,s}(\mathbb{R}^n)$. Then there exist two positive constants k_1 and α_1 such that the inequality

$$\delta(u) \geq \kappa_1 \lambda(u)^{\alpha_1} \quad (1.11)$$

holds true for any function in $D^{p,s}(\mathbb{R}^n)$.

A first result we shall use, in the direction of proving the Main Theorem, is a sort of *continuity at 0* of the asymmetry λ with respect to the deficit δ . Namely we prove, in Corollary 2.4 of Section 2, that given a sequence of functions $(u_h)_h$ such that $\delta(u_h)$ converges to 0 as h goes to infinity, then $\lambda(u_h)$ converges to 0 as well. This result is accomplished by means of the compactness Theorem 2.1, where we prove that a sequence of functions whose deficits are infinitesimal, up to be (suitably) rescaled and translated, is compact in $L^q(\mathbb{R}^n)$. After Corollary 2.4 is settled, we pass to the proof of the Main Theorem. Its proof is given in Section 3 and Section 4, each of them devoted to obtain a simplification of the class of functions we deal with. More precisely we make use of a further reduction step, aimed to prove that if the stability inequality (1.10) holds true for radial decreasing functions, it also holds for n -symmetric functions, that is functions which are symmetric with respect to n orthogonal hyperplanes. Namely we prove, in Section 3, that if there exist positive constants κ_0 and α_0 such that for any radial decreasing function $u \in D^{p,s}(\mathbb{R}^n)$ inequality (1.10) holds true, then there exist positive constants κ_1 and α_1 depending on n, p, q and s such that for any n -symmetric function $u \in D^{p,s}(\mathbb{R}^n)$ we have

$$\delta(u) \geq \kappa_1 \lambda(u)^{\alpha_1}. \quad (1.12)$$

Eventually, in Section 4, we prove that to get the stability for GNS inequality, it is not restrictive to consider only n -symmetric functions. To do this, we show the existence of two positive constants κ_2 and α_2 such that for every non-negative function $u \in D^{p,s}(\mathbb{R}^n)$ there exists an n -symmetric function \bar{u} such that the following reduction inequalities hold true

$$\lambda(u) \leq \kappa_2 \lambda(\bar{u})^{\alpha_2}, \quad \delta(\bar{u}) \leq \kappa_2 \delta(u)^{\alpha_2}. \quad (1.13)$$

It is then easy to see that combining (1.12) with the reduction inequalities (1.13) we get the claim of the Main Theorem.

2. CONTINUITY OF λ WITH RESPECT TO δ VIA A COMPACTNESS THEOREM

We devote this section to the proof of the following theorem.

Theorem 2.1. Let $(u_h)_h$ be a sequence in $D^{p,s}(\mathbb{R}^n)$ such that $\delta(u_h)$ converges to 0 as $h \rightarrow \infty$. Then there exist $(\lambda_h)_h \subset (0, +\infty)$ and $(x_h)_h \subset \mathbb{R}^n$ such that the rescaled sequence

$$w_h(x) = \tau_{\lambda_h} u_h(x - x_h) = \lambda_h^{n/q} u_h(\lambda_h(x - x_h)) \quad (2.1)$$

satisfies:

- (i) $\delta(w_h) = \delta(u_h)$; $\lambda(w_h) = \lambda(u_h)$;
- (ii) $\|w_h\|_q = \|u_h\|_q$ for $h \in \mathbb{N}$;
- (iii) there exist constants $C_0, C_1 > 0$, depending only on n, p, q, s , such that

$$\frac{1}{C_0} \leq \|\nabla w_h\|_p \leq C_0, \quad \frac{1}{C_1} \leq \|w_h\|_s \leq C_1;$$

(iv) $w_h \rightarrow w$ strongly in $L^q(\mathbb{R}^n)$ as $h \rightarrow \infty$ with $w \in D^{p,s}(\mathbb{R}^n)$.

In what follows we will always implicitly suppose that the parameters n, p, s, q and θ satisfy conditions (1.2). We begin our analysis considering the following functionals

$$G(u) = \|\nabla u\|_p^\theta \|u\|_s^{1-\theta}, \quad F(u) = \int |\nabla u|^p + \int |u|^s \quad (2.2)$$

defined for $u \in D^{p,s}(\mathbb{R}^n)$. Given $m > 0$ we define the functions

$$\psi(m) = \inf\{G(u) : \|u\|_q^q = m\}, \quad \varphi(m) = \inf\{F(u) : \|u\|_q^q = m\}. \quad (2.3)$$

Lemma 2.2. *There exists $\eta_0 = \eta_0(n, p, q, s) > 0$ with the following property. For any $u \in D^{p,s}(\mathbb{R}^n)$ there exists $\lambda > 0$ such that, if $\tau_\lambda(u) = \lambda^{n/q}u(\lambda x)$, then*

$$F(\tau_\lambda u) = \eta_0 G(u)^k \quad \text{where} \quad k = q \cdot \frac{np + ps - ns}{np + pq - ns} < q. \quad (2.4)$$

Proof. For $u \in D^{p,s}(\mathbb{R}^n)$ we have

$$\|\tau_\lambda u\|_q^q = \|u\|_q^q, \quad \|\tau_\lambda u\|_s^s = \lambda^{-n+ns/q} \|u\|_s^s \quad \text{and} \quad \|\nabla \tau_\lambda u\|_p^p = \lambda^{-n+p+\frac{np}{q}} \|\nabla u\|_p^p; \quad (2.5)$$

hence

$$F(\tau_\lambda u) = \lambda^a A + \lambda^b B = f(\lambda)$$

where $A = \|\nabla u\|_p^p$, $B = \|u\|_s^s$, $a = -n + p + np/q$ and $b = -n + ns/q$. The function f attains its minimum at

$$\lambda_m = \left(-\frac{b}{a}\right)^{\frac{1}{a-b}} \left(\frac{B}{A}\right)^{\frac{1}{a-b}}$$

with the value

$$f(\lambda_m) = \eta_0 \left(A^{\theta/p}\right)^{q\nu} \left(B^{(1-\theta)/s}\right)^{q\nu}$$

where $\eta_0 = \eta_0(n, p, s, q)$ and $\nu = \frac{np+ps-ns}{np+pq-ns}$, that is the claim of the lemma. \square

Lemma 2.3. *There exists $\alpha = \alpha(n, p, q, s) \in (0, 1)$ such that*

$$\varphi(m) = m^\alpha \varphi(1) \quad \text{for all } m > 0.$$

In particular φ is strictly super-additive in $(0, 1)$.

Proof. Let $u \in D^{p,s}(\mathbb{R}^n)$ be such that $\|u\|_q^q = m$. Let $v = u/m^{1/q}$ and set $\tau_\lambda v(x) = \lambda^{n/q}v(\lambda x)$. Setting $\lambda = m^{\frac{p-s}{np+pq-ns}}$ we get, after some calculation analogous to those in Lemma 2.2,

$$F(\tau_\lambda v) = m^{-\alpha} F(u)$$

where $\alpha = \frac{np+ps-ns}{np+pq-ns}$. If we now consider a minimizing sequence $(u_h)_h$ for $\varphi(m)$ such that $\|u_h\|_q^q = m$ and as above $v_h = u_h/m^{1/q}$, we obtain

$$\varphi(m) = \lim_{h \rightarrow \infty} F(u_h) = m^\alpha \lim_{h \rightarrow \infty} F(\tau_\lambda v_h) \geq m^\alpha \varphi(1).$$

The opposite inequality can be proved with an analogous argument considering a minimizing sequence $(v_h)_h$ for $\varphi(1)$, and setting $u_h = m^{1/q}v_h$. \square

We pass now to the proof of Theorem 2.1.

Proof of Theorem 2.1. Let φ be the function defined in (2.3). We recall the Lions's Concentration-Compactness Theorem (see [18] or [22, Theorem 4.3 and Theorem 4.8]): given a non-negative sequence $(\rho_h)_h$ in $L^1(\mathbb{R}^n)$ with fixed L^1 norm, say 1, there is a subsequence $(\rho_{h_k})_k$ which satisfies one of the following properties:

- (1) (concentration) there exists a sequence $(y_k) \subset \mathbb{R}^n$ such that for every $\varepsilon > 0$ there exists $R \in (0, \infty)$ such that $\int_{B_R(y_k)} \rho_{h_k} \geq 1 - \varepsilon$ for every $k \in \mathbb{N}$;
- (2) (vanishing) $\lim_{k \rightarrow \infty} \sup_{y \in \mathbb{R}^n} \int_{B_R(y)} \rho_{h_k} = 0$ for $0 < R < \infty$;
- (3) (dichotomy) there exist $\alpha \in (0, 1)$ and two sequences $R_h \rightarrow +\infty$ and $(y_h)_h \subset \mathbb{R}^n$ such that $\int_{B_{R_{h_k}}(y_{h_k})} \rho_{h_k} \rightarrow \alpha$ and $\int_{\mathbb{R}^n \setminus B_{2R_{h_k}}(y_{h_k})} \rho_{h_k} \rightarrow 1 - \alpha$.

Let $u_h \in D^{p,s}(\mathbb{R}^n)$ be such that $\delta(u_h) \rightarrow 0$. We can suppose that $\|u_h\|_q = 1$. Consider $w_h(x - x_h) = \tau_{\lambda_h} u_h(x - x_h)$ where $\lambda_h > 0$ is defined, for every h , as in Lemma 2.2. Thanks to formulas (2.5), each function w_h satisfies statements (i) and (ii) of the theorem. Moreover, Lemma 2.2 provides us two positive constants η_0 and k such that $F(w_h) = \eta_0 G(u_h)^k$. Since $\delta(w_h) + 1 = G(w_h)/G$, and $\delta(w_h)$ tends to 0, it follows that the sequence $F(w_h) \rightarrow G^k = \varphi(1)$ as h tends to ∞ , where φ is defined in (2.3). In particular the sequence $(w_h)_h$ must satisfy statement (iii) of the theorem. In order to prove point (iv), we apply the Concentration-Compactness Theorem to the sequence $(|w_h|^q)_h$ aiming to exclude cases (2) and (3).

If the sequence vanishes, by Hölder inequality we would get vanishing also for the sequence $(|w_h|^s)_h$, since $s < q$. It is not difficult to see that these conditions, together to the equiboundedness of $(w_h)_h$ in $D^{p,s}(\mathbb{R}^n)$, guarantee that $w_h \rightarrow 0$ strongly in L^q as $h \rightarrow \infty$ (see for instance [19] Lemma I.1). Since $\|w_h\|_q = \|u_h\|_q = 1$, we would get a contradiction. So we can exclude case (2).

The dichotomy case is more complicate and requires a longer analysis. Suppose to have dichotomy for the sequence $(|w_h|^q)_h$. Then there exist $\alpha \in (0, 1)$ and a sequence of positive numbers $R_h \rightarrow \infty$ as $h \rightarrow \infty$ such that

$$\int_{B_{R_h}} |w_h|^q \rightarrow \alpha; \quad \int_{B_{2R_h}^c} |w_h|^q \rightarrow 1 - \alpha; \quad \int_{B_{2R_h} \setminus B_{R_h}} |w_h|^q \rightarrow 0.$$

Let $f \in C_c^1(B(0, 2); [0, 1])$ such that $f = 1$ on $B(0, 1)$ and consider $f_h(x) = f(x/R_h) \in C_c^1(B(0, 2R_h); [0, 1])$. Choose also f such that $|\nabla f_h| \leq C/R_h$ for some $C > 0$. Then we have

$$\begin{aligned} F(w_h) &= \int |\nabla w_h|^p + \int |w_h|^s \\ &= \int_{B_{R_h}} |\nabla(f_h w_h)|^p + \int_{B_{2R_h}^c} |\nabla((1 - f_h)w_h)|^p + \int_{B_{2R_h} \setminus B_{R_h}} |\nabla w_h|^p \\ &\quad + \int_{B_{R_h}} |f_h w_h|^s + \int_{B_{2R_h}^c} |(1 - f_h)w_h|^s + \int_{B_{2R_h} \setminus B_{R_h}} |w_h|^s \\ &= \int |\nabla(f_h w_h)|^p + \int |\nabla[(1 - f_h)w_h]|^p \\ &\quad + \int_{B_{2R_h} \setminus B_{R_h}} \left[|\nabla w_h|^p - |\nabla(f_h w_h)|^p - |\nabla[(1 - f_h)w_h]|^p \right] \\ &\quad + \int \left[|f_h w_h|^s + |(1 - f_h)w_h|^s \right] \\ &\quad + \int_{B_{2R_h} \setminus B_{R_h}} \left[|w_h|^s - f_h^s |w_h|^s - (1 - f_h)^s |w_h|^s \right]. \end{aligned}$$

Since f_h takes values in $[0, 1]$, the last integral is non-negative. Neglecting this quantity and rearranging the terms, we get

$$\begin{aligned} F(w_h) &\geq F(f_h w_h) + F((1 - f_h)w_h) - \epsilon(h) \\ &\geq \varphi(\|f_h w_h\|_q^q) + \varphi(\|(1 - f_h)w_h\|_q^q) - \epsilon(h), \end{aligned} \quad (2.6)$$

where

$$\epsilon(h) = \int_{B_{2R_h} \setminus B_{R_h}} \left[|\nabla(f_h w_h)|^p + |\nabla[(1 - f_h)w_h]|^p - |\nabla w_h|^p \right].$$

We claim that the error $\epsilon(h)$ is estimated from above by a quantity which converges to 0 as $h \rightarrow \infty$. Indeed

$$\begin{aligned} \epsilon(h) &= \int_{B_{2R_h} \setminus B_{R_h}} \left[|\nabla(f_h w_h)|^p + |\nabla[(1 - f_h)w_h]|^p - |\nabla w_h|^p \right] \\ &= \int_{B_{2R_h} \setminus B_{R_h}} \left[|f_h \nabla w_h + w_h \nabla f_h|^p + |(1 - f_h) \nabla w_h - w_h \nabla f_h|^p - |\nabla w_h|^p \right] \\ &\leq \int_{B_{2R_h} \setminus B_{R_h}} \left[(|f_h \nabla w_h + w_h \nabla f_h| + |(1 - f_h) \nabla w_h - w_h \nabla f_h|)^p - |\nabla w_h|^p \right] \\ &\leq \int_{B_{2R_h} \setminus B_{R_h}} \left[(|\nabla w_h| + 2|w_h| |\nabla f_h|)^p - |\nabla w_h|^p \right] \\ &\leq \int_{B_{2R_h} \setminus B_{R_h}} \left[\varepsilon C_p |\nabla w_h|^p + C_\varepsilon |\nabla f_h|^p |w_h|^p \right] \\ &\leq \varepsilon C_p \sup_{h \in \mathbb{N}} \int |\nabla w_h|^p + C_\varepsilon \int_{B_{2R_h} \setminus B_{R_h}} |\nabla f_h|^p |w_h|^p. \end{aligned} \quad (2.7)$$

where the first inequality is due to the super additivity of the map $t \mapsto t^p$ on \mathbb{R}_+ , the second to the triangle inequality and the third one is the Young inequality of (suitable) parameters $\varepsilon > 0$ and C_ε . We need to estimate the quantity

$$g(h) = \int_{B_{2R_h} \setminus B_{R_h}} |\nabla f_h|^p |w_h|^p.$$

If $p > s$ then interpolating the L^p norm of the w_h 's between the L^s norm and the L^{p^*} norm and recalling that $|\nabla f_h| \leq C/R_h$, we get

$$\int_{B_{2R_h} \setminus B_{R_h}} |\nabla f_h|^p |w_h|^p \leq \frac{C^p}{R_h^p} \|w_h\|_{p^*}^{p\theta} \|w_h\|_s^{p(1-\theta)}.$$

Since we already know that w_h satisfies statement (iii) of the theorem, we get that $g(h) \rightarrow 0$ as $h \rightarrow \infty$. If $s \geq p$ we divide $g(h)$ into two terms:

$$g(h) = \int_{(B_{2R_h} \setminus B_{R_h}) \cap \{w_h \geq 1\}} |\nabla f_h|^p |w_h|^p + \int_{(B_{2R_h} \setminus B_{R_h}) \cap \{w_h < 1\}} |\nabla f_h|^p |w_h|^p = g_1(h) + g_2(h).$$

Since $q > p$ we have

$$g_1(h) \leq \frac{C^p}{R_h^p} \int_{B_{2R_h} \setminus B_{R_h}} |w_h|^q \leq \frac{C^p}{R_h^p}$$

and so $g_1(h) \rightarrow 0$ as $h \rightarrow \infty$. Moreover, by Hölder inequality of parameter p^*/p , we get

$$g_2(h) \leq \frac{C^p}{R_h^p} \left(\int_{(B_{2R_h} \setminus B_{R_h}) \cap \{w_h < 1\}} |w_h|^{p^*} \right)^{p/p^*} |(B_{2R_h} \setminus B_{R_h})|^{\frac{1}{(p/p^*)}}$$

where

$$(p/p^*)' = \frac{p/p^*}{-1 + p/p^*} = n/p.$$

Since $p^* > q$, we obtain

$$g_2(h) \leq (\omega_n(2^n - 1))^{p/n} C^p \left(\int_{(B_{2R_h} \setminus B_{R_h}) \cap \{w_h < 1\}} |w_h|^q \right)^{p/p^*}$$

where ω_n is the measure of the unit ball of \mathbb{R}^n . So also g_2 converges to 0 as $h \rightarrow \infty$. Thus, passing to the limit in (2.7), first in $h \rightarrow \infty$ and then in $\varepsilon \rightarrow 0$ we obtain that $\epsilon(h)$ can be estimated from above by an infinitesimal quantity. Since $\|w_h f_h\|_q^q$ and $\|(1 - f_h)w_h\|_q^q$ converge respectively to λ and $1 - \lambda$, we can conclude thanks to Lemma 2.3 that

$$\varphi(1) \geq \varphi(\lambda) + \varphi(1 - \lambda) > \varphi(1)$$

obtaining a contradiction.

So we can exclude also the dichotomy phenomenon. Since w_h is equibounded in $L^q(\mathbb{R}^n)$, we can consider its weak- L^q limit w . This is also a strong limit in L^q . Indeed by concentration, up to translations and since $q > 1$, we have

$$1 - \varepsilon \leq \lim_{h \rightarrow \infty} \int_{B_R} |w_h|^q = \int_{B_R} |w|^q \leq \int |w|^q \leq \liminf_{h \rightarrow \infty} \int |w_h|^q = 1.$$

This concludes the proof of the theorem. \square

Corollary 2.4. *Consider a sequence $(u_h)_h \subset D^{p,s}(\mathbb{R}^n)$ such that $\delta(u_h) \rightarrow \infty$ as $h \rightarrow 0$. Then also $\lambda(u_h) \rightarrow 0$ for $h \rightarrow \infty$.*

Proof. As a consequence of Theorem 2.1, up to pass to a subsequence and to consider a rescaling of the form $\tau_\lambda u_h(x) = \lambda^{n/q} u(\lambda x)$, we can suppose that $u_h \rightarrow u$ strongly in $L^q(\mathbb{R}^n)$. Since the map $u \mapsto \lambda(u)$ is strongly continuous in $L^q(\mathbb{R}^n)$, $\lambda(u_h)$ converges to $\lambda(u)$. Furthermore, by the semicontinuity of the deficit (with respect to the L^q -convergence), $\delta(u) = 0$. Hence u is optimal for (1.1) and $\lambda(u) = 0$. \square

3. REDUCTION TO n -SYMMETRIC FUNCTIONS

In this section we prove that inequality (1.12) holds true for n -symmetric functions (recall that a function is k -symmetric in \mathbb{R}^n , $n \geq k$, if it is symmetric with respect to k mutually orthogonal hyperplanes). We begin with a brief overview of the strategy we want to adopt. Given a function $u \in D^{p,s}(\mathbb{R}^n)$, the natural radial symmetric function to look at is its spherical rearrangement u^* (see [17, Chapter 3]). Suppose that inequality (1.10) holds true for radial symmetric decreasing functions (and so for u^*). Then by the triangle inequality we get

$$\lambda(u)^{1/q} \leq \|u - u^*\|_q + \lambda(u^*)^{1/q} \leq \|u - u^*\|_q + \kappa_0^{1/q} \delta(u^*)^{\alpha_0/q}.$$

In the direction of the proof of (1.12), we notice that by the Pólya-Szegő inequality (see for instance [2]) we have that $\delta(u^*) \leq \delta(u)$. But it is not clear if we can estimate the L^q distance between u and u^* in terms of $\delta(u)$. Indeed this turns out to be true only if a function is already n -symmetric. We shall show, in Lemma 3.1, that $\delta(u)$ estimates from above the Pólya-Szegő deficit, defined as

$$\delta_{PS}(u) = \frac{\|\nabla u\|_p - \|\nabla u^*\|_p}{\|\nabla u^*\|_p}, \quad (3.1)$$

and then, in Lemma 3.2, we prove an estimate of the L^{p^*} distance between u and u^* in terms of the L^p distance between $|\nabla u|$ and $|\nabla u^*|$.

Lemma 3.1. *There exist two positive constants δ_0 and C_0 such that for every $u \in D^{p,s}(\mathbb{R}^n)$ such that $\|u\|_q = 1$, with $\delta(u) \leq \delta_0$, up to the rescaling (2.1), we have*

$$\delta_{PS}(u) \leq C\delta(u)^{1/\theta} \quad (3.2)$$

where $\theta \in (0, 1)$ is the parameter introduced in (1.2).

Proof. By Theorem 2.1, up to rescaling it, we can suppose that u satisfies properties (i) – (iii) in (2.1). If we choose $\delta(u) \leq 1/G$, we obtain:

$$G \leq \|\nabla u^*\|_p^\theta \|u^*\|_s^{1-\theta} \leq \|\nabla u\|_p^\theta \|u\|_s^{1-\theta} \leq 1 + G. \quad (3.3)$$

Then,

$$\begin{aligned} G\delta(u) &= \left(\|\nabla u\|_p^\theta - \|\nabla u^*\|_p^\theta \right) \|u\|_s^{1-\theta} + G\delta(u^*) \\ &\geq C_1^{\theta-1} \left(\|\nabla u\|_p^\theta - \|\nabla u^*\|_p^\theta \right) + G\delta(u^*) \end{aligned} \quad (3.4)$$

where we used the fact that $\|u\|_q = 1$ (statement (ii)) and $\|u\|_s \geq C_1^{-1}$ (statement (iii)). By (3.3) there exists a positive constant c such that,

$$\|\nabla u\|_p^\theta - \|\nabla u^*\|_p^\theta \geq c \left(\|\nabla u\|_p - \|\nabla u^*\|_p \right)^\theta.$$

Now the conclusion follows from (3.4) and (3.1), with $C = c^{1/\theta}$. \square

To obtain the desired estimate of the L^p distance between u and u^* we shall use the following result, whose proof can be found in [8, Theorem 3].

Lemma 3.2. *Let $n \geq 2$, $1 < p < n$ and $z = \max\{p, 2\}$. Then there exists a positive constant C such that*

$$\int |u - u^*|^{p^*} \leq C \left(\int |u|^{p^*} \right)^{\frac{p}{n}} \left(\int |\nabla u^*|^p \right)^{\frac{z-1}{z}} \left(\int |\nabla u|^p - \int |\nabla u^*|^p \right)^{\frac{1}{z}} \quad (3.5)$$

holds for every non-negative $u \in W^{1,p}(\mathbb{R}^n)$ which is symmetric with respect to the coordinate hyperplanes.

We are now able to proceed with the proof of (1.12) when u belongs to the class of n -symmetric functions.

Theorem 3.3. *Suppose that there exist positive constants κ_0 and α_0 such that for any radial decreasing function $u \in D^{p,s}(\mathbb{R}^n)$ inequality (1.10) holds true. Then, there exist positive constants κ_1 and α_1 depending on n, p, q and s such that for any n -symmetric function $u \in D^{p,s}(\mathbb{R}^n)$ we have*

$$\delta(u) \geq \kappa_1 \lambda(u)^{\alpha_1}. \quad (3.6)$$

Proof. Since $\lambda(u) \leq 2^{q-1}$, if $\delta > 0$ and $\delta(u) \geq \min\{\delta, 1/G\}$, then we have $\lambda(u) \leq (2^{q-1}/\delta)\delta(u)$ and so (1.12) holds true with $\kappa_1 = 2^{q-1}/\delta$ and $\alpha_1 = 1$. Hence we may assume that $\delta(u) \leq \min\{\delta, 1/G\}$ for a suitably small δ . Moreover, by Theorem 2.1 we can suppose $\|u\|_q = 1$ and $\|u\|_s \in [1/C_0, C_0]$ where $C_0 \geq 1$ is a constant not depending on u . Keeping in mind these remarks, we divide the proof into two steps:

Step 1: We assume that $u \geq 0$. In this case we have, by an interpolation inequality, Lemma 3.2, and Theorem 2.1,

$$\begin{aligned}
\int |u - u^*|^q &\leq \left(\int |u - u^*|^s \right)^{(1-\theta)q/s} \left(\int |u - u^*|^{p^*} \right)^{\theta q/p^*} \\
&\leq C^{\frac{\theta q}{p^*}} 2^{\frac{(1-\theta)q(s-1)}{s}} \left(\int u^s \right) \left(\int |u|^{p^*} \right)^{\frac{p\theta q}{np^*}} \left(\int |\nabla u^*|^p \right)^{\frac{\theta q(z-1)}{p^*z}} \left(\int |\nabla u|^p - \int |\nabla u^*|^p \right)^{\frac{\theta q}{p^*z}} \\
&\leq C(n, p, q, s) \left(\int |\nabla u|^p - \int |\nabla u^*|^p \right)^\gamma.
\end{aligned} \tag{3.7}$$

where $C(n, p, q, s)$ and γ are suitable positive constants depending on n, p, s and q . Notice that we used the boundedness of the L^p norm of ∇u and of the L^s norm of u guaranteed by Theorem 2.1 (up to choose δ small enough). Moreover we exploited Lemma 3.2 and thus the assumption that $u \geq 0$. If we suppose $\delta(u) \leq \delta \leq G$, again by Theorem 2.1 we get

$$G \leq C_2 \|\nabla u^*\|_p^\theta \leq C_2 \|\nabla u\|_p^\theta = C_3 G(\delta(u) + 1) \leq C_3 G(1 + G),$$

for suitable positive constants C_2 and C_3 . Hence there exists $C_4 > 0$ such that

$$\|\nabla u\|_p^p - \|\nabla u^*\|_p^p \leq C_4 (\|\nabla u\|_p - \|\nabla u^*\|_p).$$

By the triangle inequality, estimates (3.2) and (3.7), the Pólya-Szegő inequality and the assumption on radial functions that we have as hypothesis, we can find constants C_5, C_6 and C_7 not depending on u such that

$$\begin{aligned}
\lambda(u)^q &\leq 2^{q-1} (\lambda(u^*) + \|u - u^*\|_q^q) \\
&\leq C_5 \left[\delta(u)^{\alpha_0} + \left(\int |\nabla u|^p - \int |\nabla u^*|^p \right)^\gamma \right] \\
&\leq C_6 \left[\delta(u)^{\alpha_0} + \delta(u)^{\gamma/\theta} \right] \leq C_7 \delta(u)^\xi,
\end{aligned}$$

where $\xi = \min\{\alpha_0, \gamma/\theta\}$.

Step 2: u changes sign. In this case consider the positive and the negative part of u : $u^+ = u\chi_{\{u>0\}}$ and $u^- = -u\chi_{\{u<0\}}$ (where χ_A denotes the characteristic function of the set A). By Lemma 2.2 we are provided of a positive constant λ such that

$$\eta_0 G(u)^\kappa = F(\tau_\lambda u).$$

Moreover we have that

$$F(\tau_\lambda u^\pm) \geq \inf_\mu F(\tau_\mu u^\pm) = \eta_0 G(u^\pm)^\kappa.$$

where F and G are defined in (2.2). So we get

$$\begin{aligned}
G^\kappa(\delta(u) + 1)^\kappa &= (\|\nabla u\|_p^\theta \|u\|_s^{1-\theta})^\kappa = \frac{1}{\eta_0} \left(\int |\nabla \tau_\lambda u|^p + \int |\tau_\lambda u|^s \right) \\
&= \frac{1}{\eta_0} \left(\int |\nabla \tau_\lambda u^+ + \nabla \tau_\lambda u^-|^p + |\tau_\lambda u^+ + \tau_\lambda u^-|^s \right) \\
&= \frac{1}{\eta_0} \left(\int |\nabla \tau_\lambda u^+|^p + \int |\nabla \tau_\lambda u^-|^p + \int |\tau_\lambda u^+|^s + \int |\tau_\lambda u^-|^s \right) \\
&\geq (\|\nabla u^+\|_p^\theta \|u^+\|_s^{1-\theta})^\kappa + (\|\nabla u^-\|_p^\theta \|u^-\|_s^{1-\theta})^\kappa \\
&\geq G^\kappa \left[\|u^+\|_q^\kappa + \|u^-\|_q^\kappa \right] = G^\kappa \left[\|u^+\|_q^\kappa + \|u^-\|_q^\kappa \right].
\end{aligned}$$

The last equality is due to the fact that $\tau_\lambda u^+$ and $\tau_\lambda u^-$ have disjoint supports while in the last inequality we exploited the GNS inequality. Let us set $\int (u^+)^q = t$ and $\int (u^-)^q = 1 - t$. We can suppose $t \in (0, 1)$, since u changes sign. Then the previous formula is equivalent to

$$f(t) = (t^{\kappa/q} + (1-t)^{\kappa/q})^{1/\kappa} - 1 \leq \delta(u).$$

The function f is symmetric in $[0, 1]$, vanishes at 0 and 1 and since $\kappa < q$ (by Lemma 2.2) is concave. Thus there exists $a > 0$ such that

$$f(t) \geq \frac{1}{a} \min\{t, 1-t\},$$

so that

$$\min \left\{ \int (u^+)^q, \int (u^-)^q \right\} \leq a\delta(u). \quad (3.8)$$

Suppose that the minimum in (3.8) is achieved by $\int (u^-)^q$ (being analogous the other case). Since $\delta(|u|) = \delta(u)$, we can conclude, thanks to the triangle inequality and to (3.8), that

$$\lambda(u)^{1/q} \leq \lambda(|u|)^{1/q} + \left(\int |u - |u||^q \right)^{1/q} \leq C_8(\delta(u)^{\xi/q} + \delta(u)^{1/q}) \leq \kappa_1 \delta(u)^{\alpha/q}$$

where $\alpha = \min\{\xi, 1\}$ and C_8 a positive constant not depending on u . The last inequality holds for $\delta(u) < 1$. So (3.6) holds with $\alpha_1 = \alpha/q$. \square

4. REDUCTION INEQUALITIES

The goal of this section is to prove the reduction inequalities (1.13). Namely we shall prove the following result.

Theorem 4.1. *Assume the hypotheses of Theorem 3.3. Then there exist two positive constants κ_2 and α_2 such that for every non-negative function $u \in D^{p,s}(\mathbb{R}^n)$ there exists an n -symmetric function \bar{u} such that the following reduction inequalities hold true*

$$\lambda(u) \leq \kappa_2 \lambda(\bar{u})^{\alpha_2}, \quad \delta(\bar{u}) \leq \kappa_2 \delta(u)^{\alpha_2}. \quad (4.1)$$

The proof is accomplished arguing similarly to the Sobolev case considered in [8], although some technical modifications are needed. We begin recalling that if v is an optimal function for (1.1), then any other optimal function is of the form

$$v_{a,b,x_0}(x) = av(b(x - x_0)),$$

where a and b are non-null constants and $x_0 \in \mathbb{R}^n$. We define the *relative asymmetry* of a function on an affine subspace S of \mathbb{R}^n as

$$\lambda(u|S) = \inf_{(a,b,x_0) \in \mathbb{R}^2 \times S} \left\{ \frac{\|u - v_{a,b,x_0}\|_q^q}{\|u\|_q^q} : v \text{ optimal for (1.1), } \|v_{a,b,x_0}\|_q = \|u\|_q \right\}.$$

The next Lemma shows that the infimum in the definition of the relative asymmetry is achieved.

Lemma 4.2. *Let $u \in D^{p,s}(\mathbb{R}^n)$ and S an affine space contained in \mathbb{R}^n . Then the infimum in the definition of $\lambda(u)$ and $\lambda(u|S)$ are achieved.*

Proof. Since the two cases are analogous, we show a proof just for the asymmetry. We can suppose without loss of generality that $\|u\|_q = 1$. Let us start observing that $\lambda(u) < 2$. Indeed, if v is a competitor in the definition of $\lambda(u)$, then, up to translate the center of symmetry of v , we can suppose that u and v do not have disjoint supports. Then

$$\lambda(u) \leq \int |v - u|^q = \int_{\{u>v\}} (u - v)^q + \int_{\{v>u\}} (v - u)^q < \int u^q + \int v^q = 2. \quad (4.2)$$

Let now $v_h(x) = a_h v(b_h(x - x_h))$ be a sequence of functions such that $\|v_h\|_q = 1$ and $\|u - v_h\|_q^q \rightarrow \lambda(u)$ as $h \rightarrow \infty$. We want to show that, up to subsequences, $(a_h, b_h, x_h) \rightarrow (a, b, x_0) \subset \mathbb{R}^2 \times \mathbb{R}^n$ as $h \rightarrow \infty$. We have that

$$1 = \int |v_h|^q = a_h^q \int |v(b_h(x - x_h))|^q = \frac{a_h^q}{b_h^n} \int |v|^q = \frac{a_h^q}{b_h^n} \quad (4.3)$$

that is $a_h^q = b_h^n$. Since $v \in L^q(\mathbb{R}^n)$ there exists a function $\rho(\varepsilon)$ converging to 0 as $\varepsilon \rightarrow 0$ such that for each $z \in \mathbb{R}^n$ we have

$$\int_{B(z,\varepsilon)} |v|^q \leq \rho(\varepsilon), \quad \int_{B(0,1/\varepsilon)} |v|^q \geq 1 - \rho(\varepsilon). \quad (4.4)$$

Set now $b_- = \liminf_{h \rightarrow \infty} b_h$ and $b_+ = \limsup_{h \rightarrow \infty} b_h$. We claim that $b_- > 0$ and that $b_+ < \infty$. Suppose $b_- = 0$; then, recalling that v_h is a radial function, we have

$$\begin{aligned} \int_{B(0,1/\varepsilon)} |v_h|^q &\leq \int_{B(x_h,1/\varepsilon)} |v_h|^q = \frac{a_h^q}{b_h^n} \int_{B(0,b_h/\varepsilon)} |v(y)|^q dy \\ &= \int_{B(0,b_h/\varepsilon)} |v(y)|^q dy, \end{aligned} \quad (4.5)$$

and the last quantity, up to pass to a subsequence, converges to 0 as $h \rightarrow \infty$. Thus we can suppose that, for a fixed ε and for h big enough, we have

$$\int_{B(0,1/\varepsilon)} |v_h|^q \leq \varepsilon. \quad (4.6)$$

So, thanks to (4.6) and (4.4) we have

$$\begin{aligned} \|u - v_h\|_q^q &= \int_{B(0,1/\varepsilon)} |u - v_h|^q + \int_{B(0,1/\varepsilon)^c} |u - v_h|^q \\ &\geq \left| \left(\int_{B(0,1/\varepsilon)} |u|^q \right)^{1/q} - \left(\int_{B(0,1/\varepsilon)} |v_h|^q \right)^{1/q} \right|^q \\ &\quad + \left| \left(\int_{B(0,1/\varepsilon)^c} |v_h|^q \right)^{1/q} - \left(\int_{B(0,1/\varepsilon)^c} |u|^q \right)^{1/q} \right|^q \\ &\geq [(1 - \rho(\varepsilon))^{1/q} - \varepsilon^{1/q}]^q + [(1 - \varepsilon)^{1/q} - \rho(\varepsilon)^{1/q}]^q. \end{aligned}$$

Passing to the limit in $h \rightarrow \infty$ and then in $\varepsilon \rightarrow 0$ we obtain that $\lambda(u) \geq 2$, that is a contradiction. Suppose now that $b_+ = \infty$. Then

$$\begin{aligned} \int_{B(x_h, \varepsilon)^c} |v_h|^q &= \int_{B(x_h, \varepsilon)^c} a_h^q |v(b_h(x - x_h))|^q dx = a_h^q \int_{B(0, \varepsilon)^c} |v(b_h x)|^q dx \\ &= \frac{a_h^q}{b_h^n} \int_{B(0, \varepsilon)^c} |v(z)|^q dz = \int_{B(0, b_h \varepsilon)^c} |v|^q \end{aligned} \quad (4.7)$$

and arguing as before we can suppose that $\int_{B(x_h, \varepsilon)^c} |v_h|^q \leq \varepsilon$ for h big enough. By (4.4) we get

$$\begin{aligned} \|u - v_h\|_q^q &= \int_{B(x_h, \varepsilon)} |u - v_h|^q + \int_{B(x_h, \varepsilon)^c} |u - v_h|^q \\ &\geq \left| \left(\int_{B(x_h, \varepsilon)} |v_h|^q \right)^{1/q} - \left(\int_{B(x_h, \varepsilon)} |u|^q \right)^{1/q} \right|^q \\ &\quad + \left| \left(\int_{B(x_h, \varepsilon)^c} |u|^q \right)^{1/q} - \left(\int_{B(x_h, \varepsilon)^c} |v_h|^q \right)^{1/q} \right|^q \\ &\geq (1 - \varepsilon) - \rho(\varepsilon) + (1 - \rho(\varepsilon)) - \varepsilon \end{aligned}$$

and we get, as above, a contradiction. Suppose now that $(x_h)_h$ is not bounded and extract a subsequence (not relabelled) such that $|x_h| \rightarrow \infty$. Then given $N > 0$, if h is big enough we would get $\int_{B(x_h, N)} |u|^q \leq 1/N$. If we choose N such that $\int_{B(x_h, N)} |v_h| \geq \varepsilon$ for all $h \in \mathbb{N}$, we obtain

$$\begin{aligned} \|u - v_h\|_q^q &\geq \int_{B(x_h, N)} |u - v_h|^q + \int_{B(x_h, N)^c} |u - v_h|^q \\ &\geq \left| \left(\int_{B(x_h, N)} |v_h|^q \right)^{1/q} - \left(\int_{B(x_h, N)} |u|^q \right)^{1/q} \right|^q \\ &\quad + \left| \left(\int_{B(x_h, N)^c} |u|^q \right)^{1/q} - \left(\int_{B(x_h, N)^c} |v_h|^q \right)^{1/q} \right|^q \\ &\geq \left[(1 - \varepsilon)^{1/q} - \frac{1}{N^{1/q}} \right]^q + \left[\left(1 - \frac{1}{N} \right)^{1/q} - \varepsilon^{1/q} \right]^q. \end{aligned}$$

and again we get a contradiction. \square

Clearly the asymmetry of a function estimates from below its relative asymmetry. But if we consider a n -symmetric function, also the opposite estimate is true, as shown in next lemma.

Lemma 4.3. *Let $u \in D^{p,s}(\mathbb{R}^n)$ k -symmetric with respect to k orthogonal hyperplanes and let S be the intersection of such hyperplanes. Then*

$$\lambda(u|S) \leq 3^q \lambda(u).$$

Proof. Suppose as usual that $\|u\|_q = 1$. Let $v_{a,b,x}$ a minimum for $\lambda(u)$. We consider now the orthogonal projection x_S of x on S and y the symmetric point of x_S with respect to S . Notice that since u is symmetric with respect to S , also $v_{a,b,y}$ is a minimum for $\lambda(u)$. Moreover, since the minima of the asymmetry are radial symmetric functions with decreasing profile, we have

$$\|v_{a,b,x} - v_{a,b,x_S}\|_q \leq \|v_{a,b,x} - v_{a,b,y}\|_q.$$

This observation and the triangle inequality imply that

$$\begin{aligned}\lambda(u)^{1/q} &\leq \|u - v_{a,b,x_S}\|_q \leq \|u - v_{a,b,x}\|_q + \|v_{a,b,x} - v_{a,b,x_S}\|_q \\ &= \lambda(u)^{1/q} + \|v_{a,b,x} - v_{a,b,x_S}\|_q \leq \lambda(u)^{1/q} + \|v_{a,b,x} - v_{a,b,y}\|_q \\ &\leq \lambda(u)^{1/q} + \|v_{a,b,x} - u\|_q^q + \|u - v_{a,b,y}\|_q = 3\lambda(u)^{1/q}\end{aligned}$$

and the conclusion follows. \square

The next result shows that the n -symmetry condition in the previous lemma can in some sense be relaxed.

Lemma 4.4. *There exists a constant C_0 with the following property. Consider a function $u \in L^q(\mathbb{R}^n)$, $u \geq 0$, H an hyperplane of \mathbb{R}^n and H^+ and H^- the half spaces having H as boundary. Suppose that*

$$\int_{H^+} |u|^q = \int_{H^-} |u|^q = \frac{1}{2} \int_H |u|^q,$$

then

$$\lambda(u|H) \leq C_0 \lambda(u)^{1/q}, \quad (4.8)$$

with a constant C_0 depending only on q and n . Moreover, if T_H denotes the reflection with respect to H of \mathbb{R}^n , it holds

$$\int |u \circ T_H - u|^q \leq C_0 \|u\|_q^q \lambda(u)^{1/q}. \quad (4.9)$$

Proof. Suppose without loss of generality that $\|u\|_q = 1$ and let $v_0 = v_{a,b,x_0}$ a minimum for $\lambda(u)$ centred at x_0 . Suppose moreover that $x_0 \in H^+$, being the other case analogous, and let \bar{x} the projection of x_0 on H and $\bar{v} = v_{a,b,\bar{x}}$. Then

$$\lambda(u|H) \leq \int |u - \bar{u}|^q \leq 2^{q-1} \left(\lambda(u) + \int |v_0 - \bar{v}|^q \right). \quad (4.10)$$

Consider the translated half spaces $K^\pm = H^\pm + (x_0 - \bar{x})$. Since $x_0 \in H^+$ it follows that $K^+ \subseteq H^+$ and $H^- \subseteq K^-$. We have that

$$\frac{1}{2} = \int_{K^\pm} v_0^q = \int_{H^\pm} u^q = \int_{H^\pm} \bar{v}^q$$

and

$$\int_{H^-} |v_0 - \bar{v}|^q = \int_{K^+} |v_0 - \bar{v}|^q \leq \int_{H^+} |v_0 - \bar{v}|^q;$$

hence

$$\int |v_0 - \bar{v}|^q \leq 2 \int_{H^+} |v_0 - \bar{v}|^q. \quad (4.11)$$

Since $v_0 \geq \bar{v}$ on K^+ we get that $|v_0 - \bar{v}|^q \leq v_0^q - \bar{v}^q$ on K^+ . Then

$$\begin{aligned}\int_{K^+} |v_0 - \bar{v}|^q &\leq \int_{K^+} v_0^q - \int_{K^+} \bar{v}^q = \frac{1}{2} - \int_{H^-} v_0^q \\ &\leq C \left(\|u\|_{L^q(H^-)} - \|v_0\|_{L^q(H^-)} \right) \leq C \|u - v_0\|_q = C \lambda(u)^{1/q},\end{aligned} \quad (4.12)$$

for a suitable positive constant C . Moreover

$$\begin{aligned} \int_{H^+ \setminus K^+} |v_0 - \bar{v}|^q &\leq 2^{q-1} \int_{H^+ \setminus K^+} (v_0^q + \bar{v}^q) = 2^q \int_{H^+ \setminus K^+} v_0^q \\ &= 2^q \left[\int_{H^+} v_0^q - \frac{1}{2} \right] = 2^q \left[\int_{H^+} v_0^q - \int_{H^+} u^q \right] \end{aligned}$$

and reasoning as in (4.12) we obtain

$$\int_{H^+ \setminus K^+} |v - \bar{v}|^q \leq C\lambda(u)^{1/q}. \quad (4.13)$$

Inequality (4.8) is then a consequence of (4.10), (4.11), (4.12), and (4.13). We are left to show inequality (4.9). Let \hat{u} be the optimal function in the definition of $\lambda(u|H)$. Then

$$\begin{aligned} \int_{H^\pm} |u \circ T_H - u|^q &\leq 2^{q-1} \left(\int_{H^\pm} |u \circ T_H - \hat{u}|^q + \int_{H^\pm} |u - \hat{u}|^q \right) \\ &= 2^{q-1} \int |u - \hat{u}|^q = 2^{q-1} \lambda(u|H) \leq C_0 \lambda(u)^{1/q}. \end{aligned}$$

□

Before passing to the proof of Theorem 4.1 we need another technical lemma which, roughly speaking, states that if two optimal functions for the GNS inequality are near in L^q norm, then their L^q distance on the whole \mathbb{R}^n can be estimated from above by their L^q distance on just a quarter of \mathbb{R}^n . Its proof is quite technical but it is essentially based on a Taylor expansion.

Lemma 4.5. *Let u be an optimal function for the GNS inequality of parameters s, q, p centred in 0 with $\|u\|_q = 1$, and set $u_{\alpha,z}(x) = \alpha^{n/q} u(\alpha(x-z))$ (for simplicity, $u_{1,z} = u_z$). Consider two orthogonal half spaces H and K containing the origin on their boundaries. There exist two constants $K = K(n, s, q, p) > 0$ and $\tilde{\rho} \ll 1$ such that if*

$$\int |u_{\lambda,x_0} - u_{\mu,y_0}|^q \leq \tilde{\rho},$$

then

$$\int_{H \cap K} |u_{\lambda,x_0} - u_{\mu,y_0}|^q \geq K \int |u_{\lambda,x_0} - u_{\mu,y_0}|^q. \quad (4.14)$$

Proof. Up to a rotation we can consider $H = \{e_1 = 0\}$ and $K = \{e_2 = 0\}$. So that $Q := H \cap K = \{e_1 \geq 0, e_2 \geq 0\}$. Let us define

$$T_k = \{x : 1/k < |\nabla u(x)| < M\} \quad M = \max_{\mathbb{R}^n} |\nabla u(x)|.$$

Since u is a radial function we have that each T_k is a radial set composed of a countable union of centred annuli, i.e. there exists a non-decreasing sequence of positive numbers $(r_{k,j})_j$ such that

$$T_k = \bigcup_{j \in \mathbb{N}} (B_{r_{k,j+1}} \setminus B_{r_{k,j}})$$

where B_r is the ball centred at the origin of radius r . Let now $I_k = \{j \in \mathbb{N} : r_{k,j+1} - r_{k,j} \geq 1/k\}$ and set

$$S_k = \bigcup_{j \in I_k} (B_{r_{k,j+1}} \setminus B_{r_{k,j}}).$$

We consider, for $z \in \mathbb{R}^n$, the biggest centred annulus contained in $S_k \cap (S_k + z)$:

$$\Sigma(k, z) = \{x \in S_k \cap (S_k + z) : \partial B_{|x|} \subseteq S_k \cap (S_k + z)\}.$$

Notice, that $\Sigma(k, z_1) = \Sigma(k, z_2)$ whenever $|z_1| = |z_2|$ so we may define $\Sigma(k, r) = \Sigma(k, z)$ if $|z| = r$. Clearly $\bigcup_{k \in \mathbb{N}} S_k = \mathbb{R}^n$. Moreover, since $|\nabla u|$ is continuous, we have that

$$\bigcup_{|z|>0} \Sigma(k, z) = S_k. \quad (4.15)$$

Indeed, if $x \in S_k$, there exists $r = r(x) > 0$ such that $|\nabla u|(y) > 1/k$ for every $y \in B(x, r(x))$, that is $\overline{B(x, r(x))} \subseteq S_k$. Thus it is sufficient to choose $|z| < \text{dist}(x, \partial S_k)$ and we get that $\partial B_{|x|} \subseteq S_k \cap (S_k + z)$, i.e $x \in \Sigma(k, z)$. Bearing these definitions in mind, we pass to prove inequality (4.14). Up to a change of variables we can consider just the case $\lambda = 1 + l > 1$, $\mu = 1$. For any Borel set A we have:

$$\begin{aligned} & \int_A |\lambda^{n/q} u(\lambda(x - x_0)) - u(x - y_0)|^q = \int_{A+y_0} |(1+l)^{n/q} u((1+l)(x + y_0 - x_0)) - u(x)|^q \\ &= \int_{A+y_0} \left| (1+l)^{n/q} [u(x) + \langle \nabla u(x), lx + (1+l)(y_0 - x_0) \rangle] - u(x) + R(x) \right|^q \\ &= \int_{A+y_0} \left| \langle \nabla u(x), lx + (1+l)(y_0 - x_0) \rangle + \frac{n}{q} l \langle \nabla u(x), lx + (1+l)(y_0 - x_0) \rangle + \frac{n}{q} l u(x) + R(x) \right|^q \\ &= \int_{A+y_0} \left| |\nabla u(x)| \left\langle \frac{x}{|x|}, y_0 - x_0 \right\rangle + l \left[|x| |\nabla u(x)| + \frac{n}{q} u(x) \right] + R(x) \right|^q := \int_{A+y_0} |E_{x_0, y_0, l}(x) + R(x)|^q \end{aligned} \quad (4.16)$$

where the last inequality is due to the radial symmetry of ∇u and the error term $R(x)$ is given by

$$R(x) = (1+l)^{n/q} \left[\frac{|\nabla^2 u(x)|}{|x|^2} (x \otimes x) (lx + (1+l)(y_0 - x_0))^2 \right] = O(l^2) + O(|lx + (1+l)(y_0 - x_0)|^2).$$

Notice that there exists $\tilde{\rho}_1$ such that for every $x \in S_k$

$$|R(x)| \leq \frac{1}{2} |E_{x_0, y_0, l}(x)| \quad (4.17)$$

if $|x_0| + |y_0| + l \leq \tilde{\rho}_1$, since R is an infinitesimal of higher order than $E_{x_0, y_0, l}$. We aim now to find $\tilde{\rho}$ such that, for $|x_0| + |y_0| + l \leq \tilde{\rho}$, the following chain of inequalities holds true:

$$\int |E_{x_0, y_0, l} + R|^q \leq c_1 \int_{\Sigma_{k, y_0}} |E_{x_0, y_0, l} + R|^q \leq c_2 \int_{\Sigma_{k, y_0} \cap Q} |E_{x_0, y_0, l} + R|^q \leq c_2 \int_Q |E_{x_0, y_0, l} + R|^q \quad (4.18)$$

for suitable constants c_1, c_2 and k . This would immediately imply inequality (4.14). We begin remarking that

$$\lim_{k \rightarrow \infty} \int_{S_k} |E_{x_0, y_0, l} + R|^q = \int |E_{x_0, y_0, l} + R|^q$$

so there exists \bar{k} such that for $k \geq \bar{k}$ we have

$$\int_{S_k} |E_{x_0, y_0, l} + R|^q \geq \frac{1}{2} \int |E_{x_0, y_0, l} + R|^q.$$

Moreover, in view of (4.15) we get, for any $|y_0| \leq \tilde{\rho}$ small enough,

$$\int_{\Sigma(k,y_0)} |E_{x_0,y_0,l} + R|^q \geq \frac{1}{2} \int_{S_k} |E_{x_0,y_0,l} + R|^q.$$

So the first inequality in (4.18) holds with $c_1 = 1/4$. Since the last inequality in (4.18) is trivial, we are left to prove the central one. Suppose by contradiction that there is a sequence $(x_h, y_h, l_h) \rightarrow 0 \in \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$ as $h \rightarrow \infty$ such that (4.18) does not hold. Reasoning as before we get

$$\begin{aligned} & \int_{\Sigma(k,y_h)} |E_{x_h,y_h,l_h} + R|^q \leq (3/2)^q \int_{\Sigma(k,y_h)} |E_{x_h,y_h,l_h}|^q \\ & \leq (3/2)^q \int_{\Sigma(k,y_h)} |M \left\langle \frac{x}{|x|}, y_h - x_h \right\rangle + l_h (|x| |\nabla u(x)| + (n/q)u(x))|^q \\ & \leq (3/2)^q \int_{\Sigma(k,y_h)} |M|y_h - x_h| + l_h (|x| |\nabla u(x)| + (n/q)u(x))|^q \\ & \leq (3/2)^q \int_{\Sigma(k,y_h)} |M|v_h| + c_0 l_h|^q \end{aligned} \tag{4.19}$$

where we set $v_h = y_h - x_h$ and

$$c_0 = \limsup_{h \rightarrow \infty} \max_{\Sigma(k,y_h)} |x| |\nabla u(x)| + \frac{n}{q} u(x) > 0.$$

So there exist two constants k_0, k_1 such that

$$\int_{\Sigma(k,y_h)} |E_{x_h,y_h,l_h} + R|^q \leq k_0 \int_{\Sigma(k,y_h)} |k_1 l + |v_h||^q. \tag{4.20}$$

On the other hand we have

$$\begin{aligned} & \int_{\Sigma(k,y_h) \cap Q} |E_{x_h,y_h,l_h} + R|^q \geq \frac{1}{2^q} \int_{\Sigma(k,y_h) \cap Q} |E_{x_h,y_h,l_h}|^q \\ & = \frac{1}{2^q} \int_{\Sigma(k,y_h)} \left| |\nabla u(x)| \left\langle \frac{x}{|x|}, \frac{v_h}{|v_h|} \right\rangle |v_h| + l (|x| |\nabla u(x)| + (n/q)u(x)) \right|^q. \end{aligned}$$

We have now three possible situations: $|v_h| \ll l_h$, $l_h \ll |v_h|$ or $l_h \simeq |v_h|$ as $h \rightarrow \infty$. In the first case we have, thanks to (4.20), that

$$\int_{\Sigma(k,y_h)} |E_{x_h,y_h,l_h} + R|^q \leq k_2 \int_{\Sigma(k,y_h)} l_h^q$$

for a suitable k_2 . Moreover it is easy to find positive constants k_3 and k_4 not depending on l_h and v_h such that

$$\int_{\Sigma(k,y_h) \cap Q} |E_{x_h,y_h,l_h} + R|^q \geq k_3 \int_{\Sigma(k,y_h) \cap Q} l_h^q \geq k_4 \int_{\Sigma(k,y_h)} l_h^q.$$

So in this case (4.18) holds with $c_2 = \frac{k_3}{k_4} c_1$ (or, in other terms, we get a contradiction). The second case, $l_h \ll |v_h|$ can be solved with the same argument, with the only observation that, for the estimate from above, we must further restrict the set of integration to the set (for instance) $U = \{x \in Q : |\langle x/|x|, v_0/|v_0| \rangle| \geq 1/10\}$. We are left to study the case where $|v_h| \simeq l_h$. If $\liminf_h |v_h|/l_h \geq \tilde{c} > 0$, we have that

$$\int_{\Sigma(k,y_h)} |E_{x_h,y_h,l_h} + R|^q = l_h^q \int_{\Sigma(k,y_h)} \left| c_0 \frac{l_h}{|v_h|} + M \right|^q.$$

Let us define $V = \{x \in Q : |\langle x/|x|, v_h/|v_h| \rangle| \leq \alpha\}$ where α is a constant (depending on \tilde{c}) that will be fixed later. We have

$$\begin{aligned} \int_{\Sigma(k, y_h) \cap Q} |E_{x_h, y_h, l_h} + R|^q &\geq \int_{\Sigma(k, y_h) \cap V} |E_{x_h, y_h, l_h} + R|^q \\ &\geq \kappa_0 l^q \int_{\Sigma(k, y_h) \cap V} \left| c_1 \frac{l}{|v_h|} - M\alpha \right|^q. \end{aligned}$$

Choosing α small enough, since $l_h/|v_h| \gg 0$, we can find constants k_1 and k_2 not depending on l_h and v_h such that

$$\int_{\Sigma(k, y_h) \cap Q} |E_{x_h, y_h, l_h} + R|^q \geq \kappa_1 l^q \int_{\Sigma(k, y_h) \cap V} \left| c_0 \frac{l_h}{|v_h|} + M \right|^q \geq \kappa_2 l^q \int_{\Sigma(k, y_h)} \left| c_0 \frac{l_h}{|v_h|} + M \right|^q,$$

so again (4.18) holds with $c_2 = k_2/c_1$ (thus again a contradiction). If $\liminf_h l_h/|v_h| \geq \tilde{c} > 0$, we can conclude applying an analogous argument. \square

We pass now to prove Theorem 4.1. For the sake of clearness we divide its proof into two parts. We first prove a proposition which provides us a method to pass from a generic function in $D^{p,s}(\mathbb{R}^n)$ to an $(n-1)$ -symmetric function which satisfies the reduction inequalities (4.1). Then we shall see how to obtain the last required symmetry.

Proposition 4.6. *There exists a positive constant C such that for every function $u \in D^{p,s}(\mathbb{R}^n)$ there is an $(n-1)$ -symmetric function \tilde{u} such that*

$$\lambda(u) \leq C\lambda(\tilde{u}), \quad \delta(\tilde{u}) \leq 2^{n-1}\delta(u). \quad (4.21)$$

Proof. As usual, by the homogeneity of the deficit and the asymmetry, we can consider $\|u\|_q = 1$. Moreover we can suppose that $\delta(u) < \bar{\delta}$ for an arbitrary small $\bar{\delta}$. Indeed, if $\delta(u) \geq \bar{\delta}$, let v be a radial (and so n -symmetric!) function such that $0 < \delta(v) < 2^{n-1}\bar{\delta}$. Then

$$\lambda(u) \leq 2^q = \frac{2^q}{\lambda(v)}\lambda(v) \leq \bar{C}\lambda(v), \quad \delta(v) \leq 2^{n-1}\bar{\delta} \leq 2^{n-1}\delta(u).$$

Consider, for $k = 1 \dots n$, the n hyperplanes orthogonal to the coordinate axis such that

$$\int_{H_k^+} u^q = \int_{H_k^-} u^q = \frac{1}{2}$$

where H_k^\pm are the two half spaces in which \mathbb{R}^n is divided by H_k . Denoting by T_k the reflection with respect to H_k , we define

$$u_k^+(x) = \begin{cases} u(x) & \text{if } x \in H_k^+ \\ u(T_k(x)) & \text{if } x \in H_k^- \end{cases} \quad (4.22)$$

$$u_k^-(x) = \begin{cases} u(x) & \text{if } x \in H_k^- \\ u(T_k(x)) & \text{if } x \in H_k^+ \end{cases}$$

By construction u_k^\pm are symmetric with respect to H_k . We observe now that

$$\int u^s = \frac{\int (u_k^+)^s + \int (u_k^-)^s}{2}, \quad \int |\nabla u|^p = \frac{\int |\nabla u_k^+|^p + \int |\nabla u_k^-|^p}{2},$$

and since $t \rightarrow t^{1/p}$ and $t \rightarrow t^{1/s}$ are concave functions, we have that

$$\|u\|_s \geq \frac{\|u_k^+\|_s + \|u_k^-\|_s}{2}, \quad \|\nabla u\|_p \geq \frac{\|\nabla u_k^+\|_p + \|\nabla u_k^-\|_p}{2}. \quad (4.23)$$

By the definition of $\delta(u)$ and since $(x, y) \mapsto x^\theta y^{1-\theta}$ is concave on \mathbb{R}_+^2 and strictly increasing in x and y , we get that

$$\begin{aligned} G\delta(u) &\geq \left(\frac{\|\nabla u_k^+\|_p + \|\nabla u_k^-\|_p}{2} \right)^\theta \left(\frac{\|u_k^+\|_s + \|u_k^-\|_s}{2} \right)^{1-\theta} - G \\ &\geq \frac{G}{2} \delta(u_k^+) + \frac{G}{2} \delta(u_k^-), \end{aligned}$$

and so

$$\delta(u) \geq \frac{\delta(u_k^+) + \delta(u_k^-)}{2}.$$

In particular for every $k = 1, \dots, n$

$$\max\{\delta(u_k^+), \delta(u_k^-)\} \leq 2\delta(u).$$

Let v_k^+ and v_k^- be the functions which minimize $\lambda(u_k^\pm | H_k)$. Then, by triangle inequality and Lemma 4.3 we have

$$\begin{aligned} \lambda(u) &\leq \int |u - v_k^+|^q = \int_{H_k^+} |u_k^+ - v_k^+|^q + \int_{H_k^-} |u_k^- - v_k^+|^q \\ &\leq 2^{q-1} \left(\frac{\lambda(u_k^+ | H_k) + \lambda(u_k^- | H_k)}{2} + \int_{H_k^-} |v_k^+ - v_k^-|^q \right) \\ &\leq 2^{q-2} 3^q \left(\lambda(u_k^+) + \lambda(u_k^-) + \int_{H_k^-} |v_k^+ - v_k^-|^q \right). \end{aligned}$$

We claim that if $\delta(u)$ is small enough, then for any couple of indexes $1 \leq i < j \leq n$, for $k_1 = i$ or $k_1 = j$ the following inequality holds:

$$\int_{H_{k_1}^-} |v_{k_1}^+ - v_{k_1}^-|^q \leq C \left(\int_{H_{k_1}^+} |u_{k_1}^+ - v_{k_1}^+|^q + \int_{H_{k_1}^-} |u_{k_1}^- - v_{k_1}^-|^q \right). \quad (4.24)$$

Let us show how this brings to the conclusion: by (4.23) and (4.24) we would have

$$\lambda(u) \leq C \max\{\lambda(u_{k_1}^+), \lambda(u_{k_1}^-)\}, \quad \max\{\delta(u_{k_1}^+), \delta(u_{k_1}^-)\} \leq 2\delta(u).$$

Thus we would obtain that $u_{k_1}^+$ or $u_{k_1}^-$, say $u_{k_1}^+$, is a 1-symmetric function with L^q norm equal to 1. We can now iterate this procedure exploiting two hyperplanes between the $(n-1)$ we left, obtaining a 2-symmetric function which satisfies the reductions inequalities (4.21) (with $u_{k_1}^+$ in place of u). We can continue such a construction until we have just one hyperplane left. But then we would have an $(n-1)$ -symmetric function which satisfies inequalities (4.21) that is the claim of the proposition. Thus we are only left to prove (4.24). We divide the proof of it into two further steps:

Step 1 There exists a positive constant C_0 such that for all $1 \leq i < j \leq n$, $\sigma, \tau \in \{+, -\}$ it holds

$$\int |v_i^\sigma - v_j^\tau|^q \leq C_0 \int_{H_i^\sigma \cap H_j^\tau} |v_i^\sigma - v_j^\tau|^q. \quad (4.25)$$

Step 2 (4.25) implies (4.24).

To verify (4.25) we notice that thanks to Lemma 4.5 the inequality is verified if we are able to find two positive constants ρ and C_1 such that:

$$(i) \int (v_i^\sigma)^\rho = \int (v_j^\tau)^\rho = 1;$$

- (ii) H_i^σ ed H_j^τ are two orthogonal half spaces which contains on their boundary the center of symmetry of v_i^σ e v_j^τ ;
- (iii) $\int |v_i^\sigma - v_j^\tau|^q \leq \rho$.

We need to check (i) – (iii). Clearly (i) and (ii) are true by construction. So we are left to verify (iii). We have

$$\|v_i^\sigma - v_j^\tau\|_q \leq \|v_i^\sigma - u_i^\sigma\|_q + \|u_i^\sigma - u\|_q + \|u - u_j^\tau\|_q + \|u_j^\tau - v_j^\tau\|_q. \quad (4.26)$$

Thanks to Lemma 4.4

$$\int |u_i^\sigma - u|^q = \frac{1}{2} \int |u \circ T_i - u|^q \leq C\lambda(u)^{1/q}. \quad (4.27)$$

Moreover

$$\int |v_i^\sigma - u_i^\sigma|^q \leq 2^{q-1} \left(\lambda(u|_{H_i}) + \|u - u_i^\sigma\|_p^p \right) \leq C\lambda(u)^{1/q}. \quad (4.28)$$

The same estimate holds as well for the third and the fourth addend on the right hand side of (4.26). Putting together (4.27) and (4.28) we obtain the claim of (iii) and this conclude the proof of Step 1. Let us prove now Step 2. Suppose, to fix the ideas, that $i = 1$ and $j = 2$. For $k = 1, 2$ we set

$$h_k = v_k^+ \chi_{H_k^+} + v_k^- \chi_{H_k^-}$$

where χ_A denotes the characteristic function of the set A . Thanks to (4.25),

$$\int |h_1 - h_2|^q \geq \int_{H_1^+ \cap H_2^+} |h_1 - h_2|^q = \int_{H_1^+ \cap H_2^+} |v_1^+ - v_2^+|^q \geq \frac{1}{C} \int |v_1^+ - v_2^+|^q.$$

With a similar argument, using $H_1^- \cap H_2^+$ instead of $H_1^+ \cap H_2^+$ we get

$$\int |h_1 - h_2|^q \geq \frac{1}{C} \int |v_1^- - v_2^+|^q.$$

Hence

$$\int |v_1^+ - v_1^-|^q \leq 2^q C \int |h_1 - h_2|^q. \quad (4.29)$$

Similarly we can see that

$$\int |v_2^+ - v_2^-|^q \leq 2^q C \int |h_1 - h_2|^q. \quad (4.30)$$

Furthermore we have

$$\begin{aligned}
\int |h_1 - h_2|^q &\leq 2^{q-1} \left(\int |h_1 - u|^q + \int |h_2 - u|^q \right) \\
&= 2^{q-1} \left(\int_{H_1^+} |v_1^+ - u_1^+|^q + \int_{H_1^-} |v_1^- - u_1^-|^q \right. \\
&\quad \left. + \int_{H_2^+} |v_2^+ - u_2^+|^q + \int_{H_2^-} |v_2^- - u_2^-|^q \right) \\
&\leq 2^q \max \left\{ \int_{H_1^+} |v_1^+ - u_1^+|^q + \int_{H_1^-} |v_1^- - u_1^-|^q, \right. \\
&\quad \left. \int_{H_2^+} |v_2^+ - u_2^+|^q + \int_{H_2^-} |v_2^- - u_2^-|^q \right\} \\
&= 2^q \left(\int_{H_k^+} |v_k^+ - u_k^+|^q + \int_{H_k^-} |v_k^- - u_k^-|^q \right).
\end{aligned} \tag{4.31}$$

putting together (4.29), (4.30) and (4.31) we obtain the claim of Step 2. \square

We stress that we can not symmetrize directly our function once again. A formal argument which shows a problem that may occur is the following: consider a function v such that $\delta(v) = \lambda(v) = 0$ and construct a function u as follows:

$$u(x) = v(x)\chi_{\{x_1 \geq 0\}}(x) + 2^{n/q}v(2x)\chi_{\{x_1 < 0\}}(x).$$

It follows that u is $(n-1)$ -symmetric in $H_k = \{x_k = 0\}$ for $k \neq 1$. if we try to symmetrize such function with respect to H_1 we would obtain $u_+(x) = 2^{n/q}(v(2x))$ and $u_-(x) = v(x)$. Clearly none of them satisfy the first inequality in (4.1). However we are going to see that a more refined kind of symmetrization can bypass this problem.

Proof of Theorem 4.1. Assume as usual that $\|u\|_q = 1$. We can assume, thanks to Proposition 4.6, that u is an $(n-1)$ -symmetric function and that $\delta(u) < \bar{\delta}$ with $\bar{\delta}$ arbitrarily small. Up to a rotation and a translation, we can consider u to be symmetric with respect to the coordinate axes $\{x_k = 0\}$ for $k = 2, \dots, n$ and such that

$$\int_{\{x_k > 0\}} u^q = \frac{1}{2} = \int_{\{x_k < 0\}} u^q.$$

Let u^\pm be the two symmetrizations of u with respect to the hyperplane $\{x_1 = 0\}$, constructed as in Proposition 4.6. We have that $\max\{\delta(u^+), \delta(u^-)\} \leq 2\delta(u)$. So, if $\min\{\lambda(u^+), \lambda(u^-)\} \geq C_0\lambda(u)$, we would be done. Thus we can suppose that

$$\max\{\lambda(u^+), \lambda(u^-)\} < \varepsilon\lambda(u) \tag{4.32}$$

for some constant ε to be chosen. Consider $Q = \{|x_1| \leq x_2\}$, $Q^+ = Q \cap \{x_1 > 0\}$ and $Q^- = Q \cap \{x_1 < 0\}$ and define a function \hat{u} as follows:

$$\hat{u}(x) = \begin{cases} u(x) & \text{if } x \in Q \\ u(R_1x) & \text{if } x \in R_1(Q) \\ \hat{u}(R_2x) & \text{if } x \in R_2(Q \cup R_1(Q)) \end{cases}$$

where R_1 and R_2 are the reflections with respect to $\{x_1 = x_2\}$ and $\{x_1 = -x_2\}$ respectively. The function \hat{u} satisfies all the symmetries of u with the exception of the one related to the hyperplane $\{x_2 = 0\}$, but by construction it is symmetric also with respect to $\{x_1 = \pm x_2\}$. So it is n -symmetric. It remains to show that \hat{u} satisfies the reduction inequalities (4.1). Let us start with the one which concerns the asymmetry. To this aim we denote by \hat{v} , v^+ and v^- the functions which achieve the minimum of $\lambda(\hat{u}|\{0\})$, $\lambda(u^+|\{0\})$ and $\lambda(u^-|\{0\})$ respectively. Since $\int \hat{u}^q \leq 4$, we get that

$$\begin{aligned} 3^q \lambda(\hat{u}) &\geq \lambda(\hat{u}|\{0\}) = \frac{\int |\hat{u} - \hat{v}|^q}{\int \hat{u}^q} = \frac{4}{\int \hat{u}^q} \int_Q |u - \hat{v}|^q \\ &\geq \int_{Q^+} |u^+ - \hat{v}|^q + \int_{Q^-} |u^- - \hat{v}|^q \\ &= \int_{Q^+} |u^+ - \hat{v}|^q + \int_{Q^+} |u^- - \hat{v}|^q \geq \frac{1}{2^{q-1}} \int_{Q^+} |u^+ - u^-|^q. \end{aligned}$$

The first inequality in (4.1) is then true if we can estimate $\lambda(u)$ in terms of $\int_{Q^+} |u^+ - u^-|^q$. To this aim we observe that

$$\begin{aligned} \|u^+ - u^-\|_{L^q(Q^+)} &= \frac{1}{2} \|u^+ - u^-\|_{L^q(Q)} \\ &= \frac{1}{2} \|(v^+ - v^-) - (v^+ - u^+) - (u^- - v^-)\|_{L^q(Q)} \\ &\geq \frac{1}{2} \left[\|v^+ - v^-\|_{L^q(Q)} - \|u^+ - v^+\|_{L^q(Q)} - \|u^- - v^-\|_{L^q(Q)} \right]. \end{aligned} \quad (4.33)$$

Moreover

$$\int_Q |u^\pm - v^\pm|^q \leq \int |u^\pm - v^\pm|^q = \lambda(u^\pm|\{0\}) \leq 3^q \lambda(u^\pm) \leq \varepsilon 3^q \lambda(u) \quad (4.34)$$

where we exploited Lemma 4.3 and Lemma 4.4 and the fact that u^\pm are n -symmetric functions. Thanks to (4.33) and (4.34) we obtain

$$\|u^+ - u^-\|_{L^q(Q^+)} \geq \frac{1}{2} \left[\|v^+ - v^-\|_{L^q(Q)} - 2 \left(\frac{3^q}{C(\varepsilon)} \lambda(u) \right)^{1/q} \right],$$

where $C(\varepsilon)$ is a suitable positive constant. Furthermore, always thanks to (4.34) we have

$$\begin{aligned} \lambda(u) &\leq \int |u - v^+|^q = \frac{1}{2} \int |v^+ - u^+|^q + \frac{1}{2} \int |v^+ - u^-|^q \\ &\leq \varepsilon \frac{3^q}{2} \lambda(u) + 2^{q-2} \left(\varepsilon 3^q \lambda(u) + \int |v^+ - v^-|^q \right), \end{aligned}$$

and so

$$\int |v^+ - v^-|^q \geq \frac{1}{2^{q-2}} \left[\lambda(u) - \varepsilon \frac{3^q}{2} \lambda(u) - \varepsilon 3^q \lambda(u) \right] \geq \frac{1}{2^q} \lambda(u) \quad (4.35)$$

where last inequality is justified if we choose ε small enough. Summarizing we obtained

$$\|u^+ - u^-\|_{L^q(Q^+)} \geq C_0 \lambda(u)^{1/q},$$

where C_0 is a suitable positive constant not depending on u . Let us consider now the second inequality in (4.1), that is the one concerning the deficit. Notice that we do not know if $\|\hat{u}\|_q = 1$. Nevertheless we have

$$\left| \|\hat{u}\|_{L^q(Q^+)} - (1/8)^{1/q} \right| = \left| \|u^+\|_{L^q(Q^+)} - \|v^+\|_{L^q(Q^+)} \right| \leq \|u^+ - v^+\|_{L^q(Q^+)}.$$

and, since $|s^q - t^q| \leq C_1|s - t|$ for a suitable constant C_1 if s and t are in $[0, 1]$, we get

$$\begin{aligned} \left| \int_{Q^+} \hat{u}^q - \frac{1}{8} \right| &\leq C_2 \|u^+ - v^+\|_q = C_2 \lambda (u^+ \setminus \{0\})^{1/q} \\ &\leq 3^{1/q} C_3 \lambda (u^+)^{1/q} \leq 3^{1/q} C_4 \delta (u^+)^{\alpha/q} \leq 2C_4 \delta (u)^{\alpha/q}, \end{aligned} \quad (4.36)$$

where we used Lemma 4.3, the fact that u is n -symmetric and that $\delta(u^+) \leq 2\delta(u)$ and where C_2 , C_3 and C_4 are positive constants. Analogous estimates hold on Q^- , on $U^+ = \{x_1 > 0, x_2 > 0\} \setminus Q$ and on $U^- = \{x_1 < 0, x_2 > 0\} \setminus Q$. Then, by the triangle inequality we get

$$\left| \int \hat{u}^q - 1 \right| \leq C \delta (u)^{\alpha/q} \quad (4.37)$$

for a suitable $\alpha > 0$. Let us recall the definition of the functionals F and G given in (2.2):

$$G(u) = \|\nabla u\|_p^\theta \|u\|_s^{1-\theta}, \quad F(u) = \int |\nabla u|^p + \int |u|^s. \quad (4.38)$$

By Lemma 2.2 we know that there exist two positive constants κ and η_0 such that for every u there exists $\lambda > 0$ such that $F(\tau_\lambda u) = \eta_0 G(u)^\kappa$ where $\tau_\lambda u(x) = \lambda^{n/q} u(\lambda x)$. Furthermore such λ minimizes the function $\mu \mapsto F(\tau_\mu u)$ in \mathbb{R}^+ . So we get

$$G^\kappa \|\hat{u}\|_q^\kappa (1 + \delta(\hat{u}))^\kappa = G(\hat{u})^\kappa \leq \left(\frac{1}{\eta_0} \int |\nabla \tau_\lambda \hat{u}|^p + \frac{1}{\eta_0} \int |\tau_\lambda \hat{u}|^s \right) \quad (4.39)$$

for all $\lambda > 0$. Then

$$\begin{aligned} \frac{1}{\eta_0} \left(\int |\nabla \tau_\lambda \hat{u}|^p + \int |\tau_\lambda \hat{u}|^s \right) &= \frac{4}{\eta_0} \left(\int_{Q^+ \cup Q^-} |\nabla \tau_\lambda \hat{u}|^p + \int_{Q^+ \cup Q^-} |\tau_\lambda \hat{u}|^s \right) \\ &= \frac{4}{\eta_0} \left(\int_{\{x_2 > 0\}} (|\nabla \tau_\lambda u|^p + |\tau_\lambda u|^s) - \int_{U^+ \cup U^-} (|\nabla \tau_\lambda u|^p + |\tau_\lambda u|^s) \right). \end{aligned} \quad (4.40)$$

Choosing $\lambda > 0$ such that $F(\tau_\lambda u) = \eta_0 G(u)^\kappa$, since u is symmetric with respect to $\{x_2 = 0\}$, we get

$$\begin{aligned} \int_{\{x_2 > 0\}} (|\nabla \tau_\lambda u|^p + |\tau_\lambda u|^s) &= \frac{1}{2} F(\tau_\lambda u) \\ &= \frac{\eta_0}{2} G(u)^\kappa = \frac{\eta_0 G^\kappa}{2} (\delta(u) + 1)^\kappa \\ &\leq \frac{\eta_0 G^\kappa}{2} (1 + C_5 \delta(u)), \end{aligned} \quad (4.41)$$

where the last inequality is true for $\delta(u)$ small enough. Let us consider now the function

$$v(x) = \begin{cases} \tau_\lambda u(x) & x \in U^+ \\ \tau_\lambda u(R_1 x) & x \in R_1 U^+ \\ v(S_1 x) & x \in \{x_1 < 0, x_2 > 0\} \\ v(S_2 x) & x \in \{x_2 < 0\} \end{cases}$$

where $S_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$ denotes the symmetrization map with respect to $\{x_i = 0\}$. We have

$$\begin{aligned} \int_{U^+} (|\nabla \tau_\lambda(u)|^p + |\tau_\lambda(u)|^s) &= \frac{1}{8} \left(\int (|\nabla v|^p + |v|^s) \right) \geq \frac{1}{8} \min_{\mu \in \mathbb{R}^+} F(\tau_\mu v) \\ &= \frac{\eta_0}{8} G(v)^\kappa \geq \frac{\eta_0 G^\kappa}{8} \left(\int v^q \right)^{\kappa/q} = \frac{\eta_0 G^\kappa}{8} \cdot 8^{\kappa/q} \left(\int_{U^+} u^q \right)^{\kappa/q} \quad (4.42) \\ &\geq \eta_0 G^\kappa 8^{-1+\kappa/q} \left(\frac{1}{8} - C_6 \delta(u)^\alpha \right)^{\kappa/q} \geq \eta_0 G^\kappa \left(\frac{1}{8} - C_7 \delta(u)^\alpha \right), \end{aligned}$$

for suitable constants C_6 and C_7 and for $\delta(u)$ small enough. Notice that in the last inequality we made use of (4.36). An analogous estimate can be obtained on U^- . By (4.39), (4.40), (4.41) and (4.42) we get

$$\begin{aligned} G^\kappa \|\hat{u}\|_q^\kappa (1 + \delta(\hat{u}))^\kappa &\leq \frac{4}{\eta_0} \left[\frac{\eta_0 G^\kappa}{2} (1 + C_5 \delta(u)) - 2\eta_0 G^\kappa \left(\frac{1}{8} - C_7 \delta(u) \right)^\beta \right] \\ &= G^\kappa \left[1 + C_8 \delta(u)^\alpha \right] \quad (4.43) \end{aligned}$$

for a suitable $C_8 > 0$ and where α is the minimum between 1 and β . By (4.43) and thanks to (4.37) we conclude that

$$\delta(\hat{u}) \leq \frac{1 + C_9 \delta(u)^\alpha}{1 - C_9 \delta(u)^\alpha} - 1 \leq C \delta(u)^\alpha$$

that is true again for $\delta(u)$ small enough. The conclusion follows setting $\kappa_2 = C$. □

5. ACKNOWLEDGEMENTS

The author wish to warmly thank F. Maggi for its enlightening advise during the development of this work, and L. Brasco, G. De Philippis and A. Pratelli for several useful discussions about the topic.

REFERENCES

- [1] T. Aubin, *Problèmes isopérimétriques et espaces de Sobolev*, J. Differential Geometry, **11** (1976), 573-598.
- [2] J. E. Brothers, W. P. Ziemer, *Minimal rearrangements of Sobolev functions*, J. reine angew. Math. **384** (1988), 153-179.
- [3] M. Cicalese, G. Leonardi, *A selection principle for the sharp quantitative isoperimetric inequality*, Arch. Ration. Mech. Anal., **206** (2012), 617-643.
- [4] E. A. Carlen, R. L. Frank, E. H. Lieb *Stability estimates for the lowest eigenvalue of a Schrödinger operator*, Geom. Funct. Anal., to appear (2013).
- [5] G. Bianchi, H. Egnell, *A note on the Sobolev inequality*, J. Funct. Anal., **100** (1991), 18-24.
- [6] D. Cordero-Erausquin, B. Nazaret, C. Villani, *A Mass-Transportation Approach to Sobolev and Nirenberg Inequalities*, Adv. Math., **182** (2004), 307-332.
- [7] E. A. Carlen, A. Figalli, *Stability for a GNS inequality and the log-HLS inequality, with application to the critical mass Keller-Segel equation*. Duke Math. J., **162** (2013), 579-625.
- [8] A. Cianchi, N. Fusco, F. Maggi, A. Pratelli, *The sharp Sobolev inequality in quantitative form*, J. Eur. Math. Soc., **11** (2009), 1105-1139.
- [9] M. Del Pino, J. Dolbeault, *Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions*, J. Math. Pures Appl., **81** (2002) 847-875.
- [10] L.C. Evans, R. F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, (1992).
- [11] A. Figalli, F. Maggi, A. Pratelli, *A mass transportation approach to quantitative isoperimetric inequalities*, Invent. Math., 182 (2010), no. 1, 167-211.

- [12] N. Fusco, V. Julin *A strong form of the quantitative isoperimetric inequality*, Calc. Var. Partial Differential Equations, to appear (2013).
- [13] N. Fusco, F. Maggi, A. Pratelli, *The sharp quantitative isoperimetric inequality*. Ann. of Math., **168** (2008), 941-980.
- [14] N. Fusco, F. Maggi, A. Pratelli, *The sharp quantitative Sobolev inequality for functions of bounded variation*, J. Funct. Anal., **244** (2007), 315-341.
- [15] R. R. Hall, *A quantitative isoperimetric inequality in n -dimensional*, J. Reine Angew. Math., **428** (1992), 161-176.
- [16] R. R. Hall, W. K. Hayman, A. W. Weitsman, *On asymmetry and capacity*, J. d'Analyse Math., **56** (1991), 87-123.
- [17] E. H. Lieb, M. Loss, *Analysis*, Graduate studies in Mathematics, AMS, (1996).
- [18] P.L. Lions, *The concentration-compactness principle in the Calculus of Variation. The Locally compact case, part 1*, Annales de l'I. H. P., **4** (1984), 109-145.
- [19] P.L. Lions, *The Concentration-Compactness Principle in the Calculus of Variations. The locally compact case, part 2*, Revista Matemática iberoamericana, **2** (1985), 45-121.
- [20] F. Maggi, *Sets of Finite Perimeter and Geometric Variational Problems*, Cambridge Studies in Advanced Mathematics 135, Cambridge University Press, (2012).
- [21] F. Maggi, *Some methods for studying stability in isoperimetric type problems*, Bull. Amer. Math. Society, **45** (2008), 367-408.
- [22] M. Struwe, *Variational Methods*, **34**, Springer-Verlag.
- [23] J. Serrin, M. Tang, *Uniqueness of ground states for quasilinear elliptic equations*, Indiana Univ. Math. J., **49** (2000), 897-923.
- [24] G. Talenti, *Best constants in Sobolev inequality*. Ann. Mat. Pura Appl., (1976), 353-372.

SCUOLA NORMALE SUPERIORE DI PISA, PIAZZA DEI CAVALIERI 4, 56126 PISA, ITALY
E-mail address: berardo.ruffini@sns.it