

ON TRANSVERSAL SUBMANIFOLDS AND THEIR MEASURE

VALENTINO MAGNANI, JEREMY T. TYSON, AND DAVIDE VITTONI

Abstract: We study the class of transversal submanifolds in Carnot groups. We characterize their blow-ups at transversal points and prove a negligibility theorem for their “generalized characteristic set”, with respect to the Carnot-Carathéodory Hausdorff measure. This set is made by all points of non-maximal degree. Observing that C^1 submanifolds in Carnot groups are generically transversal, the previous results prove that the “intrinsic measure” of C^1 submanifolds is generically equivalent to their Carnot-Carathéodory Hausdorff measure. As a result, the restriction of this Hausdorff measure to the submanifold can be replaced by a more manageable integral formula, that should be seen as a “sub-Riemannian mass”. Another consequence of these results is an explicit formula, only depending on the embedding of the submanifold, that computes the Carnot-Carathéodory Hausdorff dimension of C^1 transversal submanifolds.

CONTENTS

1. Introduction	2
2. Notation and preliminary results	5
3. Blow-up at transversal points	12
4. Negligibility of lower degree points in transversal submanifolds	17
5. Size of the characteristic set for $C^{1,\lambda}$ submanifolds	26
References	28

Date: July 10, 2013.

2010 Mathematics Subject Classification. Primary 53C17; Secondary 22E25, 28A78.

Key words and phrases. Stratified groups, submanifolds, Hausdorff measure.

The first author acknowledges the support of the European Project ERC AdG *GeMeThNES*.

The second author acknowledges the support of the US National Science Foundation Grants DMS-0901620 and DMS-1201875.

The third author acknowledges the support of MIUR, GNAMPA of INDAM (Italy), University of Padova, Fondazione CaRiPaRo Project “Nonlinear Partial Differential Equations: models, analysis, and control-theoretic problems” and University of Padova research project “Some analytic and differential geometric aspects in Nonlinear Control Theory, with applications to Mechanics”.

1. INTRODUCTION

A stratified group \mathbb{G} is a connected, simply connected nilpotent Lie group, whose Lie algebra \mathcal{G} has a special grading that allows for the existence of natural dilations along with a homogeneous distance that respect both dilations and group operation. A stratified group equipped with a sub-Riemannian distance is also known as Carnot group. The first developments of Geometric Analysis in Carnot groups were mainly focused on geometric properties of domains in relation with Sobolev embeddings (see, for instance, [6], [12], [16]), problems from the calculus of variations (e.g., [5], [10]), differential geometric calculus on hypersurfaces ([9]), and the structure of finite perimeter sets (a very incomplete list of references includes [1], [7], [8], [13], [14], and [20]). The preceding lists of references are far from exhaustive, representing only a small sample of the rapidly expanding literature in the field of sub-Riemannian geometric analysis.

The study of finite perimeter sets and domains naturally connects with the study of hypersurfaces and their Hausdorff measure. The fact that this measure is constructed by a fixed homogeneous distance of the group is understood. An important object in this context is the so-called \mathbb{G} -perimeter measure. It can be defined using a volume measure and a smooth left invariant metric on the horizontal subbundle of the group. This measure is equivalent, in the sense of (1.2) below, to the $(Q - 1)$ -dimensional Hausdorff measure either of the reduced boundaries in step two Carnot groups, [14], or of the topological boundaries of C^1 smooth domains in arbitrary stratified groups, [20], where Q is the Hausdorff dimension of \mathbb{G} .

The \mathbb{G} -perimeter measure for regular sets has a precise integral formula that replaces the Hausdorff measure and that does not contain the homogeneous distance. In fact, it is more manageable for minimization problems. In the development of Geometric Measure Theory on stratified groups, a natural question arises: what is the “right measure” replacing the \mathbb{G} -perimeter measure for higher codimensional sets?

In [23], a general integral formula for the “intrinsic measure” of C^1 submanifolds has been found: let Σ be a C^1 smooth submanifold of \mathbb{G} and define

$$(1.1) \quad \mu_{\Sigma}(U) = \int_{\Phi^{-1}(U)} \|(\partial_{t_1} \Phi \wedge \cdots \wedge \partial_{t_p} \Phi)_{D, \Phi(t)}\| dt$$

where $\Phi : A \rightarrow U \subset \Sigma$ is a local parametrization of Σ , A is an open set of \mathbb{R}^p and D is the degree of Σ , see Subsection 2.4 for more details. This measure yields the perimeter measure in codimension one and in several cases it is equivalent to $\mathcal{H}^D \llcorner \Sigma$ up to geometric constants, namely,

$$(1.2) \quad C^{-1} \mathcal{H}^D \llcorner \Sigma \leq \mu_{\Sigma} \leq C \mathcal{H}^D \llcorner \Sigma.$$

We stress the fact that the degree D of Σ is also equal to the Hausdorff dimension of Σ . This equivalence already appears in [23] for $C^{1,1}$ smooth submanifolds in stratified groups, under the key assumption that points of degree less than D are \mathcal{H}^D -negligible. Under this assumption, the equivalence (1.2) is a consequence of a “blow-up theorem” performed at each point of degree D , see [23, Theorem 1.1]. For more details on the notion of degree, see Subsection 2.4.

The previously mentioned \mathcal{H}^D -negligibility condition holds in many cases: for $C^{1,1}$ smooth submanifolds in two step stratified groups [22], and in the Engel group [19], for C^1 smooth non-horizontal submanifolds in all stratified groups [20, 21], and for C^1 smooth curves in all groups [18]. In all these cases, the equivalence (1.2) holds. In fact, when Σ is $C^{1,1}$ this is a consequence of the blow-up theorem of [23], while for the case of C^1 smoothness the blow-up at points of degree D is established in [21] for non-horizontal submanifolds and in [18] for all curves.

Surprisingly, for C^1 smooth submanifolds in stratified groups the equivalence (1.2) is an intriguing open question. One of the reasons behind this new difficulty is that, in higher codimension, submanifolds may belong to different classes, namely, they may have different Hausdorff dimensions, while keeping the same topological dimension. Simple examples of this phenomenon are given by the one dimensional homogeneous subgroups, that have different Hausdorff dimensions according to their degree. Clearly, analogous examples can be easily found for higher dimensional homogeneous subgroups. It is instructive to compare these cases with that of codimension one submanifolds, whose Hausdorff dimension must equal $Q - 1$.

Can we detect the “right” class of submanifolds that has the “good behaviour” of hypersurfaces, and replaces them in higher codimension? When the codimension is “low”, precisely less than the dimension of the *horizontal fibers*, this class is formed by non-horizontal submanifolds, for which (1.2) holds, [21]. In higher codimension, this class is formed by *transversal submanifolds*. A transversal submanifold is easily defined as a top-dimensional submanifold among all submanifolds having the same topological dimension p . We have a precise formula for this maximal Hausdorff dimension $D(p)$, see Section 2 for precise definitions.

In this paper, we prove that transversal submanifolds in arbitrary codimension have properties similar to those of hypersurfaces. In fact, our main result is that (1.2) holds for all C^1 smooth transversal submanifolds in arbitrary stratified groups. This follows by combining two key results: a blow-up theorem and a negligibility result, that are stated below. The estimates (1.2) show in particular that the Hausdorff dimension of C^1 smooth transversal submanifolds is equal to $D(p)$. This fact should be compared with [17, 0.6.B], where M. Gromov provides a formula for the Hausdorff dimension of generic submanifolds. Gromov also introduces the number $D_H(\Sigma)$ associated with a submanifold Σ ; this number coincides with the degree $d(\Sigma)$ introduced in Subsection 2.4, see [22, Remark 2] for a proof of this fact.

Another motivation for our study of transversal submanifolds is that C^1 smooth submanifolds are generically transversal, namely, “most” C^1 submanifolds are transversal. This suggests that these submanifolds are important in the subsequent study of higher codimensional submanifolds in Carnot groups. The fact that transversality is a generic property can be seen for instance as a simple consequence of our Lemma 2.11 and then arguing as in [21, Section 4].

The main results of our work are a “blow-up theorem” and an $\mathcal{H}^{D(p)}$ -negligibility theorem for all C^1 smooth transversal submanifolds. These theorems extend the blow-up theorem of [21] and the negligibility theorem of [20].

Theorem 1.1 (Blow-up theorem). *If Σ is a C^1 smooth submanifold of topological dimension p and $x \in \Sigma$ is transversal, then, for every compact neighbourhood F of 0 , we have*

$$(1.3) \quad F \cap \delta_{1/r}(x^{-1}\Sigma) \rightarrow F \cap \Pi_\Sigma(x) \quad \text{as } r \rightarrow 0^+$$

where the convergence is in the sense of the Hausdorff distance between compact sets and $\Pi_\Sigma(x)$ is a p -dimensional normal homogeneous subgroup of \mathbb{G} , having Hausdorff dimension equal to $D(p)$. Moreover, the following limit holds

$$(1.4) \quad \lim_{r \rightarrow 0^+} \frac{\tilde{\mu}(\Sigma \cap B(x, r))}{r^{D(p)}} = \frac{\theta_g^d((\tau_\Sigma(x))_{D(p),x})}{\|(\tau_{\Sigma,\tilde{g}}(x))_{D(p),x}\|}.$$

The proof of this theorem is given in Section 3. This section, along with Section 2, also contains the definitions of the relevant notions. It is worth to mention that in the case of $C^{1,1}$ regularity the blow-up at transversal points is already contained in [23]. In our case, where Σ is only C^1 , the approach of [23] does not apply, so we follow the method used in [18] for curves. The point here is to provide a special “weighted reparametrization” of Σ around the blow-up point, see (3.8). Our next main result is the following generalized negligibility theorem.

Theorem 1.2 (Negligibility theorem). *Let $\Sigma \subset \mathbb{G}$ be a p -dimensional C^1 submanifold and let $\Sigma_c \subset \Sigma$ denote the subset of points with degree less than $D(p)$. We have*

$$(1.5) \quad \mathcal{H}^{D(p)}(\Sigma_c) = 0.$$

We refer to Section 4 for the definition of the *generalized characteristic set* Σ_c . The proof of Theorem 1.2 relies on covering arguments and a number of technical lemmata, that aim to estimate the behaviour of the number of small balls covering the generalized characteristic set. The difficulty here is to properly translate the information on the lower degree of the points into concrete estimates on the best “local coverings” around these points, see Lemma 2.3, Lemma 4.1 and Lemma 4.2. Theorem 1.2 extends to Euclidean Lipschitz submanifolds using standard arguments (see Theorem 4.5); more precisely, for any p -dimensional Lipschitz regular submanifold $\Sigma \subset \mathbb{G}$ there holds

$$\mathcal{H}^{D(p)}(\Sigma_* \cup \Sigma_c) = 0,$$

where Σ_* is the set of points where the (Euclidean) tangent cone to Σ is not a p -dimensional plane, see Subsection 2.4. Arguing the same way, one also realizes that estimates (1.2) extend to all Euclidean Lipschitz transversal submanifolds.

The method to prove Theorem 1.2 can also be used to establish new estimates on the Carnot-Carathéodory Hausdorff dimension of Σ_c for $C^{1,\lambda}$ submanifolds in Carnot groups, where $0 < \lambda \leq 1$. These estimates show that the Carnot-Carathéodory Hausdorff dimension of Σ_c can be estimated from above by a bound smaller than $D(p)$. More precisely, we have

Theorem 1.3. *Let Σ be a p -dimensional submanifold of \mathbb{G} of class $C^{1,\lambda}$, $\lambda \in (0, 1]$. Then*

$$(1.6) \quad \begin{cases} \dim_H \Sigma_c \leq D(p) - \lambda & \text{if } \ell = \ell(p) = 1 \\ \dim_H \Sigma_c \leq D(p) - 1 & \text{if } \ell \geq 2 \text{ and } \lambda \geq \frac{1}{\ell-1} \\ \dim_H \Sigma_c \leq D(p) - \frac{\ell\lambda}{1+\lambda} & \text{if } \ell \geq 2 \text{ and } \lambda \leq \frac{1}{\ell-1} \end{cases} .$$

Both Theorems 1.2 and 1.3 generalize some results proved, in the Heisenberg group framework, in the fundamental paper [2], compare Remark 5.3. We do not know whether the estimates of Theorem 1.3 are sharp. Even in Heisenberg groups, this sharpness seems to be an interesting open question. We refer to [3] for results and open problems akin to that of estimating the size of Σ_c .

The validity of (1.2) for a large class of submanifolds makes the intrinsic measure (1.1) a reasonable notion of “sub-Riemannian mass”. This should be seen for instance in the perspective of studying special classes of isoperimetric inequalities, when either the filling current or the filling submanifold must be necessarily transversal, as it occurs for higher dimensional fillings in Heisenberg groups.

2. NOTATION AND PRELIMINARY RESULTS

2.1. Carnot groups and exponential coordinates. Let us start with a brief introduction to stratified groups; we refer to [15] for more details on the subject.

Let \mathbb{G} be a connected and simply connected Lie group with stratified Lie algebra $\mathcal{G} = V_1 \oplus \cdots \oplus V_\iota$ of step ι , satisfying the conditions $V_{i+1} = [V_1, V_i]$ for every $i \geq 1$ and $V_{\iota+1} = \{0\}$. We set

$$(2.1) \quad n_j := \dim V_j \quad \text{and} \quad m_j := n_1 + \cdots + n_j, \quad j = 1, \dots, \iota;$$

we will also use $m_0 := 0$. The degree d_j of $j \in \{1, \dots, n\}$ is defined by the condition

$$m_{d_j-1} + 1 \leq j \leq m_{d_j}.$$

We denote by n the dimension of \mathcal{G} , therefore $n = m_\iota$. We say that a basis (X_1, \dots, X_n) of \mathcal{G} is *adapted to the stratification*, or in short *adapted*, if

$$X_{m_{j-1}+1}, \dots, X_{m_j} \text{ is a basis of } V_j \text{ for any } j = 1, \dots, \iota.$$

In the sequel, we will fix a *graded metric* g on \mathbb{G} , namely, a left invariant Riemannian metric on \mathbb{G} such that the subspaces V_k are orthogonal.

Definition 2.1. An adapted basis (X_1, \dots, X_n) of \mathcal{G} that is also orthonormal with respect to a left invariant Riemannian metric is a *graded basis*.

Clearly, the Riemannian metric in the previous definition must be necessarily graded. When either an adapted or a graded basis is understood, we identify \mathbb{G} with \mathbb{R}^n by the corresponding exponential coordinates of the first kind.

We use two different ways of denoting points x of \mathbb{G} with respect to fixed exponential coordinates of the first kind adapted to a graded basis of \mathcal{G} . We use both the standard notation with “lower indices”

$$x = (x_1, \dots, x_n) \in \mathbb{R}^n$$

and the one with “upper indices”

$$x = (x^1, x^2, \dots, x^t) \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_t},$$

where clearly $x^j = (x_{m_{j-1}+1}, \dots, x_{m_j}) \in \mathbb{R}^{n_j}$ for all $j = 1, \dots, t$. By the Baker-Campbell-Hausdorff formula, the group law reads in coordinates as

$$(2.2) \quad x \cdot y = x + y + Q(x, y),$$

for a suitable polynomial function $Q : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$. Precisely, $Q = (Q_1, \dots, Q_n)$ can be written in the form

$$(2.3) \quad Q_j(x, y) = \sum_{k,h:d_k < d_j, d_h < d_j} R_j^{kh}(x, y)(x_k y_h - x_h y_k) \quad \forall j = 1, \dots, n$$

for suitable polynomial functions R_j^{kl} . It follows that for any bounded set $K \subset \mathbb{G}$ there exists $C = C(K) > 0$ such that

$$(2.4) \quad |Q(x, y)| \leq C|x||y| \quad \text{for any } x, y \in K.$$

2.1.1. Stratified groups as abstract vector spaces. To emphasize some intrinsic notions on stratified groups, while preserving the ease of using a linear structure, stratified groups can be also regarded as abstract vector spaces. In fact, connected and simply connected nilpotent Lie groups are diffeomorphic to their Lie algebra through the exponential mapping $\exp : \mathcal{G} \rightarrow \mathbb{G}$, that is an analytic diffeomorphism. As a result, we equip the Lie algebra of \mathbb{G} with a Lie group operation, given by the Baker-Campbell-Hausdorff series, that makes this Lie group isomorphic to the original \mathbb{G} . This allows us to consider \mathbb{G} as an abstract linear space, equipped with a polynomial operation and a grading $\mathbb{G} = H_1 \oplus \dots \oplus H_t$. Under this identification a graded basis (X_1, \dots, X_n) becomes an orthonormal basis of \mathbb{G} as a vector space, where $(X_{m_{j-1}+1}, \dots, X_{m_j})$ is an orthonormal basis of H_j for all $j = 1, \dots, t$.

Remark 2.2. When a stratified group \mathbb{G} is seen as an abstract vector space, equipped with a graded basis X_1, \dots, X_n , then the associated graded metric g makes this basis orthonormal. As a result, the metric g becomes the Euclidean metric with respect to the corresponding coordinates (x^1, x^2, \dots, x^t) .

2.1.2. Dilations. For every $r > 0$, a natural group automorphism $\delta_r : \mathcal{G} \rightarrow \mathcal{G}$ can be defined as the unique algebra homomorphism such that

$$\delta_r(X) := rX \quad \text{for every } X \in V_1.$$

This one parameter group of linear isomorphisms constitutes the family of the so-called *dilations* of \mathcal{G} . They canonically yield a one parameter group of dilations on \mathbb{G} and can be denoted by the same symbol. With respect to our coordinates, we have

$$\delta_r(x_1, \dots, x_j, \dots, x_n) = (rx_1, \dots, r^{d_j}x_j, \dots, r^t x_n).$$

2.1.3. Left translations. For each element $x \in \mathbb{G}$, the group operation of \mathbb{G} automatically defines the corresponding *left translation* $l_x : \mathbb{G} \rightarrow \mathbb{G}$, with $l_x(z) = xz$ for all $z \in \mathbb{G}$. Right translations r_x are defined in analogous way.

2.2. Metric facts. We will say that d is a *homogeneous distance* on \mathbb{G} if it is a continuous distance on \mathbb{G} satisfying the following conditions

$$(2.5) \quad d(zx, zy) = d(x, y) \quad \text{and} \quad d(\delta_r(x), \delta_r(y)) = rd(x, y) \quad \forall x, y, z \in \mathbb{G}, r > 0.$$

Important examples of homogeneous distances are the well known Carnot-Carathéodory distance and those constructed in [14]. It is easily seen that two homogeneous distances are always equivalent. We will denote by $B(x, r)$ and $B_E(x, r)$, respectively, the open balls of center x and radius r with respect to a (fixed) homogeneous distance d and the Euclidean distance on $\mathbb{R}^n \equiv \mathbb{G}$.

For $r > 0$ we introduce the *boxes*

$$\begin{aligned} \text{Box}(0, r) &:= \{y \in \mathbb{R}^n : |y^j| < r^j \forall j = 1, \dots, \iota\} \\ &= (-r, r)^{n_1} \times (-r^2, r^2)^{n_2} \times \dots \times (-r^\iota, r^\iota)^{n_\iota} \\ \text{Box}(x, r) &:= x \cdot \text{Box}(0, r), \quad x \in \mathbb{G}. \end{aligned}$$

By homogeneity, it is easy to observe for any homogeneous distance d there exists $C_{BB} = C_{BB}(d) \geq 1$ such that

$$(2.6) \quad \text{Box}(x, r/C_{BB}) \subset B(x, r) \subset \text{Box}(x, C_{BB}r).$$

We will also use the notation

$$\text{Box}_E^\mu(0, r) := \{y \in \mathbb{R}^\mu : |y_j| < r \forall j = 1, \dots, \mu\} = (-r, r)^\mu.$$

When given $0 < s < r$ and a linear subspace W of \mathbb{R}^μ we pose

$$\text{Box}_W^\mu(0; r, s) := \{y \in \text{Box}_E^\mu(0, r) : |\pi_W(y)| < s\},$$

where $\pi_W(y)$ is the canonical projection of y on W . If w_1, \dots, w_H is an orthonormal basis of W we clearly have

$$(2.7) \quad \{y \in \text{Box}_E^\mu(0, r) : |\langle y, w_i \rangle| < \frac{s}{\sqrt{H}} \forall i = 1, \dots, H\} \subset \text{Box}_W^\mu(0; r, s).$$

From now on, a homogeneous distance d is fixed. We will use several times the following simple fact.

Lemma 2.3. *There exists $C = C(d) > 0$ with the following property. For any fixed $r \leq 1$, $x \in B_E(0, r)$ and $j \in \{1, \dots, \iota\}$ there exists $\tilde{x} \in \mathbb{G}$ such that*

$$(2.8) \quad \tilde{x}^1 = \dots = \tilde{x}^j = 0, \quad d(x, \tilde{x}) \leq Cr^{1/j}$$

$$(2.9) \quad |\tilde{x}^h - x^h| \leq Cr^2 \quad \text{for any } h = j+1, \dots, \iota.$$

Proof. In the case $j = 1$, we define

$$\begin{aligned} \tilde{x} &= x \cdot (-x^1, 0, \dots, 0) = (x^1, \dots, x^\iota) \cdot (-x^1, 0, \dots, 0) \\ &= (0, x^2 + O(r^2), \dots, x^\iota + O(r^2)), \end{aligned}$$

where the last equality follows from (2.4) and $O(\cdot)$ is understood with respect to the Euclidean norm. By (2.6) we have

$$d(x, \tilde{x}) = d(0, (-x^1, 0, \dots, 0)) \leq C_{BB}r$$

whence (2.8) follows.

We now argue by induction on $j \geq 2$, assuming the existence of some \bar{x} such that

$$\bar{x}^1 = \dots = \bar{x}^{j-1} = 0, \quad d(x, \bar{x}) \leq Cr^{1/(j-1)} \quad \text{and} \quad |\bar{x}^h - x^h| \leq Cr^2$$

for any $h = j, \dots, \iota$. Defining $\tilde{x} = \bar{x} \cdot (0, \dots, 0, -\bar{x}^j, 0, \dots, 0)$, applying both (2.3) and (2.4) we obtain

$$\begin{aligned} \tilde{x} &= (\bar{x}^1, \dots, \bar{x}^\iota) \cdot (0, \dots, 0, -\bar{x}^j, 0, \dots, 0) \\ &= (0, \dots, 0, 0, \bar{x}^{j+1} + O(r^2), \dots, \bar{x}^\iota + O(r^2)). \end{aligned}$$

Thus, by inductive hypothesis we get $\tilde{x} = (0, \dots, 0, 0, x^{j+1} + O(r^2), \dots, x^\iota + O(r^2))$. As a result, we arrive at the following inequalities

$$\begin{aligned} d(x, \tilde{x}) &\leq d(x, \bar{x}) + d(\bar{x}, \tilde{x}) \leq Cr^{1/(j-1)} + d(0, (0, \dots, 0, -\bar{x}^j, 0, \dots, 0)) \\ &\leq Cr^{1/j} + C_{BB}|\bar{x}^j|^{1/j} = Cr^{1/j} + C_{BB}|x^j + O(r^2)|^{1/j} \leq \tilde{C}r^{1/j}. \end{aligned}$$

that complete the proof. \square

2.2.1. Hausdorff measures and coverings. For the sake of completeness, we recall the definitions of Hausdorff measures. Let $q \geq 0$ and $\delta > 0$ be fixed; we define

$$\begin{aligned} \mathcal{H}_\delta^q(E) &:= \inf \left\{ \sum_{i=1}^{\infty} (\text{diam } E_i)^q : E \subset \cup_i E_i, \text{diam } E_i < \delta \right\} \\ \mathcal{S}_\delta^q(E) &:= \inf \left\{ \sum_{i=1}^{\infty} (\text{diam } B_i)^q : E \subset \cup_i B_i, B_i = B(x_i, r_i) \text{ balls, diam } B_i < \delta \right\}. \end{aligned}$$

The q -dimensional *Hausdorff measure* of $E \subset \mathbb{G}$ is

$$\mathcal{H}^q(E) := \lim_{\delta \rightarrow 0^+} \mathcal{H}_\delta^q(E)$$

while the q -dimensional *spherical Hausdorff measure* of E is

$$\mathcal{S}^q(E) := \lim_{\delta \rightarrow 0^+} \mathcal{S}_\delta^q(E).$$

The *Hausdorff dimension* of E is

$$\dim_H E := \inf\{q : \mathcal{H}^q(E) = 0\} = \sup\{q : \mathcal{H}^q(E) = +\infty\}.$$

It is well-known that $\dim_H \mathbb{G}$ coincides with the *homogeneous dimension* $Q := n_1 + 2n_2 + \dots + \iota n_\iota$ of \mathbb{G} . The standard Euclidean Hausdorff measure on $\mathbb{G} = \mathbb{R}^n$ is denoted by $\mathcal{H}_{|\cdot|}^q$. For more information on the properties of these measures, see for instance [11, 24, 27].

We state without proof the following simple fact.

Proposition 2.4. *Let $\theta > 0$ and let $E \subset X$, where X is a metric space. If for all $\epsilon \in (0, 1)$ the set E can be covered by N_ϵ balls of radius ϵ^β with $N_\epsilon \leq C \epsilon^{-q}$ and $C > 0$ independent from ϵ , then the Hausdorff dimension of E is not greater than q/β .*

The following result, see e.g. [27, Theorem 3.3], will be useful in the sequel.

Theorem 2.5 (*5r-covering*). *Let (X, d) be a separable metric space and $E \subset X$; let $r > 0$ be fixed. Then there exists a subset $F \subset E$ at most countable such that*

$$E \subset \bigcup_{x \in F} B(x, 5r) \quad \text{and} \quad B(x, r) \cap B(x', r) = \emptyset \quad \forall x, x' \in F, x \neq x'.$$

2.3. Multi-indices, degrees and maximal dimension. We denote by I_p the set of those multi-indices $\alpha = (\alpha_1, \dots, \alpha_p) \in \{1, \dots, n\}^p$ such that $1 \leq \alpha_1 < \dots < \alpha_p \leq n$. We also set

$$d(\alpha) := d_{\alpha_1} + \dots + d_{\alpha_p}.$$

We denote by $D(p)$ the maximum integer $d(\alpha)$ when α varies in I_p . We call this number the *maximal dimension*, that is uniquely defined for any given $p \in \{1, 2, \dots, n\}$. Clearly, $D(n)$ equals the homogeneous dimension Q of \mathbb{G} . The maximal dimension can be computed in the following way. Define $\ell = \ell(p)$ by imposing

$$(2.10) \quad \begin{cases} \ell := \iota & \text{if } p \leq n_\iota \\ \sum_{j=\ell+1}^{\iota} n_j < p \leq \sum_{j=\ell}^{\iota} n_j & \text{otherwise.} \end{cases}$$

Clearly, ℓ depends on p and it can be equivalently defined by $\ell(p) := d_{n_{+1-p}}$, that represents the lowest possible degree among tangent vectors of $\text{span}\{X_{n-p+1}, \dots, X_n\}$, where (X_1, \dots, X_n) is an adapted basis of the stratified Lie algebra \mathcal{G} . It is also easy to see that

$$(2.11) \quad D(p) = \sum_{j=\ell(p)+1}^{\iota} j n_j + \ell(p) \left(p - \sum_{j=\ell(p)+1}^{\iota} n_j \right),$$

where the two summations in (2.11) have to be understood as 0 when $\ell(p) = \iota$. We also set

$$(2.12) \quad r_p := p - \sum_{j=\ell(p)+1}^{\iota} n_j \geq 1,$$

so that

$$(2.13) \quad p = r_p + n_{\ell(p)+1} + \dots + n_\iota \quad \text{and} \quad D(p) = \ell(p)r_p + (\ell(p) + 1)n_{\ell(p)+1} + \dots + \iota n_\iota.$$

It is worth noticing that $D(p) = \beta_+(p)$, where β_+ is the *upper dimension comparison function* for \mathbb{G} , introduced in [4].

2.4. Degree of submanifolds, projections and subdilations. By $\Sigma \subset \mathbb{G}$, we denote a p -dimensional Lipschitz submanifold of \mathbb{G} . We define the *singular set*

$$\Sigma_* := \{x \in \Sigma : C_x \Sigma \text{ is not a } p\text{-dimensional subspace of } \mathbb{R}^n\},$$

where $C_x \Sigma$ is the (Euclidean) tangent cone to Σ at x , i.e.,

$$C_x \Sigma := \left\{ tv \in \mathbb{R}^n : t \geq 0, v = \lim_{i \rightarrow \infty} \frac{x_i - x}{|x_i - x|} \text{ for some sequence } (x_i)_{i \in \mathbb{N}} \subset \Sigma \text{ with } x_i \rightarrow x \right\}.$$

We have the following fact.

Proposition 2.6. *For any p -dimensional Lipschitz submanifold $\Sigma \subset \mathbb{G}$, we have*

$$(2.14) \quad \mathcal{H}^{D(p)}(\Sigma_*) = 0.$$

Proof. Since $\mathcal{H}_{|\cdot|}^p(\Sigma_*) = 0$, by [4, Proposition 3.1] our claim immediately follows. \square

Given a point $x \in \Sigma \setminus \Sigma_*$, we denote by $\tau_\Sigma(x)$ its tangent vector, i.e., the p -dimensional multivector associated with the p -plane $C_x\Sigma$. We can write

$$\tau_\Sigma(x) = \sum_{\alpha \in I_p} c_\alpha X_\alpha,$$

where $X_\alpha := X_{\alpha_1} \wedge \dots \wedge X_{\alpha_p}$. We then define the *degree* $d_\Sigma(x)$ of Σ at x as

$$d_\Sigma(x) := \max\{d(\alpha) : \alpha \in I_p \text{ and } c_\alpha \neq 0\}.$$

The *degree* of Σ is $d(\Sigma) := \max\{d_\Sigma(x) : x \in \Sigma \setminus \Sigma_*\}$. Clearly, we have $d(\Sigma) \leq D(p)$. The underlying metric g on \mathbb{G} gives a natural scalar product on multivectors, whose norm will be denoted by $\|\cdot\|$. If \tilde{g} is any Riemannian metric on \mathbb{G} , then at any $x \in \mathbb{G}$ we have a canonically defined scalar product on any $\Lambda_p S_x$ where $S_x \subset T_x\mathbb{G}$ is a p -dimensional subspace. We will denote by $\|\cdot\|_{\tilde{g},x}$ its corresponding norm.

Definition 2.7. Let \tilde{g} be any Riemannian metric on \mathbb{G} , let Σ be a p -dimensional Lipschitz submanifold and let $x \in \Sigma \setminus \Sigma_*$. We define the *unit tangent p -vector* with respect to \tilde{g} as follows

$$(2.15) \quad \tau_{\Sigma,\tilde{g}}(x) = \frac{\tau_\Sigma(x)}{\|\tau_\Sigma(x)\|_{\tilde{g},x}},$$

where $\tau_\Sigma(x)$ is any tangent p -vector of Σ at x .

Dilations of \mathbb{G} canonically extend to dilations on multivectors as follows

$$(\Lambda_p \delta_r)(v_1 \wedge \dots \wedge v_p) = (\delta_r v_1) \wedge \dots \wedge (\delta_r v_p)$$

for all $v_1, \dots, v_p \in \mathbb{G}$, therefore we have

$$(\Lambda_p \delta_r)(X_\alpha) = (\Lambda_p \delta_r)(X_{\alpha_1} \wedge \dots \wedge X_{\alpha_p}) = r^{d_{\alpha_1} + \dots + d_{\alpha_p}} X_\alpha = r^{d(\alpha)} X_\alpha.$$

Definition 2.8. A p -vector $v \in \Lambda_p \mathbb{G}$ is *homogeneous* of degree $l \in \mathbb{N} \setminus \{0\}$ if $(\Lambda_p \delta_r)v = r^l v$ for all $r > 0$.

Our scalar product on $\Lambda_p(\mathbb{G})$ allows us to introduce the following canonical projections, hence homogeneous multivectors of different degrees are orthogonal.

Definition 2.9. Let $p, D \in \mathbb{N}$ be such that $1 \leq p \leq D \leq D(p)$. Let us introduce the linear subspace $\Lambda_{D,p}(\mathbb{G})$ of $\Lambda_p \mathbb{G}$ made by all homogeneous p -multivectors of degree D . With respect to the scalar product of \mathbb{G} , the following orthogonal projection

$$\pi_D : \Lambda_p(\mathbb{G}) \rightarrow \Lambda_{D,p}(\mathbb{G})$$

is uniquely defined. We say that π_D is the *projection of degree D* . If we consider a p -vector $t \in \Lambda_p S_x$ with $S_x \subset T_x\mathbb{G}$, a pointwise projection $\pi_{D,x}(t)$ is automatically

defined, taking the left translated multivector $(\Lambda_p dl_{x^{-1}})(t) \in \Lambda_p(T_0\mathbb{G})$, identifying $T_0\mathbb{G}$ with \mathbb{G} and applying π_D to this translated multivector, hence

$$\pi_{D,x}(t) = (\Lambda_p dl_x) \circ \pi_D \circ (\Lambda_p dl_{x^{-1}})(t).$$

To simplify notation, both projection π_D applied to $v \in \Lambda_p\mathbb{G}$ and $\pi_{D,x}$ applied to $w \in \Lambda_p S_x$ will be also denoted by $(v)_D$ and $(w)_{D,x}$, respectively.

Remark 2.10. The previous notions allow us to consider the “density function” with respect to \tilde{g} , defined as $\Sigma \ni x \rightarrow \|(\tau_{\Sigma,\tilde{g}}(x))_D\|$. This function naturally appears in the representation of the D -dimensional “intrinsic measure” of a submanifold.

The following lemma will be useful in the sequel. It can be proved repeating exactly the same arguments of [23, Lemma 3.1].

Lemma 2.11. *Let $\Sigma \subset \mathbb{G}$ be a p -dimensional Lipschitz submanifold and fix $x \in \Sigma \setminus \Sigma_*$. Then we can find a graded basis X_1, \dots, X_n of \mathcal{G} and a basis v_1, \dots, v_p of $T_x\Sigma$ such that, writing $v_j = \sum_{i=1}^n C_{ij}X_i(x)$, we have*

$$(2.16) \quad C := (C_{ij})_{\substack{i=1,\dots,n \\ j=1,\dots,p}} = \left[\begin{array}{c|ccc} Id_{\alpha_1} & 0 & \cdots & 0 \\ 0 & * & \cdots & * \\ \hline 0 & Id_{\alpha_2} & \cdots & 0 \\ 0 & 0 & \cdots & * \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline 0 & 0 & \cdots & Id_{\alpha_\iota} \\ 0 & 0 & \cdots & 0 \end{array} \right]$$

where α_k are integers satisfying $0 \leq \alpha_k \leq n_k$ and $\alpha_1 + \dots + \alpha_\iota = p$. The symbols 0 and * denote null and arbitrary matrices of the proper size, respectively. We have

$$(2.17) \quad d_\Sigma(x) = \sum_{k=1}^{\iota} k\alpha_k.$$

Remark 2.12. As already observed in [23, Remark 3.2], the previous lemma along with its proof are understood to hold also in the case where some α_k possibly vanishes. In this case the α_k columns of (2.16) containing Id_{α_k} and the corresponding vectors v_j^k are meant to be absent. In a similar way, if $\alpha_k = n_k$, then the null sub-matrix is missing from the corresponding diagonal box.

The integers $\alpha_1, \dots, \alpha_\iota$ of Lemma 2.11 define a “sub-grading” for a p -dimensional subspace of \mathbb{R}^n , so that in analogy with the integers m_j defined in (2.1), we set

$$(2.18) \quad \mu_0 = 0 \quad \text{and} \quad \mu_k = \sum_{l=1}^k \alpha_l \quad \text{for all} \quad k = 1, \dots, \iota.$$

This new grading allows us to define for every $j \in \{1, \dots, p\}$ the *subdegree* σ_j defined as follows

$$(2.19) \quad \sigma_j := k \quad \text{if and only if} \quad \mu_{k-1} < j \leq \mu_k \quad \text{for some} \quad k \in \{1, \dots, \iota\}.$$

The corresponding *subdilations* $\lambda_r : \mathbb{R}^p \rightarrow \mathbb{R}^p$ are defined as follows

$$\lambda_r(\xi_1, \dots, \xi_p) = (r^{\sigma_1}\xi_1, r^{\sigma_2}\xi_2, \dots, r^{\sigma_p}\xi_p) \quad \text{for all } r > 0.$$

2.5. Transversal points and transversal submanifolds. Let us fix a p -dimensional Lipschitz submanifold Σ and consider $x \in \Sigma \setminus \Sigma_*$, where Σ_* is its singular set. We say that x is *transversal* if $d_\Sigma(x)$ equals the maximal dimension $D(p)$. We say that Σ is a *transversal submanifold* if Σ contains at least one transversal point; clearly, this is equivalent to the condition $d(\Sigma) = D(p)$.

Remark 2.13. For hypersurfaces, transversal points coincide with noncharacteristic points and when $p \geq n - n_1$ a p -dimensional submanifold of \mathbb{G} is transversal if and only if it is *non-horizontal*, according to the terminology of [21].

The following corollary is an easy consequence of the fact that $X_i(0) = e_i$.

Corollary 2.14. *Under the assumptions and notations of Lemma 2.11, the point x is transversal if and only if the following conditions hold*

$$(2.20) \quad \alpha_\ell = n_\ell, \alpha_{\ell-1} = n_{\ell-1}, \dots, \alpha_{\ell+1} = n_{\ell+1}, \alpha_\ell = r_p, \alpha_{\ell-1} = 0, \dots, \alpha_1 = 0,$$

where $\ell = \ell(p)$ is defined by (2.10) and r_p is defined in (2.12). If $x = 0$ is transversal, then the vectors v_1, \dots, v_p in Lemma 2.11 constitute the columns of the matrix

$$(2.21) \quad C^0 = (C_{ij}^0) = \begin{bmatrix} * & \cdots & \cdots & \cdots & * \\ Id_{r_p} & 0 & \cdots & \cdots & 0 \\ 0 & * & \cdots & \cdots & * \\ 0 & Id_{n_{\ell+1}} & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \cdots & \vdots \\ \vdots & \ddots & \cdots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & Id_{n_\ell} \end{bmatrix}.$$

The previous corollary shows that at transversal points the associated grading given by the integers of (2.18) and (2.19) yields

$$\mu_0 = \cdots = \mu_{\ell-1} = 0, \quad \mu_\ell = r_p \quad \text{and} \quad \mu_{\ell+j} = r_p + \sum_{i=1}^j n_{\ell+i} \quad \text{for all } j = 1, \dots, \ell - \ell,$$

therefore the subdegrees are the following ones

$$(2.22) \quad \sigma_1 = \ell, \dots, \sigma_{r_p} = \ell \quad \text{and} \quad \sigma_{r_p+s+\sum_{i=1}^j n_{\ell+i}} = \sigma_{\mu_{\ell+j}+s} = \ell + j + 1$$

for all $s = 1, \dots, n_{\ell+j+1}$ and $j = 0, 1, \dots, \ell - \ell - 1$, where the term $\sum_{i=1}^j n_{\ell+i}$ in the previous formulae is meant to be zero when $j = 0$.

3. BLOW-UP AT TRANSVERSAL POINTS

This section is devoted to the proof of Theorem 1.1 stated in the Introduction. We have first to recall some more notions and fix other auxiliary objects. First of all \tilde{g} will denote any auxiliary Riemannian metric on \mathbb{G} . The corresponding Riemannian surface measure induced on a C^1 smooth submanifold $\Sigma \subset \mathbb{G}$ will be denoted by $\tilde{\mu}$.

Definition 3.1. A graded metric g on \mathbb{G} is fixed and we set $\mathbb{B} = \{x \in \mathbb{G} : d(0, x) < 1\}$, where d is a homogeneous distance. The stratified group \mathbb{G} is seen as an abstract vector space and S denotes one of its p -dimensional linear subspaces. We consider any simple p -vector τ associated to S . Then we define the *metric factor*

$$(3.1) \quad \theta_g^d(\tau) = \mathcal{H}_{|\cdot|}^p(S \cap \mathbb{B}).$$

Here $|\cdot|$ denotes the Euclidean metric on \mathbb{G} with respect to a fixed graded basis (X_1, \dots, X_n) and the sets S and \mathbb{B} are represented with respect to the associated coordinates of the first kind.

Remark 3.2. In the previous definition, any other simple p -vector $\lambda\tau$ with $\lambda \neq 0$ defines the same subspace S . Conversely, whenever a simple p -vector ζ is associated to S , that is $\zeta = \zeta_1 \wedge \dots \wedge \zeta_p$ and $(\zeta_1, \dots, \zeta_p)$ is basis of S , then $\zeta = t\tau$ for some $t \neq 0$.

Remark 3.3. The metric factor only depends on the Riemannian metric g and the homogeneous distance d . In fact, under the assumptions of Definition 3.1, let us consider another graded basis (Y_1, \dots, Y_n) with associated coordinates (y_1, \dots, y_n) of the first kind. Then the linear change of variables from these coordinates to the original coordinates (x_1, \dots, x_n) associated to (X_1, \dots, X_n) is an isometry of \mathbb{R}^n , hence the number (3.1) is preserved under the coordinates (y_i) .

About the statement of Theorem 1.1, we wish to clarify that the Lie subgroup $\Pi_\Sigma(x)$ appearing in (1.3) is also homogeneous in the sense that is closed under dilations. Furthermore, it is a p -dimensional homogeneous subgroup of \mathbb{G} of the form

$$Z \oplus H_{\ell+1} \oplus \dots \oplus H_\ell,$$

where $Z \subset H_\ell$ is a linear space of dimension r_p . The integers ℓ and r_p are defined in (2.10) and (2.12). In particular, $\Pi_\Sigma(x)$ is also a normal subgroup. The same subgroup is more conveniently defined later in the proof of Theorem 1.1, see (3.25).

Proof of Theorem 1.1. First of all, our claim allows us to assume that there exists an open neighbourhood $U \subset \mathbb{R}^p$ of the origin such that $\Psi : U \rightarrow \Sigma$ is a C^1 smooth diffeomorphism with $\Psi(0) = x$. Defining the translated submanifold $\Sigma_x = l_{x^{-1}}(\Sigma)$, we observe that

$$d_{\Sigma_x}(0) = d_\Sigma(x) = D(p) = d(\Sigma) = d(\Sigma_x).$$

We consider the translated diffeomorphism $\phi = l_{x^{-1}} \circ \Psi$, with $\phi : U \rightarrow \Sigma_x$. Taking into account Corollary 2.14, we have a graded basis of left invariant vector fields X_1, \dots, X_n and linearly independent vectors $v_1, \dots, v_p \in T_0\Sigma_x$ such that the matrix $C^0 = (C_{ij}^0)$, defined by $v_j = \sum_{i=1}^n C_{ij}^0 X_i(0) = \sum_{i=1}^n C_{ij}^0 e_i$ is given by (2.21). The vector fields X_i in our coordinates have the form

$$(3.2) \quad X_i = \sum_{l=1}^n a_i^l e_l$$

and their (nonconstant) coefficients satisfy (see e.g. [15])

$$(3.3) \quad a_i^l = \begin{cases} \delta_i^l & d_l \leq d_i \\ \text{homogeneous polynomial of degree } d_l - d_i & \text{otherwise,} \end{cases}$$

where homogeneity refers to intrinsic dilations of the group. After a linear change of variable on ϕ , we can also assume that $\partial_{t_i}\phi(0) = v_i$ for all $i = 1, \dots, p$, where each v_i is the i -th column of (2.21). Let $\pi_0 : \mathbb{R}^n \rightarrow \mathbb{R}^p$ be the projection

$$\pi_0(x_1, \dots, x_n) = (x_{m_{\ell-1}+1}, \dots, x_{m_{\ell-1}+r_p}, x_{m_{\ell}+1}, \dots, x_n).$$

Thus, $d(\pi_0 \circ \phi)(0)$ is invertible and the inverse mapping theorem provides us with new variables $y = (y_{m_{\ell-1}+1}, \dots, y_{m_{\ell-1}+r_p}, y_{m_{\ell}+1}, \dots, y_n)$ such that Σ_x is represented near the origin by $\gamma = \phi \circ (\pi_0 \circ \phi)^{-1}$, which can be written as follows

$$(\gamma_1(y), \dots, \gamma_{m_{\ell-1}}(y), y_{m_{\ell-1}+1}, \dots, y_{m_{\ell-1}+r_p}, \gamma_{m_{\ell-1}+r_p+1}(y), \dots, \gamma_{m_{\ell}}(y), y_{m_{\ell}+1}, \dots, y_n)$$

and it is defined in some smaller neighbourhood $(-c_1, c_1)^p \subset U$ for some $c_1 > 0$. Furthermore, since $d(\pi_0 \circ \phi)(0)$ is the identity mapping of \mathbb{R}^p , we get

$$(3.4) \quad (\partial_1 \gamma)(0) = v_1, \quad (\partial_2 \gamma)(0) = v_2, \quad \dots \quad (\partial_p \gamma)(0) = v_p,$$

so that we have continuous functions $C_{ij}(y)$ with $C_{ij}(0) = C_{ij}^0$ such that

$$(3.5) \quad (\partial_j \gamma)(y) = \sum_{i=1}^n C_{ij}(y) X_i(\gamma(y)) \quad \text{for all } j = 1, \dots, p.$$

Due to the structure of C^0 given in (2.21), whenever $\sigma_j = \ell$, or equivalently when $j = 1, \dots, r_p$, we have

$$(3.6) \quad C_{ij}(y) = \delta_{i-m_{\ell-1}, j} + o(1) \quad \text{for any } i \text{ such that } d_i \geq \ell.$$

When $\ell < \sigma_j \leq \iota$, or equivalently in the case $j > r_p$ and $j = \mu_{\sigma_j-1} + 1, \dots, \mu_{\sigma_j}$, we get

$$(3.7) \quad C_{ij}(y) = \begin{cases} \delta_{i-m_{\sigma_j-1}, j-\mu_{\sigma_j-1}} + o(1) & \text{if } d_i = \sigma_j \text{ or } 1 \leq i - m_{\sigma_j-1} \leq n_{\sigma_j} \\ o(1) & \text{if } d_i > \sigma_j. \end{cases}$$

Let us introduce the C^1 smooth homeomorphism $\eta : \mathbb{R}^p \rightarrow \mathbb{R}^p$ as follows

$$(3.8) \quad \eta(t) = \left(\frac{|t_1|^{\sigma_1}}{\sigma_1} \operatorname{sgn}(t_1), \dots, \frac{|t_p|^{\sigma_p}}{\sigma_p} \operatorname{sgn}(t_p) \right),$$

where its inverse mapping is given by the formula

$$\zeta(\tau) = \left(\operatorname{sgn}(\tau_1) \sqrt[\sigma_1]{\sigma_1 |\tau_1|}, \dots, \operatorname{sgn}(\tau_p) \sqrt[\sigma_p]{\sigma_p |\tau_p|} \right)$$

and all σ_j satisfy (2.22). We consider the C^1 smooth reparametrization $\Gamma(t) = \gamma(\eta(t))$ with partial derivatives

$$(3.9) \quad \partial_{t_j} \Gamma(t) = |t_j|^{\sigma_j-1} (\partial_j \gamma)(\eta(t)) = |t_j|^{\sigma_j-1} \sum_{s,i=1}^n C_{ij}(\eta(t)) a_i^s(\Gamma(t)) e_s$$

for all $j = 1, \dots, p$, where we have used both (3.2) and (3.5). We first observe that

$$(3.10) \quad \Gamma_i(t) = o(|t|^{d_i}) \quad \text{for } 1 \leq d_i < \ell.$$

In fact, we have $\gamma(0) = 0$ and $\eta(t) = O(|t|^\ell)$, hence

$$(3.11) \quad \Gamma_i(t) = \gamma_i(\eta(t)) = O(|\eta(t)|) = \begin{cases} o(|t|^{d_i}) & \text{if } d_i < \ell \\ O(|t|^\ell) & \text{if } d_i = \ell. \end{cases}$$

The main point is to prove the following rates of convergence

$$(3.12) \quad \begin{cases} \Gamma_i(t) = O(|t|^\ell) & \text{for } m_{\ell-1} < i \leq m_{\ell-1} + r_p \\ \Gamma_i(t) = o(|t|^\ell) & \text{for } m_{\ell-1} + r_p < i \leq m_\ell. \end{cases}$$

Since the first equation of (3.12) is already contained in (3.11), we have nothing to prove in the case $r_p = n_\ell$ (because (3.12) does not have the second case). Thus, we will assume that $r_p < n_\ell$ and then prove the second formula of (3.12). First of all, we apply (3.9) and compute the following partial derivatives

$$(3.13) \quad \partial_{t_j} \Gamma_i(t) = |t_j|^{\sigma_j-1} (\partial_j \gamma_i)(\eta(t)) = |t_j|^{\sigma_j-1} \sum_{k=1}^n C_{kj}(\eta(t)) a_k^i(\Gamma(t))$$

for all $j = 1, \dots, p$ and all $i = 1, \dots, n$. If $1 \leq j \leq r_p$, we rewrite the previous sum as

$$|t_j|^{\sigma_j-1} \left(\sum_{k:d_k < \ell} C_{kj}(\eta(t)) a_k^i(\Gamma(t)) + \sum_{k:d_k \geq \ell} C_{kj}(\eta(t)) a_k^i(\Gamma(t)) \right).$$

As a consequence, taking into account (3.6), it follows that

$$\begin{aligned} \partial_{t_j} \Gamma_i(t) &= |t_j|^{\ell-1} \left(a_{j+m_{\ell-1}}^i(\Gamma(t)) + \sum_{k:d_k < \ell} C_{kj}(\eta(t)) a_k^i(\Gamma(t)) + \sum_{k:d_k \geq \ell} o(1) a_k^i(\Gamma(t)) \right) \\ &= |t_j|^{\ell-1} \left(a_{j+m_{\ell-1}}^i(\Gamma(t)) + o(1) + \sum_{k:d_k < \ell} C_{kj}(\eta(t)) a_k^i(\Gamma(t)) \right). \end{aligned}$$

Since $d_k < \ell$, we have that a_k^i is a nonconstant homogenous polynomial. It follows that $a_k^i \circ \Gamma = o(1)$ and we get

$$(3.14) \quad \partial_{t_j} \Gamma_i(t) = |t_j|^{\ell-1} \left(a_{j+m_{\ell-1}}^i(\Gamma(t)) + o(1) \right)$$

Since $m_{\ell-1} + j \leq m_{\ell-1} + r_p < i \leq m_\ell$ and $d_{m_{\ell-1}+j} = d_i$, formula (3.3) implies that $a_{j+m_{\ell-1}}^i$ is the null polynomial. It follows that

$$(3.15) \quad \partial_{t_j} \Gamma_i(t) = o(|t|^{\ell-1}) \quad \text{whenever } 1 \leq j \leq r_p \text{ and } m_{\ell-1} + r_p < i \leq m_\ell.$$

If $r_p < j \leq p$, then (3.13) implies that

$$\partial_{t_j} \Gamma_i(t) = |t_j|^{\sigma_j-1} \sum_{k=1}^n C_{kj}(\eta(t)) a_k^i(\Gamma(t)) = |t_j|^{\sigma_j-1} O(1) = O(|t|^{\sigma_j-1}).$$

Since in this case $\sigma_j > \ell$, we get in particular that

$$(3.16) \quad \partial_{t_j} \Gamma_i(t) = o(|t|^{\ell-1}) \quad \text{whenever } r_p < j \leq p \text{ and } 1 \leq i \leq n.$$

Joining (3.15) with (3.16), it follows that

$$\nabla \Gamma_i(t) = o(|t|^{\ell-1}) \quad \text{for all } i = m_{\ell-1} + r_p + 1, \dots, m_\ell,$$

that proves the second equation of (3.12).

Now, we write explicitly the form of Γ as the composition $\gamma \circ \eta$. By the previous formulae for γ and η , we get

$$(3.17) \quad \Gamma(t) = \left(\Gamma_1(t), \dots, \Gamma_{m_{\ell-1}}(t), \frac{|t_1|^\ell}{\ell} \operatorname{sgn}(t_1), \dots, \frac{|t_{r_p}|^\ell}{\ell} \operatorname{sgn}(t_{r_p}), \Gamma_{m_{\ell-1}+r_p+1}(t), \dots \right. \\ \left. \dots, \Gamma_{m_\ell}(t), \frac{|t_{r_p+1}|^{\ell+1}}{\ell+1} \operatorname{sgn}(t_{r_p+1}), \dots, \frac{|t_p|^\iota}{\iota} \operatorname{sgn}(t_p) \right).$$

The new parametrization γ of Σ_x around the origin yields $\Phi : (-c_1, c_1)^p \rightarrow \Sigma$ defined as $\Phi := l_x \circ \gamma$, that is our ‘‘adapted parametrization’’ of Σ around x . Taking $r > 0$ sufficiently small, we have

$$(3.18) \quad \frac{\tilde{\mu}(\Sigma \cap B(x, r))}{r^{D(p)}} = r^{-D(p)} \int_{\Phi^{-1}(B(x, r))} \|(\partial_{y_1} \Phi \wedge \dots \wedge \partial_{y_p} \Phi)(y)\|_{\tilde{g}} dy.$$

where $\tilde{\mu}$ is the Riemannian surface measure induced by \tilde{g} on Σ . We perform the change of variable $y = \lambda_r t$, where λ_r is the subdilation of the form

$$(3.19) \quad \lambda_r(t_1, \dots, t_p) = (r^\ell t_1, \dots, r^\ell t_{r_p}, r^{\ell+1} t_{r_p+1}, \dots, r^{\ell+1} t_{r_p+n_{\ell+1}}, \dots, r^\iota t_p)$$

that yields the formula

$$(3.20) \quad \frac{\tilde{\mu}(\Sigma \cap B(x, r))}{r^{D(p)}} = \int_{\lambda_{1/r}(\Phi^{-1}(B(x, r)))} \|\partial_{y_1} \Phi(\lambda_r t) \wedge \dots \wedge \partial_{y_p} \Phi(\lambda_r t)\|_{\tilde{g}} dt.$$

The point is then to study the ‘‘behaviour’’ of the set $\lambda_{1/r}(\Phi^{-1}(B(x, r)))$ as $r \rightarrow 0^+$. To do this, we will use the formula (3.17) for Γ and the rates of convergence (3.12), taking into account the change of variables (3.8). Since $\Phi^{-1}(B(x, r)) = \gamma^{-1}(B(0, r))$, it follows that

$$(3.21) \quad \lambda_{1/r}(\Phi^{-1}(B(x, r))) = \{t \in \mathbb{R}^p : \delta_{1/r}(\gamma(\lambda_r t)) \in \mathbb{B}\},$$

where $\mathbb{B} = \{z \in \mathbb{G} : d(z, 0) < 1\}$. We observe that

$$\gamma(\lambda_r t) = \Gamma(\zeta(\lambda_r t)) = \Gamma(r \zeta(t)),$$

therefore the previous rescaled set can be written as follows

$$(3.22) \quad \lambda_{1/r}(\Phi^{-1}(B(x, r))) = \{t \in \mathbb{R}^p : \delta_{1/r}(\Gamma(r \zeta(t))) \in \mathbb{B}\}.$$

By (3.17), an element $t \in \mathbb{R}^p$ of the previous set is characterized by the property that

$$(3.23) \quad \left(\frac{\Gamma_1(r \zeta(t))}{r}, \dots, \frac{\Gamma_{m_{\ell-1}}(r \zeta(t))}{r^{\ell-1}}, t_1, \dots, t_{r_p}, \frac{\Gamma_{m_{\ell-1}+r_p+1}(r \zeta(t))}{r^\ell}, \dots \right. \\ \left. \dots, \frac{\Gamma_{m_\ell}(r \zeta(t))}{r^\ell}, t_{r_p+1}, \dots, t_p \right)$$

belongs to \mathbb{B} . This is a simple consequence of the equalities $\eta(r \zeta(t)) = \lambda_r \eta(\zeta(t)) = \lambda_r t$. We now use both (3.11) and (3.12) to conclude that the element represented in (3.23) converges to

$$(3.24) \quad \left(0, \dots, 0, t_1, \dots, t_{r_p}, 0, \dots, 0, t_{r_p+1}, \dots, t_p \right)$$

as $r \rightarrow 0^+$, uniformly with respect to t that varies in a bounded set. By standard facts on Hausdorff convergence, the previous limit implies the convergence in (1.3) where

$$(3.25) \quad \Pi_\Sigma(x) := \{z \in \mathbb{G} : z_1 = \cdots = z_{m_{\ell-1}} = z_{m_{\ell-1}+r_p+1} = \cdots = z_{m_\ell} = 0\}.$$

It can be easily seen that $\Pi_\Sigma(x)$ is the homogeneous subgroup of \mathbb{G} associated with the Lie subalgebra

$$\text{span} \{X_{m_{\ell-1}+1}, X_{m_{\ell-1}+2}, \dots, X_{m_{\ell-1}+r_p}, X_{m_\ell+1}, \dots, X_n\}.$$

Moreover, the convergence of the element represented in (3.23) to (3.24) also gives

$$(3.26) \quad \lim_{r \rightarrow 0^+} \frac{\tilde{\mu}(\Sigma \cap B(x, r))}{r^{D(p)}} = \mathcal{H}_{|\cdot|}^p(\mathbb{B} \cap S) \|(\partial_{y_1} \Phi \wedge \cdots \wedge \partial_{y_p} \Phi)(0)\|_{\tilde{g}, x}$$

where $S = \{(0, \dots, 0, t_1, \dots, t_{r_p}, 0, \dots, 0, t_{r_p+1}, \dots, t_p) \in \mathbb{R}^n : t_1, \dots, t_p \in \mathbb{R}\}$ and the metric unit ball \mathbb{B} is represented with respect to the same coordinates. Taking into account (3.4) and the matrix (2.21), we have the projection

$$\pi_{D(p), 0}((\partial_{y_1} \gamma \wedge \cdots \wedge \partial_{y_p} \gamma)(0)) = (X_{m_{\ell-1}+1} \wedge \cdots \wedge X_{m_{\ell-1}+r_p} \wedge X_{m_\ell+1} \wedge \cdots \wedge X_n)(0)$$

and the formulae $\partial_{y_j} \Phi(x) = dl_x(\partial_{y_j} \gamma(0))$ yield

$$\pi_{D(p), x}((\partial_{y_1} \Phi \wedge \cdots \wedge \partial_{y_p} \Phi)(0)) = (X_{m_{\ell-1}+1} \wedge \cdots \wedge X_{m_{\ell-1}+r_p} \wedge X_{m_\ell+1} \wedge \cdots \wedge X_n)(x).$$

We have the unit tangent p -vector

$$\tau_{\Sigma, \tilde{g}}(x) = \frac{(\partial_{y_1} \Phi \wedge \cdots \wedge \partial_{y_p} \Phi)(0)}{\|(\partial_{y_1} \Phi \wedge \cdots \wedge \partial_{y_p} \Phi)(0)\|_{\tilde{g}, x}}$$

then the previous equations for projections give

$$\|(\tau_{\Sigma, \tilde{g}}(x))_{D(p), x}\| = \frac{1}{\|(\partial_{y_1} \Phi \wedge \cdots \wedge \partial_{y_p} \Phi)(0)\|_{\tilde{g}, x}}.$$

As a result, in view of Definition 3.1, the limit (3.26) proves our last claim (1.4). \square

4. NEGLIGIBILITY OF LOWER DEGREE POINTS IN TRANSVERSAL SUBMANIFOLDS

The aim of this section is to prove Theorem 1.2 for a C^1 p -dimensional transversal submanifold $\Sigma \subset \mathbb{G}$, where we define

$$(4.1) \quad \Sigma_c := \{x \in \Sigma : d_\Sigma(x) < D(p)\}.$$

Since Σ is transversal, the subset Σ_c plays the role of a *generalized characteristic set* of Σ . Since any left translation is a diffeomorphism, for each point $x \in \Sigma$ there holds

$$(4.2) \quad T_0(x^{-1} \cdot \Sigma) = dl_{x^{-1}}(T_x \Sigma).$$

Clearly, a basis for $T_0(x^{-1} \cdot \Sigma)$ is given by

$$dl_{x^{-1}}(v_1), \dots, dl_{x^{-1}}(v_p),$$

where the vectors v_1, \dots, v_p are given by Lemma 2.11. If $v_j = \sum_i C_{ij} X_i(x)$, by the left invariance of X_i , we have

$$(4.3) \quad dl_{x^{-1}}(v_j) = \sum_{i=1}^n C_{ij} X_i(0) \quad \text{for any } j = 1, \dots, p.$$

In particular, $d_{x^{-1} \cdot \Sigma}(0) = d_\Sigma(x)$ and $0 \in (x^{-1} \cdot \Sigma)_c$ if and only if $x \in \Sigma_c$.

Taking into account (2.20), we observe that Σ_c can be written as the disjoint union

$$(4.4) \quad \Sigma_c = \Sigma_c^A \cup \Sigma_c^B,$$

where Σ_c^A and Σ_c^B are the (possibly empty) sets defined by

$$(4.5) \quad \begin{aligned} \Sigma_c^A &:= \{x \in \Sigma_c : \exists \bar{j} \geq \ell + 1 \text{ such that } \alpha_{\bar{j}} < n_{\bar{j}}\} \\ \Sigma_c^B &:= \{x \in \Sigma_c : \alpha_j = n_j \ \forall j \geq \ell + 1 \text{ and } \alpha_\ell < r_p\}. \end{aligned}$$

The integer ℓ , depending on p , is introduced in (2.10) and the nonnegative integers $\alpha_1, \dots, \alpha_\ell$ are defined in Lemma 2.11. In particular, ℓ will be used throughout this section. We notice that in the case $\ell = 1$, we must have $\alpha_\ell = r_p$, hence $\Sigma_c^B = \emptyset$.

We begin by making the further assumption that Σ is of class C^1 and such that $\Sigma \subset \phi([0, 1]^p)$ for some C^1 -regular map $\phi : [0, 1]^p \rightarrow \mathbb{G}$. By the uniform differentiability of ϕ , the boundedness of Σ and the continuity of left translations, the following statement holds: for any $\epsilon > 0$, there exists $\bar{r}_\epsilon > 0$ such that

$$(4.6) \quad \begin{aligned} &\forall r \in (0, \bar{r}_\epsilon), \\ |\langle y, w \rangle| \leq \epsilon r &\quad \forall x \in \Sigma, \ \forall y \in (x^{-1} \cdot \Sigma) \cap B_E(0, r), \\ &\quad \forall w \in (T_0(x^{-1} \cdot \Sigma))^\perp, \ |w| = 1, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the Euclidean scalar product. The orthogonal space $(T_0(x^{-1} \cdot \Sigma))^\perp$ is understood with respect to the same product. Notice that such coordinates are associated with the basis X_1, \dots, X_n given by Lemma 2.11; in particular, they depend on the chosen basepoint $x \in \Sigma$.

The proof of the negligibility stated in Theorem 1.2 stems from the following key lemmata. The proofs of these lemmata could be rather simplified; however, we present them in a form which will be helpful for some refinement provided in Section 5. In both statements, the number \bar{r}_ϵ is as in (4.6).

Lemma 4.1. *Let Σ be a C^1 submanifold such that $\Sigma \subset \phi([0, 1]^p)$ for some C^1 map $\phi : [0, 1]^p \rightarrow \mathbb{G}$; let $\theta := 1/\ell$. Then, there exists a constant $C_A = C_A(\Sigma) > 0$ such that the following property holds. For any x, ϵ, r satisfying*

$$(4.7) \quad x \in \Sigma_c^A, \quad \epsilon \in (0, 1) \quad \text{and} \quad 0 < r \leq \min\{\bar{r}_\epsilon, \epsilon^\ell\},$$

the set $(x^{-1} \cdot \Sigma) \cap B_E(0, r)$ can be covered by a family $\{B_i : i \in I\}$ of CC balls with radius r^θ such that

$$\#I \leq C_A \epsilon r^{p-\theta D(p)}.$$

Proof. From now on, the numbers C_i , with $i = 1, 2, \dots$, will denote positive constants depending only on Σ, p, \mathbb{G} and the fixed homogeneous distance d . For the reader's convenience, we divide the proof into several steps.

Step 1. By Theorem 2.5, we get a countable family $\{B(x_i, r^\theta) : i \in I\}$ such that

$$(4.8) \quad \begin{cases} x_i \in (x^{-1} \cdot \Sigma) \cap B_E(0, r) \\ (x^{-1} \cdot \Sigma) \cap B_E(0, r) \subset \bigcup_{i \in I} B(x_i, r^\theta) \\ B(x_i, r^\theta/5) \cap B(x_h, r^\theta/5) = \emptyset \text{ when } i \neq h. \end{cases}$$

We have to estimate $\#I$. By Lemma 2.3, for any $i \in I$ there exists \tilde{x}_i such that

$$(4.9) \quad \begin{aligned} \tilde{x}_i^1 = \cdots = \tilde{x}_i^\ell = 0, \quad d(x_i, \tilde{x}_i) \leq Cr^{1/\ell} = Cr^\theta, \\ |\tilde{x}_i^h - x_i^h| \leq Cr^2 \quad \text{for any } h = \ell + 1, \dots, \iota. \end{aligned}$$

Therefore, taking into account (2.6), we achieve

$$(4.10) \quad B(x_i, r^\theta) \subset B(\tilde{x}_i, (1+C)r^\theta) \subset \text{Box}(\tilde{x}_i, C_1r^\theta).$$

Let us also point out that both (4.9) and the fact that $x_i \in B_E(0, r)$ give

$$(4.11) \quad |\tilde{x}_i| \leq C_2r.$$

Step 2. Let us prove that there exists $C_3 > 0$ such that, for any $i \in I$, there holds

$$(4.12) \quad \text{Box}(\tilde{x}_i, C_1r^\theta) \subset \Omega,$$

where we have set

$$\Omega := (-C_3r^\theta, C_3r^\theta)^{n_1} \times (-C_3r^{2\theta}, C_3r^{2\theta})^{n_2} \times \cdots \times (-C_3r^{\ell\theta}, C_3r^{\ell\theta})^{n_\ell} \times \text{Box}_E^\mu(0, C_3r)$$

and $\mu := n - m_\ell$. To this aim we fix $y \in \text{Box}(0, C_1r^\theta)$, that is

$$(4.13) \quad |y^j| < (C_1r^\theta)^j \quad \forall j = 1, \dots, \iota,$$

and prove that $\tilde{x}_i \cdot y \in \Omega$. By explicit computation

$$(4.14) \quad \begin{aligned} \tilde{x}_i \cdot y &= (0, \dots, 0, \tilde{x}_i^{\ell+1}, \dots, \tilde{x}_i^\iota) \cdot (y^1, \dots, y^\iota) \\ &= (y^1, \dots, y^\ell, \tilde{x}_i^{\ell+1} + y^{\ell+1}, \tilde{x}_i^{\ell+2} + y^{\ell+2} + O(r^{1+\theta}), \dots, \tilde{x}_i^\iota + y^\iota + O(r^{1+\theta})) \end{aligned}$$

where we have used

- (2.3) for the coordinates in the layers $1, \dots, \ell + 1$;
- (2.4) for the coordinates in the layers $\ell + 2, \dots, \iota$, together with (4.11) and the fact that $|y| = O(r^\theta)$.

Here and in the sequel, all the quantities $O(\cdot)$ are uniform. From (4.14) and (4.13) it follows immediately that $\tilde{x}_i \cdot y \in \Omega$, and (4.12) follows.

Step 3. We have not used the fact that $x \in \Sigma_c^A$ yet. By definition, there exists $\bar{j} \geq \ell + 1$ such that $\alpha_{\bar{j}} < n_{\bar{j}}$. We can also assume that \bar{j} is maximum, i.e., that $\alpha_j = n_j$ for any $j > \bar{j}$; set

$$\nu := n_{\bar{j}} + n_{\bar{j}+1} + \cdots + n_\iota = n_{\bar{j}} + \alpha_{\bar{j}+1} + \cdots + \alpha_\iota.$$

The last ν rows of the matrix C given by Lemma 2.11 constitute a $\nu \times p$ matrix M of the form

$$M = \left[\begin{array}{c|c|c|c|c|c|c} 0 & \cdots & 0 & Id_{\alpha_{\bar{j}}} & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 0 & * & \cdots & * \\ \hline 0 & \cdots & 0 & 0 & Id_{n_{\bar{j}+1}} & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \hline 0 & \cdots & 0 & 0 & 0 & \cdots & Id_{n_\iota} \end{array} \right] = \left[\begin{array}{c|c|c|c|c} 0 & \cdots & 0 & Id_{\alpha_{\bar{j}}} & 0 \\ 0 & \cdots & 0 & 0 & * \\ \hline 0 & \cdots & 0 & 0 & Id_{n_{\bar{j}+1} + \cdots + n_\iota} \end{array} \right].$$

Since M has only $\alpha_{\bar{j}} + n_{\bar{j}+1} + \cdots + n_\iota < \nu$ nonzero columns, there exists a vector $z \in \mathbb{R}^\nu$ such that $|z| = 1$ and z is orthogonal to any of the columns of M . Therefore, the vector $w := (0, z) \in \mathbb{R}^n \equiv \mathbb{R}^{n-\nu} \times \mathbb{R}^\nu$ is orthogonal to any of the columns of C . By (4.2) and

(4.3), taking into account that $X_k(0) = \partial_{x_k}$, these columns generate $T_0(x^{-1} \cdot \Sigma)$. As a result, since $\bar{j} > \ell$, we are lead to the validity of the following conditions

$$(4.15) \quad \begin{cases} w \in (T_0(x^{-1} \cdot \Sigma))^\perp \\ |w| = 1 \\ w_1 = w_2 = \dots = w_{m_\ell} = 0 \end{cases} .$$

Step 4. To refine the inclusion (4.12), we will use the properties (4.15). By (4.6) one has $|\langle x_i, w \rangle| \leq \epsilon r$ for any $i \in I$. Define $w' := (w_{m_\ell+1}, w_{m_\ell+2}, \dots, w_n) \in \mathbb{R}^\mu$, where $\mu = n - m_\ell > \nu$ is the same number of Step 2. By (4.14) and (4.15), for any $y \in \text{Box}(0, C_1 r^\theta)$ we have

$$\begin{aligned} |\langle \tilde{x}_i \cdot y, w \rangle| &= |\langle (y^1, \dots, y^\ell, \tilde{x}_i^{\ell+1} + y^{\ell+1}, \tilde{x}_i^{\ell+2} + y^{\ell+2} + O(r^{1+\theta}), \dots, \tilde{x}_i^\ell + y^\ell + O(r^{1+\theta})), \\ &\quad (0, \dots, 0, w^{\ell+1}, \dots, w^\ell) \rangle| \\ &\leq |\langle (\tilde{x}_i^{\ell+1}, \dots, \tilde{x}_i^\ell), w' \rangle| + |\langle (y^{\ell+1}, \dots, y^\ell), w' \rangle| + O(r^{1+\theta}) \\ &= |\langle (x_i^{\ell+1}, \dots, x_i^\ell), w' \rangle| + O(r^{(\ell+1)\theta}) + O(r^{1+\theta}) \\ &\leq \epsilon r + O(r^{1+\theta}), \end{aligned}$$

where the second equality is justified by (4.9) and (4.13) and the last inequality follows from $(\ell + 1)\theta = 1 + \theta$. Since all the previous $O(\cdot)$ s are uniform with respect to the index i , we get

$$|\langle (\tilde{x}_i \cdot y)_\mu, w' \rangle| \leq \epsilon r + C_4 r^{1+\theta} \leq (1 + C_4) \epsilon r,$$

where $(\tilde{x}_i \cdot y)_\mu$ is the vector made by the last μ coordinates of $(\tilde{x}_i \cdot y)_\mu$ and we used the fact that, by (4.7), $r^\theta = r^{1/\ell} \leq \epsilon$. Thus, by (2.7) and (4.12) we obtain that $\text{Box}(\tilde{x}_i, C_1 r^\theta) \subset \tilde{\Omega}$, where we have set

$$\begin{aligned} \tilde{\Omega} &:= (-C_3 r^\theta, C_3 r^\theta)^{n_1} \times (-C_3 r^{2\theta}, C_3 r^{2\theta})^{n_2} \times \dots \\ &\quad \times (-C_3 r^{\ell\theta}, C_3 r^{\ell\theta})^{n_\ell} \times \text{Box}_{\text{span } w'}^\mu(0; C_3 r, C_5 \epsilon r). \end{aligned}$$

As a consequence, by (4.10) we get $B(x_i, r^\theta/5) \subset \text{Box}(\tilde{x}_i, C_1 r^\theta) \subset \tilde{\Omega}$ for all $i \in I$.

Step 5. We are ready to estimate $\#I$. The volume of $\tilde{\Omega}$ is equal to

$$a = C_6 \epsilon r^{\theta(n_1+2n_2+\dots+\ell n_\ell)+\mu} = C_6 \epsilon r^{\theta(n_1+2n_2+\dots+\ell n_\ell)+n_{\ell+1}+\dots+n_\ell},$$

while each $B(x_i, r^\theta/5)$ has volume $b = C_7 r^{\theta(n_1+2n_2+\dots+n_\ell)}$. Taking into account that the CC balls $B(x_i, r^\theta/5)$ are pairwise disjoint and contained in $\tilde{\Omega}$, we have

$$\begin{aligned} \#I \leq \frac{a}{b} &= \frac{C_6}{C_7} \epsilon r^{n_{\ell+1}+\dots+n_\ell - \theta((\ell+1)n_{\ell+1}+\dots+n_\ell)} \stackrel{(2.13)}{=} \frac{C_6}{C_7} \epsilon r^{p-r_p - \theta(D(p) - \ell r_p)} \\ &= \frac{C_6}{C_7} \epsilon r^{p - \theta D(p) + (\theta\ell - 1)r_p} \end{aligned}$$

which proves the claim and concludes the proof of the lemma. \square

While more subtle at certain points, the proof of Lemma 4.2 follows the same lines of the previous one. For the reader's benefit, we will try to make the analogies between the two proofs as evident as possible. We recall again that \bar{r}_ϵ is as in (4.6).

Lemma 4.2. *Under the assumptions of Lemma 4.1 and $\ell \geq 2$, there exists $C_B = C_B(\Sigma) > 0$ such that the following property holds. For any x, ϵ, θ, r satisfying*

$$(4.16) \quad x \in \Sigma_c^B, \quad \epsilon \in (0, 1), \quad \frac{1}{\ell} < \theta \leq \frac{1}{\ell-1} \quad \text{and} \quad 0 < r \leq \min\{\bar{r}_\epsilon, \epsilon^{1/(\ell\theta-1)}\},$$

the set $(x^{-1} \cdot \Sigma) \cap B_E(0, r)$ can be covered by a family $\{B_i : i \in I\}$ of CC balls with radius r^θ such that

$$\#I \leq C_B \epsilon^H r^{p-\theta D(p)-(\ell\theta-1)(n_\ell-r_p)},$$

where $H = H(x) := n_\ell - \alpha_\ell$ and the integers $\alpha_j = \alpha_j(x)$ are those given by Lemma 2.11.

Proof. We follow the same convention of Lemma 4.1 about the constants C_i .

Step 1. By the $5r$ -covering theorem we can cover $(x^{-1} \cdot \Sigma) \cap B_E(0, r)$ by a family of CC balls $\{B(x_i, r^\theta) : i \in I\}$ such that (4.8) holds. We have once more to estimate $\#I$. By Lemma 2.3, for any $i \in I$ there exists \tilde{x}_i such that

$$(4.17) \quad \begin{aligned} \tilde{x}_i^1 = \dots = \tilde{x}_i^{\ell-1} = 0, \quad d(x_i, \tilde{x}_i) \leq Cr^{1/(\ell-1)} \leq Cr^\theta \quad \text{and} \\ |\tilde{x}_i^h - x_i^h| \leq Cr^2 \quad \text{for any } h = \ell, \dots, \iota. \end{aligned}$$

Therefore $B(x_i, r^\theta/5) \subset B(x_i, r^\theta) \subset B(\tilde{x}_i, (1+C)r^\theta) \subset \text{Box}(\tilde{x}_i, C_8r^\theta)$. Again

$$(4.18) \quad |\tilde{x}_i| \leq C_2r.$$

Step 2. Let us prove that there exists $C_9 > 0$ such that, for any $i \in I$, there holds

$$(4.19) \quad \text{Box}(\tilde{x}_i, C_8r^\theta) \subset \Omega,$$

where now

$$(4.20) \quad \begin{aligned} \Omega := & (-C_9r^\theta, C_9r^\theta)^{n_1} \times (-C_9r^{2\theta}, C_9r^{2\theta})^{n_2} \times \dots \\ & \times (-C_9r^{(\ell-1)\theta}, C_9r^{(\ell-1)\theta})^{n_{\ell-1}} \times \text{Box}_E^\mu(0, C_9r) \end{aligned}$$

and $\mu := n - m_{\ell-1} = n_\ell + \dots + n_\iota$. As before we fix $y \in \text{Box}(0, C_8r^\theta)$,

$$(4.21) \quad |y^j| < (C_8r^\theta)^j \quad \forall j = 1, \dots, \iota$$

and prove that $\tilde{x}_i \cdot y \in \Omega$. Reasoning as in Step 2 in the proof of Lemma 4.1 we get

$$(4.22) \quad \begin{aligned} \tilde{x}_i \cdot y &= (0, \dots, 0, \tilde{x}_i^\ell, \dots, \tilde{x}_i^\iota) \cdot (y^1, \dots, y^\iota) \\ &= (y^1, \dots, y^{\ell-1}, \tilde{x}_i^\ell + y^\ell, \tilde{x}_i^{\ell+1} + y^{\ell+1} + O(r^{1+\theta}), \dots, \tilde{x}_i^\iota + y^\iota + O(r^{1+\theta})) \end{aligned}$$

where we have used (2.3), (2.4), (4.18) and the fact that $|y| = O(r^\theta)$. All the quantities $O(\cdot)$ are uniform. The inclusion (4.19) follows from (4.18), (4.22) and the fact that

$$|y^j| < (C_8r^\theta)^j = C_8^j r^{j\theta} \leq C_8^j r^{j/\ell} \leq C_8^j r \quad \forall j = \ell, \dots, \iota.$$

Step 3. Since $x \in \Sigma_c^B$ we have by definition

$$\alpha_\ell < r_p \quad \text{and} \quad \alpha_j = n_j \quad \forall j \geq \ell + 1.$$

Therefore the last μ rows of the matrix C from Lemma 2.11 constitute a $\mu \times p$ matrix M of the form

$$M = \left[\begin{array}{c|ccc|ccc} 0 & \cdots & 0 & Id_{\alpha_\ell} & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 0 & * & \cdots & * \\ \hline 0 & \cdots & 0 & 0 & Id_{n_{\ell+1}} & \cdots & 0 \\ \vdots & \cdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \hline 0 & \cdots & 0 & 0 & 0 & \cdots & Id_{n_\iota} \end{array} \right] = \left[\begin{array}{c|ccc|c} 0 & \cdots & 0 & Id_{\alpha_\ell} & 0 \\ 0 & \cdots & 0 & 0 & * \\ \hline 0 & \cdots & 0 & 0 & Id_{n_{\ell+1}+\cdots+n_\iota} \end{array} \right]$$

There are $\alpha_\ell + n_{\ell+1} + \cdots + n_\iota$ nonzero columns of M ; therefore, the columns of M span a vector subspace of \mathbb{R}^μ of dimension at most $\alpha_\ell + n_{\ell+1} + \cdots + n_\iota$. Since

$$\mu - (\alpha_\ell + n_{\ell+1} + \cdots + n_\iota) = n_\ell - \alpha_\ell = H,$$

it follows that there exist H linearly independent vectors $z_1, \dots, z_H \in \mathbb{R}^\mu$ such that $|z_k| = 1$ and z_k is orthogonal to any of the columns of M for any $k = 1, \dots, H$. In particular, the unit vectors

$$w_k := (0, z_k) \in \mathbb{R}^n \cong \mathbb{R}^{n-\mu} \times \mathbb{R}^\mu, \quad k = 1, \dots, H$$

are orthogonal to any of the columns of C , which form a basis of $T_0(x^{-1} \cdot \Sigma)$. Setting $W := \text{span}(w_1, \dots, w_H)$ we have

$$W \subset T_0(x^{-1} \cdot \Sigma)^\perp \quad \text{and} \quad \dim W = H \geq 1;$$

moreover, any vector $w \in W$ is of the form

$$(4.23) \quad w = (0, \dots, 0, w^\ell, \dots, w^\iota) = (0, w') \in \mathbb{R}^{m_{\ell-1}} \times \mathbb{R}^\mu.$$

Step 4. Again we want to refine the inclusion (4.19). By (4.6) there holds

$$|\langle x_i, w \rangle| \leq \epsilon r \quad \forall i \in I, \forall w \in W \text{ with } |w| = 1.$$

Recalling (4.22) and writing $w = (0, w') \in \mathbb{R}^{m_{\ell-1}} \times \mathbb{R}^\mu$ as in (4.23), for any $y \in \text{Box}(0, C_8 r^\theta)$ we have

$$\begin{aligned} |\langle \tilde{x}_i \cdot y, w \rangle| &= \left| \langle (y^1, \dots, y^{\ell-1}, \tilde{x}_i^\ell + y^\ell, \tilde{x}_i^{\ell+1} + y^{\ell+1} + O(r^{1+\theta}), \dots, \tilde{x}_i^\iota + y^\iota + O(r^{1+\theta})), \right. \\ &\quad \left. (0, \dots, 0, w^\ell, \dots, w^\iota) \rangle \right| \\ &\leq |\langle (\tilde{x}_i^\ell, \dots, \tilde{x}_i^\iota), w' \rangle| + |\langle (y^\ell, \dots, y^\iota), w' \rangle| + O(r^{1+\theta}) \\ &= |\langle (x_i^\ell, \dots, x_i^\iota), w' \rangle| + O(r^{\ell\theta}) + O(r^{1+\theta}) \\ &\leq \epsilon r + O(r^{\ell\theta}) + O(r^{1+\theta}) \quad \forall w \in W, |w| = 1 \end{aligned}$$

where we used (4.17) and (4.21). Since

$$\ell\theta = (\ell - 1)\theta + \theta \leq 1 + \theta,$$

we have $r^{1+\theta} \leq r^{\ell\theta}$ and thus, since all the $O(\cdot)$ s are uniform,

$$|\langle \tilde{x}_i \cdot y, w \rangle| \leq \epsilon r + C_{10} r^{\ell\theta} \leq \max\{\epsilon, C_{10} r^{\ell\theta-1}\} r \leq C_{11} \epsilon r \quad \forall w \in W, |w| = 1,$$

the last inequality following from (4.16). Using (2.7) we can then refine (4.20) to obtain

$$B(x_i, r^\theta/5) \subset \text{Box}(\tilde{x}_i, C_8 r^\theta) \subset \tilde{\Omega} \quad \forall i \in I$$

where

$$\begin{aligned} \tilde{\Omega} &:= (-C_9 r^\theta, C_9 r^\theta)^{n_1} \times (-C_9 r^{2\theta}, C_9 r^{2\theta})^{n_2} \times \dots \\ &\quad \times (-C_9 r^{(\ell-1)\theta}, C_9 r^{(\ell-1)\theta})^{n_{\ell-1}} \times \text{Box}_W^\mu(0; C_9 r, C_{11} \epsilon r). \end{aligned}$$

Step 5. We can now estimate $\#I$. Since $\dim W = H$, the volume of $\tilde{\Omega}$ is

$$a = C_{12} \epsilon^H r^{\theta(n_1+2n_2+\dots+(\ell-1)n_{\ell-1})+\mu} = C_{12} \epsilon^H r^{\theta(n_1+2n_2+\dots+(\ell-1)n_{\ell-1})+n_\ell+\dots+n_\iota},$$

while each ball $B(x_i, r^\theta/5)$ has volume $b = C_7 r^{\theta(n_1+2n_2+\dots+n_\iota)}$. Since the CC balls $B(x_i, r^\theta/5)$ are pairwise disjoint and contained in $\tilde{\Omega}$, we have

$$\begin{aligned} \#I \leq \frac{a}{b} &= \frac{C_{12}}{C_7} \epsilon^H r^{n_\ell+\dots+n_\iota-\theta(\ell n_\ell+\dots+n_\iota)} \\ &\stackrel{(2.13)}{=} \frac{C_{12}}{C_7} \epsilon^H r^{n_\ell+p-r_p-\theta[\ell(n_\ell-r_p)+\ell r_p+(\ell+1)n_{\ell+1}+\dots+n_\iota]} \\ &\stackrel{(2.13)}{=} \frac{C_{12}}{C_7} \epsilon^H r^{p+(n_\ell-r_p)-\theta[\ell(n_\ell-r_p)+D(p)]} \\ &= \frac{C_{12}}{C_7} \epsilon^H r^{p-\theta D(p)-(\ell\theta-1)(n_\ell-r_p)}, \end{aligned}$$

as claimed. \square

Lemma 4.3. *Let Σ be a C^1 submanifold such that $\Sigma \subset \phi([0, 1]^p)$ for a C^1 map $\phi : [0, 1]^p \rightarrow \mathbb{G}$. Then $\mathcal{H}^{D(p)}(\Sigma_c) = 0$.*

Proof. Clearly, it will be enough to show that

$$(4.24) \quad \mathcal{H}^{D(p)}(\Sigma_c^A) = 0 \quad \text{and} \quad \mathcal{H}^{D(p)}(\Sigma_c^B) = 0.$$

Step 1. We start by proving the first equality in (4.24); let us follow the same convention of Lemmata 4.1 and 4.2 about the constants C_i .

Let $\epsilon \in (0, 1)$ and $r \in (0, \min\{\bar{r}_\epsilon, \epsilon^\ell\}]$ be fixed. Since $(x, y) \rightarrow x^{-1}y$ is locally Lipschitz and ϕ is Lipschitz, both with respect to the Euclidean distance, we obtain $C_{13} > 0$ such that

$$(4.25) \quad \text{if } z_1, z_2 \in [0, 1]^p \text{ and } |z_1 - z_2| \leq C_{13}r, \quad \text{then } |\phi(z_1)^{-1} \cdot \phi(z_2)| < r.$$

Let us divide $[0, 1]^p$, in a standard fashion, into a family of closed subcubes of diameter not greater than $C_{13}r$; in this way there will be less than $C_{14}r^{-p}$ such subcubes. Let $(Q_j)_{j \in J}$ be the family of those subcubes with the property that

$$\phi(Q_j) \cap \Sigma_c^A \neq \emptyset$$

and fix $x_j \in \phi(Q_j) \cap \Sigma_c^A$. By (4.25) we have

$$x_j^{-1} \cdot \phi(Q_j) \subset (x_j^{-1} \cdot \Sigma) \cap B_E(0, r).$$

Writing $\theta := 1/\ell$, Lemma 4.1 ensures that $x_j^{-1} \cdot \phi(Q_j)$ can be covered by (at most) $C_A \epsilon r^{p-\theta D(p)}$ balls of radius r^θ ; by left invariance, the same holds for $\phi(Q_j)$. In particular, since $\Sigma_c^A \subset \cup_{j \in J} \phi(Q_j)$ and $\#J \leq C_{14}r^{-p}$, we have that, for any $r \in (0, \min\{\bar{r}_\epsilon, \epsilon^\ell\}]$, the set Σ_c^A can be covered by a family of CC balls with radius r^θ of cardinality controlled by $C_A C_{14} \epsilon r^{-\theta D(p)}$. Therefore

$$\mathcal{H}_{2r^{1/\ell}}^{D(p)}(\Sigma_c^A) \leq C_A C_{14} \epsilon r^{-\theta D(p)} (2r^\theta)^{D(p)} = 2^{D(p)} C_A C_{14} \epsilon$$

whence, letting $r \rightarrow 0^+$,

$$\mathcal{H}^{D(p)}(\Sigma_c^A) \leq 2^{D(p)} C_A C_{14} \epsilon.$$

The first part of (4.24) follows by letting $\epsilon \rightarrow 0$.

Step 2. Let us prove the second equality in (4.24). Let $\epsilon \in (0, 1)$ and $r \in (0, \min\{\bar{r}_\epsilon, \epsilon^{\ell-1}\}]$ be fixed; we have $\epsilon = r^\lambda$ for a suitable $\lambda = \lambda(\epsilon, r) \in (0, \frac{1}{\ell-1}]$. Define $\theta = \theta(\epsilon, r) := \frac{1+\lambda}{\ell}$ and observe that $1/\ell < \theta \leq 1/(\ell-1)$. As a result, we have

$$(4.26) \quad r^{\ell\theta-1} = r^\lambda = \epsilon;$$

in particular, $\epsilon^{1/(\ell\theta-1)} = r \leq \bar{r}_\epsilon$ (the inequality here comes from the choice of r) and the conditions in (4.16) are satisfied. As before, we divide $[0, 1]^p$ into a family of (at most) $C_{14}r^{-p}$ closed subcubes of diameter not greater than $C_{13}r$. Let $(Q_k)_{k \in K}$ be the family of those subcubes with the property that

$$\phi(Q_k) \cap \Sigma_c^B \neq \emptyset$$

and fix $x_k \in \phi(Q_k) \cap \Sigma_c^A$. By (4.25) we have again

$$x_k^{-1} \cdot \phi(Q_k) \subset (x_k^{-1} \cdot \Sigma) \cap B_E(0, r)$$

so that, by Lemma 4.2, $x_k^{-1} \cdot \phi(Q_k)$ can be covered by no more than

$$C_B \epsilon^{H(x_k)} r^{p-\theta D(p) - (\ell\theta-1)(n_\ell-r_p)}$$

balls of radius $r^\theta = \epsilon^{1/\ell} r^{1/\ell}$; by left invariance, the same holds for $\phi(Q_k)$. Notice that

$$H(x_k) = n_\ell - \alpha_\ell(x_k) \geq n_\ell - r_p + 1 \quad \forall k \in K,$$

i.e., $\phi(Q_k)$ can be covered by (at most) $C_B \epsilon^{n_\ell-r_p+1} r^{p-\theta D(p) - (\ell\theta-1)(n_\ell-r_p)}$ balls of radius r^θ . As before, this implies that

$$\#K \leq C_B C_{14} \epsilon^{n_\ell-r_p+1} r^{-\theta D(p) - (\ell\theta-1)(n_\ell-r_p)}$$

whence, using (4.26),

$$(4.27) \quad \begin{aligned} \mathcal{H}_{2r^\theta}^{D(p)}(\Sigma_c^B) &\leq C_B C_{14} \epsilon^{n_\ell-r_p+1} r^{-\theta D(p)} r^{-(\ell\theta-1)(n_\ell-r_p)} (2r^\theta)^{D(p)} \\ &= 2^{D(p)} C_B C_{14} \epsilon^{n_\ell-r_p+1} \epsilon^{-(n_\ell-r_p)} \\ &= 2^{D(p)} C_B C_{14} \epsilon. \end{aligned}$$

Observing that

$$\lim_{r \rightarrow 0^+} r^\theta = \lim_{r \rightarrow 0^+} r^{1/\ell} r^{\lambda(\epsilon, r)/\ell} = \lim_{r \rightarrow 0^+} r^{1/\ell} \epsilon^{1/\ell} = 0$$

we can let $r \rightarrow 0^+$ in (4.27) to obtain

$$\mathcal{H}^{D(p)}(\Sigma_c^B) \leq 2^{D(p)} C_B C_{14} \epsilon.$$

This proves the second equality in (4.24) and completes the proof. \square

Remark 4.4. We point out for future references the following two facts proved, respectively, in Step 1 and Step 2 of the proof of Lemma 4.3. Let $\epsilon \in (0, 1)$ be fixed and assume that Σ is a C^1 submanifold such that $\Sigma \subset \phi([0, 1]^p)$ for a C^1 map $\phi : [0, 1]^p \rightarrow \mathbb{G}$; then

$$(4.28) \quad \text{for any } r \in (0, \min\{\bar{r}_\epsilon, \epsilon^\ell\}], \text{ the set } \Sigma_c^A \text{ can be covered by a family of CC balls with radius } r^{1/\ell} \text{ of cardinality at most } C_A C_{14} \epsilon r^{-D(p)/\ell}.$$

and

$$(4.29) \quad \text{for any } r \in (0, \min\{\bar{r}_\epsilon, \epsilon^{\ell-1}\}], \text{ the set } \Sigma_c^B \text{ can be covered by a family of CC balls, with radius } \epsilon^{1/\ell} r^{1/\ell}, \text{ of cardinality at most } C_B C_{14} \epsilon^{-D(p)/\ell} r^{-D(p)/\ell}.$$

In (4.29), we used the fact that the cardinality of the involved family is controlled by

$$\begin{aligned} & C_B C_{14} \epsilon^{n_\ell - r_p + 1} r^{-\theta D(p) - (\ell\theta - 1)(n_\ell - r_p)} \\ &= C_B C_{14} \epsilon^{n_\ell - r_p + 1} \epsilon^{-D(p)/\ell} r^{-D(p)/\ell} \epsilon^{-(n_\ell - r_p)} \\ &= C_B C_{14} \epsilon^{-D(p)/\ell} r^{-D(p)/\ell}, \end{aligned}$$

where we also utilized the equalities $r^\theta = \epsilon^{1/\ell} r^{1/\ell}$ and $r^{\ell\theta - 1} = \epsilon$.

The proof of Theorem 1.2 is now at hand.

Proof of Theorem 1.2. The theorem is an easy consequence of Lemma 4.3 and a standard localization argument. \square

Actually, Theorem 1.2 can be generalized to Lipschitz p -dimensional submanifolds; recall that the singular set Σ_* was defined at the beginning of Section 2.4. Clearly, the definition of Σ_c given at (4.1) for C^1 submanifolds extends to Lipschitz submanifolds considering the subset $\Sigma \setminus \Sigma_*$ of regular points, since the pointwise degree is defined by the existence of the pointwise tangent space.

Theorem 4.5. *Let $\Sigma \subset \mathbb{G}$ be a p -dimensional Lipschitz submanifold, let Σ_* be its singular set and denote by Σ_c be the subset of points in $\Sigma \setminus \Sigma_*$ whose degree is less than $D(p)$. It follows that*

$$(4.30) \quad \mathcal{H}^{D(p)}(\Sigma_* \cup \Sigma_c) = 0.$$

Proof. By definition, Σ is locally the graph of a Euclidean Lipschitz function, hence without loss of generality, we can assume that $\Sigma \subset \phi(A)$, where ϕ is the graph function given by a Lipschitz function $f : A \rightarrow V$, $A \subset W$ is a bounded open set of W and \mathbb{G} is seen as by $W \times V$, where W and V are linear subspaces of dimensions p and $n - p$, respectively. Let $\epsilon > 0$ be arbitrarily fixed such that $0 < \epsilon < \mathcal{L}^p(A)$. By the classical Whitney's extension theorem, there exists a C^1 function $f_\epsilon : A \rightarrow V$ such that the set

$$(4.31) \quad E_\epsilon := \{z \in A : f_\epsilon(z) = f(z) \text{ and } \nabla f_\epsilon(z) = \nabla f(z)\}$$

satisfies $\mathcal{L}^p(A \setminus E_\epsilon) < \epsilon$. The graph function ϕ_ϵ associated to f_ϵ defines the C^1 submanifold $\Sigma^\epsilon := \phi_\epsilon(A)$, hence Theorem 1.2 implies that that its generalized characteristic set $\Sigma_c^\epsilon := \{x \in \Sigma^\epsilon : d_{\Sigma^\epsilon}(x) < D(p)\}$ is $\mathcal{H}^{D(p)}$ -negligible. By the conditions of (4.31), we have the inclusion $\Sigma_c \cap \phi(E_\epsilon) \subset \Sigma_c^\epsilon$, hence $\Sigma_c \cap \phi(E_\epsilon)$ is also $\mathcal{H}^{D(p)}$ -negligible. As a consequence of [4, Proposition 3.1], there exists a geometric constant $C > 0$, only depending on the diameter of $\phi(A)$ and on \mathbb{G} , such that

$$\mathcal{H}^{D(p)}(\Sigma_c) = \mathcal{H}^{D(p)}(\Sigma_c \setminus \phi(E_\epsilon)) \leq C \mathcal{H}_{| \cdot |}^p(\Sigma_c \setminus \phi(E_\epsilon)) \leq C L^p \epsilon,$$

where $L > 0$ is the Euclidean Lipschitz constant of ϕ . The arbitrary choice of ϵ implies that $\mathcal{H}^{D(p)}(\Sigma_c) = 0$ and using (2.14), the proof is accomplished. \square

5. SIZE OF THE CHARACTERISTIC SET FOR $C^{1,\lambda}$ SUBMANIFOLDS

In this section we assume that Σ is a submanifold of class $C^{1,\lambda}$ for some $\lambda \in (0, 1]$. Our aim is to refine Theorem 1.2 and obtain estimates on the Hausdorff dimension of the characteristic set Σ_c . We first assume that $\Sigma \subset \phi([0, 1]^p)$ for some map $\phi \in C^{1,\lambda}([0, 1]^p, \mathbb{G})$. Under this assumption, there exists $C = C(\Sigma) > 0$ such that

$$(5.1) \quad |\langle y, w \rangle| \leq Cr^{1+\lambda} \quad \begin{array}{l} \forall x \in \Sigma, \forall y \in (x^{-1} \cdot \Sigma) \cap B_E(0, r), \\ \forall w \in (T_0(x^{-1} \cdot \Sigma))^\perp, |w| = 1. \end{array}$$

In other words, the number \bar{r}_ϵ defined by (4.6) can be chosen to be $\bar{r}_\epsilon = (\epsilon/C)^{1/\lambda}$.

As in (4.4), we write $\Sigma_c = \Sigma_c^A \cup \Sigma_c^B$ where, following (4.5), we define

$$\begin{aligned} \Sigma_c^A &= \{x \in \Sigma_c : \exists \bar{j} \geq \ell + 1 \text{ such that } \alpha_{\bar{j}} < n_{\bar{j}}\} \\ \Sigma_c^B &= \{x \in \Sigma_c : \alpha_j = n_j \forall j \geq \ell + 1 \text{ and } \alpha_\ell < r_p\}. \end{aligned}$$

Again, if $\ell = 1$, then $\Sigma_c^B = \emptyset$.

Lemma 5.1. *Let $\Sigma \subset \mathbb{G}$ be a $C^{1,\lambda}$ submanifold such that $\Sigma \subset \phi([0, 1]^p)$ for some map $\phi \in C^{1,\lambda}([0, 1]^p, \mathbb{G})$. Then*

$$(5.2) \quad \begin{array}{ll} \dim_H \Sigma_c^A \leq D(p) - 1 & \text{if } \lambda \geq 1/\ell \\ \dim_H \Sigma_c^A \leq D(p) - \ell\lambda & \text{if } \lambda \leq 1/\ell. \end{array}$$

Proof. If $\lambda > 1/\ell$ we have

$$\min\{\bar{r}_\epsilon, \epsilon^\ell\} = \min\{(\epsilon/C)^{1/\lambda}, \epsilon^\ell\} = \epsilon^\ell$$

for any $\epsilon > 0$ small enough. We are then allowed to use (4.28) with $r := \epsilon^\ell$ and obtain that, for any $\epsilon > 0$ small enough, the set Σ_c^A can be covered by a family of balls with radius ϵ of cardinality at most $C_A C_{14} \epsilon^{-D(p)}$. By Proposition 2.4 we get

$$\dim_H \Sigma_c^A \leq D(p) - 1.$$

On the other hand, if $\lambda \leq 1/\ell$ we have

$$\min\{\bar{r}_\epsilon, \epsilon^\ell\} = \min\{(\epsilon/C)^{1/\lambda}, \epsilon^\ell\} = C_{15} \epsilon^{1/\lambda}$$

for any $\epsilon > 0$ small enough; we have utilized the usual convention on constants C_i . Using (4.28) with $r := C_{15} \epsilon^{1/\lambda}$, we get that, for any $\epsilon > 0$ small enough, the set Σ_c^A can be covered by a family of balls with radius $r^{1/\ell} = C_{16} \epsilon^{1/(\ell\lambda)}$ of cardinality at most $C_{17} \epsilon^{-D(p)/(\ell\lambda)}$. By Proposition 2.4 we get

$$\dim_H \Sigma_c^A \leq D(p) - \ell\lambda$$

and this concludes the proof. \square

Lemma 5.2. *Let $\Sigma \subset \mathbb{G}$ be a $C^{1,\lambda}$ submanifold such that $\Sigma \subset \phi([0, 1]^p)$ for some map $\phi \in C^{1,\lambda}([0, 1]^p, \mathbb{G})$; assume $\ell \geq 2$. Then*

$$(5.3) \quad \begin{array}{ll} \dim_H \Sigma_c^B \leq D(p) - 1 & \text{if } \lambda \geq \frac{1}{\ell-1} \\ \dim_H \Sigma_c^B \leq D(p) - \frac{\ell\lambda}{1+\lambda} & \text{if } \lambda \leq \frac{1}{\ell-1}. \end{array}$$

Proof. If $\lambda > 1/(\ell - 1)$ we have

$$\min\{\bar{r}_\epsilon, \epsilon^{\ell-1}\} = \min\{(\epsilon/C)^{1/\lambda}, \epsilon^{\ell-1}\} = \epsilon^{\ell-1}$$

for any $\epsilon > 0$ small enough. We are then allowed to use (4.29) with $r := \epsilon^{\ell-1}$ and obtain that, for any $\epsilon > 0$ small enough, the set Σ_c^B can be covered by a family of balls with radius $\epsilon^{1/\ell} \epsilon^{(\ell-1)/\ell} = \epsilon$ of cardinality at most

$$C_B C_{14} \epsilon \epsilon^{-D(p)/\ell} r^{-D(p)/\ell} = C_B C_{14} \epsilon^{-D(p)+1}.$$

By Proposition 2.4 we get

$$\dim_H \Sigma_c^B \leq D(p) - 1.$$

On the other hand, if $\lambda \leq 1/(\ell - 1)$ we have

$$\min\{\bar{r}_\epsilon, \epsilon^{\ell-1}\} = \min\{(\epsilon/C)^{1/\lambda}, \epsilon^{\ell-1}\} = C_{18} \epsilon^{1/\lambda}$$

for any $\epsilon > 0$ small enough. Using (4.29) with $r := C_{18} \epsilon^{1/\lambda}$, we get that, for any $\epsilon > 0$ small enough, the set Σ_c^B can be covered by a family of balls with radius $\epsilon^{1/\ell} r^{1/\ell} = C_{17} \epsilon^{(\lambda+1)/(\ell\lambda)}$ of cardinality at most

$$C_B C_{14} \epsilon \epsilon^{-D(p)/\ell} r^{-D(p)/\ell} = C_B C_{14} \epsilon^{-\frac{D(p)}{\ell}} \epsilon^{-\frac{D(p)}{\ell\lambda}} = C_B C_{14} \epsilon^{-\frac{\lambda+1}{\ell\lambda} D(p)+1}.$$

By Proposition 2.4 we get

$$\dim_H \Sigma_c^B \leq D(p) - \frac{\ell\lambda}{\lambda+1}$$

and this concludes the proof. \square

Proof of Theorem 1.3. Recalling that $\Sigma_c^B = \emptyset$ if $\ell = 1$, the proof is now an easy consequence of Lemmata 5.1 and 5.2. \square

Remark 5.3. It is interesting to analyze Theorem 1.3 when the Carnot group \mathbb{G} is the Heisenberg group \mathbb{H}^n . In this case, $\ell = \ell(p) = 1$ for all $p = 2, \dots, 2n$ and Theorem 1.3 reads as

$$(5.4) \quad \dim_H \Sigma_c \leq p + 1 - \lambda$$

for any p -dimensional submanifold $\Sigma \subset \mathbb{H}^n$ of class $C^{1,\lambda}$. These estimates coincide with the results stated in Remark 1, page 72 of [2]. In the special case $p = 1$, we have $\ell = 2$, hence Theorem 1.3 gives

$$(5.5) \quad \dim_H \Sigma_c \leq D(1) - \frac{2\lambda}{1+\lambda} = \frac{2}{1+\lambda} \leq 2 - \lambda,$$

where the last inequality is strict for all $\lambda \in (0, 1)$ and $2 - \lambda = p + 1 - \lambda$. Thus, in this special case of curves ($p = 1$), the estimates (1.6) improve that of Remark 1, page 72 of [2].

REFERENCES

- [1] L. AMBROSIO, *Some fine properties of sets of finite perimeter in Ahlfors regular metric measure spaces*, Adv. Math., **159**, 51-67, (2001)
- [2] Z. M. BALOGH, *Size of characteristic sets and functions with prescribed gradients*, J. Reine Angew. Math., **564**, 63-83, (2003)
- [3] Z. M. BALOGH, C. PINTEA & H. ROHNER, *Size of Tangencies to Non-Involutive Distributions*, Indiana Univ. Math. J. **60** (2011), no. 6, 2061–2092.
- [4] Z. M. BALOGH, J. T. TYSON & B. WARHURST, *Sub-Riemannian vs. Euclidean dimension comparison and fractal geometry on Carnot groups*, Adv. Math. **220** (2009), no. 2, 560–619
- [5] V. BARONE ADESI, F. SERRA CASSANO, D. VITTONI, *The Bernstein problem for intrinsic graphs in Heisenberg groups and calibrations*, Calc. Var. Partial Differential Equations **30**, no. 1, 17-49, (2007)
- [6] L. CAPOGNA, D. DANIELLI, N. GAROFALO, *The geometric Sobolev embedding for vector fields and the isoperimetric inequality*, Comm. Anal. Geom. **2**, n.2, 203-215, (1994)
- [7] L. CAPOGNA, N. GAROFALO, *Ahlfors type estimates for perimeter measures in Carnot-Carathéodory spaces*, J. Geom. Anal. **16**, n.3, 455-497, (2006)
- [8] D. DANIELLI, N. GAROFALO, D. M. NHIEU, *Non-doubling Ahlfors measures, Perimeter measures, and the characterization of the trace spaces of Sobolev functions in Carnot-Carathéodory spaces*, Mem. Amer. Math. Soc. **182**, n.857, (2006)
- [9] D. DANIELLI, N. GAROFALO, D. M. NHIEU, *Sub-Riemannian calculus on hypersurfaces in Carnot groups*, Adv. Math., **215**, n.1, 292-378, (2007)
- [10] D. DANIELLI, N. GAROFALO, D. M. NHIEU, *A notable family of entire intrinsic minimal graphs in the Heisenberg group which are not perimeter minimizing*, Amer. J. Math, 130 (2008) no.2, 317-339
- [11] H. FEDERER, *Geometric Measure Theory*, Springer, (1969)
- [12] B. FRANCHI, S. GALLOT, R. L. WHEEDEN, *Sobolev and isoperimetric inequalities for degenerate metrics*, Math. Ann. **300**, 557-571 (1994)
- [13] B. FRANCHI, R. SERAPIONI, F. SERRA CASSANO, *Regular hypersurfaces, intrinsic perimeter and implicit function theorem in Carnot groups*, Comm. Anal. Geom., **11**, n.5, 909-944, (2003)
- [14] B. FRANCHI, R. SERAPIONI & F. SERRA CASSANO, *On the structure of finite perimeter sets in step 2 Carnot groups*, J. Geom. An. **13**, 421-466, (2003)
- [15] G. B. FOLLAND & E. M. STEIN, *Hardy spaces on homogeneous groups*, Princeton University Press, 1982
- [16] N. GAROFALO, D. M. NHIEU, *Isoperimetric and Sobolev Inequalities for Carnot-Carathéodory Spaces and the Existence of Minimal Surfaces*, Comm. Pure Appl. Math. **49**, 1081-1144 (1996)
- [17] M. GROMOV, *Carnot-Carathéodory spaces seen from within*, in *Subriemannian Geometry*, Progress in Mathematics, **144**, edited by A. Bellaïche and J.-J. Risler, Birkhäuser Verlag, (1996)
- [18] R. KORTE, V. MAGNANI, *Measure of curves in graded groups*, Illinois J. Math., to appear
- [19] E. LE DONNE, V. MAGNANI, *Measure of submanifolds in the Engel group*, Rev. Mat. Iberoamericana, **26**, n.1, 333-346, (2010)
- [20] V. MAGNANI, *Characteristic points, rectifiability and perimeter measure on stratified groups*, J. Eur. Math. Soc., vol. 8, n.4, 585-609, (2006)
- [21] V. MAGNANI, *Non-horizontal submanifolds and coarea formula*, J. Anal. Math., **106**, 95-127, (2008)
- [22] V. MAGNANI, *Blow-up estimates at horizontal points and applications*, J. Geom. Anal. **20**, n.3, 705-722, (2010)
- [23] V. MAGNANI & D. VITTONI, *An intrinsic measure for submanifolds in stratified groups*, J. Reine Angew. Math. **619** (2008), 203–232

- [24] P. MATTILA, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, Cambridge Studies in Advanced Mathematics, **44**, Cambridge University Press, Cambridge, 1995
- [25] P. MATTILA, *Measures with unique tangent measures in metric groups*, Math. Scand. **97**, n.2, 298-308 (2005)
- [26] R. MONTI, D. MORBIDELLI, *Regular domains in homogeneous groups*, Trans. Amer. Math. Soc. **357**, n.8, 2975-3011, (2005)
- [27] L. SIMON, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, 3. Australian National University, Centre for Mathematical Analysis, Canberra, 1983. vii+272 pp.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127, PISA, ITALY

E-mail address: magnani@dm.unipi.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN ST., URBANA, IL 61801 USA

E-mail address: tyson@math.uiuc.edu

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35121 PADOVA, ITALY

E-mail address: vittone@math.unipd.it