

The energy density of martensitic thin films via dimension reduction

Lorenzo Freddi Roberto Paroni

Abstract

A variational limit defined on the space of bi-dimensional gradient Young measures is obtained from three-dimensional elasticity via dimension reduction. The obtained limit problem uniquely determines the energy density of the thin film. Our result might be used to compute the microstructure in membranes made of phase transforming material.

1 Introduction

Martensitic thin films have recently attracted much interest because of their applications in the construction of actuators, sensors and micromachines. These technological applications have generated a new interest in nonlinear membranes theories. The purpose of the present paper is to obtain, by means of a rigorous dimension reduction, the energy density of a thin film, and hence to deduce a membrane model capable to describe the microstructure observed in martensitic materials.

Phase transformations have been successfully modeled within the theory of non-linear elasticity by Ball and James [5]. Within this approach, equilibrium configurations are found by minimizing the total free energy of the body.

In the spirit of the works on dimension reduction (see for instance [1, 3]) we start by considering the total free energy

$$I_\varepsilon(y) = \int_{\Omega_\varepsilon} W(Dy) dx - \int_{\Omega_\varepsilon} \hat{f}^\varepsilon \cdot y dx$$

corresponding to a deformation y of a hyperelastic three-dimensional body which occupies the cylindrical region $\Omega_\varepsilon := \omega \times (-\varepsilon/2, \varepsilon/2)$, with Helmholtz free energy W and subject to some body forces \hat{f}^ε and boundary conditions.

Under the usual assumptions on W and \hat{f}^ε there are different topologies which ensure compactness to the family of minimizers (or quasi-minimizers). Once chosen one of these topologies, the membrane model is obtained by passing to the limit as $\varepsilon \rightarrow 0$ in an appropriate variational sense (Γ -convergence). Roughly speaking, this variational limit ensures the convergence, in such topology, of minimizers of the energy at level ε to the minimizers of the membrane problem. The obtained limit problem depends on the chosen topology.

Le Dret and Raoult [22], working with the strong convergence in L^p , with an appropriate p bigger than 1 related to the growth of W , have found a limit energy of the following form

$$J(y) = \int_{\omega} QW_0(D_\alpha y) dx_\alpha - \int_{\omega} m_3 f \cdot y dx_\alpha,$$

where QW_0 denotes the quasiconvex envelope of the function

$$W_0(\bar{F}) := \min\{W(\bar{F}|z) : z \in \mathbb{R}^3\}, \quad \bar{F} \in M^{3 \times 2}.$$

By $D_\alpha y = (D_1 y, D_2 y)$ we mean the gradient of the admissible deformation $y = y(x_\alpha)$, $x_\alpha = (x_1, x_2)$. At this introductory level it is useless to specify the meaning of $m_3 f$, for which we refer to Section 4. A similar result in the case of strings was previously obtained by Acerbi, Buttazzo and Percivale in [1]. Typically, the infimum of the total free energy I_ε of a martensitic material is not attained. The minimizing sequences shall, in general, develop fine scale oscillations, which, according to the interpretation due to Ball and James [5, 6], model the microstructure experimentally observed in specimens of phase transforming materials. On the other hand the minimum of the functional J is achieved but the limit problem, when applied to martensitic materials, does not describe the microstructure. Our aim is to derive a thin film theory capable to capture the oscillations of the minimizing sequences. Inspired by earlier works on phase transitions by Ball and James [6], James and Kinderlehrer [19], Kinderlehrer and Pedregal [20, 21], we take the variational limit in the framework of parametrized measures introduced by L.C. Young [31], which is one of the most successful tools used to characterize the oscillatory behaviour of sequences of functions. To this end we extend the three-dimensional energy functionals to the space of Young measures associating to each gradient Dy the corresponding Young measure $\mu_x = \delta_{Dy(x)}$; a similar idea was used by Paroni [26]. Since we are interested in a bi-dimensional limit, we consider the average over the thickness of the cylinders of these measures μ_x . Then, under the weak* convergence of the averaged gradient Young measures we obtain the following limit energy

$$I(\nu) = \int_{\omega} \int_{M^{3 \times 2}} W_0(\bar{F}) d\nu_{x_\alpha}(\bar{F}) dx_\alpha - \int_{\omega} m_3 f \cdot y dx_\alpha,$$

whose domain is the space of bi-dimensional gradient Young measures satisfying the appropriate boundary conditions. The functionals I and J share the same infimum value, but there is a major difference between the two functionals: while J is incapable to capture the oscillations of the minimizing sequences, the functional I characterizes the microstructure through the Young measure solution. Indeed, the minimizers of I carry more information than those of J since the minimizers of I provide a description of the microstructure and their center of mass select minimizers of J (see Section 6). The computation of the microstructure in a thin film can therefore be evaluated through the use of the functional I exactly as is done in three-dimensional elasticity, see for instance Ball and James [5, 6], and Luskin [23].

Another important feature of the limit functional I , which is missing in all the other previously obtained variational problems, and which is relevant also by the point of view of the computation of the microstructure, is that it uniquely determines the energy density W_0 of the thin film (see Section 7). The total energy of the membrane, written without using Young measures, is therefore

$$E(y) = \int_{\omega} W_0(D_\alpha y) dx_\alpha - \int_{\omega} m_3 f \cdot y dx_\alpha.$$

Thin films of martensitic material have been studied also by Bhattacharya and James [10]. These authors have considered a body characterized not only by a free energy but also by an interfacial energy; indeed they have taken a total energy of the form

$$I_{\varepsilon, \kappa}(y) = \int_{\Omega_\varepsilon} W(Dy) + \kappa |D^2 y|^2 dx - \int_{\Omega_\varepsilon} \hat{f}^\varepsilon \cdot y dx,$$

where κ is a positive constant, W has a quadratic growth and $D^2 y$ denotes the Hessian of y . The limit problem

$$J_\kappa(y, b) = \int_{\omega} W(D_\alpha y | b) + \kappa (|D_\alpha^2 y|^2 + 2|D_\alpha b|^2) dx_\alpha - \int_{\omega} m_3 f \cdot y dx_\alpha$$

was obtained by them under the weak convergence in $W^{2,2}$ of the deformation to y and the weak convergence in $W^{1,2}$ to the Cosserat vector b of the third partial derivative of a suitable rescaling of the deformation. We note that the interfacial energy plays a crucial role in order to supply a limit problem in which no quasiconvexification appears. On the other hand, it has been shown by Shu [29] that letting also $\kappa \rightarrow 0$ the variational limit coincides with that obtained by Le Dret and Raoult.

Bélik and Luskin [7, 8] have observed that the deformations with finite energy cannot have sharp interfaces between compatible variants, unless the constant κ is set equal to 0. For this reason they set the problem in the framework of functions with bounded Hessian and considered an interfacial energy proportional to the total variation of the deformation gradient.

More recently Bocea [11] has studied the limit $\mathcal{W}(y, b)$ of I_ε under the weak convergence in $W^{1,p}$ of the deformations to y and the weak convergence in L^p to b of the third partial derivative of a suitable rescaling of the deformation. Using a result of Bocea and Fonseca [12], he has proved that

$$\mathcal{W}(y, b) = \inf \int_{\Omega \times M^{3 \times 3}} W d\mu$$

where the infimum is taken among all Young measures μ generated by sequences of gradients of deformations whose third partial derivative is suitably rescaled.

We conclude this brief review by mentioning also the papers [9, 13, 28], among the wide literature on thin films, and a related paper by the authors [18] where some of the ideas used in this paper are applied to obtain the energy

density of a martensitic string starting from a three-dimensional body with a free energy that becomes infinite when the volume locally vanishes.

The paper is organized as follows. In Sections 2 and 3 are summarized well known definitions and properties of Young measures and Γ -convergence. In Section 4 we consider the three-dimensional problems and set the framework in which the Γ -limit will be taken. Moreover we briefly recall the earlier result of Le Dret and Raoult. In Section 5 various preliminary results are proved and the main coerciveness and Γ -convergence theorems are only stated. The proofs are postponed to Section 8. In Section 6 we study the relationship between our limit functional and the one obtained by Le Dret and Raoult. The energy density of the thin film is proposed in Section 7.

2 Young measures preliminaries

The main references for this section are Ball [4], Müller [24] and Pedregal [27].

Throughout the whole section Ω is an open bounded subset of \mathbb{R}^k . We denote by $M^{h \times k}$ the set of $h \times k$ real matrices which will be often identified with the Euclidean space \mathbb{R}^{hk} . By $\mathcal{M}(M^{h \times k})$ we mean the space of \mathbb{R} -valued Borel measures on $M^{h \times k}$ which can be viewed as the dual of the separable Banach space $C_0(M^{h \times k})$ under the duality

$$\langle \mu, \varphi \rangle = \int_{M^{h \times k}} \varphi d\mu.$$

We recall that a mapping $\mu : \Omega \rightarrow \mathcal{M}(M^{h \times k})$ is said to be weakly* measurable whenever the function $x \mapsto \langle \mu(x), \varphi \rangle$ is measurable for every $\varphi \in C_0(M^{h \times k})$. The space $L_w^\infty(\Omega; \mathcal{M}(M^{h \times k}))$ consists of all weakly* measurable mappings $\mu : \Omega \rightarrow \mathcal{M}(M^{h \times k})$ which are essentially bounded. This space will be endowed with the weak* topology induced by the duality with $L^1(\Omega; C_0(M^{h \times k}))$. Therefore, a sequence μ^n is weakly* converging to a limit μ if and only if

$$\int_{\Omega} \langle \mu_x^n, \varphi \rangle g(x) dx \rightarrow \int_{\Omega} \langle \mu_x, \varphi \rangle g(x) dx, \quad \forall \varphi \in C_0(M^{h \times k}), \quad \forall g \in L^1(\Omega).$$

A Young measure on Ω with target space $M^{h \times k}$ is an element ν of the space $L_w^\infty(\Omega; \mathcal{M}(M^{h \times k}))$ such that $\nu_x := \nu(x)$ is a probability measure for almost every $x \in \Omega$.

A Young measure $\mu \in L_w^\infty(\Omega; \mathcal{M}(M^{h \times k}))$ is said to be generated by the sequence of measurable functions $u^n : \Omega \rightarrow M^{h \times k}$ if

$$\delta_{u^n(\cdot)} \rightarrow \mu \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(M^{h \times k})).$$

Every Young measure is generated by some sequence of measurable functions.

The center of mass of a Young measure $\mu \in L_w^\infty(\Omega; \mathcal{M}(M^{h \times k}))$ is the function

$$x \mapsto \langle \mu_x, \text{id} \rangle = \int_{M^{h \times k}} \lambda d\mu_x(\lambda),$$

where id denotes the identity mapping.

In this paper an important role will be played by the image of a measure and by the fiber product of two Young measures.

Let (X, \mathcal{A}, μ) and (Y, \mathcal{B}) be measurable spaces and $\theta : X \rightarrow Y$. Then the image of the measure μ under the mapping θ is defined by $\theta_{\#}\mu(A) := \mu(\theta^{-1}(A))$ for every $A \in \mathcal{B}$.

The fiber product of two Young measures $\nu \in L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^m))$ and $\sigma \in L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^s))$ is the Young measure $\mu \in L_w^\infty(\Omega; \mathcal{M}(M^{m \times s}))$ usually denoted with $\mu = \nu \otimes \sigma$ and defined by

$$\mu_x = \nu_x \otimes \sigma_x,$$

where \otimes denotes the usual tensor product of measures. The following theorem of Balder and Valadier (see Valadier [30], Theorem 14) gives a sufficient condition which allows to pass to the limit in the fiber product.

Theorem 2.1 *Let ν^n and σ^n be two sequences of Young measures belonging to $L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^m))$ and $L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^s))$ respectively, such that*

$$\nu^n \rightarrow \nu \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^m))$$

while

$$\sigma^n \rightarrow \sigma = \delta_{u(\cdot)} \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^s))$$

for a suitable measurable function u . Then

$$\nu^n \otimes \sigma^n \rightarrow \nu \otimes \sigma \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(\mathbb{R}^{m+s})).$$

A large use will be done of the following theorem (see for instance Müller [24], Corollary 3.3) which states the fundamental property of Young measures.

Theorem 2.2 *Suppose that the sequence of maps $u^n : \Omega \rightarrow \mathbb{R}^h$ generates the Young measure ν . Let $f : \Omega \times \mathbb{R}^h \rightarrow \mathbb{R}$ be a Carathéodory function and assume that the negative part $f^-(x, u^n(x))$ is weakly relatively compact in $L^1(\Omega)$. Then*

$$\liminf_{k \rightarrow \infty} \int_{\Omega} f(x, u^n(x)) dx \geq \int_{\Omega} \int_{\mathbb{R}^h} f(x, \lambda) d\nu_x(\lambda) dx.$$

If, in addition, the sequence of functions $x \mapsto |f|(x, u^n(x))$ is weakly relatively compact in $L^1(\Omega)$ then

$$f(\cdot, u^n(\cdot)) \rightarrow \bar{f} \text{ weakly in } L^1(\Omega) \text{ where } \bar{f}(x) = \int_{\mathbb{R}^h} f(x, \lambda) d\nu_x(\lambda).$$

We are mainly interested into those Young measures which are generated by sequences of gradients, in the sense specified by the following definition.

Definition 2.3 *A weakly* measurable map $\nu : \Omega \rightarrow \mathcal{M}(M^{h \times k})$ is a $W^{1,p}$ -gradient Young measure if there exists a sequence of maps $u^n : \Omega \rightarrow \mathbb{R}^h$ such that $u^n \rightarrow u$ weakly in $W^{1,p}(\Omega; \mathbb{R}^h)$ and $\delta_{Du^n(\cdot)} \rightarrow \nu$ weakly* in $L_w^\infty(\Omega; M^{h \times k})$. In this case, the $W^{1,p}$ -gradient Young measure ν is said to be generated by the sequence Du^n and u is called an underlying deformation for ν .*

Gradient Young measures are in a sort of duality with quasiconvex functions.

Definition 2.4 *A function $f : M^{h \times k} \rightarrow (-\infty, +\infty]$ is said to be quasiconvex if for every open and bounded subset A of $M^{h \times k}$ with negligible boundary (with respect to the Lebesgue measure) one has*

$$\int_A f(F + D\varphi) dx \geq \int_A f(F) dx = |A|f(F) \quad \forall \varphi \in C_0^1(A; \mathbb{R}^h),$$

whenever the integral on the left hand side exists ($|A|$ denotes the Lebesgue measure of the open set A).

The following characterization of $W^{1,p}$ -gradient Young measures is due to Kinderlehrer and Pedregal [21].

Theorem 2.5 *Let $p \in [1, +\infty)$. A weakly* measurable map $\nu : \Omega \rightarrow \mathcal{M}(M^{h \times k})$ is a $W^{1,p}$ -gradient Young measure if and only if $\nu_x \geq 0$ a.e. and the following three conditions hold*

- (i) $\int_{\Omega} \int_{M^{h \times k}} |F|^p d\nu_x(F) dx < +\infty;$
- (ii) $\langle \nu_x, \text{id} \rangle = Du(x), u \in W^{1,p}(\Omega; \mathbb{R}^h);$
- (iii) $\langle \nu_x, f \rangle \geq f(\langle \nu_x, \text{id} \rangle)$ for almost every x and all quasiconvex f with $|f|(F) \leq C(|F|^p + 1)$.

The following proposition, see for instance Pedregal [27], Lemma 8.3, sometimes called Zhang's Lemma, is a variant of a result obtained by Acerbi and Fusco [2].

Proposition 2.6 *Let $p \in [1, +\infty]$ and let ν be a $W^{1,p}$ -gradient Young measure. If $u_0 \in W^{1,p}(\Omega; \mathbb{R}^h)$ is an underlying deformation for ν then there exists a sequence $u^n \in C_0^\infty(\Omega; \mathbb{R}^h)$ such that $u^n \rightarrow 0$ weakly in $W_0^{1,p}(\Omega; \mathbb{R}^h)$, $|Du^n|^p$ are equi-integrable ($|Du^n|$ equi-bounded in the case $p = +\infty$) and $D(u_0 + u^n)$ generates ν .*

3 Sequential Γ -convergence

For the purposes of the paper we need of a kind of variational convergence which allows to treat families of functionals $F_\varepsilon : X \rightarrow \overline{\mathbb{R}} = (-\infty, +\infty]$ defined on a space which may be different from the domain of the limiting functional. It is a variant of De Giorgi's Γ -convergence, and has been introduced by Anzellotti, Baldo and Percivale in [3] in order to study dimension reduction problems in mechanics. Let X be a set, let (Y, τ) be a topological space and let $q : X \rightarrow Y$.

Given a sequence of functionals $F_n : X \rightarrow \overline{\mathbb{R}}$ and a point $y \in Y$, let us denote by

$$\begin{aligned} \Gamma(q, \tau Y) \liminf_{n \rightarrow \infty} F_n(y) &:= \inf \{ \liminf_{n \rightarrow \infty} F_n(x_n) : q(x_n) \xrightarrow{\tau} y \}, \\ \Gamma(q, \tau Y) \limsup_{n \rightarrow \infty} F_n(y) &:= \inf \{ \limsup_{n \rightarrow \infty} F_n(x_n) : q(x_n) \xrightarrow{\tau} y \}, \end{aligned} \quad (3.1)$$

which are called, respectively, the sequential Γ^- -lower and the Γ^- -upper limit at the point y . In (3.1), as in the sequel, we have dropped the minus sign for notational simplicity.

Definition 3.1 *Given a sequence ε_n of positive real numbers we say that a sequence $F_{\varepsilon_n} : X \rightarrow \overline{\mathbb{R}}$ (sequentially) $\Gamma(q, \tau Y)$ -converges to a functional $F : Y \rightarrow \overline{\mathbb{R}}$ at a point $y \in Y$, and we write*

$$\Gamma(q, \tau Y) \lim_{n \rightarrow \infty} F_{\varepsilon_n}(y) = F(y)$$

if

$$\Gamma(q, \tau Y) \liminf_{n \rightarrow \infty} F_{\varepsilon_n}(y) = \Gamma(q, \tau Y) \limsup_{n \rightarrow \infty} F_{\varepsilon_n}(y) = F(y). \quad (3.2)$$

We say that the family of functionals $F_\varepsilon : X \rightarrow \overline{\mathbb{R}}$ (sequentially) $\Gamma(q, \tau Y)$ -converges to a functional $F : Y \rightarrow \overline{\mathbb{R}}$ at a point $y \in Y$, and we write

$$\Gamma(q, \tau Y) \lim_{\varepsilon \rightarrow 0} F_\varepsilon(y) = F(y) \quad (3.3)$$

if for any sequence ε_n of positive reals such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ we have that

$$\Gamma(q, \tau Y) \lim_{n \rightarrow \infty} F_{\varepsilon_n}(y) = F(y).$$

We say that a family of functionals $\Gamma(q, \tau Y)$ -converges on a set if it $\Gamma(q, \tau Y)$ -converges at every point of the set.

Remark 3.2 *Let us remark that (3.2) holds if and only if the following two conditions are satisfied*

1. for every sequence $x_n \in X$ such that $q(x_n) \xrightarrow{\tau} y$ one has

$$\liminf_{n \rightarrow \infty} F_{\varepsilon_n}(x_n) \geq F(y);$$

2. there exists a sequence $\bar{x}_n \in X$ such that $q(\bar{x}_n) \xrightarrow{\tau} y$ and

$$\lim_{n \rightarrow \infty} F_{\varepsilon_n}(\bar{x}_n) = F(y).$$

If $X = Y$ and q is the identity map, then the Γ -limits defined above are the classical De Giorgi's sequential Γ^- -limits which will be denoted by the classical notation

$$\Gamma^-(\tau X) \lim_{\varepsilon \rightarrow 0} F_\varepsilon = F.$$

Moreover, it is easy to see that, setting

$$I_\varepsilon(y) = \inf\{F_\varepsilon(x) : q(x) = y\}, \quad \varepsilon > 0$$

then (3.3) holds true if and only if

$$\Gamma^-(\tau Y) \lim_{\varepsilon \rightarrow 0} I_\varepsilon(y) = F(y).$$

Besides the notion of sequential Γ -convergence, also a more general concept of Γ -convergence, associated with a topology on X , can be introduced (see for instance Dal Maso [17]) and the two kinds of Γ -convergence coincide on first countable topological spaces. Hence when the topological space (Y, τ) is first countable, and the family F_ε is equi-coercive then the $\Gamma(q, \tau Y)$ -convergence has the variational character which is typical of Γ -convergence, that is, it ensures lower semicontinuity to the limit which, moreover, turns out to be coercive when uniform coercivity assumptions are made on the family F_ε . Moreover, roughly speaking, it preserves convergence of minima and of minimizers. This is summarized by the following statements.

Definition 3.3 *The family F_ε is said to be $(q, \tau Y)$ -equi-coercive if for every real number M there exists a τ -compact and τ -closed subset K_M of Y such that*

$$\{q(x) : F_\varepsilon(x) \leq M\} \subseteq K_M \text{ for every } \varepsilon > 0.$$

The definition above reduces to the classical one of τ -equi-coerciveness in the case $q = \text{id}$.

Proposition 3.4 *Let us assume that $\Gamma(q, \tau Y) \lim_{\varepsilon \rightarrow 0} F_\varepsilon = F$ on Y and that the family F_ε be $(q, \tau Y)$ -equi-coercive. Then we have:*

- (i) F is τ -lower semicontinuous;
- (ii) F is τ -coercive;
- (iii) if $x_\varepsilon \in X$ satisfy $\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(x_\varepsilon) = \liminf_{\varepsilon \rightarrow 0} \inf F_\varepsilon$ (e.g. if x_ε minimizes F_ε) then
 - (a) if ε_n is a sequence such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ and if $q(x_{\varepsilon_n}) \xrightarrow{\tau} y$ then y is a minimizer of F on Y and $\lim_{n \rightarrow \infty} F_{\varepsilon_n}(x_{\varepsilon_n}) = F(y)$;
 - (b) there is a sequence ε_n such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ and a minimizer y of F on Y such that $q(x_{\varepsilon_n}) \xrightarrow{\tau} y$.

Proposition 3.5 *If Y is dual of a separable Banach space, τ is the weak* topology, and $F_n : X \rightarrow \overline{\mathbb{R}}$ is $(q, \tau Y)$ -equi-coercive, then*

$$\Gamma(q, \tau Y) \lim_{n \rightarrow \infty} F_n(y) = F(y)$$

if and only if

- (i) $\liminf_{n \rightarrow \infty} F_n(x_n) \geq F(y)$ for every sequence $x_n \in X$ such that $q(x_n) \xrightarrow{\tau} y$;
- (ii) for every sequence (n_k) of positive integers there is a subsequence (n_{k_p}) and a sequence $x_p \in X$ such that

$$q(x_p) \xrightarrow{\tau} y \quad \text{and} \quad \lim_{p \rightarrow \infty} F_{n_{k_p}}(x_p) = F(y).$$

PROOF. It is an easy consequence of the weak* metrizable of compact subsets of Y and of the Urysohn property of Γ -convergence (see for instance Dal Maso [17], Chapter 8). \square

4 Setting of the problem and previous results

For all $\varepsilon > 0$, let $\Omega_\varepsilon = \{x = (x_\alpha, x_3) \in \mathbb{R}^2 \times \mathbb{R} : x_\alpha \in \omega, |x_3| < \varepsilon/2\}$, where ω is an open, bounded subset of \mathbb{R}^2 with Lipschitz boundary. We can think of Ω_ε as the reference configuration of a hyperelastic homogeneous body with stored energy density function $W : M^{3 \times 3} \rightarrow \mathbb{R}$. This function W is assumed to be continuous and to satisfy the following growth assumption

$$c(|F|^p - 1) \leq W(F) \leq C(|F|^p + 1), \quad (4.1)$$

for two constants $C \geq c > 0$. The total energy I_ε of the body is given by

$$I_\varepsilon(y) = \int_{\Omega_\varepsilon} W(Dy) dx - \int_{\Omega_\varepsilon} \hat{f}^\varepsilon \cdot y dx,$$

where the body force density \hat{f}^ε is taken in $L^q(\Omega_\varepsilon; \mathbb{R}^3)$, with $1/p + 1/q = 1$. Assuming, for the sake of simplicity, the body to be clamped on the lateral surface $\Gamma_\varepsilon := \partial\omega \times (-\varepsilon/2, \varepsilon/2)$ of Ω_ε , the equilibrium configuration will be found by minimizing the energy I_ε over the set of admissible deformations

$$\mathcal{A}_\varepsilon = \{y \in W^{1,p}(\Omega_\varepsilon; \mathbb{R}^3) : y(x) = x \text{ on } \Gamma_\varepsilon\}.$$

In order to study the behaviour of the minimizers y^ε of I_ε as $\varepsilon \rightarrow 0$, it is convenient to transform, after Ciarlet and Destuynder [15], the problems under consideration into problems over a fixed domain. To this end we define $\Omega := \Omega_1$, $\Gamma := \Gamma_1$ and a map $\theta_\varepsilon : L^p(\Omega_\varepsilon; \mathbb{R}^3) \rightarrow L^p(\Omega; \mathbb{R}^3)$ by

$$(\theta_\varepsilon y)(x) = y(x_1, x_2, \varepsilon x_3).$$

We accordingly rescale the energies by setting $I_\varepsilon^\Omega(y) = I_\varepsilon(\theta_\varepsilon^{-1}y)/\varepsilon$, which is

$$I_\varepsilon^\Omega(y) := \int_\Omega W(D_\alpha y | \frac{D_3 y}{\varepsilon}) dx - \int_\Omega f^\varepsilon \cdot y dx,$$

where we have set $f^\varepsilon := \theta_\varepsilon \hat{f}^\varepsilon$. Following Le Dret and Raoult [22] we assume that f^ε does not depend on ε and set $f := f^\varepsilon$. The admissible set over which the total energy has to be minimized becomes

$$\mathcal{A}_\varepsilon^\Omega = \{y \in W^{1,p}(\Omega; \mathbb{R}^3) : y(x) = (x_1, x_2, \varepsilon x_3) \text{ on } \Gamma\}.$$

Above and in the sequel, we identify $M^{3 \times 3}$ with $M^{3 \times 2} \times \mathbb{R}^3$, and write $(\bar{F}|z)$ for the element in $M^{3 \times 3}$ which is identified with $(\bar{F}, z) \in M^{3 \times 2} \times \mathbb{R}^3$. Moreover, $D_\alpha y$ denotes the first two columns of the gradient of y .

Within a similar framework, Le Dret and Raoult ([22], Theorem 2) have proved that, as $\varepsilon \rightarrow 0$, the family of functionals

$$J_\varepsilon(y) = \begin{cases} I_\varepsilon^\Omega(y) & \text{if } y \in \mathcal{A}_\varepsilon^\Omega \\ +\infty & \text{otherwise in } L^p(\Omega; \mathbb{R}^3) \end{cases}$$

Γ -converges in the strong topology of $L^p(\Omega; \mathbb{R}^3)$ to the functional

$$J(y) = \begin{cases} \int_\omega QW_0(D_\alpha y) dx_\alpha - \int_\omega m_3 f \cdot y dx_\alpha & \text{if } y \in W_\Gamma^{1,p}(\omega; \mathbb{R}^3) \\ +\infty & \text{otherwise in } L^p(\Omega; \mathbb{R}^3). \end{cases} \quad (4.2)$$

Here

$$W_\Gamma^{1,p}(\omega; \mathbb{R}^3) := \{y \in W^{1,p}(\omega; \mathbb{R}^3) : y(x) = (x_1, x_2, 0) \text{ on } \partial\omega\},$$

$m_3 f(x_\alpha) = \int_{-1/2}^{1/2} f(x_\alpha, x_3) dx_3$ denotes the integral mean value of f with respect to the third variable and QW_0 denotes the quasiconvex envelope of the continuous function

$$W_0(\bar{F}) := \min\{W(\bar{F}|z) : z \in \mathbb{R}^3\}, \quad \bar{F} \in M^{3 \times 2}, \quad (4.3)$$

which satisfies the same growth conditions of W . A similar result in the case of strings was previously obtained by Acerbi, Buttazzo and Percivale in [1].

With the motivation already explained in the introduction we extend the functionals I_ε^Ω to the larger space $L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$ by setting

$$I_\varepsilon^\mathcal{M}(\mu) = \begin{cases} I_\varepsilon^\Omega(y) & \text{if } \exists y \in \mathcal{A}_\varepsilon^\Omega : \mu = \delta_{Dy(\cdot)} \\ +\infty & \text{otherwise in } L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3})). \end{cases} \quad (4.4)$$

Let us remark that this functional is well defined because if $y_1, y_2 \in \mathcal{A}_\varepsilon^\Omega$ are such that $\delta_{Dy_1(x)} = \delta_{Dy_2(x)}$ then, due to the boundary condition, $y_1 = y_2$ almost everywhere.

5 The limit problem

Let us denote by $\bar{\pi}$ and π^3 the projections of $M^{3 \times 3}$ onto $M^{3 \times 2}$ and \mathbb{R}^3 , respectively, that is

$$\begin{aligned}\bar{\pi} : M^{3 \times 3} &\rightarrow M^{3 \times 2}, & \bar{\pi}(\bar{F}|z) &= \bar{F}, \\ \pi^3 : M^{3 \times 3} &\rightarrow \mathbb{R}^3, & \pi^3(\bar{F}|z) &= z,\end{aligned}$$

and by $\bar{\pi}_\# \mu_x$ and $\pi^3_\# \mu_x$ the usual image measures by the corresponding projection maps (see Section 2).

We further denote by $\mathcal{Y}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ the set of $W^{1,p}(\Omega; \mathbb{R}^3)$ -gradient Young measures with an underlying deformation y with $y(x) = (x_1, x_2, 0)$ on Γ , and by $\mathring{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ the subset of $\mathcal{Y}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ of Young measures μ with $\pi^3_\# \mu = \delta_0$, that is

$$\mathring{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3) := \{\mu \in \mathcal{Y}_\Gamma^{1,p}(\Omega; \mathbb{R}^3) : \pi^3_\# \mu = \delta_0\}.$$

The following lemma states that the sequence of functionals $I_\varepsilon^\mathcal{M}$ is equicoercive in $\mathring{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ with the weak* convergence of $L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$.

Lemma 5.1 *Let $\varepsilon_n \rightarrow 0$ and $\mu^n \in L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$ be such that*

$$\sup_n I_{\varepsilon_n}^\mathcal{M}(\mu^n) < +\infty.$$

Then there exists $\mu \in \mathring{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ and a subsequence (μ^{n_k}) such that

$$\mu^{n_k} \rightarrow \mu \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3})).$$

Moreover, denoted by y^{n_k} and y the underlying deformations of μ^{n_k} and μ which satisfy the boundary conditions $y^{n_k}(x) = (x_1, x_2, \varepsilon_{n_k} x_3)$ and $y(x) = (x_1, x_2, 0)$, respectively, we have

$$y^{n_k} \rightarrow y \text{ weakly in } W^{1,p}(\Omega; \mathbb{R}^3) \text{ and } D_3 y = 0.$$

PROOF. Since $\sup_n I_{\varepsilon_n}^\mathcal{M}(\mu^n) < +\infty$, then $\mu_x^n = \delta_{Dy^n(x)}$, $y^n \in W^{1,p}(\Omega; \mathbb{R}^3)$ and $y^n(x) = (x_1, x_2, \varepsilon_n x_3)$ on Γ . By the growth assumption on W and Poincaré inequality, we get, for every n large enough so that $\varepsilon_n < 1$

$$\begin{aligned}I_{\varepsilon_n}^\mathcal{M}(\mu^n) &= \int_\Omega W(D_\alpha y^n | \frac{D_3 y^n}{\varepsilon_n}) dx - \int_\Omega f \cdot y^n dx \\ &\geq c \|(D_\alpha y^n | \frac{D_3 y^n}{\varepsilon_n})\|_p^p - \|f\|_q \|y^n\|_p - c|\Omega| \\ &\geq c \|Dy^n\|_p^p - \|f\|_q \|y^n\|_p - c|\Omega| \\ &\geq C_1 \|y^n\|_{1,p}^p - C_2,\end{aligned}$$

where $\|\cdot\|_p$ and $\|\cdot\|_{1,p}$ denote the usual L^p and $W^{1,p}$ norms, respectively. Therefore

$$\sup_n \|y^n\|_{1,p} < +\infty \quad \text{and} \quad \sup_n \|(D_\alpha y^n | \frac{D_3 y^n}{\varepsilon_n})\|_p < +\infty$$

thus, up to subsequences,

$$y^n \rightarrow y \text{ weakly in } W^{1,p}(\Omega; \mathbb{R}^3) \quad \text{and} \quad D_3 y^n \rightarrow 0 \text{ strongly in } L^p(\Omega; \mathbb{R}^3)$$

and

$$\mu^n \rightarrow \mu \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3})).$$

It follows that $y = y(x_1, x_2)$ and $y(x) = (x_1, x_2, 0)$ on Γ and $\pi_{\#}^3 \mu = \delta_0$. Hence $\mu \in \mathring{\mathcal{Y}}_\Gamma^{1,p}$. \square

If $\mu \in \mathring{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ then it can be proved (see Lemma 5.3 below) that the center of mass of μ_x does not depend on the x_3 variable. With a subsequent example we are going to show that, in general, μ_x does depend also on x_3 . This is the main motivation which will lead us to study the Γ -convergence with respect to the weak* convergence of the averages with respect to the third variable.

Lemma 5.2 *If μ is a $W^{1,p}(\Omega; \mathbb{R}^3)$ -gradient Young measure generated by the sequence of gradients Dy^ε , and ν and γ are the Young measures generated by $D_\alpha y^\varepsilon$ and $D_3 y^\varepsilon$, respectively, then*

$$\bar{\pi}_{\#} \mu = \nu \quad \text{and} \quad \pi_{\#}^3 \mu = \gamma.$$

PROOF. The proof is a direct consequence of the definitions and of the classical integration formula with respect to an image measure. Indeed, since $D_\alpha y^\varepsilon = \bar{\pi}(Dy^\varepsilon)$ then, for every $\varphi \in C_0(M^{3 \times 2})$ and $g \in L^1(\Omega)$, we have

$$\begin{aligned} \int_{\Omega} \langle \bar{\pi}_{\#} \mu_x, \varphi \rangle g(x) dx &= \int_{\Omega} \int_{M^{3 \times 2}} \varphi(\bar{F}) d\bar{\pi}_{\#} \mu_x(\bar{F}) g(x) dx \\ &= \int_{\Omega} \int_{M^{3 \times 3}} \varphi(\bar{\pi}(F)) d\mu_x(F) g(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \varphi(\bar{\pi}(Dy^\varepsilon)) g(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \varphi(D_\alpha y^\varepsilon) g(x) dx = \int_{\Omega} \langle \nu_x, \varphi \rangle g(x) dx. \end{aligned}$$

The other relation can be proved similarly. \square

Lemma 5.3 *If $\mu \in L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$, with $\pi_{\#}^3 \mu = \delta_0$, is generated by the sequence of gradients Dy^ε with underlying deformation y , and ν is the Young measure generated by $D_\alpha y^\varepsilon$, then*

$$\mu = \nu \otimes \delta_0.$$

Moreover, the function y and the center of mass of the measures μ_x and ν_x , for almost every $x \in \Omega$, do not depend on the variable x_3 . In particular, with a small abuse of notation, we shall write

$$\langle \mu_x, \text{id} \rangle = Dy(x_\alpha), \quad \langle \nu_x, \text{id} \rangle = D_\alpha y(x_\alpha).$$

PROOF. From $\pi_{\#}^3 \mu = \delta_0$, and the previous lemma, we deduce that $D_3 y^\varepsilon$ generates the Young measure δ_0 . The first part of the statement follows by applying Theorem 2.1. Moreover, we have

$$Dy(x) = \langle \mu_x, \text{id} \rangle = \langle \nu_x \otimes \delta_0, \text{id} \rangle = \int_{M^{3 \times 3}} (\bar{F}|z) d\nu_x \otimes \delta_0(\bar{F}|z) = \langle \nu_x, \text{id} \rangle | 0,)$$

for almost every $x \in \Omega$. Thus $D_3 y = 0$, hence $y = y(x_\alpha)$ and $\langle \nu_{(x_\alpha, x_3)}, \text{id} \rangle = D_\alpha y(x_\alpha)$. \square

Example 5.4 We provide an example of a $\mu \in \mathfrak{Y}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ such that μ_x depends also on x_3 .

Let $\rho(s)$ be the 2-periodic function equaling s on $[0, 1]$ and $2-s$ on $[1, 2]$. Let $z^\varepsilon(x) := (\varepsilon x_3 \rho(x_1/\varepsilon), 0, 0)$. Then $D_3 z^\varepsilon \rightarrow 0$ in $L^\infty(\Omega; \mathbb{R}^3)$. Moreover, since

$$D_1 z_1^\varepsilon(x) = x_3 \rho'(\frac{x_1}{\varepsilon}) \text{ generates the Young measure } \frac{1}{2} \delta_{-x_3} + \frac{1}{2} \delta_{x_3},$$

we have that

$$D_\alpha z^\varepsilon \text{ generates the Young measure } \frac{1}{2} \delta_{-x_3 \bar{E}^{11}} + \frac{1}{2} \delta_{x_3 \bar{E}^{11}},$$

where \bar{E}^{11} is the matrix in $M^{3 \times 2}$ with element of place 11 equal to 1 and all the other elements equal to zero. It is now easy to check that the sequence of functions $y^\varepsilon(x) := (x_1, x_2, \varepsilon x_3) + z^\varepsilon(x) \min\{1, \text{dist}(x, \partial\Omega)/\varepsilon\}$ are in $\mathcal{A}_\varepsilon^\Omega$ and they generate a $\mu \in \mathfrak{Y}_\Gamma^{1,p}$ which depends also on x_3 .

The example above motivates the following definition.

Definition 5.5 Let X be either $M^{3 \times 2}$ or $M^{3 \times 3}$. If $\nu \in L_w^\infty(\Omega; \mathcal{M}(X))$ we define

$$\text{Av}^3 \nu : \omega \rightarrow \mathcal{M}(X),$$

the average of ν with respect to the variable x_3 , by

$$\langle \text{Av}_{x_\alpha}^3 \nu, \varphi \rangle := \langle \text{Av}^3 \nu(x_\alpha), \varphi \rangle := \int_{-1/2}^{1/2} \langle \nu_{(x_\alpha, x_3)}, \varphi \rangle dx_3,$$

for every $\varphi \in C_0(X)$.

By Fubini's theorem the map $x_\alpha \mapsto \langle \text{Av}_{x_\alpha}^3 \nu, \varphi \rangle$ is measurable, and since it is also essentially bounded, we can think of Av^3 as a mapping

$$\text{Av}^3 : L_w^\infty(\Omega; \mathcal{M}(X)) \rightarrow L_w^\infty(\omega; \mathcal{M}(X)).$$

It is easy to check that this map is continuous, which is

$$\begin{aligned} \mu^\varepsilon &\rightarrow \mu \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(X)) \\ &\downarrow \\ \text{Av}^3 \mu^\varepsilon &\rightarrow \text{Av}^3 \mu \text{ weakly* in } L_w^\infty(\omega; \mathcal{M}(X)), \end{aligned}$$

and that Av^3 maps Young measures into Young measures.

The Γ -convergence result will be made more expressive by using the average-projection mapping

$$q : L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3})) \rightarrow L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2})),$$

defined by

$$q = \bar{\pi}_\# \circ \text{Av}^3 = \text{Av}^3 \circ \bar{\pi}_\#,$$

where the commutativity of composition comes directly from the definitions. By the continuity of the average and the projection mapping, even q is continuous.

We are now in a position to state the main results of this section, whose proofs are postponed to Section 8.

We denote by $\mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$ the subset of $W^{1,p}(\omega; \mathbb{R}^3)$ -gradient Young measures with an underlying deformation y with $y(x) = (x_1, x_2, 0)$ on $\partial\omega$.

The following theorems hold under the continuity and growth assumptions made on W at the beginning of Section 4.

Theorem 5.6 *The family of energy functionals $I_\varepsilon^{\mathcal{M}}$, already defined in (4.4), is $(q, w^* L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2})))$ -equi-coercive (see Definition 3.3).*

Theorem 5.7 *Let $I_\varepsilon^{\mathcal{M}}$ be the family of functionals defined in (4.4). Then*

$$\Gamma(q, w^* L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))) \lim_{\varepsilon \rightarrow 0} I_\varepsilon^{\mathcal{M}}(\nu) = I(\nu)$$

with

$$I(\nu) = \begin{cases} \int_\omega \langle \nu_{x_\alpha}, W_0 \rangle dx_\alpha - \int_\omega m_3 f \cdot y dx_\alpha & \text{if } \nu \in \mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3) \\ +\infty & \text{otherwise in } L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2})) \end{cases} \quad (5.1)$$

where W_0 is the function already defined in (4.3), and $y \in W^{1,p}(\omega; \mathbb{R}^3)$ is the underlying deformation of ν with boundary condition $y = (x_1, x_2, 0)$ on $\partial\omega$.

By the properties of Γ -limits summarized in Proposition 3.4 we can state the following theorem which makes precise the variational character of the limit energy $I(\nu)$.

Theorem 5.8 *Let I be the functional defined in Theorem 5.7. We have:*

- (i) *I is weakly* lower semicontinuous and weakly* coercive; hence it admits minimum on the space $\mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$;*
- (ii) *if $y^\varepsilon \in W^{1,p}(\Omega; \mathbb{R}^3)$ satisfies the boundary condition $y^\varepsilon(x) = (x_1, x_2, \varepsilon x_3)$ on Γ and $\liminf_{\varepsilon \rightarrow 0} I_\varepsilon^\Omega(y^\varepsilon) = \liminf_{\varepsilon \rightarrow 0} \inf I_\varepsilon^\Omega$ (e.g. if y^ε minimizes I_ε^Ω) then*
 - (a) *if $\varepsilon_n \rightarrow 0$ and if $q(\delta_{D y^{\varepsilon_n}}) \rightarrow \nu$ weakly* in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$ then ν is a minimizer of I on $\mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$ and $\lim_{n \rightarrow \infty} I_{\varepsilon_n}^\Omega(y^{\varepsilon_n}) = I(\nu)$;*
 - (b) *there is a sequence $\varepsilon_n \rightarrow 0$ and a minimizer ν of I on $\mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$ such that $q(\delta_{D y_n^\varepsilon}) \rightarrow \nu$ weakly* in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$.*

6 Relationship with Le Dret and Raoult theory

Let J be the Γ -limit functional of Le Dret and Raoult introduced in (4.2).

Theorem 6.1 *Let $y \in L^p(\Omega; \mathbb{R}^3)$. Then*

$$J(y) = \inf I(\nu) \tag{6.1}$$

where the infimum is taken over all $\nu \in L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$ which satisfy $\langle \nu, \text{id} \rangle = D_\alpha y$ and $y(x) = (x_1, x_2, 0)$ on $\partial\omega$, with the usual convention $\inf \emptyset = +\infty$. Moreover the infimum is attained.

PROOF. Let $y \in L^p(\Omega; \mathbb{R}^3)$. Let us first of all observe that the infimum is attained as a consequence of item (i) of Theorem 5.8 and the continuity of the center of mass with respect to the weak* convergence in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$.

For convenience we denote the right hand side of (6.1) by $\tilde{I}(y)$.

Let us start by proving the inequality $J(y) \leq \tilde{I}(y)$. To this end we can assume that $\tilde{I}(y)$ be finite, that is there exists $\nu \in L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$ such that $I(\nu) = \tilde{I}(y)$ and y satisfies the boundary value problem $D_\alpha y = \langle \nu, \text{id} \rangle$, $y(x) = (x_1, x_2, 0)$ on Γ . Hence $y = y(x_\alpha)$. Then, since $I(\nu) < +\infty$, we have $\nu \in \mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$ and therefore $y \in W_\Gamma^{1,p}(\omega; \mathbb{R}^3)$. In this case

$$J(y) = \int_\omega QW_0(D_\alpha y) dx_\alpha - \int_\omega m_3 f \cdot y dx_\alpha,$$

and, for (iii) of Theorem 2.5, we have that

$$QW_0(D_\alpha y) \leq \langle \nu, W_0 \rangle,$$

which implies that $J(y) \leq I(\nu) = \tilde{I}(y)$.

In order to prove the converse inequality, it is convenient to use the fact that the involved functionals are Γ -limits. As

$$J(y) = \Gamma^-(L^p(\Omega; \mathbb{R}^3)) \lim_{\varepsilon \rightarrow 0} J_\varepsilon(y)$$

then there exist sequences $\varepsilon_n \rightarrow 0$ and $y^n \rightarrow y$ weakly in $W^{1,p}(\Omega; \mathbb{R}^3)$ such that

$$\lim_{n \rightarrow \infty} J_{\varepsilon_n}(y^n) = J(y).$$

Since the only interesting case is $J(y) < +\infty$, by coercivity and possibly passing to subsequences, we can assume that

- (Dy^n) generates a $W^{1,p}(\Omega; \mathbb{R}^3)$ -Young measure μ with underlying deformation y ,
- $(D_\alpha y^n)$ generates a Young measure ν_Ω ,
- $D_3 y^n \rightarrow D_3 y = 0$ strongly in L^p , hence $D_3 y_n$ generates the measure δ_0 .

By Lemma 5.2 follows that $\bar{\pi}_{\#}\mu = \nu_{\Omega}$ and $\pi_{\#}^3\mu = \delta_0$, so $\mu \in \mathring{\mathcal{Y}}_{\Gamma}^{1,p}(\Omega; \mathbb{R}^3)$ by definition. An application of Lemma 5.3 gives $\mu = \nu_{\Omega} \otimes \delta_0$ and $\langle \nu_{\Omega}, \text{id} \rangle = D_{\alpha}y$. Let $\mu^n = \delta_{Dy^n}$ and $\nu := q(\mu) = \text{Av}^3(\nu_{\Omega})$. Then $\nu \in \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)$ and $\langle \nu, \text{id} \rangle = D_{\alpha}y$. From the continuity of q

$$q(\mu^n) \rightarrow \nu \text{ weakly* in } L_w^{\infty}(\omega; \mathcal{M}(M^{3 \times 2}))$$

and by the Γ -convergence theorem 5.7 and the fact that $J_{\varepsilon_n}(y_n) = I_{\varepsilon_n}^{\mathcal{M}}(\mu^n)$, we obtain

$$J(y) = \lim_{n \rightarrow \infty} J_{\varepsilon_n}(y^n) = \liminf_{n \rightarrow \infty} I_{\varepsilon_n}^{\mathcal{M}}(\mu^n) \geq I(\nu) \geq \tilde{I}(y),$$

which concludes the proof. \square

The relationship between the minimizers of the two functionals I and J is made explicit in the following theorem.

Theorem 6.2 *The following properties holds for the functionals I and J .*

- (i) $\min\{J(y) : y \in W_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)\} = \min\{I(\nu) : \nu \in \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)\}$.
- (ii) *Let ν be a minimizer for I . If $y \in W_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)$ satisfies $D_{\alpha}y(x_{\alpha}) = \langle \nu_{x_{\alpha}}, \text{id} \rangle$ for almost every $x_{\alpha} \in \omega$, then y is a minimizer for J and $QW_0(D_{\alpha}y(x_{\alpha})) = \langle \nu_{x_{\alpha}}, W_0 \rangle$ for almost every $x_{\alpha} \in \omega$.*
- (iii) *Let y be a minimizer for J . If $\nu \in \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)$ satisfies $D_{\alpha}y(x_{\alpha}) = \langle \nu_{x_{\alpha}}, \text{id} \rangle$ and $QW_0(D_{\alpha}y(x_{\alpha})) = \langle \nu_{x_{\alpha}}, W_0 \rangle$ for almost every $x_{\alpha} \in \omega$, then ν is a minimizer for I .*

PROOF. It follows by noticing that J and I are the relaxed functionals of the extension by $+\infty$ of

$$E(y) = \int_{\omega} W_0(D_{\alpha}y) dx_{\alpha} - \int_{\omega} m_3 f \cdot y dx_{\alpha}$$

with respect to the strong topology of $L^p(\omega; \mathbb{R}^3)$ and the weak* topology of the space $L_w^{\infty}(\omega; \mathcal{M}(M^{3 \times 2}))$, respectively, and applying Theorem 4.4 and Corollary 4.6 of Pedregal [27]. \square

7 The energy density of the thin film

The aim of this section is to propose and motivate an energy density for the thin film. Indeed we shall deduce that W_0 and

$$E(y) = \int_{\omega} W_0(D_{\alpha}y) dx_{\alpha} - \int_{\omega} m_3 f \cdot y dx_{\alpha}$$

are the energy density and the total energy of the thin film, respectively.

To make our considerations we need to work with a general linear boundary condition $y(x) = Bx$ on Γ_ε , where B is a linear map from \mathbb{R}^3 into itself. The right Sobolev space to treat this case is

$$W_{\Gamma_B}^{1,p}(\omega; \mathbb{R}^3) := \{y \in W^{1,p}(\omega; \mathbb{R}^3) : y(x_\alpha) = \bar{B}x_\alpha \text{ on } \partial\omega\},$$

where $\bar{B} \in M^{3 \times 2}$ denotes the first two columns of the matrix representing B . It is trivial to check that in this setting the variational limit is the functional I defined in (5.1) with $\mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$ replaced, with the obvious meaning of the notation, by $\mathcal{Y}_{\Gamma_B}^{1,p}(\omega; \mathbb{R}^3)$, and the corresponding limit functional is denoted by I_B . Similarly, also the case studied by Le Dret and Raoult can be put into this framework and the limit functional is defined on $W_{\Gamma_B}^{1,p}(\omega; \mathbb{R}^3)$ and denoted by J_B .

For every continuous function $Z : M^{3 \times 2} \rightarrow \mathbb{R}$ satisfying the growth condition of order p , $c(|\bar{F}|^p - 1) \leq Z(\bar{F}) \leq C(1 + |\bar{F}|^p)$ for $0 < c < C$, we introduce the energy

$$E_Z(y) = \int_\omega Z(D_\alpha y) dx_\alpha - \int_\omega m_3 f \cdot y dx_\alpha$$

and denote by $Q[E_Z, \Gamma_B]$ the relaxed functional, that is the lower semicontinuous envelope, of E_Z on the space $W_{\Gamma_B}^{1,p}(\omega; \mathbb{R}^3)$ with respect to the weak topology. With $Y[E_Z, \Gamma_B]$ we mean the relaxation of

$$E_Z^\infty(\nu) = \begin{cases} E_Z(y) & \text{if } \nu = \delta_{D_\alpha y} : y \in W_{\Gamma_B}^{1,p}(\omega; \mathbb{R}^3) \\ +\infty & \text{otherwise in } L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2})) \end{cases}$$

with respect to the weak* topology of $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$.

As briefly noticed in the last proof, the limit functional J_B , obtained by Le Dret and Raoult, and the functional I_B , obtained by us, can be written, see Dacorogna [16] and, for instance, Murat and Tartar [25], as

$$J_B(y) = Q[E_{W_0}, \Gamma_B](y), \quad I_B(\nu) = Y[E_{W_0}, \Gamma_B](\nu).$$

Theorem 7.1 *The function W_0 is the unique continuous integrand satisfying a growth condition of order p such that $I_B = Y[E_{W_0}, \Gamma_B]$ for every linear boundary condition, which is, for every linear map B .*

PROOF. Let W_1 be one of these functions such that $I_B = Y[E_{W_1}, \Gamma_B]$ for every linear B . Then

$$\int_\omega \langle \nu_{x_\alpha}, W_0 - W_1 \rangle dx_\alpha = 0 \quad \text{for every } \nu \in \mathcal{Y}_{\Gamma_B}^{1,p}(\omega; \mathbb{R}^3),$$

and for every B . It follows that $W_0 = W_1$ by simply taking the admissible measure $\nu = \delta_{\bar{B}}$ where $\bar{B} \in M^{3 \times 2}$ are the first two columns of the matrix representing B . \square

Hence, the variational problem I determines a unique energy density W_0 . This selection is not provided by the relation $J_B = Q[W_0, \Gamma_B]$; indeed, in general, there exists an infinite number of functions W_1 such that $J_B = Q[W_0, \Gamma_B] = Q[W_1, \Gamma_B]$ for every B .

A further reason to take W_0 as the energy density of the thin film follows from the relation between $E = E_{W_0}$ and I , which from well known properties of relaxed functionals can be expressed in terms of correspondence between minimizing sequences of E and minimizers of I . In particular, minimizing sequences of E generate Young measure which are minimizers of I , and viceversa.

In Le Dret and Raoult [22] it is shown that if the energy W satisfies the principle of material frame indifference, then also W_0 is frame indifferent, moreover the symmetry relations of W_0 are derived from those of W .

8 Proofs of the main results of Section 5

Before proving the main theorems, we state some technical results.

Lemma 8.1 *Let $\nu \otimes \delta_0$ be a $W^{1,p}(\Omega; \mathbb{R}^3)$ -gradient Young measure. Then $\text{Av}^3 \nu$ is a $W^{1,p}(\omega; \mathbb{R}^3)$ -gradient Young measure, and $\langle \text{Av}_{x_\alpha}^3 \nu, \text{id} \rangle = \langle \nu_{(x_\alpha, x_3)}, \text{id} \rangle$ for almost every $x \in \Omega$. Moreover, for every $W^{1,p}(\omega; \mathbb{R}^3)$ -gradient Young measure $\bar{\mu}$, there exists a $W^{1,p}(\Omega; \mathbb{R}^3)$ -gradient Young measure $\mu \otimes \delta_0$ such that $\bar{\mu} = \text{Av}^3 \mu$.*

PROOF. To show that $\text{Av}^3 \nu$ is a $W^{1,p}(\omega; \mathbb{R}^3)$ -gradient Young measure we may use the characterization given by Kinderlehrer and Pedregal, see Theorem 2.5. Thus it suffices to show that

- (i) $\int_\omega \int_{M^{3 \times 2}} |\bar{F}|^p d\text{Av}_{x_\alpha}^3 \nu(\bar{F}) dx_\alpha < +\infty$;
- (ii) $\langle \text{Av}_{x_\alpha}^3 \nu, \text{id} \rangle = D_\alpha y(x_\alpha)$ and $y \in W^{1,p}(\omega; \mathbb{R}^3)$;
- (iii) $\langle \text{Av}_{x_\alpha}^3 \nu, f \rangle \geq f(\langle \text{Av}_{x_\alpha}^3 \nu, \text{id} \rangle)$ for almost every x_α and all quasiconvex f with $|f|(\bar{F}) \leq C(|\bar{F}|^p + 1)$.

Property (i) holds since

$$\begin{aligned} \int_\omega \int_{M^{3 \times 2}} |\bar{F}|^p d\text{Av}_{x_\alpha}^3 \nu(\bar{F}) dx_\alpha &= \int_\omega \int_{-1/2}^{1/2} \int_{M^{3 \times 2}} |\bar{F}|^p d\nu_{(x_\alpha, x_3)}(\bar{F}) dx_3 dx_\alpha \\ &= \int_\Omega \int_{M^{3 \times 2}} |(\bar{F}|z)|^p d\nu_x \otimes \delta_0(\bar{F}|z) dx < +\infty. \end{aligned}$$

Before proving property (ii) let us note that if $y \in W^{1,p}(\Omega; \mathbb{R}^3)$ is an underlying deformation for $\nu \otimes \delta_0$, then, see Lemma 5.3, $y = y(x_\alpha)$, and $\langle \nu_{(x_\alpha, x_3)}, \text{id} \rangle = D_\alpha y(x_\alpha)$. From this consideration it follows that

$$\langle \text{Av}_{x_\alpha}^3 \nu, \text{id} \rangle = \int_{-1/2}^{1/2} \langle \nu_{(x_\alpha, x_3)}, \text{id} \rangle dx_3 = D_\alpha y(x_\alpha), \quad (8.1)$$

hence property (ii) holds and $\langle \text{Av}_{x_\alpha}^3 \nu, \text{id} \rangle = \langle \nu_{(x_\alpha, x_3)}, \text{id} \rangle$. Let $f : M^{3 \times 2} \rightarrow \mathbb{R}$ be a quasiconvex function with $|f(\bar{F})| \leq C(|\bar{F}|^p + 1)$. That is, for every open and bounded set $U \subset \mathbb{R}^2$ with $|\partial U| = 0$ one has

$$\int_U f(\bar{F} + D_\alpha \varphi) dx_\alpha \geq f(\bar{F})|U|,$$

for all $\varphi \in C_0^1(U; \mathbb{R}^3)$ and all $\bar{F} \in M^{3 \times 2}$. Let us consider $f^\# : M^{3 \times 3} \rightarrow \mathbb{R}$ be defined by $f^\#(\bar{F}|z) := f(\bar{F})$. Then for every $F = (\bar{F}|z) \in M^{3 \times 3}$ and $\psi \in C_0^1(U \times (-1/2, 1/2); \mathbb{R}^3)$ we have

$$\begin{aligned} \int_{U \times (-\frac{1}{2}, \frac{1}{2})} f^\#(F + D\psi) dx &= \int_{-1/2}^{1/2} \int_U f(\bar{F} + D_\alpha \psi(x_\alpha, x_3)) dx_\alpha dx_3 \\ &\geq \int_{-1/2}^{1/2} f(\bar{F})|U| dx_3 = f^\#(\bar{F}|z)|U| = f^\#(F)|U|, \end{aligned}$$

where we have used the fact that for almost every x_3 , the function $\psi(\cdot, x_3) \in C_0^1(U, \mathbb{R}^3)$. Thus also $f^\#$ is quasiconvex. Hence, we have

$$\begin{aligned} \langle \text{Av}_{x_\alpha}^3 \nu, f \rangle &= \int_{-1/2}^{1/2} \int_{M^{3 \times 2}} f(\bar{F}) d\nu_{(x_\alpha, x_3)}(\bar{F}) dx_3 \\ &= \int_{-1/2}^{1/2} \int_{M^{3 \times 3}} f^\#(\bar{F}|z) d\nu_{(x_\alpha, x_3)} \otimes \delta_0(\bar{F}|z) dx_3 \\ &= \int_{-1/2}^{1/2} \langle \nu_{(x_\alpha, x_3)} \otimes \delta_0, f^\# \rangle dx_3 \\ &\geq \int_{-1/2}^{1/2} f^\#(\langle \nu_{(x_\alpha, x_3)} \otimes \delta_0, \text{id} \rangle) dx_3 \\ &= \int_{-1/2}^{1/2} f^\#(D_\alpha y(x_\alpha)|0) dx_3 = f(D_\alpha y(x_\alpha)) = f(\langle \text{Av}_{x_\alpha}^3 \nu, \text{id} \rangle), \end{aligned}$$

which proves property (iii). In the last equality we have used (8.1). We now prove the second part of the theorem. Let $\bar{\mu}$ be a $W^{1,p}(\omega; \mathbb{R}^3)$ -gradient Young measure. Then, obviously $\bar{\mu} \otimes \delta_0$ is a $W^{1,p}(\Omega; \mathbb{R}^3)$ -gradient Young measure and $\text{Av}^3 \bar{\mu} = \bar{\mu}$. \square

The next theorem puts into relation the spaces $\hat{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ and $\mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3)$.

Lemma 8.2 *We have*

$$\text{Av}^3 \hat{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3) = \mathcal{Y}_\Gamma^{1,p}(\omega; \mathbb{R}^3) \otimes \delta_0.$$

PROOF. Let $\mu \in \mathring{\mathcal{Y}}_{\Gamma}^{1,p}(\Omega; \mathbb{R}^3)$. Then, by Lemma 5.3, $\mu = \nu \otimes \delta_0$, and for $\varphi \in C_0(M^{3 \times 3})$ we have

$$\begin{aligned} \langle \text{Av}_{x_\alpha}^3 \mu, \varphi \rangle &= \int_{-1/2}^{1/2} \int_{M^{3 \times 3}} \varphi(\bar{F}|z) d\nu_{(x_\alpha, x_3)} \otimes \delta_0(\bar{F}|z) dx_3 \\ &= \int_{-1/2}^{1/2} \int_{M^{3 \times 2}} \varphi(\bar{F}|0) d\nu_{(x_\alpha, x_3)}(\bar{F}) dx_3 \\ &= \int_{M^{3 \times 2}} \varphi(\bar{F}|0) d\text{Av}_{x_\alpha}^3 \nu(\bar{F}) \\ &= \int_{M^{3 \times 3}} \varphi(\bar{F}|z) d\text{Av}_{x_\alpha}^3 \nu \otimes \delta_0(\bar{F}|z) \\ &= \langle (\text{Av}_{x_\alpha}^3 \nu) \otimes \delta_0, \varphi \rangle. \end{aligned}$$

Thus

$$\text{Av}^3(\nu \otimes \delta_0) = (\text{Av}^3 \nu) \otimes \delta_0,$$

and the claim of the lemma follows by an easy application of Lemma 8.1. \square

Remark 8.3 *The useful consequence of Lemma 8.2 is*

$$q(\mathring{\mathcal{Y}}_{\Gamma}^{1,p}(\Omega; \mathbb{R}^3)) = \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3).$$

Proof of Theorem 5.6

Since $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$ is dual of the space $L^1(\omega; C_0(M^{3 \times 2}))$ which is Banach and separable, the proof is a direct consequence of the following proposition.

Proposition 8.4 *If $\varepsilon_n \rightarrow 0$ and the sequence $\mu^n \in L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$ satisfy*

$$\sup_n I_{\varepsilon_n}^{\mathcal{M}}(\mu^n) < +\infty,$$

then there exists $n_k \rightarrow \infty$ and a Young measure $\nu \in L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$ such that $q(\mu^{n_k}) \rightarrow \nu$ weakly in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$. Moreover $\nu \in \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)$.*

PROOF. By Lemma 5.1 there exists $\mu \in \mathring{\mathcal{Y}}_{\Gamma}^{1,p}(\Omega; \mathbb{R}^3)$ and a subsequence μ^{n_k} such that

$$\mu^{n_k} \rightarrow \mu \text{ weakly* in } L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3})).$$

Let $\nu = q(\mu)$. By continuity of q we have that $q(\mu^{n_k}) \rightarrow \nu$ weakly* in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$. Moreover, by Remark 8.3, $\nu \in \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)$. \square

Proof of Theorem 5.7

Let ε_n be an infinitesimal sequence of positive real numbers. Since the sequence of functionals $I_{\varepsilon_n}^{\mathcal{M}}$ is $(q, w^*L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2})))$ -equi-coercive, then Proposition 3.5 applies. We start by finding a recovering sequence for a suitable subsequence $I_{\varepsilon_{n_j}}^{\mathcal{M}}$; that is, for $\nu \in \mathcal{Y}_{\Gamma}^{1,p}(\omega; \mathbb{R}^3)$ we find a sequence $y^j \in W^{1,p}(\Omega; \mathbb{R}^3)$

with $y^j(x) = (x_1, x_2, \varepsilon_{n_j} x_3)$ on Γ such that the projections on the first two columns of the averages with respect to x_3 of the associated Young measures $\mu^j = \delta_{Dy^j}$ converge to ν , weakly* in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$, and

$$\limsup_{j \rightarrow \infty} I_{\varepsilon_{n_j}}^{\mathcal{M}}(\mu^j) \leq I(\nu).$$

Let $y \in W^{1,p}(\omega; \mathbb{R}^3)$ be the underlying deformation of ν with $y = (x_1, x_2, 0)$ on Γ . By Proposition 2.6 there exists a sequence $\psi^j \in C_0^\infty(\omega; \mathbb{R}^3)$ such that, setting $v^j := y + \psi^j$, we have

$$D_\alpha v^j \text{ generates } \nu,$$

$v^j \rightarrow y$ weakly in $W^{1,p}(\omega; \mathbb{R}^3)$ and the sequence $\{|D_\alpha v^j|^p\}_j$ is equi-integrable. By a measurable selection argument (see for instance Buttazzo [14], Proposition 2.2.7), there exists a sequence of measurable functions $z^j : \omega \rightarrow \mathbb{R}^3$ such that

$$\begin{aligned} W_0(D_\alpha v^j(x_\alpha)) &= \min\{W(D_\alpha v^j(x_\alpha), z) : z \in \mathbb{R}^3\} \\ &= W(D_\alpha v^j(x_\alpha) | z^j(x_\alpha)) \text{ a.e. } x_\alpha \in \omega. \end{aligned}$$

By the growth condition from above of W we have that

$$-c \leq W_0(D_\alpha v^j) = W(D_\alpha v^j | z^j) \leq W(D_\alpha v^j | 0) \leq C(|D_\alpha v^j|^p + 1)$$

almost everywhere, and therefore the sequence $W_0(D_\alpha v^j)$ turns out to be equi-integrable. By continuity of W_0 and an application of Theorem 2.2 we get

$$\begin{aligned} \lim_{j \rightarrow \infty} \int_\Omega W(D_\alpha v^j | z^j) dx &= \lim_{j \rightarrow \infty} \int_\omega W_0(D_\alpha v^j(x_\alpha)) dx_\alpha \\ &= \int_\omega \int_{M^{3 \times 2}} W_0(\bar{F}) d\nu_{x_\alpha}(\bar{F}) dx_\alpha. \end{aligned} \quad (8.2)$$

Let us remark moreover that, by the growth assumption from below on W , even the sequence z^j turns out to be p -equi-integrable.

Let \bar{w}^j be functions in $C_0^\infty(\omega; \mathbb{R}^3)$ such that

$$\|z^j - (0, 0, 1) - \bar{w}^j\|_p < \frac{1}{j},$$

and let ε_{n_j} be a subsequence of ε_n such that

$$\|\varepsilon_{n_j} D_\alpha \bar{w}^j\|_p \rightarrow 0.$$

We define $w^j := (0, 0, 1) + \bar{w}^j$. Then $w^j = (0, 0, 1)$ on Γ , w^j is p -equi-integrable and

$$\lim_{j \rightarrow \infty} \int_\Omega [W(D_\alpha v^j | z^j) - W(D_\alpha v^j + \varepsilon_{n_j} x_3 D_\alpha w^j | w^j)] dx = 0. \quad (8.3)$$

Indeed, the sequences

$$(D_\alpha v^j | z^j) \quad \text{and} \quad (D_\alpha v^j + \varepsilon_{n_j} x_3 D_\alpha w^j | w^j)$$

generate the same Young measure and, being p -equi-integrable, the growth condition from above ensures that the sequences

$$W(D_\alpha v^j | z^j) \quad \text{and} \quad W(D_\alpha v^j + \varepsilon_{n_j} x_3 D_\alpha w^j | w^j)$$

are equi-integrable as well. Thus (8.3) follows by an application of Theorem 2.2. Let us define y^j by

$$y^j(x) := v^j(x_\alpha) + \varepsilon_{n_j} x_3 w^j(x_\alpha).$$

and observe that, using (8.3) and (8.2), we have

$$\begin{aligned} \lim_{j \rightarrow \infty} \int_{\Omega} W(D_\alpha y^j | \frac{D_3 y^j}{\varepsilon_{n_j}}) dx &= \lim_{j \rightarrow \infty} \int_{\Omega} W(D_\alpha v^j + \varepsilon_{n_j} x_3 D_\alpha w^j | w^j) dx \\ &= \lim_{j \rightarrow \infty} \int_{\Omega} W(D_\alpha v^j | z^j) dx \\ &= \int_{\omega} \int_{M^{3 \times 2}} W_0(\lambda) d\nu_{x_\alpha}(\lambda) dx_\alpha. \end{aligned}$$

Moreover, since $y^j \rightarrow \tilde{y}$ in $L^p(\Omega; \mathbb{R}^3)$ where $\tilde{y}(x_1, x_2, x_3) = y(x_1, x_2)$, we also deduce that

$$\begin{aligned} \lim_{j \rightarrow \infty} \int_{\Omega} f(x) \cdot y^j(x) dx &= \int_{\Omega} f(x) \cdot \tilde{y}(x_\alpha) dx \\ &= \int_{\omega} \int_{-1/2}^{1/2} f(x_\alpha, x_3) dx_3 \cdot y(x_\alpha) dx_\alpha. \end{aligned}$$

Thus, the first part of the proof is concluded by noting that

- $y^j = (x_1, x_2, \varepsilon_{n_j} x_3)$ on Γ ,
- $\mu^j := \delta_{D y^j} \rightarrow \nu \otimes \delta_0$ weakly* in $L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$,
- $q(\mu^j) \rightarrow q(\nu \otimes \delta_0) = \nu$ weakly* in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$ by continuity of q ,

and using Remark 8.3.

The next step consists in proving the liminf inequality for the sequence $I_{\varepsilon_n}^{\mathcal{M}}$. Let $\mu^n \in L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$ be a sequence such that $q(\mu^n)$ weakly* converges to ν in $L_w^\infty(\omega; \mathcal{M}(M^{3 \times 2}))$. We have to prove that

$$\liminf_{n \rightarrow \infty} I_{\varepsilon_n}^{\mathcal{M}}(\mu^n) \geq I(\nu).$$

Without loss of generality we may suppose that the left hand side of the equation above be finite and that the liminf be indeed a limit. Therefore $\sup_n I_{\varepsilon_n}^{\mathcal{M}}(\mu^n) < +\infty$ and, by Lemma 5.1, there exist $\mu \in \mathring{\mathcal{Y}}_\Gamma^{1,p}(\Omega; \mathbb{R}^3)$ and a subsequence μ^{n_k} such that

$$\mu^{n_k} \rightarrow \mu \quad \text{weakly* in } L_w^\infty(\Omega; \mathcal{M}(M^{3 \times 3}))$$

and the corresponding underlying deformations y^{n_k} and y which satisfy the boundary conditions $y^{n_k}(x) = (x_1, x_2, \varepsilon_{n_k} x_3)$ and $y(x) = (x_1, x_2, 0)$ are such that

$$y^{n_k} \rightarrow y \text{ weakly in } W^{1,p}(\Omega; \mathbb{R}^3) \text{ and } y = y(x_1, x_2). \quad (8.4)$$

Hence, by Remark 8.3, $\nu \in q(\overset{\circ}{\mathcal{Y}}_{\Gamma}^{1,p}(\Omega; \mathbb{R}^3)) = \mathcal{Y}^{1,p}(\omega; \mathbb{R}^3)$. Moreover,

$$\begin{aligned} \liminf_{n \rightarrow \infty} I_{\varepsilon_n}^{\mathcal{M}}(\mu^n) &= \lim_{n \rightarrow \infty} I_{\varepsilon_n}^{\mathcal{M}}(\mu^n) = \lim_{k \rightarrow \infty} I_{\varepsilon_{n_k}}^{\mathcal{M}}(\mu^{n_k}) \\ &= \lim_{k \rightarrow \infty} \left(\int_{\Omega} W(D_{\alpha} y^{n_k} | \frac{D_3 y^{n_k}}{\varepsilon_{n_k}}) dx - \int_{\Omega} f \cdot y^{n_k} dx \right) \\ &\geq \lim_{k \rightarrow \infty} \left(\int_{\Omega} W_0(D_{\alpha} y^{n_k}) dx - \int_{\Omega} f \cdot y^{n_k} dx \right) \\ &= \lim_{k \rightarrow \infty} \left(\int_{\Omega} \langle \bar{\pi}_{\#} \mu_x^{n_k}, W_0 \rangle dx - \int_{\Omega} f(x) \cdot y^{n_k}(x) dx \right) \\ &= \lim_{k \rightarrow \infty} \left(\int_{\omega} \left(\int_{-\frac{1}{2}}^{\frac{1}{2}} \langle \bar{\pi}_{\#} \mu_x^{n_k}, W_0 \rangle dx_3 \right) dx_{\alpha} - \int_{\Omega} f(x) \cdot y^{n_k} dx \right) \\ &= \lim_{k \rightarrow \infty} \left(\int_{\omega} \langle Av^3 \bar{\pi}_{\#} \mu^{n_k}, W_0 \rangle dx_{\alpha} - \int_{\Omega} f(x) \cdot y^{n_k}(x) dx \right) \\ &= \lim_{k \rightarrow \infty} \left(\int_{\omega} \langle q(\mu^{n_k})_{x_{\alpha}}, W_0 \rangle dx_{\alpha} - \int_{\Omega} f(x) \cdot y^{n_k}(x) dx \right) \\ &\geq \int_{\omega} \langle \nu_{x_{\alpha}}, W_0 \rangle dx_{\alpha} - \int_{\omega} \int_{-\frac{1}{2}}^{\frac{1}{2}} f(x_{\alpha}, x_3) dx_3 \cdot y(x_{\alpha}) dx_{\alpha} \end{aligned}$$

where in the last inequality we used (8.4) and Theorem 2.2. \square

References

- [1] E. Acerbi, G. Buttazzo, and D. Percivale, *A variational definition of the strain energy for an elastic string*, J. Elasticity **25** (1991), 137–148.
- [2] E. Acerbi and N. Fusco, *Semicontinuity problems in the calculus of variations*, Arch. Rational Mech. Anal. **86** (1984), no. 2, 125–145.
- [3] G. Anzellotti, S. Baldo, and D. Percivale, *Dimension reduction in variational problems, asymptotic development in Γ -convergence and thin structures in elasticity*, Asymptot. Anal. **9** (1994), no. 1, 61–100.
- [4] J.M. Ball, *A version of the fundamental theorem for Young measures*, PDE's and continuum models of phase transitions, Lecture Notes in Physics vol. 359, Springer, 1989, 207–215.
- [5] J.M. Ball and R.D. James, *Fine phase mixtures as minimizers of energy*, Arch. Rational Mech. Anal. **100** (1987), no. 1, 13–52.

- [6] J.M. Ball and R.D. James, *Proposed experimental tests of a theory of fine microstructure and the two well problem*, Phil. Trans. R. Soc. London A. **338** (1992), 389–450.
- [7] P. Bělík and M. Luskin, *A computational model for the indentation and phase transformation of a martensitic thin film*, J. Mech. Phys. Solids **50** (2002), no. 9, 1789–1815.
- [8] P. Bělík and M. Luskin, *A total-variation surface energy model for thin films of martensitic crystals*, Interfaces Free Bound. **4** (2002), no. 1, 71–88.
- [9] K. Bhattacharya and A. Braides, *Thin films with many small cracks*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **458** (2002), 823–840.
- [10] K. Bhattacharya and R.D. James, *A theory of thin films of martensitic materials with applications to microactuators*, J. Mech. Phys. Solids **47** (1999), no. 3, 531–576.
- [11] M. Bocea, *Young measures minimizers in the asymptotic analysis of thin films*, Preprint, 2003.
- [12] M. Bocea and I. Fonseca, *Equi-integrability results for 3D-2D dimension reduction problems*, ESAIM Control Optim. Calc. Var. **7** (2002), 443–470, (electronic).
- [13] A. Braides, I. Fonseca, and G. Francfort, *3d-2d asymptotic analysis for inhomogeneous thin films*, Indiana Univ. Math. J. **49** (2000), no. 4, 1367–1404.
- [14] G. Buttazzo, *Semicontinuity, relaxation and integral representation in the calculus of variations*, Pitman Res. Notes Math. Ser., vol. 207, Longman, 1989.
- [15] P. G. Ciarlet and P. Destuynder, *A justification of the two-dimensional linear plate model*, J. Mécanique **18** (1979), no. 2, 315–344.
- [16] B. Dacorogna, *Direct methods in the calculus of variations*, Applied Mathematical Sciences, vol. 78, Springer, 1989.
- [17] G. Dal Maso, *An introduction to Γ -convergence*, Birkhäuser, 1993.
- [18] L. Freddi and R. Paroni *A 3D-1D Young measure theory of an elastic string*, to appear in Asymptot. Anal., 2004.
- [19] R.D. James and D. Kinderlehrer, *Theory of diffusionless phase transitions*, Lecture Notes in Physics vol. 344, Springer, 1989, 51–84.
- [20] D. Kinderlehrer and P. Pedregal, *Characterizations of Young measures generated by gradients*, Arch. Rational Mech. Anal. **115** (1991), no. 4, 329–365.

- [21] D. Kinderlehrer and P. Pedregal, *Gradient Young measures generated by sequences in Sobolev spaces*, J. Geom. Analysis **4** (1994), 59–90.
- [22] H. Le Dret and A. Raoult, *The nonlinear membrane model as variational limit of nonlinear three-dimensional elasticity*, J. Math. Pures Appl. **74** (1995), no. 6, 549–578.
- [23] M. Luskin, *On the computation of crystalline microstructure*, Acta numerica, **5** (1996), 191–257.
- [24] S. Müller, *Variational models for microstructure and phase transitions.*, Calculus of variations and geometric evolution problems (Cetraro, 1996), Lecture Notes in Math. vol. 1713, Springer, 1999, 85–210.
- [25] F. Murat and L. Tartar, *Calculus of variations and homogenization*, Topics in the mathematical modelling of composite materials, Progr. Nonlinear Differential Equations Appl., vol. 31, Birkhäuser, 1997, 139–173.
- [26] R. Paroni, *From discrete to continuum problems: a Young measure approach.*, Z. angew. Math. Phys. **54** (2003), 328–348.
- [27] P. Pedregal, *Parametrized measures and variational principles*, Progress in nonlinear differential equations and their applications, Birkhäuser, 1997.
- [28] P.M. Santos and E. Zappale, *Second order analysis for thin structures*, Preprint, 2002.
- [29] Y.C. Shu, *Heterogeneous thin films of martensitic materials*, Arch. Rational Mech. Anal. **153** (2000), 39–90.
- [30] M. Valadier, *Young measures*, Methods of Nonconvex Analysis, Lecture Notes in Math. vol. 1446, Springer, 1990, 152–188.
- [31] L.C. Young, *Lectures on the calculus of variations and optimal control theory*, W. B. Saunders Co., 1969.

Lorenzo Freddi
Dipartimento di Matematica e Informatica
Università di Udine
via delle Scienze 206
33100 Udine, Italy
e-mail: freddi@dimi.uniud.it
and
Roberto Paroni
Dipartimento di Architettura e Pianificazione
Università di Sassari
Palazzo del Pou Salit
Piazza Duomo

07041 Alghero, Italy
e-mail: paroni@uniss.it