

KINETIC FORMULATION AND UNIQUENESS FOR SCALAR CONSERVATION LAWS WITH DISCONTINUOUS FLUX

GRAZIANO CRASTA, VIRGINIA DE CICCO, AND GUIDO DE PHILIPPIS

ABSTRACT. We prove a uniqueness result for BV solutions of scalar conservation laws with discontinuous flux in several space dimensions. The proof is based on the notion of kinetic solution and on a careful analysis of the entropy dissipation along the discontinuities of the flux.

1. INTRODUCTION

This paper is concerned with multidimensional scalar conservation laws with discontinuous flux, namely

$$(1) \quad u_t + \operatorname{div} \mathbf{A}(x, u) = 0, \quad (t, x) \in (0, +\infty) \times \mathbb{R}^n,$$

where the flux function $\mathbf{A}: \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ is possibly discontinuous in the first variable and satisfies some structural assumptions listed in Section 2. These type of equations have attracted a lot of attention in the last years since they naturally arise in several models (for instance models of traffic flow, flow in porous media, sedimentation processes, etc.), see [5, 18, 25] and references therein for a more detailed account on the theory.

It is well known that, even for smooth fluxes, in general the Cauchy problem

$$(2) \quad \begin{cases} u_t + \operatorname{div} \mathbf{A}(x, u) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}^n, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^n, \end{cases}$$

does not admit classical solutions. On the other hand, the notion of distributional solution is too weak in order to achieve well-posedness, in particular it does not provide uniqueness of the solution. In this context, the notion of entropy solution turns out to be the right one, as it has been shown by Vol'pert [28] in the BV setting and by Kruzkov [20] in the L^∞ framework.

The classical Kruzkov's approach [20] completely solves the problem of well-posedness in the case of smooth fluxes. Moreover, as it has been shown in recent years, using a clever change of variables and the concept of *adapted entropies*, this approach also works for a restricted class of discontinuous fluxes (see [7, 8, 11, 23, 25]).

The case of more general discontinuous fluxes in one space variable has been extensively studied (see for example [2, 5, 6, 7, 9, 16, 18, 19, 23, 24] and references therein). In particular, it has been pointed out ([2, 5, 18]) that many different admissibility criteria generate continuous semigroups of solutions, and the choice of the right criterion may depend on the physics of the problem under consideration. Indeed, in addition to

Date: September 12, 2014.

2010 Mathematics Subject Classification. Primary: 35L65; Secondary: 26A45.

Key words and phrases. Chain rule, BV functions, conservation laws with discontinuous flux.

the classical entropy criteria, one has to impose some conditions on the behavior of the solutions on the discontinuities of the flux. Roughly speaking different conditions give rise to different criteria. These different conditions are coded, for example, in the notion of *germ* (see [5]) or of well-posed Riemann solver (see [18]). One of the most studied admissibility criterion is based on the notion of vanishing viscosity solution. This will actually be the criterion that we are going to use in this paper, see Remark 3.3 below.

In spite of the intensive study concerning conservation laws with discontinuous fluxes, in the multidimensional case there are very few results available in the literature. A very general existence result has been obtained by Panov [25]. On the other hand, a well-posedness result for a restricted class of fluxes (having only one regular hypersurface of discontinuity) has been recently proved by Mitrovic [22].

In this paper we propose an entropy criterion, modeled on the ones introduced in the one-dimensional case in [6, 16, 24], which allows to prove uniqueness of *BV* entropy solutions of the Cauchy problem (2) under mild assumptions on the flux \mathbf{A} . More precisely our main result is the following:

Theorem. *Let $\mathbf{A} \in L^\infty(\mathbb{R}^n \times \mathbb{R}; \mathbb{R}^n)$ satisfies (H1)–(H7) in Section 2 below and let u_1 and u_2 be two *BV* entropy solutions of (1) (see Definition 3.2), then*

$$\int_{\mathbb{R}^n} |u_1(T, x) - u_2(T, x)| dx \leq \int_{\mathbb{R}^n} |u_1(0, x) - u_2(0, x)| dx.$$

Let us mention that our assumptions are satisfied in the particular case of fluxes of the form $\mathbf{A}(x, u) = \widehat{\mathbf{A}}(k(x), u)$ where k is in $SBV(\mathbb{R}^n; \mathbb{R}^N) \cap L^\infty(\mathbb{R}^n; \mathbb{R}^N)$ and $\widehat{\mathbf{A}}$ is smooth, see Remark 2.3.

On the other hand we do not investigate the issue of existence of *BV* entropy solutions. Let us point out that, unlike the case of fluxes with smooth dependence on x , when \mathbf{A} is discontinuous there is no general known result concerning existence of *BV* solutions, see however Remark 3.4.

Our method of proof is based on the notion of kinetic solution introduced by Lions, Perthame and Tadmor (see [21, 26, 27]). This technique has been successfully applied by Dalibard to the case of nonautonomous smooth fluxes (see [13, 14]). More precisely, we show the equivalence between the entropic and the kinetic formulation of (1). Relying on the nonautonomous chain rule for *BV* functions proved in [3] and on a careful analysis of the entropy dissipation along the discontinuities of the flux, we are then able to prove that the semigroup of solutions is contractive in the L^1 norm.

The plan of the paper is the following. In Section 2 we state the assumptions on the flux (see (H1)–(H7) below) and we discuss some preliminary results. In Section 3 we introduce our notions of entropy and kinetic solution, we prove their equivalence (see Theorem 3.9), and we state our main uniqueness result (see Theorem 3.5). In Section 4 we prove some preliminary estimates for kinetic solutions, and, finally, in Section 5 we prove the uniqueness of kinetic solutions (see Theorem 5.1).

ACKNOWLEDGMENTS. The authors would like to thank Luigi Ambrosio and Gianluca Crippa for some useful discussions, and two referees for having improved, with their comments, the presentation of the paper. The authors have been supported by the Gruppo

Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

2. ASSUMPTIONS ON THE VECTOR FIELD AND THE CHAIN RULE

In this section we first survey some useful facts about BV function that we need in the sequel, we state our main structural hypotheses on the vector field (assumptions (H1)-(H7) below) and prove some consequences of these assumptions.

2.1. BV functions. Let us start recalling our main notation and preliminary facts on BV and SBV functions. A general reference is Chapter 3 of [4], and occasionally we will give more precise references therein.

We denote by \mathcal{L}^n the Lebesgue measure in \mathbb{R}^n and by \mathcal{H}^k the k -dimensional Hausdorff measure. By Radon measure we mean a nonnegative Borel measure finite on compact sets. If μ is a Radon measure on X and ν is a Radon measure on Y we denote by $\mu \times \nu$ the the product measure on $X \times Y$, sometimes we will also write $\mu_x \times \nu_y$. Given a Borel set D we will denote by $\mu \llcorner D$ the Radon measure given by $\mu \llcorner D(B) = \mu(B \cap D)$. A set $\Sigma \subset \mathbb{R}^n$ is said to be countably \mathcal{H}^{n-1} rectifiable if \mathcal{H}^{n-1} -almost all of Σ can be covered by a sequence of C^1 hypersurfaces. Let us recall that if $\Sigma \subset \mathbb{R}^n$ is countably \mathcal{H}^{n-1} rectifiable, then $(a, b) \times \Sigma \subset \mathbb{R} \times \mathbb{R}^n$ is countably \mathcal{H}^n rectifiable and

$$\mathcal{H}^n \llcorner (a, b) \times \Sigma = \mathcal{L}^1 \llcorner (a, b) \times \mathcal{H}^{n-1} \llcorner \Sigma,$$

see [17, Theorem 3.2.23].

A function $u \in L^1(\mathbb{R}^n)$ belongs to $BV(\mathbb{R}^n)$ if its derivative in the sense of distributions is representable by a vector-valued measure $Du = (D_1u, \dots, D_nu)$ whose total variation $|Du|$ is finite, i.e.

$$\int_{\mathbb{R}^n} u \frac{\partial \phi}{\partial x_i} dx = - \int \phi D_i u \quad \forall \phi \in C_c^\infty(\mathbb{R}^n), \quad i = 1, \dots, n$$

and $|Du|(\mathbb{R}^n) < \infty$.

Approximate continuity and jump points. We say that $x \in \mathbb{R}^n$ is an approximate continuity point of u if, for some $z \in \mathbb{R}$, it holds

$$\lim_{r \downarrow 0} \int_{B_r(x)} |u(y) - z| dy = 0.$$

The number z is uniquely determined at approximate continuity points and denoted by $\tilde{u}(x)$, the so-called approximate limit of u at x . The complement of the set of approximate continuity points, the so-called singular set of u , will be denoted by S_u .

Analogously, we say that x is a jump point of u , and we write $x \in J_u$, if there exists a unit vector $\nu \in \mathbf{S}^{n-1}$ and $u^+, u^- \in \mathbb{R}$ satisfying $u^+ \neq u^-$ and

$$\lim_{r \downarrow 0} \int_{B^\pm(x, r)} |u(y) - u^\pm| dy = 0,$$

where $B^\pm(x, r) := \{y \in B_r(x) : \pm \langle y - x, \nu \rangle \geq 0\}$ are the two half balls determined by ν . At points $x \in J_u$ the triplet (u^+, u^-, ν) is uniquely determined up to a permutation of (u^+, u^-) and a change of sign of ν ; for this reason, with a slight abuse of notation, we do not emphasize the ν dependence of u^\pm and $B^\pm(x, r)$. Since we impose $u^+ \neq u^-$, it is clear that $J_u \subset S_u$, moreover, for every $u \in BV_{loc}$, $\mathcal{H}^{n-1}(S_u \setminus J_u) = 0$ and J_u is \mathcal{H}^{n-1}

countably rectifiable (see [4, Theorem 3.78]). Finally, we define the precise representative as

$$(3) \quad u^*(x) = \begin{cases} \tilde{u}(x) & x \in \mathbb{R}^n \setminus S_u \\ (u^+(x) + u^-(x))/2 & x \in J_u \\ 0 & \text{otherwise.} \end{cases}$$

Note that $\mathcal{H}^{n-1}(\{\tilde{u} \neq u^*\} \setminus J_u) = 0$, in particular, since $|Du| \ll \mathcal{H}^{n-1}$, $|\tilde{D}u|(\{\tilde{u} \neq u^*\}) = 0$.

Decomposition of the distributional derivative. For any oriented and countably \mathcal{H}^{n-1} -rectifiable set $\Sigma \subset \mathbb{R}^n$ we have

$$(4) \quad Du \llcorner \Sigma = (u^+ - u^-)\nu_\Sigma \mathcal{H}^{n-1} \llcorner \Sigma.$$

For any $u \in BV(\mathbb{R}^n)$, we can decompose Du as the sum of a diffuse part, that we shall denote $\tilde{D}u$, and a jump part, that we shall denote by $D^j u$. The diffuse part is characterized by the property that $|\tilde{D}u|(B) = 0$ whenever $\mathcal{H}^{n-1}(B)$ is finite, while the jump part is concentrated on a set σ -finite with respect to \mathcal{H}^{n-1} . The diffuse part can be then split as

$$\tilde{D}u = D^a u + D^c u$$

where $D^a u$ is the absolutely continuous part with respect to the Lebesgue measure, while $D^c u$ is the so-called Cantor part. The density of Du with respect to \mathcal{L}^n can be represented as follows

$$D^a u = \nabla u d\mathcal{L}^n,$$

where ∇u is the approximate gradient of u , see [4, Proposition 3.71 and Theorem 3.83]. Note also that $|\tilde{D}u|(\{\tilde{u} \neq u^*\}) = 0$. The jump part can be easily computed by taking $\Sigma = J_u$ (or, equivalently, S_u) in (4), namely

$$D^j u = Du \llcorner J_u = (u^+ - u^-)\nu_{J_u} \mathcal{H}^{n-1} \llcorner J_u.$$

We will say that $u \in SBV(\mathbb{R}^n)$ if $D^c u = 0$, i.e. if

$$Du = (u^+ - u^-)\nu_{J_u} \mathcal{H}^{n-1} \llcorner J_u + \nabla u d\mathcal{L}^n.$$

All these concepts and results extend, mostly arguing component by component, to vector-valued BV functions, see [4] for details.

Sets of finite perimeter and coarea formula. We will say that a measurable set E is of finite perimeter if its characteristic function χ_E belongs to $BV(\mathbb{R}^n)$. In this case we denote by

$$\partial^* E = J_{\chi_E}$$

its *reduced boundary* (note that this is slightly larger than what it is usually called reduced boundary, however it coincides with it up to a \mathcal{H}^{n-1} negligible set, see [4, Chapter 3]). Then,

$$D\chi_E = D^j \chi_E = -\nu_{\partial^* E} \mathcal{H}^{n-1} \llcorner \partial^* E,$$

in particular $\chi_E \in SBV$. We also recall the coarea formula: if $u \in BV(\mathbb{R}^n)$, then for \mathcal{L}^1 almost every $v \in \mathbb{R}$, the characteristic function of the set $\{u > v\}$ belongs to BV and we have the following equalities between measures:

$$\begin{aligned} Du &= \int_{\mathbb{R}} D\chi_{u>v} dv \\ |Du| &= \int_{\mathbb{R}} |D\chi_{\{u>v\}}| dv = \int_{\mathbb{R}} \mathcal{H}^{n-1}(\partial^*\{u > v\}) dv. \end{aligned}$$

The following lemma relates the pointwise behavior of $\chi_{\{u>v\}}$ to the pointwise behavior of u , see [15, Lemma 2.2] for the proof.

Lemma 2.1. *For any function $u: \mathbb{R}^n \rightarrow \mathbb{R}$ and any $v \in \mathbb{R}$, let $\varphi_{u,v}(x) = \chi_{\{u>v\}}(x)$. If $u \in BV(\mathbb{R}^n)$, then for \mathcal{L}^1 -a.e. $v \in \mathbb{R}$ there exists a Borel set $N_v \subset \mathbb{R}^n$, with $\mathcal{H}^{n-1}(N_v) = 0$, such that the following relation holds:*

$$\varphi_{u^\pm, v}(x) = \varphi_{u, v}^\pm(x), \quad \forall x \in \mathbb{R}^n \setminus N_v.$$

2.2. Structural assumptions on the vector field. Let $\mathbf{A} \in L^\infty(\mathbb{R}^n \times \mathbb{R}; \mathbb{R}^n)$ be such that:

- (H1) There exists a set $\mathcal{C}_{\mathbf{A}}$ with $\mathcal{L}^n(\mathcal{C}_{\mathbf{A}}) = 0$ such that $\mathbf{A}(x, \cdot) \in C^1(\mathbb{R})$ for every $x \in \mathbb{R}^n \setminus \mathcal{C}_{\mathbf{A}}$ and $\mathbf{A}(\cdot, v) \in SBV(\mathbb{R}^n)$ for every $v \in \mathbb{R}^n$.
- (H2) There exists a constant M such that

$$|\partial_v \mathbf{A}(x, v)| \leq M \quad \forall x \in \mathbb{R}^n \setminus \mathcal{C}_{\mathbf{A}}, \quad v \in \mathbb{R}.$$

- (H3) There exists a modulus of continuity ω such that

$$|\partial_v \mathbf{A}(x, u) - \partial_v \mathbf{A}(x, w)| \leq \omega(|u - w|) \quad \forall x \in \mathbb{R}^n \setminus \mathcal{C}_{\mathbf{A}}, \quad u, w \in \mathbb{R}.$$

- (H4) There exists a function $g_1 \in L^1(\mathbb{R}^n)$ such that

$$|\nabla_x \mathbf{A}(x, u) - \nabla_x \mathbf{A}(x, w)| \leq g_1(x)|u - w| \quad \forall x \in \mathbb{R}^n \setminus \mathcal{C}_{\mathbf{A}}, \quad u, w \in \mathbb{R},$$

where $\nabla_x \mathbf{A}(x, v)$ denotes the approximate gradient of the map $x \mapsto \mathbf{A}(x, v)$.

- (H5) The measure

$$\sigma := \bigvee_{u \in \mathbb{R}} |D_x \mathbf{A}(\cdot, u)|$$

satisfies $\sigma(\mathbb{R}^n) < \infty$. Here $D_x \mathbf{A}(\cdot, u)$ is the distributional gradient of the map $x \mapsto \mathbf{A}(x, u)$ (which is a measure since $\mathbf{A}(\cdot, u) \in BV$) and \bigvee denotes the least upper bound in the space of nonnegative Borel measures, see [4, Definition 1.68].

2.3. Chain rule and fine properties of \mathbf{A} . Assumptions (H1)-(H5) imply that \mathbf{A} satisfies the hypothesis of [3]. Let us summarize some consequence of this fact. If we define

$$\mathcal{N} = \left\{ x \in \mathbb{R}^n : \liminf_{r \rightarrow 0} \frac{\sigma(B_r(x))}{r^{n-1}} > 0 \right\},$$

then \mathcal{N} is a \mathcal{H}^{n-1} rectifiable set. In the sequel we shall assume that:

- (H6) $\mathcal{H}^{n-1}(\mathcal{N}) < +\infty$.

In our context, Theorem 2.2 in [3] reads as follows: For every $u \in BV(\mathbb{R}^n; \mathbb{R})$ the composite function $w(x) = \mathbf{A}(x, u(x))$ belongs to $BV(\mathbb{R}^n; \mathbb{R}^n)$ with

$$(5) \quad |Dw| \leq \sigma + M|Du|$$

and

$$(6) \quad \tilde{D}w = \nabla_x \mathbf{A}(x, \tilde{u}(x)) \mathcal{L}^n + \partial_v \mathbf{A}(x, \tilde{u}(x)) \otimes \tilde{D}u$$

$$(7) \quad D^j w = [\mathbf{A}^+(x, u^+(x)) - \mathbf{A}^-(x, u^-(x))] \otimes \nu_{J_u \cup \mathcal{N}} \mathcal{H}^{n-1} \llcorner (J_u \cup \mathcal{N}).$$

Here the functions $\mathbf{A}^\pm(x, v)$ are defined for \mathcal{H}^{n-1} almost every $x \in \mathcal{N} \cup J_u$ and every $v \in \mathbb{R}$ as

$$(8) \quad \mathbf{A}^\pm(x, v) = \lim_{r \rightarrow 0} \int_{B_r^\pm(x)} \mathbf{A}(y, v) dy,$$

in particular

$$\mathbf{A}^+(x, v) = \mathbf{A}^-(x, v) \quad \mathcal{H}^{n-1}\text{-a.e. } x \in J_u \setminus \mathcal{N} \text{ and every } v \in \mathbb{R}.$$

Note that by interchanging the derivative with the integral, applying Ascoli-Arzelà Theorem and taking into account (H2) and (H3), we can deduce as in [3, Section 3] that the map $v \mapsto \mathbf{A}^\pm(x, v)$ is C^1 for almost every $x \in \mathcal{N}$ with derivative given by $\partial_v \mathbf{A}^\pm(x, v) = (\partial_v \mathbf{A}(x, v))^\pm$ (it is part of the statement the fact that this last quantity is well defined). In particular for \mathcal{H}^{n-1} -a.e. $x \in \mathcal{N}$ and all $u, w \in \mathbb{R}$ we have

$$(9) \quad |(\partial_v \mathbf{A})^\pm(x, u) - (\partial_v \mathbf{A})^\pm(x, w)| \leq \omega(|u - w|).$$

In the same way, see [3, Section 3], for \mathcal{H}^{n-1} almost every point of $\mathbb{R}^n \setminus \mathcal{N}$ and every $v \in \mathbb{R}$ there exists the limit

$$\tilde{\mathbf{A}}(x, v) = \lim_{r \rightarrow 0} \int_{B_r(x)} \mathbf{A}(y, v) dy,$$

and the map $v \mapsto \tilde{\mathbf{A}}(x, v)$ is continuously differentiable with $\partial_v \tilde{\mathbf{A}}(x, v) = \widetilde{\partial_v \mathbf{A}}(x, v)$. Moreover for \mathcal{L}^n -a.e. x and all $u, w \in \mathbb{R}$

$$(10) \quad |\widetilde{\partial_v \mathbf{A}}(x, u) - \widetilde{\partial_v \mathbf{A}}(x, w)| \leq \omega(|u - w|).$$

We conclude this section with the following simple remark. Thanks to (H4) and the previous discussion, the functions

$$\mathbf{B}_h(x, v) = \frac{\mathbf{A}(x, v+h) - \mathbf{A}(x, v)}{h}$$

satisfy

$$(11) \quad |\mathbf{B}_h(x, v)| \leq M, \quad |D_x \mathbf{B}_h(\cdot, v)| \leq g_1(x) \mathcal{L}^n + M \mathcal{H}^{n-1} \llcorner \mathcal{N},$$

where the first inequality follows from (H2). Using (H3) one can show that $\mathbf{B}_h(x, v) \rightarrow \partial_v \mathbf{A}(x, v)$ for almost every x and every $v \in \mathbb{R}$, see [3, Section 3] for similar arguments. From (11) we deduce that $\partial_v \mathbf{A}(\cdot, v) \in SBV$. Let us now consider the decomposition

$$D_x \partial_v \mathbf{A}(\cdot, v) = \nabla_x \partial_v \mathbf{A}(\cdot, v) \mathcal{L}^n + ((\partial_v \mathbf{A})^+(\cdot, v) - (\partial_v \mathbf{A})^-(\cdot, v)) \otimes \nu_{\mathcal{N}} \mathcal{H}^{n-1} \llcorner \mathcal{N}.$$

According to the discussion below equation (8) we have $(\partial_v \mathbf{A})^\pm = \partial_v(\mathbf{A}^\pm)$ for $\mathcal{H}^{n-1} \llcorner \mathcal{N}$ almost every x and every v . Moreover, by (H4) for every v the family

$$h \mapsto \frac{\nabla_x \mathbf{A}(x, v+h) - \nabla_x \mathbf{A}(x, v)}{h}, \quad h \neq 0,$$

is weakly compact in L^1 and every one of its cluster points has to coincide with $\nabla_x \partial_v \mathbf{A}(x, v)$, i.e.

$$\nabla_x \partial_v \mathbf{A}(x, v) = w_{L^1} - \lim_{h \rightarrow 0} \frac{\nabla_x \mathbf{A}(x, v+h) - \nabla_x \mathbf{A}(x, v)}{h}.$$

Let us now define for every Borel and bounded function φ the maps

$$h_1(v) := \int \varphi(x) \nabla_x \mathbf{A}(x, v) dx, \quad h_2(v) := \int_{\mathcal{N}} \varphi(x) (\mathbf{A}^+(x, v) - \mathbf{A}^-(x, v)) \otimes \nu_{\mathcal{N}} d\mathcal{H}^{n-1}(x).$$

By the previous discussion we then see that h_1, h_2 are Lipschitz continuous and everywhere differentiable with derivatives given by

$$(12) \quad \begin{aligned} \frac{dh_1(v)}{dv} &= \int \varphi(x) \nabla_x \partial_v \mathbf{A}(x, v) dx \\ \frac{dh_2(v)}{dv} &= \int_{\mathcal{N}} \varphi(x) (\partial_v \mathbf{A}^+(x, v) - \partial_v \mathbf{A}^-(x, v)) \otimes \nu_{\mathcal{N}} d\mathcal{H}^{n-1}. \end{aligned}$$

However in the sequel we will also need the following assumption, ensuring the continuity of the map $v \rightarrow dh_1/dv$:

(H7) There exist a L^1 function g_2 and a modulus of continuity ω (which we can assume without loss of generality to be equal to the one appearing in (H3)) such that

$$|\nabla_x \partial_v \mathbf{A}(x, u) - \nabla_x \partial_v \mathbf{A}(x, w)| \leq g_2(x) \omega(|u - w|) \quad \forall u, w \in \mathbb{R}.$$

With this assumption we have

Lemma 2.2. *Let \mathbf{A} satisfy (H1)–(H7), then there exists a set $\tilde{\mathcal{C}}_{\mathbf{A}}$ with $\mathcal{L}^n(\tilde{\mathcal{C}}_{\mathbf{A}}) = 0$ such that every $x \in \mathbb{R}^n \setminus \tilde{\mathcal{C}}_{\mathbf{A}}$ is a Lebesgue point for $x \mapsto \nabla_x \mathbf{A}(x, v)$, $x \mapsto \nabla_x \widetilde{\partial_v \mathbf{A}}(x, v)$ and for any such x the map $v \mapsto \widetilde{\nabla_x \mathbf{A}}(x, v)$ is C^1 with derivative given by $\widetilde{\nabla_x \partial_v \mathbf{A}}(x, v)$*

Proof. Let $U \subset \mathbb{R}$ be a countable dense set and let

$$\tilde{\mathcal{C}}_{\mathbf{A}} = \bigcup_{u \in U} \mathbb{R}^n \setminus S_{\mathbf{A}(\cdot, u)} \cup (\mathbb{R}^n \setminus S_{g_1}) \cup (\mathbb{R}^n \setminus S_{g_2}),$$

which clearly satisfies $\mathcal{L}^n(\tilde{\mathcal{C}}_{\mathbf{A}}) = 0$. By arguing as in [3, Section 3] and using (H4) and (H7) we see that for every $x \in \mathbb{R}^n \setminus \tilde{\mathcal{C}}_{\mathbf{A}}$ the limits

$$\widetilde{\nabla_x \mathbf{A}}(x, v) = \lim_{r \rightarrow 0} \int_{B_r(x)} \nabla_x \mathbf{A}(y, v) dy \quad \widetilde{\nabla_x \partial_v \mathbf{A}}(x, v) = \lim_{r \rightarrow 0} \int_{B_r(x)} \nabla_x \partial_v \mathbf{A}(y, v) dy$$

exist for every $v \in \mathbb{R}$. By the first equality in (12) the map

$$v \mapsto h_r(v) = \int_{B_r(x)} \nabla_x \mathbf{A}(y, v) dy$$

are differentiable with derivative given by $h'_r(v) = \int_{B_r(x)} \nabla_x \partial_v \mathbf{A}(y, v) dy$. Since x is a Lebesgue point for g_2 , thanks to (H7) this is a family of equi-continuous functions in

v converging to $\widetilde{\nabla_x \partial \mathbf{A}}(x, v)$. It is now a standard argument to see that $\widetilde{\nabla_x \mathbf{A}}(x, v) = \lim_r h_r(v)$ is C^1 with derivative given by $\lim_r h_r'(v) = \nabla_x \partial_v \mathbf{A}(x, v)$. \square

Remark 2.3. Let us point out that our hypotheses include (and actually are modeled on) the case $\mathbf{A}(x, u) = \widehat{\mathbf{A}}(k(x), u)$ where $k \in SBV(\mathbb{R}^n; \mathbb{R}^N) \cap L^\infty(\mathbb{R}^n; \mathbb{R}^N)$, $\mathcal{H}^{n-1}(J_k) < +\infty$ and $\widehat{\mathbf{A}} \in C^1(\mathbb{R}^N \times \mathbb{R}, \mathbb{R}^n) \cap \text{Lip}(\mathbb{R}^N \times \mathbb{R}, \mathbb{R}^n)$.

3. FORMULATION OF THE PROBLEM

3.1. Entropic formulation. We consider the following scalar conservation law

$$(13) \quad u_t + \text{div } \mathbf{A}(x, u) = 0,$$

where $\mathbf{A}: \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ satisfies the structural assumption (H1)–(H7).

Definition 3.1 (Convex entropy pair). We say that $(S, \boldsymbol{\eta})$ is a convex entropy pair if $S \in C^2(\mathbb{R})$ is a convex function, and $\boldsymbol{\eta} = (\eta_1, \dots, \eta_n)$ is defined by

$$(14) \quad \eta_i(x, v) := \int_0^v \partial_v A_i(x, w) S'(w) dw, \quad i = 1, \dots, n.$$

In the above definition and in the sequel,

$$A_i = \mathbf{A} \cdot e_i,$$

are the components of \mathbf{A} .

Note that according to the discussion in Section 2.3, $\boldsymbol{\eta}(\cdot, v) \in SBV(\mathbb{R}^n; \mathbb{R}^n)$ for every $v \in \mathbb{R}$ and its distributional derivative is given by

$$\begin{aligned} D_x \boldsymbol{\eta}(\cdot, v) &= \left(\int_0^v \nabla_x \partial_v \mathbf{A}(x, w) S'(w) dw \right) \mathcal{L}^n \\ &\quad + \left(\int_0^v (\partial_v \mathbf{A}^+(x, w) - \partial_v \mathbf{A}^-(x, w)) S'(w) dw \right) \otimes \nu_{\mathcal{N}} d\mathcal{H}^{n-1} \llcorner \mathcal{N}. \end{aligned}$$

Definition 3.2 (Entropy solutions). A function

$$u \in C([0, T]; L^1(\mathbb{R}^n)) \cap L^\infty((0, T) \times \mathbb{R}^n) \cap L^1((0, T); BV(\mathbb{R}^n))$$

is an entropy solution of (13) if u is a solution to (13) in the sense of distributions, and there exists a (everywhere defined) Borel representative \hat{u} of u with $|\hat{u}(t, x)| \leq \|u\|_\infty$ such that, for every convex entropy pair $(S, \boldsymbol{\eta})$, one has

$$(15) \quad \begin{aligned} &\partial_t S(u) + \text{div}(\boldsymbol{\eta}(x, u)) \\ &\quad - \text{div} \boldsymbol{\eta}(x, v) \Big|_{v=\hat{u}(t, x)} + S'(\hat{u}) \text{div} \mathbf{A}(x, v) \Big|_{v=\hat{u}(t, x)} \leq 0 \end{aligned}$$

in the distributional sense. Here, by $\text{div} \mathbf{A}(x, v) \Big|_{v=\hat{u}(t, x)}$ we mean the measure whose action on a bounded and Borel function $\varphi = \varphi(t, x)$ is given by

$$(16) \quad \begin{aligned} &\sum_{i=1}^n \int_0^T dt \int_{\mathbb{R}^n} \varphi(t, x) \nabla_i A_i(x, \hat{u}(t, x)) dx \\ &\quad + \sum_{i=1}^n \int_0^T dt \int_{\mathcal{N}} \varphi(t, x) (A_i^+(x, \hat{u}(t, x)) - A_i^-(x, \hat{u}(t, x))) \nu_{\mathcal{N}}^i d\mathcal{H}^{n-1}(x), \end{aligned}$$

and the same for $\operatorname{div} \boldsymbol{\eta}(x, v)|_{v=\hat{u}(t,x)}$.

Remark 3.3. Some comments about our definition of entropy solution are in order. We shall see in Section 3.2 that, if $x \in \mathcal{N}$, then our entropy condition characterizes the (closure of the) so-called vanishing viscosity germ defined in equation (5.4) of [5] (see (24) below). This characterization has also been used by Diehl (see Condition Γ in [16]), Mitrovic [24] and Andreianov & Mitrovic [6]. In this sense, despite the appearance of a somewhat arbitrary Borel representative \hat{u} , this definition seems to be the natural extension to our general framework of the conditions cited above. We also note that, although in Definition 3.2 we require \hat{u} to be defined everywhere, it will be clear by our arguments below, that it is enough to define \hat{u} $\mathcal{L}^1 \times (\mathcal{L}^n + \sigma)$ -a.e.

Remark 3.4. By definition, if u is an entropy solution to (13), then $u(t, \cdot) \in BV(\mathbb{R}^n)$ for every t . For regular fluxes with respect to the space variables, it is well known that the entropy solution to the Cauchy problem with initial data $u_0 \in BV(\mathbb{R}^n)$ remains in BV for all times (see [20, 10]). On the other hand, it is not clear when such regularity has to be expected for entropy solutions when the flux is discontinuous in the space variables. A relevant situation for which we can expect BV -regularity of solutions is the case $n = 1$, at least for a class of fluxes widely studied in the literature (see for example [1, 18] and references therein). For instance, if we assume that the flux is piecewise constant and that, at every point of discontinuity, it satisfies an appropriate version of the crossing condition, one can show the existence of a BV solution of the Cauchy problem (see [1, Theorem 2.13] and [18, Lemma 9]).

With these definitions at hand we can now restate our main result:

Theorem 3.5. *Let $\mathbf{A} \in L^\infty(\mathbb{R}^n \times \mathbb{R}; \mathbb{R}^n)$ satisfies (H1)–(H7) and let u_1 and u_2 be two entropy solutions of (13), then*

$$(17) \quad \int_{\mathbb{R}^n} |u_1(T, x) - u_2(T, x)| dx \leq \int_{\mathbb{R}^n} |u_1(0, x) - u_2(0, x)| dx.$$

Remark 3.6. Since we are dealing with bounded solutions, one can suitably localize assumptions (H1)–(H7) in the “vertical” variable v , see for instance [3]. Furthermore one can also localize in the space variable x by just requiring for instance that $\mathbf{A}(\cdot, v)$ belongs to SBV_{loc} and similarly for u . Since this will not add any new ideas to the proof below, we leave this generalization to the interested reader. Moreover by exploiting standard techniques in the context of hyperbolic equation the following localized version of (17) can be easily obtained from the proof of Theorem 3.5:

$$\int_{B_R(0)} |u_1(T, x) - u_2(T, x)| dx \leq \int_{B_{R+VT}(0)} |u_1(0, x) - u_2(0, x)| dx \quad \forall R \geq 0,$$

where $V := \|\mathbf{A}\|_\infty$.

Since $u \in L^1((0, T); BV(\mathbb{R}^n))$, by the discussion in Section 2.3, see (5) in particular, we have that

$$\mathbf{A}(x, u(t, x)) \in L^1((0, T); BV(\mathbb{R}^n; \mathbb{R}^n)).$$

By arguing for instance as in [12, Theorem 4.3.1] we then deduce the following:

Lemma 3.7. *Let u be an entropy solution of (13), then*

$$u \in BV((0, T) \times \mathbb{R}^n).$$

Obviously we can think of $\mathbf{A}(\cdot, v)$ as a function in $SBV((0, T) \times \mathbb{R}^n; \mathbb{R}^n)$ constant in time, hence equations (13) and (7) give the following Rankine-Hugoniot conditions

$$(u^+ - u^-)\nu_t + (\mathbf{A}^+(x, u^+) - \mathbf{A}^-(x, u^-)) \cdot \nu_x = 0$$

for \mathcal{H}^n -a.e. $(t, x) \in ((0, T) \times \mathcal{N}) \cup J_u$,

where $\nu = (\nu_t, \nu_x)$ is the normal to the \mathcal{H}^n rectifiable set $((0, T) \times \mathcal{N}) \cup J_u \subset (0, T) \times \mathbb{R}^n$. In particular, since for \mathcal{H}^n almost every (t, x) in $((0, T) \times \mathcal{N}) \cap J_u$ we have $\nu = (0, \nu_{\mathcal{N}})$, we obtain

$$(18) \quad A^+(x, u^+) = A^-(x, u^-) \quad \text{for } \mathcal{H}^n\text{-a.e. } (t, x) \in ((0, T) \times \mathcal{N}) \cap J_u,$$

where we have introduced the notation

$$(19) \quad A^\pm(x, v) = \mathbf{A}^\pm(x, v) \cdot \nu_{\mathcal{N}} \quad v \in \mathbb{R}.$$

3.2. Analysis of the entropy condition on discontinuities. Let u be an entropy solution of (13). Thanks to (7) and (16), on $(0, T) \times \mathcal{N}$ the entropy inequality (15) reads as

$$(20) \quad \eta^+(x, u^+) - \eta^+(x, \hat{u}) + A^+(x, \hat{u})S'(\hat{u}) \leq \eta^-(x, u^-) - \eta^-(x, \hat{u}) + A^-(x, \hat{u})S'(\hat{u}),$$

\mathcal{H}^n almost everywhere on $(0, T) \times \mathcal{N}$ and we have dropped the dependence on (t, x) from u^\pm and \hat{u} in order to simplify the notations. Here we understand that $u^\pm = \tilde{u}$ if $x \notin J_u$ and we are using the short hand notations (19) and

$$\eta^\pm(x, v) := \boldsymbol{\eta}^\pm(x, v) \cdot \nu_{\mathcal{N}}(x).$$

Thanks to the definition of $\boldsymbol{\eta}$, (14), and (12), condition (20) can be rewritten as

$$(21) \quad A^+(x, u^+)S'(u^+) - \int_{\hat{u}}^{u^+} A^+(x, v)S''(v) dv$$

$$\leq A^-(x, u^-)S'(u^-) - \int_{\hat{u}}^{u^-} A^-(x, v)S''(v) dv.$$

By a standard approximation argument, we can plug in (21) the Kruzkov-type entropies

$$S(v) := |v - c|, \quad \eta_i(v) := \int_0^v \partial_v A_i(x, w) \text{sign}(w - c) dw,$$

where $c \in \mathbb{R}$ is a constant. We then obtain \mathcal{H}^n almost everywhere on $(0, T) \times \mathcal{N}$

$$(22) \quad A^+(x, u^+) \text{sign}(u^+ - c) - 2 \text{sign}(u^+ - \hat{u})A^+(x, c)\mathbf{1}_{(\hat{u}, u^+)}(c)$$

$$\leq A^-(x, u^-) \text{sign}(u^- - c) - 2 \text{sign}(u^- - \hat{u})A^-(x, c)\mathbf{1}_{(\hat{u}, u^-)}(c).$$

Here and in the sequel, for $a, b \in \mathbb{R}$ the symbol $\mathbf{1}_{(a, b)}$ will denote the characteristic function of the open interval $I(a, b)$ with endpoints a and b , i.e. $I(a, b) = (a, b)$ if $a < b$ or $I(a, b) = (b, a)$ if $b < a$.

We shall now derive as a consequence of (22) some inequalities which will be useful in the last part of the proof of Theorem 3.5. To this end, let us now fix a point $(t, x) \in$

$(0, T) \times \mathcal{N}$ for which (22) and (18) are valid and let $I(u^-, u^+)$ be the open interval with endpoints $u^-(t, x)$ and $u^+(t, x)$. Let us consider the following two cases:

- (a) $\hat{u} \in I(u^-, u^+)$
- (b) $\hat{u} \notin I(u^-, u^+)$.

In the case (a), taking into account the Rankine–Hugoniot condition (18), by (22) we get

$$(23) \quad [\text{sign}(u^+ - c) - \text{sign}(u^- - c)] A^+(u^+) \\ \leq 2 \text{sign}(u^+ - u^-) [A^+(c) \mathbf{1}_{(\hat{u}, u^+)}(c) + A^-(c) \mathbf{1}_{(\hat{u}, u^-)}(c)]$$

where we have dropped the dependence on x from A . This condition gives information only for $c \in I(u^-, u^+)$:

$$(24) \quad \text{sign}(u^+ - u^-) A^+(u^+) \\ \leq \text{sign}(u^+ - u^-) [A^+(c) \mathbf{1}_{(\hat{u}, u^+)}(c) + A^-(c) \mathbf{1}_{(\hat{u}, u^-)}(c)], \quad c \in I(u^-, u^+).$$

As particular cases, taking $c \nearrow \hat{u}$ and $c \searrow \hat{u}$ we get

$$\text{sign}(u^+ - u^-) A^+(u^+) \leq \text{sign}(u^+ - u^-) A^\pm(\hat{u}).$$

Similarly, in the case (b), taking again into account the Rankine–Hugoniot condition (18), by (22) we get

$$(25) \quad [\text{sign}(u^+ - c) - \text{sign}(u^- - c)] A^+(u^+) \\ \leq 2 \text{sign}(u^+ - \hat{u}) [A^+(c) \mathbf{1}_{(\hat{u}, u^+)}(c) - A^-(c) \mathbf{1}_{(\hat{u}, u^-)}(c)], \quad c \notin I(u^-, u^+).$$

We will now analyze conditions (23) and (25) in all the possible cases of different positions of u^- , u^+ , \hat{u} and c (see [24] for a similar analysis). We list all the cases for reader's convenience. First of all we remark that if $c \geq \max\{u^+, u^-, \hat{u}\}$ or $c \leq \min\{u^+, u^-, \hat{u}\}$, then, by the Rankine–Hugoniot condition (18), condition (22) does not give any information. Therefore we list all other possible cases:

Case 1: $u^+ \leq u^-$.

Subcase 1a: $u^+ \leq u^- \leq \hat{u}$

$$(26) \quad (i) \quad u^+ \leq u^- \leq c \leq \hat{u} \quad \Rightarrow \quad A^+(c) \leq A^-(c)$$

$$(27) \quad (ii) \quad u^+ \leq c \leq u^- \leq \hat{u} \quad \Rightarrow \quad A^+(c) \leq A^+(u^+).$$

Subcase 1b: $u^+ \leq \hat{u} \leq u^-$

$$(28) \quad (iii) \quad u^+ \leq \hat{u} \leq c \leq u^- \quad \Rightarrow \quad A^-(c) \leq A^-(u^-)$$

$$(29) \quad (iv) \quad u^+ \leq c \leq \hat{u} \leq u^- \quad \Rightarrow \quad A^+(c) \leq A^+(u^+).$$

Subcase 1c: $\hat{u} \leq u^+ \leq u^-$

$$(30) \quad (v) \quad \hat{u} \leq u^+ \leq c \leq u^- \quad \Rightarrow \quad A^+(c) \leq A^+(u^+)$$

$$(31) \quad (vi) \quad \hat{u} \leq c \leq u^+ \leq u^- \quad \Rightarrow \quad A^-(c) \leq A^+(c).$$

Case 2: $u^- \leq u^+$.

Subcase 2a: $u^- \leq u^+ \leq \hat{u}$

$$(32) \quad (i) \quad u^- \leq u^+ \leq c \leq \hat{u} \quad \Rightarrow \quad A^+(c) \leq A^-(c)$$

$$(33) \quad (ii) \quad u^- \leq c \leq u^+ \leq \hat{u} \quad \Rightarrow \quad A^-(u^-) \leq A^-(c).$$

Subcase 2b: $u^- \leq \hat{u} \leq u^+$

$$(34) \quad (iii) \quad u^- \leq \hat{u} \leq c \leq u^+ \quad \Rightarrow \quad A^+(u^+) \leq A^+(c)$$

$$(35) \quad (iv) \quad u^- \leq c \leq \hat{u} \leq u^+ \quad \Rightarrow \quad A^-(u^-) \leq A^-(c).$$

Subcase 2c: $\hat{u} \leq u^- \leq u^+$

$$(36) \quad (v) \quad \hat{u} \leq u^- \leq c \leq u^+ \quad \Rightarrow \quad A^+(u^+) \leq A^+(c)$$

$$(37) \quad (vi) \quad \hat{u} \leq c \leq u^- \leq u^+ \quad \Rightarrow \quad A^-(c) \leq A^+(c).$$

3.3. Kinetic formulation. Let us define the function $\chi: \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$(38) \quad \chi(v, u) := \begin{cases} 1 & \text{if } v < u, \\ 1/2 & \text{if } v = u, \\ 0 & \text{if } u < v. \end{cases}$$

Note that if $u \in BV((0, T) \times \mathbb{R}^n)$ then for almost every $v \in \mathbb{R}$ the function $(t, x) \mapsto \chi(v, u(t, x))$ belongs to SBV_{loc} .

Let \mathbf{A} satisfy conditions (H1)–(H7), and let us define

$$a_i(x, v) := \partial_v A_i(x, v) \quad \forall i = 1, \dots, n$$

$$a_{n+1}(\cdot, v) := - \sum_{i=1}^n \partial_i A_i(\cdot, v).$$

We remark that, for every $v \in \mathbb{R}$, $a_i(\cdot, v)$ is a BV function, while $a_{n+1}(\cdot, v)$ is a Radon measure, moreover according to (H5)

$$(39) \quad \sup_v |a_{n+1}(\cdot, v)|(\mathbb{R}^n) \leq \sigma(\mathbb{R}^n) < +\infty.$$

Let us denote by

$$\mathbf{a}(\cdot, v) := (a_1(\cdot, v) \mathcal{L}_x^n, \dots, a_n(\cdot, v) \mathcal{L}_x^n, a_{n+1}(\cdot, v)).$$

Note that \mathbf{a} is a Radon measure and that $\operatorname{div}_{x,v} \mathbf{a} = 0$.

Definition 3.8 (Kinetic solutions). A function

$$u \in C([0, T]; L^1(\mathbb{R}^n)) \cap L^\infty((0, T) \times \mathbb{R}^n) \cap L^1((0, T); BV(\mathbb{R}^n))$$

is a kinetic solution of (13) if u is a solution to (13) in the sense of distributions, and there exists a (everywhere defined) Borel representative \hat{u} of u with $|\hat{u}(t, x)| \leq \|u\|_\infty$ and a positive measure $m(t, x, v)$ with $m((0, T) \times \mathbb{R}^{n+1}) < +\infty$ such that the function $(t, x, v) \mapsto \chi(v, \hat{u}(t, x))$ satisfies

$$(40) \quad \partial_t \chi(v, \hat{u}(t, x)) + \operatorname{div}_{x,v} [\mathbf{a}(x, v) \chi(v, \hat{u}(t, x))] = \partial_v (m(t, x, v))$$

in the sense of distributions.

Our first results establishes the equivalence between Definitions 3.2 and 3.8.

Theorem 3.9. *Let*

$$u \in C([0, T]; L^1(\mathbb{R}^n)) \cap L^\infty((0, T) \times \mathbb{R}^n) \cap L^1((0, T); BV(\mathbb{R}^n)).$$

Then u is an entropy solution to (13) if and only if it is a kinetic solution to (13).

Proof. We divide the proof in two steps.

Step 1. Let u be a kinetic solution and let $S \in C_c^\infty(\mathbb{R})$. By testing (40) with $\phi(t)\varphi(x)S'(v)$, we then obtain

$$\begin{aligned} & \int_{(0,T) \times \mathbb{R}^n \times \mathbb{R}} \phi'(t)\varphi(x)S'(v)\chi(v, \hat{u}(t, x)) dt dx dv \\ & + \sum_{i=1}^n \int_{(0,T) \times \mathbb{R}^n \times \mathbb{R}} \phi(t)\partial_i \varphi(x)S'(v)a_i(x, v)\chi(v, \hat{u}(t, x)) dt dx dv \\ (41) \quad & + \int_{(0,T) \times \mathbb{R}^n \times \mathbb{R}} \phi(t)\varphi(x)S''(v)\chi(v, \hat{u}(t, x)) da_{n+1}(x, v) dv dt \\ & = \int_{(0,T) \times \mathbb{R}^n \times \mathbb{R}} S''(v)\varphi(x)\phi(t) dm(t, x, v). \end{aligned}$$

Now, since S is compactly supported,

$$(42) \quad \int_{\mathbb{R}} S'(v)\chi(v, \hat{u}(t, x)) dv = \int_{-\infty}^{\hat{u}(t, x)} S'(v) dv = S(\hat{u}(t, x))$$

and, for $i = 1, \dots, n$,

$$(43) \quad \int_{\mathbb{R}} S'(v)a_i(x, v)\chi(v, \hat{u}(t, x)) dv = \int_{-\infty}^{\hat{u}(t, x)} S'(v)\partial_v A_i(x, v) dv = \bar{\eta}_i(x, \hat{u}(t, x)),$$

where

$$\bar{\eta}(x, v) := \int_{-\infty}^v S'(w)\partial_v \mathbf{A}(x, w) dw = \boldsymbol{\eta}(x, v) + \int_{-\infty}^0 S'(w)\partial_v \mathbf{A}(x, w) dw$$

by the definition of $\boldsymbol{\eta}$ in (14). Moreover

$$\begin{aligned} & \int_{\mathbb{R}^n \times \mathbb{R}} \varphi(x)S''(v)\chi(v, \hat{u}(t, x)) da_{n+1}(x, v) dv \\ (44) \quad & = - \sum_{i=1}^n \int_{\mathbb{R}^n \times \mathbb{R}} \varphi(x)S''(v)\chi(v, \hat{u}(t, x)) \nabla_i A_i(x, v) dv dx \\ & \quad - \int_{\mathcal{N} \times \mathbb{R}} \varphi(x)S''(v)\chi(v, \hat{u}(t, x)) (A^+(x, v) - A^-(x, v)) dv d\mathcal{H}^{n-1}(x), \end{aligned}$$

where we have used the short hand notation (19). Now by the discussion in Section 2.3, the map $v \mapsto A^\pm(x, v)$ is C^1 for \mathcal{H}^{n-1} almost every $x \in \mathcal{N}$ with derivative given by

$v \mapsto \partial_v A^\pm(x, v)$, hence for any such x

$$\begin{aligned}
& \int_{\mathbb{R}} S''(v) \chi(v, \hat{u}(t, x)) \left(A^+(x, v) - A^-(x, v) \right) dv \\
&= \int_{-\infty}^{\hat{u}(t, x)} S''(v) \left(A^+(x, v) - A^-(x, v) \right) dv \\
(45) \quad &= S'(\hat{u}(t, x)) \left(A^+(x, \hat{u}(t, x)) - A^-(x, \hat{u}(t, x)) \right) \\
&\quad - \int_{-\infty}^{\hat{u}(t, x)} S'(v) \left(\partial_v A^+(x, v) - \partial_v A^-(x, v) \right) dv \\
&= S'(\hat{u}(t, x)) \left(A^+(x, \hat{u}(t, x)) - A^-(x, \hat{u}(t, x)) \right) - \left(\bar{\eta}^+(x, \hat{u}(t, x)) - \bar{\eta}^-(x, \hat{u}(t, x)) \right)
\end{aligned}$$

where we are using for $\bar{\eta}$ the same convention (19) used for \mathbf{A} and $\boldsymbol{\eta}$. In the same way by Lemma 2.2, for almost every $x \in \mathbb{R}^n$ the map $v \mapsto \nabla_i A_i(x, v)$ is C^1 with derivative given by $\nabla_i \partial_v A_i(x, v)$, hence for any such x

$$\begin{aligned}
& \int_{\mathbb{R}} S''(v) \chi(v, \hat{u}(t, x)) \nabla_i A_i(x, v) dv \\
(46) \quad &= S'(\hat{u}(t, x)) \nabla_i A_i(x, \hat{u}(t, x)) - \int_{-\infty}^{\hat{u}(t, x)} S'(v) \nabla_i \partial_v A_i(x, v) dv \\
&= S'(\hat{u}(t, x)) \nabla_i A_i(x, \hat{u}(t, x)) - \nabla_i \bar{\eta}_i(x, \hat{u}(t, x)).
\end{aligned}$$

Combining (41), (42), (43), (44), (45) and (46) we deduce that if u is a kinetic solution of (13), then for every function $S \in C_c^\infty(\mathbb{R})$ we have

$$\begin{aligned}
& \partial_t S(u) + \operatorname{div} (\bar{\boldsymbol{\eta}}(x, u)) \\
&\quad - \operatorname{div} \bar{\boldsymbol{\eta}}(x, v) \Big|_{v=\hat{u}} + S'(\hat{u}) \operatorname{div} \mathbf{A}(x, v) \Big|_{v=\hat{u}} = - \int S''(w) dm(\cdot, \cdot, w)
\end{aligned}$$

in the sense of distribution. We now note that $\bar{\boldsymbol{\eta}} - \boldsymbol{\eta}$ is a function of the x variable only, hence the above equation implies that

$$\begin{aligned}
& \partial_t S(u) + \operatorname{div} (\boldsymbol{\eta}(x, u)) \\
(47) \quad &\quad - \operatorname{div} \boldsymbol{\eta}(x, v) \Big|_{v=\hat{u}} + S'(\hat{u}) \operatorname{div} \mathbf{A}(x, v) \Big|_{v=\hat{u}} = - \int S''(w) dm(\cdot, \cdot, w)
\end{aligned}$$

for every $S \in C_c^\infty(\mathbb{R})$. Using the very same approximation argument of the second step of the proof of Theorem 3 in [13], we conclude that (47) holds for every convex function S of class C^2 . Since $m \geq 0$, this fact implies that u is an entropy solution of (13).

Step 2. Let u be an entropy solution of (13), let us define the distribution

$$\begin{aligned}
(48) \quad m(t, x, v) &= \partial_t \int_0^v \chi(w, u(t, x)) dw \\
&\quad + \sum_{i=1}^n \partial_i \left\{ \int_0^v a_i(x, w) \chi(w, u(t, x)) dw \right\} + a_{n+1}(x, v) \chi(v, \hat{u}(t, x)).
\end{aligned}$$

Clearly (40) is satisfied in the sense of distributions, hence to conclude that u is a kinetic solution we only have to show that m is a positive measure with $m((0, T) \times \mathbb{R}^n \times \mathbb{R}) < \infty$.

First note that by testing (48) with $\phi(t, x)\psi(v)$ with $\text{spt } \psi \cap [-\|u\|_\infty, \|u\|_\infty] = \emptyset$ and recalling that $|\hat{u}| \leq \|u\|_\infty$ we obtain that, as a distribution,

$$m(t, x, v) = \partial_t u + \text{div } \mathbf{A}(x, u) = 0 \quad \text{outside } [0, T] \times \mathbb{R}^n \times [-\|u\|_\infty, \|u\|_\infty],$$

that is $\text{spt } m \subset [0, T] \times \mathbb{R}^n \times [-\|u\|_\infty, \|u\|_\infty]$. We want to show that m is a positive measure, to this end note that by the same computations of Step 1, u satisfies (47) for every $S \in C_c^\infty(\mathbb{R})$. If $\psi(v) \geq 0$ is a positive test function, since m vanishes outside $[0, T] \times \mathbb{R}^n \times [-\|u\|_\infty, \|u\|_\infty]$ we can construct a function $S \in C_c^\infty$ which is convex on the range of \hat{u} and for which

$$\int \psi(v) dm(t, x, v) = \int S''(v) dm(t, x, v).$$

By (48) and since u is an entropy solution we then deduce that

$$\int \psi(v) dm(t, x, v) \geq 0 \quad \forall \psi \geq 0,$$

hence m is a positive measure. Moreover choosing $\phi = 1$ on $[-\|u\|_\infty, \|u\|_\infty]$ so that $S(v) = v^2/2$ on the range of \hat{u} and integrating (47) we get

$$\int_{(0, T) \times \mathbb{R}^n \times \mathbb{R}} dm(t, x, v) \leq \frac{1}{2} \int_{\mathbb{R}^n} |u(0, x)|^2 dx + \int_{(0, T) \times \mathbb{R}^n \times \mathbb{R}} d|a_{n+1}| dt.$$

By (39) we finally conclude. \square

4. PRELIMINARY ESTIMATES

In this section we shall prove some preliminary estimates for approximate solutions that will be used in Section 5.

Given an entropy solution \hat{u} , let us define the function $f(t, x, v) := \chi(v, \hat{u}(t, x))$ and let us consider a regularization of f with respect to the v variable. More precisely, let $\varphi \in C_c^\infty([-1/2, 1/2])$ be such that $\varphi(w) = \varphi(-w)$, $\varphi \geq 0$ and $\int \varphi = 1$. Let

$$\varphi_\epsilon(v) = \frac{1}{\epsilon} \varphi\left(\frac{v}{\epsilon}\right)$$

denote the standard family of mollifiers. If we define

$$f_\epsilon(t, x, v) := (f(t, x, \cdot) * \varphi_\epsilon)(v) = (\chi(\cdot, u) * \varphi_\epsilon)(v) \Big|_{u=\hat{u}(t, x)},$$

then we have the following

Proposition 4.1. *The function f_ϵ satisfies the following equation:*

$$(49) \quad \partial_t f_\epsilon(t, x, v) + \text{div}_{x, v} [\mathbf{a}(x, v) f_\epsilon(t, x, v)] = \partial_v (m_\epsilon(t, x, v)) + r_\epsilon,$$

where

$$m_\epsilon(t, x, v) := (m(t, x, \cdot) * \varphi_\epsilon)(v),$$

and the commutator

$$r_\epsilon := -\text{div}_{x, v} ((\mathbf{a}f) * \varphi_\epsilon) + \text{div}_{x, v} (\mathbf{a}f_\epsilon)$$

is a measure on $(0, T) \times \mathbb{R}^n \times \mathbb{R}$ such that

$$(50) \quad \lim_{\epsilon \rightarrow 0} |r_\epsilon|((0, T) \times \mathbb{R}^n \times (c, d)) = 0$$

for every $c, d \in \mathbb{R}$.

Proof. The fact that f_ϵ satisfies (49) is evident, hence we only have to verify the last part of the statement. To this end note let us write, with obvious notations, r_ϵ as

$$\begin{aligned} r_\epsilon &= \operatorname{div}_x(\partial_v \mathbf{A} f_\epsilon) - \operatorname{div}_x((\partial_v \mathbf{A} f)_\epsilon) \\ &\quad + \partial_v(a_{n+1} f_\epsilon) - \partial_v((a_{n+1} f)_\epsilon) := r_{1,\epsilon} + r_{2,\epsilon} \end{aligned}$$

and let us show that both $r_{1,\epsilon}$ and $r_{2,\epsilon}$ are measure satisfying (50).

• *Computation of $r_{1,\epsilon}$:*

If we test the distribution $r_{1,\epsilon}$ with functions $\phi(t, x, v) = \phi_1(t)\phi_2(x)\phi_3(v)$ and we apply Fubini Theorem we obtain

$$\begin{aligned} \langle r_{1,\epsilon}, \phi \rangle &= - \int_{\mathbb{R}} \int_0^T dv dt \phi_1(t)\phi_3(v) \int dw \varphi(w) \\ &\quad \times \int_{\mathbb{R}^n} dx D_x \phi_2(x) \cdot (\partial_v \mathbf{A}(x, v) - \partial_v \mathbf{A}(x, v - \epsilon w)) f(t, x, v - \epsilon w) \\ (51) \quad &= \int_{\mathbb{R}} \int_0^T dv dt \phi_1(t)\phi_3(v) \int dw \varphi(w) \\ &\quad \times \int_{\mathbb{R}^n} dx \phi_2(x) \operatorname{div}_x \left[(\partial_v \mathbf{A}(x, v) - \partial_v \mathbf{A}(x, v - \epsilon w)) f(t, x, v - \epsilon w) \right]. \end{aligned}$$

Recall that, by the coarea formula, for \mathcal{L}^2 almost every (v, w) the set $\{(t, x) \in (0, T) \times \mathbb{R}^n : u(t, x) > v - \epsilon w\}$ is of finite perimeter in $(0, T) \times \mathbb{R}^n$ and that denoting by $J_{v,w} = \partial^* \{(t, x) \in (0, T) \times \mathbb{R}^n : u(t, x) > v - \epsilon w\}$ we have

$$D_x f(t, x, v - \epsilon w) = D_x^j f(t, x, v - \epsilon w) = \nu_{J_{v,w}} \mathcal{H}^{n-1} \llcorner J_{v,w}.$$

Exploiting the Leibniz rule in BV , [4, Exercise 3.97], we then find that for \mathcal{L}^2 almost every (v, w)

$$\begin{aligned} &\operatorname{div} \left[(\partial_v \mathbf{A}(x, v) - \partial_v \mathbf{A}(x, v - \epsilon w)) f(t, x, v - \epsilon w) \right] \\ &= \sum_{i=1}^n (\nabla_i \partial_v A_i(x, v) - \nabla_i \partial_v A_i(x, v - \epsilon w)) f(t, x, v - \epsilon w) \mathcal{L}_t^1 \times \mathcal{L}_x^n \\ (52) \quad &+ (\partial_v A^+(x, v) - \partial_v A^+(x, v - \epsilon w)) f^+(t, x, v - \epsilon w) \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \\ &- (\partial_v A^-(x, v) - \partial_v A^-(x, v - \epsilon w)) f^-(t, x, v - \epsilon w) \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \\ &+ (\widetilde{\partial_v \mathbf{A}}(x, v) - \widetilde{\partial_v \mathbf{A}}(x, v - \epsilon w)) \cdot \nu_{J_{v,w}} \\ &\quad \times (f^+(t, x, v - \epsilon w) - f^-(t, x, v - \epsilon w)) \mathcal{H}^n \llcorner J_{v,w} \setminus ((0, T) \times \mathcal{N}), \end{aligned}$$

where we are using the notation (19). According to (H3), (H7) and (9), (10) we then obtain from (51) and (52) that $r_{1,\epsilon}$ is a measure and that

$$\begin{aligned} &|r_{1,\epsilon}|((0, T) \times \mathbb{R}^n \times (c, d)) \\ (53) \quad &\leq T\omega(\epsilon)(d - c) \|g_2\|_{L^1(\mathbb{R}^n)} + 2T\omega(\epsilon)(d - c) \mathcal{H}^{n-1}(\mathcal{N}) \\ &+ \omega(\epsilon) \int \varphi(w) dw \int_{\mathbb{R}} \mathcal{H}^n(J_{v,w}) dv. \end{aligned}$$

Clearly the first two terms on the right hand side of (53) go to zero as ϵ goes to zero. For what concerns the last one we notice that by the coarea formula

$$\begin{aligned} \int \varphi(w)dw \int_{\mathbb{R}} \mathcal{H}^n(J_{v,w})dv &\leq \int \varphi(w)dw \int_{\mathbb{R}} \mathcal{H}^n(\partial^* \{u(t,x) > v\})dv \\ &\leq |Du|((0,T) \times \mathbb{R}^n). \end{aligned}$$

Hence also the third term in right hand side of (53) goes to zero.

• *Computation of $r_{2,\epsilon}$:*

A computation similar to the previous one shows that $r_{2,\epsilon}$ is given by the following measure:

(54)

$$\begin{aligned} r_{2,\epsilon} &= \left\{ \sum_{i=1}^n \int \nabla_i \partial_v A_i(x,v) f(t,x,v-\epsilon w) \varphi(w) dw \right. \\ &\quad \left. - \sum_{i=1}^n \int (\nabla_i A_i(x,v) - \nabla_i A_i(x,v-\epsilon w)) f(t,x,v-\epsilon w) \frac{\varphi'(w)}{\epsilon} dw \right\} \mathcal{L}_t^1 \times \mathcal{L}_x^n \times \mathcal{L}_v^1 \\ &\quad + \left[\int (\partial_v A^+(x,v) - \partial_v A^-(x,v)) f(t,x,v-\epsilon w) \varphi(w) dw \right. \\ &\quad \left. - \int (A^+(x,v) - A^+(x,v-\epsilon w)) f(t,x,v-\epsilon w) \frac{\varphi'(w)}{\epsilon} dw \right. \\ &\quad \left. + \int (A^-(x,v) - A^-(x,v-\epsilon w)) f(t,x,v-\epsilon w) \frac{\varphi'(w)}{\epsilon} dw \right] \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1, \end{aligned}$$

where we are again using the convention (19). Note that by (H2) and the discussion in Section 2.3, $|A^\pm(x,v) - A^\pm(x,v-\epsilon w)| \leq M\epsilon|w|$ for \mathcal{H}^{n-1} -a.e. $x \in \mathcal{N}$ and that, by (H4), $|\nabla_i A_i(x,v) - \nabla_i A_i(x,v-\epsilon w)| \leq g_1(x)\epsilon|w|$. Hence if we can show that the term in curly brackets (respectively in square bracket) in (54) goes to zero as ϵ goes to zero for $\mathcal{L}_t^1 \times \mathcal{L}_x^n \times \mathcal{L}_v^1$ almost every (t,x,v) (respectively for $\mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1$ almost every (t,x,v)), an application of the Lebesgue dominated convergence theorem applied with respect to the measure $\mathcal{L}_t^1 \times \mathcal{L}_x^n \times \mathcal{L}_v^1$ (respectively $\mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1$) will imply that $r_{2,\epsilon}$ satisfies (50). To this end let us write

$$\begin{aligned} &\int \nabla_i \partial_v A_i(x,v) f(t,x,v-\epsilon w) \varphi(w) dw \\ &\quad - \int (\nabla_i A_i(x,v) - \nabla_i A_i(x,v-\epsilon w)) f(t,x,v-\epsilon w) \frac{\varphi'(w)}{\epsilon} dw \\ (55) \quad &= \int \left\{ \nabla_i \partial_v A_i(x,v) \varphi(w) - \nabla_i \partial_v A_i(x,v) w \varphi'(w) \right\} f(t,x,v-\epsilon w) dw \\ &\quad - \int \left\{ \frac{\nabla_i A_i(x,v) - \nabla_i A_i(x,v-\epsilon w)}{\epsilon} - w \nabla_i \partial_v A_i(x,v) \right\} f(t,x,v-\epsilon w) \varphi'(w) dw. \end{aligned}$$

Since by Lemma 2.2, the map $v \rightarrow \nabla_i A_i(x,v)$ is differentiable for almost every x with derivative given by $\nabla_i \partial_v A_i(x,v)$, by the fundamental theorem of calculus and (H7) for

every such x we can estimate the second integral in the right hand side of (55) by

$$\omega(\epsilon) g_2(x) \int |w\phi'(w)|dw,$$

which goes to zero as $\epsilon \rightarrow 0$. For what concerns the first term we notice that $f(t, x, v - \epsilon w) \rightarrow f(t, x, w)$ for almost every (t, x, v) since

$$\mathcal{L}^1 \times \mathcal{L}^n \left(\{(t, x) : \hat{u}(t, x) = v\} \right) = 0$$

for all but countably many v . Hence the first term in the right hand side of (55) goes to

$$\nabla_i \partial_v A_i(x, v) f(t, x, v) \int [\varphi(w) - w\varphi'(w)]dw = 0.$$

The term in square bracket in (54) can be treated in the same way this time using that $v \mapsto A^\pm(x, v)$ is differentiable for \mathcal{H}^{n-1} almost every $x \in \mathcal{N}$ with derivative given by $\partial_v A^\pm(x, v)$ and that

$$\mathcal{L}^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \left(\{(t, x) : \hat{u}(t, x) = v\} \right) = 0$$

for all but countably many v . □

The next step concerns the derivation of the evolution equation satisfied by $f_\epsilon^2 := (f_\epsilon)^2$. In order to simplify the notation, we define the differential operator L by

$$Lg(t, x, v) := \partial_t g(t, x, v) + \operatorname{div}_{x,v}[\mathbf{a}(x, v)g(t, x, v)].$$

Lemma 4.2. *The function f_ϵ^2 satisfies the following equation*

$$Lf_\epsilon^2 = 2f_\epsilon^* Lf_\epsilon + \mathcal{R}[f_\epsilon]$$

that is

$$(56) \quad \begin{aligned} \partial_t f_\epsilon^2(t, x, v) + \operatorname{div}_{x,v}[\mathbf{a}(x, v)f_\epsilon^2(t, x, v)] \\ = 2(f_\epsilon(t, x, v))^* [\partial_v(m_\epsilon(t, x, v)) + r_\epsilon] + \mathcal{R}[f_\epsilon] \end{aligned}$$

where f_ϵ^* is defined as in (3) and

$$(57) \quad \begin{aligned} \mathcal{R}[f_\epsilon] := & \left\{ [(\partial_v A)^+ - (\partial_v A)^-](f_\epsilon^+ - f_\epsilon)(f_\epsilon - f_\epsilon^-) \right. \\ & \left. + (f_\epsilon^+ - f_\epsilon + f_\epsilon^- - f_\epsilon)[A^+ - A^-] \partial_v f_\epsilon \right\} \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1. \end{aligned}$$

Here we are using the convention (19).

Proof. Note that $f_\epsilon(t, x, v)$ is a BV_{loc} function with respect to all its variables. By the chain rule

$$\widetilde{\partial}_t f_\epsilon^2(t, x, v) = 2f_\epsilon^*(t, x, v) \widetilde{\partial}_t f_\epsilon(t, x, v), \quad \widetilde{D}_x f_\epsilon^2(t, x, v) = 2f_\epsilon^*(t, x, v) \widetilde{D}_x f_\epsilon(t, x, v)$$

for \mathcal{H}^{n+1} almost every $(t, x, v) \in \mathbb{R}^{n+2} \setminus J_{f_\epsilon}$. Since $v \mapsto f_\epsilon(t, x, v)$ is C^1 we also have (recall that $\hat{u}(t, x)$ is everywhere defined)

$$\partial_v f_\epsilon^2(t, x, v) = 2f_\epsilon(t, x, v) \partial_v f_\epsilon(t, x, v),$$

for every (t, x, v) . Moreover

$$\chi(\cdot, u) * \varphi_\epsilon = \frac{1}{\epsilon} \int_{-\infty}^u \varphi\left(\frac{v-w}{\epsilon}\right) dw$$

is a smooth function of u . According to this $J_{f_\epsilon^2} = J_{f_\epsilon} = J_u \times \mathbb{R}$. Hence, recalling that (19) is in force,

(58)

$$\begin{aligned} & Lf_\epsilon^2 - 2(f_\epsilon)^* Lf_\epsilon \\ &= \left\{ (\partial_v A^+(f_\epsilon^2)^+ - \partial_v A^-(f_\epsilon^2)^-) - 2f_\epsilon^* (\partial_v A^+ f_\epsilon^+ - \partial_v A^- f_\epsilon^-) \right\} \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1 \\ &+ \widetilde{\partial_v \mathbf{A}} \cdot \nu_{J_{f_\epsilon}} \left\{ ((f_\epsilon^2)^+ - (f_\epsilon^2)^-) - 2f_\epsilon^* (f_\epsilon^+ - f_\epsilon^-) \right\} \mathcal{H}_{t,x,v}^{n+1} \llcorner (J_{f_\epsilon} \setminus [(0, T) \times \mathcal{N} \times \mathbb{R}]) \\ &+ \left\{ (A^+ - A^-) (2f_\epsilon - 2f_\epsilon^*) \partial_v f_\epsilon \right\} \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1 \\ &+ \left\{ ((\partial_v A)^+ - (\partial_v A)^-) (f_\epsilon - 2f_\epsilon^*) f_\epsilon \right\} \mathcal{L}_t^1 \times \mathcal{H}^{n-1} \llcorner \mathcal{N} \times \mathcal{L}_v^1, \end{aligned}$$

where we have used that $\partial_v(A^\pm) = (\partial_v A)^\pm$ for \mathcal{H}^{n-1} almost every x in \mathcal{N} and that $f_\epsilon^* = f_\epsilon$, for $\mathcal{L}^n + |\widetilde{D}_{t,x} f_\epsilon|$ almost every x , see Section 2.3. Since $(f_\epsilon^2)^\pm = (f_\epsilon^\pm)^2$ we have

$$(f_\epsilon^2)^+ - (f_\epsilon^2)^- - 2f_\epsilon^* (f_\epsilon^+ - f_\epsilon^-) = 0,$$

hence the second line in the right hand side of (58) vanishes. A simple algebraic computation now shows that (58) reduces to (57). \square

Let us now consider, for $R > 0$, the following test function $\psi_R \in C_c^\infty(\mathbb{R}^{n+1})$, $\psi_R \geq 0$ defined by

$$(59) \quad \psi_R(x, v) = \theta\left(\frac{x}{R}\right) \phi_R(v),$$

with $\theta \in C_c^\infty(\mathbb{R}^n)$, $\phi_R \in C^\infty(\mathbb{R})$, $\phi_R(v) = 1$ if $|v| \leq R$, $\phi_R(v) = 0$ if $|v| \geq R+1$, $\theta(x) = 1$ if $|x| \leq 1$ and $|\phi_R'| \leq 2$.

Lemma 4.3. *If u is a kinetic solution of (13), then*

$$\begin{aligned} (60) \quad & \lim_{R \rightarrow +\infty} \lim_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}^{n+1}} (\nu * \varphi_\epsilon) \psi_R dm_\epsilon \\ &= - \lim_{R \rightarrow +\infty} \lim_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}^{n+1}} \psi_R(x, v) d\mathcal{R}[f_\epsilon] \\ &= \int_0^T \int_{\mathcal{N}} Q(u) d\mathcal{H}^{n-1} dt, \end{aligned}$$

where (recall (19))

$$\begin{aligned} (61) \quad & Q(u) := \chi(u^-, u^+) A^+(u^-) + \chi(u^+, u^-) A^+(u^+) \\ & - \chi(u^+, u^-) A^-(u^+) - \chi(u^-, u^+) A^-(u^-) \\ & - \chi(u^+, \hat{u}) A^+(u^+) + \chi(u^+, \hat{u}) A^-(u^+) \\ & - \chi(u^-, \hat{u}) A^+(u^-) + \chi(u^-, \hat{u}) A^-(u^-), \end{aligned}$$

and

$$\nu(t, x, v) := \delta_{u+(t,x)}(v) + \delta_{u-(t,x)}(v) = -2\partial_v f^* .$$

Proof. Let $\eta_\delta \in C_c^1(0, T)$ with $\eta = 1$ on $[\delta, T - \delta]$. Since $u \in C([0, T]; L^1(\mathbb{R}^n))$, we have that $f_\epsilon \in C([0, T]; L^1(\mathbb{R}^{n+1}))$, hence by testing (56) and (49) with $\eta_\delta(t)\psi_R(x, v)$ and letting $\delta \rightarrow 0$ we obtain by standard computations

$$(62) \quad \int_{\mathbb{R}^{n+1}} [f_\epsilon^2(T, x, v) - f_\epsilon(T, x, v)] \psi_R(x, v) \, dx \, dv$$

$$(63) \quad = \int_{\mathbb{R}^{n+1}} [f_\epsilon^2(0, x, v) - f_\epsilon(0, x, v)] \psi_R(x, v) \, dx \, dv$$

$$(64) \quad + \int_0^T \int_{\mathbb{R}^{n+1}} (f_\epsilon^2 - f_\epsilon) \nabla_{x,v} \psi_R \cdot d\mathbf{a}(x, v) \, dt$$

$$(65) \quad + \int_0^T \int_{\mathbb{R}^{n+1}} [-2\partial_v f_\epsilon^*] \psi_R \, dm_\epsilon$$

$$(66) \quad + \int_0^T \int_{\mathbb{R}^{n+1}} [1 - 2f_\epsilon^*] \partial_v \psi_R \, dm_\epsilon$$

$$(67) \quad + \int_0^T \int_{\mathbb{R}^{n+1}} [2f_\epsilon^* - 1] \psi_R \, dr_\epsilon$$

$$(68) \quad + \int_0^T \int_{\mathbb{R}^{n+1}} \psi_R \, d\mathcal{R}[f_\epsilon](x, v) \, dt .$$

As ϵ goes to 0, the integral (62) tends to

$$\int_{\mathbb{R}^{n+1}} [f^2(T, x, v) - f(T, x, v)] \psi_R(x, v) \, dx \, dv = 0,$$

since $f^2 = f$. In the same way the integral (63) tends to 0. Moreover, recalling the definition of \mathbf{a} we obtain that

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^{n+1}} (f_\epsilon^2 - f_\epsilon) \nabla_{x,v} \psi_R(x, v) \cdot d\mathbf{a}(x, v) \, dt \\ &= \int_0^T \int_{\mathbb{R}^{n+1}} (f_\epsilon^2 - f_\epsilon) \nabla_x \psi_R \cdot \partial_v \mathbf{A} \, dx \, dv \, dt \\ &+ \int_0^T \int_{\mathbb{R}^{n+1}} (f_\epsilon^2 - f_\epsilon) \partial_v \psi_R \, da_{n+1} \, dt . \end{aligned}$$

We now note that, arguing as the end of the proof of Lemma 4.1, $f_\epsilon^2 \rightarrow f^2 = f$, for $\mathcal{L}_t^1 \times (\mathcal{L}^n + \sigma)_x \times \mathcal{L}_v^1$ almost every point. This and the fact that $|a_{n+1}| \ll \mathcal{L}^n + \sigma$, see (39), imply that the integral (64) tends to 0, as $\epsilon \rightarrow 0$.

Let us now we consider integral (65). We remark that

$$\partial_v f_\epsilon^* = -\frac{1}{2} \nu * \phi_\epsilon = -\frac{1}{2} [\delta_{u-(t,x)}(v) + \delta_{u+(t,x)}(v)] * \phi_\epsilon .$$

Therefore the integral (65) is equal to

$$\int_0^T \int_{\mathbb{R}^{n+1}} (\nu * \phi_\epsilon) \psi_R \, dm_\epsilon .$$

Let us now estimate the integral (66). Recalling the choice of the test function made in (59), we have that

$$\left| \int_0^T \int_{\mathbb{R}^{n+1}} m_\epsilon(t, x, v) [1 - 2f_\epsilon^*] \partial_v \psi_R(x, v) dt dx dv \right| \leq 2 m_\epsilon([0, T] \times \mathbb{R}^n \times I_R),$$

where $I_R = \{v : R \leq |v| \leq R + 1\}$. Hence, by letting $\epsilon \rightarrow 0$ and $R \rightarrow +\infty$, the integral (66) tends to 0 since $m([0, T] \times \mathbb{R}^n \times \mathbb{R}) < +\infty$.

By (50), the integral in (67) tends to 0 as $\epsilon \rightarrow 0$. Gathering all the information above, the first equality in (60) is proved.

It remains to compute

$$- \lim_{R \rightarrow +\infty} \lim_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}^{n+1}} \psi_R(x, v) d\mathcal{R}[f_\epsilon]$$

and to show that the second equality in (60) holds. We recall that the explicit form of $\mathcal{R}[f_\epsilon]$ is given in Lemma 4.2. We have that

$$\begin{aligned} & - \int_0^T \int_{\mathbb{R}^{n+1}} \psi_R(x, v) d\mathcal{R}[f_\epsilon] \\ &= - \int_0^T \int_{\mathbb{R}^{n+1}} (f_\epsilon^+ - f_\epsilon)(f_\epsilon^- - f_\epsilon)(A^+ - A^-) \partial_v \psi_R dv d\mathcal{H}^{n-1}(x) dt \\ & \quad - \int_0^T \int_{\mathbb{R}^{n+1}} [(f_\epsilon^+ - f_\epsilon) \partial_v (f_\epsilon^- - f_\epsilon) + (f_\epsilon^- - f_\epsilon) \partial_v (f_\epsilon^+ - f_\epsilon) \\ & \quad \quad + (f_\epsilon^+ - f_\epsilon + f_\epsilon^- - f_\epsilon) \partial_v f_\epsilon] (A^+ - A^-) \psi_R dv d\mathcal{H}^{n-1}(x) dt \\ &=: I_1(\epsilon, R) + I_2(\epsilon, R). \end{aligned}$$

Again, the first integral $I_1(\epsilon, R)$ tends to 0 as $\epsilon \rightarrow 0$ and $R \rightarrow +\infty$ thanks to the choice of the test function ψ_R . In order to compute the $I_2(\epsilon, R)$, let us recall that

$$\partial_v (f_\epsilon^\pm - f_\epsilon) = [\delta_{\hat{u}}(v) - \delta_{u^\pm}(v)] * \varphi_\epsilon(v), \quad \partial_v f_\epsilon = -\delta_{\hat{u}}(v) * \varphi_\epsilon(v).$$

A straightforward computation based on Lemma 4.4 below, now gives that

$$(69) \quad \lim_{R \rightarrow +\infty} \lim_{\epsilon \rightarrow 0} I_2(\epsilon, R) = \int_0^T \int_{\mathcal{N}} Q(u) d\mathcal{H}^{n-1}(x) dt,$$

where

$$\begin{aligned} (70) \quad Q(u) &:= [-\chi(u^+, \hat{u}) + \chi(u^+, u^-)] [A^+(u^+) - A^-(u^+)] \\ & \quad + \left[\frac{1}{2} - \chi(\hat{u}, u^-) + \frac{1}{2} - \chi(\hat{u}, u^+) \right] [A^+(\hat{u}) - A^-(\hat{u})] \\ & \quad + [-\chi(u^-, \hat{u}) + \chi(u^-, u^+)] [A^+(u^-) - A^-(u^-)] \\ & \quad + [-1 + \chi(\hat{u}, u^-) + \chi(\hat{u}, u^+)] [A^+(\hat{u}) - A^-(\hat{u})], \end{aligned}$$

which, after some computations is easily seen to coincide with (61). The second equality in (60) now follows from this together with (69) and (70). \square

We conclude this section with the following technical lemma that we have used in the proof of (70), for later use we state the lemma in a slightly more general setting.

Lemma 4.4. *Let $\varphi_\epsilon : \mathbb{R} \rightarrow \mathbb{R}$ be a family of symmetric mollifiers and let $h_1, h_2 : \mathbb{R} \rightarrow \mathbb{R}$ be two bounded and uniformly continuous functions, then for every $u, \hat{u} \in \mathbb{R}$*

$$(71) \quad \lim_{\epsilon \rightarrow 0} \int h_1(v) (\delta_u * \varphi_\epsilon)(v) [h_2(\cdot) \chi(\cdot, \hat{u})] * \varphi_\epsilon(v) dv = h_1(u) h_2(u) \chi(u, \hat{u}).$$

Here $\chi(u, \hat{u})$ is defined according to (38).

Proof. We start noticing that

$$\begin{aligned} I &:= \int h_1(v) (\delta_u * \varphi_\epsilon)(v) [h_2(\cdot) \chi(\cdot, \hat{u})] * \varphi_\epsilon(v) dv \\ &= \iint h_1(v) \varphi_\epsilon(v - u) h_2(w) \chi(w, \hat{u}) \varphi_\epsilon(v - w) dw dv \\ &= \iint h_1(v) \varphi_\epsilon(v - u) [h_2(w) - h_2(u)] \chi(w, \hat{u}) \varphi_\epsilon(v - w) dw dv \\ &\quad + h_2(u) \iint h_1(v) \varphi_\epsilon(v - u) \chi(w, \hat{u}) \varphi_\epsilon(v - w) dw dv \\ &=: I_1 + h_2(u) I_2. \end{aligned}$$

By exploiting the uniform continuity of h_2 we obtain for some modulus of continuity ω , that

$$(72) \quad \begin{aligned} |I_1| &\leq \iint |h_1(v)| \varphi_\epsilon(v - u) \omega(|v - w|) \varphi_\epsilon(v - w) dw dv \\ &\leq \|h_1\|_\infty \int \left(\omega(|s|) \varphi_\epsilon(s) \int \varphi_\epsilon(s + w - u) dw \right) dt \\ &\leq \|h_1\|_\infty \int \omega(|s|) \varphi_\epsilon(s) ds \end{aligned}$$

and the last integral tends to 0 as $\epsilon \rightarrow 0$. On the other hand

$$(73) \quad \lim_{\epsilon \rightarrow 0^+} I_2 = h_1(u) \chi(u, \hat{u}).$$

Indeed, from the estimate

$$\begin{aligned} |I_2 - h_1(u) \chi(u, \hat{u})| &\leq \max_{v \in [u - \epsilon, u + \epsilon]} |h_1(v) - h_1(u)| \\ &\quad + |h_1(u)| \left| \iint \varphi_\epsilon(v - u) \chi(w, \hat{u}) \varphi_\epsilon(v - w) dw dv - \chi(u, \hat{u}) \right| \end{aligned}$$

and the uniform continuity of h_1 it is enough to show that

$$(74) \quad \lim_{\epsilon \rightarrow 0^+} \iint \varphi_\epsilon(v - u) \chi(w, \hat{u}) \varphi_\epsilon(v - w) dw dv = \chi(u, \hat{u}).$$

If $u \neq \hat{u}$, then for ϵ small enough we have that $\chi(w, \hat{u}) = \chi(u, \hat{u})$ for $w \in (u - 2\epsilon, u + 2\epsilon)$, so that χ is constant in the integral. Then

$$\iint \varphi_\epsilon(v - u) \chi(w, \hat{u}) \varphi_\epsilon(v - w) dw dv = \chi(u, \hat{u}) \iint \varphi_\epsilon(v - u) \varphi_\epsilon(v - w) dw dv = \chi(u, \hat{u})$$

and (74) follows. If $u = \hat{u}$, then the integral in (74), for $\epsilon > 0$ small enough, becomes

$$\begin{aligned} \iint \varphi_\epsilon(v-u) \chi(w, u) \varphi_\epsilon(v-w) dw dv &= \int_{-\infty}^{+\infty} \varphi_\epsilon(v-u) \int_{-\infty}^u \varphi_\epsilon(v-w) dw dv \\ &= \int_{-\infty}^{+\infty} \varphi_\epsilon(v-u) \left[\frac{1}{2} + \int_v^u \varphi_\epsilon(v-w) dw \right] dv = \frac{1}{2}, \end{aligned}$$

where we have exploited that $\varphi_\epsilon(s) = \varphi_\epsilon(-s)$. Equation (74) now follows from the definition of χ , (38). Hence (71) follows from (72) and (73). \square

5. UNIQUENESS

In this Section we prove Theorem 3.5, to this end let us fix some notation. For u_1, u_2 two entropy solutions of (13), with corresponding everywhere defined Borel representatives \hat{u}_1, \hat{u}_2 , we will set $f_i(t, x, v) := \chi(v, \hat{u}_i(t, x))$ and m_i , $i = 1, 2$, for the corresponding functions and measures appearing in the kinetic formulation. We will also set

$$f_{1\epsilon}(t, x, v) := (f_1(t, x, \cdot) * \varphi_\epsilon)(v), \quad f_{2\epsilon}(t, x, v) := (f_2(t, x, \cdot) * \varphi_\epsilon)(v).$$

and

$$m_{1\epsilon}(t, x, v) := (m_1(t, x, \cdot) * \varphi_\epsilon)(v), \quad m_{2\epsilon}(t, x, v) := (m_2(t, x, \cdot) * \varphi_\epsilon)(v),$$

where φ_ϵ denotes the standard family of mollifiers.

The following theorem is the main result of this section and, by Cavalieri's principle, it immediately implies Theorem 3.5.

Theorem 5.1. *Let u_1, u_2 be two entropy solution of (13), with corresponding everywhere defined Borel representatives \hat{u}_1, \hat{u}_2 . Setting $f_i(t, x, v) := \chi(v, \hat{u}_i(t, x))$, $i = 1, 2$, we have that*

$$\int_{\mathbb{R}^{n+1}} |f_1 - f_2|(T, x, v) dx dv \leq \int_{\mathbb{R}^{n+1}} |f_1 - f_2|(0, x, v) dx dv.$$

In order to prove Theorem 5.1 we need some preliminary results concerning the interaction of two kinetic solutions:

Proposition 5.2. *With the notation above, for every test function $\psi(x, v)$ we have that*

$$\begin{aligned} \int_{\mathbb{R}^{n+1}} |f_1 - f_2|(T, x, v) \psi(x, v) dx dv &\leq \int_{\mathbb{R}^{n+1}} |f_1 - f_2|(0, x, v) \psi(x, v) dx dv \\ &\quad - \limsup_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}^{n+1}} 2\partial_v[(f_{1\epsilon} - f_{2\epsilon})^* \psi] d(m_{1\epsilon} - m_{2\epsilon})(t, x, v) \\ (75) \quad &\quad + \limsup_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}^{n+1}} \psi d\mathcal{R}[f_{1\epsilon} - f_{2\epsilon}](t, x, v) \\ &\quad - \int_0^T \int_{\mathbb{R}^{n+1}} |f_1 - f_2| \nabla_x \psi \cdot \partial_v \mathbf{A}(x, v) dx dv dt \\ &\quad + \int_0^T \int_{\mathbb{R}^{n+1}} |f_1 - f_2| \partial_v \psi da_{n+1}(x, v) dt, \end{aligned}$$

where $\mathcal{R}[f_{1\epsilon} - f_{2\epsilon}]$ is defined according to (57).

Proof. We recall that

$$Lf_{1\epsilon} = \partial_v m_{1\epsilon} + r_{1\epsilon}, \quad Lf_{2\epsilon} = \partial_v m_{2\epsilon} + r_{2\epsilon}.$$

Since $f_{1\epsilon} - f_{2\epsilon}$ is a solution of (56), by Lemma 4.2 we see that

$$\begin{aligned} & \partial_t (f_{1\epsilon} - f_{2\epsilon})^2 + \operatorname{div}_{x,v} [\mathbf{a}(f_{1\epsilon} - f_{2\epsilon})^2] \\ &= 2(f_{1\epsilon} - f_{2\epsilon})^* L(f_{1\epsilon} - f_{2\epsilon}) + \mathcal{R}[f_{1\epsilon} - f_{2\epsilon}]. \end{aligned}$$

Hence, by arguing as in Lemma 4.3, for every test function $\psi(x, v)$ we have that

$$\begin{aligned} & \int_{\mathbb{R}^{n+1}} (f_{1\epsilon} - f_{2\epsilon})^2(T, x, v) \psi(x, v) dx dv - \int_{\mathbb{R}^{n+1}} (f_{1\epsilon} - f_{2\epsilon})^2(0, x, v) \psi(x, v) dx dv \\ &= - \int_0^T \int_{\mathbb{R}^{n+1}} (f_{1\epsilon} - f_{2\epsilon})^2 \nabla_{x,v} \psi(x, v) \cdot d\mathbf{a}(x, v) dt \\ (76) \quad & + 2 \int_0^T \int_{\mathbb{R}^{n+1}} (f_{1\epsilon} - f_{2\epsilon})^* L(f_{1\epsilon} - f_{2\epsilon}) \psi(x, v) dx dv dt \\ & + \int_0^T \int_{\mathbb{R}^{n+1}} \psi(x, v) d(\mathcal{R}[f_{1\epsilon} - f_{2\epsilon}]) \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

Noticing that $(f_{1\epsilon} - f_{2\epsilon})^2 \rightarrow (f_1 - f_2)^2$, that the equality $(f_1 - f_2)^2 = |f_1 - f_2|^2$ holds for $\mathcal{L}_t^1 \times (\mathcal{L}^n + \sigma) \times \mathcal{L}_v^1$ almost every point, and that $|\mathbf{a}| \ll (\mathcal{L}^n + \sigma) \times \mathcal{L}_v^1$, we can pass to the limit in I_1 obtaining the last two integrals in the right hand side of (75). In the same way, the left hand side of (76) tends, as $\epsilon \rightarrow 0$, to

$$\int_{\mathbb{R}^{n+1}} |f_1 - f_2|(T, x, v) \psi(x, v) dx dv - \int_{\mathbb{R}^{n+1}} |f_1 - f_2|(0, x, v) \psi(x, v) dx dv.$$

Let us now consider I_2 and I_3 . Since, by Proposition 4.1,

$$Lf_{1\epsilon} = \partial_v m_{1\epsilon} + r_{1\epsilon}, \quad Lf_{2\epsilon} = \partial_v m_{2\epsilon} + r_{2\epsilon}$$

we infer that inequality (75) holds. \square

Proposition 5.3. *Let u_1, u_2 be two entropy solution of (13), with corresponding representatives \hat{u}_1, \hat{u}_2 . Setting $f_i(t, x, v) := \chi(v, \hat{u}_i(t, x))$, $i = 1, 2$, we have that*

$$\begin{aligned} (77) \quad \int_{\mathbb{R}^{n+1}} |f_1 - f_2|(T, x, v) dx dv &\leq \int_{\mathbb{R}^{n+1}} |f_1 - f_2|(0, x, v) dx dv \\ &+ \int_0^T \int_{\mathcal{N}} W(u_1, u_2) d\mathcal{H}^{n-1} dt, \end{aligned}$$

where

$$\begin{aligned} (78) \quad W(u_1, u_2) &:= A^+(u_1^+) [-2\chi(u_1^+, u_2^+) + 2\chi(u_1^-, u_2^-)] \\ &+ A^+(u_2^+) [-2\chi(u_2^+, u_1^+) + 2\chi(u_2^-, u_1^-)]. \end{aligned}$$

Proof. Let us start from inequality (75) using the test function $\psi = \psi_R$ defined in (59). It is easy to show that, as $R \rightarrow +\infty$, the last two integrals in (75) goes to 0. Hence, we only need to estimate the contribution of the two terms

$$I(\epsilon, R) := - \int_0^T \int_{\mathbb{R}^{n+1}} 2\partial_v [(f_{1\epsilon} - f_{2\epsilon})\psi_R] d(m_{1\epsilon} - m_{2\epsilon})(t, x, v)$$

and

$$H(\epsilon, R) := \int_0^T \int_{\mathbb{R}^{n+1}} \psi_R d(\mathcal{R}[f_{1\epsilon} - f_{2\epsilon}])(t, x, v)$$

for $\epsilon \rightarrow 0$ and $R \rightarrow +\infty$. We start noticing that

$$(79) \quad \begin{aligned} I(\epsilon, R) = & - \int_0^T \int_{\mathbb{R}^{n+1}} 2\partial_v [(f_{1\epsilon} - f_{2\epsilon})^*]\psi_R d(m_{1\epsilon} - m_{2\epsilon})(t, x, v) \\ & - \int_0^T \int_{\mathbb{R}^{n+1}} 2(f_{1\epsilon} - f_{2\epsilon})^* \partial_v \psi_R d(m_{1\epsilon} - m_{2\epsilon})(t, x, v), \end{aligned}$$

and, reasoning as in the final part of the proof of Lemma 4.3, the last integral goes to 0 as $\epsilon \rightarrow 0$ and $R \rightarrow +\infty$. Let us compute the first term in (79):

$$\begin{aligned} & - \int_0^T \int_{\mathbb{R}^{n+1}} 2\partial_v [(f_{1\epsilon} - f_{2\epsilon})^*]\psi_R d(m_{1\epsilon} - m_{2\epsilon}) \\ & = - \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_2^+}(v) + \delta_{u_2^-}(v) - \delta_{u_1^+}(v) - \delta_{u_1^-}(v)] * \varphi_\epsilon(v) \psi_R d(m_{1\epsilon} - m_{2\epsilon}) \\ & = \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_1^+}(v) + \delta_{u_1^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{1\epsilon} \\ & \quad + \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_2^+}(v) + \delta_{u_2^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{2\epsilon} \\ & \quad - \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_2^+}(v) + \delta_{u_2^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{1\epsilon} \\ & \quad - \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_1^+}(v) + \delta_{u_1^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{2\epsilon} \\ & =: I_1(\epsilon, R) + I_2(\epsilon, R) + I_3(\epsilon, R) + I_4(\epsilon, R). \end{aligned}$$

By Lemma 4.3,

$$\begin{aligned} & \limsup_{R \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} I_1(\epsilon, R) + I_2(\epsilon, R) \\ & \leq \int_0^T \int_{\mathcal{N}} [Q(u_1) + Q(u_2)] d\mathcal{H}^{n-1} dt, \end{aligned}$$

where $Q(u)$ is defined in (61). In order to estimate I_3 , we note that

$$\begin{aligned} I_3 = & - \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_2^+}(v) + \delta_{u_2^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{1\epsilon}(t, x, v) \\ & \leq - \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_2^+}(v) + \delta_{u_2^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{1\epsilon}^s(t, x, v), \end{aligned}$$

where $m_{1\epsilon}^s(t, \cdot, v)$ is the restriction to \mathcal{N} of the singular part of the measure $m_{1\epsilon}(t, \cdot, v)$, namely $m_{1\epsilon}^s(t, \cdot, v) = m_1^s(t, \cdot, v) * \varphi_\epsilon(v)$, with

$$m_1^s(t, x, v) = \int_0^v [(\partial_v A)^+(x, w)(\chi(w, u_1^+(t, x))) - (\partial_v A)^-(x, w)(\chi(w, u_1^-(t, x)))] d\mathcal{H}^{n-1} dw \\ - (A^+(x, v) - A^-(x, v)) \chi(v, \hat{u}_1(t, x)) d\mathcal{H}^{n-1}.$$

Hence, the contribution given by I_3 to (77) can be estimated by

$$- \limsup_{R \rightarrow +\infty} \limsup_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}^{n+1}} [\delta_{u_2^+}(v) + \delta_{u_2^-}(v)] * \varphi_\epsilon(v) \psi_R dm_{1\epsilon}^s.$$

By the explicit expression of $m_{1\epsilon}^s$ and by also taking into account Lemma 4.4, we get

$$I_3 \leq \int_0^T \int_{\mathcal{N}} \left[-\chi(u_2^+, u_1^+) A^+(u_2^+) - \chi(u_1^+, u_2^+) A^+(u_1^+) \right. \\ \left. + \chi(u_2^+, u_1^-) A^-(u_2^+) + \chi(u_1^-, u_2^+) A^-(u_1^-) \right. \\ \left. - \chi(u_2^+, \hat{u}_1) A^-(u_2^+) + \chi(u_2^+, \hat{u}_1) A^+(u_2^+) \right] d\mathcal{H}^{n-1} \\ + \int_0^T \int_{\mathcal{N}} \left[-\chi(u_2^-, u_1^+) A^+(u_2^-) - \chi(u_1^+, u_2^-) A^+(u_1^+) \right. \\ \left. + \chi(u_2^-, u_1^-) A^-(u_2^-) + \chi(u_1^-, u_2^-) A^-(u_1^-) \right. \\ \left. - \chi(u_2^-, \hat{u}_1) A^-(u_2^-) + \chi(u_2^-, \hat{u}_1) A^+(u_2^-) \right] d\mathcal{H}^{n-1}.$$

In the same way we can estimate the contribution of I_4 . The contribution of H can then be estimated with the aid of Lemma 4.4 by arguing as in the final part of the proof of Lemma 4.3. Therefore, by summing up all the estimates we obtain that (77) holds with the following choice of $W(u_1, u_2)$ where, for the reader convenience, the terms are grouped according to their provenience:

$$W(u_1, u_2) \\ = \left. \begin{aligned} & \chi(u_1^-, u_1^+) A^+(u_1^-) + \chi(u_1^+, u_1^-) A^+(u_1^+) - \chi(u_1^+, u_1^-) A^-(u_1^+) - \chi(u_1^-, u_1^+) A^-(u_1^-) \\ & - \chi(u_1^+, \hat{u}_1) A^+(u_1^+) + \chi(u_1^+, \hat{u}_1) A^-(u_1^+) - \chi(u_1^-, \hat{u}_1) A^+(u_1^-) + \chi(u_1^-, \hat{u}_1) A^-(u_1^-) \end{aligned} \right\} I_1 \\ \\ + \left. \begin{aligned} & \chi(u_2^-, u_2^+) A^+(u_2^-) + \chi(u_2^+, u_2^-) A^+(u_2^+) - \chi(u_2^+, u_2^-) A^-(u_2^+) - \chi(u_2^-, u_2^+) A^-(u_2^-) \\ & - \chi(u_2^+, \hat{u}_2) A^+(u_2^+) + \chi(u_2^+, \hat{u}_2) A^-(u_2^+) - \chi(u_2^-, \hat{u}_2) A^+(u_2^-) + \chi(u_2^-, \hat{u}_2) A^-(u_2^-) \end{aligned} \right\} I_2 \\ \\ - \left. \begin{aligned} & \chi(u_1^+, u_2^+) A^+(u_1^+) - \chi(u_1^-, u_2^+) A^+(u_1^-) - \chi(u_2^+, u_1^+) A^+(u_2^+) - \chi(u_2^+, u_1^-) A^+(u_2^+) \\ & + \chi(u_1^+, u_2^-) A^-(u_1^+) + \chi(u_1^-, u_2^-) A^-(u_1^-) + \chi(u_2^-, u_1^+) A^-(u_2^-) + \chi(u_2^-, u_1^-) A^-(u_2^-) \\ & + \chi(u_1^+, \hat{u}_2) A^+(u_1^+) - \chi(u_1^+, \hat{u}_2) A^-(u_1^+) + \chi(u_1^-, \hat{u}_2) A^+(u_1^-) - \chi(u_1^-, \hat{u}_2) A^-(u_1^-) \end{aligned} \right\} I_3$$

$$\left. \begin{aligned}
 & -\chi(u_2^+, u_1^+)A^+(u_2^+) - \chi(u_2^-, u_1^+)A^+(u_2^-) - \chi(u_1^+, u_2^+)A^+(u_1^+) - \chi(u_1^+, u_2^-)A^+(u_1^+) \\
 & + \chi(u_2^+, u_1^-)A^-(u_2^+) + \chi(u_2^-, u_1^-)A^-(u_2^-) + \chi(u_1^-, u_2^+)A^-(u_1^-) + \chi(u_1^-, u_2^-)A^-(u_1^-) \\
 & + \chi(u_2^+, \hat{u}_1)A^+(u_2^+) - \chi(u_2^+, \hat{u}_1)A^-(u_2^+) + \chi(u_2^-, \hat{u}_1)A^+(u_2^-) - \chi(u_2^-, \hat{u}_1)A^-(u_2^-)
 \end{aligned} \right\} I_4$$

$$\left. \begin{aligned}
 & -[\chi(u_1^+, u_1^-) - \chi(u_1^+, \hat{u}_1) - \chi(u_1^+, u_2^-) + \chi(u_1^+, \hat{u}_2)][A^+(u_1^+) - A^-(u_1^+)] \\
 & + [\chi(u_2^+, u_1^-) - \chi(u_2^+, \hat{u}_1) - \chi(u_2^+, u_2^-) + \chi(u_2^+, \hat{u}_2)][A^+(u_2^+) - A^-(u_2^+)] \\
 & - [\chi(u_1^-, u_1^+) - \chi(u_1^-, \hat{u}_1) - \chi(u_1^-, u_2^+) + \chi(u_1^-, \hat{u}_2)][A^+(u_1^-) - A^-(u_1^-)] \\
 & + [\chi(u_2^-, u_1^+) - \chi(u_2^-, \hat{u}_1) - \chi(u_2^-, u_2^+) + \chi(u_2^-, \hat{u}_2)][A^+(u_2^-) - A^-(u_2^-)].
 \end{aligned} \right\} H$$

We now note that all terms in $A^+(u_j^-)$ and $A^-(u_j^+)$, $j = 1, 2$, cancel out. Using the Rankine-Hugoniot conditions (18), we can sum up the terms $A^+(u_j^+)$ with $A^-(u_j^-)$, obtaining (78). \square

We are now ready to prove Theorem 5.1.

Proof of Theorem 5.1. By Proposition 5.3, it remains to prove that

$$(80) \quad A^+(u_1^+)[- \chi(u_1^+, u_2^+) + \chi(u_1^-, u_2^-)] + A^+(u_2^+)[- \chi(u_2^+, u_1^+) + \chi(u_2^-, u_1^-)] \leq 0.$$

Since u_1, u_2 are entropy solutions, recalling that

$$\chi(a, b) + \chi(b, a) = 1 \quad \forall a, b \in \mathbb{R},$$

in order to prove (80), we have to show that

$$(81) \quad W(u_1, u_2) = (\chi^+ - \chi^-) [A^+(u_2^+) - A^+(u_1^+)] \leq 0,$$

where we have set $\chi^\pm = \chi(u_1^\pm, u_2^\pm)$.

If $\text{sign}(u_2^+ - u_1^+) = \text{sign}(u_2^- - u_1^-)$, then $\chi^+ = \chi^-$, and (81) is satisfied. Hence we shall restrict our attention to the case $1 = \text{sign}(u_2^+ - u_1^+) = -\text{sign}(u_2^- - u_1^-)$, i.e.

$$(82) \quad u_1^+ < u_2^+, \quad u_2^- < u_1^-,$$

so that $\chi^+ = 1$, $\chi^- = 0$, since the case $-1 = \text{sign}(u_2^+ - u_1^+) = -\text{sign}(u_2^- - u_1^-)$ can be handled in a similar way.

Since (82) is in force in order to prove (81) we need to show that

$$(83) \quad A^+(u_2^+) - A^+(u_1^+) \leq 0 \quad \text{or, equivalently,} \quad A^-(u_2^-) - A^-(u_1^-) \leq 0$$

in each one of the following six possibilities:

$$(84) \quad u_1^+ < u_2^+ < u_2^- < u_1^-,$$

$$(85) \quad u_1^+ < u_2^- < u_2^+ < u_1^-,$$

$$(86) \quad u_2^- < u_1^+ < u_2^+ < u_1^-,$$

$$(87) \quad u_2^- < u_1^+ < u_1^- < u_2^+,$$

$$(88) \quad u_2^- < u_1^- < u_1^+ < u_2^+,$$

$$(89) \quad u_1^+ < u_2^- < u_1^- < u_2^+.$$

Here, for simplicity, we have considered only strict inequalities, but the equality cases can be handled as well. Namely, if $u_1^- = u_2^-$ or $u_1^+ = u_2^+$, then (83) trivially holds, while

the other equality cases can be proved using a continuity argument, since the quantities χ^\pm do not change.

The analysis will be the same as the one performed in [24]. For the reader's convenience, we briefly report here how to use the inequalities (26)–(37) in order to prove (83) in each of the six cases listed above. When not explicitly stated, the inequalities (26)–(37) are supposed to be used with $u = u_1$.

• Case (84). We have the following possibilities:

- $u_1^+ < u_2^+ < u_2^- < u_1^- < \hat{u}_1$: choose $c = u_2^+$ in (27).
- $u_1^+ < u_2^+ < \hat{u}_1 < u_1^-$: choose $c = u_2^+$ in (29). (We remark that the relative position of \hat{u}_1 and u_2^- is not relevant, so that this case includes the two subcases $u_1^+ < u_2^+ < \hat{u}_1 < u_2^- < u_1^-$ and $u_1^+ < u_2^+ < u_2^- < \hat{u}_1 < u_1^-$; similar considerations will apply also in some of the following cases without further reference.)
- $u_1^+ < \hat{u}_1 < u_2^- < u_1^-$: choose $c = u_2^-$ in (28).
- $\hat{u}_1 < u_1^+ < u_2^+ < u_2^- < u_1^-$: choose $c = u_2^+$ in (30).

• Case (85). We have the following possibilities:

- $u_1^+ < u_2^- < u_2^+ < u_1^- < \hat{u}_1$: choose $c = u_2^+$ in (27).
- $u_1^+ < u_2^- < u_2^+ < \hat{u}_1 < u_1^-$: choose $c = u_2^+$ in (29).
- $u_1^+ < \hat{u}_1 < u_2^- < u_2^+ < u_1^-$: choose $c = u_2^-$ in (28).
- $\hat{u}_1 < u_1^+ < u_2^- < u_2^+ < u_1^-$: choose $c = u_2^+$ in (30).
- $u_1^+ < u_2^- < \hat{u}_1 < u_2^+ < u_1^-$: this case is slightly more involved because we need to consider also the position of \hat{u}_2 . We have the following subcases:
 - (1) $u_1^+ < u_2^- < \hat{u}_1 < u_2^+ < \hat{u}_2$: apply (33) to u_2 with $c = \hat{u}_1$ and then (28) to u_1 again with $c = \hat{u}_1$, obtaining

$$A^-(u_2^-) \leq A^-(\hat{u}_1) \leq A^-(u_1^-).$$

- (2) $u_1^+ < u_2^- < \hat{u}_1 < \hat{u}_2 < u_2^+ < u_1^-$: apply (35) to u_2 with $c = \hat{u}_1$ and then (28) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^-(u_2^-) \leq A^-(\hat{u}_1) \leq A^-(u_1^-).$$

- (3) $u_1^+ < u_2^- < \hat{u}_2 < \hat{u}_1 < u_2^+ < u_1^-$: apply (34) to u_2 with $c = \hat{u}_1$ and then (29) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^+(u_2^+) \leq A^+(\hat{u}_1) \leq A^+(u_1^+).$$

- (4) $\hat{u}_2 < u_2^- < \hat{u}_1 < u_2^+ < u_1^-$: apply (36) to u_2 with $c = \hat{u}_1$ and then (29) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^+(u_2^+) \leq A^+(\hat{u}_1) \leq A^+(u_1^+).$$

• Case (86). We have the following possibilities:

- $u_2^- < u_1^+ < u_2^+ < u_1^- < \hat{u}_1$: choose $c = u_2^+$ in (27).
- $u_2^- < u_1^+ < u_2^+ < \hat{u}_1 < u_1^-$: choose $c = u_2^+$ in (29).
- $\hat{u}_1 < u_1^+ < u_2^+ < u_1^-$: choose $c = u_2^+$ in (30).
- $u_2^- < u_1^+ < \hat{u}_1 < u_2^+ < u_1^-$: this case is slightly more involved because we need to consider also the position of \hat{u}_2 . We have the following subcases:

- (1) $u_2^- < u_1^+ < \hat{u}_1 < u_2^+ < \hat{u}_2$: apply (33) to u_2 with $c = \hat{u}_1$ and then (28) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^-(u_2^-) \leq A^-(\hat{u}_1) \leq A^-(u_1^-).$$

- (2) $u_2^- < u_1^+ < \hat{u}_1 < \hat{u}_2 < u_2^+ < u_1^-$: apply (35) to u_2 with $c = \hat{u}_1$ and then (28) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^-(u_2^-) \leq A^-(\hat{u}_1) \leq A^-(u_1^-).$$

- (3) $u_2^- < \hat{u}_2 < \hat{u}_1 < u_2^+ < u_1^-$: apply (34) to u_2 with $c = \hat{u}_1$ and then (29) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^+(u_2^+) \leq A^+(\hat{u}_1) \leq A^+(u_1^+).$$

- (4) $\hat{u}_2 < u_1^+ < \hat{u}_1 < u_2^+ < u_1^-$: apply (36) to u_2 with $c = \hat{u}_1$ and then (29) to u_1 again with $c = \hat{u}_1$ obtaining

$$A^+(u_2^+) \leq A^+(\hat{u}_1) \leq A^+(u_1^+).$$

- Case (87). This case is symmetric to (85). It is enough to replace \hat{u}_1 with \hat{u}_2 and to apply (32)–(37) instead of (26)–(31) or conversely.
- Case (88). This case is symmetric to (84). It is enough to replace \hat{u}_1 with \hat{u}_2 and to apply (32)–(37) instead of (26)–(31).
- Case (89). This case is symmetric to (86). It is enough to replace \hat{u}_1 with \hat{u}_2 and to apply (32)–(37) instead of (26)–(31) or conversely. \square

REFERENCES

- [1] Adimurthi, Rajib Dutta, Shyam Sundar Ghoshal, and G. D. Veerappa Gowda, *Existence and nonexistence of TV bounds for scalar conservation laws with discontinuous flux*, Comm. Pure Appl. Math. **64** (2011), no. 1, 84–115. MR 2743877 (2012f:35334)
- [2] Adimurthi, S. Mishra, and G. D. Veerappa Gowda, *Optimal entropy solutions for conservation laws with discontinuous flux-functions*, J. Hyperbolic Differ. Equ. **2** (2005), no. 4, 783–837. MR 2195983 (2007g:35144)
- [3] L. Ambrosio, G. Crasta, V. De Cicco, and G. De Philippis, *A nonautonomous chain rule in $W^{1,p}$ and BV*, Manuscripta Math. **140** (2013), no. 3-4, 461–480. MR 3019135
- [4] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 2000. MR MR1857292 (2003a:49002)
- [5] B. Andreianov, K.H. Karlsen, and N.H. Risebro, *A theory of L^1 -dissipative solvers for scalar conservation laws with discontinuous flux*, Arch. Ration. Mech. Anal. **201** (2011), no. 1, 27–86. MR 2807133
- [6] B. Andreianov and D. Mitrovic, *Entropy conditions for scalar conservation laws with discontinuous flux revised*, to appear in Ann. Inst. H. Poincaré Anal. Non Linéaire
- [7] E. Audusse and B. Perthame, *Uniqueness for scalar conservation laws with discontinuous flux via adapted entropies*, Proc. Roy. Soc. Edinburgh Sect. A **135** (2005), no. 2, 253–265. MR MR2132749 (2006g:35174)
- [8] F. Bachmann and J. Vovelle, *Existence and uniqueness of entropy solution of scalar conservation laws with a flux function involving discontinuous coefficients*, Comm. Partial Differential Equations **31** (2006), no. 1-3, 371–395. MR MR2209759 (2008b:35170)
- [9] G.M. Coclite and N.H. Risebro, *Conservation laws with time dependent discontinuous coefficients*, SIAM J. Math. Anal. **36** (2005), no. 4, 1293–1309 (electronic). MR 2139451 (2006c:35179)

- [10] R.M. Colombo, M. Mercier, and M.D. Rosini, *Stability and total variation estimates on general scalar balance laws*, Commun. Math. Sci. **7** (2009), no. 1, 37–65. MR 2512832 (2011a:35325)
- [11] G. Crasta and V. De Cicco, *A chain rule formula in the space BV and applications to conservation laws*, SIAM J. Math. Anal. **43** (2011), no. 1, 430–456. MR 2765698
- [12] C.M. Dafermos, *Hyperbolic conservation laws in continuum physics*, third ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 325, Springer-Verlag, Berlin, 2010. MR MR2574377
- [13] A.-L. Dalibard, *Kinetic formulation for heterogeneous scalar conservation laws*, Ann. Inst. H. Poincaré Anal. Non Linéaire **23** (2006), no. 4, 475–498. MR 2245753 (2007h:35223)
- [14] A.-L. Dalibard, *Homogenization of non-linear scalar conservation laws*, Arch. Rational Mech. Anal. **192** (2009), no. 1, 117–164. MR 2481063 (2010c:35014)
- [15] V. De Cicco, N. Fusco, and A. Verde, *A chain rule formula in BV and application to lower semi-continuity*, Calc. Var. Partial Differential Equations **28** (2007), no. 4, 427–447. MR MR2293980 (2007j:49016)
- [16] S. Diehl, *A uniqueness condition for nonlinear convection-diffusion equations with discontinuous coefficients*, J. Hyperbolic Differ. Equ. **6** (2009), no. 1, 127–159. MR 2512505 (2010f:35210)
- [17] H. Federer, *Geometric measure theory*, Die Grundlehren der mathematischen Wissenschaften, Band 153, Springer-Verlag New York Inc., New York, 1969. MR MR0257325 (41 #1976)
- [18] M. Garavello, R. Natalini, B. Piccoli, and A. Terracina, *Conservation laws with discontinuous flux*, Netw. Heterog. Media **2** (2007), no. 1, 159–179 (electronic). MR MR2291816 (2008e:35126)
- [19] K. H. Karlsen, N. H. Risebro, and J. D. Towers, *L^1 stability for entropy solutions of nonlinear degenerate parabolic convection-diffusion equations with discontinuous coefficients*, Skr. K. Nor. Vidensk. Selsk. (2003), no. 3, 1–49. MR MR2024741 (2004j:35149)
- [20] S. N. Kružkov, *First order quasilinear equations with several independent variables.*, Mat. Sb. (N.S.) **81 (123)** (1970), 228–255. MR 0267257 (42 #2159)
- [21] P.-L. Lions, B. Perthame, and E. Tadmor, *A kinetic formulation of multidimensional scalar conservation laws and related equations*, J. Amer. Math. Soc. **7** (1994), no. 1, 169–191. MR 1201239 (94d:35100)
- [22] D. Mitrovic, *Existence and stability of a multidimensional scalar conservation law with discontinuous flux*, Netw. Heterog. Media **5** (2010), no. 1, 163–188. MR 2601993 (2011b:35325)
- [23] D. Mitrovic, *New entropy conditions for scalar conservation laws with discontinuous flux*, Discrete Contin. Dyn. Syst. **30** (2011), no. 4, 1191–1210. MR 2812961 (2012e:35152)
- [24] D. Mitrovic, *Proper entropy conditions for scalar conservation laws with discontinuous flux*, Tech. report, 2012.
- [25] E. Yu. Panov, *Existence and strong pre-compactness properties for entropy solutions of a first-order quasilinear equation with discontinuous flux*, Arch. Ration. Mech. Anal. **195** (2010), no. 2, 643–673. MR 2592291 (2011h:35039)
- [26] B. Perthame, *Uniqueness and error estimates in first order quasilinear conservation laws via the kinetic entropy defect measure*, J. Math. Pures Appl. (9) **77** (1998), no. 10, 1055–1064. MR 1661021 (2000e:35141)
- [27] B. Perthame, *Kinetic formulation of conservation laws*, Oxford Lecture Series in Mathematics and its Applications, vol. 21, Oxford University Press, Oxford, 2002. MR 2064166 (2005d:35005)
- [28] A.I. Vol’pert, *Spaces BV and quasilinear equations*, Mat. Sb. (N.S.) **73 (115)** (1967), 255–302. MR MR0216338 (35 #7172)

DIPARTIMENTO DI MATEMATICA “G. CASTELNUOVO”, UNIV. DI ROMA I, P.LE A. MORO 2 – I-00185
ROMA (ITALY)

E-mail address: `crasta@mat.uniroma1.it`

DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L'INGEGNERIA, UNIV. DI ROMA I, VIA A.
SCARPA 10 – I-00185 ROMA (ITALY)

E-mail address: `virginia.decicco@sbai.uniroma1.it`

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH – WINTERTHURERSTRASSE 190 – CH-8057
ZÜRICH

E-mail address: `guido.dephilippis@math.uzh.ch`