

REGULARITY PROPERTIES OF SPHERES IN HOMOGENEOUS GROUPS

ENRICO LE DONNE AND SEBASTIANO NICOLUSSI GOLO

ABSTRACT. We study left-invariant distances on Lie groups for which there exists a one-parameter family of homothetic automorphisms. The main examples are Carnot groups, in particular the Heisenberg group with the standard dilations. We are interested in criteria implying that, locally and away from the diagonal, the distance is Euclidean Lipschitz and, consequently, that the metric spheres are boundaries of Lipschitz domains in the Euclidean sense. In the first part of the paper, we consider geodesic distances. In this case, we actually prove the regularity of the distance in the more general context of sub-Finsler manifolds with no abnormal geodesics. Secondly, for general groups we identify an algebraic criterium in terms of the dilating automorphisms, which for example makes us conclude the regularity of homogeneous distances on the Heisenberg group. In such a group, we analyze in more details the geometry of metric spheres. We also provide examples of homogeneous groups where spheres presents cusps.

CONTENTS

Contents	1
1. Introduction	2
2. Preliminaries	4
2.1. Sub-Finsler structures	4
2.2. Graded groups	6
3. Regularity of sub-Finsler distances	9
3.1. About the minimal stretching	12
3.2. The End-point map is weakly* continuous	12
3.3. The differential of the End-point map is an End-point map	14
3.4. The sub-Finsler distance is Lipschitz in absence of singular geodesics	15
4. Regularity of spheres in graded groups	16

Date: September 13, 2015.

2010 Mathematics Subject Classification. 28A75, 22E25, 53C60, 53C17, 26A16.

Key words and phrases. sub-Finsler geometry, sub-Riemannian geometry, Carnot groups, Lipschitz regularity, abnormal geodesic, Heisenberg group.

S.N.G. has been supported by the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement n. 607643.

4.1. The sphere as a graph	16
4.2. An intrinsic approach	18
5. Examples	20
5.1. Three gradings on \mathbb{R}^2	20
5.2. The Heisenberg groups	23
5.3. A sub-Finsler sphere with a cusp	23
5.4. A sub-Riemannian sphere with a cusp	24
6. A closer look at the Heisenberg group	25
6.1. Proof of Proposition 6.2	26
6.2. Proof of Proposition 6.3	28
References	29

1. INTRODUCTION

The study of the asymptotic geometry of groups lead us to investigate spheres in homogeneous groups, examples of which are asymptotic cones of finitely generated nilpotent groups. A homogeneous group is a Lie group G endowed with a family of Lie group automorphisms $\{\delta_\lambda\}_{\lambda>0}$ and a left-invariant distance d for which each δ_λ multiplies the distance by λ , see Section 2.2. An algebraic characterization of these groups is known by [Sie86]. In fact, the Lie algebra \mathfrak{g} of G admits a *grading*, i.e., a decomposition $\mathfrak{g} = \bigoplus_{i>1} V_i$ such that $[V_i, V_j] \subset V_{i+j}$. For simplicity, we assume that the dilations are the ones induced by the grading. Namely, the dilation of factor λ *relative* to the grading is the one such that $(\delta_\lambda)_*(v) = \lambda^i v$ for all $v \in V_i$. Not all homogeneous metrics admit this type of dilations. We denote by 0 the neutral element of G and by \mathbb{S}_d the unit sphere at 0 for a distance d on G , i.e., $\mathbb{S}_d := \{p \in G : d(0, p) = 1\}$.

In this paper we want to exclude cusps in spheres since their presence in the asymptotic cone of a finitely generated nilpotent group may give a slower rate of convergence in the blow down, see [BLD13]. We find criteria implying that the metric spheres are boundaries of Lipschitz domains and in fact that the distance function from a point is a locally Lipschitz function with respect to a Riemannian metric.

First, we address the case where the distance d is a length distance. Thanks to a characterization of Carnot groups, see [LD15b], the group G is in this case a stratified group and d is a sub-Finsler distance. Being a *stratified group* means that the grading of \mathfrak{g} is such that the first layer V_1 generates \mathfrak{g} . Being a *sub-Finsler distance* means that there are a left-invariant subbundle $\Delta \subset TG$ and a left-invariant norm $\|\cdot\|$ on Δ such that the length induced by d of an absolutely continuous curve $\gamma : [0, 1] \rightarrow G$ is equal to $\int_0^1 \|\gamma'(t)\| dt$, where $\|\gamma'(t)\| = +\infty$ if

$\gamma'(t) \notin \Delta$. The left-invariant subbundle Δ is actually the one generated by V_1 .

In the sub-Finsler case, an obstruction to Lipschitz regularity of the sphere comes from the presence of length-minimizing curves (also called geodesics) that are not regular, in the sense that the first variation parallel to the subbundle Δ does not have maximal rank, see Definition 2.6.

Theorem 1.1. *Let G be a stratified group endowed with a sub-Finsler metric d . Let $d_0 : G \rightarrow [0, +\infty)$, $p \mapsto d(0, p)$. Let $p \in G$ be such that all geodesics from 0 to p are regular. Then, for any Riemannian metric ρ on G the function d_0 is Lipschitz with respect to ρ in some neighborhood of p .*

We will actually state and prove Theorem 1.1 in the more general setting of *sub-Finsler manifolds of constant-type norm*, see Section 2.1.

In case of homogeneity, the regularity of the distance implies also the regularity of the spheres. Hence, using Theorem 1.1 we easily get the second result for sub-Finsler homogeneous groups.

Theorem 1.2. *Let G be a stratified group endowed with a sub-Finsler metric d . Let $p \in \mathbb{S}_d$ be such that all geodesics from 0 to p are regular. Then, in smooth coordinates, the set \mathbb{S}_d is a Lipschitz graph in some neighborhood of p . In particular, if all non-constant geodesics are regular, then metric balls are Lipschitz domains.*

Notice that a ball may be a Lipschitz domain even if the distance from a point is not Lipschitz (we give an example in Remark 5.5). In Section 5 we also present examples of sub-Riemannian and sub-Finsler distances whose balls have a cusp.

At a second stage, we drop the hypothesis of d being a length distance and we present a result similar to the previous Theorem 1.2 in the context of homogeneous groups. Hereafter we denote by L_p and R_p the left and the right translations on G , respectively, and by $\bar{\delta}(p)$ the vector $\frac{d}{dt}\delta_t(p)|_{t=1} \in T_pG$, where $\{\delta_t\}_{t>0}$ are the dilations relative to a grading.

Theorem 1.3. *Let (G, d) be a homogeneous group with dilations relative to a grading, see Definition 2.12. Assume $p \in \mathbb{S}_d$ is such that*

$$(1) \quad dL_p(V_1) + dR_p(V_1) + \text{span}\{\bar{\delta}(p)\} = T_pG.$$

Then, in some neighborhood of p we have that the sphere \mathbb{S}_d is a Lipschitz graph and the distance d_0 from the identity is Lipschitz with respect to any Riemannian metric ρ .

The similarity between Theorem 1.2 and Theorem 1.3 consists in the fact that, if d is a sub-Finsler distance, then condition (1) implies that all geodesics from 0 to p are regular, see Remark 4.5.

The equality (1) or the absence of non-regular geodesics are actually quite strong conditions. However, in general we can give an upper bound for the Hausdorff dimension of spheres. In fact, if d is a homogeneous distance on a graded group of maximal degree s , then

$$(2) \quad \dim_H^\rho(G) - 1 \leq \dim_H^\rho(\mathbb{S}_d) \leq \dim_H^\rho(G) - \frac{1}{s},$$

where \dim_H^ρ is the Hausdorff dimension with respect to some (therefore any) Riemannian metric ρ . We show with Proposition 5.1 that this estimate is sharp.

In the last part of the paper, we analyze in more details an important specific example: the Heisenberg group. In this graded group we consider all possible homogeneous distances and prove that in exponential coordinates

- (i) the unit ball is a star-shaped Lipschitz domain (Proposition 6.1);
- (ii) the unit sphere is a locally Lipschitz graph with respect to the direction of the center of the group (Proposition 6.2).

We also give a method to construct homogeneous distances in the Heisenberg group with arbitrary Lipschitz regularity of the sphere. Namely, the graph of each Lipschitz function defined on the unit disk, up to adding to it a constant, is the sphere of some homogeneous distance, see Proposition 6.3. The investigation of this class of examples is meaningful in connection to Besicovitch's covering property as studied in [LDR14] and [LDR15].

The paper is organized as follows. In section 2 we will present all preliminary notions needed in the paper. We introduce sub-Finsler manifolds of constant-type norm, graded and homogeneous groups and Carnot groups. Section 3 is devoted to the proof of Theorem 1.1; first in the setting of sub-Finsler manifolds, see Theorem 3.1 proved in Section 3.4, then with a more specific result for Carnot groups, see Proposition 3.3. In Section 4 we see metric spheres as graphs over smooth spheres. Hence, we show Theorem 1.2, the inequalities (2), and Theorem 1.3. In Section 5 we present six examples: three different gradings of \mathbb{R}^2 , the Heisenberg group, a sub-Finsler sphere with a cusp and a sub-Riemannian sphere with a cusp. In Section 6 we prove stronger properties for spheres of homogeneous distances on the Heisenberg group.

2. PRELIMINARIES

2.1. Sub-Finsler structures. Let M be a manifold of dimension n . We will write TM for the tangent bundle and $\text{Vec}(M)$ for the space of smooth vector fields on M .

Definition 2.1 (Sub-Finsler structure). A *sub-Finsler structure* (of constant-type norm) of rank r on a manifold M is a triple $(\mathbb{E}, \|\cdot\|, \mathbf{f})$, where $(\mathbb{E}, \|\cdot\|)$ is a normed vector space of dimension r and $\mathbf{f} : M \times \mathbb{E} \rightarrow TM$ is a smooth bundle morphism with $\mathbf{f}(\{p\} \times \mathbb{E}) \subset T_pM$, for all $p \in M$.

Definition 2.2 (Horizontal curve). A curve $\gamma : [0, 1] \rightarrow M$ is a *horizontal curve* if it is absolutely continuous and there is $u : [0, 1] \rightarrow \mathbb{E}$ measurable, which is called a *control* of γ , such that

$$\gamma'(t) = \mathbf{f}(\gamma(t), u(t)) \quad \text{for a.e. } t \in [0, 1].$$

In this case γ is called *integral curve* of u and we write γ_u .

Definition 2.3 (Space of controls). The space of L^∞ -controls is defined as¹

$$L^\infty([0, 1]; \mathbb{E}) := \left\{ u : [0, 1] \rightarrow \mathbb{E} \text{ measurable, } \operatorname{ess\,sup}_{t \in [0, 1]} \|u(t)\| < \infty \right\}.$$

This is a Banach space with norm $\|u\|_{L^\infty} := \operatorname{ess\,sup}_{t \in [0, 1]} \|u(t)\|$.

Thanks to known results for ordinary differential equations, see [Rif14], given a control $u \in L^\infty([0, 1]; \mathbb{E})$, there is an open neighborhood $M^* \subset M \times [0, 1]$ of $M \times \{0\}$ and a map $\Phi^u : M^* \rightarrow M$, such that for all $p \in M$ the set $M^* \cap (\{p\} \times [0, 1])$ is connected and contains $(p, 0)$, and the curve $\gamma(t) := \Phi^u(p, t)$ is the unique solution to the Cauchy problem

$$\begin{cases} \gamma(0) = p \\ \gamma'(t) = \mathbf{f}(\gamma(t), u(t)) \quad \text{for a.e. } t \text{ with } (p, t) \in M^*. \end{cases}$$

The map Φ^u is called *flow* of u .

Remark 2.4. We will always assume that every $u \in L^\infty([0, 1]; \mathbb{E})$ has a flow defined on the entire interval $[0, 1]$, i.e., $M^* = M \times [0, 1]$. This happens in many cases, for example for left-invariant sub-Finsler structures on Lie groups, in particular in Carnot groups.

Definition 2.5 (End-point map). Fix $o \in M$. Define the *End-point map* with *basis point* o , $\operatorname{End}_o : L^\infty([0, 1]; \mathbb{E}) \rightarrow M$, as

$$\operatorname{End}_o(u) = \Phi^u(o, 1).$$

By standard result of ODE the map End_o is of class \mathcal{C}^1 , see [Rif14].

Definition 2.6 (Regular curves). Given $o \in M$, a control $u \in L^\infty([0, 1]; \mathbb{E})$ is said to be *regular* if it is a regular point of End_o , i.e., if $d\operatorname{End}_o(u) : L^\infty([0, 1]; \mathbb{E}) \rightarrow T_{\operatorname{End}_o(u)}M$ is surjective. A *singular control* is a control that is not regular.

¹ Among the three norms L^1 , L^2 and L^∞ for controls, we chose the latter because the unit ball in $L^1([0, 1]; \mathbb{E})$ is not weakly compact and the L^2 -space is not a Hilbert space in our contest.

Definition 2.7 (Sub-Finsler distance). The *sub-Finsler distance*, also called *Carnot-Carathéodory distance*, between two points $p, q \in M$ is

$$d(p, q) := \inf \left\{ \int_0^1 \|u(t)\| dt : u \in L^\infty([0, 1]; \mathbb{E}) \text{ with } \mathbf{End}_p(u) = q \right\}.$$

Clearly (M, d) is a metric space, even though it might happen $d(p, q) = \infty$. Let $\ell_d(\gamma)$ be the length of a curve γ with respect to d , see [AT04]. It can be proven that a curve $\gamma : [0, 1] \rightarrow (M, d)$ is Lipschitz if and only if it is horizontal and it admits a control in $L^\infty([0, 1]; \mathbb{E})$. Moreover, if γ is Lipschitz, then

$$\ell_d(\gamma) = \inf \left\{ \int_0^1 \|u(t)\| dt : u \in L^\infty([0, 1]; \mathbb{E}) \text{ control of } \gamma \right\}.$$

We will use the term *geodesic* as a synonym of *length-minimizer*.

The distance can be expressed by using the L^∞ -norm as an energy, i.e., for every $p, q \in M$

$$d(p, q) = \inf \{ \|u\|_{L^\infty} : u \in L^\infty([0, 1]; \mathbb{E}) \text{ with } \mathbf{End}_p(u) = q \}.$$

Moreover, if u realizes the infimum above, then its integral curve γ_u starting from p is a length-minimizing curve parametrized by constant velocity, i.e.,

$$d(p, q) = \|u\|_{L^\infty} = \ell_d(\gamma_u) = \|u(t)\|, \quad \text{for a.e. } t \in [0, 1].$$

Definition 2.8 (Bracket-generating condition). Let \mathcal{A} be the Lie algebra generated by the set

$$\{p \mapsto \mathbf{f}(p, X(p)) : X : M \rightarrow \mathbb{E} \text{ smooth}\} \subset \text{Vec}(M).$$

We say that the sub-Finsler structure $(\mathbb{E}, \|\cdot\|, \mathbf{f})$ on M satisfies the *bracket-generating condition* if for all $p \in M$

$$\{V(p) : V \in \mathcal{A}\} = T_p M.$$

As a consequence of the Orbit Theorem [Jur97], we have the following basic well-known fact.

Lemma 2.9. *If $(\mathbb{E}, \|\cdot\|, \mathbf{f})$ satisfies the bracket-generating condition, then the distance d induces the original topology of M and (M, d) is a locally compact and locally geodesic length space.*

By the Hopf-Rinow Theorem, see [BBI01], the assumption in Remark 2.4 implies that (M, d) is a complete, boundedly compact metric space.

2.2. Graded groups. All Lie algebras considered here are over \mathbb{R} and finite-dimensional.

Definition 2.10 (Graded group). A Lie algebra \mathfrak{g} is *graded* if it is equipped with *grading*, i.e., with a vector-space decomposition $\mathfrak{g} = \bigoplus_{i>0} V_i$, where $i > 0$ means $i \in (0, \infty)$, such that for all $i, j > 0$ it

holds $[V_i, V_j] \subset V_{i+j}$. A *graded Lie group* is a simply connected Lie group G whose Lie algebra is graded. The *maximal degree* of a graded group G is the maximum i such that $V_i \neq \{0\}$.

Graded groups are nilpotent and the exponential map $\exp : \mathfrak{g} \rightarrow G$ is a global diffeomorphism. We will denote by 0 the neutral element of G and identify $\mathfrak{g} = T_0G$.

Definition 2.11 (Dilations). In a graded group for which the Lie algebra has the grading $\mathfrak{g} = \bigoplus_{i>0} V_i$, the *dilations* relative to the grading are the group homomorphisms $\delta_\lambda : G \rightarrow G$, for $\lambda \in (0, \infty)$, such that $(\delta_\lambda)_*(v) = \lambda^i v$ for all $v \in V_i$.

In the definition above, ϕ_* denotes the Lie algebra homomorphism associated to a Lie group homomorphism ϕ , in particular, $\phi \circ \exp = \exp \circ \phi_*$. Since a graded group is simply connected, δ_λ is well defined. Notice that, for any $\lambda, \mu > 0$, $\delta_\lambda \circ \delta_\mu = \delta_{\lambda\mu}$.

Definition 2.12 (Homogeneous distances). Let G be a graded group with a dilations $\{\delta_\lambda\}_{\lambda>0}$, relative to the grading. We say that a distance d on G is *homogeneous* if it is left-invariant, i.e., for every $g, x, y \in G$ we have $d(gx, gy) = d(x, y)$, and one-homogeneous with respect to the dilations, i.e., for all $\lambda > 0$ and all $x, y \in G$ we have $d(\delta_\lambda x, \delta_\lambda y) = \lambda d(x, y)$. If d is one such a distance, then (G, d) is called *homogeneous group* (with dilations relative to the grading).

Remark 2.13. A graded group admits a homogeneous distance if and only if for $i \in (0, 1)$ we have $V_i = \{0\}$, see [HS90].

Given a homogeneous distance d , the function $p \mapsto d_0(p) := d(0, p)$ is a homogeneous norm. Here with the term *homogeneous norm* we mean a function $N : G \rightarrow [0, +\infty)$ such that for all $p, q \in G$ and all $\lambda > 0$ it holds

- (1) $N(p) = 0 \Leftrightarrow p = 0$;
- (2) $N(pq) \leq N(p) + N(q)$;
- (3) $N(p^{-1}) = N(p)$;
- (4) $N(\delta_\lambda p) = \lambda N(p)$.

In fact, homogeneous distances are in bijection with homogeneous norms on G through the formula $d(p, q) = N(p^{-1}q)$.

Homogeneous distances induce the original topology of G , see [LDR15]. Moreover, given two homogeneous distances d_1, d_2 on G , there is a constant $C > 0$ such that for all $p, q \in G$

$$(3) \quad \frac{1}{C} d_1(p, q) \leq d_2(p, q) \leq C d_1(p, q).$$

Lemma 2.14. Let G be a graded group and $0 < k_1 \leq k_2$ such that $V_i = \{0\}$ for all $i < k_1$ and all $i > k_2$. Let d be a homogeneous

distance and ρ a left-invariant Riemannian metric on G . Then there are $C, \epsilon > 0$ such that for all $p, q \in G$ with $\rho(p, q) < \epsilon$ it holds

$$(4) \quad \frac{1}{C} \rho(p, q)^{\frac{1}{k_1}} \leq d(p, q) \leq C \rho(p, q)^{\frac{1}{k_2}}.$$

In particular, the homogeneous norm d_0 is locally $\frac{1}{k_2}$ -Hölder.

Proof. We identify $G = \mathfrak{g}$ via the exponential map. So, if $p \in G$, we denote by p_i the i -th component in the decomposition $p = \sum_i p_i$ with $p_i \in V_i$. Fix a norm $|\cdot|$ on \mathfrak{g} . For any pair $(p, q) \in G \times G$ define

$$\eta(p, q) := \eta(0, p^{-1}q), \text{ where } \eta(0, p) := \max_i (|p_i|)^{\frac{1}{i}}.$$

The function η is a so-called quasi-distance, see [LDR15]. In particular, η is continuous, left-invariant and one-homogeneous with respect to the dilations δ_λ . Therefore, if d is a homogeneous distance, then there is $C > 0$ such that

$$\frac{1}{C} \eta(p, q) \leq d(p, q) \leq C \eta(p, q).$$

So, we can prove (4) only for η .

Let $C, \epsilon > 0$ be with $C\epsilon < 1$ and such that, if $\rho(0, p) < \epsilon$, then

$$(5) \quad \frac{1}{C} \rho(0, p) \leq \max_i |p_i| \leq C \rho(0, p).$$

Therefore, if $\rho(p, q) < \epsilon$, then $|(p^{-1}q)_i| \leq C \rho(p, q) < 1$ for all i and

$$(6) \quad \max_i |(p^{-1}q)_i|^{\frac{1}{k_1}} \leq \max_i (|(p^{-1}q)_i|)^{\frac{1}{i}} = \eta(p, q) \leq \max_i |(p^{-1}q)_i|^{\frac{1}{k_2}},$$

thanks to the monotonicity of the function $x \mapsto a^x$ for $0 < a < 1$. The thesis follows immediately from (5) and (6) combined. \square

Next lemma gives a characterization of sets that are the unit ball of a homogeneous distance. In this paper, we denote by $\text{int}(B)$ the interior part of a subset B . The proof of the following fact is straightforward and hence omitted.

Lemma 2.15. *Let G be a graded group with dilations δ_λ , $\lambda > 0$. A set $B \subset G$ is the unit ball with center 0 of a homogeneous distance on G if and only if B is compact, $0 \in \text{int}(B)$, $B = B^{-1}$ and*

$$(7) \quad \forall p, q \in B, \forall t \in [0, 1] \quad \delta_t(p) \delta_{1-t}(q) \in B.$$

Definition 2.16 (Stratified group). A *stratified group* is a graded group G such that its Lie algebra \mathfrak{g} is generated by the layer V_1 of the grading of \mathfrak{g} .

Notice that in a stratified group G the maximal degree s of the grading equals the nilpotency step of G and it holds $\mathfrak{g} = \bigoplus_{i=1}^s V_i$ with $[V_1, V_i] = V_{i+1}$ for all $i \in \{1, \dots, s\}$, with $V_{s+1} = \{0\}$. We also remark that all stratifications of a group G are isomorphic to each other, i.e.,

if $\mathfrak{g} = \bigoplus_{i=1}^{s'} W_i$ is a second stratification, then there is a Lie group automorphism $\phi : G \rightarrow G$ such that $\phi_*(W_i) = V_i$ for all i , see [LD15a].

In a stratified group, the map $\mathbf{f} : G \times V_1 \rightarrow TG$, $\mathbf{f}(g, v) := dL_g(v)$, is a bundle morphism with $\mathbf{f}(g, v) \in T_gG$. So, if $\|\cdot\|$ is any norm on V_1 , the triple $(V_1, \|\cdot\|, \mathbf{f})$ is a sub-Finsler structure on G . The stratified group G endowed with the corresponding sub-Finsler distance d is called *Carnot group*. Such a d is an example of a homogeneous distance on G .

Remark 2.17. As already stated, singular curves play a central role in our analysis, because they disrupt the Lipschitz regularity of the distance function. We remark that every Carnot group of nilpotency step $s \geq 3$ has singular geodesics. More precisely, there is $X \in V_1$ such that the curve $t \mapsto \exp(tX)$ is a singular geodesic. In particular, if all non-constant length-minimizing curves are regular, then the step of the group is necessarily at most 2.

3. REGULARITY OF SUB-FINSLER DISTANCES

We will prove in this section that sub-Finsler distances are Lipschitz whenever all length-minimizing curves are regular, see Theorem 3.1. Theorem 1.1 expresses this result for Carnot groups.

It is important to remind what is known in the sub-Riemannian case. A *sub-Riemannian distance* is a sub-Finsler distance whose norm on the bundle \mathbb{E} is induced by a scalar product. Rifford proved in [Rif06] that, if there are no singular length-minimizers, for all $o \in M$, not only d_o is locally Lipschitz, but also the spheres centered at o are Lipschitz hypersurfaces for almost all radii. The key points of his proof are the tools of Clarke's non-smooth calculus (see [Cla+98]) and a version of Sard's Lemma for the distance function (see [Rif04]). An exhaustive exposition of this topic can be found in [ABB15].

In Rifford's version of Sard's Lemma, one uses the fact that the L^2 norm in the Hilbert space $L^2([0, 1]; \mathbb{E})$ is smooth away from the origin. If \mathbb{E} is equipped with a generic norm, instead, the L^p norm on $L^p([0, 1]; \mathbb{E})$ with $1 \leq p \leq \infty$ may be non-smooth, hence the proof does not work in the sub-Finsler case.

The non-smoothness of the norm can be seen in another dissimilarity between sub-Riemannian and sub-Finsler distances. Sub-Riemannian distances are proven to be locally semi-concave when there are no singular length-minimizing curves. We remind that a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *semi-concave* if for each $p \in \mathbb{R}^n$ there exists a \mathcal{C}^2 function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $f \leq g$ and $f(p) = g(p)$, see [Rif14]. Semi-concavity is a stronger property than being Lipschitz. However, semi-concavity fails to hold in the sub-Finsler case. For example, the ℓ^1 -distance $d(0, (x, y)) := |x| + |y|$ on \mathbb{R}^2 is a sub-Finsler distance that is not semi-concave along the coordinate axis, although all curves are regular.

We restrict our analysis to the Lipschitz regularity of the distance function, from which we deduce regularity properties of the spheres by means of the homogeneity of Carnot groups. With this aim in view, the core of the proof of Theorem 3.1 is the bound on the point-wise Lipschitz constant (see (12) at page 16), which already appeared in the sub-Riemannian context, see [Agr09]. Our approach differs from the sub-Riemannian case for the fact that the set of optimal curves joining two points on a sub-Finsler manifold may not be compact in the $W^{1,\infty}$ topology. As an example, consider the set of all length-minimizers from $(0,0)$ to $(0,1)$ for the ℓ^∞ -distance $d(0, (x,y)) := \max\{|x|, |y|\}$ on \mathbb{R}^2 . However, we are still able to obtain a bound on the pointwise Lipschitz constant, i.e., to prove (12), by use of the weak* topology on controls.

Theorem 3.1. *Let $(\mathbb{E}, \|\cdot\|, \mathbf{f})$ be a sub-Finsler structure on M with sub-Finsler distance d . Fix o and p in M . If all the length-minimizing curves from o to p are regular, then for every Riemannian metric ρ on M there are a neighborhood U of p and $L > 0$ such that*

$$(8) \quad \forall q_1, q_2 \in U \quad d_o(q_1) - d_o(q_2) \leq L\rho(q_1, q_2).$$

The proof is presented in Section 3.4.

Remark 3.2. Theorem 3.1 can be made more quantitative. Define

$$\tau_0 := \inf \{ \tau(\text{dEnd}_o(u)) : \text{End}_o(u) = p \text{ and } \|u\|_{L^\infty} = d(o, p) \},$$

where τ the minimal stretching, which we will recall in Definition 3.4. Then, for every $L > \frac{1}{\tau_0}$, there exists a neighborhood U of p such that (8) holds. The hypothesis of regularity of all length-minimizing curves from o to p is equivalent to $\tau_0 > 0$.

In the case of Carnot groups (of step 2, see Remark 2.17), we can obtain a more global result

Proposition 3.3. *Let (G, d) be a Carnot group without non-constant singular geodesics. Then for every left-invariant Riemannian metric ρ and every neighborhood U of 0 the function $d_0 : x \mapsto d(0, x)$ is Lipschitz on $G \setminus U$. Moreover, the function $d_0^2 : x \mapsto d(0, x)^2$ is Lipschitz in a neighborhood of 0.*

Proof. Thanks to Theorem 3.1, one easily shows that there are $L > 0$ and an open neighborhood Ω of the unit sphere $\{p : d(0, p) = 1\}$ such that d_0 is L -Lipschitz on Ω .

Next, we claim that d_0 is locally L -Lipschitz on $G \setminus B_d(0, 1)$. Indeed, let $r > 0$ be such that $B_\rho(x, r) \subset \Omega$ for all $x \in S_d(0, 1)$. If $q_1, q_2 \in G \setminus B_d(0, 1)$ are such that $\rho(q_1, q_2) < r$, then there is $o \in G$ such that $d(0, q_1) = d(0, o) + d(o, q_1)$ and $d(o, q_1) = 1$, therefore

$$d_0(q_2) - d_0(q_1) \leq d(o, q_2) - d(o, q_1) \leq L\rho(o^{-1}q_2, o^{-1}q_1) = L\rho(q_2, q_1).$$

In the second step of the proof, we prove that d_0 is L -Lipschitz on $G \setminus B_d(0, 1)$. Let $p, q \in G \setminus B_d(0, 1)$ and $\gamma : [0, 1] \rightarrow G$ a ρ -length

minimizing curve from p to q . If $\text{Im}\gamma \subset G \setminus B_d(0, 1)$, then there are $0 = t_0 \leq t_1 \leq \dots \leq t_{k+1} = 1$ such that $d_0(\gamma(t_i)) - d_0(\gamma(t_{i+1})) \leq L\rho(\gamma(t_i), \gamma(t_{i+1}))$ for all i . Hence

$$\begin{aligned} d_0(p) - d_0(q) &= \sum_{i=0}^k d_0(\gamma(t_i)) - d_0(\gamma(t_{i+1})) \\ &\leq L \sum_{i=0}^k \rho(\gamma(t_i), \gamma(t_{i+1})) = L\rho(p, q). \end{aligned}$$

If instead $\text{Im}\gamma \cap B_d(0, 1) \neq \emptyset$, then there are $0 < s < t < 1$ such that $d_0(\gamma(s)) = d_0(\gamma(t)) = 1$ and $\gamma([0, s]) \subset G \setminus B_d(0, 1)$ and $\gamma([t, 1]) \subset G \setminus B_d(0, 1)$. Then

$$\begin{aligned} d_0(p) - d_0(q) &= d_0(p) - d_0(\gamma(s)) + d_0(\gamma(t)) - d_0(q) \\ &\leq L(\rho(p, \gamma(s)) + \rho(\gamma(t), q)) \leq L\rho(p, q). \end{aligned}$$

Finally, let $p, q \in G \setminus B_d(0, r)$ for $0 < r < 1$. Then $\delta_{r-1}p, \delta_{r-1}q \in G \setminus B_d(0, 1)$ and we have

$$(9) \quad d_0(p) - d_0(q) = r(d_0(\delta_{r-1}p) - d_0(\delta_{r-1}q)) \leq Lr\rho(\delta_{r-1}p, \delta_{r-1}q) \leq \frac{CL}{r}\rho(p, q),$$

where we used in the last step the fact that there exists $C > 0$ such that

$$\forall p, q \in G, \forall r \in (0, 1) \quad \rho(\delta_{r-1}p, \delta_{r-1}q) \leq Cr^{-2}\rho(p, q).$$

Now, we need to prove that d_0^2 is Lipschitz on $B_d(0, 1)$. We first claim that d_0^2 is locally $4L$ -Lipschitz on $B_d(0, 1) \setminus \{0\}$. Indeed, if $p, q \in B_d(0, 1) \setminus \{0\}$ are such that

$$\frac{1}{2} \leq \frac{d_0(p)}{d_0(q)} \leq 2,$$

then

$$0 < d_0(p) + d_0(q) \leq 4 \min\{d_0(p), d_0(q)\}.$$

Therefore, using (9),

$$\begin{aligned} d_0(p)^2 - d_0(q)^2 &= (d_0(p) + d_0(q))(d_0(p) - d_0(q)) \\ &\leq (d_0(p) + d_0(q)) \frac{CL}{\min\{d_0(p), d_0(q)\}} \rho(p, q) \\ &\leq (d_0(p) + d_0(q)) \frac{4CL}{d_0(p) + d_0(q)} \rho(p, q) = 4CL\rho(p, q). \end{aligned}$$

Finally, using again the fact that ρ is a geodesic distance, we get that d_0^2 is $4CL$ Lipschitz on $B_d(0, 1) \setminus \{0\}$ and therefore on $B_d(0, 1)$. \square

3.1. About the minimal stretching.

Definition 3.4 (Minimal Stretching). Let $(X, \|\cdot\|)$ and $(Y, \|\cdot\|)$ be normed vector spaces. We define for a continuous linear map $L : X \rightarrow Y$ the *minimal stretching*

$$\tau(L) := \inf\{\|y\| : y \in Y \setminus L(B_X(0, 1))\}$$

where $B_X(p, r) = \{q \in X : \|q - p\| < r\}$.

It is easy to prove that $\tau : L(X; Y) \rightarrow [0, +\infty)$ is continuous, where $L(X; Y)$ is the space of continuous linear mappings $X \rightarrow Y$ endowed with the operator norm.

The next proposition applies this notion to smooth functions. We omit the proof because it follows the standard arguments for the Implicit Function Theorem.

Proposition 3.5. *Let $(X, \|\cdot\|)$ and $(Y, \|\cdot\|)$ be two Banach spaces and $F : \Omega \rightarrow Y$ a \mathcal{C}^2 map, where $\Omega \subset X$ is open. Fix $\hat{x} \in \Omega$ and let $\tau_0 := \tau(dF(\hat{x})) > 0$. Then for every $C > 1$ there is $\hat{\epsilon} > 0$, such that for all $0 < \epsilon < \hat{\epsilon}$ it holds*

$$B(F(\hat{x}), \epsilon) \subset F\left(B(\hat{x}, \frac{C}{\tau_0}\epsilon)\right).$$

3.2. The End-point map is weakly* continuous. As before, let $(\mathbb{E}, \mathbf{f}, \|\cdot\|)$ be a sub-Finsler structure on a manifold M . We want to prove the following proposition.

Proposition 3.6. *Fix $o \in M$ and let $o_k \in M$ be a sequence converging to o . Let $u_k \in L^\infty([0, 1]; \mathbb{E})$ be a sequence of controls weakly* converging to $u \in L^\infty([0, 1]; \mathbb{E})$. Let γ_k (resp. γ) be the curve with control u_k (resp. u) and $\gamma_k(0) = o_k$ (resp. $\gamma(0) = o$). Then γ_k uniformly converge to γ .*

In particular, it follows that the End-point map $\mathbf{End}_o : L^\infty([0, 1]; \mathbb{E}) \rightarrow M$ is weakly* continuous.

Proof. Since the sequence u_k is bounded in $L^\infty([0, 1]; \mathbb{E})$ by the Banach–Steinhaus Theorem and the sequence o_k is bounded in (M, d) , then there is a compact set $K \subset M$ such that $\gamma_k \subset K$ for all k . Let $R > 0$ be such that $\|u_k\|_{L^\infty} \leq R$ for all $k \in \mathbb{N}$.

Thanks to the Whitney Embedding Theorem, we can assume that M is a submanifold of \mathbb{R}^N for some $N \in \mathbb{N}$. Fix a basis e_1, \dots, e_r of \mathbb{E} and define the vector fields $X_i : M \rightarrow \mathbb{R}^N$ as

$$X_i(p) := \mathbf{f}(p, e_i).$$

Since they are smooth, they are L -Lipschitz on K for some $L > 0$. We extend the vector fields $X_i : M \rightarrow \mathbb{R}^N$ to smooth functions $X_i : \mathbb{R}^N \rightarrow \mathbb{R}^N$.

Define $\eta_k : [0, 1] \rightarrow \mathbb{R}^N$ as

$$\eta_k(t) := o_k + \int_0^t u_k^i(s) X_i(\gamma(s)) ds.$$

Since $t \mapsto X_i(\gamma(t)) \in \mathbb{R}^N$ are continuous, then $u_k^i X_i(\gamma) \xrightarrow{*} u^i X_i(\gamma)$, for all $i \in \{1, \dots, r\}$. In particular, $\eta_k(t) \rightarrow \gamma(t)$ for each $t \in [0, 1]$. Moreover, since the η_k 's have uniformly bounded derivative, they are a pre-compact family of curves with respect to the topology of uniform convergence. This fact and the pointwise convergence imply that $\eta_k \rightarrow \gamma$ uniformly on $[0, 1]$.

Set $\epsilon_k := \sup_{t \in [0, 1]} |\eta_k(t) - \gamma(t)| + 2|o_k - o|$, so that $\epsilon_k \rightarrow 0$, where $|\cdot|$ is the usual norm in \mathbb{R}^N .

By Ascoli-Arzelà Theorem, the family of curves $\{\gamma_k\}_k$ is also pre-compact with respect to the uniform convergence. Hence, if we prove that the only accumulation curve of $\{\gamma_k\}_k$ is γ , then we obtain that γ_k uniformly converges to γ . So, we can assume $\gamma_k \rightarrow \xi$ uniformly for some $\xi : [0, 1] \rightarrow M$. Then we have (sums on i are hidden)

$$\begin{aligned} |\gamma_k(t) - \gamma(t)| &\leq |o_k - o| + \left| \int_0^t u_k^i(s) X_i(\gamma_k(s)) - u^i(s) X_i(\gamma(s)) ds \right| \\ &\leq |o_k - o| + \int_0^t |u_k^i(s) X_i(\gamma_k(s)) - u_k^i(s) X_i(\gamma(s))| ds + \\ &\quad + \left| \int_0^t u_k^i(s) X_i(\gamma(s)) - u^i(s) X_i(\gamma(s)) ds \right| \\ &\leq 2|o_k - o| + rRL \int_0^t |\gamma_k(s) - \gamma(s)| ds + |\eta_k(t) - \gamma(t)| \\ &\leq rRL \int_0^t |\gamma_k(s) - \gamma(s)| ds + \epsilon_k. \end{aligned}$$

Passing to the limit $k \rightarrow \infty$, we get for all $t \in [0, 1]$

$$(10) \quad |\xi(t) - \gamma(t)| \leq rRL \int_0^t |\xi(s) - \gamma(s)| ds.$$

Starting with the fact that $\|\xi - \gamma\|_{L^\infty} \leq C$ for some $C > 0$ and iterating the previous inequality, we claim that

$$|\xi(t) - \gamma(t)| \leq C \frac{(rRLt)^j}{j!} \quad \forall j \in \mathbb{N}, \forall t \in [0, 1].$$

Indeed, by induction, from (10) we get

$$|\xi(t) - \gamma(t)| \leq rRL \int_0^t C \frac{(rRL)^j}{j!} t^j ds = C \frac{(rRL)^{j+1}}{j!} \frac{t^{j+1}}{j+1}.$$

Finally, since $\lim_{j \rightarrow \infty} \frac{(rRLt)^j}{j!} = 0$, we have $|\xi(t) - \gamma(t)| = 0$ for all t . \square

3.3. The differential of the End-point map is an End-point map. The End-point map behaves like the exponential map: its differential is again an End-point map. In order to make this statement precise, we consider the case $M = \mathbb{R}^n$. Notice that we don't need any bracket-generating condition. In Corollary 3.8 we will use the results on \mathbb{R}^n to prove a statement for all manifolds.

Let $\mathbf{f} : \mathbb{R}^n \times \mathbb{E} \rightarrow \mathbb{R}^n$ be a smooth map. Given a basis e_1, \dots, e_r of \mathbb{E} , we define the vector fields $X_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$ as

$$X_i(p) := \mathbf{f}(p, e_i).$$

The differential of the End-point map with basis point 0 is the map

$$\begin{aligned} \mathbf{dEnd}_0 : L^\infty([0, 1]; \mathbb{E}) \times L^\infty([0, 1]; \mathbb{E}) &\rightarrow \mathbb{R}^n \\ (u, v) &\mapsto \mathbf{dEnd}_0(u)[v]. \end{aligned}$$

Define $Y_i, Z_i : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^n$ for $i = 1, \dots, r$ as

$$\begin{cases} Y_i(p, q) := (X_i(p), \mathbf{d}X_i(p)[q]) \\ Z_i(p, q) := (0, X_i(p)) \end{cases}$$

where $\mathbf{d}X_i(p) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the differential of X_i at p . These vector fields induce a new End-point map

$$\mathbf{End}_{00} : L^\infty([0, 1]; \mathbb{E} \times \mathbb{E}) \rightarrow \mathbb{R}^n \times \mathbb{R}^n$$

with starting point $(0, 0) \in \mathbb{R}^n \times \mathbb{R}^n$.

Proposition 3.7. *For all $u, v \in L^\infty([0, 1]; \mathbb{E})$ it holds*

$$(\mathbf{End}_0(u), \mathbf{dEnd}_0(u)[v]) = \mathbf{End}_{00}(u, v).$$

The proof is immediate, and hence omitted, once one has an explicit representation of the differential $\mathbf{dEnd}_0(u)[v]$, see [Mon02]. This result, together with Proposition 3.6, gives us the weakly* continuity of the differential of the End-point map. The next corollary is an application.

Corollary 3.8. *Let $(\mathbb{E}, \mathbf{f}, \|\cdot\|)$ be a sub-Finsler structure on a manifold M and $o \in M$. Let ρ be a Riemannian metric on M . Then the map $L^\infty([0, 1]; \mathbb{E}) \rightarrow [0, +\infty)$, $u \mapsto \tau(\mathbf{dEnd}_o(u))$ is weakly* lower semi-continuous, where τ is the minimal stretching computed with respect to the norm given by ρ .*

Proof. Let $\{u_k\}_k \subset L^\infty([0, 1]; \mathbb{E})$ be a sequence such that $u_k \xrightarrow{*} u \in L^\infty([0, 1]; \mathbb{E})$. Let γ_u be the curve with control u and starting point o . We can pull back the sub-Finsler structure from a neighborhood of γ_u to an open subset of \mathbb{R}^n via a covering map, so that we reduce the statement to the case $M = \mathbb{R}^n$.

We need to prove

$$(11) \quad \liminf_{k \rightarrow \infty} \tau(\mathbf{dEnd}_o(u_k)) \geq \tau(\mathbf{dEnd}_o(u)).$$

Set $\hat{\tau} := \tau(\text{dEnd}_0(u))$. If $\hat{\tau} = 0$, then (11) is fulfilled, so we assume $\hat{\tau} > 0$. Let $\hat{\tau} > \epsilon > 0$. Then there exists a finite-dimensional vector space $W \subset L^\infty([0, 1]; \mathbb{E})$ such that

$$B_{\text{End}_0(u)}(0, \hat{\tau} - \frac{\epsilon}{2}) \subset\subset \text{dEnd}_0(u)[B_{L^\infty}(0, 1) \cap W],$$

where, for $p \in \mathbb{R}^n$, B_p denotes a ball in $\mathbb{R}^n = T_p\mathbb{R}^n$ with respect to the norm given by ρ at p , and B_{L^∞} denotes a ball in $L^\infty([0, 1]; \mathbb{E})$ with respect to the L^∞ -norm induced by $\|\cdot\|$. Since $\dim W < \infty$ and by Propositions 3.6 and 3.7, the maps $\text{dEnd}_0(u_k)|_W$ strongly converge to $\text{dEnd}_0(u)|_W$. Moreover, the norm on $\mathbb{R}^n = T_p\mathbb{R}^n$ given by ρ continuously depends on $p \in \mathbb{R}^n$. Therefore, for k large enough we have

$$B_{\text{End}_0(u_k)}(0, \hat{\tau} - \epsilon) \subset \text{dEnd}_0(u_k)[B_{L^\infty}(0, 1) \cap W].$$

Hence

$$\liminf_{k \rightarrow \infty} \tau(\text{dEnd}_0(u_k)) \geq \hat{\tau} - \epsilon.$$

Since ϵ is arbitrary, (11) follows. \square

3.4. The sub-Finsler distance is Lipschitz in absence of singular geodesics. The proof of Theorem 3.1 is divided into the next two lemmas from which Theorem 3.1 follows.

Lemma 3.9. *Let $o, p \in M$ such that all d -minimizing curves from o to p are regular. Then there exist a compact neighborhood $K \subset M$ of p and a weakly* compact set $\mathcal{K} \subset L^\infty([0, 1]; \mathbb{E})$ such that:*

- (1) $\text{End}_o : \mathcal{K} \rightarrow K$ is onto;
- (2) $d_{CC}(o, \text{End}_o(u)) = \|u\|_{L^\infty}$ for all $u \in \mathcal{K}$;
- (3) every $u \in \mathcal{K}$ is a regular point for End_o .

Proof. For any compact neighborhood K of p , define the compact set $\mathcal{K}(K) := \{u \in L^\infty([0, 1]; \mathbb{E}) : \text{End}_o(u) \in K \text{ and } d(o, \text{End}_o(u)) = \|u\|_{L^\infty}\}$.

Since the metric d is geodesic, the End-point map $\text{End}_o : \mathcal{K}(K) \rightarrow K$ is surjective, for all K . Moreover, the second requirement holds by definition. Finally, suppose that there exist a sequence $p_k \rightarrow p$ and a sequence $u_k \in L^\infty([0, 1]; \mathbb{E})$ such that $\text{End}_o(u_k) = p_k$, $d(o, p_k) = \|u_k\|_{L^\infty}$ and u_k is a singular point for End_o , for all k . Since the sequence u_k is bounded, thanks to the Banach–Alaoglu Theorem there is $u \in L^\infty([0, 1]; \mathbb{E})$ such that, up to a subsequence, $u_k \xrightarrow{*} u$. By the continuity of End_o , we have $\text{End}_o(u) = p$. By Corollary 3.8, we have $\tau(\text{dEnd}_o(u)) \leq \liminf_k \tau(\text{dEnd}_o(u_k)) = 0$. Finally, by the lower-semicontinuity of the norm with respect to the weak* topology, we have

$$\|u\|_{L^\infty} \leq \liminf_{k \rightarrow \infty} \|u_k\|_{L^\infty} = \liminf_{k \rightarrow \infty} d(o, p_k) = d(o, p) \leq \|u\|_{L^\infty}.$$

So, u is the control of a singular length-minimizing curve from o to p , against the assumption. Therefore, there exists a neighborhood K of p such that $\mathcal{K}(K)$ contains only regular points for End_o . \square

Lemma 3.10. *Let $o \in M$, and $K \subset M$ compact. Suppose there is a weakly* compact set $\mathcal{K} \subset L^\infty([0, 1]; \mathbb{E})$ that satisfies all three properties listed in Lemma 3.9. Then for every Riemannian metric ρ on M there exists $L > 0$ such that the function $d_o : p \mapsto d(o, p)$ is locally L -Lipschitz on the interior of K .*

Proof. Let ρ be a Riemannian metric on M . We first prove that

$$(12) \quad \begin{aligned} & \exists L > 0, \forall q \in K, \exists \hat{\epsilon}_q > 0 \forall q' \in K \\ & [\rho(q, q') < \hat{\epsilon}_q \Rightarrow d_o(q') - d_o(q) \leq L\rho(q, q')]. \end{aligned}$$

Since \mathcal{K} is a weakly* compact set of regular points for End_o , then by Corollary 3.8 the function $u \mapsto \tau(\text{dEnd}_o(u))$ admits a minimum on \mathcal{K} that is strictly positive. By Proposition 3.5, there is $L > 0$ such that for every $u \in \mathcal{K}$ there is $\hat{\epsilon}_u > 0$ such that

$$(13) \quad B_\rho(\text{End}_o(u), \epsilon) \subset \text{End}_o(B_{L^\infty}(u, L\epsilon)) \quad \forall \epsilon < \hat{\epsilon}_u.$$

Let $q, q' \in K$ be such that $q = \text{End}_o(u)$ with $u \in \mathcal{K}$ and $\epsilon := \rho(q, q') < \hat{\epsilon}_u$. Then, by the inclusion (13), there is $u' \in B_{L^\infty}(u, L\epsilon)$ with $\text{End}_o(u') = q'$. So

$$d_o(q') - d_o(q) \leq \|u'\|_{L^\infty} - \|u\|_{L^\infty} \leq \|u' - u\|_{L^\infty} \leq L\epsilon = L\rho(q, q').$$

This proves the claim (12).

Finally, if p is an interior point of K , then there is a ρ -convex neighborhood U of p contained in K (see [O'N83]). So, if $q, q' \in U$, then there is a ρ -geodesic $\xi : [0, 1] \rightarrow U$ joining q to q' . Since the image of ξ is compact, there are $0 = t_1 < s_1 < t_2 < s_2 < \dots < s_{k-1} < t_k = 1$ such that $\rho(\xi(t_i), \xi(s_i)) < \hat{\epsilon}_{\xi(t_i)}$ and $\rho(\xi(s_i), \xi(t_{i+1})) < \hat{\epsilon}_{\xi(t_{i+1})}$. Therefore

$$\begin{aligned} d_o(q') - d_o(q) & \leq \sum_{i=1}^{k-1} d_o(\xi(t_{i+1})) - d_o(\xi(s_i)) + d_o(\xi(s_i)) - d_o(\xi(t_i)) \\ & \leq L \sum_{i=1}^{k-1} \rho(\xi(t_{i+1}), \xi(s_i)) + \rho(\xi(s_i), \xi(t_i)) = L\rho(q, q'). \end{aligned}$$

□

4. REGULARITY OF SPHERES IN GRADED GROUPS

This section is devoted to the proof of Theorems 1.2 and 1.3 and of the inequalities (2).

4.1. The sphere as a graph. Let (G, d) be a homogeneous group and ρ a Riemannian distance on G . Theorem 1.2 and the estimate (2) are both based on the following remark.

Remark 4.1. Let $\mathfrak{g} = \bigoplus_{i>0} V_i$ be a grading for the Lie algebra of G . Let $|\cdot|$ be the norm of a scalar product on \mathfrak{g} that makes the layers orthogonal to each other and let $S = \exp(\{v : |v| = 1\}) \subset G$. The

hypersurface S is smooth and transversal to the dilations, i.e., for all $p \in S$ we have $\frac{d}{dt}|_{t=1}\delta_t p \notin T_p S$. Define

$$\begin{aligned} \phi : S \times (0, +\infty) &\rightarrow G \setminus \{0\} \\ (p, t) &\mapsto \delta_{\frac{1}{t}} p. \end{aligned}$$

Since S is transversal to the dilations, ϕ is a diffeomorphism. Moreover, if $\Gamma := \{(p, d_0(p)) : p \in S\} \subset S \times (0, +\infty)$ is the graph of the function d_0 restricted to S , then

$$\mathbb{S}_d = \phi(\Gamma).$$

Thanks to the last remark, the estimate (2) follows from the next lemma.

Lemma 4.2. *Let $\Omega \subset \mathbb{R}^n$ be an open set and let $f : \Omega \rightarrow \mathbb{R}$ be an α -Hölder function, i.e., for all $x, y \in \Omega$ we have*

$$|f(x) - f(y)| \leq C|x - y|^\alpha,$$

for some $C > 0$, where $\alpha \in (0, 1]$. Define the graph of f as

$$\Gamma_f := \{(x, f(x)) : x \in \Omega\} \subset \mathbb{R}^{n+1}.$$

Then

$$n \leq \dim_H \Gamma_f \leq n + 1 - \alpha,$$

where \dim_H is the Hausdorff dimension. Moreover, this estimate is sharp, i.e., there exists f such that $\dim_H \Gamma_f = n + 1 - \alpha$.

The proof is straightforward by use of a simple covering argument or by an estimate of the Minkowski content of the graph. The sharpness of this result has been shown in [BU37] for the case $n = 1$. The general case, as stated here, is a simple consequence. Indeed, if $g : (0, 1) \rightarrow \mathbb{R}$ is a α -Hölder function such that $\dim_H(\Gamma_g) = 2 - \alpha$, then the graph of the function $f(x_1, \dots, x_n) := g(x_1)$ is $\Gamma_f = \Gamma_g \times (0, 1)^{n-1}$. Therefore, $\dim_H(\Gamma_f) = n + 1 - \alpha$.

In the next easy-to-prove lemma we point out that a homogenous distance is locally Lipschitz if and only if the spheres are Lipschitz graphs in the directions of the dilations.

Lemma 4.3. *Let d be a homogeneous distance on G . Let S and \mathbb{S}_d be as in Remark 4.1 and $p \in S$. Then the following conditions are equivalent:*

- (i) *Setting $\hat{p} := \delta_{d_0(p)-1}(p) \in \mathbb{S}_d$, the sphere \mathbb{S}_d is a Lipschitz graph in the direction $\delta(\hat{p})$ in some neighborhood of p ;*
- (ii) *$d_0|_S : S \rightarrow (0, +\infty)$ is Lipschitz in some neighborhood of p in S ;*
- (iii) *d_0 is Lipschitz in some neighborhood of $\delta_\lambda p$ for one, hence all, $\lambda > 0$.*

Thanks to Lemma 4.3, Theorem 1.2 is a consequence of Theorem 1.1.

4.2. An intrinsic approach. In this section we will prove Theorem 1.3. We define a *cone* in \mathbb{R}^n as

$$\text{Cone}(\alpha, h, v) := \{x \in \mathbb{R}^n : |x| \leq h \text{ and } \angle(x, v) \leq \alpha\} \subset \mathbb{R}^n,$$

where $\alpha \in [0, \pi]$, $h \in (0, +\infty]$, $v \in \mathbb{R}^n$ is the axis of the cone, and $\angle(x, v)$ is the angle between x and v . The following lemma is a simple calculus exercise and it will be used later in the proof of Theorem 1.3.

Lemma 4.4. *Let $m, k, n \in \mathbb{N}$, $p \in \mathbb{R}^m$ and $y_0 \in \mathbb{R}^k$. Let $\phi : \mathbb{R}^m \times \mathbb{R}^k \rightarrow \mathbb{R}^n$ be a smooth map such that $d(\phi_p)(y_0) : \mathbb{R}^k \rightarrow \mathbb{R}^n$ is surjective, where $\phi_x(y) := \phi(x, y)$. Let $C' \subset \mathbb{R}^k$ be a cone with axis $v' \in \mathbb{R}^k$. Then there exist a cone $C \subset \mathbb{R}^n$ with axis $d(\phi_p)(y_0)v'$ and an open neighborhood $U \subset \mathbb{R}^m$ of p such that for all $q \in U$*

$$\phi_q(y_0) + C \subset \phi_q(y_0 + C').$$

Proof of Theorem 1.3. In this proof, we consider the dilations δ_λ as defined for $\lambda \leq 0$ too, with the same definition as for $\lambda > 0$. Notice that in this way the map $G \times \mathbb{R} \rightarrow G$, $(p, \lambda) \mapsto \delta_\lambda p$, is a smooth map.

Let v_1, \dots, v_r be a basis for V_1 and set $p_i := \exp(v_i) \in G$. Up to a rescaling, we can assume $d_0(p_i) < 1$ for all i . For $p \in G$ define $\phi_p : \mathbb{R}^{2r+1} \rightarrow G$ as

$$\phi_p(u_1, \dots, u_r, s, v_1, \dots, v_r) = \delta_{u_1} p_1 \cdots \delta_{u_r} p_r \cdot \delta_s p \cdot \delta_{v_1} p_1 \cdots \delta_{v_r} p_r.$$

Let $\hat{x} \in \mathbb{R}^{2r+1}$ be the point with $u_i = 0$, $s = 1$ and $v_i = 0$, so that $\phi_p(\hat{x}) = p$. The differential of ϕ_p at \hat{x} is given by the partial derivatives

$$\begin{aligned} \frac{\partial \phi_p}{\partial u_i}(\hat{x}) &= \frac{d}{dt} \Big|_{t=0} (\delta_t p_i \cdot p) = dR_p \left(\frac{d}{dt} \Big|_{t=0} (\delta_t p_i) \right) = dR_p(v_i), \\ \frac{\partial \phi_p}{\partial s}(\hat{x}) &= \frac{d}{dt} \Big|_{t=1} (\delta_t p) = \bar{\delta}(p), \\ \frac{\partial \phi_p}{\partial v_i}(\hat{x}) &= \frac{d}{dt} \Big|_{t=0} (p \cdot \delta_t p_i) = dL_p \left(\frac{d}{dt} \Big|_{t=0} (\delta_t p_i) \right) = dL_p(v_i). \end{aligned}$$

Therefore, if $p \in \mathbb{S}_d$ is such that the condition (1) is true, then the differential $d\phi_p$ has full rank at \hat{x} , hence in a neighborhood of \hat{x} .

Define

$$\Delta := \{(u_1, \dots, u_r, s, v_1, \dots, v_r) \in \mathbb{R}^{2r+1} : s + \sum_{i=1}^r (|u_i| + |v_i|) \leq 1\}.$$

We identify G with \mathbb{R}^n through an arbitrary diffeomorphism. So, by Lemma 4.4, there is a cone C with axis $\bar{\delta}(p)$ and a neighborhood U of p such that for all $q \in U$

$$q + C \subset \phi_q(\Delta).$$

Up to restricting U , for all $q \in U$ there are cones C_q with axis $\bar{\delta}(q)$, fixed amplitude and fixed height such that $q + C_q \subset q + C$. Notice that for all $q \in \mathbb{S}_d$ we have $\phi_q(\Delta) \subset \bar{B}_d(0, 1)$ and $\phi_q(\Delta) \cap \mathbb{S}_d = \{q\}$. In particular,

for all $q \in \mathbb{S}_d \cap U$, we have $q + C_q \subset \bar{B}_d(0, 1)$ and $(q + C_q) \cap \mathbb{S}_d = \{q\}$, i.e., $\mathbb{S}_d \cap U$ is a Lipschitz graph in the direction of the dilations. Thanks to Lemma 4.3, we get that d_0 is Lipschitz in a neighborhood of p . \square

Finally, some considerations on condition (1) are due.

Remark 4.5. If (1) holds at $p \in G$ and $u \in L^\infty([0, 1]; V_1)$ is a control such that $\text{End}_0(u) = p$, then the differential $d\text{End}_0(u)$ is surjective, i.e., p is a regular value of End . Indeed, by [LD+15] (see (2.6) and (2.11) there), we have

$$dL_p(V_1) + dR_p(V_1) + \text{span}\{\bar{\delta}(p)\} \subset \text{Im}(d\text{End}_0(u)),$$

because $q \mapsto \bar{\delta}(q)$ is a contact vector field of G .

Proposition 4.6. *Let $X \in V_1$. If (1) holds for $p = \exp(X)$, then*

$$\mathfrak{g} = V_1 + [X, V_1].$$

Proof. Let X_1, \dots, X_r be a basis for V_1 and Y_1, \dots, Y_ℓ a basis for $[X, V_1]$. Let $\alpha_j^i \in \mathbb{R}$ be such that $[X, X_i] = \sum_{j=1}^\ell \alpha_j^i Y_j$. First, notice that

$$\begin{aligned} T_0G &= dL_{\exp(-X)} \left(dL_{\exp(X)}(V_1) + dR_{\exp(X)}(V_1) \right) \\ &= V_1 + dL_{\exp(-X)} \circ dR_{\exp(X)}(V_1) \\ &= V_1 + \text{Ad}_{\exp(X)}(V_1). \end{aligned}$$

Then, using the formula $\text{Ad}_{\exp(X)}(Y) = e^{\text{ad}_X}(Y) = \sum_{k=0}^\infty \frac{1}{k!} \text{ad}_X^k(Y)$, we have

$$\begin{aligned} \text{Ad}_{\exp(X)}(X_i) &= X_i + \left(\sum_{k=1}^\infty \frac{1}{k!} \text{ad}_X^{k-1}([X, X_i]) \right) \\ &= X_i + \left(\sum_{k=1}^\infty \text{ad}_X^{k-1} \left(\sum_{j=1}^\ell \alpha_j^i Y_j \right) \right) \\ &= X_i + \sum_{j=1}^\ell \alpha_j^i \left(\sum_{k=1}^\infty \text{ad}_X^{k-1}(Y_j) \right). \end{aligned}$$

It follows that $\dim(V_1 + \text{Ad}_{\exp(X)}V_1) \leq r + \ell$ and therefore $\dim \mathfrak{g} \leq r + \ell$, i.e., $\mathfrak{g} = V_1 + [X, V_1]$. \square

Proposition 4.7. *Let $Z \in V_k$, where $k > 0$ is such that $V_i = \{0\}$ for all $i > k$. If (1) holds for $p = \exp(Z)$, then*

$$\mathfrak{g} = V_1 + \text{span}\{Z\}.$$

Proof. Since $[Z, \mathfrak{g}] = \{0\}$, we have $R_p = L_p$. Moreover, $\bar{\delta}(p) = dL_p(kZ)$. So, condition (1) becomes $dL_p(V_1) + dL_p(\text{span}\{Z\}) = T_pG$. \square

In particular, if (1) holds for all $p \in G \setminus \{0\}$, then $\mathfrak{g} = V_1 \oplus V_2$ with $\dim V_2 \leq 1$ and $[X, V_1] = V_2$ for all non-zero $X \in V_1$.

5. EXAMPLES

5.1. **Three gradings on \mathbb{R}^2 .** We will present three examples of dilations on \mathbb{R}^2 . In particular we want to illustrate two applications of Theorem 1.3 and show the sharpness of the dimension estimate (2). In Remark 5.5 we give an easy example of a homogeneous distance whose unit ball is a Lipschitz domain, but the distance is not locally Lipschitz away from the diagonal.

The first and the easiest is

$$\delta_\lambda(x, y) := (\lambda x, \lambda y),$$

which gives rise to the known structure of vector space. Here, homogeneous distances are given by norms and balls are convex, hence Lipschitz domains. It's trivial to see that condition (1) holds for all $p \in \mathbb{R}^2$.

The second example is given by the dilations

$$\delta_\lambda(x, y) := (\lambda x, \lambda^2 y).$$

In this case, $\mathbb{R}^2 = V_1 \oplus V_2$ with $V_1 = \mathbb{R} \times \{0\}$ and $V_2 = \{0\} \times \mathbb{R}$, and $\bar{\delta}(x, y) = (x, 2y)$. Condition (1) holds for all $(x, y) \in \mathbb{R}^2$ with $y \neq 0$. One can actually show that, for any homogeneous metric on $(\mathbb{R}^2, \delta_\lambda)$ with closed unit ball B centered at 0, the set $I = \{x \in \mathbb{R} : (x, 0) \in B\}$ is a closed interval and there exists a function $f : I \rightarrow \mathbb{R}$ that is locally Lipschitz on the interior of I such that

$$\mathbb{S}_d \cap \{(x, y) : y \geq 0\} = \{(x, f(x)) : x \in I\}.$$

We will prove a similar statement in the Heisenberg group with an argument that applies here too, see Section 6.

The third example is given by the dilations

$$(14) \quad \delta_\lambda(x, y) := (\lambda^2 x, \lambda^2 y),$$

and it is interesting because of the next proposition.

Proposition 5.1. *There exists a homogeneous (with respect to dilations (14)) distance d on \mathbb{R}^2 whose unit sphere has Euclidean Hausdorff dimension $\frac{3}{2}$.*

Notice that $\frac{3}{2}$ is the maximal Hausdorff dimension that one gets by the estimate (2).

For proving Proposition 5.1, we need to find a set $B \subset \mathbb{R}^2$ that satisfies all four conditions listed in Lemma 2.15, in particular

$$(15) \quad \forall p, q \in B, \forall t \in [0, 1] \quad t^2 p + (1-t)^2 q \in B.$$

One easily proves the following preliminary facts.

Lemma 5.2. *Let $p, q \in \mathbb{R}^2$ and $\gamma : [0, 1] \rightarrow \mathbb{R}^2$, $\gamma(t) := t^2 p + (1-t)^2 q$.*

(1) *The curve γ is contained in the triangle of vertices $0, p, q$.*

- (2) The curve γ is an arc of the parabola passing through p and q and that is tangent to the lines $\text{span}\{p\}$ and $\text{span}\{q\}$.
- (3) If B satisfies (15) and $A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear map, then $A(B)$ satisfies (15) as well.

Lemma 5.3. For $0 < C \leq 1$, define

$$Y_C := \{(x, y) : |x| \leq 1, y \leq 1 + C\sqrt{|x|}\}.$$

Then Y_C satisfies (15).

Proof. Let $p, q \in Y_C$ and set $\gamma(t) = (\gamma_x(t), \gamma_y(t)) := t^2p + (1-t)^2q$.

If both p and q stay on one side with respect to the vertical axis, then $\gamma(t) \in Y_C$ for all $t \in [0, 1]$ thanks to the first point of Lemma 5.2 and because the two sets $Y_C \cap \{x \geq 0\}$ and $Y_C \cap \{x \leq 0\}$ are convex.

Therefore, we suppose that

$$p = (-p_x, p_y) \quad q = (q_x, q_y)$$

with $p_x, q_x > 0$. Let $t_0 \in [0, 1]$ be the unique value such that $\gamma_x(t_0) = 0$. Then the curve γ lies in the union of the two triangles with vertices $0, \gamma(0), \gamma(t_0)$ and $0, \gamma(1), \gamma(t_0)$, respectively. Therefore, γ lies in Y_C if and only if $\gamma_y(t_0) \leq 1$. Solving the equation $\gamma_x(t_0) = t_0^2(q_x - p_x) - 2q_x t_0 + q_x = 0$, one gets

$$t_0 = \frac{\sqrt{q_x}}{\sqrt{q_x} + \sqrt{p_x}}, \quad (1 - t_0) = \frac{\sqrt{p_x}}{\sqrt{q_x} + \sqrt{p_x}}.$$

From the expression of $\gamma_y(t_0) = t_0^2 p_y + (1 - t_0)^2 q_y$, we notice that, p_x and q_x fixed, the worst case is when p_y and q_y are maximal, i.e.,

$$p_y = 1 + C\sqrt{p_x}, \quad q_y = 1 + C\sqrt{q_x}.$$

Finally

$$\begin{aligned} \gamma_y(t_0) &= t_0^2 p_y + (1 - t_0)^2 q_y \\ &= \frac{q_x}{(\sqrt{q_x} + \sqrt{p_x})^2} (1 + C\sqrt{p_x}) + \frac{p_x}{(\sqrt{q_x} + \sqrt{p_x})^2} (1 + C\sqrt{q_x}) \\ &= \frac{1}{(\sqrt{q_x} + \sqrt{p_x})^2} (q_x + p_x + Cq_x\sqrt{p_x} + Cp_x\sqrt{q_x}) \\ &= 1 + \frac{-2\sqrt{p_x q_x} + Cq_x\sqrt{p_x} + Cp_x\sqrt{q_x}}{(\sqrt{q_x} + \sqrt{p_x})^2} \\ &= 1 + \sqrt{p_x q_x} \frac{-2 + C(\sqrt{q_x} + \sqrt{p_x})}{(\sqrt{q_x} + \sqrt{p_x})^2}. \end{aligned}$$

Since $-2 + C(\sqrt{q_x} + \sqrt{p_x}) \leq 0$, then we have $\gamma_y(t_0) \leq 1$, as wanted. \square

Lemma 5.4. Let $\alpha, \beta > 0$. For all $0 < \epsilon \leq \alpha$ and all $0 < C \leq \frac{\beta}{\sqrt{\alpha}}$, the set

$$Y(\epsilon, \beta, C) := \{(x, y) : |x| \leq \epsilon, y \leq \beta + C\sqrt{|x|}\}$$

satisfies (15).

Proof. Define the linear map $A(x, y) := (\alpha x, \beta y)$ and set $C' := C \frac{\sqrt{\alpha}}{\beta} \leq 1$. Then one just needs to check that

$$Y(\epsilon, \beta, C) = A(Y_{C'}) \cap \{(x, y) : |x| \leq \epsilon\},$$

where $Y_{C'}$ is defined as in the previous Lemma 5.3. \square

Proof of Proposition 5.1. First of all, let $\theta_0 > 0$ be such that for all $|\theta| \leq \theta_0$ it holds

$$(16) \quad \frac{|\theta|}{2} \leq |\cos(\frac{\pi}{2} + \theta)| = |\sin \theta| \leq 2|\theta|.$$

Moreover, let $L, m, M, C > 0$ be such that

$$\frac{L\sqrt{2}}{\sqrt{m}} \leq C \leq \frac{m}{\sqrt{2M\theta_0}}.$$

Let $f : \mathbb{R} \rightarrow (0, +\infty)$ be a function such that

$$(17) \quad \forall s, t \in \mathbb{R} \quad |f(t) - f(s)| \leq L\sqrt{|t - s|},$$

$$(18) \quad \forall t \in \mathbb{R} \quad m \leq f(t) \leq M.$$

We claim that, for $|\theta| \leq \theta_0$, we have

$$(19) \quad f\left(\frac{\pi}{2} + \theta\right) \cdot \left(\cos\left(\frac{\pi}{2} + \theta\right), \sin\left(\frac{\pi}{2} + \theta\right)\right) \in Y\left(2M\theta_0, f\left(\frac{\pi}{2}\right), C\right)$$

where $Y(2M\theta_0, f(\frac{\pi}{2}), C)$ is defined as in Lemma 5.4. Indeed, we have on one side

$$|x| := |f\left(\frac{\pi}{2} + \theta\right) \cos\left(\frac{\pi}{2} + \theta\right)| \leq M2|\theta| \leq 2M\theta_0.$$

On the other side,

$$\begin{aligned} y &:= f\left(\frac{\pi}{2} + \theta\right) \sin\left(\frac{\pi}{2} + \theta\right) \leq f\left(\frac{\pi}{2} + \theta\right) \\ &\leq f\left(\frac{\pi}{2}\right) + f\left(\frac{\pi}{2} + \theta\right) - f\left(\frac{\pi}{2}\right) \leq f\left(\frac{\pi}{2}\right) + L\sqrt{|\theta|} \\ &\leq f\left(\frac{\pi}{2}\right) + L \frac{\sqrt{2 \cos\left(\frac{\pi}{2} + \theta\right) f\left(\frac{\pi}{2} + \theta\right)}}{\sqrt{f\left(\frac{\pi}{2} + \theta\right)}} \leq f\left(\frac{\pi}{2}\right) + \frac{\sqrt{2}L}{\sqrt{m}} \sqrt{|x|} \\ &\leq f\left(\frac{\pi}{2}\right) + C\sqrt{|x|}. \end{aligned}$$

So (19) is satisfied.

Since for $\alpha := 2M\theta_0$ and $\beta := f(\frac{\pi}{2})$ we have

$$\frac{\beta}{\sqrt{\alpha}} = \frac{f(\frac{\pi}{2})}{\sqrt{2M\theta_0}} \geq \frac{m}{\sqrt{2M\theta_0}} \geq C,$$

Lemma 5.4 applies and we get that $Y(2M\theta_0, f(\frac{\pi}{2}), C)$ satisfies (15).

For any θ we set A_θ to be the counterclockwise rotation of angle θ :

$$A_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

Define the curve $\phi(t) := f(t)(\cos t, \sin t)$. Notice that $A_\theta\phi(t) = f((t - \theta) + \theta)(\cos(t + \theta), \sin(t + \theta))$ and that the function $s \mapsto f(s + \theta)$ is still satisfying both (17) and (18). So we have that, for $|t|, |s| < \frac{\theta_0}{2}$,

$$\phi\left(\frac{\pi}{2} + t\right) \in A_s[Y(2M\theta_0, f\left(\frac{\pi}{2} + s\right), C)]$$

and the set $A_s[Y(2M\theta_0, f\left(\frac{\pi}{2} + s\right), C)]$ satisfies (15).

Set

$$B := \bigcap_{|s| < \frac{\theta_0}{2}} \left(A_s[Y(2M\theta_0, f\left(\frac{\pi}{2} + s\right), C)] \cap -A_s[Y(2M\theta_0, f\left(\frac{\pi}{2} + s\right), C)] \right).$$

The set B satisfies all three conditions of Lemma 2.15, hence it is the unit ball of a homogeneous metric. Moreover,

$$\left\{ \phi\left(\frac{\pi}{2} + t\right) : |t| < \frac{\theta_0}{2} \right\} \subset \partial B.$$

The statement of Proposition 5.1 follows because there are functions $f : \mathbb{R} \rightarrow [0, +\infty)$ that satisfy (17) and (18) and such that the image of the curve ϕ has Hausdorff dimension $\frac{3}{2}$. Indeed, the image of ϕ has the same Hausdorff dimension of the graph of f , and then one uses the sharpness of Lemma 4.2. \square

Remark 5.5. Using the same arguments as in the proof of Lemma 5.3, one easily shows that the set

$$B := \{(x, y) \in \mathbb{R}^2 : |x| \leq 1, -f(-x) \leq y \leq f(x)\},$$

where

$$f(x) := \begin{cases} 1 & x \leq 0 \\ 1 + \sqrt{x} & x > 0, \end{cases}$$

is the unit ball of a homogeneous distance on \mathbb{R}^2 with dilations (14). Notice that such B is a Lipschitz domain, but the associated homogeneous distance is not Lipschitz in any neighborhood of the point $(0, 1)$, thanks to Lemma 4.3.

5.2. The Heisenberg groups. In the Heisenberg groups \mathbb{H}^n (for an introduction see [Cap+07]) condition (1) holds at every non-zero point. Therefore, balls of any homogeneous metric on \mathbb{H}^n are Lipschitz domains. We will treat more in detail the first Heisenberg group in Section 6.

5.3. A sub-Finsler sphere with a cusp. Let \mathbb{H} be the first Heisenberg group (see Section 6 for the definition). The group $G = \mathbb{H} \times \mathbb{R}$ is a stratified group with grading $(V_1 \times \mathbb{R}) \oplus V_2$, where $V_1 \oplus V_2$ is a stratification for \mathbb{H} . The line $\{0_{\mathbb{H}}\} \times \mathbb{R}$ is a singular curve in G . Moreover, it has been shown in [BLD13] that there exists a sub-Finsler distance on G whose unit sphere \mathbb{S}_d has a cusp in the intersection $\mathbb{S}_d \cap (\{0_{\mathbb{H}}\} \times \mathbb{R})$.

However, for sub-Riemannian metrics we still have balls that are Lipschitz domains, as the following Proposition 5.7 shows. But let us first recall a simple fact:

Lemma 5.6. *Let \mathbb{A} and \mathbb{B} be two stratified groups with stratifications $\bigoplus V_i$ and $\bigoplus W_i$, respectively. Endow V_1 and W_1 with a scalar product each and let d_A, d_B be the corresponding homogeneous sub-Riemannian distances.*

Then $\mathbb{A} \times \mathbb{B}$ is a Carnot group with stratification $\bigoplus_i V_i \times W_i$ and metric

$$d((a, b), (a', b')) := \sqrt{d_A(a, a')^2 + d_B(b, b')^2},$$

which is the sub-Riemannian metric generated by the scalar product on $V_1 \times W_1$ induced by the scalar products on V_1 and W_1 .

One proves this lemma by using the fact that the energy of curves on $\mathbb{A} \times \mathbb{B}$ (i.e., the integral of the squared norm of the derivative) is the sum of the energies of the two components of the curve.

Proposition 5.7. *Any homogeneous sub-Riemannian metric on $\mathbb{H} \times \mathbb{R}$ is locally Lipschitz away from the diagonal.*

Proof. First of all, we show that, up to isometry, there is only one homogeneous sub-Riemannian distance on $\mathbb{H} \times \mathbb{R}$. Let (X_1, Y_1, T_1) and (X_2, Y_2, T_2) be two basis of $V_1 \times \mathbb{R}$ that are orthonormal for two sub-Riemannian structures, respectively. We may assume $T_1, T_2 \in \{0\} \times \mathbb{R}$. Notice that $[X_i, Y_i] \notin V_1 \times \mathbb{R}$. The linear map such that $X_1 \mapsto X_2$, $Y_1 \mapsto Y_2$, $T_1 \mapsto T_2$, $[X_1, Y_1] \mapsto [X_2, Y_2]$ is an automorphism of Lie algebras and induces an isometry between the two sub-Riemannian structures.

Denoting by $d_{\mathbb{H}}$ and $d_{\mathbb{R}}$ the standard metrics on \mathbb{H} and \mathbb{R} , respectively, we prove the proposition for the product metric as in Lemma 5.6. Namely, we need to check that the function

$$(20) \quad (p, t) \mapsto d((0, 0), (p, t)) = \sqrt{d_{\mathbb{H}}(0, p)^2 + t^2}$$

is locally Lipschitz at all $(\hat{p}, \hat{t}) \neq (0, 0)$. This follows directly from Proposition 3.3. \square

5.4. A sub-Riemannian sphere with a cusp.

Proposition 5.8. *Let G be a Carnot group of step 3 endowed with a sub-Riemannian distance d_G . Then the sub-Riemannian distance d on $G \times \mathbb{R}$ given by*

$$d((p, s), (q, t)) = \sqrt{d_E(p, q)^2 + |t - s|^2}$$

has a unit sphere with a cusp at $(0_G, 1)$.

Proof. Let $\mathfrak{g} = \bigoplus_{i=1}^3 V_i$ be the Lie algebra of G and fix $Z \in V_3 \setminus \{0\}$. We identify \mathfrak{g} with G through the exponential map.

The intersection of the unit sphere in $(G \times \mathbb{R}, d)$ with the plane $\text{span}\{Z\} \times \mathbb{R}$ is given by all points (zZ, t) such that

$$(21) \quad d_G(0, zZ)^2 + t^2 = 1.$$

Since d_G is homogeneous on G , there exists $C > 0$ such that for all $z \in \mathbb{R}$

$$(22) \quad d_G(0, zZ) = C|z|^{\frac{1}{3}}.$$

Putting together (21) and (22) we obtain that this intersection consists of all the points (zZ, t) such that

$$|z| = \left(\frac{1+t}{C^2}\right)^{\frac{3}{2}} \cdot (1-t)^{\frac{3}{2}}.$$

One then easily sees that this set in \mathbb{R}^2 has a cusp at $(0, 1)$. \square

6. A CLOSER LOOK AT THE HEISENBERG GROUP

The Heisenberg group \mathbb{H} is the easiest example of a stratified group that is not Abelian and for this reason it has been studied in large extend. The most common homogeneous metrics on \mathbb{H} are the Koràny metric and the sub-Riemannian metric. Sub-Finsler metrics on \mathbb{H} arise in the study of finitely-generated groups, see [Bre14] and references therein. The geometry of spheres has been studied in [LDR14] and [DM14].

The Lie algebra \mathfrak{h} of the Heisenberg group is a three dimensional vector space $\text{span}\{X, Y, Z\}$ with a Lie bracket operation defined by the only nontrivial relation $[X, Y] = Z$.

We identify the Heisenberg group \mathbb{H} again with $\text{span}\{X, Y, Z\}$, where we define the group operation

$$p \cdot q := p + q + \frac{1}{2}[p, q] \quad \forall p, q \in \mathbb{H}.$$

Hence \mathfrak{h} is the Lie algebra of \mathbb{H} and the exponential map $\mathfrak{h} \rightarrow \mathbb{H}$ is the identity map. Notice that the inverse of an element p is $p^{-1} = -p$.

The Heisenberg algebra admits the stratification $\mathfrak{h} = V_1 \oplus V_2$ with $V_1 = \text{span}\{X, Y\}$ and $V_2 = \text{span}\{Z\}$. Denote by π the linear projection $\mathfrak{h} \rightarrow V_1$ along V_2 . Notice that this map, regarded as $\pi : (\mathbb{H}, \cdot) \rightarrow (V_1, +)$, is a group morphism.

The dilations $\delta_\lambda : \mathbb{H} \rightarrow \mathbb{H}$ are explicitly expressed by

$$\delta_\lambda(xX + yY + zZ) = x\lambda X + y\lambda Y + z\lambda^2 Z, \quad \forall \lambda > 0.$$

These are both Lie algebra automorphisms $\delta_\lambda : \mathfrak{h} \rightarrow \mathfrak{h}$ and Lie group automorphisms $\mathbb{H} \rightarrow \mathbb{H}$.

Three are the main results of this section.

Proposition 6.1. *Let $N : \mathbb{H} \rightarrow [0, +\infty)$ be a homogeneous norm. Then the unit ball*

$$B := \{p \in \mathbb{H} : N(p) \leq 1\}$$

is a star-like Lipschitz domain.

Proof. One easily shows that condition (1) holds for all $p \in \mathbb{H} \setminus \{0\}$. In order to prove that B is star-like, one first notice that if $p \in B$, then $-p \in B$, hence $\delta_t(p)\delta_{1-t}(-p) = (2t-1)p \in B$ for all $t \in [0, 1]$, and this is a straight line passing through zero. \square

Proposition 6.2. *Let N and B as in Proposition 6.1. Set $K := \pi(B) \subset V_1$. Then K is a compact, convex set with $K = -K$ and $K = \text{cl}(\text{int}(K))$, and there exists a function $f : K \rightarrow [0, +\infty)$, locally Lipschitz on $\text{int}(K)$, such that*

$$(23) \quad B = \{v + zZ : v \in K, -f(-v) \leq z \leq f(v)\}.$$

The proof is postponed to Section 6.1.

We remark that homogeneous distances and sub-Finsler homogeneous distances on \mathbb{H} have a precise relation. Indeed, if d is a homogeneous distance on \mathbb{H} , then it is easy to show that the length distance generated by d is exactly the sub-Finsler distance that has the norm on V_1 generated by the set K defined in Proposition 6.2.

Proposition 6.3. *Let $K \subset V_1$ be a compact, convex set with $-K = K$ and $0 \in \text{int}(K)$. Let $g : K \rightarrow \mathbb{R}$ be Lipschitz. Then there exists $b \in \mathbb{R}$ such that for $f := g + b$ the set B as in (23) is the unit ball of a homogeneous norm.*

The proof will appear in Section 6.2.

As a consequence of Proposition 6.3, we get the existence of homogeneous distances on \mathbb{H} that are not almost convex in the sense of [DS14]. Indeed, one can take the distance associated to $g(xX + yY) = |x|$ from Proposition 6.3.

6.1. Proof of Proposition 6.2.

Lemma 6.4. *Let $B \subset \mathbb{H}$ be an arbitrary closed set satisfying (7). If $p = v + zZ \in B$ with $v = \pi(p) \in V_1$, then $v + szZ \in B$, for all $s \in [0, 1]$. In particular,*

- (1) $\pi(B) = B \cap V_1$;
- (2) $\pi(B) \subset V_1$ is convex.

Proof. We have that for all $t \in [0, 1]$

$$B \ni \delta_t p \cdot \delta_{1-t} p = v + (t^2 + (1-t)^2)zZ.$$

Since the image of $[0, 1]$ through the map $t \mapsto (t^2 + (1-t)^2)$ is $[\frac{1}{2}, 1]$, then it follows $v + szZ \in B$ for all $s \in [\frac{1}{2}, 1]$. Iterating this process and using the closeness of B , we get $v + szZ \in B$ for all $s \in [0, 1]$. For the

last statement, take $v, w \in \pi(B) \subset B$ and notice that $tv + (1-t)w = \pi(\delta_t v \cdot \delta_{1-t} w) \in \pi(B)$. \square

Let $B = \{N \leq 1\}$ be the unit ball of a homogeneous norm and set $K := \pi(B) \subset V_1$ and $\Omega := \text{int}(K)$. First, we check that $\bar{\Omega} = K$. On the one hand, clearly we have $\bar{\Omega} \subset K$. On the other hand, if $v \in K$, then for any $t \in [0, 1)$ we have $N(\delta_t v) = tN(v) < 1$, i.e., $\delta_t v = tv \in \text{int}B \cap V \subset \Omega$. Hence $v \in \bar{\Omega}$.

If we define $f : K \rightarrow [0, +\infty)$ as $f(v) := \max\{z : v + zZ \in Q\}$, then we have (23). In order to prove that f is locally Lipschitz on Ω , we need to prove

$$(24) \quad \begin{aligned} & \forall p \in \partial B \cap \{z \geq 0\} \cap \pi^{-1}(\Omega), \\ & \exists U \ni p \text{ open, } \exists C \text{ vertical cone, s.t.} \\ & \forall q \in U \cap \partial B \text{ it holds } q + C \subset B. \end{aligned}$$

Here a *vertical cone* is a Euclidean cone with axis $-Z$ and non-empty interior.

So, fix $p \in \partial Q \cap \{z \geq 0\}$ such that $\pi(p) \in \Omega$. Define for $\theta \in \mathbb{R}$ and $\epsilon > 0$

$$v_\theta := x_\theta X + y_\theta Y := \epsilon \cos(\theta)X + \epsilon \sin(\theta)Y.$$

For $\epsilon > 0$ small enough, $\pi(p) + v_\theta \in \Omega$ for all θ . Define

$$\phi(t, \theta) := \delta_{(1-t)p} \cdot \delta_t(\pi(p) + v_\theta).$$

Clearly $\phi(t, \theta) \in B$ for $t \in [0, 1]$ and $\theta \in \mathbb{R}$, and $\phi(0, \theta) = p$ for all θ . Geometrically, $\phi([0, 1] \times \mathbb{R})$ is a ‘‘tent’’ inside B standing above the whole vertical segment from $\pi(p)$ to p . Notice that $p \neq \pi(p)$, indeed $N(p) = 1$ while $N(\pi(p)) < 1$, because $\pi(p) \in \Omega$.

We only need to prove that the curves $t \mapsto \phi(t, \theta)$ meet this vertical segment by an angle bounded away from 0. Some computations are needed: set $p = p_1 X + p_2 Y + p_3 Z$, then

$$\phi(t, \theta) = \pi(p) + tv_\theta + \left(\frac{1}{2}t(1-t)(p_1 y_\theta - p_2 x_\theta) + (1-t)^2 p_3 \right) Z.$$

We take care only of the third coordinate. Set

$$\begin{aligned} g(t) &:= \frac{1}{2}t(1-t)(p_1 y_\theta - p_2 x_\theta) + (1-t)^2 p_3 \\ &= t^2 \left(-\frac{1}{2}(p_1 y_\theta - p_2 x_\theta) + p_3 \right) + t \left(\frac{1}{2}(p_1 y_\theta - p_2 x_\theta) - 2p_3 \right) + p_3. \end{aligned}$$

Saying that the angle between the curve $t \mapsto \phi(t, \theta)$ and the vertical segment at p is uniformly greater than zero, is equivalent to give an upper bound to the derivative of g at 0 for all θ . Since

$$g'(0) = \frac{1}{2}(p_1 y_\theta - p_2 x_\theta) - 2p_3,$$

we are done.

Finally, since both ϵ and $g'(0)$ depend continuously on p , then (24) is satisfied. \square

6.2. Proof of Proposition 6.3. We consider the bilinear map $\omega : V_1 \times V_1 \rightarrow \mathbb{R}$ given by

$$\omega(v_1X + v_2Y, w_1X + w_2Y) := v_1w_2 - v_2w_1.$$

Lemma 6.5. *For any continuous function $f : K \rightarrow [0, +\infty)$, the set B as in (23) is the unit ball of a homogeneous norm on \mathbb{H} if and only if*

$$(25) \quad \forall v, w \in K \quad \forall t \in [0, 1] \\ f(tv + (1-t)w) - t^2f(v) - (1-t)^2f(w) - \frac{t(1-t)}{2}\omega(v, w) \geq 0.$$

Proof. One easily sees that $B = B^{-1}$. Notice that B is the unit ball of a homogeneous norm if and only if it satisfies (7).

\Rightarrow Assume that B satisfies (7). Then for any $v, w \in K$ we have

$$\begin{aligned} B \ni \delta_t(v + f(v)Z) \cdot \delta_{(1-t)}(w + f(w)Z) &= \\ &= tv + (1-t)w + \left(t^2f(v) + (1-t)^2f(w) + \frac{1}{2}t(1-t)\omega(v, w) \right) Z, \end{aligned}$$

hence

$$t^2f(v) + (1-t)^2f(w) + \frac{1}{2}t(1-t)\omega(v, w) \leq f(tv + (1-t)w).$$

\Leftarrow Suppose f satisfies (25). Define

$$\begin{aligned} B^+ &:= \{v + zZ : v \in K \text{ and } z \leq f(v)\}, \\ B^- &:= \{v + zZ : v \in K \text{ and } -f(-v) \leq z\}. \end{aligned}$$

We will show that both B^+ and B^- satisfy (7), from which it follows that $B = B^+ \cap B^-$ satisfies (7) as well.

So, let $v, w \in K$ and $z_1, z_2 \in \mathbb{R}$ such that $v + z_1Z, w + z_2Z \in B^+$. Then the third coordinate of $\delta_t(v + z_1Z) \cdot \delta_{(1-t)}(w + z_2Z)$ satisfies

$$\begin{aligned} t^2z_1 + (1-t)^2z_2 + \frac{1}{2}t(1-t)\omega(v, w) &\leq \\ &\leq t^2f(v) + (1-t)^2f(w) + \frac{1}{2}t(1-t)\omega(v, w) \leq f(tv + (1-t)v), \end{aligned}$$

therefore we have $\delta_t(v + z_1Z) \cdot \delta_{(1-t)}(w + z_2Z) \in B^+$ for all $t \in [0, 1]$.

The calculation for B^- is similar. \square

The verification of the next lemma is simple and therefore it is omitted.

Lemma 6.6. *Suppose that $g : K \rightarrow \mathbb{R}$ is a continuous function such that there is a constant $A \in \mathbb{R}$ with*

$$(26) \quad \forall v, w \in K, \quad \forall t \in [0, 1] \\ g(tv + (1-t)w) - t^2g(v) - (1-t)^2g(w) \geq At(1-t).$$

Then $f := g + B$ satisfies (25) with

$$B := \sup_{v,w \in K} \frac{1}{2} \left(\frac{1}{2} \omega(v, w) - A \right) = \frac{1}{4} \left(\sup_{v,w \in K} \omega(v, w) \right) - \frac{1}{2} A.$$

Lemma 6.7. *Let $g : K \rightarrow \mathbb{R}$ be L -Lipschitz. Then g satisfies (26) for*

$$A := -2L \operatorname{diam}(K) - 4 \sup_{p \in K} |g(p)|.$$

Proof. Notice that we need to show that (26) holds only for $t \in (0, 1)$ and that (26) is symmetric in t and $(1-t)$. So, it is enough to consider only the case $t \in (0, \frac{1}{2}]$:

$$\begin{aligned} & \frac{g(tv + (1-t)w) - t^2g(v) - (1-t)^2g(w)}{t(1-t)} \\ &= \frac{g(w + t(v-w)) - g(w)}{t(1-t)} + \frac{(1 - (1-t)^2)g(w)}{t(1-t)} - \frac{t}{1-t}g(v) \\ &\geq -\frac{L\|v-w\|}{1-t} + \frac{2-t}{1-t}g(w) - \frac{t}{1-t}g(v) \\ &\geq -2L \operatorname{diam}(K) - 4 \sup_{p \in K} |g(p)|. \end{aligned}$$

□

Putting together Lemmas 6.7, 6.6, and 6.5, we get Proposition 6.3.

□

REFERENCES

- [ABB15] A. A. Agrachëv, D. Barilari, and U. Boscain. *Introduction to Riemannian and sub-Riemannian geometry*. Tech. rep. 2015. URL: <http://webusers.imj-prg.fr/~daveide.barilari/Notes.php>.
- [Agr09] A. A. Agrachëv. “Any sub-Riemannian metric has points of smoothness”. In: *Dokl. Akad. Nauk* 424.3 (2009), pp. 295–298. ISSN: 0869-5652. DOI: [10.1134/S106456240901013X](https://doi.org/10.1134/S106456240901013X). URL: <http://dx.doi.org/10.1134/S106456240901013X>.
- [AT04] L. Ambrosio and P. Tilli. *Topics on analysis in metric spaces*. Vol. 25. Oxford Lecture Series in Mathematics and its Applications. Oxford University Press, Oxford, 2004, pp. viii+133. ISBN: 0-19-852938-4.
- [BBI01] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*. Providence, RI: American Mathematical Society, 2001. ISBN: 0821821296.

- [BLD13] E. Breuillard and E. Le Donne. “On the rate of convergence to the asymptotic cone for nilpotent groups and subFinsler geometry”. In: *Proc. Natl. Acad. Sci. USA* 110.48 (2013), pp. 19220–19226. ISSN: 1091-6490. DOI: [10.1073/pnas.1203854109](https://doi.org/10.1073/pnas.1203854109). URL: <http://dx.doi.org/10.1073/pnas.1203854109>.
- [Bre14] E. Breuillard. “Geometry of locally compact groups of polynomial growth and shape of large balls”. In: *Groups Geom. Dyn.* 8.3 (2014), pp. 669–732. ISSN: 1661-7207. DOI: [10.4171/GGD/244](https://doi.org/10.4171/GGD/244). URL: <http://dx.doi.org/10.4171/GGD/244>.
- [BU37] A. S. Besicovitch and H. D. Ursell. “Sets of Fractional Dimensions (V): on dimensional numbers of some continuous curves”. In: *J. London Math. Soc.* 01 (1937), pp. 18–25.
- [Cap+07] L. Capogna, D. Danielli, S. D. Pauls, and J. T. Tyson. *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*. Vol. 259. Progress in Mathematics. Birkhäuser Verlag, Basel, 2007, pp. xvi+223. ISBN: 978-3-7643-8132-5; 3-7643-8132-9.
- [Cla+98] F. H. Clarke, Y. S. Ledyaev, R. J. Stern, and P. R. Wolenski. *Nonsmooth analysis and control theory*. Vol. 178. Graduate Texts in Mathematics. Springer-Verlag, New York, 1998, pp. xiv+276. ISBN: 0-387-98336-8.
- [DM14] M. Duchin and C. Mooney. “Fine asymptotic geometry in the Heisenberg group”. In: *Indiana Univ. Math. J.* 63.3 (2014), pp. 885–916. ISSN: 0022-2518. DOI: [10.1512/iumj.2014.63.5308](https://doi.org/10.1512/iumj.2014.63.5308). URL: <http://dx.doi.org/10.1512/iumj.2014.63.5308>.
- [DS14] M. Duchin and M. Shapiro. “Rational growth in the Heisenberg group”. In: *ArXiv e-prints* (Nov. 2014). arXiv: [1411.4201](https://arxiv.org/abs/1411.4201) [[math.GR](https://arxiv.org/abs/1411.4201)].
- [HS90] W. Hebisch and A. Sikora. “A smooth subadditive homogeneous norm on a homogeneous group”. In: *Studia Math.* 96.3 (1990), pp. 231–236. ISSN: 0039-3223.
- [Jur97] V. Jurdjevic. *Geometric control theory*. Vol. 52. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1997, pp. xviii+492. ISBN: 0-521-49502-4.
- [LD+15] E. Le Donne, R. Montgomery, A. Ottazzi, P. Pansu, and D. Vittone. “Sard property for the endpoint map on some Carnot groups”. In: *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis* (2015), pp. –. ISSN: 0294-1449. DOI: [http://dx.doi.org/10.1016/j.anihpc.2015.07.004](https://doi.org/10.1016/j.anihpc.2015.07.004). URL: <http://www.sciencedirect.com/science/article/pii/S029414491500075X>.

- [LD15a] E. Le Donne. *A course on Carnot groups*. Manuscript. 2015.
- [LD15b] E. Le Donne. “A metric characterization of Carnot groups”. In: *Proc. Amer. Math. Soc.* 143.2 (2015), pp. 845–849. ISSN: 0002-9939. DOI: [10.1090/S0002-9939-2014-12244-1](https://doi.org/10.1090/S0002-9939-2014-12244-1). URL: <http://dx.doi.org/10.1090/S0002-9939-2014-12244-1>.
- [LDR14] E. Le Donne and S. Rigot. “Besicovitch Covering Property for homogeneous distances in the Heisenberg groups”. In: *ArXiv e-prints* (June 2014). arXiv: [1406.1484](https://arxiv.org/abs/1406.1484) [[math.MG](https://arxiv.org/abs/1406.1484)].
- [LDR15] E. Le Donne and S. Rigot. “Remarks about Besicovitch covering property in Carnot groups of step 3 and higher”. In: *Proc. Amer. Math. Soc.* (2015).
- [Mon02] R. Montgomery. *A tour of subriemannian geometries, their geodesics and applications*. Vol. 91. Mathematical Surveys and Monographs. Providence, RI: American Mathematical Society, 2002, pp. xx+259. ISBN: 0-8218-1391-9.
- [O’N83] B. O’Neill. *Semi-Riemannian geometry : with applications to relativity*. Pure and applied mathematics. New York: Academic Press, 1983. ISBN: 0-12-526740-1.
- [Rif04] L. Rifford. “A Morse-Sard theorem for the distance function on Riemannian manifolds”. In: *Manuscripta Math.* 113.2 (2004), pp. 251–265. ISSN: 0025-2611. DOI: [10.1007/s00229-003-0436-7](https://doi.org/10.1007/s00229-003-0436-7). URL: <http://dx.doi.org/10.1007/s00229-003-0436-7>.
- [Rif06] L. Rifford. “À propos des sphères sous-riemanniennes”. In: *Bull. Belg. Math. Soc. Simon Stevin* 13.3 (2006), pp. 521–526. ISSN: 1370-1444. URL: <http://projecteuclid.org/euclid.bbms/1161350693>.
- [Rif14] L. Rifford. *Sub-Riemannian geometry and optimal transport*. Springer Briefs in Mathematics. Springer, Cham, 2014, pp. viii+140. ISBN: 978-3-319-04803-1; 978-3-319-04804-8. DOI: [10.1007/978-3-319-04804-8](https://doi.org/10.1007/978-3-319-04804-8). URL: <http://dx.doi.org/10.1007/978-3-319-04804-8>.
- [Sie86] E. Siebert. “Contractive automorphisms on locally compact groups”. In: *Math. Z.* 191.1 (1986), pp. 73–90. ISSN: 0025-5874. DOI: [10.1007/BF01163611](https://doi.org/10.1007/BF01163611). URL: <http://dx.doi.org/10.1007/BF01163611>.