

POROSITY, DIFFERENTIABILITY AND PANSU'S THEOREM

ANDREA PINAMONTI AND GARETH SPEIGHT

ABSTRACT. We use porosity to study differentiability of Lipschitz maps on Carnot groups. Our first result states that directional derivatives of a Lipschitz function act linearly outside a σ -porous set. The second result states that irregular points of a Lipschitz function form a σ -porous set. We use these observations to give a new proof of Pansu's theorem for Lipschitz maps from a general Carnot group to a Euclidean space.

1. INTRODUCTION

A Carnot group (Definition 2.1) is a connected and simply connected Lie group whose Lie algebra admits a stratification. Carnot groups have translations, dilations, Haar measure and points are connected by horizontal curves (Definition 2.2), which are used to define the Carnot-Carathéodory distance. With so much structure, the study of analysis and geometry in Carnot groups is an active and interesting research area [1, 2, 4, 8, 13, 19, 25, 27].

The geometry of Carnot groups is highly non-trivial. For instance, any Carnot group (except for Euclidean spaces themselves) contains no subset of positive measure that is bi-Lipschitz equivalent to a subset of a Euclidean space [24]. This follows from Pansu's theorem (Theorem 2.6), a generalization of Rademacher's theorem, which asserts that Lipschitz maps between Carnot groups are differentiable almost everywhere [20, 17]. Differentiability of maps between Carnot groups is defined like that of maps between Euclidean spaces, but Euclidean translations, dilations and distances are replaced by their analogues in Carnot groups.

Many interesting geometric and analytic problems have been studied in the context of Carnot groups. For example, a geometric notion of intrinsic Lipschitz function between subgroups of a general Carnot group was introduced in [11] to study rectifiable sets [12, 18] and minimal surfaces [5, 6, 26]. Moreover, Carnot groups have been applied to study degenerate equations, control theory and potential theory [4]. More recently, they were also considered in applied mathematics, such as mathematical finance, theoretical computer science and mathematical models for neurosciences [7].

Date: March 15, 2016.

2010 Mathematics Subject Classification. 28A75, 43A80, 49Q15, 53C17.

Key words and phrases. Pansu's theorem, Porous set, Carnot group.

In this paper, we generalize to Carnot groups two statements asserting that a set of exceptional points concerning directional derivatives is σ -porous, and use them to give a new proof of Pansu's theorem for Lipschitz maps from a general Carnot group to a Euclidean space.

A set in a metric space is (upper) porous (Definition 2.7) if each of its points sees nearby relatively large holes in the set on arbitrarily small scales. A set is σ -porous if it is a countable union of porous sets. Porous sets have applications to topics like cluster set theory and differentiability of Lipschitz or convex mappings. Note there are several other types of porosity with their own applications. Properties and applications of porous sets are thoroughly discussed in the survey articles [28, 29].

Stating that an exceptional set is σ -porous is useful because σ -porous sets in metric spaces are of first category, and σ -porous sets in Euclidean spaces have Lebesgue measure zero. Proving that a set is σ -porous usually gives a stronger result than showing it is of first category or has measure zero: any topologically complete metric space without isolated points contains a closed nowhere dense set which is not σ -porous, and \mathbb{R}^n contains a closed nowhere dense set of Lebesgue measure zero which is not σ -porous [28].

Porosity has been used to study differentiability of Lipschitz mappings, even very recently. Indeed, [15] gives a version of Rademacher's theorem for Fréchet differentiability of Lipschitz mappings on Banach spaces in which porous sets are Γ -null. Roughly, a set is Γ -null if it meets typical infinite dimensional C^1 surfaces in measure zero. Applications of porosity to study differentiability in infinite dimensional Banach spaces are thoroughly discussed in the recent book [16]. In the finite dimensional setting, porosity and construction of large directional derivatives were recently used in [21] to obtain the following result: for any $n > 1$, there exists a Lebesgue null set in \mathbb{R}^n containing a point of differentiability for every Lipschitz mapping from \mathbb{R}^n to \mathbb{R}^{n-1} . Directional derivatives also played a key role in the proof of the following result in a Carnot group: there exists a measure zero set N in the Heisenberg group \mathbb{H}^n such that every Lipschitz map $f: \mathbb{H}^n \rightarrow \mathbb{R}$ is Pansu differentiable at a point of N [22].

We now briefly describe the results of this paper.

For a Lipschitz function $f: \mathbb{R}^n \rightarrow \mathbb{R}$, directional derivatives need not act linearly: $f'(x, u+v)$ may not agree with $f'(x, u) + f'(x, v)$ (see Example 3.1 for a simple example). However, [23, Theorem 2] asserts that such an implication does hold at points x outside a σ porous (even σ -directionally porous) set depending on f , even for Lipschitz maps on separable Banach spaces. We prove a similar statement in Carnot groups: directional derivatives (in horizontal directions) for Lipschitz maps act linearly outside a σ -porous set (Theorem 3.8). Note that in Carnot groups it only makes sense to consider directional derivatives in horizontal derivatives, since the composition of a Lipschitz function with a non-horizontal curve may not be Lipschitz, and consequently may fail to be differentiable. To relate directional derivatives in the different directions and hence prove Theorem 3.8, we prove and apply

Lemma 3.5 which shows how to move in a direction $U + V$ by repeatedly moving in horizontal directions U and V . Such a statement is non-trivial, since U and V are left-invariant vector fields whose direction depends upon the point where they are evaluated. Roughly, the porous sets appearing in Theorem 3.8 are sets of points at which (on some scale and to some specified accuracy) a Lipschitz function is approximated by directional derivatives in directions U and V , but is not well approximated by the sum of the directional derivatives in the direction $U + V$.

If a Lipschitz map $f: \mathbb{R}^n \rightarrow \mathbb{R}$ has all directional derivatives at a point $x \in \mathbb{R}^n$ and directional derivatives at x act linearly, then by definition f is differentiable at x . In Carnot groups we observe that existence and linear action of only directional derivatives in horizontal directions does not imply Pansu differentiability (Example 4.1). To pass from directional derivatives to Pansu differentiability, we first give another application of porosity.

A point x is a regular point of a Lipschitz function f if whenever a directional derivative of f at x in some direction exists, then that same directional derivative also controls changes in f along parallel lines with the same direction close to x (Definition 4.2). Proposition 4.3 adapts [15, Proposition 3.3] to Carnot groups, showing that the set of points at which a Lipschitz function is not regular is σ -porous.

We next prove Theorem 4.6. This states that if $f: \mathbb{G} \rightarrow \mathbb{R}$ is Lipschitz and x is a regular point at which directional derivatives exist and act linearly, then f is Pansu differentiable at x . To prove this we use Lemma 4.5, which provides a relatively short horizontal path from x to any nearby point which is a concatenation of lines in directions of a basis of V_1 . We use regularity of the point x to estimate changes along these lines using the directional derivatives at x , then linear action of directional derivatives to show that the derivative is indeed a group linear map.

As a consequence of Theorem 3.8, Proposition 4.3 and Theorem 4.6, we obtain Corollary 4.8. This asserts that existence of directional derivatives implies Pansu differentiability outside a σ -porous set. Since it is not hard to show that directional derivatives of a Lipschitz function exist almost everywhere (Lemma 4.10) and porous sets have measure zero, we obtain a new proof of Pansu's Theorem (Corollary 4.11) for mappings from Carnot groups to Euclidean spaces.

2. PRELIMINARIES

We now briefly describe the main notions and results used in the paper; the reader can consult [4] for more details.

2.1. Carnot groups. Recall that a Lie group is a smooth manifold which is also a group for which multiplication and inversion are smooth. The Lie algebra associated to a Lie group is the space of left invariant vector fields equipped with the Lie bracket $[\cdot, \cdot]$ defined on smooth functions by

$$[X, Y](f) = X(Y(f)) - Y(X(f)).$$

We also denote the direct sum of vector spaces V and W by $V \oplus W$.

Definition 2.1. A simply connected finite dimensional Lie group \mathbb{G} is said to be a *Carnot group of step s* if its Lie algebra \mathcal{G} is *stratified of step s* , this means that there exist linear subspaces V_1, \dots, V_s of \mathcal{G} such that

$$\mathcal{G} = V_1 \oplus \dots \oplus V_s$$

with

$$[V_1, V_i] = V_{i+1} \text{ if } 1 \leq i \leq s-1, \text{ and } [V_1, V_s] = \{0\}.$$

Here $[V_1, V_i] := \text{span}\{[a, b] : a \in V_1, b \in V_i\}$.

It can be shown that Definition 2.1 implies $[V_i, V_j] \subset V_{i+j}$ if $i+j \leq r$ and $[V_i, V_j] = \{0\}$ if $i+j > r$ [4, Proposition 1.1.7].

Let $m_i := \dim(V_i)$, $h_i := m_1 + \dots + m_i$ for $1 \leq i \leq s$, and $h_0 := 0$. A basis X_1, \dots, X_n of \mathcal{G} is *adapted to the stratification* if $X_{h_{i-1}+1}, \dots, X_{h_i}$ is a basis of V_i for $1 \leq i \leq s$. We define $m := m_1 = \dim(V_1)$ and note $n = h_s$. The basis X_1, \dots, X_m of V_1 induces an inner product $\omega(\cdot, \cdot)$ on V_1 for which X_1, \dots, X_m is orthonormal. We denote by $\omega(\cdot)$ the norm on V_1 induced by this inner product.

We recall the exponential map $\exp: \mathcal{G} \rightarrow \mathbb{G}$ is defined by $\exp(X) = \gamma(1)$ where $\gamma: [0, 1] \rightarrow \mathbb{G}$ is the unique solution to $\gamma'(t) = X(\gamma(t))$ and $\gamma(0) = 0$. The exponential map is a diffeomorphism between \mathbb{G} and \mathcal{G} . Using the basis X_1, \dots, X_n to identify \mathcal{G} with \mathbb{R}^n , we can identify \mathbb{G} with \mathbb{R}^n by the correspondence:

$$\exp(a_1 X_1 + \dots + a_n X_n) \in \mathbb{G} \longleftrightarrow (x_1, \dots, x_n) \in \mathbb{R}^n.$$

Using this identification of \mathbb{G} with \mathbb{R}^n , [4, Corollary 1.3.19] states that if $h_{l-1} < j \leq h_l$ for some $1 \leq l \leq s$ then:

$$X_j(x) = \partial_j + \sum_{i>h_l}^n q_{i,j}(x) \partial_i \quad i = 1, \dots, m, \quad (2.1)$$

where $q_{i,j}$ are suitable homogeneous polynomials completely determined by the group law in \mathbb{G} . Using (2.1) we can identify $X_j(x)$ with the vector $e_j + \sum_{i>h_l}^n q_{i,j}(x) e_i$ where e_j is the j 'th element of the canonical basis of \mathbb{R}^n . Denote by $p: \mathbb{R}^n \rightarrow \mathbb{R}^m$ the projection onto the first m coordinates, given by $p(x) = (x_1, \dots, x_m)$. Then $p(X_j(x))$ is independent of $x \in \mathbb{G}$, so we can unambiguously define $p(X_j) = p(X_j(x)) = e_j$ for every $x \in \mathbb{G}$ and $j = 1, \dots, m$. We extend this definition by linearity to all V_1 , in particular $p(U(x)) = p(U(y))$ for every $x, y \in \mathbb{R}^n$, $U \in V_1$.

To compute the group law in coordinates, it is possible to define a mapping $\diamond: \mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$ for which (\mathcal{G}, \diamond) is a Lie group and $\exp: (\mathcal{G}, \diamond) \rightarrow (\mathbb{G}, \cdot)$ is a group isomorphism [4, Theorem 2.2.13], in particular:

$$\exp(X) \exp(Y) = \exp(X \diamond Y) \quad \text{for all } X, Y \in \mathcal{G}. \quad (2.2)$$

The Baker-Campbell-Hausdorff formula gives a formula for \diamond :

$$X \diamond Y = X + Y + \frac{1}{2}[X, Y] + \frac{1}{12}([X, [X, Y]] + [Y, [Y, X]]) + \dots, \quad (2.3)$$

where the higher order terms are nested commutators of X and Y .

Denoting points of \mathbb{G} by $(x_1, \dots, x_n) \in \mathbb{R}^n$, the *homogeneity* $d_i \in \mathbb{N}$ of the variable x_i is defined by

$$d_j = i \quad \text{whenever } h_{i-1} + 1 \leq j \leq h_i.$$

For any $\lambda > 0$, the *dilation* $\delta_\lambda: \mathbb{G} \rightarrow \mathbb{G}$, is defined in coordinates by

$$\delta_\lambda(x_1, \dots, x_n) = (\lambda^{d_1} x_1, \dots, \lambda^{d_n} x_n)$$

and satisfies $\delta_\lambda(xy) = \delta_\lambda(x)\delta_\lambda(y)$. Using the exponential map, dilations satisfying $\exp \circ \delta_\lambda = \delta_\lambda \circ \exp$ can be defined on \mathcal{G} . These satisfy $\delta_\lambda(E) = \lambda^i E$ if $E \in V_i$ for some $1 \leq i \leq s$.

A *Haar measure* on \mathbb{G} is a non-trivial Borel measure μ on \mathbb{G} satisfying $\mu(gE) = \mu(E)$ for any $g \in \mathbb{G}$ and Borel set $E \subset \mathbb{G}$. Such a measure is unique up to scaling by a positive constant, so sets of measure zero are defined without ambiguity. Identifying \mathbb{G} with \mathbb{R}^n , any Haar measure is simply a constant multiple of n dimensional Lebesgue measure \mathcal{L}^n .

2.2. Carnot-Carathéodory distance. Recall that a curve $\gamma: [a, b] \rightarrow \mathbb{R}^n$ is *absolutely continuous* if it is differentiable almost everywhere, $\gamma' \in L^1[a, b]$ and

$$\gamma(t_2) = \gamma(t_1) + \int_{t_1}^{t_2} \gamma'(t) dt$$

whenever $t_1, t_2 \in [a, b]$.

Definition 2.2. An absolutely continuous curve $\gamma: [a, b] \rightarrow \mathbb{G}$ is *horizontal* if there exist $u_1, \dots, u_m \in L^1[a, b]$ such that

$$\gamma'(t) = \sum_{j=1}^m u_j(t) X_j(\gamma(t))$$

for almost every $t \in [a, b]$. We define the horizontal length of such a curve γ by:

$$L(\gamma) = \int_a^b |u(t)| dt,$$

where $u = (u_1, \dots, u_m)$ and $|\cdot|$ denotes the Euclidean norm on \mathbb{R}^m .

The Chow-Rashevskii Theorem asserts that any two points of \mathbb{G} can be connected by horizontal curves [4, Theorem 9.1.3].

Definition 2.3. The *Carnot-Carathéodory distance* d on \mathbb{G} is defined by:

$$d(x, y) = \inf\{L(\gamma) : \gamma: [0, 1] \rightarrow \mathbb{G} \text{ horizontal joining } x \text{ to } y\}$$

for $x, y \in \mathbb{G}$.

It is well-known that the Carnot-Carathéodory distance satisfies the relations $d(zx, zy) = d(x, y)$ and $d(\delta_r(x), \delta_r(y)) = rd(x, y)$ for $x, y, z \in \mathbb{G}$ and $r > 0$. It induces on \mathbb{G} the same topology as the Euclidean distance but is not bi-Lipschitz equivalent to the Euclidean distance. It follows from the definition of the exponential map that $d(\exp(tE)) \leq |t|\omega(E)$ whenever $t \in \mathbb{R}$ and $E \in V_1$.

A *homogeneous norm* on \mathbb{G} is a continuous function $D: \mathbb{G} \rightarrow [0, \infty)$ such that $D(\delta_\lambda(x)) = \lambda D(x)$ for every $\lambda > 0$ and $x \in \mathbb{G}$, and $D(x) > 0$ if and only if $x \neq 0$. We will mostly use the homogeneous norm $d(x) := d(x, 0)$, but it is also useful to consider the homogeneous norm given by the explicit formula:

$$\|x\| := \left(\sum_{i=1}^s |x^i|^{2r!/i} \right)^{\frac{1}{2s!}} \quad (2.4)$$

where $x = (x^1, \dots, x^s) \in \mathbb{R}^m \times \dots \times \mathbb{R}^{m_s} = \mathbb{R}^n$ and $|x^i|$ denotes the Euclidean norm on \mathbb{R}^{m_i} . By [4, Proposition 5.1.4] there exists $c > 0$ such that

$$c^{-1}\|x\| \leq d(x) \leq c\|x\| \quad \text{for every } x \in \mathbb{G}. \quad (2.5)$$

2.3. Directional derivatives and Pansu differentiability. If $x \in \mathbb{G}$ and $E \in V_1$ is horizontal then the map $t \mapsto x \exp(tE)$ is Lipschitz. Consequently if $f: \mathbb{G} \rightarrow \mathbb{R}$ is Lipschitz then the composition $t \mapsto f(x \exp(tE))$ is a Lipschitz mapping from \mathbb{R} to itself, hence differentiable almost everywhere. Thus it makes sense to define directional derivatives of Lipschitz maps $f: \mathbb{G} \rightarrow \mathbb{R}$ in horizontal directions. Note, however, that if $E \in \mathcal{G} \setminus V_1$ then the composition $t \mapsto f(x \exp(tE))$ may not be Lipschitz. In this paper we only consider directional derivatives in horizontal directions, as in [22].

Definition 2.4. Suppose $f: \mathbb{G} \rightarrow \mathbb{R}$, $x \in \mathbb{G}$ and $E \in V_1$. We say that f is *differentiable at x in direction E* if the limit

$$Ef(x) = \lim_{t \rightarrow 0} \frac{f(x \exp(tE)) - f(x)}{t}$$

exists.

Suppose $f: \mathbb{G} \rightarrow \mathbb{R}$, $x \in \mathbb{G}$ and $X_j f(x)$ exists for every $1 \leq j \leq m$. Then we define the *horizontal gradient* of f at x by

$$\nabla_H f(x) := \sum_{i=1}^m (X_i f(x)) X_i(x),$$

which can be represented in coordinates by $(X_1 f(x), \dots, X_m f(x)) \in \mathbb{R}^m$.

Let $\tilde{\mathbb{G}}$ be a Carnot group with distance \tilde{d} and dilations $\tilde{\delta}_r$.

Definition 2.5. A map $L: \mathbb{G} \rightarrow \tilde{\mathbb{G}}$ is *group linear* if $L(xy) = L(x)L(y)$ and $L(\delta_r(x)) = \tilde{\delta}_r(L(x))$ whenever $x, y \in \mathbb{G}$ and $r > 0$.

A map $f: \mathbb{G} \rightarrow \tilde{\mathbb{G}}$ is *Pansu differentiable* at $x_0 \in \mathbb{G}$ if there exists a group linear map $L: \mathbb{G} \rightarrow \tilde{\mathbb{G}}$ such that

$$\lim_{h \rightarrow 0} \frac{\tilde{d}(f(x)^{-1}f(xh), L(h))}{d(h)} = 0.$$

If such a map L exists then it is unique and we denote it by df .

The following fundamental result generalizes Rademacher's theorem to Carnot groups and is due to Pansu [20].

Theorem 2.6 (Pansu's Theorem). *Let $f: \mathbb{G} \rightarrow \tilde{\mathbb{G}}$ be a Lipschitz map. Then f is Pansu differentiable almost everywhere.*

We fix a Carnot group \mathbb{G} and an adapted basis X_1, \dots, X_n of \mathcal{G} , with corresponding inner product norm ω , throughout this paper.

2.4. Porous sets. We now define porous sets and σ -porous sets; for more information see the extensive survey articles [28, 29]. Intuitively, a set is porous if every point of the set sees relatively large holes in the set on arbitrarily small scales. We denote by $B(x, r)$ the open ball of centre x and radius $r > 0$ in a metric space.

Definition 2.7. Let (M, ρ) be a metric space, $E \subset M$ and $a \in M$. We say that E is *porous at a* if there exist $\lambda > 0$ and a sequence $x_n \rightarrow a$ such that

$$B(x_n, \lambda\rho(a, x_n)) \cap E = \emptyset$$

for every $n \in \mathbb{N}$.

A set E is *porous* if it is porous at each point $a \in E$ with λ independent of a . A set is *σ -porous* if it is a countable union of porous sets.

Porous sets in Carnot groups have measure zero. This follows from the fact that Haar measure on Carnot groups is Ahlfors regular, hence doubling, so the Lebesgue differentiation theorem applies [14, Theorem 1.8]. If a porous set had positive measure then it would have a Lebesgue density point, which is impossible due to the presence of relatively large holes, which have relatively large measure, on arbitrarily small scales.

3. DIRECTIONAL DERIVATIVES ACTING LINEARLY

Even for a Lipschitz function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ with all directional derivatives $f'(x, v)$ at a point $x \in \mathbb{R}^2$, directional derivatives need not act linearly: $f'(x, a_1v_1 + a_2v_2) \neq a_1f'(x, v_1) + a_2f'(x, v_2)$ in general. Consequently, existence of directional derivatives alone does not suffice for differentiability. We illustrate this with a simple example.

Example 3.1. Define $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ by $f(x, y) = \min(x, y)$. Then f is a Lipschitz function with all directional derivatives at $(0, 0)$. However, f is not differentiable at $(0, 0)$, since:

$$f'((0, 0), (1, 1)) = 1 \neq 0 = \frac{\partial f}{\partial x}(0, 0) + \frac{\partial f}{\partial y}(0, 0).$$

If x lies outside a σ -porous set depending on f , then directional derivatives do act linearly: if $f'(x, v_1)$ and $f'(x, v_2)$ exist then $f'(x, a_1v_1 + a_2v_2)$ exists and is equal to $a_1f'(x, v_1) + a_2f'(x, v_2)$. Such a statement holds even for Lipschitz maps from a separable Banach space X to a Banach space Y [23, Theorem 2]. In this section we prove an analogue in Carnot groups, showing that directional derivatives act linearly outside a σ -porous set (Theorem 3.8).

3.1. Geometric lemmas. We now give several results describing the geometry of Carnot groups, essential for our study of porosity and differentiability. Before proving the first geometric lemma, we state an estimate for the norm of a commutator of group elements [10, Lemma 2.13]. Recall that \mathbb{G} is a Carnot group of step s , $h_s := n$ and the homogeneous norm $\|\cdot\|$ was defined in (2.4). The following lemma is stated using a different homogeneous norm in [10]. However the desired statement follows from [10, Lemma 2.13] because any two homogeneous norms differ by a constant multiplicative factor.

Lemma 3.2. *There is a constant $C > 0$ such that*

$$\|x^{-1}yx\| \leq C \left(\|y\| + \|x\|^{\frac{1}{s}} \|y\|^{\frac{s-1}{s}} + \|x\|^{\frac{s-1}{s}} \|y\|^{\frac{1}{s}} \right) \quad \text{for } x, y \in \mathbb{G}.$$

The first geometric lemma bounds the Carnot-Carathéodory distance between points obtained by flowing from two nearby points in the direction of the same horizontal vector field.

Lemma 3.3. *Suppose $\lambda \in (0, 1)$ and $t \in (-1, 1)$. Let $x, y \in \mathbb{G}$ satisfy $d(x, y) \leq \lambda|t|$ and $U \in V_1$. Then, there exists $C = C_1 > 0$ such that*

$$d(x \exp(tU), y \exp(tU)) \leq C\lambda^{1/s}|t| \max(1, \omega(U)).$$

Proof. Throughout the proof C will denote a positive constant depending only on \mathbb{G} and possibly different from line to line. Using equation (2.5), we can estimate with the equivalent homogeneous norm $\|\cdot\|$ instead of $d(\cdot)$. Without loss of generality assume $y = 0$ and $s \geq 2$. Indeed, if $s = 1$ then $\mathbb{G} \cong \mathbb{R}^n$ and

$$\|\exp(tU)^{-1}x \exp(tU)\| = \|x\| \leq C\lambda|t|$$

so the statement follows. Now suppose $s \geq 2$. Using Lemma 3.2 and the bound $\|x\| \leq C\lambda|t|$ we have:

$$\begin{aligned} \|\exp(tU)^{-1}x \exp(tU)\| &\leq C \left(\|x\| + \|x\|^{\frac{1}{s}} \|\exp(tU)\|^{\frac{s-1}{s}} + \|x\|^{\frac{s-1}{s}} \|\exp(tU)\|^{\frac{1}{s}} \right) \\ &= C \left(\|x\| + \|x\|^{\frac{1}{s}} |t|^{\frac{s-1}{s}} \omega(U)^{\frac{s-1}{s}} + \|x\|^{\frac{s-1}{s}} |t|^{\frac{1}{s}} \omega(U)^{\frac{1}{s}} \right) \\ &\leq C|t| \left(\lambda + \lambda^{\frac{1}{s}} \omega(U)^{\frac{s-1}{s}} + \lambda^{\frac{s-1}{s}} \omega(U)^{\frac{1}{s}} \right) \\ &\leq C|t| \lambda^{\frac{1}{s}} \left(1 + \omega(U)^{\frac{s-1}{s}} + \omega(U)^{\frac{1}{s}} \right), \end{aligned}$$

where in the last inequality we used $\lambda \in (0, 1)$ and $s \geq 2$. The conclusion easily follows. \square

We next recall [4, Lemma 19.1.4], as summarized in [4, page 717]. Keeping in mind (2.2), the lemma describes how to move in the direction of a Lie bracket of many directions, by moving repeatedly backwards and forwards in the individual directions. Recall that an iterated bracket is of length k with entries Y_1, \dots, Y_q if it is of the form $[Y_{j_1}, \dots, [Y_{j_{k-1}}, Y_{j_k}] \dots]$ for some multi-index $(j_1, \dots, j_k) \in \{1, \dots, q\}^k$.

Lemma 3.4. *Suppose $Y_1, \dots, Y_q \in \mathcal{G}$. Then we can write*

$$[Y_q, \dots, [Y_2, Y_1] \dots] = Y_{j_1} \diamond \dots \diamond Y_{j_{c(q)}} + R(Y_1, \dots, Y_q),$$

where $c(q)$ is an integer depending only on q , and

$$Y_{j_1}, \dots, Y_{j_{c(q)}} \in \{\pm Y_1, \dots, \pm Y_q\}.$$

The remainder term $R(Y_1, \dots, Y_q)$ is a linear combination of brackets of length $\geq q + 1$ of the vector fields Y_1, \dots, Y_q .

The next lemma shows how to move in a direction $U + V$ by repeatedly making increments in directions U and V . Such a statement will be useful when we want to relate directional derivatives in direction $U + V$ to directional derivatives in direction U and in direction V .

Lemma 3.5. *There is a constant $C = C_2 > 0$ for which the following holds. For any pair $U, V \in V_1$, there exist $U_1, \dots, U_N \in \{U, V\}$ and $\rho_1, \dots, \rho_N \in \mathbb{R}$ with $|\rho_i| \leq C$ and $N \leq C$ such that*

$$\exp(U + V) = \exp(\rho_1 U_1) \exp(\rho_2 U_2) \cdots \exp(\rho_N U_N).$$

Proof. We prove the lemma by induction on the step of \mathbb{G} . Throughout the proof, C denotes a constant depending only on \mathbb{G} , which may vary from line to line. If \mathbb{G} has step one the statement is clear: simply take $N = 2$, $U_1 = U$, $U_2 = V$ and use the equality:

$$\exp(U + V) = \exp(U) \exp(V),$$

which follows from the Baker-Campbell-Hausdorff formula (2.2) and (2.3).

Suppose $s > 1$ and the lemma holds for Carnot groups of step $s - 1$. Let \mathbb{G} be a Carnot group of step s with Lie algebra $\mathcal{G} = V_1 \oplus \dots \oplus V_s$ and adapted basis X_1, \dots, X_n of \mathcal{G} . Let $q = h_{s-1}$, then X_1, \dots, X_q is a basis of $V_1 \oplus \dots \oplus V_{s-1}$ and X_{q+1}, \dots, X_n is a basis of V_s .

Claim 3.6. *There exists a Carnot group \mathbb{H} of step $s - 1$ whose Lie algebra $\mathcal{H} = W_1 \oplus \dots \oplus W_{s-1}$ is spanned by an adapted basis Y_1, \dots, Y_q , with Lie bracket defined as follows:*

$$[Y_i, Y_j]_{\mathbb{H}} = \sum_{k=1}^q a_k Y_k \quad \text{if} \quad [X_i, X_j]_{\mathbb{G}} = \sum_{k=1}^n a_k X_k. \quad (3.1)$$

Proof. Fix a q -dimensional vector space \mathcal{H} and a basis Y_1, \dots, Y_q of \mathcal{H} . For $1 \leq i \leq s - 1$ define:

$$W_i = \text{span}\{Y_j : X_j \in V_i\}.$$

Then $\mathcal{H} = W_1 \oplus \cdots \oplus W_{s-1}$ and $W_{s-1} \neq 0$. Define $[\cdot, \cdot]_{\mathbb{H}}: \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ using (3.1) and bilinearity. The vector space \mathcal{H} equipped with $[\cdot, \cdot]_{\mathbb{H}}$ is a Lie algebra, so there exists a simply connected Lie group \mathbb{H} with Lie algebra \mathcal{H} [4, Theorem 2.2.14]. Using (3.1), one can check:

$$[W_1, W_i]_{\mathbb{H}} = W_{i+1} \text{ for } 1 \leq i \leq s-2, \text{ and } [W_1, W_{s-1}]_{\mathbb{H}} = \{0\}.$$

Hence \mathbb{H} is a Carnot group of step $s-1$. \square

Using (3.1), we can choose a Lie algebra homomorphism $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ satisfying $\varphi(X_i) = Y_i$ for $1 \leq i \leq q$ and $\varphi(X_i) = 0$ if $q < i \leq n$. The kernel of φ is V_s . We may choose a Lie group homomorphism $F: \mathbb{G} \rightarrow \mathbb{H}$ satisfying $dF = \varphi$ [4, Remark 2.1.60].

Suppose $U, V \in V_1$. Then $dF(U), dF(V) \in W_1$. Since \mathbb{H} has step $s-1$, we may apply our inductive hypothesis to \mathbb{H} and use linearity of dF to write:

$$\exp_{\mathbb{H}}(dF(U+V)) = \exp_{\mathbb{H}}(\rho_1 dF(U_1)) \cdots \exp_{\mathbb{H}}(\rho_N dF(U_N)), \quad (3.2)$$

with $U_i \in \{U, V\}$, $|\rho_i| \leq C$ and $N \leq C$. Here C is a constant depending only on \mathbb{H} , hence C is determined by \mathbb{G} . Since F is a Lie group homomorphism, the equality $\exp_{\mathbb{H}} \circ dF = F \circ \exp_{\mathbb{G}}$ holds. Hence (3.2) implies:

$$\begin{aligned} F(\exp_{\mathbb{G}}(U+V)) &= F(\exp_{\mathbb{G}}(\rho_1 U_1)) \cdots F(\exp_{\mathbb{G}}(\rho_N U_N)) \\ &= F(\exp_{\mathbb{G}}(\rho_1 U_1) \cdots \exp_{\mathbb{G}}(\rho_N U_N)). \end{aligned} \quad (3.3)$$

Claim 3.7. *There exists $Z \in V_s$ such that*

$$\exp_{\mathbb{G}}(U+V+Z) = \exp_{\mathbb{G}}(\rho_1 U_1) \cdots \exp_{\mathbb{G}}(\rho_N U_N). \quad (3.4)$$

Furthermore, $Z = \eta_1 Z_1 + \dots + \eta_p Z_p$ with $|\eta_i| \leq C$ and $p \leq C$ for a constant C determined by \mathbb{G} . Each term Z_i is a Lie bracket of length s of U and V .

Proof. The map $\exp_{\mathbb{G}}$ is a diffeomorphism, so necessarily there exists $Z \in \mathcal{G}$ such that (3.4) holds. Combining (3.3) and (3.4) gives:

$$F(\exp_{\mathbb{G}}(U+V+Z)) = F(\exp_{\mathbb{G}}(U+V)).$$

Hence:

$$\exp_{\mathbb{H}}(dF(U) + dF(V) + dF(Z)) = \exp_{\mathbb{H}}(dF(U) + dF(V)).$$

Since $\exp_{\mathbb{H}}$ is a diffeomorphism this implies

$$dF(U) + dF(V) + dF(Z) = dF(U) + dF(V),$$

so $dF(Z) = 0$. The kernel of dF is V_s , so $Z \in V_s$. By (3.4), the fact that $\exp_{\mathbb{G}}$ is invertible and (2.2) we obtain:

$$\begin{aligned} Z &= \exp_{\mathbb{G}}^{-1}(\exp_{\mathbb{G}}(\rho_1 U_1) \cdots \exp_{\mathbb{G}}(\rho_N U_N)) - U - V \\ &= (\rho_1 U_1) \diamond \cdots \diamond (\rho_N U_N) - U - V. \end{aligned} \quad (3.5)$$

Using the Baker-Campbell-Hausdorff formula (2.3) we can write:

$$Z = J_1 + \dots + J_s, \quad (3.6)$$

where each J_i is a linear combination of Lie brackets of length i of U and V . The number of terms in each linear combination is bounded by a constant

depending on \mathbb{G} . The scalars in these linear combinations are polynomials in the numbers ρ_i , with bounded degree and coefficients. Since the numbers ρ_i are uniformly bounded, the scalars are also bounded by a constant depending on \mathbb{G} . Note $Z \in V_s$ implies $J_i = 0$ for every $1 \leq i \leq s-1$. Hence $Z = J_s$ is of the desired form. \square

Any of the Z_i in Claim 3.7 can be written in the form:

$$[E_s, \dots [E_2, E_1] \dots]_{\mathbb{G}} \in V_s, \quad E_i \in \{U, V\}. \quad (3.7)$$

Given such a Lie bracket, use Lemma 3.4 to find $j_1, \dots, j_{c(s)}$ such that

$$[E_s, \dots [E_2, E_1] \dots]_{\mathbb{G}} = E_{j_1} \diamond \dots \diamond E_{j_{c(s)}} + R(E_1, \dots, E_s),$$

where $c(s)$ is an integer depending only on s , and

$$E_{j_1}, \dots, E_{j_{c(s)}} \in \{\pm U, \pm V\}.$$

The remainder term $R(E_1, \dots, E_s)$ is a linear combination of brackets of height $\geq s+1$ of the vector fields E_1, \dots, E_s . Since \mathbb{G} has step s , this implies $R = 0$. Consequently:

$$[E_s, \dots [E_2, E_1] \dots]_{\mathbb{G}} = E_{j_1} \diamond \dots \diamond E_{j_{c(s)}}.$$

Using (2.2), this implies:

$$\exp_{\mathbb{G}}([E_s, \dots [E_2, E_1] \dots]_{\mathbb{G}}) = \exp_{\mathbb{G}}(E_{j_1}) \cdots \exp_{\mathbb{G}}(E_{j_{c(s)}}), \quad (3.8)$$

where $c(s) \leq C$.

Claim 3.7 states that $Z = \eta_1 Z_1 + \dots + \eta_p Z_p$ with $|\eta_i| \leq C$, $p \leq C$ and each Z_i of the form in (3.7). Since $Z_i \in V_s$ for $1 \leq i \leq p$ we have $[Z_i, Z_j]_{\mathbb{G}} = 0$, so:

$$\exp_{\mathbb{G}}(Z) = \exp_{\mathbb{G}}(\eta_1 Z_1) \cdots \exp_{\mathbb{G}}(\eta_p Z_p).$$

Hence we may use (3.8) for each Z_i and combine the resulting expressions to write:

$$\exp_{\mathbb{G}}(Z) = \exp_{\mathbb{G}}(\delta_1 Q_1) \cdots \exp_{\mathbb{G}}(\delta_M Q_M), \quad (3.9)$$

with $M \leq C$, $|\delta_i| \leq C$ and $Q_i \in \{U, V\}$. Combining (3.4) and (3.9), we deduce:

$$\begin{aligned} & \exp_{\mathbb{G}}(\rho_1 U_1) \cdots \exp_{\mathbb{G}}(\rho_N U_N) \exp_{\mathbb{G}}(-\delta_M Q_M) \cdots \exp_{\mathbb{G}}(-\delta_1 Q_1) \\ &= \exp_{\mathbb{G}}(U + V + Z) \exp_{\mathbb{G}}(-Z) \\ &= \exp_{\mathbb{G}}(U + V). \end{aligned} \quad (3.10)$$

The second equality follows from the fact $Z \in V_s$ so $[U + V + Z, -Z]_{\mathbb{G}} = 0$ and the Baker-Campbell-Hausdorff formula (2.3) simplifies. Notice that $|\rho_i|, |\delta_i|, N, M \leq C$ and $U_i, Q_i \in \{U, V\}$, so (3.10) gives the required representation holds for $\exp(U + V)$. Hence the lemma holds for \mathbb{G} which has step s . This completes the inductive step, proving the lemma. \square

3.2. Geometry to differentiability. We now use Lemma 3.3 and Lemma 3.5 to show directional derivatives act linearly outside a σ -porous set.

Theorem 3.8. *Suppose $f: \mathbb{G} \rightarrow \mathbb{R}$ is Lipschitz. Then there is a σ -porous set $A \subset \mathbb{G}$ such that directional derivatives act linearly at every point $x \in \mathbb{G} \setminus A$, namely the following implication holds: if $E_1 f(x)$ and $E_2 f(x)$ exist for some $E_1, E_2 \in V_1$, then $(a_1 E_1 + a_2 E_2) f(x)$ exists and*

$$(a_1 E_1 + a_2 E_2) f(x) = a_1 E_1 f(x) + a_2 E_2 f(x)$$

for all $a_1, a_2 \in \mathbb{R}$.

It follows directly from the definition that if a directional derivative $E f(x)$ exists then $(sE) f(x)$ exists and is equal to $s(E f(x))$ for any $s \in \mathbb{R}$. Hence to prove Theorem 3.8 we may assume $a_1 = a_2 = 1$.

Recall the constant C_2 from Lemma 3.5. Given $U, V \in V_1 \setminus \{0\}$, $y, z \in \mathbb{R}$ and $\varepsilon, \delta > 0$, let $A(U, V, y, z, \varepsilon, \delta)$ be the set of $x \in \mathbb{G}$ such that for all $|t| < \delta$:

$$|f(x \exp(tU)) - f(x) - ty| \leq \varepsilon |t|, \quad (3.11)$$

$$|f(x \exp(tV)) - f(x) - tz| \leq \varepsilon |t|, \quad (3.12)$$

and there exist arbitrarily small t for which:

$$|f(x \exp(tU + tV)) - f(x) - t(y + z)| > 2\varepsilon C_2 |t|. \quad (3.13)$$

Fix $x \in A(U, V, y, z, \varepsilon, \delta)$. Use Lemma 3.5 to choose $U_1, \dots, U_N \in \{U, V\}$ and $\rho_1, \dots, \rho_N \in \mathbb{R}$ with $|\rho_i| \leq C_2$ and $N \leq C_2$ such that

$$\exp(U + V) = \exp(\rho_1 U_1) \cdots \exp(\rho_N U_N). \quad (3.14)$$

Choose

$$|t| < \min \left(\delta, \frac{1}{(1 + \omega(U) + \omega(V)) C_2} \right) \quad (3.15)$$

satisfying (3.13). Dilating both sides of (3.14) by the factor t gives:

$$\exp(tU + tV) = \exp(t\rho_1 U_1) \cdots \exp(t\rho_N U_N). \quad (3.16)$$

Let $x_0 = x$. For $1 \leq i \leq N$ let $x_i = x_{i-1} \exp(t\rho_i U_i)$, $y_i = y$ if $U_i = U$ and $y_i = z$ if $U_i = V$. Note that $x_N = x \exp(tU + tV)$ by (3.16).

Claim 3.9. *If U and V are linearly independent then the following statements hold:*

- (1) $\sum_{i=1}^N \rho_i y_i = y + z$,
- (2) there exists $0 \leq i \leq N - 1$ such that

$$|f(x_{i+1}) - f(x_i) - t\rho_{i+1} y_{i+1}| > 2\varepsilon |t|.$$

Proof. We first verify (1). Define:

$$I_U = \{i : U_i = U\} \text{ and } I_V = \{i : U_i = V\}.$$

Recall that $p: \mathbb{G} \rightarrow \mathbb{R}^m$ is the projection onto the first m coordinates. The equality (3.14) implies

$$\begin{aligned} p(U) + p(V) &= \rho_1 p(U_1) + \dots + \rho_N p(U_N) \\ &= \left(\sum_{i \in I_U} \rho_i \right) p(U) + \left(\sum_{i \in I_V} \rho_i \right) p(V). \end{aligned}$$

Since $U, V \in V_1$ and are linearly independent, $p(U)$ and $p(V)$ are linearly independent. Hence:

$$\sum_{i \in I_U} \rho_i = \sum_{i \in I_V} \rho_i = 1.$$

Consequently:

$$\begin{aligned} \sum_{i=1}^N \rho_i y_i &= \left(\sum_{i \in I_U} \rho_i \right) y + \left(\sum_{i \in I_V} \rho_i \right) z \\ &= y + z, \end{aligned}$$

which proves (1).

Next suppose (2) fails. Then:

$$|f(x_{i+1}) - f(x_i) - t\rho_{i+1}y_{i+1}| \leq 2\varepsilon|t| \text{ for } 0 \leq i \leq N-1. \quad (3.17)$$

We estimate using (1), (3.17) and $N \leq C_2$:

$$\begin{aligned} |f(x \exp(tU + tV)) - f(x) - t(y + z)| &= \left| f(x_N) - f(x) - t \sum_{i=1}^N \rho_i y_i \right| \\ &= \left| \sum_{i=1}^N (f(x_i) - f(x_{i-1}) - t\rho_i y_i) \right| \\ &\leq \sum_{i=1}^N |f(x_i) - f(x_{i-1}) - t\rho_i y_i| \\ &\leq 2\varepsilon C_2 |t|, \end{aligned}$$

This estimate violates our choice of t satisfying (3.13). This proves (2). \square

Claim 3.10. *The set $A(U, V, y, z, \varepsilon, \delta)$ is porous for each choice of the parameters $U, V, y, z, \varepsilon, \delta$.*

Proof. We may assume that U and V are linearly independent, otherwise $A(U, V, y, z, \varepsilon, \delta) = \emptyset$ and the claim is trivial. Use Claim 3.9(2) and the equality $x_{i+1} = x_i \exp(t\rho_{i+1}U_{i+1})$ to choose $0 \leq i \leq N-1$ satisfying:

$$|f(x_i \exp(t\rho_{i+1}U_{i+1})) - f(x_i) - t\rho_{i+1}y_{i+1}| > 2\varepsilon|t|. \quad (3.18)$$

Recall the constant $C_1 > 0$ defined in Lemma 3.3. Let

$$\begin{aligned} \Lambda &= 2\text{Lip}(f)C_1C_2 \max(\omega(U), \omega(V)), \\ \lambda &= \min \left(\frac{\varepsilon^s}{\Lambda^s}, \frac{\varepsilon}{2\text{Lip}(f)}, 1 \right), \end{aligned} \quad (3.19)$$

and

$$r = \min(\lambda\omega(U), \lambda\omega(V), \lambda)|t|. \quad (3.20)$$

Fix $q \in B(x_i, r)$. We will apply Lemma 3.3 with parameters:

- λ as defined in (3.19),
- t replaced by $t\rho_{i+1}\omega(U_{i+1})$,
- x replaced by x_i ,
- y replaced by q ,
- U replaced by $U_{i+1}/\omega(U_{i+1})$.

To see the hypotheses of Lemma 3.3 are satisfied, notice:

- (3.15) and the bound $|\rho_{i+1}| \leq C_2$ implies $|t\rho_{i+1}\omega(U_{i+1})| < 1$,
- $0 < \lambda < 1$ is clear from the definition in (3.19),
- $d(x_i, q) \leq \lambda|t|$ using $q \in B(x_i, r)$ and the definition of r in (3.20).

Also note $\omega(U_{i+1}/\omega(U_{i+1})) \leq 1$ and recall $|\rho_{i+1}| \leq C_2$. Applying Lemma 3.3 gives:

$$\begin{aligned} d(x_i \exp(t\rho_{i+1}U_{i+1}), q \exp(t\rho_{i+1}U_{i+1})) &\leq C_1\lambda^{1/s}|t\rho_{i+1}|\omega(U_{i+1}) \\ &\leq C_1C_2|t|\omega(U_{i+1})\lambda^{1/s} \\ &\leq C_1C_2|t|\omega(U_{i+1})\varepsilon/\Lambda \\ &\leq \varepsilon|t|/(2\text{Lip}(f)). \end{aligned}$$

Hence:

$$|f(x_i \exp(t\rho_{i+1}U_{i+1})) - f(q \exp(t\rho_{i+1}U_{i+1}))| \leq \varepsilon|t|/2. \quad (3.21)$$

Since $q \in B(x_i, r)$ we can also estimate:

$$|f(x_i) - f(q)| \leq \text{Lip}(f)r \leq \text{Lip}(f)\lambda|t| \leq \varepsilon|t|/2. \quad (3.22)$$

Applying (3.18) together with (3.21) and (3.22) gives:

$$|f(q \exp(t\rho_{i+1}U_{i+1})) - f(q) - t\rho_{i+1}y_{i+1}| > \varepsilon|t|. \quad (3.23)$$

Recall that either $U_{i+1} = U$ and $y_{i+1} = y$ or $U_{i+1} = V$ and $y_{i+1} = z$. Hence (3.23) shows that (3.11) or (3.12) fails with x replaced by q . Consequently $q \notin A(U, V, y, z, \varepsilon, \delta)$. Since q was an arbitrary member of $B(x_i, r)$, we deduce:

$$B(x_i, r) \cap A(U, V, y, z, \varepsilon, \delta) = \emptyset.$$

Since $N \leq C_2$ and $|\rho_i| \leq C_2$, we can estimate as follows:

$$\begin{aligned} d(x, x_i) &\leq d(x_0, x_1) + \dots + d(x_{i-1}, x_i) \\ &\leq \omega(t\rho_1U_1) + \dots + \omega(t\rho_NU_N) \\ &\leq |t||\rho_1|\omega(U_1) + \dots + |t||\rho_N|\omega(U_N) \\ &\leq C_2^2 \max(\omega(U), \omega(V))|t|. \end{aligned}$$

Since $r = \min(\lambda\omega(U), \lambda\omega(V), \lambda)|t|$, $B(x_i, r)$ is a relatively large hole in $A(U, V, y, z, \varepsilon, \delta)$ close to x . Since t can be chosen arbitrarily small, it follows that $A(U, V, y, z, \varepsilon, \delta)$ is porous at x . The point x was chosen arbitrarily from $A(U, V, y, z, \varepsilon, \delta)$, so the claim is proven. \square

Proof of Theorem 3.8. Fix a countable dense set $W \subset V_1$. Let A be the countable union of sets $A(U, V, y, z, \varepsilon, \delta)$, with $U, V \in W$ and $y, z, \varepsilon, \delta \in \mathbb{Q}$ such that $\varepsilon, \delta > 0$. Clearly A is σ -porous.

Suppose $x \in \mathbb{G} \setminus A$ and $E_1 f(x), E_2 f(x)$ exist. Choose rational $\varepsilon > 0$ and rational $\delta > 0$ such that for all $|t| < \delta$:

$$|f(x \exp(tE_1)) - f(x) - tE_1 f(x)| \leq \varepsilon|t|/2$$

and

$$|f(x \exp(tE_2)) - f(x) - tE_2 f(x)| \leq \varepsilon|t|/2.$$

Fix $y, z \in \mathbb{Q}$ such that $|y - E_1 f(x)| < \varepsilon/4$ and $|z - E_2 f(x)| < \varepsilon/4$. Choose $F_1, F_2 \in W$ such that

$$\begin{aligned} d(\exp(E_1), \exp(F_1)) &< \varepsilon/4 \text{Lip}(f), \\ d(\exp(E_2), \exp(F_2)) &< \varepsilon/4 \text{Lip}(f), \end{aligned}$$

and

$$d(\exp(E_1 + E_2), \exp(F_1 + F_2)) \leq \varepsilon.$$

Analogues of (3.11) and (3.12) hold with E_1 replaced by F_1 and E_2 replaced by F_2 . For the first one notice that for any $|t| < \delta$:

$$\begin{aligned} |f(x \exp tF_1) - f(x) - ty| &\leq |f(x \exp tE_1) - f(x) - tE_1 f(x)| \\ &\quad + |f(x \exp tE_1) - f(x \exp tF_1)| \\ &\quad + |ty - tE_1 f(x)| \\ &\leq \varepsilon|t|/2 + \text{Lip}(f)|t|d(\exp(E_1), \exp(F_1)) \\ &\quad + |t||y - E_1 f(x)| \\ &\leq \varepsilon|t|. \end{aligned}$$

The second estimate is similar. Since $x \in \mathbb{G} \setminus A$, necessarily $x \notin A(F_1, F_2, y, z, \varepsilon, \delta)$. However, (3.11) and (3.12) hold, so necessarily (3.13) fails. Hence, for all sufficiently small t ,

$$|f(x \exp(tF_1 + tF_2)) - f(x) - t(y + z)| \leq 2\varepsilon C_1 |t|.$$

This implies that for sufficiently small t :

$$\begin{aligned} &|f(x \exp(tE_1 + tE_2)) - f(x) - t(E_1 f(x) + E_2 f(x))| \\ &\leq |f(x \exp(tF_1 + tF_2)) - f(x) - t(y + z)| \\ &\quad + |t| \text{Lip}(f) d(\exp(E_1 + E_2), \exp(F_1 + F_2)) \\ &\quad + |t||E_1 f(x) - y| + |t||E_2 f(x) - z| \\ &\leq 2\varepsilon C_1 |t| + |t| \text{Lip}(f) \varepsilon + |t| \varepsilon / 2 \\ &\leq (2C_1 + \text{Lip}(f) + 1/2) \varepsilon |t|. \end{aligned}$$

This implies that $(E_1 + E_2)f(x)$ exists and is equal to $E_1 f(x) + E_2 f(x)$, which proves the theorem. \square

4. REGULARITY AND DIFFERENTIABILITY

For Lipschitz maps between Euclidean spaces, existence and linear action of directional derivatives implies differentiability (by definition). The following example shows that for a Lipschitz map $f: \mathbb{G} \rightarrow \mathbb{R}$, existence and linear action of directional derivatives (in only horizontal directions) does not suffice for Pansu differentiability.

Example 4.1. Let $\mathbb{H}^1 \equiv \mathbb{R}^3$ be the first Heisenberg group. The Heisenberg group is the simplest example of non trivial Carnot group of step 2. The group law has the form

$$(x, y, t)(x', y', t') = \left(x + x', y + y', t + t' + \frac{1}{2}(xy' - x'y) \right).$$

The homogeneous norm defined in (2.4) reads as

$$\|(x, y, t)\| = ((x^2 + y^2)^2 + t^2)^{1/4}, \quad (x, y, t) \in \mathbb{H}^1$$

and $\tilde{d}(a, b) = \|a^{-1}b\|$ defines a metric on \mathbb{H}^1 . Let

$$A = (\mathbb{R}^2 \times \{0\}) \cup (\{(0, 0)\} \times \mathbb{R}) \subset \mathbb{H}^1,$$

equipped with the restriction of the metric d on \mathbb{H}^1 . Define $f: A \rightarrow \mathbb{R}$ by $f(x, y, 0) = 0$ and $f(0, 0, t) = \sqrt{|t|}$. We claim that f is Lipschitz with respect to restriction of the metric \tilde{d} . Clearly, it suffices to prove that there exists $C > 0$ such that

$$|f(x, y, 0) - f(0, 0, t)| \leq C\tilde{d}((x, y, 0), (0, 0, t)) \quad \text{for all } x, y, t \in \mathbb{R} \quad (4.1)$$

and

$$|f(0, 0, t) - f(0, 0, s)| \leq C\tilde{d}((0, 0, t), (0, 0, s)) \quad \text{for all } t, s \in \mathbb{R} \quad (4.2)$$

Using the definition of the group law and \tilde{d} :

$$\tilde{d}((x, y, 0), (0, 0, t)) = \|(-x, -y, t)\| \geq c(\|(x, y)\|_{\mathbb{R}^2} + \sqrt{|t|})$$

and

$$|f(x, y, 0) - f(0, 0, t)| = \sqrt{|t|} \leq \frac{1}{c}\tilde{d}((x, y, 0), (0, 0, t))$$

so (4.1) follows. Moreover,

$$|f(0, 0, t) - f(0, 0, s)| = \left| \sqrt{|t|} - \sqrt{|s|} \right| \leq \sqrt{|t-s|} = \tilde{d}((0, 0, t), (0, 0, s)).$$

By the classical McShane extension theorem (see for example [14, Theorem 6.2]), f admits an extension to a Lipschitz function $f: \mathbb{H}^1 \rightarrow \mathbb{R}$, note that by (2.5), f is Lipschitz also with respect to the Carnot-Carathéodory distance d . Clearly $Ef(0, 0, 0) = 0$ for all $E \in V_1$, so f has all directional derivatives and they act linearly at $(0, 0, 0)$. Indeed, recalling that $\exp(tE) \in \mathbb{R}^2 \times \{0\}$, we have:

$$Ef(0, 0, 0) = \lim_{t \rightarrow 0} \frac{f(\exp(tE)) - f(0, 0, 0)}{t} = 0.$$

However, f is not Pansu differentiable at $(0, 0, 0)$. Indeed:

$$\lim_{t \rightarrow 0} \frac{|f(0, 0, t^2) - f(0, 0, 0)|}{d((0, 0, t^2), (0, 0, 0))} = \frac{|t|}{|t|} = 1 > 0.$$

In this section we adapt [15] to define regular points of Lipschitz functions on Carnot groups and show that the set of irregular points of a Lipschitz function is σ -porous (Proposition 4.3). We then show that outside another σ -porous set, existence of directional derivatives (in horizontal directions) implies Pansu differentiability (Theorem 4.6 and Corollary 4.8). This then yields a new proof of Pansu's theorem (Corollary 4.11).

4.1. Regular points. We adapt [15, Definition 3.1] from Banach spaces to define regular points for Lipschitz maps on Carnot groups. Intuitively, x is a regular point of f if whenever $E \in V_1$ and $Ef(x)$ exists, then $Ef(x)$ controls changes in f along all lines close to x in direction E .

Definition 4.2. Suppose $f: \mathbb{G} \rightarrow \mathbb{R}$. We say that $x \in \mathbb{G}$ is a *regular point* of f if for every $E \in V_1$ for which $Ef(x)$ exists,

$$\lim_{t \rightarrow 0} \frac{f(x\delta_t(u) \exp(tE)) - f(x\delta_t(u))}{t} = Ef(x)$$

uniformly for $d(u) \leq 1$. A point is *irregular* if it is not regular.

We adapt [15, Proposition 3.3] from Banach spaces to show that irregular points of a Lipschitz function form a σ -porous set.

Proposition 4.3. *Let $f: \mathbb{G} \rightarrow \mathbb{R}$ be Lipschitz. Then the set of irregular points of f is σ -porous.*

Proof. Let \mathcal{J} be a countable dense subset of V_1 . For $p, q \in \mathbb{N}$, $E \in \mathcal{J}$ and $w \in \mathbb{Q}$, let $A_{p,q,E,w}$ be the set of $x \in \mathbb{G}$ such that

$$|f(x \exp tE) - f(x) - tw| \leq |t|/p \text{ for } |t| < 1/q \quad (4.3)$$

and

$$\limsup_{t \rightarrow 0} \sup_{d(u) \leq 1} \left| \frac{f(x\delta_t(u) \exp(tE)) - f(x\delta_t(u))}{t} - w \right| > 2/p.$$

Fix p, q, E, w and $x \in A_{p,q,E,w}$. Then there are arbitrarily small $|t| < 1/q$ such that for some $u \in \mathbb{G}$ with $d(u) \leq 1$, depending on t ,

$$|f(x\delta_t(u) \exp(tE)) - f(x\delta_t(u)) - tw| > 2|t|/p. \quad (4.4)$$

Let

$$\lambda = \min \left(\frac{1}{(2C_1 p \text{Lip}(f)(1 + \omega(E)))^s}, \frac{1}{2p \text{Lip}(f)}, 1 \right),$$

where C_1 is the constant from Lemma 3.3. Fix $|t| < 1/q$ and $u \in \mathbb{G}$ with $d(u) \leq 1$ satisfying (4.4). Suppose $y \in B(x\delta_t(u), \lambda|t|)$. We may apply

Lemma 3.3, with U replaced by E and x replaced by $x\delta_t(u)$, to obtain:

$$\begin{aligned} |f(y \exp(tE)) - f(x\delta_t(u) \exp(tE))| &\leq \text{Lip}(f)d(y \exp(tE), x\delta_t(u) \exp(tE)) \\ &\leq \text{Lip}(f)C_1\lambda^{1/s}|t| \max(1, \omega(E)) \\ &\leq |t|/2p. \end{aligned}$$

Hence, using also (4.4) and the assumption $d(y, x\delta_t(u)) < \lambda|t|$,

$$\begin{aligned} |f(y \exp(tE)) - f(y) - tw| &\geq |f(x\delta_t(u) \exp(tE)) - f(x\delta_t(u)) - tw| \\ &\quad - |f(y \exp(tE)) - f(x\delta_t(u) \exp(tE))| \\ &\quad - |f(x\delta_t(u)) - f(y)| \\ &> |t|/p. \end{aligned}$$

Since $|t| < 1/q$ the previous estimate contradicts (4.3), so $y \notin A_{p,q,E,w}$. To summarise:

$$d(x, x\delta_t(u)) = d(\delta_t(u)) \leq |t|$$

and we have shown:

$$B(x\delta_t(u), \lambda|t|) \cap A_{p,q,E,w} = \emptyset.$$

Since $|t|$ could be chosen arbitrarily small, this shows $A_{p,q,E,w}$ is porous. Every irregular point of f belongs to one of the countable collection of porous sets $A_{p,q,E,w}$, where $p, q \in \mathbb{N}$, $E \in \mathcal{J}$ and $w \in \mathbb{Q}$. Hence the set of irregular points of f is σ -porous, as desired. \square

4.2. Differentiability and Pansu's theorem. To pass from directional derivatives to Pansu's theorem, we use a lemma showing how to join arbitrary points by following directions from our chosen basis X_1, \dots, X_m of V_1 . To prove our desired statement we first quote the following lemma [4, Theorem 19.2.1].

Lemma 4.4. *Let $\mathcal{Z} = \{Z_1, \dots, Z_m\}$ be a basis for V_1 and fix a homogeneous norm ρ on \mathbb{G} . Then there exist constants $M \in \mathbb{N}$ (depending only on \mathbb{G}) and $c_0 > 0$ (depending on \mathbb{G} , ρ and \mathcal{Z}) for which the following holds.*

For every $x \in \mathbb{G}$, there exist $x_1, \dots, x_M \in \exp(V_1)$ with the following properties:

- $x = x_1 \cdots x_M$,
- $\rho(x_j) \leq c_0\rho(x)$ for all $j = 1, \dots, M$,
- for every $j = 1, \dots, M$, there exist $t_j \in \mathbb{R}$ and $i_j \in \{1, \dots, m\}$ such that $x_j = \exp(t_j Z_{i_j})$.

Lemma 4.5. *There are constants $M \in \mathbb{N}$ and $Q > 0$ such that the following holds for every point $h \in \mathbb{G}$. For $1 \leq j \leq M$, there exist $t_j \geq 0$ and $E_j \in \{\pm X_1, \dots, \pm X_m\}$ such that:*

- (1) $\sum_{j=1}^M t_j \leq Qd(h)$,
- (2) $h = \exp(t_1 E_1) \cdots \exp(t_M E_M)$.

Proof. Apply Lemma 4.4 with the basis $\mathcal{Z} = \{X_1, \dots, X_m\}$ of V_1 and homogeneous norm $x \mapsto d(x)$ on \mathbb{G} . We obtain $M \in \mathbb{N}$ such that for every $h \in \mathbb{G}$, there exist $t_j \in \mathbb{R}$ and $i_j \in \{1, \dots, m\}$ for $1 \leq j \leq M$, such that

$$h = \exp(t_1 X_{i_1}) \cdots \exp(t_M X_{i_M}), \quad (4.5)$$

and

$$d(\exp t_j X_{i_j}) \leq c_0 d(h). \quad (4.6)$$

Since $\omega(X_{i_j}) = 1$ for every j , (4.6) implies $|t_j| \leq c_0 d(h)$ for $1 \leq j \leq M$. Hence:

$$\sum_{j=1}^M |t_j| \leq M c_0 d(h). \quad (4.7)$$

For each $1 \leq j \leq M$ choose $E_j = X_{i_j}$ if $t_j \geq 0$, or choose $E_j = -X_{i_j}$ and replace t_j by $-t_j$ if $t_j < 0$. After such a replacement we have $t_j \geq 0$ for every j , (4.5) still holds giving the desired equality (2), and (4.7) gives (1). \square

We now prove a pointwise differentiability result, whose hypotheses combine the various conditions we have investigated so far.

Theorem 4.6. *Suppose $f: \mathbb{G} \rightarrow \mathbb{R}$ is Lipschitz and $x \in \mathbb{G}$ has the following properties:*

- $X_1 f(x), \dots, X_m f(x)$ exist,
- directional derivatives act linearly at x ,
- x is a regular point of f .

Then f is Pansu differentiable at the point x with group linear derivative $L(\cdot) = \langle p(\cdot), \nabla_H f(x) \rangle$, where $p: \mathbb{G} \rightarrow \mathbb{R}^m$ is projection onto the first m coordinates.

Proof. First we remark that $h \mapsto \langle p(h), \nabla_H f(x) \rangle$ is group linear. Indeed, the group operation is Euclidean in the first m coordinates so:

$$\begin{aligned} \langle p(h_1 h_2), \nabla_H f(x) \rangle &= \langle p(h_1) + p(h_2), \nabla_H f(x) \rangle \\ &= \langle p(h_1), \nabla_H f(x) \rangle + \langle p(h_2), \nabla_H f(x) \rangle, \end{aligned}$$

and

$$\langle p(\delta_r(h)), \nabla_H f(x) \rangle = \langle r p(h), \nabla_H f(x) \rangle = r \langle p(h), \nabla_H f(x) \rangle$$

for any $h_1, h_2 \in \mathbb{G}$ and $r > 0$.

We may assume $x = 0$. Fix M and Q as in Lemma 4.5 and let $\varepsilon > 0$. Since $X_1 f(x), \dots, X_m f(x)$ exist and directional derivatives act linearly at x , necessarily $E f(x)$ exists for every $E \in V_1$. Since 0 is a regular point of f , for each $E \in V_1$ we can find $\delta > 0$ such that if $0 < |t| < \delta$ and $d(u) \leq 1$ then:

$$|f(\delta_t(u) \exp(tE)) - f(\delta_t(u)) - t E f(0)| \leq \varepsilon |t|. \quad (4.8)$$

Since $\mathcal{J} := \{E \in V_1 : \omega(E) \leq 1\}$ is compact, we may additionally choose $\delta > 0$ so that (4.8) holds uniformly if $0 < |t| < \delta$, $d(u) \leq 1$ and $E \in \mathcal{J}$.

Fix $h \in \mathbb{G} \setminus \{0\}$ with $d(h) \leq \delta/Q$. Use Lemma 4.5 to choose $t_i \geq 0$ and $E_i \in \{\pm X_1, \dots, \pm X_m\}$ for $1 \leq i \leq M$ such that

$$\sum_{j=1}^M t_j \leq Qd(h) \quad (4.9)$$

and

$$h = \exp(t_1 E_1) \cdots \exp(t_M E_M). \quad (4.10)$$

Let $x_1 = 0$ and $x_{i+1} = x_i \exp(t_i E_i)$ for $1 \leq i \leq M$. Notice $x_{M+1} = h$ follows from (4.10).

Claim 4.7. *For each $1 \leq i \leq M$:*

$$|f(x_{i+1}) - f(x_i) - t_i E_i f(0)| \leq Q\varepsilon d(h).$$

Proof. Fix $1 \leq i \leq M$ and let $t = Qd(h)$. We check that there exist $u \in \mathbb{G}$ with $d(u) \leq 1$ and $E \in V_1$ with $\omega(E) \leq 1$ such that $x_i = \delta_t(u)$ and $t_i E_i = tE$.

We begin by solving $x_i = \delta_t(u)$. Notice

$$x_i = \exp(t_1 E_1) \cdots \exp(t_{i-1} E_{i-1}).$$

Using the definition of x_i , $\omega(E_j) \leq 1$ and (4.9), we can estimate $d(x_i)$ as follows:

$$\begin{aligned} d(x_i) &\leq \sum_{j=1}^{i-1} d(x_j, x_{j+1}) = \sum_{j=1}^{i-1} d(\exp(t_j E_j)) \\ &\leq \sum_{j=1}^{i-1} t_j \omega(E_j) \\ &\leq Qd(h), \end{aligned}$$

Since $t = Qd(h)$, we can choose $u \in \mathbb{G}$ with $d(u) \leq 1$ such that $x_i = \delta_t(u)$.

Since $h \neq 0$, we have $t \neq 0$. Let $E = (t_i/t)E_i$. Since $\omega(E_i) = 1$ and (4.9) implies $t_i \leq Qd(h)$, we have

$$\omega(E) = t_i/t = t_i/Qd(h) \leq 1.$$

Hence $E \in V_1$ solves $tE = t_i E_i$ with $\omega(E) \leq 1$ as desired.

Since:

- $x_{i+1} = x_i \exp(t_i E_i)$,
- $x_i = \delta_t(u)$ and $t_i E_i = tE$,
- $t = Qd(h) < \delta$,

we may apply inequality (4.8) to obtain:

$$|f(x_{i+1}) - f(x_i) - t_i E_i f(0)| \leq Q\varepsilon d(h).$$

This proves the claim. □

The group operation in the first m coordinates is Euclidean, (4.10) states $\exp(t_1 E_1) \cdots \exp(t_M E_M) = h$, and $\exp(\theta E) = \theta E(0)$ whenever $\theta \in \mathbb{R}$ and $E \in V_1$. Hence:

$$(t_1 E_1(0)) \cdots (t_M E_M(0)) = h.$$

Since $p(X_i(0))$ is the i 'th standard basis vector of \mathbb{R}^m in coordinates, we deduce:

$$p \left(\sum_{i=1}^M t_i E_i(0) \right) = p \left(\sum_{i=1}^m h_i X_i(0) \right).$$

Hence $\sum_{i=1}^M t_i E_i$ and $\sum_{i=1}^m h_i X_i$ are two horizontal vectors whose projections in \mathbb{R}^m are equal at 0. Consequently:

$$\sum_{i=1}^M t_i E_i = \sum_{i=1}^m h_i X_i.$$

Since directional derivatives act linearly at 0, we deduce:

$$\sum_{i=1}^M t_i E_i f(0) = \sum_{i=1}^m h_i X_i f(0).$$

We then estimate as follows:

$$\begin{aligned} |f(h) - f(0) - \langle p(h), \nabla_H f(0) \rangle| &= \left| \sum_{i=1}^M (f(x_{i+1}) - f(x_i)) - \sum_{i=1}^m h_i X_i f(0) \right| \\ &= \left| \sum_{i=1}^M (f(x_{i+1}) - f(x_i)) - \sum_{i=1}^M t_i E_i f(0) \right| \\ &\leq \sum_{i=1}^M |f(x_{i+1}) - f(x_i) - t_i E_i f(0)| \\ &\leq MQ\varepsilon d(h). \end{aligned}$$

Hence for $d(h) \leq \delta/Q$ we have:

$$|f(h) - f(0) - \langle p(h), \nabla_H f(0) \rangle| \leq MQ\varepsilon d(h).$$

This proves that f is Pansu differentiable at x with the desired derivative. \square

We can combine Theorem 4.6 with Theorem 3.8 and Proposition 4.3 to obtain the following application of porous sets.

Corollary 4.8. *Let $f: \mathbb{G} \rightarrow \mathbb{R}$ be Lipschitz. Then there exists a σ -porous set P for which the following implication holds at all points $x \notin P$: if $X_i f(x)$ exists for every $1 \leq i \leq m$, then f is Pansu differentiable at x .*

Proof. Use Theorem 3.8 to choose a σ -porous set A , outside which directional derivatives act linearly. Let B be the set of irregular points of f defined in Definition 4.2, which Proposition 4.3 states is σ -porous. Clearly the set $A \cup B$ is σ -porous. Suppose $X_i f(x)$ exists for all $1 \leq i \leq m$ and

$x \notin A \cup B$. Then Theorem 4.6 asserts that f is Pansu differentiable at x , proving the desired implication. \square

We will use existence of directional derivatives and Corollary 4.8 to prove Pansu's theorem. To construct many directional derivatives we first need the following well-known fact, for example see [17, Lemma 3.3].

Lemma 4.9. *Let Z_1, Z_2 be two subspaces of \mathcal{G} with $Z_1 \oplus Z_2 = \mathcal{G}$ and $\dim(Z_1) = 1$. Then there are open neighborhoods of the origin $\Omega_1 \subset Z_1$, $\Omega_2 \subset Z_2$ and an open neighborhood of the identity $U \subset \mathbb{G}$ such that the map*

$$\phi: \Omega_2 \times \Omega_1 \rightarrow U$$

defined by

$$\phi(a, z) = \exp(a) \exp(z)$$

is a diffeomorphism.

Lemma 4.10. *Suppose $f: \mathbb{G} \rightarrow \mathbb{R}$ is Lipschitz and $E \in V_1$. Then the directional derivative $Ef(x)$ exists for almost every $x \in \mathbb{G}$.*

Proof. Let $Z_1 \subset \mathcal{G}$ be the subspace spanned by E and let Z_2 be the subspace of \mathcal{G} for which $\mathcal{G} = Z_1 \oplus Z_2$. Let Ω_1, Ω_2, U and ϕ be given by Lemma 4.9. Choose $\delta > 0$ so that if $|t| < \delta$ then $tE \in \Omega_1$. Notice that all these quantities are independent of the map f .

For each fixed $a \in \Omega_2$, $F_a: (-\delta, \delta) \rightarrow \mathbb{R}$ defined by $F_a(t) = f(\phi(a, tE))$ is a composition of Lipschitz functions, hence Lipschitz and therefore differentiable almost everywhere. Note $\exp((t+h)E) = \exp(tE) \exp(hE)$ follows from (2.3) and $[E, E] = 0$. Hence for $t \in (-\delta, \delta)$:

$$\begin{aligned} \frac{|F_a(t+h) - F_a(t)|}{h} &= \frac{|f(\phi(a, (t+h)E)) - f(\phi(a, tE))|}{h} \\ &= \frac{|f(\phi(a, tE) \exp(hE)) - f(\phi(a, tE))|}{h}. \end{aligned}$$

Using Definition 2.4 and existence of $F'_a(t)$ for almost every $t \in (-\delta, \delta)$, we deduce that $Ef(\phi(a, tE))$ exists and is equal to $F'_a(t)$ for almost every $t \in (-\delta, \delta)$.

Using the basis X_1, \dots, X_n of \mathcal{G} , \mathcal{G} is identified with \mathbb{R}^n so equipped with Lebesgue measure. Viewing ϕ as a map between subsets of \mathbb{R}^n , Fubini's theorem implies that $Ef(\phi(Z))$ exists for almost every $Z \in \Omega_2 \times \Omega_1$. Since ϕ is a diffeomorphism, it preserves sets of measure zero. Hence $Ef(x)$ exists for almost every $x \in \phi(\Omega_2 \times \Omega_1) = U$. This holds for every Lipschitz map f , with the set U independent of f .

For any $R > 0$, let $g_R(x) = f(\delta_R(x))$. It is easy to show that $Eg_R(x)$ exists if and only if $Ef(\delta_R(x))$ exists, in which case $Eg_R(x) = R(Ef(\delta_R(x)))$. Since dilations preserve sets of measure zero, $Ef(x)$ exists if and only if $Eg_R(\delta_{\frac{1}{R}}(x))$ exists. Suppose f is a Lipschitz function for which $Ef(x)$ fails to exist at points x in a set of positive measure. Then we can find a compact set K of positive measure such that $Ef(x)$ does not exist for any $x \in K$.

Now fix $R > 0$ such that $\delta_{\frac{1}{R}}(K) \subset U$. Then g_R is a Lipschitz function and $Eg_R(y)$ does not exist for points $y \in \delta_{\frac{1}{R}}(K) \subset U$ and $\delta_{\frac{1}{R}}(K)$ has positive measure. This contradicts what we showed above with \bar{f} replaced by g_R . Hence $Ef(x)$ exists for almost every $x \in \mathbb{G}$, for any Lipschitz function f . \square

We now use Lemma 4.10 and Corollary 4.8 to obtain Pansu's theorem for mappings from \mathbb{G} to a Euclidean space.

Corollary 4.11 (Pansu's Theorem). *Suppose $f: \mathbb{G} \rightarrow \mathbb{R}^N$ is Lipschitz. Then f is Pansu differentiable almost everywhere.*

Proof. First assume $N = 1$. Using Lemma 4.10, we can find a measure zero set $A \subset \mathbb{G}$ such that $X_i f(x)$ exists for $1 \leq i \leq m$ and $x \in \mathbb{G} \setminus A$. Using Corollary 4.8, choose a σ -porous set P such that $x \notin P$ and existence of $X_i f(x)$ for $1 \leq i \leq m$ together imply that f is Pansu differentiable at x . Since porous sets have measure zero, the set $A \cup P$ has measure zero. The function f is Pansu differentiable outside $A \cup P$, consequently Pansu differentiable almost everywhere.

Now assume $N > 1$ and let $f = (f_1, \dots, f_N): \mathbb{G} \rightarrow \mathbb{R}^N$ be Lipschitz. Clearly $f_i: \mathbb{G} \rightarrow \mathbb{R}$ is Lipschitz for all $1 \leq i \leq N$, therefore there exists a null set $B \subset \mathbb{G}$ so that every component f_i is Pansu differentiable in $\mathbb{G} \setminus B$. Let $L: \mathbb{G} \rightarrow \mathbb{R}^N$ be the group linear map defined by $L = (L_1, \dots, L_N)$ where L_i is the Pansu derivative of f_i . For every $x \in \mathbb{G} \setminus B$ we have:

$$\lim_{h \rightarrow 0} \frac{|f(xh) - f(x) - L(h)|}{d(h)} = \left(\sum_{i=1}^N \lim_{h \rightarrow 0} \frac{(f_i(xh) - f_i(x) - L_i(h))^2}{d(h)^2} \right)^{\frac{1}{2}} = 0.$$

Hence f is Pansu differentiable almost everywhere. \square

REFERENCES

- [1] Agrachev, A., Barilari, D., Boscain, U.: *Introduction to Riemannian and sub-Riemannian geometry (from hamiltonian viewpoint)*, notes available at <http://webusers.imj-prg.fr/~davide.barilari/>.
- [2] Bellaïche, A.: *The tangent space in sub-Riemannian geometry*, Journal of Mathematical Sciences, 83(4) (1994), 461–476.
- [3] Bate, D., Speight, G.: *Differentiability, porosity and doubling in metric measure spaces*, Proceedings of the American Mathematical Society 141 (2013), 971–985.
- [4] Bonfiglioli, A., Lanconelli, E., Uguzzoni, F.: *Stratified Lie groups and potential theory for their sub-laplacians*, Springer Monographs in Mathematics 26, New York, Springer-Verlag (2007).
- [5] Citti, C., Manfredini, M., Pinamonti, A., Serra Cassano, F.: *Smooth approximation for intrinsic Lipschitz functions in the Heisenberg group*, Calc. Var. Partial Differential Equations 49(3–4) (2014), 1279–1308.

- [6] Citti, C., Manfredini, M., Pinamonti, A., Serra Cassano, F.: *Poincaré-type inequality for Lipschitz continuous vector fields*, J. Math. Pures Appl. (9) 105(3) (2016), 265–292.
- [7] Sarti, A., Citti, G., Petitot, J.: *The symplectic structure of the primary visual cortex*, Biol. Cybernet. 98(1) (2008), 33–48.
- [8] Capogna, L., Danielli, D., Pauls, S., Tyson, J.: *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*, Birkhauser, Progress in Mathematics, 259 (2007).
- [9] Folland, G.B., Stein, E.: *Hardy spaces on homogeneous groups*, Princeton University Press (1982).
- [10] Franchi, B., Serapioni, R.: *Intrinsic Lipschitz graphs within Carnot groups*, Journal of Geometric Analysis, DOI: 10.1007/s12220-015-9615-5.
- [11] Franchi, B., Serapioni, R., Serra Cassano, F.: *Differentiability of intrinsic Lipschitz functions within Heisenberg groups*, J. Geom. Anal. 21(4) (2011), 1044–1084.
- [12] Franchi, B., Serapioni, R., Serra Cassano, F.: *Rectifiability and perimeter in the Heisenberg group*, Math. Ann. 321 (2001), 479–531.
- [13] Gromov, M.: *Carnot-Carathéodory spaces seen from within*, Progress in Mathematics, 144 (1996), 79–323.
- [14] Heinonen, J.: *Lectures on analysis on metric spaces*, Universitext. Springer-Verlag, New York (2001).
- [15] Lindenstrauss, J., Preiss, D.: *On Fréchet differentiability of Lipschitz maps between Banach spaces*, Annals of Mathematics 157 (2003), 257–288.
- [16] Lindenstrauss, J., Preiss, D., Tiser, J.: *Fréchet differentiability of Lipschitz functions and porous sets in Banach spaces*, Annals of Mathematics Studies 179, Princeton University Press (2012).
- [17] Magnani, V.: *Differentiability and area formula on stratified Lie groups*, Houston J. Math. 27(2) (2001), 297–323.
- [18] Magnani, V.: *Towards differential calculus in stratified groups*, J. Aust. Math. Soc. 95(1) (2013), 76–128.
- [19] Montgomery, R.: *A tour of subriemannian geometries, their geodesics and applications*, American Mathematical Society, Mathematical Surveys and Monographs, 91 (2006).
- [20] Pansu, P.: *Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un*, Annals of Mathematics 129(1) (1989), 1–60.
- [21] Preiss, D., Speight, G.: *Differentiability of Lipschitz functions in Lebesgue null sets*, Inventiones Mathematicae 199(2) (2015), 517–559.
- [22] Pinamonti, A., Speight, G.: *A measure zero universal differentiability set in the Heisenberg group*, preprint available at arXiv:1505.07986.

- [23] Preiss, D., Zajicek, L.: *Directional derivatives of Lipschitz functions*, Israel Journal of Mathematics 125 (2001), 1–27.
- [24] Semmes, S.: *On the nonexistence of bi-Lipschitz parameterizations and geometric problems about A_∞ -weights*, Revista Matemática Iberoamericana, 12(2) (1996), 337–410.
- [25] Serra Cassano, F.: *Some topics of geometric measure theory in Carnot groups*, Volume I, EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, to appear.
- [26] Serra Cassano, F., Vittone, D.: *Graphs of bounded variation, existence and local boundedness of non-parametric minimal surfaces in the Heisenberg group*, Adv. Calc. Var. 7(4) (2014), 409–492.
- [27] Vittone, D.: *The regularity problem for sub-Riemannian geodesics*, CRM Series, 17, Ed. Norm., Pisa (2014), 193–226. Notes available online at <http://cvgmt.sns.it/paper/2416/>.
- [28] Zajicek, L.: *Porosity and σ -porosity*, Real Analysis Exchange 13(2) (1987/1988), 314–350.
- [29] Zajicek, L.: *On σ -porous sets in abstract spaces*, Abstract and Applied Analysis 2005(5), 509–534.

(Andrea Pinamonti) UNIVERSITY OF BOLOGNA, ITALY
E-mail address: Andrea.Pinamonti@gmail.com

(Gareth Speight) UNIVERSITY OF CINCINNATI, UNITED STATES
E-mail address: Gareth.Speight@uc.edu