

Trace and extension theorems for functions of bounded variation *

Lukáš Malý Nageswari Shanmugalingam Marie Snipes

November 7, 2015

Abstract

In this paper we show that every L^1 -integrable function on $\partial\Omega$ can be obtained as the trace of a function of bounded variation in Ω whenever Ω is a domain with regular boundary $\partial\Omega$ in a doubling metric measure space. In particular, the trace class of $BV(\Omega)$ is $L^1(\partial\Omega)$ provided that Ω supports a 1-Poincaré inequality. We also construct a bounded linear extension from a Besov class of functions on $\partial\Omega$ to $BV(\Omega)$.

1 Overview

A class of problems in analysis, called boundary value problems, is the group of problems that seek to find solutions to a given equation in a domain, subject to a prescribed condition on the behavior of the function at the boundary of the domain. A wide category of such problems deal with the Dirichlet boundary conditions; in such problems one wishes to prescribe the trace value of the solution at the boundary of the domain. Given a domain Ω , it is therefore natural to ask which functions f defined on $\partial\Omega$ can be extended to functions F (of some specified regularity) on the interior of Ω .

***2010 Mathematics Subject Classification:** Primary 46E35; Secondary 26A45, 26B30, 30L99, 31E05.

Keywords: Besov, BV, metric measure space, co-dimension 1 Hausdorff measure, trace, extension, Whitney cover.

More specifically, what class of functions on the boundary can be realized as the traces of the functions F in the preceding question?

The model problem that motivates our study is the problem of finding least gradient functions from the class of functions of bounded variation (BV), with prescribed boundary data, see [4, 25]. Therefore the regularity of the extended function inside the domain is BV regularity.

The paper [4] first studied the trace and extension problem for functions of bounded variation in Euclidean Lipschitz domains. It was shown there that the trace functions of BV functions on the domain lie in the L^1 -class of the boundary. In contrast, the work [13] demonstrated that every L^1 -function on the boundary of a Euclidean half-space (and hence boundaries of Lipschitz domains) has a BV extension to the half-space. Together, these two results indicate that the trace class of BV functions on a Euclidean Lipschitz domain is the L^1 -class of its boundary.

In the metric setting, a version of the Dirichlet problem associated with BV functions was considered in [15], but their notion of trace required that the BV function be defined on a larger domain. In [23] this requirement was dispensed with for domains whose boundaries are more regular (Euclidean Lipschitz domains satisfy this regularity condition). In [23] it was shown that if in addition the domain supports a 1-Poincaré inequality, then the trace of a BV function on the domain lies in a suitable L^1 -class of the boundary, thus providing an analog of the results of [4] in the metric setting. The recent work [30] gave an analog of the extension result of [13] for Lipschitz domains in Carnot–Carathéodory spaces, which indicated that it is possible to identify the trace class of BV functions in more general metric measure spaces. The goal of this paper is to provide such an identification, by adapting the technique of [13] to the metric setting.

In this paper Ω denotes a domain in a metric measure space (X, d, μ) . The natural measure on Ω is the restriction of μ to Ω . The measure we consider on the boundary $\partial\Omega$ is the co-dimension 1 Hausdorff measure $\mathcal{H} := \mathcal{H}|_{\partial\Omega}$ (see (1.9) below). The function spaces related to $\partial\Omega$ will have norms computed using the measure \mathcal{H} , and this being understood, we will not explicitly mention the measure in the notation representing these function spaces.

We now state the two main theorems of this paper. In what follows, $T : BV(\Omega) \rightarrow L^1(\partial\Omega)$ is the trace operator as constructed in [23], see (2.17).

thm:main1

Theorem 1.1. *Let Ω be a bounded domain in X that satisfies the density condition (1.14) and the co-dimension 1 Ahlfors regularity (1.15). Then there*

is a bounded linear extension operator $E : B_{1,1}^0(\partial\Omega) \rightarrow BV(\Omega)$ such that $T \circ E$ is the identity operator on $B_{1,1}^0(\partial\Omega)$.

thm:main2

Theorem 1.2. *With Ω a bounded domain in X that satisfies the density condition (1.14) and the co-dimension 1 Ahlfors regularity (1.15), there is a nonlinear bounded extension operator $\text{Ext} : L^1(\partial\Omega) \rightarrow BV(\Omega)$ such that $T \circ \text{Ext}$ is the identity operator on $L^1(\partial\Omega)$.*

The extension from $L^1(\partial\Omega)$ to $BV(\Omega)$ cannot in general be linear; this is not an artifact of our proof, see [27, 28].

Combining the above results with those of [23] we obtain the following identification of the trace class of $BV(\Omega)$.

Corollary 1.3. *Let X support a 1-Poincaré inequality. With Ω a bounded domain in X that satisfies the density condition (1.14), the co-dimension 1 Ahlfors regularity (1.15), and 1-Poincaré inequality, we have that the trace class of $BV(\Omega)$ is $L^1(\partial\Omega)$.*

For clarity, we note that the statements of Theorems 1.1 and 1.2 do not require the domains or the ambient metric measure space to support any Poincaré inequality, which allows the domains to have interior cusps or slits. Thus, even in the Euclidean setting our methods give rise to new results, as the results of [13] and [30] are in the setting of Lipschitz domains.

A related problem is to investigate the extensions of functions from a domain Ω to the whole space. See [5, 8, 17, 22, 24].

Acknowledgement: The research of the first author was supported by the Knut and Alice Wallenberg Foundation (Sweden), and the research of the second author was partially supported by the NSF grant DMS-1500440 (U.S.A.). Majority of the research for this paper was conducted during the visit of the third author to the University of Cincinnati; she wishes to thank that institution for its kind hospitality.

1.4 Notation and definitions

In this section (Z, d, ν) denotes a metric measure space with ν a Radon measure. We say that ν is *doubling* if there is a constant C_D such that for each $z \in Z$ and $r > 0$,

$$0 < \nu(B(z, 2r)) \leq C_D \nu(B(z, r)) < \infty.$$

Given a Lipschitz function f on a subset $A \subset Z$, we set

$$\text{LIP}(f, A) := \sup_{x, y \in A : x \neq y} \frac{|f(x) - f(y)|}{d(x, y)}.$$

When x is a point in the interior of $A \subset Z$, we set

$$\text{Lip } f(x) := \limsup_{y \rightarrow x} \frac{|f(y) - f(x)|}{d(y, x)}.$$

In what follows, given a metric measure space (Z, d, ν) , the space $L^1_{\text{loc}}(Z)$ consists of functions on Z that are integrable on *bounded* subsets of Z . We follow [26] to define the function class $BV(Z)$. The space $BV(Z)$ of functions of bounded variation consists of functions in $L^1(Z)$ that also have finite total variation on Z . The total variation of a function on a metric measure space is measured using upper gradients; the notion of upper gradients, first formulated in [20] (with the terminology “very weak gradients”), plays the role of $|\nabla u|$ in the metric setting where no natural distributional derivative structure exists. A Borel function $g : Z \rightarrow [0, \infty]$ is an upper gradient of $u : Z \rightarrow \mathbb{R} \cup \{\pm\infty\}$ if the following inequality holds for all (rectifiable) curves $\gamma : [a, b] \rightarrow Z$, (denoting $x = \gamma(a)$ and $y = \gamma(b)$),

$$|u(y) - u(x)| \leq \int_{\gamma} g \, ds$$

whenever $u(x)$ and $u(y)$ are both finite, and $\int_{\gamma} g \, ds = \infty$ otherwise. For each function u as above, we set $I(u : Z)$ to be the infimum of the quantity $\int_Z g \, d\nu$ over all upper gradients (in Z) g of u .

rem:lip-upp1

Remark 1.5. We note here that if u is a (locally) Lipschitz function on Z , then $\text{Lip } u$ is an upper gradient of u ; see for example [19]. We refer the interested reader to [6, 18] for more on upper gradients.

The *total variation* of the function $u \in L^1_{\text{loc}}(Z)$ is given by

$$\|Du\|(Z) := \inf \left\{ \liminf_{i \rightarrow \infty} I(u_i : Z) : u_i \in \text{Lip}_{\text{loc}}(Z), u_i \rightarrow u \text{ in } L^1_{\text{loc}}(Z) \right\}.$$

rem:lip-upp2

Remark 1.6. From Remark 1.5 we know that if u is a locally Lipschitz continuous function on Z , then $\|Du\|(Z) \leq \int_Z \text{Lip } u \, d\mu$.

For each open set $U \subset Z$ we can set $\|Du\|(U)$ similarly:

$$\|Du\|(U) := \inf \left\{ \liminf_{i \rightarrow \infty} I(u_i : U) : u_i \in \text{Lip}_{\text{loc}}(U), u_i \rightarrow u \text{ in } L^1_{\text{loc}}(U) \right\}.$$

It was shown in [26] that if $\|Du\|(Z)$ is finite, then $U \mapsto \|Du\|(U)$ is the restriction of a Radon measure to open sets of Z . We use $\|Du\|$ to also denote this Radon measure.

Definition 1.7. The space $BV(Z)$ of functions of bounded variation is equipped with the norm

$$\|u\|_{BV(Z)} := \|u\|_{L^1(Z)} + \|Du\|(Z).$$

This definition of BV agrees with the standard notion of BV functions in the Euclidean setting, see [2, 12, 31]. See also [3] for more on the BV class in the metric setting.

We say that a measurable set $E \subset Z$ is of *finite perimeter* if $\chi_E \in BV(Z)$. It follows from [26] that the superlevel set $E_t := \{z \in Z : u(z) > t\}$ has finite perimeter for almost every $t \in \mathbb{R}$ and that the coarea formula

$$\|Du\|(A) = \int_{\mathbb{R}} \|D\chi_{E_t}\|(A) dt$$

holds true whenever $A \subset Z$ is a Borel set.

Definition 1.8 (cf. [1]). A metric space Z supports a 1-Poincaré inequality if there exist positive constants λ and C such that for all balls $B \subset Z$ and all $u \in L^1_{\text{loc}}(Z)$,

$$\int_B |u - u_B| d\nu \leq C \text{rad}(B) \frac{\|Du\|(\lambda B)}{\nu(\lambda B)}.$$

Here and in the rest of the paper, f_A denotes the *integral mean* of a function $f \in L^0(Z)$ over a measurable set $A \subset Z$ of finite positive measure, defined as

$$f_A = \int_A f d\nu = \frac{1}{\nu(A)} \int_A f d\nu$$

whenever the integral on the right-hand side exists, not necessarily finite though. Furthermore, given a ball $B = B(x, r) \subset Z$ and $\lambda > 0$, the symbol λB denotes the inflated ball $B(x, \lambda r)$.

Given $A \subset Z$, we define its *co-dimension 1 Hausdorff measure* $\mathcal{H}(A)$ by

$$\mathcal{H}(A) = \lim_{\delta \rightarrow 0^+} \inf \left\{ \sum_i \frac{\nu(B_i)}{\text{rad}(B_i)} : B_i \text{ balls in } Z, \text{rad}(B_i) < \delta, A \subset \bigcup_i B_i \right\}. \quad (1.9)$$

eq:deff-mathcal

It was shown in [1] that if ν is doubling and supports a 1-Poincaré inequality, then there is a constant $C \geq 1$ such that whenever $E \subset Z$ is of finite perimeter,

$$C^{-1} \mathcal{H}(\partial_m E) \leq \|D\chi_E\|(Z) \leq C \mathcal{H}(\partial_m E),$$

where $\partial_m E$ is the *measure-theoretic boundary* of E . It consists of those points $z \in Z$ for which

$$\limsup_{r \rightarrow 0^+} \frac{\mu(B(z, r) \cap E)}{\mu(B(z, r))} > 0 \quad \text{and} \quad \limsup_{r \rightarrow 0^+} \frac{\mu(B(z, r) \setminus E)}{\mu(B(z, r))} > 0.$$

We next turn our attention to the definition of other function spaces to be considered in this paper. The Besov classes, much studied in the Euclidean setting, made their first appearance in the metric setting in [9] and were explored further in [14].

Definition 1.10. For a fixed $R > 0$, the *Besov space* $B_{1,1}^\theta(Z)$ of smoothness $\theta \in [0, 1]$ consists of functions of finite Besov norm that is given by

$$\|u\|_{B_{1,1}^\theta(Z)} = \|u\|_{L^1(Z)} + \int_0^R \int_Z \int_{B(x,t)} |u(y) - u(x)| d\nu(y) d\nu(x) \frac{dt}{t^{1+\theta}}. \quad (1.11)$$

eq:Besov

We will show that the function class $B_{1,1}^\theta(Z)$ is in fact independent of the choice of $R \in (0, \infty)$, see Lemma 3.3 below.

The following *fractional John–Nirenberg space* was first generalized to the metric measure space setting in [17]. In the Euclidean setting it was first studied in [10] and [11], but the case $\theta = 0$ in the Euclidean setting appeared in the earlier work of John and Nirenberg [21]. The fractional John–Nirenberg space $A_{1,\tau}^\theta(Z)$, where $\theta \in [0, 1]$ is its smoothness and $\tau \geq 1$ the dilation factor, is defined via its norm

$$\|u\|_{A_{1,\tau}^\theta(Z)} = \|u\|_{L^1(Z)} + \sup_{\mathcal{B}_\tau} \sum_{B \in \mathcal{B}_\tau} \frac{1}{\text{rad}(B)^\theta} \int_{\tau B} |u - u_{\tau B}| d\nu, \quad (1.12)$$

eq:JN

where the supremum is taken over all collections \mathcal{B}_τ of balls in Z of radius at most R/τ such that $\tau B_1 \cap \tau B_2$ is empty whenever $B_1, B_2 \in \mathcal{B}_\tau$ with $B_1 \neq B_2$. The class $A_{1,\tau}^\theta(Z)$ is also independent of the exact choice of $R \in (0, \infty)$.

1.13 Standing assumptions

Throughout this paper (X, d, μ) is a metric measure space, with μ a Borel regular measure. We assume that X is complete and that μ is doubling on X . Furthermore, $\Omega \subset X$ is a bounded domain and there is a constant $C \geq 1$ such that for all $x \in \partial\Omega$, $z \in \Omega$, and $0 < r \leq \text{diam}(\Omega)$, we have

$$\mu(B(z, r) \cap \Omega) \geq C^{-1} \mu(B(z, r)), \quad (1.14) \quad \boxed{\text{density}}$$

and

$$C^{-1} \frac{\mu(B(x, r))}{r} \leq \mathcal{H}(B(x, r) \cap \partial\Omega) \leq C \frac{\mu(B(x, r))}{r}. \quad (1.15) \quad \boxed{\text{boundary-Ahlfors-regularity}}$$

The property of satisfying (1.15) will be called *Ahlfors codimension 1 regularity* of $\partial\Omega$.

Throughout the paper C represents various constants that depend solely on the doubling constant, constants related to the Poincaré inequality, and the constants related to (1.14) and (1.15). The precise value of C is not of interest to us at this time, and its value may differ in each occurrence. Given expressions a and b , we say that $a \approx b$ if there is a constant $C \geq 1$ such that $C^{-1}a \leq b \leq Ca$.

2 Bounded linear extension from Besov class to BV class: proof of Theorem 1.1

ec:B110-extension

2.1 Whitney cover and partition of unity

The following theorem from [18, Section 4.1] gives the existence of a Whitney *covering* of an open subset Ω of a doubling metric space X by balls whose radii are comparable to their distance from the boundary, see also [7].

Theorem 2.2. *Let $\Omega \subsetneq X$ be open. Then there exists a countable collection $\mathcal{W}_\Omega = \{B(p_{j,i}, r_{j,i}) = B_{j,i}\}$ of balls in Ω so that*

- $\bigcup_{j,i} B_{j,i} = \Omega$,
- $\sum_{j,i} \chi_{B(p_{j,i}, 2r_{j,i})} \leq 2C_D^5$,
- $2^{j-1} < r_{j,i} \leq 2^j$ for all i ,
- and so that $r_{j,i} = \frac{1}{8} \text{dist}(p_{j,i}, X \setminus \Omega)$.

Here the constant C_D is the doubling constant of the measure μ .

Since the radii of the balls are sufficiently small, we have that $2B_i \subset \Omega$. By the boundedness of Ω there is a largest exponent j that occurs in the cover; we denote this exponent by j_0 . Hence $-j \in \mathbb{N} \cup \{0, \dots, -j_0\}$. Note that 2^{j_0} is comparable to $\text{diam}(\Omega)$. One wishing to keep track of the relationships between various constants should therefore keep in mind that the constants that depend on j_0 then depend on $\text{diam}(\Omega)$.

We also note that no ball in level j intersects a ball in level $j+2$. This follows by the reverse triangle inequality $d(p_{j,i}, p_{j+2,k}) \geq 2^{j+4} - 2^{j+3} = 2^{j+3}$ and the bounds on the radii: $2^{j-1} < r_{j,i} \leq 2^j$ and $2^{j+1} < r_{j+2,k} \leq 2^{j+2}$.

As in [18, Section 4.1], there is a Lipschitz partition of unity $\{\varphi_{j,i}\}$ subordinate to the Whitney decomposition \mathcal{W}_Ω , that is, $\sum_{j,i} \varphi_{j,i} \equiv \chi_\Omega$ and for every ball $B_{j,i} \in \mathcal{W}_\Omega$, we have that $\chi_{B_{j,i}} \leq \varphi_{j,i} \leq \chi_{2B_{j,i}}$ and $\varphi_{j,i}$ is $C/r_{j,i}$ -Lipschitz continuous.

`ssec:BesovExt`

2.3 An extension of Besov functions

Suppose that $f : \partial\Omega \rightarrow \mathbb{R}$ is a function in $B_{1,1}^0(\partial\Omega)$. We want to define a function $F : \Omega \rightarrow \mathbb{R}$ whose trace is the original function f on $\partial\Omega$.

Consider the center of the Whitney ball $p_{j,i} \in \Omega$ and choose a closest point $q_{j,i} \in \partial\Omega$. Define $U_{j,i} := B(q_{j,i}, r_{j,i}) \cap \partial\Omega$. We set $a_{j,i} := \int_{U_{j,i}} f(y) d\mathcal{H}(y)$. Then for $x \in \Omega$ set

$$F(x) := \sum_{j,i} a_{j,i} \varphi_{j,i}.$$

In subsequent results in this section we will show that $F \in BV(\Omega)$. From the following proposition and Remark 1.6 we obtain the desired bound for $\|DF\|(\Omega)$.

`prop:extnBounds`

Proposition 2.4. *Given $\Omega \subset X$ and $f \in B_{1,1}^0(\partial\Omega)$, there exists $C > 0$ such that*

$$\int_{\Omega} \text{Lip } F d\mu \leq C \|f\|_{B_{1,1}^0(\partial\Omega)}.$$

Proof. Fix a ball $B_{\ell,m} \in \mathcal{W}_\Omega$, and fix a point $x \in B_{\ell,m}$. For all $y \in B_{\ell,m}$,

$$\begin{aligned} |F(y) - F(x)| &= \left| \sum_{j,i} a_{j,i} (\varphi_{j,i}(y) - \varphi_{j,i}(x)) \right| \\ &= \left| \sum_{j,i} (a_{j,i} - a_{\ell,m}) (\varphi_{j,i}(y) - \varphi_{j,i}(x)) \right| \leq \sum_{\substack{j,i \text{ s.t.} \\ 2B_{j,i} \cap B_{\ell,m} \neq \emptyset}} |a_{j,i} - a_{\ell,m}| \frac{C}{r_{j,i}} d(y, x). \end{aligned}$$

The last inequality in the above sequence follows from the Lipschitz constant of $\varphi_{j,i}$. Rearranging and noting that if the balls intersect then $|j - \ell| \leq 1$, we see that

$$\frac{|F(y) - F(x)|}{d(y, x)} \leq \frac{C}{r_{\ell, m}} \sum_{\substack{j, i \text{ s.t.} \\ 2B_{j, i} \cap B_{\ell, m} \neq \emptyset}} |a_{j, i} - a_{\ell, m}|.$$

Hence, we want to bound terms of the form $|a_{j, i} - a_{\ell, m}|$:

$$\begin{aligned} |a_{j, i} - a_{\ell, m}| &= \left| \int_{U_{j, i}} f(z) d\mathcal{H}(z) - \int_{U_{\ell, m}} f(z) d\mathcal{H}(z) \right| \\ &= \left| \int_{U_{j, i}} \int_{U_{\ell, m}} (f(z) - f(w)) d\mathcal{H}(w) d\mathcal{H}(z) \right| \\ &\leq \int_{U_{j, i}} \int_{U_{\ell, m}} |f(z) - f(w)| d\mathcal{H}(w) d\mathcal{H}(z) \\ &\leq \frac{1}{\mathcal{H}(U_{j, i}) \mathcal{H}(U_{\ell, m})} \int_{U_{j, i}} \int_{U_{\ell, m}} |f(z) - f(w)| d\mathcal{H}(w) d\mathcal{H}(z) \\ &\leq \frac{C}{\mathcal{H}(U_{\ell, m}^*) \mathcal{H}(U_{\ell, m}^*)} \int_{U_{j, i}} \int_{U_{\ell, m}} |f(z) - f(w)| d\mathcal{H}(w) d\mathcal{H}(z) \quad (2.5) \quad \boxed{\text{ineq: expandballs1}} \\ &\leq \frac{C}{\mathcal{H}(U_{\ell, m}^*) \mathcal{H}(U_{\ell, m}^*)} \int_{U_{\ell, m}^*} \int_{U_{\ell, m}^*} |f(z) - f(w)| d\mathcal{H}(w) d\mathcal{H}(z) \\ &= C \int_{U_{\ell, m}^*} \int_{U_{\ell, m}^*} |f(z) - f(w)| d\mathcal{H}(w) d\mathcal{H}(z), \end{aligned}$$

where $U_{\ell, m}^*$ denotes the expanded subset of the boundary:

$$U_{\ell, m}^* := B(q_{\ell, m}, 2^6 r_{\ell, m}) \cap \partial\Omega. \quad (2.6) \quad \boxed{\text{eq: expandedsubset}}$$

By the doubling property of X , the boundary regularity condition on $\partial\Omega$, and the definition of codimension-1 Hausdorff measure, we have

$$\mathcal{H}(U_{\ell, m}^*) \leq C \mathcal{H}(U_{\ell, m}),$$

which gave inequality (2.5). The above estimates together with the bounded

overlap of the Whitney balls yield the following inequality:

$$\begin{aligned} \text{Lip } F(x) &= \limsup_{y \rightarrow x} \frac{|F(y) - F(x)|}{d(y, x)} \\ &\leq \frac{C}{r_{\ell, m}} \int_{U_{\ell, m}^*} \int_{U_{\ell, m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \end{aligned} \quad (2.7)$$

eq:pointwise-Lip-

for $x \in B_{\ell, m}$. From (2.7) and (1.15) we see that

$$\begin{aligned} \int_{\Omega} \text{Lip } F(x) \, d\mu(x) &\leq \sum_{\ell, m} \int_{B_{\ell, m}} \text{Lip } F(x) \, d\mu(x) \\ &\leq \sum_{\ell, m} \mu(B_{\ell, m}) \frac{C}{r_{\ell, m}} \int_{U_{\ell, m}^*} \int_{U_{\ell, m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\leq C \sum_{\ell, m} \mathcal{H}(U_{\ell, m}) \int_{U_{\ell, m}^*} \int_{U_{\ell, m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\leq C \sum_{\ell=-\infty}^{j_0} \sum_m \int_{U_{\ell, m}^*} \int_{U_{\ell, m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\leq C \sum_{\ell=-\infty}^{j_0} \int_{\partial\Omega} \int_{B(z, 2^{7+\ell})} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z). \end{aligned}$$

Here the last inequality follows from the uniformly bounded overlap of the balls $U_{\ell, m}^*$ for each ℓ . Without loss of generality, we may choose $R = 2^{j_0+7}$ in the definition of the Besov norm (1.11). Note that $R \approx \text{diam}(\Omega)$ then. The following estimate (cf. the proof of [14, Theorem 5.2]) concludes the proof:

$$\begin{aligned} &\sum_{\ell=-\infty}^{j_0} \int_{\partial\Omega} \int_{B(z, 2^{7+\ell})} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\approx \int_{t=0}^{2^{j_0+7}} \int_{\partial\Omega} \int_{B(z, t)} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \frac{dt}{t} \quad (2.8) \\ &\leq C \|f\|_{B_{1,1}^0(\partial\Omega)}. \end{aligned}$$

eq:BesovEquivSum

□

We will use the extension constructed in this section in formulating a nonlinear bounded extension from $L^1(\partial\Omega, \mathcal{H})$ to $BV(\Omega)$ in the subsequent sections. There we will need the following estimates for the integral of the gradient and the function on layers of Ω .

layer-est-grad

Lemma 2.9. For $0 \leq \rho_1 < \rho_2 < \text{diam}(\Omega)/2$, set

$$\Omega(\rho_1, \rho_2) := \{x \in \Omega : \rho_1 \leq \text{dist}(x, X \setminus \Omega) < \rho_2\}. \quad (2.10)$$

eq:def-OmRhoRho

Let $\mathcal{J}(\rho_1, \rho_2)$ be the collection of all $\ell \in \mathbb{Z}$ such that there is some $m \in \mathbb{N}$ with $2B_{\ell,m} \cap \Omega(\rho_1, \rho_2)$ non-empty. Then

$$\int_{\Omega(\rho_1, \rho_2)} \text{Lip } F \, d\mu \leq C \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \int_{\partial\Omega} \int_{B(z, 2^{7+\ell})} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z).$$

Proof. For each $\ell \in \mathcal{J}(\rho_1, \rho_2)$ let $\mathcal{I}(\ell)$ denote the collection of all $m \in \mathbb{N}$ for which $2B_{\ell,m} \cap \Omega(\rho_1, \rho_2)$ is non-empty. Then by (2.7) and (1.15),

$$\begin{aligned} \int_{\Omega(\rho_1, \rho_2)} \text{Lip } F \, d\mu &\leq \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}(\ell)} \int_{B_{\ell,m}} \text{Lip } F \, d\mu \\ &\leq C \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}(\ell)} \frac{\mu(B_{\ell,m})}{r_{\ell,m}^\ell} \int_{U_{\ell,m}^*} \int_{U_{\ell,m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\leq C \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}(\ell)} \mathcal{H}(U_{\ell,m}) \int_{U_{\ell,m}^*} \int_{U_{\ell,m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &= C \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}(\ell)} \int_{U_{\ell,m}^*} \int_{U_{\ell,m}^*} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\leq C \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}(\ell)} \int_{U_{\ell,m}^*} \int_{B(z, 2^{7+\ell})} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ &\leq C \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} \int_{\partial\Omega} \int_{B(z, 2^{7+\ell})} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z). \quad \square \end{aligned}$$

ayer-est-grad-lip

Corollary 2.11. Using the notation of Lemma 2.9, we have that

$$\int_{\Omega(\rho_1, \rho_2)} \text{Lip } F \, d\mu \leq C \rho_2 \mathcal{H}(\partial\Omega) \text{LIP}(f, \partial\Omega)$$

whenever f is Lipschitz on $\partial\Omega$.

Proof. For a fixed $\ell \in \mathbb{Z}$, we can estimate

$$\begin{aligned} \int_{\partial\Omega} \int_{B(z, 2^{7+\ell})} |f(z) - f(w)| \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ \leq \int_{\partial\Omega} \int_{B(z, 2^{7+\ell})} \text{LIP}(f, \partial\Omega) \, d(z, w) \, d\mathcal{H}(w) \, d\mathcal{H}(z) \\ \leq C \mathcal{H}(\partial\Omega) \text{LIP}(f, \partial\Omega) 2^{7+\ell}. \end{aligned}$$

Therefore,

$$\int_{\Omega(\rho_1, \rho_2)} \text{Lip } F \, d\mu \leq C \mathcal{H}(\partial\Omega) \text{LIP}(f, \partial\Omega) \sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} 2^\ell.$$

Every ball $B = B(p, r) \in \mathcal{I}(\ell)$ satisfies $2^{\ell-1} < r \leq 2^\ell$ and $\text{dist}(p, X \setminus \Omega) = 8r$. There is $C \geq 1$ such that $C^{-1}\rho_1 \leq 2^\ell \leq C\rho_2$ whenever $\ell \in \mathcal{J}(\rho_1, \rho_2)$. Thus, $\sum_{\ell \in \mathcal{J}(\rho_1, \rho_2)} 2^\ell \leq C\rho_2$. \square

We next turn our attention to the L^1 -estimates for F .

em:L1-est_Whitney

Lemma 2.12. *There exists $C > 0$ such that*

$$\int_{\Omega} |F| \, d\mu \leq C \text{diam}(\Omega) \|f\|_{L^1(\partial\Omega)}.$$

Proof. We first consider a fixed ball $B_{\ell, m}$ from the Whitney cover. Then

$$\begin{aligned} \int_{B_{\ell, m}} |F(x)| \, d\mu(x) &= \int_{B_{\ell, m}} \left| \sum_{j, i} \int_{U_{j, i}} f(y) \, d\mathcal{H}(y) \varphi_{j, i}(x) \right| \, d\mu(x) \\ &\leq \int_{B_{\ell, m}} \sum_{j, i} \left| \int_{U_{j, i}} f(y) \, d\mathcal{H}(y) \right| \varphi_{j, i}(x) \, d\mu(x) \\ &= \int_{B_{\ell, m}} \sum_{\substack{j, i \text{ s.t.} \\ 2B_{j, i} \cap B_{\ell, m} \neq \emptyset}} \left| \int_{U_{j, i}} f(y) \, d\mathcal{H}(y) \right| \varphi_{j, i}(x) \, d\mu(x). \end{aligned}$$

Recall that if $2B_{j, i} \cap B_{\ell, m} \neq \emptyset$, then $|j - \ell| \leq 1$, so $\mathcal{H}(U_{j, i}) \approx \mathcal{H}(U_{\ell, m}^*)$. Also, for $U_{\ell, m}^*$ as defined in equation (2.6), $U_{j, i} \subset U_{\ell, m}^*$. Furthermore, by the construction of the Whitney decomposition, each point is in a fixed number of dilated Whitney balls $2B_{j, i}$. Hence,

$$\begin{aligned} \int_{B_{\ell, m}} \sum_{\substack{j, i \text{ s.t.} \\ 2B_{j, i} \cap B_{\ell, m} \neq \emptyset}} \left| \int_{U_{j, i}} f(y) \, d\mathcal{H}(y) \right| \varphi_{j, i}(x) \, d\mu(x) \\ \leq C \int_{B_{\ell, m}} \int_{U_{\ell, m}^*} |f(y)| \, d\mathcal{H}(y) \, d\mu(x) \leq C \mu(B_{\ell, m}) \int_{U_{\ell, m}^*} |f(y)| \, d\mathcal{H}(y). \end{aligned}$$

In view of (1.15), we obtain that

$$\int_{B_{\ell,m}} |F(x)| d\mu(x) \leq C r_{\ell,m} \int_{U_{\ell,m}^*} |f(y)| d\mathcal{H}(y). \quad (2.13) \quad \boxed{\text{eq:Ball-F-est}}$$

Summing up and noting that $\Omega = \bigcup_{\ell,m} B_{\ell,m}$, we have

$$\begin{aligned} \int_{\Omega} |F| d\mu &\leq C \sum_{\ell=-\infty}^{j_0} \sum_m r_{\ell,m} \int_{U_{\ell,m}^*} |f| d\mathcal{H} \leq C \sum_{\ell=-\infty}^{j_0} 2^\ell \sum_m \int_{U_{\ell,m}^*} |f| d\mathcal{H} \\ &\leq C \sum_{\ell=-\infty}^{j_0} 2^\ell \int_{\partial\Omega} |f| d\mathcal{H} \leq C \text{diam}(\Omega) \|f\|_{L^1(\partial\Omega)}. \quad \square \end{aligned}$$

We now aim to obtain an analog of Lemma 2.9 for the L^1 -norm of F on the layer $\Omega(\rho_1, \rho_2)$.

t-Fn-boundaryball

Lemma 2.14. *Let $z \in \partial\Omega$ and $r \in (0, \text{diam}(\Omega)/2)$. Then,*

$$\int_{B(z,r) \cap \Omega(\rho_1, \rho_2)} |F| d\mu \leq C \min\{r, \rho_2\} \int_{B(z, 2^8 r) \cap \partial\Omega} |f| d\mathcal{H}$$

whenever $0 \leq \rho_1 < \min\{r, \rho_2\}$ and $\rho_2 < \text{diam}(\Omega)/2$.

Proof. Since $B(z, r) \cap \Omega(\rho_1, \rho_2) = B(z, r) \cap \Omega(\rho_1, \min\{r, \rho_2\})$, we do not lose any generality by assuming that $\rho_2 \leq r$. Similarly as in the proof of Lemma 2.9, we set $\mathcal{J}'(\rho_1, \rho_2)$ to be the collection of all ℓ for which there is some m such that $2B_{\ell,m} \cap \Omega(\rho_1, \rho_2) \cap B(z, r)$ is non-empty, and for each $\ell \in \mathcal{J}'(\rho_1, \rho_2)$ we set $\mathcal{I}'(\ell)$ to be the collection of all $m \in \mathbb{N}$ for which $2B_{\ell,m} \cap \Omega(\rho_1, \rho_2) \cap B(z, r)$ is non-empty. Then by (2.13),

$$\begin{aligned} \int_{B(z,r) \cap \Omega(\rho_1, \rho_2)} |F| d\mu &\leq \sum_{\ell \in \mathcal{J}'(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}'(\ell)} \int_{B_{\ell,m}} |F| d\mu \\ &\leq C \sum_{\ell \in \mathcal{J}'(\rho_1, \rho_2)} \sum_{m \in \mathcal{I}'(\ell)} r_{\ell,m} \int_{U_{\ell,m}^*} |f| d\mathcal{H}. \end{aligned}$$

The triangle inequality yields that

$$d(z, q_{\ell,m}) \leq d(z, p_{\ell,m}) + d(p_{\ell,m}, q_{\ell,m}) \leq 2d(z, p_{\ell,m}) \leq 2(r + 2r_{\ell,m}),$$

where $B_{\ell,m} = B(p_{\ell,m}, r_{\ell,m})$ and $U_{\ell,m} = B(q_{\ell,m}, r_{\ell,m}) \cap \partial\Omega$ with $q_{\ell,m} \in \partial\Omega$ being a boundary point lying closest to $p_{\ell,m}$. Moreover, $8r_{\ell,m} = \text{dist}(p_{\ell,m}, X \setminus \Omega) \leq d(p_{\ell,m}, z) \leq r + 2r_{\ell,m}$. Hence, $r_{\ell,m} \leq \frac{1}{6}r$. Consequently, $d(z, q_{\ell,m}) \leq \frac{8}{3}r$ and $U_{\ell,m} \subset B(z, (\frac{8}{3} + \frac{1}{6})r)$. Thus, $U_{\ell,m}^* \subset B(z, 2^8 r)$ and

$$\begin{aligned} \int_{B(z,r) \cap \Omega(\rho_1, \rho_2)} |F| d\mu &\leq C \sum_{\ell \in \mathcal{J}'(\rho_1, \rho_2)} 2^\ell \int_{B(z, 2^8 r) \cap \partial\Omega} |f| d\mathcal{H} \\ &\leq C \rho_2 \int_{B(z, 2^8 r) \cap \partial\Omega} |f| d\mathcal{H}, \end{aligned}$$

where the last inequality can be verified as follows: Every ball $B = B(p, r) \in \mathcal{I}'(\ell)$ satisfies $2^{\ell-1} < r \leq 2^\ell$ and $\text{dist}(p, X \setminus \Omega) = 8r$. There is $C \geq 1$ such that $C^{-1}\rho_1 \leq 2^\ell \leq C\rho_2$ whenever $\ell \in \mathcal{J}'(\rho_1, \rho_2)$. Thus, $\sum_{\ell \in \mathcal{J}'(\rho_1, \rho_2)} 2^\ell \leq C\rho_2$. \square

By covering $\partial\Omega$ by balls of radii r , whose overlap is bounded, we obtain the following corollary.

layer-est-Fn

Corollary 2.15. *With the notation of Lemma 2.9, we have*

$$\int_{\Omega(\rho_1, \rho_2)} |F| d\mu \leq C \rho_2 \int_{\partial\Omega} |f| d\mathcal{H}.$$

2.16 Trace of extension is the identity mapping

From the above lemma we know that given a function $f \in B_{1,1}^0(\partial\Omega)$ the corresponding function F is in the class $N^{1,1}(\Omega) \subset BV(\Omega)$, where $N^{1,1}(\Omega)$ is a Newtonian class introduced in [29]. The mapping $f \mapsto F$ is denoted by the operator $E : B_{1,1}^0(\partial\Omega) \rightarrow BV(\Omega)$. This operator is bounded by Proposition 2.4 and Lemma 2.12, and it is linear by construction.

We now wish to show that the trace of F returns the original function f , i.e., $T \circ E$ is the identity function on $B_{1,1}^0(\partial\Omega)$. It was shown in [23] that if Ω satisfies our standing assumptions and supports a 1-Poincaré inequality, then for each $u \in BV(\Omega)$ and for \mathcal{H} -a.e. $z \in \partial\Omega$ there is a number $Tu(z) \in \mathbb{R}$ such that

$$\limsup_{r \rightarrow 0^+} \int_{B(z,r) \cap \Omega} |u(y) - Tu(z)| d\mu(y) = 0. \quad (2.17) \quad \text{def of trace}$$

The map $u \mapsto Tu$ is called the *trace* of $BV(\Omega)$. Moreover, $Tu \in L^1(\partial\Omega)$ provided that $u \in BV(\Omega)$.

Note also that $B_{1,1}^0(\partial\Omega) \subset L^1(\partial\Omega)$ and the inclusion is strict in general, which is shown in Example 3.11 below. Further properties of the Besov classes are explored in Section 3.

For the sake of clarity, let us explicitly point out that the following lemma shows that the BV extension of a function of the Besov class $B_{1,1}^0(\partial\Omega)$, as constructed above, has a well-defined trace even though no Poincaré inequality for Ω or for X is assumed.

trace-Besov

Lemma 2.18. *For f and F as above, and for \mathcal{H} -a.e. $z \in \partial\Omega$,*

$$\lim_{r \rightarrow 0^+} \int_{B(z,r) \cap \Omega} |F(x) - f(z)| d\mu(x) = 0.$$

That is, $TEf(z)$ exists for \mathcal{H} -a.e. $z \in \partial\Omega$.

Proof. Since $f \in B_{1,1}^0(\partial\Omega) \subset L^1(\partial\Omega)$, we know by the doubling property of $\mathcal{H}|_{\partial\Omega}$ that \mathcal{H} -a.e. $z \in \partial\Omega$ is a Lebesgue point of f . Let z be such a point. Since $\sum_{j,i} \varphi_{j,i} = \chi_\Omega$, we have

$$\begin{aligned} \int_{B(z,r) \cap \Omega} |F - f(z)| d\mu &= \int_{B(z,r) \cap \Omega} \left| \sum_{j,i} \left(\int_{U_{j,i}} f d\mathcal{H} \right) \varphi_{j,i}(x) - f(z) \right| d\mu(x) \\ &= \int_{B(z,r) \cap \Omega} \left| \sum_{j,i} \left(\int_{U_{j,i}} (f - f(z)) d\mathcal{H} \right) \varphi_{j,i}(x) \right| d\mu(x) \\ &\leq \int_{B(z,r) \cap \Omega} \sum_{j,i} \left(\int_{U_{j,i}} |f - f(z)| d\mathcal{H} \right) \varphi_{j,i}(x) d\mu(x). \end{aligned}$$

By the properties of the Whitney covering \mathcal{W}_Ω ,

$$\begin{aligned} &\int_{B(z,r) \cap \Omega} \sum_{j,i} \left(\int_{U_{j,i}} |f - f(z)| d\mathcal{H} \right) \varphi_{j,i}(x) d\mu(x) \\ &\leq \sum_{\substack{\ell, m \text{ s.t.} \\ 2B_{\ell,m} \cap B(z,r) \neq \emptyset}} \int_{B_{\ell,m}} \sum_{j,i} \left(\int_{U_{j,i}} |f - f(z)| d\mathcal{H} \right) \varphi_{j,i}(x) d\mu(x) \\ &\leq \sum_{\substack{\ell, m \text{ s.t.} \\ 2B_{\ell,m} \cap B(z,r) \neq \emptyset}} \int_{B_{\ell,m}} \sum_{\substack{j,i \text{ s.t.} \\ 2B_{j,i} \cap B_{\ell,m} \neq \emptyset}} \left(\int_{U_{j,i}} |f - f(z)| d\mathcal{H} \right) d\mu(x). \end{aligned}$$

If $2B_{j,i} \cap B_{\ell,m}$ is non-empty, then $U_{j,i} \subset U_{\ell,m}^*$ and $\mathcal{H}(U_{j,i}) \approx \mathcal{H}(U_{\ell,m}^*)$. Therefore by (1.15) we have

$$\begin{aligned}
& \int_{B(z,r) \cap \Omega} \sum_{j,i} \left(\int_{U_{j,i}} |f - f(z)| d\mathcal{H} \right) \varphi_{j,i}(x) d\mu(x) \\
& \leq C \sum_{\substack{\ell,m \text{ s.t.} \\ 2B_{\ell,m} \cap B(z,r) \neq \emptyset}} \left(\int_{U_{\ell,m}^*} |f - f(z)| d\mathcal{H} \right) \mu(B_{\ell,m}) \\
& \leq C \sum_{\substack{\ell,m \text{ s.t.} \\ 2B_{\ell,m} \cap B(z,r) \neq \emptyset}} r_{\ell,m} \int_{U_{\ell,m}^*} |f - f(z)| d\mathcal{H}.
\end{aligned}$$

Let $\mathcal{J}(B(z,r))$ denote the collection of all $\ell \in \mathbb{Z}$ for which there is some $m \in \mathbb{N}$ such that $2B_{\ell,m} \cap B(z,r)$ is non-empty. For each $\ell \in \mathcal{J}(B(z,r))$, set $\mathcal{I}(\ell)$ to be the collection of all $m \in \mathbb{N}$ for which $2B_{\ell,m} \cap B(z,r)$ is non-empty. Then,

$$\begin{aligned}
& \int_{B(z,r) \cap \Omega} \sum_{j,i} \left(\int_{U_{j,i}} |f - f(z)| d\mathcal{H} \right) \varphi_{j,i}(x) d\mu(x) \\
& \leq C \sum_{\ell \in \mathcal{J}(B(z,r))} 2^\ell \sum_{m \in \mathcal{I}(\ell)} \int_{U_{\ell,m}^*} |f - f(z)| d\mathcal{H} \\
& \leq C \sum_{\ell \in \mathcal{J}(B(z,r))} 2^\ell \int_{B(z,2^7 r) \cap \partial\Omega} |f - f(z)| d\mathcal{H} \\
& \leq C r \int_{B(z,2^7 r) \cap \partial\Omega} |f - f(z)| d\mathcal{H}.
\end{aligned}$$

In the above, we used the fact that $\sum_{\ell \in \mathcal{J}(B(z,r))} 2^\ell \approx r$, since only the indices $\ell \in \mathbb{Z}$ for which $2^\ell \approx \text{dist}(B_{\ell,m}, X \setminus \Omega) \leq r$ are allowed to be in $\mathcal{J}(B(z,r))$. From the fact that z is a Lebesgue point of f , we now have

$$\begin{aligned}
\int_{B(z,r) \cap \Omega} |F - f(z)| d\mu & \leq C \frac{r}{\mu(B(z,r) \cap \Omega)} \int_{B(z,2^7 r) \cap \partial\Omega} |f - f(z)| d\mathcal{H} \\
& \leq C \int_{B(z,2^7 r) \cap \partial\Omega} |f - f(z)| d\mu \rightarrow 0 \text{ as } r \rightarrow 0^+.
\end{aligned}$$

This completes the proof of the lemma. \square

3 Comparison of $B_{1,1}^0(\partial\Omega)$ and other function spaces

sec:spaces

We now wish to show that $B_{1,1}^0(\partial\Omega)$ has more interesting functions than mere constant functions. What functions are in $B_{1,1}^0(\partial\Omega)$? Since the results of this section deal with function spaces based on more general doubling metric measure spaces, we consider the underlying metric measure space (Z, d, ν) . The other function spaces include L^1 , BV, and the fractional John–Nirenberg spaces as well as the class of Lipschitz functions.

Let $Z = (Z, d, \nu)$ be a metric space endowed with a doubling measure. In applications in this paper, Z will be $\partial\Omega \subset X$ and ν will be the Hausdorff co-dimension 1 measure $\mathcal{H}|_{\partial\Omega}$.

3.1 Preliminary results

cor:ballcovering

Lemma 3.2. *For every $\tau > 3$, there is $C = C(C_D, \tau) \geq 1$ such that for every $r > 0$ there is an at most countable set of points $\{x_j\}_j \subset Z$ (alternatively, $\{x_j\}_j \subset \Omega$, where $\Omega \subset Z$ is arbitrary) such that*

- $B(x_j, r) \cap B(x_k, r) = \emptyset$ whenever $j \neq k$;
- $Z = \bigcup_j B(x_j, \tau r)$ (alternatively, $\Omega \subset \bigcup_j B(x_j, \tau r)$);
- $\sum_j \chi_{B(x_j, \tau r)} \leq C$.

The above lemma is widely known to experts in the field, but we were unable to find it in current literature; hence we provide a sketch of its proof.

Proof. An application of Zorn’s lemma or [18, Lemma 4.1.12] gives a countable set $A \subset Z$ such that for distinct points $x, y \in A$ we have $d(x, y) \geq r$, and for each $z \in Z$ there is some $x \in A$ such that $d(z, x) < r$. The countable collection $\{B(x, r) : x \in A\}$ can be seen to satisfy the requirements set forth in the lemma because of the doubling property of ν . \square

lem:BesovIndepR

Lemma 3.3. *Let $f \in L^1(Z)$. Then,*

$$\int_0^r \int_Z \int_{B(x,t)} |f(y) - f(x)| d\nu(y) d\nu(x) \frac{dt}{t^{1+\theta}} < \infty \quad \text{if and only if}$$

$$\int_0^R \int_Z \int_{B(x,t)} |f(y) - f(x)| d\nu(y) d\nu(x) \frac{dt}{t^{1+\theta}} < \infty,$$

where $0 < r < R < \infty$. If $\theta > 0$, then the equivalence holds true even for $R = \infty$.

Proof. By the triangle inequality, we obtain for $t > 0$ that

$$\begin{aligned} \int_Z \int_{B(x,t)} |f(y) - f(x)| d\nu(y) d\nu(x) &\leq \int_Z \int_{B(x,t)} (|f(y)| + |f(x)|) d\nu(y) d\nu(x) \\ &= \int_Z \left(|f(x)| + \int_{B(x,t)} |f(y)| d\nu(y) \right) d\nu(x) \\ &= \|f\|_{L^1(Z)} + \int_Z \int_{B(x,t)} |f(y)| d\nu(y) d\nu(x). \end{aligned}$$

The Fubini theorem and the doubling condition then yield

$$\begin{aligned} \int_Z \int_{B(x,t)} |f(y)| d\nu(y) d\nu(x) &\approx \int_{Z \times Z} \frac{|f(y)| \chi_{(0,t)}(d(x,y))}{\nu(B(y,t))} d(\nu \times \nu)(x, y) \\ &= \int_Z |f(y)| \int_{B(y,t)} d\nu(x) d\nu(y) = \|f\|_{L^1(Z)}. \end{aligned}$$

For $r > 0$ set

$$I(r) := \int_0^r \int_Z \int_{B(x,t)} |f(y) - f(x)| d\nu(y) d\nu(x) \frac{dt}{t^{1+\theta}}.$$

Then,

$$\begin{aligned} I(R) &= I(r) + \int_r^R \int_Z \int_{B(x,t)} |f(y) - f(x)| d\nu(y) d\nu(x) \frac{dt}{t^{1+\theta}} \\ &\leq I(r) + C \int_r^R \|f\|_{L^1(Z)} \frac{dt}{t^{1+\theta}} = I(r) + C \|f\|_{L^1(Z)} \left(\frac{1}{r^\theta} - \frac{1}{R^\theta} \right), \end{aligned}$$

with obvious modification for $\theta = 0$. □

lem:BesovEquiv

Lemma 3.4. *Let $R \leq 2 \operatorname{diam}(Z)$ and $\theta \in [0, 1]$. Then,*

$$I(R) \approx \int_Z \int_{B(x,R)} \frac{|f(y) - f(x)|}{\nu(B(x, d(x,y))) d(x,y)^\theta} d\nu(y) d\nu(x).$$

Proof. The equivalence follows from the Fubini theorem, see also [14, Theorem 5.2]. □

Lemma 3.5. *There is a constant $C \geq 1$ and there are collections of balls \mathcal{B}^k , $k = 0, 1, \dots$, such that*

$$C^{-1}I(1) \leq \sum_{k=0}^{\infty} \sum_{B \in \mathcal{B}^k} \frac{1}{\text{rad}(B)^\theta} \int_B |f - f_B| d\nu \leq CI(4).$$

Moreover, $\text{rad}(B) \approx 2^{-k}$ whenever $B \in \mathcal{B}^k$, and the balls within each collection \mathcal{B}^k have bounded overlap (also after inflation by a given factor $\tau \geq 1$).

Proof. By Lemma 3.2, there is a constant $C = C(C_D, \tau) \geq 1$ and collections of balls $\tilde{\mathcal{B}}^k$, $k = 0, 1, \dots$, such that $\text{rad}(B) = 2^{-k}$ for every $B \in \tilde{\mathcal{B}}^k$ and $1 \leq \sum_{B \in \tilde{\mathcal{B}}^k} \chi_{2\tau B}(x) \leq C$ for all $x \in Z$. Then, by (2.8),

$$\begin{aligned} I(1) &\leq \sum_{k=0}^{\infty} 2^{k\theta} \sum_{B \in \tilde{\mathcal{B}}^k} \int_B \int_{B(x, 2^{-k})} |f(y) - f(x)| d\nu(y) d\nu(x) \\ &\leq C \sum_{k=0}^{\infty} 2^{k\theta} \sum_{B \in \tilde{\mathcal{B}}^k} \int_B \int_{2B} |f(y) - f(x)| d\nu(y) d\nu(x) \\ &\leq C \sum_{k=0}^{\infty} \sum_{B \in \tilde{\mathcal{B}}^k} \frac{1}{(2^{-k})^\theta} \int_{2B} \int_{2B} |f(y) - f(x)| d\nu(y) d\nu(x) \\ &\leq C \sum_{k=0}^{\infty} \sum_{B \in \tilde{\mathcal{B}}^k} \frac{1}{\text{rad}(B)^\theta} \int_{2B} |f - f_{2B}| d\nu. \end{aligned}$$

Thus, we choose $\mathcal{B}^k = \{2B : B \in \tilde{\mathcal{B}}^k\}$, $k = 0, 1, \dots$, to conclude the proof of the first inequality.

The proof of the second inequality follows analogous steps backwards. Recall that $\text{rad}(B) = 2^{1-k}$ whenever $B \in \mathcal{B}^k$. Thus,

$$\begin{aligned} &\sum_{k=0}^{\infty} \sum_{B \in \mathcal{B}^k} \frac{1}{\text{rad}(B)^\theta} \int_B |f - f_B| d\nu \\ &\approx \sum_{k=0}^{\infty} \sum_{B \in \mathcal{B}^k} 2^{k\theta} \int_B \int_B |f(y) - f(x)| d\nu(y) d\nu(x) \\ &\leq C \sum_{k=0}^{\infty} 2^{k\theta} \sum_{B \in \mathcal{B}^k} \int_B \int_{B(x, 2^{2-k})} |f(y) - f(x)| d\nu(y) d\nu(x) \\ &\leq C \sum_{k=0}^{\infty} 2^{k\theta} \int_Z \int_{B(x, 2^{2-k})} |f(y) - f(x)| d\nu(y) d\nu(x), \end{aligned}$$

where we used the fact that the balls have uniformly bounded overlap within each collection \mathcal{B}^k . \square

Remark 3.6. Let $0 \leq \theta < \eta \leq 1$. Then, $\|u\|_{B_{1,1}^\theta} \leq C(1 + R^{\eta-\theta})\|u\|_{B_{1,1}^\eta}$ and $\|u\|_{A_{1,\tau}^\theta} \leq C(1 + R^{\eta-\theta})\|u\|_{A_{1,\tau}^\eta}$ for any $\tau \geq 1$.

3.7 Comparison of function spaces with $B_{1,1}^\theta(Z)$

Proposition 3.8. Let $\theta \in [0, 1]$ and $\tau \geq 1$ be arbitrary. Then there is a constant $C \geq 1$, which depends on θ and τ , such that

$$\|u\|_{A_{1,\tau}^\theta} \leq C\|u\|_{B_{1,1}^\theta}.$$

Proof. Let \mathcal{B}_τ be a fixed collection of non-overlapping balls in Z of radius at most R/τ . Then,

$$\begin{aligned} \sum_{B \in \mathcal{B}_\tau} \frac{1}{\text{rad}(B)^\theta} \int_{\tau B} |u - u_{\tau B}| d\nu &\approx \sum_{B \in \mathcal{B}_\tau} \frac{1}{(\tau \text{rad}(B))^\theta} \int_{\tau B} \int_{\tau B} |u(x) - u(y)| d\nu(y) d\nu(x) \\ &= \int_Z \sum_{B \in \mathcal{B}_\tau} \frac{\chi_{\tau B}(x)}{(\tau \text{rad}(B))^\theta} \int_{\tau B} |u(x) - u(y)| d\nu(y) d\nu(x) \\ &\leq \int_Z \sum_{B \in \mathcal{B}_\tau} \chi_{\tau B}(x) \int_{\tau B} \frac{|u(x) - u(y)|}{\nu(B(x, d(x, y)))d(x, y)^\theta} d\nu(y) d\nu(x) \\ &\leq \int_Z \int_{B(x, 2R)} \frac{|u(x) - u(y)|}{\nu(B(x, d(x, y)))d(x, y)^\theta} d\nu(y) d\nu(x) \approx I(2R), \end{aligned}$$

where we used that ν is doubling and $B(x, d(x, y)) \subset 3\tau B$ for all $x, y \in \tau B$. Taking supremum over all collections of balls concludes the proof. \square

Proposition 3.9. Assume that Z is bounded. Let $0 \leq \theta < \eta \leq 1$ and $\tau \geq 1$. Then, $A_{1,\tau}^\eta(Z) \subset B_{1,1}^\theta(Z)$.

Proof. We use the characterization of Besov functions from Lemma 3.5.

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{B \in \mathcal{B}^k} \frac{1}{\text{rad}(B)^\theta} \int_B |f - f_B| d\nu &\leq C \sum_{k=0}^{\infty} \sum_{B \in \mathcal{B}^k} \frac{2^{k(\theta-\eta)}}{\text{rad}(B)^\eta} \int_B |f - f_B| d\nu \leq C \sum_{k=0}^{\infty} 2^{k(\theta-\eta)} \|f\|_{A_{1,\tau}^\eta(Z)}. \quad \square \end{aligned}$$

Lemma 3.10. *For every $\tau \geq 1$, we have $L^1(Z) = A_{1,\tau}^0(Z)$.*

Proof. Let \mathcal{B}_τ be a fixed collection of balls in Z that remain pairwise disjoint after being inflated τ -times. Then, the triangle inequality yields that

$$\begin{aligned} \sum_{B \in \mathcal{B}_\tau} \frac{1}{\text{rad}(B)^0} \int_{\tau B} |u - u_{\tau B}| d\nu &\approx \sum_{B \in \mathcal{B}_\tau} \left(\int_{\tau B} |u| + \left| \int_{\tau B} u d\nu \right| \right) d\nu \\ &= \sum_{B \in \mathcal{B}_\tau} 2 \int_{\tau B} |u| d\nu \leq 2 \int_Z |u| d\nu. \end{aligned}$$

Hence, $\|u\|_{A_{1,\tau}^0(Z)} \leq 3\|u\|_{L^1(Z)}$.

Conversely, $\|u\|_{A_{1,\tau}^0(Z)} \geq \|u\|_{L^1(Z)}$ by the definition (1.12). \square

The following example shows that the inclusion $B_{1,1}^0(Z) \subset A_{1,\tau}^0(Z)$ may, in general, be strict.

exa:Bes0!=L1

Example 3.11. Let

$$f(x) = \sum_{j=1}^{\infty} \chi_{[1/(j+1), 1/j)}(x) u(4^j x), \quad x \in (0, 1),$$

where u is the 1-periodic extension of $\chi_{[0, 1/2)}$. Obviously, $f \in L^\infty(0, 1)$. Hence, $f \in L^1(0, 1) = A_{1,\tau}^0(0, 1)$. On the other hand, $u \notin B_{1,1}^0(0, 1)$, which we are about to show.

We will use the characterization of $B_{1,1}^0(0, 1)$ from Lemma 3.5. There, we may choose $\mathcal{B}^k = \{(l2^{-k}, (l+2)2^{-k}) : l = 0, 1, \dots, 2^k - 2\}$ to get $\int_B |f - f_B| \approx 1$ whenever $B \subset (0, 1/j)$ and $B \in \mathcal{B}^k$ for some $k \leq j$. Then,

$$\sum_{k=0}^{\infty} \sum_{B \in \mathcal{B}^k} \int_B |f - f_B| \geq C^{-1} \sum_{k=0}^{\infty} \sum_{\substack{B \in \mathcal{B}^k \\ B \subset (0, k^{-1})}} |B| \approx \sum_{k=0}^{\infty} \frac{1}{k} = \infty.$$

Next, we will provide a family of examples that show that $BV(Z)$ is, in general, strictly smaller space than $B_{1,1}^\theta(Z)$ for every $\theta \in [0, 1)$.

Example 3.12. Let $\alpha \in (\theta, 1)$. Then, the Weierstrass function

$$u_\alpha(x) = \sum_{k=1}^{\infty} \frac{\cos(2^k \pi x)}{2^{k\alpha}}, \quad x \in [0, 1],$$

is α -Hölder continuous but nowhere differentiable in $[0, 1]$ by Hardy [16]. Hence, $u_\alpha \notin BV[0, 1]$ as it would have been differentiable a.e. otherwise. Since $\alpha > \theta$, we have $\mathcal{C}^{0,\alpha}[0, 1] \subset B_{1,1}^\theta[0, 1]$ by [14, Lemma 6.2].

In conclusion, we have now proved the following theorem.

Theorem 3.13. *Let $\tau \geq 1$ and $\theta \in (0, 1]$ be arbitrary. Then,*

$$L^1(Z) = A_{1,\tau}^0(Z) \supset B_{1,1}^0(Z) \supset A_{1,\tau}^\theta(Z) \supset A_{1,\tau}^1(Z) \subset BV(Z),$$

where all but the last of the inclusions are strict in general. Furthermore, Lipschitz functions on Z belong to $B_{1,1}^0(Z)$.

We know from [17, Theorem 1.1] that $A_{1,\tau}^1(Z) \subset BV(Z)$. Note however that $A_{1,\tau}^1(Z) = BV(Z)$ holds by [17, Corollary 1.3] whenever Z supports a 1-Poincaré inequality.

4 Extension theorem for L^1 boundary data: proof of Theorem 1.2

sec:l1-extension

Given an L^1 -function on $\partial\Omega$, we will construct its BV extension in Ω using the linear extension operator for $B_{1,1}^0(\partial\Omega)$ boundary data. Observe however that the mapping $f \in L^1(\partial\Omega) \mapsto F \in BV(\Omega)$ will be nonlinear, which is not surprising in view of [27].

Instead of constructing the extension using a Whitney decomposition of Ω , we will set up a sequence of layers inside Ω whose widths depend not only on their distance from $X \setminus \Omega$, but also on the function itself (more accurately, on the choice of the sequence of Lipschitz approximations of the function in L^1 -class). Using a partition of unity subordinate to these layers, we will glue together BV extensions (from Theorem 1.1) of Lipschitz functions on $\partial\Omega$ that approximate the boundary data in $L^1(\partial\Omega)$. Roughly speaking, the closer the layer lies to $X \setminus \Omega$, the better we need the approximating Lipschitz data to be. The core idea of such a construction can be traced back to Gagliardo [13] who discussed extending $L^1(\mathbb{R}^{n-1})$ functions to $W^{1,1}(\mathbb{R}_+^n)$.

First, we approximate f in $L^1(\partial\Omega)$ by a sequence of Lipschitz continuous functions $\{f_k\}_{k=1}^\infty$ such that $\|f_{k+1} - f_k\|_{L^1(\partial\Omega)} \leq 2^{2-k}\|f\|_{L^1(\partial\Omega)}$. Note that this requirement of rate of convergence of f_k to f also ensures that $f_k \rightarrow f$ pointwise \mathcal{H} -a.e. in $\partial\Omega$. For technical reasons, we choose $f_1 \equiv 0$.

Next, we choose a decreasing sequence of real numbers $\{\rho_k\}_{k=1}^\infty$ such that:

- $\rho_1 \leq \text{diam}(\Omega)/2$;
- $0 < \rho_{k+1} \leq \rho_k/2$;
- $\sum_k \rho_k \text{LIP}(f_{k+1}, \partial\Omega) \leq C\|f\|_{L^1(\partial\Omega)}$.

These will now be used to define layers in Ω . Let

$$\psi_k(x) = \max\left\{0, \min\left\{1, \frac{\rho_k - \text{dist}(x, X \setminus \Omega)}{\rho_k - \rho_{k+1}}\right\}\right\}, \quad x \in \Omega.$$

Then, the sequence of functions $\{\psi_{k-1} - \psi_k : k = 2, 3, \dots\}$ serves as a partition of unity in $\Omega(0, \rho_2)$ subordinate to the system of layers given by $\{\Omega(\rho_{k+1}, \rho_{k-1}) : k = 2, 3, \dots\}$.

Recall that Lipschitz continuous functions lie in the Besov class $B_{1,1}^0$. Thus, we can apply the linear extension operator $E : B_{1,1}^0(\partial\Omega) \rightarrow BV(\Omega)$, whose properties were established in Section 2, to define the extension of $f \in L^1(\partial\Omega)$ by extending its Lipschitz approximations in layers, i.e.,

$$\begin{aligned} F(x) &:= \sum_{k=2}^{\infty} (\psi_{k-1}(x) - \psi_k(x)) E f_k(x) \\ &= \sum_{k=1}^{\infty} \psi_k(x) (E f_{k+1}(x) - E f_k(x)), \quad x \in \Omega. \end{aligned} \tag{4.1} \quad \boxed{\text{eq:L1-ext}}$$

The following result shows that the above extension is in the class $BV(\Omega)$ with appropriate norm bounds (see Remark 1.6).

prop:trace-L1

Proposition 4.2. *Given $f \in L^1(\partial\Omega)$, the extension defined by (4.1) satisfies*

$$\begin{aligned} \|F\|_{L^1(\Omega)} &\leq C \text{diam}(\Omega) \|f\|_{L^1(\partial\Omega)} \quad \text{and} \\ \|\text{Lip } F\|_{L^1(\Omega)} &\leq C(1 + \mathcal{H}(\Omega)) \|f\|_{L^1(\partial\Omega)}. \end{aligned}$$

Proof. Corollary 2.15 allows us to obtain the desired L^1 estimate for F . Since the extension on $B_{1,1}^0(\partial\Omega)$ is linear, we have that $E f_{k+1} - E f_k = E(f_{k+1} - f_k)$. Therefore,

$$\begin{aligned} \|F\|_{L^1(\Omega)} &\leq \sum_{k=1}^{\infty} \|\psi_k E(f_{k+1} - f_k)\|_{L^1(\Omega)} \leq \sum_{k=1}^{\infty} \|E(f_{k+1} - f_k)\|_{L^1(\Omega(0, \rho_k))} \\ &\leq C \sum_{k=1}^{\infty} \rho_k \|f_{k+1} - f_k\|_{L^1(\partial\Omega)} \leq C \rho_1 \|f\|_{L^1(\partial\Omega)} \leq C \text{diam}(\Omega) \|f\|_{L^1(\partial\Omega)}. \end{aligned}$$

In order to obtain the L^1 estimate for $\text{Lip } F$, we first apply the product rule for locally Lipschitz functions, which yields that

$$\begin{aligned} \text{Lip } F &= \sum_{k=1}^{\infty} (|E(f_{k+1} - f_k)| \text{Lip } \psi_k + \psi_k \text{Lip}(E(f_{k+1} - f_k))) \\ &\leq \sum_{k=1}^{\infty} \left(\frac{|E(f_{k+1} - f_k)| \chi_{\Omega(\rho_{k+1}, \rho_k)}}{\rho_k - \rho_{k+1}} + \chi_{\Omega(0, \rho_k)} \text{Lip}(E(f_{k+1} - f_k)) \right) \end{aligned}$$

It follows from Corollary 2.15 that

$$\begin{aligned} \sum_{k=1}^{\infty} \left\| \frac{E(f_{k+1} - f_k)}{\rho_k - \rho_{k+1}} \right\|_{L^1(\Omega(\rho_{k+1}, \rho_k))} &\leq C \sum_{k=1}^{\infty} \frac{\rho_k}{\rho_k - \rho_{k+1}} \|f_{k+1} - f_k\|_{L^1(\partial\Omega)} \\ &\leq C \sum_{k=1}^{\infty} \|f_{k+1} - f_k\|_{L^1(\partial\Omega)} \leq C \|f\|_{L^1(\partial\Omega)}. \end{aligned}$$

Next, we apply Corollary 2.11 to see that

$$\begin{aligned} \sum_{k=1}^{\infty} \|\text{Lip } E(f_{k+1} - f_k)\|_{L^1(\Omega(0, \rho_k))} &\leq C \sum_{k=1}^{\infty} \rho_k \mathcal{H}(\partial\Omega) \text{LIP}(f_{k+1} - f_k, \partial\Omega) \\ &\leq C \mathcal{H}(\partial\Omega) \sum_{k=1}^{\infty} \rho_k (\text{LIP}(f_{k+1}, \partial\Omega) + \text{LIP}(f_k, \partial\Omega)) \\ &\leq C \mathcal{H}(\partial\Omega) \|f\|_{L^1(\partial\Omega)}, \end{aligned}$$

where we used the defining properties of $\{\rho_k\}_{k=1}^{\infty}$ to obtain the ultimate inequality. Altogether, $\|\text{Lip } F\|_{L^1(\Omega)} \leq C(1 + \mathcal{H}(\partial\Omega))\|f\|_{L^1(\partial\Omega)}$. \square

subsec-L1Trace

4.3 Trace of the extended functions

In this section we complete the proof of Theorem 1.2 by showing that the trace of the extended function yields the original function back.

Proposition 4.4. *Let $F \in BV(\Omega)$ be the extension of $f \in L^1(\partial\Omega)$ as constructed in (4.1). Then,*

$$\lim_{r \rightarrow 0} \int_{B(z, r) \cap \Omega} |F - f(z)| d\mu = 0$$

for \mathcal{H} -a.e. $z \in \partial\Omega$.

Proof. Let E_0 be the collection of all $z \in \partial\Omega$ for which $\lim_k f_k(z) = f(z)$, and for $k \in \mathbb{N}$ let E_k be the collection of all $z \in \partial\Omega$ for which $TEf_k(z) = f_k(z)$ exists. Lemma 2.18 yields that $\mathcal{H}(\partial\Omega \setminus \bigcap_{k=0}^{\infty} E_k) = 0$. We define also an auxiliary sequence $\{F_n\}_{n=1}^{\infty}$ of functions approximating F by

$$F_n = \sum_{k=2}^n (\psi_{k-1} - \psi_k) E f_k + \sum_{k=n+1}^{\infty} (\psi_{k-1} - \psi_k) E f_n, \quad n \in \mathbb{N}.$$

It can be shown that $F_n \rightarrow F$ in $BV(\Omega)$, but we will not need this fact here. Note that $F_n = E f_n$ in $\Omega(0, \rho_n)$ and hence the trace of F_n exists on $\partial\Omega$ and coincides with the trace of $E f_n$, i.e., with f_n .

Fix a point $z \in \bigcap_{k=0}^{\infty} E_k$ and let $\varepsilon > 0$. Then, we can find $j \in \mathbb{N}$ such that $|f_k(z) - f(z)| < \varepsilon$ for every $k \geq j$. Next, we choose $k_0 > j$ such that $R := \rho_{k_0}$ satisfies:

- $R \text{LIP}(f_j, \partial\Omega) < \varepsilon$;
- $\int_{B(z,r)} |F_j - f_j(z)| d\mu < \varepsilon$ for every $r < R$;
- $\sum_{k=k_0}^{\infty} \rho_k \text{LIP}(f_{k+1}, \partial\Omega) < \varepsilon$.

For every $r \in (0, \rho_{k_0+1}) \subset (0, R/2)$, we can then estimate

$$\begin{aligned} \int_{B(z,r) \cap \Omega} |F - f(z)| d\mu &\leq \int_{B(z,r) \cap \Omega} |F - F_j| d\mu + \int_{B(z,r) \cap \Omega} |F_j - f_j(z)| d\mu + |f_j(z) - f(z)| \\ &\leq \int_{B(z,r) \cap \Omega} |F - F_j| d\mu + 2\varepsilon. \end{aligned} \tag{4.5} \quad \boxed{\text{eq:L1-trace}}$$

For such r , choose $k_r > k_0$ such that $\rho_{k_r+1} \leq r < \rho_{k_r}$. Then,

$$\begin{aligned} \int_{B(z,r) \cap \Omega} |F - F_j| d\mu &\leq \sum_{k=k_r}^{\infty} \int_{B(z,r) \cap \Omega} (\psi_{k-1} - \psi_k) |E(f_k - f_j)| d\mu \\ &\leq \sum_{k=k_r}^{\infty} \int_{B(z,r) \cap \Omega(\rho_{k+1}, \rho_{k-1})} |E(f_k - f_j)| d\mu \\ &\leq C \sum_{k=k_r}^{\infty} \min\{r, \rho_{k-1}\} \int_{B(z, 2^8 r) \cap \partial\Omega} |f_k - f_j| d\mathcal{H} \end{aligned} \tag{4.6} \quad \boxed{\text{eq:L1-trace-A}}$$

by Lemma 2.14. In the last inequality above, we used the fact that when $k = k_r$, we must have $B(z, r) \cap \Omega(\rho_{k_r+1}, \rho_{k_r-1}) = B(z, r) \cap \Omega(\rho_{k_r+1}, r)$ by the choice of $r < \rho_{k_r}$.

Let us, for the sake of brevity, write $U_r = B(z, 2^8 r) \cap \partial\Omega$. As $f_k - f_j$ is Lipschitz continuous, we have by the choice of j , and the fact that $k \geq j$,

$$\begin{aligned} \int_{U_r} |f_k - f_j| d\mathcal{H} &\leq \int_{U_r} |f_k - f_j - (f_k(z) - f_j(z))| d\mathcal{H} + |f_k(z) - f_j(z)| \mathcal{H}(U_r) \\ &\leq Cr \mathcal{H}(U_r) \text{LIP}(f_k - f_j, U_r) + 2\varepsilon \mathcal{H}(U_r). \end{aligned} \tag{4.7} \quad \boxed{\text{eq:L1-trace-B}}$$

Observe that $r \mathcal{H}(U_r) \approx \mu(B(z, r))$ by (1.14), (1.15), and the doubling condition for μ . Note that $\sum_{k=k_r}^{\infty} \rho_{k-1} \leq C \rho_{k_r-1} \leq CR$. Combining this with (4.6) and (4.7) gives us that

$$\begin{aligned} \int_{B(z, r) \cap \Omega} |F - F_j| d\mu &\leq \sum_{k=k_r}^{\infty} C \rho_{k-1} \mu(B(z, r)) (\text{LIP}(f_k, \partial\Omega) + \text{LIP}(f_j, \partial\Omega)) \\ &\quad + 2\varepsilon \mu(B(z, r)) \sum_{k=k_r}^{\infty} \frac{\min\{r, \rho_{k-1}\}}{r} \\ &\leq C \mu(B(z, r)) \left(\sum_{k=k_0}^{\infty} (\rho_k \text{LIP}(f_{k+1}, \partial\Omega)) + R \text{LIP}(f_j, \partial\Omega) + \varepsilon \right) \\ &\leq C \mu(B(z, r)) \varepsilon. \end{aligned}$$

Plugging this estimate into (4.5) completes the proof. \square

4.8 Summary

In conclusion, we have shown that every function in $L^1(\partial\Omega)$ has an extension to $BV(\Omega)$ in such a way that the trace of the extension returns the original function. This extension is nonlinear, but it is bounded. In a preceding section we demonstrated that there is a bounded linear extension from the subclass $B_{1,1}^0(\partial\Omega)$ to $BV(\Omega)$. Note that $B_{1,1}^0(\partial\Omega)$, containing all the Lipschitz functions on $\partial\Omega$, must necessarily be dense in $L^1(\partial\Omega)$. It therefore follows that the extension from $L^1(\partial\Omega)$ to $BV(\Omega)$ cannot be continuous on $L^1(\partial\Omega)$ (see [27] for the fact that in general any extension from $L^1(\partial\Omega)$ to $BV(\Omega)$ cannot be both bounded and linear).

References

- [A1] [1] L. Ambrosio, *Fine properties of sets of finite perimeter in doubling metric measure spaces*, Calculus of variations, nonsmooth analysis and related topics. Set-Valued Anal. **10** (2002), no. 2–3, 111–128.
- [AFP] [2] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems.*, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [AMP] [3] L. Ambrosio, M. Miranda, Jr., and D. Pallara, *Special functions of bounded variation in doubling metric measure spaces*, Calculus of variations: topics from the mathematical heritage of E. De Giorgi, 1–45, Quad. Mat., **14**, Dept. Math., Seconda Univ. Napoli, Caserta, 2004.
- [AnzGia] [4] G. Anzelotti and M. Giaquinta, *Funzioni BV e tracce*, Rendiconti del Seminario Matematico della Università di Padova, **60** (1978), 1–21. http://www.numdam.org/item?id=RSMUP_1978__60__1_0
- [BM] [5] A. Baldi and F. Montefalcone, *A note on the extension of BV functions in metric measure spaces*, J. Math. Anal. Appl. **340** (2008), no. 1, 197–208.
- [BB] [6] A. Björn and J. Björn, *Nonlinear potential theory on metric spaces*, EMS Tracts in Mathematics, 17. European Mathematical Society (EMS), Zürich, 2011. xii+403 pp.
- [BBS] [7] A. Björn, J. Björn, and N. Shanmugalingam, *Sobolev extensions of Hölder continuous and characteristic functions on metric spaces*, Canad. J. Math. **59** (2007), no. 6, 1135–1153.
- [BS] [8] A. Björn, J. Björn, and N. Shanmugalingam, *Poincaré inequalities, uniform domains and extension properties for Newton–Sobolev functions in metric spaces*, J. Math. Anal. Appl. **332** (2007), no. 1, 190–208.
- [BourPaj] [9] M. Bourdon and H. Pajot, *Cohomologie l_p et espaces de Besov*, J. Reine Angew. Math. **558** (2003), 85–108.
- [BBM] [10] J. Bourgain, H. Brezis, and P. Mironescu, *Another look at Sobolev spaces*, Optimal Control and Partial Differential Equations, in honour of Professor A. Bensoussan’s 60th birthday, J. L. Menaldi et al. (eds.), IOS Press, Amsterdam (2001), 439–455.

- [Brez] [11] H. Brezis, *How to recognize constant functions. A connection with Sobolev spaces*, Uspekhi Mat. Nauk **57** (2002), no. 4, 59–74 (in Russian); English transl.: Russian Math. Surveys **57** (2002), 693–708.
- [EvaG92] [12] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions* Studies in Advanced Mathematics series, CRC Press, Boca Raton, 1992, viii+268 pp.
- [Gag] [13] E. Gagliardo, *Caratterizzazioni delle tracce sulla frontiera relative ad alcune classi di funzioni in n variabili*, Ren. Sem. Mat. Univ. Padova, **27** (1957) 284–305.
- [GKS] [14] A. Gogatishvili, P. Koskela, and N. Shanmugalingam, *Interpolation properties of Besov spaces defined on metric spaces*, Math. Nachr., **283** no. 2 (2010) 215–231.
- [HaKiLa] [15] H. Hakkarainen, J. Kinnunen, and P. Lahti, *Regularity of minimizers of the area functional in metric spaces*, Adv. Calc. Var. **8** (2015), no. 1, 55–68.
- [Har] [16] G. H. Hardy, *Weierstrass’s non-differentiable function*, Trans. Amer. Math. Soc. **17** (1916), 301–325.
- [HKT] [17] T. Heikkinen, P. Koskela, and H. Tuominen, *Sobolev-type spaces from generalized Poincaré inequalities*, Studia Math. **181** (2007), no. 1, 1–16.
- [HKST] [18] J. Heinonen, P. Koskela, N. Shanmugalingam, and J. Tyson, *Sobolev spaces on metric measure spaces: an approach based on upper gradients*, New Mathematical Monographs **27**, Cambridge University Press, 2015. xi+448 pp.
- [Hei01] [19] J. Heinonen, *Lectures on analysis on metric spaces*, Universitext. Springer-Verlag, New York, 2001. x+141 pp.
- [HK] [20] J. Heinonen and P. Koskela, *Quasiconformal maps in metric spaces with controlled geometry*, Acta Math. **181** (1998), no. 1, 1–61.
- [JN] [21] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426.
- [L] [22] P. Lahti, *Extensions and traces of functions of bounded variation on metric spaces*, J. Math. Anal. Appl. **423** (2015), no. 1, 521–537.

- [LS] [23] P. Lahti and N. Shanmugalingam, *Trace theorems for functions of bounded variation in metric setting*, in preparation. Preprint available at [arXiv:1507.07006](https://arxiv.org/abs/1507.07006).
- [Maz] [24] V. G. Maz'ya, *Sobolev spaces*, Translated from the Russian by T. O. Shaposhnikova. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1985. xix+486 pp.
- [M-Sr] [25] M. Miranda Sr., *Comportamento delle successioni convergenti di frontiere minimali*, Rend. Sem. Mat. Univ. Padova, **38** (1967), 238–257.
- [M] [26] M. Miranda Jr., *Functions of bounded variation on “good” metric spaces*, J. Math. Pures Appl. (9) **82** (2003), no. 8, 975–1004.
- [Pee] [27] J. Peetre, *A counterexample connected with Gagliardo’s trace theorem*, Comment. Math., **2** (1979) 277–282.
- [PW] [28] A. Pełczyński and M. Wojciechowski, *Sobolev spaces in several variables in L^1 -type norms are not isomorphic to Banach lattices*, Ark. Mat., **40** (2002) 363–382.
- [S] [29] N. Shanmugalingam, *Newtonian spaces: An extension of Sobolev spaces to metric measure spaces*, Rev. Mat. Iberoamericana **16**(2) (2000), 243–279.
- [Vit] [30] D. Vittone, *Lipschitz surfaces, perimeter and trace theorems for BV functions in Carnot–Carathéodory spaces*, to appear in Ann. Sc. Norm. Super. Pisa Cl. Sci.
- [Zie89] [31] W. P. Ziemer, *Weakly differentiable functions. Sobolev spaces and functions of bounded variation*, Graduate Texts in Mathematics, 120. Springer-Verlag, New York, 1989.

Address:

L. Malý: Department of Mathematical Sciences, University of Cincinnati, Cincinnati, OH 45221-0025, U.S.A.

malyls@ucmail.uc.edu

N. Shanmugalingam: Department of Mathematical Sciences, University of Cincinnati, Cincinnati, OH 45221-0025, U.S.A.

shanmun@ucmail.uc.edu

M. Snipes: Department of Mathematics, Kenyon College, Gambier, OH 43022, U.S.A.

snipesm@kenyon.edu