

An extremal eigenvalue problem for the Wentzell-Laplace operator

M. Dambrine, D. Kateb, J. Lamboley

September 26, 2014

Abstract: We consider the question of giving an upper bound for the first nontrivial eigenvalue of the Wentzell-Laplace operator of a domain Ω , involving only geometrical informations. We provide such an upper bound, by generalizing Brock’s inequality concerning Steklov eigenvalues, and we conjecture that balls maximize the Wentzell eigenvalue, in a suitable class of domains, which would improve our bound. To support this conjecture, we prove that balls are critical domains for the Wentzell eigenvalue, in any dimension, and that they are local maximizers in dimension 2 and 3, using an order two sensitivity analysis. We also provide some numerical evidence.

Keywords: Wentzell eigenvalues, eigenvalue estimates, Shape optimization, Shape derivatives, Stability, Quantitative isoperimetric inequality.

AMS Subject Classification: *Primary* 35P15; *Secondary* 49K20, 49K40.

Contents

1	Introduction	2
2	Upper bound for $\lambda_{1,\beta}$	8
2.1	Preliminary remarks and results.	8
2.2	Proof of Theorem 1.1	10
2.3	On the sharpness of the upper bounds.	14
3	First order shape calculus	15
3.1	Notations and preliminary result for shape deformation	16
3.2	Regularity of the eigenfunctions and eigenvalues with respect to the parameter	18
3.3	Shape derivative of simple eigenvalues of the Wentzell-Laplace problem	22
3.4	Shape derivative of multiple eigenvalues of the Wentzell-Laplace problem	22
3.4.1	The general result	22
3.4.2	The case of balls	23
3.5	Numerical illustrations	25
4	Testing if the ball is a local maximum for λ_1: second order arguments	26
4.1	Construction of the matrix E of the second derivatives	27
4.2	Computation of the trace	28

A	Some classical results on tangential differential calculus	31
A.1	Notations and definitions. Preliminary results	31
A.2	A commutation lemma	32
B	Spherical harmonics	33
C	Intermediate results for the second shape derivative matrix	35
D	Computing u'	41
D.1	Explicit resolution of (58) to compute \tilde{u}_j	42
D.1.1	Explicit representation of \tilde{u}_j in the case $d = 2$	42
D.1.2	Explicit representation of \tilde{u}_j in the case $d = 3$	43
D.2	The explicit expression of the trace of $E^{(1)}$	44
E	Shape Derivatives of Steklov and Laplace-Beltrami eigenvalues problem	46

1 Introduction

Background. Let $d \geq 2$ and Ω be a bounded domain in \mathbb{R}^d (i.e a bounded connected open set) supposed to be sufficiently smooth (of class C^3), and we denote Δ_τ the Laplace-Beltrami operator on $\partial\Omega$. Motivated by generalized impedance boundary conditions, we consider the eigenvalue problem for Wentzell boundary conditions

$$\begin{cases} -\Delta u = 0 & \text{in } \Omega \\ -\beta\Delta_\tau u + \partial_n u = \lambda u & \text{on } \partial\Omega \end{cases} \quad (1)$$

where β is a given real number and ∂_n denotes the outward unit normal derivative.

The coefficient β appears as a surface diffusion coefficient arising in a passage to the limit in the thickness of the boundary layer for coated object (see [22, 1, 16]). A general derivation of Wentzell boundary conditions can be found in [15]. The coefficient can be either positive or negative. We first consider the case $\beta \geq 0$ where the obtained boundary value problem is coercive.

This problem couples surface and volume effects through the Steklov eigenvalue problem in Ω with the Laplace-Beltrami eigenvalue problem on $\partial\Omega$. Let us recall some known facts about these two problems. The **Steklov eigenvalue problem** consists in solving

$$\begin{cases} \Delta u = 0 & \text{in } \Omega \\ \partial_n u = \lambda^S u & \text{on } \partial\Omega \end{cases} \quad (2)$$

It has a discrete spectrum consisting in a sequence

$$\lambda_0^S(\Omega) = 0 < \lambda_1^S(\Omega) \leq \lambda_2^S(\Omega) \dots \rightarrow +\infty$$

where the λ^S are called Steklov eigenvalues. Brock-Weinstock inequality states that λ_1^S is maximized by the ball among all open sets of fixed volume $|\Omega|$. It was first proved in the case $d = 2$ by Weinstock and extended by Brock to any dimension in [6] (Weinstock inequality is slightly stronger but restricted to simply-connected domains: he proved indeed that the disk maximizes λ_1^S among simply-connected sets of given perimeter). A quantitative form of this inequality was recently obtained by Brasco, De Philippis and Ruffini who proved in [5] that

$$\lambda_1^S(\Omega) \leq \lambda_1^S(B) \left[1 - \delta_d \left(\frac{|\Omega \Delta B(x_{\partial\Omega})|}{|\Omega|} \right)^2 \right],$$

where δ_d is an explicit nonnegative constant depending only on d , $x_{\partial\Omega}$ is the center of mass of $\partial\Omega$ and $B(x_{\partial\Omega})$ is the ball centered in $x_{\partial\Omega}$ with volume $|\Omega|$ ¹. Let us emphasize that no additional topological assumption is needed.

It is well-known that the **spectrum of the Laplace-Beltrami operator** on $\partial\Omega$, that is numbers λ such that the equation $-\Delta_\tau u = \lambda u$ on $\partial\Omega$ has nontrivial solutions, is also discrete and satisfies:

$$\lambda_0^{LB}(\partial\Omega) = 0 < \lambda_1^{LB}(\partial\Omega) \leq \lambda_2^{LB}(\partial\Omega) \dots \rightarrow +\infty$$

Again, one can ask if λ_1^{LB} takes its maximal value on the euclidean sphere, among hypersurfaces of fixed $(d-1)$ -dimensional volume. Here, the answer is more complicated than for the Steklov problem. It depends on both the topology of the surface and the dimension. In [19], Hersch gave a positive answer if $d = 3$ for surfaces homomorphic to the euclidean sphere. In the cases $d > 3$ or without topological restriction, the answer is negative (see [3, 10, 11], and Section 2.1 for the 2-dimensional case)).

When $\beta \geq 0$, the **spectrum of the Laplacian with Wentzell conditions** consists in an increasing countable sequence of eigenvalues

$$\lambda_{0,\beta}(\Omega) = 0 < \lambda_{1,\beta}(\Omega) \leq \lambda_{2,\beta}(\Omega) \dots \rightarrow +\infty \quad (3)$$

with corresponding real orthonormal (in $L^2(\partial\Omega)$) eigenfunctions u_0, u_1, u_2, \dots . As in the previous cases, the first eigenvalue is zero with constants as corresponding eigenfunctions. As usual, we adopt the convention that each eigenvalue is repeated according to its multiplicity. Hence, the first eigenvalue of interest is $\lambda_{1,\beta}$. A variational characterization of the eigenvalues is available: we introduce the Hilbert space

$$H(\Omega) = \{u \in H^1(\Omega), \text{Tr}_{\partial\Omega}(u) \in H^1(\partial\Omega)\},$$

where $\text{Tr}_{\partial\Omega}$ is the trace operator, and we define on $H(\Omega)$ the two bilinear forms

$$A_\beta(u, v) = \int_\Omega \nabla u \cdot \nabla v \, dx + \beta \int_{\partial\Omega} \nabla_\tau u \cdot \nabla_\tau v \, d\sigma, \quad B(u, v) = \int_{\partial\Omega} uv, \quad (4)$$

where ∇_τ is the tangential gradient. Since we assume β is nonnegative, the two bilinear forms are positive and the variational characterization for the k -th eigenvalue is

$$\lambda_{k,\beta}(\Omega) = \min \left\{ \frac{A_\beta(v, v)}{B(v, v)}, v \in H(\Omega), \int_{\partial\Omega} v u_i = 0, i = 0, \dots, k-1 \right\} \quad (5)$$

In particular, when $k = 1$, the minimum is taken over the functions orthogonal to the eigenfunctions associated to $\lambda_{0,\beta} = 0$, i.e constant functions. To describe this spectrum, one can notice that the eigenvalue problem can be rewritten purely on $\partial\Omega$ as:

$$-\beta \Delta_\tau u + \text{D}u = \lambda u$$

where D denotes the Dirichlet-to-Neumann map, that is a selfadjoint, positive pseudodifferential operator of order one. Therefore, this problem can be seen as a compact perturbation of the usual Laplace-Beltrami operator. This point of view was used in [4] and justify that high order eigenvalues of the Laplace-Wentzell problem look like those of the Laplace-Beltrami operator.

¹The results in [5] are stated with the Fraenkel asymmetry, meaning that the previous inequality is stated for the ball B of volume $|\Omega|$ that minimizes $|\Omega \Delta B|$, but from the proof (see [5, Section 5]) we can conclude that the ball $B(x_{\partial\Omega})$ of volume $|\Omega|$ and such that $\int_{\partial\Omega} (x - x_{\partial\Omega}) d\sigma = 0$ is in fact valid as well.

However, we are interested in this work, in studying low order eigenvalues and more precisely in giving an upper bound for the second eigenvalue $\lambda_{1,\beta}$ involving only geometrical informations. Please remark that we are not seeking for lower bound, because even with very strong geometrical assumption, there is none. Indeed, a consequence of our results is that

$$\inf \{ \lambda_{1,\beta}(\Omega), \Omega \text{ convex}, |\Omega| = m \} = 0 \quad (6)$$

for any value of $\beta \geq 0$ and $m \geq 0$, see Remark 2.5. An important remark at this point is that the bilinear form A_β is not homogeneous with respect to dilatation of the domain. Therefore, the volume of Ω plays a crucial role in $\lambda_{1,\beta}$. As a surface term appears also in A_β (corresponding to the Laplace-Beltrami operator), the perimeter of Ω (i.e. the volume of $\partial\Omega$) should also play a crucial role.

Notice that when $\beta = 0$ we retrieve the Steklov eigenvalues, and we recover the Laplace-Beltrami eigenvalues by considering $\frac{1}{\beta}\lambda_{1,\beta}$ and letting β go to $+\infty$, see Section 2.1.

Note also that the close but distinct eigenvalue problem

$$\begin{cases} -\Delta u = \lambda u & \text{in } \Omega \\ \Delta u + \alpha \partial_n u + \gamma u = 0 & \text{on } \partial\Omega \end{cases} \quad (7)$$

was considered by J.B. Kennedy in [21]. He transforms this problem into a Robin type problem to prove a Faber-Krahn type inequality when the constants α, γ are non negative: the ball is the best possible domain among those of given volume.

The results of the paper. We first apply the strategy of F. Brock for the Steklov eigenvalue problem to the Wentzell eigenvalue problem and obtain a first upper bound of $\lambda_{1,\beta}(\Omega)$ in terms of purely geometric quantities (we actually provide a refined version, using [5]):

Theorem 1.1 *Let Ω a smooth set such that $\int_{\partial\Omega} x = 0$. Let $\Lambda[\Omega]$ be the spectral radius of the symmetric and positive semidefinite matrix $P(\Omega) = (p_{ij})_{i,j=1,\dots,d}$ defined as*

$$p_{ij} = \int_{\partial\Omega} (\delta_{ij} - n_i n_j), \quad (8)$$

where \mathbf{n} is the outward normal vector to $\partial\Omega$. Then if $\beta \geq 0$, one has:

$$S(\Omega) := \sum_{i=1}^d \frac{1}{\lambda_{i,\beta}(\Omega)} \geq \frac{\int_{\partial\Omega} |x|^2}{|\Omega| + \beta \Lambda[\Omega]} \geq \frac{d \omega_d^{-1/d} |\Omega|^{\frac{d+1}{d}}}{|\Omega| + \beta \Lambda[\Omega]} \left[1 + \gamma_d \left(\frac{|\Omega \Delta B|}{|B|} \right)^2 \right]. \quad (9)$$

where

$$\gamma_d = \frac{d+1}{d} \frac{2^{1/d} - 1}{4}, \quad (10)$$

$\omega_d = |B_1|$ and B is the ball of volume $|\Omega|$ and with the same center of mass than $\partial\Omega$. Equality holds in (9) if Ω is a ball.

A trivial consequence of Theorem 1.1 is the following upper bound for $\lambda_{1,\beta}(\Omega)$.

Corollary 1.2 *With the same notations as in Theorem 1.1, if $\beta \geq 0$, it holds:*

$$\lambda_{1,\beta}(\Omega) \leq d \frac{|\Omega| + \beta \Lambda[\Omega]}{\int_{\partial\Omega} |x|^2} \leq \frac{|\Omega| + \beta \Lambda[\Omega]}{\omega_d^{-1/d} |\Omega|^{\frac{d+1}{d}} \left[1 + \gamma_d \left(\frac{|\Omega \Delta B|}{|B|} \right)^2 \right]}. \quad (11)$$

where B and γ_d are as in Theorem 1.1. Equality holds in (11) if Ω is a ball.

Note that the method used for the Wentzell eigenvalue problem also applies for the Laplace-Beltrami case and provides an upper bound for λ_1^{LB} without any topological assumption on Ω .

Theorem 1.3 *With the same notations as in Theorem 1.1, it holds*

$$S^{LB}(\partial\Omega) := \sum_{i=1}^d \frac{1}{\lambda_i^{LB}(\partial\Omega)} \geq \frac{\int_{\partial\Omega} |x|^2}{\Lambda[\Omega]} \geq \frac{d\omega_d^{-1/d} |\Omega|^{\frac{d+1}{d}}}{\Lambda[\Omega]} \left[1 + \gamma_d \left(\frac{|\Omega \Delta B|}{|B|} \right)^2 \right]. \quad (12)$$

and

$$\lambda_1^{LB}(\partial\Omega) \leq d \frac{\Lambda[\Omega]}{\int_{\partial\Omega} |x|^2} \leq \frac{\Lambda[\Omega]}{\omega_d^{-1/d} |\Omega|^{\frac{d+1}{d}} \left[1 + \gamma_d \left(\frac{|\Omega \Delta B|}{|B|} \right)^2 \right]}. \quad (13)$$

Equality holds in (12) and (13) if Ω is a ball.

It is expected in this type of extremal eigenvalue problem that ball are maximizers. We are not able to fully justify the natural following conjecture:

Conjecture: The ball maximizes the first non-trivial Wentzell-Laplace eigenvalue among smooth open sets of given volume and which are homeomorphic to the ball.

The topological restriction is motivated by the limit case $\beta \rightarrow +\infty$ as we noticed before (see also Section 2.1). In Section 2.2, we observe that the intermediate bound in (11) has both its numerator and denominator that are minimized by the ball, under volume constraint, so there is a competition. In Section 2.3 we observe that in fact, the ball does not minimize this bound in general (see Figure 2.3). Therefore, we can not deduce from this bound the maximality of balls (though it might work for certain values of β and the volume constraint). About the upper bound (11), we show that it is larger than $\lambda_{1,\beta}(B)$ for every $\beta > 0$ (with equality for the ball) and hence again does not implies that balls are maximizing $\lambda_{1,\beta}$. To check if balls are relevant candidates for maximizers in our case, we then turn our attention to a shape sensitivity analysis of $\lambda_{1,\beta}$.

Therefore, we first wonder if the ball is a critical shape in any dimension. With respect to shape sensitivity, the main difficulty is to handle multiple eigenvalues which leads to a nonsmooth dependency of $\lambda_{1,\beta}$ with respect to Ω . However, for a fixed deformation field $\mathbf{V} \in W^{3,\infty}(\Omega, \mathbb{R}^d)$, along the transport of Ω by $T_t = I + t\mathbf{V}$, we prove the existence of smooth branches of eigenvalues and eigenfunctions associated to the subspace generated by the group of eigenvalues and provide a characterization of the derivative along the branches: $\lambda_{1,\beta}$ is then the minimum value among these d smooth branches.

Theorem 1.4 *We distinguish the case of simple and multiple eigenvalue.*

- If $\lambda = \lambda_{k,\beta}(\Omega)$ is a simple eigenvalue of the Wentzell problem, then the application $t \mapsto \lambda(t) = \lambda_{k,\beta}(\Omega_t)$ (where $\Omega_t = (I + t\mathbf{V})(\Omega)$) is differentiable and the derivative at $t = 0$ is

$$\lambda'(0) = \int_{\partial\Omega} V_n \left(|\nabla_\tau u|^2 - |\partial_n u|^2 - \lambda H |u_0|^2 + \beta (H I_d - 2D^2b) \nabla_\tau u \cdot \nabla_\tau u \right) d\sigma.$$

where u is the normalized eigenfunction associated to λ , D^2b is the Hessian of the signed distance function (see (48)), $H = \text{Tr}(D^2b)$ is the mean curvature of $\partial\Omega$, I_d is the identity matrix of size

d , and $V_n = \mathbf{V} \cdot \mathbf{n}_{\partial\Omega}$ is the normal component of the deformation. Moreover, the shape derivative u' at $t = 0$ of the eigenfunction satisfies

$$\begin{cases} \Delta u' = 0 & \text{in } \Omega, \\ -\beta \Delta_\tau u' + \partial_n u' - \lambda u' = \beta \Delta_\tau (V_n \partial_n u) + \beta \operatorname{div}_\tau (V_n (H I_d - 2D^2 b) \nabla_\tau u) \\ \quad + \operatorname{div}_\tau (V_n \nabla_\tau u) - \lambda' u + \lambda V_n (\partial_n u + H u) & \text{on } \partial\Omega. \end{cases} \quad (14)$$

- Let λ be a multiple eigenvalue of order $m \geq 2$. Let $(u_k)_{k=1, \dots, m}$ denote the eigenfunctions associated to λ . Then there exist m functions $t \mapsto \lambda_{k,\beta}(t)$, $k = 1, \dots, m$ defined in a neighborhood of 0 such that

- $\lambda_{k,\beta}(0) = \lambda$,
- for every t in a neighborhood of 0, $\lambda_{k,\beta}(t)$ is an eigenvalue of $\Omega_t = (I + t\mathbf{V})(\Omega)$,
- the functions $t \mapsto \lambda_{k,\beta}(t)$, $k = 1, \dots, m$ admit derivatives and their values at 0 are the eigenvalues of the $m \times m$ matrix $M = M_\Omega(V_n)$ of entries (M_{ij}) defined by

$$M_{ij} = \int_{\partial\Omega} V_n \left(\nabla_\tau u_i \cdot \nabla_\tau u_j - \partial_n u_i \partial_n u_j - \lambda H u_i u_j + \beta (H I_d - 2D^2 b) \nabla_\tau u_i \cdot \nabla_\tau u_j \right) d\sigma.$$

Notice that in the notations above and contrary to (3), the functions $\lambda_k(t)$ are no longer ordered. As a byproduct of this result, notice that we can write the corresponding shape derivatives for the Steklov and Laplace-Beltrami eigenvalue problem (see Appendix E). Another consequence of this result, regarding our conjecture, is that we are able to check that balls are critical shapes for $\lambda_{1,\beta}$ by computing the trace of the previously defined matrix $M = M_B$ (recall that $\lambda_{1,\beta}(B)$ is an eigenvalue of multiplicity d the dimension). But first, we make a short remark about the notion of volume preserving deformation:

Remark 1.5 *In the next results and in many places in the paper, we will consider volume preserving smooth deformations of domains, that is to say $\Omega_t = T_t(\Omega)$ where $t \mapsto T_t$ satisfies:*

- $T_0 = Id$,
- for every t near 0, T_t is a $W^{3,\infty}$ -diffeomorphism from Ω onto its image $\Omega_t = T_t(\Omega)$,
- the application $t \mapsto T_t$ is real-analytic near $t = 0$.
- for every t near 0, $|\Omega_t| = |\Omega|$.

More generally, it can be sufficient to assume that the volume is preserved at the first or the second order, depending on whether we are interested in first or second order conditions. For example, if one considers $T_t = I + t\mathbf{V}$ the vector field \mathbf{V} is said to be volume preserving at first order if it satisfies $\int_{\partial\Omega} V_n d\sigma = 0$; indeed for $\Omega_t = (I + t\mathbf{V})(\Omega)$, we have $\frac{d}{dt}|_{t=0} |\Omega_t| = \int_{\partial\Omega} V_n d\sigma$.

When dealing with second order considerations as in Theorem 1.7, we need that the volume is preserved at the second order, so T_t is volume preserving at second order if

$$\frac{d^2}{dt^2} |\Omega_t|_{t=0} = \int_{\partial\Omega} (W + V_n \partial_n V_n + H V_n^2) d\sigma = 0,$$

where $\mathbf{V} = \frac{1}{t}(T_t - I)$, V_n is the value at $t = 0$ of $\mathbf{V} \cdot \mathbf{n}_{\partial\Omega_t}$, and W denotes the derivative of $\mathbf{V} \cdot \mathbf{n}_{\partial\Omega_t}$ with respect to t at $t = 0$.

Proposition 1.6 Any ball B is a critical shape for $\lambda_{1,\beta}$ with volume constraint, in the sense that for every volume preserving deformations \mathbf{V} ,

$$\text{Tr}(M_B(V_n)) = \sum_{k=1}^d \lambda'_{k,\beta}(0) = 0,$$

where $(t \mapsto \lambda_{k,\beta}(t))_{k=1\dots d}$ are defined in Theorem 1.4.

In particular, $0 \in \partial\lambda_{1,\beta}(B; V_n) := [\inf_{i=1\dots d} \lambda'_{i,\beta}(0), \sup_{i=1\dots d} \lambda'_{i,\beta}(0)]$ the directional subdifferential associated to the first non trivial eigenvalue.

Moreover, this subdifferential reduces to $\{0\}$ if V_n is orthogonal to spherical harmonics of order two: in other words, in that case, the directional derivative exists in the usual sense and vanishes.

Two situations can now occur: either the subdifferential in direction V_n is not reduced to $\{0\}$ and then one can deduce from the previous statement that B locally maximizes $\lambda_{1,\beta}$ along $t \mapsto B_t$ (see for example (c) and (d) in Figure 5), or the subdifferential in direction V_n is $\{0\}$ and then this first order shape calculus does not allow us to conclude that the ball is a local maximizer of $\lambda_{1,\beta}$. Hence, for the directions V_n in \mathcal{H} defined as the **Hilbert space generated by spherical harmonics of order greater or equal to three**, we now consider the second order analysis to wonder if the ball satisfies the second order necessary condition of optimality, and obtain the following result in dimension two and three.

Theorem 1.7 Let B be a ball of radius R in \mathbb{R}^2 or \mathbb{R}^3 and $t \mapsto B_t = T_t(B)$ a second order volume preserving deformation. $\lambda_{1,\beta}(B)$ is an eigenvalue of multiplicity d the dimension, and we denote $t \mapsto \lambda_{k,\beta}(t)$, $k = 1, \dots, d$ the branches obtained in Theorem 1.4.

Then the functions $t \mapsto \lambda_{k,\beta}(t)$, $k = 1, \dots, d$ admit a second derivative and their values at 0 are the eigenvalues of the $d \times d$ matrix $E = E_B(V_n)$ defined in Section 4. Moreover, there exists a nonnegative number $\mu(= \mu(\beta))$ independent of the radius R such that

$$\text{Tr}(E_B(V_n)) = \sum_{k=1}^d \lambda''_{k,\beta}(0) \leq -\mu K(R) \int_{\partial B} (|\nabla_\tau V_n|^2 + |V_n|^2) d\sigma = -\mu K(R) \|V_n\|_{H^1(\partial B)}^2.$$

holds for or all $V_n \in \mathcal{H}$, with $K(R) = \frac{d}{R^{2+d}\omega_{d-1}}$.

As a consequence of Proposition 1.6 and Theorem 1.7, we have the result:

Corollary 1.8 If B is a ball in \mathbb{R}^2 or \mathbb{R}^3 , and $t \mapsto T_t \in W^{3,\infty}(B, \mathbb{R}^d)$ a smooth (second order) volume preserving deformation, then

$$\lambda_{1,\beta}(B) \geq \lambda_{1,\beta}(T_t(B)), \quad \text{for } t \text{ small enough.}$$

Plan of the paper. The paper is organized as follows: in section 2, we prove Theorem 1.1 by adapting the strategy of Brock and present some numerical tests to illustrate the sharpness of the upper bound. The first order shape analysis is presented in section 3, while the second order shape analysis is presented in section 4. The background material for shape calculus and the proofs of technical intermediary results are postponed to the annexes.

2 Upper bound for $\lambda_{1,\beta}$

2.1 Preliminary remarks and results.

Let us start by a few remarks on the proofs in the two limit cases $\beta \rightarrow +\infty$ (that is the Laplace-Beltrami eigenvalue problem), and $\beta = 0$ (that is the Steklov eigenvalue problem).

On the Laplace-Beltrami case:

The case $d = 2$ is trivial: it suffices to argue on each connected component of $\partial\Omega$. We introduce $\gamma : [0, L]$ a parametrization by the arclength of a connected component Γ of $\partial\Omega$, then for any $u \in H^1(\partial\Omega)$, the Rayleigh quotient can be written as

$$\frac{\int_{\Gamma} |\nabla_{\tau} u|^2}{\int_{\Gamma} u^2} = \frac{\int_0^L [(u \circ \gamma)']^2}{\int_0^L (u \circ \gamma)^2}.$$

Hence, the $\lambda_1^{LB}(\Gamma)$ is nothing but the infimum of $\|u'\|_{L^2(0,L)}^2$ among periodic functions u with 0 mean value and $\|u\|_{L^2(0,L)} = 1$, that is to say $4\pi^2/L^2$. It is a decreasing function of the length of the connected component of the boundary. Then, if Ω is simply connected, combined with the isoperimetric inequality, the previous computations leads to $\lambda_1^{LB}(\partial\Omega) \leq \lambda_1^{LB}(\partial B)$ where B is a disk of same area than Ω .

Moreover, if $\partial\Omega$ has more than one connected component, then $\lambda_1^{LB} = 0$ since the multiplicity of 0 as eigenvalue is at least the number of connected component. To check that claim, it suffices to check that the functions taking the value 1 on one of the connected component and 0 elsewhere are independent eigenfunctions associated to the eigenvalue 0. We conclude that in dimension 2, $\lambda_1^{LB}(\partial\Omega) \leq \lambda_1^{LB}(\partial B)$, where B is a disk of same area than Ω .

The case $d = 3$ is more complex. There is a classical result of J. Hersch [19]: if $\Omega \subset \mathbb{R}^3$ is homeomorphic to the ball, then

$$\lambda_1^{LB}(\partial\Omega) \leq \lambda_1^{LB}(\partial B), \text{ for all } \Omega \text{ such that } |\partial\Omega| = |\partial B|. \quad (15)$$

We first extend Hersch statement to domains of same volume by a classic homogeneity argument.

Lemma 2.1 *If $\Omega \subset \mathbb{R}^3$ is homeomorphic to the ball, then*

$$\lambda_1^{LB}(\partial\Omega) \leq \lambda_1^{LB}(\partial B) \text{ if } |\Omega| = |B|.$$

Proof of Lemma 2.1.:

One easily checks that $\Omega \mapsto \lambda_1^{LB}(\partial\Omega)$ is homogeneous of degree -2 , so $\Omega \mapsto \lambda_1^{LB}(\Omega)|\partial\Omega|^{2/(d-1)}$ is homogeneous of degree 0. Then we get from Hersch's inequality (15), that

$$\lambda_1^{LB}(\partial\Omega)|\partial\Omega|^{\frac{2}{d-1}} \leq \lambda_1^{LB}(\partial B)|\partial B|^{\frac{2}{d-1}}, \text{ for all } \Omega \text{ such that } |\partial\Omega| = |\partial B|. \quad (16)$$

Thanks to the invariance by translation of λ_1^{LB} and the perimeter, and using the 0-homogeneity of the previous product, we get that the previous inequality is in fact valid for any ball B and any domain Ω . We combine with the isoperimetric inequality

$$\frac{|\partial B|^{\frac{d}{d-1}}}{|B|} \leq \frac{|\partial\Omega|^{\frac{d}{d-1}}}{|\Omega|}$$

to conclude. ■

On the Steklov case:

In the general case $\beta \geq 0$, we will adapt the original Brock's proof; the main tool is an isoperimetric inequality for the moment of inertia of the boundary $\partial\Omega$ with respect to the origin. The general form of the weighted isoperimetric inequality due to F. Betta, F. Brock, A. Mercaldo and M.R. Posteraro [2] is:

Lemma 2.2 *Let $\Omega \subset \mathbb{R}^d$ be an open set and let f be a continuous, nonnegative and nondecreasing function defined on $[0, \infty]$. Moreover, we suppose that*

$$t \mapsto \left(f(t^{\frac{1}{d}}) - f(0) \right) t^{1-\frac{1}{d}} \text{ is convex for } t \geq 0$$

Then

$$\int_{\partial\Omega} f(|x|) d\sigma \geq f(R) |\partial B_R|, \tag{17}$$

where B_R is the ball centered at the origin such that $|B_R| = |\Omega|$.

Let us remark that the function $t \mapsto t^p$ satisfies the assumptions of the lemma as soon as $p \geq 1$ and in particular for $p = 2$. In that case and in order to prove a refinement of Brock's inequality, L. Brasco, G. De Philippis and B. Ruffini established a qualitative refinement of this inequality (Theorem B of [5]):

Lemma 2.3 *There exists an explicit dimensional constant γ_d such that for every bounded, open Lipschitz set Ω in \mathbb{R}^d ,*

$$\int_{\partial\Omega} |x|^2 d\sigma \geq R^2 |\partial B_R| \left[1 + \gamma_d \left(\frac{|\Omega \Delta B_R|}{|B_R|} \right)^2 \right], \tag{18}$$

where B_R is the ball centered at the origin such that $|B_R| = |\Omega|$ and γ_d is the constant defined in (10).

On the Wentzell case:

An important remark for the sequel is the particular case when Ω is a ball B_R of radius R . The eigenspace corresponding to $\lambda_{1,\beta}$ is d -dimensional: it consists to the restrictions on the sphere S_R^{d-1} of the linear functions in \mathbb{R}^d spanned by the coordinates functions. It follows, from the theory of spherical harmonic functions that

$$\lambda_{1,\beta}(B_R) = \lambda_{2,\beta}(B_R) = \dots = \lambda_{d,\beta}(B_R) = \frac{(d-1)\beta + R}{R^2}. \tag{19}$$

The Laplace-Beltrami operator on ∂B_R and the Steklov operator also are diagonal on the basis of spherical harmonics, hence

$$\lambda_{1,\beta}(B_R) = \lambda_1^S(B_R) + \beta \lambda_1^{LB}(\partial B_R),$$

and more generally the eigenvalue associated to spherical harmonics of order l is

$$\lambda_{(l)}(B_R) = \frac{l(l+d-2)\beta + R}{R^2}. \tag{20}$$

But, this situation is specific to the ball: indeed, in general we only have the inequality

$$\lambda_{1,\beta}(\Omega) \geq \lambda_1^S(\Omega) + \beta \lambda_1^{LB}(\Omega).$$

Moreover, we can easily prove that for any smooth Ω , $\lim_{\beta \rightarrow \infty} \frac{1}{\beta} \lambda_{1,\beta}(\Omega) = \lambda_1^{LB}(\Omega)$: indeed, we have a first trivial inequality $\frac{1}{\beta} \lambda_{1,\beta}(\Omega) \geq \lambda_1^{LB}(\Omega)$ for any $\beta \geq 0$, and using the variational formulation (5), we obtain $\forall v \in H(\Omega)$ with the addition condition $\int_{\partial\Omega} v = 0$,

$$\overline{\lim}_{\beta \rightarrow \infty} \frac{1}{\beta} \lambda_{1,\beta}(\Omega) \leq \overline{\lim}_{\beta \rightarrow \infty} \frac{\frac{1}{\beta} \int_{\Omega} |\nabla v|^2 + \int_{\partial\Omega} |\nabla_{\tau} v|^2}{\int_{\partial\Omega} v^2} = \frac{\int_{\partial\Omega} |\nabla_{\tau} v|^2}{\int_{\partial\Omega} v^2}$$

which leads to the result.

For example if $d = 3$, combining Brock's inequality and Lemma 2.1, we obtain that the right-hand side in the previous inequality is maximized by the ball, among domains of given volume and homeomorphic to the ball. Unfortunately, this is not enough to obtain that balls are maximizing the Wentzell eigenvalue.

So in order to obtain an estimate of $\lambda_{1,\beta}$, we look into the strategies used for the extremal problems, which are the Steklov ($\beta = 0$) and the Laplace-Beltrami ($\beta \rightarrow +\infty$) cases. The strategies of Brock and Hersch for those cases are actually close but distinct: they use the coordinate functions as test functions in the Rayleigh quotient characterization of eigenvalues. In the case of the Laplace-Beltrami operator though, J. Hersch had an additional step: he first transports the surface $\partial\Omega$ on the sphere by a conformal mapping, and use the conformal invariance of the Dirichlet energy for 2-dimensional surfaces. In the following, we choose to follow the ideas of Brock. This allows to obtain an estimate with no assumption on the topology or the dimension of the domain. Indeed, the above mentioned phenomenon of decoupling between the different connected components does not appear in the Steklov case, due to the volume term, and in fact Brock's result is valid for every (smooth enough) domain. The same volume term appears in the Wentzell case and the approach of Brock is then the natural one. However, one expects from these topological considerations that it will not provide an optimal result.

2.2 Proof of Theorem 1.1

Our strategy to prove Theorem 1.1 is to use the following characterization for the inverse trace of eigenvalues (stated by J. Hersch in [18] and proved by G. Hile and Z. Xu in [20])

$$\sum_{i=1}^d \frac{1}{\lambda_{i,\beta}} = \max_{v_1, \dots, v_d} \sum_{i=1}^d \frac{B(v_i, v_i)}{A_{\beta}(v_i, v_i)}, \quad (21)$$

where the functions $(v_i)_{i=1, \dots, d}$ are non zero functions that are B -orthogonal to the constants and pairwise A_{β} -orthogonal.

Before proving Theorem 1.1, we now present some preliminary results.

Lemma 2.4 *The matrix $P[\Omega]$ defined by (8) is symmetric, positive definite. Its spectral radius $\Lambda[\Omega]$ satisfies*

$$(d-1)|\partial\Omega| \geq \Lambda[\Omega] \geq \frac{d-1}{d} |\partial\Omega|. \quad (22)$$

In particular, among sets of given volume, the spectral radius is minimal for the ball.

Proof: The matrix $P(\Omega)$ is symmetric by definition. For $\mathbf{y} = (y_1, \dots, y_d) \in \mathbb{R}^d$ with $\mathbf{y} \neq \mathbf{0}$, we check that

$$\sum_{i,j=1}^d y_i(\delta_{ij} - \mathbf{n}_i \mathbf{n}_j) y_j = \mathbf{y}^T \mathbf{y} - (\mathbf{y}^T \mathbf{n})^2 \geq 0$$

by Cauchy-Schwarz inequality. By integration over $\partial\Omega$, $P[\Omega]$ is positive semidefinite. Assume, by contradiction, that P is not definite: then there is a vector $\mathbf{y} \neq \mathbf{0}$ such that

$$0 = \sum_{i,j=1}^d y_i \left(\int_{\partial\Omega} (\delta_{ij} - \mathbf{n}_i \mathbf{n}_j) \right) y_j = \int_{\partial\Omega} (\mathbf{y}^T \mathbf{y} - (\mathbf{y}^T \mathbf{n})^2).$$

The equality case of Cauchy-Schwarz inequality $\mathbf{y}^T \mathbf{y} - (\mathbf{y}^T \mathbf{n})^2 = 0$ is therefore satisfied everywhere on $\partial\Omega$, this holds if and only if \mathbf{y} and \mathbf{n} are colinear. Hence, \mathbf{n} is constant on $\partial\Omega$ which contradicts the boundedness of Ω .

The matrix $P[\Omega]$ has positive eigenvalues. Their sum is the trace $\text{Tr}(P[\Omega])$, hence

$$\text{Tr}(P[\Omega]) \geq \Lambda[\Omega] \geq \frac{\text{Tr}(P[\Omega])}{d} \quad \text{with} \quad \text{Tr}(P[\Omega]) = \sum_{i=1}^d \int_{\partial\Omega} (1 - \mathbf{n}_i^2) = (d-1)|\partial\Omega|.$$

Therefore

$$(d-1)|\partial\Omega| \geq \Lambda[\Omega] \geq \frac{(d-1)}{d} |\partial\Omega| \geq \frac{(d-1)}{d} |\partial B|.$$

The last inequality is obtained by the usual isoperimetric inequality and assuming B is a ball such that $|\Omega| = |B|$. Let us compute $\Lambda[B]$. From the invariance by rotation of the ball, there exists a real number a such that $P[B] = aI_d$. In others words, we have

$$\int_{\partial B} \mathbf{n}_i \mathbf{n}_j = 0, \quad i \neq j \quad \text{and} \quad \int_{\partial B} (1 - \mathbf{n}_i^2) = \int_{\partial B} (1 - \mathbf{n}_1^2), \quad i = 1, \dots, d.$$

The real number a is determined using the trace of the matrix: we obtain that $d \Lambda[B] = (d-1)|\partial B|$, and so $\Lambda(\Omega) \geq \Lambda(B)$. ■

Remark 2.5 *The inequalities in (22) are sharp. The lower bound is reached when Ω is a ball and the upper bound is the limit of the collapsing stadium S_ε (union of a rectangle and two half-disks) of unit area and width ε when ε tends to 0: one checks by an explicit elementary calculus that:*

$$|\partial S_\varepsilon| = \frac{2}{\varepsilon} + \frac{\pi\varepsilon}{2} \quad \text{while} \quad \Lambda[S_\varepsilon] = \frac{2}{\varepsilon}.$$

This example is also useful to prove (6): indeed, we easily prove

$$\int_{\partial S_\varepsilon} |x|^2 \geq \frac{\alpha}{\varepsilon^3},$$

where α is a universal constant, so using (11), we obtain (6) for $d = 2$ and $m = 1$. The other cases can be handled similarly.

Proof of Theorem 1.1:

We first translate and rotate coordinates x_i , $i = 1, 2, \dots, d$ such that

$$\forall i \neq j \in \llbracket 1, d \rrbracket^2, \quad \int_{\partial\Omega} x_i = 0 \text{ and } \int_{\partial\Omega} x_i x_j = 0.$$

We now construct a family which is pairwise A_β -orthogonal, and B -orthogonal to \mathbb{R} . We consider a collection of a family of functions w_1, w_2, \dots, w_d in the vector space spanned by the coordinates functions: there is a matrix C such that

$$w_i = \sum_{j=1}^d c_{ij} x_j, \quad i \in \llbracket 1, d \rrbracket.$$

Brock used directly the coordinate functions to deal with A_0 . Here, we need an A_β -orthogonal family, hence the matrix C will be chosen to that end. Since the coordinates functions are L^2 orthogonal to the constants, each w_i is L^2 -orthogonal to the constants (that is to say the eigenfunctions associated to the smallest eigenvalue $\lambda_0 = 0$).

Let us compute $A_\beta(w_i, w_j)$. First, we get $\nabla w_i = (c_{i1}, c_{i2}, \dots, c_{id})^T$ then

$$\int_{\Omega} \nabla w_i \cdot \nabla w_j = \int_{\Omega} \sum_{k,m=1}^d c_{ik} c_{jm} = |\Omega| (CC^T)_{ij}.$$

To compute the second term of the sum occurring in A_β , we recall that

$$\nabla_\tau w_i \cdot \nabla_\tau w_j = \nabla w_i \cdot \nabla w_j - (\nabla w_i \cdot \mathbf{n})(\nabla w_j \cdot \mathbf{n}_j).$$

We therefore get

$$\begin{aligned} \int_{\partial\Omega} \nabla_\tau w_i \cdot \nabla_\tau w_i &= \int_{\partial\Omega} \left[\sum_{k=1}^d c_{ik} c_{jk} - \left(\sum_{k=1}^d c_{ik} \mathbf{n}_k \right) \cdot \left(\sum_{k=1}^d c_{jk} \mathbf{n}_k \right) \right] \\ &= \int_{\partial\Omega} \left[\sum_{k=1}^d c_{ik} c_{jk} - \sum_{k,l=1}^d c_{ik} c_{jl} \mathbf{n}_k \cdot \mathbf{n}_l \right]. \end{aligned}$$

We introduce $P[\Omega]$ the matrix defined in (8) to get

$$\int_{\partial\Omega} \nabla_\tau w_i \cdot \nabla_\tau w_j = \sum_{k,m} c_{ik} p_{km} c_{jm} = (CP[\Omega]C^T)_{ij}.$$

Gathering all the terms, it comes that

$$A_\beta(w_i, w_j) = |\Omega| (CC^T)_{ij} + \beta (CP[\Omega]C^T)_{ij} \quad (23)$$

Since $P[\Omega]$ is a real symmetric matrix, we can choose an orthogonal matrix C such that $CP[\Omega]C^T$ is diagonal. Hence, $CC^T = I$ and finally w_i and w_j are A_β -orthogonal if $i \neq j$ while

$$A_\beta(w_i, w_i) = |\Omega| + \beta (CP[\Omega]C^T)_{ii} \leq |\Omega| + \beta \Lambda[\Omega]. \quad (24)$$

and we can apply Hile and Xu's inequality (see [20]).

Since by assumption

$$\int_{\partial\Omega} x_i x_j = 0$$

when $i \neq j$, it comes that

$$B(w_i, w_i) = \sum_{k=1}^d c_{ik}^2 \int_{\partial\Omega} x_k^2$$

and then

$$S(\Omega) = \sum_{i=1}^d \frac{1}{\lambda_{i,\beta}(\Omega)} \geq \frac{\sum_{i=1}^d \sum_{k=1}^d c_{ik}^2 \int_{\partial\Omega} x_k^2}{|\Omega| + \beta\Lambda[\Omega]} = \frac{\sum_{k=1}^d \left(\int_{\partial\Omega} x_k^2 \right) \sum_{i=1}^d c_{ik}^2}{|\Omega| + \beta\Lambda[\Omega]} = \frac{\int_{\partial\Omega} |x|^2}{|\Omega| + \beta\Lambda[\Omega]},$$

which is the first part of the result. Then using first the isoperimetric weighted inequality (17) for $p = 2$, we get

$$\int_{\partial\Omega} |x|^2 \geq R^2 |\partial B_R|,$$

and so

$$\frac{\int_{\partial\Omega} |x|^2}{|\Omega| + \beta\Lambda[\Omega]} \geq \frac{R^2 |\partial B_R|}{|\Omega| + \beta\Lambda[\Omega]} = \frac{R^2}{\frac{|B_R|}{|\partial B_R|} + \frac{\beta\Lambda[\Omega]}{|\partial B_R|}}.$$

If $\Omega = B_R$, we know that $d|B_R| = R|\partial B_R|$ and then

$$\frac{R^2}{\frac{|B_R|}{|\partial B_R|} + \frac{\beta\Lambda[B_R]}{|\partial B_R|}} = \frac{R^2}{\frac{R}{d} + \beta \frac{d-1}{d}} = \frac{d}{\lambda_{1,\beta}(B_R)},$$

and prove the equality case. By the quantitative version of the isoperimetric inequality for the moment of inertia of $\partial\Omega$ with respect to the origin (18), we also get the precise version:

$$\frac{\int_{\partial\Omega} |x|^2}{|\Omega| + \beta\Lambda[\Omega]} \geq \frac{R^2 |\partial B_R|}{|\Omega| + \beta\Lambda[\Omega]} \left[1 + \gamma_d \left(\frac{|\Omega \Delta B_R|}{|B_R|} \right)^2 \right].$$

Using the definition of R and $|\Omega| = |B_R|$, we obtain $R^2 |\partial B_R| = d\omega_d^{-1/d} |\Omega|^{\frac{d+1}{d}}$ and the desired inequality. \blacksquare

Proof of Corollary 1.2:

Since $\lambda_{1,\beta}(\Omega) \leq \lambda_{i,\beta}(\Omega)$ for $i = 1, \dots, d$, we get

$$\lambda_{1,\beta}(\Omega) \leq \frac{d}{S(\Omega)} \leq d \frac{|\Omega| + \beta\Lambda[\Omega]}{\int_{\partial\Omega} |x|^2} \leq \frac{d}{1 + \gamma_d \left(\frac{|\Omega \Delta B_R|}{|B_R|} \right)^2} \frac{|\Omega| + \beta\Lambda[\Omega]}{d\omega_d^{-1/d} |\Omega|^{\frac{d+1}{d}}}.$$

Proof of Theorem 1.3:

It is a direct adaptation of the previous proof to the Laplace-Beltrami case: it suffices to replace the bilinear form $A_\beta(u, v)$ by $A(u, v) = \int_\Omega \nabla u \cdot \nabla v$. Then Equation (24) becomes $A(w_i, w_i) = (CP[\Omega]C^T)_{ii} \leq \Lambda[\Omega]$ and the conclusion follows. \blacksquare

2.3 On the sharpness of the upper bounds.

Testing the sharpness. Let us denote $M_1(\Omega)$ the upper bound (11). In order to emphasize the improvement to the inequality of Brasco, De Philippis and Ruffini, we also plot the rougher upper bound

$$M_3(\Omega) = \frac{|\Omega| + \beta\Lambda[\Omega]}{\omega_d^{-1/d}|\Omega|^{\frac{d+1}{d}}} = d \frac{|\Omega| + \beta\Lambda[\Omega]}{R^2|\partial B_R|}.$$

It is clear from the bound of $\Lambda[\Omega]$ stated in (22) that

$$\lambda_{1,\beta}(B_R) = M_1(B_R) \leq M_2(\Omega) = \frac{d}{\left(1 + \gamma_d \frac{|\Omega\Delta B_R|}{|B_R|}\right)^2} \frac{|\Omega| + \beta\Lambda[\Omega]}{R^2|\partial B_R|}.$$

We also plot the shaper bound

$$M_1(\Omega) = d \frac{|\Omega| + \beta\Lambda[\Omega]}{\int_{\partial\Omega} |x|^2}.$$

This inequality means that proving that balls are maximizers would be strictly better than (11). Let us illustrate this fact with some numerical illustrations. We compute $\lambda_{1,\beta}(\Omega)$ and $M_i(\Omega)$ ($i = 1, 2$) for several parametrized families of plane domains when $\beta = 1$. In Figure 1(a), we present the case of ellipses of area π (their semiaxis are e^t and e^{-t} , t is in abscissa) while in Figure 1(b) and 1(d) we present the case of the star-shaped domains Ω_t defined in polar coordinate by $r(\theta) = a(t)(2 + \cos(k\theta))$ where $a(t)$ is a constant chosen such that $|\Omega_t| = \pi$.

From these graphs, it seems that the upper bounds $M_i(\Omega)$ lack of precision when Ω is far from a ball and that the maximality of balls is possible and would improve the upper bound given in Corollary 1.2.

Some numerical tests. It is natural to wonder if the ball have the largest $\lambda_{1,\beta}$ among all the domains of same volume that are homeomorphic to the ball. This question cannot be solved with estimate (11), as shows Figure 1(a). Therefore, to conclude this section, we would like to present some numerical experiments in favor of such property.

Let us start by computing the value of $\lambda_{1,\beta}(\Omega)$ when Ω is an ellipse of fixed volume. We present here the results of our numerical computations for $\beta \in \{0.1, 1, 5, 10\}$ when $|\Omega| = \pi$. Then when the volume of Ω is 4π . In both figures, the abscissa stands for the eccentricity of the ellipse. It seems that the ball maximizes $\lambda_{1,\beta}$ among ellipses of fixed area.

Let us show some computations in dimension three. We consider families of ellipsoids with semi-axes defined by $(\exp(\alpha_i t))_{i=1,2,3}$ where $\alpha_1 + \alpha_2 + \alpha_3 = 0$ to insure the volume constraint. The ball B corresponds to $t = 0$. We remind that in this case, $\lambda_{1,\beta}(B)$ has multiplicity 3 at the sphere, we then have plotted the three corresponding eigenvalues in two cases: first for the family such that $\alpha = (2, -0.8, -1.2)$ in Figure 3(a), then for $\alpha = (2, -1, -1)$ in Figure 3(b). In the last case, the defined ellipsoids are of revolution and we observe that in this particular case $\lambda_{3,\beta} \approx \lambda_{4,\beta}$. One can wonder if it is really the case.

Let $E(a, b)$ be an ellipsoid of volume $4\pi/3$ where a is the larger semiaxis and b the middle one. We now show in Figure 4 the surfaces $z = \lambda_{i,\beta}(E(a, b))$ where $i = 1, 2, 3$. The pictures have been obtained by interpolation after the computations of the eigenvalues on 2700 ellipsoids. Again one can attest that the ball seems to maximize $\lambda_{1,\beta}$ among ellipsoids.




Figure 1: Comparison of $\lambda_{1,\beta}(\Omega)$ and $M_i(\Omega)$. Here $\lambda_{1,\beta}(B_1) = 2$.




Figure 2: $\lambda_{1,\beta}(\Omega)$ when Ω is an ellipse of volume $|\Omega|$

3 First order shape calculus

In order to go one step further, we adopt a shape optimization point of view and prove in this section that the ball is a critical point. The main difficulty here is that the eigenvalue $\lambda_{1,\beta}(B)$ has multiplicity the dimension of the ambient space. We need some technical material on shape derivative and tangential calculus on manifold to justify the results stated in this section; to simplify the reading




Figure 3: $(\lambda_{1,\beta}(\Omega_t), \lambda_{2,\beta}(\Omega_t), \lambda_{3,\beta}(\Omega_t))$ when Ω_t is a parametrized ellipsoid of volume $4\pi/3$




Figure 4: $(\lambda_{1,\beta}(\Omega), \lambda_{2,\beta}(\Omega), \lambda_{3,\beta}(\Omega))$ when $\Omega = E(a,b)$ is an ellipsoid of volume $4\pi/3$

of this work, we postpone these reminders in Appendix A.

Let us emphasize that from this point we do not make the assumption $\beta \geq 0$, and therefore all the results of this section and the following are valid for any $\beta \in \mathbb{R}$. Though from now on we drop the notation β in $\lambda_{1,\beta}$ since there is no possible confusion anymore.

3.1 Notations and preliminary result for shape deformation

We adopt the formalism of Hadamard's shape calculus and consider the map $t \mapsto T_t = I + t\mathbf{V}$ where $\mathbf{V} \in W^{3,\infty}(\Omega, \mathbb{R}^d)$ and t is small enough. We denote by

$$\Omega_t = T_t(\Omega) = \{x + t\mathbf{V}(x), x \in \Omega\}.$$

Remark 3.1 *More generally the results and computations from this section are valid if $t \mapsto T_t$ satisfies:*

- $T_0 = Id$,
- for every t near 0, T_t is a $W^{3,\infty}$ -diffeomorphism from Ω onto its image $\Omega_t = T_t(\Omega)$.
- The application $t \mapsto T_t$ is real-analytic near $t = 0$.

We need to introduce the surface jacobian ω_t defined as

$$\omega_t(x) = \det(DT_t(x)) \|(DT_t(x)^T)^{-1}\mathbf{n}(x)\|,$$

and the functions

$$A_t(x) = (DT_t(x))^{-1}(DT_t(x)^T)^{-1}, \quad \tilde{A}_t(x) = \det(DT_t(x))A_t(x), \quad C_t(x) = \omega_t(x)A_t(x).$$

We have to study the transport of the considered eigenvalue problem on the deformed domain Ω_t . To that end, we first rewrite the deformed equation on the fixed domain Ω and its boundary $\partial\Omega$: we have to describe how are transported the Laplace-Beltrami and the Dirichlet-to-Neumann operators.

Transport of the Dirichlet-to-Neumann map. Let us consider the Dirichlet-to-Neumann operator defined on its natural space $D_t : H^{1/2}(\partial\Omega_t) \rightarrow H^{-1/2}(\partial\Omega_t)$. It maps a function ϕ_t in $H^{1/2}(\partial\Omega_t)$ onto the normal derivative of its harmonic expansion in Ω_t , that is to say $D_t(\phi_t) = \partial_{\mathbf{n}_t} u_t$, where u^t solves the boundary values problem:

$$\begin{cases} -\Delta u_t = 0 & \text{in } \Omega_t, \\ u_t = \phi_t & \text{on } \partial\Omega_t. \end{cases} \quad (25)$$

To compute the quantity \mathcal{D}_t such that $\mathcal{D}_t(\phi_t \circ T_t) = [D_t(\phi_t)] \circ T_t$, we transport the boundary value problem (25) back on the domain Ω . In others words, \mathcal{D}_t makes the following diagram commutative:

$$\begin{array}{ccc} H^{1/2}(\partial\Omega_t) & \xrightarrow{T_t} & H^{1/2}(\partial\Omega) \\ \mathcal{D}_t \downarrow & & \downarrow \mathcal{D}_t \\ H^{-1/2}(\partial\Omega_t) & \xrightarrow{T_t} & H^{-1/2}(\partial\Omega) \end{array}$$

To be more precise, we have the following result proved in [12].

Lemma 3.2 *Given $\psi \in H^{1/2}(\partial\Omega)$, we denote v^t the solution of the boundary value problem*

$$\begin{cases} -\operatorname{div}(\tilde{A}_t \nabla v^t) = 0 & \text{in } \Omega, \\ v^t = \psi & \text{on } \partial\Omega. \end{cases} \quad (26)$$

and then define $\mathcal{D}_t \psi \in H^{-1/2}(\partial\Omega)$ as:

$$\mathcal{D}_t \psi : f \in H^{1/2}(\partial\Omega) \mapsto \int_{\Omega} \tilde{A}_t(x) \nabla v^t(x) \cdot \nabla E(f)(x) dx,$$

where E is a continuous extension operator from $H^{1/2}(\partial\Omega)$ to $H^1(\Omega)$. Then the relation

$$(D_t \varphi) \circ T_t = \mathcal{D}_t [\varphi \circ T_t] \quad (27)$$

holds for all functions $\varphi \in H^{1/2}(\Omega_t)$.

Setting $u^t = u_t \circ T_t$, we check from the variational formulation, that the function u^t is the unique solution of the transported boundary value problem:

$$\begin{cases} -\operatorname{div}(\tilde{A}_t \nabla u^t) &= 0 & \text{in } \Omega, \\ u^t &= \phi_t \circ T_t & \text{on } \partial\Omega. \end{cases} \quad (28)$$

Hence, setting $y = T_t(x)$, $x \in \Omega$ we get formally

$$D_t(\phi_t)(y) = \nabla u_t(y) \cdot \mathbf{n}_t(y) = (DT_t(x)^T)^{-1} \nabla v_t(x) \cdot \frac{(DT_t(x)^T)^{-1} \mathbf{n}(x)}{\|(DT_t(x)^T)^{-1} \mathbf{n}(x)\|} = \frac{A_t(x) \mathbf{n}(x) \cdot \nabla u^t(x)}{\|(DT_t(x)^T)^{-1} \mathbf{n}(x)\|}.$$

Here again, we can give a sense to the co-normal derivative $A_t \mathbf{n} \cdot \nabla u^t$ thanks to the boundary value problem (28): this quantity is defined in a weak sense as the previous Dirichlet-to-Neumann operator D_t .

Transport of the Laplace-Beltrami operator. We recall now the expression of the transported Laplace-Beltrami operator, relying on the relation

$$\forall \varphi \in H^2(\partial\Omega_t), \quad (\Delta_\tau \varphi) \circ T_t = \frac{1}{\omega_t(x)} \operatorname{div}_\tau (C_t(x) \nabla_\tau (\varphi \circ T_t)(x)) \quad \text{on } \partial\Omega. \quad (29)$$

Let us denote by \mathcal{L}_t the operator defined as

$$\begin{aligned} \mathcal{L}_t[\varphi \circ T_t](x) &= \\ &= \frac{1}{\omega_t(x)} \operatorname{div}_\tau \left\{ C_t(x) \nabla [\varphi \circ T_t](x) - \frac{C_t(x) \nabla [\varphi \circ T_t](x) \cdot \mathbf{n}(x)}{A_t(x) \mathbf{n}(x) \cdot \mathbf{n}(x)} A_t(x) \mathbf{n}(x) \right\} \end{aligned} \quad (30)$$

for $\varphi \in H^{5/2}(\Omega_t)$. In [12], we show the following lemma:

Lemma 3.3 *The identity*

$$[\Delta_\tau \varphi] \circ T_t = \mathcal{L}_t[\varphi \circ T_t] \quad (31)$$

holds for all functions φ belonging to $H^{5/2}(\Omega_t)$.

3.2 Regularity of the eigenfunctions and eigenvalues with respect to the parameter

The section is a slight variation of a theorem due to Ortega and Zuazua on the existence and regularity of eigenvalues and associated eigenfunctions in the case of Stokes system [24]. The difficulty comes from the possible multiple eigenvalues. The main result is, for a fixed deformation field $\mathbf{V} \in W^{3,\infty}(\Omega, \mathbb{R}^d)$, the existence of smooth branches of eigenvalue. In other words, the eigenvalues are not regular when sorted in the increasing order, but can be locally relabeled around the multiple point in order to remain smooth. The restriction is that this labeling depends on the deformation field \mathbf{V} hence one cannot hope to prove Fréchet-differentiability.

Theorem 3.4 *Let Ω be an open smooth bounded domain of \mathbb{R}^d . Assume that λ is an eigenvalue of multiplicity m of the Wentzell-Laplace operator. We suppose that $T_t = I + t\mathbf{V}$ for some $\mathbf{V} \in W^{3,\infty}(\Omega, \mathbb{R})^d$ and denote $\Omega_t = T_t(\Omega)$. Then there exists m real-valued continuous functions $t \mapsto \lambda_i(t)$, $i = 1, 2, \dots, m$ and m functions $t \mapsto u_i^t \in H^{\frac{5}{2}}(\Omega)$ such that the following properties hold*

1. $\lambda_i(0) = \lambda$, $i = 1, \dots, m$,
2. the functions $t \mapsto \lambda_i(t)$ and $t \mapsto u_i^t$, $i = 1, 2, \dots, m$ are analytic in a neighborhood of $t = 0$.
3. The functions $u_{i,t}$ defined by $u_{i,t} \circ T_t = u_i^t$ are normalized eigenfunctions associated to $\lambda_i(t)$ on the moving domain Ω_t . If one considers K compact subset such that $K \subset \Omega_t$ for all t small enough, then $t \mapsto u_{i,t}|_K$ is also an analytic function of t in a neighborhood of $t = 0$.
4. Let $I \subset \mathbb{R}$ be an interval such that \bar{I} contains only the eigenvalue λ of the Wentzell problem of multiplicity m . Then there exists a neighborhood of $t = 0$ such that $\lambda_i(t)$ $i = 1, \dots, m$ are the only eigenvalues of Ω_t which belongs to I .

Proof: Let λ be an eigenvalue of multiplicity m and let u_1, \dots, u_m the orthonormal eigenfunctions associated to λ . Let $(\lambda(t), u_t)$ be an eigenpair satisfying

$$(P_t) \begin{cases} -\Delta u_t & = 0 & \text{in } \Omega_t, \\ -\beta \Delta_\tau u_t + \partial_{n_t} u_t & = \lambda(t) u_t & \text{on } \partial \Omega_t. \end{cases}$$

Setting $u^t = u_t \circ T_t$, Lemma 3.2 (transport of the Dirichlet-to-Neumann map) and 3.3 (transport of the Laplace-Beltrami operator) show that the system (P_t) above is equivalent to the following equation set on the boundary

$$(-\beta \mathcal{L}_t + \mathcal{D}_t) u^t = \lambda(t) \omega_t u^t \text{ on } \partial \Omega. \quad (32)$$

Consider the operator $S(t)$ defined on $H^{3/2}(\partial \Omega)$ by

$$v \mapsto S(t)v = -\beta \mathcal{L}_t v + \mathcal{D}_t v \quad (33)$$

From their expressions computed for example in [17, Section 5-2] and the regularity assumption on T_t , all the operators C_t , A_t and ω_t are analytic in a neighborhood of $t = 0$. Since $\det(DT_t) > 0$ for t small enough, we deduce that all the expressions involved in C_t , \mathcal{L}_t and \mathcal{D}_t are analytic in a neighborhood of $t = 0$. This enables us to conclude that $S(t)$ is also analytic in a neighborhood of zero.

To show that the eigenvalues and the corresponding eigenfunctions are analytic in a neighborhood of zero, we apply the Lyapunov-Schmidt reduction in order to treat a problem on a finite dimensional space, namely the kernel of $S(0) - \lambda I$. To that end, we rewrite the problem (P_t) on the fixed domain $\partial \Omega$ as

$$S(t)(u^t) - \lambda(t) \omega_t u^t = 0.$$

From the decomposition

$$(S(0) - \lambda)(u^t) = \left[(S(0) - S(t)) + [(\lambda(t) - \lambda) \omega_t + \lambda(\omega_t - 1)] \right] u^t,$$

u^t is solution of the equation

$$(S(0) - \lambda)(u^t) = W(t, \lambda(t) - \lambda) u^t, \quad (34)$$

where we have set $R(t) = S(0) - S(t) + \lambda(\omega_t - 1)$ and $W(t, \alpha) = R(t) + \alpha \omega_t I$. From the Lyapunov-Schmidt Theorem (see [24, Lemma 3-2, p. 999]), we obtain that $S(0) - \lambda$ has a right inverse operator denoted by K . Hence the equation above implies that $u^t = KW(t, \lambda(t) - \lambda) u^t + \psi_t$ where $\psi_t \in \text{Ker}(S(0) - \lambda)$, i.e $\psi_t = \sum_{k=1}^m c_k(t) \phi_k$ where (ϕ_k) is a basis of $\text{Ker}(S(0) - \lambda)$. Notice that $I -$

$KW(t, \lambda(t) - \lambda)$ is invertible on $\text{Ker}(S(0) - \lambda I)$, the inverse of his operator restricted to this kernel will be denoted by $(I - KW(t, \lambda(t) - \lambda))^{-1}$ so that

$$u^t = (I - KW(t, \lambda(t) - \lambda))^{-1} \psi_t.$$

From (34), $W(t, \lambda(t) - \lambda)u^t$ belongs to $\text{Im}(S(0) - \lambda) = \text{Ker}^\perp(S(0) - \lambda)$ since $S(0)$ is a Fredholm selfadjoint operator, and then

$$\sum_{k=1}^m c_k(t) \langle W(t, \lambda(t) - \lambda)(I - KW(t, \lambda(t) - \lambda))^{-1} \phi_k, \phi_i \rangle = 0, \quad i = 1, 2, \dots, m, \quad (35)$$

where $\langle \cdot, \cdot \rangle$ denote the scalar product of $L^2(\partial\Omega)$. This shows that a vector of coefficients $C = (c_j)_{j=1, \dots, m} \neq 0$ is a solution if and only if the determinant of the $m \times m$ matrix $M(t, \lambda(t) - \lambda)$ with entries

$$M(t, \alpha)_{i,j} = \langle W(t, \alpha)(I - KW(t, \alpha))^{-1} \phi_j, \phi_i \rangle$$

satisfies

$$\det(M(t, \lambda(t) - \lambda)) = 0.$$

Hence $\lambda(t)$ is an eigenvalue of our problem if and only if $\det(M(t, \lambda(t) - \lambda)) = 0$. Note that $t \mapsto M(t, \lambda(t))$ is analytic around $t = 0$.

For small values of t the operator $(I - KW(t, \alpha))^{-1}$ is well defined since $I - KW(0, 0) = I$ and $t \mapsto (I - KW(t, \alpha))^{-1}$ is analytic around $t = 0$. On the other hand, if $\det M(t, \alpha) = 0$ then (35) has a nontrivial solution $c_1(t), \dots, c_m(t)$ and this means that $\lambda(t) = \lambda + \alpha$ is an eigenvalue of (P_t) .

We focus now on $\det M(t, \alpha)$ for $\alpha \in \mathbb{R}$. From the fact that $W(0, \alpha) = \alpha I$, it comes that for sufficiently small values of α , the operator $I - KW(0, \alpha)$ is invertible on $\text{Ker}(S(0) - \lambda I)$ and from the Von Neumann expansion we write

$$\langle W(0, \alpha)(I - KW(0, \alpha))^{-1} \phi_i, \phi_j \rangle = \alpha \left[\delta_{ij} + \sum_{k=1}^{\infty} \alpha^k \langle K^k \phi_i, \phi_j \rangle \right];$$

hence

$$\det(M(0, \alpha)) = \alpha^m + \sum_{i=1}^{\infty} \beta_i \alpha^{m+i} = \alpha^m \left(1 + \sum_{i=1}^{\infty} \beta_i \alpha^i \right).$$

Since $\det(M(0, \alpha)) \neq 0$ is the restriction on $t = 0$ of $\det(M(t, \alpha))$, we deduce from the Weierstrass preparation theorem that there is neighborhood of $(0, 0)$ such that $\det(M(t, \alpha))$ is uniquely representable as

$$\det(M(t, \alpha)) = P_m(t, \alpha)h(t, \alpha)$$

where

$$P_m(t, \alpha) = \alpha^m + \sum_{k=1}^m a_k(t) \alpha^{m-k}$$

and where

$$h(t, \alpha) \neq 0.$$

Furthermore, the coefficients $a_k(t)$, $k = 1, \dots, m$ are real and analytic in a neighborhood of $t = 0$. Then $\det(M(t, \alpha)) = 0$ if and only if $P_m(t, \alpha) = 0$. If $\alpha_k(t)$, $k = 1, \dots, m$ are the real roots of the polynomial, we take $\lambda_1(t) = \lambda + \alpha_1(t)$ if $\alpha_1(t)$ is not identically equal to zero.

We now have to find the $(m - 1)$ other branches $\lambda_i(t)$ and the corresponding eigenfunction $u_{i,t}$ for $i = 2, \dots, m$. We use the idea of the deflation method by considering the operator

$$S_2(t) = S(t) - \lambda_1 P_1(t)$$

where P_1 is the orthogonal projection on the subspace spanned by $u_{1,t}$. At $t = 0$, we obtain

$$S_2(0)u_j = S(0)u_j - \lambda \delta_{1j} u_j$$

in other terms $S_2(0)u_j = \lambda u_j$, $j = 2, \dots, m$ while $S_2(0)u_1 = 0$. This shows that λ is an eigenvalue of multiplicity $m - 1$ of $S_2(0)$ with eigenvalues u_2, \dots, u_m . One can show that these functions are the only linearly independent eigenfunctions associated to λ . Now we can apply the same recipe used before to the operator S_2 instead of S . We then get a branch $\lambda_2(t)$ such that $t \mapsto \lambda_2(t)$ is analytic in a neighborhood of $t = 0$. Iterating the process, we get at the end the $m - 1$ branches $\lambda_i(t)$, $i = 1, \dots, m$ such that each branch is analytic in a neighborhood of $t = 0$ and m corresponding eigenfunctions forming an orthonormal set of functions in $H^{\frac{3}{2}}(\partial\Omega_t)$.

The proof of the last item follows the same lines than the proof of Ortega and Zuazua for the Stokes system, see [24]. \blacksquare

Theorem 3.5 *With the notations of Theorem 3.4, if $t \mapsto (\lambda(t), u_t)$ is one of the smooth eigenpair path $(\lambda_i(t), u_{i,t})$ of Ω_t for the Wentzell problem, then the shape derivative $u' = (\partial_t u_t)|_{t=0}$ of the eigenfunction satisfies*

$$\begin{aligned} \Delta u' &= 0 \text{ in } \Omega, \\ -\beta \Delta_\tau u' + \partial_n u' - \lambda u' &= \beta \Delta_\tau (V_n \partial_n u) - \beta \operatorname{div}_\tau (V_n (2D^2 b - H I_d) \nabla_\tau u) \\ &\quad + \operatorname{div}_\tau (V_n \nabla_\tau u) - \lambda'(0)u + \lambda V_n (\partial_n u + H u) \text{ on } \partial\Omega. \end{aligned} \quad (36)$$

Proof: The fact that u' is harmonic inside the domain is trivial. To derive the boundary condition satisfied by u' , we use a test function ϕ_t defined on $\partial\Omega_t$ with $\partial_n \phi_t = 0$ as used in the proof of Lemma 3.2 and 3.3 in [12]. We get the following weak formulation valid for all t small enough:

$$\int_{\partial\Omega_t} \beta \nabla_\tau u(t, x) \cdot \nabla_\tau \phi_t \, d\sigma_t + \int_{\partial\Omega_t} \partial_{n_t} u(t, x) \phi_t \, d\sigma_t - \lambda(t) \int_{\partial\Omega_t} u(t, x) \phi_t \, d\sigma_t = 0.$$

We take the derivative with respect to t and get at $t = 0$:

$$\beta \frac{d}{dt} \left(\int_{\partial\Omega_t} \nabla_\tau u(t, x) \cdot \nabla_\tau \phi_t \, d\sigma_t \right) \Big|_{t=0} + \frac{d}{dt} \left(\int_{\partial\Omega_t} \partial_{n_t} u(t, x) \phi_t \, d\sigma_t \right) \Big|_{t=0} = \frac{d}{dt} \left(\lambda(t) \int_{\partial\Omega_t} u(t, x) \phi_t(x) \, d\sigma_t \right) \Big|_{t=0}.$$

From [14] and [7], we get

$$\frac{d}{dt} \left(\int_{\partial\Omega_t} \nabla_\tau u(t, x) \cdot \nabla_\tau \phi_t \, d\sigma_t \right) \Big|_{t=0} = \int_{\partial\Omega} \left(-\Delta_\tau u' - \Delta_\tau (V_n \partial_n u) + \operatorname{div}_\tau ((2D^2 b - H I_d) \nabla_\tau u) \right) \phi \, d\sigma.$$

After some lengthy but straightforward computations we also obtain

$$\frac{d}{dt} \left(\int_{\partial\Omega_t} \partial_{n_t} u \phi_t \, d\sigma_t \right) \Big|_{t=0} = \int_{\partial\Omega} \partial_n u' \phi \, d\sigma - \int_{\partial\Omega} \nabla_\tau V_n \cdot \nabla_\tau u \phi \, d\sigma + \int_{\partial\Omega} V_n (\partial_n u + H u) \phi \, d\sigma$$

and

$$\begin{aligned} \frac{d}{dt} \left(\int_{\partial\Omega_t} \lambda(t) u_t \phi_t \, d\sigma_t \right) \Big|_{t=0} &= \lambda'(0) \int_{\partial\Omega} u \phi \, d\sigma \\ &\quad + \lambda \int_{\partial\Omega} u' \phi \, d\sigma + \lambda \int_{\partial\Omega} \partial_n u \phi \, d\sigma + \lambda \int_{\partial\Omega} H u \phi \, d\sigma. \end{aligned}$$

To end the proof of this second point, it suffices to gather the relations. \blacksquare

3.3 Shape derivative of simple eigenvalues of the Wentzell-Laplace problem

Let λ be a simple eigenvalue of the Wentzell-Laplace equation (1) and let u be the corresponding normalized eigenfunction. We give in this subsection the explicit formula for the shape derivative of the eigenvalue of the Wentzell-Laplace operator associated to (1).

On $\Omega_t = (I + t\mathbf{V})(\Omega)$ with t small, there is a unique eigenvalue $\lambda(t)$ near λ which is an analytic function with respect of the parameter t . The associated eigenfunction $u_t(x) = u(t, x)$ is solution of the problem (1). The shape derivative denoted u' is the partial derivative $\partial_t u(t, x)$ evaluated at $t = 0$ and solves (36). Let us deduce the analytic expression of $\lambda'(0)$:

Theorem 3.6 *If (λ, u) is an eigenpair (with u normalized) for the Wentzell problem with the additional assumption that λ is simple then the application $t \rightarrow \lambda(t)$ is analytic and its derivative at $t = 0$ is*

$$\lambda'(0) = \int_{\partial\Omega} V_n \left(|\nabla_\tau u|^2 - |\partial_n u|^2 - \lambda H |u|^2 + \beta(H I_d - 2D^2 b) \nabla_\tau u \cdot \nabla_\tau u \right) d\sigma.$$

Proof: We start with the result of Theorem 3.5. Let us multiply the two sides of (36) the boundary condition satisfied by u' by the eigenfunction u and integrate over the boundary $\partial\Omega$:

$$\begin{aligned} 0 &= \int_{\partial\Omega} v'(-\beta\Delta_\tau u + \partial_n u - \lambda u) d\sigma + \int_{\partial\Omega} V_n \partial_n u (-\beta\Delta_\tau u) d\sigma \\ &\quad + \int_{\partial\Omega} \beta V_n (H I_d - 2D^2 b) \nabla_\tau u \cdot \nabla_\tau u d\sigma + \int_{\partial\Omega} V_n |\nabla_\tau u|^2 - \lambda'(0) \int_{\partial\Omega} |u|^2 \\ &\quad - \lambda \int_{\partial\Omega} V_n (u \partial_n u + H |u|^2) d\sigma. \end{aligned}$$

Using the boundary condition satisfied by the eigenfunction: $-\beta\Delta_\tau u + \partial_n u - \lambda u = 0$, it follows that

$$\begin{aligned} 0 &= \int_{\partial\Omega} V_n \partial_n u (\lambda u - \partial_n u) d\sigma + \int_{\partial\Omega} \beta V_n (H I_d - 2D^2 b) \nabla_\tau u \cdot \nabla_\tau u d\sigma \\ &\quad + \int_{\partial\Omega} V_n \nabla_\tau |u|^2 - \lambda'(0) \int_{\partial\Omega} |u|^2 - \lambda \int_{\partial\Omega} V_n (u \partial_n u + H |u|^2) d\sigma. \end{aligned}$$

and the normalization condition $\int_{\partial\Omega} u^2 d\sigma = 1$ implies

$$\begin{aligned} \lambda'(0) &= - \int_{\partial\Omega} V_n |\partial_n u|^2 d\sigma + \int_{\partial\Omega} \beta V_n (H I_d - 2D^2 b) \nabla_\tau u \cdot \nabla_\tau u d\sigma \\ &\quad + \int_{\partial\Omega} V_n |\nabla_\tau u|^2 - \lambda \int_{\partial\Omega} V_n H |u|^2 d\sigma. \end{aligned}$$

■

3.4 Shape derivative of multiple eigenvalues of the Wentzell-Laplace problem

3.4.1 The general result

We suppose that λ is an eigenvalue of multiplicity m . For smooth deformation $t \mapsto \Omega_t$, there will be m eigenvalues close to λ (counting their multiplicities) for small values of t . We know that such a multiple eigenvalue is no longer differentiable in the classical sense. We are then led to compute the directional derivative of $t \mapsto \lambda_i(t)$ at $t = 0$ where $\lambda_i(t), j = 1, \dots, m$ are given by Theorem 3.4. This is the second part of Theorem 1.4 that we recall here:

Theorem 3.7 *Let λ be a multiple eigenvalue of order $m \geq 2$. Then each $t \mapsto \lambda_i(t)$ for $i \in \llbracket 1, d \rrbracket$ given by Theorem 3.4 has a derivative near 0, and the values of $(\lambda'_i(0))_{i \in \llbracket 1, d \rrbracket}$ are the eigenvalues of the matrix $M(V_n) = (M_{jk})_{1 \leq j, k \leq m}$ defined by*

$$M_{jk} = \int_{\partial\Omega} V_n \left(\nabla_\tau u_j \cdot \nabla_\tau u_k - \partial_n u_j \partial_n u_k - \lambda H u_j u_k + \beta (H I_d - 2D^2 b) \nabla_\tau u_j \cdot \nabla_\tau u_k \right) d\sigma. \quad (37)$$

Proof of Theorem 3.7:

Let $t \mapsto (u(t, x), \lambda(t) = \lambda(\Omega_t))$ a smooth path of eigenpair of the Laplace-Wentzell problem, so that it satisfies

$$\begin{cases} \Delta u(t, x) = 0 & \text{in } \Omega_t \\ -\beta \Delta_\tau u(t, x) + \partial_n u(t, x) = \lambda(t) u(t, x) & \text{on } \partial\Omega_t. \end{cases}$$

We have proved that $u' = \partial_t u(0, x)$ is harmonic in Ω and satisfies the boundary condition (36) on $\partial\Omega$. We use the decomposition of $u = u(0, x)$ as

$$u = \sum_{j=1}^m c_j u_j$$

for some $c = (c_1, c_2, \dots, c_m)^T \neq 0$. Multiplying the two sides equation of (36) by u_k , we get after some integration by parts the eigenvalue equation

$$\lambda'(0)c = Mc$$

where $M = (M_{jk})_{1 \leq i, j \leq m}$ is defined by (37). From this, we deduce that the set of derivatives $(\lambda'_i(0))_{i \in \llbracket 1, d \rrbracket}$ is exactly the set of eigenvalues of the matrix M , which achieves the proof of Theorem 3.7. \blacksquare

3.4.2 The case of balls

We consider now the case where the domain is a ball of radius R . The problem is invariant under translation. In order to remove the invariance, we fix the center of mass of the boundary of the domain, as in Section 2.

The coordinates functions x_i are eigenfunctions of the Wentzell-Laplace operator, so we get

$$\lambda = \frac{\beta(d-1) + R}{R^2}, \text{ and } u_i(x) = \frac{x_i}{\|x_i\|_{L^2(\partial B_R)}} = \frac{x_i}{\sqrt{\omega_d R^{d+1}}}.$$

Corollary 3.8 *Let $\Omega = B_R$ be a ball of radius R , λ_1 its first non-trivial eigenvalue, which is of multiplicity d . The shape derivatives of the maps $t \mapsto \lambda_i(t)$, $i = 1, \dots, d$ given by Theorem 3.4 are the eigenvalues of the matrix $M_{B_R}(V_n) = (M_{jk})_{j, k=1, \dots, d}$ defined by*

$$M_{jk} = \frac{\delta_{jk}}{\omega_d R^{d+1}} \left(1 + \beta \frac{d-3}{R} \right) \int_{\partial B_R} V_n - C(d, R) \int_{\partial B_R} V_n x_j x_k d\sigma \quad (38)$$

where $C(d, R) = \frac{(d+1)(1+\beta \frac{d-2}{R})}{\omega_d R^{d+3}}$.

Proof of Corollary 3.8:

We use (37). On one hand we check the geometric quantities:

$$H = \frac{d-1}{R}, \quad D^2b(x) = \frac{1}{R}I_d - \frac{1}{R^3}(x_i x_j)_{i,j}$$

so since $\nabla_\tau u_j, \nabla_\tau u_k$ are in the tangent space of ∂B_R , we obtain that

$$(HI_d - 2D^2b(x))\nabla_\tau u_j \cdot \nabla_\tau u_k = \frac{d-3}{R}\nabla_\tau u_j \cdot \nabla_\tau u_k$$

and on the other hand:

$$\partial_n u_j = \frac{x_i}{R\sqrt{\omega_d R^{d+1}}} \quad \nabla_\tau u_j \cdot \nabla_\tau u_k = \frac{1}{\omega_d R^{1+d}} \left(\delta_{jk} - \frac{x_j x_k}{R^2} \right)$$

Therefore, the matrix $M = M_{B_R}$ has the following entries

$$\begin{aligned} M_{jk} &= \frac{1}{\omega_d R^{d+1}} \int_{\partial B_R} V_n \left[\left(\delta_{jk} - \frac{x_j x_k}{R^2} \right) - \frac{x_j x_k}{R^2} - \lambda \frac{d-1}{R} x_j x_k + \beta \frac{d-3}{R} \left(\delta_{jk} - \frac{x_j x_k}{R^2} \right) \right] d\sigma \\ &= \frac{\delta_{jk}}{\omega_d R^{d+1}} \left(1 + \beta \frac{d-3}{R} \right) \int_{\partial B_R} V_n - \left[\frac{d+1 + \beta \frac{(d-1)^2 + d - 3}{R}}{\omega_d R^{d+3}} \right] \int_{\partial B_R} V_n x_j x_k d\sigma. \end{aligned}$$

This leads to the result since $(d-1)^2 + d - 3 = (d+1)(d-2)$. ■

From this formula, we deduce a first interesting result:

Proposition 3.9 *If V is a volume preserving deformation, then the following statements are equivalent:*

- (i) V_n is orthogonal (in $L^2(\partial B_R)$) to homogeneous harmonic polynomials of degree 2,
- (ii) $M_{B_R}(V_n) = 0$.

Proof of Proposition 3.9:

We denote \mathcal{H}_2 the space of homogeneous harmonic polynomials of degree 2 (therefore we use here a slightly different notation than in Section 4). Let us suppose that $M(V_n) = 0$; this means that

$$\int_{\partial B_R} V_n x_j x_k d\sigma = 0, \text{ for all } j, k = 1, \dots, d, \text{ and in particular } V_n \text{ is orthogonal to } \mathcal{H}_2.$$

If we assume now that V_n is orthogonal to \mathcal{H}_2 , using that

$$\mathcal{H}_2 = \text{span} \{x_j x_k, j \neq k \in \{1, \dots, d\}, x_1^2 - x_j^2, j = 2, \dots, d\}.$$

and moreover that $\int_{\partial B_R} V_n = 0$, we obtain

$$d \int_{\partial B_R} V_n x_1^2 = \sum_{j=2}^d \int_{\partial B_R} V_n (x_1^2 - x_j^2) + \int_{\partial B_R} \sum_{j=1}^d x_j^2 = 0,$$

and therefore

$$\int_{\partial B_R} V_n x_j^2 = \int_{\partial B_R} V_n (x_j^2 - x_1^2) = 0,$$

which concludes the proof. ■

In the case where $M_{B_R}(V_n) \neq 0$, we compute the trace of the matrix $M_{B_R}(V_n)$ to obtain information on its eigenvalues.

Proposition 3.10 *When Ω is a ball of radius R , then*

$$\text{Tr}(M_{B_R}(V_n)) = 0 \tag{39}$$

for all volume preserving deformations.

Proof of Proposition 3.10:

It comes that

$$\text{Tr}(M_{B_R}(V_n)) = -C(d, R) \int_{\partial B_R} \sum_{j=1}^d x_j^2 V_n d\sigma = -C(d, R) \sum_{j=1}^d x_j^2 \int_{\partial B_R} V_n d\sigma = 0$$

since we are concerned with deformations preserving the volume. ■

As a consequence of Proposition 3.9 and Proposition 3.10, there is the following alternative: either the only eigenvalue of $M(V_n)$ is 0, or $M(V_n)$ has at least one nonnegative and one nonpositive eigenvalue. Each $t \mapsto \lambda_i(t)$ given by Theorem 3.4 has a directional derivative at $t = 0$ denoted by $\lambda'_i(0)$. We then define, as usual [8], $\partial\lambda_1$ the subgradient of λ_1 by $\partial\lambda_1 = [\inf_{i=1\dots d} \lambda'_i(0), \sup_{i=1\dots d} \lambda'_i(0)]$. With this notation, $0 \in \partial\lambda_1$ and we say the ball is a critical shape.

3.5 Numerical illustrations

In order to illustrate Proposition 3.10, we consider the two dimensional case and consider perturbations of the disk given in polar coordinates by

$$\rho_t(\theta) = R + tf(\theta)$$

where f has zero mean value.

In Figure 5, the computations are made in the case $R = 1$ and $\beta = 10$, the deformation parameter t appears in the abscissa.

In both collection of figures, we can see the derivatives of the second and third eigenvalues vanish at the ball in every case except when $f(\theta) = \cos(2\theta)$, where the regular lines cross, leading to a really non differentiable second eigenvalue. This is coherent with Proposition 3.9. Let us explicit the case $V_n = R^2 \cos 2\theta$, where we are led to compute the eigenvalues of the following symmetric matrix

$$M = -\frac{3}{\pi R} \begin{pmatrix} \int_0^{2\pi} \cos 2\theta \cos^2 \theta d\theta & 0 \\ 0 & \int_0^{2\pi} \cos 2\theta \sin^2 \theta d\theta \end{pmatrix}$$

whose eigenvalues are $\alpha_1 = -\frac{3}{2R}$ and $\alpha_2 = \frac{3}{2R}$.




Figure 5: $\lambda_1(\Omega)$ and $\lambda_2(\Omega)$ in the direction of $f(\theta)$ - $|B_R| = \pi$, $\beta = 10$.

4 Testing if the ball is a local maximum for λ_1 : second order arguments

We know that any ball is a critical point for volume preserving deformations. Therefore, if the subgradient $\partial\lambda_1(B; V_n) \neq \{0\}$, then the ball is a local maximizer. It remains to deal with the case where all the eigenvalues of $M_B(V_n)$ are 0; this case corresponds to V_n orthogonal to the harmonics of order two. Then, we aim at proving that the second derivative of λ_1 along at least one of the smooth branches is nonpositive.

The necessary order two conditions of optimality are: the second derivative of the Lagrangian should be non positive on the subspace orthogonal to the space generated by the gradient of the volume constraint. We compute:

$$\text{Vol}'(0) = \int_{\partial B_R} V_n \tag{40}$$

Hence $\text{Vol}'(0) = 0$ if and only if $V_n \in (\mathcal{H}_0)^\perp$ where \mathcal{H}_k denotes the linear space of spherical harmonics of order k . Due to the previous remarks, we hence consider deformation field in the hilbertian space \mathcal{H} spanned by all the spherical harmonics of order $l \in I = \mathbb{N} \setminus \{0, 2\}$. The normal component of such a field is orthogonal to spherical harmonics of order 0 and 2.

The goal of this section is to present the different steps for the computations. We will characterize the matrix E whose eigenvalues are the second order derivatives of the smooth branches of eigenvalues. It turns out that this computation is hard even in the case of a ball. Nevertheless, the computation of $\text{Tr}(E)$ is much simpler than the individual computations of the entries. In order to prove that the ball is a local maximum of λ_1 , it suffices to prove that its trace is nonpositive: therefore at least one smooth branch of eigenvalues has a nonpositive second order derivative.

In this section, we consider deformations preserving the volume at second order and not only at first order. Hence, we cannot consider deformation T_t of type $I + tV$ with V independent of t and

introduce deformations S_t that are the flow at time t of a vector field V (see also Remark 1.5). Notice that $S_t = I + tV + o(t)$ so that $T_t - S_t = o(t)$ and first order shape derivatives are unchanged. In particular, one has

$$\frac{d^2}{dt^2} \text{Vol}(S_t(\Omega)) = \int_{\partial\Omega(t)} \left(\frac{\partial}{\partial t}(V_{n(t)}) + V_{n(t)} \frac{\partial}{\partial n(t)}(V_{n(t)}) + HV_{n(t)}^2 \right) d\sigma$$

and the volume preservation at second order means that

$$\left(\frac{d^2}{dt^2} \text{Vol}(S_t(\Omega)) \right)_{|t=0} = \int_{\partial\Omega} \left(\frac{\partial}{\partial t}(V_{n(t)}) + V_{n(t)} \frac{\partial}{\partial n(t)}(V_{n(t)}) + HV_{n(t)}^2 \right)_{|t=0} d\sigma = 0. \quad (41)$$

4.1 Construction of the matrix E of the second derivatives

Let $(u(t, x), \lambda(t) = \lambda(\Omega_t))$ be an eigenpair of the Laplace-Wentzell problem, that is to say solves

$$\begin{cases} \Delta u(t, x) = 0 & \text{in } \Omega_t \\ -\beta \Delta_\tau u(t, x) + \partial_n u(t, x) = \lambda(t)u(t, x) & \text{on } \partial\Omega_t \end{cases}$$

We use the decomposition of $u = u(0, x)$ in the basis of eigenfunctions:

$$u = \sum_{j=1}^d c_j u_j$$

for some c_1, c_2, \dots, c_d not all zero. We have shown that the vector $c = (c_1, c_2, \dots, c_d)^T$ is solution of

$$\lambda'(0)c = M(V_n)c$$

where the matrix $M(V_n) = (M_{jk})_{1 \leq i, j \leq d}$ is defined by (37).

To compute the second derivative at $t = 0$, one has to compute the first shape derivative $u'(x) = u'(0, x)$. Fredholm's alternative insures the existence of a unique harmonic function \tilde{u}_j orthogonal to the eigenfunctions u_1, u_2, \dots, u_d and satisfying on $\partial\Omega$ the boundary condition

$$\begin{aligned} -\beta \Delta_\tau \tilde{u}_j + \partial_n \tilde{u}_j - \lambda \tilde{u}_j &= \beta \left[\Delta_\tau [V_n \partial_n u_j] + \text{div}_\tau [V_n (HI_d - 2D^2 b) \cdot \nabla_\tau u_j] \right] \\ &\quad + \text{div}_\tau [V_n \nabla_\tau u_j] + \lambda' u_j + \lambda V_n (\partial_n u_j + H u_j). \end{aligned} \quad (42)$$

It follows that

$$u' = \sum_{j=1}^d \tilde{c}_j u_j + \sum_{j=1}^d c_j \tilde{u}_j \quad (43)$$

for some c_j, \tilde{c}_j when $j = 1, \dots, d$. We point out that the (c_j) are the same coefficients as the decomposition of u in the basis (u_j) .

The strategy is straightforward : we have to consider the equation satisfied by u' on the boundary $\partial\Omega$ and take its shape derivative again. A first look to the second derivative shows that we will encounter three operators :

- the first contains only u'' and its expression is the following

$$E^{(0)} = -\beta \Delta u'' + \partial_n u'' - \lambda u''$$

- concerning the term in u' and $\lambda' = 0$ we have

$$\begin{aligned} E^{(1)} &= -2\beta\Delta_\tau(V_n\partial_n u') - 2\operatorname{div}_\tau(V_n(I + \beta\mathcal{A})\nabla_\tau u') \\ &\quad - 2\left[\lambda' u' + \lambda V_n(\partial_n u' + H u')\right] \end{aligned}$$

where $\mathcal{A} = HI - 2D^2b$ is the deviatoric part of the curvature tensor.

- The remaining term is $E^{(2)}$ contains only u ; we give a more explicit expression below.

Green-Riemann identity tells us that $\langle E^{(0)}, u_i \rangle = \langle u'', -\beta\Delta_\tau u_i + \partial_n u_i - \lambda u_i \rangle = 0$, $i = 1, \dots, d$. This means that the term $E^{(0)}$ will have no influence in the determination of the second derivative of the eigenvalue. We will focus only on $E^{(1)}$ and $E^{(2)}$.

Construction of $E^{(2)}$:

The computations are very technical. We need first to use a test function ϕ which is the restriction of a test function Φ defined on a tubular neighborhood of the boundary such that its normal derivative on $\partial\Omega$ is zero. This kind of extension is well discussed in the book [13] of Delfour-Zolesio. Taking the shape derivative of the boundary condition (36) (in the multiple case) we need to compute

$$\begin{aligned} \left. \left(\frac{d}{dt} \int_{\partial\Omega_t} V_n \nabla_\tau u \cdot \nabla_\tau \phi \, d\sigma_t \right) \right|_{t=0} &= \langle A^{(1)} u', \phi \rangle + \langle A^{(2)} u, \phi \rangle, \\ \beta \left. \left(\frac{d}{dt} \int_{\partial\Omega_t} \mathcal{A}(t) V_n \nabla_\tau u \cdot \nabla_\tau \phi \, d\sigma_t \right) \right|_{t=0} &= \langle B^{(1)} u', \phi \rangle + \langle B^{(2)} u, \phi \rangle, \\ - \left. \frac{d}{dt} \left(\int_{\partial\Omega_t} \left[\lambda' u + \lambda(u' + V_n \partial_n u + V_n H u) \right] \phi \, d\sigma_t \right) \right|_{t=0} &= \langle C^{(0)} u'', \phi \rangle + \langle C^{(1)} u', \phi \rangle + \langle C^{(2)} u, \phi \rangle, \\ \beta \left. \left(\frac{d}{dt} \int_{\partial\Omega_t} \nabla_\tau (V_n \nabla_\tau \partial_n u) \cdot \nabla_\tau \phi \, d\sigma_t \right) \right|_{t=0} &= \langle D^{(1)} u', \phi \rangle + \langle D^{(2)} u, \phi \rangle. \end{aligned}$$

The remaining $E^{(2)}$ containing only u is then given by

$$E^{(2)} = A^{(2)}u + B^{(2)}u + C^{(2)}u + D^{(2)}u.$$

For an operator L involved in $E^{(i)}$, $i = 1, 2, 3$ we denote by $(L_{ij})_{i,j=1,\dots,d}$ the matrix of L in the basis of the eigenvalues. After calculations (see also Remark D.1 in the Appendix), we get the following linear equation

$$(\lambda'' I - E)c + 2(-M(V_n) + \lambda' I)\tilde{c} = 0$$

(corresponding to the second derivation) together with

$$(-M(V_n) + \lambda' I)c = 0.$$

(corresponding to the first derivation) where the matrix $E = (E_{ij})$ is split into $E = E^{(1)} + E^{(2)}$ where the terms involving u' are gathered in $E^{(1)}$ and the terms involving u are gathered in $E^{(2)}$.

4.2 Computation of the trace

Since the direct computations of the eigenvalues are difficult, we restrict ourselves to the cases $d = 2$ or $d = 3$, and we will focus on the trace of E and prove that $\operatorname{Tr}(E)$ is nonpositive. We start with the trace of $E^{(2)}$:

Lemma 4.1 Assume $d \in \{2, 3\}$. With $K(R) = \frac{d}{R^{2+d}\omega_{d-1}}$, we have

$$\mathrm{Tr}(E^{(2)}) = -(d\beta + R)RK(R) \int_{\partial B_R} |\nabla_\tau V_n|^2 d\sigma - K(R) \int_{\partial B_R} V_n^2 d\sigma. \quad (44)$$

for all deformations preserving volume and such that V_n is orthogonal to spherical harmonics of order two.

Proof: The computation of $E^{(2)}$ is done in the Appendix C, and to obtain the result, we sum all the traces given by Lemmas C.1, C.2, C.3 and C.4. \blacksquare

Concerning $\mathrm{Tr}(E^{(1)})$, we start with the following Lemma which is straightforward (see also Remark D.1):

Lemma 4.2 We have

$$\mathrm{Tr}(E^{(1)}) = 2 \int_{\partial\Omega} V_n \sum_{j=1}^d \left(-\partial_n \tilde{u}_j \partial_n u_j - H\lambda \tilde{u}_j u_j + (I + \beta(HI_d - 2D^2b)) \nabla_\tau \tilde{u}_j \cdot \nabla_\tau u_j \right) d\sigma. \quad (45)$$

holds for all deformations preserving volumes such that V_n is orthogonal to spherical harmonics of order two.

From this result we deduce the following, which is proved in Appendix D:

Proposition 4.3 Assume $d = 3$ and set $\alpha = \frac{\beta}{R}$. We denote $Y_l^m, m = -l, \dots, m$ any spherical harmonic of order $l \in I$. If

$$V_n = \sum_{l \in I} R^l \left(\sum_{m=-l}^l v_{l,m} Y_l^m \right),$$

then

$$\mathrm{Tr}(E^{(1)}) = -K(R) \left(\sum_{l \in I} [A_{l,\alpha} + B_{l,\alpha}] R^{2l+1} \sum_{m=-l}^l |v_{l,m}|^2 \right)$$

where

$$A_{l,\alpha} = \frac{l}{2l+1} \frac{l+2}{l-2} (4\alpha + 2l) \frac{1 + \alpha(3-l)}{1 + \alpha(l+1)} \text{ and } B_{l,\alpha} = \frac{l+1}{l} \frac{l-1}{l} (4\alpha + 2) \frac{1 + \alpha(4+l)}{1 + \alpha(3+l)}.$$

Since $\mathrm{Tr}E = \mathrm{Tr}(E^{(1)}) + \mathrm{Tr}(E^{(2)})$, we will then deduce the following result

Proposition 4.4 Assume $d \in \{2, 3\}$. Then there exists a nonnegative constant μ such that

$$\mathrm{Tr}(E) \leq -K(R)\mu \int_{\partial B_R} |\nabla_\tau V_n|^2 + |V_n|^2 d\sigma.$$

holds for all preserving volume deformations such that V_n is orthogonal to \mathcal{H} .

Proof: We distinguish the case $d = 2$ and $d = 3$.

The case $d = 2$. Let us compute the trace of the matrix E . Gathering all the results of Lemma 4.1 with the computations of Appendix D concerning the trace of the different matrices involved in the matrix E , we obtain the following formula: when

$$V_n = \sum_{l \in I} \frac{R^l}{\sqrt{\pi}} \left(v_1^{(l)} \cos l\theta + v_2^{(l)} \sin l\theta \right), \quad l \in I,$$

we have

$$\text{Tr}(E) = -K(R) \sum_{l \in I} G(\alpha, l) (l^2 + 1) R^{2l+1} ((v_1^{(l)})^2 + (v_2^{(l)})^2). \quad (46)$$

where

$$G(\alpha, l) = \frac{(l^2 - 1) (2 + l^2 + 2\alpha^2(l - 2)l^2 + \alpha(l - 2)(l^2 + 2))}{2(1 + l^2) (l - 2)l(1 + \alpha l)}.$$

Let us remark that $G(\alpha, 1) = 0$. This could have been guessed since the Wentzell eigenvalues are translation invariance: we recall that, denoting Bar the center of mass of the boundary, we have

$$\text{Bar}'(0) = \int_{\partial B_R} x V_n$$

so that deformations orthogonal to spherical harmonics of order 1 preserve at first order the center of mass. A close look to the fraction G shows that it has no pole for $\alpha > 0$ and $l \geq 3$, that it is nonnegative for $l > 2$ and that $G(l, \alpha) \rightarrow 1$ when $l \rightarrow +\infty$; then there is a nonnegative constant μ such that for all $l \geq 3$, $\mu \leq G(l, \alpha)$. This gives

$$\text{Tr}(E) \leq -K(R)\mu \int_{\partial B_R} |\nabla_\tau V_n|^2 + |V_n|^2 d\sigma.$$

The case $d = 3$. The strategy is the same, and we use again Lemma 4.1 and the detailed computations from Section D.2: we get for $l \in I$:

$$V_n = \sum_{l \in I} R^l \sum_{p=-l}^l v_p^{(l)} Y_l^p,$$

$$\text{Tr}(E) = -K(R) \sum_{l \in I} F(\alpha, l) (l(l+1) + 1) R^{2l+1} \sum_{p=-l}^l (v_p^{(l)})^2.$$

where $F(\alpha, l)$ is the fraction

$$F(\alpha, l) = \frac{(l-1) \sum_{m=0}^3 P_m(l) \alpha^m}{(l(l+1) + 1) l (1 + \alpha(l+1)) (2l+1) (l-2) (1 + \alpha(l+3))},$$

and where the polynomial P_m are defined as

$$\begin{aligned} P_0(X) &= 2X^4 + 5X^3 + 16X^2 - 8, \\ P_1(X) &= 4X^5 + 18X^4 + 40X^3 + 68X^2 - 28X - 56, \\ P_2(X) &= 2X^6 + 21X^5 + 42X^4 + 35X^3 + 16X - 112, \\ P_3(X) &= 8X^6 + 18X^5 + 24X^4 - 68X^3 - 144X^2 - 112X - 64. \end{aligned}$$

Let us remark that $F(\alpha, 1) = 0$ for the same reason than in dimension two. By Descartes's rule of signs, the polynomials P_m have at most one positive root. Since $P_m(0) < 0$ and $P_m(2) > 0$ for $m = 0, \dots, 3$, P_m has exactly one positive root which is in $[0, 2]$. Since $l > 2$, there exists a nonnegative constant μ such that for all $k \geq 3$, $\mu \leq F(k, \alpha)$ and

$$\mathrm{Tr}(E) \leq -K(R) \mu \int_{\partial B_R} |\nabla_\tau V_n|^2 + |V_n|^2 d\sigma.$$

■

A Some classical results on tangential differential calculus

We recall some facts about tangential operators acting on functions defined on $\partial\Omega$. The formulas involve the extensions of functions and the differential calculus becomes easier since we will use the classical euclidean differential calculus in a neighborhood of $\partial\Omega$. The canonical extension will be provided thanks to the oriented distance and the orthogonal projection on the tangent plane. For more details, the interested reader will consult the book [13] of M. Delfour and J.P. Zolesio from which we borrowed the necessary material.

A.1 Notations and definitions. Preliminary results

We recall some essential notations and definitions that are needed for the computations of shape derivatives. Given a smooth function $f : \partial\Omega \mapsto \mathbb{R}$, we define its tangential gradient $\nabla_\tau f$ as

$$\nabla_\tau f = \nabla \tilde{f} - \nabla \tilde{f} \cdot \mathbf{n} \mathbf{n} \quad (47)$$

where \tilde{f} is any extension of f in a tubular neighborhood of $\partial\Omega$. An extension is easily obtained when $\partial\Omega$ is smooth. The tangential gradient does not depends on the extension.

It is also useful to define the tangential gradient as the normal projection of $\nabla \tilde{f}$ to the tangent hyperplane of $\partial\Omega$; in other words

$$\nabla_\tau f = \nabla \tilde{f} - n \otimes n \nabla \tilde{f}, \text{ on } \partial\Omega.$$

We also need the definition of the tangential divergence : for a tensor u , we define the surface divergence as

$$\mathrm{div}_\tau u = \mathrm{Tr}(\nabla_\tau u)$$

For regular functions we define the surface Laplacian or Laplace-Beltrami operator as

$$\Delta_\tau f := \mathrm{div}_\tau (\nabla_\tau f).$$

We recall the definition of the oriented distance $b_{\partial\Omega}$:

$$b_{\partial\Omega}(x) = \begin{cases} d_\Omega(x) & \text{for } x \in \mathbb{R}^d \setminus \overline{\Omega} \\ -d_\Omega(x) & \text{for } x \in \Omega, \end{cases} \quad (48)$$

where the notation d_Ω stands for the distance function for a subset $\Omega \subset \mathbb{R}^d$:

$$d_\Omega(x) = \inf_{y \in \Omega} |x - y|$$

We shall sometimes write b instead of $b_{\partial\Omega}$; its gradient is an extension of the normal vector field \mathbf{n} in a neighborhood of $\partial\Omega$.

Let D^2b be the Weingarten operator with entries $(\nabla_\tau)_i n_j$ where n_j is the j -th component of \mathbf{n} . The normal vector is known to be in the kernel of D^2b , while the other eigenfunctions are tangential with the corresponding eigenvalues given by the principal curvatures of $\partial\Omega$.

Let κ_i , $i = 1, \dots, d-1$ be the non zero eigenvalues of D^2b . We define the mean curvature H as

$$H = \sum_{i=1}^{d-1} \kappa_i = \text{Tr}(D^2b) = \Delta b, \text{ on } \partial\Omega. \quad (49)$$

An important result about the normal derivative of this quantities is:

Proposition A.1 *Suppose that the boundary $\partial\Omega$ is of class C^3 . Then the normal derivative of the mean curvature H is*

$$\partial_n H = - \sum_{i=1}^{d-1} \kappa_i^2. \quad (50)$$

Other known identities: we denote \mathbf{x} the identity function. We have

$$\begin{aligned} -\Delta_\tau \mathbf{x} &= H \mathbf{n} \\ \text{div}_\tau \mathbf{n} &= H \mathbf{n} \end{aligned}$$

Tangential integral formula: Given two functions f (scalar) and \mathbf{v} smooth enough, we have

$$\int_{\partial\Omega} f \text{div}_\tau \mathbf{v} + \int_{\partial\Omega} \nabla_\tau f \cdot \mathbf{v} = \int_{\partial\Omega} H f \mathbf{v} \cdot \mathbf{n}$$

Shape derivative of the main curvature H and of the normal \mathbf{n} in the direction of a velocity \mathbf{V} :

Proposition A.2 *Let a surface $\partial\Omega$ be of class C^2 . The shape derivatives of the normal \mathbf{n} and of the mean curvature H in the direction of the velocity vector \mathbf{V} are*

$$\begin{aligned} \mathbf{n}' &= -\nabla_\tau V_n \\ H' &= -\Delta_\tau V_n \end{aligned} \quad (51)$$

where $V_n = \langle \mathbf{V}, \mathbf{n} \rangle$ denotes the normal component of the vector deformation \mathbf{V} .

A.2 A commutation lemma

Here f and g are two smooth functions defined on \mathcal{U} a neighborhood of $\partial\Omega$; the notation b stands for the oriented distance. Recall that its gradient is an extension of the normal field \mathbf{n} on $\partial\Omega$.

Proposition A.3 *We have*

$$\partial_n(\nabla_\tau f \cdot \nabla_\tau g) + 2(D^2b \nabla_\tau f) \cdot \nabla_\tau g = \nabla_\tau(\partial_n f) \cdot \nabla_\tau g + \nabla_\tau(\partial_n g) \cdot \nabla_\tau f \quad (52)$$

Proof: A straightforward computation gives

$$\partial_n(\nabla f \cdot \nabla g) = (D^2 f \nabla g) \cdot \mathbf{n} + (D^2 g \nabla f) \cdot \mathbf{n}$$

and

$$\begin{aligned}\nabla(\partial_{\mathbf{n}}f).\nabla g &= \nabla(\nabla f.\mathbf{n}).\nabla g \\ &= (D^2f\mathbf{n}).\nabla g + (D^2b\nabla f).\nabla g\end{aligned}$$

hence

$$\begin{aligned}\nabla(\partial_{\mathbf{n}}f).\nabla g + \nabla(\partial_{\mathbf{n}}g).\nabla f &= 2(D^2b\nabla f).\nabla g + (D^2f\mathbf{n}).\nabla g + (D^2g\mathbf{n}).\nabla f \\ &= 2(D^2b\nabla f).\nabla g + \partial_{\mathbf{n}}(\nabla f.\nabla g)\end{aligned}$$

We use now the decomposition of ∇ into its normal and tangential components and the well known identity $D^2b\mathbf{n}.\mathbf{n} = 0$. We get

$$\begin{aligned}\nabla_{\tau}(\partial_{\mathbf{n}}f).\nabla_{\tau}g + \nabla_{\tau}(\partial_{\mathbf{n}}g).\nabla_{\tau}f &+ \frac{\partial^2 f}{\partial n^2} \frac{\partial g}{\partial n} + \frac{\partial^2 g}{\partial n^2} \frac{\partial f}{\partial n} = \\ &2(D^2b\nabla_{\tau}f).\nabla_{\tau}g + \partial_{\mathbf{n}}(\nabla_{\tau}f.\nabla_{\tau}g) + \frac{\partial^2 f}{\partial n^2} \frac{\partial g}{\partial n} + \frac{\partial^2 g}{\partial n^2} \frac{\partial f}{\partial n}\end{aligned}\quad (53)$$

hence

$$\nabla_{\tau}(\partial_{\mathbf{n}}f).\nabla_{\tau}g + \nabla_{\tau}(\partial_{\mathbf{n}}g).\nabla_{\tau}f = 2(D^2b\nabla_{\tau}f).\nabla_{\tau}g + \partial_{\mathbf{n}}(\nabla_{\tau}f.\nabla_{\tau}g)$$

■

B Spherical harmonics

In order to explicit the shape hessian under consideration, a useful tool is the surface spherical harmonics defined as the restriction to the surface of the unit sphere of harmonic polynomials in the special case $d = 3$. We recall here facts from [25, pages 139-141]. Spherical Harmonics are defined as restrictions of homogeneous harmonic polynomials to the unit sphere. The spherical harmonics are said of order k when the harmonic homogeneous polynomial is of degree k . We denote by \mathcal{H}_k the space of spherical harmonics of degree k . We show that is also the eigenspace of the Laplace-Beltrami operator on the unit sphere associated with the eigenvalue $k(k+1)$. Its dimension is

$$d_k = 2k + 1.$$

Let $(Y_k^l)_{-k \leq l \leq k}$ be an orthonormal basis of \mathcal{H}_k with respect to the $L^2(\partial B_1)$ scalar product. The $(\mathcal{H}_k)_{k \in \mathbb{N}}$ spans a vector space dense in $L^2(\partial B_1)$ and the family $(Y_k^l)_{k \in \mathbb{N}, -k \leq l \leq k}$ is a Hilbert basis of $L^2(\partial B_1)$. To be more precise, if $f \in L^2(\partial B_1)$, then there exists a unique representation

$$f = \sum_{k=0}^{\infty} \mathbf{Y}_k$$

where the series converge to f in the L^2 norm and

$$\mathbf{Y}_k = \sum_{l=-k}^k b_k^l Y_k^l \in \mathcal{H}_k$$

If $x = (x_1, x_2, x_3) \in \mathbb{R}^3$, it is natural to use on a sphere the spherical coordinates (r, θ, ϕ) where r is the radius and θ and ϕ are the Euler angles. The spherical harmonic Y_k^l is defined with the Euler angles (θ, ϕ) as

$$Y_k^l = (-1)^l \sqrt{\left[\frac{k + \frac{1}{2}(k-l)!}{2\pi (k+l)!} \right]} e^{il\phi} \mathbb{P}_k^l(\cos \theta), \quad -k \leq l \leq k.$$

where the polynomial \mathbb{P}_k^l is the associated Legendre polynomial. The formula giving the explicit form of these polynomials can be found in the book of Nedelec [23, page 24].

When $k \neq k'$, we have also the orthogonality property

$$\int_{\partial B_1} \mathbf{Y}_k \mathbf{Y}_{k'} d\sigma = 0$$

when $\mathbf{Y}_k \in \mathcal{H}_k$ and $\mathbf{Y}_{k'} \in \mathcal{H}_{k'}$. An homogeneity argument shows that any function φ in $L^2(\partial B_R)$ can be decomposed as the Fourier series:

$$\varphi(x) = \sum_{k=0}^{\infty} R^k \left(\sum_{l=-k}^k \alpha_{k,l}(\varphi) Y_k^l \left(\frac{x}{|x|} \right) \right), \quad \text{for } |x| = R.$$

Then, by construction, the function u defined by

$$u(x) = \sum_{k=0}^{\infty} |x|^k \left(\sum_{l=1}^{d_k} \alpha_{k,l}(\varphi) Y_k^l \left(\frac{x}{|x|} \right) \right), \quad \text{for } |x| \leq R,$$

is harmonic in B_R and satisfies $u = \varphi$ on ∂B_R .

We recall now some results about the integration of three spherical harmonics, they will enable us to estimate $\text{Tr}(E)$ in dimension three. When we integrate three spherical harmonics, we use coefficients called Clebsch-Gordon coefficients or Wigner-3j coefficients. The Wigner-3j coefficients are mostly used; they are related to Clebsch-Gordon coefficients via some known formula that the interested reader will find in the book of Cohen-Tannoudji and al [9, Tome 2, Annex B].

The first general result concerns the product of two spherical harmonics; it is given by the following proposition

Proposition B.1 *Given $l_1, l_2 > 0$ two natural integers and $-l_1 \leq m_1 \leq l_1$, $-l_2 \leq m_2 \leq l_2$, we have*

$$Y_{l_1}^{m_1} Y_{l_2}^{m_2} = (-1)^{m_1+m_2} \sum_{L=|l_1-l_2|}^{l_1+l_2} \sqrt{\frac{(2l_1+1)(2l_2+1)(2L+1)}{4\pi}} \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & L \\ m_1 & m_2 & -m_1-m_2 \end{pmatrix} Y_L^{m_1+m_2},$$

where $\begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} l_1 & l_2 & L \\ m_1 & m_2 & -m_1-m_2 \end{pmatrix}$ are the Wigner-3j symbols.

The second result concerns the integration of three spherical harmonics.

Proposition B.2 *We have:*

$$\int_{\partial B_1} Y_{l_1}^{m_1} Y_{l_2}^{m_2} Y_{l_3}^{m_3} = \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{4\pi}} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix}.$$

In particular it holds

Proposition B.3 *Let l be a natural integer and m an integer. We have*

1. If $-l \leq m \leq l$ then

$$\int_{\partial B_1} Y_l^m Y_0^0 \overline{Y_l^m} = \sqrt{\frac{1}{4\pi}},$$

and

$$\int_{\partial B_1} Y_l^m Y_1^1 \overline{Y_{l-1}^{m+1}} = -\sqrt{\frac{3}{8\pi}} \sqrt{\frac{(l-m)(l-m-1)}{(2l+1)(2l-1)}}.$$

2. If $-l-1 \leq m \leq l+1$ then

$$\int_{\partial B_1} Y_l^m Y_1^0 \overline{Y_{l+1}^m} = \sqrt{\frac{3}{4\pi}} \sqrt{\frac{(l+m+1)(l-m+1)}{(2l+1)(2l+3)}},$$

3. If $-l-2 \leq m \leq l$ then

$$\int_{\partial B_1} Y_l^m Y_1^1 \overline{Y_{l+1}^{m+1}} = \sqrt{\frac{3}{8\pi}} \sqrt{\frac{(l+m+1)(l+m+2)}{(2l+1)(2l+3)}},$$

C Intermediate results for the second shape derivative matrix

We need to construct the matrix associated to the second shape derivative. To that end, we have to compute the explicit formula for all the shape derivatives of order one involved in the formula giving λ' (see Theorem 3.6). In this appendix, we focus on the term $E^{(2)}$ introduced in Section 4.1. Since these computations are very technical, we only give the main line and the used arguments, omitting a couple of details. In the following lines, we denote by $H(t)$ the mean curvature associated to the boundary of Ω_t and $\mathcal{A}(t)$ the deviatoric part defined on $\partial\Omega_t$ as

$$\mathcal{A}(t) = H(t)I - 2D^2b(t)$$

(see [13] for the terminology).

In order to deal with the weak formulation on the boundary $\partial\Omega_t$, we will make use of a test function ϕ which is the restriction of a test function Φ defined on a tubular neighborhood of the boundary such that its normal derivative is zero. This kind of extension is well discussed in the book [13] of Delfour-Zolesio.

In this differentiation, nineteen terms arise and we introduce some notations to study them separately. For all function test $\phi \in H^1(\partial\Omega)$, we will need in the sequel the following quantities:

$$\begin{aligned} A(u, u', \phi) &= \left(\frac{d}{dt} \int_{\partial\Omega_t} V_n \nabla_\tau u \cdot \nabla_\tau \phi d\sigma_t \right) \Big|_{t=0}, \\ B(u, u', \phi) &= \beta \left(\frac{d}{dt} \int_{\partial\Omega_t} \mathcal{A}(t) V_n \nabla_\tau u \cdot \nabla_\tau \phi d\sigma_t \right) \Big|_{t=0}, \\ C(u, u', u'', \phi) &= -\frac{d}{dt} \left(\int_{\partial\Omega_t} [\lambda' u + \lambda(u' + V_n \partial_n u + V_n H u)] \phi d\sigma_t \right) \Big|_{t=0}, \\ D(u, u', \phi) &= \beta \left(\frac{d}{dt} \int_{\partial\Omega_t} \nabla_\tau (V_n \partial_n u) \cdot \nabla_\tau \phi d\sigma_t \right) \Big|_{t=0}. \end{aligned}$$

We will now study independently each term A, B, C and D , when $\Omega = B_R \subset \mathbb{R}^2$ or \mathbb{R}^3 , and $t \mapsto \Omega_t$ is volume preserving.

Study of $D(u, u', \phi)$. First, we denote

$$W = \frac{d}{dt} (\mathbf{V} \cdot \mathbf{n}_{\Omega_t})|_{t=0}.$$

From the derivative formula of boundary integrals, we know that we have to compute three main terms: the first corresponding to the shape derivative, the second concerns the normal derivative of the integrand and the third is related to the term related to the mean curvature H . The first term is

$$\begin{aligned} & \beta \left(\int_{\partial\Omega_t} \frac{d}{dt} [\nabla_\tau(V_n \partial_n u) \cdot \nabla_\tau \phi] d\sigma_t \right) \Big|_{t=0} \\ &= \beta \left(\int_{\partial B_R} \nabla_\tau(V_n \cdot \partial_n u' - V_n \nabla_\tau u \cdot \nabla_\tau V_n) \cdot \nabla_\tau \phi d\sigma + \int_{\partial B_R} \nabla_\tau(V_n' \cdot \partial_n u) \cdot \nabla_\tau \phi d\sigma \right) \\ & \quad + \beta \int_{\partial B_R} \partial_n(V_n \partial_n u) \nabla_\tau V_n \cdot \nabla_\tau \phi d\sigma \\ &= -\beta \int_{\partial B_R} \Delta_\tau(V_n \cdot \partial_n u') \phi d\sigma + \beta \int_{\partial B_R} \nabla_\tau(V_n' \cdot \partial_n u) \cdot \nabla_\tau \phi d\sigma \\ & \quad + \beta \int_{\partial B_R} \partial_n(V_n \partial_n u) \nabla_\tau V_n \cdot \nabla_\tau \phi d\sigma + \beta \int_{\partial B_R} \Delta_\tau(V_n \nabla_\tau u \cdot \nabla_\tau V_n) \phi d\sigma. \end{aligned}$$

The third term is

$$\beta \int_{\partial B_R} H V_n \nabla_\tau(V_n \partial_n u) \cdot \nabla_\tau \phi d\sigma = -\beta \int_{\partial B_R} \operatorname{div}_\tau \left(H V_n \nabla_\tau(V_n \partial_n u) \right) \phi d\sigma.$$

We focus now on the second term. We have

$$\begin{aligned} & \beta \int_{\partial B_R} V_n \partial_n [\nabla_\tau(V_n \partial_n u) \cdot \nabla_\tau \phi] d\sigma \\ &= \beta \int_{\partial B_R} V_n \nabla_\tau [\partial_n(V_n \partial_n u)] \cdot \nabla_\tau \phi d\sigma - 2\beta \int_{\partial B_R} V_n (D^2 b \nabla_\tau [V_n \partial_n u]) \cdot \nabla_\tau \phi d\sigma \\ &= \beta \left(\int_{\partial B_R} V_n \nabla_\tau [\partial_n V_n \partial_n u] \cdot \nabla_\tau \phi d\sigma - 2 \int_{\partial B_R} V_n (D^2 b \nabla_\tau [V_n \partial_n u]) \cdot \nabla_\tau \phi d\sigma \right) \\ &= -\beta \int_{\partial B_R} \operatorname{div}_\tau [V_n \nabla_\tau [\partial_n u \partial_n V_n] - 2V_n D^2 b \nabla_\tau [V_n \partial_n u]] \phi d\sigma. \end{aligned}$$

We expand $D(u, \phi)$ into a sum $\langle D^{(1)}u', \phi \rangle + \langle D^{(2)}u, \phi \rangle$. For $D^{(2)}$, we will set $D^{(2)} = \sum_{k=1}^3 D^{(2,k)}$ where

$$\begin{aligned} \langle D^{(1)}u', \phi \rangle &= \beta \int_{\partial B_R} \nabla_\tau [V_n \cdot \partial_n u'] \cdot \nabla_\tau \phi d\sigma = -\beta \int_{\partial B_R} \Delta_\tau [V_n \partial_n u'] \phi d\sigma \\ \langle D^{(2,1)}u, \phi \rangle &= \beta \left[\int_{\partial B_R} (-\Delta_\tau [W \partial_n u] \phi d\sigma - \int_{\partial B_R} \operatorname{div}_\tau [V_n \partial_n V_n \nabla_\tau [\partial_n u]] \phi d\sigma, \right. \\ & \quad \left. - \int_{\partial B_R} \operatorname{div}_\tau [H V_n \nabla_\tau (V_n \partial_n u)] \phi d\sigma \right], \\ \langle D^{(2,2)}u, \phi \rangle &= -\beta \int_{\partial B_R} \operatorname{div}_\tau [\partial_n u \partial_n V_n \nabla_\tau V_n] \phi d\sigma + \beta \int_{\partial B_R} \Delta_\tau [V_n \partial_n u \nabla_\tau V_n] \phi d\sigma, \\ \langle D^{(2,3)}u, \phi \rangle &= 2\beta \int_{\partial B_R} \operatorname{div}_\tau [V_n D^2 b \cdot \nabla_\tau [V_n \partial_n u]] \phi d\sigma. \end{aligned}$$

We denote $D^{(1)}$ and $D^{(2,k)}$, $k = 1, 2, 3$ the matrices whose elements are defined by

$$D_{ij}^{(1)} = \langle D^{(1)} \tilde{u}_i, u_j \rangle, \text{ and } D_{ij}^{(2,k)} = \langle D^{(2,k)} u_i, u_j \rangle, \quad i, j = 1, 2, \dots, d.$$

We give a result concerning the traces of the matrices.

Lemma C.1 *We have*

$$\text{Tr}(D^{(2,1)}) = \text{Tr}(D^{(2,2)}) = 0 \text{ and } \text{Tr}(D^{(2,3)}) = -\frac{2\beta(d-1)K(R)}{R} \int_{\partial B_R} V_n^2 \, d\sigma,$$

with the normalization constant $K(R) = \frac{d}{R^{2+d\omega_d}}$.

Proof of Lemma C.1:

We have

$$\begin{aligned} \text{Tr}(D^{(2,1)}) &= \beta \left[\int_{\partial B_R} -\Delta_\tau \left(W \sum_{i=1}^d \partial_n u_i \right) u_i \, d\sigma - \int_{\partial B_R} \text{div}_\tau \left(V_n \partial_n V_n \sum_{i=1}^d \nabla_\tau (\partial_n u_i) \right) u_i \, d\sigma \right. \\ &\quad \left. - \int_{\partial B_R} \sum_{i=1}^d \text{div}_\tau \left(H V_n \nabla_\tau (V_n \partial_n u_i) \right) u_i \, d\sigma \right] \\ &= \beta \int_{\partial B_R} V_n' (d-1) \sum_{i=1}^d |\partial_n u_i|^2 \frac{d\sigma}{R} + \beta \int_{\partial B_R} V_n \partial_n V_n \sum_{i=1}^d \nabla_\tau (\partial_n u_i) \cdot \nabla_\tau u_i \, d\sigma \\ &\quad + \beta \int_{\partial B_R} H V_n^2 \sum_{i=1}^d \nabla_\tau (\partial_n u_i) \cdot \nabla_\tau u_i \, d\sigma + \beta \int_{\partial B_R} H \sum_{i=1}^d \partial_n u_i V_n \nabla_\tau V_n \cdot \nabla_\tau u_i \, d\sigma \end{aligned} \quad (54)$$

Combining the two facts (coming from algebraic properties of spherical harmonics, see Appendix B),

$$(d-1) \sum_{i=1}^d \frac{|\partial_n u_i|^2}{R} = \sum_{i=1}^d \nabla_\tau (\partial_n u_i) \cdot \nabla_\tau u_i = \frac{d(d-1)}{R^{2+d\omega_d}} = (d-1)K(R). \quad (55)$$

and

$$\int_{\partial B_R} V_n \sum_{i=1}^d \partial_n u_i \nabla_\tau V_n \cdot \nabla_\tau u_i = 0, \quad (56)$$

we get

$$\text{Tr}(D^{(2,1)}) = (d-1) \sum_{i=1}^d |\partial_n u_i|^2 \int_{\partial B_R} \left(W + V_n \partial_n V_n + H V_n^2 \right) \frac{d\sigma}{R}.$$

Since we assumed the deformation to be volume preserving up to the second order (41), we have $\text{Tr}(D^{(2,1)}) = 0$. The same strategy applies for $\text{Tr}(D^{(2,2)})$.

We focus now on $\text{Tr}(D^{(2,3)})$. We first expand the second term in the definition of $D^{(4)}$:

$$\begin{aligned} \text{Tr}(D^{(2,3)}) &= \beta \sum_{i=1}^d \int_{\partial B_R} V_n \partial_n u_i \nabla_\tau [\partial_n V_n] \cdot \nabla_\tau u_i - 2V_n \partial_n u_i D^2 b \nabla_\tau V_n \cdot \nabla_\tau u_i \, d\sigma \\ &\quad - \beta \sum_{i=1}^d \int_{\partial B_R} 2V_n^2 D^2 b \nabla_\tau (\partial_n u_i) \cdot \nabla_\tau u_i \, d\sigma. \end{aligned}$$

We follow the same argument thanks to the relations (55)-(56) and the fact

$$\sum_{i=1}^d D^2 b \nabla_\tau (\partial_n u_i) \cdot \nabla_\tau u_i = \frac{(d-1)K(R)}{R}$$

on the sphere. Recall that on the sphere $D^2 b = I_d/R$ when restricted to the tangent space. \blacksquare

Study of $B(u, u', \phi)$. In the same manner, we begin to compute the derivative of the integrand:

$$\begin{aligned} \frac{d}{dt} \left(\mathcal{A}(t) V_n \nabla_\tau u \cdot \nabla_\tau \phi \right) \Big|_{t=0} = \\ \mathcal{A}' V_n \nabla_\tau u \cdot \nabla_\tau \phi + \mathcal{A} V_n' \nabla_\tau u \cdot \nabla_\tau \phi + \mathcal{A} V_n \nabla_\tau u' \cdot \nabla_\tau \phi - \mathcal{A} V_n \partial_n u \nabla_\tau V_n \cdot \nabla_\tau \phi. \end{aligned}$$

Denote $\mathcal{A} = (a_{ij})_{1 \leq i, j \leq d}$ and $\tilde{\mathcal{A}} = (\partial_n a_{ij})_{1 \leq i, j \leq d}$. Thanks to lemma A.3, we get

$$V_n \partial_n \left(V_n \mathcal{A} \cdot \nabla_\tau u \cdot \nabla_\tau \phi \right) = V_n^2 \tilde{\mathcal{A}} \cdot \nabla_\tau u \cdot \nabla_\tau \phi + V_n \partial_n V_n \mathcal{A} \cdot \nabla_\tau u \cdot \nabla_\tau \phi + V_n^2 \mathcal{A} \partial_n \left[\nabla_\tau u \cdot \nabla_\tau \phi \right].$$

From the relation

$$\begin{aligned} \beta \frac{d}{dt} \int_{\partial \Omega_t} \mathcal{A}(t) V_n \nabla_{\partial \Omega_t} u \cdot \nabla_{\partial \Omega_t} \phi \, d\sigma_t \Big|_{t=0} = \int_{\partial B_R} \frac{d}{dt} \left(\mathcal{A}(t) V_n \nabla_{\partial \Omega_t} u \cdot \nabla_{\partial \Omega_t} \phi \right) \Big|_{t=0} \, d\sigma \\ + \int_{\partial B_R} V_n \partial_n \left(\mathcal{A} V_n \nabla_\tau u \cdot \nabla_\tau \phi \right) \, d\sigma + \int_{\partial B_R} H V_n^2 \mathcal{A} \nabla_t u \cdot \nabla_\tau \phi \, d\sigma_t, \end{aligned}$$

we gather all the terms and obtain $B(u, \phi) = \langle B^{(1)} u', \phi \rangle + \langle B^{(2)} u', \phi \rangle$; we then set

$$\langle B^{(2)} u, \phi \rangle = \sum_{i=1}^4 \langle B^{(2,i)} u, \phi \rangle,$$

where

$$\begin{aligned} \langle B^{(2,1)} u', \phi \rangle &= -\beta \int_{\partial B_R} \operatorname{div}_\tau [V_n \mathcal{A} \cdot \nabla_\tau u'] \phi \, d\sigma, \\ \langle B^{(2,1)} u, \phi \rangle &= -\beta \int_{\partial B_R} \operatorname{div}_\tau [(W + H V_n^2 + V_n \partial_n V_n) \mathcal{A} \cdot \nabla_\tau u] \phi \, d\sigma, \\ \langle B^{(2,2)} u, \phi \rangle &= -\beta \int_{\partial B_R} \operatorname{div}_\tau [\partial_n u V_n \mathcal{A} \cdot \nabla_\tau V_n] \phi \, d\sigma, \\ \langle B^{(2,3)} u, \phi \rangle &= -\beta \int_{\partial B_R} \operatorname{div}_\tau [V_n \mathcal{A}' \cdot \nabla_\tau u] \phi \, d\sigma, \\ \langle B^{(2,4)} u, \phi \rangle &= \beta \int_{\partial B_R} V_n^2 \partial_n [\mathcal{A} \cdot \nabla_\tau u \cdot \nabla_\tau \phi] \, d\sigma. \end{aligned}$$

We get

$$\begin{aligned} B^{(2,4)} u, \phi &= \beta \int_{\partial B_R} V_n^2 (\partial_n [\mathcal{A}] \cdot \nabla_\tau u \cdot \nabla_\tau \phi) \, d\sigma + \beta \int_{\partial B_R} V_n^2 \mathcal{A} \cdot \nabla_\tau \partial_n u \cdot \nabla_\tau \phi \, d\sigma \\ &\quad - \beta \int_{\partial \Omega} 2(D^2 b \mathcal{A}) \cdot \nabla_\tau u \cdot \nabla_\tau \phi \, d\sigma \\ &= -\beta \int_{\partial B_R} \operatorname{div}_\tau \left[V_n^2 \left(\tilde{\mathcal{A}} \cdot \nabla_\tau u + \mathcal{A} \cdot \nabla_\tau [\partial_n u] \right) - 2D^2 b \mathcal{A} \cdot \nabla_\tau u \right] \phi \, d\sigma \end{aligned}$$

Let $B^{(2,k)}$ $k = 1, 2, 3, 4$ denote the respective matrices associated to the operator with respect to the basis of eigenvectors. We have the following result:

Lemma C.2 *We have*

$$\mathrm{Tr}\left(\sum_{i=1}^4 B^{(2,i)}\right) = -\beta(d-1)RK(R) \int_{\partial B_R} |\nabla_\tau V_n|^2 d\sigma + 2\frac{\beta K(R)}{R} \int_{\partial B_R} V_n^2 d\sigma.$$

Proof of Lemma C.2:

Using the same arguments as before, we prove easily that $\mathrm{Tr}(B^{(2,1)}) = \mathrm{Tr}(B^{(2,2)}) = 0$.

For the other terms, above all we have to focus on the term

$$\mathrm{Tr}(B^{(2,3)}) = \beta \int_{\partial B_R} V_n \sum_{i=1}^d (\mathcal{A}' \cdot \nabla_\tau u_i) \cdot \nabla_\tau u_i d\sigma.$$

We have, thanks to the expression of shape derivation of the normal vector and of the mean curvature given in Proposition A.2:

$$\mathcal{A}(t) = H(t) - 2D^2b(t) \Rightarrow \mathcal{A}' = -\Delta_\tau V_n + 2D(\nabla_\tau V_n);$$

then

$$\begin{aligned} \mathrm{Tr}(B^{(2,3)}) &= \beta \int_{\partial B_R} V_n \sum_{i=1}^d (\mathcal{A}' \cdot \nabla_\tau u_i) \cdot \nabla_\tau u_i d\sigma \\ &= -\beta \int_{\partial B_R} V_n \Delta_\tau V_n \sum_{i=1}^d |\nabla_\tau u_i|^2 d\sigma + 2\beta \int_{\partial B_R} V_n \sum_{i=1}^d \left[D(\nabla_\tau V_n) \cdot \nabla_\tau u_i \right] \cdot \nabla_\tau u_i d\sigma \\ &= -\beta \int_{\partial B_R} V_n \Delta_\tau V_n \sum_{i=1}^d |\nabla_\tau u_i|^2 d\sigma + 2\beta \int_{\partial B_R} V_n \sum_{i=1}^d \left[D_\tau(\nabla_\tau V_n) \cdot \nabla_\tau u_i \right] \cdot \nabla_\tau u_i d\sigma \\ &= -\beta \int_{\partial B_R} V_n \Delta_\tau V_n \sum_{i=1}^d |\nabla_\tau u_i|^2 d\sigma + 2\beta \int_{\partial B_R} V_n \sum_{i=1}^d \left[D_\tau^2 V_n \cdot \nabla_\tau u_i \right] \cdot \nabla_\tau u_i d\sigma \\ &= -\beta \int_{\partial B_R} V_n \Delta_\tau V_n \sum_{i=1}^d |\nabla_\tau u_i|^2 d\sigma + 2\beta \int_{\partial B_R} V_n \mathrm{Tr}(D_\tau^2 V_n) \sum_{i=1}^d |\nabla_\tau u_i|^2 d\sigma \end{aligned}$$

Since $\mathrm{Tr}(D_\tau^2 V_n) = \Delta_\tau V_n$, and since $\sum_{i=1}^d |\nabla_\tau u_i|^2 = RK(R)$, on ∂B_R we get

$$\mathrm{Tr}(B^{(2,3)}) = \beta \int_{\partial B_R} V_n \Delta_\tau V_n \sum_{i=1}^d |\nabla_\tau u_i|^2 d\sigma = \beta(d-1)RK(R) \int_{\partial B_R} V_n \Delta_\tau V_n d\sigma.$$

Concerning $\mathrm{Tr}(B^{(2,4)})$, we have to distinguish the case $d = 2$ from the case $d = 3$. If $d = 3$ then $\mathcal{A} = 0$; this implies that $\mathrm{Tr}(B^{(2,4)})$ is reduced to

$$\mathrm{Tr}(B^{(2,4)}) = (d-1)K(R) \frac{\beta}{R} \int_{\partial B_R} V_n^2 d\sigma.$$

If $d = 2$, then $\mathcal{A} + \tilde{\mathcal{A}}$ is a null matrix and this leads to

$$\begin{aligned} \mathrm{Tr}(B^{(2,4)}) &= 2\beta \int_{\partial B_R} V_n^2 \sum_{i=1}^d D^2b \cdot \nabla_\tau u_i \cdot \nabla_\tau u_i d\sigma \\ &= 2K(R) \frac{\beta}{R} \int_{\partial B_R} V_n^2 d\sigma. \end{aligned}$$

Then for $d = 2, 3$ we get

$$\text{Tr}(B^{(2,4)}) = 2\beta \frac{K(R)}{R};$$

■

Study of $A(u, u', \phi)$. We have

$$\begin{aligned} \frac{d}{dt} \left(\int_{\partial\Omega_t} V_n \nabla_\tau u \cdot \nabla_\tau \phi d\sigma_t \right) \Big|_{t=0} &= \int_{\partial B_R} W \nabla_\tau u \cdot \nabla_\tau \phi d\sigma + \int_{\partial B_R} V_n \nabla_\tau u' \cdot \nabla_\tau \phi d\sigma \\ &+ \int_{\partial B_R} V_n \nabla_\tau V_n \cdot \left[\partial_n u \nabla_\tau \phi + \partial_n \phi \nabla_\tau u \right] + \left(V_n \partial_n [V_n \nabla_\tau u \cdot \nabla_\tau \phi] + H V_n^2 \nabla_\tau u \cdot \nabla_\tau \phi \right) d\sigma. \end{aligned}$$

Since $\partial_n \phi = 0$, it comes that

$$\begin{aligned} \int_{\partial B_R} V_n \nabla_\tau V_n \cdot \left[\partial_n u \nabla_\tau \phi + \partial_n \phi \nabla_\tau u \right] d\sigma &= -\frac{1}{2} \int_{\partial B_R} V_n^2 \left[\partial_n u \Delta_\tau \phi + \nabla_\tau [\partial_n u] \cdot \nabla_\tau \phi \right] \\ &= -\frac{1}{2} \int_{\partial B_R} V_n^2 \left(\partial_n [\nabla_\tau u \cdot \nabla_\tau \phi] + 2D^2 b \nabla_\tau u \cdot \nabla_\tau \phi \right) d\sigma - \frac{1}{2} \int_{\partial B_R} V_n^2 \partial_n u \Delta_\tau \phi d\sigma \end{aligned}$$

Hence, gathering the equivalent terms we get

$$\begin{aligned} \frac{d}{dt} \int_{\partial\Omega_t} V_n \nabla_\tau u \cdot \nabla_\tau \phi d\sigma_t \Big|_{t=0} &= \int_{\partial B_R} W \nabla_\tau u' \cdot \nabla_\tau \phi d\sigma + \int_{\partial B_R} V_n \nabla_\tau u' \cdot \nabla_\tau \phi d\sigma \\ &- \frac{1}{2} \int_{\partial B_R} \Delta_\tau [V_n^2 \partial_n u] \phi - \partial_n (V_n^2 \nabla_\tau u \cdot \nabla_\tau \phi) d\sigma + \int_{\partial B_R} (H I_d - D^2 b) V_n^2 \nabla_\tau u \cdot \nabla_\tau \phi d\sigma. \end{aligned}$$

We split these terms into $A(u, \phi) = \langle A^{(1)} u', \phi \rangle + \langle A^{(2)} u, \phi \rangle$. As before, we set $\langle A^{(2)} u, \phi \rangle = \sum_{i=1}^3 \langle A^{(i)} u, \phi \rangle$ where

$$\begin{aligned} \langle A^{(1)} u', \phi \rangle &= \int_{\partial B_R} -\text{div}_\tau [V_n \nabla_\tau u'] \phi d\sigma, \\ \langle A^{(2,1)} u, \phi \rangle &= \int_{\partial B_R} -\text{div}_\tau [(W + H V_n^2 + V_n \partial_n V_n) \nabla_\tau u] \phi d\sigma, \\ \langle A^{(2,2)} u, \phi \rangle &= \int_{\partial B_R} \text{div}_\tau [\partial_n u V_n \nabla_\tau V_n] \phi d\sigma, \\ \langle A^{(2,3)} u, \phi \rangle &= \int_{\partial B_R} \text{div}_\tau [V_n^2 (2D^2 b \nabla_\tau u - \nabla_\tau (\partial_n u))] \phi d\sigma. \end{aligned}$$

We have

Lemma C.3 *We have*

$$\text{Tr}(A^{(2,1)}) = 0, \quad \text{Tr}(A^{(2,2)}) = 0 \quad \text{and} \quad \text{Tr}(A^{(2,3)}) = -K(R) \int_{\partial B_R} V_n^2 d\sigma.$$

The proof of Lemma C.3 follows the lines of the proof of Lemma C.2.

Study of $C(u, u', u'', \phi)$. We decompose $C(u, u', u'', \phi)$ as follows:

$$C(u, \phi) = \langle C^{(0)}u'', \phi \rangle + \langle C^{(1)}u', \phi \rangle + \langle C^{(2)}u, \phi \rangle$$

with $\langle C^{(2)}u, \phi \rangle = \sum_{i=3}^6 \langle C^{(2,i)}u, \phi \rangle$ where

$$\begin{aligned} \langle C^{(0)}u'', \phi \rangle &= -\lambda \int_{\partial B_R} u'' \phi \, d\sigma \\ \langle C^{(1)}u', \phi \rangle &= -2 \int_{\partial B_R} (\lambda' u' + \lambda V_n (\partial_n u' + H u')) \phi \, d\sigma \\ \langle C^{(2,1)}u, \phi \rangle &= -\lambda'' \int_{\partial B_R} u \phi - \lambda' \int_{\partial B_R} V_n \partial_n u \phi \, d\sigma \\ \langle C^{(2,2)}u, \phi \rangle &= -\lambda \int_{\partial B_R} (W + V_n \partial_n V_n + H V_n^2) (\partial_n u + H u) \phi \, d\sigma \\ \langle C^{(2,3)}u, \phi \rangle &= -\lambda \int_{\partial B_R} V_n \left(-\nabla_\tau V_n \cdot \nabla_\tau u + H' u \right) \phi \, d\sigma \\ &= \lambda \int_{\partial B_R} V_n \left(\nabla_\tau V_n \cdot \nabla_\tau u + \Delta_\tau V_n u \right) \phi \, d\sigma \\ \langle C^{(2,4)}u, \phi \rangle &= -\lambda \int_{\partial B_R} V_n^2 \left(\partial_n^2 u - u \sum_{i=1}^{d-1} \kappa_i^2 + H \partial_n u \right) \phi \, d\sigma \\ &= 0. \end{aligned}$$

Denoting by $(C^{(2,j)})$, $j = 1, 2, 3, 4$ the matrices associated to the linear operators $C^{(2,p)}$, $p = 1, 2, 3, 4$ in the basis of eigenvectors, we get:

Lemma C.4 *We have*

$$\sum_{j=1}^4 \text{Tr}(C^{(2,j)}) = \lambda R^3 K(R) \int_{\partial B_R} V_n \Delta_\tau V_n \, d\sigma = -((d-1)\beta + R) R K(R) \int_{\partial B_R} |\nabla_\tau V_n|^2 \, d\sigma.$$

Proof of Lemma C.4:

The proof is straightforward and obeys to the same arguments used before. The only non null trace concerns the factor in $-H' = \Delta_\tau V_n$. \blacksquare

D Computing u'

In this section, we focus on the computation of the trace of $E^{(1)}$ introduced in Section 4.1. We recall that $t \mapsto (\lambda(t), u(t, \cdot))$ is solution of

$$\begin{aligned} \Delta u &= 0 & \text{in } T_t(B_R), \\ -\beta \Delta_\tau u + \partial_n u - \lambda(t)u &= 0 & \text{on } \partial T_t(B_R). \end{aligned} \tag{57}$$

To compute the second derivative, one must know $u' = u'(0)$. For the reader convenience, we recall the problem (36) solved by u' .

$$\begin{aligned} \Delta u' &= 0 \text{ in } B_R, \\ -\beta \Delta_\tau u' + \partial_n u' - \lambda v' &= \beta \Delta_\tau (V_n \partial_n u) - \beta \text{div}_\tau (V_n (2D^2 b - H I_d) \nabla_\tau u) \\ &\quad + \text{div}_\tau (V_n \nabla_\tau u) - \lambda' u + \lambda V_n (\partial_n u + H u) \text{ on } \partial B_R. \end{aligned}$$

First, Fredholm's alternative insures the existence of a unique harmonic function \tilde{u}_j orthogonal to the eigenfunctions u_1, u_2, \dots, u_d and satisfying on ∂B_R the boundary condition

$$\begin{aligned} -\beta \Delta_\tau \tilde{u}_j + \partial_n \tilde{u}_j - \lambda \tilde{u}_j &= \beta \left[\Delta_\tau [V_n \partial_n u_j] + \operatorname{div}_\tau [V_n (HI_d - 2D^2 b) \cdot \nabla_\tau u_j] \right] \\ &\quad + \operatorname{div}_\tau [V_n \nabla_\tau u_j] + \lambda' u_j + \lambda V_n (\partial_n u_j + H u_j). \end{aligned} \quad (58)$$

It follows that

$$u' = \sum_{j=1}^d \tilde{c}_j u_j + \sum_{j=1}^m c_j \tilde{u}_j \quad (59)$$

for some c_j, \tilde{c}_j when $j = 1, \dots, d$. We point out that the (c_j) are the same coefficients as the decomposition of u in the basis (u_j) of the eigenspace associated to λ : $u = c_1 u_1 + \dots + c_d u_d$.

Remark D.1 *We recall that we only need the terms \tilde{u}_j : we inject this decomposition of u' in $E^{(1)}$:*

$$\begin{aligned} E^{(1)} \phi &= -2 \sum_{j=1}^d \tilde{c}_j \left[\int_{\partial B_R} V_n \partial_n u_j \partial_n \phi \, d\sigma + 2 \frac{R + \beta(d-3)}{R} \int_{\partial B_R} V_n \nabla_\tau u_j \cdot \nabla_\tau \phi \, d\sigma \right] \\ &\quad - 2 \sum_{j=1}^m c_j \left[\int_{\partial B_R} V_n \partial_n \tilde{u}_j \partial_n \phi \, d\sigma + 2 \frac{R + \beta(d-3)}{R} \int_{\partial B_R} V_n \nabla_\tau \tilde{u}_j \cdot \nabla_\tau \phi \, d\sigma \right. \\ &\quad \left. - 2\lambda \int_{\partial B_R} V_n H u_j \phi \, d\sigma - 2\lambda \int_{\partial B_R} V_n H \tilde{u}_j \phi \, d\sigma \right]. \end{aligned}$$

By construction the first sum cancels and we simply get

$$E_{jk}^{(1)} = 2 \int_{\partial \Omega} V_n \left(-\partial_n \tilde{u}_j \partial_n u_k - H \lambda \tilde{u}_j u_k + (I + \beta (HI_d - 2D^2 b)) \nabla_\tau \tilde{u}_j \cdot \nabla_\tau u_k \right) d\sigma$$

D.1 Explicit resolution of (58) to compute \tilde{u}_j

Let us now compute \tilde{u}_j solution of (58). This step consists in technical computations. For the completeness of the presentation, we present the case of dimension three, we will then simply state the results in dimension two. From now on, we do not consider the case $d \geq 4$ for technical reasons.

D.1.1 Explicit representation of \tilde{u}_j in the case $d = 2$.

We illustrate the computation of the elements \tilde{u}_i , $i = 1, 2$ in the case $d = 2$. The eigenfunctions are the normalized coordinates functions that is (u_1, u_2) given by

$$u_1(r, \theta) = r \frac{\cos \theta}{\sqrt{\pi R^3}} \text{ and } u_2(r, \theta) = r \frac{\sin \theta}{\sqrt{\pi R^3}}.$$

We have

Lemma D.2 *Let V be a deformation of normal component $V_n = R^k (v_1^{(k)} \cos k\theta + v_2^{(k)} \sin k\theta)$, then*

$$\begin{aligned} \tilde{u}_1(r, \theta) &= \frac{r^{k+1}}{2\sqrt{\pi R^{\frac{7}{2}}}} \frac{1-k}{k} \left[v_1^{(k)} \cos(k+1)\theta + v_2^{(k)} \sin(k+1)\theta \right] \\ &\quad + \frac{r^{k-1}}{2\sqrt{\pi R^{\frac{3}{2}}}} \frac{1+k}{k-2} \left[\frac{\beta(2-k) + R}{k\beta + R} \right] \left[v_1^{(k)} \cos(k-1)\theta + v_2^{(k)} \sin(k-1)\theta \right] \end{aligned} \quad (60)$$

and

$$\begin{aligned}\tilde{u}_2(r, \theta) &= \frac{r^{k+1}}{2\sqrt{\pi}R^{\frac{7}{2}}} \frac{1-k}{k} \left[-v_2^{(k)} \cos(k+1)\theta + v_1^k \sin(k+1)\theta \right] \\ &\quad + \frac{r^{k-1}}{2\sqrt{\pi}R^{\frac{3}{2}}} \frac{1+k}{k-2} \left[\frac{\beta(2-k)+R}{k\beta+R} \right] \left[v_2^{(k)} \cos(k-1)\theta - v_1^k \sin(k-1)\theta \right]\end{aligned}\tag{61}$$

In order to justify these formulae, one has to compute a, b, c, d the coefficients

$$\tilde{u}_j = a^{(k)} \cos(k+1)\theta + b^{(k)} \sin(k+1)\theta + c^{(k)} \cos(k-1)\theta + d^{(k)} \sin(k-1)\theta$$

such that \tilde{u}_j satisfies (58) with $u_i = \frac{x_i}{\|x_i\|_{L^2(\partial B_R)}}$. We left the tedious computations to the reader.

D.1.2 Explicit representation of \tilde{u}_j in the case $d = 3$

We begin with the case where $V_n = r^l Y_l^m$ and $\varphi_p = r Y_1^p$ where $-l \leq m \leq l$ and $-1 \leq p \leq 1$. We introduce the coefficients:

$$C_{l-1,p}^{(l,1,m,p)} = (-1)^{m+p} \sqrt{\frac{3(2l-1)(2l+1)}{4\pi}} \begin{pmatrix} l & 1 & l-1 \\ m & p & -m-p \end{pmatrix} \begin{pmatrix} l & 1 & l-1 \\ 0 & 0 & 0 \end{pmatrix},$$

and

$$C_{l+1,p}^{(l,1,m,p)} = (-1)^{m+p} \sqrt{\frac{3(2l+1)(2l+3)}{4\pi}} \begin{pmatrix} l & 1 & l+1 \\ m & p & -m-p \end{pmatrix} \begin{pmatrix} l & 1 & l+1 \\ 0 & 0 & 0 \end{pmatrix},$$

where we use the Wigner $3j$ symbol and Clebsch-Gordan coefficients. We set $\alpha = \beta/R$ in order to obtain an adimensional constant.

Lemma D.3 *Let $l \neq 0$ be a natural integer and let $-l \leq m \leq l$. Let $V_n = r^l Y_l^m$ and $u_p = r Y_1^p$ where $-1 \leq p \leq 1$. The unique solution of (58) that is orthogonal to $\text{Span}(Y_1^{-1}, Y_1^0, Y_1^1)$ is given by*

$$\tilde{u}_p = a_{l-1,p,\alpha}^{(l,1,m,p)} r^{l-1} Y_{l-1}^{m+p} + a_{l+1,p,\alpha}^{(l,1,m,p)} \frac{r^{l+1}}{R^2} Y_{l+1}^{m+p}$$

where

$$a_{l-1,p,\alpha}^{(l,1,m,p)} = \frac{l+2}{l-2} \frac{1+\alpha(3-l)}{1+\alpha(1+l)} C_{l-1,p}^{(l,1,m,p)} \quad \text{and} \quad a_{l+1,p,\alpha}^{(l,1,m,p)} = \frac{l-1}{l} \frac{1+\alpha(4+l)}{1+\alpha(3+l)} C_{l+1,p}^{(l,1,m,p)}.$$

Proof of Lemma D.3:

We first decompose the right hand side of (58) into the basis of spherical harmonics. Taking into account that

$$\begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix} = 0$$

whenever (l_1, l_2, L) satisfies the triangular inequality and $l_1 + l_2 + L$ is odd, we get

$$\beta V_n \partial_n u_p = \beta R^l Y_l^m Y_1^p = \beta R^l \left[C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} + C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right]$$

and then

$$\beta \Delta_\tau (V_n \partial_n u_p) = \alpha R^{l-1} \left[l(1-l) C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} - (l+1)(l+2) C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right].$$

We also have

$$\begin{aligned}
\nabla_\tau V_n \cdot \nabla_\tau u_p &= \frac{1}{2} [\Delta_\tau (V_n u_p) - V_n \Delta_\tau u_p - u_p \Delta_\tau V_n] \\
&= \frac{R^{l-1}}{2} \left[l(1-l) C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} - (l+1)(l+2) C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right. \\
&\quad + 2 C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} + 2 C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \\
&\quad \left. + l(l+1) C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} + l(l+1) C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right] \\
&= R^{l-1} \left[(l+1) C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} - l C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right].
\end{aligned}$$

Since $\operatorname{div}_\tau V_n \nabla_\tau u_p = \nabla_\tau V_n \cdot \nabla_\tau u_p + V_n \Delta_\tau u_p$, it comes

$$\operatorname{div}_\tau V_n \nabla_\tau u_p = R^{l-1} \left[(l-1) C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} - (l+2) C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right].$$

Hence, gathering the various terms in the right hand side of (58), we see that \tilde{u}_p is solution of

$$\begin{aligned}
-\beta \Delta_\tau \tilde{u}_p + \partial_n \tilde{u}_p - \lambda_2 \tilde{u}_p &= \\
R^{l-1} \left[(l+2)(1+\alpha(3-l)) C_{l-1,p}^{(l,1,m,p)} Y_{l-1}^{m+p} + (1-l)(1+\alpha(4+l)) C_{l+1,p}^{(l,1,m,p)} Y_{l+1}^{m+p} \right].
\end{aligned}$$

After identification, we obtain:

$$\tilde{u}_p = a_{l-1,p,\alpha}^{(l,1,m,p)} r^{l-1} Y_{l-1}^{m+p} + a_{l+1,p,\alpha}^{(l,1,m,p)} \frac{r^{l+1}}{R^2} Y_{l+1}^{m+p},$$

where the coefficients $a_{l\pm 1,p,\alpha}^{(l,1,m,p)}$ are defined in Lemma D.3. ■

As a corollary, we deduce the general case for V_n .

Corollary D.4 *If*

$$V_n = \sum_{l=2}^{\infty} r^l \sum_{m=-l}^l v_{l,m} Y_l^m \text{ and } u_p = \sum_{p=-1}^1 \alpha_p Y_1^p,$$

then

$$\tilde{u}_p = \sum_{l=2}^{\infty} \sum_{m=-l}^l \sum_{p=-1}^1 \alpha_p v_{l,m} \left[a_{l-1,p,\alpha}^{(l,1,m,p)} r^{l-1} Y_{l-1}^{m+p} + a_{l+1,p,\alpha}^{(l,1,m,p)} \frac{r^{l+1}}{R^2} Y_{l+1}^{m+p} \right].$$

D.2 The explicit expression of the trace of $E^{(1)}$

We leave the tedious but easy computations of the case $d = 2$ to the reader; the obtained result is written in (46). We focus here on the much more technical case $d = 3$.

We set $u_j = K(R)(\alpha_{-1}^j Y_1^{-1} + \alpha_0^j Y_1^0 + \alpha_1^j Y_1^1)$ for $1 \leq j \leq 3$ where

$$\begin{aligned}
\alpha_{-1}^1 &= 1/\sqrt{2}, & \alpha_0^1 &= 0, & \alpha_1^1 &= 1/\sqrt{2}, \\
\alpha_{-1}^2 &= 0, & \alpha_0^2 &= 1, & \alpha_1^2 &= 0, \\
\alpha_{-1}^3 &= -i/\sqrt{2}, & \alpha_0^3 &= 0, & \alpha_1^3 &= i/\sqrt{2}.
\end{aligned}$$

On the sphere in dimension 3, the deviatoric part of the curvature cancels and the entries of $E^{(1)}$ are

$$\mathrm{Tr}(E^{(1)}) = \sum_{j=1}^3 E_{jj}^{(1)} \text{ where } E_{jj}^{(1)} = \int_{\partial\Omega} V_n \left(-\partial_n \tilde{u}_j \partial_n u_j - H \lambda \tilde{u}_j u_j + \nabla_\tau \tilde{u}_j \cdot \nabla_\tau u_j \right) d\sigma,$$

where each \tilde{u}_j corresponding to u_j is computed thanks to Corollary D.4.

We first state a technical result to perform this summation. We postpone its proof to the end of the section.

Lemma D.5 *Let $V_n = R^l Y_l^m$, $-l \leq m \leq l$ and*

$$\psi = r Y_1^p$$

for $-1 \leq p \leq 1$. Let m' and p' be integer such that $-l \leq m' \leq l$ and $-1 \leq p' \leq 1$ and suppose

$$\tilde{\psi} = a r^{l-1} Y_{l-1}^{m'+p'} + b \frac{r^{l+1}}{R^2} Y_{l+1}^{m'+p'}.$$

Then

$$\begin{aligned} & \int_{\partial B_R} V_n \left(-\partial_n \tilde{\psi} \partial_n \psi - H \lambda \tilde{\psi} \psi + \nabla_\tau \tilde{\psi} \cdot \nabla_\tau \psi \right) d\sigma \\ &= -a (4\alpha + 2l) R^{2l-1} \int_{\partial B_1} Y_{l-1}^{m'+p'} Y_l^m Y_1^p - b (4\alpha + 2) R^{2l-1} \int_{\partial B_1} Y_{l+1}^{m'+p'} Y_l^m Y_1^p. \end{aligned}$$

As a consequence, we get for $j = 1, 2, 3$

$$\begin{aligned} E_{jj}^{(1)} = & -K(R) R^{2l+1} \left[(4\alpha + 2l) \frac{l+2}{l-2} \frac{1+\alpha(3+l)}{1+\alpha(1+l)} \sum_{m=-l}^l \sum_{p=-1}^1 |\alpha_p^j|^2 |v_{l,m}|^2 \left(\int_{\partial B_1} \overline{Y_{l-1}^{m+p}} Y_l^m Y_1^p \right)^2 \right. \\ & \left. + (4\alpha + 2) \frac{l-1}{l} \frac{1+\alpha(4+l)}{1+\alpha(3+l)} \sum_{m=-l}^l \sum_{p=-1}^1 |\alpha_p^j|^2 |v_{l,m}|^2 \left(\int_{\partial B_1} \overline{Y_{l+1}^{m+p}} Y_l^m Y_1^p \right)^2 \right]. \end{aligned}$$

We are now in position to prove Proposition 4.3 concerning the trace of $E^{(1)}$ in dimension $d = 3$.

Proof of Proposition 4.3:

We have to sum the $E_{jj}^{(1)}$ obtained before the statement of Proposition 4.3. By the normalization condition $\sum_j |\alpha_p^j|^2 = 1$, our main task is to compute the sum over $p = -1, 0, 1$ of the integrals involving three spherical harmonics. The values of this type of integral is recalled in Propositions B.2 and B.3. Elementary computations then give

$$\sum_{m=-l}^l \sum_{p=-1}^1 \left(\int_{\partial B_1} \overline{Y_{l-1}^{m+p}} Y_l^m Y_1^p \right)^2 = \frac{3}{4\pi} \frac{l}{2l+1} \text{ and } \sum_{m=-l}^l \sum_{p=-1}^1 \left(\int_{\partial B_1} \overline{Y_{l+1}^{m+p}} Y_l^m Y_1^p \right)^2 = \frac{3}{4\pi} \frac{l+1}{2l+1}.$$

■

Proof of Lemma D.5:

We compute:

$$\begin{aligned} -V_n \partial_n \tilde{\psi} \partial_n \psi &= -R^{2l-1} \left[a(l-1) Y_{l-1}^{m'+p'} + b(l+1) Y_{l+1}^{m'+p'} \right] Y_l^m Y_1^p, \\ -\lambda H V_n \tilde{\psi} \psi &= -R^{2l-1} (4\alpha + 2) \left[a Y_{l-1}^{m'+p'} + b Y_{l+1}^{m'+p'} \right] Y_l^m Y_1^p, \end{aligned}$$

We have also

$$\begin{aligned} \int_{\partial B_R} V_n \nabla_\tau \tilde{\psi} \cdot \nabla_\tau \psi &= \frac{1}{2} \int_{\partial B_R} V_n \left[\Delta_\tau (\tilde{\psi} \psi) - \psi \Delta_\tau \tilde{\psi} - \tilde{\psi} \Delta_\tau \psi \right] \\ &= -\frac{1}{2} l(l+1) R^{2l-1} \int_{\partial B_1} (a Y_{l-1}^{m'+p'} + b Y_{l+1}^{m'+p'}) Y_l^m Y_1^p \\ &\quad + R^{2l+1} \int_{\partial B_1} (a Y_{l-1}^{m'+p'} + b Y_{l+1}^{m'+p'}) Y_l^m Y_1^p \\ &\quad + \frac{1}{2} R^{2l-1} \int_{\partial B_1} \left[a l(l-1) Y_{l-1}^{m'+p'} + b (l+1)(l+2) Y_{l+1}^{m'+p'} \right] Y_l^m Y_1^p \\ &= R^{2l-1} \int_{\partial B_1} \left[a (l-1) Y_{l-1}^{m'+p'} + b (l+2) Y_{l+1}^{m'+p'} \right] Y_l^m Y_1^p. \end{aligned}$$

We obtain the result by summing the three terms. ■

E Shape Derivatives of Steklov and Laplace-Beltrami eigenvalues problem

The following result is obtained by taking $\beta = 0$ in Theorem 1.4.

Theorem E.1 [Steklov eigenvalues] *We distinguish the case of simple and multiple eigenvalue.*

- If $\lambda = \lambda_k(\Omega)$ is a simple eigenvalue of the Steklov problem and u an associated eigenfunction, then the application $t \rightarrow \lambda(t) = \lambda_k((I + t\mathbf{V})(\Omega))$ is differentiable and the derivative at $t = 0$ is

$$\lambda'(0) = \int_{\partial\Omega} V_n \left(|\nabla_\tau u|^2 - |\partial_n u|^2 - \lambda H |u|^2 \right) d\sigma.$$

The shape derivative u' of the eigenfunction satisfies

$$\begin{aligned} \Delta u' &= 0 \text{ in } \Omega, \\ \partial_n u' - \lambda u' &= \operatorname{div}_\tau (V_n \nabla_\tau u) - \lambda'(0) u + \lambda V_n (\partial_n u + H u) \text{ on } \partial\Omega. \end{aligned}$$

- Let λ be a multiple eigenvalue of order $m \geq 2$. Let (u_j) for $1 \leq j \leq m$ denote the eigenfunctions associated to λ . Then there exists m functions $t \mapsto \lambda_k(t)$, $k = 1, \dots, m$ defined in a neighborhood of 0 such that

- $\lambda_k(0) = \lambda$,
- for every t in a neighborhood of 0, $\lambda_k(t)$ is an Steklov eigenvalue of $\Omega_t = (I + t\mathbf{V})(\Omega)$,

- the functions $t \mapsto \lambda_k(t), k = 1, \dots, m$ admit derivatives which are the eigenvalues of the $m \times m$ matrix $M = M_\Omega(V_n)$ of entries (M_{ij}) defined by

$$M_{jk} = \int_{\partial\Omega} V_n \left(-\partial_n u_j \partial_n u_k - H \lambda u_j u_k + \nabla_\tau u_j \cdot \nabla_\tau u_k \right) d\sigma.$$

The following result is obtain by taking $\beta \rightarrow +\infty$ in Theorem 1.4.

Theorem E.2 [Laplace-Beltrami eigenvalues] *We distinguish the case of simple and multiple eigenvalue.*

- If $\lambda = \lambda_k(\Omega)$ is a simple eigenvalue of the Laplace-Beltrami problem and u an associated eigenfunction, then the application $t \rightarrow \lambda(t) = \lambda_k((I + t\mathbf{V})(\Omega))$ is differentiable and the derivative at $t = 0$ is

$$\lambda'(0) = \int_{\partial\Omega} V_n \left((H I_d - 2D^2b) \nabla_\tau u \cdot \nabla_\tau u \right) d\sigma.$$

The shape derivative v' of the eigenfunction satisfies

$$\begin{aligned} \Delta u' &= 0 \text{ in } \Omega, \\ -\Delta_\tau u' &= \Delta_\tau(V_n \partial_n u) - \operatorname{div}_\tau(V_n(2D^2b - H I_d) \nabla_\tau u) - \lambda'(0)u \text{ on } \partial\Omega. \end{aligned}$$

- Let λ be a multiple eigenvalue of order $m \geq 2$. Let (u_j) for $1 \leq j \leq m$ denote the eigenfunctions associated to λ . Then there exists m functions $t \mapsto \lambda_k(t), k = 1, \dots, m$ defined in a neighborhood of 0 such that
 - $\lambda_k(0) = \lambda$,
 - for every t in a neighborhood of 0, $\lambda_k(t)$ is a Laplace-Beltrami eigenvalue of $\Omega_t = (I + t\mathbf{V})(\Omega)$,
 - the functions $t \mapsto \lambda_k(t), k = 1, \dots, m$ admit derivatives which are the eigenvalues of the $m \times m$ matrix $M = M_\Omega(V_n)$ of entries (M_{ij}) defined by

$$M_{jk} = \int_{\partial\Omega} V_n \left((H I_d - 2D^2b) \nabla_\tau u_i \cdot \nabla_\tau u_j \right) d\sigma.$$

Acknowledgements. Part of the work was supported by the project Projet ANR-12-BS01-0007 OPTIFORM financed by the French Agence Nationale de la Recherche (ANR). We would also like to thank the anonymous reviewer for its careful reading of the previous version of the manuscript, that helped to improve and clarify the paper.

References

- [1] A. Bendali and K. Lemrabet. The effect of a thin coating on the scattering of a time-harmonic wave for the Helmholtz equation. *SIAM J. Appl. Math.*, 56(6):1664–1693, 1996.
- [2] M. F. Betta, F. Brock, A. Mercaldo, and M. R. Posteraro. A weighted isoperimetric inequality and applications to symmetrization. *J. Inequal. Appl.*, 4(3):215–240, 1999.
- [3] D. D. Bleecker. The spectrum of a Riemannian manifold with a unit Killing vector field. *Trans. Amer. Math. Soc.*, 275(1):409–416, 1983.

- [4] V. Bonnaillie-Noël, M. Dambrine, F. Hérau, and G. Vial. On generalized Ventcel's type boundary conditions for Laplace operator in a bounded domain. *SIAM J. Math. Anal.*, 42(2):931–945, 2010.
- [5] L. Brasco, G. De Philippis, and B. Ruffini. Spectral optimization for the Stekloff-Laplacian: the stability issue. *J. Funct. Anal.*, 262(11):4675–4710, 2012.
- [6] F. Brock. An isoperimetric inequality for eigenvalues of the Stekloff problem. *ZAMM Z. Angew. Math. Mech.*, 81(1):69–71, 2001.
- [7] F. Caubet, M. Dambrine, and D. Kateb. Shape optimization methods for the inverse obstacle problem with generalized impedance boundary conditions. *Inverse Problems*, 29(11), 2013.
- [8] F. H. Clarke. *Optimization and nonsmooth analysis*. Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons, Inc., New York, 1983. A Wiley-Interscience Publication.
- [9] C. Cohen Tannoudji, B. Diu, and F. Laloe. *Mécanique Quantique*. Hermann, Paris, 1997.
- [10] B. Colbois and J. Dodziuk. Riemannian metrics with large λ_1 . *Proc. Amer. Math. Soc.*, 122(3):905–906, 1994.
- [11] B. Colbois, E. B. Dryden, and A. El Soufi. Bounding the eigenvalues of the Laplace-Beltrami operator on compact submanifolds. *Bull. Lond. Math. Soc.*, 42(1):96–108, 2010.
- [12] M. Dambrine and D. Kateb. Persistency of wellposedness of Ventcel's boundary value problem under shape deformations. *J. Math. Anal. Appl.*, 394(1):129–138, 2012.
- [13] M. C. Delfour and J.-P. Zolésio. *Shapes and geometries*, volume 4 of *Advances in Design and Control*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001. Analysis, differential calculus, and optimization.
- [14] F. R. Desaint and J.-P. Zolésio. Manifold derivative in the Laplace-Beltrami equation. *J. Funct. Anal.*, 151(1):234–269, 1997.
- [15] G. R. Goldstein. Derivation and physical interpretation of general boundary conditions. *Adv. Differential Equations*, 11(4):457–480, 2006.
- [16] H. Haddar, P. Joly, and H.-M. Nguyen. Generalized impedance boundary conditions for scattering by strongly absorbing obstacles: the scalar case. *Math. Models Methods Appl. Sci.*, 15(8):1273–1300, 2005.
- [17] A. Henrot and M. Pierre. *Variation et optimisation de formes*, volume 48 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer, Berlin, 2005. Une analyse géométrique. [A geometric analysis].
- [18] J. Hersch. Caractérisation variationnelle d'une somme de valeurs propres consécutives; généralisation d'inégalités de Pólya-Schiffer et de Weyl. *C. R. Acad. Sci. Paris*, 252:1714–1716, 1961.
- [19] J. Hersch. Quatre propriétés isopérimétriques de membranes sphériques homogènes. *C. R. Acad. Sci. Paris Sér. A-B*, 270:A1645–A1648, 1970.

- [20] G. N. Hile and Z. Y. Xu. Inequalities for sums of reciprocals of eigenvalues. *J. Math. Anal. Appl.*, 180(2):412–430, 1993.
- [21] J. Kennedy. A Faber-Krahn inequality for the Laplacian with generalised Wentzell boundary conditions. *J. Evol. Equ.*, 8(3):557–582, 2008.
- [22] K. Lemrabet and D. Teniou. Vibrations d’une plaque mince avec raidisseur sur le bord. *Maghreb Math. Rev.*, 2(1):27–41, 1992.
- [23] J.-C. Nédélec. *Acoustic and electromagnetic equations*, volume 144 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2001. Integral representations for harmonic problems.
- [24] J. H. Ortega and E. Zuazua. Generic simplicity of the eigenvalues of the Stokes system in two space dimensions. *Adv. Differential Equations*, 6(8):987–1023, 2001.
- [25] E. M. Stein and G. Weiss. *Introduction to Fourier analysis on Euclidean spaces*. Princeton University Press, Princeton, N.J., 1971. Princeton Mathematical Series, No. 32.

Marc Dambrine

Université de Pau et des Pays de l’Adour

E-mail: marc.dambrine@univ-pau.fr

http://web.univ-pau.fr/~mdambrin/Marc_Dambrine/Home.html

Djalil Kateb

Université de Technologie de Compiègne.

E-mail: djalil.kateb@utc.fr

<http://www.lmac.utc.fr/membres/kateb>

Jimmy Lamboley

Université Paris-Dauphine

E-mail: lamboley@math.cnrs.fr

<https://www.ceremade.dauphine.fr/~lamboley/>