

LOGARITHMIC SOBOLEV INEQUALITY REVISITED

HOAI-MINH NGUYEN AND MARCO SQUASSINA

ABSTRACT. We provide a new characterization of the logarithmic Sobolev inequality.

1. INTRODUCTION

The classical Sobolev inequality translates information about the derivatives of a function into information about the size of the function itself. Precisely, for a function u with square summable gradient in dimension N one obtains that u is $2N/(N-2)$ -summable, that is a gain in summability which depends on N and which tends to deteriorate as $N \rightarrow \infty$. On the other hand, since the middle fifties, people have started looking at possible replacements of the Sobolev inequality in order to provide an improvement in the summability *independent* of the dimension N , which can be done in terms of integrability properties of $u^2 \log u^2$. This was firstly done by Stam [23] who proved the logarithmic Sobolev inequality with Gauss measure $d\mathcal{G}$

$$\int_{\mathbb{R}^N} u^2 \log \frac{u^2}{\|u\|_{2,d\mathcal{G}}^2} d\mathcal{G} \leq \frac{1}{\pi} \int_{\mathbb{R}^N} |\nabla u|^2 d\mathcal{G}, \quad d\mathcal{G} = e^{-\pi|x|^2} dx.$$

The formula was originally discovered in quantum field theory in order to handle estimates which are uniform in the space dimension, for systems with a large number of variables. A different proof and further insight was obtained by Gross in [17]. See also the work of Adams and Clarke [1] for an elementary proof of the previous inequality. These properties are widely used in statistical mechanics, quantum field theory and differential geometry. A variant of the logarithmic Sobolev inequality with Gauss measure is given by the following one parameter family of euclidean inequalities [18, Theorem 8.14]

$$\int_{\mathbb{R}^N} u^2 \log \frac{u^2}{\|u\|_2^2} dx + N(1 + \log a)\|u\|_2^2 \leq \frac{a^2}{\pi} \int_{\mathbb{R}^N} |\nabla u|^2 dx.$$

for any $u \in H^1(\mathbb{R}^N)$ and $a > 0$. A version of this inequality for fractional Sobolev spaces $H^s(\mathbb{R}^N)$ can be found in [13]. Recently some new characterization of the Sobolev spaces were provided in [4, 19, 21] (see also [2, 3, 5–9, 20]) in terms of the following family of nonlocal functionals

$$I_\delta(u) := \int \int_{\{|u(y)-u(x)|>\delta\}} \frac{\delta^2}{|x-y|^{N+2}} dx dy, \quad \delta > 0,$$

where u is a measurable function on \mathbb{R}^N . In particular, if $N \geq 3$ and $I_\delta(u) < \infty$ for some $\delta > 0$, then in [21] it was proved that

$$(1.1) \quad \int_{\{|u|>\lambda_N\delta\}} |u|^{2N/(N-2)} dx \leq C_N I_\delta(u)^{N/(N-2)},$$

2010 *Mathematics Subject Classification.* 46E35, 28D20, 82B10, 49A50.

Key words and phrases. Nonlocal functionals, logarithmic Sobolev inequality, entropy.

The second author is member of *Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* (GNAMPA) of the *Istituto Nazionale di Alta Matematica* (INdAM).

for some positive constants C_N and λ_N . This is a sort of nonlocal improvement of the classical Sobolev inequality and it is also possible to show that in the singular limit $\delta \searrow 0$ one recovers the classical Sobolev result, since I_δ converges to the Dirichlet energy up to a normalization constant. The aim of this note is to remark that in this context also a logarithmic type estimate holds. Thus we have the summability gain independent of N can be controlled in terms of $I_\delta(u)$.

More precisely, we have the following

Theorem 1.1. *Let $u \in L^2(\mathbb{R}^N)$ ($N \geq 3$). There is a positive constant C_N such that*

$$\int_{\mathbb{R}^N} \frac{u^2}{\|u\|_2^2} \log \frac{u^2}{\|u\|_2^2} dx + \frac{N}{2} \log \|u\|_2^2 \leq \frac{N}{2} \log \left(C_N \delta^{\frac{4}{N}} \|u\|_2^{\frac{2N-4}{N}} + C_N I_\delta(u) \right),$$

for all $\delta > 0$. In particular, if $u \in L^2(\mathbb{R}^N)$ is such that $I_\delta(u) < \infty$ for some $\delta > 0$, then

$$(1.2) \quad \int_{\mathbb{R}^N} u^2 \log u^2 dx < +\infty.$$

Proof. By a simple normalization argument, we may reduce the assertion to proving that

$$(1.3) \quad \int_{\mathbb{R}^N} u^2 \log u^2 dx \leq \frac{N}{2} \log \left(C_N \delta^{\frac{4}{N}} + C_N I_\delta(u) \right), \quad \text{for all } \delta > 0,$$

for any $u \in L^2(\mathbb{R}^N)$ such that $\|u\|_2 = 1$. Considering the normalized outer measure

$$\mu(E) := \int_E u^2(x) dx, \quad \mu(\mathbb{R}^N) = 1,$$

and using Jensen's inequality for concave nonlinearities and with measure μ , we have

$$(1.4) \quad \begin{aligned} \log \left(\int_{\mathbb{R}^N} |u|^{\frac{2N}{N-2}} dx \right) &= \log \left(\int_{\mathbb{R}^N} |u|^{\frac{4}{N-2}} d\mu \right) \geq \int_{\mathbb{R}^N} \log |u|^{\frac{4}{N-2}} d\mu \\ &= \frac{2}{N-2} \int_{\mathbb{R}^N} u^2 \log u^2 dx. \end{aligned}$$

On the other hand, applying (1.1), we derive that, for all $\delta > 0$,

$$\frac{2}{N-2} \int_{\mathbb{R}^N} u^2 \log u^2 dx \leq \log \left(D_N \delta^{\frac{4}{N-2}} + C_N I_\delta(u)^{\frac{N}{N-2}} \right),$$

for some positive constant D_N , which implies (1.3). Here we used the fact that

$$\int_{\{|u| \leq \lambda_N \delta\}} |u|^{\frac{2N}{N-2}} dx \leq \lambda_N^{\frac{4}{N-2}} \delta^{\frac{4}{N-2}},$$

since $\int_{\mathbb{R}^N} u^2 dx = 1$. □

Defining a notion of *entropy* as typical in statistical mechanics:

$$\text{Ent}_\mu(f) := \int_{\mathbb{R}^N} \frac{f}{\|f\|_{1,\mu}} \log \frac{f}{\|f\|_{1,\mu}} d\mu + \frac{N}{2} \log \|f\|_{1,\mu}, \quad f \geq 0, \quad \|f\|_{1,\mu} := \int f d\mu,$$

the conclusion of the previous results reads as

$$u \in L^2(\mathbb{R}^N), \exists \delta > 0 : I_\delta(u) < +\infty \implies \text{Ent}_{\mathcal{L}^N}(u^2) < +\infty.$$

Remark 1.2 (Logarithmic NLS). If $u \in H^1(\mathbb{R}^N)$, then the results of [19] show that

$$(1.5) \quad \lim_{\delta \searrow 0} I_\delta(u) = Q_N \int_{\mathbb{R}^N} |\nabla u|^2 dx,$$

for some constant $Q_N > 0$. Hence, passing to the limit as $\delta \searrow 0$ in the inequality of Theorem 1.1 one recovers classical forms of the logarithmic inequality. The logarithmic Schrödinger equation

$$(1.6) \quad i\partial_t \phi + \Delta \phi + \phi \log |\phi|^2 = 0, \quad \phi : [0, \infty) \times \mathbb{R}^N \rightarrow \mathbb{C}, \quad N \geq 3,$$

admits applications to quantum mechanics, quantum optics, transport and diffusion phenomena, theory of superfluidity and Bose-Einstein condensation (see [25] and [10–12]). The *standing waves* solutions of (1.6) solve the following semi-linear elliptic problem

$$(1.7) \quad -\Delta u + \omega u = u \log u^2, \quad u \in H^1(\mathbb{R}^N).$$

These equations were recently investigated in [14, 24]. From a variational point of view, the search of solutions to (1.7) can be associated with the study of critical points (in a nonsmooth sense) of the lower semi-continuous functional $J : H^1(\mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$J(u) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx + \frac{\omega + 1}{2} \int_{\mathbb{R}^N} u^2 dx - \frac{1}{2} \int_{\mathbb{R}^N} u^2 \log u^2 dx,$$

which is well defined by the logarithmic Sobolev inequality. Due to Theorem 1.1 and (1.5), one could handle a kind of *nonlocal approximations* of (1.7), formally defined for $\delta > 0$ by

$$I'_\delta(u) + \omega u = u \log u^2,$$

which are associated with the energy functional $J_\delta : H^1(\mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$J_\delta(u) = I_\delta(u) + \frac{\omega + 1}{2} \int_{\mathbb{R}^N} u^2 dx - \frac{1}{2} \int_{\mathbb{R}^N} u^2 \log u^2 dx.$$

Since there holds $I_\delta(u) \leq C_N \int_{\mathbb{R}^N} |\nabla u|^2 dx$ for all $\delta > 0$ and $u \in H^1(\mathbb{R}^N)$ (cf. [19, Theorem 2]) the energy functional J_δ is well defined, for every $\delta > 0$.

Remark 1.3 (Magnetic case). If $A : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is locally bounded and $u : \mathbb{R}^N \rightarrow \mathbb{C}$, we set

$$\Psi_u(x, y) := e^{i(x-y) \cdot A\left(\frac{x+y}{2}\right)} u(y), \quad x, y \in \mathbb{R}^N.$$

It was observed in [15] that the following *Diamagnetic inequality* holds

$$\left| |u(x)| - |u(y)| \right| \leq \left| \Psi_u(x, x) - \Psi_u(x, y) \right|, \quad \text{for a.e. } x, y \in \mathbb{R}^N.$$

In turn, by defining

$$I_\delta^A(u) := \int_{\{|\Psi_u(x, y) - \Psi_u(x, x)| > \delta\}} \frac{\delta^2}{|x - y|^{N+2}} dx dy,$$

we have

$$(1.8) \quad I_\delta(|u|) \leq I_\delta^A(u), \quad \text{for all } \delta > 0 \text{ and all measurable } u : \mathbb{R}^N \rightarrow \mathbb{C}.$$

Then, Theorem 1.1 yields the following *Magnetic logarithmic Sobolev inequality*. For $u \in L^2(\mathbb{R}^N)$, there is a positive constant C_N such that

$$\int_{\mathbb{R}^N} \frac{|u|^2}{\|u\|_2^2} \log \frac{|u|^2}{\|u\|_2^2} dx + \frac{N}{2} \log \|u\|_2^2 \leq \frac{N}{2} \log \left(C_N \delta^{\frac{4}{N}} \|u\|_2^{\frac{2N-4}{N}} + C_N I_\delta^A(u) \right).$$

Notice that, since $I_\delta(|u|) \approx \|\nabla|u|\|_2^2$ as $\delta \searrow 0$ [19] and $I_\delta^A(u) \approx \|\nabla u - iAu\|_2^2$ as $\delta \searrow 0$ [22], from inequality (1.8) one recovers $\|\nabla|u|\|_2 \leq \|\nabla u - iAu\|_2$ which follows from the well-know diamagnetic inequality for the gradients $|\nabla|u|| \leq |\nabla u - iAu|$, see [18].

As a companion to Theorem 1.1, we also have the following

Theorem 1.4. *Let $u \in L^2(\mathbb{R}^N)$ ($N \geq 3$). Assume that there exists a non-decreasing function $F : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $F(ts) \leq t^\beta F(s)$ for any $s, t \geq 0$ and some $\beta > 0$ and*

$$(1.9) \quad \int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+2}} dx dy < +\infty.$$

Then there exists a positive constant $C_{N,F}$ such that

$$\int_{\mathbb{R}^N} \frac{u^2}{\|u\|_2^2} \log \frac{u^2}{\|u\|_2^2} dx + \frac{N}{2} \log \|u\|_2^\beta \leq \frac{N}{2} \log \left(C_{N,F} \|u\|_2^\beta + C_{N,F} \int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+2}} dx dy \right),$$

In particular, condition (1.2) holds.

Proof. Consider the statement when $\|u\|_2 = 1$. In light of inequality (1.4), since by [21, Proposition 6] there exists $C_N > 0$ and $\lambda_N > 0$ such that

$$(1.10) \quad \int_{\{|u| > \lambda_N F(1/2)\}} |u|^{2N/(N-2)} dx \leq C_N \left(\frac{1}{F(1/2)} \int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+2}} dx dy \right)^{N/(N-2)},$$

by arguing as in the previous proof, we get

$$\frac{2}{N-2} \int_{\mathbb{R}^N} u^2 \log u^2 \leq \log \left(D_{N,F} + D_{N,F} \left(\int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+2}} dx dy \right)^{N/(N-2)} \right),$$

where we used the fact that

$$\int_{\{|u| \leq \lambda_N F(1/2)\}} |u|^{\frac{2N}{N-2}} dx \leq \lambda_N^{\frac{4}{N-2}} F(1/2)^{\frac{4}{N-2}},$$

since $\int_{\mathbb{R}^N} u^2 dx = 1$. Then, we get

$$\int_{\mathbb{R}^N} u^2 \log u^2 \leq \frac{N}{2} \log \left(C_{N,F} + C_{N,F} \int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+2}} dx dy \right).$$

In the general case, using the sub-homogeneity condition on F yields

$$\int_{\mathbb{R}^N} \frac{u^2}{\|u\|_2^2} \log \frac{u^2}{\|u\|_2^2} \leq \frac{N}{2} \log \left(C_{N,F} + \frac{C_{N,F}}{\|u\|_2^\beta} \int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+2}} dx dy \right),$$

which yields the desired conclusion. \square

Remark 1.5 ($L^p(\mathbb{R}^N)$ -version). If $p > 1$ and $N > p$, one has a variant of (1.4), namely

$$(1.11) \quad \log \left(\int_{\mathbb{R}^N} |u|^{\frac{Np}{N-p}} dx \right) \geq \frac{p}{N-p} \int_{\mathbb{R}^N} |u|^p \log |u|^p dx.$$

Then, by arguing as in the proofs of Theorems 1.1 and 1.4 with

$$(1.12) \quad u \mapsto \int \int_{\{|u(y) - u(x)| > \delta\}} \frac{\delta^p}{|x - y|^{N+p}} dx dy, \quad u \mapsto \int_{\mathbb{R}^{2N}} \frac{F(|u(x) - u(y)|)}{|x - y|^{N+p}} dx dy,$$

in place of $I_\delta(u)$ and (1.9) respectively, it is possible to get corresponding log-Sobolev inequalities as for the case $p = 2$, via the results of [21]. In particular, if $u \in L^p(\mathbb{R}^N)$ and the functionals in (1.12) are finite at u for some $\delta > 0$, then

$$\int_{\mathbb{R}^N} |u|^p \log |u|^p dx < +\infty.$$

The Euclidean logarithmic Sobolev inequalities for the p -case have been intensively studied, see e.g. the work of Del Pino and Dolbeault [16] and the references therein.

REFERENCES

- [1] R.A. Adams, F.H. Clarke, *Gross's logarithmic Sobolev inequality: a simple proof*, Amer. J. Math. **101** (1979), 1265–1269. [1](#)
- [2] J. Bourgain, H. Brezis, P. Mironescu, *Another look at Sobolev spaces*, in *Optimal Control and Partial Differential Equations. A Volume in Honor of Professor Alain Bensoussan's 60th Birthday* (eds. J. L. Menaldi, E. Rofman and A. Sulem), IOS Press, Amsterdam, 2001, 439–455. [1](#)
- [3] J. Bourgain, H. Brezis, P. Mironescu, *Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications*, J. Anal. Math. **87** (2002), 77–101. [1](#)
- [4] J. Bourgain and H-M. Nguyen, *A new characterization of Sobolev spaces*, C. R. Acad. Sci. Paris **343** (2006), 75–80. [1](#)
- [5] H. Brezis, *How to recognize constant functions. Connections with Sobolev spaces*, Russian Mathematical Surveys **57** (2002), 693–708. [1](#)
- [6] H. Brezis, *New approximations of the total variation and filters in imaging*, Rend Accad. Lincei **26** (2015), 223–240. [1](#)
- [7] H. Brezis, H-M. Nguyen, *Non-local functionals related to the total variation and connections with Image Processing*, preprint. <http://arxiv.org/abs/1608.08204> [1](#)
- [8] H. Brezis, H-M. Nguyen, *The BBM formula revisited*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **27** (2016), 515–533. [1](#)
- [9] H. Brezis, H-M. Nguyen, *Two subtle convex nonlocal approximations of the BV-norm*, Nonlinear Anal. **137** (2016), 222–245. [1](#)
- [10] T. Cazenave, *Stable solutions of the logarithmic Schrödinger equation*, Nonlinear Anal. **7** (1983), 1127–1140. [3](#)
- [11] T. Cazenave, *An introduction to nonlinear Schrödinger equations*, Textos de Métodos Matemáticos **26**, Universidade Federal do Rio de Janeiro 1996. [3](#)
- [12] T. Cazenave, A. Haraux, *Équations d'évolution avec non linéarité logarithmique*, Ann. Fac. Sci. Toulouse Math. **2** (1980), 21–51. [3](#)
- [13] A. Cotsoioli, N. Tavoularis, *On logarithmic Sobolev inequalities for higher order fractional derivatives*, Comptes Rendus Acad. Sci. Paris **340** (2005), 205–208. [1](#)
- [14] P. d'Avenia, E. Montefusco, M. Squassina, *On the logarithmic Schrödinger equation*, Commun. Contemp. Math. **16** (2014), 1350032, 15 pp. [3](#)
- [15] P. d'Avenia, M. Squassina, *Ground states for fractional magnetic operators*, ESAIM COCV, to appear. [3](#)
- [16] M. Del Pino, J. Dolbeault, *The optimal euclidean L^p -Sobolev logarithmic inequality*, J. Funct. Anal. **197** (2003), 151–161. [4](#)
- [17] L. Gross, *Logarithmic Sobolev Inequalities*, Amer. J. Math. **97** (1975), 1061–1083. [1](#)
- [18] E. Lieb and M. Loss, *Analysis*, Graduate Studies in Mathematics **14**, AMS, 2001. [1](#), [3](#)
- [19] H-M. Nguyen, *Some new characterizations of Sobolev spaces*, J. Funct. Anal. **237** (2006), 689–720. [1](#), [2](#), [3](#)
- [20] H-M. Nguyen, *Further characterizations of Sobolev spaces*, J. Eur. Math. Soc. **10** (2008), 191–229. [1](#)
- [21] H-M. Nguyen, *Some inequalities related to Sobolev norms*, Calc. Var. PDEs **41** (2011), 483–509. [1](#), [4](#)
- [22] H-M. Nguyen, A. Pinamonti, M. Squassina, E. Vecchi, *A new characterization of magnetic Sobolev spaces*, in preparation. [3](#)
- [23] A.J. Stam, *Some inequalities satisfied by the quantities of information of Fisher and Shannon*, Information and Control **2** (1959), 101–112. [1](#)
- [24] W.C. Troy, *Uniqueness of positive ground state solutions of the logarithmic Schrödinger equation*, Arch. Ration. Mech. Anal. **222** (2016), 1581–1600. [3](#)
- [25] K.G. Zloshchastiev, *Logarithmic nonlinearity in theories of quantum gravity: origin of time and observational consequences*, Grav. Cosmol. **16** (2010), 288–297. [3](#)

(H.-M. Nguyen) DEPARTMENT OF MATHEMATICS
 EPFL SB CAMA
 STATION 8 CH-1015 LAUSANNE, SWITZERLAND
E-mail address: hoai-minh.nguyen@epfl.ch

(M. Squassina) DIPARTIMENTO DI MATEMATICA E FISICA
 UNIVERSITÀ CATTOLICA DEL SACRO CUORE
 VIA DEI MUSEI 41, I-25121 BRESCIA, ITALY
E-mail address: marco.squassina@unicatt.it