

OPTIMAL TRANSPORT WITH COULOMB COST AND THE SEMICLASSICAL LIMIT OF DENSITY FUNCTIONAL THEORY

UGO BINDINI AND LUIGI DE PASCALE

ABSTRACT. We present some progress in the direction of determining the semiclassical limit of the Levy-Lieb or Hohenberg-Kohn universal functional in Density Functional Theory for Coulomb systems. In particular we give a proof of the fact that for Bosonic systems with an arbitrary number of particles the limit is the multimarginal optimal transport problem with Coulomb cost and that the same holds for Fermionic systems with 2 or 3 particles. Comparisons with previous results are reported. The approach is based on some techniques from the optimal transportation theory.

1. INTRODUCTION AND PRELIMINARY RESULTS

The ground state of a system of N electrons interacting through Coulomb forces between each others and with M nuclei is described by one of the following minimal values (ground values)

$$E_h^S = \min_S T_h(\psi) + V_{ee}(\psi) - V_{ne}(\psi), \quad (1.1)$$

or

$$E_h^A = \min_A T_h(\psi) + V_{ee}(\psi) - V_{ne}(\psi), \quad (1.2)$$

and by the corresponding minimizers (ground states).

The minimization domains are the following sets of wave functions

$$\mathcal{S} = \{\psi \in H^1((\mathbb{R}^d \times \mathbb{Z}_2)^N) : \int |\psi|^2 dz = 1, \text{ for all permutations } \sigma \text{ of } N \text{ points} \\ \psi((x_{\sigma(1)}, \alpha_{\sigma(1)}), \dots, (x_{\sigma(N)}, \alpha_{\sigma(N)})) = \psi((x_1, \alpha_1), \dots, (x_N, \alpha_N))\},$$

$$\mathcal{A} = \{\psi \in H^1((\mathbb{R}^d \times \mathbb{Z}_2)^N) : \int |\psi|^2 dz = 1, \text{ for all permutations } \sigma \text{ of } N \text{ points} \\ \psi((x_{\sigma(1)}, \alpha_{\sigma(1)}), \dots, (x_{\sigma(N)}, \alpha_{\sigma(N)})) = \text{sign}(\sigma)\psi((x_1, \alpha_1), \dots, (x_N, \alpha_N))\},$$

where we adopted the common notation

$$\int f(z) dz := \int f(z_1, \dots, z_N) dz_1 \dots dz_N := \sum_{\alpha_1, \dots, \alpha_N=0,1} \int_{\mathbb{R}^d} f(x_1, \alpha_1; \dots; x_N, \alpha_N) dx_1 \dots dx_N.$$

The three terms in the functionals are: the kinetic energy

$$T_h(\psi) = \frac{\hbar^2}{2m} \int |\nabla \psi|^2(z) dz,$$

the electron-electron interaction energy

$$V_{ee}(\psi) = \int \sum_{1 \leq i < j \leq N} \frac{|\psi|^2(z)}{|x_i - x_j|} dz$$

Date: July 25, 2017.

2010 Mathematics Subject Classification. 49J45, 49N15, 49K30.

Key words and phrases. Density Functional Theory, Multimarginal optimal transportation, Monge-Kantorovich problem, duality theory, Coulomb cost.

and the nuclei-electrons interaction energy

$$V_{ne}(\psi) = \int \sum_{1 \leq i < j \leq N} \sum_{k=1}^M \frac{|\psi|^2(z) Z_k}{|x_i - N_k|} dz.$$

The values E_h^S and E_h^A represents, respectively, the ground value for a Bosonic and Fermionic system of particles with N electrons and M nuclei. The N_k are the positions of the nuclei, while the couple (x_i, α_i) is the position-spin of the i -th electron. In the usual Born interpretation, $|\psi|^2$ is the probability distribution of the N electrons and, according with the indistinguishability principle, it is invariant with respect to permutations of the N (x_i, α_i) variables.

Computing the ground values above amounts to solving a Schrödinger equation in \mathbb{R}^{dN} and the numerical cost scales exponentially with N . The Density Functional Theory (DFT from now on) is an alternative introduced in the late sixties by Hohenberg, Kohn and Sham. However the desire to describe the system in term of a different variable is much older and we may consider the Thomas-Fermi model as a precursor of this theory.

We associate to every wave function ψ a probability density on \mathbb{R}^d defined as follows ¹:

$$\rho_\psi(x) := \sum_{\alpha_1, \dots, \alpha_N=0,1} \int_{\mathbb{R}^{d(N-1)}} |\psi|^2(x, \alpha_1; \dots; x_N, \alpha_N) dx_2 \dots dx_N.$$

The map which associates ρ to ψ will be denoted by $\psi \downarrow \rho_\psi$ and ρ will be called single electron density or electronic density. If we wish to describe the system in terms of ρ , the problems above should be reformulated as a minimization of suitable energies with respect to ρ which, no matter how big N is, is always a probability measure in \mathbb{R}^d . The exact image of the map $\psi \downarrow \rho_\psi$ was characterized by Lieb [26] who proved that the image of both \mathcal{S} and \mathcal{A} is

$$H = \{\rho \mid 0 \leq \rho, \int \rho = 1, \sqrt{\rho} \in H^1(\mathbb{R}^d)\}.$$

For $\rho \in H$ we introduce the Levy-Lieb functional [25, 26] also known as Hohenberg-Kohn functional²

$$F_h^S(\rho) = \inf_{\psi \in \mathcal{S}, \psi \downarrow \rho} \{T_h(\psi) + V_{ee}(\psi)\}, \quad (1.3)$$

$$F_h^A(\rho) = \inf_{\psi \in \mathcal{A}, \psi \downarrow \rho} \{T_h(\psi) + V_{ee}(\psi)\}. \quad (1.4)$$

Then the problems above can be reformulated as follows:

$$E_h^S = \min_{\rho \in H} F_h^S(\rho) + N \int v(x) \rho(x) dx,$$

and

$$E_h^A = \min_{\rho \in H} F_h^A(\rho) + N \int v(x) \rho(x) dx,$$

where we denoted

$$v(x) = - \sum_{k=1}^M \frac{Z_k}{|x - N_k|}.$$

This paper deals with the semiclassical limit of the two functionals $F_h^S(\rho)$ and $F_h^A(\rho)$ and more precisely it concerns the relations of the multimarginal optimal transport theory with these semiclassical limits. The ties between the DFT for Coulomb systems and Optimal transport appeared first in [5, 11] and by now have revealed to be a precious tool for the

¹In the usual definition the integral in the definition of ρ_ψ is also multiplied by a factor N , but here we prefer to deal with probability measures.

²These functionals are sometimes denoted by $F_{HK}(\psi)$. For the sake of a lighter notation, here we will not report the HK .

understanding. To detail better the results, let us shortly introduce the multimarginal optimal transport problem of interest here.

For every $\rho \in H$ consider the set

$$\Pi(\rho) := \{P \in \mathcal{P}(\mathbb{R}^{dN}) \mid \pi_{\#}^i P = \rho, i = 1, \dots, N\},$$

and then the multimarginal optimal transport problem with Coulomb cost

$$C(\rho) := \min_{P \in \Pi(\rho)} \int \sum_{1 \leq i < j \leq N} \frac{1}{|x_i - x_j|} dP. \quad (1.5)$$

Duality for (1.5) and some properties of the functional $C(\rho)$ have been studied in [14, 6]. The structure of 1-dimensional minimizers has been investigated in [9].

The results on optimal transportation needed in this paper will be reported in the next section.

In this paper we will prove that

Theorem 1.1. *For all $\rho \in H$ and $d, N \in \mathbb{N}$,*

$$F_h^S(\rho) \xrightarrow{h \rightarrow 0} C(\rho),$$

and that

Theorem 1.2. *For all $\rho \in H$, $d = 1, 2, 3, 4$, $N = 2, 3$*

$$F_h^A(\rho) \xrightarrow{h \rightarrow 0} C(\rho).$$

The same problem was previously studied in [11]. It is easy to see that the proof presented in that paper adapts to prove Theorem 1.1 for any N . In this case our contribution consists in a more direct use of optimal transport techniques with the consequent simplifications. The convergence 1.2 was proved in [11] for $N = 2$. Here we extend the result to $N = 3$ and we are able to enlarge the class of approximating antisymmetric wave functions also for $N = 2$.

Remark 1.3. Although the usual description is limited to the physical dimension $d = 3$, here we explored also other dimensions in the hope to shed some light on the problems which are still open.

Since the functionals appearing in Theorems 1.1 and 1.2 above are all expressed as minimal values, the natural tool to deal with their convergence is the Γ -convergence which also we shortly introduce in the next section.

2. TOOLS: Γ -CONVERGENCE AND MULTIMARGINAL OPTIMAL TRANSPORT

2.1. Definition of Γ -convergence and basic results. A crucial tool that we will use throughout this paper is Γ -convergence. All the details can be found, for instance, in Braides's book [4] or in the classical book by Dal Maso [13]. In what follows, (X, d) is a metric space or a topological space equipped with a convergence.

Definition 2.1. Let $(F_n)_n$ be a sequence of functions $X \mapsto \bar{\mathbb{R}}$. We say that $(F_n)_n$ Γ -converges to F and we write $F_n \xrightarrow[n]{\Gamma} F$ if for any $x \in X$ we have

- for any sequence $(x_n)_n$ of X converging to x

$$\liminf_n F_n(x_n) \geq F(x) \quad (\Gamma\text{-liminf inequality});$$

- there exists a sequence $(x_n)_n$ converging to x and such that

$$\limsup_n F_n(x_n) \leq F(x) \quad (\Gamma\text{-limsup inequality}).$$

This definition is actually equivalent to the following equalities for any $x \in X$:

$$F(x) = \inf \left\{ \liminf_n F_n(x_n) : x_n \rightarrow x \right\} = \inf \left\{ \limsup_n F_n(x_n) : x_n \rightarrow x \right\}$$

The function $x \mapsto \inf \left\{ \liminf_n F_n(x_n) : x_n \rightarrow x \right\}$ is called Γ -liminf of the sequence $(F_n)_n$ and the other one its Γ -limsup. A useful result is the following (which for instance implies that a constant sequence of functions does not Γ -converge to itself in general).

Proposition 2.2. *The Γ -liminf and the Γ -limsup of a sequence of functions $(F_n)_n$ are both lower semi-continuous on X .*

The main interest of Γ -convergence resides in its consequences in terms of convergence of minima:

Theorem 2.3. *Let $(F_n)_n$ be a sequence of functions $X \rightarrow \bar{\mathbb{R}}$ and assume that $F_n \xrightarrow[n]{\Gamma} F$. Assume moreover that there exists a compact and non-empty subset K of X such that*

$$\forall n \in N, \inf_X F_n = \inf_K F_n$$

(we say that $(F_n)_n$ is equi-mildly coercive on X). Then F admits a minimum on X and the sequence $(\inf_X F_n)_n$ converges to $\min F$. Moreover, if $(x_n)_n$ is a sequence of X such that

$$\lim_n F_n(x_n) = \lim_n (\inf_X F_n)$$

and if $(x_{\phi(n)})_n$ is a subsequence of $(x_n)_n$ having a limit x , then $F(x) = \inf_X F$.

2.2. Multimarginal optimal transportation and composition of optimal transport plans. In this subsection we present some basic results about multimarginal optimal transportation with Coulomb cost.

An element P of $\Pi(\rho)$ is commonly called a transport plan for ρ .

If $\rho \in \mathcal{P}(\mathbb{R}^d)$, define

$$\mu_\rho(t) = \sup_{x \in \mathbb{R}^d} \rho(B(x, t)).$$

Definition 2.4. The *concentration* of ρ is

$$\mu_\rho = \lim_{t \rightarrow 0} \mu_\rho(t).$$

Note that $\rho \in H$ implies $\mu_\rho = 0$, since $\sqrt{\rho} \in H^1$ implies $\rho \in L^1$.

It is commonly assumed that, if $\mu_\rho < 1/N$, then $\mathcal{C}(\rho) < +\infty$. Here we provide a simple argument adapted to the particular case $\rho \in H$:

Proposition 2.5. *If $p \geq \frac{2d}{2d-1}$ and $\rho \in L^1(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$, then $\mathcal{C}(\rho) < +\infty$.*

Proof. Consider the transport plan $P(x_1, \dots, x_N) = \rho(x_1) \cdots \rho(x_N)$; then,

$$\begin{aligned} C(P) &= \int c(X) dP(X) = \sum_{1 \leq i < j \leq N} \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{\rho(x_i) \rho(x_j)}{|x_i - x_j|} dx_i dx_j \\ &\leq C(d, p) \|\rho\|_p^2, \end{aligned}$$

where the last inequality follows from the Hardy-Littlewood-Sobolev and Hölder inequalities. \square

Definition 2.6. If $\alpha > 0$, let

$$D_\alpha = \left\{ X \in \mathbb{R}^{Nd} \text{ s.t. } \exists i \neq j \text{ with } |x_i - x_j| < \alpha \right\}$$

be an open strip around

$$D_0 = \left\{ X \in \mathbb{R}^{Nd} \text{ s.t. } \exists i \neq j \text{ with } x_i = x_j \right\},$$

which is the set where the cost function c is singular.

Note that if a transport plan P has the property that $P(D_\alpha) = 0$ for some $\alpha > 0$ then $C(P)$ is finite. The converse is not true in general, i.e. there may well be transport plans of finite cost whose supports have distance 0 from the set D_0 .

However, this is not the case if P is optimal, as recently proved in [6]:

Theorem 2.7. *Let $\rho \in \mathcal{P}(\mathbb{R}^d)$ with $\mu_\rho < \frac{1}{N(N-1)^2}$, with $C(\rho) < +\infty$, and let β be such that*

$$\mu_\rho(\beta) < \frac{1}{N(N-1)^2}.$$

If $P \in \Pi(\rho)$ is an optimal plan for the problem (1.5), then $P|_{D_\alpha} = 0$ for every α such that

$$\alpha < \frac{2\beta}{N^2(N-1)}.$$

3. A Γ -CONVERGENCE RESULT AND PROOFS OF THE MAIN THEOREMS

It is useful to reformulate the different variational problems so that they have a common domain. For every $\psi \in \mathcal{S}$ or $\psi \in \mathcal{A}$ there is a natural transport plan $P_\psi \in \rho_\psi$ defined by

$$P_\psi = \sum_{\alpha_1, \dots, \alpha_N} |\psi|^2(x_1, \alpha_1; \dots; x_N, \alpha_N).$$

Then for every $\rho \in H$ define $\mathcal{F}_h^{\mathcal{S}}, \mathcal{F}_h^{\mathcal{A}} : \Pi(\rho) \rightarrow \mathbb{R}^+ \cup \{+\infty\}$ as follows:

$$\mathcal{F}_h^{\mathcal{S}}(P) = \begin{cases} T_h(\psi) + V_{ee}(\psi) & \text{if } P = P_\psi \text{ for some } \psi \in \mathcal{S} \\ +\infty & \text{otherwise,} \end{cases}$$

and

$$\mathcal{F}_h^{\mathcal{A}}(P) = \begin{cases} T_h(\psi) + V_{ee}(\psi) & \text{if } P = P_\psi \text{ for some } \psi \in \mathcal{A} \\ +\infty & \text{otherwise.} \end{cases}$$

It follows that

$$F_h^{\mathcal{S}}(\rho) = \min_{\Pi(\rho)} \mathcal{F}_h^{\mathcal{S}}(P), \quad (3.1)$$

and

$$F_h^{\mathcal{A}}(\rho) = \min_{\Pi(\rho)} \mathcal{F}_h^{\mathcal{A}}(P). \quad (3.2)$$

Concerning the optimal transport problem we only need to incorporate the symmetry constraint in the transport functional. For every $\sigma \in \mathfrak{S}^N$ permutation of $\{1, \dots, N\}$ and every $P \in \mathcal{P}(\mathbb{R}^{Nd})$ we consider $\sigma_\# P$ the image measure via σ of the measure P , where with a little abuse of notations we have denoted by

$$\sigma(x_1, \dots, x_N) := (x_{\sigma(1)}, \dots, x_{\sigma(N)}).$$

For every $P \in \Pi(\rho)$ we may consider the measure

$$\tilde{P} := \frac{1}{N!} \sum_{\sigma \in \mathfrak{S}^N} \sigma_\# P.$$

We have that $\tilde{P} \in \Pi(\rho)$ and, since the cost is also permutation invariant the transport cost of \tilde{P} is the same as the cost of P . We say that P is symmetric if $\sigma_\# P = P$ for every $\sigma \in \mathfrak{S}^N$. Then, for example, the measure \tilde{P} above is symmetric. Define, then

Definition 3.1.

$$\mathfrak{C}_{\mathcal{S}}(P) = \begin{cases} \int cdP & \text{if } P \text{ is symmetric,} \\ +\infty & \text{otherwise.} \end{cases}$$

We will omit the \mathcal{S} and only use $\mathfrak{C}(P)$ if symmetry is not required.

By the previous discussion

$$C(\rho) = \min_{\Pi(\rho)} \mathfrak{C}_{\mathcal{S}}(P). \quad (3.3)$$

Then the common domain of minimization of $\mathcal{F}_h^{\mathcal{S}}$, $\mathcal{F}_h^{\mathcal{A}}$ and $\mathfrak{C}_{\mathcal{S}}$ is $\Pi(\rho)$ which we consider embedded in the space of probability measures \mathcal{P} equipped with the tight convergence.

Definition 3.2. A generalized sequence of wave functions $\{\psi_h\}$ converges to a transport plan P if $P_{\psi_h} \rightharpoonup P$.

We will prove

Theorem 3.3. For every $\rho \in H$ the functionals $\mathcal{F}_h^{\mathcal{S}}$ are mildly equicoercive and

$$\mathcal{F}_h^{\mathcal{S}} \xrightarrow{\Gamma} \mathfrak{C}_{\mathcal{S}},$$

with respect to the tight convergence of measures.

Theorem 3.4. For every $\rho \in H$ the functionals $\mathcal{F}_h^{\mathcal{A}}$ are mildly equicoercive and for $d = 2, 3, 4$ and $N = 2, 3$,

$$\mathcal{F}_h^{\mathcal{A}} \xrightarrow{\Gamma} \mathfrak{C}_{\mathcal{S}},$$

with respect to the tight convergence of measures.

The proofs of Theorems 3.3 and 3.4 above coincide for a large part and will constitute the rest of this section.

3.1. Equicoerciveness.

Lemma 3.5. $\Pi(\rho)$ is compact with respect to the weak convergence.

Proof. First we prove that $\Pi(\rho)$ is tight. In fact, for $\varepsilon > 0$ let $K \subseteq \mathbb{R}^d$ compact such that

$$\int_{K^c} \rho(x) dx \leq \varepsilon.$$

Observe that

$$(K^N)^c = \left(K^c \times \mathbb{R}^{d(N-1)} \right) \cup \left(\mathbb{R}^d \times K^c \times \mathbb{R}^{d(N-2)} \right) \cup \dots \cup \left(\mathbb{R}^{d(N-1)} \times K^c \right),$$

and for every $P \in \Pi(\rho)$

$$\int_{(K^N)^c} dP(X) \leq N\varepsilon$$

and K^N is compact. By Prokhorov's Theorem we deduce that $\Pi(\rho)$ is relatively compact. However, if $P_n \rightharpoonup P$ and $P_n \in \Pi(\rho)$, for every function $\phi(x_j) \in C_b(\mathbb{R}^{Nd})$ depending only on the j -th variable one has

$$\int_{\mathbb{R}^{Nd}} \phi(x_j) dP_n(X) = \int_{\mathbb{R}^d} \phi(x) \rho(x) dx$$

which implies

$$\int_{\mathbb{R}^d} \phi(x) \rho(x) dx = \int_{\mathbb{R}^{Nd}} \phi(x_j) dP(X) = \int_{\mathbb{R}^d} \phi(x) d\pi_{\#}^j(P)(x).$$

Since ϕ was arbitrary, $\pi_{\#}^j(P)(x) = \rho(x)$, and hence $P \in \Pi(\rho)$. \square

3.2. Γ -lim inf inequality.

Proposition 3.6. *Let $\rho \in H$ and let P a probability measure on \mathbb{R}^{Nd} . If $\{\psi_h\}_h \subseteq \mathcal{S}(\rho)$ (or $\mathcal{A}(\rho)$) is a generalized sequence such that $P_{\psi_h} \rightharpoonup P$. Then*

- (i) P is symmetric,
- (ii) $\liminf_{h \rightarrow 0} \mathcal{F}_h^{\mathcal{S}}(P_{\psi_h}) \geq \mathfrak{C}_{\mathcal{S}}(P)$.

Proof. It is easy to see that invariance with respect to permutations is a closed condition in the space of probability measures. For the inequality: $\liminf_{h \rightarrow 0} \mathcal{F}_h^{\mathcal{S}}(P_{\psi_h}) = \liminf_{h \rightarrow 0} T_h(\psi_h) + V_{ee}(\psi_h) \geq \liminf_{h \rightarrow 0} V_{ee}(\psi_h) \geq \mathfrak{C}_{\mathcal{S}}(P)$. \square

3.3. An approximation procedure for $P \in \Pi(\rho)$.

Proposition 3.7. *Let $\rho \in H$, and $P \in \Pi(\rho)$ be a transport plan such that*

$$P|_{D_\alpha} = 0 \tag{3.4}$$

for some $\alpha > 0$.

Then there exists a family of plans $\{P_\varepsilon\}_{\varepsilon > 0}$ such that:

- (i) for every $\varepsilon > 0$, $P_\varepsilon \in \Pi(\rho)$ and is absolutely continuous with respect to the Lebesgue measure, with density given by $\varphi_\varepsilon^2(X)$, where φ_ε is a suitable H^1 function;
- (ii) $P_\varepsilon \rightharpoonup P$ as $\varepsilon \rightarrow 0$;
- (iii) $\limsup_{\varepsilon \rightarrow 0} \mathfrak{C}(P_\varepsilon) \leq \mathfrak{C}(P)$;
- (iv) the kinetic energy of φ_ε is explicitly controlled:

$$\int |\nabla \varphi_\varepsilon(X)|^2 dX \leq N \left(\|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\varepsilon^2} \right)$$

for a suitable constant $K > 0$.

The proof is made of several steps.

We start regularizing by convolution. The plans we obtain are regular but do not have the same marginals as P . We use the standard mollifiers $\eta: \mathbb{R}^d \rightarrow \mathbb{R}$ as

$$\eta(u) = \begin{cases} ke^{-\frac{1}{1-|u|^2}} & |u| < 1 \\ 0 & |u| \geq 1 \end{cases}$$

where $k = k(d) > 0$ is a suitable constant, which depends only on the dimension d , such that $\int_{\mathbb{R}^d} \eta(u) du = 1$. Set also

$$\eta_\varepsilon(u) = \frac{1}{\varepsilon^d} \eta\left(\frac{u}{\varepsilon}\right).$$

The next Lemma is well known and it is a useful tool to estimate some L^2 norms which will appear later.

Lemma 3.8. *There exists a constant $K = K(d) > 0$, depending only on the dimension d , such that*

$$\int_{B(0,\varepsilon)} \frac{|\nabla \eta_\varepsilon(u)|^2}{\eta_\varepsilon(u)} du = \frac{K}{\varepsilon^2}.$$

Proof. Point out that

$$\nabla \eta(u) = \begin{cases} \frac{-2ku}{(1-|u|^2)^2} e^{-\frac{1}{1-|u|^2}} & |u| < 1 \\ 0 & |u| \geq 1 \end{cases}$$

and

$$\nabla \eta_\varepsilon(u) = \frac{1}{\varepsilon^{d+1}} \nabla \eta\left(\frac{u}{\varepsilon}\right).$$

Now

$$\begin{aligned} \int_{B(0,\varepsilon)} \frac{|\nabla\eta_\varepsilon(u)|^2}{\eta_\varepsilon(u)} du &= \int_{B(0,\varepsilon)} \frac{1}{\varepsilon^{d+2}} \frac{|\nabla\eta(u/\varepsilon)|^2}{\eta(u/\varepsilon)} du = \frac{1}{\varepsilon^2} \int_{B(0,1)} \frac{|\nabla\eta(v)|^2}{\eta(v)} dv \\ &= \frac{1}{\varepsilon^2} \int_{B(0,1)} \frac{4k|v|^2}{(1-|v|^2)^4} e^{-\frac{1}{1-|v|^2}} dv = \frac{K(d)}{\varepsilon^2}. \end{aligned}$$

□

Now we define the function

$$H_\varepsilon(Y - X) = \prod_{i=1}^N \eta_\varepsilon(y_i - x_i).$$

and use it as mollifier to regularize the transport plan P defining

$$\tilde{P}_\varepsilon(Y) = \int H_\varepsilon(Y - X) dP(X).$$

Note that the marginals of \tilde{P}_ε^3 are different from ρ , but may be written explicitly:

$$\begin{aligned} \rho_\varepsilon(y) &= \int_{\mathbb{R}^{(N-1)d}} \tilde{P}_\varepsilon(y, y_2, \dots, y_N) dy_2 \dots dy_N \\ &= \int_{\mathbb{R}^{(N-1)d}} \int \eta_\varepsilon(y - x_1) \eta_\varepsilon(y_2 - x_2) \dots \eta_\varepsilon(y_N - x_N) dP(X) dy_2 \dots dy_N \\ &= \int \eta_\varepsilon(y - x_1) dP(X) = \int_{\mathbb{R}^d} \rho(x) \eta_\varepsilon(y - x) dx = (\rho * \eta_\varepsilon)(y) \end{aligned}$$

Lemma 3.9. *Let α be as in the statement of Proposition 3.7 and let $Y \in \mathbb{R}^{Nd}$ be such that $|y_i - y_j| < \alpha/2$ for some $i \neq j$, and $\varepsilon < \alpha/4$, then $\tilde{P}_\varepsilon(Y) = 0$.*

Proof. Note that

$$\tilde{P}_\varepsilon(Y) = \int H_\varepsilon(Y - X) \prod_{j=1}^N \chi_{B(y_j, \varepsilon)}(x_j) dP(X).$$

If Y and ε are as in the statement, and $x_i \in B(y_i, \varepsilon)$, $x_j \in B(y_j, \varepsilon)$, then

$$|x_i - x_j| \leq |x_i - y_i| + |y_i - y_j| + |y_j - x_j| \leq \alpha.$$

The thesis follows from (3.4).

□

Define now

$$\tilde{\varphi}_\varepsilon(Y) = \sqrt{\tilde{P}_\varepsilon(Y)}.$$

Lemma 3.10. *For every $\varepsilon > 0$, $\tilde{\varphi}_\varepsilon \in H^1(\mathbb{R}^{Nd})$.*

Proof. Fix $\varepsilon > 0$. Clearly $\tilde{\varphi}_\varepsilon$ is L^2 , since

$$\begin{aligned} \int \tilde{\varphi}_\varepsilon^2(Y) dY &= \int \tilde{P}_\varepsilon(Y) dY = \iint H_\varepsilon(Y - X) dP(X) dY \\ &= \iint H_\varepsilon(Y - X) dY dP(X) = \int dP(X) = 1. \end{aligned}$$

³As usual we mean the marginals of $\tilde{P}_\varepsilon(Y) dY$.

Now we estimate $|\nabla\tilde{\varphi}_\varepsilon|^2$. Using the Cauchy-Schwarz inequality,

$$\begin{aligned} \left| \nabla\tilde{P}_\varepsilon(Y) \right| &\leq \int |\nabla H_\varepsilon(Y - X)| dP(X) \\ &\leq \sqrt{\int \frac{|\nabla H_\varepsilon(Y - X)|^2}{H_\varepsilon(Y - X)} dP(X)} \sqrt{\int H_\varepsilon(Y - X) dP(X)} \\ &= \sqrt{\int \frac{|\nabla H_\varepsilon(Y - X)|^2}{H_\varepsilon(Y - X)} dP(X)} \sqrt{\tilde{P}_\varepsilon(Y)} \end{aligned}$$

where the first integral is extended to the set where the integrand is defined, namely $|x_j - y_j| < \varepsilon \forall j$. Therefore, with the same convention,

$$\begin{aligned} \int |\nabla\tilde{\varphi}_\varepsilon(Y)|^2 dY &= \frac{1}{4} \int \frac{|\nabla\tilde{P}_\varepsilon(Y)|^2}{\tilde{P}_\varepsilon(Y)} dY \leq \frac{1}{4} \iint \frac{|\nabla H_\varepsilon(Y - X)|^2}{H_\varepsilon(Y - X)} dP(X) dY \\ &= \frac{1}{4} \sum_{j=1}^N \iint \frac{|\nabla_j H_\varepsilon(Y - X)|^2}{H_\varepsilon(Y - X)} dY dP(X) \\ &\leq \frac{1}{4} \sum_{j=1}^N \iint \frac{|\nabla\eta_\varepsilon(y_j - x_j)|^2}{\eta_\varepsilon(y_j - x_j)} dy_j dP(X) \\ &= \frac{N}{4} \int \frac{|\nabla\eta_\varepsilon(u)|^2}{\eta_\varepsilon(u)} du = \frac{KN}{4\varepsilon^2}. \end{aligned}$$

□

In the next step we introduce a natural technique to get back the original marginals ρ without losing too much regularity. This technique is original and different from the one presented in [11]. We point out that, in a different context, this construction may well be generalized to a plan with different marginals ρ^1, \dots, ρ^N .

The construction fits in the general scheme of composition of transport plans as presented in [1].

For $x, y \in \mathbb{R}^d$ define

$$\gamma_\varepsilon(x, y) := \frac{\rho(x)\eta_\varepsilon(y - x)}{\rho_\varepsilon(y)}$$

with the convention that it is zero if $\rho_\varepsilon(y) = 0$. The two variables function $\rho(x)\eta_\varepsilon(y - x)$ is the key point of the construction, since it has the following property that links the different marginals:

$$\int_{\mathbb{R}^d} \rho(x)\eta_\varepsilon(y - x) dx = \rho_\varepsilon(y), \quad \int_{\mathbb{R}^d} \rho(x)\eta_\varepsilon(y - x) dy = \rho(x).$$

To simplify the notation we set

$$\begin{aligned} m_\varepsilon(X, Y) &= \prod_{i=1}^N \gamma_\varepsilon(x_i, y_i), \\ m_\varepsilon^j(X, Y) &= \prod_{\substack{i=1 \\ i \neq j}}^N \gamma_\varepsilon(x_i, y_i). \end{aligned}$$

Note that

$$\int_{\mathbb{R}^d} \gamma_\varepsilon(x, y) dx = \chi_{\rho_\varepsilon > 0}(y).$$

Now set

$$\Gamma_\varepsilon(X, Y) := \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon(X, Y)$$

and observe that

$$\int \Gamma_\varepsilon(X, Y) dX = \tilde{\varphi}_\varepsilon^2(Y) \prod_{i=1}^N \chi_{\rho_\varepsilon > 0}(y_i) = \tilde{\varphi}_\varepsilon^2(Y) = \tilde{P}_\varepsilon(Y)$$

where we used the following remark, which will be implicit from now on:

Remark 3.11. If Y is such that $\rho_\varepsilon(y_i) = 0$, then

$$0 = \rho_\varepsilon(y_i) = \int_{\mathbb{R}^{(N-1)d}} \tilde{\varphi}_\varepsilon(y_1, y_2, \dots, y_n)^2 dy_1 \dots \hat{d}y_i \dots dy_N$$

and hence $\tilde{\varphi}_\varepsilon(Y) = 0$.

Define

$$P_\varepsilon(X) = \int \Gamma_\varepsilon(X, Y) dY$$

and calculate the marginals of P_ε to get

$$\begin{aligned} \int_{\mathbb{R}^{(N-1)d}} P_\varepsilon(X) dx_2 \dots dx_N &= \int_{\mathbb{R}^{(N-1)d}} \int \Gamma_\varepsilon(X, Y) dY dx_2 \dots dx_N \\ &= \int \tilde{\varphi}_\varepsilon^2(Y) \gamma_\varepsilon(x_1, y_1) dY \\ &= \int_{\mathbb{R}^d} \rho_\varepsilon(y_1) \gamma_\varepsilon(x_1, y_1) dy_1 \\ &= \rho(x_1) \end{aligned}$$

and similarly also the other $N - 1$ marginals are equal to ρ .

Lemma 3.12. *Let α be as in the statement of Proposition 3.4. If $|x_i - x_j| < \alpha/4$ for some $i \neq j$, and $\varepsilon < \alpha/8$, then $P_\varepsilon(X) = 0$.*

Proof. Fix X and ε as in the statement, and suppose $m_\varepsilon(X, Y) > 0$. Then necessarily $|y_i - x_i| < \varepsilon$ and $|y_j - x_j| < \varepsilon$, so that

$$|y_i - y_j| \leq |y_i - x_i| + |x_i - x_j| + |x_j - y_j| \leq \alpha/2.$$

From Lemma 3.9 it follows that $\tilde{\varphi}_\varepsilon^2(Y) = 0$. □

We define now the function $\varphi_\varepsilon(X) = \sqrt{P_\varepsilon(X)}$, and proceed to estimate its kinetic energy.

Estimate for the kinetic energy. Calculate first the gradient with respect to x_j of P_ε :

$$\begin{aligned} \nabla_j P_\varepsilon(X) &= \nabla_j \int \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon^j(X, Y) \frac{\rho(x_j) \eta_\varepsilon(y_j - x_j)}{\rho_\varepsilon(y_j)} dY \\ &= \int \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon^j(X, Y) \frac{\nabla \rho(x_j) \eta_\varepsilon(y_j - x_j)}{\rho_\varepsilon(y_j)} dY \\ &\quad - \int \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon^j(X, Y) \frac{\rho(x_j) \nabla \eta_\varepsilon(y_j - x_j)}{\rho_\varepsilon(y_j)} dY \\ &= A(X) + B(X). \end{aligned}$$

We define for simplicity

$$J(X) = \int \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon^j(X, Y) \frac{\eta_\varepsilon(y_j - x_j)}{\rho_\varepsilon(y_j)} dY$$

so that, for example,

$$P_\varepsilon(X) = \rho(x_j) J(X),$$

$$A(X) = \nabla \rho(x_j) J(X).$$

Now

$$\nabla_j \varphi_\varepsilon(X) = \nabla_j \sqrt{P_\varepsilon(X)} = \frac{\nabla_j P_\varepsilon(X)}{2\sqrt{P_\varepsilon(X)}} = \frac{1}{2\sqrt{P_\varepsilon(X)}} [A(X) + B(X)]$$

and we estimate the square of the L^2 norm of every term.

$$\begin{aligned} \int \frac{|A(X)|^2}{P_\varepsilon(X)} dX &= \int \frac{|\nabla \rho(x_j)|^2}{\rho(x_j)} J(X) dX \\ &= \iint \frac{|\nabla \rho(x_j)|^2}{\rho(x_j)} \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon^j(X, Y) \frac{\eta_\varepsilon(y_j - x_j)}{\rho_\varepsilon(y_j)} dY dX \\ &= \int_{\mathbb{R}^d} \left(\int \frac{|\nabla \rho(x_j)|^2}{\rho(x_j)} \tilde{\varphi}_\varepsilon^2(Y) \frac{\eta_\varepsilon(y_j - x_j)}{\rho_\varepsilon(y_j)} dY \right) dx_j \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|\nabla \rho(x_j)|^2}{\rho(x_j)} \eta_\varepsilon(y_j - x_j) dy_j dx_j \\ &= \int_{\mathbb{R}^d} \frac{|\nabla \rho(x_j)|^2}{\rho(x_j)} dx_j \\ &\leq 4 \|\sqrt{\rho}\|_{H^1}^2. \end{aligned}$$

By Cauchy-Schwarz inequality,

$$|B(X)|^2 \leq \rho(x_j) J(X) \cdot \int \tilde{\varphi}_\varepsilon(Y)^2 m_\varepsilon^j(X, Y) \frac{\rho(x_j) |\nabla \eta_\varepsilon(y_j - x_j)|^2}{\rho_\varepsilon(y_j) \eta_\varepsilon(y_j - x_j)} dY.$$

(Here the integral is extended to the region where $\eta_\varepsilon(y_j - x_j) > 0$.) Therefore, with the same convention,

$$\begin{aligned} \int \frac{|B(X)|^2}{P_\varepsilon(X)} dX &\leq \iint \tilde{\varphi}_\varepsilon(Y)^2 m_\varepsilon^j(X, Y) \frac{\rho(x_j) |\nabla \eta_\varepsilon(y_j - x_j)|^2}{\rho_\varepsilon(y_j) \eta_\varepsilon(y_j - x_j)} dY dX \\ &= \int_{\mathbb{R}^d} \int \tilde{\varphi}_\varepsilon(Y)^2 \frac{\rho(x_j) |\nabla \eta_\varepsilon(y_j - x_j)|^2}{\rho_\varepsilon(y_j) \eta_\varepsilon(y_j - x_j)} dY dx_j \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\rho(x_j) |\nabla \eta_\varepsilon(y_j - x_j)|^2}{\eta_\varepsilon(y_j - x_j)} dy_j dx_j = \int_{\mathbb{R}^d} \frac{|\nabla \eta_\varepsilon(y)|^2}{\eta_\varepsilon(y)} dy = \frac{K}{\varepsilon^2}. \end{aligned}$$

Moreover,

$$\begin{aligned} \int \frac{A(X) \cdot B(X)}{4P_\varepsilon(X)} dX &= -\frac{1}{4} \iint \frac{\nabla \rho(x_j) \cdot \nabla \eta_\varepsilon(x_j - y_j)}{\rho_\varepsilon(y_j)} \tilde{\varphi}_\varepsilon^2(Y) m_\varepsilon^j(X, Y) dY dX \\ &= -\frac{1}{4} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \nabla \rho(x_j) \cdot \nabla \eta_\varepsilon(x_j - y_j) dx_j dy_j \\ &= -\frac{1}{4} \left(\int_{\mathbb{R}^d} \nabla \rho(x) dx \right) \cdot \left(\int_{\mathbb{R}^d} \nabla \eta_\varepsilon(z) dz \right) \end{aligned}$$

and the second factor is equal to 0, as is easy to see integrating in spherical coordinates.

Thus

$$\int |\nabla_j \varphi_\varepsilon(X)|^2 dX \leq \frac{1}{4} \int \frac{|A(X)|^2 + |B(X)|^2}{P_\varepsilon(X)} dX \leq \|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\varepsilon^2},$$

and by summation over j

$$\int |\nabla \varphi_\varepsilon(X)|^2 dX \leq N \left(\|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\varepsilon^2} \right).$$

Here we prove (ii). First we give the following lemma, which specifies that, for $\varepsilon \rightarrow 0$, the mass of \tilde{P}_ε is concentrated near the mass of P .

Lemma 3.13. *Suppose $R, \delta > 0$ are such that*

$$\int_{|X|>R} dP(X) \leq \delta;$$

then, if $\varepsilon\sqrt{N} < R$,

$$\int_{|Y|>2R} \tilde{P}_\varepsilon(Y) dY \leq \delta.$$

Proof.

$$\begin{aligned} \int_{Y>2R} \tilde{P}_\varepsilon(Y) dY &= \iint_{|Y|>2R} H_\varepsilon(Y-X) dP(X) dY \\ &= \iint_{|Y|>2R \cap |X|>R} H_\varepsilon(Y-X) dP(X) dY \\ &\leq \iint_{|X|>R} H_\varepsilon(Y-X) dY dP(X) = \int_{|X|>R} dP(X) \leq \delta \end{aligned}$$

since, where $|X| \leq R$, one has $|Y-X| \geq R > \varepsilon\sqrt{N}$, and hence there exists i such that $|x_i - y_i| > \varepsilon$. \square

Next, to prove that $P_\varepsilon \rightarrow P$, we interpolate by \tilde{P}_ε in between:

Lemma 3.14. $\tilde{P}_\varepsilon \rightarrow P$.

Proof. Let $\phi \in C_b(\mathbb{R}^{Nd})$;

$$\begin{aligned} \left| \int \phi(Y) d\tilde{P}_\varepsilon(Y) - \int \phi(X) dP(X) \right| &= \left| \iint [\phi(Y) - \phi(X)] H_\varepsilon(Y-X) dP(X) dY \right| \\ &\leq \iint |\phi(Y) - \phi(X)| H_\varepsilon(Y-X) dP(X) dY. \end{aligned}$$

Given $\delta > 0$, let R be such that the hypothesis of Lemma 3.13 holds. We divide $\mathbb{R}^{Nd} \times \mathbb{R}^{Nd}$ in three disjoint regions:

$$E_1 = \{|X| > R\} \quad E_2 = \{|X| \leq R, |Y| \leq 2R\} \quad E_3 = \{|X| \leq R, |Y| > 2R\}.$$

As before, if $\varepsilon\sqrt{N} < R$, on E_3 one has $H_\varepsilon(X-Y) \equiv 0$.

$$\begin{aligned} \iint_{E_1} |\phi(Y) - \phi(X)| H_\varepsilon(Y-X) dP(X) dY &\leq 2\|\phi\|_\infty \iint_{E_1} H_\varepsilon(Y-X) dP(X) dY \\ &\leq 2\delta\|\phi\|_\infty. \end{aligned}$$

On the other hand, E_2 is compact; take ε_0 such that $|X-Y| \leq \varepsilon_0$ implies $|\phi(X) - \phi(Y)| \leq \delta$. If $\varepsilon\sqrt{N} \leq \varepsilon_0$ we get

$$\begin{aligned} \iint_{E_2} |\phi(Y) - \phi(X)| H_\varepsilon(Y-X) dP(X) dY &\leq \delta \iint_{E_2} H_\varepsilon(Y-X) dP(X) dY \\ &\leq \delta \iint H_\varepsilon(Y-X) dP(X) dY = \delta. \end{aligned}$$

\square

Lemma 3.15. $P_\varepsilon \rightharpoonup P$.

Proof. Let $\phi \in C_b(\mathbb{R}^{Nd})$. Using the fact that $\tilde{P}_\varepsilon \rightharpoonup P$ (Lemma 3.14), it is left to estimate

$$\begin{aligned} \left| \int \phi(X) P_\varepsilon(X) dX - \int \phi(Y) \tilde{P}_\varepsilon(Y) dY \right| &= \left| \iint [\phi(X) - \phi(Y)] \Gamma_\varepsilon(X, Y) dX dY \right| \\ &\leq \iint |\phi(X) - \phi(Y)| \Gamma_\varepsilon(X, Y) dX dY \end{aligned}$$

As in the proof of Lemma 3.14, given $\delta > 0$ let R be such that the hypothesis of Lemma 3.13 holds. We divide $\mathbb{R}^{Nd} \times \mathbb{R}^{Nd}$ in three disjoint regions:

$$E_1 = \{|Y| > 2R\} \quad E_2 = \{|Y| \leq 2R, |X| \leq 3R\} \quad E_3 = \{|Y| \leq 2R, |X| > 3R\}.$$

If $\varepsilon\sqrt{N} < R$, as before, on E_3 the integral is zero since $\Gamma_\varepsilon(X, Y) \equiv 0$ there. Thanks to Lemma 3.13,

$$\begin{aligned} \iint_{E_1} |\phi(X) - \phi(Y)| \Gamma_\varepsilon(X, Y) dX dY &\leq 2\|\phi\|_\infty \iint_{E_1} \Gamma_\varepsilon(X, Y) dX dY \\ &= 2\|\phi\|_\infty \int_{\{|Y| > 2R\}} \tilde{P}_\varepsilon(Y) dY \leq 2\delta\|\phi\|_\infty \end{aligned}$$

Exactly as before, using that E_2 is compact and ϕ is absolutely continuous the thesis follows. \square

It is left to prove (iii). The cost function c is not continuous neither bounded. However, recall Lemma 3.12, it is bounded on the complement of $D_{\alpha/4}$. With this in mind, consider the function $v: \mathbb{R}^{2d} \rightarrow \mathbb{R}$ defined as

$$v(x, y) = \begin{cases} \frac{1}{|x - y|} & \text{if } |x - y| \geq \alpha/4 \\ 4/\alpha & \text{elsewhere.} \end{cases}$$

and set

$$\bar{c}(X) = \sum_{1 \leq i < j \leq N} v(x_i, x_j).$$

Clearly $\bar{c}(X) \leq c(X)$, and \bar{c} is continuous (sum of continuous functions), bounded by $\binom{N}{2} \frac{4}{\alpha}$; moreover, thanks to the property (3.4) and Lemma 3.12,

$$\int \bar{c}(X) dP(X) = \int c(X) dP(X), \quad \int \bar{c}(X) P_\varepsilon(X) dX = \int c(X) P_\varepsilon(X) dX.$$

We can conclude the estimate as follows:

$$\limsup_{\varepsilon \rightarrow 0} \mathfrak{C}(P_\varepsilon) = \limsup_{\varepsilon \rightarrow 0} \int \bar{c}(X) P_\varepsilon(X) dX = \int \bar{c}(X) dP(X) = \int c(X) dP(X). \quad \square$$

Proposition 3.7 may be extended to all plans. Since we will make use of Th. 2.7 we limit ourself to the case of symmetric transport plans. The same proof for non symmetric plans requires a version of Th. 2.7 for plans with different marginals. The extension is contained, for example, in [3].

Proposition 3.16. *Let $P \in \Pi(\rho)$ a symmetric plan (not necessarily satisfying the property (3.4)). Then there exists a family of plans $\{P_\varepsilon\}_{\varepsilon > 0}$ such that:*

- (i) for every $\varepsilon > 0$, $P_\varepsilon \in \Pi(\rho)$ and is absolutely continuous with respect to the Lebesgue measure, with density given by $\varphi_\varepsilon^2(X)$, where φ_ε is a suitable H^1 function;
- (ii) $P_\varepsilon \rightharpoonup P$ as $\varepsilon \rightarrow 0$;

(iii) $\limsup_{\varepsilon \rightarrow 0} \mathfrak{C}_S(P_\varepsilon) \leq \mathfrak{C}_S(P)$;

(iv) the kinetic energy of φ_ε is explicitly controlled:

$$\int |\nabla \varphi_\varepsilon(X)|^2 dX \leq N \left(\|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\varepsilon^2} \right)$$

for a suitable constant $K > 0$.

Proof. The proof of (i), (ii) and (iv) holds in general since it does not require (3.4). And, in fact, carefully following the constructions in Propositions 3.7 one may observe that if P is permutations invariant then the approximating P_ε have the same property. Then we only need to prove (iii) and so if $\mathfrak{C}_S(P) = \infty$ there is nothing to prove. Suppose $\mathfrak{C}_S(P) = K < \infty$. Let $r > 0$ be a parameter, and split

$$P = Q_r + P|_{D_r}.$$

Let σ_r be the marginals of Q_r , and $\tilde{\rho}_r$ those of $P|_{D_r}$; clearly $\sigma_r + \tilde{\rho}_r = \rho$. Since $\rho \in L^1 \cap L^{\frac{d}{d-2}}$ by Sobolev embedding, and $\tilde{\rho}_r \leq \rho$ pointwise, we have $\tilde{\rho}_r \in L^1 \cap L^{\frac{d}{d-2}}$. Although $\tilde{\rho}_r$ needs not to be a probability measure on \mathbb{R}^d , we can suppose there exists $\lambda_r > 0$ such that

$$\int_{\mathbb{R}^d} \tilde{\rho}_r(x) dx = \frac{1}{\lambda_r} < 1,$$

otherwise $P|_{D_r} = 0$ and we get the result directly by Proposition 3.7. Let now \tilde{P}_r be a symmetric optimal transport plan in $\Pi(\lambda_r \tilde{\rho}_r)$, and define

$$P_r = Q_r + \frac{\tilde{P}_r}{\lambda_r},$$

which lies in $\Pi(\rho)$. On the one hand we have the following

Lemma 3.17. $P_r \rightarrow P$.

Proof. Recall that $\mathfrak{C}(P)$ is finite to get

$$K = \mathfrak{C}(P) \geq \mathfrak{C}(P|_{D_r}) \geq \frac{1}{r} P(D_r),$$

and *a fortiori* for \tilde{P}_r/λ_r due to the optimality. Hence

$$\lim_{r \rightarrow 0} P(D_r) = \lim_{r \rightarrow 0} \frac{\tilde{P}_r(D_r)}{\lambda_r} = 0.$$

Take $f \in C_b(\mathbb{R}^{Nd})$, and estimate

$$\begin{aligned} \left| \int f(X) dP_r(X) - \int f(X) dP(X) \right| &= \left| \int f(X) dP|_{D_r}(X) - \frac{1}{\lambda_r} \int f(X) d\tilde{P}_r(X) \right| \\ &\leq \|f\|_\infty \left(P(D_r) + \frac{\tilde{P}_r(D_r)}{\lambda_r} \right) \rightarrow 0 \end{aligned}$$

as $r \rightarrow 0$. □

On the other hand,

$$\mathfrak{C}_S(P_r) = \mathfrak{C}_S(Q_r) + \frac{\mathfrak{C}_S(\tilde{P}_r)}{\lambda_r} \leq \mathfrak{C}_S(Q_r) + \mathfrak{C}_S(P|_{D_r}) = \mathfrak{C}_S(P),$$

thus

$$\limsup_{r \rightarrow 0} \mathfrak{C}_S(P_r) \leq \mathfrak{C}_S(P).$$

Thanks to Proposition 2.5 $\mathfrak{C}_S(\tilde{P}_r)$ is finite, and by Theorem 2.7 there exists $\alpha = \alpha(r) > 0$ such that \tilde{P}_r is supported outside D_α .⁴ Recall now Proposition 3.7 to find $\varphi_{\varepsilon,r}$ weakly converging to P_r as $\varepsilon \rightarrow 0$, with

$$\int |\nabla \varphi_{\varepsilon,r}(X)|^2 dX \leq N \left(\|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\varepsilon^2} \right)$$

and

$$\limsup_{\varepsilon \rightarrow 0} \mathfrak{C}_S(|\varphi_{\varepsilon,r}|^2) = \mathfrak{C}_S(P_r).$$

It suffices now to take $\{\varphi_{r,r}\}_{r>0}$ to conclude. In fact, given $\delta > 0$, let R be such that

$$\int_{\{|X|>R\}} dP_r(X) \leq \delta.$$

Note that R may be chose independent from r , since the marginals of P_r are all equal to ρ , and we may choose $K \subseteq \mathbb{R}^d$ compact such that

$$\int_K \rho(x) dx \leq \frac{\delta}{N},$$

and R sufficiently large such that $K^N \subseteq B(0, R)_{\mathbb{R}^{Nd}}$.

Now to prove weak convergence take $\phi \in C_b(\mathbb{R}^{Nd})$ and proceed as in the previous paragraph to estimate

$$\left| \int \phi(X) \varphi_{r,r}^2(X) dX - \int \phi(X) dP_r(X) \right|,$$

using in addition Lemma 3.17 to estimate

$$\left| \int \phi(X) dP_r(X) - \int \phi(X) dP(X) \right|.$$

□

3.4. Constructing wave-functions. A wave-function depends on N space-spin variables. In the previous subsections we worked mainly in \mathbb{R}^{Nd} , since we were considering transport plans in $\Pi(\rho)$. To introduce the spin we will separate the spin dependence as follows: for every s binary string of length N we consider the function $\psi_s(x_1, \dots, x_N) = \psi(x_1, s_1, \dots, x_N, s_N)$, then we describe ψ as a 2^N -dimensional vector

$$\boldsymbol{\psi}(z_1, \dots, z_N) = (\psi_s(X))_{s \in S}.$$

As an example, if $N = 2$ we would have

$$\boldsymbol{\psi}(z_1, z_2) = \begin{pmatrix} \psi_{00}(x_1, x_2) \\ \psi_{01}(x_1, x_2) \\ \psi_{10}(x_1, x_2) \\ \psi_{11}(x_1, x_2) \end{pmatrix},$$

and for $N = 3$ a wave-function would be represented as

$$\boldsymbol{\psi}(z_1, z_2, z_3) = \begin{pmatrix} \psi_{000}(x_1, x_2, x_3) \\ \psi_{001}(x_1, x_2, x_3) \\ \psi_{010}(x_1, x_2, x_3) \\ \psi_{011}(x_1, x_2, x_3) \\ \psi_{100}(x_1, x_2, x_3) \\ \psi_{101}(x_1, x_2, x_3) \\ \psi_{110}(x_1, x_2, x_3) \\ \psi_{111}(x_1, x_2, x_3) \end{pmatrix}.$$

⁴Observe that $\alpha(r)$ may be chose decreasing as $r \rightarrow 0$, as follows from Theorem 2.7.

Note that now the density $|\psi(X)|^2$ is simply the square of the Euclidean norm of the vector ψ , and the same holds for $\nabla\psi$, once we set

$$\nabla\psi(z_1, \dots, z_N) = (\nabla\psi_s(X))_{s \in S}.$$

Let us take now a fermionic (i.e., antisymmetric) wave-function ψ , and consider a spin state $s = (s_1, \dots, s_N)$. If $i < j$ are such that $s_i = s_j$, consider $\sigma = (i j) \in \mathfrak{S}_N$ to get

$$\begin{aligned} \psi_s(x_1, \dots, x_N) &= \text{sgn}(\sigma)\psi_{\sigma(s)}(x_{\sigma(1)}, \dots, x_{\sigma(N)}) \\ &= -\psi_s(x_1, \dots, x_j, \dots, x_i, \dots, x_N). \end{aligned}$$

Hence we get the following

Remark 3.18. If ψ is fermionic and s is a spin state, ψ_s is *separately* antisymmetric with respect to the spatial variables such that $s_j = 0$, and with respect to the spatial variables such that $s_j = 1$.

Consider now two spin states s and s' with the same number of ones and zeroes. Then ψ_s and $\psi_{s'}$ are related: taking $\sigma \in \mathfrak{S}_N$ such that $\sigma(s) = s'$, we get

$$\psi_{s'}(x_{\sigma(1)}, \dots, x_{\sigma(N)}) = \text{sgn}(\sigma)\psi_s(x_1, \dots, x_N).$$

These observations will be used in the following.

3.5. Fermionic wave-functions with given density. Suppose we have $\alpha > 0$ and a symmetric function $\psi(x_1, \dots, x_N) \geq 0$ of H^1 with the property that

$$\psi(X) = 0 \text{ if } |x_i - x_j| < \alpha \text{ for some } i \neq j. \quad (3.5)$$

We wonder if there exists a fermionic wave-function ψ such that

$$\sum_{s \in S} |\psi(x_1, s_1, \dots, x_N, s_N)|^2 = \psi^2(X),$$

and

$$\|\psi\|_{H^1} \leq C\|\psi\|_{H^1}$$

for a suitable constant C . In fact, we managed to prove the following

Proposition 3.19. *For $N = 2, 3$, $d = 3, 4$, given $\psi \in H^1$ symmetric, with $\psi|_{D_\alpha} = 0$ for some $\alpha > 0$, there exists ψ fermionic such that*

$$\sum_{s \in S} |\psi_s(X)|^2 = \psi^2(X)$$

and

$$\sum_{s \in S} |\nabla\psi_s(X)|^2 \leq |\nabla\psi(X)|^2 + \frac{C}{\alpha^2}\psi^2(X).$$

In [11], the following ψ is given as a wave-function:

$$\begin{aligned} \psi_{00} &= 0 \\ \psi_{01} &= \psi(x_1, x_2) \\ \psi_{10} &= -\psi(x_1, x_2) \\ \psi_{11} &= 0, \end{aligned}$$

which is in fact fermionic with bounded kinetic energy. Note, however, that this construction cannot work for a larger number of particles. Indeed, if a binary string s has length $N \geq 3$, then there are at least two ones, or two zeros, on places $i \neq j$ – thus the corresponding function ψ_s must change sign for a suitable flip of the variables (namely, $x_i \mapsto x_j$, $x_j \mapsto x_i$). We will exhibit a wave-function ψ (with square density ψ^2) such that $\psi_{10} = \psi_{01} = 0$. This forces ψ_{00} and ψ_{11} to be different from zero and antisymmetric. We remark also that, for this kind of construction, the condition (3.5) is “morally necessary”.

3.6. Construction for $N = 2$, $d = 3$. The variable X will be expanded as $X = (x, y) = (x_1, x_2, x_3; y_1, y_2, y_3)$. Set $r = \alpha/\sqrt{3}$, and take $\xi = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ a primitive cubic root of 1. The key point is to choose two auxiliary C^∞ functions $a, b: \mathbb{R} \rightarrow \mathbb{R}$ such that

- (i) $a^2 + b^2 = 1$;
- (ii) b is symmetric, $b(t) = 0$ if $|t| \geq r$;
- (iii) a is antisymmetric, $a(t) = -1$ if $t \leq -r$, $a(t) = 1$ if $t \geq r$;
- (iv) $|a'|, |b'| \leq k/r$.

Note that the constant $k > 1$ may be chosen arbitrarily close to 1. To shorten the notation we set $a_j = a(x_j - y_j)$, $b_j = b(x_j - y_j)$ for $j = 1, 2, 3$. Now define

$$g_1(x, y) = \frac{1}{\sqrt{3}}(a_1 + b_1 a_2 + b_1 b_2 a_3)$$

$$g_\xi(x, y) = \frac{\sqrt{2}}{\sqrt{3}}(a_1 + \xi b_1 a_2 + \xi^2 b_1 b_2 a_3).$$

By direct computation one sees that $|g_1|^2 + |g_\xi|^2 = a_1^2 + b_1^2 a_2^2 + b_1^2 b_2^2 a_3^2$, since clearly $|a_1 + \bar{\xi} b_1 a_2 + \bar{\xi}^2 b_1 b_2 a_3| = |a_1 + \xi b_1 a_2 + \xi^2 b_1 b_2 a_3|$. Next we define the wave-function

$$\begin{aligned} \psi_{00}(x, y) &= g_1(x, y)\psi(x, y) \\ \psi_{01}(x, y) &= 0 \\ \psi_{10}(x, y) &= 0 \\ \psi_{11}(x, y) &= g_\xi(x, y)\psi(x, y). \end{aligned}$$

The following equality is crucial in the construction

$$\psi^2(x, y) b_1^2 b_2^2 a_3^2 = \psi^2(x, y) b_1^2 b_2^2. \quad (3.6)$$

This holds because where $|a_3|^2 = 1$, i.e. where $|x_3 - y_3| \geq r$, the equality holds. It also holds where $b_1 = 0$ or $b_2 = 0$, i.e. where $|x_1 - y_1| \geq r$ or $|x_2 - y_2| \geq r$. It is left the region where $|x_j - y_j| \leq r$ for every $j = 1, 2, 3$, but there it holds

$$|x - y| = \sqrt{|x_1 - y_1|^2 + |x_2 - y_2|^2 + |x_3 - y_3|^2} \leq r\sqrt{3} = \alpha,$$

and hence the equality holds because $\psi^2(x, y) = 0$.

Now one can compute

$$\begin{aligned} |\psi_{00}|^2 + |\psi_{11}|^2 &= \left(|g_1(x, y)|^2 + |g_\xi(x, y)|^2 \right) \psi^2(x, y) \\ &= (a_1^2 + b_1^2 a_2^2 + b_1^2 b_2^2 a_3^2) \psi^2(x, y) \\ &= (a_1^2 + b_1^2 a_2^2 + b_1^2 b_2^2) \psi^2(x, y) \\ &= (a_1^2 + b_1^2) \psi^2(x, y) = \psi^2(x, y). \end{aligned}$$

Next come the estimates for the derivatives. Since

$$\begin{aligned} \nabla \psi_{00}(x, y) &= \psi(x, y) \nabla g_1(x, y) + g_1(x, y) \nabla \psi(x, y) \\ \nabla \psi_{11}(x, y) &= \psi(x, y) \nabla g_\xi(x, y) + g_\xi(x, y) \nabla \psi(x, y), \end{aligned}$$

it follows

$$\begin{aligned} |\nabla \psi(x, y)|^2 &= |\nabla \psi(x, y)|^2 + \left(|\nabla g_1(x, y)|^2 + |\nabla g_\xi(x, y)|^2 \right) \psi^2(x, y) \\ &\quad + \psi(x, y) \nabla \psi(x, y) \cdot v(x, y) \end{aligned}$$

where

$$v(x, y) = 2g_1(x, y)\nabla g_1(x, y) + g_{\bar{\xi}}(x, y)\nabla g_{\bar{\xi}}(x, y) + g_{\xi}(x, y)\nabla g_{\xi}(x, y).$$

We claim that $\psi(x, y)v(x, y) = 0$. Again by direct computation one gets

$$v = 6[a_1\nabla a_1 + b_1a_2\nabla(b_1a_2) + b_1b_2a_3\nabla(b_1b_2a_3)].$$

Since the next steps work for general d , we group the result in the following

Lemma 3.20. *Let a_j, b_j be defined as before and evaluated in the point $x - y$. Then*

$$[a_1\nabla a_1 + (b_1a_2)\nabla(b_1a_2) + \cdots + (b_1b_2 \cdots b_{d-1}a_d)\nabla(b_1b_2 \cdots b_{d-1}a_d)]\psi(x, y) = 0.$$

Proof. Observe that

$$\begin{aligned} \nabla a_1(x, y) &= (a'(x-y), 0, \dots, 0, -a'(x-y), 0, \dots, 0) \\ \nabla b_1(x, y) &= (b'(x-y), 0, \dots, 0, -b'(x-y), 0, \dots, 0) \end{aligned} \quad (3.7)$$

and similarly for the other gradients. Moreover, from $a^2 + b^2 = 1$ it follows $aa' + bb' = 0$, while $\psi b_1b_2 \cdots b_{d-1}\nabla a_d = 0$ and $\psi b_1b_2 \cdots b_{d-1}a_d^2 = \psi b_1b_2 \cdots b_d$ for the same reason as in claim 3.6. Hence we have

$$\begin{aligned} &[a_1\nabla a_1 + (b_1a_2)\nabla(b_1a_2) + \cdots + (b_1b_2 \cdots b_{d-1}a_d)\nabla(b_1b_2 \cdots b_{d-1}a_d)]\psi \\ &= [a_1\nabla a_1 + (b_1a_2)\nabla(b_1a_2) + \cdots + (b_1b_2 \cdots b_{d-1})\nabla(b_1b_2 \cdots b_{d-1})]\psi. \end{aligned}$$

A ‘‘chain reaction’’ is now generated by the following formula, valid for every $k \geq 1$:

$$\begin{aligned} &(b_1 \cdots b_k a_{k+1})\nabla(b_1 \cdots b_k a_{k+1}) + (b_1 \cdots b_k b_{k+1})\nabla(b_1 \cdots b_k b_{k+1}) \\ &= (b_1 \cdots b_k a_{k+1}^2)\nabla(b_1 \cdots b_k) + (b_1^2 \cdots b_k^2) a_{k+1} \nabla a_{k+1} \\ &\quad + (b_1 \cdots b_k b_{k+1}^2)\nabla(b_1 \cdots b_k) + (b_1^2 \cdots b_k^2) b_{k+1} \nabla b_{k+1} \\ &= (b_1 \cdots b_k)\nabla(b_1 \cdots b_k). \end{aligned}$$

□

It is left to estimate $(|\nabla g_1(x, y)|^2 + |\nabla g_{\xi}(x, y)|^2) \psi(x, y)$. Again we compute directly

$$|\nabla g_1(x, y)|^2 + |\nabla g_{\xi}(x, y)|^2 = 3 \left(|\nabla a_1|^2 + |\nabla(b_1a_2)|^2 + |\nabla(b_1b_2a_3)|^2 \right).$$

Note, however, that because of (3.7) we have $\nabla a_i \cdot \nabla b_j = 0$ and $\nabla b_i \cdot \nabla b_j = 0$ if $i \neq j$. Therefore,

$$\begin{aligned} |\nabla g_1(x, y)|^2 + |\nabla g_{\xi}(x, y)|^2 &= |\nabla a_1|^2 + b_1^2 |\nabla a_2|^2 + a_2^2 |\nabla b_1|^2 + b_2^2 a_3^2 |\nabla b_1|^2 \\ &\quad + b_1^2 a_3^2 |\nabla b_2|^2 + b_1^2 b_2^2 |\nabla a_3|^2. \end{aligned}$$

and, using again the idea of claim (3.6),

$$\begin{aligned} &(|\nabla g_1(x, y)|^2 + |\nabla g_{\xi}(x, y)|^2) \psi^2(x, y) \\ &= \left(|\nabla a_1|^2 + b_1^2 |\nabla a_2|^2 + a_2^2 |\nabla b_1|^2 + b_2^2 |\nabla b_1|^2 + b_1^2 |\nabla b_2|^2 \right) \psi^2(x, y) \\ &= \left(|\nabla a_1|^2 + |\nabla b_1|^2 + b_1^2 \left(|\nabla a_2|^2 + |\nabla b_2|^2 \right) \right) \psi^2(x, y) \\ &\leq \frac{8k^2}{r^2} \psi^2(x, y) = \frac{24k^2}{\alpha^2} \psi^2(x, y). \end{aligned}$$

3.7. Construction for $N = 3$, $d = 3$. In this case, let the variable be $X = (x, y, z) = (x_1, x_2, x_3; y_1, y_2, y_3; z_1, z_2, z_3)$, and define as before

$$a_j(x, y) = a(x_j - y_j) \quad b_1(x, y) = b(x_1 - y_1)$$

for $j = 2, 3$. As in the case $N = 2$, define also

$$\begin{aligned} g_1(x, y) &= a_1(x, y) + b_1(x, y)a_2(x, y) + b_1(x, y)b_2(x, y)a_3(x, y) \\ g_\xi(x, y) &= a_1(x, y) + \xi b_1(x, y)a_2(x, y) + \bar{\xi} b_1(x, y)b_2(x, y)a_3(x, y). \end{aligned}$$

Now comes the definition of the wave-function:

$$\begin{aligned} \psi_{000}(x, y, z) &= 0 \\ \psi_{001}(x, y, z) &= \frac{1}{3}g_1(x, y)\psi(x, y, z) \\ \psi_{010}(x, y, z) &= -\frac{1}{3}g_1(x, z)\psi(x, y, z) \\ \psi_{011}(x, y, z) &= \frac{\sqrt{2}}{3}g_\xi(y, z)\psi(x, y, z) \\ \psi_{100}(x, y, z) &= \frac{1}{3}g_1(y, z)\psi(x, y, z) \\ \psi_{101}(x, y, z) &= -\frac{\sqrt{2}}{3}g_\xi(x, z)\psi(x, y, z) \\ \psi_{110}(x, y, z) &= \frac{\sqrt{2}}{3}g_\xi(x, y)\psi(x, y, z) \\ \psi_{111}(x, y, z) &= 0. \end{aligned}$$

It is quite easy to see that ψ is indeed fermionic. The fact that

$$\sum_{s \in S} |\psi_s(x, y, z)|^2 = \psi^2(x, y, z)$$

is proved exactly in the same way as for $N = 2$, and also the gradient estimates, considering the couples ψ_{001} - ψ_{110} , ψ_{010} - ψ_{101} and ψ_{100} - ψ_{011} . Hence we proved the Proposition 3.19, with $C = 24k^2$ for $k > 1$ arbitrary.

3.8. The case $d = 4$. A very similar construction may be done for $d = 4$. In the case $N = 2$ one simply chooses

$$\begin{aligned} g_1(x, y) &= \frac{1}{\sqrt{2}}(a_1 + ib_1a_2 + b_1b_2a_3 + ib_1b_2b_3a_4) \\ g_2(x, y) &= \frac{1}{\sqrt{2}}(a_1 + ib_1a_2 - b_1b_2a_3 - ib_1b_2b_3a_4). \end{aligned}$$

and

$$\begin{aligned} \psi_{00}(x, y) &= g_1(x, y)\psi(x, y) \\ \psi_{01}(x, y) &= 0 \\ \psi_{10}(x, y) &= 0 \\ \psi_{11}(x, y) &= g_2(x, y)\psi(x, y). \end{aligned}$$

It is easy to verify that $|g_1(x, y)|^2 + |g_2(x, y)|^2 = a_1^2 + b_1^2a_2^2 + b_1^2b_2^2a_3^2 + b_1^2b_2^2b_3^2a_4^2$, and proceeding as before

$$\left(|g_1(x, y)|^2 + |g_2(x, y)|^2 \right) \psi^2(x, y) = \psi^2(x, y).$$

To estimate the derivatives, note that

$$\begin{aligned} 2\Re(\bar{g}_1 \nabla g_1) &= a_1 \nabla a_1 + a_1 \nabla(b_1 b_2 a_3) + b_1 a_2 \nabla(b_1 a_2) + b_1 a_2 \nabla(b_1 b_2 b_3 a_4) \\ &\quad + b_1 b_2 a_3 \nabla a_1 + b_1 b_2 a_3 \nabla b_1 b_2 a_3 + b_1 b_2 b_3 a_4 \nabla(b_1 a_2) \\ &\quad + b_1 b_2 b_3 a_4 \nabla(b_1 b_2 b_3 a_4) \\ 2\Re(\bar{g}_2 \nabla g_2) &= a_1 \nabla a_1 - a_1 \nabla(b_1 b_2 a_3) + b_1 a_2 \nabla(b_1 a_2) - b_1 a_2 \nabla(b_1 b_2 b_3 a_4) \\ &\quad - b_1 b_2 a_3 \nabla a_1 + b_1 b_2 a_3 \nabla b_1 b_2 a_3 - b_1 b_2 b_3 a_4 \nabla(b_1 a_2) \\ &\quad + b_1 b_2 b_3 a_4 \nabla(b_1 b_2 b_3 a_4) \end{aligned}$$

This yields, using Lemma 3.20,

$$\begin{aligned} &(2\Re(\bar{g}_1 \nabla g_1) + 2\Re(\bar{g}_2 \nabla g_2)) \psi \\ &= 2[a_1 \nabla a_1 + b_1 a_2 \nabla(b_1 a_2) + b_1 b_2 a_3 \nabla(b_1 b_2 a_3) + b_1 b_2 b_3 a_4 \nabla(b_1 b_2 b_3 a_4)] \psi \\ &= 0. \end{aligned}$$

Therefore,

$$|\nabla \psi_{00}(x, y)|^2 + |\nabla \psi_{00}(x, y)|^2 = |\nabla \psi|^2 + \left(|\nabla g_1(x, y)|^2 + |\nabla g_2(x, y)|^2 \right) \psi^2$$

and we conclude with the estimate

$$\begin{aligned} &\left(|\nabla g_1|^2 + |\nabla g_2|^2 \right) \psi^2 \\ &= \left(|\nabla a_1|^2 + b_1^2 |\nabla a_2|^2 + a_2^2 |\nabla b_1|^2 + b_2^2 a_3^2 |\nabla b_1|^2 + b_1^2 a_3^2 |\nabla b_2|^2 \right. \\ &\quad \left. + b_1^2 b_2^2 |\nabla a_3|^2 + b_2^2 b_3^2 |\nabla b_1|^2 + b_1^2 b_3^2 |\nabla b_2|^2 + b_1^2 b_2^2 |\nabla b_3|^2 \right) \psi^2 \\ &= \left[|\nabla a_1|^2 + |\nabla b_1|^2 + b_1^2 \left(|\nabla a_2|^2 + |\nabla b_2|^2 \right) \right. \\ &\quad \left. + b_1^2 b_2^2 \left(|\nabla a_3|^2 + |\nabla b_3|^2 \right) \right] \psi^2 \\ &\leq \frac{12k^2}{\alpha^2} \psi^2 = \frac{36k^2}{\alpha^2} \psi^2, \end{aligned}$$

which shows that in this case C can be chosen $36k^2$ for $k > 1$ arbitrary.

For 3 particles it suffices to repeat the construction of Subsection 4.5 in order to obtain Proposition 3.19 with $C = 36k^2$.

3.9. Γ – lim sup-inequality. Finally we get the Γ -lim sup inequality, and thus the entire proof, both in the symmetric and the antisymmetric case.

Bosonic case. We can complete the proof of Theorem 3.3:

Proof. of Th. 3.3 We proved equicoerciveness in Subsection 3.1 and, for all $P \in \Pi(\rho)$, in Subsection 3.2 we already proved the Γ -lim inf inequality. Thus it is left to find a family of bosonic wave-functions ψ_{\hbar} such that $P_{\psi_{\hbar}} \rightharpoonup P$ and

$$\limsup_{\hbar \rightarrow 0} \{T_{\hbar}(\psi_{\hbar}) + V_{ee}(\psi_{\hbar})\} \leq \mathfrak{C}_{\mathcal{S}}(P).$$

Define $\varepsilon(\hbar) = \sqrt{\hbar}$, and set

$$\begin{aligned} \psi_{\hbar}(x_1, s_1, \dots, x_N, s_N) &= \begin{cases} \psi_{\varepsilon(\hbar)}(x_1, \dots, x_N) & \text{if } s_1 = \dots = s_N = 0 \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{pmatrix} \psi_{\varepsilon(\hbar)}(x_1, \dots, x_N) \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \end{aligned}$$

where ψ_{ε} are given by Proposition 3.7.

These wave-functions are clearly bosonic, and satisfy $|\boldsymbol{\psi}_{\hbar}(X)|^2 = \psi_{\varepsilon(\hbar)}^2(X)$, $|\nabla \boldsymbol{\psi}_{\hbar}(X)|^2 = |\nabla \psi_{\varepsilon(\hbar)}(X)|^2$, so that

$$\begin{aligned} T_{\hbar}(\boldsymbol{\psi}_{\hbar}) &= \frac{\hbar^2}{2} \int |\nabla \psi_{\varepsilon(\hbar)}(X)|^2 dX \leq \frac{N\hbar^2}{2} \left(\|\rho\|_{H^2}^2 + \frac{K}{4\varepsilon(\hbar)^2} \right) \\ &= \frac{N\hbar^2}{2} \|\rho\|_{H^2}^2 + \frac{KN\hbar}{8}, \end{aligned}$$

while

$$V_{ee}(\boldsymbol{\psi}_{\hbar}) = \int c(X) \psi_{\varepsilon(\hbar)}^2(X) dX = \mathfrak{C}_{\mathcal{S}}(P_{\varepsilon(\hbar)}).$$

Now the thesis follows from Proposition 3.7, since

$$\limsup_{\hbar \rightarrow 0} \{T_{\hbar}(\boldsymbol{\psi}_{\hbar}) + V_{ee}(\boldsymbol{\psi}_{\hbar})\} = \limsup_{\hbar \rightarrow 0} V_{ee}(\boldsymbol{\psi}_{\hbar}) \leq \mathfrak{C}_{\mathcal{S}}(P).$$

□

Fermionic case. We can complete the proof of Theorem 3.4:

Proof. of Th. 3.4 We proved equicoerciveness in Subsection 3.1 and, for all $P \in \Pi(\rho)$, in Subsection 3.2, we already proved the Γ -lim inf inequality. Thus it is left to find a family of fermionic wave-functions $\boldsymbol{\psi}_{\hbar}$ such that $P_{\boldsymbol{\psi}_{\hbar}} \rightarrow P$ and

$$\limsup_{\hbar \rightarrow 0} \{T_{\hbar}(\boldsymbol{\psi}_{\hbar}) + V_{ee}(\boldsymbol{\psi}_{\hbar})\} \leq \mathfrak{C}_{\mathcal{S}}(P).$$

Consider a sequence of functions $\{\psi_{\varepsilon}\}$ as in the thesis of Proposition 3.7. Recall that ψ_{ε} is supported outside $D_{\alpha(\varepsilon)}$, where $\alpha(\varepsilon) \searrow 0$ as $\varepsilon \rightarrow 0$ – hence there exists α^{-1} in a right neighbourhood of 0. We may then consider a corresponding family of wave-functions $\{\psi_{\varepsilon}\}$ given by Proposition 3.19. Define

$$\varepsilon(\hbar) = \max \left\{ \alpha^{-1}(\sqrt{\hbar}), \sqrt{\hbar} \right\},$$

and observe that $\varepsilon(\hbar) \rightarrow 0$ as $\hbar \rightarrow 0$. We take $\boldsymbol{\psi}_{\hbar} = \psi_{\varepsilon(\hbar)}$ as a recovery sequence.

It is easy to estimate the kinetic energy:

$$\begin{aligned} T_{\hbar}(\boldsymbol{\psi}_{\varepsilon(\hbar)}) &= \frac{\hbar^2}{2} \int |\nabla \psi_{\varepsilon(\hbar)}(X)|^2 dX \\ &\leq \frac{\hbar^2}{2} \left\{ \int |\nabla \psi_{\varepsilon(\hbar)}(X)|^2 dX + \frac{C}{\alpha^2(\varepsilon(\hbar))} \int \psi_{\varepsilon(\hbar)}^2(X) dX \right\} \\ &\leq \frac{\hbar^2}{2} \left\{ N \|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\varepsilon^2(\hbar)} + \frac{C}{\hbar} \right\} \\ &\leq \frac{\hbar^2}{2} \left\{ N \|\sqrt{\rho}\|_{H^1}^2 + \frac{K}{4\hbar} + \frac{C}{\hbar} \right\} \end{aligned}$$

which tends to 0 as $\hbar \rightarrow 0$. On the other hand, with the notation of Proposition 3.7,

$$V_{ee}(\boldsymbol{\psi}_{\hbar}) = \int c(X) \psi_{\varepsilon(\hbar)}^2(X) dX = \mathfrak{C}_{\mathcal{S}}(P_{\varepsilon(\hbar)}).$$

Now the thesis follows, since

$$\limsup_{\hbar \rightarrow 0} \{T_{\hbar}(\boldsymbol{\psi}_{\hbar}) + V_{ee}(\boldsymbol{\psi}_{\hbar})\} = \limsup_{\hbar \rightarrow 0} V_{ee}(\boldsymbol{\psi}_{\hbar}) \leq \mathfrak{C}_{\mathcal{S}}(P).$$

□

3.10. Conclusions. The Γ -convergence result of the previous section allow us to prove Theorems 1.1 and 1.2:

Proof. of Th.1.1. Thanks to the formulations (3.1) and (3.3) the thesis follows from Theorem 2.3, applied to the functionals \mathcal{F}_h^S and \mathcal{C}_S . We proved the Γ -convergence and equicoercivity in Th. 3.3. \square

Proof. of Th.1.2 Thanks to the formulations (3.2) and (3.3) the thesis follows from Theorem 2.3, applied to the functionals \mathcal{F}_h^A and \mathcal{C}_S . We proved the Γ -convergence and equicoercivity in Th. 3.4. \square

ACKNOWLEDGEMENT

The research of the second author is part of the project 2010A2TFX2 *Calcolo delle Variazioni* funded by the Italian Ministry of Research and is partially financed by the “Fondi di ricerca di ateneo” of the University of Pisa and by the INDAM-GNAMPA.

ADDENDUM

After this paper was accepted, two new papers appeared [24, 12]. Both papers contain a proof of the convergence for general N . The approach is different.

REFERENCES

- [1] L. AMBROSIO, N. GIGLI and G. SAVARÉ, *Gradient Flows in Metric Spaces and in the Space of Probability Measures*, Birkhäuser Verlag, Basel – Boston – Berlin
- [2] M. Beiglböck, C. Léonard, W. Schachermayer. A general duality theorem for the Monge–Kantorovich transport problem. *Studia Math.*, 209:2, 2012.
- [3] U. Bindini. Γ -convergence and Optimal Transportation in Density Functional Theory. Master Thesis, University of Pisa, November 2016. Available at <https://etd.adm.unipi.it/t/etd-09242016-113718/>
- [4] A. Braides. *Gamma-convergence for Beginners*, volume 22. Oxford University Press Oxford, 2002.
- [5] G. Buttazzo, L. De Pascale, and P. Gori-Giorgi. Optimal-transport formulation of electronic density-functional theory. *Physical Review A*, 85(6):062502, 2012.
- [6] G. Buttazzo, T. Champion, and L. De Pascale, Continuity and estimates for multimarginal optimal transportation problems with singular costs, *Applied Mathematics & Optimization*, on-line first, 2017.
- [7] G. Carlier. On a class of multidimensional optimal transportation problems. *J. Convex Anal.*, 10(2):517–530, 2003.
- [8] G. Carlier, B. Nazaret. Optimal transportation for the determinant. *ESAIM Control Optim. Calc. Var.*, 14(04):678–698, 2008.
- [9] M. Colombo, L. De Pascale, S. Di Marino. Multimarginal optimal transport maps for 1-dimensional repulsive costs. *Canad. J. Math*, 67(2):350–368, 2015.
- [10] M. Colombo, S. Di Marino. Equality between Monge and Kantorovich multimarginal problems with Coulomb cost. *Ann. Mat. Pura Appl.*, 194 (2):307–320, 2015.
- [11] C. Cotar, G. Friesecke, C. Klüppelberg. Density functional theory and optimal transportation with Coulomb cost. *Comm. Pure Appl. Math.*, 66(4):548–599, 2013.
- [12] C. Cotar, G. Friesecke, C. Klüppelberg. Smoothing of transport plans with fixed marginals and rigorous semiclassical limit of the Hohenberg-Kohn functional. Preprint: [arXiv:1706.05676v1](https://arxiv.org/abs/1706.05676v1) [math-ph]
- [13] G. Dal Maso. *An introduction to Γ -convergence*. Springer, 1993.
- [14] L. De Pascale, Optimal transport with Coulomb cost. Approximation and duality. *ESAIM Math. Model. Numer. Anal.*, 49(6):1643–1657, 2015.
- [15] G. Friesecke, C.B. Mendl, B. Pass, C. Cotar, C. Klüppelberg. N-density representability and the optimal transport limit of the Hohenberg-Kohn functional. *The Journal of chemical physics*, 139(16):164109, 2013.
- [16] W. Gangbo, A. Swiech. Optimal maps for the multidimensional Monge-Kantorovich problem. *Comm. Pure Appl. Math.*, 51(1):23–45, 1998.
- [17] N. Ghoussoub, A. Moameni. A self-dual polar factorization for vector fields. *Comm. Pure Appl. Math.*, 66 (6) (2013), 905–933.
- [18] P. Gori-Giorgi, M. Seidl. Density functional theory for strongly-interacting electrons: perspectives for physics and chemistry. *Physical Chemistry Chemical Physics*, 12(43):14405–14419, 2010.

- [19] P. Gori-Giorgi, M. Seidl, G. Vignale. Density-functional theory for strongly interacting electrons. *Physical review letters*, 103(16):166402, 2009.
- [20] H. Heinich. Problème de Monge pour n probabilités. *C. R. Math. Acad. Sci. Paris*, 334(9):793–795, 2002.
- [21] P. Hohenberg, W. Kohn. Inhomogeneous electron gas. *Physical review*, 136(3B):B864, 1964.
- [22] H.G. Kellerer. Duality theorems for marginal problems. *Probab. Theory Related Fields*, 67(4):399–432, 1984.
- [23] W. Kohn, L.J. Sham. Self-consistent equations including exchange and correlation effects. *Physical Review*, 140(4A):A1133, 1965.
- [24] M. Lewin. Semi-classical limit of the Levy-Lieb functional in Density Functional Theory. Preprint, [arXiv:1706.02199](https://arxiv.org/abs/1706.02199) [math-ph]
- [25] M. Levy. Universal variational functionals of electron densities, first-order density matrices, and natural spin-orbitals and solution of the v -representability problem. *Proceedings of the National Academy of Sciences*, 76(12):6062–6065, 1979.
- [26] E.H. Lieb. Density functionals for coulomb systems. *International journal of quantum chemistry*, 24(3):243–277, 1983.
- [27] C.B. Mendl, L. Lin. Kantorovich dual solution for strictly correlated electrons in atoms and molecules. *Physical Review B*, 87(12):125106, 2013.
- [28] B. Pass. Uniqueness and monge solutions in the multimarginal optimal transportation problem. *SIAM J. Math. Anal.*, 43(6):2758–2775, 2011.
- [29] B. Pass. On the local structure of optimal measures in the multi-marginal optimal transportation problem. *Calc. Var. Partial Differential Equations*, 43(3-4):529–536, 2012.
- [30] S.T. Rachev, L. Rüschendorf. *Mass transportation problems. Vol. I: Theory*. Probability and its Applications (New York). Springer-Verlag, New York, 1998.
- [31] M. Seidl. Strong-interaction limit of density-functional theory. *Physical Review A*, 60(6):4387, 1999.
- [32] M. Seidl, P. Gori-Giorgi, A. Savin. Strictly correlated electrons in density-functional theory: A general formulation with applications to spherical densities. *Physical Review A*, 75(4):042511, 2007.
- [33] M. Seidl, J.P. Perdew, M. Levy. Strictly correlated electrons in density-functional theory. *Physical Review A*, 59(1):51, 1999.

Ugo Bindini: Scuola Normale Superiore,
Piazza dei Cavalieri 5, 56127 Pisa - ITALY
ugo.bindini@sns.it

Luigi De Pascale: Dipartimento di Matematica e Informatica, Università di Firenze
Viale Morgagni 67/A , 50134 Firenze - ITALY
luigi.depascale@unifi.it
<http://web.math.unifi.it/~depascal/>