

UNIQUENESS OF SOLUTIONS IN MEAN FIELD GAMES WITH SEVERAL POPULATIONS AND NEUMANN CONDITIONS

MARTINO BARDI AND MARCO CIRANT

ABSTRACT. We study the uniqueness of solutions to systems of PDEs arising in Mean Field Games with several populations of agents and Neumann boundary conditions. The main assumption requires the smallness of some data, e.g., the length of the time horizon. This complements the existence results for MFG models of segregation phenomena introduced by the authors and Achdou. An application to robust Mean Field Games is also given.

CONTENTS

1.	Introduction	1
2.	The uniqueness theorem	3
2.1.	The main result	3
2.2.	Examples and remarks	7
3.	Special cases and applications	9
3.1.	Well-posedness of segregation models	10
3.2.	Well-posedness of robust Mean Field Games	11
4.	Appendix: a Comparison Principle	11
	References	12

1. INTRODUCTION

The systems of partial differential equations associated to finite-horizon Mean Field Games (briefly, MFGs) with N populations of agents have the form

$$(1.1) \quad \begin{cases} -\partial_t v_k - \Delta v_k + H_k(x, Dv_k) = F_k(x, m(t, \cdot)), & \text{in } (0, T) \times \Omega, \\ \partial_t m_k - \Delta m_k - \operatorname{div}(D_p H_k(x, Dv_k) m_k) = 0 & \text{in } (0, T) \times \Omega, \\ v_k(T, x) = G_k(x, m(T, \cdot)), \quad m_k(0, x) = m_{0,k}(x) & \text{in } \Omega, \quad k = 1, \dots, N, \end{cases}$$

where the unknown m is a vector of probability densities on Ω , F_k and G_k are function of this vector and represent the running and terminal costs of a representative agent of the k -population, and v_k is the value function of this agent. The first N equations are parabolic of Hamilton-Jacobi-Bellman type and backward in time with a terminal condition, the second N equations are parabolic of Kolmogorov-Fokker-Planck type and forward in time with an initial condition. If the state space $\Omega \subseteq \mathbb{R}^d$ is not all \mathbb{R}^d , boundary conditions must also be imposed. In most of the

Date: April 30, 2018.

Key words and phrases. Mean Field Games, multi-populations, uniqueness, Neumann boundary conditions, robust Mean Field Games.

The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). They are partially supported by the research projects "Mean-Field Games and Nonlinear PDEs" of the University of Padova and "Nonlinear Partial Differential Equations: Asymptotic Problems and Mean-Field Games" of the Fondazione CaRiPaRo.

theory of MFGs they are periodic, which are the easiest to handle, here we will consider instead Neumann conditions, i.e.,

$$(1.2) \quad \partial_n v_k = 0, \quad \partial_n m_k + m_k D_p H_k(x, Dv_k) \cdot n = 0 \quad \text{on } (0, T) \times \partial\Omega.$$

There is a large literature on the existence of solutions for these equations, especially in the case of a single population $N = 1$, beginning with the pioneering papers of Lasry and Lions [27, 28, 29] and Huang, Caines and Malhame [23, 24, 25], see the lecture notes [10, 22], the books [19, 9], the survey [20], and the references therein. Systems with several populations, $N > 1$, were treated with Neumann conditions in [12, 15] for the stationary case and in [1] in the evolutive case, with periodic conditions in [4, 11].

Uniqueness of solutions is a much more delicate issue. For one population Lasry and Lions [27, 28, 29] discovered a monotonicity condition on the costs F and G that together with the convexity in p of the Hamiltonian $H(x, p)$ implies the uniqueness of classical solutions. It reads

$$(1.3) \quad \int_{\mathbb{R}} (F(x, \mu) - F(x, \nu)) d(\mu - \nu)(x) > 0, \quad \text{if } \mu \neq \nu$$

and it means that a representative agent prefers the regions of the state space that are less crowded. This is a restrictive condition that is satisfied in some models and not in others. When it fails, non-uniqueness may arise: this was first observed in the stationary case by Lasry and Lions [29] and other counterexamples were shown by [21, 3, 6, 18]. The need of a condition such as (1.3) for having uniqueness for finite-horizon MFGs was discussed at length in [31], and some explicit examples of non-uniqueness appeared very recently in [8], [14], and in [5] that presents also a probabilistic proof and references on other examples obtained by the probabilistic approach.

For multi-population problems, $N > 1$, there are extensions of the monotonicity condition (1.3) in [12, 5] and they are even more restrictive: they impose not only aversion to crowd within each population, but also that the costs due to this effect dominate the costs due to the interactions with the other populations. This is not the case in the multi-population models of segregation in urban settlements proposed in [1] following the ideas of the Nobel Prize Thomas Schelling [34]. There the interactions between two different populations are the main cause of the dynamics, and in fact examples of multiple solutions were shown in [1] and [15] for the stationary case and in [5] for the evolutive one. Therefore a different criterion giving uniqueness in some cases is particularly desirable when $N > 1$.

A second regime for uniqueness was introduced in a lecture of P.L. Lions on January 9th, 2009 [31]: it occurs if the length T of the time horizon is short enough. To our knowledge Lions' original argument did not appear in print. For finite state MFGs, uniqueness for short time was proved by Gomes, Mohr, and Souza [16] as part of their study of the large population limit. For continuous state, an existence and uniqueness result under a "small data" condition was given in [25] for Linear-Quadratic-Gaussian MFGs using a contraction mapping argument to solve the associated system of Riccati differential equations, and similar arguments were used for different classes of linear-quadratic problems in [36, 32]. The well-posedness when $H(x, Dv) - F(x, m)$ is replaced by $\varepsilon \mathcal{H}(x, Dv, m)$ with ε small is studied in [2], and another result for small Hamiltonian is in [35] for nonconvex H .

Very recently the first author and Fischer [5] revived Lions' argument to show that the smoothness of the Hamiltonian is the crucial property to have small-time uniqueness without monotonicity of the costs and convexity of H , and gave an example of non-uniqueness for all $T > 0$ and $H(x, p) = |p|$. The uniqueness theorem for small data in [5] holds for $N = 1$ and $\Omega = \mathbb{R}^d$ with conditions on the behaviour of the solutions at infinity.

In the present paper we focus instead on $N \geq 1$ and Neumann boundary conditions, which is the setting of the MFG models of segregation in [1]. The new difficulties arise from the boundary conditions, that require different methods for some estimates, especially on the L^∞ norm of the densities m_k . Our first uniqueness result assumes a suitable smoothness of the Hamiltonians H_k , but neither convexity nor growth conditions, and that the costs F_k, G_k are Lipschitz in L^2 with respect to the measure m , with no monotonicity. The smallness condition on the data depends on the range of the spacial gradient of the solutions v_k , unless $D_p H_k$ are bounded and globally

Lipschitz for all k . Then we complement such result with some a priori gradient estimates on v_k , under an additional quadratic growth condition on H_k and some more regularity of the costs, and get a $\bar{T} > 0$ depending only on the data such that there is uniqueness for all horizons $T \leq \bar{T}$. Finally, we give sufficient conditions ensuring both existence and uniqueness for the system (1.1) with the boundary conditions (1.2), as well as for some robust MFGs considered in [7, 32], which are interesting examples with nonconvex Hamiltonian.

We mention that in the stationary case, uniqueness up to (space) translation may hold without (1.3) in force. A special class of MFG on \mathbb{R}^d enjoying such a feature has been identified in [13].

The paper is organised as follows. Section 2 contains the main result about uniqueness for small data, possibly depending on gradient bounds on the solutions. Section 3 gives further sufficient conditions depending only on the data for uniqueness and existence of solutions. The Appendix 4 recalls a comparison principle for HJB equations with Neumann conditions.

2. THE UNIQUENESS THEOREM

Consider the MFG system for N populations

$$(2.1) \quad \begin{cases} -\partial_t v_k - \Delta v_k + H_k(x, Dv_k) = F_k(x, m(t, \cdot)), & \text{in } (0, T) \times \Omega, \\ \partial_t m_k - \Delta m_k - \operatorname{div}(D_p H_k(x, Dv_k) m_k) = 0 & \text{in } (0, T) \times \Omega, \\ \partial_n v_k = 0, \partial_n m_k + m_k D_p H_k(x, Dv_k) \cdot n = 0 & \text{on } (0, T) \times \partial\Omega, \\ v_k(T, x) = G_k(x, m(T, \cdot)), m_k(0, x) = m_{0,k}(x) & \text{in } \Omega \end{cases}$$

where $k = 1, \dots, N$, Dv_k denotes the gradient of the k -th component v_k of the unknown v with respect to the space variables, Δ is the Laplacian with respect to the space variables x , $D_p H_k$ is the gradient of the Hamiltonian of the k -th population with respect to the moment variable, $\Omega \subseteq \mathbb{R}^d$ is a bounded open set with boundary $\partial\Omega$ of class $C^{2,\beta}$ for some $\beta > 0$, and $n(x)$ is its exterior normal at x . The components m_k of the unknown vector m are bounded densities of probability measures on Ω , i.e., m lives in

$$\mathcal{P}_N(\Omega) := \left\{ \mu = (\mu_1, \dots, \mu_N) \in L^\infty(\Omega)^N : \mu_k \geq 0, \int_\Omega \mu_k(x) dx = 1 \right\}.$$

F and G represent, respectively, the running and terminal cost of the MFG

$$F : \bar{\Omega} \times \mathcal{P}_N(\Omega) \rightarrow \mathbb{R}^N, \quad G : \bar{\Omega} \times \mathcal{P}_N(\Omega) \rightarrow \mathbb{R}^N.$$

By classical solutions we will mean functions of (t, x) of class C^1 in t and C^2 in x in $[0, T] \times \bar{\Omega}$.

2.1. The main result. Our main assumptions are the smoothness of the Hamiltonians and a Lipschitz continuity of the costs in the norm $\|\cdot\|_2$ of $L^2(\Omega)^N$ that we state next. We consider $H_k : \bar{\Omega} \times \mathbb{R}^d \rightarrow \mathbb{R}$ continuous and satisfying

$$(2.2) \quad D_p H_k(x, p) \text{ is continuous and locally Lipschitz in } p \text{ uniformly in } x \in \bar{\Omega}.$$

We will assume F, G satisfy, for all μ, ν ,

$$(2.3) \quad \|F(\cdot, \mu) - F(\cdot, \nu)\|_2^2 \leq L_F \|\mu - \nu\|_2^2,$$

$$(2.4) \quad \|DG(\cdot, \mu) - DG(\cdot, \nu)\|_2^2 \leq L_G \|\mu - \nu\|_2^2,$$

Theorem 2.1. *Assume (2.2), (2.3), (2.4), $m_0 \in \mathcal{P}_N(\Omega)$, and $(\bar{v}, \bar{m}), (\bar{v}, \bar{m})$ are two classical solutions of (2.1). Denote*

$$(2.5) \quad \mathcal{C} := \operatorname{co}\{D\bar{v}(t, x), D\bar{v}(t, x) : (t, x) \in (0, T) \times \Omega\},$$

$$(2.6) \quad C_H := \max_{k=1, \dots, N} \sup_{x \in \Omega, p \in \mathcal{C}} |D_p H_k(x, p)|,$$

$$(2.6) \quad \bar{C}_H := \max_{k=1, \dots, N} \sup_{x \in \Omega, p, q \in \mathcal{C}} \frac{|D_p H_k(x, p) - D_p H_k(x, q)|}{|p - q|}$$

Then there exists a function Ψ of $T, L_F, L_G, C_H, \bar{C}_H, N$ and $\max_k \|m_{0,k}\|_\infty$ (depending also on Ω), such that the inequality $\Psi < 1$ implies $\tilde{v}(t, \cdot) = \bar{v}(t, \cdot)$ and $\tilde{m}(t, \cdot) = \bar{m}(t, \cdot)$ for all $t \in [0, T]$, and $\Psi < 1$ holds if either T , or \bar{C}_H , or the pair L_F, L_G is small enough.

For the the proof we need two auxiliary results.

Proposition 2.2. *There are constants $r > 1$ and $C > 0$ depending only on d and Ω such that*

$$(2.7) \quad \|m_k\|_{L^\infty((0,T)\times\Omega)} \leq C[1 + \|m_{0,k}\|_\infty + (1+T)\|D_p H_k(\cdot, Dv_k)\|_{L^\infty((0,T)\times\Omega)}]^r, \quad k = 1, \dots, N.$$

Proof. Step 1. We aim at proving that for any $q \in [1, (d+2)/(d+1))$ there exists a constant \bar{C} depending only on d, q and Ω such that any positive classical solution φ of the backward heat equation

$$\begin{cases} -\partial_t \varphi - \Delta \varphi = 0 & \text{on } (0, t) \times \Omega \\ \partial_n \varphi = 0 & \text{on } (0, t) \times \partial\Omega \\ \int_\Omega \varphi(t, x) dx = 1, \end{cases}$$

satisfies

$$\|\nabla \varphi\|_{L^q((0,t)\times\Omega)} \leq \bar{C}(1+t)^{1/q}.$$

We follow the strategy presented in [19, Section 5]. Note first that $\int_\Omega \varphi(s, x) dx = 1$ for all $s \in (0, t)$, by integrating by parts the equation and using the boundary conditions. We proceed in the case $d \geq 3$; if $d = 1$ or $d = 2$, one argues in a similar way (see the discussion below). Let $\alpha \in (0, 1)$ to be chosen later; multiplying the equation by $\alpha \varphi^{\alpha-1}$ and integrating by parts yield for all $s \in (0, t)$

$$\int_\Omega |\nabla \varphi^{\alpha/2}(s, x)|^2 dx = \frac{\alpha}{4(\alpha-1)} \partial_t \int_\Omega \varphi^\alpha(s, x) dx.$$

Integrating in time and using the fact that $\int_\Omega \varphi(s, x) dx = 1$ give

$$(2.8) \quad \int_0^t \int_\Omega |\nabla \varphi^{\alpha/2}|^2 dx ds = \frac{\alpha}{4(1-\alpha)} \int_\Omega \varphi^\alpha(0, x) dx - \frac{\alpha}{4(1-\alpha)} \int_\Omega \varphi^\alpha(t, x) dx \leq c_1,$$

where c_1 depends on d and Ω (the positive constants c_2, c_3, \dots used in the sequel will have the same dependance).

We now exploit the continuous embedding of $W^{1,2}(\Omega)$ into $L^{\frac{2d}{d-2}}(\Omega)$; the adaption of this proof to the cases $d = 1, 2$ is straightforward, as the injection of $W^{1,2}(\Omega)$ is into $L^p(\Omega)$ for all $p \geq 1$. Hence, for all $s \in (0, t)$, by Hölder and Sobolev inequalities

$$\begin{aligned} \int_\Omega \varphi^{\alpha+\frac{2}{d}}(s, x) dx &\leq \left(\int_\Omega \varphi(s, x) dx \right)^{\frac{2}{d}} \left(\int_\Omega \varphi^{\frac{\alpha}{2} \frac{2d}{d-2}}(s, x) dx \right)^{\frac{d-2}{d}} \\ &\leq c_2 \left(\int_\Omega |\nabla \varphi^{\alpha/2}|^2 dx + \int_\Omega \varphi^\alpha dx \right) \leq c_2 \left(\int_\Omega |\nabla \varphi^{\alpha/2}|^2 dx + 1 + |\Omega| \right), \end{aligned}$$

so

$$(2.9) \quad \int_0^t \int_\Omega \varphi^{\alpha+\frac{2}{d}} dx ds \leq c_3 \left(\int_0^t \int_\Omega |\nabla \varphi^{\alpha/2}|^2 dx ds + t \right).$$

Finally, since $q < (d+2)/(d+1)$, we may choose $\alpha \in (0, 1)$ such that

$$q \frac{2-\alpha}{2-q} = \alpha + \frac{2}{d},$$

and therefore, by the identity $\nabla \varphi^{\alpha/2} = \frac{\alpha}{2} \varphi^{\frac{\alpha-2}{2}} \nabla \varphi$ and Young's inequality

$$\begin{aligned} \int_0^t \int_\Omega |\nabla \varphi|^q dx ds &= \left(\frac{2}{\alpha} \right)^q \int_0^t \int_\Omega |\nabla \varphi^{\alpha/2}|^q \varphi^{q \frac{2-\alpha}{2}} dx ds \leq \\ &c_4 \left(\int_0^t \int_\Omega |\nabla \varphi^{\alpha/2}|^2 dx ds + \int_0^t \int_\Omega \varphi^{q \frac{2-\alpha}{2}} dx ds \right) \leq c_4(c_1 + c_3(c_1 + t)), \end{aligned}$$

in view of (2.8) and (2.9), and the desired estimate follows.

Step 2. Fix $t \in (0, T)$ and $1 < q < (d+2)/(d+1)$. Let φ_0 be any non-negative smooth function on Ω such that $\partial_n \varphi_0 = 0$ on $\partial\Omega$ and $\int_\Omega \varphi_0(x) dx = 1$. Let φ be the solution of the backward heat equation

$$\begin{cases} -\partial_t \varphi - \Delta \varphi = 0 & \text{on } (0, t) \times \Omega \\ \partial_n \varphi = 0 & \text{on } (0, t) \times \partial\Omega \\ \varphi(t, x) = \varphi_0(x) & \text{on } \Omega. \end{cases}$$

Note that φ is positive on $(0, t) \times \Omega$ by the strong maximum principle. Multiply the KFP equation in (2.1), integrate by parts and use the boundary conditions for m_k to get

$$\int_0^t \int_\Omega \partial_t m_k \varphi + \nabla m_k \cdot \nabla \varphi + D_p H_k(x, Dv_k) \cdot \nabla \varphi m_k dx ds = 0.$$

Integrating again by parts (in space-time) yields

$$\int_\Omega m_k(t, x) \varphi_0(x) = \int_\Omega m_k(0, x) \varphi_0(x) - \int_0^t \int_\Omega D_p H_k(x, Dv_k) \cdot \nabla \varphi m_k dx ds,$$

using the equation and the boundary condition for φ . Hence,

$$\begin{aligned} \int_\Omega m_k(t, x) \varphi_0(x) &\leq \|m_{k,0}\|_\infty + \|D_p H_k(\cdot, Dv_k)\|_{L^\infty((0,t) \times \Omega)} \int_0^t \int_\Omega |\nabla \varphi| |m_k| dx ds, \\ &\leq \|m_{k,0}\|_\infty + \bar{C}(1+t)^{1/q} \|D_p H_k(\cdot, Dv_k)\|_{L^\infty((0,t) \times \Omega)} \|m_k\|_{L^{q'}((0,t) \times \Omega)} \end{aligned}$$

by Step 1. By the arbitrariness of φ_0 , one obtains

$$\|m_k(t, \cdot)\|_\infty \leq \|m_{k,0}\|_\infty + \bar{C}(1+t)^{1/q} \|D_p H_k(\cdot, Dv_k)\|_{L^\infty((0,t) \times \Omega)} \|m_k\|_{L^{q'}((0,t) \times \Omega)},$$

and since

$$\|m_k\|_{L^{q'}((0,t) \times \Omega)} \leq \left(\int_0^t \|m_k(s, \cdot)\|_\infty^{q'-1} \int_\Omega m_k(s, x) dx ds \right)^{1/q'} \leq \|m_k\|_{L^\infty((0,t) \times \Omega)}^{1/q} t^{1/q'},$$

we have

$$\|m_k(t, \cdot)\|_\infty \leq \|m_{k,0}\|_\infty + \bar{C}(1+t) \|D_p H_k(\cdot, Dv_k)\|_{L^\infty((0,t) \times \Omega)} \|m_k\|_{L^\infty((0,t) \times \Omega)}^{1/q}.$$

Passing to the supremum on $t \in (0, T)$, we conclude (r in the statement can be chosen to be q'). \square

Lemma 2.3 (A mean-value theorem). *Let $\mathcal{K} \subseteq \mathbb{R}^d$, $f : \bar{\Omega} \times \mathcal{K} \rightarrow \mathbb{R}^d$ be continuous and Lipschitz continuous in the second entry with constant L , uniformly in the first. Then there exists a measurable matrix-valued function $M(\cdot, \cdot, \cdot)$ such that*

$$(2.10) \quad f(x, p) - f(x, q) = M(x, p, q)(p - q), \quad |M(x, p, q)| \leq L, \quad \forall x \in \bar{\Omega}, p, q \in \mathcal{K}.$$

Proof. Mollify f in the variables p and get a sequence f_n converging to f locally uniformly and with Jacobian matrix satisfying $\|D_p f_n\|_\infty \leq L$. Since f_n is C^1 in p the standard mean-value theorem gives

$$(2.11) \quad f_n(x, p) - f_n(x, q) = \int_0^1 Df_n(x, q + s(p - q))(p - q) ds =: M_n(x, p, q)(p - q),$$

and $|M_n| \leq L$. We define the matrix M componentwise by setting

$$M(x, p, q)_{ij} := \liminf_n M_n(x, p, q)_{ij}, \quad i, j = 1, \dots, d,$$

so that it is measurable in (x, p, q) and satisfies $|M(x, p, q)| \leq L$. Now we take the \liminf_n in the i -th component of the identity (2.11) and get the i -th component of the desired identity (2.10). \square

Proof of Thm. 2.1. Step 1. First observe that, by the regularity of the solutions, $C_H < +\infty$ and $\bar{C}_H < +\infty$. We set

$$v := \tilde{v} - \bar{v}, \quad m := \tilde{m} - \bar{m}, \quad B_k(t, x) := \int_0^1 D_p H_k(x, D\bar{v}(t, x) + s(D\tilde{v} - D\bar{v})(t, x)) ds$$

and observe that $|B_k| \leq C_H$ for all k and v_k satisfies

$$(2.12) \quad \begin{cases} -\partial_t v_k + B_k(t, x) \cdot Dv_k = \Delta v_k + F_k(x, \tilde{m}(t)) - F_k(x, \bar{m}(t)) & \text{in } (0, T) \times \Omega \\ \partial_n v_k = 0 & \text{on } (0, T) \times \partial\Omega, \quad v_k(T, x) = G_k(x, \tilde{m}(T)) - G_k(x, \bar{m}(T)). \end{cases}$$

Step 2. By the divergence theorem and the boundary conditions we compute

$$\begin{aligned} - \int_t^T \int_{\Omega} \partial_t v_k \Delta v_k ds &= \int_t^T \frac{d}{dt} \int_{\Omega} \frac{|Dv_k|^2}{2} dx ds - \int_t^T \int_{\partial\Omega} \partial_t v_k Dv_k \cdot n d\sigma \\ &= \frac{1}{2} \|Dv_k(T, \cdot)\|_2^2 - \frac{1}{2} \|Dv_k(t, \cdot)\|_2^2. \end{aligned}$$

Now we set

$$\bar{F}(t, x) := F(x, \bar{m}) - F(x, \tilde{m}), \quad \bar{G}(t, x) := G(x, \bar{m}) - G(x, \tilde{m}),$$

multiply the PDE in (2.12) by Δv_k , integrate, use the terminal condition in (2.12) and estimate

$$\begin{aligned} \frac{1}{2} \|Dv_k(t, \cdot)\|_2^2 + \int_t^T \|\Delta v_k(s, \cdot)\|_2^2 &\leq \frac{1}{2} \|D\bar{G}(T, \cdot)\|_2^2 + \\ \|B_k\|_{\infty} \int_t^T \left(\frac{1}{2\varepsilon} \|Dv_k(s, \cdot)\|_2^2 + \frac{\varepsilon}{2} \|\Delta v_k(s, \cdot)\|_2^2 \right) ds &+ \int_t^T \left(\frac{1}{2\varepsilon} \|\bar{F}(s, \cdot)\|_2^2 + \frac{\varepsilon}{2} \|\Delta v_k(s, \cdot)\|_2^2 \right) ds. \end{aligned}$$

Next we choose ε such that $1 = (\|B_k\|_{\infty} + 1)\varepsilon/2$ and use the assumptions (2.4) and (2.3) to get

$$\|Dv_k(t, \cdot)\|_2^2 \leq L_G \|m(T, \cdot)\|_2^2 + \int_t^T \frac{L_F}{\varepsilon} \|m(s, \cdot)\|_2^2 ds + \frac{\|B_k\|_{\infty}}{\varepsilon} \int_t^T \|Dv_k(s, \cdot)\|_2^2 ds.$$

Then Gronwall inequality gives, for $c_o := (\|B_k\|_{\infty} + 1)/2 = 1/\varepsilon$ and for all $0 \leq t \leq T$,

$$(2.13) \quad \|Dv_k(t, \cdot)\|_2^2 \leq \left(L_G \|m(T, \cdot)\|_2^2 + c_o L_F \int_t^T \|m(s, \cdot)\|_2^2 ds \right) e^{c_o \|B_k\|_{\infty} T}.$$

Step 3. In order to write a PDE solved by m we apply Lemma 2.3 to $D_p H_k : \bar{\Omega} \times \mathcal{C} \rightarrow \mathbb{R}^d$, which is Lipschitz in p by the assumption in (2.6), and get a matrix M_k such that

$$D_p H_k(x, D\bar{v}_k) - D_p H_k(x, D\check{v}_k) = M_k(x, D\bar{v}_k, D\check{v}_k)(D\bar{v}_k - D\check{v}_k),$$

with $|M_k| \leq \bar{C}_H$. Now define

$$\tilde{B}_k(t, x) := D_p H_k(x, D\bar{v}_k), \quad A_k(t, x) := \tilde{m}_k M(x, D\bar{v}_k, D\check{v}_k), \quad \tilde{F}_k(t, x) := A_k(t, x)(D\bar{v}_k - D\check{v}_k).$$

Then m_k satisfies

$$(2.14) \quad \begin{cases} \partial_t m_k - \operatorname{div}(\tilde{B}_k m_k) = \Delta m_k + \operatorname{div} \tilde{F}_k & \text{in } (0, T) \times \mathbb{R}^d, \\ \partial_n m_k + (m_k \tilde{B}_k + \tilde{F}_k) \cdot n = 0 & \text{on } (0, T) \times \partial\Omega, \quad m_k(0, x) = 0. \end{cases}$$

with $|\tilde{B}_k| \leq C_H$ and $|A_k| \leq \mathcal{M} \bar{C}_H$ by the assumption (2.6), where

$$\mathcal{M} := \max_k C[1 + \|m_{0,k}\|_{\infty} + (1+T)\|D_p H_k(\cdot, Dv_k)\|_{L^{\infty}((0,T) \times \Omega)}]^r$$

is the upper bound on m_k given by Proposition 2.2 (where C depends only on the set Ω).

Step 4. We multiply the PDE in (2.14) by m_k and integrate by parts to get

$$\begin{aligned} 0 &= \int_0^t \frac{d}{dt} \int_{\Omega} \frac{m_k^2}{2} dx ds + \int_0^t \int_{\Omega} |Dm_k|^2 dx ds - \int_0^t \int_{\partial\Omega} m_k Dm_k \cdot n d\sigma ds \\ &\quad + \int_0^t \int_{\Omega} m_k \tilde{B}_k \cdot Dm_k dx ds - \int_0^t \int_{\partial\Omega} m_k^2 \tilde{B}_k \cdot n d\sigma ds \\ &\quad + \int_0^t \int_{\Omega} \tilde{F}_k \cdot Dm_k dx ds - \int_0^t \int_{\partial\Omega} m_k \tilde{F}_k \cdot n d\sigma ds. \end{aligned}$$

By the initial and boundary conditions in (2.14) we obtain

$$\begin{aligned} \frac{1}{2} \|m_k(t, \cdot)\|_2^2 + \int_0^t \|Dm_k(s, \cdot)\|_2^2 ds &= - \int_0^t \int_\Omega \left(m_k \tilde{B}_k + \tilde{F}_k \right) \cdot Dm_k dx ds \leq \\ &= \frac{1}{2\varepsilon} \int_0^t \|\tilde{F}_k(s, \cdot)\|_2^2 ds + \frac{\|\tilde{B}_k\|_\infty}{2\varepsilon} \int_0^t \|m_k(s, \cdot)\|_2^2 ds + \varepsilon \frac{\|\tilde{B}_k\|_\infty + 1}{2} \int_0^t \|Dm_k(s, \cdot)\|_2^2 ds, \end{aligned}$$

and with the choice $\varepsilon = 2/(\|\tilde{B}\|_\infty + 1) =: 1/c_1$

$$\|m_k(t, \cdot)\|_2^2 \leq c_1 \int_0^t \|\tilde{F}_k(s, \cdot)\|_2^2 ds + c_1 \|\tilde{B}_k\|_\infty \int_0^t \|m(s, \cdot)\|_2^2 ds.$$

Then Gronwall inequality and the definition of \tilde{F}_k give, for all $0 \leq t \leq T$,

$$(2.15) \quad \|m_k(t, \cdot)\|_2^2 \leq c_1 e^{c_1 \|\tilde{B}_k\|_\infty T} \|A_k\|_\infty^2 \int_0^t \|Dv_k(s, \cdot)\|_2^2 ds.$$

Step 5. Now we set

$$\phi(t) := \|Dv(t, \cdot)\|_2^2 = \sum_{k=1}^N \|Dv_k(t, \cdot)\|_2^2$$

and assume w.l.o.g. $C_H \geq 1$, so that $c_o, c_1 \leq C_H$. By combining (2.13) and (2.15) we get

$$\begin{aligned} \phi(t) &\leq N e^{C_H^2 T} \left(L_G \|m(T, \cdot)\|_2^2 + C_H L_F \int_t^T \|m(s, \cdot)\|_2^2 ds \right) \\ &\leq \bar{C}_H^2 C \left(L_G \int_0^T \phi(s) ds + C_H L_F \int_t^T \int_0^\tau \phi(s) ds d\tau \right), \quad C := N C_H e^{C_H^4 T^2} \mathcal{M}^2. \end{aligned}$$

Then $\Phi := \sup_{0 \leq t \leq T} \phi(t)$ satisfies

$$\Phi \leq \Phi \Psi, \quad \Psi := T \bar{C}_H^2 C (L_G + L_F C_H T / 2),$$

which implies $\Phi = 0$ if $\Psi < 1$. Therefore under such condition we conclude that $D\tilde{v}_k(t, x) = D\bar{v}_k(t, x)$ for all k, x and $0 \leq t \leq T$. By the uniqueness of solution for the KFP equation (e.g., Thm. I.2.2, p. 15 of [26]) we deduce $\tilde{m} = \bar{m}$ and then, by the Comparison Principle for the HJB equation in the Appendix, $\tilde{v} = \bar{v}$.

Finally, it is clear that Ψ can be made less than 1 by choosing either T , or \bar{C}_H , or both L_G and L_F small enough. \square

2.2. Examples and remarks.

Example 2.1. *Integral costs.* Consider F_k and G_k of the form

$$F_k(x, \mu) = F_o \left(x, \int_\Omega K(x, y) \mu(y) dy \right), \quad G_k(x, \mu) = g_1(x) \int_\Omega \bar{K}(x, y) \cdot \mu(y) dy + g_2(x)$$

with $F_o : \bar{\Omega} \times \mathbb{R}^N \rightarrow \mathbb{R}$ measurable and Lipschitz in the second variable uniformly in the first, whereas K is an $N \times N$ matrix with components in $L^2(\Omega \times \Omega)$. Then F_k satisfies (2.3). About G_k we assume $g_1, g_2 \in C^1(\Omega)$, Dg_1 bounded, the vector \bar{K} and its Jacobian $D_x \bar{K}$ with components in $L^2(\Omega \times \Omega)$. Then it satisfies (2.4). Of course all the data F_o, K, \bar{K}, g_i are allowed to change with the index $k = 1, \dots, N$.

Example 2.2. *Local costs.* Take $G_k = G_k(x)$ independent of $m(T)$ and F_k of the form $F_k(x, \mu) = F_k^l(x, \mu(x))$ with $F_k^l : \bar{\Omega} \times [0, +\infty)^N \rightarrow \mathbb{R}$ measurable and Lipschitz in the second variable uniformly in the first. Then F_k satisfies (2.3).

Example 2.3. *Costs depending on the moments.* The mean value of the density μ , $M(\mu) = \int_\Omega y \mu(y) dy$, and all its moments $\int_\Omega y^j \mu(y) dy$, $j = 2, 3, \dots$, are Lipschitz in L^2 by Example 2.1. Then any F_k (resp., G_k) depending on μ only via these quantities satisfies (2.3) (resp., (2.4)) if it is Lipschitz with respect to them uniformly with respect to x .

Example 2.4. Convex Hamiltonians. The usual Hamiltonians in MFGs are those arising from classical Calculus of Variations, e.g., $H_k(x, p) = b_k(x)(c_k + |p|^2)^{\beta_k/2}$, which satisfies the assumption (2.2) if $b_k \in C(\Omega)$ and either $c_k > 0$ or $c_k = 0$ and $\beta_k \geq 2$.

A related class of Hamiltonians are those of Bellman type associated to nonlinear systems, affine in the control $\alpha \in \mathbb{R}^d$,

$$(2.16) \quad H_k(x, p) := \sup_{\alpha} \{-(f_k(x) + g_k(x)\alpha) \cdot p - L_k(x, \alpha)\} = -f_k(x) \cdot p + L_k^*(x, -g_k(x)^T p),$$

where f_k is a Lipschitz vector field, g_k a Lipschitz square matrix, $L_k(x, \alpha)$ is the running cost of using the control α (adding to $F_k(x, m)$ in the cost functional of a representative player), and $L_k^*(x, \cdot)$ is its convex conjugate with respect to α . In this case one can check the assumption (2.2) on an explicit expression of L_k^* . For instance, if $L_k(x, \alpha) = |\alpha|^\gamma/\gamma$ then

$$H_k(x, p) = -f_k(x) \cdot p + \frac{\gamma-1}{\gamma} |g_k(x)^T p|^{\gamma/(\gamma-1)},$$

which satisfies (2.2) if $\gamma \leq 2$.

Example 2.5. Nonconvex Hamiltonians. Two-person 0-sum differential games give rise to the Isaacs Hamiltonians, which are defined in a way similar to (2.16) but as the inf-sup over two sets of controls. A motivation for considering these Hamiltonians in MFGs is proposed in [35]. A relevant example is the case of robust control, or nonlinear H_∞ control, studied in connection with MFGs by [7, 32] (see also the references therein). In this class of problems a deterministic disturbance $\sigma(x)\beta$ affects the control system (σ is a Lipschitz square matrix) and a worst case analysis is performed by assuming that $\beta \in \mathbb{R}^d$ is the control of an adversary who wishes to maximise the cost functional of the representative agent; a term $-\delta|\beta|^2/2$, with $\delta > 0$ is added to the running cost to penalise the energy of the disturbance. The Hamiltonian for robust control then becomes

$$(2.17) \quad H_k^{(r)}(x, p) := H_k(x, p) + \inf_{\beta} \{-\sigma(x)\beta + \delta|\beta|^2/2\} \cdot p = H_k(x, p) - \frac{|\sigma(x)^T p|^2}{2\delta},$$

which is the sum of the convex H_k of the previous example and a concave function of p . Clearly it satisfies the condition (2.2) if and only if H_k does.

Remark 2.1. Continuous dependence on data. Our proof of uniqueness can be adapted to show the Lipschitz dependence of solutions on some data. For instance, in Theorem 2.1 we may assume that $\bar{m}(0, x) = \bar{m}_0(x)$ and $\tilde{m}(0, x) = \tilde{m}_0(x)$, with $\bar{m}_0, \tilde{m}_0 \in \mathcal{P}_N(\Omega)$. Then a simple variant of the proof allows to estimate

$$\|\tilde{m}(t, \cdot) - \bar{m}(t, \cdot)\|_2^2 \leq \frac{C}{\delta} \|\tilde{m}_0 - \bar{m}_0\|_2^2$$

where $0 < \delta \leq 1 - \Psi$ and C depends on the same quantities as Ψ . A similar estimate holds for $\|D\tilde{v}(t, \cdot) - D\bar{v}(t, \cdot)\|_2^2$. Under some further assumptions on the costs F and G one can also use results on the HJB equation to obtain the continuous dependence of v itself upon the initial data m_0 . More precise results on continuous dependence of solutions with respect to data will be given elsewhere.

Remark 2.2. The statement of Theorem 2.1 holds with the same proof for solutions \mathbb{Z}^d -periodic in the space variable x in the case that F_k and G_k are \mathbb{Z}^d -periodic in x and without Neumann boundary conditions. In such case of periodic boundary conditions a uniqueness result for short T was presented by Lions in [31] for $N = 1$, regularizing running cost F , and for terminal cost G independent of $m(T)$. He used estimates in L^1 norm for m and in L^∞ norm for Dv , instead of the L^2 norms we used here in (2.13) and (2.15). See also [5] for the case of a single population.

Remark 2.3. The constants C_H and \bar{C}_H in the theorem depend only on the data of (2.1) if H_k and $D_p H_k$ are globally Lipschitz in p , uniformly in x , for all k . In this case the smallness condition $\Psi < 1$ does not depend on the solutions \tilde{v}, \bar{v} . In the next section we reach the same conclusion for much more general Hamiltonians H_k under some mild additional conditions on the costs F_k, G_k .

Remark 2.4. If the volatility is different among the populations the terms $\Delta v_k, \Delta m_k$ in (2.1) are replaced, respectively, by $\nu_k \Delta v_k$ and $\nu_k \Delta m_k$. If the constants ν_k are all positive, the theorem remains true with the function Ψ now depending also on ν_1, \dots, ν_N and minor changes in the proof. The case of volatility depending on x leads to operators of the form $\text{trace}(\sigma_k(x)\sigma_k^T(x)D^2v_k)$ in the the HJB equations and their adjoints in the KFP equations. This can also be treated, with some additional work in the proof, if such operators are uniformly elliptic, i.e., the minimal eigenvalue of the matrix $\sigma_k(x)\sigma_k^T(x)$ is bounded away from 0 for $x \in \bar{\Omega}$.

Remark 2.5. The $C^{2,\beta}$ regularity of $\partial\Omega$ can be weakened in Theorem 2.1. Here we used, e.g., Theorem IV.5.3 of [26] to produce a smooth test function φ in the proof of Proposition 2.2. However, we could work instead with a weak solution of the backward heat equation, which exists, for instance, if $\partial\Omega \in C^{1,\beta}$ by Theorem 6.49 of [30], or if it is "piecewise smooth" by Theorem III.5.1 in [26].

3. SPECIAL CASES AND APPLICATIONS

The function Ψ of Theorem 2.1 may depend on the solutions \tilde{v}, \bar{v} if the Hamiltonians H_k are not globally Lipschitz or they have unbounded second derivatives, because the constants C_H, \bar{C}_H may depend on the range of $D\tilde{v}$ and $D\bar{v}$. Under some further assumptions we can estimate these quantities and therefore get a uniqueness result where the function Ψ depends only on the data of the problem (2.1). The additional assumptions are

$$(3.1) \quad |F_k(x, \mu)| \leq C_F, \quad |G_k(x, \mu)| \leq C_G, \quad \forall x \in \bar{\Omega}, \mu \in \mathcal{P}_N(\Omega), k = 1, \dots, N,$$

$$(3.2) \quad |H_k(x, p)| \leq \alpha(1 + |p|^2), \quad |D_p H_k(x, p)|(1 + |p|) \leq \alpha(1 + |p|^2), \quad \forall x, p, k,$$

and $x \rightarrow G(x, \mu)$ of class C^2 , for all $\mu \in \mathcal{P}_N(\Omega)$, with

$$(3.3) \quad \|DG_k(\cdot, \mu)\|_\infty + \|D^2G_k(\cdot, \mu)\|_\infty \leq C'_G, \forall k.$$

Corollary 3.1. *Assume (2.2), (2.3), (2.4), $m_0 \in \mathcal{P}_N(\Omega)$, (3.1), (3.2), and (3.3). Then there exists $\bar{T} > 0$ such that for all $T \in (0, \bar{T}]$ can be at most one classical solution of (2.1).*

Proof. By Assumption (3.1) the functions $\pm(C_G + t(C_F + \alpha))$ are, respectively, a super- and a subsolution of the HJB equation in (2.1) with homogeneous Neumann condition and terminal condition G_k , for any k and m . Then the Comparison Principle in the Appendix gives for any solution of (2.1) the estimate

$$|v_k(t, x)| \leq C_G + TC_F, \quad \forall (t, x) \in [0, T] \times \bar{\Omega}, k = 1, \dots, N.$$

Now we can use an estimate of Theorem V.7.2, p. 486 of [26], stating that there is a constant K , depending only on $\max |v_k|, \alpha, C'_G$, and $\partial\Omega$, such that

$$|Dv_k(t, x)| \leq K, \quad \forall (t, x) \in [0, T] \times \bar{\Omega}, k = 1, \dots, N.$$

Then the constant C_H in (2.5) is bounded by $C'_H := \alpha(1 + K^2)/(1 + K)$, and \bar{C}_H defined by (2.6) can be estimated by

$$\bar{C}'_H := \max_{k=1, \dots, N} \sup_{x \in \Omega, |p|, |q| \leq K} \frac{|D_p H_k(x, p) - D_p H_k(x, q)|}{|p - q|}.$$

Now Theorem 2.1 gives the conclusion. \square

Remark 3.1. The constant \bar{T} in the Corollary depends only on $L_F, L_G, N, \alpha, C_F, C_G, C'_G, \max_k \|m_{0,k}\|_\infty, \Omega$ and the constants C'_H, \bar{C}'_H built in the proof. A similar results holds if, instead of T small, we assume L_F and L_G suitably small.

Example 3.1. *Costs satisfying the assumptions.* The nonlocal cost F_k and G_k of Example 2.1 satisfy Assumption (3.1) if, for instance, K, \bar{K} , and g_i are bounded and F_o is continuous.

The Assumption (3.3) is verified if $g_1, g_2 \in C^2(\bar{\Omega})$ and $|D_x^2 \bar{K}(x, y)| + |D_x^2 \bar{K}(x, y)| \leq C$ for all x, y .

For the local cost F_k of Example 2.2, (3.1) holds if F_k^l is bounded.

3.1. Well-posedness of segregation models. Next we combine this uniqueness result with an existence theorem for models of urban settlements and residential choice proposed in [1]. We take for simplicity

$$(3.4) \quad N = 2, \quad G_k \equiv 0, \quad H_k(x, p) = h_k(x, |p|).$$

We endow $\mathcal{P}_2(\Omega)$ with the Kantorovitch-Rubinstein distance and strengthen condition (3.1) to

$$(3.5) \quad (F_1, F_2) : \bar{\Omega} \times \mathcal{P}_2(\Omega) \rightarrow \mathbb{R}^2 \text{ continuous and with bounded range in } C^{1,\beta}(\bar{\Omega}),$$

for some $\beta > 0$. We also assume a compatibility condition and further regularity on m_0 :

$$(3.6) \quad \partial_n m_{0,k} = 0 \text{ on } \partial\Omega, \quad m_{0,k} \in C^{2,\beta}(\bar{\Omega}), \quad k = 1, 2.$$

Corollary 3.2. *Assume (2.2), (2.3), (3.2), (3.4), (3.5), (3.6), and $H_k \in C^1(\bar{\Omega} \times \mathbb{R}^d)$. Then there exists $\bar{T} > 0$ such that for all $T \in (0, \bar{T}]$ there exists a unique classical solution of (2.1).*

Proof. The existence of a solution (for any T) follows from Theorem 12 of [1]. Let us only note that, by (3.4), $D_p H_k(x, p) = \partial_{|p|} h_k(x, |p|) p / |p|$, and then the compatibility condition in (3.6) and the Neumann condition for v_k imply also the compatibility condition

$$(3.7) \quad \partial_n m_{0,k} + m_{0,k} D_p H_k(x, Dv_k(0, x)) \cdot n = 0 \quad \forall x \in \partial\Omega.$$

The uniqueness of the solution for small T follows from Corollary 3.1. \square

Remark 3.2. Here the constant \bar{T} depends on $L_F, \alpha, C_F, \max_k \|m_{0,k}\|_\infty, \Omega$, and the constants C'_H, \bar{C}'_H built in the proof of Corollary 3.1. The solution m and Dv depend in a Lipschitz way from the initial condition m_0 , as explained in Remark 2.1.

Example 3.2. *Costs of Schelling type.* Let $K_k : \bar{\Omega} \times \bar{\Omega} \rightarrow \mathbb{R}$ be Lipschitz and such that, for some $U(x)$ neighborhood of x , $K_k(x, y) = 1$ for $y \in U(x)$ and $K_k(x, y) = 0$ for y out of a small neighborhood of $U(x)$. Then

$$N_k(x, \mu_k) := \int_{\Omega} K_k(x, y) \mu_k(y) dy$$

represents the amount of population k around x . The cost functional for the k -th population introduced in [1] and inspired by the studies on segregation of T. Schelling [34] is of the form

$$F_k(x, \mu_1, \mu_2) := \left(\frac{N_k(x, \mu_k)}{N_k(x, \mu_k) + N_{3-k}(x, \mu_{3-k}) + \eta} - a_k \right)^-,$$

where $(\)^-$ denotes the negative part and $\eta > 0$ is very small. It means that if the ratio of the k -th population with respect to the total population in the neighborhood of x is above the threshold a_k , then a representative agent of this population is happy because his cost is 0, whereas below the threshold the agent incurs in a cost and therefore he wants to move from the neighborhood. These costs fall within Example 3.1 and satisfy (2.3) and (3.1). Moreover $F_k : \Omega \times \mathcal{P}_2(\Omega) \rightarrow \mathbb{R}$ is Lipschitz.

To meet the assumptions of Corollary 3.2 we assume the kernel K is of class C^2 in x and we approximate the negative part $(\)^-$ with a smooth function, e.g.,

$$\varphi_\varepsilon(r) := \frac{\sqrt{r^2 + \varepsilon^2} - r}{2},$$

for a small $\varepsilon > 0$. Then the cost functionals

$$F_k^\varepsilon(x, \mu_1, \mu_2) := \varphi_\varepsilon \left(\frac{N_k(x, \mu_k)}{N_k(x, \mu_k) + N_{3-k}(x, \mu_{3-k}) + \eta} - a_k \right)$$

satisfy also (3.5).

Example 3.3. *Hamiltonians.* Typical examples are either $H_k(x, p) = b_k(x)|p|^2$, with $b_k \in C(\bar{\Omega})$, or

$$H_k(x, p) = b_k(x)(1 + |p|^2)^{\beta_k/2}, \quad 0 < \beta_k \leq 2.$$

They satisfy (2.2) and (3.2), moreover they are in $C^1(\bar{\Omega} \times \mathbb{R}^d)$ if $b_k \in C^1(\bar{\Omega})$.

Remark 3.3. In the last Corollary 3.2 the simplifying assumption $G_k \equiv 0$ can be dropped and replaced with $G_k : \bar{\Omega} \times \mathcal{P}_2(\Omega) \rightarrow \mathbb{R}$ continuous, with bounded range in $C^{2,\beta}(\bar{\Omega})$, and satisfying (2.4). Then (3.3) holds and the constant \bar{T} depends also on L_G, C_G , and C'_G . Examples of such terminal costs can be given along the lines of Examples 2.1, 3.1, and 3.2.

3.2. Well-posedness of robust Mean Field Games. For simplicity we limit ourselves to a single population of agents, so $N = 1$ and we drop the subscripts k . The representative agent has the dynamics in \mathbb{R}^d

$$dX_s = (f(X_s) + g(X_s)\alpha_s + \sigma(X_s)\beta_s) ds + dW_s,$$

where f is a C^1 vector field in $\bar{\Omega}$, g and σ are C^1 scalar functions in $\bar{\Omega}$, W_s is a d -dimensional Brownian motion, α_s, β_s take values in \mathbb{R}^d and are, respectively, the control of the agent and a disturbance affecting the system. The cost functional is (for $\delta > 0$)

$$\mathbb{E} \left[\int_0^T \left(F(X_s, m(s, \cdot)) + \frac{|\alpha_s|^2}{2} - \delta \frac{|\beta_s|^2}{2} \right) ds + G(X_T, m(T, \cdot)) \right]$$

that the agent wants to minimise whereas the disturbance, modeled as a second player in a 2-person 0-sum game, wants to maximise. This leads to the Hamiltonian

$$(3.8) \quad H(x, p) = -f(x) \cdot p + g^2(x) \frac{|p|^2}{2} - \sigma^2(x) \frac{|p|^2}{2\delta}.$$

Note that here $g(x)$ and $\sigma(x)$ are scalars, different from Examples 2.4 and 2.5. On the costs we assume

$$(3.9) \quad F, G : \bar{\Omega} \times \mathcal{P}_1(\Omega) \rightarrow \mathbb{R} \text{ continuous with bounded range, resp., in } C^{1,\beta}(\bar{\Omega}) \text{ and } C^{2,\beta}(\bar{\Omega})$$

for some $\beta > 0$. The compatibility condition and regularity on m_0 now are

$$(3.10) \quad \partial_n m_0 - m_0 f \cdot n = 0 \text{ on } \partial\Omega, \quad m_0 \in C^{2,\beta}(\bar{\Omega}).$$

Corollary 3.3. *Assume $N = 1$ with the Hamiltonian defined by (3.8), (2.3), (2.4), (3.9), and (3.10). Then for all $T > 0$ there is a classical solution of (2.1), and there exists $\bar{T} > 0$ such that for all $T \in (0, \bar{T}]$ such solution is unique.*

Proof. The existence of a solution follows from Theorem 12 of [1]. In fact, $H \in C^1(\bar{\Omega} \times \mathbb{R}^d)$ and it has quadratic growth. Moreover

$$D_p H(x, p) = -f(x) + g^2(x)p - \frac{\sigma^2(x)}{\delta}p,$$

and then the compatibility condition in (3.10) and the Neumann condition for v imply again the compatibility condition (3.7).

The uniqueness of the solution for small T follows from Corollary 3.1, since H satisfies also (2.2). \square

Remark 3.4. Also here the solution m and Dv depend in a Lipschitz way from the initial condition m_0 , as explained in Remark 2.1.

Remark 3.5. Our example of robust MFG is different from the one in [7]. In that paper the state space is $\Omega = \mathbb{R}$, one-dimensional without boundary, the control system is linear in the state X_s , and the volatility is σX_s instead of 1, for some positive constant σ , so the parabolic operators in the HJB and KFP equations of (2.1) are degenerate at the origin. The well-posedness of the MFG system of PDEs in [7] is an open problem.

4. APPENDIX: A COMPARISON PRINCIPLE

The next result is known but we give its elementary proof for lack of a precise reference.

Proposition 4.1. *Assume $\Omega \subseteq \mathbb{R}^d$ is bounded with C^2 boundary, $H : \bar{\Omega} \times \mathbb{R}^d$ is of class C^1 with respect to p , and $u, v : [0, T] \times \bar{\Omega} \rightarrow \mathbb{R}$ are C^1 in t and C^2 in x and satisfy*

$$\begin{cases} -\partial_t u - \Delta u + H(x, Du) \leq -\partial_t v - \Delta v + H(x, Dv), & \text{in } (0, T) \times \Omega, \\ \partial_n u \leq \partial_n v, & \text{on } (0, T) \times \partial\Omega, \\ u(T, x) \leq v(T, x) & \text{in } \Omega. \end{cases}$$

Then $u \leq v$ in $[0, T] \times \bar{\Omega}$.

Proof. Let us assume first that

$$-\partial_t(u - v) - \Delta(u - v) + H(x, Du) - H(x, Dv) < 0 \quad \text{in } [0, T] \times \Omega,$$

$\partial_n(u - v) < 0$ on $[0, T] \times \partial\Omega$, and $(u - v)(T, x) \leq \delta$. Then the maximum of $u - v$ can be attained only at $t = T$, which implies $u - v \leq \delta$ in $[0, T] \times \bar{\Omega}$.

Now take $g \in C^2(\bar{\Omega})$ such that $Dg(x) = n(x)$ for all $x \in \partial\Omega$ and define

$$v_\varepsilon(t, x) := v(t, x) + \varepsilon(T - t)C + \varepsilon g(x).$$

Then $\partial_n(u - v_\varepsilon) = \partial_n(u - v) - \varepsilon < 0$ and $(u - v_\varepsilon)(T, x) \leq \varepsilon \|g\|_\infty$. Moreover, by Taylor's formula, for some q with $|q| \leq \|Dg\|_\infty$,

$$\begin{aligned} & -\partial_t(u - v_\varepsilon) - \Delta(u - v_\varepsilon) + H(x, Du) - H(x, Dv_\varepsilon) = \\ & -\partial_t(u - v) - \Delta(u - v) + H(x, Du) - H(x, Dv) - \varepsilon(C - \Delta g + D_p H(x, q) \cdot Dg) < -\varepsilon \end{aligned}$$

if C is chosen large enough. Then

$$u \leq v_\varepsilon + \varepsilon \|g\|_\infty \leq v + \varepsilon(TC + 2\|g\|_\infty)$$

and we conclude by letting $\varepsilon \rightarrow 0$. □

Remark 4.1. The result remains true if $\partial\Omega$ is merely C^1 and satisfies an interior sphere condition. This can be proved in a less direct way by linearizing the inequality for $u - v$ and then using the parabolic Strong Maximum Principle and the parabolic version of Hopf's Lemma for linear equations (see, e.g., [33]).

REFERENCES

- [1] Y. Achdou, M. Bardi, M. Cirant: Mean Field Games models of segregation, *Math. Models Methods Appl. Sci.* 27 (2017), 75–113,
- [2] D.M. Ambrose: Strong solutions for time-dependent mean field games with non-separable Hamiltonians, *J. Math. Pures Appl.* (9) 113 (2018), 141–154.
- [3] M. Bardi: Explicit solutions of some Linear-Quadratic Mean Field Games, *Netw. Heterog. Media* 7 (2012), 243–261
- [4] M. Bardi, E. Feleqi: Nonlinear elliptic systems and mean field games, *NoDEA Nonlinear Differential Equations Appl.* 23 (2016), 23–44
- [5] M. Bardi, M. Fischer: On non-uniqueness and uniqueness of solutions in some finite-horizon Mean Field Games, preprint arXiv 1707.00628, to appear in *ESAIM Control Optim. Calc. Var.*
- [6] M. Bardi, F.S. Priuli: Linear-Quadratic N -person and Mean-Field Games with Ergodic Cost, *SIAM J. Control Optim.* 52 (2014), 3022–3052.
- [7] D. Bauso, H. Tembine, T. Basar: Robust mean field games. *Dyn. Games Appl.* 6 (2016), 277–303
- [8] A. Briani, P. Cardaliaguet: Stable solutions in potential mean field game systems, *NoDEA Nonlinear Differential Equations Appl.* 25 (2018), no. 1, Art. 1, 26 pp.
- [9] R. Carmona, F. Delarue: *Probabilistic Theory of Mean Field Games with Applications I - II*, Springer 2018.
- [10] P. Cardaliaguet: Notes on Mean Field Games (from P-L. Lions' lectures at Collège de France), 2010.
- [11] P. Cardaliaguet, A. Porretta, D. Tonon: A segregation problem in multi-population Mean Field Games, *Annals of I.S.D.G.*, 15 (2017), pp 49–70.
- [12] M. Cirant: Multi-population mean field games systems with Neumann boundary conditions, *J. Math. Pures Appl.* 103 (2015), 1294–1315.
- [13] M. Cirant: Stationary focusing Mean-Field Games, *Comm. Partial Differential Equations.* 41(2016) 1324–1346.
- [14] M. Cirant, D. Tonon: Time-dependent focusing Mean-Field Games: the sub-critical case, preprint arXiv:1704.04014, to appear in *J. Dynam. Differential Equations*

- [15] M. Cirant, G. Verzini: Bifurcation and segregation in quadratic two-populations mean field games systems, *ESAIM Control Optim. Calc. Var.* 23 (2017), 1145–1177.
- [16] D.A. Gomes, J. Mohr, and R.R. Souza. Continuous time finite state mean field games. *Appl. Math. Optim.* 68 (2013), 99–143.
- [17] D. Gomes, L. Nurbekyan and E. Pimentel: Economic models and mean-field games theory, IMPA Mathematical Publications, Instituto Nacional de Matemática Pura e Aplicada, Rio de Janeiro, 2015.
- [18] D. Gomes, L. Nurbekyan and M. Prazeres: One-dimensional stationary mean-field games with local coupling, *Dyn. Games Appl.* 8 (2018), 315–351.
- [19] D. Gomes, E. Pimentel, and V. Voskanyan, *Regularity theory for mean-field game systems*, Springer 2016.
- [20] D. Gomes, J. Saude: Mean field games models: a brief survey. *Dyn. Games Appl.* 4 (2014), 110–154
- [21] O. Guéant, A reference case for mean field games models, *J. Math. Pures Appl.* (9), 92 (2009), 276–294.
- [22] O. Guéant, J.-M. Lasry, P.-L. Lions: Mean field games and applications, in "Paris-Princeton Lectures on Mathematical Finance 2010" (eds. R. A. Carmona, et al.), *Lecture Notes in Math.*, 2003, Springer, Berlin, (2011), 205–266.
- [23] M. Huang, R.P. Malhamé and P.E. Caines, Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle, *Commun. Inf. Syst.* 6 (2006) 221–251.
- [24] M. Huang, P.E. Caines and R.P. Malhamé, Large-population cost-coupled LQG problems with nonuniform agents: individual-mass behavior and decentralized ϵ -Nash equilibria, *IEEE Trans. Automat. Control* 52 (2007) 1560–1571.
- [25] M. Huang, P.E. Caines and R.P. Malhamé, An invariance principle in large population stochastic dynamic games. *J. Syst. Sci. Complex.* 20 (2007), 162–172.
- [26] O.A. Ladyzenskaja, V.A. Solonnikov, N.N. Uralceva: Linear and quasilinear equations of parabolic type. *Translations of Mathematical Monographs*, Vol. 23 American Mathematical Society, Providence, R.I. 1968
- [27] J.-M. Lasry, P.-L. Lions: Jeux champ moyen. I. Le cas stationnaire, *C. R. Math. Acad. Sci. Paris* 343 (2006), 619–625
- [28] J.-M. Lasry, P.-L. Lions: Jeux champ moyen. II. Horizon fini et controle optimal, *C. R. Math. Acad. Sci. Paris* 343 (2006), 679–684.
- [29] J.-M. Lasry, P.-L. Lions: Mean field games, *Jpn. J. Math.* 2 (2007), 229–260.
- [30] G.M. Lieberman: Second order parabolic differential equations. World Scientific Publishing Co., Inc., River Edge, NJ, 1996
- [31] P.-L. Lions: Lectures at Collège de France 2008-9.
- [32] J. Moon and T. Başar. Linear quadratic risk-sensitive and robust mean field games. *IEEE Trans. Automat. Control*, 62(3):1062–1077, 2016.
- [33] M.H. Protter, H.F. Weinberger: Maximum principles in differential equations. Prentice-Hall, Inc., Englewood Cliffs, N.J. 1967.
- [34] T. C. Schelling: *Micromotives and Macrobehavior*, Norton, 1978
- [35] H.V. Tran: A note on nonconvex Mean Field Games, preprint 2016, arXiv:1612.04725
- [36] B.-C. Wang and J.-F. Zhang. Mean field games for large-population multiagent systems with Markov jump parameters. *SIAM J. Control Optim.*, 50(4):2308–2334, 2012.

DEPARTMENT OF MATHEMATICS "T. LEVI-CIVITA", UNIVERSITY OF PADOVA, VIA TRIESTE 63, 35121 PADOVA, ITALY

E-mail address: bardi@math.unipd.it, cirant@math.unipd.it