

ON THE SINGULAR LOCAL LIMIT FOR CONSERVATION LAWS WITH NONLOCAL FLUXES

MARIA COLOMBO, GIANLUCA CRIPPA, AND LAURA V. SPINOLO

ABSTRACT. We give an answer to a question posed in [2], which can be loosely speaking formulated as follows. Consider a family of continuity equations where the velocity field is given by the convolution of the solution with a regular kernel. In the singular limit where the convolution kernel is replaced by a Dirac delta, one formally recovers a conservation law: can we rigorously justify this formal limit? We exhibit counter-examples showing that, despite numerical evidence suggesting a positive answer, one in general does not have convergence of the solutions. We also show that the answer is positive if we consider viscous perturbations of the nonlocal equations. In this case, in the singular local limit the solutions converge to the solution of the viscous conservation law.

KEYWORDS: nonlocal conservation law, nonlocal continuity equation, singular limit, local limit.

MSC (2010): 35L65.

1. INTRODUCTION AND MAIN RESULTS

We are concerned with the so-called nonlocal continuity equation (or nonlocal conservation law)

$$(1.1) \quad \partial_t w + \operatorname{div} [w b(w * \eta)] = 0.$$

In the previous expression, $b : \mathbb{R} \rightarrow \mathbb{R}^d$ is a Lipschitz continuous vector field, $w : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ is the unknown and div denotes the divergence computed with respect to the space variable only. The symbol $*$ denotes the convolution computed with respect to the space variable only and η is a convolution kernel satisfying

$$(1.2) \quad \eta : \mathbb{R}^d \rightarrow \mathbb{R}, \quad \eta \in C_c^\infty(\mathbb{R}^d), \quad \eta(x) = 0 \text{ if } |x| \geq 1, \quad \eta \geq 0, \quad \int_{\mathbb{R}^d} \eta(x) dx = 1.$$

In recent years, conservation laws involving nonlocal terms have been extensively studied owing to their applications to models for sedimentation [3], pedestrian [7] and vehicular [5] traffic, and others. We refer to the recent paper [5] for a more extended discussion and a more complete list of references. Here we only mention that the basic idea underpinning the use of equations like (1.1) in traffic models is, very loosely speaking, the following. The unknown w represents the density of pedestrians or cars and b their velocity. The nonlocal term $w * \eta$ appears since one postulates that pedestrians or drivers tune their velocity depending on the density of pedestrians or cars surrounding them.

In the present work we investigate a question posed by Amorim, R. Colombo and Teixeira in [2]. To precisely state the question, we consider the family of Cauchy problems

$$(1.3) \quad \begin{cases} \partial_t u_\varepsilon + \operatorname{div} [u_\varepsilon b(u_\varepsilon * \eta_\varepsilon)] = 0 \\ u_\varepsilon(0, x) = \bar{u}(x), \end{cases}$$

where b is as before a Lipschitz continuous vector field, ε is a positive parameter and \bar{u} is a summable and bounded initial datum. Assume that the family of convolution kernels η_ε is obtained from η by setting

$$(1.4) \quad \eta_\varepsilon(x) := \frac{1}{\varepsilon^d} \eta\left(\frac{x}{\varepsilon}\right), \quad 0 < \varepsilon \leq 1,$$

in such a way that when $\varepsilon \rightarrow 0^+$ the family η_ε converges weakly-* in the sense of measures to the Dirac delta. This implies that, when $\varepsilon \rightarrow 0^+$, the Cauchy problem (1.3) *formally* reduces to a scalar

conservation law

$$(1.5) \quad \begin{cases} \partial_t u + \operatorname{div} [ub(u)] = 0 \\ u(0, x) = \bar{u}(x). \end{cases}$$

The by now classical theory by Kruřkov [15] provides global existence and uniqueness results for so-called *entropy admissible* solutions. We refer to [11] for the definition and an extended discussion concerning entropy solution of conservation laws. The question posed in [2] can be formulated as follows.

Question 1. Can we rigorously justify the singular limit from (1.3) to (1.5)? In other words, does u_ε converge to the entropy admissible u , in a suitable topology?

Some remarks are here in order. First, Question 1 is motivated by numerical experiments. Indeed, in [2, § 3.3] the authors exhibit numerical evidence suggesting that there should be convergence. Second, to the best of our knowledge, the only previous analytical result concerning Question 1 is due to Zumbrun [17] and states that the answer to Question 1 is positive provided that the limit entropy solution u is smooth and the convolution kernel is even, i.e. $\eta(x) = \eta(-x)$ (see [17, Proposition 4.1] for a more precise statement). Third, even in the case $d = 1$, $b(u_\varepsilon) = u_\varepsilon$, establishing weak compactness of the family $\{u_\varepsilon\}$ is not a priori sufficient to establish convergence. Indeed, one needs strong convergence (or some more refined argument) to pass to the limit in the nonlinear term $u_\varepsilon u_\varepsilon * \eta_\varepsilon$. Fourth, similar questions show up when considering equations in transport form instead of in continuity form as in (1.3) and (1.5) (see for instance [4]); the analysis in such a case shares some similarities with that in the present paper and we plan to address it in future work.

In this paper we exhibit explicit counter-examples showing that the answer to Question 1 is, in general, negative. Also, we show that the answer is positive if we add to the right hand side of the first line of both (1.3) and (1.5) a viscous term. As we explain below, this is relevant in connection with the numerical analysis of the singular limit from (1.3) to (1.5).

We now describe our results more precisely. Our counter-examples can be summarized as follows:

- In § 5.1 we exhibit a counter-example showing that, in general, u_ε does not converge to the entropy admissible solution u weakly in L^p or weakly* in L^∞ . The example uses a family of even convolution kernels. See Lemma 5.1 for the precise statement. A drawback of this example is that the initial datum \bar{u} changes sign. This is not completely satisfactory in view of the applications, where the unknown typically represents a density.
- In § 5.2 we exhibit a counter-example with a nonnegative initial datum where we show that u_ε does not converge to u weakly in L^p or weakly* in L^∞ . See Lemma 5.2 for the precise statement. A drawback of this example is that we have to use “completely asymmetric” convolution kernels, namely we assume that $\eta(x) = 0$ for every $x > 0$. Note that this is consistent with numerical experiments provided in [2, § 3.2] and [5, § 5], where “completely asymmetric” kernels are connected with highly oscillatory behaviors of the solution.
- In § 5.3 we exhibit a counter-example involving a nonnegative initial datum and a family of even convolution kernels. In this counter-example we show that for every $\delta > 0$ the family u_ε does not converge to u strongly in $L^{1+\delta}$. See Lemma 5.5 for the precise statement.

As mentioned before, we manage to establish positive results by adding to the first line of (1.3) and (1.5) a second order perturbation. More precisely, we consider the family of Cauchy problems

$$(1.6) \quad \begin{cases} \partial_t u_{\varepsilon\nu} + \operatorname{div} [u_{\varepsilon\nu} b(u_{\varepsilon\nu} * \eta_\varepsilon)] = \nu \Delta u_{\varepsilon\nu} \\ u_{\varepsilon\nu}(0, x) = \bar{u}(x), \end{cases}$$

which depends on two parameters $\varepsilon > 0$ and $\nu > 0$. When $\varepsilon \rightarrow 0^+$ and ν is fixed, the family of Cauchy problems (1.6) *formally* reduces to

$$(1.7) \quad \begin{cases} \partial_t u_\nu + \operatorname{div} [u_\nu b(u_\nu)] = \nu \Delta u_\nu \\ u_\nu(0, x) = \bar{u}(x). \end{cases}$$

On the other hand, when $\nu \rightarrow 0^+$ and ε is fixed, the family of Cauchy problems formally reduces to (1.3), while (1.7) reduces to (1.5) (see (1.9) below for a scheme). The reason why we consider the viscous approximations (1.6), (1.7) is the following. As mentioned before, Question 1 is motivated by the numerical evidence exhibited in [2]. The numerical tests showing convergence are obtained by using a Lax-Friedrichs type scheme involving some so-called *numerical viscosity*, as it typical of many numerical schemes for conservation laws (see the book by LeVeque [16] for an extended introduction). Very loosely speaking, the numerical viscosity consists of finite differences terms that mimic a second order operator like the Laplacian. For this reason, the analysis of the viscous approximation (1.6), (1.7) may provide some insight in the understanding of the numerical tests.

Our main result involving the singular limit from (1.6) to (1.7) is the following.

Theorem 1.1. *Let b be a Lipschitz continuous function, $\bar{u} \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$, $\nu > 0$ and p such that*

$$(1.8) \quad 2 \leq p < \infty, \quad p > d.$$

Let $u_{\varepsilon\nu}$ and u_ν be the solutions of (1.6) and (1.7) starting from \bar{u} , respectively. Then

$$u_{\varepsilon\nu} \rightarrow u_\nu \quad \text{strongly in } L_{\text{loc}}^\infty([0, +\infty[; L^p(\mathbb{R}^d)).$$

Some remarks are here in order. First, the Cauchy problem (1.6) has a unique weak solution, see Theorem 2.1 in § 2 for the precise statement. Second, in the case $d = 1$, $b(u) = u$, $p = 2$, $\bar{u} \in W^{1,\infty}(\mathbb{R})$, Theorem 1.1 was established by Calderoni and Pulvirenti [6]. The main novelties of Theorem 1.1 with respect to the analysis in [6] can be summarized as follows:

- We provide a completely different proof. Indeed, in [6] the authors explicitly compute the equations satisfied by the Fourier transforms $\hat{u}_{\varepsilon\nu}$ and \hat{u}_ν and use them to control the L^2 norm of the difference. The proof explicitly uses the fact that $b(u) = u$ and the regularity of the initial datum.
- On the other hand, our argument is based on a-priori estimates obtained by extensively using energy estimates and the Duhamel representation formula. We first establish Theorem 3.1 in the case when the initial datum \bar{u} is regular. Next, we introduce a careful perturbation argument and we establish the proof in the general case. Our argument is fairly robust, it applies to general velocity fields b , to equations in several space dimensions, and to rough initial data, and provides more quantitative estimates, see Remark 1.3 below.

As a further remark, we explicitly point out that Theorem 1.1 requires neither symmetry conditions on the convolution kernels η_ε nor sign conditions on the initial datum \bar{u} .

Finally, we discuss the vanishing viscosity limit from (1.6) to (1.3). Our result is the following.

Proposition 1.2. *Under the assumptions of Theorem 1.1, let $u_{\varepsilon\nu}$ and u_ε satisfy (1.6) and (1.3), respectively. For every $\varepsilon > 0$, we have that $u_{\varepsilon\nu} \xrightarrow{*} u_\varepsilon$ weakly* in $L_{\text{loc}}^\infty([0, +\infty[\times\mathbb{R}^d)$ as $\nu \rightarrow 0^+$.*

Note that in the statement of Proposition 1.2 the parameter $\varepsilon > 0$ is fixed and hence the weak* convergence suffices to pass to the limit in the equation, owing to the regularizing effect of the convolution. Also, note that at the local level the vanishing viscosity limit from (1.7) to (1.5) is established in the work by Kruřkov [15].

The take-home message obtained by combining the counter-examples, Theorem 1.1, Kruřkov's Theorem and Proposition 1.2 can be therefore represented as follows:

$$(1.9) \quad \begin{array}{ccc} \partial_t u_{\varepsilon\nu} + \operatorname{div} [u_{\varepsilon\nu} b(u_{\varepsilon\nu} * \eta_\varepsilon)] = \nu \Delta u_{\varepsilon\nu} & \xrightarrow[\text{Theorem 1.1}]{\varepsilon \rightarrow 0^+} & \partial_t u_\nu + \operatorname{div} [u_\nu b(u_\nu)] = \nu \Delta u_\nu \\ \downarrow \nu \rightarrow 0^+ \text{ Proposition 1.2} & & \downarrow \nu \rightarrow 0^+ \text{ Kruřkov's Theorem} \\ \partial_t u_\varepsilon + \operatorname{div} [u_\varepsilon b(u_\varepsilon * \eta_\varepsilon)] = 0 & \xrightarrow[\text{False in general}]{\varepsilon \rightarrow 0^+} & \partial_t u + \operatorname{div} [u b(u)] = 0 \end{array}$$

To conclude, we make a remark concerning the “diagonal” convergence, which can be tracked explicitly in the case of regular initial data.

Remark 1.3. Under the assumptions of Theorem 1.1, let $u_{\varepsilon\nu}$ satisfy (1.6), let u be the Kruřkov entropy admissible solution of (1.5), and fix p satisfying (1.8). Combining Kruřkov's Theorem with Theorem 1.1 and by a diagonal argument we infer that there is a sequence (ε_n, ν_n) such that $\varepsilon_n \rightarrow 0^+$, $\nu_n \rightarrow 0^+$ and $u_{\varepsilon_n \nu_n} \rightarrow u$ strongly in $L_{\text{loc}}^\infty([0, +\infty[; L^p(\mathbb{R}^d))$, as $n \rightarrow +\infty$. In the case when the initial datum is sufficiently regular, namely $\bar{u} \in W^{1,p}(\mathbb{R}^d)$, we explicitly determine a coupling $\varepsilon \leq e^{-C\nu^{-\beta}}$ (for constants $C > 0$ and $\beta > 0$ specified later) under which the above diagonal convergence holds true (see Theorem 3.1 below).

Paper outline. The paper is organized as follows. In § 2 we establish well-posedness of the Cauchy problem (1.6), we slightly extend known well-posedness results for (1.3) and we establish Proposition 1.2. In § 3 we establish Theorem 1.1 under the additional assumption that the initial datum \bar{u} is regular. In § 4 we complete the proof of Theorem 1.1 and in § 5 we discuss the counter-examples to the nonlocal to local limit from (1.3) to (1.5).

Notation. For the readers' convenience, we recall here the main notation used in the present paper.

We denote by $C(a_1, \dots, a_N)$ a constant only depending on the quantities a_1, \dots, a_N . Its precise value can vary from occurrence to occurrence.

General mathematical symbols.

- $f * g$: the convolution of the functions f and g , computed with respect to the variable x only.
- $\text{div } f$: the divergence of the vector field f , computed with respect to the x variable only.
- $\mathbf{1}_E$: the characteristic function of the measurable set E .
- $|E|$: the Lebesgue measure of the measurable set E .
- L^p : the Lebesgue space $L^p(\mathbb{R}^d)$, $p \in [1, +\infty]$.
- $\|\cdot\|_{L^p}$: the standard norm in $L^p(\mathbb{R}^d)$.

Symbols introduced in the present paper.

- b : the vector field satisfying (1.10).
- L : the Lipschitz constant in (1.10).
- η, η_ε : the convolution kernel in (1.2) and (1.4).
- u : the entropy solution of the conservation law (1.5).
- u_ε : the solution of the nonlocal nonviscous problem (1.3).
- u_ν : the solution of the local viscous problem (1.7).
- $u_{\varepsilon\nu}$: the solution of the nonlocal nonviscous problem (1.6).
- G, G_ν : the heat kernel in (3.3) and (3.5).
- $S_t^{\varepsilon\nu}, S_t^\nu$: the semigroups defined in (4.1).

Remark 1.4. Consider the Lipschitz continuous function $b : \mathbb{R} \rightarrow \mathbb{R}^d$ in (1.3), (1.5), (1.6) and (1.7). We can assume, with no loss of generality, that $b(0) = 0$. Indeed, assume that this is not the case and that $b(0) = \xi \neq 0$. Assume furthermore that the function $u_{\varepsilon\nu}$ satisfies (1.6), then we can set

$$\tilde{u}_{\varepsilon\nu}(t, x) := u_{\varepsilon\nu}(t, x - \xi t),$$

and obtain that $\tilde{u}_{\varepsilon\nu}$ satisfies

$$\partial_t \tilde{u}_{\varepsilon\nu} + \text{div} \left[\tilde{u}_{\varepsilon\nu} \tilde{b}(\tilde{u}_{\varepsilon\nu} * \eta_\varepsilon) \right] = \nu \Delta \tilde{u}_{\varepsilon\nu} \quad \text{where} \quad \tilde{b}(\tilde{u}_{\varepsilon\nu} * \eta_\varepsilon) := b(\tilde{u}_{\varepsilon\nu} * \eta_\varepsilon) - \xi.$$

For this reason in the following we assume that b satisfies

$$(1.10) \quad b(0) = 0, \quad |b(x) - b(y)| \leq L|x - y| \text{ for every } x, y \in \mathbb{R}.$$

Remark 1.5. Theorem 1.1 states that $u_{\varepsilon\nu} \rightarrow u_\nu$ strongly in $L_{\text{loc}}^\infty([0, +\infty[; L^p)$, hence to establish the thesis it suffices to prove that, for every $T > 0$, $u_{\varepsilon\nu} \rightarrow u_\nu$ strongly in $L^\infty([0, T]; L^p)$. A similar remark applies to Proposition 1.2 and to the other positive results, which are all local in time. To simplify the notation, in the following we take $T = 1$.

2. PRELIMINARY RESULTS: WELL-POSEDNESS OF THE VISCOUS AND NONVISCOUS CAUCHY PROBLEM WITH NONLOCAL FLUXES

This section is organized as follows: in § 2.1 we establish well-posedness of the nonlocal viscous Cauchy problem (1.6) by relying on a fixed point argument. In § 2.2 we establish a uniqueness result for the nonlocal conservation law (1.3) that slightly extends previous results in [2, 8, 10]. Finally, in § 2.3 we establish the proof of the nonlocal vanishing viscosity result stated in Proposition 1.2.

2.1. Well-posedness of the viscous Cauchy problem with a nonlocal flux. We establish the following well-posedness result.

Theorem 2.1. *Let $\bar{u} \in L^1 \cap L^\infty(\mathbb{R}^d)$ and b a vector field satisfying (1.10). Then the nonlocal viscous Cauchy problem (1.6) has a distributional solution $u_{\varepsilon\nu}$, unique in the class (2.1)-(2.2), that satisfies*

$$(2.1) \quad \|u_{\varepsilon\nu}(t, \cdot)\|_{L^1} \leq \|\bar{u}\|_{L^1}, \quad \|u_{\varepsilon\nu}(t, \cdot)\|_{L^\infty} \leq C(\|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{L^1}, \|\nabla\eta\|_{L^\infty}, L, d, \varepsilon), \quad \text{for every } t \in [0, 1],$$

$$(2.2) \quad \partial_t u_{\varepsilon\nu} \in L^2([0, 1]; H^{-1}(\mathbb{R}^d)), \quad u_{\varepsilon\nu} \in L^2([0, 1]; H^1(\mathbb{R}^d)).$$

Remark 2.2. The function $u_{\varepsilon\nu}$ is in principle only defined for a.e. (t, x) . However, the regularity (2.2) implies that, up to changing $u_{\varepsilon\nu}$ in a set of measure 0 in $[0, 1] \times \mathbb{R}^d$, we can assume that $u_{\varepsilon\nu} \in C^0([0, 1]; L^2(\mathbb{R}^d))$. In the following, we always consider this L^2 -continuous representative; in this way the function $u_{\varepsilon\nu}$ is well-defined for every t and the estimates (2.1) hold for every t .

Proof of Theorem 2.1. To simplify the notation, let $\varepsilon = 1$, $\nu = 1$ and consider the Cauchy problem

$$(2.3) \quad \begin{cases} \partial_t v + \operatorname{div} [vb(v * \eta)] = \Delta v \\ v(0, x) = \bar{u}(x). \end{cases}$$

The proof straightforwardly extends to the general case and relies on a classical fixed point argument that we sketch below.

STEP 1: we introduce the functional setting. We fix a constant $0 < \tau < 1$, to be determined in the following, and we define the set X by setting

$$(2.4) \quad X := \{z \in C^0([0, \tau]; L^2(\mathbb{R}^d)) : \|z(t, \cdot)\|_{L^1} \leq \|\bar{u}\|_{L^1} \forall t \in [0, \tau]\}.$$

We fix a function $\zeta \in X$ and we consider the Cauchy problem

$$(2.5) \quad \begin{cases} \partial_t z + \operatorname{div} [zb(\zeta * \eta)] = \Delta z \\ z(0, x) = \bar{u}(x). \end{cases}$$

Since ζ is now fixed, the equation at the first line of the above system is a standard linear parabolic equation with smooth coefficients. By using classical methods for evolution equations (see for instance [14, § 7]) one can show that (2.5) has a unique solution satisfying

$$\partial_t z \in L^2([0, \tau]; H^{-1}(\mathbb{R}^d)), \quad z \in L^2([0, \tau]; H^1(\mathbb{R}^d)),$$

which implies that (up to re-defining z on a negligible set of times) $z \in C^0([0, \tau]; L^2(\mathbb{R}^d))$. In the following, we always identify z and its L^2 -continuous representative, in such a way that $z(t, \cdot)$ is well-defined for every $t > 0$. We define the map T by setting $T(\zeta) = z$, where z is the solution of (2.5).

STEP 2: we show that the map T defined as in STEP 1 attains values in X . We fix a regular function $\beta : \mathbb{R} \rightarrow \mathbb{R}$ and by multiplying the equation at the first line of (2.5) times $\beta'(z)$ we get

$$(2.6) \quad \partial_t [\beta(z)] + \operatorname{div} [b(\zeta * \eta)\beta(z)] + \operatorname{div} [b(\zeta * \eta)](z\beta'(z) - \beta(z)) = \operatorname{div} [\nabla z \beta'(z)] - \beta''(z)|\nabla z|^2.$$

We point out that by (1.10) and (2.4)

$$|\operatorname{div} [b(\zeta * \eta)]| \leq C(L, d) \|\nabla\eta\|_{L^\infty} \|\bar{u}\|_{L^1}.$$

By space-time integrating (2.6) we get

$$(2.7) \quad \int_{\mathbb{R}^d} \beta(z)(t, \cdot) dx - \int_{\mathbb{R}^d} \beta(\bar{u}) dx + \int_0^t \int_{\mathbb{R}^d} \beta''(z) |\nabla z|^2 dx ds \\ \leq C(L, d) \|\bar{u}\|_{L^1} \|\nabla \eta\|_{L^\infty} \int_0^t \int_{\mathbb{R}^d} |z\beta'(z) - \beta(z)| dx ds, \quad \text{for every } t \in [0, \tau].$$

By applying (2.7) with $\beta(z) = z^2$ and using the Grönwall Lemma we get that for every $t \in [0, \tau]$

$$(2.8) \quad \|z(t, \cdot)\|_{L^2} \leq C(L, d, \|\bar{u}\|_{L^1}, \|\nabla \eta\|_{L^\infty}) \|\bar{u}\|_{L^2}.$$

Also, by using (2.7) and choosing a suitable approximation of $\beta(z) = |z|$, we get

$$(2.9) \quad \int_{\mathbb{R}^d} |z|(t, \cdot) dx - \int_{\mathbb{R}^d} |\bar{u}| dx \leq 0.$$

This implies that the solution of (2.5), i.e. $T(\zeta)$, belongs to the set X defined as in (2.4).

STEP 3: we show that the map T defined as in STEP 1 is a contraction provided that τ is sufficiently small. We fix $\zeta_1, \zeta_2 \in X$ and we term $z_1 = T(\zeta_1)$ and $z_2 = T(\zeta_2)$. First, we point out that owing to the Young Inequality

$$(2.10) \quad \|b(\zeta_1 * \eta) - b(\zeta_2 * \eta)\|_{L^\infty} \leq L \|(\zeta_1 - \zeta_2) * \eta\|_{L^\infty} \leq L \|(\zeta_1 - \zeta_2)\|_{C^0([0, \tau]; L^2)} \|\eta\|_{L^2},$$

for every $t \in [0, \tau]$. By subtracting the equation for z_2 from the equation for z_1 we get

$$\partial_t [z_1 - z_2] + \operatorname{div} \left[[z_1 - z_2] b(\zeta_1 * \eta) + z_2 [b(\zeta_1 * \eta) - b(\zeta_2 * \eta)] \right] = \Delta [z_1 - z_2].$$

By arguing as in STEP 2 and recalling that $z_2 \equiv z_1$ at $t = 0$, we arrive at

$$(2.11) \quad \int_{\mathbb{R}^d} |z_1 - z_2|^2(t, \cdot) dx \leq C(L, d, \|\bar{u}\|_{L^1}, \|\nabla \eta\|_{L^\infty}) \int_0^t \int_{\mathbb{R}^d} |z_1 - z_2|^2(s, \cdot) dx ds \\ + 2 \left| \int_0^t \int_{\mathbb{R}^d} \operatorname{div} \left[z_2 [b(\zeta_1 * \eta) - b(\zeta_2 * \eta)] \right] (z_1 - z_2) dx ds \right| - 2 \int_0^t \int_{\mathbb{R}^d} |\nabla [z_1 - z_2]|^2 dx ds.$$

Next, we point out that

$$\left| \int_0^t \int_{\mathbb{R}^d} \operatorname{div} \left[z_2 [b(\zeta_1 * \eta) - b(\zeta_2 * \eta)] \right] (z_1 - z_2) dx ds \right| = \left| \int_0^t \int_{\mathbb{R}^d} z_2 [b(\zeta_1 * \eta) - b(\zeta_2 * \eta)] \cdot \nabla [z_1 - z_2] dx ds \right| \\ \leq \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} z_2^2 |b(\zeta_1 * \eta) - b(\zeta_2 * \eta)|^2 dx ds + \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} |\nabla [z_1 - z_2]|^2 dx ds.$$

To control the first term in the right hand side of the above expression we combine (2.8) and (2.10). By plugging the above inequality into (2.11) we then arrive at

$$\int_{\mathbb{R}^d} |z_1 - z_2|^2(t, \cdot) dx \leq C(L, d, \|\bar{u}\|_{L^1}, \eta) \left[\int_0^t \int_{\mathbb{R}^d} |z_1 - z_2|^2(s, \cdot) dx ds + \tau \|\bar{u}\|_{L^2}^2 \|(\zeta_1 - \zeta_2)\|_{C^0([0, \tau]; L^2)}^2 \right]$$

and owing to the Grönwall Lemma and recalling that $z_1 = T(\zeta_1)$, $z_2 = T(\zeta_2)$ this implies that T is a contraction provided that τ is sufficiently small. To establish existence and uniqueness on the interval $[0, 1]$ we iterate the above argument a finite number of times.

STEP 4: we establish the L^∞ estimate. We recall (2.5), we set

$$\Xi := \|\operatorname{div}[b(\zeta * \eta)]\|_{L^\infty}$$

and we point out that the solution z of the Cauchy problem (2.5) satisfies

$$(2.12) \quad \|z(t, \cdot)\|_{L^\infty} \leq \|\bar{u}\|_{L^\infty} \exp(\Xi t), \quad \text{for every } t.$$

The proof of the above estimate is standard, and can be found for instance in [9, Lemma 3.4]. By construction, the solution of (2.3) satisfies (2.5) provided that $\zeta = z$. If this is the case, by (2.9)

$$\Xi = \|\operatorname{div}[b(z * \eta)]\|_{L^\infty} \leq C(L, d) \|z\|_{L^1} \|\nabla \eta\|_{L^\infty} \leq C(L, d) \|\bar{u}\|_{L^1} \|\nabla \eta\|_{L^\infty}$$

and owing to (2.12) this establishes the L^∞ estimate in (2.1). \square

2.2. Well-posedness of the Cauchy problem for a continuity equation with nonlocal flux. In this section we establish an existence and uniqueness result that slightly extends the well-posedness result in [2] (see also [8]).

Proposition 2.3. *Assume that b and η satisfy (1.10) and (1.2), respectively, and that $\bar{u} \in L^1 \cap L^\infty$. Then the Cauchy problem (1.3) has a distributional solution that satisfies*

$$u_\varepsilon \in L^\infty_{\text{loc}}([0, +\infty[; L^\infty) \cap C^0([0, +\infty[; L^1).$$

Also, the solution is unique in the class of locally bounded, distributional solutions.

In the following we identify u_ε and its L^1 strongly continuous representative. The existence part of Proposition 2.3 is a consequence of the analysis in [2, § 2] (see also [8, 10]). The relatively new part is the uniqueness: indeed, in [2] uniqueness is established in a more restrictive class. More precisely, in [8] it is shown that there is a unique solution u , in the sense of Kruřkov [15], of the conservation law

$$(2.13) \quad \begin{cases} \partial_t u_\varepsilon + \operatorname{div} [u_\varepsilon g_\varepsilon] = 0 \\ u_\varepsilon(0, x) = \bar{u}(x) \end{cases}$$

provided that the function g_ε is given by $g_\varepsilon := b(u_\varepsilon * \eta_\varepsilon)$ (see [2, Definition 2.1]). On the other hand, Proposition 2.3 states the uniqueness of locally bounded distributional solutions.

Proof of Proposition 2.3, uniqueness. Let u_ε be a distributional solution of (1.3). Then u_ε is a distributional solution of (2.13). Next, we observe that the first line of (2.13) is a continuity equation with a regular in space coefficient g_ε . Every locally bounded distributional solution of (2.13) is therefore renormalized, meaning that for every $\beta \in C^1(\mathbb{R})$ we have that $\beta(u)$ is a distributional solution of

$$(2.14) \quad \partial_t [\beta(u)] + \operatorname{div} [\beta(u)g_\varepsilon] + \operatorname{div} g_\varepsilon [\beta'(u)u - \beta(u)] = 0.$$

This is for instance an application (in a very easy case) of the DiPerna-Lions-Ambrosio theory, see [1, 13]. Equation (2.14) implies that, up to redefining u_ε in a negligible set, $u_\varepsilon \in C^0([0, +\infty[; L^1)$: this can be proved by arguing as in the proof of Corollary 3.14 in [12]. By (2.14), u_ε is a solution, in the sense of Kruřkov, of the conservation law (2.13). Since by the analysis in [8, § 2] distributional solutions of (1.3) that are also Kruřkov solutions of (2.13) are unique, this concludes the proof of Proposition 2.3. \square

2.3. Proof of Proposition 1.2. Let $\varepsilon > 0$. We consider $u_{\varepsilon\nu}$ satisfying (1.6) and we recall the L^∞ estimate in (2.1). We fix a sequence ν_n and a function $u_\varepsilon \in L^\infty([0, 1] \times \mathbb{R}^d)$ such that

$$(2.15) \quad u_{\varepsilon\nu_n} \xrightarrow{*} u_\varepsilon \text{ weakly}^* \text{ in } L^\infty([0, 1] \times \mathbb{R}^d) \text{ as } \nu_n \rightarrow 0^+.$$

We claim that u_ε is a distributional solution of (1.3). To take the limit in the distributional formulation of (1.6) and prove this claim, it is enough to show that

$$(2.16) \quad u_{\varepsilon\nu_n} * \eta_\varepsilon \rightarrow u_\varepsilon * \eta_\varepsilon \text{ strongly in } L^1_{\text{loc}}([0, 1] \times \mathbb{R}^d).$$

If the claim is true, since bounded, distributional solutions of (1.3) are unique by Proposition 2.3, the whole family $u_{\varepsilon\nu_n}$ converges to u_ε weakly* in $L^\infty([0, 1] \times \mathbb{R}^d)$, proving Proposition 1.2.

To show (2.16) we point out first that by (2.1) for every $t \in [0, 1]$

$$(2.17) \quad \|[u_{\varepsilon\nu} * \eta_\varepsilon](t, \cdot)\|_{L^\infty} + \|\nabla[u_{\varepsilon\nu} * \eta_\varepsilon](t, \cdot)\|_{L^\infty} \leq \|u_{\varepsilon\nu}(t, \cdot)\|_{L^\infty} \|\eta_\varepsilon\|_{W^{1,1}} \leq C(\|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{L^1}, \eta, L, d, \varepsilon).$$

The time derivative of $u_{\varepsilon\nu} * \eta_\varepsilon$ is obtained by convolving every term in (1.6) with η_ε , that is

$$\partial_t [u_{\varepsilon\nu} * \eta_\varepsilon](t, x) = -\operatorname{div} [\eta_\varepsilon * (u_{\varepsilon\nu} b(u_{\varepsilon\nu}))] + \nu u_{\varepsilon\nu} * \Delta \eta_\varepsilon.$$

By using (1.10), (1.4) and (2.1) we conclude that

$$(2.18) \quad \|\partial_t [u_{\varepsilon\nu} * \eta_\varepsilon](t, \cdot)\|_{L^\infty} \leq \|\nabla \eta_\varepsilon\|_{L^\infty} \|u_{\varepsilon\nu} b(u_{\varepsilon\nu})\|_{L^1} + \nu \|u_{\varepsilon\nu}\|_{L^1} \|\Delta \eta_\varepsilon\|_{L^\infty} \leq C(\|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{L^1}, \eta, L, d, \varepsilon).$$

Finally, we combine (2.17) and (2.18) and we apply the Ascoli-Arzelà Theorem: there is a continuous function w such that, up to subsequences (that we do not re-label) $u_{\varepsilon\nu_n} * \eta_\varepsilon \rightarrow w$ uniformly on compact sets of $[0, 1] \times \mathbb{R}^d$. For any $\phi \in C_c^\infty([0, 1] \times \mathbb{R}^d)$, terming $\check{\eta}_\varepsilon(z) := \eta_\varepsilon(-z)$ and by (2.15), we have

$$\begin{aligned} \int_0^1 \int_{\mathbb{R}^d} \phi(t, x) w(t, x) dx dt &= \lim_{n \rightarrow \infty} \int_0^1 \int_{\mathbb{R}^d} \phi[u_{\varepsilon\nu_n} * \eta_\varepsilon] dx dt = \lim_{n \rightarrow \infty} \int_0^1 \int_{\mathbb{R}^d} u_{\varepsilon\nu_n} [\phi * \check{\eta}_\varepsilon] dy dt \\ &= \int_0^1 \int_{\mathbb{R}^d} u_\varepsilon [\phi * \check{\eta}_\varepsilon] dy dt = \int_0^1 \int_{\mathbb{R}^d} \phi[u_\varepsilon * \eta_\varepsilon] dx dt. \end{aligned}$$

By the arbitrariness of ϕ we deduce that $w = u_\varepsilon * \eta_\varepsilon$ a.e. in $[0, 1] \times \mathbb{R}^d$ and hence we prove (2.16). \square

3. CONVERGENCE OF THE NONLOCAL VISCOUS APPROXIMATION FOR REGULAR DATA

In this section we establish the nonlocal to local limit asserted in Theorem 1.1 assuming more restrictive conditions on the initial data. This intermediate result is pivotal to the proof of Theorem 1.1.

Theorem 3.1. *Fix $0 < \nu < 1/4$. Assume that b satisfies (1.10) and η_ε satisfies (1.2) and (1.4). Let p satisfy (1.8), let $\beta = (p + d)/(p - d)$ and assume that $\bar{u} \in L^1 \cap L^\infty \cap W^{1,p}(\mathbb{R}^d)$. Let $u_{\varepsilon\nu}$ and u_ν be the solutions of the Cauchy problems (1.6) and (1.7), respectively. Then there exists $C := C(d, p, L, \|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{W^{1,p}})$ such that, if $\varepsilon \leq e^{-C\nu^{-\beta}}$, we have*

$$(3.1) \quad \|u_{\varepsilon\nu}(t, \cdot) - u_\nu(t, \cdot)\|_{L^p} \leq \varepsilon e^{C\nu^{-\beta}}, \quad \text{for every } t \in [0, 1].$$

This, in particular, implies that $u_{\varepsilon\nu} \rightarrow u_\nu$ strongly in $L_{\text{loc}}^\infty([0, +\infty[; L^p(\mathbb{R}^d))$ as $\varepsilon \rightarrow 0^+$.

To establish Theorem 3.1 we introduce the function

$$(3.2) \quad z_\varepsilon := u_{\varepsilon\nu} - u_\nu.$$

Note that, to simplify the notation, we do not explicitly indicate the dependence of z_ε on ν . Next, we compute the equation satisfied by z_ε and we perform careful a-priori estimates by extensively using the Duhamel representation formula.

The proof of Theorem 3.1 is organized as follows: in § 3.1 we review some basic results concerning viscous conservation laws. In § 3.2 we provide the proof of Theorem 3.1 by establishing precise a-priori estimates on the growth rate of the function z_ε in (3.2).

3.1. Preliminary results. In § 3.1.1 we recall some basic results about heat kernels, in § 3.1.2 we go over some a-priori estimates on solutions of viscous conservation laws that we need in the following.

3.1.1. Heat kernels. We recall some basic properties of the heat kernel $G :]0, +\infty[\times \mathbb{R}^d \rightarrow \mathbb{R}$

$$(3.3) \quad G(t, x) := C(d) \frac{1}{t^{d/2}} \exp\left(-\frac{|x|^2}{4t}\right).$$

The normalization constant $C(d)$ is chosen in such a way that $\|G(t, \cdot)\|_{L^1(\mathbb{R}^d)} = 1$, for every $t > 0$. Since

$$|\nabla G(t, x)| = C(d) \frac{|x|}{t^{d/2+1}} \exp\left(-\frac{|x|^2}{4t}\right),$$

by using spherical coordinates and by making the change of variables $\rho' = \rho/2t^{1/2}$ we get

$$\|\nabla G(t, \cdot)\|_{L^q(\mathbb{R}^d)} = C(d, q) \left(\int_0^\infty \rho^{d-1} \frac{\rho^q}{t^{q(d/2+1)}} e^{-\frac{q\rho^2}{4t}} d\rho \right)^{1/q} = C(d, q) t^{\frac{d-q(d+1)}{2q}} = C(d, q) t^\alpha.$$

For later use we have set

$$(3.4) \quad \alpha := \frac{d - q(d+1)}{2q}$$

in the formula above. Given $\nu > 0$, we introduce the kernel

$$(3.5) \quad G_\nu(t, x) := G(\nu t, x) = \frac{1}{\nu^d} G\left(\frac{t}{\nu}, \frac{x}{\nu}\right),$$

which is the fundamental solution of the equation $\partial_t u = \nu \Delta u$ and satisfies

$$(3.6) \quad \|G_\nu(t, \cdot)\|_{L^1} = 1, \quad \|\nabla G_\nu(t, \cdot)\|_{L^q} = \|\nabla G(\nu t, \cdot)\|_{L^q} = C(d, q)(\nu t)^\alpha.$$

3.1.2. A priori estimates on solutions of a viscous conservation law. The following lemma collects some classical a-priori estimates we need in the following.

Lemma 3.2. *Let $\nu \in (0, 1)$. Assume b satisfies (1.10), $\bar{u} \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$. The solution of the Cauchy problem (1.7) satisfies:*

$$(3.7) \quad \|u_\nu(t, \cdot)\|_{L^p} \leq \|\bar{u}\|_{L^p}, \quad \text{for every } t \in [0, 1] \text{ and every } p \in [1, +\infty].$$

Let u_ν and w_ν be the two solutions corresponding to the data \bar{u} and \bar{w} , respectively.

Then we have the following stability estimate: for every $p \in [1, +\infty]$,

$$(3.8) \quad \|u_\nu(t, \cdot) - w_\nu(t, \cdot)\|_{L^p} \leq e^{C(d,p,L,\|\bar{u}\|_{L^\infty},\|\bar{w}\|_{L^\infty})\nu^{-1}} \|\bar{u} - \bar{w}\|_{L^p}, \quad \text{for every } t \in [0, 1].$$

If we also require $\bar{u} \in W^{1,p}(\mathbb{R}^d)$, then we have

$$(3.9) \quad \|\nabla u_\nu(t, \cdot)\|_{L^p} \leq e^{C(d,p,L,\|\bar{u}\|_{L^\infty})\nu^{-1}} \|\nabla \bar{u}\|_{L^p}, \quad \text{for every } t \in [0, 1].$$

Remark 3.3. Note that [11, Lemma 6.3.3] implies that the function u_ν (which a priori is only defined for a.e. (t, x)) has a representative such that the function $t \mapsto u(t, \cdot)$ is continuous from $[0, 1]$ to L^1 endowed with the strong topology. Here and in the following, we always identify u_ν and its L^1 -continuous representative.

Proof of Lemma 3.2. When $p = \infty$, the estimate (3.7) is a maximum principle, which is a classical result [11, § VI]. The result for $p \in [1, +\infty[$ is also classical, but for the sake of completeness we provide a sketch of the proof. We rewrite the equation at the first line of (1.7) in the quasi-linear form

$$\partial_t u_\nu + f'(u_\nu) \operatorname{div} u_\nu = \nu \Delta u_\nu, \quad \text{where} \quad f(u) := ub(u).$$

We set $\beta(u) := |u|^p$ and we multiply the above equation by $\beta'(u_\nu)$. We arrive at

$$\partial_t [\beta(u_\nu)] + f'(u_\nu) \beta'(u_\nu) \operatorname{div} u_\nu = \nu \Delta [\beta(u_\nu)] - \nu \beta''(u_\nu) |\nabla u_\nu|^2.$$

We fix a function $h : \mathbb{R} \rightarrow \mathbb{R}^d$ satisfying $h' = f' \beta'$ and we rewrite the above equation as

$$\partial_t [\beta(u_\nu)] + \operatorname{div} [h(u_\nu)] = \nu \Delta [\beta(u_\nu)] - \nu \beta''(u_\nu) |\nabla u_\nu|^2.$$

Next, we integrate with respect to x and use the convexity of the function β : we get

$$\frac{d}{dt} \int_{\mathbb{R}^d} \beta(u_\nu) dx \leq 0,$$

which implies (3.7). To prove (3.8), we take the difference between the equation (1.7) for u_ν and w_ν

$$\partial_t (u_\nu - w_\nu) + \operatorname{div} ((u_\nu - w_\nu) b(u_\nu) + w_\nu (b(u_\nu) - b(w_\nu))) = \nu \Delta (u_\nu - w_\nu).$$

Multiplying by $p(u_\nu - w_\nu) |u_\nu - w_\nu|^{p-2}$ the previous equation and integrating in space we have

$$\partial_t \|u_\nu - w_\nu\|_{L^p}^p = \int_{\mathbb{R}^d} p(u_\nu - w_\nu) |u_\nu - w_\nu|^{p-2} \left[-\operatorname{div} ((u_\nu - w_\nu) b(u_\nu) + w_\nu (b(u_\nu) - b(w_\nu))) + \nu \Delta (u_\nu - w_\nu) \right].$$

Integrating by parts, using assumptions (1.10) on b and (3.7) with $p = \infty$ we get

$$\begin{aligned} \partial_t \|u_\nu - w_\nu\|_{L^p}^p &\leq C(p) \int_{\mathbb{R}^d} |\nabla (u_\nu - w_\nu)| |u_\nu - w_\nu|^{p-1} |b(u_\nu)| + |u_\nu - w_\nu|^{p-2} |\nabla (u_\nu - w_\nu)| |w_\nu| |b(u_\nu) - b(w_\nu)| \\ &\quad - \nu \int_{\mathbb{R}^d} |\nabla (u_\nu - w_\nu)|^2 |u_\nu - w_\nu|^{p-2} \\ &\leq C_0 \int_{\mathbb{R}^d} |\nabla (u_\nu - w_\nu)| |u_\nu - w_\nu|^{p-1} - \nu \int_{\mathbb{R}^d} |\nabla (u_\nu - w_\nu)|^2 |u_\nu - w_\nu|^{p-2}, \end{aligned}$$

for a suitable constant C_0 . By Young inequality we have

$$\int_{\mathbb{R}^d} |\nabla(u_\nu - w_\nu)| |u_\nu - w_\nu|^{p-1} \leq \frac{\nu}{2C_0} \int_{\mathbb{R}^d} |\nabla(u_\nu - w_\nu)|^2 |u_\nu - w_\nu|^{p-2} + \frac{C_0}{2\nu} \int_{\mathbb{R}^d} |u_\nu - w_\nu|^p,$$

which implies

$$\partial_t \|u_\nu - w_\nu\|_{L^p}^p \leq C\nu^{-1} \|u_\nu - w_\nu\|_{L^p}^p.$$

The Grönwall lemma allows to conclude the validity of (3.8).

By the characterization of Sobolev functions in terms of finite differences (notice that for $p = 1$ it would involve functions of bounded variation, but we know a priori that $\nabla u_\nu \in L^2$ for every $t \in [0, 1]$), we have

$$\|\nabla u_\nu(t, \cdot)\|_{L^p} \leq C \sup_{h \in \mathbb{R}^d \setminus \{0\}} \frac{1}{|h|} \|u_\nu(t, \cdot) - u_\nu(t, \cdot + h)\|_{L^p}, \quad \text{for every } t \in [0, 1].$$

Applying the stability (3.8) to \bar{u} and $\bar{u}(\cdot + h)$ we estimate the right-hand side

$$\|\nabla u_\nu(t, \cdot)\|_{L^p} \leq e^{C(d,L,\|\bar{u}\|_{L^\infty})\nu^{-1}} \sup_{h \in \mathbb{R}^d \setminus \{0\}} \frac{1}{|h|} \|\bar{u}(\cdot) - \bar{u}(\cdot + h)\|_{L^p} \leq e^{C(d,L,\|\bar{u}\|_{L^\infty})\nu^{-1}} \|\nabla \bar{u}\|_{L^p}$$

for every $t \in [0, 1]$. This proves (3.9). \square

Lemma 3.4. *Assume that b satisfies (1.10) and that u_ν satisfies (3.9). Assume furthermore that the convolution kernel η_ε satisfies (1.2) and (1.4). Then for every $p \in [1, +\infty[$ we have*

$$(3.10) \quad \|u_\nu(t, \cdot) - \eta_\varepsilon * u_\nu(t, \cdot)\|_{L^p} \leq \varepsilon e^{C(d,p,L,\|\bar{u}\|_{L^\infty})\nu^{-1}} \|\nabla \bar{u}\|_{L^p}, \quad \text{for every } t \in [0, 1].$$

Proof. By Jensen's inequality applied with respect to the probability measure $\eta_\varepsilon dx$ and by the finite differences characterization of Sobolev functions, we get

$$\begin{aligned} \|u_\nu(t, \cdot) - \eta_\varepsilon * u_\nu(t, \cdot)\|_{L^p}^p &\stackrel{(1.2),(1.4)}{=} \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} [u_\nu(t, x-y) - u_\nu(t, x)] \eta_\varepsilon(y) dy \right|^p dx \\ &\leq \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |u_\nu(t, x-y) - u_\nu(t, x)|^p \eta_\varepsilon(y) dy dx \\ &\leq \|\nabla u_\nu(t, \cdot)\|_{L^p(\mathbb{R}^d)}^p \int_{\mathbb{R}^d} \eta_\varepsilon(y) |y|^p dy \\ &\stackrel{(3.9)}{\leq} \varepsilon^p e^{C(d,p,L,\|\bar{u}\|_{L^\infty})\nu^{-1}} \|\nabla \bar{u}\|_{L^p(\mathbb{R}^d)}^p \int_{\mathbb{R}^d} \eta_\varepsilon(y) \left(\frac{|y|}{\varepsilon}\right)^p dy. \end{aligned}$$

Since η_ε is supported where $|y| \leq \varepsilon$, the last integrand in the right-hand side is estimated by $\|\eta_\varepsilon\|_{L^1} = 1$, which concludes the proof of (3.10). \square

3.2. Proof of Theorem 3.1. First, we recall that $u_{\varepsilon\nu}$ is the solution of (1.6) and u_ν is the solution of (1.7) and we define z_ε as in (3.2). Note that z_ε satisfies the equation

$$\partial_t z_\varepsilon + \operatorname{div} \mathcal{T}_\varepsilon = \nu \Delta z_\varepsilon$$

where, thanks to (3.2), the term \mathcal{T}_ε is given by

$$(3.11) \quad \begin{aligned} \mathcal{T}_\varepsilon &:= u_{\varepsilon\nu} b(u_{\varepsilon\nu} * \eta_\varepsilon) - u_\nu b(u_\nu) = u_{\varepsilon\nu} [b(u_{\varepsilon\nu} * \eta_\varepsilon) - b(u_\nu)] + [u_{\varepsilon\nu} - u_\nu] b(u_\nu) \\ &= [z_\varepsilon + u_\nu] [b([z_\varepsilon + u_\nu] * \eta_\varepsilon) - b(u_\nu)] + z_\varepsilon b(u_\nu). \end{aligned}$$

We now proceed as follows: in § 3.2.1 we establish some a-priori estimates on z_ε , which are the key point in the proof, and in § 3.2.2 we conclude the proof of Theorem 3.1.

3.2.1. *A-priori estimates on z_ε .* We establish a-priori estimates on the solution of the Cauchy problem

$$(3.12) \quad \begin{cases} \partial_t z_\varepsilon + \operatorname{div} \mathcal{T}_\varepsilon = \nu \Delta z_\varepsilon \\ z_\varepsilon(0, x) = z_0(x). \end{cases}$$

Lemma 3.5. *Let b satisfy (1.10), $0 < \varepsilon, \nu \leq 1/4$, η_ε as in (1.2) and (1.4), $z_0 \in L^p(\mathbb{R}^d)$ with p as in (1.8), and $\beta = (p+d)/(p-d)$. Assume furthermore that the function u_ν satisfies (3.7) and (3.10). Then there exist $c_0 := c_0(d, p, L, \|\bar{u}\|_{L^\infty}) > 0$ and $\tau_0 := \tau_0(d, p, L, \|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{W^{1,p}}) > 0$, such that if*

$$(3.13) \quad \|z_0\|_{L^p} \leq 1/4, \quad \varepsilon \leq e^{-c_0\nu^{-1}}/4$$

the solution of the Cauchy problem (3.12) starting from z_0 satisfies

$$(3.14) \quad \|z_\varepsilon(t, \cdot)\|_{L^p} \leq 2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon], \quad \text{for every } t \in [0, \tau_0\nu^\beta].$$

Proof. Let c_0 be the maximum between the constant in Lemma 3.4 and 1. Set

$$(3.15) \quad \tau := \sup \{t \in [0, 1] : \|z_\varepsilon(s, \cdot)\|_{L^p} \leq 2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon], \text{ for every } s \in [0, t]\}.$$

Owing to Remarks 2.2 and 3.3 and to (2.1) and (3.7), the functions $u_{\varepsilon\nu}$ and u_ν are continuous from $[0, +\infty[$ to L^p . Hence, the function $\|z_\varepsilon(t, \cdot)\|_{L^p}$ is continuous and $\tau > 0$. Moreover, (3.15) implies

$$(3.16) \quad \|z_\varepsilon(\tau, \cdot)\|_{L^p} = 2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon].$$

We represent the solution of the Cauchy problem (3.12) by the Duhamel Principle as

$$z_\varepsilon(\tau, \cdot) = G_\nu(\tau, \cdot) * z_0 - \int_0^\tau \int_{\mathbb{R}^d} \nabla G_\nu(\tau - s, \cdot - y) \cdot \mathcal{T}_\varepsilon(s, y) dy ds$$

where G_ν denotes the heat kernel (3.5). We apply (3.6) and the Bochner and Young Theorems to get

$$(3.17) \quad \begin{aligned} \|z_\varepsilon(\tau, \cdot)\|_{L^p} &\leq \|G_\nu(\tau, \cdot)\|_{L^1} \|z_0\|_{L^p} + \int_0^\tau \left\| \int_{\mathbb{R}^d} \nabla G_\nu(\tau - s, \cdot) \cdot \mathcal{T}_\varepsilon(s, \cdot) \right\|_{L^p} ds \\ &\leq \|z_0\|_{L^p} + \int_0^\tau \|\nabla G_\nu(\tau - s, \cdot)\|_{L^q} \|\mathcal{T}_\varepsilon(s, \cdot)\|_{L^{p/2}} ds, \end{aligned}$$

noting that $p/2 \geq 1$ owing to (1.8), and setting $q := p/(p-1)$. Only in the rest of this proof we denote by C any constant that only depends on $d, p, L, \|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{W^{1,p}}$. For every $s \in [0, \tau]$, by (3.11), the Hölder inequality and (1.10) we have

$$\begin{aligned} \|\mathcal{T}_\varepsilon(s, \cdot)\|_{L^{p/2}} &\leq (\|z_\varepsilon\|_{L^p} + \|u_\nu\|_{L^p}) \|b([z_\varepsilon + u_\nu] * \eta_\varepsilon) - b(u_\nu)\|_{L^p} + \|z_\varepsilon\|_{L^p} \|b(u_\nu)\|_{L^p} \\ &\leq C(\|z_\varepsilon\|_{L^p} + \|u_\nu\|_{L^p}) \|([z_\varepsilon + u_\nu] * \eta_\varepsilon) - u_\nu\|_{L^p} + C\|z_\varepsilon\|_{L^p} \|u_\nu\|_{L^p} \end{aligned}$$

(all functions at the right hand side are evaluated at time s). By the Young inequality we have $\|z_\varepsilon * \eta_\varepsilon\|_{L^p} \leq \|z_\varepsilon\|_{L^p}$, and applying also (3.10) we get

$$(3.18) \quad \|\mathcal{T}_\varepsilon(s, \cdot)\|_{L^{p/2}} \leq C(\|z_\varepsilon(s, \cdot)\|_{L^p} + \|u_\nu(s, \cdot)\|_{L^p}) [\|z_\varepsilon(s, \cdot)\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon].$$

We recall that $s \leq \tau$ and so (3.14) holds. Also, we recall (3.4) and (3.6) and we point out that $\alpha = -(d+p)/(2p) \in (-1, 0)$ by (1.8). Using this and (3.18), we go back to (3.17) to get

$$(3.19) \quad \begin{aligned} \|z_\varepsilon(\tau, \cdot)\|_{L^p} &\leq \|z_0\|_{L^p} + C\nu^\alpha [2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] + 1] [2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] + e^{c_0\nu^{-1}}\varepsilon] \tau^{\alpha+1} \\ &\leq \|z_0\|_{L^p} + 6C\nu^\alpha [\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] \tau^{\alpha+1}, \end{aligned}$$

where in the last inequality we used (3.13) to show that $2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] + 1 \leq 2$. By comparing (3.19) with (3.16) we arrive at

$$2[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] \leq \|z_0\|_{L^p} + 6C\nu^\alpha [\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] \tau^{\alpha+1},$$

which implies

$$[\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] \leq 6C\nu^\alpha [\|z_0\|_{L^p} + e^{c_0\nu^{-1}}\varepsilon] \tau^{\alpha+1} \implies (6C)^{-1} \nu^{-\alpha} \leq \tau^{\alpha+1}.$$

This gives a lower bound on τ and concludes the proof of the Lemma by choosing $\tau_0 = (6C)^{-\frac{1}{\alpha+1}}$. \square

3.2.2. *Conclusion of the proof of Theorem 3.1.* Let $0 < \nu < 1/4$, and let τ_0 and c_0 be as in the statement of Lemma 3.5. Let $m := \text{int}((\tau_0\nu^\beta)^{-1}) + 1$, where $\text{int}(\cdot)$ denotes the integer part, and let $\varepsilon \leq e^{-2c_0\nu^{-1}}4^{-m-1}$. This is implied for instance by

$$\varepsilon \leq e^{-c\nu^{-\beta}} \quad \text{for } c := c(d, p, L, \|\bar{u}\|_{L^\infty}, \|\bar{u}\|_{W^{1,p}}) > 0.$$

We show by induction that for every $i = 1, \dots, m$

$$(3.20) \quad \|z_\varepsilon(t, \cdot)\|_{L^p} \leq e^{c_0\nu^{-1}}\varepsilon 4^i \quad \text{for every } t \in [(i-1)\tau_0\nu^\beta, i\tau_0\nu^\beta].$$

Indeed, by (1.6), (1.7) and (3.2) we have $z_\varepsilon(0, x) \equiv 0$; we apply estimate (3.14) on $[0, \tau_0\nu^\beta]$ to get (3.20) with $i = 1$. If the statement holds true for i , we have that $\|z_\varepsilon(i\tau_0\nu^\beta, \cdot)\|_{L^p} \leq e^{c_0\nu^{-1}}\varepsilon 4^i \leq e^{-c_0\nu^{-1}}/4$; to get the statement for $i + 1$, we apply estimate (3.14) with $z_0 := z_\varepsilon(i\tau_0\nu^\beta, \cdot)$, obtaining

$$\|z_\varepsilon(t, \cdot)\|_{L^p} \leq 2e^{c_0\nu^{-1}}[\varepsilon 4^i + \varepsilon] \leq 4^{i+1}e^{c_0\nu^{-1}}\varepsilon \quad \text{for every } t \in [i\tau_0\nu^\beta, (i+1)\tau_0\nu^\beta].$$

This establishes (3.1) and concludes the proof of Theorem 3.1. \square

4. PROOF OF THEOREM 1.1

We first explain the basic ideas of the proof. We need some notation: we term

$$(4.1) \quad S_t^{\varepsilon\nu} : L^1 \cap L^\infty \times [0, +\infty[\rightarrow L^1 \cap L^\infty, \quad S_t^\nu : L^1 \cap L^\infty \times [0, +\infty[\rightarrow L^1 \cap L^\infty$$

the semigroup of solutions of the equations at the first line of (1.6) and (1.7), respectively. In other words, $u_{\varepsilon\nu}(t, \cdot) = S_t^{\varepsilon\nu}\bar{u}$ and $u_\nu(t, \cdot) = S_t^\nu\bar{u}$. Next, we fix $\mathbf{d} \in L^1 \cap L^\infty$ and a regularity parameter $0 < \lambda < 1$ and we decompose \mathbf{d} as

$$(4.2) \quad \mathbf{d} := \mathbf{d}_r + \mathbf{d}_s, \quad \text{where} \quad \mathbf{d}_r := \mathbf{d} * \rho_\lambda, \quad \mathbf{d}_s = \mathbf{d} - \mathbf{d} * \rho_\lambda.$$

In the previous expression ρ_λ is a given standard family of convolution kernels, obtained by setting $\rho_\lambda(x) := \lambda^{-d}\rho(x/\lambda)$ for a standard (i.e., smooth, positive, radial, compactly supported, and with unit integral) convolution kernel ρ , with $\|\rho\|_{C^1(\mathbb{R}^d)} \leq C(d)$.

Note that \mathbf{d}_r is regular, and hence we can apply Theorem 3.1 to show that $S_t^{\varepsilon\nu}\mathbf{d}_r$ converges to $S_t^\nu\mathbf{d}_r$, with a convergence rate that deteriorates when $\lambda \rightarrow 0^+$. Also, we can choose the regularizing parameter λ in such a way that $\mathbf{d}_s = \mathbf{d} - \mathbf{d}_r$ is small. The basic point in the proof of Theorem 1.1 is then establishing a uniform control on the growth of $\|S_t^{\varepsilon\nu}\mathbf{d} - S_t^{\varepsilon\nu}\mathbf{d}_r\|_{L^p}$. This is done in § 4.1 below. Next, in § 4.2 we establish some stability estimates and in § 4.3 we conclude the proof by using an iteration argument.

4.1. **Perturbations estimates.** We begin by establishing some perturbation estimates.

Lemma 4.1. *Fix p satisfying (1.8), $\mathbf{d} \in L^1 \cap L^\infty$, and let \mathbf{d}_r and \mathbf{d}_s be as in (4.2). Assume that*

$$(4.3) \quad \|\mathbf{d}_s\|_{L^p} \leq \delta \leq 1,$$

$$(4.4) \quad \|\mathbf{d}\|_{L^p} \leq D, \quad \|\mathbf{d}\|_{L^\infty} \leq B$$

for some positive constants $D > 0$, $B > 0$. Then there are constants $\bar{\varepsilon}(d, p, L, B, D, \nu, \lambda)$ and $\sigma = \sigma(d, p, L, D, \nu)$ such that, if $\varepsilon \leq \bar{\varepsilon}$, then

$$\|S_t^{\varepsilon\nu}\mathbf{d} - S_t^{\varepsilon\nu}\mathbf{d}_r\|_{L^p} \leq 2\delta, \quad \text{for every } t \in [0, \sigma].$$

Proof. We set

$$(4.5) \quad v_\varepsilon := S_t^{\varepsilon\nu}\mathbf{d} - S_t^{\varepsilon\nu}\mathbf{d}_r$$

and we point out that v_ε is a solution of the Cauchy problem

$$\begin{cases} \partial_t v_\varepsilon + \text{div} [v_\varepsilon b(S_t^{\varepsilon\nu}\mathbf{d} * \eta_\varepsilon) + S_t^{\varepsilon\nu}\mathbf{d}_r (b(S_t^{\varepsilon\nu}\mathbf{d} * \eta_\varepsilon) - b(S_t^{\varepsilon\nu}\mathbf{d}_r * \eta_\varepsilon))] = \nu \Delta v_\varepsilon \\ v_\varepsilon(0, x) = \mathbf{d}_s(x). \end{cases}$$

We introduce σ by setting

$$\sigma := \sup \{t \in [0, 1] : \|v_\varepsilon(s, \cdot)\|_{L^p} \leq 2\delta \text{ for every } s \in [0, t]\}.$$

Note that, if $\sigma < 1$, we have

$$(4.6) \quad \|v_\varepsilon(\sigma, \cdot)\|_{L^p} = 2\delta.$$

We now provide a lower bound on σ . By using the Duhamel representation formula we get

$$v_\varepsilon(t, \cdot) = G_\nu(t, \cdot) * \mathbf{d}_s - \int_0^t \int_{\mathbb{R}^d} \nabla G_\nu(t-s, \cdot - y) \cdot \left[v_\varepsilon b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon) + S_t^{\varepsilon\nu} \mathbf{d}_r [b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon) - b(S_t^{\varepsilon\nu} \mathbf{d}_r * \eta_\varepsilon)] \right] (s, y) dy ds.$$

We fix $q := p/(p-1)$ and $\alpha > -1$ given by (3.4). Applying the Bochner and Young Theorems we get

$$(4.7) \quad \begin{aligned} \|v_\varepsilon(t, \cdot)\|_{L^p} &\leq \|G_\nu(t, \cdot)\|_{L^1} \|\mathbf{d}_s\|_{L^p} \\ &\quad + \int_0^t \|\nabla G_\nu(t-s, \cdot)\|_{L^q} \left[\|v_\varepsilon b(S_t^\varepsilon \mathbf{d} * \eta_\varepsilon)(s, \cdot) + S_t^\varepsilon \mathbf{d}_r [b(S_t^\varepsilon \mathbf{d} * \eta_\varepsilon) - b(S_t^\varepsilon \mathbf{d}_r * \eta_\varepsilon)](s, \cdot)\|_{L^{p/2}} \right] ds \\ &\stackrel{(3.6), (4.3)}{\leq} \delta + C(d, p, \nu) \int_0^t (t-s)^\alpha \left[\|v_\varepsilon b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)(s, \cdot)\|_{L^{p/2}} \right. \\ &\quad \left. + \|S_t^{\varepsilon\nu} \mathbf{d}_r [b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon) - b(S_t^{\varepsilon\nu} \mathbf{d}_r * \eta_\varepsilon)](s, \cdot)\|_{L^{p/2}} \right] ds. \end{aligned}$$

Next, by the Hölder inequality, (1.10), (4.5) and the Young inequality to get

$$(4.8) \quad \begin{aligned} &\|v_\varepsilon b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)\|_{L^{p/2}} + \|S_t^{\varepsilon\nu} \mathbf{d}_r [b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon) - b(S_t^{\varepsilon\nu} \mathbf{d}_r * \eta_\varepsilon)]\|_{L^{p/2}} \\ &\leq \|v_\varepsilon\|_{L^p} \|b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)\|_{L^p} + \|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \|b(S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon) - b(S_t^{\varepsilon\nu} \mathbf{d}_r * \eta_\varepsilon)\|_{L^p} \\ &\leq L \|v_\varepsilon\|_{L^p} \|S_t^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon\|_{L^p} + L \|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \|v_\varepsilon * \eta_\varepsilon\|_{L^p} \\ &\leq L \|v_\varepsilon\|_{L^p} \|S_t^{\varepsilon\nu} \mathbf{d}\|_{L^p} + L \|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \|v_\varepsilon\|_{L^p} \\ &\leq L \|v_\varepsilon\|_{L^p} [\|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} + \|v_\varepsilon\|_{L^p}] + L \|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \|v_\varepsilon\|_{L^p} \\ &\leq 2L \|v_\varepsilon\|_{L^p} [\|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} + \|v_\varepsilon\|_{L^p}]. \end{aligned}$$

We recall the definition (4.2) of \mathbf{d}_r and we point out that \mathbf{d}_r is smooth and henceforth satisfies the hypotheses of Theorem 3.1. By applying (3.1) and (3.7), we get

$$(4.9) \quad \|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \leq \|S_t^{\varepsilon\nu} \mathbf{d}_r - S_t^\nu \mathbf{d}_r\|_{L^p} + \|S_t^\nu \mathbf{d}_r\|_{L^p} \leq C(d, p, L, \|\mathbf{d}_r\|_{L^\infty}, \|\mathbf{d}_r\|_{W^{1,p}}, \nu) \varepsilon + \|\mathbf{d}_r\|_{L^p}.$$

Since $\mathbf{d}_r = \mathbf{d} * \rho_\lambda$, by (4.4) we have

$$\|\mathbf{d}_r\|_{L^\infty} \leq \|\mathbf{d}\|_{L^\infty} \leq B, \quad \|\mathbf{d}_r\|_{L^p} \leq \|\mathbf{d}\|_{L^p} \leq D, \quad \|\nabla \mathbf{d}_r\|_{L^p} \leq \|\mathbf{d}\|_{L^p} \|\nabla \rho_\lambda\|_{L^1} \leq C(d, D, \lambda)$$

and hence (4.9) implies

$$\|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \leq C(d, p, L, B, D, \nu, \lambda) \varepsilon + D,$$

so if $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \nu, \lambda)$ is sufficiently small, then

$$(4.10) \quad \|S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \leq 3D/2.$$

We combine (4.6), (4.7), (4.8) and (4.10) and we get that

$$\begin{aligned} \|v_\varepsilon(\sigma, \cdot)\|_{L^p} &= 2\delta \leq \delta + C(d, L, p, \nu) \int_0^\sigma (\sigma - s)^\alpha \left[\|v_\varepsilon\|_{L^p} [D + \|v_\varepsilon\|_{L^p}] \right] (s, \cdot) ds \\ &\stackrel{s \leq \sigma}{\leq} \delta + C(d, L, p, \nu) \int_0^\sigma (\sigma - s)^\alpha \left[2\delta [D + 2\delta] \right] (s, \cdot) ds = \left[1 + C(d, L, p, \nu) \sigma^{\alpha+1} [D + 2\delta] \right] \delta \\ &\stackrel{\delta \leq 1}{\leq} \left[1 + C(d, L, p, \nu) \sigma^{\alpha+1} [D + 2] \right] \delta. \end{aligned}$$

The above chain of inequalities implies that $1 \leq C(d, L, p, \nu)\sigma^{\alpha+1}[D+2]$ and this provides a lower bound on σ that only depends on d, p, L, ν and D . \square

4.2. Stability estimates. We now establish a conditional stability estimate.

Lemma 4.2. *Fix $\mathbf{d}_1, \mathbf{d}_2 \in L^1 \cap L^\infty$ and p satisfying (1.8) and assume there are constants $F > 0$ and $T > 0$ such that*

$$(4.11) \quad \|S_t^{\varepsilon\nu} \mathbf{d}_1\|_{L^p}, \|S_t^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \leq F, \quad \text{for every } t \in [0, T].$$

Then there is a threshold $\varpi = \varpi(L, F, d, p, \nu) \in]0, T]$ such that

$$\|S_t^{\varepsilon\nu} \mathbf{d}_1 - S_t^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \leq 2\|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p}, \quad \text{for every } t \in [0, \varpi].$$

Proof. We use the Duhamel Representation Formula and get

$$\begin{aligned} S_t^{\varepsilon\nu} \mathbf{d}_1 - S_t^{\varepsilon\nu} \mathbf{d}_2 &= [\mathbf{d}_1 - \mathbf{d}_2] * G_\nu(t, \cdot) \\ &\quad - \int_0^t \int_{\mathbb{R}^d} \nabla G_\nu(t-s, \cdot - y) [S_s^{\varepsilon\nu} \mathbf{d}_1 b(S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon) - S_s^{\varepsilon\nu} \mathbf{d}_2 b(S_s^{\varepsilon\nu} \mathbf{d}_2 * \eta_\varepsilon)](y) dy ds, \end{aligned}$$

which owing to the Bochner and Young Theorems implies

$$(4.12) \quad \begin{aligned} \|S_t^{\varepsilon\nu} \mathbf{d}_1 - S_t^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} &\leq \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} \|G_\nu(t, \cdot)\|_{L^1} \\ &\quad + \int_0^t \|\nabla G_\nu(t-s, \cdot)\|_{L^q} \|S_s^{\varepsilon\nu} \mathbf{d}_1 b(S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon) - S_s^{\varepsilon\nu} \mathbf{d}_2 b(S_s^{\varepsilon\nu} \mathbf{d}_2 * \eta_\varepsilon)\|_{L^{p/2}} ds \end{aligned}$$

provided $q := p/(p-1)$. By the Hölder inequality we get

$$(4.13) \quad \begin{aligned} &\|S_s^{\varepsilon\nu} \mathbf{d}_1 b(S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon) - S_s^{\varepsilon\nu} \mathbf{d}_2 b(S_s^{\varepsilon\nu} \mathbf{d}_2 * \eta_\varepsilon)\|_{L^{p/2}} \\ &\leq \|S_s^{\varepsilon\nu} \mathbf{d}_1 b(S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon) - S_s^{\varepsilon\nu} \mathbf{d}_2 b(S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon)\|_{L^{p/2}} + \|S_s^{\varepsilon\nu} \mathbf{d}_2 b(S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon) - S_s^{\varepsilon\nu} \mathbf{d}_2 b(S_s^{\varepsilon\nu} \mathbf{d}_2 * \eta_\varepsilon)\|_{L^{p/2}} \\ &\stackrel{\text{Hölder, (1.10)}}{\leq} L \|S_s^{\varepsilon\nu} \mathbf{d}_1 - S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \|S_s^{\varepsilon\nu} \mathbf{d}_1 * \eta_\varepsilon\|_{L^p} + L \|S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \| [S_s^{\varepsilon\nu} \mathbf{d}_1 - S_s^{\varepsilon\nu} \mathbf{d}_2] * \eta_\varepsilon \|_{L^p} \\ &\stackrel{\text{Young, (1.2)}}{\leq} L \|S_s^{\varepsilon\nu} \mathbf{d}_1 - S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \|S_s^{\varepsilon\nu} \mathbf{d}_1\|_{L^p} + L \|S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \|S_s^{\varepsilon\nu} \mathbf{d}_1 - S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \\ &\stackrel{(4.11)}{\leq} C(L, F) \|S_s^{\varepsilon\nu} \mathbf{d}_1 - S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p}. \end{aligned}$$

We now introduce the value ϖ by setting

$$\varpi := \sup \{t \in [0, 1] : \|S_s^{\varepsilon\nu} \mathbf{d}_1 - S_s^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \leq 2\|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} \text{ for every } s \in [0, t]\}.$$

Note that

$$(4.14) \quad \|S_\varpi^{\varepsilon\nu} \mathbf{d}_1 - S_\varpi^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} = 2\|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p}.$$

Also, by combining (4.12), (3.6) and (4.13) we get that

$$(4.15) \quad \begin{aligned} \|S_\varpi^{\varepsilon\nu} \mathbf{d}_1 - S_\varpi^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} &\leq \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} + C(L, F) \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} \int_0^\varpi \|\nabla G_\nu(\varpi - s, \cdot)\|_{L^q} ds \\ &\stackrel{(3.6), (3.4)}{\leq} \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} + C(L, F, p, d, \nu) \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} \int_0^\varpi (\varpi - s)^\alpha ds \\ &\leq \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} [1 + C(L, F, p, d, \nu) \varpi^{\alpha+1}]. \end{aligned}$$

By comparing (4.15) with (4.14) we get

$$2\|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p} \leq [1 + C(L, F, d, p, \nu) \varpi^{\alpha+1}] \|\mathbf{d}_1 - \mathbf{d}_2\|_{L^p}$$

and this provides a lower bound on ϖ . \square

We conclude this paragraph by establishing a uniform a-priori estimate on the growth of \mathbf{d} .

Lemma 4.3. *Assume that $\mathbf{d} \in L^\infty \cap L^1$ and that*

$$\|\mathbf{d}\|_{L^p} \leq Q.$$

Then there is a constant $\theta = \theta(d, p, L, Q) > 0$ such that

$$\|S_t^{\varepsilon\nu} \mathbf{d}\|_{L^p} \leq 2Q, \quad \text{for every } t \in [0, \theta].$$

Proof. We set

$$\theta := \sup \{t \in [0, 1] : \|S_s^{\varepsilon\nu} \mathbf{d}\|_{L^p} \leq 2Q, \text{ for every } s \in [0, t]\}$$

and we point out that

$$(4.16) \quad \|S_\theta^{\varepsilon\nu} \mathbf{d}\|_{L^p} = 2Q.$$

To establish a lower bound on θ we use the Duhamel representation formula. We have

$$S_\theta^{\varepsilon\nu} \mathbf{d} = \mathbf{d} * G_\nu(\theta, \cdot) - \int_0^\theta \int_{\mathbb{R}^d} \nabla G_\nu(\theta - s, \cdot - y) \cdot [S_s^{\varepsilon\nu} \mathbf{d} b(S_s^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)](y) dy ds.$$

We use the Bochner and Young Theorems and we get

$$\|S_\theta^{\varepsilon\nu} \mathbf{d}\|_{L^p} \leq \|\mathbf{d}\|_{L^p} + \int_0^\theta \|\nabla G_\nu(\theta - s, \cdot)\|_{L^q} \|S_s^{\varepsilon\nu} \mathbf{d} b(S_s^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)\|_{L^{p/2}} ds,$$

provided $q := p/(p-1)$. Next, by Hölder inequality, (1.10), and since $s \leq \theta$ we infer that

$$\|S_s^{\varepsilon\nu} \mathbf{d} b(S_s^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)\|_{L^{p/2}} \leq \|S_s^{\varepsilon\nu} \mathbf{d}\|_{L^p} \|b(S_s^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon)\|_{L^p} \leq L \|S_s^{\varepsilon\nu} \mathbf{d}\|_{L^p} \|S_s^{\varepsilon\nu} \mathbf{d} * \eta_\varepsilon\|_{L^p} \leq L \|S_s^{\varepsilon\nu} \mathbf{d}\|_{L^p}^2 \leq 4LQ^2.$$

We let $\alpha > -1$ be as in (3.4). By (3.6) and the above inequalities we infer that

$$\|S_\theta^{\varepsilon\nu} \mathbf{d}\|_{L^p} \leq Q + C(d, p, L)\theta^{\alpha+1}Q^2$$

and by comparing the above inequality with (4.16) we establish a lower bound on θ . \square

4.3. Conclusion of the proof of Theorem 1.1. We first introduce some notation. First, we fix a parameter $0 < h < 1$. We set

$$D := \|\bar{u}\|_{L^p}, \quad B := \|\bar{u}\|_{L^\infty}, \quad F := 4D, \quad Q := 2D$$

and choose a threshold $\xi = \xi(d, p, L, D, \nu, Q)$ in such a way that

$$\xi := \min\{\sigma, \varpi, \theta\},$$

where σ , ϖ and θ are as in the statement of Lemma 4.1, Lemma 4.2 and Lemma 4.3, respectively.

STEP 1: we choose $\mathbf{d} := \bar{u}$ and the regularity parameter λ in (4.2) (depending only on p , \bar{u} and h) in such a way that

$$\|\mathbf{d}_s\|_{L^p} = \|\mathbf{d} - \mathbf{d} * \rho_\lambda\|_{L^p} \leq h < 1.$$

We establish convergence on the interval $[0, \xi]$. First we decompose \bar{u} as in (4.2). Note that

$$\|\mathbf{d}_r\|_{L^p} \leq \|\mathbf{d}\|_{L^p} \leq D, \quad \|\mathbf{d}_r\|_{L^\infty} \leq \|\mathbf{d}\|_{L^\infty} \leq B.$$

Next, we fix $t \in [0, \xi]$ and we introduce the following decomposition:

$$(4.17) \quad \|S_t^{\varepsilon\nu} \mathbf{d} - S_t^\nu \mathbf{d}\|_{L^p} \leq \|S_t^{\varepsilon\nu} \mathbf{d} - S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} + \|S_t^{\varepsilon\nu} \mathbf{d}_r - S_t^\nu \mathbf{d}_r\|_{L^p} + \|S_t^\nu \mathbf{d}_r - S_t^\nu \mathbf{d}\|_{L^p} =: T_1 + T_2 + T_3.$$

To control the term T_1 , we apply Lemma 4.1 and we infer that, if $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \nu, \lambda)$, then

$$(4.18) \quad \|S_t^{\varepsilon\nu} \mathbf{d} - S_t^{\varepsilon\nu} \mathbf{d}_r\|_{L^p} \leq 2h.$$

To control the term T_2 , we apply Theorem 3.1. First, we point out that

$$\nabla \mathbf{d}_r = \mathbf{d} * \nabla \rho_\lambda \implies \|\nabla \mathbf{d}_r\|_{L^p} \leq \|\bar{u}\|_{L^p} \|\nabla \rho_\lambda\|_{L^1} = C(d, D, \lambda).$$

By applying Theorem 3.1 we arrive at

$$(4.19) \quad \|S_t^{\varepsilon\nu} \mathbf{d}_r - S_t^\nu \mathbf{d}_r\|_{L^p} \leq C(d, p, L, B, D, \nu, \lambda) \varepsilon \leq h$$

provided that $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \lambda, \nu, h)$. Finally, to control the term T_3 we apply (3.8) and we get

$$(4.20) \quad \|S_t^\nu \mathbf{d}_r - S_t^\nu \mathbf{d}\|_{L^p} \leq C(d, p, L, B, \nu) \|\mathbf{d}_r - \mathbf{d}\|_{L^p} \leq C(d, p, L, B, \nu) h.$$

By combining (4.18), (4.19) and (4.20) with (4.17) we eventually get that

$$(4.21) \quad \|S_t^{\varepsilon\nu} \mathbf{d} - S_t^\nu \mathbf{d}\|_{L^p} \leq C(d, p, L, B, \nu) h$$

provided that $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \lambda, \nu, h)$.

STEP 2: we establish convergence on the interval $[\xi, 2\xi]$. First, we fix $t \in [0, \xi]$ and we introduce the following decomposition:

$$(4.22) \quad \begin{aligned} \|S_{t+\xi}^{\varepsilon\nu} \bar{u} - S_{t+\xi}^\nu \bar{u}\|_{L^p} &= \|S_t^{\varepsilon\nu} S_\xi^{\varepsilon\nu} \bar{u} - S_t^\nu S_\xi^\nu \bar{u}\|_{L^p} \\ &\leq \|S_t^{\varepsilon\nu} S_\xi^{\varepsilon\nu} \bar{u} - S_t^{\varepsilon\nu} S_\xi^\nu \bar{u}\|_{L^p} + \|S_t^{\varepsilon\nu} S_\xi^\nu \bar{u} - S_t^\nu S_\xi^\nu \bar{u}\|_{L^p} =: S_1 + S_2. \end{aligned}$$

To control the term S_1 we apply Lemma 4.2. First, we set

$$\mathbf{d}_1 := S_\xi^{\varepsilon\nu} \bar{u}, \quad \mathbf{d}_2 := S_\xi^\nu \bar{u}$$

and we recall that $F = 4\|\bar{u}\|_{L^p}$. Now we want to show that (4.11) holds true: we do this by applying Lemma 4.3. First, we check that

$$(4.23) \quad \|S_t^{\varepsilon\nu} \mathbf{d}_2\|_{L^p} \leq F.$$

We recall that $Q = 2\|\bar{u}\|_{L^p}$ and we point out that, owing to (3.7), we have

$$\|\mathbf{d}_2\|_{L^p} \leq \|\bar{u}\|_{L^p} \leq Q.$$

By applying Lemma 4.3, we get (4.23). Next, by (4.21) and (3.7) we point out that

$$\|\mathbf{d}_1\|_{L^p} \leq \|S_\xi^{\varepsilon\nu} \bar{u} - S_\xi^\nu \bar{u}\|_{L^p} + \|S_\xi^\nu \bar{u}\|_{L^p} \leq C(d, p, L, B, \nu) h + \|\bar{u}\|_{L^p} \leq 2\|\bar{u}\|_{L^p} = Q$$

provided that h is sufficiently small. By applying Lemma 4.3, we get $\|S_t^{\varepsilon\nu} \mathbf{d}_1\|_{L^p} \leq F$ and by recalling (4.23) we conclude that (4.11) is satisfied. By applying Lemma 4.2 we conclude that

$$S_1 \stackrel{(4.22)}{=} \|S_t^{\varepsilon\nu} S_\xi^{\varepsilon\nu} \bar{u} - S_t^{\varepsilon\nu} S_\xi^\nu \bar{u}\|_{L^p} \leq 2\|S_\xi^{\varepsilon\nu} \bar{u} - S_\xi^\nu \bar{u}\|_{L^p} \stackrel{(4.21)}{\leq} C(d, p, L, B, \nu) h,$$

provided that $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \lambda, \nu, h)$.

We now control S_2 , the second term in (4.22). We set $\mathbf{d} := S_\xi^\nu \bar{u}$ and we point out that

$$\|S_\xi^\nu \bar{u}\|_{L^p} \leq D, \quad \|S_\xi^\nu \bar{u}\|_{L^\infty} \leq B$$

owing to (3.7). By applying the same argument as in STEP 1 we conclude that

$$S_2 = \|S_t^{\varepsilon\nu} S_\xi^\nu \bar{u} - S_t^\nu S_\xi^\nu \bar{u}\|_{L^p} \leq C(d, p, L, B, \nu) h$$

provided that $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \lambda, \nu, h)$. By recalling (4.22), this establishes the convergence on the interval $[\xi, 2\xi]$.

STEP 3: by iterating the argument at STEP 2 a finite number of times we can prove that

$$\|S_t^{\varepsilon\nu} \bar{u} - S_t^\nu \bar{u}\|_{L^p} \leq C(d, p, L, B, \nu) h, \quad \text{for every } t \in [0, 1]$$

provided that $\varepsilon \leq \bar{\varepsilon}(d, p, L, B, D, \lambda, \nu, h)$. This establishes the strong L^p convergence and concludes the proof of Theorem 1.1. \square

5. COUNTER-EXAMPLES

In this section we focus on the family of Cauchy problems in one space dimension

$$(5.1) \quad \begin{cases} \partial_t u_\varepsilon + \partial_x [u_\varepsilon u_\varepsilon * \eta_\varepsilon] = 0 \\ u_\varepsilon(0, \cdot) = \bar{u}, \end{cases}$$

which is exactly (1.3) in the case when $d = 1$ and $b(u) = u$. When $\varepsilon \rightarrow 0^+$, the Cauchy problem in (5.1) *formally* reduces to the Cauchy problem for the Burgers' equation

$$(5.2) \quad \begin{cases} \partial_t u + \partial_x [u^2] = 0 \\ u(0, \cdot) = \bar{u}. \end{cases}$$

In this section we provide three explicit counter-examples showing that, in general, u_ε does not converge to the entropy admissible solution u .

5.1. A counter-example with sign-changing data and symmetric kernels. We begin by stating and proving our first counter-example.

Lemma 5.1. *Assume that η_ε satisfies (1.2) and (1.4) and that η is an even function, namely $\eta(x) = \eta(-x)$ for every x . Assume furthermore that the initial datum $\bar{u} \in BV(\mathbb{R})$ is an odd function, namely $\bar{u}(x) = -\bar{u}(-x)$ for a.e. x , and such that*

$$(5.3) \quad \bar{u}(x) := \begin{cases} 1 & -1 < x < 0 \\ -1 & 0 < x < 1 \\ 0 & |x| > 2. \end{cases}$$

Let u_ε be the solution of (5.1) and u be the entropy admissible solution of (5.2). Then

$$(5.4) \quad \int_{-\infty}^0 u(t, x) dx < \int_{-\infty}^0 \bar{u}(x) dx = \int_{-\infty}^0 u_\varepsilon(t, x) dx, \quad \text{for every } t \in [0, 1/4[.$$

In particular, the family $\{u_\varepsilon\}_{\varepsilon > 0}$ does not converge to u , not even in the weak topology of L^p , $p \geq 1$, in the weak* topology of L^∞ , or up to subsequences.

The precise meaning of the last statement is the following: for every $p \geq 1$ and $T > 0$ the statement “there is a sequence ε_k such that $\varepsilon_k \rightarrow 0^+$ and $u_{\varepsilon_k} \rightharpoonup u$ in $L^p([0, T] \times \mathbb{R})$ ” is false; the statement “there is a sequence ε_k such that $\varepsilon_k \rightarrow 0^+$ and $u_{\varepsilon_k} \overset{*}{\rightharpoonup} u$ in $L^\infty([0, T] \times \mathbb{R})$ ” is also false. The basic idea underpinning Lemma 5.1 is, very loosely speaking, the following: one can show that for t small enough, the entropy admissible solution of the Cauchy problem (5.2), (5.3) has a steady shock at $x = 0$ between the values 1 (on the left) and -1 (on the right). By using the formal computation

$$\frac{d}{dt} \int_{-\infty}^0 u(t, x) dx \stackrel{(5.2)}{=} - \int_{-\infty}^0 \partial_x [u^2](t, x) dx = -u^2(0^-) = -1 < 0$$

we infer the first inequality in (5.4). On the other hand, we can show that the solution u_ε of (5.1), (5.3) is odd. Since the function η_ε is even, this implies that $u_\varepsilon * \eta_\varepsilon = 0$ at $x = 0$ and hence that

$$\frac{d}{dt} \int_{-\infty}^0 u_\varepsilon(t, x) dx \stackrel{(5.1)}{=} - \int_{-\infty}^0 \partial_x [u_\varepsilon u_\varepsilon * \eta_\varepsilon](t, x) dx = 0,$$

which in turn implies the equality in (5.4). By (5.4) and doing some more work one can eventually rule out weak convergence. We now give the rigorous proof of Lemma 5.1.

Proof of Lemma 5.1. We proceed according to the following steps.

STEP 1: we investigate the structure of the entropy solution u . First, we collect some properties of u :

- a) $u \in C^0([0, +\infty[; L^1(\mathbb{R}))$.
- b) Since $\|\bar{u}\|_{L^\infty} \leq 1$, then by the maximum principle $\|u(t, \cdot)\|_{L^\infty} \leq 1$ for every $t \geq 0$.
- c) Since $\bar{u} \in BV(\mathbb{R})$, then $u(t, \cdot) \in BV(\mathbb{R})$ for every $t \geq 0$.

d) A 0-speed shock is created at $t = 0$ at the origin $x = 0$. Owing to the finite propagation speed, this shock will survive for some time. More precisely, we have

$$(5.5) \quad u(t, x) = \begin{cases} 1 & \text{for a.e } x \in]-1/2, 0[\\ -1 & \text{for a.e } x \in]0, 1/2[\end{cases} \quad \text{for every } t \in [0, 1/4].$$

e) Owing to the finite propagation speed and to the fact that $\bar{u} = 0$ if $|x| > 2$, we have $u(t, x) = 0$ for a.e. $|x| \geq 3$ and for every $t \in [0, 1/4]$.

We now want to show that

$$(5.6) \quad \int_{-4}^0 u(1/4, x) dx = \int_{-4}^0 \bar{u}(x) dx - \frac{1}{4}.$$

We can *formally* obtain (5.6) by pointing out that

$$\frac{d}{dt} \int_{-4}^0 u(t, x) dx \stackrel{(5.2)}{=} - \int_{-4}^0 \partial_x [u^2](t, x) dx = -u^2(t, 0^-) + u^2(t, -4) \stackrel{d), e)}{=} -1$$

and by integrating with respect to time. We now sketch a rigorous argument to justify (5.6). First, we point out that u is a distributional solution of (5.2), which amounts to say that

$$(5.7) \quad \int_0^{+\infty} \int_{\mathbb{R}} u \partial_t \varphi \, dx dt + \int_0^{+\infty} \int_{\mathbb{R}} u^2 \partial_x \varphi \, dx dt + \int_{\mathbb{R}} \varphi(0, \cdot) \bar{u} \, dx = 0$$

for every $\varphi \in C_c^\infty(\mathbb{R}^2)$. We now introduce the sequence of functions $\{\chi_n\} \subseteq C_c^\infty(\mathbb{R})$ such that

$$(5.8) \quad \chi_n(x) = \begin{cases} 1 & -4 + 1/n \leq x \leq -1/n \\ 0 & x \leq -4 \text{ or } x \geq 0. \end{cases}$$

As a matter of fact, χ_n is an approximation of the characteristic function of $[-4, 0]$. We fix an arbitrary $\theta \in C_c^\infty(]0, 1/4[)$, we plug $\varphi_n(t, x) := \chi_n(x)\theta(t)$ as a test function in (5.7) and we point out that

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbb{R}} u^2 \partial_x \varphi_n \, dx dt &= \int_0^{1/4} \theta(t) \int_{\mathbb{R}} u^2(t, x) \chi_n'(x) \, dx dt \\ &\stackrel{(5.8)}{=} \int_0^{1/4} \theta(t) \int_{-4}^{-4+1/n} u^2(t, x) \chi_n'(x) \, dx dt + \int_0^{1/4} \theta(t) \int_{-1/n}^0 u^2(t, x) \chi_n'(x) \, dx dt \\ &\stackrel{(5.5), e)}{=} \int_0^{1/4} \theta(t) \int_{-4}^{-4+1/n} 0 \cdot \chi_n'(x) \, dx dt + \int_0^{1/4} \theta(t) \int_{-1/n}^0 1 \cdot \chi_n'(x) \, dx dt \\ &\stackrel{(5.8)}{=} \int_0^{1/4} \theta(t) (-1) \, dt. \end{aligned}$$

Next, we let $n \rightarrow +\infty$ in the other term in (5.7) and we eventually arrive at

$$\int_0^{1/4} \theta'(t) \int_{-4}^0 u(t, x) \, dx \, dt + \int_0^{1/4} \theta(t) (-1) \, dt = 0.$$

Owing to the arbitrariness of θ , this implies that the continuous function

$$(5.9) \quad t \mapsto \int_{-4}^0 u(t, x) \, dx$$

has distributional derivative equal to -1 . This implies that the above function is actually absolutely continuous and, owing to the Fundamental Theorem of Calculus, we get (5.6).

Since we will need it in the following, we also point that, since the map in (5.9) is continuous, then (5.6) implies that there is $h > 0$ such that

$$\int_{1/4-h}^{1/4+h} \int_{-4}^0 u(t, x) \, dx dt \leq \int_{1/4-h}^{1/4+h} \left(\int_{-4}^0 \bar{u} \, dx - \frac{1}{8} \right) dt = 2h \int_{-4}^0 \bar{u} \, dx - \frac{h}{4}.$$

In other words, if we define E by setting

$$(5.10) \quad E := \{(t, x) : t \in [1/4 - h, 1/4 + h], x \in [-4, 0]\}$$

and we denote by $\mathbf{1}_E$ the characteristic function of E , then

$$(5.11) \quad \int_0^\infty \int_{\mathbb{R}} \mathbf{1}_E u \, dx dt \leq 2h \int_{-4}^0 \bar{u} \, dx - \frac{h}{4}.$$

STEP 2: we show that the distributional solution u_ε of (5.1) is odd, namely that, for a.e. $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$, $u(t, x) = -u(t, -x)$. We set $v_\varepsilon(t, x) := -u_\varepsilon(t, -x)$. If we can prove that v_ε is also a distributional solution of the Cauchy problem (5.1), then by the uniqueness part of Proposition 2.3 we get that for every $t \geq 0$ it holds $v_\varepsilon(t, x) = u_\varepsilon(t, x)$ for a.e. x , namely that u_ε is an odd function.

To show that v_ε is a distributional solution of (5.1), we first observe that by using the fact that η_ε is even and making the change of variables $z = -y$ we get

$$(5.12) \quad (v_\varepsilon * \eta_\varepsilon)(t, x) = - \int_{\mathbb{R}} u_\varepsilon(t, -x + y) \eta_\varepsilon(y) dy = - \int_{\mathbb{R}} u_\varepsilon(t, -x - z) \eta_\varepsilon(z) dz = -(u_\varepsilon * \eta_\varepsilon)(t, -x).$$

Next, we fix $\varphi \in C_c^\infty(\mathbb{R}^2)$, we set $\phi(t, x) := -\varphi(t, -x)$ and we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbb{R}} v_\varepsilon \partial_t \varphi \, dx dt + \int_0^{+\infty} \int_{\mathbb{R}} v_\varepsilon (v_\varepsilon * \eta_\varepsilon) \partial_x \varphi \, dx dt + \int_{\mathbb{R}} \varphi(0, \cdot) \bar{u} \, dx = [z = -x] \\ & = \int_0^{+\infty} \int_{\mathbb{R}} (-u_\varepsilon) (-\partial_t \phi) \, dz dt + \int_0^{+\infty} \int_{\mathbb{R}} (-u_\varepsilon) (-u_\varepsilon * \eta_\varepsilon) (\partial_x \phi) \, dz dt + \int_{\mathbb{R}} (-\phi(0, \cdot)) (-\bar{u}) \, dz = 0. \end{aligned}$$

To establish the last equality we have used the fact that u_ε is a distributional solution of (5.1). The above chain of equalities states that v_ε is a distributional solution of (5.1) and hence concludes STEP 2.

STEP 3: we show that

$$(5.13) \quad u_\varepsilon(t, x) = 0, \quad \text{for a.e. } |x| \geq 2 \text{ and every } t \geq 0 \text{ and } \varepsilon > 0.$$

We note that u_ε is a distributional solution of the Cauchy problem

$$(5.14) \quad \begin{cases} \partial_t u_\varepsilon + \partial_x [u_\varepsilon g_\varepsilon] = 0 \\ u_\varepsilon(0, \cdot) = \bar{u} \end{cases}$$

provided that the vector field g_ε is defined as $g_\varepsilon(t, x) := u_\varepsilon * \eta_\varepsilon$. Since the vector field g_ε is smooth, then we can apply the method of characteristics. We term $X(t, x)$ the characteristic curve solving the Cauchy problem

$$(5.15) \quad \begin{cases} \frac{dX}{dt} = g_\varepsilon(t, X) \\ X(0, x) = x. \end{cases}$$

Recall that u_ε is an odd function by STEP 2. Since η_ε is an even function by assumption, by arguing as in the chain of equalities (5.12) we obtain that $u_\varepsilon * \eta_\varepsilon$ is an odd function. Since it is also smooth, we eventually conclude that

$$(5.16) \quad g_\varepsilon(t, 0) = u_\varepsilon * \eta_\varepsilon(t, 0) = 0, \quad \text{for every } t \geq 0.$$

This means that $X(t, 0) \equiv 0$ and, since (5.15) has a unique solution, implies that the characteristic curves *cannot cross* the t axis. Since $\bar{u}(x) \geq 0$ if $x \leq 0$ and $\bar{u}(x) \leq 0$ if $x \geq 0$, this in turn implies that

$$(5.17) \quad \text{for every } t \geq 0, u_\varepsilon(t, x) \geq 0 \text{ for a.e. } x < 0 \text{ and } u_\varepsilon(t, x) \leq 0 \text{ for a.e. } x > 0.$$

This implies that, if $x \leq -2$, then $x + \varepsilon \leq 0$ and hence

$$(5.18) \quad g_\varepsilon(t, x) = u_\varepsilon * \eta_\varepsilon(t, x) = \int_{x-\varepsilon}^{x+\varepsilon} u_\varepsilon(y) \eta_\varepsilon(x-y) dy \stackrel{(1.10), (5.17)}{\geq} 0.$$

If $x_1 < x_2$, then $X(t, x_1) < X(t, x_2)$ for every $t \geq 0$ (to see this, we use again the fact that the solution of (5.15) is unique). By recalling (5.18), this implies that

$$x \geq -2 \implies X(t, x) \geq -2 \text{ for every } t \geq 0$$

and hence that

$$\text{for every } t \geq 0, X(t, x) < -2 \implies x < -2.$$

Since $\bar{u}(x) = 0$ for a.e. $x \leq 2$, this eventually implies that $u_\varepsilon(t, x) = 0$ for every $x \leq -2$. Since the function u_ε is odd, this establishes (5.13).

STEP 4: we conclude the proof. Recall that the set E is defined as in (5.10) for a suitable h and assume that we have shown that

$$(5.19) \quad \int_0^{+\infty} \int_{\mathbb{R}} \mathbf{1}_E u_\varepsilon dx dt = 2h \int_{-4}^0 \bar{u} dx.$$

Since the function $\mathbf{1}_E \in L^p(\mathbb{R}^+ \times \mathbb{R})$, for every $p \in [1, +\infty]$, then by comparing (5.19) and (5.11) we rule out the possibility that u_ε converges weakly or weakly* to u . We are thus left with establishing (5.19). To this end, we first use the formal computation

$$(5.20) \quad \frac{d}{dt} \int_{-4}^0 u_\varepsilon(t, x) dx \stackrel{(5.1)}{=} - \int_{-4}^0 \partial_x [u_\varepsilon(u_\varepsilon * \eta_\varepsilon)](t, x) dx = u_\varepsilon(u_\varepsilon * \eta_\varepsilon)(t, -4) - u_\varepsilon(u_\varepsilon * \eta_\varepsilon)(t, 0) \stackrel{(5.13), (5.16)}{=} 0.$$

This implies that

$$\int_0^{+\infty} \int_{\mathbb{R}} \mathbf{1}_E u_\varepsilon dx dt = \int_{1/4-h}^{1/4+h} \int_{-4}^0 u_\varepsilon dx dt = \int_{1/4-h}^{1/4+h} \int_{-4}^0 \bar{u}(x) dx dt = 2h \int_{-4}^0 \bar{u}(x) dx,$$

namely (5.19). To provide a rigorous justification of (5.20) one can argue as in STEP 1. This concludes the proof of the lemma. \square

5.2. A counter-example with positive data and asymmetric kernels. This paragraph aims at establishing the following lemma, which rules out also the possibility that u_{ε_k} weakly converges to a distributional, not necessarily entropy admissible, solution of (5.2).

Lemma 5.2. *Assume that η_ε satisfies (1.2) and (1.4) and moreover that*

$$(5.21) \quad \eta(x) = 0, \quad \text{for every } x \geq 0.$$

Let \bar{u} be given by

$$(5.22) \quad \bar{u}(x) = \begin{cases} 1 & -1 < x < 0 \\ 0 & \text{otherwise.} \end{cases}$$

Let u_ε be the solution of the Cauchy problem (5.1), (5.22) and u be the entropy admissible solution of (5.2), (5.22). Then

- (1) *the family of distributional solutions $\{u_\varepsilon\}_{\varepsilon>0}$ does not converge to u , not even in the weak topology of L^p , $p \geq 1$, in the weak* topology of L^∞ , or up to subsequences;*
- (2) *more in general, any weak limit w of a subsequence of $\{u_\varepsilon\}_{\varepsilon>0}$ (in the weak topology of L^p , $p \geq 1$, in the weak* topology of L^∞) cannot be a L^2_{loc} distributional (not necessarily entropy admissible) solution of (5.2).*

The basic idea underpinning Lemma 5.2 is, very loosely speaking, the following. Owing to (5.21), the convolution $u_\varepsilon * \eta_\varepsilon$ evaluated at the point x only depends on the values of u_ε on the right hand side of x . Owing to the particular structure of the initial datum \bar{u} this implies that $u_\varepsilon * \eta_\varepsilon(0, 0) = 0$ and hence that the characteristic line of the velocity field $u_\varepsilon * \eta_\varepsilon$ starting at $x = 0$ has zero initial speed. Then, one can show that the speed is identically zero: this implies that the characteristic lines coming from the half line $x < 0$ cannot cross the axis $x = 0$, and hence that no mass can enter the half line $x > 0$. In conclusion, $u_\varepsilon(t, x) = 0$ for a.e. $x > 0$. Notice that this last equality could be shown

also by noticing that the approximating sequence in the construction of u_ε in [10, § 5] enjoys the same property.

On the other hand, the entropy admissible solution of (5.2), (5.22) is explicit and not identically 0 for $x > 0$. With some more work, one can show that any *distributional* solution of (5.2), (5.22) is not identically 0 for $x > 0$. This allows to rule out weak convergence to a distributional solution. We now make the previous argument rigorous.

Lemma 5.3. *Assume that η and η_ε satisfy (1.2), (1.4) and (5.21) and let \bar{u} be as in (5.22). Then*

$$(5.23) \quad \text{for every } t \geq 0, u_\varepsilon(t, x) = 0 \text{ for a.e. } x < -1 \text{ and a.e. } x > 0.$$

Proof. We argue according to the following steps.

STEP 1: we show that $u_\varepsilon(t, x) = 0$ for a.e. $x < -1$. We use the method of characteristics: note that u_ε is a distributional solution of the continuity equation (5.14) provided the vector field g_ε is given by $g_\varepsilon := u_\varepsilon * \eta_\varepsilon$. Since $\bar{u} \geq 0$, then $g_\varepsilon(t, x) \geq 0$ for every (t, x) . This implies that, for every $t \geq 0$ and every $x < -1$, the characteristic line $Y_t(s, x)$ solving the (backward) Cauchy problem

$$\begin{cases} \frac{dY_t}{ds} = g_\varepsilon(s, Y_t) \\ Y_t(t, x) = x \end{cases}$$

satisfies $Y_t(0, x) < -1$ and hence $\bar{u}(Y_t(0, x)) = 0$. Since the value 0 is propagated along the characteristic lines of the continuity equation, then $u_\varepsilon(t, x) = 0$.

STEP 2: we regard again u_ε as the solution of the continuity equation (5.14) and we term X the characteristic line solving the (forward) Cauchy problem (5.15). We claim that

$$(5.24) \quad X(t, x) = x \quad \text{for every } t \geq 0, x \geq 0.$$

Indeed, by the spatial smoothness of the vector field $u_\varepsilon * \eta_\varepsilon$, the characteristic lines “cannot cross” the curve $X(t, 0)$; in particular for any $t > 0$ and $x > X(t, 0)$ we have $Y_t(0, x) > 0$. Hence

$$(5.25) \quad u_\varepsilon(t, x) = 0 \quad \text{for any } t > 0, x \geq X(t, 0).$$

Since η_ε satisfies (1.2) and (1.4), for any $x \in \mathbb{R}$ the quantity $u_\varepsilon * \eta_\varepsilon(x)$ is an average, weighted with η_ε , of the values of $u_\varepsilon(t, \cdot)$ on the right of x . From (5.25), we deduce that

$$(5.26) \quad u_\varepsilon * \eta_\varepsilon(t, x) = 0 \quad \text{for any } t > 0, x \geq X(t, 0).$$

From (5.26) applied to $x = X(t, 0)$ and (5.15) with $x = 0$, we deduce that $X(t, 0) = 0$ for any $t > 0$; applying again (5.26) with this further information, we deduce (5.24).

Since the value 0 is propagated along the characteristic lines of the continuity equation, which in turn are constant for any $x \geq 0$ thanks to (5.24), we have shown that $u_\varepsilon(t, x) = 0$ for any $x \geq 0$, concluding the proof of (5.23). \square

Proof of Lemma 5.2(1). First, we point out that, if \bar{u} is given by (5.22), then the entropy admissible solution of the Cauchy problem (5.2) is

$$(5.27) \quad u(t, x) = \begin{cases} 0 & x \leq -1 \text{ or } x \geq t \\ \frac{x+1}{2t} & -1 \leq x \leq 2t-1 \\ 1 & 2t-1 \leq x \leq t, \end{cases} \quad \text{for a.e. } (t, x) \in [0, 1] \times \mathbb{R}.$$

Assume by contradiction there is a sequence $\{\varepsilon_k\}$ such that u_{ε_k} weakly converges to u . We use as a test function the characteristic function of the set $E := [0, 1/2] \times [0, 1]$. Since

$$\int_{\mathbb{R}^+ \times \mathbb{R}} u_{\varepsilon_k} \mathbf{1}_E dx dt \stackrel{\text{Lemma 5.3}}{=} 0, \quad \int_{\mathbb{R}^+ \times \mathbb{R}} u \mathbf{1}_E dx dt \stackrel{(5.27)}{=} \int_0^{1/2} \int_0^t 1 dx dt = \frac{1}{8},$$

then we find a contradiction. \square

The proof of Lemma 5.2(2) is based on the following result, which could be generalized to Young measure solutions of the Cauchy problem (5.2) (we refer to [11] for an extended discussion on Young measures and their applications to nonlinear conservation laws).

Lemma 5.4. *Let $a, b \in \mathbb{R}$, $a < b$, and let $u \in L^2_{\text{loc}}([0, 1] \times \mathbb{R})$ be a nonnegative, distributional solution of the Cauchy problem (5.2) compactly supported in $[0, 1] \times (a, b)$. Then the baricenter of u is a nondecreasing function and*

$$(5.28) \quad \int_a^b x u(t, x) dx \geq \left(\int_a^b \bar{u} \right)^2 t + \int_a^b x \bar{u}(x) dx.$$

The proof of Lemma 5.2(2) straightforwardly follows from Lemma 5.4. Indeed, any nonnegative distributional solution u of the Cauchy problem (5.2) starting from \bar{u} in (5.22) cannot satisfy

$$u(t, x) = 0 \text{ for a.e. } t \in [0, 1], x \in (-\infty, -1) \cup (0, \infty),$$

because otherwise it would contradict (5.28) for $a = -1 - \sigma, b = \sigma$ (σ arbitrarily small) and any $t > 1/2$. Hence we find a contradiction with (5.23) as in the proof of Lemma 5.2(1).

Proof of Lemma 5.4. The conservation law (5.2) implies

$$\int_a^b u(t, x) dx = \int_{\mathbb{R}} u(t, x) dx = \int_a^b \bar{u}(x) dx \quad \text{for a.e. } t > 0.$$

We perform some *formal* computations, which can be made rigorous by arguing as in the proof of Lemma 5.1: by (5.2), the Jensen inequality, and the previous equality, we have

$$\begin{aligned} \frac{d}{dt} \int_a^b x u(t, x) dx &= \int_a^b x \partial_t u(t, x) dx = - \int_a^b x \partial_x [u^2](t, x) dx \\ &= \left[-x u^2(t, x) \right]_{x=b}^{x=a} + \int_a^b u^2(t, x) dx \geq \frac{1}{b-a} \left(\int_a^b u(t, x) dx \right)^2. \end{aligned}$$

Integrating in time, we get (5.28). \square

5.3. A counter-example with positive data and symmetric kernels. We now establish the following result.

Lemma 5.5. *Assume that η and η_ε are as in (1.10) and (1.4), respectively, and that η is an even function. Let u denote the entropy admissible solution of (5.2), (5.22) and u_ε the solution of (5.1), (5.22). Then for every $\delta > 0$, the family u_ε does not converge to u strongly in $L^{1+\delta}$, not even up to subsequences. More precisely,*

$$(5.29) \quad \forall t > 0, \nexists \{\varepsilon_k\}, \varepsilon_k \rightarrow 0^+ \text{ such that } u_{\varepsilon_k}(t, \cdot) \rightarrow u(t, \cdot) \text{ strongly in } L^{1+\delta}.$$

Note that (5.29) rules out the possibility that u_ε converges to u in $L^{1+\delta}([0, 1] \times \mathbb{R})$: indeed, if this were true then, up to subsequences, $u_\varepsilon(t, \cdot) \rightarrow u(t, \cdot)$ in $L^{1+\delta}$ for a.e. t , and this is ruled out by (5.29).

The basic idea underpinning Lemma 5.5 is the following. We introduce the entropy function

$$\mathcal{E}(u) := \int_{\mathbb{R}} u \ln u \, dx,$$

where by a slight abuse of notation we have continuously extended the function $u \ln u$ with value 0 for $u = 0$. By using the formal computation (5.35), one gets that

$$\frac{d}{dt} \mathcal{E}(u_\varepsilon) = 0$$

if u_ε is a nonnegative solution of (5.1). On the other hand, the function $u \ln u$ is convex and hence $\mathcal{E}(u)$ is non increasing for nonnegative entropy admissible solutions of (5.2). In particular, if the initial datum is as in (5.22), then $\mathcal{E}(u)$ is strictly decreasing. After some more work this allows us to rule out the strong convergence of u_ε to u .

The precise argument requires some preliminary results.

Lemma 5.6. *Fix $\delta > 0$ and assume that $\{v_k\} \subseteq L^{1+\delta}$ satisfies $v_k \rightarrow v$ in $L^{1+\delta}$, for some compactly supported $v \in L^{1+\delta}$. Then*

$$(5.30) \quad \int_{\mathbb{R}} v \ln v \, dx \geq \limsup_{k \rightarrow +\infty} \int_{\mathbb{R}} v_k \ln v_k \, dx.$$

Proof. Let $\Omega \subset \mathbb{R}$ be a compact set s.t. $v = 0$ a.e. in $\mathbb{R} \setminus \Omega$. Up to a (not relabelled) subsequence, we can assume that the lim sup in the right-hand side in (5.30) is a limit. Since $v_k \rightarrow v$ in $L^{1+\delta}$, up to a further subsequence, we can assume that v_k converges pointwise to v a.e. in \mathbb{R} and that there exists a function $h \in L^{1+\delta}(\mathbb{R})$ such that $h \geq |v_k|$ a.e. for any $k \in \mathbb{N}$. We observe that

$$(5.31) \quad |w \ln w| \leq C(\delta)(\mathbf{1}_{\{w < 1\}} + \mathbf{1}_{\{w \geq 1\}}|w|^{1+\delta}), \quad \text{for every } w \geq 0.$$

Since the function $s \rightarrow s \ln s$ is negative for $s < 1$, by (5.31), and since $v_k \rightarrow v$ in $L^{1+\delta}(\mathbb{R} \setminus \Omega)$ we have

$$\limsup_{k \rightarrow \infty} \int_{\mathbb{R} \setminus \Omega} v_k \ln v_k \, dx \leq \limsup_{k \rightarrow \infty} \int_{\mathbb{R} \setminus \Omega} v_k \ln v_k \mathbf{1}_{v_k \geq 1} \, dx \leq C(\delta) \lim_{k \rightarrow \infty} \int_{\mathbb{R} \setminus \Omega} |v_k|^{1+\delta} \, dx = 0.$$

Since the functions $v_k \ln v_k$ converge pointwise to $v \ln v$ as $k \rightarrow \infty$ and the convergence is dominated by $C(\delta)(1 + |h|^{1+\delta}) \in L^1(\Omega)$, we have

$$\int_{\Omega} v \ln v \, dx = \lim_{k \rightarrow \infty} \int_{\Omega} v_k \ln v_k \, dx \geq \lim_{k \rightarrow \infty} \int_{\Omega} v_k \ln v_k \, dx + \limsup_{k \rightarrow \infty} \int_{\mathbb{R} \setminus \Omega} v_k \ln v_k \, dx = \limsup_{k \rightarrow \infty} \int_{\mathbb{R}} v_k \ln v_k \, dx,$$

which proves (5.30). \square

Lemma 5.7. *Let u_ε be the solution of (5.2) and assume that η_ε satisfies (1.10) and (1.4) and that η is an even function. Assume that $\bar{u} \in L^\infty$ is a compactly supported function satisfying $\bar{u} \geq 0$ and*

$$\int_{\mathbb{R}} \bar{u} \ln \bar{u} \, dx < +\infty.$$

Then for every $t \geq 0$ the distributional solution satisfies the following properties: $u_\varepsilon(t, \cdot) \geq 0$ and

$$(5.32) \quad \int_{\mathbb{R}} u_\varepsilon \ln u_\varepsilon(t, \cdot) \, dx = \int_{\mathbb{R}} \bar{u} \ln \bar{u} \, dx.$$

Proof. First, we point out that one can check by direct computation that, for every $a, b \in L^2(\mathbb{R})$, $c \in C_c^\infty(\mathbb{R})$, we have

$$(5.33) \quad \int_{\mathbb{R}} a(b * c) \, dx = \int_{\mathbb{R}} (a * \check{c})b \, dx,$$

where $\check{c}(x) = c(-x)$. We apply the above formula with $a = b = u_\varepsilon(t, \cdot)$ and $c = \eta'_\varepsilon$. Since η_ε is an even function, then the derivative η'_ε is an odd function and hence $\check{\eta}'_\varepsilon = -\eta'_\varepsilon$. We then obtain

$$\int_{\mathbb{R}} u_\varepsilon \partial_x [u_\varepsilon * \eta_\varepsilon](t, \cdot) \, dx = \int_{\mathbb{R}} u_\varepsilon (u_\varepsilon * \eta'_\varepsilon)(t, \cdot) \, dx \stackrel{(5.33)}{=} - \int_{\mathbb{R}} (u_\varepsilon * \eta'_\varepsilon) u_\varepsilon(t, \cdot) \, dx = - \int_{\mathbb{R}} u_\varepsilon \partial_x [u_\varepsilon * \eta_\varepsilon](t, \cdot) \, dx,$$

which implies that

$$(5.34) \quad \int_{\mathbb{R}} u_\varepsilon \partial_x [u_\varepsilon * \eta_\varepsilon](t, \cdot) \, dx = 0.$$

We can then establish (5.32) by using the following (formal) computation:

$$(5.35) \quad \begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} u_\varepsilon \ln u_\varepsilon(t, \cdot) \, dx &= \int_{\mathbb{R}} (1 + \ln u_\varepsilon) \partial_t u_\varepsilon(t, \cdot) \, dx \stackrel{(5.2)}{=} - \int_{\mathbb{R}} (1 + \ln u_\varepsilon) \partial_x [u_\varepsilon (u_\varepsilon * \eta_\varepsilon)](t, \cdot) \, dx \\ &= \int_{\mathbb{R}} \partial_x u_\varepsilon \frac{1}{u_\varepsilon} u_\varepsilon (u_\varepsilon * \eta_\varepsilon)(t, \cdot) \, dx = \int_{\mathbb{R}} \partial_x u_\varepsilon (u_\varepsilon * \eta_\varepsilon)(t, \cdot) \, dx = - \int_{\mathbb{R}} u_\varepsilon \partial_x [u_\varepsilon * \eta_\varepsilon](t, \cdot) \, dx \stackrel{(5.34)}{=} 0. \end{aligned}$$

To make the above argument rigorous, we recall that u_ε is the solution of the Cauchy problem (5.14), where the velocity field $g_\varepsilon = u_\varepsilon * \eta_\varepsilon$ is smooth. By the renormalization property, for every $\beta \in C^1$ we have (2.14). This implies that

$$(5.36) \quad \int_{\mathbb{R}} \beta(u_\varepsilon(t, \cdot)) dx = \int_{\mathbb{R}} \beta(\bar{u}) dx - \int_0^t \int_{\mathbb{R}} \partial_x [u_\varepsilon * \eta_\varepsilon] [u_\varepsilon \beta'(u_\varepsilon) - \beta(u_\varepsilon)] dx ds.$$

We construct a sequence of functions $\beta_n : \mathbb{R}^+ \rightarrow \mathbb{R}$ by setting

$$\beta_n(v) := \int_0^v \left[1 + \ln \left(\xi + \frac{1}{n} \right) \right] d\xi.$$

Note that

$$\begin{aligned} \beta_n(v) &\rightarrow \int_0^v [1 + \ln \xi] d\xi = v \ln v, \quad \text{for every } v \geq 0, \text{ as } n \rightarrow +\infty, \\ v\beta'_n(v) - \beta_n(v) &\rightarrow v, \quad \text{for every } v \geq 0, \text{ as } n \rightarrow +\infty. \end{aligned}$$

By testing the inequality (5.36) with β_n and passing to the limit for $n \rightarrow +\infty$ we obtain

$$\int_{\mathbb{R}} u_\varepsilon \ln u_\varepsilon dx = \int_{\mathbb{R}} \bar{u} \ln \bar{u} dx - \int_0^t \int_{\mathbb{R}} \partial_x [u_\varepsilon * \eta_\varepsilon] u_\varepsilon ds dx \stackrel{(5.34)}{=} \int_{\mathbb{R}} \bar{u} \ln \bar{u} dx.$$

This concludes the proof of the lemma. \square

Proof of Lemma 5.5. If \bar{u} is given by (5.22), then the entropy admissible solution u can be explicitly computed and is given by (5.27). Note that \bar{u} only attains the values 0 and 1, whereas if $t \in (0, 1]$ then u attains values between 0 and 1. Therefore

$$(5.37) \quad \int_{\mathbb{R}} u(t, x) \ln u(t, x) dx < 0 = \int_{\mathbb{R}} \bar{u} \ln \bar{u} dx \quad \text{for any } t \in (0, 1].$$

Owing to Lemma 5.7, for every $\varepsilon > 0$ and $t \geq 0$ we have

$$(5.38) \quad \int_{\mathbb{R}} u_\varepsilon(t, x) \ln u_\varepsilon(t, x) dx = 0.$$

Assume by contradiction that there is a sequence $\varepsilon_k \rightarrow 0^+$ and a time $t > 0$ such that $u_{\varepsilon_k}(t, \cdot) \rightarrow u(t, \cdot)$ strongly in $L^{1+\delta}(\mathbb{R})$. We apply Lemma 5.6 with $v_k := u_{\varepsilon_k}(t, \cdot)$, $v := u(t, \cdot)$. By combining (5.38) and (5.30) we get

$$\int_{\mathbb{R}} u(t, x) \ln u(t, x) dx \geq 0,$$

which contradicts the second inequality in (5.37). This concludes the proof of the lemma. \square

6. ACKNOWLEDGMENTS

The authors wish to thank Stefano Bianchini, Rinaldo Colombo, Elio Marconi, Mario Pulvirenti, Giuseppe Savaré and Giuseppe Toscani for valuable discussions. In particular, Lemma 5.4 is due to Elio Marconi and the equality (5.35) was pointed out to us by Giuseppe Savaré. GC is partially supported by the Swiss National Science Foundation grant 200020_156112 and by the ERC Starting Grant 676675 FLIRT. LVS is a member of the GNAMPA group of INDAM and of the PRIN National Project ‘‘Hyperbolic Systems of Conservation Laws and Fluid Dynamics: Analysis and Applications’’. Part of this work was done when LVS and MC were visiting the University of Basel, and its kind hospitality is gratefully acknowledged.

REFERENCES

- [1] L. Ambrosio. Transport equation and Cauchy problem for BV vector fields. *Invent. Math.*, 158(2):227–260, 2004.
- [2] P. Amorim, R. M. Colombo, and A. Teixeira. On the numerical integration of scalar nonlocal conservation laws. *ESAIM Math. Model. Numer. Anal.*, 49(1):19–37, 2015.
- [3] F. Betancourt, R. Bürger, K. H. Karlsen, and E. M. Tory. On nonlocal conservation laws modelling sedimentation. *Nonlinearity*, 24(3):855–885, 2011.
- [4] H. S. Bhat and R. C. Fetecau. A Hamiltonian regularization of the Burgers equation. *J. Nonlinear Sci.*, 16(6):615–638, 2006.
- [5] S. Blandin and P. Goatin. Well-posedness of a conservation law with non-local flux arising in traffic flow modeling. *Numer. Math.*, 132(2):217–241, 2016.
- [6] P. Calderoni and M. Pulvirenti. Propagation of chaos for Burgers’ equation. *Ann. Inst. H. Poincaré Sect. A (N.S.)*, 39(1):85–97, 1983.
- [7] R. M. Colombo, M. Garavello, and M. Lécureux-Mercier. A class of nonlocal models for pedestrian traffic. *Math. Models Methods Appl. Sci.*, 22(4):1150023, 34, 2012.
- [8] R. M. Colombo, M. Herty, and M. Mercier. Control of the continuity equation with a non local flow. *ESAIM Control Optim. Calc. Var.*, 17(2):353–379, 2011.
- [9] G. Crippa, C. Donadello, and L. V. Spinolo. A note on the initial-boundary value problem for continuity equations with rough coefficients. In *Hyperbolic problems: theory, numerics and applications*, volume 8 of *AIMS Series on Applied Mathematics*, pages 957–966. 2014.
- [10] G. Crippa and M. Lécureux-Mercier. Existence and uniqueness of measure solutions for a system of continuity equations with non-local flow. *NoDEA Nonlinear Differential Equations Appl.*, 20(3):523–537, 2013.
- [11] C. M. Dafermos. *Hyperbolic conservation laws in continuum physics*, volume 325 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, fourth edition, 2016.
- [12] C. De Lellis. Notes on hyperbolic systems of conservation laws and transport equations. In *Handbook of differential equations: evolutionary equations. Vol. III*, Handb. Differ. Equ., pages 277–382. Elsevier/North-Holland, Amsterdam, 2007.
- [13] R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, 98(3):511–547, 1989.
- [14] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [15] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81 (123):228–255, 1970.
- [16] R. J. LeVeque. *Finite volume methods for hyperbolic problems*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2002.
- [17] K. Zumbrun. On a nonlocal dispersive equation modeling particle suspensions. *Quart. Appl. Math.*, 57(3):573–600, 1999.

M.C. INTITUTE FOR THEORETICAL STUDIES, ETH ZÜRICH, CLAUSIUSSTRASSE 47, CH-8092 ZÜRICH, SWITZERLAND,
 INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, CH-8057 ZÜRICH, SWITZERLAND.
E-mail address: maria.colombo@math.uzh.ch

G.C. DEPARTEMENT MATHEMATIK UND INFORMATIK, UNIVERSITÄT BASEL, SPIEGELGASSE 1, CH-4051 BASEL,
 SWITZERLAND.
E-mail address: gianluca.crippa@unibas.ch

L.V.S. IMATI-CNR, VIA FERRATA 1, I-27100 PAVIA, ITALY.
E-mail address: spinolo@imati.cnr.it