

A TWO-PHASE PROBLEM WITH ROBIN CONDITIONS ON THE FREE BOUNDARY

SERENA GUARINO LO BIANCO, DOMENICO ANGELO LA MANNA, BOZHIDAR VELICHKOV

ABSTRACT. We study for the first time a two-phase free boundary problem in which the solution satisfies a Robin boundary condition. We consider the case in which the solution is continuous across the free boundary and we prove an existence and a regularity result for minimizers of the associated variational problem. Finally, in the appendix, we give an example of a class of Steiner symmetric minimizers.

1. INTRODUCTION

For a fixed a constant $\beta > 0$ and a smooth bounded open set $D \subset \mathbb{R}^d$, $d \geq 2$, we consider the functional

$$J_\beta(u, \Omega) = \int_D |\nabla u|^2 dx + \beta \int_{\partial^* \Omega} u^2 d\mathcal{H}^{d-1},$$

defined on the couples (u, Ω) , where $u \in H^1(D)$, $\Omega \subset \mathbb{R}^d$ is a set of finite perimeter in the sense of De Giorgi (see Section 2) and $\partial^* \Omega$ denotes the reduced boundary of Ω (see Section 2). Recall that, when Ω is smooth, $\partial^* \Omega$ is the topological boundary of Ω .

In this paper we study the existence and the regularity of minimizers of the functional J_β among all couples (u, Ω) , which are fixed outside the domain D . Precisely, throughout the paper, we fix a set $E \subset \mathbb{R}^d$ of finite perimeter, a constants $m > 0$ and a function

$$v \in H_{loc}^1(\mathbb{R}^d) \quad \text{such that} \quad v \geq m \quad \text{in} \quad \mathbb{R}^d \quad \text{and} \quad \int_{\partial^* E} v^2 d\mathcal{H}^{d-1} < +\infty ;$$

we define the admissible sets

$$\mathcal{V} = \{u \in H_{loc}^1(\mathbb{R}^d) : u - v \in H_0^1(D)\},$$

$$\mathcal{E} = \{\Omega \subset \mathbb{R}^d : \text{Per}(\Omega) < +\infty \text{ and } \Omega = E \text{ in } \mathbb{R}^d \setminus D\},$$

and we consider the variational minimization problem

$$\min \{J_\beta(u, \Omega) : u \in \mathcal{V}, \Omega \in \mathcal{E}\}. \tag{1.1}$$

Our main result is the following.

Theorem 1.1 (Existence and regularity of minimizers). *Let $\beta > 0$, $D \subset \mathbb{R}^d$, v , E , \mathcal{V} and \mathcal{E} be as above. Then the following holds.*

- (i) *There exists a solution $(u, \Omega) \in \mathcal{V} \times \mathcal{E}$ to the variational problem (1.1).*
- (ii) *For every solution (u, Ω) of (1.1), u is Hölder continuous and bounded from below by a strictly positive constant in D .*
- (iii) *If (u, Ω) is a solution to (1.1), then the free boundary $\partial\Omega \cap D$ can be decomposed as the disjoint union of a regular part $\text{Reg}(\partial\Omega)$ and a singular part $\text{Sing}(\partial\Omega)$, where :*
 - *$\text{Reg}(\partial\Omega)$ is a C^∞ hypersurface and a relatively open subset of $\partial\Omega$, and the function u is C^∞ smooth on $\text{Reg}(\partial\Omega)$;*
 - *$\text{Sing}(\partial\Omega)$ is a closed set, which is empty if $d \leq 7$, discrete if $d = 8$, and of Hausdorff dimension $d - 8$, if $d > 8$.*

Remark 1.2. We notice that if (u, Ω) is a solution to (1.1), then u is harmonic in the interior of Ω and $D \setminus \Omega$. Thus, as a consequence of Theorem 1.1 (iii), in a neighborhood of a regular point $x_0 \in \text{Reg}(\partial\Omega)$, the functions $u : \bar{\Omega} \rightarrow \mathbb{R}$ and $u : D \setminus \bar{\Omega} \rightarrow \mathbb{R}$ are C^∞ up to the free boundary $\partial\Omega$.

1.1. Outline of the proof and organization of the paper. The main difficulty in the proof of Theorem 1.1 is to prove the existence of a minimizing couple (u, Ω) and to show that the function u is Hölder continuous and bounded from below by a strictly positive constant in D . The almost-minimality of the solutions is proved in Theorem 5.1. Finally, in the Appendix, we give examples of minimizers in domains D symmetric with respect to the hyperplane $\{x_d = 0\}$.

1.1.1. Existence. The existence of a solution (u, Ω) and the regularity of u (Hölder regularity and non-degeneracy) are treated simulatenously. The reason is that if (u_n, Ω_n) is a minimizing sequence for (1.1), then in order to get the compactness of Ω_n , we need a uniform bound (from above) on the perimeter $\text{Per}(\Omega_n)$, for which we need the functions u_n to be bounded from below by a strictly positive constant. Now, notice that we cannot simply replace u_n by $u_n \vee \varepsilon$, for some $\varepsilon > 0$; this is due to the fact that the second term in J_β is increasing in u :

$$\int_{\partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} \leq \int_{\partial^* \Omega_n} (\varepsilon \vee u_n)^2 d\mathcal{H}^{d-1}.$$

Thus, we select a minimizing sequence which is in some sense optimal. Precisely, we take (u_n, Ω_n) to be solution of the auxiliary problem

$$\min \left\{ J_\beta(u, \Omega) : u \in \mathcal{V}, \Omega \in \mathcal{E}, u \geq \frac{1}{n} \text{ in } D \right\}, \quad (1.2)$$

for which the existence of an optimal set is much easier (see Section 3, Proposition 3.1). Still, we do not have a uniform (independent from n) bound from below for the functions u_n , so we still miss the uniform bound on the perimeter of Ω_n .

On the other hand, we are able to prove that the sequence u_n is uniformly Hölder continuous in D (see Section 3, Lemma 3.5). This allows to extract a subsequence u_n that converges locally uniformly in D to a non-negative Hölder continuous function $u_\infty : D \rightarrow \mathbb{R}$ (see Section 4). Now, on each of the sets $\{u_\infty > t\}$, $t > 0$, the sequence Ω_n has uniformly bounded perimeter. This allows to extract a subsequence Ω_n that converges pointwise almost-everywhere on $\{u_\infty > 0\}$ to some Ω_∞ . Thus, we have constructed our candidate for a solution: $(u_\infty, \Omega_\infty)$.

In order to prove that $(u_\infty, \Omega_\infty)$ is an admissible competitor in (1.1), we need to show that Ω_∞ has finite perimeter. We do this in Section 4. We first use the optimality of (u_n, Ω_n) to prove that $(u_\infty, \Omega_\infty)$ is optimal when compared to a special class of competitors. This optimality condition can be written as (we refer to Lemma 4.1 for the precise statement) :

$$J_\beta(u_\infty, \Omega_\infty) \leq J_\beta(u_t, \Omega_t) \quad \text{where} \quad u_t = u_\infty \vee t \quad \text{and} \quad \Omega_t = \Omega_\infty \cup \{u_\infty \leq t\}, \quad (1.3)$$

for any $t > 0$. Next, from this special optimality condition we deduce that the function u_∞ is bounded from below by a strictly positive constant (see Proposition 4.2). From this, in Section 4, we deduce that Ω_∞ has finite perimeter in \mathbb{R}^d and that the couple $(u_\infty, \Omega_\infty)$ is a solution to (1.1).

1.1.2. Hölder continuity and non-degeneracy of u . Let now (u, Ω) be any solution of (1.1). In order to prove the Hölder continuity and the non-degeneracy of u it is sufficient to exploit some of the estimates that we already used to prove the existence. Indeed, we can test the optimality of (u, Ω) with the competitors from (1.3). Thus, for $t > 0$ small enough, we have

$$J_\beta(u, \Omega) \leq J_\beta(u_t, \Omega_t) \quad \text{where} \quad u_t = u \vee t \quad \text{and} \quad \Omega_t = \Omega \cup \{u \leq t\}. \quad (1.4)$$

In particular,

$$\begin{aligned} \int_D |\nabla u|^2 dx + \beta \int_{\partial^* \Omega} u^2 &\leq \int_D |\nabla(u \vee t)|^2 dx + \beta \int_{\partial^*(\Omega \cup \{u < t\})} u^2 \\ &\leq \int_D |\nabla(u \vee t)|^2 dx + \beta t^2 \text{Per}(\{u < t\}) + \beta \int_{\{u > t\} \cap \partial^* \Omega} u^2, \end{aligned}$$

which proves that u satisfies the optimality condition (4.1) from Lemma 4.1:

$$\int_{\{u < t\}} |\nabla u|^2 dx \leq \beta t^2 \text{Per}(\{u < t\}). \quad (1.5)$$

Now, applying Proposition 4.2, we get that u is bounded from below by a strictly positive constant in D . Finally, Proposition 3.5 gives that u is Hölder continuous in D . This proves Theorem 1.1 (ii).

1.1.3. Regularity of the free boundary. In order to prove the regularity of the free boundary (Theorem 1.1 (iii)), we use the Hölder continuity and the non-degeneracy of u to show that a solution Ω is an almost-minimizer of the perimeter. We do this in Theorem 5.1. Now, from the classical regularity theory for almost-minimizers of the perimeter (see [11]), we obtain that (inside D) the free boundary $\partial\Omega$ can be decomposed into a $C^{1,\alpha}$ -regular part $\text{Reg}(\partial\Omega)$ and a (possibly empty) singular part of Hausdorff dimension smaller than $d - 8$.

Finally, in Theorem 5.2, we prove the C^∞ regularity of $\text{Reg}(\partial\Omega)$. In order to do so, we first show (see Lemma 5.3) that in a neighborhood of a regular point x_0 , the restrictions u_+ and u_- of u on Ω and $D \setminus \Omega$ are solutions of the following transmission problem:

$$\begin{cases} \Delta u_+ = 0 & \text{in } \Omega, \\ \Delta u_- = 0 & \text{in } D \setminus \bar{\Omega}, \\ u_+ = u_- = u & \text{on } \partial\Omega, \\ \frac{\partial u_+}{\partial \nu_\Omega} - \frac{\partial u_-}{\partial \nu_\Omega} + 2\beta u = 0 & \text{on } \partial\Omega, \end{cases}$$

where ν_Ω is the normal derivative to $\partial\Omega$. Now, using the recent results [4] and [6], we get that u_+ and u_- are as regular as the free boundary $\partial\Omega$ (see Lemma 5.4). On the other hand, using variations of u along smooth vector fields, we obtain that $\text{Reg}(\partial\Omega)$ solves an equation of the form

$$\text{“Mean curvature of } \partial\Omega\text{”} = F(\nabla u_+, \nabla u_-, u_\pm) \quad \text{on } \partial\Omega,$$

where F is an explicit (rational) function of ∇u_\pm and u . In particular, this implies that $\partial\Omega$ gains one more derivative with respect to u , that is, $u \in C^{k,\alpha} \Rightarrow \partial\Omega \in C^{k+1,\alpha}$. Thus, by a bootstrap argument, we get that the regular part of the free boundary is C^∞ .

1.2. On the non-degeneracy of the solutions. We notice that the competitors (u_t, Ω_t) in (1.3) are the two-phase analogue of the ones used by Caffarelli and Kriventsov in [3], where the authors study a one-phase version of (1.1). Nevertheless, the functional in [3] involves the measure of Ω , which means that the optimality condition there corresponds to

$$J_\beta(u, \Omega) + \bar{C} |\Omega \cap \{u \leq t\}| \leq J_\beta(u_t, \Omega_t) \quad \text{where } u_t = u \vee t \quad \text{and } \Omega_t = \Omega \setminus \{u \leq t\},$$

where $\bar{C} > 0$. The presence of the constant \bar{C} allows to prove the bound from below by using a differential inequality for a suitably chosen function $f(t)$, which is given in terms of u and $\{u < t\}$ (see Proposition 4.2 and [3, Theorem 3.2]). In Proposition 4.2, we exploit the same idea, but since we do not have the constant \bar{C} , we can only conclude that $f(t) \geq \varepsilon t$ (which is not in contradiction with the fact that $f(t)$ is defined for every $t > 0$). So, we continue, and we use this lower bound to obtain a bound of the form

$$c \leq \beta^{1/2} \text{Per}(\{u < t\})^{1/2} |\{u < t\}|^{1/2} \quad \text{for every } t > 0, \quad (1.6)$$

where $u := u_\infty$ and c is a constant depending on β and d . Then, we notice that this entails

$$c \leq \beta^{3/4} \text{Per}(\{u < t\})^{1/4} |\{u < t\}|^{3/4} \quad \text{for every } t > 0.$$

and we use an iteration procedure to get that

$$c \leq \beta^{1-1/2^n} \text{Per}(\{u < t\})^{1/2^n} |\{u < t\}|^{1-1/2^n} \quad \text{for every } t > 0.$$

Passing to the limit as $n \rightarrow \infty$, we get that if u is not bounded away from zero, then

$$c \leq \beta |\{u < t\}| \leq \beta |D| \quad \text{for every } t > 0. \quad (1.7)$$

Now, this means that the measure of the zero-set $|\{u = 0\}|$ is bounded from below. Thus, using again the optimality of u , we get that (1.6) holds with an arbitrary small $\varepsilon > 0$ in place of β , we get that ‘

$$c \leq \varepsilon |\{u < t\}| \quad \text{for every } t > 0,$$

which is impossible.

A similar non-degeneracy result was proved by Bucur and Giacomini in [1] by a De Giorgi iteration scheme¹. Precisely, one can prove that any solution to (1.1) satisfies the optimality condition from [1, Remark 3.7]. Thus, [1, Theorem 3.5] also holds for the solutions of (1.1). Conversely, the argument from 4.2 can be applied to the minimizers from [1] to obtain the bound from below from [1, Theorem 3.5].

1.3. One-phase and two-phase problems with Robin boundary conditions. The problem (1.1) is the first instance of a two-phase free boundary problem with Robin boundary conditions. Precisely, we notice that if Ω is a fixed set with smooth boundary and if u minimizes the functional $J_\beta(\cdot, \Omega)$ in $H^1(D)$, then the functions

$$u_+ := u \quad \text{on } \bar{\Omega} \quad \text{and} \quad u_- := u \quad \text{on } D \setminus \Omega,$$

are harmonic respectively in Ω and $D \setminus \bar{\Omega}$, and satisfy the following conditions on $\partial\Omega \cap D$:

$$u_+ = u_- \quad \text{and} \quad \left(\frac{\partial u_+}{\partial \nu_+} + \frac{\beta}{2} u_+ \right) + \left(\frac{\partial u_-}{\partial \nu_-} + \frac{\beta}{2} u_- \right) = 0 \quad \text{on } \partial\Omega \cap D, \quad (1.8)$$

where ν_+ and ν_- are the exterior and the interior normals to $\partial\Omega$. Notice that (1.8) is a two-phase counterpart of the one-phase problem

$$\Delta u = 0 \quad \text{in } \Omega, \quad \frac{\partial u}{\partial \nu} + \beta u = 0 \quad \text{on } \partial\Omega \cap D, \quad (1.9)$$

which was studied by Bucur-Luckhaus in [2] and Caffarelli-Kriventsov in [3]. As explained in [3], the Robin condition in (1.9) naturally arises in the physical situation in which the heat diffuses freely in Ω , the temperature is set to be zero on the surface $\partial\Omega$, which is separated from the interior of Ω by an infinitesimal insulator. The two-phase problem (1.8) also may be interpreted in this way, in this case the heat diffuses freely both inside Ω and outside, in $D \setminus \bar{\Omega}$; the temperature is set to be zero on the surface $\partial\Omega$, which is insulated from both sides; the continuity of the temperature means that the heat transfer is allowed also across $\partial\Omega$, which happens for instance if the surface $\partial\Omega$ is replaced by a very thin (infinitesimal) net.

Even if the problems in [2]-[3] and in the present paper lead to the free boundary conditions of the same type, the techniques are completely different. For instance, the problem studied in [2]-[3] is a free discontinuity problem as the function u jumps from positive in Ω to zero in $D \setminus \bar{\Omega}$. Thus, the corresponding variational minimization problem can be naturally stated in the class of SBV functions, which clearly influences both the existence and the regularity techniques; roughly speaking, the existence is obtained through a compactness theorem in the SBV class, while the regularity relies on techniques related to the Mumford-Shah functional.

¹We are grateful to the anonymous referee for bringing to our attention the reference [1].

In our case, the problem can be stated for the functions (u_1, u_2) with disjoint supports ($u_1 u_2 = 0$ almost-everywhere in D) which satisfy the following constraints: the sum $u_1 + u_2$ should be a Sobolev function (this corresponds to the continuity condition in (1.8)); u_1^2 and u_2^2 are SBV functions whose jump sets are contained in the boundary of the positivity sets $\{u_1 > 0\}$ and $\{u_2 > 0\}$. Now, it is reasonable to expect that an existence result can be proved also in this class, but then, in order to prove that a solution to (1.1) exists, one should show that u_1 and u_2 are of the form $u_1 = u \mathbb{1}_\Omega$ and $u_2 = u \mathbb{1}_{D \setminus \Omega}$ for a set of finite perimeter $\Omega \subset \mathbb{R}^d$, u being the sum $u_1 + u_2$. Summarizing, working in the class of SBV functions would allow to state (1.1) in a weaker form, but it doesn't seem to be a shortcut to the existence of a solution (of (1.1)) as it will require the analysis of the jump sets of the optimal couples in the SBV class. Thus, we prefer not to rely on the advanced compactness results for SBV functions, but to prove the existence of a solution from scratch.

Finally, as explained in Section 1.1, once we know that an optimal couple (u, Ω) exists, and that u is non-degenerate and Hölder continuous, the regularity of the free boundary $\partial\Omega$ follows immediately since the set Ω becomes an almost-minimizer of the perimeter.

2. PRELIMINARIES

2.1. Sets of finite perimeter. Let $A \subset \mathbb{R}^d$ be an open set in \mathbb{R}^d . We recall that the set $E \subset \mathbb{R}^d$ is said to have a *finite perimeter in A* if

$$\text{Per}(E, A) = \sup \left\{ \int_A \text{div } \xi(x) \, dx : \xi \in C_c^1(A; \mathbb{R}^d), \sup_{x \in \mathbb{R}^d} |\xi(x)| \leq 1 \right\}, \quad (2.1)$$

is finite. We say that E has a *locally finite perimeter in A*, if for every open set $B \subset \mathbb{R}^d$ such that $\overline{B} \subset A$, we have that $\text{Per}(E, B) < \infty$. We say that E is of *finite perimeter* if

$$\text{Per}(E) := \text{Per}(E, \mathbb{R}^d) < +\infty.$$

By the De Giorgi structure theorem (see for instance [10, Theorem II.4.9]), if the set $E \subset \mathbb{R}^d$ has locally finite perimeter in A , then there is a set $\partial^* E \subset A \cap \partial E$ called *reduced boundary* such that

$$\text{Per}(E, B) = \mathcal{H}^{d-1}(B \cap \partial^* E) \quad \text{for every set } B \subset\subset A,$$

where \mathcal{H}^{d-1} is the $(d-1)$ -dimensional Hausdorff measure in \mathbb{R}^d . Moreover, there is a \mathcal{H}^{d-1} -measurable map $\nu_E : \partial^* E \rightarrow \mathbb{R}^d$, called *generalized normal* such that $|\nu_E| = 1$ and

$$\int_E \text{div } \xi(x) \, dx = \int_{\partial^* E} \nu_E \cdot \xi \, d\mathcal{H}^{d-1} \quad \text{for every } \xi \in C_c^1(A; \mathbb{R}^d).$$

2.2. Capacity and traces of Sobolev functions. We define the capacity (or the 2-capacity) of a set $E \subset \mathbb{R}^d$ as

$$\text{cap}(E) = \inf \left\{ \|u\|_{H^1(\mathbb{R}^d)}^2 : u \in H^1(\mathbb{R}^d), u \geq 1 \text{ in a neighborhood of } E \right\}.$$

Suppose now that $d \geq 3$. It is well-known that the sets of zero capacity have zero $d-1$ dimensional Hausdorff measure (see for instance [7, Section 4.7.2, Theorem 4]), that is,

$$\text{If } \text{cap}(E) = 0, \quad \text{then } \mathcal{H}^{d-1}(E) = 0.$$

Moreover, the Sobolev functions are defined up to a set of zero capacity (i.e. *quasi-everywhere*), that is, if $A \subset \mathbb{R}^d$ is an open set and $u \in H^1(A)$, then there is a set $\mathcal{N}_u \subset \mathbb{R}^d$ such that $\text{cap}(\mathcal{N}_u) = 0$ and

$$u(x_0) = \lim_{r \rightarrow 0} \frac{1}{|B_r|} \int_{B_r(x_0)} u(x) \, dx \quad \text{for every } x_0 \in A \setminus \mathcal{N}_u.$$

Moreover, for every function $u \in H^1(A)$ there is a sequence $u_n \in C^\infty(A) \cap H^1(A)$ and a set $\mathcal{N} \subset A$ of zero capacity such that:

- u_n converges to u strongly in $H^1(A)$;

- $u(x) = \lim_{n \rightarrow \infty} u_n(x)$ for every $x \in A \setminus (\mathcal{N} \cup \mathcal{N}_u)$.

In particular, if $E \subset \mathbb{R}^d$ is a set of locally finite perimeter in the open set $A \subset \mathbb{R}^d$ and if $u \in H^1(A)$, then the function u^2 is defined \mathcal{H}^{d-1} -almost everywhere on $\partial^* E$ and is \mathcal{H}^{d-1} -measurable on $\partial^* E$. Thus, the integral

$$\mathcal{I}(u, E) := \int_{A \cap \partial^* E} u^2 d\mathcal{H}^{d-1} \quad \text{is well-defined.}$$

As a consequence of the discussion above, we have the following proposition.

Proposition 2.1. *Suppose that D is a smooth bounded open set in \mathbb{R}^d and that $u \in H^1(D)$ is a Sobolev function. Then, there is a set $\mathcal{N}_u \subset D$ such that $\mathcal{H}^{d-1}(\mathcal{N}_u) = 0$ and*

$$u(x_0) = \lim_{r \rightarrow 0} \frac{1}{|B_r|} \int_{B_r(x_0)} u(x) dx \quad \text{for every } x_0 \in D \setminus \mathcal{N}_u.$$

Moreover, if $E \subset \mathbb{R}^d$ is a set of locally finite perimeter in \mathbb{R}^d , then the function $u : \partial^* E \cap D \rightarrow \mathbb{R}$ is defined \mathcal{H}^{d-1} -almost everywhere and is \mathcal{H}^{d-1} -measurable on $\partial^* E \cap D$. In particular, the integral $\mathcal{I}(u, D)$ is well-defined.

Remark 2.2. In the case $d = 2$, (2.1) still holds. In fact, it is sufficient to notice that if $u \in H^1(D)$, then $u \in W^{1,p}(D)$ for any $1 < p < 2$. In particular, it is sufficient to results from [7], this time in the space $W^{1,p}(D)$, for p close to 2.

In the next subsection, we will go through the main properties of this functional, which we will need in the proof of Theorem 1.1.

2.3. Properties of the functional \mathcal{I} . We first notice that we can use an integration by parts to write \mathcal{I} as in (2.1).

Lemma 2.3. *Let $E \subset \mathbb{R}^d$ be a set of locally finite perimeter in the open set $A \subset \mathbb{R}^d$ and let $u \in H^1(A)$ be locally bounded in A . Then, the following holds.*

- (i) *For every $\xi \in C_c^1(A; \mathbb{R}^d)$ we have*

$$\int_{A \cap \partial^* E} (\xi \cdot \nu_E) u^2 d\mathcal{H}^{d-1} = \int_A \operatorname{div}(u^2 \xi) dx. \quad (2.2)$$

- (ii) *We have the formula*

$$\int_{A \cap \partial^* E} u^2 d\mathcal{H}^{d-1} = \sup \left\{ \int_{A \cap \partial^* E} (\xi \cdot \nu_E) u^2 d\mathcal{H}^{d-1} : \xi \in C_c^1(A; \mathbb{R}^d), |\xi| \leq 1 \right\}. \quad (2.3)$$

Proof. The first claim follows by a classical approximation argument with functions of the form $\phi_n * u$, where ϕ_n is a sequence of mollifiers. In order to prove claim (ii), we notice that

$$\int_{A \cap \partial^* E} u^2 d\mathcal{H}^{d-1} \leq \sup \left\{ \int_{A \cap \partial^* E} (\xi \cdot \nu_E) u^2 d\mathcal{H}^{d-1} : \xi \in C_c^1(A; \mathbb{R}^d), |\xi| \leq 1 \right\}.$$

Thus, it is sufficient to find a sequence $\xi_n \in C_c^1(A; \mathbb{R}^d)$, $|\xi_n| \leq 1$, such that

$$\int_{A \cap \partial^* E} u^2 d\mathcal{H}^{d-1} = \lim_{n \rightarrow \infty} \int_{A \cap \partial^* E} (\xi_n \cdot \nu_E) u^2 d\mathcal{H}^{d-1}.$$

Let A_n be a sequence of open sets such that $A_n \subset \subset A$ and $\mathbb{1}_{A_n} \rightarrow \mathbb{1}_A$. Then

$$\int_{A \cap \partial^* E} u^2 d\mathcal{H}^{d-1} = \lim_{n \rightarrow \infty} \int_{A_n \cap \partial^* E} u^2 d\mathcal{H}^{d-1}.$$

Setting $M_n = \sup_{A_n} u^2$, we can find $\xi_n \in C_c^1(A; \mathbb{R}^d)$ such that $|\xi_n| \leq 1$, and

$$0 \leq \operatorname{Per}(E, A_n) - \int_{\partial^* E} (\xi_n \cdot \nu_E) d\mathcal{H}^{d-1} \leq \frac{1}{nM_n}.$$

In particular, this implies that

$$0 \leq \int_{A_n \cap \partial^* E} u^2 d\mathcal{H}^{d-1} - \int_{A_n \cap \partial^* E} (\xi_n \cdot \nu_E) u^2 d\mathcal{H}^{d-1} \leq \frac{1}{n},$$

which concludes the proof. \square

Lemma 2.4 (Main semicontinuity lemma). *Suppose that $A \subset \mathbb{R}^d$ is a bounded open set and that $h : A \rightarrow \mathbb{R}$ is a non-negative function in $L^1(A)$. Let $u_n \in H^1(A)$ be a sequence of functions and $\Omega_n \subset \mathbb{R}^d$ be a sequence of sets of locally finite perimeter in A such that:*

- (a) $0 \leq u_n \leq h$ in A , for every $n \in \mathbb{N}$;
- (b) there is a function $u_\infty \in H^1(A)$ such that u_n converges to u_∞ weakly in $H^1(A)$ and pointwise almost-everywhere in A ;
- (c) there is a set $\Omega_\infty \subset \mathbb{R}^d$ of locally finite perimeter in A such that the sequence of characteristic functions $\mathbb{1}_{\Omega_n}$ converges to $\mathbb{1}_{\Omega_\infty}$ pointwise almost-everywhere in A .

Then,

$$\int_{A \cap \partial^* \Omega_\infty} u_\infty^2 d\mathcal{H}^{d-1} \leq \liminf_{n \rightarrow \infty} \int_{A \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1}. \quad (2.4)$$

Proof. Notice that, for every $u \in H^1(A)$ and every set of finite perimeter Ω , we have

$$\int_{A \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1} = \sup \left\{ \int_{A \cap \partial^* \Omega} (\xi \cdot \nu_\Omega) u^2 d\mathcal{H}^{d-1} : \xi \in C_c^1(A; \mathbb{R}^d), |\xi| \leq 1 \right\},$$

where ν_Ω denotes the exterior normal to $\partial^* \Omega$. We use the notation

$$\nu_n := \nu_{\Omega_n} \quad \text{and} \quad \nu_\infty := \nu_{\Omega_\infty}.$$

Let now $\xi \in C_c^1(A; \mathbb{R}^d)$, $|\xi| \leq 1$ be fixed. By the divergence theorem, we have

$$\begin{aligned} \liminf_{n \rightarrow \infty} \int_{A \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} &\geq \liminf_{n \rightarrow \infty} \int_{A \cap \Omega_n} \operatorname{div}(u_n^2 \xi) dx = \liminf_{n \rightarrow \infty} \int_A \mathbb{1}_{\Omega_n} \operatorname{div}(u_n^2 \xi) dx \\ &= \liminf_{n \rightarrow \infty} \int_A \left(2(u_n \mathbb{1}_{\Omega_n} \xi) \cdot \nabla u_n + (u_n \mathbb{1}_{\Omega_n}) (u_n \operatorname{div} \xi) \right) dx \\ &= \int_A \left(2(u_\infty \mathbb{1}_{\Omega_\infty} \xi) \cdot \nabla u_\infty + (u_\infty \mathbb{1}_{\Omega_\infty}) (u_\infty \operatorname{div} \xi) \right) dx \\ &= \int_{A \cap \partial^* \Omega_\infty} \operatorname{div}(u_\infty^2 \xi) dx = \int_{A \cap \partial^* \Omega_\infty} (\xi \cdot \nu_\infty) u_\infty^2 d\mathcal{H}^{d-1}, \end{aligned}$$

where in order to pass to the limit we used that the sequence $u_n \mathbb{1}_{\Omega_n}$ converges strongly in $L_{loc}^2(A)$ to $u_\infty \mathbb{1}_{\Omega_\infty}$, as a consequence of the fact that it converges pointwise a.e. and is bounded by h . Now, taking the supremum over ξ , we get (2.4). \square

3. A FAMILY OF APPROXIMATING PROBLEMS

We use the notations $D, \beta, E, v, \mathcal{E}, \mathcal{V}$ from Section 1. Moreover, we fix a constant

$$\varepsilon \in [0, m),$$

where m is the lower bound of the function v , and we consider the auxiliary problem

$$\min \{ J_\beta(\Omega, u) : \Omega \in \mathcal{E}, u \in \mathcal{V}, u \geq \varepsilon \text{ in } \mathbb{R}^d \}. \quad (3.1)$$

Proposition 3.1 (Existence of a solution). *Let \mathcal{E} and \mathcal{V} be as above. Then, for every $0 < \varepsilon < m$, there is a solution to the problem (3.1).*

Proof. Let (u_n, Ω_n) be a minimizing sequence for (3.1). Since

$$\int_D |\nabla u_n|^2 dx + \int_{\partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} = J_\beta(u_n, \Omega_n) \leq J_\beta(v, E),$$

for every $n \in \mathbb{N}$, we have

$$\int_D |\nabla u_n|^2 dx \leq J_\beta(v, E) \quad \text{and} \quad \text{Per}(\Omega_n) \leq \frac{1}{\beta \varepsilon^2} J_\beta(v, E).$$

Thus, there are subsequences u_n and Ω_n such that:

- u_n converges strongly in $L^2(D)$, weakly in $H^1(D)$ and pointwise almost-everywhere to a function $u_\infty \in H^1(D)$;
- $\mathbb{1}_{\Omega_n}$ converges to $\mathbb{1}_{\Omega_\infty}$ strongly in $L^1(D)$ and pointwise almost-everywhere.

Moreover, we can assume that $u_n \leq h$ on D , where h is the harmonic function :

$$\Delta h = 0 \quad \text{in } D, \quad h - v \in H_0^1(D).$$

Indeed, we have

$$\begin{aligned} \int_D |\nabla u_n|^2 dx &= \int_D |\nabla(u_n \wedge h)|^2 dx + \int_D |\nabla(u_n \vee h)|^2 dx - \int_D |\nabla h|^2 dx \geq \int_D |\nabla(u_n \wedge h)|^2 dx, \\ \text{and} \quad \int_{\partial^* \Omega} u_n^2 d\mathcal{H}^{d-1} &\geq \int_{\partial^* \Omega} (u_n \wedge h)^2 d\mathcal{H}^{d-1}, \end{aligned}$$

which gives that

$$J_\beta(u_n \wedge h, \Omega_n) \leq J_\beta(u_n, \Omega_n).$$

On the other hand, we have that

$$J_\beta(u_n \vee 0, \Omega_n) \leq J_\beta(u_n, \Omega_n).$$

Thus, we can assume that $0 \leq u_n \leq h$, for every $n \in \mathbb{N}$, and so the hypotheses of Lemma 2.4 are satisfied, which means that (2.4) holds. Moreover, by the semicontinuity of the H^1 norm we have

$$\int_D |\nabla u_\infty|^2 dx \leq \liminf_{n \rightarrow \infty} \int_D |\nabla u_n|^2 dx,$$

which finally implies that

$$J_\beta(u_\infty, \Omega_\infty) \leq \liminf_{n \rightarrow \infty} J_\beta(u_n, \Omega_n). \quad \square$$

Lemma 3.2 (Subharmonicity of the solutions). *Let $m > 0$, $\beta > 0$ and $\varepsilon \in [0, m)$ be fixed. Let the function $u_\varepsilon \in H^1(D)$ and the set of finite perimeter Ω_ε be such that the couple $(u_\varepsilon, \Omega_\varepsilon)$ is a solution to the problem (3.1). Then u_ε is subharmonic in D and there is a positive Radon measure μ_ε such that*

$$-\int_D \nabla u_\varepsilon \cdot \nabla \varphi dx = \int_D \varphi d\mu_\varepsilon \quad \text{for every } \varphi \in H_0^1(D).$$

Remark 3.3. μ_ε is the distributional Laplacian of u_ε . We will use the notation $\mu_\varepsilon = \Delta u_\varepsilon$.

Proof. Let $\varphi \leq u_\varepsilon$ be a function in $H^1(D)$ such that $\varphi = u_\varepsilon$ on ∂D . Then, testing the optimality of $(u_\varepsilon, \Omega_\varepsilon)$ with $(\varphi \vee \varepsilon, \Omega_\varepsilon)$ and using the fact that $u_\varepsilon \geq \varphi \vee \varepsilon$, we get

$$\begin{aligned} \int_D |\nabla \varphi|^2 dx &\geq \int_D |\nabla(\varphi \vee \varepsilon)|^2 dx \\ &\geq \int_D |\nabla u_\varepsilon|^2 dx + \int_{\partial^* \Omega_\varepsilon} u_\varepsilon^2 d\mathcal{H}^{d-1} - \int_{\partial^* \Omega_\varepsilon} (\varphi \vee \varepsilon)^2 d\mathcal{H}^{d-1} \geq \int_D |\nabla u_\varepsilon|^2 dx, \end{aligned}$$

which concludes the proof. \square

We will next show that the family of solutions $\{u_\varepsilon\}_{\varepsilon \in (0, m)}$ is uniformly Hölder continuous. We will use the following lemma, which can be proved in several different ways. Here, we give a short proof based on the mean-value formula for subharmonic functions. Similar argument was used to prove the Lipschitz continuity of the solutions to some free boundary problems (see for instance [12] and the references therein).

Lemma 3.4 (A general condition for the Hölder continuity). *Let D be a bounded open set in \mathbb{R}^d and let $h \in L_{loc}^\infty(D)$. Suppose that $u \in H^1(D)$ is such that*

- (a) $0 \leq u \leq h$ in D ;
- (b) u is subharmonic in D ;
- (c) there are constants $K > 0$ and $\alpha \in [0, 1)$ such that

$$\Delta u(B_r(x_0)) \leq K r^{d-1-\alpha} \quad \text{for every } x_0 \in \overline{D}_\delta \quad \text{and every } 0 < r < \frac{\delta}{2}, \quad (3.2)$$

where $\delta > 0$ and

$$D_\delta := \{x \in D : \text{dist}(x, \partial D) > \delta\}. \quad (3.3)$$

Then, there is a constant C depending on δ , h , α and K such that

$$|u(x) - u(y)| \leq C|x - y|^{\frac{1-\alpha}{2-\alpha}} \quad \text{for every } x, y \in \overline{D}_\delta.$$

Proof. We first notice that the following formula is true for every subharmonic function $u \in H^1(D)$ and for every $x_0 \in D$ and $0 < s < t < \text{dist}(x_0, \partial D)$.

$$\int_{\partial B_t(x_0)} u d\mathcal{H}^{d-1} - \int_{\partial B_s(x_0)} u d\mathcal{H}^{d-1} = \frac{1}{d\omega_d} \int_s^t r^{1-d} \Delta u(B_r(x_0)) dr.$$

In particular, the function

$$r \mapsto \int_{\partial B_r(x_0)} u d\mathcal{H}^{d-1},$$

is monotone and we can define the function u pointwise *everywhere* as

$$u(x_0) := \lim_{r \rightarrow 0} \int_{\partial B_r(x_0)} u d\mathcal{H}^{d-1}.$$

As a consequence, for every $R < \text{dist}(x_0, \partial D)$, we have

$$\int_{\partial B_R(x_0)} u d\mathcal{H}^{d-1} - u(x_0) = \frac{1}{d\omega_d} \int_0^R r^{1-d} \Delta u(B_r(x_0)) dr.$$

Now, applying (3.2), and integrating in r , we get that if $x_0 \in \overline{D}_\delta$ and $R < \frac{\delta}{2}$, then

$$0 \leq \int_{\partial B_R(x_0)} u d\mathcal{H}^{d-1} - u(x_0) \leq C R^{1-\alpha} \quad \text{where} \quad C := \frac{K}{d\omega_d(1-\alpha)}, \quad (3.4)$$

which, by the subharmonicity of u , implies

$$0 \leq \int_{B_R(x_0)} u dx - u(x_0) \leq C R^{1-\alpha}. \quad (3.5)$$

Let now $x_0, y_0 \in \overline{D}_\delta$ be such that

$$|x_0 - y_0| \leq 1 \quad \text{and} \quad R := |x_0 - y_0|^\gamma \leq \frac{\delta}{4},$$

where $\gamma \in (0, 1)$ will be chosen later.

Now, since $B_R(x_0) \subset B_{R+|x_0-y_0|}(y_0) \subset B_{2R}(y_0) \subset D$, we can estimate

$$\begin{aligned} u(x_0) - u(y_0) &\leq \int_{B_R(x_0)} u(x) dx - u(y_0) \leq \frac{(R + |x_0 - y_0|)^d}{R^d} \int_{B_{R+|x_0-y_0|}(y_0)} u(x) dx - u(y_0) \\ &= (1 + |x_0 - y_0|^{1-\gamma})^d \int_{B_{R+|x_0-y_0|}(y_0)} u(x) dx - u(y_0) \\ &\leq \left(1 + d2^{d-1}|x_0 - y_0|^{1-\gamma}\right) \int_{B_{R+|x_0-y_0|}(y_0)} u(x) dx - u(y_0), \end{aligned}$$

where in the last inequality we used that $|x_0 - y_0|^{1-\gamma} \leq 1$. Now, using (3.5), we get

$$\begin{aligned} u(x_0) - u(y_0) &\leq \left(\int_{B_{R+|x_0-y_0|}(y_0)} u(x) dx - u(y_0) \right) + d2^{d-1}|x_0 - y_0|^{1-\gamma} \|u\|_{L^\infty(B_{2R}(y_0))} \\ &\leq C(R + |x_0 - y_0|)^{1-\alpha} + d2^{d-1}|x_0 - y_0|^{1-\gamma} \|h\|_{L^\infty(B_{2R}(y_0))} \\ &\leq \left(2C + d2^{d-1}M_{\delta/2} \right) |x_0 - y_0|^{1-\gamma}, \end{aligned}$$

where $M_{\delta/2}$ is the maximum of h on the set $\overline{D}_{\delta/2}$ and where we choose $\gamma = \frac{1}{2-\alpha}$, which implies that $\gamma(1-\alpha) = 1-\gamma$ and $1-\gamma = \frac{1-\alpha}{2-\alpha}$. \square

Proposition 3.5 (Hölder continuity of the solution). *Let $m > 0$, $\beta > 0$ and $\varepsilon \in [0, m]$ be fixed. Let the function $u_\varepsilon \in H^1(D)$ and the set of finite perimeter Ω_ε be such that the couple $(u_\varepsilon, \Omega_\varepsilon)$ is a solution to the problem (3.1) with some $v \in H^1(D)$ and $E \subset \mathbb{R}^d$. Then, for every $\delta > 0$, there is a constant C depending on D , δ and v (but not on ε) such that*

$$|u_\varepsilon(x) - u_\varepsilon(y)| \leq C|x - y|^{1/3} \quad \text{for every } x, y \in D_\delta.$$

Proof. By Lemma 3.2, we have that u_ε is subharmonic and, in particular, $0 \leq u_\varepsilon \leq h$ in D , where h is the harmonic extension of v in D . Thus, it is sufficient to prove that (3.2) holds. Let $x_0 \in \overline{D}_\delta$ and $R \leq \frac{\delta}{2}$. Let $\varphi \in C_c^\infty(B_{3R/2}(x_0))$ be such that

$$\varphi = 1 \quad \text{on } B_R(x_0), \quad |\nabla \varphi| \leq \frac{3}{R} \quad \text{in } B_{3R/2}(x_0).$$

Now, we test the optimality of $(u_\varepsilon, \Omega_\varepsilon)$ with $(\tilde{u}_\varepsilon, \tilde{\Omega}_\varepsilon)$, where

$$\tilde{u}_\varepsilon = u_\varepsilon + R^{1/2}\varphi \quad \text{and} \quad \tilde{\Omega}_\varepsilon = \Omega_\varepsilon \cup B_{3R/2}(x_0).$$

Thus, we get

$$\int_D |\nabla u_\varepsilon|^2 dx + \beta \int_{\partial^* \Omega_\varepsilon} u_\varepsilon^2 d\mathcal{H}^{d-1} \leq \int_D |\nabla \tilde{u}_\varepsilon|^2 dx + \beta \int_{\partial^* \tilde{\Omega}_\varepsilon} \tilde{u}_\varepsilon^2 d\mathcal{H}^{d-1}$$

and

$$\begin{aligned} \int_{\partial^* \tilde{\Omega}_\varepsilon} \tilde{u}_\varepsilon^2 d\mathcal{H}^{d-1} &\leq \int_{\partial^* \Omega_\varepsilon \setminus B_{3R/2}(x_0)} u_\varepsilon^2 d\mathcal{H}^{d-1} + \int_{\partial B_{3R/2}(x_0)} u_\varepsilon^2 d\mathcal{H}^{d-1} \\ &\leq \int_{\partial^* \Omega_\varepsilon} u_\varepsilon^2 d\mathcal{H}^{d-1} + C_d R^{d-1} M_{\delta/4}^2, \end{aligned}$$

where $M_\rho := \sup \{h(x) : x \in \overline{D}_\rho\}$. Thus, we obtain

$$2R^{1/2} \int_{B_{3R/2}(x_0)} -\nabla u_\varepsilon \cdot \nabla \varphi dx \leq R \int_{B_{3R/2}(x_0)} |\nabla \varphi|^2 dx + \beta C_d R^{d-1} M_{\delta/4}^2 \leq C_d (1 + \beta M_{\delta/4}^2) R^{d-1},$$

which implies that

$$\Delta u_\varepsilon(B_R(x_0)) \leq C_d (1 + \beta M_{\delta/4}^2) R^{d-3/2},$$

which concludes the proof of (3.2) with $\alpha = 1/2$. \square

4. EXISTENCE OF AN OPTIMAL SET

4.1. Definition of (u_0, Ω_0) . Now, for any $\varepsilon \in (0, m)$, we consider the solution $(u_\varepsilon, \Omega_\varepsilon)$ of (3.1). As a consequence of Proposition (3.5), we can find a sequence $\varepsilon_n \rightarrow 0$ and a function $u_0 \in H^1(D) \cap C^{0,1/3}(D)$ such that :

- u_{ε_n} converges to u_0 uniformly on every set D_δ , $\delta > 0$, where D_δ is defined in (3.3);
- u_{ε_n} converges to u_0 strongly in $L^2(D)$;
- u_{ε_n} converges to u_0 weakly in $H^1(D)$.

Our aim in this section is to show that u_0 is a solution to (1.1).

The construction of Ω_0 is more delicate. First, we fix $t > 0$ and $\delta > 0$ and we notice that the perimeter of Ω_{ε_n} is bounded on the open set $\{u_0 > t\} \cap D_\delta$. Indeed, the uniform convergence of u_{ε_n} to u_0 implies that, for n large enough ($n \geq N_{t,\delta}$, for some fixed $N_{t,\delta} \in \mathbb{N}$),

$$u_{\varepsilon_n} \geq \frac{t}{2} \quad \text{on} \quad D_\delta \cap \{u_0 > t\}.$$

Thus, we have

$$J_\beta(v, E) \geq \beta \int_{D_\delta \cap \{u_0 > t\} \cap \partial^* \Omega_{\varepsilon_n}} u_{\varepsilon_n}^2 d\mathcal{H}^{d-1} \geq \frac{\beta t^2}{2} \text{Per}(\Omega_{\varepsilon_n}; D_\delta \cap \{u_0 > t\}).$$

Now, if we choose t such that $\text{Per}(\{u_0 > t\}) < \infty$ (which, by the co-area formula, is true for almost-every $t > 0$), then we have that

$$\text{Per}(\Omega_{\varepsilon_n} \cap \{u_0 > t\} \cap D_\delta) \leq C_{t,\delta} \quad \text{for every} \quad n \geq N_{t,\delta},$$

for some constant $C_{t,\delta} > 0$. Now, since all the sets $\Omega_{\varepsilon_n} \cap \{u_0 > t\} \cap D_\delta$ are contained in D and have uniformly bounded perimeter, we can find a set Ω_0 and a subsequence for which

$$\mathbb{1}_{\Omega_{\varepsilon_n} \cap \{u_0 > t\} \cap D_\delta}(x) \rightarrow \mathbb{1}_{\Omega_0 \cap \{u_0 > t\} \cap D_\delta}(x) \quad \text{for almost-every} \quad x \in D.$$

Thus, by a diagonal sequence argument, we can extract a subsequence of ε_n (still denoted by ε_n) and we can define the set $\Omega_0 \subset \mathbb{R}^d$ as the pointwise limit

$$\mathbb{1}_{\Omega_0}(x) = \lim_{n \rightarrow \infty} \mathbb{1}_{\Omega_{\varepsilon_n} \cap \{u_0 > 0\}}(x) \quad \text{for almost-every} \quad x \in \{u_0 > 0\},$$

and we notice that, by construction, $\Omega_0 \subset \{u_0 > 0\}$. Notice that, we do not know a priori that Ω_0 has finite perimeter. We only know that

$$\text{Per}(\Omega_0 \cap \{u_0 > t\} \cap D_\delta) < \infty \quad \text{for every} \quad \delta > 0 \quad \text{and almost-every} \quad t > 0.$$

which means that $\Omega_0 \cap \{u_0 > t\}$ has locally finite perimeter in D for almost-every $t > 0$.

4.2. An optimality condition. As pointed out above, we do not know if the couple (u_0, Ω_0) is even an admissible competitor for (1.1) (we need to show that $\Omega_0 \in \mathcal{E}$). Nevertheless, we can still prove that it satisfies a suitable optimality condition.

Lemma 4.1 (The optimality condition at the limit). *Let u_0 and Ω_0 be as in Section 4.1. Then, for almost-every $t > 0$, we have*

$$\int_{\{u_0 < t\}} |\nabla u_0|^2 dx \leq \beta t^2 \text{Per}(\{u_0 < t\}). \quad (4.1)$$

Proof. Let now $t > 0$ be fixed and such that the set $\{u_0 < t\}$ has finite perimeter. Then, for n large enough, we can use the couple $(u_0 \vee t, \Omega_0 \cup \{u_0 < t\})$ to test the optimality of $(u_{\varepsilon_n}, \Omega_{\varepsilon_n})$. Notice that the set $\Omega_0 \cup \{u_0 < t\}$ has finite perimeter for a.e $t \in (0, m)$, as observed in the previous section. For the sake of simplicity, we write $u_{\varepsilon_n} = u_n$, $\Omega_{\varepsilon_n} = \Omega_n$, $u_0 = u$ and $\Omega_0 = \Omega$. Thus, we have

$$\begin{aligned} & \int_D |\nabla u_n|^2 dx + \beta \int_{\{u > t\} \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} \\ & \leq \int_D |\nabla u_n|^2 dx + \beta \int_{\partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} \\ & \leq \int_D |\nabla(u \vee t)|^2 dx + \beta \int_{\partial^*(\Omega \cup \{u < t\})} u^2 d\mathcal{H}^{d-1} \\ & \leq \int_D |\nabla(u \vee t)|^2 dx + \beta t^2 \text{Per}(\{u < t\}) + \beta \int_{\{u > t\} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1}. \end{aligned} \quad (4.2)$$

Now, by the weak convergence of u_n to u , we get that

$$\int_D |\nabla u|^2 dx \leq \liminf_{n \rightarrow \infty} \int_D |\nabla u_n|^2 dx.$$

On the other hand, setting $U_{t,\delta}$ to be the open set

$$U_{t,\delta} = \mathbb{R}^d \setminus (\overline{D}_\delta \cap \{u \leq t\}),$$

for some fixed $\delta > 0$, and applying Lemma 2.4, we have that

$$\int_{U_{t,\delta} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1} \leq \liminf_{n \rightarrow \infty} \int_{U_{t,\delta} \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} \leq \liminf_{n \rightarrow \infty} \int_{\{u > t\} \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1}.$$

Taking the limit as $\delta \rightarrow 0$, by the monotone convergence theorem, we get that

$$\lim_{\delta \rightarrow 0} \int_{U_{t,\delta} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1} = \int_{(\mathbb{R}^d \setminus (D \cap \{u \leq t\})) \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1}$$

Now since

$u(x) = h(x)$ for quasi-every $x \in \mathbb{R}^d \setminus D$ and for \mathcal{H}^{d-1} -almost-every $x \in \mathbb{R}^d \setminus D$, and since $h \geq m > t$ on ∂D , we have that

$$\int_{(\mathbb{R}^d \setminus (D \cap \{u \leq t\})) \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1} = \int_{\{u > t\} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1}. \quad (4.3)$$

Thus, we get that

$$\int_{\{u > t\} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1} \leq \liminf_{n \rightarrow \infty} \int_{D \cap \{u > t\} \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1}. \quad (4.4)$$

Now, using (4.4) and (4.2), we obtain

$$\begin{aligned} \int_D |\nabla u|^2 dx + \beta \int_{\{u > t\} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1} \\ \leq \liminf_{n \rightarrow \infty} \int_D |\nabla u_n|^2 dx + \beta \int_{\{u > t\} \cap \partial^* \Omega_n} u_n^2 d\mathcal{H}^{d-1} \\ \leq \int_D |\nabla(u \vee t)|^2 dx + \beta t^2 \text{Per}(\{u < t\}) + \beta \int_{\{u > t\} \cap \partial^* \Omega} u^2 d\mathcal{H}^{d-1}, \end{aligned}$$

which gives (4.1). \square

4.3. Non-degeneracy. The crucial observation in this section is that the functions u satisfying the optimality condition (4.1) are non-degenerate in the sense of the following proposition.

Proposition 4.2 (Non-degeneracy). *Let $\beta > 0$, $m > 0$, D be a bounded open set of \mathbb{R}^d and $u \in H^1(D)$ be a non-negative function in D such that $u \geq m$ on ∂D . Let $\Omega \subset D$ be a set of finite perimeter in D . Suppose that u and Ω satisfy the optimality condition*

$$\int_{\Omega_t} |\nabla u|^2 dx \leq \beta t^2 \text{Per}(\Omega_t) \quad \text{where} \quad \Omega_t = \{u \leq t\}, \quad (4.5)$$

for almost-every $t \in (0, m)$. Then, $|\Omega_t| = 0$ for some $t > 0$.

Proof. By contradiction, suppose that

$$|\Omega_t| > 0 \quad \text{for every} \quad t > 0.$$

Let $t \in (0, m)$ be fixed. By the co-area formula, the Cauchy-Schwartz inequality and the optimality condition (4.5), we get

$$\int_{\Omega_t} |\nabla u| = \int_0^t \text{Per}(\Omega_s) ds \leq \left(\int_{\Omega_t} |\nabla u|^2 \right)^{1/2} |\Omega_t|^{1/2} \leq t \beta^{1/2} \text{Per}(\Omega_t)^{1/2} |\Omega_t|^{1/2}. \quad (4.6)$$

We now set

$$f(t) := \int_0^t \text{Per}(\Omega_s) ds = \int_{\Omega_t} |\nabla u| dx .$$

Using (4.6), we will estimate $f(t)$ from below.

Step 1. *Non-degeneracy of f .* By the isoperimetric inequality and the estimate (4.6), there is a dimensional constant C_d such that

$$\int_0^t \text{Per}(\Omega_s) ds \leq t\beta^{1/2}C_d \text{Per}(\Omega_t)^{\frac{2d-1}{2d-2}} .$$

Using the definition of f , we can re-write this inequality as

$$f(t)^{\frac{2d-2}{2d-1}} \leq t^{\frac{2d-2}{2d-1}} (\beta^{1/2}C_d)^{\frac{2d-2}{2d-1}} f'(t) .$$

After rearranging the terms and integrating from 0 to t , we obtain

$$f(t)^{\frac{1}{2d-1}} - f(0)^{\frac{1}{2d-1}} \geq \frac{t^{\frac{1}{2d-1}}}{(\beta^{1/2}C_d)^{\frac{2d-2}{2d-1}}} .$$

Now, since u is non-negative in D , we have that $f(0) = 0$. Thus

$$f(t) \geq \frac{t}{(\beta^{1/2}C_d)^{2d-2}} .$$

Setting

$$C = (\beta C_d)^{1-d} , \tag{4.7}$$

we obtain the lower bound

$$f(t) \geq Ct .$$

In particular, as a consequence of (4.6), we get that

$$C \leq \beta^{1/2} \text{Per}(\Omega_t)^{1/2} |\Omega_t|^{1/2} . \tag{4.8}$$

Step 2. *Non-degeneracy of $|\Omega_t|$.* Let $\alpha \in (0, 1)$ be fixed. Then, we have that

$$\begin{aligned} \int_0^t \text{Per}(\Omega_s)^\alpha |\Omega_s|^{1-\alpha} ds &\leq \left(\int_0^t \text{Per}(\Omega_s) ds \right)^\alpha \left(\int_0^t |\Omega_s| ds \right)^{1-\alpha} \\ &\leq \left(t\beta^{1/2} \text{Per}(\Omega_t)^{1/2} |\Omega_t|^{1/2} \right)^\alpha \left(t|\Omega_t| \right)^{1-\alpha} = t\beta^{\alpha/2} \text{Per}(\Omega_t)^{\alpha/2} |\Omega_t|^{1-\alpha/2} . \end{aligned}$$

Thus, we obtain that for fixed $T \in (0, m)$ and $C > 0$, the following implication holds :

$$\begin{aligned} \text{If } C &\leq \text{Per}(\Omega_t)^\alpha |\Omega_t|^{1-\alpha} \text{ for every } t \in (0, T), \\ \text{then } C &\leq \beta^{\alpha/2} \text{Per}(\Omega_t)^{\alpha/2} |\Omega_t|^{1-\alpha/2} \text{ for every } t \in (0, T). \end{aligned} \tag{4.9}$$

We claim that, for every $n \geq 1$ and every $t \in (0, m)$, we have the inequality

$$C \leq \beta^{1-1/2^n} \text{Per}(\Omega_t)^{1/2^n} |\Omega_t|^{1-1/2^n} . \tag{4.10}$$

In order to prove (4.10), we argue by induction on n . When $n = 1$, (4.10) is precisely (4.8). In order to prove that the claim (4.10) for $n \in \mathbb{N}$ implies the same claim for $n + 1$, we apply (4.9) for $\alpha = 2^{-n}$, $n \in \mathbb{N}$, which gives precisely (4.10) with $n + 1$. This concludes the proof of (4.10). Next, passing to the limit as $n \rightarrow \infty$, we obtain that

$$C \leq \beta |\Omega_t| \text{ for every } t \in (0, T),$$

where C is given by (4.7). Thus, there is a dimensional constant $C_d > 0$ such that

$$\beta^{-d} C_d \leq |\Omega_t| \text{ for every } t \in [0, m). \tag{4.11}$$

Step 3. *Conclusion.* We now notice that

$$\lim_{t \rightarrow 0} |\Omega_t| = |\Omega_0| > 0 .$$

Thus, for every $\varepsilon > 0$, there is T_ε such that for all $t \in (0, T_\varepsilon)$ we have

$$\int_{\Omega_t} |\nabla u| = \int_0^t \text{Per}(\Omega_s) ds \leq \left(\int_{\Omega_t} |\nabla u|^2 \right)^{1/2} |\Omega_t \setminus \Omega_0|^{1/2} \leq t\varepsilon^{1/2} \text{Per}(\Omega_t)^{1/2} |\Omega_t|^{1/2}. \quad (4.12)$$

Now, repeating the argument from Step 1 and Step 2, we get that (4.11) should hold with ε in place of β . Since $\varepsilon > 0$ is arbitrary, this is a contradiction. \square

4.4. Existence of a solution. We are now in position to prove that the couple (u_0, Ω_0) , constructed in Section 4.1, is a solution to (1.1).

Proposition 4.3 (Existence of a solution). *There is a dimensional constant $C_d > 0$ such that if D is a bounded open set of \mathbb{R}^d and $\beta > 0$ is a given positive constant, then the following holds. For every set $E \subset \mathbb{R}^d$ of finite perimeter and every $v \in H^1(\mathbb{R}^d)$ satisfying*

$$v \geq m \quad \text{on } D \quad \text{for some constant } m > 0,$$

there is a solution (u, Ω) of the problem (1.1).

Proof. Let (u_0, Ω_0) be as in Section 4.1. Then, by Lemma 4.1, (u_0, Ω_0) satisfies the optimality condition (4.5). Now, by Proposition 4.2 we get that $u_0 \geq t$ in D , for some $t > 0$. In particular, Ω_0 has finite perimeter in D . Precisely, for every $\delta > 0$, we have

$$\begin{aligned} \text{Per}(\Omega_0; D_\delta) &\leq \liminf_{n \rightarrow \infty} \text{Per}(\Omega_{\varepsilon_n}; D_\delta) \leq \frac{4}{t^2} \liminf_{n \rightarrow \infty} \int_{D_\delta \cap \partial^* \Omega_{\varepsilon_n}} u_{\varepsilon_n}^2 d\mathcal{H}^{d-1} \\ &\leq \frac{4}{\beta t^2} \liminf_{n \rightarrow \infty} J_\beta(u_{\varepsilon_n}, \Omega_{\varepsilon_n}) \leq \frac{4}{\beta t^2} J_\beta(v, E). \end{aligned}$$

Passing to the limit as $\delta \rightarrow 0$, we get

$$\text{Per}(\Omega_0; D) \leq \frac{4}{\beta t^2} J_\beta(v, E).$$

In particular, this implies that Ω_0 is a set of finite perimeter in \mathbb{R}^d . Indeed,

$$\begin{aligned} \text{Per}(\Omega_0) &\leq \text{Per}(\Omega_0; D) + 2\text{Per}(D) + \text{Per}(\Omega_0; \mathbb{R}^d \setminus \overline{D}) \\ &\leq \frac{4}{\beta t^2} J_\beta(v, E) + 2\text{Per}(D) + \text{Per}(E; \mathbb{R}^d \setminus \overline{D}). \end{aligned}$$

Thus, the couple (u_0, Ω_0) is admissible in (1.1); it now remains to prove that it is optimal. Let $\tilde{u} \in H^1(D)$ be non-negative on D and such that $u - v \in H_0^1(D)$. Let $\tilde{\Omega} \subset \mathbb{R}^d$ be a set of finite perimeter such that $\tilde{\Omega} = E$ on $\mathbb{R}^d \setminus D$. It is sufficient to prove that

$$J_\beta(u_0, \Omega_0) \leq J_\beta(\tilde{u}, \tilde{\Omega}).$$

Let $\varepsilon > 0$ be fixed. We now use the couple $(\tilde{u} \vee \varepsilon, \tilde{\Omega})$ to test the optimality of $(u_{\varepsilon_n}, \Omega_{\varepsilon_n})$:

$$J_\beta(u_{\varepsilon_n}, \Omega_{\varepsilon_n}) \leq J_\beta(\tilde{u} \vee \varepsilon, \tilde{\Omega}).$$

Passing to the limit as $\varepsilon \rightarrow 0$, we get

$$J_\beta(u_{\varepsilon_n}, \Omega_{\varepsilon_n}) \leq J_\beta(\tilde{u}, \tilde{\Omega}).$$

Now, Lemma 2.4 and the semicontinuity of the H^1 norm gives that $J_\beta(u_0, \Omega_0) \leq J_\beta(\tilde{u}, \tilde{\Omega})$, which concludes the proof. \square

5. REGULARITY OF THE FREE BOUNDARY

In this section, we prove the regularity of the free boundary. In Theorem 5.1, we prove that the solutions of (1.1) are almost-minimizers for the perimeter in D . As a consequence, $\partial\Omega$ can be decomposed into a regular and a singular part and that the regular part is $C^{1,\alpha}$ manifold. Then, in Theorem 5.2, we prove that the regular part of the free boundary is C^∞ smooth.

Theorem 5.1. *Let (u, Ω) be a solution to (1.1). there is a constant $C > 0$ such that Ω is an almost-minimizer of the perimeter in the following sense:*

$$\text{Per}(\Omega; B_r(x_0)) \leq (1 + Cr^{1/3}) \text{Per}(\Omega'; B_r(x_0)),$$

for every ball $B_r(x_0) \subset D$ and every set $\Omega' \subset \mathbb{R}^d$ such that $\Omega = \Omega'$ outside $B_r(x_0)$.

In particular, the free boundary $\partial\Omega \cap D$ can be decomposed as the disjoint union of a regular part $\text{Reg}(\partial\Omega)$ and a singular part $\text{Sing}(\partial\Omega)$, where

- (i) $\text{Reg}(\partial\Omega)$ is a relatively open subset of $\partial\Omega$ and is a $C^{1,\alpha}$ smooth manifold;
- (ii) $\text{Sing}(\partial\Omega)$ is a closed set, which is empty if $d \leq 7$, discrete if $d = 8$, and of Hausdorff dimension $d - 8$, if $d > 8$.

Proof. We first notice that by Lemma 3.4, $u \in C^{0,1/3}(D)$. Let $\delta > 0$, $x_0 \in D_\delta$ and $r < \delta/2$. We consider a set $\Omega' \subset \mathbb{R}^d$ such that $\Omega' \Delta \Omega \subset\subset B_r(x_0)$. Testing the optimality of (u, Ω) against (u, Ω') we get that

$$\int_{B_r(x_0) \cap \partial^* \Omega} u^2 d\mathcal{H}^{n-1} \leq \int_{B_r(x_0) \cap \partial^* \Omega'} u^2 d\mathcal{H}^{n-1},$$

which implies that

$$\left(\min_{B_r(x_0)} u^2 \right) \text{Per}(\Omega; B_r(x_0)) \leq \left(\max_{B_r(x_0)} u^2 \right) \text{Per}(\Omega'; B_r(x_0)).$$

By regularity of u , we have that

$$\max_{B_r(x_0)} u^2 \leq \min_{B_r(x_0)} u^2 + Cr^{1/3} \leq \left(\min_{B_r(x_0)} u^2 \right) \left(1 + \frac{C}{t} r^{1/3} \right),$$

where in the second inequality, we used that $u \geq t > 0$. Thus, we obtain

$$\text{Per}(\Omega; B_r(x_0)) \leq \left(1 + \frac{C}{t} r^{1/3} \right) \text{Per}(\Omega'; B_r(x_0)),$$

which proves that Ω is an almost-minimizer of the perimeter in D . \square

We next prove that regular part the free boundary $\text{Reg}(\partial\Omega)$ is C^∞ .

Theorem 5.2. *Let (u, Ω) be a solution to (1.1). Let*

$$D \cap \partial\Omega = \text{Reg}(\partial\Omega) \cup \text{Sing}(\partial\Omega)$$

be the decomposition of the free boundary given by Theorem 5.1. Then, in a neighborhood of any point $x_0 \in \text{Reg}(\partial\Omega)$, $\partial\Omega$ is C^∞ -regular and the function u is C^∞ on $\partial\Omega$.

Proof. We fix a point $x_0 \in \text{Reg}(\partial\Omega)$. Without loss of generality, we can assume that $x_0 = 0$.

Step 1. Notation. For any $x \in \mathbb{R}^d$, we use the notation $x = (x', x_d)$, where $x' \in \mathbb{R}^{d-1}$ and $x_d \in \mathbb{R}$. By the $C^{1,\alpha}$ regularity of $\text{Reg}(\partial\Omega)$, in a neighborhood of the origin $B' \times (-\varepsilon, \varepsilon) \subset \mathbb{R}^{d-1} \times \mathbb{R}$, $\partial\Omega$ is the graph of a $C^{1,\alpha}$ regular function $\eta : B' \rightarrow \mathbb{R}$, where B' is a ball in \mathbb{R}^{d-1} ; the set Ω coincides with the subgraph of η in a neighborhood of the origin:

$$B' \times (-\varepsilon, \varepsilon) \cap \Omega = \{(x', x_d) \in B' \times (-\varepsilon, \varepsilon) : x_d < \eta(x')\}.$$

and the exterior normal ν_Ω is given by

$$\nu_\Omega = \frac{(-\nabla_{x'} \eta, 1)}{\sqrt{1 + |\nabla_{x'} \eta|^2}}, \quad (5.1)$$

where $\nabla_{x'}\eta$ is the gradient of η in the first $d-1$ variables. Let u_+ and u_- be the restrictions of u on the sets $\bar{\Omega}$ and $D \setminus \Omega$; since u is continuous across $\partial\Omega$, we have $u_+ = u_-$ on $\partial\Omega$. Moreover, we write the gradients of u_+ and u_- as

$$\nabla u_{\pm} = (\nabla_{x'}u_{\pm}, \partial_{x_d}u_{\pm}) \in \mathbb{R}^{d-1} \times \mathbb{R}.$$

Step 2. *Transmission condition and $C^{1,\alpha}$ regularity of u .* In Lemma 5.3, we keep fixed and free boundary $\partial\Omega$ and we use vertical perturbations of the function u to obtain a Robin-type transmission condition on $\partial\Omega$. We notice that the recent results [4] and [6] imply the $C^{1,\alpha}$ -regularity of u_+ and u_- , up to the boundary $\partial\Omega$. Thus, the gradient is well-defined and the transmission conditions (5.2) hold in the classical sense.

Step 3. *Optimality condition and $C^{2,\alpha}$ regularity of $\text{Reg}(\partial\Omega)$.* In Lemma 5.5 we perform variations of the optimal set to find the geometric equation solved by $\partial\Omega$. Precisely, we find that the curvature of the optimal set solves an equation of the form

$$\text{“Mean curvature of } \partial\Omega\text{”} = F(\nabla u_+, \nabla u_-, u_{\pm}) \quad \text{on } \partial\Omega.$$

In particular, this implies that if u is $C^{k,\alpha}$, for some $k \geq 1$, then $\partial\Omega$ is $C^{k+1,\alpha}$.

Step 4. *Bootstrap and C^∞ regularity of $\partial\Omega$.* In Lemma 5.4 we use the recent results of [6] to show that if the boundary $\partial\Omega$ is $C^{k,\alpha}$ for some $k \geq 2$, then the solutions u_+ and u_- are also $C^{k,\alpha}$ regular up to the boundary $\partial\Omega$. Finally, applying this result (Lemma 5.4) and the result from the previous step (Lemma 5.5), we get that $\partial\Omega$ is C^∞ . \square

Lemma 5.3 (Robin and continuity conditions on $\partial\Omega$). *Suppose that $\partial\Omega$ is $C^{1,\alpha}$ regular in the neighborhood of the origin. Let $\eta : B' \rightarrow \mathbb{R}$, u_+ and u_- be as above. Then, for every $x' \in B'$ we have*

$$\begin{cases} \nabla_{x'}\eta \cdot \nabla_{x'}u_+ - \nabla_{x'}\eta \cdot \nabla_{x'}u_- = -(\partial_{x_d}u_+ - \partial_{x_d}u_-)|\nabla_{x'}\eta|^2 \\ \sqrt{1 + |\nabla_{x'}\eta|^2}(\partial_{x_d}u_+ - \partial_{x_d}u_-) + \beta u = 0, \end{cases} \quad (5.2)$$

where u_+ , u_- and their partial derivatives are calculated in $(x', \eta(x')) \in \partial\Omega$.

Proof. Let $\phi \in C_c^\infty(D)$ be a smooth function supported in $B' \times (-\varepsilon, \varepsilon)$. Then, the optimality of u gives that

$$\begin{aligned} 0 &= \left. \frac{\partial}{\partial t} \right|_{t=0} J_\beta(u + t\phi, \Omega) = \int_{D \setminus \partial\Omega} 2\nabla u \cdot \nabla \phi \, dx + \beta \int_{\partial\Omega} 2u\phi \, d\mathcal{H}^{d-1} \\ &= \int_{\partial\Omega} 2(\nu_\Omega \cdot \nabla u_+ - \nu_\Omega \cdot \nabla u_- + \beta u)\phi \, d\mathcal{H}^{d-1}, \end{aligned}$$

where in the last inequality we integrated by parts u_+ in Ω and u_- in $D \setminus \Omega$. Since ϕ is arbitrary we get that u satisfies the Robin-type condition on $\partial\Omega$

$$\nu_\Omega \cdot \nabla u_+ - \nu_\Omega \cdot \nabla u_- + \beta u \quad \text{on } \partial\Omega. \quad (5.3)$$

Now, using (5.1), we can re-write this as

$$(-\nabla_{x'}\eta \cdot \nabla_{x'}u_+ + \partial_{x_d}u_+) - (-\nabla_{x'}\eta \cdot \nabla_{x'}u_- + \partial_{x_d}u_-) + \beta u \sqrt{1 + |\nabla_{x'}\eta|^2} = 0. \quad (5.4)$$

On the other hand u is continuous across $\partial\Omega$. This means that

$$\nabla_{x'}u_+(x', \eta(x')) + \partial_{x_d}u_+(x', \eta(x'))\nabla_{x'}\eta = \nabla_{x'}u_-(x', \eta(x')) + \partial_{x_d}u_-(x', \eta(x'))\nabla_{x'}\eta.$$

Multiplying by $\nabla_{x'}\eta$, we get

$$\nabla_{x'}\eta \cdot \nabla_{x'}u_+ + \partial_{x_d}u_+|\nabla_{x'}\eta|^2 = \nabla_{x'}\eta \cdot \nabla_{x'}u_- + \partial_{x_d}u_-|\nabla_{x'}\eta|^2, \quad (5.5)$$

where u_+ , u_- and their partial derivatives are calculated in $(x', \eta(x'))$. Putting together (5.4) and (5.5), we get (5.2). \square

Lemma 5.4 (Smooth boundary \Rightarrow smooth function). *Let (u, Ω) be a solution of (1.1). Suppose that, in a neighborhood of zero, $\partial\Omega$ is $C^{k,\alpha}$ -regular for some $k \geq 1$. Then, in a neighborhood of the origin, the functions u_+ and u_- are $C^{k,\alpha}$ up to the boundary $\partial\Omega$.*

Proof. We argue by induction. The case $k = 1$ follows by [6]. We suppose that $k \geq 2$ and that the claim holds for $k - 1$. Suppose that $\partial\Omega$ is the graph of $\eta : B' \rightarrow \mathbb{R}$, $\eta \in C^{k,\alpha}(B')$, and consider the functions

$$v_+(x', x_d) := u_+(x', x_d + \eta(x')) \quad \text{and} \quad v_-(x', x_d) := u_-(x', x_d + \eta(x')),$$

defined on the half-space $\{x_d \geq 0\}$. We set

$$A_\eta = \begin{pmatrix} \mathcal{N}_{d-1} & -(\nabla_{x'}\eta)^t \\ -\nabla_{x'}\eta & |\nabla_{x'}\eta|^2 \end{pmatrix},$$

where \mathcal{N}_{d-1} is the null $(d-1) \times (d-1)$ matrix and we notice that A_η has $C^{k-1,\alpha}$ regular coefficients. Now, since u_+ and u_- are harmonic in Ω and $D \setminus \bar{\Omega}$, we have that v_+ and v_- are solutions to the transmission problem

$$\begin{cases} -\operatorname{div}((Id + A_\eta)\nabla v_+) = 0 & \text{in } \{x_d > 0\} \\ -\operatorname{div}((Id + A_\eta)\nabla v_-) = 0 & \text{in } \{x_d < 0\} \\ v_+ = v_- & \text{on } \{x_d = 0\} \\ \partial_{x_d}v_+ - \partial_{x_d}v_- + \frac{\beta}{2\sqrt{1 + |\nabla_{x'}\eta|^2}}(v_+ + v_-) = 0 & \text{on } \{x_d = 0\}. \end{cases}$$

We now fix $k - 1$ directions i_1, \dots, i_{k-1} , $i_j \neq d$ for every j , and we consider the functions

$$w_+ := \partial_{i_1}\partial_{i_2}\dots\partial_{i_{k-1}}v_+ \quad \text{and} \quad w_- := \partial_{i_1}\partial_{i_2}\dots\partial_{i_{k-1}}v_-.$$

We notice that, in $\{x_d > 0\}$ and $\{x_d < 0\}$ the functions w_+ and w_- are solutions to

$$-\operatorname{div}((Id + A_\eta)\nabla w_\pm) + \sum_{I,J} \operatorname{div}(\partial_I A_\eta \partial_J \nabla u_\pm) = 0,$$

where the sum is over all multiindices I and J such that the sets I and J are disjoint subsets of $\{i_1, i_2, \dots, i_{k-1}\}$, $I \cup J = \{i_1, i_2, \dots, i_{k-1}\}$ and I is non-empty. In particular, using that $A_\eta \in C^{k-1,\alpha}$ and $\nabla u \in C^{k-2,\alpha}$ (since by hypothesis $u_\pm \in C^{k-1,\alpha}$), we get that w_\pm solve

$$-\operatorname{div}((Id + A_\eta)\nabla w_\pm) + \operatorname{div}(F_\pm) = 0 \quad \text{in } \{\pm x_d > 0\},$$

where F_+ and F_- are $C^{0,\alpha}$ continuous functions (depending on i_1, \dots, i_k). On the other hand, on the boundary $\{x_d = 0\}$ we have that $w_+ = w_-$ and

$$\partial_{x_d}w_+ - \partial_{x_d}w_- + \partial_{i_1}\partial_{i_2}\dots\partial_{i_{k-1}}\left(\frac{\beta(u_+ + u_-)}{2\sqrt{1 + |\nabla_{x'}\eta|^2}}\right) = 0 \quad \text{on } \{x_d = 0\}.$$

Reasoning as above, we notice that this condition can be written as

$$\partial_{x_d}w_+ - \partial_{x_d}w_- = g \quad \text{on } \{x_d = 0\},$$

where g is a $C^{0,\alpha}$ function. Now, applying [6, Theorem 1.2], we get that w_+ and w_- are $C^{1,\alpha}$ regular up to the boundary $\{x_d = 0\}$. Thus, the trace $u_+ = u_-$ is $C^{k,\alpha}$ smooth on $\{x_d = 0\}$. Finally, the classical Schauder estimates give that u_+ and u_- are $C^{k,\alpha}$ on $\{x_d \geq 0\}$ and $\{x_d \leq 0\}$, respectively. \square

Lemma 5.5 (Smooth function \Rightarrow smooth boundary). *Let (u, Ω) be a solution of (1.1). Suppose that, in a neighborhood of zero, $\partial\Omega$ is $C^{1,\alpha}$ -regular and that the functions u_+ and u_- are $C^{k,\alpha}$ up to the boundary $\partial\Omega$, for some $k \geq 1$. Then, $\partial\Omega$ is $C^{k+1,\alpha}$ -regular in a neighborhood of zero.*

Proof. Let $\xi \in C_c^\infty(D; \mathbb{R}^d)$ be a given vector field with compact support in D and let Ψ_t be the function

$$\Psi_t(x) = x + t\xi(x) \quad \text{for every } x \in D.$$

Then, for t small enough, $\Psi_t : D \rightarrow D$ is a diffeomorphism and setting $\Phi_t := \Psi_t^{-1}$, the function $u_t := u \circ \Phi_t$ is well-defined and belongs to $H^1(D)$; the function $t \mapsto \int_D |\nabla u_t|^2 dx$ is differentiable at $t = 0$ and

$$\frac{\partial}{\partial t} \Big|_{t=0} \int_D |\nabla u_t|^2 dx = \int_D (-2\nabla u D\xi \cdot \nabla u + |\nabla u|^2 \operatorname{div} \xi) dx.$$

It is immediate to check that

$$-2\nabla u D\xi \cdot \nabla u + |\nabla u|^2 \operatorname{div} \xi = \operatorname{div}(|\nabla u|^2 \xi - 2(\xi \cdot \nabla u) \nabla u) \quad \text{in } D \setminus \partial\Omega.$$

We now take ξ to be smooth outside $\partial\Omega$ and such that

$$\xi = \phi \nu_\Omega \quad \text{on } \partial\Omega,$$

where ν_Ω is the exterior normal to $\partial\Omega$ and $\phi : \partial\Omega \rightarrow \mathbb{R}$ is continuous and with compact support. Integrating by parts, we get

$$\begin{aligned} \frac{\partial}{\partial t} \Big|_{t=0} \int_D |\nabla u_t|^2 dx &= \int_{\partial\Omega} \left(|\nabla u_+|^2 (\xi \cdot \nu_\Omega) - 2(\xi \cdot \nabla u_+) (\nu_\Omega \cdot \nabla u_+) \right) d\mathcal{H}^{d-1} \\ &\quad - \int_{\partial\Omega} \left(|\nabla u_-|^2 (\xi \cdot \nu_\Omega) - 2(\xi \cdot \nabla u_-) (\nu_\Omega \cdot \nabla u_-) \right) d\mathcal{H}^{d-1}, \end{aligned}$$

where $u_+ := u$ on Ω , and $u_- := u$ on $D \setminus \bar{\Omega}$. Now, if

$$\xi = \phi e_d \quad \text{and} \quad \nu_\Omega = \frac{(-\nabla_{x'} \eta, 1)}{\sqrt{1 + |\nabla_{x'} \eta|^2}},$$

then

$$\begin{aligned} &\sqrt{1 + |\nabla_{x'} \eta|^2} \left(|\nabla u_+|^2 (\xi \cdot \nu_\Omega) - 2(\xi \cdot \nabla u_+) (\nu_\Omega \cdot \nabla u_+) \right) \\ &\quad - \sqrt{1 + |\nabla_{x'} \eta|^2} \left(|\nabla u_-|^2 (\xi \cdot \nu_\Omega) - 2(\xi \cdot \nabla u_-) (\nu_\Omega \cdot \nabla u_-) \right) \\ &= \phi \left(|\nabla u_+|^2 - |\nabla u_-|^2 \right) \\ &\quad - 2\phi \left(\partial_{x_d} u_+ (-\nabla_{x'} \eta \cdot \nabla_{x'} u_+ + \partial_{x_d} u_+) - \partial_{x_d} u_- (-\nabla_{x'} \eta \cdot \nabla_{x'} u_- + \partial_{x_d} u_-) \right). \end{aligned}$$

We now suppose that $x_0 \in \operatorname{Reg}(\partial\Omega)$ and that $\partial\Omega$ is the graph of the $(C^{1,\alpha})$ function $\eta : B' \rightarrow \mathbb{R}$, where B' is a ball in \mathbb{R}^{d-1} . Taking $\xi = e_d \phi$ and $\Omega_t = \Phi_t(\Omega)$, we have

$$\begin{aligned} \frac{\partial}{\partial t} \Big|_{t=0} \int_{\partial\Omega_t} u_t^2 d\mathcal{H}^{d-1} &= \frac{\partial}{\partial t} \Big|_{t=0} \int_{B'} u^2(x', \eta(x')) \sqrt{1 + |\nabla_{x'} \eta + t \nabla_{x'} \phi|^2} dx' \\ &= \int_{B'} \frac{u^2(x', \eta(x'))}{\sqrt{1 + |\nabla_{x'} \eta|^2}} \nabla_{x'} \eta \cdot \nabla_{x'} \phi dx' \\ &= \int_{B'} u^2(x', \eta(x')) H(x', \eta(x')) \phi(x') dx' \\ &\quad - 2 \int_{B'} \phi(x') u(x', \eta(x')) \frac{(\nabla_{x'} u + \partial_{x_d} u \nabla_{x'} \eta) \cdot \nabla_{x'} \eta}{\sqrt{1 + |\nabla_{x'} \eta|^2}} dx'. \end{aligned}$$

In particular, combining these two computations and using the optimality of (u, Ω) , we get

$$\begin{aligned} 0 &= \frac{\partial}{\partial t} \Big|_{t=0} J_\beta(u_t, \Omega_t) = \int_{B'} \beta u^2 H(x') \phi(x') dx' \\ &\quad + \int_{B'} \left(|\nabla u_+|^2 - |\nabla u_-|^2 \right) \phi(x') dx' \\ &\quad - \int_{B'} 2 \left(1 + |\nabla_{x'} \eta|^2 \right) \left((\partial_{x_d} u_+)^2 - (\partial_{x_d} u_-)^2 \right) \phi(x') dx' \end{aligned}$$

Since ϕ is arbitrary, we obtain that η is a solution of the problem

$$-\operatorname{div}_{x'} \left(\frac{\nabla_{x'} \eta}{\sqrt{1 + |\nabla_{x'} \eta|^2}} \right) = f(x') \quad \text{in } B',$$

where

$$f(x') = \frac{1}{\beta u^2(x', \eta(x'))} \left[(|\nabla u_+|^2 - |\nabla u_-|^2) - 2(1 + |\nabla_{x'} \eta|^2) \left((\partial_{x_d} u_+)^2 - (\partial_{x_d} u_-)^2 \right) \right],$$

and all the derivatives of u_+ and u_- are calculated at $(x', \eta(x'))$. Since the right-hand side f is $C^{k-1, \alpha}$ regular, we get that η is $C^{k+1, \alpha}$ regular. This concludes the proof. \square

APPENDIX A. EXAMPLES OF MINIMIZERS

In this section, we use a calibration argument to prove that if $E = \{x_d > 0\}$ and $v \equiv 1$, then in any Steiner symmetric set $D \subset \mathbb{R}^d$, the solution (Ω, u) is unique, u is even with respect to the hyperplane $\{x_d = 0\}$ and Ω is precisely the half-space E . Our main result is the following.

Proposition A.1. *Let D be an open set, Steiner symmetric with respect to the hyperplane $\{x_d = 0\}$. Let E be the half-ball $E = B \cap \{x_d > 0\}$, for some large ball B containing D , and let $v \equiv 1$. Then there is a unique solution (u, Ω) to (1.1), where $\Omega = E$, u is positive and even with respect to $\{x_d = 0\}$ and solves the equation*

$$\Delta u = 0 \quad \text{in } \{x_d > 0\} \cap D, \quad \partial_{x_d} u = \frac{1}{2} \beta u \quad \text{on } D \cap \{x_d = 0\}. \quad (\text{A.1})$$

Proof. Let $\tilde{u} \in \mathcal{V}$ and $\tilde{\Omega} \in \mathcal{E}$ be given. We will prove that

$$J_\beta(u, \Omega) \leq J_\beta(\tilde{u}, \tilde{\Omega}),$$

with an equality, if and only if, $(u, \Omega) = (\tilde{u}, \tilde{\Omega})$. First, we notice that, since $J_\beta(1 \wedge \tilde{u} \vee 0, \tilde{\Omega}) \leq J_\beta(\tilde{u}, \tilde{\Omega})$, we can suppose that $0 \leq \tilde{u} \leq 1$. We then write \tilde{u} as $\tilde{u} = 1 - \varphi$ for some $\varphi \in H_0^1(D)$ such that $0 \leq \varphi \leq 1$ and we define the function $\tilde{u}_* = 1 - \varphi_*$, where $\varphi_* \in H_0^1(D)$ is the Steiner symmetrization of φ . We will show that

$$J_\beta(\tilde{u}_*, \Omega) \leq J_\beta(\tilde{u}, \tilde{\Omega}) \quad (\text{A.2})$$

Indeed, it is well-known that the Steiner symmetrization decreases the Dirichlet energy:

$$\int_D |\nabla \tilde{u}_*|^2 dx = \int_D |\nabla \varphi_*|^2 dx \leq \int_D |\nabla \varphi|^2 dx = \int_D |\nabla \tilde{u}|^2 dx.$$

In order to estimate also the second term of the energy J_β , we use a calibration-type argument. We first notice that, by construction, along every line orthogonal to $\{x_d = 0\}$, the symmetrized function achieves its maximum in zero. Precisely

$$\varphi(x', x_d) \leq \sup_{x_d} \varphi(x', x_d) = \varphi_*(x', 0).$$

Thus, by the definition of \tilde{u}_* , we have

$$\begin{aligned} \int_{B \cap \partial \tilde{\Omega}} \tilde{u}^2(x', x_d) d\mathcal{H}^{d-1} &\geq \int_{B \cap \partial \tilde{\Omega}} \tilde{u}_*^2(x', 0) d\mathcal{H}^{d-1} \geq \int_{B \cap \partial \tilde{\Omega}} \tilde{u}_*^2(x', 0) \nu_{\tilde{\Omega}} \cdot e_d d\mathcal{H}^{d-1} \\ &= \int_{B \cap \partial \Omega} \tilde{u}_*^2(x', 0) \nu_{\Omega} \cdot (-e_d) d\mathcal{H}^{d-1} + \int_{\Omega \Delta \tilde{\Omega}} \operatorname{div}(\tilde{u}_*^2(x', 0) e_d) dx, \end{aligned}$$

where in order to get the last equality we used the divergence theorem in $\Omega \Delta \tilde{\Omega}$. Now, we notice that $\operatorname{div}(\tilde{u}_*^2(x', 0) e_d) = 0$ and that $\nu_{\Omega} = -e_d$. Thus, we get

$$\int_{B \cap \partial \tilde{\Omega}} \tilde{u}^2(x', x_d) d\mathcal{H}^{d-1} \geq \int_{B \cap \partial \Omega} \tilde{u}_*^2 d\mathcal{H}^{d-1},$$

which concludes the proof of (A.2). Finally, we notice that the problem

$$\min \left\{ J_\beta(u, \Omega) : u \in H^1(D \cap \{x_d > 0\}), \quad u = 1 \text{ on } \partial D \cap \{x_d > 0\} \right\},$$

has a unique solution u , which is Steiner symmetric, non-negative and satisfies (A.1). \square

ACKNOWLEDGMENTS

The first author was partially supported by PRIN 2017 *Nonlinear Differential Problems via Variational, Topological and Set-valued Methods* (Grant 2017AYM8XW) and the INdAM-GNAMPA project 2020 “Problemi di ottimizzazione con vincoli via trasporto ottimo e incertezza”. The second author was partially supported by the Academy of Finland grant 314227. The third author has been partially supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement VAREG, No. 853404).

REFERENCES

- [1] D. Bucur, A. Giacomini. *Shape optimization problems with Robin conditions on the free boundary..* Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), no. 6, 1539–1568.
- [2] D. Bucur, S. Luckhaus. *Monotonicity formula and regularity for general free discontinuity problems.* Arch. Rat. Mech. Anal. **211** (2014), 489–511.
- [3] L. A. Caffarelli, D. Kriventsov. *A free boundary problem related to thermal insulation.* Comm. PDE **41** (7) (2016), 1149–1182.
- [4] L. A. Caffarelli, M. Soria-Carro, P. R. Stinga. *Regularity for $C^{1,\alpha}$ interface transmission problems.* ArXiv:2004.07322 (2020).
- [5] Ennio De Giorgi. *Frontiere orientate di misura minima.* Seminario di Matematica della Scuola Normale Superiore di Pisa, 1960-61. Editrice Tecnica Scientifica, Pisa (1961).
- [6] H. Dong. *A simple proof of regularity for $C^{1,\alpha}$ interface transmission problems.* <https://arxiv.org/abs/2004.09365> (2020).
- [7] L. C. Evans, R. F. Gariepy. *Measure theory and fine properties of functions.* Boca Raton, FL: CRC Press, 2nd revised edition (2015).
- [8] E. Giusti. *Minimal surfaces and functions of bounded variations.* Birkhäuser, Boston (1984).
- [9] D. Kriventsov. *A free boundary problem related to thermal insulation: flat implies smooth.* Calc. Var. PDE **58** (2019).
- [10] F. Maggi. *Sets of finite perimeter and geometric variational problems: an introduction to Geometric Measure Theory.* Cambridge University Press **135** (2012).
- [11] I. TAMANINI: *Regularity results for almost minimal hyperurfaces in \mathbb{R}^n .* Quaderni del Dipartimento di Matematica dell’ Università di Lecce. (1984).
- [12] B. Velichkov. *Regularity of the one-phase free boundaries.* Lecture notes 2019. Available at <http://cvgmt.sns.it/paper/4367/>

SERENA GUARINO LO BIANCO, UNIVERSITÀ DEGLI STUDI DI NAPOLI “FEDERICO II”, DIPARTIMENTO DI AGRARIA, VIA UNIVERSITÀ 100, 80055 PORTICI (NA), ITALIA.

Email address: `serena.guarinolobianco@unina.it`

DOMENICO ANGELO LA MANNA, UNIVERSITY OF JYVASKYLÄ, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35 (MAD) FI-40014, FINLAND

Email address: `domenicolamanna@hotmail.it`

BOZHIDAR VELICHKOV:

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA
LARGO BRUNO PONTECORVO, 5, 56127 PISA - ITALY

Email address: `bozhidar.velichkov@unipi.it`