

Quasistatic limit of a dynamic viscoelastic model with memory

Gianni Dal Maso and Francesco Sapio

Abstract. We study the behaviour of the solutions to a dynamic evolution problem for a viscoelastic model with long memory, when the rate of change of the data tends to zero. We prove that a suitably rescaled version of the solutions converges to the solution of the corresponding stationary problem.

Keywords: evolution problems with memory, linear second order hyperbolic systems, dynamic mechanics, elastodynamics, viscoelasticity.

MSC 2010: 35B25, 35B40, 35L53, 35Q74, 74H20, 74D05.

1. Introduction

The most common models of viscoelasticity with long memory, such as the Maxwell model (see [4], [5], [6], [15]), lead to a dynamic evolution governed by a system of partial differential equations of the form

$$\ddot{u}(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})eu(t)) + \int_{-\infty}^t \frac{1}{\beta} e^{-\frac{t-\tau}{\beta}} \operatorname{div}(\mathbb{B}eu(\tau)) d\tau = \ell(t) \quad (1.1)$$

in Ω for $t \in [0, T]$, where $\Omega \subset \mathbb{R}^d$ is the reference configuration, $[0, T]$ is the time interval, $u(t)$ and $eu(t)$ are the displacement at time t and the symmetric part of its gradient, \mathbb{A} and \mathbb{B} are the elasticity and viscosity tensors, $\beta > 0$ is a material constant, and $\ell(t)$ is the external load at time t . This system is complemented by boundary and initial conditions

$$u(t) = z(t) \quad \text{on } \partial\Omega \text{ for } t \in [0, T], \quad (1.2)$$

$$u(t) = u_{in}(t) \quad \text{in } \Omega \text{ for } t \in (-\infty, 0], \quad (1.3)$$

where z and u_{in} are prescribed functions, the latter representing the history of the displacement for $t \leq 0$. Existence and uniqueness for (1.1)–(1.3) can be found in [3].

In this paper we study the quasistatic limit of the solutions to this problem, i.e., the limit of these solutions when the rate of change of the data

tends to zero. More precisely, given a small parameter $\varepsilon > 0$, we consider the solution u^ε of (1.1)–(1.3) corresponding to $\ell(\varepsilon t)$, $z(\varepsilon t)$, and $u_{in}(\varepsilon t)$. To study the asymptotic behaviour of u^ε as $\varepsilon \rightarrow 0^+$ it is convenient to introduce the rescaled solution $u_\varepsilon(t) := u^\varepsilon(t/\varepsilon)$, which turns out to be the solution of the system

$$\varepsilon^2 \ddot{u}_\varepsilon(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})e u_\varepsilon(t)) + \int_{-\infty}^t \frac{1}{\beta \varepsilon} e^{-\frac{t-\tau}{\beta \varepsilon}} \operatorname{div}(\mathbb{B} e u_\varepsilon(\tau)) d\tau = \ell(t) \quad (1.4)$$

in Ω for $t \in [0, T]$, with boundary and initial conditions (1.2) and (1.3).

Under different assumptions on $\ell(t)$, $z(t)$, and $u_{in}(t)$ we prove (Theorems 3.6 and 3.7) that $u_\varepsilon(t)$ converges, as $\varepsilon \rightarrow 0^+$, to the solution $u_0(t)$ of the stationary problem

$$-\operatorname{div}(\mathbb{A} e u_0(t)) = \ell(t) \quad \text{in } \Omega \text{ for } t \in [0, T], \quad (1.5)$$

with boundary condition (1.2).

By using just the energy-dissipation inequality, it is not difficult to prove a similar result for the Kelvin-Voigt model, in which the viscosity term

$$-\operatorname{div}(\mathbb{B} e \dot{u}(t)) + \int_{-\infty}^t \frac{1}{\beta} e^{-\frac{t-\tau}{\beta}} \operatorname{div}(\mathbb{B} e \dot{u}(\tau)) d\tau \quad (1.6)$$

is replaced by $-\operatorname{div}(\mathbb{B} e \dot{u}(t))$. On the other hand, in the case of the equation of elastodynamics without damping terms, i.e., when $\mathbb{B} = 0$, by using the Fourier decomposition with respect to the eigenfunctions of the operator $-\operatorname{div}(\mathbb{A} e u)$, we can easily see that the convergence of u_ε to u_0 does not hold in general. The purpose of this paper is to prove that the non-local damping term (1.6) is enough to obtain the convergence of the solutions of the evolution problems to the solution of the stationary problem.

Our result can be considered in the framework of the study of the quasistatic limits, i.e. the convergence of the solutions to second order evolution equations with rescaled times towards the solutions to the corresponding stationary equations. Similar problems in finite dimension have been studied in [1, 10, 13, 7]. A special case involving the wave equations on time-dependent intervals in dimension one has been studied in [9, 12]. The main novelty of our problem is the non-local form of the damping term, given by (1.6).

The main tools to prove our results are two different estimates (Lemmas 3.8 and 5.2), related to the energy-dissipation balance (2.25) and to the elliptic system (5.17) obtained from (1.4) via Laplace Transform. After a precise statement of all assumptions, more details on the line of proof will be given after Theorem 3.7.

2. Hypotheses and statement of the problem

Let d be a positive integer and let $\Omega \subset \mathbb{R}^d$ be a bounded open set with Lipschitz boundary. We use standard notation for Lebesgue and Sobolev spaces.

Let $\mathbb{R}_{sym}^{d \times d}$ be the space of all symmetric $d \times d$ matrices. For convenience we set

$$\begin{aligned} H &:= L^2(\Omega; \mathbb{R}^d), \quad \tilde{H} := L^2(\Omega; \mathbb{R}_{sym}^{d \times d}), \\ V &:= H^1(\Omega; \mathbb{R}^d), \quad V_0 := H_0^1(\Omega; \mathbb{R}^d), \quad V'_0 := H^{-1}(\Omega; \mathbb{R}^d), \end{aligned} \quad (2.1)$$

and we always identify the dual of H with H itself. The symbols (\cdot, \cdot) and $\|\cdot\|$ denote the scalar product and the norm in H or in \tilde{H} , according to the context. The duality product between V'_0 and V_0 is denoted by $\langle \cdot, \cdot \rangle$. Given $u \in V$, its strain eu is defined as the symmetric part of its gradient, i.e., $eu := \frac{1}{2}(\nabla u + \nabla u^T)$, where ∇u is the Jacobian matrix, whose components are $(\nabla u)_{ij} := \partial_j u_i$ for $i, j = 1, \dots, d$.

Under these assumptions, the Second Korn Inequality (see, e.g., [11, Theorem 2.4]) states that there exists a positive constant $C_K = C_K(\Omega)$ such that

$$\|\nabla u\| \leq C_K (\|u\|^2 + \|eu\|^2)^{1/2} \quad \text{for every } u \in V. \quad (2.2)$$

Moreover, there exists a positive constant $C_P = C_P(\Omega)$ such that the following Korn-Poincaré Inequality holds (see, e.g., [11, Theorem 2.7]):

$$\|u\| \leq C_P \|eu\| \quad \text{for every } u \in V_0. \quad (2.3)$$

Thanks to (2.2) we can use on the space V the equivalent norm

$$\|u\|_V := (\|u\|^2 + \|eu\|^2)^{1/2} \quad \text{for every } u \in V.$$

Let $\mathcal{L}(\mathbb{R}_{sym}^{d \times d}; \mathbb{R}_{sym}^{d \times d})$ be the space of all linear operators from $\mathbb{R}_{sym}^{d \times d}$ into itself. We assume that the elasticity and viscosity tensors \mathbb{A} and \mathbb{B} , which depend on the variable $x \in \Omega$, satisfy the following assumptions:

$$\mathbb{A}, \mathbb{B} \in L^\infty(\Omega; \mathcal{L}(\mathbb{R}_{sym}^{d \times d}; \mathbb{R}_{sym}^{d \times d})), \quad (2.4)$$

and for a.e. $x \in \Omega$

$$\begin{aligned} \mathbb{A}(x)\xi_1 \cdot \xi_2 &= \xi_1 \cdot \mathbb{A}(x)\xi_2, \\ \mathbb{B}(x)\xi_1 \cdot \xi_2 &= \xi_1 \cdot \mathbb{B}(x)\xi_2 \end{aligned} \quad \text{for every } \xi_1, \xi_2 \in \mathbb{R}_{sym}^{d \times d}, \quad (2.5)$$

$$\begin{aligned} c_{\mathbb{A}}|\xi|^2 &\leq \mathbb{A}(x)\xi \cdot \xi \leq C_{\mathbb{A}}|\xi|^2, \\ c_{\mathbb{B}}|\xi|^2 &\leq \mathbb{B}(x)\xi \cdot \xi \leq C_{\mathbb{B}}|\xi|^2 \end{aligned} \quad \text{for every } \xi \in \mathbb{R}_{sym}^{d \times d}, \quad (2.6)$$

where $c_{\mathbb{A}}$, $c_{\mathbb{B}}$, $C_{\mathbb{A}}$, and $C_{\mathbb{B}}$ are positive constants independent of x , and the dot denotes the Euclidean scalar product of matrices.

Let us fix $T > 0$ and $\beta > 0$. To give a precise meaning to the notion of solution to problem (1.2)–(1.4) we introduce the function spaces

$$\begin{aligned} \mathcal{V} &:= L^2(0, T; V) \cap H^1(0, T; H) \cap H^2(0, T; V'_0), \\ \mathcal{V}_0 &:= L^2(0, T; V_0) \cap H^1(0, T; H) \cap H^2(0, T; V'_0), \\ \mathcal{V}_{loc} &:= L^2_{loc}(-\infty, T; V) \cap H^1_{loc}(-\infty, T; H) \cap H^2_{loc}(-\infty, T; V'_0). \end{aligned}$$

Remark 2.1. By the Sobolev Embedding Theorem, if $u \in \mathcal{V}$ (resp. $u \in \mathcal{V}_{loc}$), then

$$\begin{aligned} u &\in C^0([0, T]; H) \cap C^1([0, T]; V'_0), \\ (\text{resp. } u &\in C^0((-\infty, T); H) \cap C^1((-\infty, T); V'_0)). \end{aligned}$$

We study problem (1.2)–(1.4) with ℓ , z , and u_{in} depending on ε . Let us consider $\varepsilon > 0$ and

$$f_\varepsilon \in L^2(0, T; H), \quad g_\varepsilon \in H^1(0, T; V'_0), \quad z_\varepsilon \in H^2(0, T; H) \cap H^1(0, T; V), \quad (2.7)$$

$$u_{\varepsilon, in} \in C^0((-\infty, T); H) \cap C^1((-\infty, T); V'_0) \text{ such that}$$

$$u_{\varepsilon, in}(0) \in V, \quad u_{\varepsilon, in}(0) - z_\varepsilon(0) \in V_0, \\ \dot{u}_{\varepsilon, in}(0) \in H, \quad \int_{-\infty}^0 \frac{1}{\beta\varepsilon} e^{\frac{\tau}{\beta\varepsilon}} \|u_{\varepsilon, in}(\tau)\|_V d\tau < +\infty. \quad (2.8)$$

The notion of solution to (1.2)–(1.4) is made precise by the following definition.

Definition 2.2. We say that u_ε is a solution to the viscoelastic dynamic system (1.2)–(1.4), with forcing term $\ell = f_\varepsilon + g_\varepsilon$, boundary condition z_ε , and initial condition $u_{\varepsilon, in}$, if

$$u_\varepsilon \in \mathcal{V}_{loc} \quad \text{and} \quad u_\varepsilon - z_\varepsilon \in \mathcal{V}_0, \quad (2.9a)$$

$$\varepsilon^2 \ddot{u}_\varepsilon(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})e u_\varepsilon(t)) + \int_{-\infty}^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e u_\varepsilon(\tau)) d\tau \\ = f_\varepsilon(t) + g_\varepsilon(t) \quad \text{for a.e. } t \in [0, T], \quad (2.9b)$$

$$u_\varepsilon(t) = u_{\varepsilon, in}(t) \quad \text{for every } t \in (-\infty, 0]. \quad (2.9c)$$

In the next remark we shall see that (2.9) can be reduced to the following problem starting from 0:

$$u_\varepsilon \in \mathcal{V} \quad \text{and} \quad u_\varepsilon - z_\varepsilon \in \mathcal{V}_0, \quad (2.10a)$$

$$\varepsilon^2 \ddot{u}_\varepsilon(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})e u_\varepsilon(t)) + \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e u_\varepsilon(\tau)) d\tau \\ = \varphi_\varepsilon(t) + \gamma_\varepsilon(t) \quad \text{for a.e. } t \in [0, T], \quad (2.10b)$$

$$u_\varepsilon(0) = u_\varepsilon^0 \text{ in } H \quad \text{and} \quad \dot{u}_\varepsilon(0) = u_\varepsilon^1 \text{ in } V'_0, \quad (2.10c)$$

with $\varphi_\varepsilon \in L^2(0, T; H)$, $\gamma_\varepsilon \in H^1(0, T; V'_0)$, $u_\varepsilon^0 \in V$, $u_\varepsilon^0 - z_\varepsilon(0) \in V_0$, $u_\varepsilon^1 \in H$.

Remark 2.3. It is easy to see that u_ε is a solution according to Definition 2.2 if and only if its restriction to $[0, T]$, still denoted by u_ε , solves (2.10) with

$$\varphi_\varepsilon = f_\varepsilon, \quad \gamma_\varepsilon = g_\varepsilon - p_\varepsilon, \quad u_\varepsilon^0 = u_{\varepsilon, in}(0), \quad u_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0), \quad (2.11)$$

where

$$p_\varepsilon(t) := e^{-\frac{t}{\beta\varepsilon}} g_\varepsilon^0 \quad \text{with} \quad g_\varepsilon^0 := \int_{-\infty}^0 \frac{1}{\beta\varepsilon} e^{\frac{\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e u_{\varepsilon, in}(\tau)) d\tau. \quad (2.12)$$

To solve problem (2.10) it is enough to study the corresponding problem with homogeneous boundary condition:

$$v_\varepsilon \in \mathcal{V}_0, \quad (2.13a)$$

$$\varepsilon^2 \ddot{v}_\varepsilon(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})e v_\varepsilon(t)) + \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e v_\varepsilon(\tau)) d\tau$$

$$= h_\varepsilon(t) + \ell_\varepsilon(t) \quad \text{for a.e. } t \in [0, T], \quad (2.13b)$$

$$v_\varepsilon(0) = v_\varepsilon^0 \text{ in } H \quad \text{and} \quad \dot{v}_\varepsilon(0) = v_\varepsilon^1 \text{ in } V_0', \quad (2.13c)$$

with

$$h_\varepsilon \in L^2(0, T; H), \quad \ell_\varepsilon \in H^1(0, T; V_0'), \quad v_\varepsilon^0 \in V_0, \quad v_\varepsilon^1 \in H. \quad (2.14)$$

Remark 2.4. The function u_ε is a solution to (2.10) if and only if $v_\varepsilon = u_\varepsilon - z_\varepsilon$ solves (2.13) with

$$\begin{aligned} h_\varepsilon(t) &= \varphi_\varepsilon(t) - \varepsilon^2 \ddot{z}_\varepsilon(t), \\ \ell_\varepsilon(t) &= \gamma_\varepsilon(t) + \operatorname{div}((\mathbb{A} + \mathbb{B})e z_\varepsilon(t)) - \int_0^t \frac{1}{\beta \varepsilon} e^{-\frac{t-\tau}{\beta \varepsilon}} \operatorname{div}(\mathbb{B} e z_\varepsilon(\tau)) d\tau, \\ v_\varepsilon^0 &= u_\varepsilon^0 - z_\varepsilon(0), \quad v_\varepsilon^1 = u_\varepsilon^1 - \dot{z}_\varepsilon(0), \end{aligned} \quad (2.15)$$

Therefore, existence and uniqueness for (2.13) imply existence and uniqueness for (2.10).

Remark 2.5. In [3] problem (2.13) has been studied with initial conditions taken in the sense of interpolation spaces. Given two Hilbert spaces X and Y , the symbol $[X, Y]_\theta$ denotes the interpolation space between X and Y of exponent $\theta \in (0, 1)$. Thanks to [8, Theorem 3.1] we have the following inclusions:

$$\begin{aligned} L^2(0, T; V_0) \cap H^1(0, T; H) &\subset C^0([0, T]; V_0^{\frac{1}{2}}) \\ L^2(0, T; H) \cap H^1(0, T; V_0') &\subset C^0([0, T]; V_0^{-\frac{1}{2}}), \end{aligned}$$

where $V_0^{\frac{1}{2}} := [V_0, H]_{\frac{1}{2}}$ and $V_0^{-\frac{1}{2}} := [H, V_0']_{\frac{1}{2}}$. Consequently

$$\mathcal{V}_0 \subset C^0([0, T]; V_0^{\frac{1}{2}}) \cap C^1([0, T]; V_0^{-\frac{1}{2}}).$$

Therefore, the initial conditions in (2.13) are satisfied also in the stronger sense

$$v_\varepsilon(0) = v_\varepsilon^0 \text{ in } V_0^{\frac{1}{2}} \quad \text{and} \quad \dot{v}_\varepsilon(0) = v_\varepsilon^1 \text{ in } V_0^{-\frac{1}{2}}. \quad (2.16)$$

The following proposition provides the main properties of the solutions. We recall that, if X is a Banach space, $C_w^0([0, T]; X)$ denotes the space of all weakly continuous functions from $[0, T]$ to X , namely, the vector space of all functions $u: [0, T] \rightarrow X$ such that for every $x' \in X'$ the function $t \mapsto \langle x', u(t) \rangle$ is continuous from $[0, T]$ to \mathbb{R} .

Proposition 2.6. *Given $\varepsilon > 0$, assume (2.7) and (2.8). Then there exists a unique solution u_ε to the viscoelastic dynamic system (2.9). Moreover, it satisfies*

$$u_\varepsilon \in C^0([0, T]; V) \cap C^1([0, T]; H). \quad (2.17)$$

Proof. By Remarks 2.3 and 2.4 it is enough to prove the theorem for (2.13). Existence and uniqueness are proved in [3], taking into account Remark 2.5 about the equivalence between the initial conditions in the sense of (2.13) and (2.16).

After an integration by parts with respect to time, it easy to see that the weak formulation (2.13) is equivalent to the following one:

$$\begin{aligned} & -\varepsilon^2 \int_0^T (\dot{v}_\varepsilon(t), \dot{\varphi}(t)) dt + \int_0^T ((\mathbb{A} + \mathbb{B})ev_\varepsilon(t) - \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \mathbb{B}ev_\varepsilon(\tau), e\varphi(t)) dt \\ & = \int_0^T [(h_\varepsilon(t), \varphi(t)) + \langle \ell_\varepsilon(t), \varphi(t) \rangle] dt \quad \text{for every } \varphi \in C_c^\infty(0, T; V). \end{aligned} \quad (2.18)$$

In [14], in a more general context, it has been proved that if v_ε satisfies (2.18) and the initial conditions in the sense of (2.13), then it satisfies also

$$\begin{aligned} v_\varepsilon & \in C_w^0([0, T]; V) \quad \text{and} \quad \dot{v}_\varepsilon \in C_w^0([0, T]; H), \\ \lim_{t \rightarrow 0^+} \|v_\varepsilon(t) - v_\varepsilon^0\|_V & = 0 \quad \text{and} \quad \lim_{t \rightarrow 0^+} \|\dot{v}_\varepsilon(t) - v_\varepsilon^1\| = 0. \end{aligned} \quad (2.19)$$

We fix $s \in [0, T]$. We want to prove

$$\lim_{t \rightarrow s^+} \|v_\varepsilon(t) - v_\varepsilon(s)\|_V = 0 \quad \text{and} \quad \lim_{t \rightarrow s^+} \|\dot{v}_\varepsilon(t) - \dot{v}_\varepsilon(s)\| = 0. \quad (2.20)$$

Thanks to the theory developed in [3] there exists a unique $\tilde{v}_\varepsilon \in L^2(s, T; V_0) \cap H^1(s, T; H) \cap H^2(s, T; V_0')$ such that

$$\begin{aligned} & \varepsilon^2 \ddot{\tilde{v}}_\varepsilon(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})e\tilde{v}_\varepsilon(t)) + \int_s^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e\tilde{v}_\varepsilon(\tau)) d\tau \\ & = h_\varepsilon(t) + \ell_\varepsilon(t) - \int_0^s \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_\varepsilon(\tau)) d\tau \quad \text{for a.e. } t \in [s, T], \end{aligned} \quad (2.21)$$

$$\lim_{t \rightarrow s^+} \|\tilde{v}_\varepsilon(t) - v_\varepsilon(s)\| = 0 \quad \text{and} \quad \lim_{t \rightarrow s^+} \|\dot{\tilde{v}}_\varepsilon(t) - \dot{v}_\varepsilon(s)\|_{V_0'} = 0. \quad (2.22)$$

By the results in [14] the function \tilde{v}_ε satisfies also

$$\lim_{t \rightarrow s^+} \|\tilde{v}_\varepsilon(t) - v_\varepsilon(s)\|_V = 0 \quad \text{and} \quad \lim_{t \rightarrow s^+} \|\dot{\tilde{v}}_\varepsilon(t) - \dot{v}_\varepsilon(s)\| = 0. \quad (2.23)$$

Since clearly v_ε satisfies (2.21) and (2.22), by uniqueness we have $\tilde{v}_\varepsilon(t) = v_\varepsilon(t)$ for every $t \in [s, T]$. In particular, from (2.23) we deduce that (2.20) holds.

To complete the proof we need the following proposition about the energy-dissipation balance, where \tilde{H} is defined by (2.1), and $\mathcal{W}_\varepsilon(t)$ represents the work done in the interval $[0, t]$.

Proposition 2.7. *Given $\varepsilon > 0$, we assume (2.14). Let v_ε be the solution to (2.13) and let $w_\varepsilon: [0, T] \rightarrow \tilde{H}$ be defined by*

$$w_\varepsilon(t) := e^{-\frac{t}{\beta\varepsilon}} \int_0^t \frac{1}{\beta\varepsilon} e^{\frac{\tau}{\beta\varepsilon}} ev_\varepsilon(\tau) d\tau \quad \text{for every } t \in [0, T]. \quad (2.24)$$

Then $w_\varepsilon \in H^1(0, T; \tilde{H})$ and the following energy-dissipation balance holds for every $t \in [0, T]$:

$$\frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon(t)\|^2 + \frac{1}{2} (\mathbb{A}ev_\varepsilon(t), ev_\varepsilon(t))$$

$$\begin{aligned}
 & + \frac{1}{2}(\mathbb{B}(ev_\varepsilon(t) - w_\varepsilon(t)), ev_\varepsilon(t) - w_\varepsilon(t)) + \beta\varepsilon \int_0^t (\mathbb{B}\dot{w}_\varepsilon(\tau), \dot{w}_\varepsilon(\tau))d\tau \\
 & = \frac{\varepsilon^2}{2}\|v_\varepsilon^1\|^2 + \frac{1}{2}((\mathbb{A} + \mathbb{B})ev_\varepsilon^0, ev_\varepsilon^0) + \mathcal{W}_\varepsilon(t),
 \end{aligned} \tag{2.25}$$

where

$$\begin{aligned}
 \mathcal{W}_\varepsilon(t) & := \langle \ell_\varepsilon(t), v_\varepsilon(t) \rangle - \langle \ell_\varepsilon(0), v_\varepsilon^0 \rangle \\
 & \quad + \int_0^t (h_\varepsilon(\tau), \dot{v}_\varepsilon(\tau))d\tau - \int_0^t \langle \dot{\ell}_\varepsilon(\tau), v_\varepsilon(\tau) \rangle d\tau.
 \end{aligned}$$

Proof. It is convenient to extend the data of our problem to the interval $[0, 2T]$ by setting

$$h_\varepsilon(t) := 0 \quad \text{and} \quad \ell_\varepsilon(t) := \ell_\varepsilon(T) \quad \text{for every } t \in (T, 2T].$$

It is clear that $h_\varepsilon \in L^2(0, 2T; H)$ and $\ell_\varepsilon \in H^1(0, 2T; V_0')$. By uniqueness of the solution to (2.13), the solution on $[0, 2T]$ is an extension of v_ε , still denoted by v_ε . We also consider the extension of w_ε on $[0, 2T]$ defined by (2.24).

Since $ev_\varepsilon \in L^2(0, 2T; \tilde{H})$, it follows from (2.24) that $w_\varepsilon \in H^1(0, 2T; \tilde{H})$, and

$$\beta\varepsilon\dot{w}_\varepsilon(t) = ev_\varepsilon(t) - w_\varepsilon(t) \quad \text{for a.e. } t \in [0, 2T]. \tag{2.26}$$

Thanks to (2.20) in $[0, 2T]$ and (2.26) there exists a representative of \dot{w}_ε such that

$$\lim_{t \rightarrow s^+} \|\dot{w}_\varepsilon(t) - \dot{w}_\varepsilon(s)\| = 0 \quad \text{for every } s \in [0, 2T]. \tag{2.27}$$

Moreover, since v_ε satisfies (2.13) in $[0, 2T]$, we have for a.e. $t \in [0, 2T]$

$$\varepsilon^2\ddot{v}_\varepsilon(t) - \operatorname{div}(\mathbb{A}ev_\varepsilon(t)) - \operatorname{div}(\mathbb{B}(ev_\varepsilon(t) - w_\varepsilon(t))) = h_\varepsilon(t) + \ell_\varepsilon(t). \tag{2.28}$$

Multiplying (2.26) and (2.28) by $\psi \in \tilde{H}$ and $\varphi \in V_0$, respectively, and then integrating over Ω and adding the results, for a.e. $t \in [0, 2T]$ we get

$$\begin{aligned}
 \varepsilon^2\langle \ddot{v}_\varepsilon(t), \varphi \rangle + (\mathbb{A}ev_\varepsilon(t), e\varphi) + (\mathbb{B}(ev_\varepsilon(t) - w_\varepsilon(t)), e\varphi - \psi) \\
 + \beta\varepsilon(\mathbb{B}\dot{w}_\varepsilon(t), \psi) = (h_\varepsilon(t), \varphi) + \langle \ell_\varepsilon(t), \varphi \rangle.
 \end{aligned} \tag{2.29}$$

Given a function r from $[0, 2T]$ into a Banach space X , for every $\eta > 0$ we define the sum and the difference $\sigma^\eta r, \delta^\eta r: [0, 2T - \eta] \rightarrow X$ by

$$\sigma^\eta r(t) := r(t + \eta) + r(t) \quad \text{and} \quad \delta^\eta r(t) := r(t + \eta) - r(t).$$

For a.e. $t \in [0, 2T - \eta]$ we have $\sigma^\eta v_\varepsilon(t), \delta^\eta v_\varepsilon(t) \in V_0$ and $\sigma^\eta w_\varepsilon(t), \delta^\eta w_\varepsilon(t) \in \tilde{H}$. For a.e. $t \in [0, 2T - \eta]$ we use (2.29) first at time t and then at time $t + \eta$, with $\varphi := \delta^\eta v_\varepsilon(t)$ and $\psi := \delta^\eta w_\varepsilon(t)$. By summing the two expressions and then integrating in time on the interval $[0, t]$ we get

$$\int_0^t [\varepsilon^2 K_\eta(\tau) + A_\eta(\tau) + B_\eta(\tau) + \varepsilon D_\eta(\tau)]d\tau = \int_0^t W_\eta(\tau)d\tau, \tag{2.30}$$

where for a.e. $\tau \in [0, 2T - \eta]$

$$K_\eta(\tau) := \langle \sigma^\eta \ddot{v}_\varepsilon(\tau), \delta^\eta v_\varepsilon(\tau) \rangle,$$

$$\begin{aligned}
A_\eta(\tau) &:= (\mathbb{A} \sigma^\eta ev_\varepsilon(\tau), \delta^\eta ev_\varepsilon(\tau)), \\
B_\eta(\tau) &:= (\mathbb{B}(\sigma^\eta ev_\varepsilon(\tau) - \sigma^\eta w_\varepsilon(\tau)), \delta^\eta ev_\varepsilon(\tau) - \delta^\eta w_\varepsilon(\tau)), \\
D_\eta(\tau) &:= \beta(\mathbb{B} \sigma^\eta \dot{w}_\varepsilon(\tau), \delta^\eta w_\varepsilon(\tau)), \\
W_\eta(\tau) &:= (\sigma^\eta h_\varepsilon(\tau), \delta^\eta v_\varepsilon(\tau)) + \langle \sigma^\eta \ell_\varepsilon(\tau), \delta^\eta v_\varepsilon(\tau) \rangle.
\end{aligned}$$

An integration by parts in time gives

$$\begin{aligned}
\int_0^t K_\eta(\tau) d\tau &= (\sigma^\eta \dot{v}_\varepsilon(t), \delta^\eta v_\varepsilon(t)) - (\sigma^\eta \dot{v}_\varepsilon(0), \delta^\eta v_\varepsilon(0)) \\
&\quad - \int_0^t (\sigma^\eta \dot{v}_\varepsilon(\tau), \delta^\eta \dot{v}_\varepsilon(\tau)) d\tau \\
&= \int_t^{t+\eta} (\sigma^\eta \dot{v}_\varepsilon(t), \dot{v}_\varepsilon(\tau)) d\tau - \int_0^\eta (\sigma^\eta \dot{v}_\varepsilon(0), \dot{v}_\varepsilon(\tau)) d\tau \\
&\quad - \int_0^t \|\dot{v}_\varepsilon(\tau + h)\|^2 d\tau + \int_0^t \|\dot{v}_\varepsilon(\tau)\|^2 d\tau \\
&= \int_t^{t+\eta} [(\sigma^\eta \dot{v}_\varepsilon(t), \dot{v}_\varepsilon(\tau)) - \|\dot{v}_\varepsilon(\tau)\|^2] d\tau \\
&\quad - \int_0^\eta [(\sigma^\eta \dot{v}_\varepsilon(0), \dot{v}_\varepsilon(\tau)) - \|\dot{v}_\varepsilon(\tau)\|^2] d\tau. \tag{2.31}
\end{aligned}$$

Moreover

$$\int_0^t A_\eta(\tau) d\tau = \int_t^{t+\eta} (\mathbb{A} ev_\varepsilon(\tau), ev_\varepsilon(\tau)) d\tau - \int_0^\eta (\mathbb{A} ev_\varepsilon(\tau), ev_\varepsilon(\tau)) d\tau, \tag{2.32}$$

$$\begin{aligned}
\int_0^t B_\eta(\tau) d\tau &= \int_t^{t+\eta} (\mathbb{B}(ev_\varepsilon(\tau) - w_\varepsilon(\tau)), ev_\varepsilon(\tau) - w_\varepsilon(\tau)) d\tau \\
&\quad - \int_0^\eta (\mathbb{B}(ev_\varepsilon(\tau) - w_\varepsilon(\tau)), ev_\varepsilon(\tau) - w_\varepsilon(\tau)) d\tau, \tag{2.33}
\end{aligned}$$

$$\int_0^t D_\eta(\tau) d\tau = \beta \int_0^t \int_\tau^{\tau+\eta} (\mathbb{B} \sigma^\eta \dot{w}_\varepsilon(\tau), \dot{w}_\varepsilon(s)) ds d\tau, \tag{2.34}$$

$$\begin{aligned}
\int_0^t W_\eta(\tau) d\tau &= \int_0^t \int_\tau^{\tau+\eta} (\sigma^\eta h_\varepsilon(\tau), \dot{v}_\varepsilon(s)) ds d\tau + \int_{t-\eta}^t \langle \sigma^\eta \ell_\varepsilon(\tau), v_\varepsilon(\tau + \eta) \rangle d\tau \\
&\quad - \int_\eta^t \int_{\tau-\eta}^{\tau+\eta} \langle \dot{\ell}_\varepsilon(s), v_\varepsilon(\tau) \rangle ds d\tau - \int_0^\eta \langle \sigma^\eta \ell_\varepsilon(\tau), v_\varepsilon(\tau) \rangle d\tau. \tag{2.35}
\end{aligned}$$

We now divide by η all terms of (2.31)–(2.35). Observing that

$$\sigma^\eta h_\varepsilon \xrightarrow[\eta \rightarrow 0^+]{} 2h_\varepsilon \quad \text{strongly in } L^2(0, T; H),$$

$$\int_0^t \left\| \int_\tau^{\tau+\eta} \dot{v}_\varepsilon(s) ds - \dot{v}_\varepsilon(\tau) \right\|^2 d\tau \xrightarrow[\eta \rightarrow 0^+]{} 0,$$

$$\int_\eta^t \left\| \int_{\tau-\eta}^{\tau+\eta} \dot{\ell}_\varepsilon(s) ds - \dot{\ell}_\varepsilon(\tau) \right\|_{V_0'}^2 d\tau \xrightarrow[\eta \rightarrow 0^+]{} 0,$$

thanks to (2.20) in $[0, 2T)$ and (2.27), we can pass to the limit as $\eta \rightarrow 0^+$, and from (2.30) we obtain that (2.25) is satisfied for every $t \in [0, T]$. \square

Proof of Proposition 2.6 (Continuation). Now we want to prove (2.17). By using (2.25), for every $t \in [0, T]$ we can write

$$\begin{aligned} & \frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon(t)\|^2 + \frac{1}{2}((\mathbb{A} + \mathbb{B})ev_\varepsilon(t), ev_\varepsilon(t)) \\ &= \frac{\varepsilon^2}{2} \|v_\varepsilon^1\|^2 + \frac{1}{2}((\mathbb{A} + \mathbb{B})ev_\varepsilon^0, ev_\varepsilon^0) + \mathcal{W}_\varepsilon(t) \\ & - \frac{1}{2}(\mathbb{B}w_\varepsilon(t), w_\varepsilon(t)) + (\mathbb{B}ev_\varepsilon(t), w_\varepsilon(t)) - \beta \varepsilon \int_0^t (\mathbb{B}\dot{w}_\varepsilon(\tau), \dot{w}_\varepsilon(\tau)) d\tau. \end{aligned} \quad (2.36)$$

Let $\Psi_\varepsilon: [0, T] \rightarrow [0, +\infty)$ be defined by

$$\Psi_\varepsilon(t) := \frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon(t)\|^2 + \frac{1}{2}((\mathbb{A} + \mathbb{B})ev_\varepsilon(t), ev_\varepsilon(t));$$

since $w_\varepsilon \in C^0([0, T]; \tilde{H})$, thanks to (2.19) and (2.36) we have $\Psi_\varepsilon \in C^0([0, T])$.

Now we fix $t \in [0, T]$. Given a sequence $\{t_k\}_k \subset [0, T]$ such that $t_k \rightarrow t$ as $k \rightarrow +\infty$, we define

$$\mathcal{E}_k := \frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon(t_k) - \dot{v}_\varepsilon(t)\|^2 + \frac{1}{2}((\mathbb{A} + \mathbb{B})(ev_\varepsilon(t_k) - ev_\varepsilon(t)), ev_\varepsilon(t_k) - ev_\varepsilon(t)).$$

By elementary computations we have

$$\mathcal{E}_k = \Psi_\varepsilon(t_k) + \Psi_\varepsilon(t) - \varepsilon^2(\dot{v}_\varepsilon(t_k), \dot{v}_\varepsilon(t)) - ((\mathbb{A} + \mathbb{B})ev_\varepsilon(t_k), ev_\varepsilon(t)),$$

therefore, by (2.3) and (2.6) there exists a positive constant $C = C(\mathbb{A}, \mathbb{B}, \Omega)$ such that

$$\begin{aligned} & \varepsilon^2 \|\dot{v}_\varepsilon(t_k) - \dot{v}_\varepsilon(t)\|^2 + \|v_\varepsilon(t_k) - v_\varepsilon(t)\|_V^2 \\ & \leq C \left(\Psi_\varepsilon(t_k) + \Psi_\varepsilon(t) - \varepsilon^2(\dot{v}_\varepsilon(t_k), \dot{v}_\varepsilon(t)) - ((\mathbb{A} + \mathbb{B})ev_\varepsilon(t_k), ev_\varepsilon(t)) \right). \end{aligned}$$

The right-hand side of the previous inequality tends to 0 as $k \rightarrow +\infty$ because of (2.19) and the continuity of Ψ_ε . Since $z_\varepsilon \in C^0([0, T]; V)$, by (2.7), and $u_\varepsilon = v_\varepsilon + z_\varepsilon$, we obtain (2.17). \square

3. Statement of the main results

In this section we present the main results about the convergence, as $\varepsilon \rightarrow 0^+$, of the solutions u_ε . We assume the following hypotheses on the dependence on $\varepsilon > 0$ of our data:

(H1) $\{f_\varepsilon\}_\varepsilon \subset L^2(0, T; H)$ and $\{g_\varepsilon\}_\varepsilon \subset H^1(0, T; V'_0)$ such that

$$\begin{aligned} f_\varepsilon & \xrightarrow{\varepsilon \rightarrow 0^+} f \quad \text{strongly in } L^2(0, T; H), \\ g_\varepsilon & \xrightarrow{\varepsilon \rightarrow 0^+} g \quad \text{strongly in } W^{1,1}(0, T; V'_0); \end{aligned}$$

(H2) $\{z_\varepsilon\}_\varepsilon \subset H^2(0, T; H) \cap H^1(0, T; V)$ such that

$$z_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} z \quad \text{strongly in } W^{2,1}(0, T; H) \cap W^{1,1}(0, T; V);$$

(H3) $\{u_{\varepsilon, in}\}_{\varepsilon} \subset C^0((-\infty, 0]; V) \cap C^1((-\infty, 0]; H)$, $u_{in} \in C^0((-\infty, 0]; V)$, and there exist $a > 0$ such that

$$\begin{aligned} u_{\varepsilon, in} &\xrightarrow{\varepsilon \rightarrow 0^+} u_{in} && \text{strongly in } C^0([-a, 0]; V), \\ \varepsilon \dot{u}_{\varepsilon, in} &\xrightarrow{\varepsilon \rightarrow 0^+} 0 && \text{strongly in } C^0([-a, 0]; H), \\ \int_{-\infty}^{-a} \frac{1}{\beta \varepsilon} e^{\frac{\tau}{\beta \varepsilon}} \|u_{\varepsilon, in}(\tau)\|_V d\tau &\xrightarrow{\varepsilon \rightarrow 0^+} 0, \\ \int_{-\infty}^{-a} \frac{1}{\beta \varepsilon} e^{\frac{\tau}{\beta \varepsilon}} \|u_{in}(\tau)\|_V d\tau &\xrightarrow{\varepsilon \rightarrow 0^+} 0. \end{aligned}$$

Remark 3.1. Let $u_{\varepsilon}^0 = u_{\varepsilon, in}(0)$, $u_{\varepsilon}^1 = \dot{u}_{\varepsilon, in}(0)$, and $u^0 = u_{in}(0)$. Hypothesis (H3) implies

$$u_{\varepsilon}^0 \xrightarrow{\varepsilon \rightarrow 0^+} u^0 \quad \text{strongly in } V \quad \text{and} \quad \varepsilon u_{\varepsilon}^1 \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } H.$$

Our purpose is to show that the solutions u_{ε} converge, as $\varepsilon \rightarrow 0^+$, to the solution u_0 of the stationary problem (1.5) with boundary condition (1.2). The notion of solution to this problem is the usual one:

$$\begin{cases} u_0(t) \in V, & u_0(t) - z(t) \in V_0, & \text{for a.e. } t \in [0, T], \\ -\operatorname{div}(\mathbb{A} e u_0(t)) = f(t) + g(t) & & \text{for a.e. } t \in [0, T]. \end{cases} \quad (3.1)$$

Remark 3.2. The existence and uniqueness of a solution u_0 to (3.1) follows easily from the Lax-Milgram Lemma. Since $f + g \in L^2(0, T; V'_0)$, the estimate for the solution implies also $u_0 \in L^2(0, T; V)$.

We shall sometimes use the corresponding problem with homogeneous boundary conditions:

$$\begin{cases} v_0(t) \in V_0 & \text{for a.e. } t \in [0, T], \\ -\operatorname{div}(\mathbb{A} e v_0(t)) = h(t) + \ell(t) & \text{for a.e. } t \in [0, T], \end{cases} \quad (3.2)$$

with $h \in L^2(0, T; H)$ and $\ell \in H^1(0, T; V'_0)$.

Remark 3.3. The function u_0 is a solution to (3.1) if and only if $v_0 = u_0 - z$ is a solution to (3.2) with

$$h(t) = f(t) \quad \text{and} \quad \ell(t) = g(t) + \operatorname{div}(\mathbb{A} e z(t)).$$

The following lemma will be used to prove the regularity with respect to time of the solution to (3.1).

Lemma 3.4. *Let $m \in \mathbb{N}$ and $p \in [1, +\infty)$. If $f = 0$, $g \in W^{m,p}(0, T; V'_0)$, and $z \in W^{m,p}(0, T; V)$, then the solution u_0 to problem (3.1) satisfies $u_0 \in W^{m,p}(0, T; V)$.*

Proof. By Remark 3.3 it is enough to consider the case $z = 0$. Let $R : V'_0 \rightarrow V_0$ be the resolvent operator defined as follows:

$$R(\psi) = \varphi \iff \begin{cases} \varphi \in V_0, \\ -\operatorname{div}(\mathbb{A} e \varphi) = \psi. \end{cases}$$

Since $u_0(t) = R(g(t))$, the conclusion follows from the continuity of the linear operator R . \square

Remark 3.5. If $f = 0$, since $g \in W^{1,1}(0, T; V'_0)$ and $z \in W^{1,1}(0, T; V)$, we can apply Lemma 3.4 to obtain that the solution u_0 to (3.1) belongs to $W^{1,1}(0, T; V)$, hence $u_0 \in C^0([0, T]; V)$.

In the final statement of the next theorem, besides (H1)–(H3) we assume $f_\varepsilon = 0$ and the following compatibility condition: there exists an extension of g (still denoted by g) such that

$$g \in W^{1,1}(-a, T; V'_0) \quad \text{and} \quad -\operatorname{div}(\mathbb{A}eu_{in}(t)) = g(t) \quad \text{for } t \in [-a, 0]. \quad (3.3)$$

The meaning of (3.3) is that $u_{in}(t)$ is in equilibrium with external loads for $t \in [-a, 0]$. This condition must be required if we want to obtain uniform convergence of u_ε to u_0 also near $t = 0$.

We are now in a position to state the main results of this paper.

Theorem 3.6. *Let us assume (H1)–(H3). Let u_ε be the solution to the viscoelastic dynamic system (2.9) and let u_0 be the solution to the stationary problem (3.1). Then*

$$u_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} u_0 \quad \text{strongly in } L^2(0, T; V), \quad (3.4)$$

$$\varepsilon \dot{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; H). \quad (3.5)$$

If, in addition, $f_\varepsilon = 0$ for every $\varepsilon > 0$, then

$$u_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} u_0 \quad \text{strongly in } L^\infty(\eta, T; V) \text{ for every } \eta \in (0, T), \quad (3.6)$$

$$\varepsilon \dot{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(\eta, T; H) \text{ for every } \eta \in (0, T). \quad (3.7)$$

If $f_\varepsilon = 0$ for every $\varepsilon > 0$ and the compatibility condition (3.3) holds, then we have also

$$u_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} u_0 \quad \text{strongly in } L^\infty(0, T; V), \quad (3.8)$$

$$\varepsilon \dot{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(0, T; H). \quad (3.9)$$

In the case of solutions to problems (2.10) we have the following results, assuming that

$$u_\varepsilon^0 \xrightarrow{\varepsilon \rightarrow 0^+} u^0 \quad \text{strongly in } V \quad \text{and} \quad \varepsilon u_\varepsilon^1 \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } H. \quad (3.10)$$

Theorem 3.7. *Let us assume (H1), (H2), and (3.10). Let u_ε be the solution to the viscoelastic dynamic system (2.10), with $\varphi_\varepsilon = f_\varepsilon$ and $\gamma_\varepsilon = g_\varepsilon$, and let u_0 be the solution to the stationary problem (3.1). Then (3.4) and (3.5) hold. Moreover, if $f_\varepsilon = 0$ for every $\varepsilon > 0$, then (3.6) and (3.7) hold.*

Theorems 3.6 and 3.7 will be proved in several steps. First, we prove (3.8) and (3.9) when $f_\varepsilon = 0$ and the compatibility condition (3.3) holds (Theorem 4.1). For $g \in H^2(0, T; V'_0)$ the proof is based on the estimate in Lemma 3.8 below, which is derived from the energy-dissipation balance (2.25).

The general case is obtained by an approximation argument based on the same estimate.

Next, we prove that (3.4) holds for the solutions of (2.10) if $\gamma_\varepsilon = \gamma = 0$, $z_\varepsilon = 0$, $u_\varepsilon^0 = 0$, and $u_\varepsilon^1 = 0$ (Proposition 6.1). The proof is obtained by means of a careful estimate of the solutions to the elliptic system (5.17) obtained from (2.13) via Laplace Transform (Section 5). Under the general assumptions (H1), (H2), and (3.10) the same result is deduced from the previous one by an approximation argument based again on Lemma 3.8 below.

Then, (3.5) is obtained from (3.4) using a suitable test function in (2.10) (Theorem 6.3). A further approximation argument gives (3.4) and (3.5) under the assumptions (H1), (H2), and (H3) (Theorem 6.4).

Finally, if $f_\varepsilon = 0$, we obtain (3.6) and (3.7) from (3.4) and (3.5) (Lemma 7.1), concluding the proof of Theorems 3.6 and 3.7.

The following lemma, derived from the energy-dissipation balance (2.25) of Proposition 2.7, will be frequently used to approximate the solutions of (2.13) by means of solutions corresponding to more regular data.

Lemma 3.8. *Given $\varepsilon > 0$, $\varphi_\varepsilon \in L^2(0, T; H)$, $\ell_\varepsilon \in H^1(0, T; V'_0)$, $v_\varepsilon^0 \in V_0$, and $v_\varepsilon^1 \in H$, let v_ε be the solution to (2.13) with $h_\varepsilon = \varepsilon\varphi_\varepsilon$. Then there exists a positive constant $C_E = C_E(\mathbb{A}, \mathbb{B}, \Omega, T)$, independent of ε , such that*

$$\begin{aligned} \varepsilon^2 \|\dot{v}_\varepsilon\|_{L^\infty(0, T; H)}^2 + \|v_\varepsilon\|_{L^\infty(0, T; V)}^2 \\ \leq C_E \left(\varepsilon^2 \|v_\varepsilon^1\|^2 + \|v_\varepsilon^0\|_V^2 + \|\varphi_\varepsilon\|_{L^1(0, T; H)}^2 + \|\ell_\varepsilon\|_{W^{1,1}(0, T; V'_0)}^2 \right). \end{aligned}$$

Proof. By the energy-dissipation balance (2.25) proved in Proposition 2.7 and by (2.3) and (2.6) there exists a positive constant $C = C(\mathbb{A}, \mathbb{B}, \Omega)$ such that for every $t \in [0, T]$ we have

$$\varepsilon^2 \|\dot{v}_\varepsilon(t)\|^2 + \|v_\varepsilon(t)\|_V^2 \leq C \left(\varepsilon^2 \|v_\varepsilon^1\|^2 + \|v_\varepsilon^0\|_V^2 + \mathcal{W}_\varepsilon(t) \right), \quad (3.11)$$

where the work is now defined by

$$\begin{aligned} \mathcal{W}_\varepsilon(t) &= \langle \ell_\varepsilon(t), v_\varepsilon(t) \rangle - \langle \ell_\varepsilon(0), v_\varepsilon^0 \rangle \\ &\quad - \int_0^t \langle \dot{\ell}_\varepsilon(\tau), v_\varepsilon(\tau) \rangle d\tau + \int_0^t (\varphi_\varepsilon(\tau), \varepsilon \dot{v}_\varepsilon(\tau)) d\tau. \end{aligned} \quad (3.12)$$

Let $K_\varepsilon := \varepsilon \|\dot{v}_\varepsilon(t)\|_{L^\infty(0, T; H)}$ and $E_\varepsilon := \|v_\varepsilon(t)\|_{L^\infty(0, T; V)}$, which are finite by (2.17). Thanks to (3.11) and (3.12) for every $t \in [0, T]$ we get

$$\begin{aligned} \varepsilon^2 \|\dot{v}_\varepsilon(t)\|^2 + \|v_\varepsilon(t)\|_V^2 \\ \leq C \left(\varepsilon^2 \|v_\varepsilon^1\|^2 + \|v_\varepsilon^0\|_V^2 + \left(3 + \frac{2}{T}\right) \|\ell_\varepsilon\|_{W^{1,1}(0, T; V'_0)} E_\varepsilon + \|\varphi_\varepsilon\|_{L^1(0, T; H)} K_\varepsilon \right). \end{aligned}$$

By passing to the supremum with respect to t and using the Young Inequality we can find a positive constant $C_E = C_E(\mathbb{A}, \mathbb{B}, \Omega, T)$ such that

$$K_\varepsilon^2 + E_\varepsilon^2 \leq C_E \left(\varepsilon^2 \|v_\varepsilon^1\|^2 + \|v_\varepsilon^0\|_V^2 + \|\varphi_\varepsilon\|_{L^1(0, T; H)}^2 + \|\ell_\varepsilon\|_{W^{1,1}(0, T; V'_0)}^2 \right),$$

which concludes the proof. \square

In the proof of Theorem 3.6 we shall use the following lemma, which ensure that it is enough to consider the case $z_\varepsilon = 0$ and $z = 0$.

Lemma 3.9. *If Theorem 3.6 holds when $z_\varepsilon = 0$ for every $\varepsilon > 0$, then it holds for arbitrary $\{z_\varepsilon\}_\varepsilon$ and z satisfying (H2).*

Proof. It is not restrictive to assume $\operatorname{div}(\mathbb{B}ez_\varepsilon(0)) = \operatorname{div}(\mathbb{B}ez(0)) = 0$. Indeed, if this is not the case, we can consider the solutions z_ε^0 and z^0 to the stationary problems

$$\begin{cases} z_\varepsilon^0 \in V_0, \\ -\operatorname{div}(\mathbb{B}ez_\varepsilon^0) = \operatorname{div}(\mathbb{B}ez_\varepsilon(0)), \end{cases} \quad \text{and} \quad \begin{cases} z^0 \in V_0, \\ -\operatorname{div}(\mathbb{B}ez^0) = \operatorname{div}(\mathbb{B}ez(0)), \end{cases}$$

and we can replace $z_\varepsilon(t)$ and $z(t)$ by $\tilde{z}_\varepsilon(t) := z_\varepsilon(t) + z_\varepsilon^0$ and $\tilde{z}(t) := z(t) + z^0$. It is clear that $\operatorname{div}(\mathbb{B}e\tilde{z}_\varepsilon(0)) = \operatorname{div}(\mathbb{B}e\tilde{z}(0)) = 0$ and that problems (2.9) and (3.1) do not change passing from z_ε and z to \tilde{z}_ε and \tilde{z} .

Let $\psi_\varepsilon, \psi: [0, T] \rightarrow V'_0$ be the functions defined by

$$\psi_\varepsilon(t) := \begin{cases} 0 & \text{if } t \in (-\infty, 0), \\ \operatorname{div}(\mathbb{B}ez_\varepsilon(t)) & \text{if } t \in [0, T], \\ \operatorname{div}(\mathbb{B}ez_\varepsilon(T)) & \text{if } t \in (T, +\infty), \end{cases} \quad (3.13)$$

$$\psi(t) := \begin{cases} 0 & \text{if } t \in (-\infty, 0), \\ \operatorname{div}(\mathbb{B}ez(t)) & \text{if } t \in [0, T], \\ \operatorname{div}(\mathbb{B}ez(T)) & \text{if } t \in (T, +\infty). \end{cases} \quad (3.14)$$

Since

$$\operatorname{div}(\mathbb{B}ez_\varepsilon(0)) = \operatorname{div}(\mathbb{B}ez(0)) = 0, \quad z_\varepsilon \in H^1(0, T; V), \quad z \in W^{1,1}(0, T; V),$$

we obtain $\psi_\varepsilon \in H^1_{loc}(\mathbb{R}; V'_0)$ and $\psi \in W^{1,1}_{loc}(\mathbb{R}; V'_0)$. Moreover, thanks to (H2) we have

$$\psi_\varepsilon \xrightarrow[\varepsilon \rightarrow 0^+]{} \psi \quad \text{strongly in } W^{1,1}_{loc}(\mathbb{R}; V'_0). \quad (3.15)$$

Since u_ε is the solution to (2.9), by Remark 2.3 it solves (2.10) with $\gamma_\varepsilon = g_\varepsilon - p_\varepsilon$ and initial conditions defined by (2.11), where p_ε is defined by (2.12). By Remark 2.4 the function $v_\varepsilon = u_\varepsilon - z_\varepsilon$ is the solution to (2.13) with

$$\begin{aligned} h_\varepsilon(t) &= f_\varepsilon(t) - \varepsilon^2 \ddot{z}_\varepsilon(t), \\ \ell_\varepsilon(t) &= g_\varepsilon(t) - p_\varepsilon(t) + \operatorname{div}((\mathbb{A} + \mathbb{B})ez_\varepsilon(t)) \\ &\quad - \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ez_\varepsilon(\tau)) d\tau, \end{aligned} \quad (3.16)$$

and initial conditions v_ε^0 and v_ε^1 defined by (2.15). We define the family of convolution kernels $\{\rho_\varepsilon\}_\varepsilon \subset L^1(\mathbb{R})$ by

$$\rho_\varepsilon(t) := \begin{cases} \frac{1}{\beta\varepsilon} e^{-\frac{t}{\beta\varepsilon}} & \text{if } t \in [0, +\infty), \\ 0 & \text{if } t \in (-\infty, 0), \end{cases} \quad (3.17)$$

and notice that, by (3.14) and (3.13), the integral in (3.16) coincides with $(\rho_\varepsilon * \psi_\varepsilon)(t)$, hence

$$\ell_\varepsilon(t) = g_\varepsilon(t) - p_\varepsilon(t) + \operatorname{div}(\mathbb{A}ez_\varepsilon(t)) + \psi_\varepsilon(t) - (\rho_\varepsilon * \psi_\varepsilon)(t) \quad \text{for every } t \in [0, T].$$

By Remark 3.3 the function $v_0 = u_0 - z$ is the solution to (3.2) with $h = f$ and $\ell = g + \operatorname{div}(\mathbb{A}ez)$. By the definition of v_ε and v_0 it is clear that to prove the theorem it is enough to show that the conclusions of Theorem 3.6 holds for v_ε and v_0 . To this aim, we introduce the solution \tilde{v}_ε to (2.13) with $h_\varepsilon = f_\varepsilon$, $\ell_\varepsilon = g_\varepsilon - p_\varepsilon + \operatorname{div}(\mathbb{A}ez_\varepsilon)$, and $v_\varepsilon^0, v_\varepsilon^1$ defined by (2.15). Then the function $\bar{v}_\varepsilon := v_\varepsilon - \tilde{v}_\varepsilon$ satisfies (2.13) with $h_\varepsilon = -\varepsilon^2 \ddot{z}_\varepsilon$, $\ell_\varepsilon = \psi_\varepsilon - \rho_\varepsilon * \psi_\varepsilon$, and homogeneous initial conditions. By Lemma 3.8 we can write

$$\begin{aligned} \varepsilon^2 \|\ddot{v}_\varepsilon\|_{L^\infty(0,T;H)}^2 + \|\bar{v}_\varepsilon\|_{L^\infty(0,T;V)}^2 \\ \leq C_E \left(\varepsilon^2 \|\ddot{z}_\varepsilon\|_{L^1(0,T;H)}^2 + \|\psi_\varepsilon - \rho_\varepsilon * \psi_\varepsilon\|_{W^{1,1}(0,T;V'_0)}^2 \right). \end{aligned} \quad (3.18)$$

By (3.15) and by classical results on convolutions we obtain

$$\psi_\varepsilon - \rho_\varepsilon * \psi_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } W^{1,1}(0,T;V'_0).$$

Since $\{\ddot{z}_\varepsilon\}_\varepsilon$ is bounded in $L^1(0,T;H)$ by (H2), from (3.18) we deduce

$$v_\varepsilon - \tilde{v}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(0,T;V), \quad (3.19)$$

$$\varepsilon(\dot{v}_\varepsilon - \dot{\tilde{v}}_\varepsilon) \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(0,T;H). \quad (3.20)$$

By Remark 2.3 the function \tilde{v}_ε is the solution to (2.9) with g_ε replaced by $g_\varepsilon + \operatorname{div}(\mathbb{A}ez_\varepsilon)$ and $z_\varepsilon = 0$. Thanks to (H1) and (H2) we have

$$g_\varepsilon + \operatorname{div}(\mathbb{A}ez_\varepsilon) \xrightarrow{\varepsilon \rightarrow 0^+} g + \operatorname{div}(\mathbb{A}ez) \quad \text{strongly in } W^{1,1}(0,T;V'_0).$$

Since by hypothesis, Theorem 3.6 holds in the case of homogeneous boundary condition, its conclusions are valid for \tilde{v}_ε and v_0 . Thanks to (3.19) and (3.20) the same results hold for v_ε and v_0 . This concludes the proof. \square

In a similar way we can prove the following result.

Lemma 3.10. *If Theorem 3.7 holds when $z_\varepsilon = 0$ for every $\varepsilon > 0$, then it holds for arbitrary $\{z_\varepsilon\}_\varepsilon$ and z satisfying (H2).*

4. The uniform convergence

In this section we shall prove (3.8) and (3.9) of Theorem 3.6 under the compatibility condition (3.3).

Theorem 4.1. *Let us assume (H1)–(H3), the compatibility condition (3.3), and $f_\varepsilon = 0$ for every $\varepsilon > 0$. Let u_ε be the solution to the viscoelastic dynamic system (2.9) and let u_0 be the solution to the stationary problem (3.1), with $f = 0$. Then (3.8) and (3.9) hold.*

To prove the theorem we need the following lemma, which gives the result when g is more regular.

Lemma 4.2. *Under the assumptions of Theorem 4.1, if $g \in H^2(0, T; V'_0)$, then (3.8) and (3.9) hold.*

Proof. Thanks to Lemma 3.9 we can suppose $z = 0$ and $z_\varepsilon = 0$ for every $\varepsilon > 0$. Let p_ε be defined by (2.11). Since u_ε is the solution to (2.9), thanks to Remark 2.3 it solves (2.13) with $h_\varepsilon = 0$, $\ell_\varepsilon = g_\varepsilon - p_\varepsilon$, $v_\varepsilon^0 = u_{\varepsilon, in}(0)$, and $v_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0)$. We fix $b > a > 0$ and we extend the function g in (3.3) to $(-\infty, T)$ in such a way $g \in W^{1,1}(-\infty, T; V'_0)$ and $g(t) = 0$ for every $t \in (-\infty, -b]$. Since $z = 0$ we can extend u_0 by solving the following problem:

$$\begin{cases} u_0(t) \in V_0 & \text{for every } t \in (-\infty, T], \\ -\operatorname{div}(\mathbb{A}eu_0(t)) = g(t) & \text{for every } t \in (-\infty, T]. \end{cases}$$

We observe that $u_0 = 0$ on $(-\infty, -b]$ and $u_0 = u_{in}$ on $[-a, 0]$ by the compatibility condition (3.3).

Assume $g \in H^2(0, T; V'_0)$. By Lemma 3.4 (with $z = 0$) we have $u_0 \in H^2(0, T; V)$, hence by (3.1) we get for a.e. $t \in [0, T]$

$$\begin{aligned} \varepsilon^2 \ddot{u}_0(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})eu_0(t)) + \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}eu_0(\tau)) d\tau \\ = \varepsilon^2 \ddot{u}_0(t) + g(t) - \operatorname{div}(\mathbb{B}eu_0(t)) + (\rho_\varepsilon * \operatorname{div}(\mathbb{B}eu_0))(t) - \tilde{p}_\varepsilon(t), \end{aligned} \quad (4.1)$$

where ρ_ε is defined by (3.17) and

$$\tilde{p}_\varepsilon(t) := e^{-\frac{t}{\beta\varepsilon}} \tilde{g}_\varepsilon^0 \quad \text{with} \quad \tilde{g}_\varepsilon^0 := \int_{-b}^0 \frac{1}{\beta\varepsilon} e^{\frac{\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}eu_0(\tau)) d\tau.$$

Let $q_\varepsilon := g_\varepsilon - g + \operatorname{div}(\mathbb{B}eu_0) - (\rho_\varepsilon * \operatorname{div}(\mathbb{B}eu_0)) - p_\varepsilon + \tilde{p}_\varepsilon$. By (4.1) the function $\tilde{u}_\varepsilon := u_\varepsilon - u_0$ satisfies (2.13) with $h_\varepsilon = -\varepsilon^2 \ddot{u}_0$, $\ell_\varepsilon = q_\varepsilon$, $v_\varepsilon^0 = u_{\varepsilon, in}(0) - u_0(0)$, and $v_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0) - \dot{u}_0(0)$.

Since $g \in W^{1,1}(-\infty, T; V'_0)$ and $g = 0$ on $(-\infty, -b]$, by Lemma 3.4 we obtain $u_0 \in W^{1,1}(-\infty, T; V)$ and therefore $\operatorname{div}(\mathbb{B}eu_0) \in W^{1,1}(-\infty, T; V'_0)$. Then the properties of convolutions imply

$$\rho_\varepsilon * \operatorname{div}(\mathbb{B}eu_0) \xrightarrow{\varepsilon \rightarrow 0^+} \operatorname{div}(\mathbb{B}eu_0) \quad \text{strongly in } W^{1,1}(-\infty, T; V'_0). \quad (4.2)$$

As we have already observed, by the compatibility condition (3.3) we have $u_0 = u_{in}$ on $[-a, 0]$, hence

$$\begin{aligned} \|\tilde{g}_\varepsilon^0 - g_\varepsilon^0\|_{V'_0} &\leq \int_{-\infty}^{-a} \frac{1}{\beta\varepsilon} e^{\frac{\tau}{\beta\varepsilon}} (\|\operatorname{div}(\mathbb{B}(eu_{\varepsilon, in}(\tau)))\|_{V'_0} + \|\operatorname{div}(\mathbb{B}(eu_0(\tau)))\|_{V'_0}) d\tau \\ &\quad + \|\operatorname{div}(\mathbb{B}(eu_{\varepsilon, in} - eu_{in}))\|_{L^\infty(-a, 0; V'_0)}. \end{aligned}$$

Thanks to (H3) we obtain $\tilde{g}_\varepsilon^0 - g_\varepsilon^0 \rightarrow 0$ strongly in V'_0 as $\varepsilon \rightarrow 0^+$. Hence

$$\tilde{p}_\varepsilon - p_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } W^{1,1}(0, T; V'_0). \quad (4.3)$$

By (H1), (4.2), and (4.3) we have

$$q_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } W^{1,1}(0, T; V'_0). \quad (4.4)$$

Since $u_0(0) = u_{in}(0)$, (H3) gives

$$u_{\varepsilon,in}^0 := u_{\varepsilon,in}(0) - u_0(0) \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } V, \quad (4.5)$$

$$\varepsilon u_{\varepsilon,in}^1 := \varepsilon(\dot{u}_{\varepsilon,in}(0) - \dot{u}_0(0)) \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } H. \quad (4.6)$$

By using Lemma 3.8 we get

$$\begin{aligned} & \varepsilon^2 \|\dot{u}_\varepsilon\|_{L^\infty(0,T;H)}^2 + \|\bar{u}_\varepsilon\|_{L^\infty(0,T;V)}^2 \\ & \leq C_E \left(\varepsilon^2 \|u_{\varepsilon,in}^1\|^2 + \|u_{\varepsilon,in}^0\|_V^2 + \varepsilon^2 \|\ddot{u}_0\|_{L^1(0,T;H)}^2 + \|q_\varepsilon\|_{W^{1,1}(0,T;V'_0)}^2 \right), \end{aligned}$$

therefore thanks to (4.4), (4.5), and (4.6) we obtain the conclusion. \square

In the proof of Theorems 4.1, 6.2, and 6.4 we shall use the following density result.

Lemma 4.3. *Let X, Y be two Hilbert spaces such that $X \hookrightarrow Y$ continuously, with X dense in Y . Then for every $m, n \in \mathbb{N}$ with $m \leq n$, and $p \in [1, 2]$ the space $H^n(0, T; X)$ is dense in $W^{m,p}(0, T; Y)$.*

Proof. Since every simple function with values in Y can be approximated by simple functions with values in X , it is easy to see that $L^2(0, T; X)$ is dense in $L^p(0, T; Y)$.

To prove the result for $m = 1$ we fix $u \in W^{1,p}(0, T; Y)$. By the density of $L^2(0, T; X)$ in $L^p(0, T; Y)$ we can find a sequence $\{\psi_k\}_k \subset L^2(0, T; X)$ such that $\psi_k \rightarrow \dot{u}$ strongly in $L^p(0, T; Y)$ as $k \rightarrow +\infty$. By the density of X in Y there exists $\{u_k^0\}_k \subset X$ such that $u_k^0 \rightarrow u(0)$ strongly in Y as $k \rightarrow +\infty$. Now we define

$$u_k(t) := \int_0^t \psi_k(\tau) d\tau + u_k^0.$$

It is easy to see that $\{u_k\}_k \subset H^1(0, T; X)$ and

$$u_k \xrightarrow{k \rightarrow +\infty} u \quad \text{strongly in } W^{1,p}(0, T; Y).$$

Arguing by induction we can prove that for every integer $m \geq 0$ the space $H^m(0, T; X)$ is dense in $W^{m,p}(0, T; Y)$. Since $H^n(0, T; X)$ is dense in $H^m(0, T; X)$, the conclusion follows. \square

We are now in a position to deduce Theorem 4.1 from Lemma 4.2 by means of an approximation argument.

Proof of Theorem 4.1. Thanks to Lemma 3.9 we can suppose $z = 0$ and $z_\varepsilon = 0$ for every $\varepsilon > 0$. We fix $\delta > 0$. By Lemma 4.3 there exists a function $\psi \in H^2(0, T; V'_0)$ such that

$$\|\psi - g\|_{W^{1,1}(0,T;V'_0)} < \delta. \quad (4.7)$$

By (H1) there exists a positive number $\varepsilon_0 = \varepsilon_0(\delta)$ such that

$$\|\psi - g_\varepsilon\|_{W^{1,1}(0,T;V'_0)} < \delta \quad \text{for every } \varepsilon \in (0, \varepsilon_0). \quad (4.8)$$

Let p_ε be defined by (2.12). Since u_ε is the solution to (2.9) with $f_\varepsilon = 0$ and $z_\varepsilon = 0$, thanks to Remark 2.3 it solves (2.13) with $h_\varepsilon = 0$, $\ell_\varepsilon = g_\varepsilon - p_\varepsilon$,

$v_\varepsilon^0 = u_{\varepsilon, in}(0)$, and $v_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0)$. Moreover, let \tilde{u}_ε be solution to (2.13) with $h_\varepsilon = 0$, $\ell_\varepsilon = \psi - p_\varepsilon$, $v_\varepsilon^0 = u_{\varepsilon, in}(0)$, and $v_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0)$, and let \tilde{u}_0 be the solution to (3.2) with $h = 0$ and $\ell = \psi$. Thanks to Remark 2.3 the function \tilde{u}_ε is the solution to (2.9) with $f_\varepsilon = 0$, $g_\varepsilon = \psi$, and $z_\varepsilon = 0$, hence by Lemma 4.2 we have

$$\tilde{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} \tilde{u}_0 \quad \text{strongly in } L^\infty(0, T; V), \quad (4.9)$$

$$\varepsilon \dot{\tilde{u}}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(0, T; H). \quad (4.10)$$

We now consider the functions $\bar{u}_0 := \tilde{u}_0 - u_0$ and $\bar{u}_\varepsilon := \tilde{u}_\varepsilon - u_\varepsilon$. Since \bar{u}_0 is the solution to (3.2), with $h = 0$ and $\ell = \psi - g$, by the Lax-Milgram Lemma we get

$$\|\bar{u}_0\|_{L^\infty(0, T; V)} \leq \frac{C_P^2 + 1}{c_A} \|\psi - g\|_{L^\infty(0, T; V'_0)} \leq \frac{C_P^2 + 1}{c_A} \left(1 + \frac{1}{T}\right) \|\psi - g\|_{W^{1,1}(0, T; V'_0)}. \quad (4.11)$$

Moreover, since \bar{u}_ε is the solution to (2.13), with $h_\varepsilon = 0$, $\ell_\varepsilon = \psi - g_\varepsilon$, $v_\varepsilon^0 = 0$, and $v_\varepsilon^1 = 0$, thanks to Lemma 3.8 we get

$$\varepsilon^2 \|\dot{\bar{u}}_\varepsilon\|_{L^\infty(0, T; H)}^2 + \|\bar{u}_\varepsilon\|_{L^\infty(0, T; V)}^2 \leq C_E \|\psi - g_\varepsilon\|_{W^{1,1}(0, T; V'_0)}^2. \quad (4.12)$$

By combining (4.7), (4.8), (4.11), and (4.12), we can find a positive constant $C = C(\mathbb{A}, \mathbb{B}, \Omega, T)$ such that

$$\varepsilon \|\dot{\bar{u}}_\varepsilon\|_{L^\infty(0, T; H)} + \|\bar{u}_\varepsilon\|_{L^\infty(0, T; V)} + \|\bar{u}_0\|_{L^\infty(0, T; V)} \leq C\delta \quad (4.13)$$

for every $\varepsilon \in (0, \varepsilon_0)$. Since

$$\begin{aligned} \|u_\varepsilon - u_0\|_{L^\infty(0, T; V)} &\leq \|\bar{u}_\varepsilon\|_{L^\infty(0, T; V)} + \|\tilde{u}_\varepsilon - \tilde{u}_0\|_{L^\infty(0, T; V)} + \|\bar{u}_0\|_{L^\infty(0, T; V)}, \\ \varepsilon \|\dot{u}_\varepsilon\|_{L^\infty(0, T; H)} &\leq \varepsilon \|\dot{\bar{u}}_\varepsilon\|_{L^\infty(0, T; H)} + \varepsilon \|\dot{\tilde{u}}_\varepsilon\|_{L^\infty(0, T; H)}, \end{aligned}$$

by (4.9), (4.10), and (4.13) we have

$$\limsup_{\varepsilon \rightarrow 0^+} \|u_\varepsilon - u_0\|_{L^\infty(0, T; V)} \leq C\delta \quad \text{and} \quad \limsup_{\varepsilon \rightarrow 0^+} \|\varepsilon \dot{u}_\varepsilon\|_{L^\infty(0, T; H)} \leq C\delta.$$

The conclusion follows from the arbitrariness of $\delta > 0$. \square

5. Use of the Laplace Transform

In this section we shall use the Laplace Transform to prepare the proof of the convergence, as $\varepsilon \rightarrow 0^+$, of the solutions of the problems

$$v_\varepsilon \in \mathcal{V}_0, \quad (5.1a)$$

$$\begin{aligned} \varepsilon^2 \ddot{v}_\varepsilon(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})ev_\varepsilon(t)) + \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_\varepsilon(\tau)) d\tau &= h_\varepsilon(t) \\ \text{for a.e. } t \in [0, T], \end{aligned} \quad (5.1b)$$

$$v_\varepsilon(0) = 0 \quad \text{in } H \quad \text{and} \quad \dot{v}_\varepsilon(0) = 0 \quad \text{in } V'_0, \quad (5.1c)$$

to the solution v_0 of the problem

$$\begin{cases} v_0(t) \in V_0 & \text{for a.e. } t \in [0, T], \\ -\operatorname{div}(\mathbb{A} \epsilon v_0(t)) = h(t) & \text{for a.e. } t \in [0, T], \end{cases} \quad (5.2)$$

when $\{h_\varepsilon\}_\varepsilon \subset L^2(0, T; H)$, $h \in L^2(0, T; H)$, and

$$h_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} h \quad \text{strongly in } L^2(0, T; H), \quad (5.3)$$

This partial result will be the starting point for the proof of the convergence in $L^2(0, T; V)$ under the general assumptions of Theorem 3.6.

5.1. The Laplace Transform for functions with values in Hilbert spaces

Given a complex Hilbert space X , let $r \in L^1_{loc}(0, +\infty; X)$ be a function such that

$$\int_0^{+\infty} e^{-\alpha t} \|r(t)\|_X dt < +\infty \quad \text{for every } \alpha > 0, \quad (5.4)$$

and let $\mathbb{C}_+ := \{s \in \mathbb{C} : \Re(s) > 0\}$. The Laplace Transform of r is the function $\hat{r} : \mathbb{C}_+ \rightarrow X$ defined by

$$\hat{r}(s) := \int_0^{+\infty} e^{-st} r(t) dt \quad \text{for every } s \in \mathbb{C}_+. \quad (5.5)$$

Besides \hat{r} , we shall also use the notation $\mathcal{L}(r)$, which is sometimes written as $\mathcal{L}_t(r(t))$, with dummy variable t . In the particular case $r \in L^\infty(0, +\infty; X)$ we have

$$\|\hat{r}(s)\|_X \leq \frac{1}{s_1} \|r\|_{L^\infty(0, +\infty; X)} \quad \text{for every } s = s_1 + is_2 \in \mathbb{C}_+, \text{ with } s_1, s_2 \in \mathbb{R}.$$

There is a close connection between the Laplace Transform and the Fourier Transform, defined for every $\rho \in L^1(\mathbb{R}; X)$ as the function $\mathcal{F}(\rho) \in L^\infty(\mathbb{R}; X)$ given by

$$\mathcal{F}(\rho)(\xi) = \int_{-\infty}^{+\infty} e^{-i\xi t} \rho(t) dt \quad \text{for every } \xi \in \mathbb{R}. \quad (5.6)$$

For $\mathcal{F}(\rho)$ we use also the notation $\mathcal{F}_t(\rho(t))$ with dummy variable t . For the main properties of the Fourier and Laplace Transforms of functions with values in Hilbert spaces we refer to [2].

We extend the function r satisfying (5.4) by setting $r(t) = 0$ for every $t < 0$. By (5.5) and (5.6) we have

$$\mathcal{L}_t(r(t))(s) = \mathcal{F}_t(e^{-s_1 t} r(t))(s_2) \quad \text{for every } s = s_1 + is_2 \in \mathbb{C}_+, s_1, s_2 \in \mathbb{R}.$$

We remark that the Laplace Transform commutes with linear transformations, as shown in the following proposition (see, for instance [2, Proposition 1.6.2]).

Proposition 5.1. *Let X and Y be two complex Hilbert spaces, let $r \in L^1_{loc}(0, +\infty; X)$, and let T be a continuous linear operator from X to Y . Then $T \circ r \in L^1_{loc}(0, +\infty; Y)$. If in addition, r satisfies (5.4), then the same property holds also for $T \circ r$, with X replaced by Y , and $\mathcal{L}(T \circ r)(s) = (T \circ \hat{r})(s)$ for every $s \in \mathbb{C}_+$.*

Now we consider the Inverse Laplace Transform. Let $R : \mathbb{C}_+ \rightarrow X$ be a function. Suppose that there exists $r \in L^1_{loc}(0, +\infty; X)$ such that (5.4) holds and $\mathcal{L}(r)(s) = R(s)$ for every $s \in \mathbb{C}_+$. In this case we say that r is the Inverse Laplace Transform of R , and we use the notation $r = \mathcal{L}^{-1}(R)$ or $r = \mathcal{L}_s^{-1}(R(s))$ with dummy variable s . It can be proven that r is uniquely determined up to a negligible set (see [2, Theorem 1.7.3]). Moreover, r can be obtained by the Bromwich Integral Formula:

$$r(t) = \mathcal{L}^{-1}(R)(t) = \frac{e^{s_1 t}}{2\pi} \lim_{k \rightarrow +\infty} \int_{-k}^k e^{is_2 t} R(s_1 + is_2) ds_2, \quad (5.7)$$

where s_1 is an arbitrary positive number. Clearly (5.7) can be expressed in terms of the Inverse Fourier Transform, namely

$$r(t) = \mathcal{L}_s^{-1}(R(s))(t) = e^{s_1 t} \mathcal{F}_{s_2}^{-1}(R(s_1 + is_2))(t), \quad (5.8)$$

where $\mathcal{F}_{s_2}^{-1}(R(s_1 + is_2))$ denotes the Inverse Fourier Transform with respect to the variable s_2 .

To use the Laplace Transform, we extend our problems from the interval $[0, T]$ to $[0, +\infty)$. To do this, we extend the functions h_ε and h , introduced in (5.3), by setting them equal to zero in $(T, +\infty)$, and we consider the solution to (5.1) in $[0, +\infty)$, which we still denote v_ε . Moreover, we consider the solution to (5.2) in $[0, +\infty)$, which we still denote v_0 . Notice that, thanks to the choice of the extension we have

$$h_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} h \quad \text{strongly } L^2(0, +\infty; H).$$

By Proposition 3.8 and by using the equality $h_\varepsilon = 0$ on $(T, +\infty)$, we get

$$v_\varepsilon \in L^\infty(0, +\infty; V_0) \quad \text{and} \quad \dot{v}_\varepsilon \in L^\infty(0, +\infty; H). \quad (5.9)$$

Since $h \in L^2(0, T; H)$ and $h = 0$ on $(T, +\infty)$, by means of standard estimates for the solution to (5.2) we obtain

$$v_0 \in L^2(0, +\infty; V_0) \quad \text{and} \quad v_0 = 0 \quad \text{on } (T, +\infty). \quad (5.10)$$

From (2.4), (5.1), and (5.9) we can deduce

$$\ddot{v}_\varepsilon \in L^2(0, T; V'_0) \cap L^\infty(T, +\infty; V'_0). \quad (5.11)$$

To study our problem by means of the Laplace Transform we introduce the complexification of the Hilbert spaces H , V_0 , and V'_0 defined by

$$\hat{H} := L^2(\Omega; \mathbb{C}^d), \quad \hat{V}_0 := H^1(\Omega; \mathbb{C}^d), \quad \hat{V}'_0 := H^{-1}(\Omega; \mathbb{C}^d).$$

The symbols (\cdot, \cdot) and $\|\cdot\|$ denote the hermitian product and the norm in \hat{H} or in other complex L^2 spaces. For every $s \in \mathbb{C}_+$ the Laplace Transforms $\hat{h}_\varepsilon(s)$ and $\hat{h}(s)$ of h_ε and h in \hat{H} are well defined. Thanks to (5.9) and (5.10) the Laplace Transforms $\hat{v}_\varepsilon(s)$ and $\hat{v}_0(s)$ in \hat{V}_0 are well defined for every $s \in \mathbb{C}_+$. By (5.11) the Laplace Transform $\hat{\dot{v}}_\varepsilon(s)$ of $\dot{v}_\varepsilon(s)$ in \hat{V}'_0 is well defined for every

$s \in \mathbb{C}_+$. Using (5.9) we can integrate by parts two times in the integral which defines \hat{v}_ε and, since $v_\varepsilon(0) = 0$ and $\dot{v}_\varepsilon(0) = 0$, we obtain

$$\hat{v}_\varepsilon(s) = s^2 \hat{v}_\varepsilon(s) \quad \text{for every } s \in \mathbb{C}_+. \quad (5.12)$$

By considering the operators $S_\mathbb{A}, S_\mathbb{B} : \hat{V}_0 \rightarrow \hat{V}'_0$ defined by

$$S_\mathbb{A}(\psi) := -\operatorname{div}(\mathbb{A}e\psi) \quad \text{and} \quad S_\mathbb{B}(\psi) := -\operatorname{div}(\mathbb{B}e\psi),$$

we can rephrase (5.1) and (5.2) for a.e. $t \in [0, +\infty)$ as equalities in \hat{V}'_0 :

$$\varepsilon^2 \ddot{v}_\varepsilon(t) = S_\mathbb{B} \left(\int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} v_\varepsilon(\tau) d\tau \right) - (S_\mathbb{A} + S_\mathbb{B})(v_\varepsilon(t)) + h_\varepsilon(t), \quad (5.13)$$

$$S_\mathbb{A}(v_0(t)) = h(t). \quad (5.14)$$

Now we want to consider the Laplace Transforms, in the sense of \hat{V}'_0 , of both sides of these equations. By Proposition 5.1 we can say

$$\mathcal{L}(S_\mathbb{A}(v_\varepsilon)) = S_\mathbb{A}(\hat{v}_\varepsilon), \quad \mathcal{L}(S_\mathbb{B}(v_\varepsilon)) = S_\mathbb{B}(\hat{v}_\varepsilon), \quad \mathcal{L}(S_\mathbb{A}(v_0)) = S_\mathbb{A}(\hat{v}_0), \quad (5.15)$$

where \hat{v}_ε and \hat{v}_0 are the Laplace Transforms of v_ε and v_0 , respectively, in the sense of \hat{V}_0 . Moreover, since we have

$$\sup_{t \in [0, +\infty)} \left\| \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} v_\varepsilon(\tau) d\tau \right\|_{V_0} \leq \|v_\varepsilon\|_{L^\infty(0, +\infty; V_0)},$$

this integral admits Laplace Transform in the sense of \hat{V}_0 , which for every $s \in \mathbb{C}_+$ satisfies

$$\mathcal{L}_t \left(\int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} v_\varepsilon(\tau) d\tau \right) (s) = \frac{1}{\beta\varepsilon s + 1} \hat{v}_\varepsilon(s).$$

Hence, by using Proposition 5.1 again, we obtain

$$\mathcal{L}_t \left(S_\mathbb{B} \left(\int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} v_\varepsilon(\tau) d\tau \right) \right) (s) = \frac{1}{\beta\varepsilon s + 1} S_\mathbb{B}(\hat{v}_\varepsilon(s)). \quad (5.16)$$

5.2. Properties of the Laplace Transform of the solutions

Thanks to (5.12), (5.15), and (5.16) we can apply the Laplace Transform to both sides of (5.13) and (5.14) to deduce for every $s \in \mathbb{C}_+$ the following equalities in \hat{V}'_0 :

$$\varepsilon^2 s^2 \hat{v}_\varepsilon(s) - \operatorname{div}((\mathbb{A} + \mathbb{B})e\hat{v}_\varepsilon(s)) + \frac{1}{\beta\varepsilon s + 1} \operatorname{div}(\mathbb{B}e\hat{v}_\varepsilon(s)) = \hat{h}_\varepsilon(s), \quad (5.17)$$

$$-\operatorname{div}(\mathbb{A}e\hat{v}_0(s)) = \hat{h}(s). \quad (5.18)$$

Our purpose is to prove that for every $s_1 > 0$ we have

$$\int_{-\infty}^{+\infty} \|\hat{v}_\varepsilon(s_1 + is_2) - \hat{v}_0(s_1 + is_2)\|_{V_0}^2 ds_2 \xrightarrow{\varepsilon \rightarrow 0^+} 0. \quad (5.19)$$

To prove (5.19) we need two lemmas. In the first one we deduce from (5.17) an estimate for $\hat{v}_\varepsilon(s)$, which is used in the second lemma to prove a convergence result for $\hat{v}_\varepsilon(s)$.

Lemma 5.2. *For every $s \in \mathbb{C}_+$ there exists a positive constant $M(s)$ such that*

$$\|\hat{v}_\varepsilon(s)\|_{\hat{V}_0} \leq M(s) \|\hat{h}_\varepsilon(s)\| \quad \text{for every } \varepsilon \in (0, 1). \quad (5.20)$$

Proof. We fix $\varepsilon \in (0, 1)$ and for every $s \in \mathbb{C}_+$ we define the operator $S_\varepsilon(s): \hat{V}_0 \rightarrow \hat{V}'_0$ in the following way:

$$S_\varepsilon(s)(\psi) := \varepsilon^2 s^2 \psi - \operatorname{div}((\mathbb{A} + \mathbb{B})e\psi) + \frac{1}{\beta\varepsilon s + 1} \operatorname{div}(\mathbb{B}e\psi) \quad \text{for every } \psi \in \hat{V}_0.$$

Since $S_\varepsilon(s)(\hat{v}_\varepsilon(s)) = \hat{h}_\varepsilon(s)$ by (5.17), the Lax-Milgram Lemma, together with the Korn-Poincaré Inequality (2.3), implies (5.20) if we can show that for every $s \in \mathbb{C}_+$ there exists a positive constant $K(s)$, independent of ε , such that

$$|\langle S_\varepsilon(s)(\psi), \psi \rangle| \geq c_\mathbb{A} K(s) \|e\psi\|^2 \quad \text{for every } \psi \in \hat{V}_0, \quad (5.21)$$

where

$$|\langle S_\varepsilon(s)(\psi), \psi \rangle| = \frac{|(\beta\varepsilon^3 s^3 + \varepsilon^2 s^2) \|\psi\|^2 + \beta\varepsilon s((\mathbb{A} + \mathbb{B})e\psi, e\psi) + (\mathbb{A}e\psi, e\psi)|}{|\beta\varepsilon s + 1|}.$$

We can suppose $\psi \in \hat{V}_0 \setminus \{0\}$, otherwise the inequality is trivially satisfied, and we set

$$a := \frac{(\mathbb{A}e\psi, e\psi)}{\|\psi\|^2} \quad \text{and} \quad b := \frac{((\mathbb{A} + \mathbb{B})e\psi, e\psi)}{\|\psi\|^2},$$

which satisfy, thanks to the Korn-Poincaré Inequality (2.3) and to (2.4)–(2.6), the following relations

$$a \geq \frac{c_\mathbb{A} \|e\psi\|^2}{\|\psi\|^2} \geq \frac{c_\mathbb{A}}{C_P^2} =: a_0, \quad b \geq \frac{(c_\mathbb{A} + c_\mathbb{B}) \|e\psi\|^2}{\|\psi\|^2} \geq \frac{c_\mathbb{A} + c_\mathbb{B}}{C_P^2} =: b_0, \quad (5.22)$$

$$a \leq c_0 a \leq b \leq c_1 a,$$

where $c_0 := 1 + \frac{c_\mathbb{B}}{C_\mathbb{A}}$ and $c_1 := 1 + \frac{C_\mathbb{B}}{C_\mathbb{A}}$. Therefore, to prove (5.21) it is enough to obtain

$$\left| \frac{\beta\varepsilon^3 s^3 + \varepsilon^2 s^2 + \beta b \varepsilon s + a}{\beta\varepsilon s + 1} \right| \geq K(s) a \quad \text{for every } s \in \mathbb{C}_+. \quad (5.23)$$

For simplicity of notation we set $z = \varepsilon s$ and we consider two cases.

Case $b > \frac{2}{3\beta^2}$. In this situation, thanks to (A.2) and (A.3) we know that the polynomial $\beta z^3 + z^2 + \beta b z + a$ has one real root z_0 and two complex and conjugate ones w and \bar{w} . Therefore, thanks to Lemmas A.1 and A.2, we can write

$$\begin{aligned} \left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| &= \left| \frac{\beta(z - z_0)(z - w)(z - \bar{w})}{\beta z + 1} \right| \\ &\geq \left| \frac{\beta(z - z_0)}{\beta z + 1} \right| |\Re(w)| |\Im(w)| \\ &= \left| \frac{\beta(z - z_0)}{\beta z + 1} \right| |\Re(w)| \sqrt{3|\Re(w)|^2 + \frac{2}{\beta} \Re(w) + b} \end{aligned}$$

$$\begin{aligned}
&\geq \left| \frac{\beta(z - z_0)}{\beta z + 1} \right| \alpha \sqrt{b - \frac{1}{3\beta^2}} \\
&\geq \left| \frac{\beta(z - z_0)}{\beta z + 1} \right| \alpha \sqrt{\frac{b}{2}} \geq \frac{\alpha}{\sqrt{3}} \left| \frac{z}{\beta z + 1} \right|, \tag{5.24}
\end{aligned}$$

where in the last inequality we used $z_0 < 0$.

If $a \leq 2|z|^2$, then $|z| \geq \frac{a}{2|z|}$ and, thanks to (5.24), we deduce

$$\left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| \geq \frac{\alpha}{2\sqrt{3}} \frac{a}{|z(\beta z + 1)|}. \tag{5.25}$$

For $a > 2|z|^2$ we have

$$\frac{1}{a} \left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| = \left| \frac{z^2}{a} + \frac{\beta b z + a}{a(\beta z + 1)} \right| \geq \left| \frac{\beta b z + a}{a(\beta z + 1)} \right| - \frac{1}{2},$$

and, by writing $z = x + iy$, we obtain

$$\left| \frac{\beta b z + a}{a(\beta z + 1)} \right| = \left| \frac{\beta b x + a + i\beta b y}{\beta a x + a + i\beta a y} \right| = \sqrt{\frac{(\beta b x + a)^2 + \beta^2 b^2 y^2}{(\beta a x + a)^2 + \beta^2 a^2 y^2}} \geq 1,$$

which implies

$$\left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| \geq \frac{a}{2}. \tag{5.26}$$

By (5.25) and (5.26) in the case $b > \frac{2}{3\beta^2}$ we conclude

$$\left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| \geq \min \left\{ \frac{1}{2}, \frac{\alpha}{2\sqrt{3}} \frac{1}{|z(\beta z + 1)|} \right\} a. \tag{5.27}$$

Case $b_0 \leq b \leq \frac{2}{3\beta^2}$. In this case, thanks to (5.22), we have $a_0 \leq a \leq \frac{2}{3\beta^2}$. We define

$$R := \sqrt{\frac{2(2 + c_1)}{3\beta^2}}.$$

Then for $z \in \mathbb{C}_+$, with $|z| > R$, we get

$$\begin{aligned}
\frac{1}{a} \left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| &= \left| \frac{z^2}{a} + \frac{\beta b z + a}{a(\beta z + 1)} \right| \\
&\geq \frac{3\beta^2 |z|^2}{2} - \frac{b}{a} \left| \frac{\beta z}{\beta z + 1} \right| - \frac{1}{|\beta z + 1|} \geq 1, \tag{5.28}
\end{aligned}$$

where we used the inequalities $|\beta z| \leq |\beta z + 1|$ and $1 \leq |\beta z + 1|$.

To deal with the case $z \in \mathbb{C}_+$, with $|z| \leq R$, we define

$$I := \{a, b \in \mathbb{R} : b_0 \leq b \leq \frac{2}{3\beta^2}, \quad a_0 \leq a \leq \frac{2}{3\beta^2}\},$$

$$\gamma := \min \left\{ \left| \frac{\beta z^3 + z^2 + \beta b z + a}{a(\beta z + 1)} \right| : \Re(z) \geq 0, \quad |z| \leq R, \quad a, b \in I \right\},$$

and we claim $\gamma > 0$. Indeed the function under examination is continuous with respect to (z, a, b) , and by Lemma A.1 it does not vanish in the compact

set considered in the minimum problem. By using also (5.28) we conclude that for $b_0 \leq b \leq \frac{2}{3\beta^2}$ we have

$$\left| \frac{\beta z^3 + z^2 + \beta b z + a}{\beta z + 1} \right| \geq \min\{\gamma, 1\}a. \quad (5.29)$$

for every $z \in \mathbb{C}_+$ and every a satisfying (5.22). Since $\varepsilon \in (0, 1)$ we have

$$\frac{1}{|\varepsilon s(\beta \varepsilon s + 1)|} \geq \frac{1}{|s(\beta s + 1)|},$$

therefore, by setting

$$K(s) := \min \left\{ \frac{1}{2}, \frac{\alpha}{2\sqrt{3}} \frac{1}{|s(\beta s + 1)|}, \gamma \right\},$$

from (5.27) and (5.29) we obtain (5.23), which concludes the proof. \square

5.3. Convergence of the Laplace Transform of the solutions

We begin by proving the pointwise convergence.

Lemma 5.3. *For every $s \in \mathbb{C}_+$ we have*

$$\hat{v}_\varepsilon(s) \xrightarrow{\varepsilon \rightarrow 0^+} \hat{v}_0(s) \quad \text{strongly in } \hat{V}_0.$$

Proof. Thanks to (5.3) and to the Hölder Inequality for every $s \in \mathbb{C}_+$ we get

$$\begin{aligned} \|\hat{h}_\varepsilon(s) - \hat{h}(s)\| &\leq \int_0^{+\infty} e^{-\Re(s)t} \|h_\varepsilon(t) - h(t)\| dt \\ &\leq \frac{1}{\sqrt{2\Re(s)}} \|h_\varepsilon - h\|_{L^2(0,T;H)} \xrightarrow{\varepsilon \rightarrow 0^+} 0. \end{aligned} \quad (5.30)$$

Consequently, thanks to Lemma 5.2, for every $s \in \mathbb{C}_+$ there exist two constants $\bar{M}(s) > 0$ and $\varepsilon(s) \in (0, 1)$ such that

$$\|\hat{v}_\varepsilon(s)\|_{\hat{V}_0} \leq \bar{M}(s) \quad \text{for every } \varepsilon \in (0, \varepsilon(s)). \quad (5.31)$$

By (5.31) we can say that for every $s \in \mathbb{C}_+$ there exist a sequence $\varepsilon_j \rightarrow 0^+$ and $v^*(s) \in \hat{V}_0$ such that

$$\hat{v}_{\varepsilon_j}(s) \xrightarrow{j \rightarrow +\infty} v^*(s) \quad \text{weakly in } \hat{V}_0. \quad (5.32)$$

Thanks to (2.5) and (5.32) for every $\psi \in \hat{V}_0$ we deduce

$$\begin{aligned} ((\mathbb{A} + \mathbb{B})e\hat{v}_{\varepsilon_j}(s), e\psi) &\xrightarrow{j \rightarrow +\infty} ((\mathbb{A} + \mathbb{B})ev^*(s), e\psi), \\ |\varepsilon_j^2 s^2 (\hat{v}_{\varepsilon_j}(s), \psi)| &\leq \varepsilon_j^2 |s|^2 \bar{M}(s) \|\psi\| \xrightarrow{j \rightarrow +\infty} 0, \\ \left| \frac{1}{\beta \varepsilon_j s + 1} (\mathbb{B}e\hat{v}_{\varepsilon_j}(s), e\psi) - (\mathbb{B}ev^*(s), e\psi) \right| \\ &\leq |(\mathbb{B}(e\hat{v}_{\varepsilon_j}(s) - ev^*(s)), e\psi)| + \frac{\beta \varepsilon_j |s|}{|\beta \varepsilon_j s + 1|} |(\mathbb{B}e\hat{v}_{\varepsilon_j}(s), e\psi)| \\ &\leq |(e\hat{v}_{\varepsilon_j}(s) - ev^*(s), \mathbb{B}e\psi)| + \beta \varepsilon_j |s| C_{\mathbb{B}} \bar{M}(s) \|e\psi\| \xrightarrow{j \rightarrow +\infty} 0. \end{aligned}$$

Therefore by (5.30) we have

$$\begin{cases} v^*(s) \in \hat{V}_0, \\ -\operatorname{div}(\mathbb{A}ev^*(s)) = \hat{h}(s). \end{cases} \quad (5.33)$$

Since, by (5.18), $\hat{v}_0(s)$ is a solution to (5.33), by uniqueness $v^*(s) = \hat{v}_0(s)$. Moreover, since the limit does not depend on the subsequence, the whole sequence satisfies

$$\hat{v}_\varepsilon(s) \xrightarrow{\varepsilon \rightarrow 0^+} \hat{v}_0(s) \quad \text{weakly in } \hat{V}_0 \text{ for every } s \in \mathbb{C}_+. \quad (5.34)$$

To prove the strong convergence we use $\hat{v}_\varepsilon(s)$ and $\hat{v}_0(s)$ as test function in (5.17) and (5.18), respectively. By subtracting the two equalities, we obtain

$$\begin{aligned} & (\mathbb{A}e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s)) - (\mathbb{A}e\hat{v}_0(s), e\hat{v}_0(s)) \\ &= (\hat{h}_\varepsilon(s), \hat{v}_\varepsilon(s)) - (\hat{h}(s), \hat{v}_0(s)) - \varepsilon^2 s^2 \|\hat{v}_\varepsilon(s)\|^2 - \frac{\beta \varepsilon s}{\beta \varepsilon s + 1} (\mathbb{B}e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s)), \end{aligned}$$

from which we deduce

$$\begin{aligned} & |(\mathbb{A}e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s)) - (\mathbb{A}e\hat{v}_0(s), e\hat{v}_0(s))| \\ & \leq |(\hat{h}_\varepsilon(s), \hat{v}_\varepsilon(s)) - (\hat{h}(s), \hat{v}_0(s))| + \varepsilon^2 |s|^2 \|\hat{v}_\varepsilon(s)\|^2 + \beta \varepsilon |s| C_{\mathbb{B}} \|e\hat{v}_\varepsilon(s)\|^2. \end{aligned}$$

By using again (5.30), (5.31), and (5.34), we can deduce

$$\lim_{\varepsilon \rightarrow 0^+} (\mathbb{A}e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s)) = (\mathbb{A}e\hat{v}_0(s), e\hat{v}_0(s)) \quad \text{for every } s \in \mathbb{C}_+. \quad (5.35)$$

Thanks to the coerciveness assumption (2.6), the conclusion follows from the weak convergence (5.34) together with (5.35). \square

Now we are in a position to prove the following result about the convergence in the space L^2 on the lines $\{s_1 + is_2 : s_2 \in \mathbb{R}\}$.

Proposition 5.4. *The functions \hat{v}_ε and \hat{v}_0 satisfy (5.19).*

Proof. For every $s \in \mathbb{C}_+$, by using $\hat{v}_\varepsilon(s)$ as test function in (5.17) we obtain

$$\begin{aligned} & \left(\beta \varepsilon^3 s^3 + \varepsilon^2 s^2 + \beta \frac{((\mathbb{A} + \mathbb{B})e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s))}{\|\hat{v}_\varepsilon(s)\|^2} \varepsilon s + \frac{(\mathbb{A}e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s))}{\|\hat{v}_\varepsilon(s)\|^2} \right) \|\hat{v}_\varepsilon(s)\|^2 \\ &= (\beta \varepsilon s + 1)(\hat{h}_\varepsilon(s), \hat{v}_\varepsilon(s)). \end{aligned} \quad (5.36)$$

As before, we set

$$a := \frac{(\mathbb{A}e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s))}{\|\hat{v}_\varepsilon(s)\|^2} \quad \text{and} \quad b := \frac{((\mathbb{A} + \mathbb{B})e\hat{v}_\varepsilon(s), e\hat{v}_\varepsilon(s))}{\|\hat{v}_\varepsilon(s)\|^2}, \quad (5.37)$$

and we observe that (5.22) holds. Therefore, thanks to (5.24), Lemma A.1, and (5.29) we can deduce

$$\begin{aligned} \left| \frac{\beta \varepsilon^3 s^3 + \varepsilon^2 s^2 + \beta b \varepsilon s + a}{\beta \varepsilon s + 1} \right| &\geq \left| \frac{\beta(\varepsilon s - z_0)}{\beta \varepsilon s + 1} \right| \alpha \sqrt{\frac{b}{2}} \\ &\geq \beta |z_0| \alpha \sqrt{\frac{a}{2}} \geq \frac{\beta \alpha^2}{\sqrt{2}} \sqrt{a} \quad \text{for } b > \frac{2}{3\beta^2}, \end{aligned}$$

$$\left| \frac{\beta \varepsilon^3 s^3 + \varepsilon^2 s^2 + \beta b \varepsilon s + a}{\beta \varepsilon s + 1} \right| \geq \min\{\gamma, 1\} a \geq \min\{\gamma, 1\} \sqrt{a_0} \sqrt{a} \quad \text{for } b \leq \frac{2}{3\beta^2},$$

where in the first line we used the inequality $|z_0(\beta \varepsilon s + 1)| \leq |\varepsilon s - z_0|$ for every $s \in \mathbb{C}_+$, which follows from the condition $z_0 < 0$.

As a consequence of these inequalities and of (5.36) there exists a positive constant $C = C(\alpha, \beta, \gamma, a_0)$ such that for every $s \in \mathbb{C}_+$ we have

$$\|\hat{v}_\varepsilon(s)\|^2 = \left| \frac{\beta \varepsilon s + 1}{\beta \varepsilon^3 s^3 + \varepsilon^2 s^2 + \beta b \varepsilon s + a} \right| |(\hat{h}_\varepsilon(s), \hat{v}_\varepsilon(s))| \leq \frac{C}{\sqrt{a}} \|\hat{h}_\varepsilon(s)\| \|\hat{v}_\varepsilon(s)\|.$$

Therefore, by using (5.37) and the coerciveness assumption (2.6), we can write

$$\sqrt{c_A} \|e \hat{v}_\varepsilon(s)\| \leq \sqrt{(\mathbb{A} e \hat{v}_\varepsilon(s), e \hat{v}_\varepsilon(s))} \leq C \|\hat{h}_\varepsilon(s)\|,$$

from which, recalling (2.3), we deduce

$$\|\hat{v}_\varepsilon(s)\|_{\hat{V}_0} \leq (C_P + 1) \frac{C}{\sqrt{c_A}} \|\hat{h}_\varepsilon(s)\| \quad \text{for every } s \in \mathbb{C}_+. \quad (5.38)$$

By extending the function h_ε to $(-\infty, 0)$ with value 0, we can write

$$\hat{h}_\varepsilon(s) = \int_0^{+\infty} e^{-st} h_\varepsilon(t) dt = \int_{-\infty}^{+\infty} e^{-st} h_\varepsilon(t) dt = \mathcal{F}_t(e^{-s_1 t} h_\varepsilon(t))(s_2).$$

Since for every $s = s_1 + i s_2 \in \mathbb{C}_+$ the function $t \mapsto e^{-s_1 t} h_\varepsilon(t)$ belongs to $L^2(\mathbb{R}; H)$, by the properties of the Fourier Transform we deduce that $s_2 \mapsto \hat{h}_\varepsilon(s_1 + i s_2)$ belongs to $L^2(\mathbb{R}; \hat{H})$ for every $\varepsilon > 0$. Moreover, by using (5.3) and the Plancherel Theorem, we can write

$$\begin{aligned} \int_{-\infty}^{+\infty} \|\hat{h}_\varepsilon(s_1 + i s_2) - \hat{h}(s_1 + i s_2)\|^2 ds_2 \\ &= \int_{-\infty}^{+\infty} \|\mathcal{F}_t(e^{-s_1 t} (h_\varepsilon(t) - h(t)))(s_2)\|^2 ds_2 \\ &= \int_{-\infty}^{+\infty} \|e^{-s_1 t} (h_\varepsilon(t) - h(t))\|^2 dt \\ &\leq \int_0^T \|h_\varepsilon(t) - h(t)\|^2 dt \xrightarrow{\varepsilon \rightarrow 0^+} 0. \end{aligned} \quad (5.39)$$

Since $\hat{v}_\varepsilon(s) \rightarrow \hat{v}_0(s)$ strongly in \hat{V}_0 by Lemma 5.3 and $\hat{h}_\varepsilon(s) \rightarrow \hat{h}(s)$ strongly in \hat{H} by (5.30), thanks to (5.38) and (5.39) we can apply the Generalized Dominated Convergence Theorem to get the conclusion. \square

6. L^2 convergence

In this section we shall prove (3.4) and (3.5) under the assumptions of Theorems 3.6 and 3.7. We begin by proving the following partial result.

Proposition 6.1. *Let $\{h_\varepsilon\}_\varepsilon \subset L^2(0, T; H)$ and $h \in L^2(0, T; H)$ be such that (5.3) holds. Let v_ε and v_0 be the solutions to problems (5.1) and (5.2). Then*

$$v_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} v_0 \quad \text{strongly in } L^2(0, T; V).$$

Proof. By the Plancherel Theorem we deduce from (5.8) and Proposition 5.4 that for every $s_1 > 0$

$$\begin{aligned} \|v_\varepsilon - v_0\|_{L^2(0, T; V)}^2 &= \int_0^T \|v_\varepsilon(t) - v_0(t)\|_V^2 dt = \int_0^T \|\mathcal{L}^{-1}(\hat{v}_\varepsilon - \hat{v}_0)(t)\|_V^2 dt \\ &\leq e^{s_1 T} \int_{-\infty}^{+\infty} \|\mathcal{F}_{s_2}^{-1}(\hat{v}_\varepsilon(s_1 + is_2) - \hat{v}_0(s_1 + is_2))(t)\|_V^2 ds_2 \\ &= e^{s_1 T} \int_{-\infty}^{+\infty} \|\hat{v}_\varepsilon(s_1 + is_2) - \hat{v}_0(s_1 + is_2)\|_{V_0}^2 ds_2 \xrightarrow{\varepsilon \rightarrow 0^+} 0, \end{aligned}$$

which concludes the proof. \square

Theorem 6.2. *Let us assume (H1), (H2), and (3.10). Let u_ε be the solution to the viscoelastic dynamic system (2.10), with $\varphi_\varepsilon = f_\varepsilon$ and $\gamma_\varepsilon = g_\varepsilon$, and let u_0 be the solution to the stationary problem (3.1). Then (3.4) holds.*

Proof. Thanks to Lemma 3.10 it is enough to prove the theorem in the case $z = 0$ and $z_\varepsilon = 0$ for every $\varepsilon > 0$. We divide the proof into two steps.

Step 1. The case $u_\varepsilon^1 = 0$. We reduce the problem to the case of homogeneous initial conditions by considering the functions

$$v_\varepsilon(t) := u_\varepsilon(t) - u_\varepsilon^0 \quad \text{and} \quad v_0(t) := u_0(t) - u^0 \quad \text{for a.e. } t \in [0, T]. \quad (6.1)$$

Let us define

$$q_\varepsilon(t) := g_\varepsilon(t) + \operatorname{div}(\mathbb{A}eu_\varepsilon^0) + e^{-\frac{t}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}eu_\varepsilon^0) \quad \text{for every } t \in [0, T], \quad (6.2)$$

$$q(t) := g(t) + \operatorname{div}(\mathbb{A}eu^0) \quad \text{for every } t \in [0, T]. \quad (6.3)$$

Since $u_\varepsilon^1 = 0$, it is easy to see that v_ε satisfies (2.13) with $h_\varepsilon = f_\varepsilon$, $\ell_\varepsilon = q_\varepsilon$, $v_\varepsilon^0 = 0$, and $v_\varepsilon^1 = 0$, while v_0 satisfies (3.2) with $h = f$ and $\ell = q$. By (3.10) and (6.1), to prove (3.4) it is enough to show that

$$v_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} v_0 \quad \text{strongly in } L^2(0, T; V). \quad (6.4)$$

In order to apply Proposition 6.1, we approximate the forcing terms of the problems for v_ε and v_0 by means of functions in $H^1(0, T; H)$ and we consider the corresponding solutions \tilde{v}_ε and \tilde{v}_0 , for which Proposition 6.1 yields $\tilde{v}_\varepsilon \rightarrow \tilde{v}_0$ strongly in $L^2(0, T; V)$ as $\varepsilon \rightarrow 0^+$. Finally we show that $\|\tilde{v}_\varepsilon - v_\varepsilon\|_{L^2(0, T; V)}$ and $\|\tilde{v}_0 - v_0\|_{L^2(0, T; V)}$ are small uniformly with respect to ε , and this leads to the proof of (6.4).

Let us fix $\delta > 0$. Thanks to the density of H in V'_0 and to Lemma 4.2 we can find $\psi \in H^1(0, T; H)$ and $h_\mathbb{A}^0, h_\mathbb{B}^0 \in H$ such that

$$\|\psi - g\|_{W^{1,1}(0, T; V'_0)} < \delta, \quad (6.5)$$

$$\|h_\mathbb{A}^0 - \operatorname{div}(\mathbb{A}eu^0)\|_{V'_0} < \delta, \quad \|h_\mathbb{B}^0 - \operatorname{div}(\mathbb{B}eu^0)\|_{V'_0} < \delta. \quad (6.6)$$

Thanks to (H1) and (3.10) there exist $\varepsilon_0 = \varepsilon_0(\delta) \in (0, \frac{1}{\beta})$ such that for every $\varepsilon \in (0, \varepsilon_0)$ we have

$$\|\psi - g_\varepsilon\|_{W^{1,1}(0,T;V'_0)} < \delta, \quad (6.7)$$

$$\|h_{\mathbb{A}}^0 - \operatorname{div}(\mathbb{A}eu_\varepsilon^0)\|_{V'_0} < \delta, \quad \|h_{\mathbb{B}}^0 - \operatorname{div}(\mathbb{B}eu_\varepsilon^0)\|_{V'_0} < \delta. \quad (6.8)$$

Let $\varphi_\varepsilon, \varphi: [0, T] \rightarrow H$ be defined for every $t \in [0, T]$ by

$$\varphi_\varepsilon(t) := \psi(t) + h_{\mathbb{A}}^0 + e^{-\frac{t}{\beta\varepsilon}} h_{\mathbb{B}}^0 \quad \text{and} \quad \varphi(t) := \psi(t) + h_{\mathbb{A}}^0. \quad (6.9)$$

By (6.2), (6.3), (6.5), (6.6), (6.7), (6.8), and (6.9) for every $\varepsilon \in (0, \varepsilon_0)$ we obtain

$$\begin{aligned} \|\varphi_\varepsilon - q_\varepsilon\|_{W^{1,1}(0,T;V'_0)} &\leq \|\psi - g_\varepsilon\|_{W^{1,1}(0,T;V'_0)} + T\|h_{\mathbb{A}}^0 - \operatorname{div}(\mathbb{A}eu_\varepsilon^0)\|_{V'_0} \\ &\quad + (\beta\varepsilon + 1)\|h_{\mathbb{B}}^0 - \operatorname{div}(\mathbb{B}eu_\varepsilon^0)\|_{V'_0} \leq (3 + T)\delta, \end{aligned} \quad (6.10)$$

$$\begin{aligned} \|\varphi - q\|_{L^\infty(0,T;V'_0)} &\leq (1 + \frac{1}{T})\|\psi - g\|_{W^{1,1}(0,T;V'_0)} + \|h_{\mathbb{A}}^0 - \operatorname{div}(\mathbb{A}eu_\varepsilon^0)\|_{V'_0} \\ &\leq (2 + \frac{1}{T})\delta. \end{aligned} \quad (6.11)$$

Since $t \mapsto e^{-\frac{t}{\beta\varepsilon}} \psi_{\mathbb{B}}^0$ converges to 0 strongly in $L^2(0, T; H)$ as $\varepsilon \rightarrow 0^+$, by (6.9) we have

$$\varphi_\varepsilon \xrightarrow[\varepsilon \rightarrow 0^+]{} \varphi \quad \text{strongly in } L^2(0, T; H). \quad (6.12)$$

Let \tilde{v}_ε be the solution to (5.1) with $h_\varepsilon = f_\varepsilon + \varphi_\varepsilon$ and let \tilde{v}_0 be the solution to (5.2) with $h = f + \varphi$. By (H1) and (6.12) we have

$$f_\varepsilon + \varphi_\varepsilon \xrightarrow[\varepsilon \rightarrow 0^+]{} f + \varphi \quad \text{strongly in } L^2(0, T; H),$$

hence Proposition 6.1 yields

$$\tilde{v}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0^+]{} \tilde{v}_0 \quad \text{strongly in } L^2(0, T; V). \quad (6.13)$$

To estimate the difference $\tilde{v}_\varepsilon - v_\varepsilon$ we observe that it solves (2.13) with $h_\varepsilon = 0$, $\ell_\varepsilon = \varphi_\varepsilon - q_\varepsilon$, $v_\varepsilon^0 = 0$, and $v_\varepsilon^1 = 0$. Therefore, by Lemma 3.8 we have

$$\|\tilde{v}_\varepsilon - v_\varepsilon\|_{L^2(0,T;V)} \leq \sqrt{C_E T} \|\varphi_\varepsilon - q_\varepsilon\|_{W^{1,1}(0,T;V'_0)}. \quad (6.14)$$

To estimate the difference $\tilde{v}_0 - v_0$ we observe that it solves (3.2) with $h = 0$ and $\ell = \varphi - q$. Therefore by the Lax-Milgram Lemma we obtain

$$\|\tilde{v}_0 - v_0\|_{L^2(0,T;V)} \leq \frac{\sqrt{T}(C_P^2 + 1)}{c_A} \|\varphi - q\|_{L^\infty(0,T;V'_0)}. \quad (6.15)$$

By (6.10), (6.11), (6.14), and (6.15) there exists a positive constant $C = C(\mathbb{A}, \mathbb{B}, \Omega, T)$ such that

$$\|\tilde{v}_\varepsilon - v_\varepsilon\|_{L^2(0,T;V)} + \|\tilde{v}_0 - v_0\|_{L^2(0,T;V)} \leq C\delta,$$

hence

$$\begin{aligned} \|v_\varepsilon - v_0\|_{L^2(0,T;V)} &\leq \|v_\varepsilon - \tilde{v}_\varepsilon\|_{L^2(0,T;V)} + \|\tilde{v}_\varepsilon - \tilde{v}_0\|_{L^2(0,T;V)} + \|\tilde{v}_0 - v_0\|_{L^2(0,T;V)} \\ &\leq \|\tilde{v}_\varepsilon - \tilde{v}_0\|_{L^2(0,T;V)} + C\delta. \end{aligned}$$

This inequality, together with (6.13), gives

$$\limsup_{\varepsilon \rightarrow 0^+} \|v_\varepsilon - v_0\|_{L^2(0,T;V)} \leq C\delta.$$

By the arbitrariness of $\delta > 0$ we obtain (6.4), which concludes the proof of Step 1.

Step 2. The general case. Let \tilde{u}_ε be the solution to (2.13) with $h_\varepsilon = f_\varepsilon$, $\ell_\varepsilon = g_\varepsilon$, $v_\varepsilon^0 = u_\varepsilon^0$, and $v_\varepsilon^1 = 0$. By Step 1

$$\tilde{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} u_0 \quad \text{strongly in } L^2(0,T;V). \quad (6.16)$$

The function $u_\varepsilon - \tilde{u}_\varepsilon$ is the solution to (2.13) with all data equal to 0 except v_ε^1 , which is now equal to u_ε^1 . Therefore, Lemma 3.8 and (3.10) yield

$$\|u_\varepsilon - \tilde{u}_\varepsilon\|_{L^\infty(0,T;V)} \leq \varepsilon \sqrt{C_E} \|u_\varepsilon^1\| \xrightarrow{\varepsilon \rightarrow 0^+} 0,$$

which, together with (6.16), gives (3.4). \square

In the following theorem, under the assumptions of Theorem 3.7 we deduce (3.5) from (3.4).

Theorem 6.3. *Let us assume (H1), (H2), and (3.10). Let u_ε be the solution to the viscoelastic dynamic system (2.10), with $\varphi_\varepsilon = f_\varepsilon$ and $\gamma_\varepsilon = g_\varepsilon$, and let u_0 be the solution to the stationary problem (3.1). Then (3.5) holds.*

Proof. Thanks to Lemma 3.10 we can suppose $z = 0$ and $z_\varepsilon = 0$ for every $\varepsilon > 0$. It is convenient to extend the data of our problem to the interval $[0, 2T]$ by setting for every $t \in (T, 2T]$

$$f_\varepsilon(t) := 0, \quad f(t) := 0, \quad g_\varepsilon(t) := g_\varepsilon(T), \quad g(t) := g(T).$$

Since (H1) holds, it is clear that $\{f_\varepsilon\}_\varepsilon \subset L^2(0, 2T; H)$, $\{g_\varepsilon\}_\varepsilon \in H^1(0, 2T; V'_0)$,

$$f_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} f \quad \text{strongly in } L^2(0, 2T; H) \quad (6.17)$$

$$g_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} g \quad \text{strongly in } W^{1,1}(0, 2T; V'_0). \quad (6.18)$$

Moreover, the solution to (2.10) on $[0, 2T]$ with the extended data is an extension of u_ε , which is still denoted by u_ε . Similarly, the solution to (3.1) on $[0, 2T]$ is still denoted by u_0 . Since (6.17) and (6.18) hold, Theorem 6.2 gives

$$u_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} u_0 \quad \text{strongly in } L^2(0, 2T; V). \quad (6.19)$$

We further extend u_ε to \mathbb{R} by setting $u_\varepsilon(t) = 0$ for every $t \in \mathbb{R} \setminus [0, 2T]$, and we define

$$w_\varepsilon(t) := \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} eu_\varepsilon(\tau) d\tau = (\rho_\varepsilon * eu_\varepsilon)(t) \quad \text{for every } t \in \mathbb{R},$$

where ρ_ε is as in (3.17). By the properties of convolutions and (6.19) we get

$$eu_\varepsilon - w_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(\mathbb{R}; \tilde{H}). \quad (6.20)$$

Thanks to (6.19) and (6.20), by using (2.10) and (3.1) we obtain

$$\varepsilon^2 \ddot{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, 2T; V'_0). \quad (6.21)$$

Since

$$\varepsilon^2 \dot{u}_\varepsilon(t) = \varepsilon^2 u_\varepsilon^1 + \varepsilon^2 \int_0^t \ddot{u}_\varepsilon(\tau) d\tau \quad \text{for every } t \in [0, 2T],$$

(3.10) and (6.21) imply

$$\varepsilon^2 \dot{u}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, 2T; V'_0). \quad (6.22)$$

By (6.19) and (6.22) there exists a sequence $\varepsilon_j \rightarrow 0^+$ such that for a.e. $t \in [0, 2T]$ we have

$$u_{\varepsilon_j}(t) \xrightarrow{j \rightarrow +\infty} u_0(t) \quad \text{strongly in } V, \quad (6.23)$$

$$\varepsilon_j^2 \dot{u}_{\varepsilon_j}(t) \xrightarrow{j \rightarrow +\infty} 0 \quad \text{strongly in } V'_0. \quad (6.24)$$

We choose $T_0 \in (T, 2T)$ such that (6.23) and (6.24) hold at $t = T_0$. This implies

$$\varepsilon_j^2 (\dot{u}_{\varepsilon_j}(T_0), u_{\varepsilon_j}(T_0)) = \langle \varepsilon_j^2 \dot{u}_{\varepsilon_j}(T_0), u_{\varepsilon_j}(T_0) \rangle \xrightarrow{j \rightarrow +\infty} 0. \quad (6.25)$$

Since $z_\varepsilon = 0$ for a.e. $t \in [0, T_0]$ we can use $u_\varepsilon(t) \in V_0$ as test function in (2.10). Then we integrate by parts in time on the interval $(0, T_0)$ to obtain

$$\begin{aligned} & -\varepsilon_j^2 \int_0^{T_0} \|\dot{u}_{\varepsilon_j}(t)\|^2 dt + \int_0^{T_0} (\mathbb{A} e u_{\varepsilon_j}(t), e u_{\varepsilon_j}(t)) dt \\ & + \int_0^{T_0} (\mathbb{B} (e u_{\varepsilon_j}(t) - w_{\varepsilon_j}(t)), e u_{\varepsilon_j}(t)) dt = \int_0^{T_0} (f_{\varepsilon_j}(t), u_{\varepsilon_j}(t)) dt \\ & + \int_0^{T_0} \langle g_{\varepsilon_j}(t), u_{\varepsilon_j}(t) \rangle dt - \varepsilon_j^2 (\dot{u}_{\varepsilon_j}(T_0), u_{\varepsilon_j}(T_0)) + \varepsilon_j^2 (u_{\varepsilon_j}^0, u_{\varepsilon_j}^1). \end{aligned}$$

Thanks to (3.1), (3.10), (6.17), (6.19), (6.20), and (6.25) the first term on the left-hand side of the previous equation tends to 0 as $j \rightarrow +\infty$. Since $T_0 > T$ we have

$$\varepsilon_j^2 \int_0^T \|\dot{u}_{\varepsilon_j}(t)\|^2 dt \xrightarrow{j \rightarrow +\infty} 0.$$

By the arbitrariness of the sequence $\{\varepsilon_j\}_j$ we have

$$\varepsilon^2 \int_0^T \|\dot{u}_\varepsilon(t)\|^2 dt \xrightarrow{\varepsilon \rightarrow 0^+} 0,$$

which concludes the proof. \square

We now use Theorems 6.2 and 6.3 to obtain (3.4) and (3.5) under the assumptions of Theorem 3.6.

Theorem 6.4. *Let us assume (H1)–(H3). Let u_ε be the solution to the viscoelastic dynamic system (2.9) and let u_0 be the solution to the stationary problem (3.1). Then (3.4) and (3.5) hold.*

Proof. Thanks to Lemma 3.9 we can suppose $z = 0$ and $z_\varepsilon = 0$ for every $\varepsilon > 0$. Let p_ε be defined by (2.12). Since $z_\varepsilon = 0$, by Remark 2.3 the function u_ε solves (2.13) with $h_\varepsilon = f_\varepsilon$, $\ell_\varepsilon = g_\varepsilon - p_\varepsilon$, $v_\varepsilon^0 = u_{\varepsilon, in}(0)$, and $v_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0)$. To obtain (3.4) and (3.5) we cannot apply Theorems 6.2 and 6.3 directly, because $\{p_\varepsilon\}_\varepsilon$ does not converge to 0 in $W^{1,1}(0, T; V'_0)$ as $\varepsilon \rightarrow 0^+$ and, in general, $p_\varepsilon \notin L^2(0, T; H)$.

To overcome this difficulty we construct a family $\{q_\varepsilon\}_\varepsilon \subset H^1(0, T; H)$ such that $\|q_\varepsilon - p_\varepsilon\|_{W^{1,1}(0, T; V'_0)}$ is uniformly small and $q_\varepsilon \rightarrow 0$ strongly in $L^2(0, T; H)$ as $\varepsilon \rightarrow 0^+$. Then we can apply Theorems 6.2 and 6.3 to the solutions v_ε to (2.13) with p_ε replaced by q_ε , obtaining that $v_\varepsilon \rightarrow u_0$ strongly in $L^2(0, T; V)$ and $\varepsilon \dot{v}_\varepsilon \rightarrow 0$ strongly in $L^2(0, T; H)$. Finally, we show that $\|v_\varepsilon - u_\varepsilon\|_{L^2(0, T; V)}$ and $\varepsilon \|\dot{v}_\varepsilon - \dot{u}_\varepsilon\|_{L^2(0, T; H)}$ are small uniformly with respect to ε , and this leads to the proof of (3.4) and (3.5).

To construct q_ε we consider g_ε^0 introduced in (2.12) and we define

$$\tilde{g}_\varepsilon^0 := \int_{-\infty}^0 \frac{1}{\beta \varepsilon} e^{\frac{\tau}{\beta \varepsilon}} \operatorname{div}(\mathbb{B} e u_{in}(\tau)) d\tau = (\rho_\varepsilon * \operatorname{div}(\mathbb{B} e u_{in})) (0),$$

where ρ_ε is the convolution kernel in (3.17). By (H3) we have $\operatorname{div}(\mathbb{B} e u_{in}) \in C^0((-\infty, 0]; V'_0)$, hence the properties of convolutions imply

$$\tilde{g}_\varepsilon^0 \xrightarrow{\varepsilon \rightarrow 0^+} g^0 := \operatorname{div}(\mathbb{B} e u_{in}(0)) \quad \text{strongly in } V'_0. \quad (6.26)$$

Since

$$\begin{aligned} \|g_\varepsilon^0 - \tilde{g}_\varepsilon^0\|_{V'_0} &\leq \int_{-\infty}^{-a} \frac{1}{\beta \varepsilon} e^{\frac{\tau}{\beta \varepsilon}} (\|\operatorname{div}(\mathbb{B}(e u_{\varepsilon, in}(\tau)))\|_{V'_0} + \|\operatorname{div}(\mathbb{B}(e u_{in}(\tau)))\|_{V'_0}) d\tau \\ &\quad + \|\operatorname{div}(\mathbb{B}(e u_{\varepsilon, in} - e u_{in}))\|_{L^\infty(-a, 0; V'_0)}, \end{aligned}$$

thanks to (H3) we have $g_\varepsilon^0 - \tilde{g}_\varepsilon^0 \rightarrow 0$ strongly in V'_0 as $\varepsilon \rightarrow 0^+$, hence (6.26) implies

$$g_\varepsilon^0 \xrightarrow{\varepsilon \rightarrow 0^+} g^0 \quad \text{strongly in } V'_0. \quad (6.27)$$

Let us fix $\delta > 0$. By the density of H in V'_0 we can find $h^0 \in H$ such that $\|h^0 - g^0\|_{V'_0} < \delta$. By (6.27) there exists $\varepsilon_0 = \varepsilon_0(\delta) \in (0, \frac{1}{\beta})$ such that

$$\|h^0 - g_\varepsilon^0\|_{V'_0} < \delta \quad \text{for every } \varepsilon \in (0, \varepsilon_0). \quad (6.28)$$

Let $q_\varepsilon \in H^1(0, T; H)$ be defined by $q_\varepsilon(t) := e^{-\frac{t}{\beta \varepsilon}} h^0$ for every $t \in [0, T]$. Then

$$q_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; H). \quad (6.29)$$

Since $p_\varepsilon(t) = e^{-\frac{t}{\beta \varepsilon}} g_\varepsilon^0$, by (6.28) we have also

$$\|q_\varepsilon - p_\varepsilon\|_{W^{1,1}(0, T; V'_0)} \leq (\beta \varepsilon + 1) \|h^0 - g_\varepsilon^0\|_{V'_0} \leq 2\delta \quad \text{for every } \varepsilon \in (0, \varepsilon_0). \quad (6.30)$$

Let v_ε be the solution to (2.13) with $h_\varepsilon = f_\varepsilon - q_\varepsilon$, $\ell_\varepsilon = g_\varepsilon$, $v_\varepsilon^0 = u_{\varepsilon, in}(0)$, and $v_\varepsilon^1 = \dot{u}_{\varepsilon, in}(0)$. By (H1) and (6.29) we have

$$\begin{aligned} f_\varepsilon - q_\varepsilon &\xrightarrow{\varepsilon \rightarrow 0^+} f \quad \text{strongly in } L^2(0, T; H) \\ g_\varepsilon &\xrightarrow{\varepsilon \rightarrow 0^+} g \quad \text{strongly in } W^{1,1}(0, T; V'_0). \end{aligned}$$

By (H3) we have

$$\begin{aligned} u_{\varepsilon, in}(0) &\xrightarrow{\varepsilon \rightarrow 0^+} u_{in}(0) \quad \text{strongly in } V \\ \varepsilon \dot{u}_{\varepsilon, in}(0) &\xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } H. \end{aligned}$$

Therefore we can apply Theorems 6.2 and 6.3 to obtain

$$v_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0^+} u_0 \quad \text{strongly in } L^2(0, T; V) \quad (6.31)$$

$$\varepsilon \dot{v}_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; H). \quad (6.32)$$

To estimate the difference $v_{\varepsilon} - u_{\varepsilon}$ we observe that it solves (2.13) with $h_{\varepsilon} = 0$, $\ell_{\varepsilon} = p_{\varepsilon} - q_{\varepsilon}$, $v_{\varepsilon}^0 = 0$, and $v_{\varepsilon}^1 = 0$. Therefore, by Lemma 3.8 and (6.30) we have

$$\varepsilon^2 \|\dot{v}_{\varepsilon} - \dot{u}_{\varepsilon}\|_{L^2(0, T; H)}^2 + \|v_{\varepsilon} - u_{\varepsilon}\|_{L^2(0, T; V)}^2 \leq C_E \|q_{\varepsilon} - p_{\varepsilon}\|_{W^{1,1}(0, T; V'_0)}^2 \leq 4C_E \delta^2. \quad (6.33)$$

Since by (6.33)

$$\begin{aligned} \|u_{\varepsilon} - u_0\|_{L^2(0, T; V)} &\leq \|u_{\varepsilon} - v_{\varepsilon}\|_{L^2(0, T; V)} + \|v_{\varepsilon} - u_0\|_{L^2(0, T; V)} \\ &\leq \|v_{\varepsilon} - u_0\|_{L^2(0, T; V)} + 2\sqrt{C_E} \delta, \\ \varepsilon \|\dot{u}_{\varepsilon}\|_{L^2(0, T; H)} &\leq \varepsilon \|\dot{u}_{\varepsilon} - \dot{v}_{\varepsilon}\|_{L^2(0, T; H)} + \varepsilon \|\dot{v}_{\varepsilon}\|_{L^2(0, T; H)} \\ &\leq \varepsilon \|\dot{v}_{\varepsilon}\|_{L^2(0, T; H)} + 2\sqrt{C_E} \delta, \end{aligned}$$

thanks to (6.31) and (6.32) we have

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0^+} \|u_{\varepsilon} - u_0\|_{L^2(0, T; V)} &\leq 2\sqrt{C_E} \delta, \\ \limsup_{\varepsilon \rightarrow 0^+} \varepsilon \|\dot{u}_{\varepsilon}\|_{L^2(0, T; H)} &\leq 2\sqrt{C_E} \delta. \end{aligned}$$

By the arbitrariness of $\delta > 0$ we obtain (3.4) and (3.5), which concludes the proof. \square

7. The local uniform convergence

In this section we shall prove (3.6) and (3.7) under the assumptions of Theorems 3.6 and 3.7. The proof is based on the following lemma.

Lemma 7.1. *Let $\{\ell_{\varepsilon}\}_{\varepsilon} \subset H^1(0, T; V'_0)$ and $\ell \in W^{1,1}(0, T; V'_0)$ be such that*

$$\ell_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0^+} \ell \quad \text{strongly in } W^{1,1}(\eta, T; V'_0) \text{ for every } \eta \in (0, T). \quad (7.1)$$

Let v_{ε} be a solution to the viscoelastic dynamic system (2.13) with $h_{\varepsilon} = 0$ and arbitrary initial data. Moreover, let v_0 be the solution to the stationary problem (3.2) with $h = 0$. We assume that

$$v_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0^+} v_0 \quad \text{strongly in } L^2(0, T; V), \quad (7.2)$$

$$\varepsilon \dot{v}_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; H). \quad (7.3)$$

Then

$$v_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} v_0 \quad \text{strongly in } L^\infty(\eta, T; V) \text{ for every } \eta \in (0, T), \quad (7.4)$$

$$\varepsilon \dot{v}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(\eta, T; H) \text{ for every } \eta \in (0, T). \quad (7.5)$$

Proof. We divide the proof into two steps.

Step 1. Let us assume $\ell_\varepsilon = \ell \in H^2(0, T; V'_0)$ for every $\varepsilon > 0$. By Lemma 3.4 (with $z = 0$) we have $v_0 \in H^2(0, T; V)$, hence recalling (3.2) we get for a.e. $t \in [0, T]$

$$\begin{aligned} \varepsilon^2 \ddot{v}_0(t) - \operatorname{div}((\mathbb{A} + \mathbb{B})ev_0(t)) + \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_0(\tau)) d\tau \\ = \varepsilon^2 \ddot{v}_0(t) + \ell(t) - \operatorname{div}(\mathbb{B}ev_0(t)) + \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_0(\tau)) d\tau. \end{aligned} \quad (7.6)$$

Now we define $\bar{v}_\varepsilon := v_\varepsilon - v_0$ and observe that by (7.2) and (7.3) we have

$$\bar{v}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; V), \quad (7.7)$$

$$\varepsilon \dot{\bar{v}}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; H). \quad (7.8)$$

Let us consider

$$q_\varepsilon(t) := \operatorname{div}(\mathbb{B}ev_0(t)) - \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_0(\tau)) d\tau.$$

Since v_ε satisfies (2.13) with $h_\varepsilon = 0$, by (7.6) the function \bar{v}_ε satisfies (2.13) with $h_\varepsilon = -\varepsilon^2 \ddot{v}_0$ and $\ell_\varepsilon = q_\varepsilon$. After two integrations by parts in time we deduce

$$\begin{aligned} \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_0(\tau)) d\tau &= \operatorname{div}(\mathbb{B}ev_0(t)) - e^{-\frac{t}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}ev_0(0)) \\ &\quad - \beta\varepsilon \operatorname{div}(\mathbb{B}e\dot{v}_0(t)) + \beta\varepsilon e^{-\frac{t}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e\dot{v}_0(0)) + \beta\varepsilon \int_0^t e^{-\frac{t-\tau}{\beta\varepsilon}} \operatorname{div}(\mathbb{B}e\ddot{v}_0(\tau)) d\tau, \end{aligned}$$

hence

$$q_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } W^{1,1}(\eta, T; V'_0) \text{ for every } \eta \in (0, T). \quad (7.9)$$

Now we fix $\delta \in (0, T)$, and we consider $\eta \in (0, \delta)$ and $\zeta \in (\eta, \delta)$. We define the family of functions $\{\bar{w}_\varepsilon\}_\varepsilon \subset H^1(0, T; \tilde{H})$ by

$$\bar{w}_\varepsilon(t) := \int_0^t \frac{1}{\beta\varepsilon} e^{-\frac{t-\tau}{\beta\varepsilon}} e\bar{v}_\varepsilon(\tau) d\tau = (\rho_\varepsilon * e\bar{v}_\varepsilon)(t) \quad \text{for every } t \in [0, T],$$

where ρ_ε is defined by (3.17) and \bar{v}_ε is extended to \mathbb{R} by setting $\bar{v}_\varepsilon(t) = 0$ on $\mathbb{R} \setminus [0, T]$. By properties of convolutions we have

$$e\bar{v}_\varepsilon - \bar{w}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^2(0, T; \tilde{H}). \quad (7.10)$$

By the energy-dissipation balance (2.25) of Proposition 2.7, for every $t \in [\eta, T]$ and $s \in (\eta, \zeta)$ we can write

$$\begin{aligned} & \frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon(t)\|^2 + \frac{1}{2} (\mathbb{A} e \bar{v}_\varepsilon(t), e \bar{v}_\varepsilon(t)) + \frac{1}{2} (\mathbb{B} (e \bar{v}_\varepsilon(t) - \bar{w}_\varepsilon(t), e \bar{v}_\varepsilon(t) - \bar{w}_\varepsilon(t)) \\ & + \beta \varepsilon \int_s^t (\mathbb{B} \dot{w}_\varepsilon(\tau), \dot{w}_\varepsilon(\tau)) d\tau = \frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon(s)\|^2 + \frac{1}{2} (\mathbb{A} e \bar{v}_\varepsilon(s), e \bar{v}_\varepsilon(s)) \\ & + \frac{1}{2} (\mathbb{B} (e \bar{v}_\varepsilon(s) - \bar{w}_\varepsilon(s), e \bar{v}_\varepsilon(s) - \bar{w}_\varepsilon(s)) + \mathcal{W}_\varepsilon(t, s), \end{aligned} \quad (7.11)$$

where the work is defined by

$$\begin{aligned} \mathcal{W}_\varepsilon(t, s) &= \langle q_\varepsilon(t), \bar{v}_\varepsilon(t) \rangle - \langle q_\varepsilon(s), \bar{v}_\varepsilon(s) \rangle \\ &\quad - \int_s^t \langle \dot{q}_\varepsilon(\tau), \bar{v}_\varepsilon(\tau) \rangle d\tau - \varepsilon \int_s^t \langle \ddot{v}_0(\tau), \varepsilon \dot{v}_\varepsilon(\tau) \rangle d\tau. \end{aligned}$$

Now we take the mean value with respect to s of all terms of (7.11) on (η, ζ) , and we pass to the supremum with respect to t on $[\eta, T]$. Thanks to (2.3) and (2.6) we deduce

$$\begin{aligned} & \frac{\varepsilon^2}{2} \|\dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)}^2 + \frac{C_{\mathbb{A}}}{2(C_P^2 + 1)} \|\bar{v}_\varepsilon\|_{L^\infty(\eta, T; V)}^2 \\ & \leq \frac{\varepsilon^2}{2} \int_\eta^\zeta \|\dot{v}_\varepsilon(s)\|^2 ds + \frac{C_{\mathbb{A}}}{2} \int_\eta^\zeta \|\bar{v}_\varepsilon(s)\|_V^2 ds \\ & \quad + \frac{C_{\mathbb{B}}}{2} \int_\eta^\zeta \|e \bar{v}_\varepsilon(s) - \bar{w}_\varepsilon(s)\|^2 ds + \int_\eta^\zeta \sup_{t \in [\eta, T]} |\mathcal{W}_\varepsilon(t, s)| ds. \end{aligned} \quad (7.12)$$

Notice that for every $s \in (\eta, \zeta)$ we have

$$\begin{aligned} \sup_{t \in [\eta, T]} |\mathcal{W}_\varepsilon(t, s)| &\leq (2\|q_\varepsilon\|_{L^\infty(\eta, T; V'_0)} + \|\dot{q}_\varepsilon\|_{L^1(\eta, T; V'_0)}) \|\bar{v}_\varepsilon\|_{L^\infty(\eta, T; V)} \\ &\quad + \varepsilon \|\ddot{v}_0\|_{L^1(\eta, T; H)} \|\varepsilon \dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)} \\ &\leq (3 + \frac{2}{T}) \|q_\varepsilon\|_{W^{1,1}(\eta, T; V'_0)} \|\bar{v}_\varepsilon\|_{L^\infty(\eta, T; V)} \\ &\quad + \varepsilon \|\ddot{v}_0\|_{L^1(\eta, T; H)} \|\varepsilon \dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)}, \end{aligned}$$

hence thanks to the Young Inequality and (7.12) there exists a positive constant $C = C(\mathbb{A}, \mathbb{B}, \Omega, T)$ such that

$$\begin{aligned} & \varepsilon^2 \|\dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)}^2 + \|\bar{v}_\varepsilon\|_{L^\infty(\eta, T; V)}^2 \leq C \left(\varepsilon^2 \int_\eta^\zeta \|\dot{v}_\varepsilon(s)\|^2 ds + \int_\eta^\zeta \|\bar{v}_\varepsilon(s)\|_V^2 ds \right. \\ & \quad \left. + \int_\eta^\zeta \|e \bar{v}_\varepsilon(s) - \bar{w}_\varepsilon(s)\|^2 ds + \|q_\varepsilon\|_{W^{1,1}(\eta, T; V'_0)}^2 + \varepsilon^2 \|\ddot{v}_0\|_{L^1(\eta, T; H)}^2 \right). \end{aligned} \quad (7.13)$$

By passing to the limit in (7.13) as $\varepsilon \rightarrow 0^+$, thanks to (7.7), (7.8), (7.9), and (7.10) we obtain

$$\varepsilon \|\dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)} + \|\bar{v}_\varepsilon\|_{L^\infty(\eta, T; V)} \xrightarrow{\varepsilon \rightarrow 0^+} 0,$$

which, by the definition of \bar{v}_ε , concludes the proof of (7.4) and (7.5) in the case $\ell \in H^2(0, T; V'_0)$.

Step 2. In the general case $\ell \in W^{1,1}(0, T; V'_0)$ we use an approximation argument. Given $\delta > 0$, by Lemma 4.3 there exists a function $\psi \in H^2(0, T; H)$ such that

$$\|\psi - \ell\|_{W^{1,1}(0, T; V'_0)} < \delta. \quad (7.14)$$

Thanks to (7.1) for every $\sigma \in (0, T)$ there exists a positive number $\varepsilon_0 = \varepsilon_0(\delta, \sigma)$ such that

$$\|\psi - \ell_\varepsilon\|_{W^{1,1}(\sigma, T; V'_0)} < \delta \quad \text{for every } \varepsilon \in (0, \varepsilon_0). \quad (7.15)$$

Let \tilde{v}_ε be the solution to (2.13) in the interval $[\sigma, T]$ with $h_\varepsilon = 0$, $\ell_\varepsilon = \psi$, $\tilde{v}_\varepsilon(\sigma) = v_\varepsilon(\sigma)$, and $\dot{\tilde{v}}_\varepsilon(\sigma) = \dot{v}_\varepsilon(\sigma)$, and let \tilde{v}_0 be the solution to (3.2) in the interval $[0, T]$ with $h = 0$ and $\ell = \psi$. By applying Step 1 in the interval $[\sigma, T]$ we obtain

$$\tilde{v}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} \tilde{v}_0 \quad \text{strongly in } L^\infty(\eta, T; V) \text{ for every } \eta \in (\sigma, T), \quad (7.16)$$

$$\varepsilon \dot{\tilde{v}}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{strongly in } L^\infty(\eta, T; H) \text{ for every } \eta \in (\sigma, T). \quad (7.17)$$

We set $\bar{v}_0 := \tilde{v}_0 - v_0$ and $\bar{v}_\varepsilon := \tilde{v}_\varepsilon - v_\varepsilon$. We observe that \bar{v}_0 is the solution to (3.2) with $h = 0$ and ℓ replaced by $\psi - \ell$, hence by the Lax-Milgram Lemma we get

$$\|\bar{v}_0\|_{L^\infty(0, T; V)} \leq \frac{C_P^2 + 1}{c_A} \|\psi - \ell\|_{L^\infty(0, T; V'_0)} \leq \frac{C_P^2 + 1}{c_A} \left(1 + \frac{1}{T}\right) \|\psi - \ell\|_{W^{1,1}(0, T; V'_0)}. \quad (7.18)$$

Moreover, \bar{v}_ε is the solution to (2.13) in the interval $[\sigma, T]$ with $h_\varepsilon = 0$, ℓ_ε replaced by $\psi - \ell_\varepsilon$, and homogeneous initial conditions. Thanks to Lemma 3.8 we obtain

$$\varepsilon \|\dot{\bar{v}}_\varepsilon\|_{L^\infty(\sigma, T; H)}^2 + \|\bar{v}_\varepsilon\|_{L^\infty(\sigma, T; V)}^2 \leq C_E \|\psi - \ell_\varepsilon\|_{W^{1,1}(\sigma, T; V'_0)}^2. \quad (7.19)$$

By combining (7.14), (7.15), (7.18), and (7.19), we can find a positive constant $C = C(\mathbb{A}, \mathbb{B}, \Omega, T)$ such that

$$\varepsilon \|\dot{\bar{v}}_\varepsilon\|_{L^\infty(\sigma, T; H)} + \|\bar{v}_\varepsilon\|_{L^\infty(\sigma, T; V)} + \|\bar{v}_0\|_{L^\infty(\sigma, T; V)} \leq C\delta. \quad (7.20)$$

Since for every $\eta \in (\sigma, T)$ we have

$$\begin{aligned} \|v_\varepsilon - v_0\|_{L^\infty(\eta, T; V)} &\leq \|\bar{v}_\varepsilon\|_{L^\infty(\eta, T; V)} + \|\tilde{v}_\varepsilon - \tilde{v}_0\|_{L^\infty(\eta, T; V)} + \|\bar{v}_0\|_{L^\infty(\eta, T; V)}, \\ \varepsilon \|\dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)} &\leq \varepsilon \|\dot{\tilde{v}}_\varepsilon\|_{L^\infty(\eta, T; H)} + \varepsilon \|\dot{\bar{v}}_\varepsilon\|_{L^\infty(\eta, T; H)}, \end{aligned}$$

thanks to (7.16), (7.17), and (7.20) we obtain

$$\limsup_{\varepsilon \rightarrow 0^+} \|v_\varepsilon - v_0\|_{L^\infty(\eta, T; V)} \leq C\delta \quad \text{and} \quad \limsup_{\varepsilon \rightarrow 0^+} \varepsilon \|\dot{v}_\varepsilon\|_{L^\infty(\eta, T; H)} \leq C\delta,$$

for every $\eta \in (\sigma, T)$. By the arbitrariness of $\delta > 0$ and $\sigma > 0$ we conclude. \square

Now we are in a position to prove (3.6) and (3.7).

Theorem 7.2. *Let us assume (H1), (H2), (3.10), and $f_\varepsilon = 0$ for every $\varepsilon > 0$. Let u_ε be the solution to the viscoelastic dynamic system (2.10), with $\varphi_\varepsilon = 0$ and $\gamma_\varepsilon = g_\varepsilon$, and let u_0 be the solution to the stationary problem (3.1), with $f = 0$. Then (3.6) and (3.7) hold.*

Proof. By Theorems 6.2 and 6.3 we obtain (3.4) and (3.5). Since $f_\varepsilon = 0$ and $g_\varepsilon \rightarrow g$ strongly in $W^{1,1}(0, T; V'_0)$ as $\varepsilon \rightarrow 0^+$ by (H1), we can apply Lemma 7.1 to conclude. \square

Theorem 7.3. *Let us assume (H1)–(H3) and $f_\varepsilon = 0$ for every $\varepsilon > 0$. Let u_ε be the solution to the viscoelastic dynamic system (2.9) and let u_0 be the solution to the stationary problem (3.1), with $f = 0$. Then (3.6) and (3.7) hold.*

Proof. Thanks to Lemma 3.9 we can suppose $z = 0$ and $z_\varepsilon = 0$ for every $\varepsilon > 0$. By Theorem 6.4 we obtain (3.4) and (3.5). Since u_ε is a solution to (2.9) with $f_\varepsilon = 0$, by Remark 2.3 it solves (2.13) with $h_\varepsilon = 0$ and $\ell_\varepsilon = g_\varepsilon - p_\varepsilon$, where p_ε is defined by (2.12). Since

$$g_\varepsilon - p_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} g \quad \text{strongly in } W^{1,1}(\eta, T; V'_0) \text{ for every } \eta \in (0, T),$$

we can apply Lemma 7.1 to conclude. \square

Finally we can prove Theorems 3.6 and 3.7.

Proof of Theorem 3.6. It is enough to combine Theorems 4.1, 6.4, and 7.3. \square

Proof of Theorem 3.7. It is enough to combine Theorems 6.2, 6.3, and 7.2. \square

Appendix A.

Throughout this section we fix $a_0 > 0$, $b_0 > 0$, and $c_1 \geq c_0 > 1$. For every a, b with

$$c_0 a \leq b \leq c_1 a, \quad b \geq b_0, \quad a \geq a_0, \quad (\text{A.1})$$

we consider the polynomial $p(z) := \beta z^3 + z^2 + \beta b z + a$ depending on the complex variable z . The following result about the roots of this polynomial is used in the proof of Lemma 5.2 and Proposition 5.4.

Lemma A.1. *There exists a positive constant $\alpha = \alpha(\beta, a_0, b_0, c_0, c_1)$ such that, for every $a, b \in \mathbb{R}$ satisfying (A.1), the roots of the polynomial p have real parts in the interval $(-\frac{1}{\beta}, -\alpha)$.*

Proof. Let us set $z := x + iy$ with $x, y \in \mathbb{R}$. Then $p(z) = 0$ if and only if

$$\begin{cases} \beta x^3 + x^2 + \beta b x - (3\beta x + 1)y^2 + a = 0, \\ y(-\beta y^2 + 3\beta x^2 + 2x + \beta b) = 0, \end{cases}$$

from which we derive

$$\begin{cases} q(x) := \beta x^3 + x^2 + \beta b x + a = 0, \\ y = 0, \end{cases} \quad (\text{A.2})$$

$$\begin{cases} r(x) := 8\beta x^3 + 8x^2 + 2\left(\frac{1}{\beta} + \beta b\right)x + b - a = 0, \\ y^2 = 3x^2 + \frac{2}{\beta}x + b. \end{cases} \quad (\text{A.3})$$

By recalling $a > 0$ and $b - a \geq (c_0 - 1)a > 0$, for every $x \geq 0$ we have $q(x) > 0$ and $r(x) > 0$, and so the real part of the roots cannot be positive or zero. Moreover, since for every $x \leq -\frac{1}{\beta}$ we have $\beta x^3 + x^2 \leq 0$, we obtain

$$\begin{aligned} q(x) &\leq -b + a \leq (1 - c_0)a < 0 \\ r(x) &\leq b - a - 2\left(\frac{1}{\beta^2} + b\right) = -b - a - \frac{2}{\beta^2} < 0, \end{aligned}$$

which imply that the real part of the roots does not belong to $(-\infty, -\frac{1}{\beta}]$. Therefore, by calling $z_1, z_2, z_3 \in \mathbb{C}$ the three roots of the polynomial p , we can say

$$\Re(z_i) \in (-\frac{1}{\beta}, 0) \quad \text{for } i = 1, 2, 3. \quad (\text{A.4})$$

Case 1: there is only one real root. In this case by (A.3) there exists a unique $x_1 \in (-\frac{1}{\beta}, 0)$ which satisfies $r(x_1) = 0$ and $3x_1^2 + \frac{2}{\beta}x_1 + b > 0$. Indeed by setting $y_1 := \sqrt{3x_1^2 + \frac{2}{\beta}x_1 + b}$ we obtain that $x_1 + iy_1$ and $x_1 - iy_1$ are two distinct non-real roots of p . Since

$$\begin{aligned} r(-\frac{1}{2\beta}) &= -\frac{1}{\beta^2} + \frac{2}{\beta^2} - \frac{1}{\beta^2} - b + b - a = -a < 0, \\ r(-\frac{\beta(b-a)}{2(b\beta^2+1)}) &= \frac{\beta^2(b-a)^2((a+b)\beta^2+2)}{(b\beta^2+1)^3} > 0, \end{aligned}$$

then $x_1 \in (-\frac{1}{2\beta}, -\frac{\beta(b-a)}{2(b\beta^2+1)})$. Moreover

$$\begin{aligned} q(-\frac{1}{\beta}) &= -\frac{1}{\beta^2} + \frac{1}{\beta^2} - b + a = -b + a < 0, \\ q(-\frac{a}{\beta b}) &= -\frac{a^3}{b^3\beta^2} + \frac{a^2}{b^2\beta^2} - a + a = \frac{a^2(b-a)}{b^3\beta^2} > 0, \end{aligned}$$

hence there exists $x_0 \in (-\frac{1}{\beta}, -\frac{a}{\beta b})$ such that $q(x_0) = 0$. As a consequence of this, $(x_0, 0)$ satisfies the system in (A.2), which implies that x_0 is the real root of p , hence we have $\Re(z_i) \in (-\frac{1}{\beta}, \max\{-\frac{a}{\beta b}, -\frac{\beta(b-a)}{2(b\beta^2+1)}\})$. Thanks to (A.1) we can say $-\frac{a}{\beta b} \leq -\frac{1}{c_1\beta}$ and $-\frac{\beta(b-a)}{2(b\beta^2+1)} \leq \frac{\beta(1-c_0)a}{2(c_1a\beta^2+1)} \leq \frac{\beta(1-c_0)a_0}{2(c_1a_0\beta^2+1)}$, where in the last inequality we use the decreasing property of the function $a \mapsto \frac{\beta(1-c_0)a}{2(c_1a\beta^2+1)}$. This implies

$$\Re(z_i) \in (-\frac{1}{\beta}, \max\{-\frac{1}{c_1\beta}, \frac{\beta(1-c_0)a_0}{2(c_1a_0\beta^2+1)}\}) \quad \text{for } i = 1, 2, 3. \quad (\text{A.5})$$

Case 2: there are only real roots. In this case we have $b \leq \frac{1}{3\beta^2}$, otherwise $q'(x) > 0$ for every $x \in \mathbb{R}$, which forces p to have also non-real roots. Thanks to (A.1) we have also $a < b \leq \frac{1}{3\beta^2}$. By setting $\tilde{b}_0 := 1 - \sqrt{1 - 3b_0\beta^2}$, we can write

$$-\tilde{b}_0 a_0 \beta \geq -\tilde{b}_0 a \beta \geq -(1 - \sqrt{1 - 3b_0\beta^2})a\beta > \frac{-1 + \sqrt{1 - 3b_0\beta^2}}{3\beta} > -\frac{1}{\beta},$$

which implies

$$q'(x) > 0 \quad \text{for every } x \in [-\tilde{b}_0 a_0 \beta, +\infty). \quad (\text{A.6})$$

Since

$$\begin{aligned} q(-\tilde{b}_0 a_0 \beta) &\geq \beta^2 \tilde{b}_0^2 a_0^2 (1 - \beta^2 \tilde{b}_0 a_0) + a_0 (1 - \beta^2 \tilde{b}_0 b) \\ &> a_0 (1 + \beta^2 \tilde{b}_0^2 a_0) (1 - \beta^2 \tilde{b}_0 b) > 0, \end{aligned}$$

thanks to (A.1), (A.4), and (A.6) we get

$$\Re(z_i) \in (-\frac{1}{\beta}, -\tilde{b}_0 a_0 \beta), \quad \text{for } i = 1, 2, 3. \quad (\text{A.7})$$

By combining (A.5) and (A.7), we obtain the conclusion with

$$\alpha := \min\{\tilde{b}_0 a_0 \beta, \frac{1}{c_1 \beta}, \frac{\beta(c_0 - 1)a_0}{2(c_1 a_0 \beta^2 + 1)}\}.$$

□

The following easy estimate is used in the proof of Lemma 5.2.

Lemma A.2. *For every $z, w \in \mathbb{C}$ with $\Re(z) > 0$ and $\Re(w) < 0$ the following inequality holds:*

$$|(z - w)(z - \bar{w})| \geq |\Re(w)| |\Im(w)|.$$

Proof. Without loss of generality we can suppose $\Im(w) > 0$, otherwise we exchange the role of w with \bar{w} . If $\Im(z) > 0$, then

$$\begin{aligned} |z - w| &\geq |\Re(z - w)| = |\Re(z) + \Re(-w)| = \Re(z) + \Re(-w) \geq |\Re(w)|, \\ |z - \bar{w}| &\geq |\Im(z - \bar{w})| = |\Im(z) + \Im(w)| = \Im(z) + \Im(w) \geq |\Im(w)|, \end{aligned}$$

which give the conclusion in this case. If $\Im(z) < 0$, then

$$\begin{aligned} |z - w| &\geq |\Im(z - w)| = |-\Im(-z) - \Im(w)| = \Im(-z) + \Im(w) \geq |\Im(w)|, \\ |z - \bar{w}| &\geq |\Re(z - \bar{w})| = |\Re(z) + \Re(-w)| = \Re(z) + \Re(-w) \geq |\Re(w)|, \end{aligned}$$

which conclude the proof. □

ACKNOWLEDGEMENTS. This paper is based on work supported by the National Research Project (PRIN 2017) "Variational Methods for Stationary and Evolution Problems with Singularities and Interfaces", funded by the Italian Ministry of University and Research. The authors are members of the *Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* (GNAMPA) of the *Istituto Nazionale di Alta Matematica* (INdAM).

References

- [1] V. AGOSTINIANI: Second order approximations of quasistatic evolution problems in finite dimension, *Discrete Contin. Dyn. Syst.* **32** (2010), 1125–1167.
- [2] W. ARENDT, C. BATTY, M. HIEBER, F. NEUBRANDER: Vector-valued Laplace Transforms and Cauchy Problems, Monographs in Mathematics **96**. Birkhäuser, Basel, 2001.
- [3] C. DAFERMOS: An abstract Volterra equation with applications to linear viscoelasticity, *Journal of Differential Equations* **7** (1970), 554–569.

- [4] R. DAUTRAY AND J.L. LIONS: Mathematical Analysis and Numerical Methods for Science and Technology, Volume 1 Physical Origins and Classical Methods, Original French edition published by Masson, S.A., Paris, 1984
- [5] M. FABRIZIO, C. GIORGI, V. PATA: A New Approach to Equations with Memory, *Arch. Rational Mech. Anal.* **198** (2010), 189–232.
- [6] M. FABRIZIO AND A. MORRO: Mathematical Problems in Linear Viscoelasticity. SIAM Studies in Applied Mathematics **12** 2001.
- [7] P. GIDONI AND F. RIVA: A vanishing inertia analysis for finite dimensional rate-independent systems with nonautonomous dissipation and an application to soft crawlers, *Calc. Var. Partial Differential Equations* **60** (2021) Paper 191.
- [8] J.L. LIONS AND E. MAGENES: Non-Homogeneous Boundary-Value Problems and Applications. Vol. 1 **181**, Springer-Verlag, 1972.
- [9] G. LAZZARONI AND L. NARDINI: On the quasistatic limit of dynamic evolutions for a peeling test in dimension one, *J. Nonlinear Sci.* **28** (2018), 269–304.
- [10] L. NARDINI: A note on the convergence of singularly perturbed second order potential-type equations, *J. Dynam. Differential Equations* **29** (2017), 783–707.
- [11] O.A. OLEINIK, A.S. SHAMAEV, AND G.A. YOSIFIAN: Mathematical problems in elasticity and homogenization. Studies in Mathematics and its Applications, 26. North-Holland Publishing Co., Amsterdam, 1992.
- [12] F. RIVA: On the approximation of quasistatic evolutions for the debonding of a thin film via vanishing inertia and viscosity, *J. Nonlinear Sci.* **30** (2020), 903–951.
- [13] G. SCILLA AND F. SOLOMBRINO: A variational approach to the quasistatic limit of viscous dynamic evolutions in finite dimension, *J. Differential Equations* **267** (2019), 6216–6264.
- [14] F. SAPIO: A dynamic model for viscoelasticity in domains with time-dependent cracks. Preprint SISSA 14/2020/MATE.
- [15] L.I. SLEPYAN: Models and phenomena in fracture mechanics, Foundations of Engineering Mechanics. Springer-Verlag, Berlin, 2002.

Gianni Dal Maso

(Gianni Dal Maso) SISSA, via Bonomea 265, 34136 Trieste, Italy

e-mail: dalmaso@sissa.it

Francesco Sapiro

(Francesco Sapiro) SISSA, via Bonomea 265, 34136 Trieste, Italy

e-mail: fsapiro@sissa.it