

# A BMO-type characterization of higher order Sobolev spaces

Serena Guarino Lo Bianco<sup>1</sup> and Roberta Schiattarella<sup>2\*†</sup>

<sup>1</sup>Dipartimento di Agraria, Università di Napoli “Federico II”, via  
Università 100, Portici (NA), 80055, Italy.

<sup>2</sup>Dipartimento di Matematica e Applicazioni “R. Caccioppoli”,  
Università di Napoli “Federico II”, Via Cintia, Napoli, 80126, Italy.

\*Corresponding author(s). E-mail(s): [roberta.schiattarella@unina.it](mailto:roberta.schiattarella@unina.it);  
Contributing authors: [serena.guarinolobianco@unina.it](mailto:serena.guarinolobianco@unina.it);

†These authors contributed equally to this work.

## Abstract

We obtain a new characterization of the higher Sobolev space  $W^{m,p}(\mathbb{R}^n)$ ,  $m \in \mathbb{N}$  and  $p \in (1, +\infty)$  and of the space  $BV^m$ , the space of functions of higher order bounded variation. The characterizations are in term of BMO-type seminorms. The results unify and substantially extend previous results in [16] and [13].

**Keywords:** Higher order Sobolev spaces, Higher order bounded variation, BMO-type seminorms

**MSC Classification:** 46E35 , 26B30

## 1 Introduction

Let  $W_{\text{loc}}^{m,p}(\mathbb{R}^n)$  ( $m \in \mathbb{N}$ ,  $1 \leq p < \infty$ ), denote the Sobolev space of functions belonging to  $L_{\text{loc}}^p(\mathbb{R}^n)$  whose distribution derivatives up to order  $m$  belong to  $L_{\text{loc}}^p(\mathbb{R}^n)$ .

In [4], the Authors studied a characterization of  $W^{m,p}$  based on J. Bourgain, H. Brezis and P. Mironescu’s approach introduced in [6] (see also [8]). In particular they prove that if  $f \in W^{m-1,p}(\Omega)$ ,  $1 < p < \infty$  and  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^n$

## 2 A BMO-type characterization of higher order Sobolev spaces

then  $f$  belongs to  $W^{m,p}(\Omega)$  if and only if,

$$\liminf_{\varepsilon \rightarrow 0} \int_{\Omega} \int_{\Omega} \frac{|R^{m-1}f(x,y)|^p}{|x-y|^{mp}} \rho_{\varepsilon}(|x-y|) dx dy < \infty \quad (1)$$

where  $\rho_{\varepsilon}$ , with  $\varepsilon > 0$ , are radial mollifiers and  $R^{m-1}f$  is the Taylor  $(m-1)$  remainder of  $f$ . For  $p = 1$ , the condition (1) describes  $BV^m$ .

Here we say that a  $W^{m-1,1}(\Omega)$  is of  $m$ -th order bounded variation  $BV^m$  if its  $m$ -th order partial derivatives in the sense of distributions are finite Radon measures. Spaces of this kind have been studied in [11] as applications in mathematical imaging in the setting of isotropic and anisotropic variants of the TV-model (see also [14]).

Another characterization of  $W^{m,p}$ ,  $1 < p < \infty$ , ( $BV^m$  for  $p = 1$ ) formulated in terms of the  $m$ -th differences has been presented in [5].

In this article we are concerned with a characterization of  $W^{m,p}$   $1 < p < \infty$ , ( $BV^m$  for  $p = 1$ ) as the limit of certain BMO-type seminorms similar to the one introduced by J. Bourgain, H. Brezis, P. Mironescu in [7].

In [16] the Authors showed that a function  $f \in L^p_{\text{loc}}(\mathbb{R}^n)$  belongs to the Sobolev space  $W^{1,p}_{\text{loc}}(\mathbb{R}^n)$ ,  $1 < p < +\infty$ , if and only if

$$\lim_{\varepsilon \rightarrow 0^+} K(\varepsilon, 1, p) < +\infty$$

where

$$K_{\varepsilon}(f, 1, p) := \varepsilon^{n-p} \sup_{\mathcal{G}_{\varepsilon}} \sum_{Q' \in \mathcal{G}_{\varepsilon}} \int_{Q'} \left| f(x) - \int_{Q'} f \right|^p dx, \quad (2)$$

and the supremum on the right hand side is taken over all families  $\mathcal{G}_{\varepsilon}$  of disjoint  $\varepsilon$ -cubes  $Q' = Q'(x_0, \varepsilon)$  of side length  $\varepsilon$ , centered in  $x_0$ , with arbitrary orientation. Moreover, if  $f \in W^{1,p}_{\text{loc}}(\mathbb{R}^n)$  and  $p \geq 1$  then

$$\lim_{\varepsilon \rightarrow 0^+} K_{\varepsilon}(f, 1, p) = \gamma(n, p) \int_{\mathbb{R}^n} |\nabla f|^p dx \quad (3)$$

where

$$\gamma(n, p) := \max_{\nu \in \mathbb{S}^{n-1}} \int_Q |x \cdot \nu|^p dx \quad (4)$$

where  $Q = \left(-\frac{1}{2}, \frac{1}{2}\right)^n$ .

Following some ideas in [1], an analogous representation formula is obtained for the total variation of  $SBV$  functions in [15] (see also [12]). For related results see also [10], [13].

Here, given a function  $f \in W^{m-1,p}_{\text{loc}}(\mathbb{R}^n)$ ,  $p \geq 1$ , for any  $\varepsilon > 0$ , we consider

$$K_{\varepsilon}(f, m, p) := \varepsilon^{n-mp} \sup_{\mathcal{G}_{\varepsilon}} \sum_{Q' \in \mathcal{G}_{\varepsilon}} \int_{Q'} |f(x) - P^{m-1}_{Q'}[f](x)|^p dx,$$

where the families  $\mathcal{G}_\varepsilon$  are as above and  $P_{Q'}^{m-1}[f]$  is the polynomial of degree  $m - 1$  centered at  $x_0$ , given by

$$P_{Q'}^{m-1}[f](x) = \sum_{|\alpha| \leq m-1} (x - x_0)^\alpha \int_{Q'} (D^\alpha f)(s) ds. \quad (5)$$

In particular, for  $m = 1$  and  $m = 2$  we have:

$$P_{Q'}^0[f](x) = \int_{Q'} f; \quad P_{Q'}^1[f](x) = \int_{Q'} f + \sum_{i=1}^n (x_i - x_{0_i}) \left( \int_{Q'} \frac{\partial f}{\partial y_i}(y) dy \right).$$

Our main Theorem reads as follows:

**Theorem 1** *Let  $p > 1$  and  $f \in W_{loc}^{m-1,p}(\mathbb{R}^n)$ , then*

$$|\nabla^m f| \in L_{loc}^p(\mathbb{R}^n) \iff \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, p) < \infty. \quad (6)$$

*Moreover, if  $f \in W_{loc}^{m,p}(\mathbb{R}^n)$  and  $p \geq 1$  we have also*

$$\lim_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, p) = \beta(n, m, p) \int_{\mathbb{R}^n} |\nabla^m f|^p dx. \quad (7)$$

The constant in (7) is given by

$$\beta(n, m, p) := \max_{v \in \mathbb{S}^{N-1}} \left( \frac{1}{m!} \right)^p \int_Q \left| v \cdot x^m - \int_Q v \cdot y^m dy \right|^p dx. \quad (8)$$

where  $N = n^m$  and we refer to Section 2 for the notation.

Note that this Theorem is exactly an extension of Theorem 2.2 in [16] to the higher order case; indeed, in the case  $m = 1$ , since  $\int_Q x \cdot v dx = 0$ , the constant  $\beta(n, 1, p)$  coincides with the one defined in (4).

A drawback of the formula (7) is that one does not recover the function in  $BV^m$ . However, we are able to show that it is possible to characterize the functions in  $BV^m(\mathbb{R}^n)$  as the functions  $f \in W_{loc}^{m-1,1}(\mathbb{R}^n)$  such that  $\limsup_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, 1) < +\infty$ .

## 2 Notation and preliminaries

We denote by  $Q = \left(-\frac{1}{2}, \frac{1}{2}\right)^n \subset \mathbb{R}^n$  the unit cube with faces parallel to coordinate axes in  $\mathbb{R}^n$ . For any  $z \in \mathbb{R}^n$  and  $\varepsilon > 0$  we denote by  $Q_\varepsilon(z) = z + \varepsilon Q$  the cube of sidelength  $\varepsilon$  centered in  $z$ .

For  $m, n \geq 1$ , we denote by  $N_j = n^{m-j}$  for  $j = 0, \dots, m$ . Given  $v \in \mathbb{R}^{N_0}$  we denotes its components by  $v_{i_1, \dots, i_k, \dots, i_m}$  with  $i_k = 1 \dots n$ . Taking  $x \in \mathbb{R}^n$ ,  $x = (x_{i_k})_{i_k \in \{1, \dots, n\}}$  we define the product  $v \cdot x$  as the element of  $\mathbb{R}^{N_1}$  given by

$$(v \cdot x)_{i_1, \dots, i_{k-1}, i_{k+1}, \dots, i_m} = \sum_{i_k=1}^n v_{i_1, i_2, \dots, i_m} x_{i_k}.$$

#### 4 A BMO-type characterization of higher order Sobolev spaces

The product of  $v \in R^{N_0}$  and  $m$  times the vector  $x \in \mathbb{R}^n$ ,  $v \cdot x \cdot x \cdots x$  is an element of  $\mathbb{R}^{N_m} = \mathbb{R}$  and it is denoted for brevity by  $v \cdot x^m$ .

For a multi-index  $\alpha = (\alpha_1, \dots, \alpha_n)$ ,  $\alpha_i \geq 0$  and a point  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ , we denote by

$$x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}$$

the monomial of degree  $|\alpha| = \sum_{i=1}^n \alpha_i$ .

In the same way,

$$D^\alpha u = \frac{\partial^{|\alpha|} u}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}$$

is a weak partial derivative of order  $|\alpha|$ .

Sometimes, we use the convention that  $D^0 u = u$ . Moreover, let  $\nabla^m u$  be a vector with the components  $D^\alpha u$ ,  $|\alpha| = m$ .

### 2.1 The Sobolev space $W^{m,p}$

**Definition 1** Let  $\Omega \subset \mathbb{R}^n$  be an open set, let  $m \in \mathbb{N}$ , and let  $1 \leq p < \infty$ . The Sobolev space  $W^{m,p}(\Omega)$  is the space of all functions  $u \in L^p(\Omega)$  which admit  $\alpha$ -th weak derivative  $D^\alpha u$  in  $L^p(\Omega)$  for every  $\alpha \in \mathbb{N}^n$  with  $1 \leq |\alpha| \leq m$ .

The space  $W^{m,p}(\Omega)$  is endowed with the norm

$$\|u\|_{W^{m,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \sum_{1 \leq |\alpha| \leq m} \|D^\alpha u\|_{L^p(\Omega)}$$

**Definition 2** Let  $\Omega \subset \mathbb{R}^n$  be an open set, let  $m \in \mathbb{N}$ , and let  $1 \leq p < \infty$ . The homogeneous Sobolev space  $\dot{W}^{m,p}(\Omega)$  is the space of all functions  $u \in L^1_{loc}(\Omega)$  whose  $\alpha$ -th weak derivative  $D^\alpha u$  belongs to  $L^p(\Omega)$  for every  $\alpha \in \mathbb{N}^n$  with  $|\alpha| = m$ .

Note that the inclusion

$$W^{m,p}(\Omega) \subseteq \dot{W}^{m,p}(\Omega)$$

holds. Moreover, as a consequence of Poincaré's inequality for sufficiently regular domains of finite measure the spaces  $\dot{W}^{m,p}(\Omega)$  and  $W^{m,p}(\Omega)$  actually coincide.

The space  $\dot{W}^{m,p}(\Omega)$  is equipped with the seminorm

$$|u|_{\dot{W}^{m,p}(\Omega)} = \|\nabla^m u\|_{L^p(\Omega)}.$$

Sometimes we will also use the equivalent seminorm  $u \mapsto \sum_{|\alpha|=m} \|D^\alpha u\|_{L^p(\Omega)}$ .

The equivalence of the norm permit to have a useful density result as in [18, Remark 11.28]. Indeed, if  $u \in \dot{W}^{m,p}(\Omega)$  then for every  $\sigma > 0$  there exists  $v \in C^\infty(\Omega) \cap \dot{W}^{m,p}(\Omega)$  such that  $\|u - v\|_{W^{m,p}(\Omega)} \leq \sigma$ .

Let  $\Omega \subset \mathbb{R}^n$  be an open bounded set and let  $E \subset \Omega$  be a Lebesgue measurable set with finite positive measure. Let  $1 \leq p \leq +\infty$  and let  $m \in \mathbb{N}$ . Then, for every

$u \in W^{m,p}(\Omega)$ , there exists a polynomial  $P_E^{m-1}[u]$  of degree  $m - 1$  such that for every multi-index  $\alpha \in \mathbb{R}^n$  with  $0 \leq |\alpha| \leq m - 1$  (see [18, Exercise 13.26]),

$$\int_E \left( D^\alpha u(x) - D^\alpha P_E^{m-1}[u](x) \right) dx = 0. \quad (9)$$

**Theorem 2** (Poincarè inequality in  $W^{m,p}$  [18, Theorem 13.27]) *Let  $m \in \mathbb{N}$ , let  $1 \leq p < +\infty$  and let  $\Omega \subset \mathbb{R}^n$  be an open, bounded and convex set. Then there exists a positive constant  $C = C(m, n, p, \Omega) > 0$  such that,*

$$\sum_{k=0}^{m-1} \|\nabla^k (u - P_\Omega^{m-1}[u])\|_{L^p(\Omega)} \leq C \|\nabla^m u\|_{L^p(\Omega)},$$

for every  $u \in W^{m,p}$  and for every  $k = 0, \dots, m - 1$ .

Notice that for  $m = 1$  the previous Theorem is the classical Poincarè inequality and the polynomial  $P_\Omega[u]$  is the mean of  $u$  over  $\Omega$ . In particular, if  $u \in W^{m,p}(Q')$  with  $Q' = Q'(x_0, \varepsilon)$ , then there exists a unique polynomial  $P_{Q'}^{m-1}[u]$  of degree  $m - 1$  such that (9) holds and there exists a constant  $C = C(n, m, p)$  such that

$$\int_{Q'} |u - P_{Q'}^{m-1}[u]|^p \leq C \varepsilon^{mp} \int_{Q'} |\nabla^m u|^p. \quad (10)$$

Next, we consider the Sobolev–Gagliardo–Nirenberg’s embedding in  $W^{m,p}$  (see Lemma 2.1 in [19]).

Let  $n > mp$ ,  $1 \leq p < \frac{n}{m}$ . Let  $u \in W^{m,p}(Q')$  with  $Q' = Q'(x_0, \varepsilon)$ . Then there exists a unique polynomial  $P_{Q'}^{m-1}[u]$  of degree  $m - 1$  such that (9) holds and there exists a constant  $C = C(n, m, p)$  such that

$$\left( \int_{Q'} |u - P_{Q'}^{m-1}[u]|^{p^*} \right)^{\frac{1}{p^*}} \leq C \left( \frac{1}{\varepsilon^{n-mp}} \int_{Q'} |\nabla^m u|^p \right)^{\frac{1}{p}} \quad (11)$$

where  $p^* = \frac{np}{n-mp}$ .

Moreover, the following easy properties of  $P_\Omega[u]$  holds:

- Linearity:

$$P_\Omega[u](x) + P_\Omega[v](x) = P_\Omega[u + v](x).$$

- Scaling:

$$P_{\varepsilon\Omega}[u](\varepsilon x) = P_\Omega[u_\varepsilon](x),$$

where  $u_\varepsilon(x) := u(\varepsilon x)$ .

We write

$$T_y^m u(x) = \sum_{|\alpha| \leq m} D^\alpha u(y) \frac{(x-y)^\alpha}{\alpha!}$$

for the Taylor polynomial of order  $m$  and

$$R^m u(x, y) = u(x) - T_y^m u(x)$$

for the Taylor remainder of order  $m$ .

## 2.2 Functions of higher-order bounded variation

Let  $\Omega \subset \mathbb{R}^n$  be an open set. A function  $u \in L^1(\Omega)$  is of bounded variation (for short  $u \in BV(\Omega)$ ) if  $u$  has a distributional gradient in form of a Radon measure of finite total mass and write

$$|\nabla u|(\Omega) = \sup \left\{ u \operatorname{div} \varphi : \varphi \in C_0^1(\Omega), \|\varphi\|_{L^\infty} \leq 1 \right\}.$$

We define

$$BV^m(\Omega) = \{u \in W^{m-1,1}(\Omega), \nabla^{m-1}u \in BV(\Omega, S^{m-1}(\mathbb{R}^n))\}$$

the space of (real valued) functions of  $m$ -th order bounded variation, i.e. the set of all functions, whose distributional gradients up to order  $m-1$  are represented through 1-integrable tensor-valued functions and whose  $m$ -th distributional gradient is a tensor-valued Radon measure of finite total variation. Here  $S^k(\mathbb{R}^n)$  denotes the set of all symmetric tensors of order  $k$  with real components, which is naturally isomorphic to the set of all  $k$ -linear symmetric maps  $(\mathbb{R}^n)^k \rightarrow \mathbb{R}$  (see [11]).

It becomes a Banach space with the norm

$$\|u\|_{BV^m(\Omega)} = \|u\|_{W^{m-1,1}(\Omega)} + |\nabla^m u|(\Omega).$$

Here the total variation of  $\nabla^{m-1}u$  is denoted by  $|\nabla^m u|(\Omega)$  and defined by

$$|\nabla^m u|(\Omega) = \sup \left( \sum_{\alpha_1, \dots, \alpha_m=1}^n \int_{\Omega} D_{\alpha_1, \dots, \alpha_{m-1}} u \cdot \partial_{\alpha_m} \varphi_{\alpha_1, \dots, \alpha_m} dx \right),$$

where the supremum is taken over all  $\varphi \in C_0^1(\Omega, \mathbb{R}^n)$  with  $\|\varphi\|_{\infty} = 1$ .

Obviously,  $W^{m,1}(\Omega)$  is a subspace of  $BV^m(\Omega)$ .

The definition of  $BV^m$  generalizes that of the classical space of functions of bounded variation and many results about  $BV$  can be obtained in  $BV^m$  similarly (see [17]). We recall a higher-order variant of the famous Poincaré inequality, which will be useful throughout the sequel:

**Theorem 3** (Poincaré inequality in  $BV^m$  [14, Lemma 2.2]) *Let  $\Omega \subset \mathbb{R}^n$  be an open and bounded subset with Lipschitz boundary,  $m \in \mathbb{N}$ ,  $1 \leq p < \infty$ . Then there exist a constant  $C > 0$ , depending only on  $\Omega$ ,  $m$  and  $n$  such that for all  $u \in BV^m(\Omega)$*

$$\|u\|_{BV^m(\Omega)} \leq C |\nabla^m u|(\Omega).$$

In particular, the following version of Poincaré's inequality holds.

Let  $u \in BV^m(Q')$  with  $Q' = Q'(x_0, \varepsilon)$ , then there exists a unique polynomial  $P_{Q'}^{m-1}[u]$  of degree  $m-1$  such that (9) holds and there exists a constant  $C = C(n, m)$

such that

$$\int_{Q'} |u - P_{Q'}^{m-1}[u]| \leq C \varepsilon^m |\nabla^m u|(Q') \quad (12)$$

By the nature of its definition, the space  $BV^m$  inherits the Poincaré-Wirtinger inequality which can be proved exactly as the corresponding first order result.

Let  $n > m$ ,  $u \in BV^m(Q')$  with  $Q' = Q'(x_0, \varepsilon)$ . Then there exists a unique polynomial  $P_{Q'}^{m-1}[u]$  of degree  $m - 1$  such that (9) holds and there exists a constant  $C = C(n, m)$  such that

$$\left( \int_{Q'} |u - P_{Q'}^{m-1}[u]|^{\frac{n}{n-m}} \right)^{\frac{n-m}{n}} \leq C \frac{1}{\varepsilon^{n-m}} |\nabla^m u|(Q'). \quad (13)$$

We end this subsection with a higher-order variant of the compactness result in  $BV$  (Theorem 3.23 in [3]).

**Proposition 4** (Compactness result in  $BV^m$  [17, Lemma 2.1]) *Let  $\Omega \subset \mathbb{R}^n$  be open and bounded with Lipschitz boundary, and let  $(u_k)_{k=1}^\infty$  be a sequence of  $BV^m$  functions such that*

$$\|u_k\|_{BV^m(\Omega)} \leq M$$

*for some constant  $M > 0$ . Then there is a subsequence  $(u_{k_l})_{l=1}^\infty$  and a function  $u \in BV^m(\Omega)$  such that*

$$\|u - u_{k_l}\|_{W^{m-1,1}(\Omega)} \rightarrow 0 \text{ for } l \rightarrow \infty \text{ and } \|u\|_{BV^m(\Omega)} \leq M.$$

## 2.3 Other useful inequalities

The following tools will be useful in the sequel.

Given  $\delta \in (0, 1)$ , from the convexity of the function  $t \rightarrow |t|^p$  we get for every  $a, b \in \mathbb{R}$

$$|a + b|^p = \left| \frac{1}{(1 + \delta)}(1 + \delta)a + \frac{\delta}{1 + \delta} \frac{1 + \delta}{\delta} b \right|^p \leq (1 + \delta)^p |a|^p + \frac{(1 + \delta)^p}{\delta^p} |b|^p \quad (14)$$

Taking into account (14), we also obtain the following pointwise inequality

$$|a - b|^p \geq \frac{1}{(1 + \delta)^p} |a|^p - \frac{1}{\delta^p} |b|^p \quad (15)$$

for every  $a, b \in \mathbb{R}$ . Given  $\xi, \eta \in \mathbb{R}^n$  it holds

$$\left| |\xi|^p - |\eta|^p \right| \leq p (|\xi| + |\eta|)^{p-1} |\xi - \eta| \quad (16)$$

and, given  $\xi, \eta \in \mathbb{R}^n \setminus \{0\}$  it holds

$$\left| \frac{\xi}{|\xi|} - \frac{\eta}{|\eta|} \right| \leq 2 \frac{|\xi - \eta|}{|\xi|}. \quad (17)$$

## 2.4 The local version of the functional $K_\varepsilon(f, m, p)$

We define the following local counterpart of (2) which will be use in Step 3 of proof of Theorem 1

$$K_\varepsilon(f, m, p, \Omega) = \varepsilon^{n-mp} \sup_{\mathcal{G}_\varepsilon} \sum_{Q' \in \mathcal{G}_\varepsilon} \int_{Q'} |f(x) - P_{Q'}^{m-1}[f](x)|^p dx, \quad (18)$$

where the supremum on the right hand side is taken over all families  $\mathcal{G}_\varepsilon$  of disjoint open cubes of sidelength  $\varepsilon$  and arbitrary orientation contained in  $\Omega$ .

This quantity is strictly related to the  $L^p$  norm of  $\nabla^m f$ . Indeed, for  $p < \frac{n}{m}$  with  $p^\star = \frac{np}{n-mp}$ , by using Hölder inequality, we have

$$\|f\|_{L^p(Q)} \leq \|f\|_{L^{p^\star}(Q)} \quad (19)$$

Thus, there exists a constant  $C$  depending only on  $Q, m, p$  such that for  $Q' = \varepsilon Q + x_0$ , by (19) and (11), we get

$$\varepsilon^{n-mp} \int_{Q'} |f(x) - P_{Q'}^{m-1}[f]|^p dx \leq C \int_{Q'} |\nabla^m f|^p. \quad (20)$$

Summing over all sets  $Q'$  in  $\mathcal{G}_\varepsilon$ , we obtain

$$\varepsilon^{n-mp} \sum_{Q' \in \mathcal{G}_\varepsilon} \int_{Q'} |f(x) - P_{Q'}^{m-1}[f]|^p dx \leq C \|\nabla^m f\|_{L^p(\Omega)}^p$$

and therefore

$$K_\varepsilon(f, m, p, \Omega) \leq C \|\nabla^m f\|_{L^p(\Omega)}^p.$$

We conclude this subsection, by observing that if  $\bar{v} \in \mathbb{S}^{N-1}$  is a vector maximizing the integral in (8),  $x_0 \in \mathbb{R}^n$  and  $Q_\eta(x_0)$  is a cube of side length  $\eta$  with center in  $x_0$  then

$$\frac{1}{(m!)^p} \int_{Q_\eta(x_0)} \left| (x - x_0)^m \cdot \bar{v} - \int_{Q_\eta(x_0)} (y - x_0)^m \cdot \bar{v} dy \right|^p dx = \beta(n, m, p) \cdot \eta^{n+mp}. \quad (21)$$

## 3 The case $m = 2$

In this section we deal with the case  $m = 2$ . In this case it is easier to make some explicit computations. Moreover we give an estimates on the constant  $\beta(n, 2, p)$  in terms of the Laplacian of the function  $f \in W^{2,p}$ .

We prove the following

**Proposition 5** *Let  $f \in W^{2,p}$  and  $\beta(n, 2, p)$  as in (8). Then the following estimate from below holds true*

$$\beta(n, 2, p) \geq C_{n,p} |\Delta f(0)|^p. \quad (22)$$



First, by virtue of (9), it is possible to characterize  $P_\Omega[u]$  for  $m = 2$ . Fixed  $x_0 \in \Omega$ , a generic polynomial of degree 1 centered in  $x_0$  is given by

$$P_\Omega^1[u](x) = \langle a, x - x_0 \rangle + b, \quad a \in \mathbb{R}^n, b \in \mathbb{R}.$$

By (9) with  $|\alpha| = 0$ , we have

$$b|\Omega| = \int_\Omega (u(x) - \langle a, x - x_0 \rangle) dx$$

which implies

$$b = \int_\Omega (u(x) - \langle a, x - x_0 \rangle) dx.$$

Moreover, for every  $i = 1, \dots, n$ , again (9) for  $|\alpha| = 1$  gives

$$a_i = \int_\Omega \frac{\partial u}{\partial x_i}(x) dx$$

and we write

$$a = \int_\Omega \nabla u(x) dx.$$

Then the polynomial  $P_\Omega^1(u)$  is

$$P_\Omega^1[u](x) = \int_\Omega \left( u(y) - \left\langle \int_\Omega \nabla u, y \right\rangle \right) dy + \left\langle \int_\Omega \nabla u(y) dy, x - x_0 \right\rangle \quad (23)$$

where, with a slight abuse of notation, we mean

$$\left\langle \int_\Omega \nabla u(y) dy, x - x_0 \right\rangle = \sum_{j=1}^n (x_j - x_{0j}) \int_\Omega \frac{\partial u}{\partial y_j}(y) dy.$$

*Remark 1* We observe that if  $\Omega$  is symmetric with respect to  $x_0$ , the polynomial  $P_\Omega^1[u]$  has a simpler form, indeed

$$\int_\Omega \left\langle \int_\Omega \nabla u, y \right\rangle dy = 0,$$

and then

$$P_\Omega^1[u](x) = \int_\Omega u(y) dy + \left\langle \int_\Omega \nabla u(y) dy, x - x_0 \right\rangle \quad (24)$$

*Proof of Proposition 5* We observe that when  $m = 2$ ,  $p \geq 1$ , (8) reads as

$$\beta(n, 2, p) := \max_{v \in \mathbb{S}^{n^2-1}} \frac{1}{4} \int_Q \left| v \cdot x^2 - \int_Q v \cdot y^2 dy \right|^p dx. \quad (25)$$

In this case  $v \cdot x^2$  can equivalently be write as

$$\langle Ax, x \rangle$$

10 *A BMO-type characterization of higher order Sobolev spaces*

where  $A \in \mathcal{M}(n)$  is a matrix  $n \times n$  and  $\langle \cdot, \cdot \rangle$  denote the usual scalar product in  $\mathbb{R}^n$ .

It is worth to remark that

$$\beta(n, 2, p) \geq \frac{1}{2^p} \int_Q \left| \langle \nabla^2 f(0)x, x \rangle - \int_Q \langle \nabla^2 f(0)y, y \rangle \right|^p dx. \tag{26}$$

Firstly we observe that denoting by  $e_i$  the canonical basis of  $\mathbb{R}^n$ , by  $O \in O(n)$  an orthogonal matrix and by  $\mathcal{R} \in SO(n)$  a rotation around the origin taking  $O^{-1}(Q)$  into  $Q$  we have

$$\int_{O^{-1}(Q)} y_i^2 dy = \int_{O^{-1}(Q)} (y \cdot e_i)^2 dy = \int_{\mathcal{R} \circ O^{-1}(Q)} (\mathcal{R}w \cdot e_i)^2 dw = \int_Q (w \cdot \mathcal{R}^{-1} e_i)^2 dy = \frac{1}{12}$$

Moreover, given  $A \in \mathcal{S}(n)$  a symmetric matrix there exist  $O \in O(n)$  and  $D \in \mathcal{D}(n)$  such that  $A = ODO^{-1}$ . Thus we have

$$\begin{aligned} \int_Q \langle Az, z \rangle dz &= \int_Q \langle (ODO^{-1})z, z \rangle dz = \int_Q \langle (DO^{-1})z, O^{-1}z \rangle dz = \int_{O^{-1}(Q)} \langle Dy, y \rangle dy \\ &= \int_{O^{-1}(Q)} \sum_{i=1}^n \lambda_i y_i^2 dy = \sum_{i=1}^n \lambda_i \int_{O^{-1}(Q)} y_i^2 dy = \frac{1}{12} \sum_{i=1}^n \lambda_i \end{aligned} \tag{27}$$

Then we can estimate from below  $\beta(n, 2, p)$  using (26) and (27), proving (22).

Indeed, setting  $\nabla^2 f(0) = A$  we have

$$\int_Q \langle Ax, x \rangle = \frac{\Delta f(0)}{12}.$$

Moreover setting  $\bar{y} = \min y_i$ , we have

$$\begin{aligned} \frac{1}{2^p} \int_Q \left| \langle Ax, x \rangle - \int_Q \langle Ay, y \rangle \right|^p dx &= \frac{1}{2^p} \int_Q \left| \langle (DO^{-1})x, O^{-1}x \rangle - \frac{\Delta f(0)}{12} \right|^p dx \\ &= \frac{1}{2^p} \int_Q \left| \sum \lambda_i y_i^2 - \frac{\Delta f(0)}{12} \right|^p dx \geq \frac{1}{2^p} \int_{O^{-1}(Q)} \left| \sum \lambda_i \bar{y}^2 - \frac{\Delta f(0)}{12} \right|^p dx \\ &= \frac{1}{2^p} |\Delta f(0)|^p \int_{O^{-1}(Q)} \left| \bar{y} - \frac{1}{12} \right|^p dx = C_{n,p} |\Delta f(0)|^p. \end{aligned} \tag{28}$$

□

## 4 A characterization of $W^{m,p}$

*Proof of Theorem 1* We divide the proof in three steps, proving first the limsup and liminf inequalities in (7) and then the validity of (6).

As a starting point we fix a bounded open set  $\Omega \subset \mathbb{R}^n$  and  $f \in W^{m,p}(\Omega)$ . Given  $\sigma > 0$ , there exists a function  $g \in C_c^\infty(\Omega)$  such that  $\|f - g\|_{W^{m,p}(\Omega)} < \sigma$  and we choose  $\varepsilon > 0$  such that

$$|\nabla^m g(x) - \nabla^m g(y)| \leq \sigma, \quad \forall x, y, \quad |x - y| \leq \frac{\sqrt{n}\varepsilon}{2} \tag{29}$$

Let us take now a family  $\mathcal{G}_\varepsilon$  of disjoint open cubes  $Q'$  of side length  $\varepsilon$  and arbitrary orientation and let us denote by  $\mathcal{G}'_\varepsilon$  the subfamily of  $\mathcal{G}_\varepsilon$  made by all cubes  $Q' \in \mathcal{G}_\varepsilon$  such that  $Q' \subset \Omega$ .

### Step1 (limsup inequality).

We are going to show that

$$\limsup_{\varepsilon \rightarrow 0^+} K_\varepsilon(f, m, p) \leq \beta(n, m, p) \int_{\mathbb{R}^n} |\nabla^m f|^p dx.$$

We may assume, without loss of generality, that  $|\nabla^m f| \in L^p(\Omega)$ . Using (14) and the linearity of  $P_{Q'}^{m-1}[f]$ , for any  $Q' \in \mathcal{G}'_\varepsilon$  we have:

$$\int_{Q'} |f - P_{Q'}^{m-1}[f]|^p dx \leq (1 + \delta)^p \int_{Q'} |g - P_{Q'}^{m-1}[g]|^p dx + M_\delta \int_{Q'} |(f - g) - P_{Q'}^{m-1}[f - g]|^p dx \quad (30)$$

where  $M_\delta = (1 + \delta)^p / \delta^p$ .

We recall the notation in Section 2, so denoting by  $x_0$  the center of the cube  $Q'$  and for all  $x \in Q'$  we write

$$g(x) = T_{x_0}^m g(x) + R^m g(x, x_0),$$

where  $|R^m g(x, y)| < (n^{\frac{m}{2}} \sigma \varepsilon^m) / 2^m = C_1 \sigma \varepsilon^m$ .

We now estimate the two terms in (30). Let us focus on the first addendum: using again (14) we have

$$\begin{aligned} & \int_{Q'} |g - P_{Q'}^{m-1}[g]|^p dx \\ &= \int_{Q'} \left| \frac{1}{m!} \nabla^m g(x_0) \cdot (x - x_0)^m + R^m g(x, x_0) - \left[ \int_{Q'} \frac{1}{m!} \nabla^m g(x_0) \cdot (y - x_0)^m dy + \int_{Q'} R^m g(y, x_0) dy \right] \right|^p dx \\ &\leq (1 + \delta)^p \frac{1}{(m!)^p} \int_{Q'} \left| \nabla^m g(x_0) \cdot (x - x_0)^m - \int_{Q'} \nabla^m g(x_0) \cdot (y - x_0)^m dy \right|^p dx + 2^p M_\delta \int_{Q'} |R^m g(x, x_0)|^p dx \\ &\leq (1 + \delta)^p \beta(n, m, p) \varepsilon^{mp} |\nabla^m g(x_0)|^p + C_2 M_\delta \sigma^p \varepsilon^{mp}. \end{aligned} \quad (31)$$

Moreover, applying again (14) and (29) we have

$$|\nabla^m g(x_0)|^p \leq (1 + \delta)^p \int_{Q'} |\nabla^m g(x)|^p dx + C_3 M_\delta \sigma^p.$$

Hence

$$\int_{Q'} |g - P_{Q'}^{m-1}[g]|^p dx \leq \beta(n, m, p) (1 + \delta)^{2p} \varepsilon^{mp} \int_{Q'} |\nabla^m g(x)|^p dx + C_4 M_\delta \varepsilon^{mp} \sigma^p. \quad (32)$$

Let us focus now on the second addendum in (30). By Poincaré inequality in  $W^{m,p}$  (see Theorem 2), we have

$$\int_{Q'} |(f - g) - P_{Q'}^{m-1}[f - g]|^p dx \leq C_p \varepsilon^{mp-n} \int_{Q'} |\nabla^m (f - g)|^p dx \quad (33)$$

where  $C_p$  is the Poincaré constant for cubes.

Observe now that  $\#\{\mathcal{G}'_\varepsilon\} \leq \varepsilon^{-n} |\Omega|$  and set  $\Omega_\varepsilon = \{x \in \Omega : \text{dist}(x, \partial\Omega) > \varepsilon \sqrt{n}\}$ . Using (30), (32) and (33) we have

$$\begin{aligned} & \varepsilon^{n-mp} \sum_{Q' \in \mathcal{G}'_\varepsilon} \int_{Q'} |f - P_{Q'}^{m-1}[f]|^p dx \\ &\leq \varepsilon^{n-mp} \sum_{Q' \in \mathcal{G}'_\varepsilon} \int_{Q'} |f - P_{Q'}^{m-1}[f]|^p dx + C_6 \sum_{Q' \in \mathcal{G}'_\varepsilon \setminus \mathcal{G}'_\varepsilon} \int_{Q'} |\nabla^m f|^p \\ &\leq (1 + \delta)^p \varepsilon^{n-mp} \sum_{Q' \in \mathcal{G}'_\varepsilon} \int_{Q'} |g - P_{Q'}^{m-1}[g]|^p dx + C_p M_\delta \int_{\Omega} |\nabla^m (f - g)|^p + C_6 \int_{\mathbb{R}^n \setminus \Omega_\varepsilon} |\nabla^m f|^p dx \\ &\leq (1 + \delta)^{3p} \beta(n, m, p) \sum_{Q' \in \mathcal{G}'_\varepsilon} \int_{Q'} |\nabla^m g(x)|^p dx + C_4 M_\delta \varepsilon^n \sigma^p + C_p M_\delta \sigma^p + C_6 \int_{\mathbb{R}^n \setminus \Omega_\varepsilon} |\nabla^m f|^p dx \\ &\leq (1 + \delta)^{3p} \beta(n, m, p) \int_{\Omega} |\nabla^m f(x)|^p dx + C_4 M_\delta \varepsilon^n \sigma^p + C_p M_\delta \sigma^p + C_6 \int_{\mathbb{R}^n \setminus \Omega_\varepsilon} |\nabla^m f|^p dx \end{aligned} \quad (34)$$

12 *A BMO-type characterization of higher order Sobolev spaces*

where the constants depend only on  $n$ ,  $p$  and  $|\Omega|$ . Then, taking the supremum over all the families of cubes  $\mathcal{G}_\varepsilon$ , and then letting first  $\varepsilon \rightarrow 0^+$ ,  $\sigma \rightarrow 0$ ,  $\delta \rightarrow 0$  and  $\Omega \uparrow \mathbb{R}^n$  we conclude.

**Step2 (liminf inequality).** We fix  $\Omega \subset \mathbb{R}^n$ , we assume again that  $f \in W_{loc}^{m,p}(\Omega)$  and we fix  $\sigma > 0$  and  $g \in C_c^\infty(\Omega)$  as in the previous Step. We prove that

$$\liminf_{\varepsilon \rightarrow 0^+} K_\varepsilon(f, m, p) \geq \beta(n, m, p) \int_{\mathbb{R}^n} |\nabla^m f|^p dx. \quad (35)$$

So, for  $\eta \in (0, 1)$  we consider the set

$$U_\eta = \{x \in \Omega : |\nabla^m g(x)| > \eta\}$$

With a clever use of Lemma 2.95 of [3] (as in Proposition 3.6 of [15]) it is possible to find  $k$  sufficiently small pairwise disjoint open sets  $S_j \subset \mathbb{S}^{N-1}$  covering  $\mathbb{S}^{N-1}$ . Precisely,

$$\bigcup_{j=1}^k \bar{S}_j = \mathbb{S}^{N-1}$$

$$\text{diam } S_j < \eta \text{ for all } j = 1 \dots k$$

$$\left| \bigcup_{j=1}^k \left\{ x \in U_\eta : \frac{\nabla^m g(x)}{|\nabla^m g(x)|} \in \partial S_j \right\} \right| = 0.$$

For all  $j = 1, \dots, k$  we denote

$$A_j = \left\{ x \in U_\eta : \frac{\nabla^m g(x)}{|\nabla^m g(x)|} \in S_j \right\},$$

which are open sets with the property

$$\left| U_\eta \setminus \bigcup_{j=1}^k A_j \right| = 0. \quad (36)$$

For  $\varepsilon > 0$  we consider the family  $\mathcal{F}_\varepsilon$  of all open cubes with faces parallel to the coordinate planes, side length  $\varepsilon$ , centered at all points of the form  $\varepsilon v$ , with  $v \in \mathbb{Z}^n$ . Then for all  $j = 1, \dots, k$  we choose  $M_j \in S_j$  and we denote by  $R_j \in SO(n)$  a rotation that takes  $e_1$  into  $M_j$ .

Note that in this way, denoting by  $x'$  the center of the cube  $Q' \in \mathcal{F}_\varepsilon$ , we have (see (21)),

$$\frac{1}{(m!)^p} \int_{R_j(Q')} \left| (x - x')^m \cdot \bar{v} - \int_{R_j(Q')} (y - x')^m \cdot \bar{v} dy \right|^p dx = \beta(n, m, p) \cdot \varepsilon^{n+mp}.$$

For all  $j = 1, \dots, k$  we denote by  $R_j(Q_{h,j})$ ,  $Q_{h,j} \in \mathcal{F}_\varepsilon$ ,  $h = 1, \dots, m_j$ , the elements of  $\mathcal{G}_\varepsilon$  contained in  $A_j$ . By (36) there exists  $\varepsilon(\sigma, \eta)$  such that if  $\varepsilon < \varepsilon(\sigma, \eta)$  then

$$\left| U_\eta \setminus \bigcup_{j=1}^k \bigcup_{h=1}^{m_j} \mathcal{R}_j(Q_{h,j}) \right| \leq \eta^p.$$

We denote by  $x_{h,j}$  the center of the cube  $\mathcal{R}_j(Q_{h,j})$  and we argue as in Step 1. Indeed we have

$$\begin{aligned}
& \int_{R_j(Q_{h,j})} \left| g - P_{R_j(Q_{h,j})}^{m-1}[g] \right|^p dx \\
& \geq \frac{1}{(1+\delta)^p} \frac{1}{(m!)^p} \int_{R_j(Q_{h,j})} \left| \nabla^m g(x_{h,j}) \cdot (x - x_{h,j})^m - \int_{R_j(Q_{h,j})} \nabla^m g(x_{h,j}) \cdot (x - x_{h,j})^m \right|^p dx \\
& \quad - \frac{2^p}{\delta^p} \int_{R_j(Q_{h,j})} |R^m g(x, x_{h,j})|^p dx \\
& \geq \frac{1}{(1+\delta)^{2p}} \frac{|\nabla^m g(x_{h,j})|^p}{(m!)^p} \int_{R_j(Q_{h,j})} \left| M_j \cdot (x - x_{h,j})^m - \int_{R_j(Q_{h,j})} M_j \cdot (x - x_{h,j})^m \right|^p dx \\
& \quad - \frac{2^p}{\delta^p} \frac{|\nabla^m g(x_{h,j})|^p}{(m!)^p} \int_{R_j(Q_{h,j})} \left| (\nabla^m g(x_{h,j}) - M_j) \cdot (x - x_{h,j})^m - \int_{R_j(Q_{h,j})} (\nabla^m g(x_{h,j}) - M_j) \cdot (x - x_{h,j})^m \right|^p dx \\
& \quad - C_7 \frac{\sigma^p \varepsilon^{mp}}{\delta^p} \\
& \geq \frac{\varepsilon^{mp} \beta(n, m, p) |\nabla^m g(x_{h,j})|^p}{(1+\delta)^{2p}} - \frac{C_8 \eta^p \varepsilon^{mp}}{\delta^p} \|\nabla^m g\|_{L^\infty}^p - C_7 \frac{\sigma^p \varepsilon^{mp}}{\delta^p}.
\end{aligned} \tag{37}$$

Now, adding on  $j$  and  $h$  the previous inequality, recalling (36), we have

$$\begin{aligned}
& \varepsilon^{n-mp} \sum_{R_j(Q_{h,j}) \in \mathcal{G}_\varepsilon} \int_{R_j(Q_{h,j})} \left| g - P_{Q'}^{m-1}[g] \right|^p dx \\
& \geq \varepsilon^{n-mp} \sum_{j=1}^k \sum_{h=1}^{m_j} \frac{\varepsilon^{mp} \beta(n, m, p) |\nabla^m g(x_{h,j})|^p}{(1+\delta)^{2p}} - \frac{C_8 \eta^p \varepsilon^{mp}}{\delta^p} \|\nabla^m g\|_{L^\infty}^p - C_7 \frac{\sigma^p \varepsilon^{mp}}{\delta^p} \\
& \geq \frac{\beta(n, m, p)}{(1+\delta)^{3p}} \sum_{j=1}^k \sum_{h=1}^{m_j} \int_{R_j(Q_{h,j})} |\nabla^m g|^p - \frac{C_8 \eta^p \varepsilon^n}{\delta^p} \|\nabla^m g\|_{L^\infty}^p - C_7 \frac{\sigma^p \varepsilon^n}{\delta^p} \\
& \geq \frac{\beta(n, m, p)}{(1+\delta)^{3p}} \int_{\Omega} |\nabla^m g|^p - \frac{C_7 \eta^p}{\delta^p} (1 + \|\nabla^m g\|_{L^\infty}^p) - C \frac{\sigma^p}{\delta^p} \\
& \geq \frac{\beta(n, m, p)}{(1+\delta)^{4p}} \int_{\Omega} |\nabla^m f|^p - \frac{C_7 \eta^p}{\delta^p} (1 + \|\nabla^m g\|_{L^\infty}^p) - C \frac{\sigma^p}{\delta^p},
\end{aligned}$$

where the constants may change from line to line and depend only on  $p$ ,  $n$  and  $|\Omega|$ . We conclude choosing  $\eta$  small enough and consequently  $\varepsilon$  small,

$$\begin{aligned}
& \varepsilon^{n-mp} \sum_{Q' \in \mathcal{G}_\varepsilon} \int_{Q'} \left| f - P_{Q'}^{m-1}[f] \right|^p dx \\
& \geq \frac{1}{(1+\delta)^p} \varepsilon^{n-mp} \sum_{Q' \in \mathcal{G}_\varepsilon} \int_{Q'} \left| g - P_{Q'}^{m-1}[g] \right|^p dx - \frac{1}{\delta^p} \int_{\Omega} |\nabla^m(f - g)|^p \\
& \geq \frac{\beta(n, m, p)}{(1+\delta)^{5p}} \int_{\Omega} |\nabla^m f|^p - \frac{C \sigma^p}{\delta^p},
\end{aligned}$$

where again  $C$  may change from line to line and depend on  $p$ ,  $n$  and  $|\Omega|$ . To conclude we take the supremum over all the families  $\mathcal{G}_\varepsilon$  and let first  $\varepsilon \rightarrow 0$ ,  $\sigma \rightarrow 0$ ,  $\delta \rightarrow 0$  and  $\Omega \uparrow \mathbb{R}^n$ , proving (35).

**Step3 (proof of (6))** Now let  $p > 1$ ,  $f \in W_{\text{loc}}^{m-1,p}(\mathbb{R}^n)$  and  $\liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, p) < \infty$ . We fix  $\sigma > 0$ ,  $\Omega \subset \mathbb{R}^n$  and observe that there exist  $r > 0$  and a finite family of pairwise disjoint

open cubes  $Q(x_i, r)$  such that

$$\left| \Omega \setminus \bigcup_{i=1}^m Q(x_i, r) \right| < \sigma. \tag{38}$$

$$|\nabla^m f(x) - \nabla^m f(y)| < \sigma \tag{39}$$

Moreover we fix  $0 < \varepsilon < r$  and we set  $f_\varepsilon(x) = (\varrho_\varepsilon * f)(x)$ , where  $\varrho$  is a standard mollifier with compact support in the unit cube  $Q$  and  $\varrho_\varepsilon(x) = \varepsilon^{-n}\varrho(x/\varepsilon)$ .

For every  $Q(x_i, r)$  we consider a family  $\mathcal{H}_\varepsilon$  of pairwise disjoint cubes  $Q_j = z_j + \varepsilon Q \subset Q(x_i, r)$ , for  $j = 1, \dots, k$ .

We compute now

$$\begin{aligned} |\nabla^m f_\varepsilon(z_j)|^p &= \left| \int_{\mathbb{R}^n} f(y) \nabla^m \rho_\varepsilon(z_j - y) dy \right|^p = \left| \int_{\mathbb{R}^n} (f(y) - P_{Q_j}^{m-1}[f](y)) \nabla^m \rho_\varepsilon(z_j - y) dy \right|^p \\ &\leq \varepsilon^{(-m-n)p+n} \int_{Q_j} |f(y) - P_{Q_j}^{m-1}[f](y)|^p dy = \varepsilon^{-mp} \int_{Q_j} |f(y) - P_{Q_j}^{m-1}[f](y)|^p dy. \end{aligned}$$

Moreover, by (29) and (14), we have

$$|\nabla^m f_\varepsilon(z_j)|^p \geq \frac{1}{1 + \delta} \varepsilon^{-n} \int_{Q_j} |\nabla^m f_\varepsilon(x)|^p dx - \frac{C}{\delta^p} \sigma^p$$

Then

$$\frac{1}{1 + \delta} \int_{Q_j} |\nabla^m f_\varepsilon(x)|^p dx \leq \varepsilon^{n-mp} \int_{Q_j} |f(y) - P_{Q_j}^{m-1}[f](y)|^p dy + \frac{C}{\delta^p} \sigma^p \varepsilon^n.$$

Summing up all the cubes in  $\mathcal{H}_\varepsilon$ , we obtain

$$\begin{aligned} &\frac{1}{1 + \delta} \int_{Q(x_i, r)} |\nabla^m f_\varepsilon(x)|^p dx \\ &\leq \frac{1}{1 + \delta} \sum_{j=1}^k \int_{Q_j} |\nabla^m f_\varepsilon(x)|^p dx \leq \varepsilon^{n-mp} \sum_{j=1}^k \int_{Q_j} |f(y) - P_{Q_j}^{m-1}[f](y)|^p dy + \frac{C}{\delta^p} \sigma^p \varepsilon^n \\ &\leq \varepsilon^{n-mp} \sum_{j=1}^k \int_{Q_j} |f(y) - P_{Q_j}^{m-1}[f](y)|^p dy + \frac{C}{\delta^p} \sigma^p r^n, \tag{40} \end{aligned}$$

where the last inequality follows since  $ke^n \leq r^n$ . Taking the supremum with respect to all families  $\mathcal{H}_\varepsilon$  and the liminf with respect to  $\varepsilon$ , we have

$$\frac{1}{1 + \delta} \int_{Q(x_i, r)} |\nabla^m f(x)|^p dx \leq \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, p, Q(x_i, r)) + \frac{C}{\delta^p} \sigma^p r^n.$$

Summing up with respect to  $i$  and using (38) we have

$$\frac{1}{1 + \delta} \int_{\Omega} |\nabla^m f(x)|^p dx \leq \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, p, \Omega) + \frac{C}{\delta^p} \sigma^p |\Omega|.$$

Letting  $\sigma \rightarrow 0, \delta \rightarrow 0$  and  $\Omega \uparrow \mathbb{R}^n$ , we conclude. □

*Remark 2* We observe that Theorem 7 hold also in an open set  $\Omega$  with the same proof replacing  $K_\varepsilon(f, m, p)$  by the quantity  $K_\varepsilon(f, m, p, \Omega)$  defined in (18).

**Corollary 6** *Let  $p > 1, n > mp, p^* = \frac{np}{n-mp}, \Omega \subset \mathbb{R}^n$  and  $\mathcal{G}_\varepsilon$  a pairwise disjoint family of translations  $Q'$  of  $\varepsilon Q$  contained in  $\Omega$ . Then, the following three statements are equivalent:*

i)  $f \in W^{m,p}(\Omega)$ ;

ii)

$$\sup_{\mathcal{G}_\varepsilon} \sum_{Q' \in \mathcal{G}_\varepsilon} \varepsilon^{n-mp} \int_{Q'} |f - P_{Q'}^{m-1}[f]|^p < +\infty;$$

iii)

$$\sup_{\mathcal{G}_\varepsilon} \sum_{Q' \in \mathcal{G}_\varepsilon} \|f - P_{Q'}^{m-1}[f]\|_{L^{p^*}(Q')}^p < +\infty$$

*Proof* In this proof the constant  $C$  may change from line to line.

We prove that *iii*)  $\Rightarrow$  *ii*). By Hölder's inequality it holds

$$\varepsilon^{n-mp} \int_{Q'} |f - P_{Q'}^{m-1}[f]|^p dx \leq \frac{\varepsilon^{n-mp}}{\varepsilon^n} \left( \int_{Q'} |f - P_{Q'}^{m-1}[f]|^{\frac{np}{n-mp}} \right)^{\frac{n-mp}{n}} |Q'|^{\frac{mp}{n}} = \|f - P_{Q'}^{m-1}[f]\|_{L^{p^*}(Q')}^p. \quad (41)$$

Summing over all sets  $Q'$  in  $\mathcal{G}_\varepsilon$  and passing to the supremum, we conclude.

We prove that *i*)  $\Rightarrow$  *iii*). Using the Sobolev-Gagliardo-Nirenberg inequality (11), we obtain that there exists a constant  $C = C(n, m, p)$  such that

$$\|f - P_{Q'}^{m-1}[f]\|_{L^{p^*}(Q')} \leq C \|\nabla^m f\|_{L^p}. \quad (42)$$

Summing over  $Q'$  in  $\mathcal{G}_\varepsilon$  and passing to the supremum over all families  $G_\varepsilon$  the proof is completed.

The equivalence *i*)  $\Leftrightarrow$  *ii*) is proved in [9]. □

## 5 A characterization of higher order bounded variation

In this section we deal with the case  $p = 1$ . This case is not included in Theorem 1 since (6) hold only for  $p > 1$ .

The case  $m = 1$  was treated in [16]. They proved that (see Proposition 2.4 of [16]) if  $f \in L^1_{loc}(\mathbb{R}^n)$  then

$$f \in BV(\mathbb{R}^n) \iff \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, 1, 1) < +\infty \quad (43)$$

Precisely, they prove that for  $f \in L^1_{loc}(\mathbb{R}^n)$  it holds

$$\frac{1}{4} |\nabla f|(\mathbb{R}^n) \leq \liminf_{\varepsilon \rightarrow 0^+} K_\varepsilon(f, 1, 1) \leq \limsup_{\varepsilon \rightarrow 0^+} K_\varepsilon(f, 1, 1) \leq \frac{1}{2} |\nabla f|(\mathbb{R}^n),$$

where the total variation of  $f$  in  $\Omega \subset \mathbb{R}^n$ , possibly equal to  $+\infty$ , is defined by setting

$$|\nabla f|(\Omega) := \sup \left\{ \int_{\Omega} f(x) \operatorname{div} \varphi(x) dx \quad : \quad \varphi \in C^1_c(\Omega), \|\varphi\|_\infty \leq 1 \right\}$$

We prove a similar characterization for the case  $m > 1$ . Now an equivalence similar to (43) involve the space  $BV^m(\mathbb{R}^n)$  of functions of  $m$ -th order bounded variation (see Section 2).

Precisely, we prove the following

**Proposition 7** *Let  $f \in W_{loc}^{m-1,1}(\mathbb{R}^n)$ . Then*

$$f \in BV^m(\mathbb{R}^n) \iff \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, 1) < +\infty$$

*Moreover, there is a positive constants  $C$ , independent of  $f$ , such that*

$$|\nabla^m f|(\mathbb{R}^n) \leq \liminf_{\varepsilon \rightarrow 0^+} K_\varepsilon(f, m, 1) \leq \limsup_{\varepsilon \rightarrow 0^+} K_\varepsilon(f, m, 1) \leq C|\nabla^m f|(\mathbb{R}^n). \quad (44)$$

*Proof* To prove the first inequality in (44) we argue as in Step 3 of Theorem 1. In particular, we have

$$\frac{1}{1+\delta} \int_{Q(x_i, r)} |\nabla^m f_\varepsilon(x)| dx \leq \varepsilon^{n-m} \sum_{j=1}^k \int_{Q_j} |f(y) - P_{Q_j}^{m-1}[f](y)| dy + \frac{C}{\delta} \sigma r^n, \quad (45)$$

Taking the supremum with respect to all families  $\mathcal{H}_\varepsilon$  and the liminf with respect to  $\varepsilon$ , we have

$$\frac{1}{1+\delta} \liminf_{\varepsilon \rightarrow 0} \int_{Q(x_i, r)} |\nabla^m f_\varepsilon(x)| dx \leq \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, Q(x_i, r)) + \frac{C}{\delta} \sigma r^n.$$

By the compactness in  $BV^m$  (Proposition 4), we get

$$\frac{1}{1+\delta} |\nabla^m f|(Q(x_i, r)) dx \leq \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, Q(x_i, r)) + \frac{C}{\delta} \sigma r^n$$

Summing up with respect to  $i$  and using (38) we obtain

$$\frac{1}{1+\delta} |\nabla^m f|(\Omega) dx \leq \liminf_{\varepsilon \rightarrow 0} K_\varepsilon(f, m, \Omega) + \frac{C}{\delta} \sigma r^n |\Omega|$$

We conclude letting  $\sigma \rightarrow 0$ ,  $\delta \rightarrow 0$ ,  $\Omega \uparrow \mathbb{R}^n$ .

In order to prove the estimate from above in (44), it is sufficient to apply the Poincaré inequality in  $BV^m$  (see Section 2). □

**Corollary 8** *Let  $n > m$ ,  $1^* = \frac{n}{n-m}$ ,  $\Omega \subset \mathbb{R}^n$  and  $\mathcal{G}_\varepsilon$  is any pairwise disjoint family of translations  $Q'$  of  $\varepsilon Q$  contained in  $\Omega$ . Then, the following three statements are equivalent:*

i)  $f \in BV^m(\Omega)$ ;

ii)

$$\sup_{\mathcal{G}_\varepsilon} \sum_{Q' \in \mathcal{G}_\varepsilon} \varepsilon^{n-m} \int_{Q'} |f - P_{Q'}^{m-1}[f]| < +\infty$$

iii)

$$\sup_{\mathcal{G}_\varepsilon} \sum_{Q' \in \mathcal{G}_\varepsilon} \|f - P_{Q'}^{m-1}[f]\|_{L^{1^*}(Q')} < +\infty$$

*Proof* We prove that iii)  $\Rightarrow$  ii). By Hölder's inequality it holds

$$\varepsilon^{n-m} \int_{Q'} |f - P_{Q'}^{m-1}[f]| dx \leq \|f - P_{Q'}^{m-1}[f]\|_{L^{1^*}(Q')}. \quad (46)$$



The conclusion follows by summing over all sets  $Q'$  in  $\mathcal{G}_\varepsilon$ .

We prove that  $i) \Rightarrow iii)$ . By using (13) there exists a constant  $C = C(n,m)$  such that

$$\|f - P_{Q'}^{m-1}[f]\|_{L^{1^*}(Q')} \leq C \|\nabla^m f\|_{L^p}(Q') \quad (47)$$

The conclusion follows again by summing over all sets  $Q'$  in  $\mathcal{G}_\varepsilon$ .

The equivalence  $i) \Leftrightarrow ii)$  is proved in [9]. □

## Declarations

- Funding (The authors are members of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of INdAM. The research of S.G.L.B. has been funded by PRIN Project 2017AYM8XW and the research of R.S. has been funded by PRIN Project 2017JFFHSH.)
- Conflict of interest/Competing interests (There is no conflict of interest)
- Ethics approval (Not applicable)
- Consent to participate (Not applicable)
- Consent for publication (The Authors consent for publication)
- Availability of data and materials (Not applicable)
- Code availability (Not applicable)
- Authors' contributions (The Authors contributed equally to this work.)

## References

- [1] L. AMBROSIO, J. BOURGAIN, H. BREZIS, A. FIGALLI, *BMO-type norms related to the perimeter of sets*, Comm. Pure Appl. Math., 69 (2016), 1062–1086.
- [2] L. AMBROSIO, G. COMI, *Anisotropic Surface Measures as Limits of Volume Fractions*. Current Research in Nonlinear Analysis 135 (2018), 1-32.
- [3] L. AMBROSIO, N. FUSCO, D. PALLARA, *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [4] B. BOJARSKI, L. IHNATSYEVA, J. KINNUNEN, *How to recognize polynomials in higher order Sobolev spaces*, Math. Scand. 112 (2013), 161-181.
- [5] R. Borghol, *Some properties of Sobolev spaces*, Asymptot. Anal. 51 (2007), no.3-4, 303-318.
- [6] J. BOURGAIN, H. BREZIS, P. MIRONESCU, *Another look at Sobolev spaces*, Optimal Control and Partial Differential Equations, IOS Press, Amsterdam 2001, 439–455.
- [7] J. BOURGAIN, H. BREZIS, P. MIRONESCU, *A new function space and applications*, Journal of the EMS, 17 (2015), 2083–2101.

18 *A BMO-type characterization of higher order Sobolev spaces*

- [8] H. BREZIS, *How to recognize constant functions. A connection with Sobolev spaces.*, (Russian) Uspekhi Mat. Nauk 57 (2002), no. 4(346), 59–74; translation in Russian Math. Surveys 57 (2002), no. 4, 693–708.
- [9] A. BRUDNYI, Y. BRUDNYI, *On the Banach structure of multivariate BV spaces*, Dissertationes Math. 548 (2020), 52 pp.
- [10] G. DI FRATTA, A. FIORENZA, *BMO-type seminorms from Escher-type tessellations*, J. Funct. Anal. 279 (2020), 108556.
- [11] F. DEMENGEL, R. TEMAM, *Convex functions of a measure and applications*, Indiana University Mathematical Journal, vol. 33 n. 5 (1984), pp. 673-709.
- [12] F. FARRONI, N. FUSCO, S. GUARINO LO BIANCO, R. SCHIATTARELLA, *A formula for the anisotropic total variation of BV functions*, Journal of Functional Analysis 278 (9) (2020), 108451 .
- [13] F. FARRONI, S. GUARINO LO BIANCO, R. SCHIATTARELLA, *BMO-type seminorms generating Sobolev functions*, J. Math. Anal. Appl. 491 (2020), no.1, 124298, 15pp.
- [14] M. FUCHS, J. MULLER, *A higher order TV-type variational problem related to the denoising and inpainting of images*, Nonlinear Anal. 154 (2017), 122-147.
- [15] N. FUSCO, G. MOSCARIELLO, C. SBORDONE, *A formula for the total variation of SBV functions*. J. Funct. Anal. 270 (2016), no. 1, 419-446.
- [16] N. FUSCO, G. MOSCARIELLO, C. SBORDONE, *BMO-type seminorms and Sobolev functions*. ESAIM Control Optim. Calc. Var. 24 (2018), no. 2, 835-847.
- [17] R.L. JERRARD, H.M. SONER, *Functions of bounded higher variation*, Indiana Univ. Math. J., 51 (2002), pp. 645-677, 10.1512/iumj.2002.51.2229
- [18] G. LEONI *A first course in Sobolev spaces*. Second Edition. Graduate Studies in Mathematics, 181. American Mathematical Society, Providence, RI (2017).
- [19] MARCUS, *Exceptional sets with respect to Lebesgue differentiation of functions in Sobolev spaces*, Ann. Sc. Norm. Sup. Pisa, Cl. Sc., 1, no 1-2 (1974), 113-130.