

COARSE-GRAINING OF A DISCRETE MODEL FOR EDGE DISLOCATIONS IN THE REGULAR TRIANGULAR LATTICE

R. ALICANDRO, L. DE LUCA, G. LAZZARONI, M. PALOMBARO, AND M. PONSIGLIONE

ABSTRACT. We consider a discrete model of planar elasticity where the particles, in the reference configuration, sit on a regular triangular lattice and interact through nearest neighbor pairwise potentials, with bonds modeled as linearized elastic springs. Within this framework we introduce plastic slip fields, whose discrete circulation around each triangle detects the possible presence of an edge dislocation.

We provide a Γ -convergence analysis, as the lattice spacing tends to zero, of the elastic energy induced by edge dislocations in the energy regime corresponding to a finite number of geometrically necessary dislocations.

Keywords: Dislocations, Topological singularities, Plasticity, Discrete to continuum limits, Gamma-convergence.

MSC2020: 74C05, 58K45, 70G75, 74G65, 49J45.

CONTENTS

Introduction	1
1. The model and the main result	3
2. The continuum case	5
3. Proof of the main result	8
3.1. Preliminary results	8
3.2. Proof of Theorem 1.1	10
4. Degeneracies of the model and possible kinematic constraints	15
References	16

INTRODUCTION

Dislocations are line defects in the periodic structure of crystals and are considered the main microscopic mechanism of plastic flow. Idealized straight dislocations are classified into two types, *edge* and *screw*, while in real crystals dislocations are actually curved lines of mixed type [15, 14]. From a mathematical viewpoint, dislocations can be seen as topological line singularities around which the elastic strain has non trivial circulation given by a vector of the underlying lattice and referred to as *Burgers vector*.

Here we focus on planar elasticity, where the relevant dislocations are of edge nature and can be seen as topological point singularities of the elastic strain. Specifically, we study the discrete elastic energy induced by a finite system of edge dislocations in a finite portion $\Omega \cap \varepsilon\mathbb{T}$ of the regular triangular lattice $\varepsilon\mathbb{T}$, where ε is the lattice spacing.

As customary in the linearized framework, we adopt the additive decomposition of the *discrete deformation gradient* du , defined on pairs of nearest neighbors, into an elastic and a plastic part. The latter is represented by an additional variable σ , referred to as *slip*, defined on pairs of nearest neighbors and taking values in the set of lattice vectors. In this way we identify dislocations as points around which the discrete circulation of the plastic slip σ , or, equivalently, the discrete circulation of $du - \sigma$, is non trivial (see [1, 8, 4] for related models for screw dislocations and topological singularities in the XY model). Such a procedure agrees with the formalism of the eigenstrains considered in [3].

We focus on the simple case where nearest neighbors interact through linearized pairwise potentials; specifically, in absence of singularities, the energy functional is defined by

$$G_\varepsilon(u) := \frac{1}{2\varepsilon^2} \sum_{\substack{i,j \in \Omega \cap \varepsilon\mathbb{T} \\ |i-j|=\varepsilon}} (du(i,j) \cdot (j-i))^2.$$

The above energy can be formally derived by linearization of suitable nonlinear frame invariant functionals (see Remark 1.2); further rigorous linearization results in terms of Γ -convergence are provided in [2, 6, 19]. In the limit as $\varepsilon \rightarrow 0$ the energy $\frac{1}{\varepsilon^2}G_\varepsilon$ approximates, up to a pre-factor, the continuum isotropic elastic energy with Lamé parameters both equal to 1 (see Remark 3.1).

In presence of a plastic slip field σ , the energy functional should depend only on the elastic strain $du - \sigma$, and hence the resulting elastic energy reads as

$$F_\varepsilon(u, \sigma) := \frac{1}{2\varepsilon^2} \sum_{\substack{i,j \in \Omega \cap \varepsilon\mathbb{T} \\ |i-j|=\varepsilon}} ((du(i,j) - \sigma(i,j)) \cdot (j-i))^2.$$

It is well known that the energy induced by an isolated edge dislocation, both in the discrete and in the continuum setting, is of order $\varepsilon^2 |\log \varepsilon|$, while short dipoles of opposite dislocations induce a much smaller energy of order ε^2 . The aim of the present paper is to determine the asymptotic behavior (as $\varepsilon \rightarrow 0$) of F_ε in the energy regime $\varepsilon^2 |\log \varepsilon|$. This corresponds to a finite distribution of *geometrically necessary edge dislocations*, i.e., to the superposition of a finite number of isolated dislocations plus clusters of singularities with total Burgers vector equal to zero (usually referred to as *statistically stored dislocations*).

Our results are obtained within the rigorous formalism of Γ -convergence and consist in a compactness property for the dislocation measures and in the derivation of the effective limit energy induced by dislocations. Specifically, our analysis shows that the discrete dislocation density generated by the plastic slip σ converges, in the sense of flat convergence (1.7), to a finite sum of Dirac deltas $\sum_{k=1}^K b^k \delta_{x^k}$ with $b^k \in \mathbb{T}$. The effective energy (namely the Γ -limit) “counts” the limiting singularities with a coefficient $\varphi(b^k)$ given by the explicit formula (2.13). These results rely on a similar analysis performed in [11, 9] in a *semi-discrete* setting within the so-called core radius approach.

We restrict our analysis to configurations of plastic slips inducing dislocations with minimal mutual distance larger than or equal to $\sqrt{3}\varepsilon$. Roughly speaking, this means that two neighboring triangles cannot both contain a dislocation. Such a mild separation assumption (MS) guarantees that each dislocation induces a core energy of order ε^2 ; the latter estimate is essential also in the core radius approach, and represents the starting point in the so-called ball construction technique [18], which is devised to provide sharp lower bounds. This is not a mere technical assumption; indeed, removing it, one can exhibit unphysical configurations with zero stored elastic energy where all triangles contain a dislocation (see Remark 3.5). In this respect, it seems that our linearized model (without assuming (MS)) fails to describe the core energy stored in a single triangle and induced by the presence of a dislocation. On the other hand, under the mild separation assumption (MS), each dislocation is surrounded by an annulus of “elastically deformed” triangles where a finite amount of energy is stored, as it follows from an application of Korn’s inequality.

Although our model provides a good description of “non-pathological” dislocation configurations, it exhibits some degeneracies, due to the discrete linearized framework relying on a reference configuration, as well as to the presence of the slip variable (see Section 4). A natural way to rule out such degeneracies could be to include kinematic constraints on the slip fields mimicking (in a discrete framework) pure shear/deviatoric stress conditions that are typically assumed in (continuum) elasto-plasticity. In this respect, in Section 4, we discuss a possible constraint on the slip fields, based on the (formal) linearization of the volume preserving condition on dislocation-free ε -triangles; we remark that our Γ -convergence analysis still holds true under such a constraint.

Finally, some comments are in order. First, we do believe that a similar discrete-to-continuum analysis could be developed in the $\varepsilon^2 |\log \varepsilon|^2$ regime, which corresponds to a limit diffused distribution of dislocations and to an effective energy accounting for both elastic and plastic effects

(see [12] for an analogous result within the core-radius approach). Second, it would be interesting to look at the coarse-graining for the energy F_ε in even higher regimes, such as that of the grain boundaries (see the recent papers [13, 16, 10]). Third, a challenging issue would be to deal directly with nonlinear models built on the deformed configurations. An intriguing intermediate attempt in this direction could be to consider *hybrid models* combining the mathematical efficiency of linearized theories together with the mechanical understanding of dislocations in terms of interpenetrating pentagon-heptagon pairs usually observed in the deformed configurations.

Notation. For every $m, n \in \mathbb{N}$ and for every matrix $M \in \mathbb{R}^{m \times n}$, we denote by $M^* \in \mathbb{R}^{n \times m}$ the transpose matrix of M , i.e., such that $M_{ij}^* = M_{ji}$ for every $i = 1, \dots, n$ and $j = 1, \dots, m$. In what follows the elements (x_1, x_2) of \mathbb{R}^2 will be also identified (whenever it is convenient) with column vectors $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = (x_1 \ x_2)^*$. We denote by $\{e_1, e_2\}$ the canonical basis of \mathbb{R}^2 , where $e_1 = (1, 0)$ and $e_2 = (0, 1)$. Given two vectors $a = (a_1, a_2)$, $b = (b_1, b_2) \in \mathbb{R}^2$, we set $a \wedge b := a_1 b_2 - a_2 b_1$; moreover, we set $a^\perp := (-a_2, a_1)$.

For every open set U with ∂U smooth and for every $s \in \partial U$, we denote by $\tau(s)$ the tangent vector to ∂U at s defined as $\tau(s) = n^\perp(s)$, where $n(s)$ denotes the outer normal unitary vector to ∂U at s .

For every $p \in \mathbb{R}^2$ and for every $0 < r < R$ we define the annulus $A_{r,R}(p) := B_R(p) \setminus \overline{B_r}(p)$, where, for every $\rho > 0$, $B_\rho(p)$ and $\overline{B}_\rho(p)$ denote the open and the closed ball centered at p with radius ρ , respectively.

Moreover, for every bounded open set $A \subset \mathbb{R}^2$ and for every $\beta \in L^2(A; \mathbb{R}^{2 \times 2})$ the symbol $\text{Curl } \beta$ denotes the row-by-row distributional curl of β ; formally,

$$\text{Curl } \beta = (\partial_{x_1} \beta_{12} - \partial_{x_2} \beta_{11}, \partial_{x_1} \beta_{22} - \partial_{x_2} \beta_{21}).$$

Analogously, the symbol $\text{Div } \beta$ will denote the row-by-row distributional divergence of β , formally given by

$$(\text{Div } \beta)_i = \partial_{x_1} \beta_{i1} + \partial_{x_2} \beta_{i2}, \quad i \in \{1, 2\}.$$

Finally, $\mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ denotes the class of \mathbb{R}^2 valued Radon measures on \mathbb{R}^2 .

1. THE MODEL AND THE MAIN RESULT

In this section we introduce our model and state the main result.

Reference lattice. We set $\nu := \frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2$ and $\eta := -\frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2$. Let $\mathbb{T} := \text{span}_{\mathbb{Z}}\{e_1, \nu\}$ and set

$$T^+ := \text{conv}\{0, e_1, \nu\} \quad \text{and} \quad T^- := \text{conv}\{0, e_1, -\eta\},$$

where, for every $a, b, c \in \mathbb{R}^2$, the set $\text{conv}\{a, b, c\}$ denotes the convex envelope of the points a, b, c , i.e., the (closed) triangle with vertices at a, b, c . For every $\varepsilon > 0$ we denote by \mathcal{T}_ε the family of the triangles T_ε of the form $i + \varepsilon T^\pm$, with $i \in \varepsilon \mathbb{T}$.

Let $\Omega \subset \mathbb{R}^2$ be a bounded open set with Lipschitz continuous boundary, representing the domain of definition of the relevant fields in the model. For every $\varepsilon > 0$, we set

$$\mathcal{T}_\varepsilon(\Omega) := \{T_\varepsilon \in \mathcal{T}_\varepsilon : T_\varepsilon \subset \Omega\}$$

and we define $\Omega_\varepsilon := \bigcup_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} T_\varepsilon$. Moreover, we set $\Omega_\varepsilon^0 := \Omega_\varepsilon \cap \varepsilon \mathbb{T}$ and we denote by Ω_ε^1 the family of nearest neighbor bonds in Ω_ε , i.e., $\Omega_\varepsilon^1 := \{(i, j) \in \Omega_\varepsilon^0 \times \Omega_\varepsilon^0 : |i - j| = \varepsilon\}$. Trivially, $(i, j) \in \Omega_\varepsilon^1$ if and only if $(j, i) \in \Omega_\varepsilon^1$.

In the following we will generalize the notation introduced above to general subsets of \mathbb{R}^2 (not necessarily open). In particular, for every triangle $T_\varepsilon \in \mathcal{T}_\varepsilon$, we have

$$(T_\varepsilon)_\varepsilon^1 = \{(i, j) \in (T_\varepsilon \cap \varepsilon \mathbb{T}) \times (T_\varepsilon \cap \varepsilon \mathbb{T}) : i \neq j\}.$$

For every map $V : (T_\varepsilon)_\varepsilon^1 \rightarrow \mathbb{R}^2$, we define the *discrete circulation* of V on the “boundary of T_ε ” as

$$(1.1) \quad dV(T_\varepsilon) := V(i, j) + V(j, k) + V(k, i),$$

where (i, j, k) is a triple of counter-clockwise oriented vertices of T_ε .

Displacement, plastic slip, dislocation measure. We denote by $\mathcal{D}_\varepsilon(\Omega)$ the class of displacements $u : \Omega_\varepsilon^0 \rightarrow \mathbb{R}^2$; moreover, for every $u \in \mathcal{D}_\varepsilon(\Omega)$ we define the *discrete gradient* $du : \Omega_\varepsilon^1 \rightarrow \mathbb{R}^2$ of u as $du(i, j) := u(j) - u(i)$ for every $(i, j) \in \Omega_\varepsilon^1$.

We define the class of *plastic slips* $\mathcal{S}_\varepsilon(\Omega)$ as

$$\mathcal{S}_\varepsilon(\Omega) := \left\{ \sigma : \Omega_\varepsilon^1 \rightarrow \varepsilon\mathbb{T} : \sigma(i, j) = -\sigma(j, i) \text{ for every } (i, j) \in \Omega_\varepsilon^1 \right\}.$$

For every $\sigma \in \mathcal{S}_\varepsilon(\Omega)$ we define the discrete dislocation measure associated to σ as

$$\mu[\sigma] := - \sum_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} d\sigma(T_\varepsilon) \delta_{x_{T_\varepsilon}},$$

where $d\sigma$ is defined in (1.1) and x_{T_ε} denotes the barycenter of the triangle T_ε . Notice that for every $u \in \mathcal{D}_\varepsilon(\Omega)$ and for every $\sigma \in \mathcal{S}_\varepsilon(\Omega)$ it holds

$$\mu[\sigma] = \sum_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} d(du - \sigma)(T_\varepsilon) \delta_{x_{T_\varepsilon}}.$$

The class of *admissible dislocation measures*, denoted by $X_\varepsilon(\Omega)$, is the family of measures of the form $\mu = \sum_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} b(T_\varepsilon) \delta_{x_{T_\varepsilon}}$ with $b(T_\varepsilon) \in \varepsilon\mathbb{T}$ and satisfying the following *mild separation* property:

- (MS) for every $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ with $\mu(T_\varepsilon) \neq 0$, we have
 $\partial T_\varepsilon \cap \partial \Omega_\varepsilon = \emptyset$ and $\mu(T'_\varepsilon) = 0$ for every $T'_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ with $\partial T'_\varepsilon \cap \partial T_\varepsilon \neq \emptyset$.

Finally, we set

$$X(\Omega) := \left\{ \mu = \sum_{k=1}^K b^k \delta_{x^k} : K \in \mathbb{N}, b^k \in \mathbb{T}, x^k \in \Omega \right\}.$$

The energy functional and the main result. We are now in a position to define the energy functionals $F_\varepsilon : \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega) \rightarrow [0, +\infty)$ as

$$(1.2) \quad F_\varepsilon(u, \sigma) := \frac{1}{2\varepsilon^2} \sum_{(i, j) \in \Omega_\varepsilon^1} [(du(i, j) - \sigma(i, j)) \cdot (j - i)]^2,$$

where \cdot denotes the standard scalar product in \mathbb{R}^2 .

We will consider also localized versions of the functional $F_\varepsilon(u, \sigma)$ in (1.2). More specifically, for every set $A \subset \mathbb{R}^2$, we define $F_\varepsilon(\cdot, \cdot; A) : \mathcal{D}_\varepsilon(A) \times \mathcal{S}_\varepsilon(A) \rightarrow [0, +\infty)$ as

$$(1.3) \quad F_\varepsilon(u, \sigma; A) := \frac{1}{2\varepsilon^2} \sum_{(i, j) \in A_\varepsilon^1} [(du(i, j) - \sigma(i, j)) \cdot (j - i)]^2,$$

so that $F_\varepsilon(u, \sigma; \Omega) = F_\varepsilon(u, \sigma)$.

Since in our analysis the relevant parameter is the dislocation measure μ associated to σ , we let the energy functionals depend only on (the admissible measures) μ , by defining $\mathcal{F}_\varepsilon : X_\varepsilon(\Omega) \rightarrow [0, +\infty)$ as

$$\mathcal{F}_\varepsilon(\mu) := \inf_{\substack{(u, \sigma) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega) \\ \mu[\sigma] = \mu}} F_\varepsilon(u, \sigma).$$

For every $b \in \mathbb{T}$, we set

$$(1.4) \quad \varphi(b) := \frac{1}{3\pi} \min \left\{ \sum_{i=1}^3 |z_i| : z_1, z_2, z_3 \in \mathbb{Z}, \quad b = z_1 e_1 + z_2 \nu + z_3 \eta \right\}.$$

Our main result is the following theorem.

Theorem 1.1. *The following Γ -convergence result holds true.*

- (i) (*Compactness*) Let $\{\mu_\varepsilon\}_\varepsilon \subset \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ be such that $\mu_\varepsilon \in X_\varepsilon(\Omega)$ for every $\varepsilon > 0$. If $\mathcal{F}_\varepsilon(\mu_\varepsilon) \leq C\varepsilon^2 |\log \varepsilon|$, then, up to a subsequence, $\frac{\mu_\varepsilon}{\varepsilon} \xrightarrow{\text{flat}} \mu$ (as $\varepsilon \rightarrow 0$) for some $\mu \in X(\Omega)$.

(ii) (Γ -liminf inequality) For every $\mu = \sum_{k=1}^K b^k \delta_{x^k} \in X(\Omega)$ and for every $\{\mu_\varepsilon\}_\varepsilon \subset \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ with $\mu_\varepsilon \in X_\varepsilon(\Omega)$ for every $\varepsilon > 0$ and such that $\frac{\mu_\varepsilon}{\varepsilon} \xrightarrow{\text{flat}} \mu$ (as $\varepsilon \rightarrow 0$) it holds

$$(1.5) \quad \liminf_{\varepsilon \rightarrow 0} \frac{\mathcal{F}_\varepsilon(\mu_\varepsilon)}{\varepsilon^2 |\log \varepsilon|} \geq \frac{\sqrt{3}}{2} \sum_{k=1}^K \varphi(b^k).$$

(iii) (Γ -limsup inequality) For every $\mu = \sum_{k=1}^K b^k \delta_{x^k} \in X(\Omega)$ there exists $\{\mu_\varepsilon\}_\varepsilon \subset \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ such that $\mu_\varepsilon \in X_\varepsilon(\Omega)$ for every $\varepsilon > 0$, $\frac{\mu_\varepsilon}{\varepsilon} \xrightarrow{\text{flat}} \mu$ (as $\varepsilon \rightarrow 0$) and

$$(1.6) \quad \limsup_{\varepsilon \rightarrow 0} \frac{\mathcal{F}_\varepsilon(\mu_\varepsilon)}{\varepsilon^2 |\log \varepsilon|} \leq \frac{\sqrt{3}}{2} \sum_{k=1}^K \varphi(b^k).$$

The convergence appearing in Theorem 1.1 is the *flat convergence*, that is the convergence with respect to the *flat norm*, defined by

$$(1.7) \quad \|\mu\|_{\text{flat}} := \sup_{\substack{\phi \in C_c^{0,1}(\Omega) \\ \|\phi\|_{C^{0,1}} \leq 1}} \left| \int_{\Omega} \phi \, d\mu \right|, \quad \text{for every } \mu \in \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2),$$

where $C^{0,1}(\Omega)$ is the space of Lipschitz continuous functions endowed with the norm

$$\|\phi\|_{C^{0,1}} := \sup_{x \in \Omega} |\phi(x)| + \sup_{\substack{x, y \in \Omega \\ x \neq y}} \frac{|\phi(x) - \phi(y)|}{|x - y|},$$

and $C_c^{0,1}(\Omega)$ is the subspace of $C^{0,1}$ functions compactly supported in Ω .

Remark 1.2. We illustrate how to formally derive (1.2) from a nonlinear frame invariant model with nearest neighbor pairwise interaction potentials, in the case where no dislocation is present, i.e., $\sigma \equiv 0$. We assume that, given a deformation $v: \Omega_\varepsilon^0 \rightarrow \mathbb{R}^2$, the total interaction energy is

$$E_\varepsilon(v) = \varepsilon^2 \sum_{(i,j) \in \Omega_\varepsilon^1} \psi \left(\left| \frac{dv(i,j)}{\varepsilon} \right| \right),$$

where ψ is a C^2 function such that $\psi(1) = \psi'(1) = 0$ and $\psi''(1) > 0$, so that the identity is an equilibrium configuration. Expressing the energy in terms of the displacement $u(x) = \frac{v(x) - x}{\delta}$, with $\delta > 0$, and rescaling by δ^2 , the total interaction energy reads as

$$E_\varepsilon^\delta(x + \delta u) = \frac{\varepsilon^2}{\delta^2} \sum_{(i,j) \in \Omega_\varepsilon^1} \psi \left(\left| \frac{j-i}{\varepsilon} + \delta \frac{du(i,j)}{\varepsilon} \right| \right).$$

A second order Taylor expansion of $E_\varepsilon^\delta(x + \delta u)$ with respect to δ about the point $\delta = 0$ gives

$$E_\varepsilon^\delta(x + \delta u) = F_\varepsilon(u) + o_\delta(1),$$

where

$$F_\varepsilon(u) = \frac{1}{2\varepsilon^2} \psi''(1) \sum_{(i,j) \in \Omega_\varepsilon^1} [du(i,j) \cdot (j-i)]^2,$$

which coincides with (1.2) up to a multiplicative constant when $\sigma \equiv 0$. We refer to [13] for a formal derivation of (1.2) starting from a nonlinear model with discrete plastic slips.

2. THE CONTINUUM CASE

In order to prove Theorem 1.1 we will use the corresponding result in the continuum setting [9], that we briefly recall here. Let A be an open bounded subset of \mathbb{R}^2 with Lipschitz continuous boundary. For every $\mu \in X(A)$ we set

$$(2.1) \quad A_\varepsilon(\mu) := A \setminus \bigcup_{x \in \text{supp}(\mu)} \bar{B}_\varepsilon(x)$$

and we define the set $\mathcal{AS}_\varepsilon(\mu)$ of admissible strains associated to μ as

$$(2.2) \quad \mathcal{AS}_\varepsilon(\mu) := \left\{ \beta \in L^2(A_\varepsilon(\mu); \mathbb{R}^{2 \times 2}) : \text{Curl } \beta = 0 \text{ in } A_\varepsilon(\mu), \right. \\ \left. \int_{\partial U} \beta \tau \, d\mathcal{H}^1 = \mu(U) \text{ for every open set } U \subset A \text{ with } \partial U \subset A_\varepsilon(\mu) \text{ smooth} \right\}.$$

Let \mathbb{C} be an elasticity tensor, i.e., a linear operator from $\mathbb{R}^{2 \times 2}$ into itself satisfying the following property: There exist two constants $0 < c_1 < c_2$ such that

$$c_1 |\beta^{\text{sym}}|^2 \leq \frac{1}{2} \mathbb{C} \beta : \beta \leq c_2 |\beta^{\text{sym}}|^2 \quad \text{for every } \beta \in \mathbb{R}^{2 \times 2},$$

where $\beta^{\text{sym}} := \frac{1}{2}(\beta + \beta^*)$.

The elastic energy of a field $\beta \in \mathcal{AS}_\varepsilon(\mu)$ in the body A is given by

$$E_\varepsilon(\beta; A_\varepsilon(\mu)) := \frac{1}{2} \int_{A_\varepsilon(\mu)} \mathbb{C} \beta : \beta \, dx,$$

and the energy induced by the dislocation distribution μ in the body A is defined by

$$(2.3) \quad \mathcal{E}_\varepsilon(\mu; A) := \inf_{\beta \in \mathcal{AS}_\varepsilon(\mu)} E_\varepsilon(\beta; A_\varepsilon(\mu)) + |\mu|(A), \quad \text{for every } \mu \in X(A).$$

In (2.3), the first addendum on the right hand side is the elastic energy induced by the dislocation measure μ , whereas the second addendum plays the role of a plastic core energy.

In order to introduce the self-energy of an edge dislocation, for every $b \in \mathbb{T}$ we first define the strain field $\beta_{\mathbb{R}^2}^{b, \mathbb{C}}$ satisfying the circulation condition

$$\text{Curl } \beta = b \delta_0 \text{ in } \mathbb{R}^2$$

and the equilibrium equation

$$\text{Div } \mathbb{C} \beta = 0 \text{ in } \mathbb{R}^2.$$

As shown in [9], the strain field $\beta_{\mathbb{R}^2}^{b, \mathbb{C}}$ is given, in polar coordinates, by

$$(2.4) \quad \beta_{\mathbb{R}^2}^{b, \mathbb{C}}(\rho, \theta) := \frac{1}{\rho} (f^{b, \mathbb{C}}(\theta) \otimes (-\sin \theta, \cos \theta) + g^{b, \mathbb{C}} \otimes (\cos \theta, \sin \theta)),$$

where the constant $g^{b, \mathbb{C}} \in \mathbb{R}^2$ and the function $f^{b, \mathbb{C}} \in C^0([0, 2\pi]; \mathbb{R}^2)$, with $f(0) = f(2\pi)$ and $\int_0^{2\pi} f^{b, \mathbb{C}}(\omega) \, d\omega = b$, are uniquely determined by the vector b and the tensor \mathbb{C} .

The corresponding displacement $u_{\mathbb{R}^2}^{b, \mathbb{C}}$ (i.e., such that $\nabla u_{\mathbb{R}^2}^{b, \mathbb{C}} = \beta_{\mathbb{R}^2}^{b, \mathbb{C}}$) is computed explicitly in the literature (see, for instance, [5, formula 4.1.25]) and, in polar coordinates, is given by

$$(2.5) \quad u_{\mathbb{R}^2}^{b, \mathbb{C}}(\rho, \theta) = F^{b, \mathbb{C}}(\theta) + g^{b, \mathbb{C}} \log \rho,$$

where $F^{b, \mathbb{C}}(\theta) = \int_0^\theta f^{b, \mathbb{C}}(\omega) \, d\omega$, for $\theta \in [0, 2\pi)$. Note that the displacement above is uniquely determined up to a constant.

For every $b \in \mathbb{T}$ we set

$$(2.6) \quad \psi^{\mathbb{C}}(b) := \int_0^{2\pi} \frac{1}{2} \mathbb{C} \Gamma^{b, \mathbb{C}}(\theta) : \Gamma^{b, \mathbb{C}}(\theta) \, d\theta = \frac{1}{|\log r|} \int_{B_1 \setminus \overline{B}_r} \frac{1}{2} \mathbb{C} \beta_{\mathbb{R}^2}^{b, \mathbb{C}} : \beta_{\mathbb{R}^2}^{b, \mathbb{C}} \, dx, \quad 0 < r < 1,$$

where we have set $\Gamma^{b, \mathbb{C}}(\theta) := (f^{b, \mathbb{C}}(\theta) \otimes (-\sin \theta, \cos \theta) + g^{b, \mathbb{C}} \otimes (\cos \theta, \sin \theta))$ for every $\theta \in [0, 2\pi]$. Finally, for every $b \in \mathbb{T}$ we define

$$(2.7) \quad \varphi^{\mathbb{C}}(b) := \min \left\{ \sum_{i=1}^N |z_i| \psi^{\mathbb{C}}(b_i) : z_i \in \mathbb{Z}, b_i \in \mathbb{T}, N \in \mathbb{N}, \sum_{i=1}^N z_i b_i = b \right\}.$$

The following result is a slight variant of [9, Theorem 4].

Theorem 2.1. *The following Γ -convergence result holds true.*

(i) (Compactness) Let $\{\mu_\varepsilon\}_\varepsilon \subset X(A)$ and let $\{\beta_\varepsilon\}_\varepsilon$ be a sequence of fields with $\beta_\varepsilon \in \mathcal{AS}_\varepsilon(\mu_\varepsilon)$ (for every $\varepsilon > 0$) such that

$$(2.8) \quad E_\varepsilon(\beta_\varepsilon; A_\varepsilon(\mu_\varepsilon)) + |\mu_\varepsilon|(A) \leq M |\log \varepsilon|;$$

then, up to a subsequence, $\mu_\varepsilon \xrightarrow{\text{flat}} \mu$ (as $\varepsilon \rightarrow 0$) for some $\mu \in X(A)$.

(ii) (Γ -liminf inequality) For every $\mu = \sum_{k=1}^K b^k \delta_{x^k} \in X(A)$, for every $\{\mu_\varepsilon\}_\varepsilon \subset X(A)$ with $\mu_\varepsilon \xrightarrow{\text{flat}} \mu$ (as $\varepsilon \rightarrow 0$) and $|\mu_\varepsilon|(A) \leq C |\log \varepsilon|$ and for every sequence of fields $\{\beta_\varepsilon\}_\varepsilon$ with $\beta_\varepsilon \in \mathcal{AS}_\varepsilon(\mu_\varepsilon)$ (for every $\varepsilon > 0$) it holds

$$(2.9) \quad \liminf_{\varepsilon \rightarrow 0} \frac{E_\varepsilon(\beta_\varepsilon; A_\varepsilon(\mu_\varepsilon))}{|\log \varepsilon|} \geq \sum_{k=1}^K \varphi^{\mathbb{C}}(b^k);$$

(iii) (Γ -limsup inequality) For every $\mu = \sum_{k=1}^K b^k \delta_{x^k} \in X(A)$ there exists a sequence of measures $\{\mu_\varepsilon\}_\varepsilon \subset X(A)$ such that $\mu_\varepsilon \xrightarrow{\text{flat}} \mu$ (as $\varepsilon \rightarrow 0$) and

$$(2.10) \quad \limsup_{\varepsilon \rightarrow 0} \frac{\mathcal{E}_\varepsilon(\mu_\varepsilon; A)}{|\log \varepsilon|} \leq \sum_{k=1}^K \varphi^{\mathbb{C}}(b^k).$$

The proof of Theorem 2.1 can be obtained by arguing verbatim as in the proof of [9, Theorem 4]. In fact, the minor differences between Theorem 2.1 and [9, Theorem 4] are the following. First, in [9] the admissible strains β should satisfy the condition

$$\int_{A_\varepsilon(\mu)} (\beta - \beta^*) dx = 0.$$

Such a condition can be always enforced in view of the invariance of the elastic energy with respect to translations. Second, in [9, Theorem 4] the compactness property is stated enforcing (2.8) only for the optimal β_ε . Third, in [9, Theorem 4], the lower bound (2.9) is provided for the functional \mathcal{E}_ε ; there, the assumption $|\mu_\varepsilon|(A) \leq C |\log \varepsilon|$ is automatically satisfied by sequences with equibounded energy. However, assuming $|\mu_\varepsilon|(A) \leq C |\log \varepsilon|$, the same proof provides the same lower bound for $\frac{1}{|\log \varepsilon|} E_\varepsilon(\beta_\varepsilon; A_\varepsilon(\mu_\varepsilon))$.

Now, we specialize the functions $\psi^{\mathbb{C}}$ and $\varphi^{\mathbb{C}}$ to the particular isotropic case we deal with in the discrete-to-continuum limit. More specifically, let \mathbb{C} be the isotropic elasticity tensor with Lamé parameters both equal to 1, i.e.,

$$(2.11) \quad \mathbb{C}\beta : \beta := |\text{tr } \beta|^2 + 2|\beta^{\text{sym}}|^2,$$

By [7, formulas (3.3) and (3.4)], for the specific choice of \mathbb{C} in (2.11) we have that the constant $g^{b,\mathbb{C}}$ and the function $f^{b,\mathbb{C}}$ in (2.4) are given by

$$(2.12) \quad \begin{aligned} g^{b,\mathbb{C}} &= -\frac{1}{6\pi} b^\perp, \\ f^{b,\mathbb{C}}(\theta) &= \frac{1}{2\pi} b - \frac{1}{3\pi} (-b_1 \cos(2\theta) - b_2 \sin(2\theta), b_2 \cos(2\theta) - b_1 \sin(2\theta)); \end{aligned}$$

therefore, by straightforward computations, in this case

$$\psi(b) := \psi^{\mathbb{C}}(b) = \frac{1}{3\pi} |b|^2, \quad b \in \mathbb{T},$$

and hence, by (2.7),

$$(2.13) \quad \begin{aligned} \varphi(b) = \varphi^{\mathbb{C}}(b) &= \frac{1}{3\pi} \min \left\{ \sum_{i=1}^N |z_i| |b_i|^2 : z_i \in \mathbb{Z}, b_i \in \mathbb{T}, N \in \mathbb{N}, \sum_{i=1}^N z_i b_i = b \right\} \\ &= \frac{1}{3\pi} \min \left\{ \sum_{i=1}^3 |z_i| : z_1, z_2, z_3 \in \mathbb{Z}, b = z_1 e_1 + z_2 \nu + z_3 \eta \right\}, \end{aligned}$$

which is exactly (1.4).

3. PROOF OF THE MAIN RESULT

This section is devoted to the proof of Theorem 1.1. In Subsection 3.1 below we collect some auxiliary results that will be instrumental to prove Theorem 1.1.

3.1. Preliminary results. We start by deriving the continuum isotropic elasticity tensor associated to our discrete functional.

Remark 3.1. Let $\varepsilon > 0$ and let $T_\varepsilon \in \mathcal{T}_\varepsilon$. Let moreover $u : T_\varepsilon \cap \varepsilon\mathbb{T} \rightarrow \mathbb{R}^2$ and $\sigma : (T_\varepsilon)_\varepsilon^1 \rightarrow \varepsilon\mathbb{T}$ be such that $\sigma(i, j) = -\sigma(j, i)$ for every $(i, j) \in (T_\varepsilon)_\varepsilon^1$. If $\beta \in \mathbb{R}^{2 \times 2}$ satisfies

$$\beta(j - i) = du(i, j) - \sigma(i, j) \quad \text{for every } (i, j) \in (T_\varepsilon)_\varepsilon^1,$$

then, by straightforward computations, we have that

$$(3.1) \quad F_\varepsilon(u, \sigma; T_\varepsilon) = \varepsilon^2 \left(|e_1^* \beta e_1|^2 + |\nu^* \beta \nu|^2 + |\eta^* \beta \eta|^2 \right) = \varepsilon^2 \frac{3}{8} \mathbb{C} \beta : \beta = \frac{\sqrt{3}}{2} \int_{T_\varepsilon} \mathbb{C} \beta : \beta \, dx,$$

where $F_\varepsilon(\cdot, \cdot; T_\varepsilon)$ is defined in (1.3) and \mathbb{C} is given in (2.11).

In the next lemma we construct, far from the singularities, a (continuous) strain field β that is compatible with a given distribution of dislocations and whose (continuous) energy behaves like the discrete energy F_ε . Such an estimate, together with a bound on the total variation, will allow us to deduce the Γ -liminf inequality in Theorem 1.1(ii) directly from the analogous statement in the continuous setting (Theorem 2.1(ii)).

Lemma 3.2. *Let $\varepsilon > 0$ and let $(u, \sigma) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega)$. Let moreover*

$$(3.2) \quad \mathcal{K}_\varepsilon := \{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega) : \mu[\sigma](T_\varepsilon) = 0\}, \quad K_\varepsilon := \bigcup_{T_\varepsilon \in \mathcal{K}_\varepsilon} T_\varepsilon.$$

Then, there exists a piecewise constant (namely, constant on each triangle $T_\varepsilon \in \mathcal{K}_\varepsilon$) field $\beta^{u, \sigma, \mathcal{K}_\varepsilon} \in L^2(K_\varepsilon; \mathbb{R}^{2 \times 2})$ such that:

- (a) $du(i, j) - \sigma(i, j) = \beta_{|T_\varepsilon}^{u, \sigma, \mathcal{K}_\varepsilon}(j - i)$ for every $i, j \in T_\varepsilon$ with $T_\varepsilon \in \mathcal{K}_\varepsilon$;
- (b) $F_\varepsilon(u, \sigma; K_\varepsilon) = \frac{\sqrt{3}}{2} \int_{K_\varepsilon} \frac{1}{2} \mathbb{C} \beta^{u, \sigma, \mathcal{K}_\varepsilon} : \beta^{u, \sigma, \mathcal{K}_\varepsilon} \, dx + \frac{1}{4\varepsilon^2} \sum_{\substack{(i, j) \in K_\varepsilon^1 \\ i, j \in \partial K_\varepsilon}} |(du(i, j) - \sigma(i, j)) \cdot (j - i)|^2$,
where F_ε is defined in (1.3) and \mathbb{C} is defined in (2.11);
- (c) $\text{Curl } \beta^{u, \sigma, \mathcal{K}_\varepsilon} = 0$ in K_ε and $\int_{\partial U} \beta^{u, \sigma, \mathcal{K}_\varepsilon} \tau \, d\mathcal{H}^1 = \mu[\sigma](U)$ for every smooth open set $U \subset \Omega$ such that $\partial U \subset K_\varepsilon$.

Proof. Let $T_\varepsilon = \text{conv}\{i, j, k\} \in \mathcal{K}_\varepsilon$ and assume that the triple (i, j, k) is counterclockwise oriented. Let $v_{T_\varepsilon} \in \mathcal{D}_\varepsilon(T_\varepsilon)$ be defined by

$$v_{T_\varepsilon}(i) = 0, \quad v_{T_\varepsilon}(j) = du(i, j) - \sigma(i, j), \quad v_{T_\varepsilon}(k) = v_{T_\varepsilon}(j) + du(j, k) - \sigma(j, k),$$

and notice that the discrete gradient dv_{T_ε} of v_{T_ε} agrees with $du - \sigma$. For every $T_\varepsilon \in \mathcal{K}_\varepsilon$ let $\tilde{v}_{T_\varepsilon} : T_\varepsilon \rightarrow \mathbb{R}^2$ be the piecewise affine interpolation of v_{T_ε} and let $\beta^{u, \sigma, \mathcal{K}_\varepsilon} : K_\varepsilon \rightarrow \mathbb{R}^{2 \times 2}$ be the map defined by

$$\beta^{u, \sigma, \mathcal{K}_\varepsilon} := \sum_{T_\varepsilon \in \mathcal{K}_\varepsilon} D\tilde{v}_{T_\varepsilon} \chi_{T_\varepsilon},$$

where $D\tilde{v}_{T_\varepsilon}$ denotes the (continuous) gradient of the function $\tilde{v}_{T_\varepsilon}$. Clearly $\beta^{u, \sigma, \mathcal{K}_\varepsilon} \in L^2(K_\varepsilon; \mathbb{R}^{2 \times 2})$. Moreover, by construction, we have that property (a) is satisfied; furthermore, since for every $T_\varepsilon \in \mathcal{K}_\varepsilon$ and for every $(i, j) \in T_\varepsilon^1$ it holds

$$\beta_{|T_\varepsilon}^{u, \sigma, \mathcal{K}_\varepsilon}(j - i) = du(i, j) - \sigma(i, j),$$

by Remark 3.1 we get

$$\begin{aligned} F_\varepsilon(u, \sigma; K_\varepsilon) &= \frac{1}{2} \sum_{T_\varepsilon \in \mathcal{K}_\varepsilon} F_\varepsilon(u, \sigma; T_\varepsilon) + \frac{1}{4\varepsilon^2} \sum_{\substack{(i,j) \in K_\varepsilon^1 \\ i,j \in \partial K_\varepsilon}} |(\mathrm{d}u(i, j) - \sigma(i, j)) \cdot (j - i)|^2 \\ &= \frac{\sqrt{3}}{2} \int_{K_\varepsilon} \frac{1}{2} \mathbb{C} \beta^{u, \sigma, \mathcal{K}_\varepsilon} : \beta^{u, \sigma, \mathcal{K}_\varepsilon} \, \mathrm{d}x + \frac{1}{4\varepsilon^2} \sum_{\substack{(i,j) \in K_\varepsilon^1 \\ i,j \in \partial K_\varepsilon}} |(\mathrm{d}u(i, j) - \sigma(i, j)) \cdot (j - i)|^2, \end{aligned}$$

i.e., property (b).

Finally, we prove that also (c) is satisfied. To this end, we notice that if T_ε^1 and T_ε^2 are two triangles sharing one edge (i, j) , then, by construction, $\beta_{|T_\varepsilon^1}^{u, \sigma, \mathcal{K}_\varepsilon}(j - i) = \beta_{|T_\varepsilon^2}^{u, \sigma, \mathcal{K}_\varepsilon}(j - i)$ so that $\mathrm{Curl} \beta^{u, \sigma, \mathcal{K}_\varepsilon} = 0$ on K_ε . Now, since for every $T_\varepsilon \in \mathcal{K}_\varepsilon$,

$$\int_{\partial T_\varepsilon} \beta^{u, \sigma, \mathcal{K}_\varepsilon} \tau \, \mathrm{d}\mathcal{H}^1 = 0,$$

by using Stokes' Theorem, for every smooth open set $U \subset \Omega$ such that $\partial U \subset K_\varepsilon$, we have

$$\int_{\partial U} \beta^{u, \sigma, \mathcal{K}_\varepsilon} \tau \, \mathrm{d}\mathcal{H}^1 = \int_{U \cap \partial K_\varepsilon} \beta^{u, \sigma, \mathcal{K}_\varepsilon} \tau \, \mathrm{d}\mathcal{H}^1 = - \sum_{T_\varepsilon \subset U} \mathrm{d}\sigma(T_\varepsilon) = \mu[\sigma](U),$$

which concludes the proof of property (c) and of the whole lemma. \square

The following result allows to extend the field β constructed in Lemma 3.2 above up to the boundary of Ω .

Lemma 3.3. *Let $\varepsilon > 0$ and let $(u, \sigma) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega)$ be such that $\mu[\sigma] \in X_\varepsilon(\Omega)$; let moreover \mathcal{K}_ε and K_ε be defined as in (3.2) and let $\beta = \beta^{u, \sigma, \mathcal{K}_\varepsilon}$ be the field provided by Lemma 3.2. Then, there exists a field $\hat{\beta} = \hat{\beta}^{u, \sigma, \mathcal{K}_\varepsilon} \in L^2(K_\varepsilon \cup (\Omega \setminus \Omega_\varepsilon); \mathbb{R}^{2 \times 2})$ such that*

- (i) $\hat{\beta} = \beta$ in K_ε ;
- (ii) $\mathrm{Curl} \hat{\beta} = 0$ (in the sense of distributions);
- (iii) $\int_{K_\varepsilon \cup (\Omega \setminus \Omega_\varepsilon)} \mathbb{C} \hat{\beta} : \hat{\beta} \, \mathrm{d}x \leq C \int_{K_\varepsilon} \mathbb{C} \beta : \beta \, \mathrm{d}x$, for some constant C independent of ε .

Proof. It is enough to notice that, in view of (MS), each of the triangles $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ touching $\partial \Omega_\varepsilon$ satisfies $\mu[\sigma](T_\varepsilon) = 0$. Therefore, in order to construct a field $\hat{\beta}$ satisfying properties (i), (ii), and (iii), it is enough to extend locally by reflection the field β provided by Lemma 3.2. \square

We conclude this subsection showing how assumption (MS) allows to estimate the total variation of the dislocation measure μ with the elastic energy $\mathcal{F}_\varepsilon(\mu)$.

Lemma 3.4. *There exists $\bar{C} > 0$ such that $|\frac{\mu}{\varepsilon}|(\Omega) \leq \bar{C} \frac{1}{\varepsilon^2} \mathcal{F}_\varepsilon(\mu)$ for every $\varepsilon > 0$ and for every $\mu \in X_\varepsilon(\Omega)$.*

Proof. Let $\varepsilon > 0$ and $\mu \in X_\varepsilon(\Omega)$ be fixed. Let moreover $(u, \sigma) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega)$ with $\mu[\sigma] = \mu$. For every $p \in \mathrm{supp} \mu$ let $\mathcal{K}_\varepsilon^p$ be the set of the triangles $T_\varepsilon \neq T_\varepsilon^p$ sharing at least one vertex with the triangle T_ε^p centered at p and let $K_\varepsilon^p := \bigcup_{T_\varepsilon \in \mathcal{K}_\varepsilon^p} T_\varepsilon$. By the very definition of $X_\varepsilon(\Omega)$, we have that $\mu(T_\varepsilon) = 0$ for every $T_\varepsilon \in \bigcup_{p \in \mathrm{supp} \mu} \mathcal{K}_\varepsilon^p$. Let $\beta^{u, \sigma, \mathcal{K}_\varepsilon^p}$ be the map defined in Lemma 3.2 above. Recalling the notation for the annulus $A_{r, R}(p)$, for every $p \in \mathrm{supp} \mu$ we set

$$S_\varepsilon^p := \frac{1}{|A_{\frac{\sqrt{3}}{3}\varepsilon, \frac{5\sqrt{3}}{12}\varepsilon}(p)|} \int_{A_{\frac{\sqrt{3}}{3}\varepsilon, \frac{5\sqrt{3}}{12}\varepsilon}(p)} (\beta^{u, \sigma, \mathcal{K}_\varepsilon^p})^{\mathrm{skew}} \, \mathrm{d}y,$$

where, for every $\beta \in \mathbb{R}^{2 \times 2}$, we have set $\beta^{\text{skew}} := \frac{1}{2}(\beta - \beta^*)$. By Korn's inequality, there exists a constant κ (independent of ε and p) such that

$$\begin{aligned} \int_{A_{\frac{\sqrt{3}}{3}\varepsilon, \frac{5\sqrt{3}}{12}\varepsilon}(p)} |(\beta^{u, \sigma, \mathcal{K}_\varepsilon^p})^{\text{sym}}|^2 dx &\geq \kappa \int_{A_{\frac{\sqrt{3}}{3}\varepsilon, \frac{5\sqrt{3}}{12}\varepsilon}(p)} |\beta^{u, \sigma, \mathcal{K}_\varepsilon^p} - S_\varepsilon^p|^2 dx \\ &\geq \kappa \int_{\frac{\sqrt{3}}{3}\varepsilon}^{\frac{5\sqrt{3}}{12}\varepsilon} \int_{\partial B_\rho(p)} |(\beta^{u, \sigma, \mathcal{K}_\varepsilon^p} - S_\varepsilon^p) \tau|^2 d\mathcal{H}^1 d\rho \\ &\geq \frac{\kappa}{2\pi} \log 2 |\mu(p)|^2 = \frac{\kappa}{2\pi} \log 2 \varepsilon^2 \left| \frac{\mu}{\varepsilon}(p) \right|^2 \geq \frac{\kappa}{2\pi} \log 2 \varepsilon^2 \left| \frac{\mu}{\varepsilon}(p) \right|, \end{aligned}$$

where the second inequality follows by Fubini theorem, estimating $|\beta^{u, \sigma, \mathcal{K}_\varepsilon^p} - S_\varepsilon^p|$ from below by its tangential part, and the third one is a consequence of Jensen's inequality using Lemma 3.2(c) together with

$$\int_{\partial B_\rho(p)} S_\varepsilon^p \tau d\mathcal{H}^1 = 0.$$

By assumption (MS), Lemma 3.2(b) and the very definition of \mathbb{C} in (2.11), using that the annuli $A_{\frac{\sqrt{3}}{3}\varepsilon, \frac{5\sqrt{3}}{12}\varepsilon}(p)$ are pairwise disjoint and contained in K_ε^p , we thus deduce that

$$\begin{aligned} F_\varepsilon(u, \sigma) &\geq \frac{\sqrt{3}}{2} \sum_{p \in \text{supp } \mu} \frac{1}{2} \int_{K_\varepsilon^p} \mathbb{C} \beta^{u, \sigma, \mathcal{K}_\varepsilon^p} : \beta^{u, \sigma, \mathcal{K}_\varepsilon^p} dx \geq C \sum_{p \in \text{supp } \mu} \int_{A_{\frac{\sqrt{3}}{3}\varepsilon, \frac{5\sqrt{3}}{12}\varepsilon}(p)} |(\beta^{u, \sigma, \mathcal{K}_\varepsilon^p})^{\text{sym}}|^2 dx \\ &\geq C \varepsilon^2 \left| \frac{\mu}{\varepsilon} \right|(\Omega), \end{aligned}$$

which, taking the infimum over the pairs $(u, \sigma) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega)$ with $\mu[\sigma] = \mu$, provides the claim for $\tilde{C} = \frac{1}{C}$. \square

Remark 3.5. Without the assumption (MS), Lemma 3.4 does not hold. As an example, for every $\varepsilon > 0$, let $u_\varepsilon \equiv 0$ and

$$\sigma_\varepsilon(i, j) := \begin{cases} +\sqrt{3}\varepsilon e_2 & \text{if } j = i + \varepsilon e_1, \\ -\sqrt{3}\varepsilon e_2 & \text{if } j = i - \varepsilon e_1, \\ 0 & \text{elsewhere in } \Omega_\varepsilon^1. \end{cases}$$

Trivially, $|\frac{\mu[\sigma_\varepsilon]}{\varepsilon}|(\Omega) \sim \frac{1}{\varepsilon^2}$, $\|\frac{\mu[\sigma_\varepsilon]}{\varepsilon}\|_{\text{flat}} \sim \frac{1}{\varepsilon}$, and $\mathcal{F}_\varepsilon(\mu[\sigma_\varepsilon]) = F_\varepsilon(u_\varepsilon, \sigma_\varepsilon) \equiv 0$.

3.2. Proof of Theorem 1.1. We are now in a position to prove Theorem 1.1.

Proof of Theorem 1.1(i). For every $\varepsilon > 0$, let $(u_\varepsilon, \sigma_\varepsilon) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega)$ with $\mu[\sigma_\varepsilon] = \mu_\varepsilon$ be such that

$$(3.3) \quad F_\varepsilon(u_\varepsilon, \sigma_\varepsilon) \leq 2\mathcal{F}_\varepsilon(\mu_\varepsilon).$$

We let

$$(3.4) \quad \mathcal{K}_\varepsilon := \{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega) : \mu_\varepsilon(T_\varepsilon) = 0\} \quad K_\varepsilon := \bigcup_{T_\varepsilon \in \mathcal{K}_\varepsilon} T_\varepsilon,$$

and we set $\beta_\varepsilon := \beta^{u_\varepsilon, \sigma_\varepsilon, \mathcal{K}_\varepsilon}$ and $\hat{\beta}_\varepsilon := \hat{\beta}^{u_\varepsilon, \sigma_\varepsilon, \mathcal{K}_\varepsilon}$, where $\beta^{u_\varepsilon, \sigma_\varepsilon, \mathcal{K}_\varepsilon}$ and $\hat{\beta}^{u_\varepsilon, \sigma_\varepsilon, \mathcal{K}_\varepsilon}$ are the fields provided by Lemmas 3.2 and 3.3, respectively. Furthermore, we set $\tilde{\mu}_\varepsilon := \frac{\mu_\varepsilon}{\varepsilon}$ and $\tilde{\beta}_\varepsilon := \frac{\hat{\beta}_\varepsilon}{\varepsilon}$. Recalling the definition of $\Omega_\varepsilon(\tilde{\mu}_\varepsilon)$ in (2.1) and using, in order of appearance, Lemma 3.3(iii), Lemma 3.2(b), (3.3) and the energy bound, we have

$$(3.5) \quad \begin{aligned} \frac{1}{2} \int_{\Omega_\varepsilon(\tilde{\mu}_\varepsilon)} \mathbb{C} \tilde{\beta}_\varepsilon : \tilde{\beta}_\varepsilon dx &= \frac{1}{2\varepsilon^2} \int_{\Omega_\varepsilon(\tilde{\mu}_\varepsilon)} \mathbb{C} \hat{\beta}_\varepsilon : \hat{\beta}_\varepsilon dx \leq \frac{C}{2\varepsilon^2} \int_{K_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon dx \\ &\leq \frac{C}{\varepsilon^2} F_\varepsilon(u_\varepsilon, \sigma_\varepsilon) \leq \frac{C}{\varepsilon^2} \mathcal{F}_\varepsilon(\mu_\varepsilon) \leq C |\log \varepsilon|; \end{aligned}$$

moreover, by Lemma 3.4 and by the energy bound, we have

$$(3.6) \quad |\tilde{\mu}_\varepsilon|(\Omega) \leq C |\log \varepsilon|.$$




FIGURE 1. Geometric construction used in the proof of Theorem 1.1(iii). The position of each dislocation is approximated with a point $x^{k,\varepsilon}$ sitting on the barycenter of a triangle of $\mathcal{T}_\varepsilon(\Omega)$. The approximate displacement is the restriction to the lattice points of a continuum displacement having a jump on each half-line $\gamma^{k,\varepsilon}$. The sets $S_{k,\varepsilon}^1$ defined in the proof contain bonds across the jumps: for example, the points i , $i \pm \varepsilon \eta$ displayed in the picture are such that both bonds $(i, i - \varepsilon \eta)$, $(i, i + \varepsilon \eta)$ belong to the set $S_{k,\varepsilon}^1$.

In view of Lemma 3.3(ii) and of Lemma 3.2(c), we have $\tilde{\beta}_\varepsilon \in \mathcal{AS}_\varepsilon(\tilde{\mu}_\varepsilon)$ for every $\varepsilon > 0$ (with $A = \Omega$ in (2.2)), so that, by (3.5) and (3.6), applying Theorem 2.1(i) with $\mu_\varepsilon = \tilde{\mu}_\varepsilon$ and $\beta_\varepsilon = \tilde{\beta}_\varepsilon$, we deduce the claim. \square

Proof of Theorem 1.1(ii). We can assume without loss of generality that $\mathcal{F}_\varepsilon(\mu_\varepsilon) \leq C\varepsilon^2 |\log \varepsilon|$, which in view of Lemma 3.4, implies that $|\frac{\mu_\varepsilon}{\varepsilon}|(\Omega) \leq C |\log \varepsilon|$.

Let $A \subset\subset \Omega$ be an open bounded subset of \mathbb{R}^2 with Lipschitz continuous boundary such that $\text{supp } \mu \subset A$. Let $\varepsilon > 0$ small enough such that $A \subset \Omega_\varepsilon$. Let moreover $(u_\varepsilon, \sigma_\varepsilon) \in \mathcal{D}_\varepsilon(\Omega) \times \mathcal{S}_\varepsilon(\Omega)$ with $\mu[\sigma_\varepsilon] = \mu_\varepsilon$ be such that

$$(3.7) \quad \mathcal{F}_\varepsilon(\mu_\varepsilon) \leq F_\varepsilon(u_\varepsilon, \sigma_\varepsilon) + \varepsilon^2,$$

and let \mathcal{K}_ε be defined by (3.4); we set $\beta_\varepsilon := \beta^{u_\varepsilon, \sigma_\varepsilon, \mathcal{K}_\varepsilon}$, where $\beta^{u_\varepsilon, \sigma_\varepsilon, \mathcal{K}_\varepsilon}$ is the field provided by Lemma 3.2, $\tilde{\mu}_\varepsilon := \frac{\mu_\varepsilon}{\varepsilon}$ and $\tilde{\beta}_\varepsilon := \frac{\beta_\varepsilon}{\varepsilon}$. By (3.7) and Lemma 3.2 we have that $\tilde{\beta}_\varepsilon \in \mathcal{AS}_\varepsilon(\tilde{\mu}_\varepsilon)$ (for every $\varepsilon > 0$) and

$$\mathcal{F}_\varepsilon(\mu_\varepsilon) \geq F_\varepsilon(u_\varepsilon, \sigma_\varepsilon) - \varepsilon^2 \geq \frac{\sqrt{3}}{2} \varepsilon^2 E_\varepsilon(\tilde{\beta}_\varepsilon; A_\varepsilon(\tilde{\mu}_\varepsilon)) - \varepsilon^2,$$

whence the claim follows by applying Theorem 2.1(ii) with $\mu_\varepsilon = \tilde{\mu}_\varepsilon$ and $\beta_\varepsilon = \tilde{\beta}_\varepsilon$. \square

Proof of Theorem 1.1(iii). Let $\mu = \sum_{k=1}^K b^k \delta_{x^k} \in X(\Omega)$. By standard density arguments in Γ -convergence we can assume that $\varphi(b^k) = \psi(b^k)$ for every $k = 1, \dots, K$, i.e., that $|b^k| = 1$ for every $k = 1, \dots, K$. Let $\varepsilon \mathbb{T}_+^v(\Omega)$ be the set of the barycenters of the triangles $T_\varepsilon^+ = i + \varepsilon T^+ \in \mathcal{T}_\varepsilon(\Omega)$.

For every $k = 1, \dots, K$ and for every $\varepsilon > 0$, let $x^{k,\varepsilon} \in \varepsilon \mathbb{T}_+^v(\Omega)$ be such that

$$|x^{k,\varepsilon} - x^k| = \min\{|y - x^k| : y \in \varepsilon \mathbb{T}_+^v(\Omega)\}.$$

For every $k = 1, \dots, K$ set (see Figure 1)

$$\gamma^{k,\varepsilon} := \begin{cases} \{x^{k,\varepsilon} + \lambda e_1 : \lambda \geq 0\} & \text{if } b^k = \pm e_1, \\ \{x^{k,\varepsilon} + \lambda \nu : \lambda \geq 0\} & \text{if } b^k = \pm \nu, \\ \{x^{k,\varepsilon} + \lambda \eta : \lambda \geq 0\} & \text{if } b^k = \pm \eta. \end{cases}$$

Let $\bar{u}^{k,\varepsilon} \in C^2(\mathbb{R}^2 \setminus \gamma^{k,\varepsilon}; \mathbb{R}^2)$ be a function satisfying the following properties,

$$(3.8) \quad S_{\bar{u}^{k,\varepsilon}} = \gamma^{k,\varepsilon}, \quad [\bar{u}^{k,\varepsilon}] = b^k \text{ on } \gamma^{k,\varepsilon}, \quad \nabla \bar{u}^{k,\varepsilon}(\cdot) = \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(\cdot - x^{k,\varepsilon}),$$

Therefore, (3.9) is proved once provided that for every fixed $k, l = 1, \dots, K$ with $l \neq k$ it holds

$$(3.10) \quad \limsup_{\varepsilon \rightarrow 0} \frac{1}{2\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_{\frac{1}{\varepsilon}}^1} [(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j)) \cdot (j-i)]^2 = \frac{\sqrt{3}}{2} \psi(b^k),$$

$$(3.11) \quad \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_{\frac{1}{\varepsilon}}^1} [(du^{l,\varepsilon}(i,j) - \sigma^{l,\varepsilon}(i,j)) \cdot (j-i)]^2 = 0,$$

$$(3.12) \quad \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_{\frac{1}{\varepsilon}}^1} [(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j)) \cdot (j-i)] \\ [(du^{l,\varepsilon}(i,j) - \sigma^{l,\varepsilon}(i,j)) \cdot (j-i)] = 0,$$

$$(3.13) \quad \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in \Omega_{\varepsilon}^1 \setminus (\bigcup_{l=1}^K B_r(x^{l,\varepsilon}))_{\frac{1}{\varepsilon}}^1} [(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j)) \cdot (j-i)]^2 = 0.$$

Now, by the very definition of $\beta_{\mathbb{R}^2}^{b,\mathbb{C}}$ in (2.4), for every $b \in \mathbb{T}$, we have that there exists a universal constant $C > 0$ such that

$$(3.14) \quad |\beta_{\mathbb{R}^2}^{b,\mathbb{C}}(\rho, \theta)| \leq \frac{C}{\rho}, \quad |\nabla \beta_{\mathbb{R}^2}^{b,\mathbb{C}}(\rho, \theta)| \leq \frac{C}{\rho^2};$$

moreover, by the very definition of $u^{k,\varepsilon}$ and $\sigma^{k,\varepsilon}$,

$$(3.15) \quad \frac{du^{k,\varepsilon}(i,j)}{\varepsilon} - \frac{\sigma^{k,\varepsilon}(i,j)}{\varepsilon} = \int_0^1 \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(i + t(j-i) - x^{k,\varepsilon})(j-i) dt,$$

for every $k = 1, \dots, K$, for every $T_{\varepsilon} \in \mathcal{T}_{\varepsilon}$, and for every $(i,j) \in (T_{\varepsilon})_{\frac{1}{\varepsilon}}^1$. As a consequence, for every $k = 1, \dots, K$, for every $T_{\varepsilon} \in \mathcal{T}_{\varepsilon}$ with $\mu_{\varepsilon}(T_{\varepsilon}) = 0$, for every $(i,j) \in (T_{\varepsilon})_{\frac{1}{\varepsilon}}^1$, and for every $x \in T_{\varepsilon}$, we get

$$(3.16) \quad \frac{1}{\varepsilon^2} \left[(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j)) \cdot (j-i) \right]^2 = \left[\left(\frac{du^{k,\varepsilon}(i,j)}{\varepsilon} - \frac{\sigma^{k,\varepsilon}(i,j)}{\varepsilon} \right) \cdot (j-i) \right]^2 \\ = \varepsilon^4 \left[\int_0^1 \left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(i + t(j-i) - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) dt \right]^2 \\ \leq \varepsilon^4 \int_0^1 \left[\left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(i + t(j-i) - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) \right]^2 dt \\ = \varepsilon^4 \left[\left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(x - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) \right]^2 \\ + \varepsilon^4 \int_0^1 \left(\left[\left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(i + t(j-i) - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) \right]^2 \right. \\ \left. - \left[\left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(x - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) \right]^2 \right) dt,$$

where the second equality follows from (3.15), and the inequality is a consequence of Jensen's inequality. Now, by (3.14) for every $k = 1, \dots, K$ and for every $\xi \in \{\pm e_1, \pm \nu, \pm \eta\}$ it holds

$$|\nabla([\xi^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}} \xi]^2)(x - x^{k,\varepsilon})| \leq \frac{C}{|x - x^{k,\varepsilon}|^3} \quad \text{for every } x \neq x^{k,\varepsilon}.$$

In particular, for every $k = 1, \dots, K$, for every $T_{\varepsilon} \in \mathcal{T}_{\varepsilon}$ with $\mu_{\varepsilon}(T_{\varepsilon}) = 0$, for every $(i,j) \in (T_{\varepsilon})_{\frac{1}{\varepsilon}}^1$, and for every $x \in T_{\varepsilon}$,

$$\left[\left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(i + t(j-i) - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) \right]^2 - \left[\left(\frac{j-i}{\varepsilon} \right)^* \beta_{\mathbb{R}^2}^{b^k,\mathbb{C}}(x - x^{k,\varepsilon}) \left(\frac{j-i}{\varepsilon} \right) \right]^2 \leq \frac{C \varepsilon}{|x - x^{k,\varepsilon}|^3}.$$

Hence, for every $T_\varepsilon \in \mathcal{T}_\varepsilon$ with $\mu_\varepsilon(T_\varepsilon) = 0$, by (3.16) and integrating over T_ε , we get

$$\begin{aligned}
& F_\varepsilon(u^{k,\varepsilon}, \sigma^{k,\varepsilon}; T_\varepsilon) \\
& \leq \frac{4}{\sqrt{3}} \varepsilon^2 \int_{T_\varepsilon} \left(|e_1^* \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) e_1|^2 + |\nu^* \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) \nu|^2 + |\eta^* \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) \eta|^2 \right) dx \\
(3.17) \quad & + C \varepsilon^3 \int_{T_\varepsilon} \frac{1}{|x - x^{k,\varepsilon}|^3} dx \\
& = \sqrt{3} \varepsilon^2 \int_{T_\varepsilon} \frac{1}{2} \mathbb{C} \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) : \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) dx + C \varepsilon^3 \int_{T_\varepsilon} \frac{1}{|x - x^{k,\varepsilon}|^3} dx,
\end{aligned}$$

where the equality follows by the second equality in (3.1) with β replaced by $\beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon})$. Now, denoting by $T_\varepsilon^{x^{k,\varepsilon}} \in \mathcal{T}_\varepsilon(\Omega)$ the triangle centered at $x^{k,\varepsilon}$, by (3.14) and (3.15) for every $k = 1, \dots, K$ we get

$$F_\varepsilon(u^{k,\varepsilon}, \sigma^{k,\varepsilon}; T_\varepsilon^{x^{k,\varepsilon}}) \leq C \varepsilon^2,$$

for some constant C independent of ε . Moreover, recalling the notation previously introduced for the annuli, an integration in polar coordinates shows that

$$\int_{A_{\delta, 2r}(x^{k,\varepsilon})} \frac{1}{|x - x^{k,\varepsilon}|^3} dx \leq \frac{C}{\delta},$$

for every $0 < \delta < 2r$. Therefore, by (3.17) and (2.6), for every $k = 1, \dots, K$ and for $c > 0$ small enough (as for instance $c < \frac{\sqrt{3}}{6}$) we obtain

$$\begin{aligned}
& \frac{1}{2\varepsilon^2} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_\varepsilon^{\frac{1}{2}}} \left[\left(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j) \right) \cdot (j - i) \right]^2 \\
(3.18) \quad & \leq \frac{1}{2} \sum_{T_\varepsilon \in \mathcal{T}_\varepsilon(B_{2r}(x^{k,\varepsilon}))} F_\varepsilon(u^{k,\varepsilon}, \sigma^{k,\varepsilon}; T_\varepsilon) \\
& \leq \frac{\sqrt{3}}{2} \varepsilon^2 \int_{A_{c\varepsilon, 2r}(x^{k,\varepsilon})} \frac{1}{2} \mathbb{C} \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) : \beta_{\mathbb{R}^2}^{b^k, \mathbb{C}}(x - x^{k,\varepsilon}) dx + C \varepsilon^2 \\
& \leq \varepsilon^2 |\log \varepsilon| \frac{\sqrt{3}}{2} \psi(b^k) + C \varepsilon^2,
\end{aligned}$$

where the constant C changes from line to line; this proves (3.10).

Moreover, by (3.17) together with (3.14), for every $k, l = 1, \dots, K$ with $l \neq k$, we have

$$\begin{aligned}
& \frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_\varepsilon^{\frac{1}{2}}} \left[(du^{l,\varepsilon}(i,j) - \sigma^{l,\varepsilon}(i,j)) \cdot (j - i) \right]^2 \\
(3.19) \quad & \leq \frac{1}{\varepsilon^2 |\log \varepsilon|} \sum_{T_\varepsilon \in \mathcal{T}_\varepsilon(B_{2r}(x^{k,\varepsilon}))} F_\varepsilon(u^{l,\varepsilon}, \sigma^{l,\varepsilon}; T_\varepsilon) \\
& \leq \frac{C}{|\log \varepsilon|} \left(\int_{B_{2r}(x^{k,\varepsilon})} \mathbb{C} \beta_{\mathbb{R}^2}^{b^l, \mathbb{C}}(x - x^{l,\varepsilon}) : \beta_{\mathbb{R}^2}^{b^l, \mathbb{C}}(x - x^{l,\varepsilon}) dx + \varepsilon \int_{B_{2r}(x^{k,\varepsilon})} \frac{1}{|x - x^{l,\varepsilon}|^3} dx \right) \\
& \leq \frac{C_r}{|\log \varepsilon|} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,
\end{aligned}$$

where C_r depends only on r . Now, by (3.19) we obtain (3.11). Analogously, one can prove that also (3.13) holds true.

Finally, by the Hölder inequality, using (3.18) and (3.19), for every $k, l = 1, \dots, K$ with $l \neq k$ we get

$$\begin{aligned} & \frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_\varepsilon^1} [(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j)) \cdot (j-i)] [(du^{l,\varepsilon}(i,j) - \sigma^{l,\varepsilon}(i,j)) \cdot (j-i)] \\ & \leq \left(\frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{k,\varepsilon}))_\varepsilon^1} [(du^{k,\varepsilon}(i,j) - \sigma^{k,\varepsilon}(i,j)) \cdot (j-i)]^2 \right)^{\frac{1}{2}} \\ & \quad \left(\frac{1}{\varepsilon^4 |\log \varepsilon|} \sum_{(i,j) \in (B_{2r}(x^{l,\varepsilon}))_\varepsilon^1} [(du^{l,\varepsilon}(i,j) - \sigma^{l,\varepsilon}(i,j)) \cdot (j-i)]^2 \right)^{\frac{1}{2}} \\ & \leq \frac{C}{\sqrt{|\log \varepsilon|}} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

which proves (3.12) and concludes the proof of the Γ -limsup inequality. \square

4. DEGENERACIES OF THE MODEL AND POSSIBLE KINEMATIC CONSTRAINTS

We highlight that the functional (1.2) exhibits some degeneracies that are typical of linearized energies as well as of discrete models built on a reference configuration; the presence of the plastic slip field σ actually makes the system even less rigid and produces even more unphysical configurations.

As an example, let $\Omega := [-1, 1]^2$ and define $u_\varepsilon^\pm \in \mathcal{D}_\varepsilon(\Omega)$ as

$$(4.1) \quad u_\varepsilon^\pm(i) := \begin{cases} \pm \varepsilon \nu & \text{if } i_2 \leq 0 \\ 0 & \text{elsewhere in } \Omega_\varepsilon^0. \end{cases}$$

Notice that u_ε^- can be interpreted as a microscopic crack, whereas u_ε^+ can be seen as an unphysical crack which produces interpenetration of matter generating the superposition of two lines of atoms. It is easy to check that there exist $\sigma_\varepsilon^\pm \in \mathcal{S}_\varepsilon(\Omega)$ such that $F_\varepsilon(u_\varepsilon^\pm, \sigma_\varepsilon^\pm) = 0$. Moreover, up to slightly modifying u_ε^- , one can construct a function $u_\varepsilon \in \mathcal{D}_\varepsilon(\Omega)$ such that the crack does not disconnect Ω and has two ending points which in our model are classified as dislocations with opposite Burgers vector. In particular, what in our model is identified as a plastic slip, in some cases should be rather understood as a microscopic crack opening.

Furthermore, we observe that the energy in (1.2) is invariant with respect to integer dilations of the lattice. In order to see this, it is enough to fix $\lambda \in \mathbb{Z}$ and to take the displacement u_ε^λ and the slip $\sigma_\varepsilon^\lambda$ defined by

$$(4.2) \quad u_\varepsilon^\lambda(i) := \lambda \varepsilon i \quad \sigma_\varepsilon^\lambda(i, j) := \varepsilon \lambda (j - i).$$

Trivially, $du_\varepsilon^\lambda(i, j) \equiv \sigma_\varepsilon^\lambda(i, j)$ and hence $F_\varepsilon(u_\varepsilon^\lambda, \sigma_\varepsilon^\lambda) \equiv 0$.

In view of the degeneracies pointed out above, one may ask whether additional suitable kinematic constraints on the plastic slips could prevent such unphysical behaviors. In the present framework, it seems natural to incorporate linearized constraints on the plastic slips, mimicking deviatoric/pure shear volume preserving conditions on each ε -triangle $T_\varepsilon \in \mathcal{T}_\varepsilon$.

Let us discuss now a possible way to introduce such a kinematic constraint in our model. In order to (formally) linearize the nonlinear volume preserving constraint, consider deformation gradients $\text{Id} + \delta z$ close to the identity, where $z \in \mathbb{R}^{2 \times 2}$ represents a plastic deformation gradient, $\delta > 0$ is a small parameter with respect to which the linearization is performed and Id is the identity matrix in $\mathbb{R}^{2 \times 2}$. Enforcing that the deformed configuration $(\text{Id} + \delta z)(T_\varepsilon)$ has the same (oriented) area of T_ε we obtain

$$[(k-i) + \delta z(k-i)] \wedge [(j-i) + \delta z(j-i)] = (k-i) \wedge (j-i),$$

for every triple (i, j, k) of counterclockwise oriented vertices of T_ε . Neglecting lower order terms, this yields

$$(4.3) \quad z(k-i) \wedge (j-i) - z(j-i) \wedge (k-i) = 0,$$

for every triple (i, j, k) as above. Summing (4.3) over two triples of the type (i, j, k) , (j, k, i) , one can see that condition (4.3) is equivalent to the well known trace free constraint on z ; we refer the interested reader to [17] for a rigorous derivation via Γ -convergence of such a kind of trace free constraints starting from continuous nonlinear models in plasticity, relying on multiplicative rather than additive decompositions of the deformation gradient.

We now rewrite (4.3) in terms of the (linear) slip field σ , which is, by construction, given by

$$(4.4) \quad \sigma(i, j) = z(j-i) \quad \text{for every bond } (i, j) \text{ of the triangle } T_\varepsilon.$$

Condition (4.3) becomes

$$(4.5) \quad \sigma(i, k) \wedge (j-i) - \sigma(i, j) \wedge (k-i) = 0,$$

for every triple (i, j, k) of counterclockwise oriented vertices of T_ε . As a consequence of (4.4), we have that $\mu[\sigma](T_\varepsilon) = 0$, and hence (4.5) is justified by the reasoning above only on dislocation-free triangles. On the other hand, one can easily check that the only condition (4.5) (assumed for every triple (i, j, k) of counterclockwise oriented vertices of T_ε) implies that $\mu[\sigma](T_\varepsilon) = 0$. Summarizing, in our framework, a reasonable linearization of the volume preserving constraint seems to be provided by condition (4.5) assumed on all dislocation-free ε -triangles T_ε .

Notice that (4.5) rules out the degeneracies pointed out by the examples above. More precisely, one can check that, given a displacement as in (4.1) or (4.2), the minimum energy in a dislocation-free triangle among all slip fields fulfilling (4.5) is strictly positive. On the other hand (MS) guarantees that dislocation-free triangles are necessarily present. Finally, our Γ -convergence result still holds true under the additional constraint (4.5), which is satisfied by the recovery sequence constructed in the proof of Theorem 1.1(iii).

REFERENCES

- [1] Alicandro, R., De Luca, L., Garroni, A., Ponsiglione: Metastability and dynamics of discrete topological singularities in two dimensions: A Γ -convergence approach. *Arch. Rational Mech. Anal.* **214** (2014), 269–330.
- [2] Alicandro, R., Lazzaroni, G., Palombaro, M.: Derivation of linear elasticity for a general class of atomistic energies. *SIAM J. Math. Anal.* **53** (2021), 5060–5093.
- [3] Ariza, M. P.; Ortiz, M.: Discrete crystal elasticity and discrete dislocations in crystals. *Arch. Rational Mech. Anal.* **178** (2005), 149–226.
- [4] Bach, A., Cicalese, M., Kreutz, L., Orlando, G.: The antiferromagnetic XY model on the triangular lattice: topological singularities. To appear in *Indiana Univ. Math. J.*
- [5] Bacon, D.J., Barnett, D.M., Scattergood, R.O.: Anisotropic continuum theory of lattice defects. *Progress Mater. Sci.* **23** (1978), 51–262.
- [6] Braides, A., Solci, M., Vitali, E.: A derivation of linear elastic energies from pair-interaction atomistic systems. *Netw. Heterog. Media* **2** (2007), 551–567.
- [7] Cermelli, P., Leoni, G.: Renormalized energies and forces on dislocations. *SIAM J. Math. Anal.* **37** (2005), 1131–1160.
- [8] De Luca, L: Γ -convergence analysis for discrete topological singularities: The anisotropic triangular lattice and the long range interaction energy. *Asymptot. Anal.* **96** (2016), 185–221.
- [9] De Luca, L., Garroni, A., Ponsiglione, M.: Γ -convergence analysis of systems of edge dislocations: the self energy regime. *Arch. Rational Mech. Anal.* **206** (2012), 885–910.
- [10] Fanzon, S., Palombaro, M., Ponsiglione, M.: Derivation of Linearised Polycrystals from a Two-Dimensional System of Edge Dislocations. *SIAM J. Math. Anal.* **51** (2019), 3956–3981.
- [11] Garroni, A., Leoni, G., Ponsiglione, M.: Gradient theory for plasticity via homogenization of discrete dislocations. *J. Eur. Math. Soc.* **12** (2010), 1231–1266.
- [12] Ginster, J.: Plasticity as the Γ -Limit of a Two-dimensional Dislocation Energy: The Critical Regime Without the Assumption of Well-Separateness, *Arch. Rational Mech. Anal.* **233** (2019), 1253–1288.
- [13] Giuliani, A., Theil, F.: Long range order in atomistic models for solids, to appear in *J. Eur. Math. Soc.*
- [14] Hirth, J.P., Lothe, J.: *Theory of Dislocations*, Krieger Publishing Company, Malabar, Florida, 1982.
- [15] Hull, D., Bacon, D.J.: *Introduction to dislocations*, Butterworth-Heinemann, 2011.
- [16] Lauteri, G., Luckhaus, S.: An energy estimate for dislocation configurations and the emergence of Cosserat-type structures in metal plasticity. arXiv:1608.06155, 2016.
- [17] Mielke, A., Stefanelli, U.: Linearized plasticity is the evolutionary Gamma-limit of finite plasticity. *J. Eur. Math. Soc.* **15** (2013), 923–948.
- [18] Sandier, E., Serfaty, S.: *Vortices in the Magnetic Ginzburg-Landau Model*, Progress in Nonlinear Differential Equations and Their Applications, vol. 70, Birkhäuser Boston, Boston (MA), 2007.
- [19] Schmidt, B.: On the derivation of linear elasticity from atomistic models. *Netw. Heterog. Media* **4** (2009), 789–812.

(Roberto Alicandro) DIEI, UNIVERSITÀ DI CASSINO E DEL LAZIO MERIDIONALE, VIA DI BIASIO 43, 03043 CASSINO (FR), ITALY

E-mail address, R. Alicandro: alicandr@unicas.it

(Lucia De Luca (Corresponding author)) ISTITUTO PER LE APPLICAZIONI DEL CALCOLO “M. PICONE”, IAC-CNR, VIA DEI TAURINI 19, 00185 ROMA, ITALY

E-mail address, L. De Luca: lucia.deluca@cnr.it

(Giuliano Lazzaroni) DIMAI, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY

E-mail address, G. Lazzaroni: giuliano.lazzaroni@unifi.it

(Mariapia Palombaro) DISIM, UNIVERSITÀ DELL’AQUILA, VIA VETOIO, 67100 L’AQUILA, ITALY

E-mail address, M. Palombaro: mariapia.palombaro@univaq.it

(Marcello Ponsiglione) DIP. DI MATEMATICA, UNIV. ROMA-I “LA SAPIENZA”, PIAZZALE ALDO MORO 5, 00185 ROMA, ITALY

E-mail address, M. Ponsiglione: ponsigli@mat.uniroma1.it