

EMERGENCE OF WULFF-CRYSTALS FROM ATOMISTIC SYSTEMS ON THE FCC AND HCP LATTICES

MARCO CICALESSE, LEONARD KREUTZ, AND GIAN PAOLO LEONARDI

ABSTRACT. We consider a system of N hard spheres sitting on the nodes of either the FCC or HCP lattice and interacting via a sticky-disk potential. As N tends to infinity (continuum limit), assuming the interaction energy does not exceed that of the ground-state by more than $N^{2/3}$ (surface scaling), we obtain the variational coarse grained model by Γ -convergence. More precisely, we prove that the continuum limit energies are of perimeter type and we compute explicitly their Wulff shapes. Our analysis shows that crystallization on FCC is preferred to that on HCP for N large enough. The method is based on integral representation and concentration-compactness results that we prove for general periodic lattices in any dimension.

1. INTRODUCTION

A fundamental problem in crystallography is to understand why ensembles of large number of atoms arrange themselves into crystals at low temperatures. From the mathematical point of view, proving that equilibrium configurations of certain phenomenological interaction energies exhibit these structures is referred to as the crystallization problem [8].


At zero temperature the internal energy of a configuration of atoms is expected to be solely governed by its geometric arrangement. Within the framework of molecular mechanics [1, 24, 31], one identifies each ensemble of atoms with its atomic positions $X = \{x_1, \dots, x_N\} \subset \mathbb{R}^3$ and associates to it a configurational energy of the form

$$\mathcal{E}(X) := \frac{1}{2} \sum_{i \neq j} V(|x_i - x_j|),$$

where $V: \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ is an empirical pair interaction potential (the factor $\frac{1}{2}$ accounts for double counting). Such potentials are typically repulsive at short distances and attractive at large distances. While clustering is favored by long range attraction, the density of a cluster cannot get too large due to short-range repulsion.

Notably, even under simplifying assumptions on the interaction potentials, the mathematical literature on rigorous crystallization results is scarce. In fact, for finite N , only results in one and two space dimensions are available. For example, if V is of Lennard–Jones type, crystallization has been proved only in one space dimension [26]. In higher space dimensions only partial results are available. Most notably, in [18, 20, 39] it has been proven that crystalline structures have optimal bulk energy scaling. In two dimensions, only results for (some variants of) the sticky

Key words and phrases. Wulff shape, isoperimetric inequality, crystallization, Γ -convergence, lattice systems, concentration-compactness.

FIGURE 1. The sticky disc interaction potential V .

disc potential (see Fig. 1)

$$V(r) := \begin{cases} +\infty & \text{if } r < 1, \\ -1 & \text{if } r = 1, \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

are available [16, 28, 30, 36]. More recently, crystallization results have been proved for ionic compounds [21, 22] and carbon structures in [35]. The potential given in (1) models the atoms as hard spheres that interact exactly when two of them are tangent. In \mathbb{R}^n the kissing number $k(n)$ is the highest number of n -dimensional spheres of radius $\frac{1}{2}$ which are tangent to a given sphere of the same size. It is well known that $k(2) = 6$ and $k(3) = 12$, see [37]. For a given configuration of non-overlapping equal balls centered at $X = \{x_1, \dots, x_N\} \subset \mathbb{R}^n$, $N \in \mathbb{N} \cup \{+\infty\}$ the coordination number of $x \in X$ is the number of spheres centered at $y \in X \setminus \{x\}$ and tangent to the one centered at x . In two dimensions there is a unique (up to a rigid motion) configuration made of infinitely many particles such that all atoms have as coordination number the kissing number. Such a set X is the triangular lattice with lattice spacing one. In three dimensions the problem is much more intricate. In fact, there exist infinitely many configurations with constant coordination number equal to $k(3)$. An infinite class of configurations can be obtained by stacking in an appropriate way layers of triangular lattices. A remarkable result by Hales [27] shows that all such structures solve Kepler's conjecture which is to say that they have the maximal packing density in \mathbb{R}^3 . Two notable cases of the aforementioned structures are the face-centered cubic lattice \mathcal{L}_{FCC} and the hexagonal closed-packed lattice \mathcal{L}_{HCP} (see (7)–(9) for their precise definition) which are the most prevalent among the crystalline arrangements in the periodic table of elements.

In this paper we want to investigate already crystallized configurations, i.e. configurations $X \subset \mathcal{L}$ where $\mathcal{L} = \mathcal{L}_{\text{FCC}}$ or $\mathcal{L} = \mathcal{L}_{\text{HCP}}$. For such $X = \{x_1, \dots, x_N\} \subset \mathcal{L}$, fixing the lattice spacing to be 1, we have

$$\mathcal{E}(X) = \frac{1}{2} \sum_{i \neq j} V(|x_i - x_j|) = - \sum_{i=1}^N \#(\mathcal{N}(x_i) \cap X), \text{ where } \mathcal{N}(x) = \{y \in \mathcal{L} : |x - y| = 1\}.$$

As described above the minimal energy per atom is $-k(3) = -12$. Further information on \mathcal{E} as N grows can be obtained by referring it to the minimal energy per atom and calculating the *excess energy* $E_N(X)$ defined below. More precisely, in Theorem 2.3, we carry out a rigorous

variational asymptotic expansion (see [10]) of $\mathcal{E}(X)$, by considering

$$E_N(X) = N^{-2/3} (\mathcal{E}(X) + 12N) = N^{-2/3} \sum_{i=1}^N (12 - \#(\mathcal{N}(x) \cap X)) \quad (2)$$

and calculating its Γ -limit [9, 13] as N tends to infinity. This analysis has been done in two dimensions for configurations confined to the triangular lattice [7] as well as without any confinement assumption [23]. Note that, the scaling factor $N^{-2/3}$ is used in order to keep the energy bounded as the number of atoms grows. In fact, given a low energy configuration of N atoms, the number of those contributing to the energy scales like $N^{2/3}$ for N large. By associating to each configuration its rescaled empirical measure

$$\mu_N(X) := \frac{1}{N} \sum_{i=1}^N \delta_{N^{-1/3}x_i},$$

we show in Theorem 2.3 i) that the sequence of rescaled energies (2) is equi-coercive with respect to the weak*-convergence of the associated empirical measures. In Theorem 2.3 ii), iii) we exploit integral representation theorems [2, 3, 5] to show that the limit energy is finite on the set of measures $\mu = \sqrt{2}\mathcal{L}^3 \llcorner V$, where $V \subset \mathbb{R}^3$ is a set of finite perimeter, on which the energy takes the form

$$E_{\mathcal{L}}(\mu) := \int_{\partial^*V} \varphi_{\mathcal{L}}(\nu) d\mathcal{H}^2. \quad (3)$$

Here, ∂^*V denotes the reduced boundary of the set V , $\nu(x)$ denotes its unit outer normal at the point $x \in \partial^*V$ and $\varphi_{\mathcal{L}}$ is an anisotropic surface energy density depending on the underlying lattice \mathcal{L} . In the case of multi-lattices, like the HCP-lattice, this integral representation result has not yet been proven in the literature. We defer to Section 5 for a proof of this result whose main ingredient is the integral representation theorem in [3]. Furthermore, in the same section we prove general compactness and concentration lemmata that ensure the convergence of the rescaled empirical measures of minimizers of the discrete problem (2) to the Wulff shape (up to a constant density factor) of the associated limiting anisotropic perimeter energy (21). Such kind of result was previously known only in two dimensions [7]. Its extension to higher dimensions, see Lemma 5.16, requires more refined tools from geometric measure theory that, to the best of our knowledge, are exploited in this setting here for the first time. The main body of this work lies in the calculation of the surface energy density $\varphi_{\mathcal{L}}: \mathbb{R}^3 \rightarrow [0, +\infty)$ both for the FCC and the HCP lattices. Here, we take advantage of a recently proved finite cell formula [11]. Finally, for both lattices, we solve the associated isoperimetric problem [19]

$$m_{\mathcal{L}} := \min \left\{ \int_{\partial^*V} \varphi_{\mathcal{L}}(\nu) d\mathcal{H}^2 : |V| = 1 \right\} \quad (4)$$

by calculating the (up to translation unique) set realizing the minimum in (4), also known as the Wulff shape [40]. We show that $m_{\text{FCC}} < m_{\text{HCP}}$ which also implies (since Γ -convergence and coercivity implies the convergence of minimum values) that, for large number of atoms, crystallization on the face-centered cubic lattice is preferred to that on the hexagonal-closed packed lattice. We finally mention [6] for some preliminary computations on the Wulff shape of the FCC and HCP.

In contrast to the uniqueness of the Wulff crystal in the continuum setting, minimizers to the discrete isoperimetric problem [29] are non-unique [17]. Over the last years there has been a remarkable interest in establishing fluctuation estimates between different minimizers, i.e., estimating (several notions of) distances between different minimizers. Maximal fluctuation

estimates between two minimizers have been first conjectured in [7] in the case of the crystallization on the triangular lattice and have been later proved in [38] and [15]. The same estimates have been proved in [33, 21] and [14, 22] for the square and the honeycomb lattices, respectively. A general approach linking the quantitative anisotropic isoperimetric inequality to such fluctuation estimates has been set up in [12] by two of the authors. In dimensions larger than two these fluctuation estimates have been only established for the cubic lattice in [32] and for \mathbb{Z}^d in [34]. In order to establish the aforementioned fluctuation estimates, however, an understanding of the limiting macroscopic Wulff shape is essential. Since the present work yields these shapes for the FCC and HCP lattices, it is our opinion that it may be considered an indispensable first step to prove fluctuation estimates also for such lattices.

The article is structured as follows. In Section 2, we introduce the necessary mathematical preliminaries, the model, and the main results. In Section 3 we prove Proposition 2.4 and 2.5, by calculating the surface energy density as well as the Wulff crystal associated to both the FCC and the HCP lattices. In Section 4 we prove the main Γ -convergence Theorem 2.3. The latter is a consequence of a more general theory for discrete perimeter energies on general periodic lattices developed in Section 5.

2. SETTING AND NOTATION

Given a set of vectors $V \subset \mathbb{R}^n$ we denote by $\text{span}_{\mathbb{Z}}V$ the set of finite linear combinations of elements of V with coefficients in \mathbb{Z} . We denote by \mathfrak{M} the collection of all Lebesgue measurable subsets of \mathbb{R}^n . Given $A \in \mathfrak{M}$ we denote by $|V|$ its n -dimensional Lebesgue measure, i.e., $|V| = \mathcal{L}^n(V)$, and \mathcal{H}^n its n -dimensional Hausdorff measure. Given a countable set X , we denote by $\#X$ the cardinality of X . Given $a, b \in \mathbb{R}^n$ we denote by $\langle a, b \rangle$ their scalar product. We denote by \mathbb{S}^{n-1} the set of unitary vectors in \mathbb{R}^n . For any $\nu \in \mathbb{S}^{n-1}$ let $\{\nu_1, \dots, \nu_n = \nu\}$ be an orthonormal basis of \mathbb{R}^n , and let $Q^\nu := \{x \in \mathbb{R}^n : |\langle x, \nu_i \rangle| < 1/2, i = 1, \dots, n\}$ be a unit cube centered at the origin with faces parallel and orthogonal to ν . For $T > 0$ and $x \in \mathbb{R}^n$ we set $Q_T^\nu(x) = x + TQ^\nu$ and we write $Q_T^\nu = Q_T^\nu(0)$. For $r > 0$ and $x \in \mathbb{R}^n$ we denote by $B_r(x)$ the n -dimensional Euclidean ball of radius r centered at x (for $x = 0$ we write B_r in place of $B_r(0)$) and we set $\omega_n = |B_1(x)|$. For $r > 0$ and $A \subset \mathbb{R}^n$ we set $(A)_r = B_r + A$. Given $A \subset \mathbb{R}^n$ open, we define the set of positive Radon measures by $\mathcal{M}_+(A)$. We say that $\{\mu_k\}_k \subset \mathcal{M}_+(A)$ converges to $\mu \in \mathcal{M}_+(A)$ with respect to the weak star topology and we write $\mu_k \xrightarrow{*} \mu$ if

$$\lim_{k \rightarrow \infty} \int_A \varphi d\mu_k = \int_A \varphi d\mu \text{ for all } \varphi \in C_c(A).$$

we denote by $BV(A)$ the space of functions of bounded variation in A and we denote by $BV_{\text{loc}}(A) = \{u : u \in BV(K) \text{ for all } K \subset\subset A, K \text{ open}\}$. Given a function $u \in BV(A)$ we use the notation of [4] for the jump set $J(u)$ and the measure theoretic normal $\nu_u : J(u) \rightarrow \mathbb{S}^{n-1}$. For $V \subset A$, $V \in \mathfrak{M}$ we denote the relative perimeter of V in A by

$$\text{Per}(V, A) = \sup \left\{ \int_V \text{div } v \, dx : v \in C_c^\infty(A; \mathbb{R}^n), \|v\|_\infty \leq 1 \right\}.$$

For $\nu \in \mathbb{S}^{n-1}$ we set

$$u_\nu(i) := \begin{cases} 1 & \text{if } \langle x, \nu \rangle \geq 0; \\ 0 & \text{otherwise.} \end{cases} \quad (5)$$

In Section 2–Section 4 we set $n = 3$.




FIGURE 2. On the left: The FCC-lattice. On the right: The HCP-lattice. Pairs of points at distance one are connected via the dashed lines.

Definition of HCP and FCC lattices. In the following we define the *face-centered cubic lattice* (short FCC-lattice) and the *hexagonal closed-packed lattice* (short HCP-lattice). To this end, we introduce the vectors

$$b_1 := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad b_2 := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad b_3 := \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \quad (6)$$

and

$$e_1 := \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad e_2 := \frac{1}{2} \begin{pmatrix} 1 \\ \sqrt{3} \\ 0 \end{pmatrix}, \quad e_3 := \frac{2}{3}\sqrt{6} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad v_1 := \frac{1}{3}(e_1 + e_2) + \frac{1}{2}e_3. \quad (7)$$

We define the FCC-lattice as

$$\mathcal{L}_{\text{FCC}} := \text{span}_{\mathbb{Z}} \{b_1, b_2, b_3\} \quad (8)$$

and the HCP-lattice by

$$\mathcal{L}_{\text{HCP}} := \text{span}_{\mathbb{Z}} \{e_1, e_2, e_3\} \cup (\text{span}_{\mathbb{Z}} \{e_1, e_2, e_3\} + v_1). \quad (9)$$

The two lattices are illustrated in Figure 2. We shall write \mathcal{L} to generically denote one of the two lattices defined above. We define the *neighborhood* of a point $x \in \mathcal{L}_{\text{FCC}}$ as the set

$$\mathcal{N}_{\text{FCC}}(x) := \{\pm b_1, \pm b_2, \pm b_3, \pm(b_1 - b_2), \pm(b_1 - b_3), \pm(b_2 - b_3)\} + x. \quad (10)$$

Similarly, for a point $x \in \mathcal{L}_{\text{HCP}}$ we define its *neighborhood* as follows: if $x \in \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\}$ then

$$\mathcal{N}_{\text{HCP}}(x) := \{\pm e_1, \pm e_2, \pm(e_1 - e_2), v_1, v_1 - e_1, v_1 - e_2, v_1 - e_3, v_1 - e_1 - e_3, v_1 - e_2 - e_3\} + x, \quad (11)$$

while if $x \in \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\} + v_1$ then

$$\mathcal{N}_{\text{HCP}}(x) := \{\pm e_1, \pm e_2, \pm(e_1 - e_2), -v_1, e_1 - v_1, e_2 - v_1, e_3 - v_1, e_1 + e_3 - v_1, e_2 + e_3 - v_1\} + x. \quad (12)$$

Note that $\mathcal{N}_{\text{FCC}}(x) = \mathcal{N}_{\text{FCC}}(0) + x$ for all $x \in \mathcal{L}_{\text{FCC}}$, while this is no more the case for $x \in \mathcal{L}_{\text{HCP}}$. Also for \mathcal{N} we omit the subscript if we do not need to distinguish between FCC and HCP. It is

straightforward to check that for all $x, y \in \mathcal{L}$,

$$x \in \mathcal{N}(y) \iff |x - y| = 1.$$

Given \mathcal{L} we define the *Voronoi cell* of $x \in \mathcal{L}$ (with respect to \mathcal{L}) by

$$\mathcal{V}_{\mathcal{L}}(x) := \{y \in \mathbb{R}^3 : |y - x| \leq |y - z| \text{ for all } z \in \mathcal{L}\}. \quad (13)$$

Accordingly, given $\varepsilon > 0$ we write $\mathcal{V}_{\varepsilon\mathcal{L}}(x)$ for the Voronoi cell centered at $x \in \varepsilon\mathcal{L}$ with respect to the scaled lattice $\varepsilon\mathcal{L}$. Given $X \subset \varepsilon\mathcal{L}$ we say that $y \in \mathcal{N}_{\varepsilon}(x)$ if and only if $\varepsilon^{-1}y \in \mathcal{N}(\varepsilon^{-1}x)$.

Definition of the Energy. Given $X \subset \mathcal{L}$ and $A \subset \mathbb{R}^3$ we define the configurational energy of X localized to the set A as

$$E_{\mathcal{L}}(X, A) := \sum_{x \in X \cap A} (12 - \#\mathcal{N}(x) \cap X).$$

In the formula above we can interpret the set X as the occupancy of the crystal \mathcal{L} , i.e., the set of those nodes of \mathcal{L} occupied by atoms. The quantity $12 - \#\mathcal{N}_{\mathcal{L}}(x) \cap X$ is also known as the valence of the point x with respect to X , i.e., the number of neighbours missing in X in order to have a neighbourhood of maximal cardinality (the number 12 in the formula). Note that we can also rewrite the energy as

$$E_{\mathcal{L}}(X, A) = \frac{1}{2} \sum_{x \in \mathcal{L} \cap A} \sum_{y \in \mathcal{L}} c(x, y) |\chi_X(y) - \chi_X(x)|,$$

where

$$c(x, y) = \begin{cases} 1 & \text{if } y \in \mathcal{N}(x), \\ 0 & \text{otherwise.} \end{cases} \quad (14)$$

Periodicity of the interaction coefficients. By definition

$$\mathcal{L}_{\text{FCC}} = \mathcal{L}_{\text{FCC}} + b_1 = \mathcal{L}_{\text{FCC}} + b_2 = \mathcal{L}_{\text{FCC}} + b_3.$$

As a consequence of that, for any $x, y \in \mathcal{L}_{\text{FCC}}$ it holds that

$$c(x + b_1, y + b_1) = c(x + b_2, y + b_2) = c(x + b_3, y + b_3) = c(x, y).$$

According to the last two equalities, we say that the lattice \mathcal{L}_{FCC} as well as the interaction coefficients of its configurational energy are periodic with periodicity cell

$$T_{\text{FCC}} = \{\lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3 : \lambda_i \in [0, 1)\}, \quad (15)$$

or simply that they are T_{FCC} -periodic. Similarly, we observe that \mathcal{L}_{HCP} and its interaction coefficients are T_{HCP} -periodic, where the periodicity cell is defined as

$$T_{\text{HCP}} = \{\lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3 : \lambda_i \in [0, 1)\}. \quad (16)$$

Surface scaling of the configurational energy. For $\varepsilon > 0$ such that $\varepsilon^3 \#X \rightarrow 1$ as $\varepsilon \rightarrow 0$ we consider the following family of scaled energies

$$G_{\mathcal{L}, \varepsilon}(X) := \varepsilon^2 \sum_{x \in X} (12 - \#\mathcal{N}_{\varepsilon}(x) \cap X).$$

Note that, modeling the points in $X \subset \varepsilon\mathcal{L}$ as hard-spheres of ε diameter, the quantity $\varepsilon^3 \#X$ is of order one (according to the scaling assumption above) and proportional to the volume of the union of the spheres in X . Hence, the scaling factor ε^2 in the energy functional turns out to be

a surface scaling. We also define the rescaled empirical measures associated to the configuration X as

$$\mu_\varepsilon := \varepsilon^3 \sum_{x \in X} \delta_x. \quad (17)$$

Upon identifying $X \subset \varepsilon\mathcal{L}$ with its empirical measure μ_ε , we can regard these energies to be defined on $\mathcal{M}_+(\mathbb{R}^3)$ by setting

$$E_{\mathcal{L},\varepsilon}(\mu) := \begin{cases} G_{\mathcal{L},\varepsilon}(X) & \text{if } \mu = \mu_\varepsilon \text{ given in (17) for some } X \subset \varepsilon\mathcal{L}, \\ +\infty & \text{otherwise.} \end{cases} \quad (18)$$

The coarse grained continuum energy. For \mathcal{L} we define the *homogenized surface energy density* $\varphi_{\mathcal{L}}: \mathbb{R}^3 \rightarrow [0, +\infty]$ as the convex positively homogeneous function of degree one such that for all $\nu \in \mathbb{S}^2$ we have

$$\varphi_{\mathcal{L}}(\nu) := \lim_{T \rightarrow +\infty} \frac{1}{T^2} \inf \{ E_{\mathcal{L}}(X, Q_T^\nu) : X \subset \mathcal{L}, \chi_X(i) = u_\nu(i) \text{ for } i \in \mathcal{L} \setminus Q_{T-3}^\nu \}, \quad (19)$$

where u_ν is given by (5).

In order to be able to apply [11, Proposition 2.6] and eventually obtain an alternative representation of $\varphi_{\mathcal{L}}$ (up to a coordinate transformation and reparametrization of the interaction coefficients), we define for $u: \mathcal{L} \rightarrow \mathbb{R}$, $A \subset \mathbb{R}^3$ the energy

$$F_{\mathcal{L}}(u, A) := \frac{1}{2} \sum_{x \in \mathcal{L} \cap A} \sum_{y \in \mathcal{L}} c(x, y) |u(y) - u(x)|.$$

We are now in position to state [11, Proposition 2.6].

Proposition 2.1. *Let $c(x, y)$ be as in (14). Then*

$$\varphi_{\mathcal{L}}(\nu) = \frac{1}{|T_{\mathcal{L}}|} \inf \{ F_{\mathcal{L}}(u, T_{\mathcal{L}}) : u: \mathcal{L} \rightarrow \mathbb{R}, u(\cdot) - \langle \nu, \cdot \rangle \text{ is } T_{\mathcal{L}}\text{-periodic} \}. \quad (20)$$

With the definition of surface energy density at hand we can define the *coarse-grained continuum energy* $E_{\mathcal{L}}: \mathcal{M}_+(\mathbb{R}^3) \rightarrow [0, +\infty]$ as

$$E_{\mathcal{L}}(\mu) := \begin{cases} \int_{\partial^* V} \varphi_{\mathcal{L}}(\nu) d\mathcal{H}^2 & \text{if } \mu = \sqrt{2}\mathcal{L}^3 \llcorner V, \chi_V \in BV_{\text{loc}}(\mathbb{R}^3), \\ +\infty & \text{otherwise.} \end{cases} \quad (21)$$

with $\varphi_{\mathcal{L}}$ given by (19). Here, $\partial^* V$ denotes the reduced boundary of the set V , ν its outer normal and \mathcal{H}^2 , as noted at the beginning of this section, stands for the 2-dimensional Hausdorff measure in \mathbb{R}^3 (cf. [4], Chapters 2.8 and 3.5).

In what follows we say that F_ε Γ -converges to F if for all sequences $\{\varepsilon_j\}_j$ converging to 0 we have $\Gamma\text{-}\lim_j F_{\varepsilon_j} = F$.

The Wulff Crystal. In this section we calculate the Wulff crystals of the coarse grained FCC and HCP lattices. To the best of our knowledge, this is the first time that such a calculation has been carried out in a rigorous analytical way. In what follows we introduce the notion of Wulff shape in the general case of \mathbb{R}^n . While in the rest of this section we limit ourselves to the case $n = 3$, in Section 5 we consider general n .

Given $\varphi: \mathbb{R}^n \rightarrow [0, +\infty)$ convex, non-degenerate, (i.e. there exist $0 < c < C$ such that $c \leq \varphi(\nu) \leq C$ for all $\nu \in \mathbb{S}^{n-1}$) positively homogeneous of degree one, we define the Wulff set of φ by

$$W_\varphi := \{\zeta \in \mathbb{R}^n: \langle \zeta, \nu \rangle \leq \varphi(\nu) \text{ for all } \nu \in \mathbb{S}^{n-1}\}. \quad (22)$$

Thanks to the anisotropic isoperimetric inequality (cf. [19]), we have that W_φ is the unique (up to rigid motions) minimizer of

$$\min \left\{ \int_{\partial^* A} \varphi(\nu) \, d\mathcal{H}^{n-1}: |A| = |W_\varphi| \right\}.$$

Given $\lambda > 0$ we set $W_\lambda = \left(\frac{\lambda}{|W_\varphi|}\right)^{1/n} W_\varphi$ so that $|W_\lambda| = \lambda$ and, by scaling, it solves the minimum problem above among all sets $A \subset \mathbb{R}^n$ with $|A| = \lambda$. We recall here that, defining $\varphi^\circ: \mathbb{R}^n \rightarrow [0, +\infty)$ by

$$\varphi^\circ(\xi) = \sup_{\nu \in \mathbb{S}^{n-1}} \frac{\langle \nu, \xi \rangle}{\varphi(\nu)},$$

it holds that $W_\varphi = \{\varphi^\circ \leq 1\}$.

Definition 2.2. Let (X, τ) be a topological space and let $F_k: X \rightarrow [0, +\infty]$. For $x \in X$ we set

$$\Gamma\text{-lim sup}_{k \rightarrow +\infty} F_k(x) = \inf \left\{ \limsup_{k \rightarrow +\infty} F_k(x_k): x_k \xrightarrow{\tau} x \right\}$$

and

$$\Gamma\text{-lim inf}_{k \rightarrow +\infty} F_k(x) = \inf \left\{ \liminf_{k \rightarrow +\infty} F_k(x_k): x_k \xrightarrow{\tau} x \right\}.$$

If there exists $F: X \rightarrow [0, +\infty]$ such that

$$F(x) = \Gamma\text{-lim sup}_{k \rightarrow +\infty} F_k(x) = \Gamma\text{-lim inf}_{k \rightarrow +\infty} F_k(x),$$

we say that F_k Γ -converges with respect to τ to F and we write

$$F(x) = \Gamma\text{-lim}_{k \rightarrow +\infty} F_k(x).$$

If we have $(F_\varepsilon)_{\varepsilon > 0}: X \rightarrow [0, +\infty]$ we say that F_ε Γ -converges with respect to τ to F if F_{ε_k} Γ -converges with respect to τ to F for all $\varepsilon_k \rightarrow 0$.

The following variational coarse-graining result is proved in Section 4.

Theorem 2.3. *Let $\varepsilon \rightarrow 0$, and let $E_{\mathcal{L}, \varepsilon}$ and $E_{\mathcal{L}}$ be the energy functionals defined in (18) and (21), respectively.*

i) (Compactness) *Let $\{\mu_\varepsilon\}_\varepsilon \subset \mathcal{M}_+(\mathbb{R}^3)$ be such that*

$$\sup_{\varepsilon > 0} E_{\mathcal{L}, \varepsilon}(\mu_\varepsilon) < +\infty.$$

Then there exists $V \subset \mathbb{R}^3$ such that $\chi_V \in BV_{\text{loc}}(\mathbb{R}^3)$, $\mu = \sqrt{2}\mathcal{L}^3 \llcorner V$, and a subsequence (not relabeled) such that $\mu_\varepsilon \xrightarrow{} \mu$. Furthermore, if μ_ε is such that*

$$E_{\mathcal{L}, \varepsilon}(\mu_\varepsilon) = \inf_{\nu \in \mathcal{M}_+(\mathbb{R}^3): |\nu|(\mathbb{R}^3) = \varepsilon^3 n_\varepsilon} E_{\mathcal{L}, \varepsilon}(\nu),$$

with $\varepsilon^3 n_\varepsilon \rightarrow \sqrt{2}v$, then $\mu = \sqrt{2}\mathcal{L}^3 \llcorner W_{\varphi_{\mathcal{L}}}^v$, where $W_{\varphi_{\mathcal{L}}}^v = \lambda W_{\varphi_{\mathcal{L}}}$ (defined in (22)) for $\lambda > 0$ such that $|W_{\varphi_{\mathcal{L}}}^v| = v$.

ii) (*Limsup inequality*) Let $\mu \in \mathcal{M}_+(\mathbb{R}^3)$ be such that $\mu_\varepsilon \xrightarrow{*} \mu$. Then

$$E_{\mathcal{L}}(\mu) \leq \liminf_{\varepsilon \rightarrow 0} E_{\mathcal{L},\varepsilon}(\mu_\varepsilon).$$

iii) (*Limsup inequality*) Let $\mu \in \mathcal{M}_+(\mathbb{R}^3)$. Then there exists $\{\mu_\varepsilon\}_\varepsilon \subset \mathcal{M}_+(\mathbb{R}^3)$ such that $\mu_\varepsilon \xrightarrow{*} \mu$ and

$$E_{\mathcal{L}}(\mu) \geq \liminf_{\varepsilon \rightarrow 0} E_{\mathcal{L},\varepsilon}(\mu_\varepsilon).$$




FIGURE 3. The Wulff Crystal of the FCC-lattice on the left and HCP-lattice on the right.




FIGURE 4. The sublevel set $\{\varphi_{\text{FCC}} \leq 1\}$ on the left and the sublevel set $\{\varphi_{\text{HCP}} \leq 1\}$ on the right.

Explicit formula of the surface energy densities. Taking advantage of the representation formula (20) stated in Proposition (2.1), we provide the explicit formulas of the surface energy density $\varphi_{\mathcal{L}_{\text{FCC}}}$ and $\varphi_{\mathcal{L}_{\text{HCP}}}$. Their sublevel sets are depicted in Figure 4. With the two explicit formulas at hand we can calculate the polar functions of both densities, the associated Wulff shapes and the surface energy per unit volume of both the FCC and HCP crystals. In order not to overburden the reader with notation, we write φ_{FCC} and φ_{HCP} for $\varphi_{\mathcal{L}_{\text{FCC}}}$ and $\varphi_{\mathcal{L}_{\text{HCP}}}$ as well as W_{FCC} and W_{HCP} instead of $W_{\varphi_{\mathcal{L}_{\text{FCC}}}}$ and $W_{\varphi_{\mathcal{L}_{\text{HCP}}}}$. W_{FCC} and W_{HCP} are depicted in Figure 3.

Proposition 2.4. *The following formulas hold true.*

$$\varphi_{\text{FCC}}(\nu) = |\nu_1 + \nu_2| + |\nu_1 + \nu_3| + |\nu_2 + \nu_3| + |\nu_1 - \nu_2| + |\nu_1 - \nu_3| + |\nu_2 - \nu_3|, \quad (23)$$

and

$$\varphi_{\text{FCC}}^\circ(\zeta) = \max \left\{ \frac{1}{4} \|\zeta\|_\infty, \frac{1}{6} \|\zeta\|_1 \right\}. \quad (24)$$

In particular, W_{FCC} is a truncated octahedron and its surface energy per unit volume is

$$|W_{\text{FCC}}|^{-2/3} \int_{\partial^* W_{\text{FCC}}} \varphi_{\text{FCC}}(\nu) \, d\mathcal{H}^2 = 3 \cdot 2^2 \cdot 2^{2/3}. \quad (25)$$

Proposition 2.5. *The following formulas hold true.*

$$\begin{aligned} \varphi_{\text{HCP}}(\nu) &= \sqrt{2} (|\langle e_1, \nu \rangle| + |\langle e_2, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle|) + \frac{1}{\sqrt{2}} |\langle e_3, \nu \rangle| \\ &+ \sqrt{2} \max \{ |\langle e_1, \nu \rangle|, |\langle e_2, \nu \rangle|, |\langle e_3, \nu \rangle|, |\langle e_1 - e_2, \nu \rangle| \}, \end{aligned} \quad (26)$$

and

$$\begin{aligned} \varphi_{\text{HCP}}^\circ(\zeta) &= \max \left\{ \frac{2}{7\sqrt{2}} \left(|\zeta_1| + \frac{1}{\sqrt{3}} |\zeta_2| + \frac{3}{2\sqrt{6}} |\zeta_3| \right), \frac{1}{2\sqrt{3}} |\zeta_3|, \right. \\ &\left. \frac{2}{3\sqrt{6}} |\zeta_2|, \frac{4}{7\sqrt{6}} |\zeta_2| + \frac{3}{14\sqrt{3}} |\zeta_3|, \frac{1}{3\sqrt{2}} \left(|\zeta_1| + \frac{1}{\sqrt{3}} |\zeta_2| \right) \right\}. \end{aligned} \quad (27)$$

In particular, W_{HCP} is a truncated elongated hexagonal bipyramid and its surface energy per unit volume is

$$|W_{\text{HCP}}|^{-2/3} \int_{\partial^* W_{\text{HCP}}} \varphi_{\text{HCP}}(\nu) \, d\mathcal{H}^2 = 3 \cdot 2^{2/3} \cdot 65^{1/3}. \quad (28)$$

3. PROOF OF PROPOSITION 2.4 AND PROPOSITION 2.5

In this section we prove Proposition 2.4 and Proposition 2.5. To this end, we use Proposition 2.1 to note that $\varphi_{\mathcal{L}}$ is given by (20).

Proof of Proposition 2.4. We divide the proof into several steps. First, we calculate φ_{FCC} . Then, we calculate $\varphi_{\text{FCC}}^\circ$. Lastly, we calculate (25). Recall (6).

Step 1. (Calculation of φ_{FCC}) We make use of Proposition 2.1 in order to calculate φ_{FCC} . First of all, owing to (15), we note that

$$|T_{\text{FCC}}| = \frac{1}{3} \sqrt{6} \cdot \frac{1}{2} \sqrt{3} = \frac{1}{2} \sqrt{2}. \quad (29)$$

Given $u: \mathcal{L}_{\text{FCC}} \rightarrow \mathbb{R}$ such that $u(\cdot) - \langle \nu, \cdot \rangle$ is T_{FCC} -periodic we have that $u(x + b_i) = u(x) + \langle b_i, \nu \rangle$ for all $i = 1, 2, 3$. Therefore, u is an affine function of the form $u(x) = \langle x, \nu \rangle + c$, $x \in \mathcal{L}_{\text{FCC}}$ for some $c \in \mathbb{R}$. Lastly, note that $\mathcal{L}_{\text{FCC}} \cap T_{\text{FCC}} = \{0\}$. Using (20) and (29), we obtain

$$\varphi_{\text{FCC}}(\nu) = \frac{1}{2}\sqrt{2} \sum_{\xi \in \mathcal{N}_{\text{FCC}}} |u(\xi) - u(0)| = \frac{1}{2}\sqrt{2} \sum_{\xi \in \mathcal{N}_{\text{FCC}}} |\langle \xi, \nu \rangle|.$$

Employing now (10), we obtain (23).

Step 2.(Calculation of $\varphi_{\text{FCC}}^\circ$) Let G be the isometry group on \mathbb{R}^3 whose elements $g \in G$ are the linear isometries $g: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $g(\nu_1, \nu_2, \nu_3) = (b_1\nu_{\pi_1}, b_2\nu_{\pi_2}, b_3\nu_{\pi_3})$ where π is a permutation on $\{1, 2, 3\}$ and $b_i \in \{-1, 1\}$. Since $\varphi_{\text{FCC}}(g(\nu)) = \varphi_{\text{FCC}}(\nu)$ for all $g \in G$, $\nu \in \mathbb{R}^3$, we infer that

$$\varphi_{\text{FCC}}^\circ(\zeta) := \max_{\substack{\nu \in \mathbb{R}^3 \\ \varphi_{\text{FCC}}(\nu) \leq 1}} \langle \zeta, \nu \rangle = \max_{\substack{\nu \in \mathbb{R}^3 \\ \varphi_{\text{FCC}}(g^{-1}(\nu)) \leq 1}} \langle \zeta, g^{-1}(\nu) \rangle = \max_{\substack{\nu \in \mathbb{R}^3 \\ \varphi_{\text{FCC}}(\nu) \leq 1}} \langle g(\zeta), \nu \rangle = \varphi_{\text{FCC}}^\circ(g(\zeta)),$$

also relying on the property $g^T = g^{-1}$. Therefore, we can assume that $0 \leq \zeta_1 \leq \zeta_2 \leq \zeta_3$. Thus, if we want to maximize $\langle \zeta, \nu \rangle$ under the condition $\varphi_{\text{FCC}}(\nu) \leq 1$, we can as well assume that $0 \leq \nu_1 \leq \nu_2 \leq \nu_3$, so that condition $\varphi_{\text{FCC}}(\nu) \leq 1$ becomes equivalent to

$$4\nu_3 + 2\nu_2 \leq 1.$$

Therefore, noting that any linear function attains its maximum at the extreme points of a convex set and consulting Figure 5, we obtain

$$\begin{aligned} \max_{\substack{0 \leq \nu_1 \leq \nu_2 \leq \nu_3 \\ 4\nu_3 + 2\nu_2 \leq 1}} \zeta_1\nu_1 + \zeta_2\nu_2 + \zeta_3\nu_3 &= \max_{\substack{0 \leq \nu_2 \leq \nu_3 \\ 4\nu_3 + 2\nu_2 \leq 1}} (\zeta_1 + \zeta_2)\nu_2 + \zeta_3\nu_3 = \max \left\{ \frac{1}{4}\zeta_3, \frac{1}{6}(\zeta_1 + \zeta_2 + \zeta_3) \right\} \\ &= \max \left\{ \frac{1}{4}\|\zeta\|_\infty, \frac{1}{6}\|\zeta\|_1 \right\}. \end{aligned}$$

This is the desired formula (24) and concludes Step 2.




FIGURE 5. The set $\{0 \leq \nu_2 \leq \nu_3\} \cap \{4\nu_3 + 2\nu_2 \leq 1\}$ depicted in gray.

Step 3.(Calculation of (25)) Note that the set $W_{\varphi_{\text{FCC}}}$ is the intersection of a cube $\|\zeta\|_\infty \leq 4$ with an octahedron $\|\zeta\|_1 \leq 6$, see Fig. 3. Its boundary has 6 square faces, where $\nu = \pm(1, 0, 0)$ (resp. $\pm(0, 1, 0)$ or $\pm(0, 0, 1)$) and 8 hexagonal faces, where $\nu = \frac{1}{\sqrt{3}}(\pm 1 \pm 1 \pm 1)$. First, we consider the set where $\nu = (1, 0, 0)$, the other cases where $\varphi_{\text{FCC}}^\circ(\zeta) = \frac{1}{4}\|\zeta\|_\infty = 1$ contributing with the same value. The square is given by

$$S_1^+ = \{(4, \zeta_2, \zeta_3) : |\zeta_2| + |\zeta_3| \leq 2\} = \left\{ \frac{1}{4}\|\zeta\|_\infty = \frac{1}{4}\zeta_1 = 1 \right\} \cap \left\{ \frac{1}{6}\|\zeta\|_1 \leq 1 \right\}.$$

Therefore, $\mathcal{H}^2(S_1^+) = 8$ and $\varphi_{\text{FCC}}((1, 0, 0)) = 4$. Similarly, we obtain the same measure and value of φ_{FCC} for the other squares S_1^-, S_2^\pm, S_3^\pm , where ν is (up to sign) one of the coordinate unit vectors. Hence,

$$\sum_{i=1}^3 \int_{S_i^+} \varphi_{\text{FCC}}(\nu) d\mathcal{H}^2 + \sum_{i=1}^3 \int_{S_i^-} \varphi_{\text{FCC}}(\nu) d\mathcal{H}^2 = 6 \cdot 8 \cdot 4 = 3 \cdot 2^6. \quad (30)$$

Next, we consider the contribution of a hexagon. We consider the hexagon contained in the set $\zeta_i \geq 0$ for all i . Here, we have $\nu = \frac{1}{\sqrt{3}}(1, 1, 1)$ and $\varphi_{\text{FCC}}(\nu) = 2\sqrt{3}$. The 6 sides of the hexagon have all side-length $2\sqrt{2}$. To see this, there are sides of the form $(4, 2-t, t), t \in [0, 2]$ or $(4-t, 0, 2+t), t \in [0, 2]$ and their permutations (up to identifying t with $2-t$ in the first case and $4-t$ and $2+t$ in the second case). An equilateral hexagon H of side-length $2\sqrt{2}$ satisfies $\mathcal{H}^2(H) = 12\sqrt{3}$. Labeling the hexagons by $H_i, i = 0, \dots, 7$, we obtain

$$\sum_{i=0}^7 \int_{H_i} \varphi_{\text{FCC}}(\nu) d\mathcal{H}^2 = 8 \cdot \mathcal{H}^2(H_i) \cdot \varphi_{\text{FCC}}\left(\frac{1}{\sqrt{3}}(\pm 1, \pm 1, \pm 1)\right) = 8 \cdot 12\sqrt{3} \cdot 2\sqrt{3} = 3^2 \cdot 2^6. \quad (31)$$

Using (30) and (31), we obtain

$$\int_{\partial W_{\text{FCC}}} \varphi_{\text{FCC}}(\nu) d\mathcal{H}^2 = 3 \cdot 2^6 + 3^2 \cdot 2^6 = 3 \cdot 2^8. \quad (32)$$

Let $C := \{\zeta \in \mathbb{R}^3 : \zeta_i \geq 0 \text{ for all } i = 1, 2, 3 \text{ and } \frac{1}{4}\|\zeta\|_\infty \geq \frac{1}{6}\|\zeta\|_1\}$ and $C^c := \{\zeta \in \mathbb{R}^3 : \zeta_i \geq 0 \text{ for all } i = 1, 2, 3 \text{ and } \frac{1}{4}\|\zeta\|_\infty < \frac{1}{6}\|\zeta\|_1\}$. We split the calculation of the volume $W \cap \{\zeta \in \mathbb{R}^3 : \zeta_i \geq 0 \text{ for all } i\}$ into the set $C \cap W_{\text{FCC}}$ and $C^c \cap W_{\text{FCC}}$. Noting that on this set $|\nabla \varphi_{\text{FCC}}^\circ(\zeta)| = \frac{1}{4} \mathcal{L}^3$ -a.e. on C , due to the coarea-formula, we have

$$\begin{aligned} |\{C \cap W_{\text{FCC}}\}| &= 4 \int_{C \cap W_{\text{FCC}}} |\nabla \varphi_{\text{FCC}}^\circ(\zeta)| d\zeta = 4 \int_0^1 \mathcal{H}^2(C \cap \{\varphi_{\text{FCC}}^\circ(\zeta) = s\}) ds \\ &= \int_0^1 4 \cdot s^2 \cdot 6 ds = 8. \end{aligned}$$

Here we used that, $C \cap \{\varphi_{\text{FCC}}^\circ(\zeta) = s\} = s(S_1^+ \cup S_2^+ \cup S_3^+) \cap \{\zeta_i \geq 0\}$ and the scaling properties of the 2-dimensional Hausdorff-measure. On the other hand, using that $|\nabla \varphi_{\text{FCC}}^\circ(\zeta)| = \frac{\sqrt{3}}{6} \mathcal{L}^3$ -a.e. on C^c , we have

$$\begin{aligned} |\{C^c \cap W_{\text{FCC}}\}| &= 2\sqrt{3} \int_{C^c \cap W_{\text{FCC}}} |\nabla \varphi_{\text{FCC}}^\circ(\zeta)| d\zeta = 2\sqrt{3} \int_0^1 \mathcal{H}^2(C^c \cap \{\varphi_{\text{FCC}}^\circ(\zeta) = s\}) ds \\ &= 2\sqrt{3} \int_0^1 s^2 \cdot 12\sqrt{3} ds = 3 \cdot 2^3. \end{aligned}$$

Taking into account also the sets $\{\pm \zeta_i \geq 0\}$, we obtain

$$|W_{\text{FCC}}| = 8(8 + 3 \cdot 2^3) = 2^8.$$

Now, this together with (32) yields (25). \square

Proof of Proposition 2.5. We divide the proof into several steps. First, we calculate φ_{HCP} . Then, we calculate $\varphi_{\text{HCP}}^\circ$. Lastly, we calculate (28).

Step 1.(Calculation of φ_{HCP}) We make use of Proposition 2.1 in order to calculate φ_{HCP} . First of all, due to (16), note that

$$|T_{\text{HCP}}| = \frac{2}{3}\sqrt{6} \cdot \frac{1}{2}\sqrt{3} = \sqrt{2}. \quad (33)$$

Given $u: \mathcal{L}_{\text{FCC}} \rightarrow \mathbb{R}$ such that $u(\cdot) - \langle \nu, \cdot \rangle$ is T_{HCP} -periodic we have that $u(x + e_i) = u(x) + \langle e_i, \nu \rangle$ for all $i = 1, 2, 3$ and $\mathcal{L}_{\text{HCP}} \cap T_{\text{HCP}} = \{0, v_1\}$. Hence, there exist $c_1, c_2 \in \mathbb{R}$ such that

$$u(x) = \begin{cases} \langle x, \nu \rangle + c_1 & x \in \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\}; \\ \langle x_0, \nu \rangle + c_2 & x = x_0 + v_1, \text{ with } x_0 \in \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\}. \end{cases}$$

Setting $c_2 - c_1 = t$, recalling (11) and (12), we therefore obtain

$$F_{\mathcal{L}_{\text{HCP}}}(u, T_{\text{HCP}}) = 2(|\langle e_1, \nu \rangle| + |\langle e_2, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle|) + |t| + |t - \langle e_1, \nu \rangle| + |t - \langle e_2, \nu \rangle| \\ + |t - \langle e_3, \nu \rangle| + |t - \langle e_3 + e_1, \nu \rangle| + |t - \langle e_3 + e_2, \nu \rangle|.$$

Employing Proposition 2.1 and (33), we have

$$\varphi_{\text{HCP}}(\nu) = \sqrt{2}(|\langle e_1, \nu \rangle| + |\langle e_2, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle|) + \frac{1}{\sqrt{2}} \min_{t \in \mathbb{R}} g_{\nu}(t), \quad (34)$$

where

$$g_{\nu}(t) := |t| + |t - \langle e_1, \nu \rangle| + |t - \langle e_2, \nu \rangle| + |t - \langle e_3, \nu \rangle| + |t - \langle e_3 + e_1, \nu \rangle| + |t - \langle e_3 + e_2, \nu \rangle|.$$

Next, we show that

$$\min_{t \in \mathbb{R}} g_{\nu}(t) = |\langle e_3, \nu \rangle| + 2 \max\{|\langle e_1, \nu \rangle|, |\langle e_2, \nu \rangle|, |\langle e_1 - e_2, \nu \rangle|, |\langle e_3, \nu \rangle|\}. \quad (35)$$

Note that if (35) is shown, (26) is proven and Step 1 is concluded. In order to prove (35), we first note that $g_{\nu}(t)$ is a piecewise affine function such that $g_{\nu}(t) \rightarrow +\infty$ as $|t| \rightarrow +\infty$. Hence, it attains its minimum at a point of non-differentiability. The function g_{ν} is not differentiable for $t \in \{0, \langle e_1, \nu \rangle, \langle e_2, \nu \rangle, \langle e_3, \nu \rangle, \langle e_3 + e_1, \nu \rangle, \langle e_3 + e_2, \nu \rangle\}$ and therefore

$$\min_{t \in \mathbb{R}} g_{\nu}(t) = |\langle e_3, \nu \rangle| + \min\{f_k(\nu) : k \in \{0, \dots, 5\}\},$$

where

$$\begin{aligned} f_0(\nu) &= |\langle e_1, \nu \rangle| + |\langle e_2, \nu \rangle| + |\langle e_3 + e_1, \nu \rangle| + |\langle e_3 + e_2, \nu \rangle|, \\ f_1(\nu) &= |\langle e_1, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle| + |\langle e_3 - e_1, \nu \rangle| + |\langle e_3 + e_2 - e_1, \nu \rangle|, \\ f_2(\nu) &= |\langle e_2, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle| + |\langle e_3 - e_2, \nu \rangle| + |\langle e_3 + e_1 - e_2, \nu \rangle|, \\ f_3(\nu) &= |\langle e_1, \nu \rangle| + |\langle e_2, \nu \rangle| + |\langle e_3 - e_1, \nu \rangle| + |\langle e_3 - e_2, \nu \rangle|, \\ f_4(\nu) &= |\langle e_1, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle| + |\langle e_3 + e_1, \nu \rangle| + |\langle e_3 + e_1 - e_2, \nu \rangle|, \\ f_5(\nu) &= |\langle e_2, \nu \rangle| + |\langle e_1 - e_2, \nu \rangle| + |\langle e_3 + e_2, \nu \rangle| + |\langle e_3 + e_2 - e_1, \nu \rangle|. \end{aligned}$$

It is easy to see that

$$\min_{t \in \mathbb{R}} g_{\nu}(t) = |\langle e_3, \nu \rangle| + \min\{|\langle e_1, R_k \nu \rangle| + |\langle e_2, R_k \nu \rangle| + |\langle e_3 + e_1, R_k \nu \rangle| \\ + |\langle e_3 + e_2, R_k \nu \rangle| : k \in \{0, \dots, 5\}\},$$

where R_k is the rotation of angle $k\pi/3$ around the x_3 -axis. Noting also that $\min_{t \in \mathbb{R}} g_{\nu}(t) = \min_{t \in \mathbb{R}} g_{-\nu}(t)$, it is not restrictive to assume that $\langle e_1, \nu \rangle \geq 0, \langle e_2, \nu \rangle \geq 0, \langle e_3, \nu \rangle \geq 0$. We only

consider the case, where $\langle e_1, \nu \rangle \geq \langle e_2, \nu \rangle \geq \langle e_3, \nu \rangle \geq 0$, the other being dealt with in a similar fashion. In this case we have $f_0(\nu) \geq f_3(\nu)$, $f_4(\nu) \geq f_5(\nu)$, and

$$\begin{aligned} f_1(\nu) &= \langle e_1, \nu \rangle + \langle e_1 - e_2, \nu \rangle + \langle e_1 - e_3, \nu \rangle + |\langle e_3 + e_2 - e_1, \nu \rangle| \\ &= 2\langle e_1, \nu \rangle + \langle e_1 - e_2 - e_3, \nu \rangle + |\langle e_3 + e_2 - e_1, \nu \rangle| \geq 2\langle e_1, \nu \rangle; \\ f_2(\nu) &= \langle e_2, \nu \rangle + \langle e_1 - e_2, \nu \rangle + \langle e_2 - e_3, \nu \rangle + \langle e_3 + e_1 - e_2, \nu \rangle = 2\langle e_1, \nu \rangle; \\ f_3(\nu) &= \langle e_1, \nu \rangle + \langle e_2, \nu \rangle + \langle e_1 - e_3, \nu \rangle + \langle e_2 - e_3, \nu \rangle = 2\langle e_1, \nu \rangle + 2\langle e_2, \nu \rangle - 2\langle e_3, \nu \rangle \geq 2\langle e_1, \nu \rangle; \\ f_4(\nu) &= \langle e_1, \nu \rangle + \langle e_2, \nu \rangle + \langle e_1 - e_3, \nu \rangle + |\langle e_3 + e_2 - e_1, \nu \rangle| \\ &= 2\langle e_1, \nu \rangle + \langle e_3 + e_2 - e_1, \nu \rangle + |\langle e_3 + e_2 - e_1, \nu \rangle| \geq 2\langle e_1, \nu \rangle. \end{aligned}$$

Hence, we see that (35) holds true. This together with (34) establishes (26) and concludes Step 1.

Step 2.(Calculation of $\varphi_{\text{HCP}}^\circ$) In order to calculate $\varphi_{\text{HCP}}^\circ$, we exploit the symmetries of $\varphi_{\text{HCP}}^\circ$. Note that

$$\varphi_{\text{HCP}} \left(- \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} \right) = \varphi_{\text{HCP}} \left(\begin{pmatrix} \nu_1 \\ \nu_2 \\ -\nu_3 \end{pmatrix} \right) = \varphi_{\text{HCP}} \left(\begin{pmatrix} -\nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} \right) = \varphi_{\text{HCP}} \left(\begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} \right). \quad (36)$$

Given $\zeta \in \mathbb{R}^3$ we can find $R = T_1^{\alpha_1} \circ T_2^{\alpha_2} \circ T_3^{\alpha_3}$, $\alpha_i \in \{0, 1\}$ such that $(R\zeta)_i \geq 0$ for all i . Thus,

$$\begin{aligned} \varphi_{\text{HCP}}^\circ(\zeta) &= \max_{\varphi_{\text{HCP}}(\nu) \leq 1} \langle \nu, \zeta \rangle = \max_{\varphi_{\text{HCP}}(\nu) \leq 1} \langle R\nu, R\zeta \rangle \\ &= \max_{\varphi_{\text{HCP}}(R^{-1}\nu) \leq 1} \langle \nu, R\zeta \rangle = \max_{\varphi_{\text{HCP}}(\nu) \leq 1} \langle \nu, R\zeta \rangle = \varphi_{\text{HCP}}^\circ(R\zeta). \end{aligned} \quad (37)$$

It therefore suffices to calculate $\varphi_{\text{HCP}}^\circ$ for $\zeta \in \mathbb{R}^3$ such that $\zeta_i \geq 0$. This together with (36) implies that if $\nu = (\nu_1, \nu_2, \nu_3)$ is such that $\varphi_{\text{HCP}}(\nu) \leq 1$ and

$$\langle \nu, \zeta \rangle = \max_{\varphi_{\text{HCP}}(\nu) \leq 1} \langle \nu, \zeta \rangle,$$

then $\nu_i \geq 0$ for all i . Additionally, a maximizer ν can be chosen such that φ_{HCP} is not differentiable at ν . Therefore, there are the following cases to consider:

- (a) $\langle e_1 - e_2, \nu \rangle = 0$;
- (b) $\langle e_1 - e_3, \nu \rangle = 0, \langle e_1 - e_2, \nu \rangle \geq 0$;
- (c) $\langle e_2 - e_3, \nu \rangle = 0, \langle e_3 - e_1, \nu \rangle \geq 0$;
- (d) $\langle e_3, \nu \rangle = 0$;
- (e) $\langle e_1, \nu \rangle = 0$.

Here, we point out that the points on the boundary $\nu_2 = 0$ are excluded as possible maximum points by arguing in the following way: If there were a point ν such that $\nu_2 = 0$, then $\varphi_{\text{HCP}}(\nu)$ would either be differentiable and thus ν would not be a maximum point or ν would satisfy one of the cases (a)-(e).

Maximum of case (a). Since $\langle e_1 - e_2, \nu \rangle = 0$, we have $\nu_1 = \sqrt{3}\nu_2$. Hence, $\nu = (t, \frac{1}{\sqrt{3}}t, s)$ for some $t, s \geq 0$. Now, using (26), we have

$$\varphi_{\text{HCP}}(\nu) = \sqrt{2} \left(2\nu_1 + \frac{1}{3}\sqrt{6}\nu_3 + \max\{\nu_1, \frac{2}{3}\sqrt{6}\nu_3\} \right).$$

(a.1) $t \geq \frac{2}{3}\sqrt{6}s$: Since the maximum is attained for $\varphi_{\text{HCP}}(\nu) = 1$, we have $t = \frac{1}{3\sqrt{2}} - \frac{1}{9}\sqrt{6}s$. Now, $t \geq 0$ together with $t \geq \frac{2}{3}\sqrt{6}s$ implies $0 \leq s \leq \frac{3}{14\sqrt{3}}$. Noting that

$$\langle \nu, \zeta \rangle = t \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right) + s\zeta_3 = \left(\frac{1}{3\sqrt{2}} - \frac{1}{9}\sqrt{6}s \right) \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right) + s\zeta_3,$$

we obtain

$$\max_{\nu \text{ sat. (a.1)}} \langle \nu, \zeta \rangle = \max \left\{ \frac{2}{7\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 + \frac{3}{2\sqrt{6}}\zeta_3 \right), \frac{1}{3\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right) \right\}. \quad (38)$$

(a.2) $t \leq \frac{2}{3}\sqrt{6}s$: Using $\varphi_{\text{HCP}}(\nu) = 1$, we obtain $t = \frac{1}{2\sqrt{2}} - \frac{\sqrt{6}}{2}s$. Now, $t \geq 0$ together with $t \leq \frac{2}{3}\sqrt{6}s$ implies $\frac{3}{14\sqrt{3}} \leq s \leq \frac{1}{2\sqrt{3}}$. Noting that

$$\langle \nu, \zeta \rangle = t \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right) + s\zeta_3 = \left(\frac{1}{3\sqrt{2}} - \frac{1}{9}\sqrt{6}s \right) \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right) + s\zeta_3,$$

we obtain

$$\max_{\nu \text{ sat. (a.2)}} \langle \nu, \zeta \rangle = \max \left\{ \frac{2}{7\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 + \frac{3}{2\sqrt{6}}\zeta_3 \right), \frac{1}{2\sqrt{3}}\zeta_3 \right\}. \quad (39)$$

Maximum of case (b). Since $\langle e_1 - e_3, \nu \rangle = 0$, we have $\nu_1 = \frac{2}{3}\sqrt{6}\nu_3$. Hence, $\nu = (t, s, \frac{3}{2\sqrt{6}}t)$ for some $t, s \geq 0$. Now using (26), we have

$$\varphi_{\text{HCP}}(\nu) = \frac{7}{2}\sqrt{2}\nu_1.$$

Hence, since the maximum is attained for $\varphi_{\text{HCP}}(\nu) = 1$, we have $\nu_1 = \frac{2}{7\sqrt{2}}$. Additionally, since $\langle e_1 - e_2, \nu \rangle \geq 0$, we have $\nu_2 \leq \frac{2}{7\sqrt{6}}$, and due to the form of ν , we have $\nu_3 = \frac{3}{14\sqrt{3}}$. This implies

$$\max_{\nu \text{ sat. (b)}} \langle \nu, \zeta \rangle = \frac{2}{7\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 + \frac{3}{2\sqrt{6}}\zeta_3 \right). \quad (40)$$

Maximum of case (c). Since $\langle e_2 - e_3, \nu \rangle = 0$, we have $\frac{1}{2}\nu_1 + \frac{1}{2}\sqrt{3}\nu_2 = \frac{2}{3}\sqrt{6}\nu_3$. Now using (26), we have

$$\varphi_{\text{HCP}}(\nu) = \frac{7}{2}\sqrt{2}\langle e_3, \nu \rangle = \frac{14}{3}\sqrt{3}\nu_3.$$

Hence, since the maximum is attained for $\varphi_{\text{HCP}}(\nu) = 1$, we have $\nu_3 = \frac{3}{14\sqrt{3}}$. Additionally, since $\langle e_3 - e_1, \nu \rangle \geq 0$, we have $\nu_1 \leq \frac{2}{7\sqrt{2}}$. Due to the form of $\langle e_2 - e_3, \nu \rangle = 0$, we have $\nu_2 = \frac{4}{7\sqrt{6}} - \frac{1}{\sqrt{3}}\nu_1$. Note that $\nu_2 \geq 0$ for all $0 \leq \nu_1 \leq \frac{2}{7\sqrt{2}}$. Therefore

$$\langle \nu, \zeta \rangle = \nu_1\zeta_1 + \left(\frac{4}{7\sqrt{6}} - \frac{1}{\sqrt{3}}\nu_1 \right) \zeta_2 + \frac{3}{14\sqrt{3}}\zeta_3.$$

This implies

$$\max_{\nu \text{ sat. (c)}} \langle \nu, \zeta \rangle = \max \left\{ \frac{4}{7\sqrt{6}}\zeta_2 + \frac{3}{14\sqrt{3}}\zeta_3, \frac{2}{7\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 + \frac{3}{2\sqrt{6}}\zeta_3 \right) \right\}. \quad (41)$$

Maximum of case (d). We have $\nu_3 = 0$ and therefore

$$\varphi_{\text{HCP}}(\nu) = \sqrt{2}(\langle e_1, \nu \rangle + \langle e_2, \nu \rangle + |\langle e_1 - e_2, \nu \rangle| + \max\{\langle e_1, \nu \rangle, \langle e_2, \nu \rangle\})$$

We distinguish to cases

- (d.1) $\langle e_1 - e_2, \nu \rangle \geq 0$;
- (d.2) $\langle e_1 - e_2, \nu \rangle \leq 0$.

Maximum of case (d.1). In the case $\langle e_1 - e_2, \nu \rangle \geq 0$ we have $\varphi_{\text{HCP}}(\nu) = 3\sqrt{2}\nu_1$ and therefore, since $\varphi_{\text{HCP}}(\nu) = 1$, $\nu_1 = \frac{1}{3\sqrt{2}}$. The inequality $\langle e_1 - e_2, \nu \rangle \geq 0$ implies that $0 \leq \nu_2 \leq \frac{1}{\sqrt{3}}\nu_1 = \frac{1}{3\sqrt{6}}$. Hence,

$$\max_{\nu \text{ sat. (d.1)}} \langle \nu, \zeta \rangle = \frac{1}{3\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right). \quad (42)$$

Maximum of case (d.2). In the case $\langle e_1 - e_2, \nu \rangle \leq 0$ we have

$$\varphi_{\text{HCP}}(\nu) = 3\sqrt{2}\langle e_2, \nu \rangle = \sqrt{2} \left(\frac{3}{2}\nu_1 + \frac{3}{2}\sqrt{3}\nu_2 \right).$$

This, together with $\varphi_{\text{HCP}}(\nu) = 1$, implies, $\nu_1 = \frac{2}{3\sqrt{2}} - \sqrt{3}\nu_2$ and therefore $\nu_2 \leq \frac{2}{3\sqrt{6}}$. Additionally, since $\langle e_1 - e_2 \rangle \leq 0$, we have $\frac{1}{3\sqrt{6}} \leq \nu_2$. Therefore,

$$\langle \nu, \zeta \rangle = \nu_1\zeta_1 + \nu_2\zeta_2 = \left(\frac{3}{2\sqrt{2}} - \sqrt{3}\nu_2 \right) \zeta_1 + \nu_2\zeta_2.$$

This implies

$$\max_{\nu \text{ sat. (d.2)}} \langle \nu, \zeta \rangle = \max \left\{ \frac{2}{3\sqrt{6}}\zeta_2, \frac{1}{3\sqrt{2}} \left(\zeta_1 + \frac{1}{\sqrt{3}}\zeta_2 \right) \right\}. \quad (43)$$

Maximum of case (e). In the case $\nu_1 = 0$ we have

$$\varphi_{\text{HCP}}(\nu) = \sqrt{2} \left(\sqrt{3}\nu_2 + \frac{1}{3}\sqrt{6}\nu_3 + \max \left\{ \frac{\sqrt{3}}{2}\nu_2, \frac{2}{3}\sqrt{6}\nu_3 \right\} \right).$$

We distinguish between two cases:

- (e.1) $\langle e_2, \nu \rangle \geq \langle e_3, \nu \rangle$;
- (e.2) $\langle e_2, \nu \rangle \leq \langle e_3, \nu \rangle$.

Maximum of case (e.1). In this case, we have

$$\varphi_{\text{HCP}}(\nu) = \sqrt{2} \left(\frac{3}{2}\sqrt{3}\nu_2 + \frac{1}{3}\sqrt{6}\nu_3 \right).$$

Therefore, since $\varphi_{\text{HCP}}(\nu) = 1$, we have $\nu_2 = \frac{2}{3\sqrt{6}} - \frac{2}{9}\sqrt{2}\nu_3$. Hence, $\nu_3 \leq \frac{3}{2\sqrt{3}}$. Additionally, since $\langle e_2 - e_3, \nu \rangle \geq 0$, we have $\nu_3 \leq \frac{3}{14\sqrt{3}}$. Therefore,

$$\langle \nu, \zeta \rangle = \nu_2\zeta_2 + \nu_3\zeta_3 = \left(\frac{2}{3\sqrt{6}} - \frac{2}{9}\sqrt{2}\nu_3 \right) \zeta_2 + \nu_3\zeta_3.$$

Hence,

$$\max_{\nu \text{ sat. (e.1)}} \langle \nu, \zeta \rangle = \max \left\{ \frac{2}{3\sqrt{6}}\zeta_2, \frac{4}{7\sqrt{6}}\zeta_2 + \frac{3}{14\sqrt{3}}\zeta_3 \right\}. \quad (44)$$

Maximum of case (e.2). In this case, we have

$$\varphi_{\text{HCP}}(\nu) = \sqrt{2} \left(\sqrt{3}\nu_2 + \sqrt{6}\nu_3 \right).$$

Therefore, since $\varphi_{\text{HCP}}(\nu) = 1$, we have $\nu_2 = \frac{1}{\sqrt{6}} - \sqrt{2}\nu_3$. Hence, $\nu_3 \leq \frac{1}{2\sqrt{3}}$. Additionally, since $\langle e_2 - e_3, \nu \rangle \leq 0$, we have $\nu_3 \geq \frac{3}{14\sqrt{3}}$. Therefore,

$$\langle \nu, \zeta \rangle = \nu_2\zeta_2 + \nu_3\zeta_3 = \left(\frac{1}{3\sqrt{6}} - \sqrt{2}\nu_3 \right) \zeta_2 + \nu_3\zeta_3.$$

Hence,

$$\max_{\nu \text{ sat. (e.1)}} \langle \nu, \zeta \rangle = \max \left\{ \frac{1}{2\sqrt{3}}\zeta_3, \frac{4}{7\sqrt{6}}\zeta_2 + \frac{3}{14\sqrt{3}}\zeta_3 \right\}. \quad (45)$$

Exploiting (38)–(45), and (37), we obtain (27). This concludes Step 2.

Step 3. (Calculation of (28)) In order to calculate (28), we split the calculation of $\partial^* W_{\text{HCP}} = \{\varphi_{\text{HCP}}^\circ(\zeta) = 1\}$ into different sets, where the maximum of $\varphi_{\text{HCP}}^\circ$ is attained. We consider the following cases

- (a) $A_a := \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{1}{2\sqrt{3}}|\zeta_3| = 1\}$;
- (b) $A_b := \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{2}{3\sqrt{6}}|\zeta_2| = 1\}$;
- (c) $A_c := \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{4}{7\sqrt{6}}|\zeta_2| + \frac{3}{14\sqrt{3}}|\zeta_3| = 1\}$;
- (d) $A_d := \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{1}{3\sqrt{2}}(|\zeta_1| + \frac{1}{\sqrt{3}}|\zeta_2|) = 1\}$;
- (e) $A_e := \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{2}{7\sqrt{2}}(|\zeta_1| + \frac{1}{\sqrt{3}}|\zeta_2| + \frac{3}{2\sqrt{6}}|\zeta_3|) = 1\}$.

In each of the cases, one can determine the area, shape and normal of the set, by invoking the condition that the maximum is attained $\varphi_{\text{HCP}}^\circ$ for the respective function and therefore all the other functions f in the definition of $\varphi_{\text{HCP}}^\circ$ satisfy $f \leq 1$. In the following, we only collect the results, since the calculations are elementary (but very long).

Calculations for case (a). In this case, we see that $\nu = (0, 0, \pm 1)$ \mathcal{H}^2 -a.e., since this set is contained in the level set of the function $|\zeta_3| = c$ for some $c > 0$. Additionally, we see that the set is a union of two hexagons of side length $2\sqrt{2}$. Therefore, for each of the two hexagons H_i we have $\mathcal{H}^2(H_i) = 12\sqrt{3}$. Furthermore, $\varphi_{\text{HCP}}(\nu) = 2\sqrt{3}$. Hence

$$\int_{A_a} \varphi_{\text{HCP}}(\nu) d\mathcal{H}^2 = 2 \cdot 12\sqrt{3} \cdot 2\sqrt{3} = 2^4 \cdot 3^2. \quad (46)$$

Calculations for case (b). In this case, we see that $\nu = (0, \pm 1, 0)$ \mathcal{H}^2 -a.e., since this set is contained in the level set of the function $|\zeta_2| = c$ for some $c > 0$. Additionally, we see that the set is a union of two rectangles with side lengths $3\sqrt{2}$ and $\frac{4}{3}\sqrt{3}$. Therefore, for each of the two rectangles S_i we have $\mathcal{H}^2(S_i) = 4\sqrt{6}$. Furthermore, $\varphi_{\text{HCP}}(\nu) = \frac{3}{2}\sqrt{6}$. Hence

$$\int_{A_b} \varphi_{\text{HCP}}(\nu) d\mathcal{H}^2 = 2 \cdot 4\sqrt{6} \cdot \frac{3}{2}\sqrt{6} = 2^3 \cdot 3^2. \quad (47)$$

Calculations for case (c). In this case, we see that $\nu = (3/41)^{1/2}(0, \pm 8/\sqrt{6}, \pm \sqrt{3})$ \mathcal{H}^2 -a.e., since this set is contained in the level set of the function $\frac{4}{7\sqrt{6}}|\zeta_2| + \frac{3}{14\sqrt{3}}|\zeta_3| = c$ for some $c > 0$. Additionally, we see that the set is a union of four trapezoids with height $(41/6)^{1/2}$ and two parallel sides of lengths $3\sqrt{2}$ and $2\sqrt{2}$. Therefore, for each of the four trapezoids T_i we have $\mathcal{H}^2(T_i) = \frac{5}{2}(\frac{41}{3})^{1/2}$. Furthermore, $\varphi_{\text{HCP}}(\nu) = 14(\frac{3}{41})^{1/2}$. Hence

$$\int_{A_c} \varphi_{\text{HCP}}(\nu) d\mathcal{H}^2 = 4 \cdot \frac{5}{2} \left(\frac{41}{3}\right)^{1/2} \cdot 14 \left(\frac{3}{41}\right)^{1/2} = 2^2 \cdot 5 \cdot 7. \quad (48)$$

Calculations for case (d). In this case, we see that $\nu = \frac{1}{2}(\pm\sqrt{3}, \pm 1, 0)$ \mathcal{H}^2 -a.e., since this set is contained in the level set of the function $|\zeta_1| + \frac{1}{\sqrt{3}}|\zeta_2| = c$ for some $c > 0$. Additionally, we see that the set is a union of four rectangles with side length $3\sqrt{2}$ and $\frac{4}{3}\sqrt{3}$. Therefore, for each of

the four rectangles R_i we have $\mathcal{H}^2(R_i) = 4\sqrt{6}$. Furthermore, $\varphi_{\text{HCP}}(\nu) = \frac{3}{2}\sqrt{6}$. Hence

$$\int_{A_d} \varphi_{\text{HCP}}(\nu) \, d\mathcal{H}^2 = 4 \cdot 4\sqrt{6} \cdot \frac{3}{2}\sqrt{6} = 2^4 \cdot 3^2. \quad (49)$$

Calculations for case (e). In this case, we see that $\nu = 2(6/41)^{1/2}(\pm 1, \pm \frac{1}{\sqrt{3}}, \pm \frac{3}{2\sqrt{6}})$ \mathcal{H}^2 -a.e., since this set is contained in the level set of the function $|\zeta_1| + \frac{1}{\sqrt{3}}|\zeta_2| + \frac{3}{2\sqrt{6}}|\zeta_3| = c$ for some $c > 0$. Additionally, we see that the set is a union of eight trapezoids with height $(41/6)^{1/2}$ and two parallel sides of lengths $3\sqrt{2}$ and $2\sqrt{2}$. Therefore, for each of the eight trapezoids Z_i we have $\mathcal{H}^2(Z_i) = \frac{5}{2}(\frac{41}{3})^{1/2}$. Furthermore, $\varphi_{\text{HCP}}(\nu) = 14(\frac{3}{41})^{1/2}$. Hence

$$\int_{A_e} \varphi_{\text{HCP}}(\nu) \, d\mathcal{H}^2 = 8 \cdot \frac{5}{2} \left(\frac{41}{3}\right)^{1/2} \cdot 14 \left(\frac{3}{41}\right)^{1/2} = 2^3 \cdot 5 \cdot 7. \quad (50)$$

Taking into account (46)–(50), we obtain

$$\int_{\partial^* W_{\text{HCP}}} \varphi_{\text{HCP}}(\nu) \, d\mathcal{H}^2 = 2^5 \cdot 3^2 + 2^3 \cdot 3^2 + 2^2 \cdot 5 \cdot 7 + 2^4 \cdot 3^2 + 2^3 \cdot 5 \cdot 7 = 780. \quad (51)$$

Next, we need to calculate $|W_{\text{HCP}}|$, since $W_{\text{HCP}} = \{\varphi_{\text{HCP}}^\circ \leq 1\} \cap (C_a \cup C_b \cup C_c \cup C_d \cup C_e)$, where

$$\begin{aligned} C_a &= \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{1}{2\sqrt{3}}|\zeta_3|\}, \\ C_b &:= \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{2}{3\sqrt{6}}|\zeta_2|\}, \\ C_c &:= \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{4}{7\sqrt{6}}|\zeta_2| + \frac{3}{14\sqrt{3}}|\zeta_3|\}, \\ C_d &:= \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{1}{3\sqrt{2}}(|\zeta_1| + \frac{1}{\sqrt{3}}|\zeta_2|)\}, \\ C_e &:= \{\zeta \in \mathbb{R}^3 : \varphi_{\text{HCP}}^\circ(\zeta) = \frac{2}{7\sqrt{2}}(|\zeta_1| + \frac{1}{\sqrt{3}}|\zeta_2| + \frac{3}{2\sqrt{6}}|\zeta_3|)\}. \end{aligned}$$

Note that $\mathcal{H}^2(C_\alpha \cap \{\varphi_{\text{HCP}}^\circ(\zeta) = s\}) = s^2 \mathcal{H}^2(A_\alpha)$ for all $\alpha \in \{a, b, c, d, e\}$. In the set C_a we have that $|\nabla \varphi_{\text{HCP}}^\circ(\zeta)| = \frac{1}{2\sqrt{3}} \mathcal{L}^3$ -a.e.. Due to the coarea formula, we have

$$\begin{aligned} |C_a \cap W_{\text{HCP}}| &= 2\sqrt{3} \int_{C_a \cap W_{\text{HCP}}} |\nabla \varphi_{\text{HCP}}^\circ(\zeta)| \, d\zeta \\ &= 2\sqrt{3} \int_0^1 \mathcal{H}^2(C_a \cap \{\varphi_{\text{HCP}}^\circ(\zeta) = s\}) \, ds = \frac{2}{3}\sqrt{3} \mathcal{H}^2(A_a) = 2^4 \cdot 3. \end{aligned} \quad (52)$$

In the set C_b , we have that $|\nabla \varphi_{\text{HCP}}^\circ(\zeta)| = \frac{2}{3\sqrt{6}} \mathcal{L}^3$ -a.e.. Due to the coarea formula, we have

$$\begin{aligned} |C_b \cap W_{\text{HCP}}| &= \frac{3}{2}\sqrt{6} \int_{C_b \cap W_{\text{HCP}}} |\nabla \varphi_{\text{HCP}}^\circ(\zeta)| \, d\zeta \\ &= \frac{3}{2}\sqrt{6} \int_0^1 \mathcal{H}^2(C_b \cap \{\varphi_{\text{HCP}}^\circ(\zeta) = s\}) \, ds = \frac{1}{2}\sqrt{6} \mathcal{H}^2(A_b) = 2^3 \cdot 3. \end{aligned} \quad (53)$$

In the set C_c , we have that $|\nabla\varphi_{\text{HCP}}^\circ(\zeta)| = \frac{1}{14}(41/3)^{1/2} \mathcal{L}^3$ -a.e.. Due to the coarea formula, we have

$$\begin{aligned} |C_c \cap W_{\text{HCP}}| &= 14 \left(\frac{3}{41}\right)^{1/2} \int_{C_c \cap W_{\text{HCP}}} |\nabla\varphi_{\text{HCP}}^\circ(\zeta)| \, d\zeta \\ &= 14 \left(\frac{3}{41}\right)^{1/2} \int_0^1 \mathcal{H}^2(C_c \cap \{\varphi_{\text{HCP}}^\circ(\zeta) = s\}) \, ds \\ &= \frac{1}{3} 14 \left(\frac{3}{41}\right)^{1/2} \mathcal{H}^2(A_c) = \frac{2^2 \cdot 5 \cdot 7}{3}. \end{aligned} \quad (54)$$

In the set C_d , we have that $|\nabla\varphi_{\text{HCP}}^\circ(\zeta)| = \frac{2}{3\sqrt{6}} \mathcal{L}^3$ -a.e.. Due to the coarea formula, we have

$$\begin{aligned} |C_d \cap W_{\text{HCP}}| &= \frac{3}{2} \sqrt{6} \int_{C_d \cap W_{\text{HCP}}} |\nabla\varphi_{\text{HCP}}^\circ(\zeta)| \, d\zeta \\ &= \frac{3}{2} \sqrt{6} \int_0^1 \mathcal{H}^2(C_d \cap \{\varphi_{\text{HCP}}^\circ(\zeta) = s\}) \, ds \\ &= \frac{1}{2} \sqrt{6} \mathcal{H}^2(A_d) = 2^4 \cdot 3. \end{aligned} \quad (55)$$

In the set C_e , we have that $|\nabla\varphi_{\text{HCP}}^\circ(\zeta)| = \frac{1}{14}(41/3)^{1/2} \mathcal{L}^3$ -a.e.. Due to the coarea formula, we have

$$\begin{aligned} |C_e \cap W_{\text{HCP}}| &= 14 \left(\frac{3}{41}\right)^{1/2} \int_{C_e \cap W_{\text{HCP}}} |\nabla\varphi_{\text{HCP}}^\circ(\zeta)| \, d\zeta \\ &= 14 \left(\frac{3}{41}\right)^{1/2} \int_0^1 \mathcal{H}^2(C_e \cap \{\varphi_{\text{HCP}}^\circ(\zeta) = s\}) \, ds \\ &= \frac{1}{3} 14 \left(\frac{3}{41}\right)^{1/2} \mathcal{H}^2(A_e) = \frac{2^3 \cdot 5 \cdot 7}{3}. \end{aligned} \quad (56)$$

Using (52)–(56), we obtain $|W_{\text{HCP}}| = 260$. This together with (51) yields (28). \square

4. Γ -CONVERGENCE ANALYSIS ON THE FCC AND HCP LATTICES

In this section we prove Theorem 2.3. In order to prove the compactness statement, we provide some preliminary lemmata about the shape of the Voronoi cells of the FCC-lattice as well as the HCP-lattice (see Figure 6). In what follows we use the notation $\mathcal{N}_{\text{FCC}} = \mathcal{N}_{\mathcal{L}_{\text{FCC}}}(0)$ and $\mathcal{N}_{\text{HCP}} = \mathcal{N}_{\mathcal{L}_{\text{HCP}}}(0)$.

Lemma 4.1. (*Voronoi cell in the FCC-lattice*) *Let us take $x \in \mathcal{L}_{\text{FCC}}$. Then*

$$\mathcal{V}_{\mathcal{L}_{\text{FCC}}}(x) = x + V_{\text{FCC}}, \quad \text{where } V_{\text{FCC}} := \left\{ y \in \mathbb{R}^3 : \max_{b \in \mathcal{N}_{\text{FCC}}} \langle b, y \rangle \leq \frac{1}{2} \right\}. \quad (57)$$

Given $b_0 \in \mathcal{N}_{\text{FCC}}$ the face

$$S_{b_0} := \left\{ y \in \mathbb{R}^3 : \max_{b \in \mathcal{N}_{\text{FCC}}} \langle b, y \rangle = \langle b_0, y \rangle = \frac{1}{2} \right\} \quad (58)$$

is a rhombus with $\mathcal{H}^2(S_{b_0}) = \frac{1}{4}\sqrt{2}$. Moreover, for each $b_0 \in \mathcal{N}_{\text{FCC}}$ the face S_{b_0} of $\mathcal{V}_{\text{FCC}}(0)$ is shared with the Voronoi cell $\mathcal{V}_{\text{FCC}}(b_0)$. Lastly, we have $|\mathcal{V}_{\text{FCC}}(x)| = \frac{1}{2}\sqrt{2}$ for all $x \in \mathcal{L}_{\text{FCC}}$.




FIGURE 6. Left: the Voronoi cell V_{FCC} of the FCC lattice. Right: the Voronoi cell V_{HCP} of the HCP lattice.

Lemma 4.2. (*Voronoi cell in the HCP-lattice*) *Let us take $x \in \mathcal{L}_{\text{HCP}}$. Then*

$$\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(x) = \begin{cases} x + V_{\text{HCP}} & \text{if } x \in \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\}, \\ x - V_{\text{HCP}} & \text{if } x \in (v_1 + \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\}), \end{cases} \quad (59)$$

where

$$V_{\text{HCP}} := \left\{ y \in \mathbb{R}^3 : \max_{b \in \mathcal{N}_{\text{HCP}}} \langle b, y \rangle \leq \frac{1}{2} \right\}.$$

For $b_0 \in \mathcal{N}_{\text{HCP}}$ we set

$$S_{b_0} := \left\{ y \in \mathbb{R}^3 : \max_{b \in \mathcal{N}_{\text{HCP}}} \langle b, y \rangle = \langle b_0, y \rangle = \frac{1}{2} \right\}. \quad (60)$$

If $b_0 \in \{\pm e_1, \pm e_2, \pm(e_1 - e_2)\}$ the face S_{b_0} is a trapezoid of area $\frac{1}{4}\sqrt{2}$. If $b_0 \in \{v_1, v_1 - e_1, v_1 - e_2, v_1 - e_3, v_1 - e_1 - e_3, v_1 - e_2 - e_3\}$ the face S_{b_0} is a rhombus of area $\frac{1}{8}\sqrt{6}$. Moreover, for each $b_0 \in \mathcal{N}_{\text{HCP}}$ the face S_{b_0} is shared with the Voronoi cell $\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(b_0)$. Lastly, we have $|\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(x)| = \frac{1}{2}\sqrt{2}$ for all $x \in \mathcal{L}_{\text{HCP}}$.

Proof of Lemma 4.1. We split the proof of the lemma into four steps. First, we prove (57). In the second step, we show that each face is a rhombus and calculate its area. Lastly, we show that each neighboring Voronoi cell $V_{\text{FCC}}(b)$, $b \in \mathcal{N}_{\text{FCC}}$ shares one face with the Voronoi cell $V_{\text{FCC}}(0)$.

Step 1.(Proof of (57)) To check (57), since \mathcal{L}_{FCC} is a Bravais-lattice (see (8)), it suffices to consider the case $x = 0$. Let $\mathcal{V}_{\mathcal{L}_{\text{FCC}}}(0)$ denote the Voronoi cell of \mathcal{L}_{FCC} at $x = 0$ defined according to (13).

Step 1.1.($\mathcal{V}_{\mathcal{L}_{\text{FCC}}}(0) \subset V_{\text{FCC}}$) Let $y \in \mathcal{V}_{\mathcal{L}_{\text{FCC}}}(0)$. By the very definition of Voronoi cell we have that for all $b \in \mathcal{N}_{\text{FCC}}$ it holds $|y| \leq |y - b|$. Noting that $|b| = 1$ for all $b \in \mathcal{N}_{\text{FCC}} \subset \mathcal{L}_{\text{FCC}}$, we have

$$|y| \leq |y - b| \iff |y|^2 \leq |y - b|^2 = |y|^2 - 2\langle b, y \rangle + |b|^2 \iff \langle b, y \rangle \leq \frac{1}{2},$$

that is the inclusion $\mathcal{V}_{\mathcal{L}_{\text{FCC}}}(0) \subset V_{\text{FCC}}$.

Step 1.2.($V_{\text{FCC}} \subset \mathcal{V}_{\mathcal{L}_{\text{FCC}}}(0)$) We show that for $y \in V_{\text{FCC}}$ we have $|y| \leq |y - z|$ for all $z \in \mathcal{L}_{\text{FCC}}$.

This is equivalent to

$$y \in V_{\text{FCC}} \implies \langle y, z \rangle \leq \frac{1}{2}|z|^2 \quad \text{for all } z \in \mathcal{L}_{\text{FCC}}. \quad (61)$$

We first observe that if $z \in \mathcal{N}_{\text{FCC}}$, (61) is trivial since $|z| = 1$. Next, we prove (61) for all $z \in \mathcal{L}_{\text{FCC}} \setminus \mathcal{N}_{\text{FCC}}$. We distinguish two cases:

- (a) $z = \lambda_1 b_j + \lambda_2 b_k$ $\lambda_1, \lambda_2 \in \mathbb{Z}$, $j, k \in \{1, 2, 3\}$, $j \neq k$;
- (b) $z = \lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3$ $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{Z}$.

Proof in case (a). We only show the statement for $z = \lambda_1 b_1 + \lambda_2 b_2$ for $\lambda_1, \lambda_2 \in \mathbb{Z}$, the cases with any other combination of two vectors being analogous. If $\lambda_1 \lambda_2 \geq 0$, since $\langle b_1, b_2 \rangle \geq 0$, we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, \lambda_1 b_1 + \lambda_2 b_2 \rangle \leq \frac{1}{2}|\lambda_1 b_1|^2 + \frac{1}{2}|\lambda_2 b_2|^2 \\ &= \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2|^2 - \lambda_1 \lambda_2 \langle b_1, b_2 \rangle \leq \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2|^2 = \frac{1}{2}|z|^2. \end{aligned}$$

On the other hand, if $\lambda_1 \lambda_2 \leq 0$ and without loss of generality $|\lambda_1| \leq |\lambda_2|$, noting that $b_1 - b_2 \in \mathcal{N}_{\text{FCC}}$, we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, \lambda_1 b_1 + \lambda_2 b_2 \rangle = \langle y, (\lambda_2 + \lambda_1) b_2 + \lambda_1 (b_1 - b_2) \rangle \leq \frac{1}{2}|(\lambda_2 + \lambda_1) b_2|^2 + \frac{1}{2}|\lambda_1 (b_1 - b_2)|^2 \\ &= \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2|^2 - \lambda_1 (\lambda_2 + \lambda_1) \langle (b_1 - b_2), b_2 \rangle \leq \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2|^2. \end{aligned}$$

Here, the last inequality follows, since $|b_1| = |b_2|$ and therefore $\lambda_1 (\lambda_2 + \lambda_1) \langle (b_1 - b_2), b_2 \rangle \geq 0$. This concludes case (a).

Proof in case (b). We now show that (57) holds true in the case of $b = \lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3$ with $\lambda_i \in \mathbb{Z}$. We restrict to the case $\lambda_1 \geq 0, \lambda_2 \geq 0$ and $\lambda_3 \leq 0$, since if all λ_i are of the same sign, (57) can be deduced from the fact that it holds true for $b \in \mathcal{N}_{\text{FCC}}$ and the fact that $\langle b_j, b_k \rangle \geq 0$. Without loss of generality, we assume $|\lambda_2| \leq |\lambda_3|$. Hence, observing that $b_2 - b_3 \in \mathcal{N}_{\text{FCC}}$, we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, \lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3 \rangle = \langle y, \lambda_1 b_1 + (\lambda_3 + \lambda_2) b_3 + \lambda_2 (b_2 - b_3) \rangle \\ &\leq \frac{1}{2}|\lambda_1 b_1 + (\lambda_3 + \lambda_2) b_3|^2 + \frac{1}{2}|\lambda_2 (b_2 - b_3)|^2 \\ &= \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3|^2 - \langle (\lambda_1 b_1 + (\lambda_3 + \lambda_2) b_3), \lambda_2 (b_2 - b_3) \rangle \\ &= \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3|^2 - (\lambda_3 + \lambda_2) \lambda_2 \langle b_2 - b_3, b_3 \rangle \leq \frac{1}{2}|\lambda_1 b_1 + \lambda_2 b_2 + \lambda_3 b_3|^2. \end{aligned}$$

Here, the last inequality follows from $|b_2| = |b_3|$ and $\lambda_3 + \lambda_2 \leq 0$ whereas the equality in the last line is due to $\langle b_1, b_2 \rangle = \langle b_1, b_3 \rangle = \langle b_2, b_3 \rangle$. This concludes case (b) and with that Step 1.2.

Step 2. (The faces of the Voronoi cell) To show that each face of the Voronoi cell V_{FCC} is a rhombus with area $\frac{1}{4}\sqrt{2}$ we first exploit its symmetries. Let $i \in \{1, 2, 3\}$ and let $T_i: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear mapping that flips the i -th entry, i.e.

$$(T_i x)_j = \begin{cases} -x_i & \text{if } i = j, \\ x_j & \text{otherwise.} \end{cases}$$

We observe that

$$T_i \mathcal{N}_{\text{FCC}} = \{T_i b: b \in \mathcal{N}_{\text{FCC}}\} = \mathcal{N}_{\text{FCC}}, \quad \text{for all } i \in \{1, 2, 3\}.$$

Moreover, given a permutation $\pi \in S_3$ we have that

$$\pi \mathcal{N}_{\text{FCC}} = \{\pi b : b \in \mathcal{N}_{\text{FCC}}\} = \mathcal{N}_{\text{FCC}}.$$

It therefore suffices to restrict only to the case in which the vector b_0 agrees with the vector $b_1 \in \mathcal{N}_{\text{FCC}}$. We claim that this face has corners given by

$$c_1 = \left(\frac{1}{2}\sqrt{2}, 0, 0 \right), c_2 = \left(0, \frac{1}{2}\sqrt{2}, 0 \right), c_3 = \frac{1}{4} \left(\sqrt{2}, \sqrt{2}, \sqrt{2} \right), c_4 = \frac{1}{4} \left(\sqrt{2}, \sqrt{2}, -\sqrt{2} \right). \quad (62)$$

Note that, if this were true then it is easy to see that S_{b_0} is a rhombus and $\mathcal{H}^2(S_{b_0}) = \frac{1}{4}\sqrt{2}$. It remains to prove (62). Let us denote by y a corner of S_{b_0} . We can assume that $y_1, y_2 \geq 0$. Were this not the case, then there could be $b' \in \mathcal{N}_{\text{FCC}}$ such that $\langle b', y \rangle > \langle b, y \rangle$, thus contradicting the definition of S_{b_0} in (58). If $y_1 = 0$ (or $y_2 = 0$), then $y_2 = \frac{1}{2}\sqrt{2}$ (resp. $y_1 = \frac{1}{2}\sqrt{2}$) and since, $\langle b', y \rangle \leq \frac{1}{2}$ for all $b' \in \mathcal{N}_{\text{FCC}}$ we have $y_3 = 0$. Hence, we find the two corners with coordinates $(\frac{1}{2}\sqrt{2}, 0, 0)$ and $(0, \frac{1}{2}\sqrt{2}, 0)$. Now, if $y_1 > 0$ and $y_2 > 0$, then assuming that $y_3 \geq 0$ we have that the corner is equal to $\langle b_1, y \rangle = \langle b_2, y \rangle = \langle b_3, y \rangle = \frac{1}{2}\sqrt{2}$. Thus, necessarily $y_1 = y_2 = y_3 = \frac{1}{4}\sqrt{2}$. If instead $y_3 < 0$, then the corner is equal to $\langle b_1, y \rangle = \langle b_2, y \rangle = \langle b_1 - b_3, y \rangle = \frac{1}{2}\sqrt{2}$ which implies $y_1 = y_2 = -y_3 = \frac{1}{4}\sqrt{2}$. Hence (62) holds true and this concludes Step 2.

Step 3.(Neighbors share faces) We want to show that for each $b_0 \in \mathcal{N}_{\text{FCC}}$ we have that the face S_{b_0} of $V_{\text{FCC}}(0)$ is shared with the Voronoi cell $V_{\text{FCC}}(b_0)$. By the symmetries shown in Step 2 it suffices to prove this statement only for $b_0 = b_1$. Using (62) we see that the corners of the face S_{b_0} of the Voronoi cell $V_{\text{FCC}}(0)$ coincide with the corners of the face $S_{-b_0} + b_0$ of the Voronoi cell $V_{\text{FCC}}(b_0)$.

Step 4.(Volume of the Voronoi cell) In order to calculate the volume of the Voronoi cell we note that \mathcal{L}_{FCC} is a Bravais-lattice with spanning vectors b_1, b_2, b_3 . Since, the Voronoi cells of all the points are the same, it suffices to calculate the fraction of points per unit volume. This, then gives also the volume per point. Since, the Voronoi cells are space filling the volume per point is equal to the volume of each Voronoi cell. Due to (15) we have that

$$|T_{\text{FCC}}| = \frac{1}{2}\sqrt{2}.$$

Furthermore, we have that

$$\bigcup_{x \in \mathcal{L}_{\text{FCC}}} (x + T_{\text{FCC}}) = \mathbb{R}^3, \text{ and } \mathcal{L}_{\text{FCC}} \cap T_{\text{FCC}} = \{0\}.$$

Hence, each points of the lattice occupies a volume $|T_{\text{FCC}}| = \frac{1}{2}\sqrt{2}$ and the volume of the Voronoi cell must be the same. This concludes Step 3 and thus the proof of the lemma. \square

Proof of Lemma 4.2. We split the proof of the lemma into four steps. First, we prove (59). In the second step, we show that 6 of the faces are rhombi, the 6 other faces are trapezoids, and we calculate the area of each face. Lastly, given $x \in \mathcal{L}_{\text{HCP}}$, we show that each neighboring Voronoi cell $\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(y), y \in \mathcal{N}_{\text{HCP}}(x)$ shares a face with the Voronoi cell $V_{\text{HCP}}(x)$.

Step 1.(Shape of the Voronoi cell) The purpose of this step is to prove (59). Here, we only show this equality in the case that $x = 0$, the case $x \neq 0$ being treated in a similar fashion.

Step 1.1.($\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(0) \subset V_{\text{HCP}}$) Given $y \in \mathcal{V}_{\mathcal{L}_{\text{HCP}}}(0)$ we have that $|y| \leq |y - b|$. Now, noting that $|b| = 1$ for all $b \in \mathcal{N}_{\text{HCP}} \subset \mathcal{L}_{\text{HCP}}$, we have

$$|y|^2 \leq |y - b|^2 = |y|^2 - 2\langle y, b \rangle + |b|^2 \iff \langle b, y \rangle \leq \frac{1}{2}.$$

This concludes Step 1.1.

Step 1.2. ($V_{\text{HCP}} \subset \mathcal{V}_{\mathcal{L}_{\text{HCP}}}(0)$) We show that for $y \in V_{\text{HCP}}$ we have $|y| \leq |y - z|$, for all $z \in \mathcal{L}_{\text{HCP}}$. This is equivalent to

$$y \in V_{\text{HCP}} \implies \langle y, z \rangle \leq \frac{1}{2}|z|^2 \quad \text{for all } z \in \mathcal{L}_{\text{HCP}}. \quad (63)$$

Since, $|b| = 1$ for all $b \in \mathcal{N}_{\text{HCP}}$ (63) is true for all $b \in \mathcal{N}_{\text{HCP}}$. Next, we prove (63) for all $z \in \mathcal{L}_{\text{HCP}} \setminus \mathcal{N}_{\text{HCP}}$. We distinguish several cases:

- (a) $z = \lambda_1 e_1 + \lambda_2 e_2, \lambda_1, \lambda_2 \in \mathbb{Z}$;
- (b) $z = \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3, \lambda_1, \lambda_2, \lambda_3 \in \mathbb{Z}$;
- (c) $z = v_1 + \lambda_1 e_1 + \lambda_2 e_2, \lambda_1, \lambda_2 \in \mathbb{Z}$;
- (d) $z = v_1 + \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3, \lambda_1, \lambda_2, \lambda_3 \in \mathbb{Z}$;

Proof in case (a). If $\lambda_1, \lambda_2 \geq 0$, using that $\langle e_1, e_2 \rangle \geq 0$, we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, \lambda_1 e_1 + \lambda_2 e_2 \rangle \leq \frac{1}{2}|\lambda_1 e_1|^2 + \frac{1}{2}|\lambda_2 e_2|^2 \\ &= \frac{1}{2}|\lambda_1 e_1 + \lambda_2 e_2|^2 - \lambda_1 \lambda_2 \langle e_1, e_2 \rangle \leq \frac{1}{2}|\lambda_1 e_1 + \lambda_2 e_2|^2 = \frac{1}{2}|z|^2. \end{aligned}$$

On the other hand, if $\lambda_1 \lambda_2 \leq 0$ and without loss of generality $\lambda_1 \geq |\lambda_2| \geq 0$, noting that $e_2 - e_1 \in \mathcal{N}_{\text{HCP}}$, we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, \lambda_1 e_1 + \lambda_2 e_2 \rangle = \langle y, \lambda_2 (e_2 - e_1) + (\lambda_1 + \lambda_2) e_1 \rangle \leq \frac{1}{2}|\lambda_2 (e_2 - e_1)|^2 + \frac{1}{2}|(\lambda_2 + \lambda_1) e_1|^2 \\ &= \frac{1}{2}|\lambda_1 e_1 + \lambda_2 e_2|^2 - \lambda_2 (\lambda_1 + \lambda_2) \langle e_2 - e_1, e_1 \rangle \leq \frac{1}{2}|\lambda_1 e_1 + \lambda_2 e_2|^2 = \frac{1}{2}|z|^2. \end{aligned}$$

Here, the last inequality follows, since $\lambda_2 \leq 0 \leq \lambda_1 + \lambda_2$ and $\langle e_2 - e_1, e_1 \rangle \leq 0$. This concludes case (a).

Proof in case (b). We first show that $\langle y, e_3 \rangle \leq \frac{1}{2}|e_3|^2$. Using that $v_1, v_1 - e_1, v_1 - e_2 \in \mathcal{N}_{\text{HCP}}$, that $3v_1 - e_1 - e_2 = \frac{3}{2}e_3$ we have

$$\begin{aligned} \langle y, e_3 \rangle &= \frac{2}{3} \langle y, v_1 + v_1 - e_1 + v_1 - e_2 \rangle \\ &\leq \frac{1}{3} \left(\left| \frac{1}{3}(e_1 + e_2) \right|^2 + \left| \frac{1}{3}(e_2 - 2e_1) \right|^2 + \left| \frac{1}{3}(e_1 - 2e_2) \right|^2 \right) + \left| \frac{1}{2}e_3 \right|^2 \leq \frac{1}{2}|e_3|^2. \end{aligned}$$

Here, the last inequality follows by calculating the norms of $e_1 + e_2, e_1 - 2e_2, e_2 - 2e_1$ and e_3 by using (7). Note that now, the case of $z = \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3$ follows from case (a) using that $\langle e_3, e_1 \rangle = \langle e_3, e_2 \rangle = 0$.

Proof of case (c). Let $z = v_1 + \lambda_1 e_1 + \lambda_2 e_2$. If $\lambda_1, \lambda_2 \geq 0$ we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, v_1 + \lambda_1 e_1 + \lambda_2 e_2 \rangle \leq \frac{1}{2}|v_1|^2 + \frac{1}{2}|\lambda_1 e_1|^2 + \frac{1}{2}|\lambda_2 e_2|^2 \leq \frac{1}{2}|v_1|^2 + \frac{1}{2}|\lambda_1 e_1 + \lambda_2 e_2|^2 \\ &= \frac{1}{2}|v_1 + \lambda_1 e_1 + \lambda_2 e_2|^2 - \langle v_1, \lambda_1 e_1 + \lambda_2 e_2 \rangle \leq \frac{1}{2}|v_1 + \lambda_1 e_1 + \lambda_2 e_2|^2. \end{aligned}$$

The second inequality uses that $\langle e_1, e_2 \rangle \geq 0$ and the last inequality uses that $\langle v_1, e_1 \rangle, \langle v_1, e_2 \rangle \geq 0$. Now assume that $\lambda_1 \geq 0, \lambda_2 < 0$. Then, since $\langle (v_1 - e_2), e_1 \rangle = 0$ and $\langle v_1 - e_2, e_2 \rangle \leq 0$, again exploiting that $v_1 - e_2 \in \mathcal{N}_{\text{HCP}}$ it holds that

$$\begin{aligned} \langle y, z \rangle &= \langle y, (v_1 - e_2) + \lambda_1 e_1 + (\lambda_2 + 1)e_2 \rangle \leq \frac{1}{2}|v_1 - e_2|^2 + \frac{1}{2}|\lambda_1 e_1 + (\lambda_2 + 1)e_2|^2 \\ &= \frac{1}{2}|v_1 + \lambda_1 e_1 + \lambda_2 e_2|^2 - \langle v_1 - e_2, \lambda_1 e_1 + (\lambda_2 + 1)e_2 \rangle \leq \frac{1}{2}|v_1 + \lambda_1 e_1 + \lambda_2 e_2|^2 = \frac{1}{2}|z|^2. \end{aligned}$$

The case where $\lambda_1 < 0, \lambda_2 \geq 0$ (resp. $\lambda_1, \lambda_2 < 0$) is being treated in a similar fashion by replacing $v_1 - e_2$ with $v_1 - e_1$ (resp. $v_1 - e_1 - e_2$).

Proof of case (d). Here, we only treat the case of $z = v_1 + \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3, \lambda_3 \geq 0$. Since $\langle v_1 + \lambda_1 e_1 + \lambda_2 e_2, e_3 \rangle \geq 0$, we have

$$\begin{aligned} \langle y, z \rangle &= \langle y, v_1 + \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3 \rangle \leq \frac{1}{2} |v_1 + \lambda_1 e_1 + \lambda_2 e_2|^2 + \frac{1}{2} |\lambda_3 e_3|^2 \\ &= \frac{1}{2} |v_1 + \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3|^2 - \langle v_1 + \lambda_1 e_1 + \lambda_2 e_2, e_3 \rangle \leq \frac{1}{2} |v_1 + \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3|^2 = \frac{1}{2} |z|^2. \end{aligned}$$

The case of $\lambda_3 < 0$ follows by replacing v_1 with $v_1 - e_3$ in the last two cases (c) and (d). This concludes Step 1.2 and, together with Step 1.1, shows (59).

Step 2.(The faces of the Voronoi cell) In order to calculate the faces of V_{HCP} we use (59) and exploit its symmetries. We note that if $R \in SO(3)$ is any rotation of integer multiples of $2\pi/3$ around the x_3 -axis we have that

$$R\mathcal{N}_{\text{HCP}} = \{Rb : b \in \mathcal{N}_{\text{HCP}}\} = \mathcal{N}_{\text{HCP}}. \quad (64)$$

Moreover, if $T_3 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the reflection with respect to the (x_1, x_2) -plane, i.e.

$$(T_3 x)_j := \begin{cases} x_j & j = 1, 2, \\ -x_3 & j = 3, \end{cases} \quad (65)$$

we have that

$$T_3 \mathcal{N}_{\text{HCP}} = \{T_3 b : b \in \mathcal{N}_{\text{HCP}}\} = \mathcal{N}_{\text{HCP}}. \quad (66)$$

Exploiting (64) and (66), it suffices to find the corners of S_{b_0} in (60) for

$$(a) \quad b_0 = e_1, \quad (b) \quad b_0 = -e_1, \quad (c) \quad b_0 = v_1.$$

Corners in case (a). We claim that in the case of $b_0 = e_1$ that the corners of S_{b_0} are given by the points

$$\begin{aligned} c_1 &= \left(\frac{1}{2}, \frac{1}{6}\sqrt{3}, \frac{1}{12}\sqrt{6} \right), \quad c_2 = \left(\frac{1}{2}, \frac{1}{6}\sqrt{3}, -\frac{1}{12}\sqrt{6} \right), \\ c_3 &= \left(\frac{1}{2}, -\frac{1}{6}\sqrt{3}, \frac{1}{6}\sqrt{6} \right), \quad c_4 = \left(\frac{1}{2}, -\frac{1}{6}\sqrt{3}, -\frac{1}{6}\sqrt{6} \right). \end{aligned} \quad (67)$$

In particular, the face S_{b_0} is a trapezoid with two bases of length $\frac{1}{6}\sqrt{6}, \frac{1}{3}\sqrt{6}$ and height $\frac{1}{3}\sqrt{3}$. Hence, $\mathcal{H}^2(S_{b_0}) = \frac{1}{4}\sqrt{2}$. It remains to prove (67). Let $y \in S_{b_0}$ be a corner. Due to (66), we can assume that $y_3 \geq 0$, since the other corners are just found by applying the mapping T_3 (see (65)) to the corners with positive coordinates. By the definition of S_{b_0} we have that $\langle y, e_1 \rangle \geq \langle y, e_1 - e_2 \rangle$ which is equivalent to $\langle y, e_2 \rangle \geq 0$. Now, if $\langle y, e_2 \rangle > 0$, then y is given by $\langle y, e_1 \rangle = \langle y, e_2 \rangle = \langle y, v_1 \rangle = \frac{1}{2}$. This linear system has a unique solution given by $c_1 = (\frac{1}{2}, \frac{1}{6}\sqrt{3}, \frac{1}{12}\sqrt{6})$. On the other hand, if $\langle y, e_2 \rangle = 0$, then y is given by $\langle y, e_2 \rangle = 0, \langle y, e_1 \rangle = \langle y, v_1 \rangle = \frac{1}{2}$. The unique solution of this linear system is given by $c_3 = (\frac{1}{2}, -\frac{1}{6}\sqrt{3}, \frac{1}{6}\sqrt{6})$. This shows (67) and concludes case (a).

Corners in case (b). We claim that in the case of $b_0 = -e_1$ that the corners of S_{b_0} are given by the points

$$\begin{aligned} c_1 &= \left(-\frac{1}{2}, \frac{1}{6}\sqrt{3}, \frac{1}{12}\sqrt{6} \right), \quad c_2 = \left(-\frac{1}{2}, \frac{1}{6}\sqrt{3}, -\frac{1}{12}\sqrt{6} \right), \\ c_3 &= \left(-\frac{1}{2}, -\frac{1}{6}\sqrt{3}, \frac{1}{6}\sqrt{6} \right), \quad c_4 = \left(-\frac{1}{2}, -\frac{1}{6}\sqrt{3}, -\frac{1}{6}\sqrt{6} \right). \end{aligned} \quad (68)$$

In particular, the face S_{b_0} is a trapezoid with two bases of length $\frac{1}{6}\sqrt{6}$, $\frac{1}{3}\sqrt{6}$ and height $\frac{1}{3}\sqrt{3}$. Hence, $\mathcal{H}^2(S_{b_0}) = \frac{1}{4}\sqrt{2}$. It remains to prove (68). Let $y \in S_{b_0}$ be a corner. Due to (66), as in case (a), we can assume that $y_3 \geq 0$. By the definition of S_{b_0} we have that $\langle y, -e_1 \rangle \geq \langle y, e_2 - e_1 \rangle$ which is equivalent to $\langle y, e_2 \rangle \leq 0$. Now, if $\langle y, e_2 \rangle = 0$, then y is given by $\langle y, e_2 \rangle = 0$, $\langle y, v_1 - e_1 \rangle = \langle y, -e_1 \rangle = \frac{1}{2}$. We see that the unique solution of this linear system is given by $c_1 = (-\frac{1}{2}, \frac{1}{6}\sqrt{3}, \frac{1}{12}\sqrt{6})$. On the other hand, if $\langle y, e_2 \rangle < 0$, then y is given by $\langle y, v_1 \rangle = 0$, $\langle y, -e_1 \rangle = \langle y, -e_2 \rangle = \frac{1}{2}$. The unique solution is now given by $c_3 = (-\frac{1}{2}, -\frac{1}{6}\sqrt{3}, \frac{1}{6}\sqrt{6})$. This shows (68) and concludes case (b).

Corners in case (c). We claim that in the case of $b_0 = v_1$ that the corners of S_{b_0} are given by the points

$$\begin{aligned} c_1 &= \left(\frac{1}{2}, \frac{1}{6}\sqrt{3}, \frac{1}{12}\sqrt{6} \right), c_2 = \left(0, 0, \frac{1}{4}\sqrt{6} \right), \\ c_3 &= \left(0, \frac{1}{3}\sqrt{3}, \frac{1}{6}\sqrt{6} \right), c_4 = \left(\frac{1}{2}, -\frac{1}{6}\sqrt{3}, \frac{1}{6}\sqrt{6} \right). \end{aligned} \quad (69)$$

In particular, the face S_{b_0} is a rhombus. Hence, $\mathcal{H}^2(S_{b_0}) = \frac{1}{8}\sqrt{6}$. It remains to prove (67). Let $y \in S_{b_0}$ be a corner. By the definition of S_{b_0} we have that $\langle y, v_1 \rangle \geq \langle y, v_1 - e_1 \rangle, \langle y, v_1 - e_2 \rangle$ which is equivalent to $\langle y, e_1 \rangle, \langle y, e_2 \rangle \geq 0$. Now if, $\langle y, e_2 \rangle > 0$ then the corner solves the linear system $\langle y, e_1 \rangle = \langle y, e_2 \rangle = \langle y, v_1 \rangle = \frac{1}{2}$. Its unique solution is $c_1 = (\frac{1}{2}, \frac{1}{6}\sqrt{3}, \frac{1}{12}\sqrt{6})$. On the other hand if $\langle y, e_2 \rangle = 0$, then the corners are given by those y such that $\langle y, e_2 \rangle = 0, \langle y, e_1 \rangle = \langle y, v_1 \rangle = \frac{1}{2}$ or $\langle y, e_1 \rangle = \langle y, e_2 \rangle = 0, \langle y, v_1 \rangle = \frac{1}{2}$. These points have coordinates $c_2 = (\frac{1}{2}, -\frac{1}{6}\sqrt{3}, \frac{1}{6}\sqrt{6})$ and $c_3 = (0, 0, \frac{1}{4}\sqrt{6})$. Finally, if $\langle y, e_1 \rangle = 0$ and $\langle y, e_2 \rangle > 0$, then y is obtained by solving $\langle y, e_1 \rangle = 0, \langle y, e_2 \rangle = \langle y, v_1 \rangle = \frac{1}{2}$. Hence it has coordinates $c_4 = (0, \frac{\sqrt{3}}{3}, \frac{\sqrt{6}}{6})$. This proves (69) and concludes Step 2.

Step 3.(Neighbors share faces) We want to show that for each $b_0 \in \mathcal{N}_{\text{HCP}}$ we have that the face $\overline{S_{b_0}}$ of $\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(0)$ is shared with the Voronoi cell $\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(b_0)$. By Step 1 we have that $\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(0) = V_{\text{HCP}}$ and $\mathcal{V}_{\mathcal{L}_{\text{HCP}}}(b_0) = b_0 - V_{\text{HCP}}$. Hence, they share the side $\langle y, b_0 \rangle = \frac{1}{2} = \langle b_0 - y, b_0 \rangle$.

Step 4.(Volume of the Voronoi cell) In order to calculate the volume of the Voronoi cell we note that \mathcal{L}_{HCP} is periodic with respect to the vectors e_1, e_2, e_3 . Since the Voronoi cells of all the points occupy the same volume, it suffices to calculate the fraction of points per unit volume. The inverse of this number is the volume per point. Since the Voronoi cells are space filling the volume per point is equal to the volume of each Voronoi cell. Due to (16) we have that

$$|T_{\text{HCP}}| = \sqrt{2}.$$

Furthermore, we have that

$$\bigcup_{x \in \text{span}_{\mathbb{Z}}\{e_1, e_2, e_3\}} (x + T_{\text{HCP}}) = \mathbb{R}^3, \text{ and } \mathcal{L}_{\text{HCP}} \cap T_{\text{HCP}} = \{0, v_1\}.$$

Hence, the volume per point is $\frac{1}{2}|T_{\text{HCP}}| = \frac{1}{2}\sqrt{2}$ and it agrees with the volume of the Voronoi cell. This concludes Step 4 and thus the proof of the lemma. \square

We are now in the position to prove Theorem 2.3.

Proof of Theorem 2.3. All the statements are consequences of Proposition 5.10, Lemma 5.11, Theorem 5.14 and Lemma 5.16 once we show that \mathcal{L}_{FCC} and \mathcal{L}_{HCP} are periodic admissible sets (according to Definition 5.1 and Definition 5.8) and we observe that, due to Lemma 4.1 and Lemma 4.2, $\mathcal{N}_{\text{FCC}}(x) = \mathcal{NN}(x)$ (in the sense of Definition 5.2) as well as $\mathcal{N}_{\text{HCP}}(x) = \mathcal{NN}(x)$

in the respective cases. We first show that both lattices are admissible sets. Let us first observe that

$$(T_{\text{FCC}} + x) \cap \mathcal{L}_{\text{FCC}} \neq \emptyset \quad \text{and} \quad (T_{\text{HCP}} + x) \cap \mathcal{L}_{\text{HCP}} \neq \emptyset \quad \text{for all } x \in \mathbb{R}^3.$$

Therefore, (L1) is satisfied for both \mathcal{L}_{FCC} and \mathcal{L}_{HCP} with

$$R := \max\{\text{diam}(T_{\text{FCC}}), \text{diam}(T_{\text{HCP}})\} \leq \max\left\{\sum_{k=1}^3 |e_k|, \sum_{k=1}^3 |b_k|\right\} < +\infty,$$

where we recalled Definition 15 and Definition 16. On the other hand, (L2) is satisfied with $r = 1$, see the discussion at the definition of the FCC and HCP lattice in Section 2. Concerning periodicity: We observe that for all $z = (z_1, z_2, z_3) \in \mathbb{Z}^3$ we have

$$\mathcal{L}_{\text{FCC}} = \mathcal{L}_{\text{FCC}} + \sum_{k=1}^3 z_k b_k, \quad \text{and} \quad \mathcal{L}_{\text{HCP}} = \mathcal{L}_{\text{HCP}} + \sum_{k=1}^3 z_k e_k$$

and thus both \mathcal{L}_{FCC} and \mathcal{L}_{HCP} are periodic according to Definition 5.8. The statement follows by Theorem 5.14 with $c_{nn}(x) = 1$. \square

5. GENERAL PERIODIC LATTICES

This section deals with integral representation and concentrated-compactness properties of energies defined on general periodic lattices.

Definition 5.1. Let $\Sigma \subset \mathbb{R}^n$ be a countable set of points in \mathbb{R}^n . We call Σ an *admissible* set of points if the following two conditions hold:

- (L1) There exists $R > 0$ such that $\inf_{x \in \mathbb{R}^n} \#(\Sigma \cap B_R(x)) \geq 1$;
- (L2) There exists $r > 0$ such that $\text{dist}(x, \Sigma \setminus \{x\}) \geq r$ for all $x \in \Sigma$.

Definition 5.2. We define the Voronoi cell of $x \in \Sigma$ as

$$\mathcal{V}(x) := \{z \in \mathbb{R}^n : |x - z| \leq |y - z| \text{ for all } y \in \Sigma\}. \quad (70)$$

The set of nearest neighbors of Σ is defined by

$$\mathcal{NN}(\Sigma) := \{(x, y) \in \Sigma \times \Sigma : \mathcal{H}^{n-1}(\mathcal{V}(x) \cap \mathcal{V}(y)) > 0\},$$

We set $\mathcal{NN}(x) = \{y \in \Sigma : (x, y) \in \mathcal{NN}(\Sigma)\}$. Given $\varepsilon > 0$ we denote by $\varepsilon\Sigma := \{\varepsilon x : x \in \Sigma\}$ and for $x \in \varepsilon\Sigma$ we set $\mathcal{V}_\varepsilon(x) = \varepsilon\mathcal{V}(\varepsilon^{-1}x)$ the Voronoi cell of $x \in \varepsilon\Sigma$, and $\mathcal{NN}_\varepsilon(x) = \{y \in \varepsilon\Sigma : \varepsilon^{-1}(x, y) \in \mathcal{NN}(\Sigma)\}$ the set of nearest neighbors of x in $\varepsilon\Sigma$.

We now define for $u : \Sigma \rightarrow \{0, 1\}$ the two energy functionals given by

$$F_\varepsilon(u, A) := \sum_{\substack{(x, y) \in \mathcal{NN}(\Sigma) \\ \varepsilon x \in A}} \varepsilon^{n-1} c_{nn}(x - y) |u(\varepsilon x) - u(\varepsilon y)| \quad (71)$$

and

$$\hat{F}_\varepsilon(u, A) := \sum_{\substack{(x, y) \in \mathcal{NN}(\Sigma) \\ \varepsilon x, \varepsilon y \in A}} \varepsilon^{n-1} c_{nn}(x - y) |u(\varepsilon x) - u(\varepsilon y)|, \quad (72)$$

where $c_{nn} : \mathbb{R}^n \rightarrow [0, +\infty]$ satisfies

$$C^{-1} \leq c_{nn}(x) \leq C, \quad \text{for all } x \in \mathbb{R}^n. \quad (73)$$

When $A = \mathbb{R}^n$ we omit the dependence on it and write $F_\varepsilon(u) = F_\varepsilon(u, \mathbb{R}^n)$ and $\hat{F}_\varepsilon(u) = \hat{F}_\varepsilon(u, \mathbb{R}^n)$.

Remark 5.3. (Difference between F_ε and \hat{F}_ε) We want to point out the difference between F_ε and \hat{F}_ε : In the formula defining F_ε the sum is taken over all $(x, y) \in \mathcal{NN}(\Sigma)$ such that $\varepsilon x \in A$. Instead in the case of \hat{F}_ε the sum takes only those $(x, y) \in \mathcal{NN}(\Sigma)$ such that both $\varepsilon x \in A$ and $\varepsilon y \in A$. The functional $F_\varepsilon(u, \cdot)$ is an additive set function on disjoint sets, i.e., given $A, B \subset \mathbb{R}^n$ such that $A \cap B = \emptyset$, we have

$$F_\varepsilon(u, A \cup B) = F_\varepsilon(u, A) + F_\varepsilon(u, B),$$

whereas $\hat{F}_\varepsilon(u, \cdot)$ is only super-additive on disjoint sets. Our Γ -convergence result will be stated for the functional F_ε . The reason for us to introduce \hat{F}_ε is that our proof will use the integral representation result proven in [3], see Theorem 5.7. However, we will show later on that the Γ -convergence of F_ε is equivalent to that of \hat{F}_ε .

Given $X \subset \varepsilon\Sigma$ we write with a slight abuse of notation

$$F_\varepsilon(X, A) = F_\varepsilon(\chi_X, A).$$

Note that, due to (73), we have

$$C^{-1} \sum_{x \in X} \varepsilon^{n-1} \#(\mathcal{NN}_\varepsilon(x) \setminus X) \leq F_\varepsilon(X, A) \leq C \sum_{x \in X} \varepsilon^{n-1} \#(\mathcal{NN}_\varepsilon(x) \setminus X).$$

Hypothesis (73) corresponds to [3, Hypothesis 1] in the case that, according to the notation in [3, Equation (5.23)], $c_{nn}^\varepsilon(x, y) = c_{nn}(x - y)$ and $c_{lr}^\varepsilon(x, y) = 0$. It is worth observing that in [3] a more general class of functionals was investigated, namely those for which also certain long-range interactions between points in Σ contribute to the energy, i.e., $c_{lr}(x, y) \neq 0$. For the sake of exposition and simplicity, here we consider the case $c_{lr} = 0$, that is the energy accounts only for the nearest neighbor interactions. However, with some more involved multi-scale constructions, all the statements below extend to the more general case where also long-range interactions are considered.

Definition 5.4. Given $X \subset \varepsilon\mathcal{L}$ we define the rescaled empirical measures associated to X as

$$\mu_\varepsilon = \varepsilon^n \sum_{x \in X} \delta_x. \quad (74)$$

Furthermore, recalling (70), we define

$$V_\varepsilon(X) := \bigcup_{x \in X} V_\varepsilon(x). \quad (75)$$

Henceforth, we drop the dependence on X and simply write V_ε . Given $A \subset \mathbb{R}^n$ open with $\partial A \in \text{Lip}$, with slight abuse of notation we define $F_\varepsilon: \mathcal{M}_+(A) \rightarrow [0, +\infty]$ (similarly $\hat{F}_\varepsilon: \mathcal{M}_+(A) \rightarrow [0, +\infty]$) by

$$F_\varepsilon(\mu, A) = \begin{cases} F_\varepsilon(X, A) & \mu \text{ is given by (74) for some } X \subset \varepsilon\mathcal{L}; \\ +\infty & \text{otherwise.} \end{cases}$$

Additionally, we define $F_\varepsilon: L_{\text{loc}}^1(A) \rightarrow [0, +\infty]$ (similarly $\hat{F}_\varepsilon: L_{\text{loc}}^1(A) \rightarrow [0, +\infty]$) by

$$F_\varepsilon(u, A) = \begin{cases} F_\varepsilon(X, A) & u = \chi_{V_\varepsilon} \text{ and } V_\varepsilon \text{ is given by (75) for some } X \subset \varepsilon\mathcal{L}; \\ +\infty & \text{otherwise.} \end{cases}$$

It is necessary for us to introduce two different domains of definition for the extended functional F_ε , since we want to make use of [3, Theorem 5.5]. As it will turn out the two types of extension are equivalent, cf. Lemma 5.11 and Corollary 5.12.

Let $(\Omega, \mathbb{P}, \mathcal{F})$ be a probability space. Hereafter we recall some definitions from [3] (Definition 5.1 and Definition 5.4):

Definition 5.5. We say that a family $(\tau_z)_{z \in \mathbb{Z}^n}$, $\tau_z: \Omega \rightarrow \Omega$, is an *additive group action* on Ω if

$$\tau_{z_1+z_2} = \tau_{z_2} \circ \tau_{z_1} \quad \text{for all } z_1, z_2 \in \mathbb{Z}^n.$$

Such an additive group action is called *measure preserving* if

$$\mathbb{P}(\tau_z B) = \mathbb{P}(B) \quad \text{for all } B \in \mathcal{F}, z \in \mathbb{Z}^n.$$

If in addition, for all $B \in \mathcal{F}$ we have

$$(\tau_z(B) = B \quad \text{for all } z \in \mathbb{Z}^n) \implies \mathbb{P}(B) \in \{0, 1\},$$

then $(\tau_z)_{z \in \mathbb{Z}^n}$ is called *ergodic*.

Definition 5.6. A random variable $\mathcal{L}: \Omega \rightarrow (\mathbb{R}^n)^{\mathbb{Z}^n}$, $\omega \mapsto \mathcal{L}(\omega) = \{\mathcal{L}(\omega)(i)\}_{i \in \mathbb{Z}^n}$ is called a stochastic lattice. We say that \mathcal{L} is admissible if $\mathcal{L}(\omega)$ is admissible in the sense of Definition 5.1 and the constants r, R can be chosen independent of ω \mathbb{P} -almost surely. The stochastic lattice \mathcal{L} is said to be *stationary* if there exists a measure preserving group action $(\tau_z)_{z \in \mathbb{Z}^n}$ on Ω such that, for \mathbb{P} -almost every $\omega \in \Omega$, $\mathcal{L}(\tau_z \omega) = \mathcal{L}(\omega) + z$. If in addition $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then \mathcal{L} is called ergodic, too.

We now state a simplified version of [3, Theorem 5.5] which is enough for our purposes.

Theorem 5.7. (*Stochastic homogenization of spin systems*) Let \mathcal{L} be a stationary and ergodic stochastic lattice and let \hat{F}_ε be defined by (72). Let $A \subset \mathbb{R}^n$ be open and bounded with $\partial A \in \text{Lip}$. For \mathbb{P} -almost every ω the functionals $F_\varepsilon(\omega)$ Γ -converge with respect to the strong $L^1(A)$ -topology to the functional $F_{\text{hom}}: L^1(A) \rightarrow [0, +\infty]$ defined by

$$F_{\text{hom}}(u, A) := \begin{cases} \int_{J(u) \cap A} \varphi_{\text{hom}}(\nu_u) d\mathcal{H}^{n-1} & \text{if } u \in BV(A; \{0, 1\}), \\ +\infty & \text{otherwise.} \end{cases}$$

The function $\varphi_{\text{hom}}: \mathbb{R}^n \rightarrow [0, +\infty]$ is given by

$$\varphi_{\text{hom}}(\nu) = \lim_{T \rightarrow +\infty} \frac{1}{T^{n-1}} \int_{\Omega} \inf \{ F(u, Q_T^\nu) : u: \mathcal{L}(\omega) \rightarrow \{0, 1\}, \\ u(i) = \nu(i) \text{ for } i \in \mathcal{L}(\omega) \setminus Q_{T-l_T}^\nu \} d\mathbb{P}(\omega),$$

where $l_T \rightarrow +\infty$ and $l_T/T \rightarrow 0$ as $T \rightarrow +\infty$.

Definition 5.8. Let $\mathcal{L} \subset \mathbb{R}^n$ be an admissible set of points. We say that \mathcal{L} is periodic if there exists a basis $\{e_1, \dots, e_n\} \subset \mathbb{R}^n$ such that

$$\mathcal{L} + e_k = \mathcal{L} \quad \text{for all } k = 1, \dots, n.$$

We denote by

$$Q := \left\{ \sum_{k=1}^n \lambda_k e_k : 0 \leq \lambda_k < 1 \right\}$$

the periodicity cell of \mathcal{L} . We set

$$\rho := \frac{\#(\mathcal{L} \cap Q)}{|Q|}. \tag{76}$$

In the following we assume, up to a change of coordinates that $\{e_1, \dots, e_n\}$ is the standard orthonormal basis of \mathbb{R}^n .

We collect the following general properties of periodic admissible set of points.

Lemma 5.9. *(Properties of periodic admissible sets) Let \mathcal{L} be a periodic admissible set of points. The following holds true:*

- (i) $B_{r/2}(x) \subset \mathcal{V}(x) \subset B_R(x)$ for all $x \in \mathcal{L}$;
- (ii) $\mathcal{NN}(x) \subset B_{2R}(x)$ for all $x \in \mathcal{L}$;
- (iii) There exists $C = C(n, r, R) \in (0, +\infty)$ such that $\sup_{x \in \mathcal{L}} \#\mathcal{NN}(x) \leq C$. In particular, $\partial\mathcal{V}(x)$ is made out of finitely many $(n-1)$ -dimensional polyhedral faces;
- (iv) There exists $C = C_{\mathcal{L}} > 0$ such that for all $X \subset \varepsilon\mathcal{L}$ and $A \subset \mathbb{R}^n$ there holds

$$F_\varepsilon(X, A) \leq C \text{Per}(V_\varepsilon, (A)_{R\varepsilon}) \text{ and } \text{Per}(V_\varepsilon, A) \leq CF_\varepsilon(X, (A)_{R\varepsilon}).$$

Proof. Apart from (iv) all of these facts are classical. We collect their proof here for completeness.

proof of (i), (ii): Let $x \in \mathcal{L}$. The inclusion $B_{r/2}(x) \subset \mathcal{V}(x)$ follows from (L2) since for all $y \in B_{r/2}(x)$ and $z \in \mathcal{L} \setminus \{x\}$

$$|z - y| \geq |x - z| - |x - y| \geq r - r/2 = r/2 \geq |x - y|.$$

As for the inclusion $\mathcal{V}(x) \subset B_R(x)$ assume that there exists $y \in \mathcal{V}(x) \setminus B_R(x)$. We have for all $z \in \mathcal{L} \setminus \{x\}$

$$|y - z| \geq |y - x| \geq R.$$

This implies that $B_R(y) \cap \mathcal{L} = \emptyset$ contradicting (L1). Finally $\mathcal{NN}(x) \subset B_{2R}(x)$ since for $y \in \mathcal{NN}(x)$ we have that $\mathcal{V}(x) \cap \mathcal{V}(y) \neq \emptyset$ which implies $B_R(x) \cap B_R(y) \neq \emptyset$.

proof of (iii): Due to (i) and (ii) we have that $B_{r/2}(y) \cap B_{r/2}(z) = \emptyset$, $y, z \in \mathcal{NN}(x)$, $y \neq z$ and $B_{r/2}(y) \subset B_{2R+r}(x)$ for all $y \in \mathcal{NN}(x)$. Therefore

$$\omega_n \left(\frac{r}{2}\right)^n \#\mathcal{NN}(x) = \bigcup_{y \in \mathcal{NN}(x)} |B_{r/2}(y)| \leq |B_{2R+r}(x)| \leq \omega_n (2R+r)^n$$

and thus the claim follows with $C = (2 + 4R/r)^n$.

proof of (iv): The desired inequalities follow from the following observation: Given $x \in X$, we have that

$$\mathcal{H}^{n-1}(\mathcal{V}_\varepsilon(x) \cap \partial V_\varepsilon) > 0 \iff \mathcal{NN}_\varepsilon(x) \setminus X \neq \emptyset. \quad (77)$$

Additionally, we note that, for $x \in X$ such that $\mathcal{NN}_\varepsilon(x) \setminus X \neq \emptyset$, there exists $C > 0$ such that

$$C^{-1}\varepsilon^{n-1} \leq \mathcal{H}^{n-1}(\mathcal{V}_\varepsilon(x) \cap \partial V_\varepsilon) \leq C\varepsilon^{n-1}. \quad (78)$$

Now, summing over all $x \in X \cap A$ and noting that each Voronoi cell intersects only a finite number of other Voronoi cells, using (77), (78), (i), and (iii), we obtain

$$\begin{aligned} F_\varepsilon(X, A) &= \sum_{x \in X \cap A} \varepsilon^{n-1} \#\mathcal{NN}_\varepsilon(x) \setminus X \leq C \sum_{x \in X \cap A} \mathcal{H}^{n-1}(\mathcal{V}_\varepsilon(x) \cap \partial V_\varepsilon) \\ &\leq C \mathcal{H}^{n-1}(\partial V_\varepsilon \cap (A)_{R\varepsilon}) = C \text{Per}(V_\varepsilon, (A)_{R\varepsilon}). \end{aligned}$$

This yields the first inequality in (iv). On the other hand, owing to (i) we have $\partial V_\varepsilon \cap A \subset \bigcup_{x \in X \cap (A)_{R\varepsilon}} (\mathcal{V}_\varepsilon(x) \cap \partial V_\varepsilon)$, and thus, by (77) and (78),

$$\begin{aligned} \text{Per}(V_\varepsilon, A) &= \mathcal{H}^{n-1}(\partial V_\varepsilon \cap A) \leq \sum_{x \in X \cap (A)_{R\varepsilon}} \mathcal{H}^{n-1}(\mathcal{V}_\varepsilon(x) \cap \partial V_\varepsilon) \\ &\leq C\varepsilon^{n-1} \sum_{x \in X \cap (A)_{R\varepsilon}} \#(\mathcal{NN}_\varepsilon(x) \setminus X) = CF_\varepsilon(X, (A)_{R\varepsilon}). \end{aligned}$$

This shows the second inequality in (iv) and concludes the proof. \square

Proposition 5.10. (*Compactness of the piecewise-constant interpolants*) *Let \mathcal{L} be an admissible periodic set of points and F_ε defined in (71) with \mathcal{L} in place of Σ . Let $A \subset \mathbb{R}^n$ be open and let $\{X_\varepsilon\}_\varepsilon \subset \varepsilon\mathcal{L}$ be such that*

$$\sup_{\varepsilon > 0} F_\varepsilon(X_\varepsilon, A) < +\infty.$$

Then there exists a set of finite perimeter $V \subset A$ and a subsequence (not relabeled) such that $\chi_{V_\varepsilon} \rightarrow \chi_V$ with respect to the strong $L^1_{\text{loc}}(A)$ -topology.

Proof. Let X_ε be as above and let $A' \subset \subset A$ such that $(A')_{R\varepsilon} \subset A$, and $\partial A' \in \text{Lip}$. We observe, due to the second inequality of Lemma 5.9(iv),

$$\text{Per}(V_\varepsilon, A') \leq CF_\varepsilon(X_\varepsilon, (A')_{R\varepsilon}) \leq CF_\varepsilon(X_\varepsilon, A) \leq C.$$

Therefore

$$\|\chi_{V_\varepsilon}\|_{L^1(A')} + |D\chi_{V_\varepsilon}|(A') \leq C(|A'| + \text{Per}(V_\varepsilon, A')) \leq C.$$

We use [4, Theorem 3.39] to deduce that there exists a subsequence (depending on A') and a set of finite perimeter V such that $\chi_{V_\varepsilon} \rightarrow \chi_V$ in $L^1(A')$. By a diagonal argument on a sequence $A'_k \uparrow A$ as $k \rightarrow +\infty$, we obtain the claim. \square

Lemma 5.11. (*Equivalence of convergences*) *Let \mathcal{L} be an admissible periodic set of points and F_ε defined in (71) with \mathcal{L} in place of Σ . Let $A \subset \mathbb{R}^n$ be open and let $V \subset A$ be a set of finite perimeter and let $\{X_\varepsilon\}_\varepsilon \subset \varepsilon\mathcal{L}$ for each $\varepsilon > 0$ be such that*

$$\sup_{\varepsilon > 0} F_\varepsilon(X_\varepsilon, A) < +\infty. \quad (79)$$

Then, setting μ_ε and V_ε as in (74) and (75), the following are equivalent:

- (i) $\mu_\varepsilon \xrightarrow{*} \mu$ with respect to the weak star topology of measures and $\mu = \rho \mathcal{L}^n \llcorner V$.
- (ii) $\chi_{V_\varepsilon} \rightarrow \chi_V$ with respect to the strong $L^1_{\text{loc}}(A)$ -topology.

Proof. We proceed in two steps. First, we construct a sequence of auxiliary measure ν_ε and show that its weak*-convergence is equivalent to the weak*-convergence of the sequence of measures μ_ε . Then, for this sequence of measures we show that its weak*-convergence is equivalent to (ii). Step 1. (Construction of the auxiliary measure) Let $\{X_\varepsilon\}_\varepsilon$ be as in the assumptions of the lemma and let $v \in C_c(A)$ such that $\text{supp } v \subset \subset A$. We assume that $\varepsilon > 0$ is small enough such that for all $k \in \varepsilon\mathbb{Z}^n$ there holds

$$Q_{(3+R)\varepsilon}(k) \cap \text{supp } v \neq \emptyset \implies Q_{(3+R)\varepsilon}(k) \subset \subset A. \quad (80)$$

Fix $R_0 > 0$ such that $\text{supp } v \subset B_{R_0}$. Since v is uniformly continuous, it admits a modulus of continuity $\omega = \omega_v: [0, +\infty) \rightarrow [0, +\infty)$, i.e., an increasing function such that $\omega(0) = 0$ and

$$|v(x) - v(y)| \leq \omega(|x - y|) \text{ for all } x, y \in \mathbb{R}^n. \quad (81)$$

We set

$$\begin{aligned}\mathcal{I}_\varepsilon^{\text{full}} &:= \{k \in \varepsilon\mathbb{Z}^d : Q_{(3+R)\varepsilon}(k) \cap X = Q_{(3+R)\varepsilon}(k) \cap \varepsilon\mathcal{L}\}, \\ \mathcal{I}_\varepsilon^{\text{empty}} &:= \{k \in \varepsilon\mathbb{Z}^d : Q_{(3+R)\varepsilon}(k) \cap X = \emptyset\},\end{aligned}$$

and $\mathcal{I}_\varepsilon^{\text{bad}} := \{k \in \varepsilon\mathbb{Z}^n : Q_{(3+R)\varepsilon}(k) \cap \text{supp } v \neq \emptyset\} \setminus (\mathcal{I}_\varepsilon^{\text{full}} \cup \mathcal{I}_\varepsilon^{\text{empty}})$. We now set

$$\nu_\varepsilon := \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \varepsilon^n \#(\mathcal{L} \cap Q) \delta_k.$$

Our goal is to show that

$$\mu_\varepsilon \xrightarrow{*} \mu \quad \text{in the sense of measures} \quad \iff \quad \nu_\varepsilon \xrightarrow{*} \mu \quad \text{in the sense of measures.} \quad (82)$$

First of all we claim that

$$\#\mathcal{I}_\varepsilon^{\text{bad}} \leq C\varepsilon^{1-n}. \quad (83)$$

In fact, due to its definition and (80), we have that for all $k \in \mathcal{I}_\varepsilon^{\text{bad}}$ there exists $x \in X \cap Q_{(3+R)\varepsilon}(k) \subset A$ such that $\#(\mathcal{N}\mathcal{N}(x) \setminus X) \neq \emptyset$. Therefore, since for $k \in \varepsilon\mathbb{Z}^n$ fixed $Q_{(3+R)\varepsilon}(k) \cap Q_{(3+R)\varepsilon}(j) \neq \emptyset$ for only finitely many j independent of ε , we have

$$\varepsilon^{n-1} \#\mathcal{I}_\varepsilon^{\text{bad}} \leq \sum_{k \in \mathcal{I}_\varepsilon^{\text{bad}}} \sum_{x \in Q_{(3+R)\varepsilon}(k) \cap X} \varepsilon^{n-1} \#(\mathcal{N}\mathcal{N}(x) \setminus X) \leq CF_\varepsilon(X_\varepsilon, A).$$

Using (79) yields (83). Let now $k \in \mathcal{I}_\varepsilon^{\text{full}}$. Then

$$\begin{aligned}\int_{Q_\varepsilon(k)} v(x) d\nu_\varepsilon(x) &= v(k) \varepsilon^n \#(\mathcal{L} \cap Q) = \sum_{x \in Q_\varepsilon(k) \cap \varepsilon\mathcal{L}} \varepsilon^n v(k) \\ &= \sum_{x \in Q_\varepsilon(k) \cap X} \varepsilon^n v(x) + \sum_{x \in Q_\varepsilon(k) \cap \varepsilon\mathcal{L}} \varepsilon^n (v(k) - v(x)) \\ &= \int_{Q_\varepsilon(k)} v(x) d\mu_\varepsilon(x) + \sum_{x \in Q_\varepsilon(k) \cap \varepsilon\mathcal{L}} \varepsilon^n (v(k) - v(x)).\end{aligned}$$

Thus,

$$\left| \int_{Q_\varepsilon(k)} v(x) d(\nu_\varepsilon - \mu_\varepsilon)(x) \right| \leq \sum_{x \in Q_\varepsilon(k) \cap \varepsilon\mathcal{L}} \varepsilon^n |v(k) - v(x)| \leq \omega(\varepsilon\sqrt{n}) |Q_\varepsilon(k)|$$

and, recalling that $\text{supp } v \subset B_{R_0}$ and (81), we have for $\varepsilon > 0$ small enough

$$\sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \left| \int_{Q_\varepsilon(k)} v(x) d(\nu_\varepsilon - \mu_\varepsilon)(x) \right| \leq \omega(\varepsilon\sqrt{n}) |B_{2R_0}|. \quad (84)$$

Noting that both $|\mu_\varepsilon|(Q_\varepsilon(k))$ and $|\nu_\varepsilon|(Q_\varepsilon(k))$ are bounded above by $\varepsilon^n \#(\mathcal{L} \cap Q) \leq C\varepsilon^n$ for all $k \in \varepsilon\mathbb{Z}^n$, using (83), we observe

$$\sum_{k \in \mathcal{I}_\varepsilon^{\text{bad}}} \left| \int_{Q_\varepsilon(k)} v(x) d(\nu_\varepsilon - \mu_\varepsilon)(x) \right| \leq 2\|v\|_\infty \#\mathcal{I}_\varepsilon^{\text{bad}} \varepsilon^n \#(\mathcal{L} \cap Q) \leq C\varepsilon \|v\|_\infty. \quad (85)$$

Therefore, noting that $\mu_\varepsilon \lfloor_{Q_\varepsilon(k)} = \nu_\varepsilon \lfloor_{Q_\varepsilon(k)}$ for $k \in \mathcal{I}_\varepsilon^{\text{empty}}$, using (84) and (85), we obtain

$$\left| \int_{\mathbb{R}^n} v(x) d(\nu_\varepsilon - \mu_\varepsilon)(x) \right| \leq \omega(\varepsilon\sqrt{n}) |B_{2R}| + C\varepsilon \|v\|_\infty.$$

This shows (82).

Step 2.(Equivalence of convergence) We now prove that

$$\chi_{V_\varepsilon} \rightarrow \chi_V \text{ in } L^1_{\text{loc}}(A) \iff \nu_\varepsilon \xrightarrow{*} \mu \text{ in the sense of measures.}$$

First of all, recalling ρ defined in (76), we note

$$\begin{aligned} \int_{\mathbb{R}^n} v \, d\nu_\varepsilon &= \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \varepsilon^n \#(\mathcal{L} \cap Q) v(k) = \frac{\#(\mathcal{L} \cap Q)}{|Q|} \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} |Q_\varepsilon(k)| v(k) \\ &= \rho \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \int_{Q_\varepsilon(k)} v(y) \, dy + \rho \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \int_{Q_\varepsilon(k)} (v(k) - v(y)) \, dy. \end{aligned} \quad (86)$$

Now, due to (81), we have for $\varepsilon > 0$ small enough

$$\left| \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \int_{Q_\varepsilon(k)} (v(k) - v(y)) \, dy \right| \leq \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \int_{Q_\varepsilon(k)} |v(k) - v(y)| \, dy \leq C\omega(\varepsilon\sqrt{n})|B_{2R_0}|. \quad (87)$$

Note that, by Lemma 5.9(i), we have

$$\rho \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \int_{Q_\varepsilon(k)} v(y) \, dy = \rho \sum_{k \in \mathcal{I}_\varepsilon^{\text{full}}} \int_{Q_\varepsilon(k) \cap V_\varepsilon} v(y) \, dy \quad (88)$$

and also $V_\varepsilon \cap Q_\varepsilon(k) = \emptyset$ for $k \in \mathcal{I}_\varepsilon^{\text{empty}}$. Note that by (83) we have

$$\sum_{k \in \mathcal{I}_\varepsilon^{\text{bad}}} \int_{Q_\varepsilon(k)} |v(y)| \, dy \leq \varepsilon^n \# \mathcal{I}_\varepsilon^{\text{bad}} \|v\|_\infty \leq C\varepsilon \|v\|_\infty. \quad (89)$$

Due to (86)–(89), we obtain that

$$\rho \chi_{V_\varepsilon} \xrightarrow{*} \mu \text{ in the sense of measures} \iff \nu_\varepsilon \xrightarrow{*} \mu \text{ in the sense of measures.}$$

Now clearly (ii) implies (i), since the $L^1_{\text{loc}}(A)$ convergence of the characteristic functions implies their weak* convergence as measures. As for the implication (i) to (ii) we proceed as follows. Let $V \subset A$ be a set of finite perimeter and assume that $\mu_\varepsilon \xrightarrow{*} \mu$ and $\mu = \rho \mathcal{L}^n \llcorner V$. By Step 1 this is equivalent to $\nu_\varepsilon \xrightarrow{*} \mu$, hence to $\rho \chi_{V_\varepsilon} \xrightarrow{*} \rho \chi_V$ in the sense of measures (which is to say that $\chi_{V_\varepsilon} \xrightarrow{*} \chi_V$). Take now an arbitrary subsequence (not relabeled) of $\{X_\varepsilon\}_\varepsilon$. We show that there exists a further subsequence (again not relabeled) such that $\chi_{V_\varepsilon} \rightarrow \chi_V$ with respect to the strong $L^1_{\text{loc}}(A)$ -topology. Since $L^1_{\text{loc}}(A)$ -topology satisfies the Urysohn property this implies the claim. By the compactness statement in Proposition 5.10 we have that there exists a set of finite perimeter $V' \subset \mathbb{R}^n$ and a further subsequence $\{X_{\varepsilon_k}\}_k \subset \{X_\varepsilon\}_\varepsilon$ such that $\chi_{V_{\varepsilon_k}} \rightarrow \chi_{V'}$ with respect to the strong $L^1_{\text{loc}}(A)$ -topology. Since this implies their weak* convergence as measures and we already know that the whole sequence converges to χ_V we deduce $V = V'$ which implies the claim and concludes the proof of the lemma. \square

Corollary 5.12. (Equivalence of Γ -convergence) *Let \mathcal{L} be an admissible periodic set of points and let F_ε be defined in (71) with \mathcal{L} in place of Σ . Let $A \subset \mathbb{R}^n$ be a bounded open set such that $\partial A \in \text{Lip}$. Then the following statements are equivalent:*

- (i) $F_\varepsilon: \mathcal{M}_+(A) \rightarrow [0, +\infty]$ Γ -converges with respect to the weak* convergence of measures to the functional $F_{\text{hom}}: \mathcal{M}_+(A) \rightarrow [0, +\infty]$ defined as

$$F_{\text{hom}}(\mu, A) := \begin{cases} \int_{\partial^* V \cap A} \varphi_{\text{hom}}(\nu) \, d\mathcal{H}^{n-1} & \text{if } \mu = \rho \mathcal{L}^n \llcorner V; \\ +\infty & \text{otherwise.} \end{cases}$$

(ii) $F_\varepsilon: L^1_{\text{loc}}(A) \rightarrow [0, +\infty]$ Γ -converges with respect to strong $L^1_{\text{loc}}(A)$ -topology to the functional $F_{\text{hom}}: L^1_{\text{loc}}(A) \rightarrow [0, +\infty]$ defined as

$$F_{\text{hom}}(u, A) := \begin{cases} \int_{\partial^* V \cap A} \varphi_{\text{hom}}(\nu) \, d\mathcal{H}^{n-1} & \text{if } u = \chi_V \text{ and } \chi_V \in BV(A); \\ +\infty & \text{otherwise.} \end{cases}$$

Proof. The statement follows directly from Lemma 5.11. \square

Remark 5.13. The analogous statements are true for \hat{F}_ε as well.

Theorem 5.14. (Γ -convergence for periodic admissible lattices) *Let \mathcal{L} be an admissible periodic set of points and let F_ε be defined by (71) with \mathcal{L} in place of Σ . Let $A \subset \mathbb{R}^n$ be bounded open set with $\partial A \in \text{Lip}$ or $A = \mathbb{R}^n$. The functionals F_ε Γ -converge with respect to weak* convergence of measures to the functional $F_{\text{hom}}: \mathcal{M}_+(A) \rightarrow [0, +\infty]$ defined by*

$$F_{\text{hom}}(\mu, A) := \begin{cases} \int_{\partial^* V \cap A} \varphi_{\text{hom}}(\nu) \, d\mathcal{H}^{n-1} & \text{if } \mu = \rho \mathcal{L}^n \llcorner V; \\ +\infty & \text{otherwise.} \end{cases}$$

The function $\varphi_{\text{hom}}: \mathbb{R}^n \rightarrow [0, +\infty]$ is given by

$$\varphi_{\text{hom}}(\nu) = \lim_{T \rightarrow +\infty} \frac{1}{T^{n-1}} \inf \{ F(X, Q_T^\nu) : X \subset \mathcal{L}, \chi_X(i) = u_\nu(i) \text{ for } i \in \mathcal{L} \setminus Q_{T-l_T}^\nu \},$$

where $l_T \rightarrow +\infty$ and $l_T/T \rightarrow 0$ as $T \rightarrow +\infty$.

Proof. Step 1. (Probabilistic setup) We exploit the integral representation result (Theorem 5.7) to obtain the specific form of the Γ -limit. We fix $(\Omega, \mathcal{F}, \mathbb{P}) = (\{0\}, \{\{0\}, \emptyset\}, \delta_0)$ to be a probability space and a trivial additive and ergodic group action (see Definition 5.1 in [3]) $\tau_z: \{0\} \rightarrow \{0\}$, $z \in \mathbb{Z}^3$ given by $\tau_z(0) = 0$. With respect to this group action $\mathcal{L}(0) = \mathcal{L}$ is an admissible stationary and ergodic stochastic lattice according to Definition 5.6. In fact, since \mathcal{L} is periodic according to Definition 5.8, for all $z = (z_1, z_2, z_3) \in \mathbb{Z}^3$ we have

$$\mathcal{L}(0) = \mathcal{L}, \quad \mathcal{L}(\tau_z(0)) = \mathcal{L}(0) = \mathcal{L}(0) + \sum_{k=1}^n z_k e_k.$$

Therefore, all conditions of Theorem 5.7 are satisfied. This shows that for $\hat{F}_\varepsilon: L^1_{\text{loc}}(A) \rightarrow [0, +\infty]$ we have

$$\Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \hat{F}_\varepsilon(\chi_V, A) = F_{\text{hom}}(\chi_V, A)$$

for all $A \subset \mathbb{R}^n$, $\partial A \in \text{Lip}$, $\chi_V \in BV(A, \{0, 1\})$. Note that, by Corollary 5.12, this is equivalent to saying that for $\hat{F}_\varepsilon: \mathcal{M}_+(A) \rightarrow [0, +\infty]$ we have

$$\Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \hat{F}_\varepsilon(\mu, A) = F_{\text{hom}}(\mu, A).$$

This concludes Step 1.

Step 2. (Γ -convergence of F_ε) We use the Γ -convergence of \hat{F}_ε obtained in Step 1 in order to prove the Γ -convergence of F_ε . Let us first prove the result for $A \subset \mathbb{R}^n$ open and bounded with $\partial A \in \text{Lip}$. We note that

$$\hat{F}_\varepsilon(X, A) \leq F_\varepsilon(X, A)$$

and therefore for all $\mu_\varepsilon \xrightarrow{*} \mu$, we have

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(\mu_\varepsilon, A) \geq \liminf_{\varepsilon \rightarrow 0} \hat{F}_\varepsilon(\mu_\varepsilon, A) \geq F_{\text{hom}}(\mu, A).$$

Therefore $\Gamma\text{-lim inf}_{\varepsilon \rightarrow 0} F_\varepsilon(\mu, A) \geq F_{\text{hom}}(\mu, A)$. Next, we show the Limsup-inequality. Let $V \subset A$ be such that

$$F_{\text{hom}}(\rho\chi_V, A) < +\infty \quad (90)$$

since otherwise there is nothing to prove. Using Lemma 5.9(iv), we obtain that $\text{Per}(V, A) < +\infty$, i.e., V is a set of finite perimeter. In particular, by [25, Theorem 1.3] (noting that $\partial A \in \text{Lip}$ implies that A is an extension domain), there exists an extension $V_\varepsilon \subset \mathbb{R}^n$ such that

- (i) $V_\varepsilon \cap A = V \cap A$ up to a set of zero \mathcal{L}^n -measure;
- (ii) $\text{Per}(V_\varepsilon, \mathbb{R}^n) \leq C\text{Per}(V, A)$;
- (iii) $\mathcal{H}^{d-1}(\partial^* V_\varepsilon \cap \partial A) = 0$.

Now (ii) together with (90) implies, again by Lemma 5.9(iv), that

$$F_{\text{hom}}(\rho\chi_{V_\varepsilon}, \mathbb{R}^n) = \int_{\partial^* V_\varepsilon} \varphi_{\text{hom}}(\nu) d\mathcal{H}^{n-1} < +\infty.$$

Fix $\delta > 0$ and let $\mu_\varepsilon^\delta \rightarrow \rho\chi_{V_\varepsilon}$ weakly* in $\mathcal{M}_+((A)_\delta)$ be such that

$$\limsup_{\varepsilon \rightarrow 0} \hat{F}_\varepsilon(\mu_\varepsilon^\delta, (A)_\delta) \leq F_{\text{hom}}(\rho\chi_{V_\varepsilon}, (A)_\delta).$$

Note that, due to Lemma 5.9(ii), for every $\varepsilon < \delta/(2R)$ we have that

$$F_\varepsilon(X, A) \leq \hat{F}_\varepsilon(X, (A)_\delta).$$

Therefore, recalling Definition 2.2, we obtain,

$$\Gamma\text{-lim sup}_{\varepsilon \rightarrow 0} F_\varepsilon(\rho\chi_V, A) \leq \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(\mu_\varepsilon^\delta, A) \leq \limsup_{\varepsilon \rightarrow 0} \hat{F}_\varepsilon(\mu_\varepsilon^\delta, (A)_\delta) \leq F_{\text{hom}}(\rho\chi_{V_\varepsilon}, (A)_\delta).$$

Sending $\delta \rightarrow 0$ we obtain

$$\Gamma\text{-lim sup}_{\varepsilon \rightarrow 0} F_\varepsilon(\rho\chi_V, A) \leq F_{\text{hom}}(\rho\chi_{V_\varepsilon}, \bar{A}) = F_{\text{hom}}(\rho\chi_{V_\varepsilon}, A) = F_{\text{hom}}(\rho\chi_V, A),$$

where the last equality follows by properties (i) and (iii) of V_ε . This shows the desired integral representation for all $A \subset \mathbb{R}^n$ such that $\partial A \in \text{Lip}$.

Step 3.(Integral representation on unbounded sets) It remains to prove the integral representation of the Γ -limit for \mathbb{R}^n . The Liminf-inequality follows by monotonicity since for all $R > 0$ and $X \subset \varepsilon\mathcal{L}$ we have

$$F_\varepsilon(X, B_R) \leq F_\varepsilon(X)$$

and therefore, given $\mu_\varepsilon \rightarrow \rho\chi_V$ weakly* in $\mathcal{M}_+(\mathbb{R}^n)$, we have

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(\mu_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(\mu_\varepsilon, B_R) \geq F_{\text{hom}}(\rho\chi_V, B_R).$$

The claim follows by taking the supremum over $R > 0$. We now turn our attention to the Limsup inequality. We can assume without loss of generality that $V \subset \mathbb{R}^n$ is a set of finite perimeter and

$$C\text{Per}(V, \mathbb{R}^n) \leq \int_{\partial^* V} \varphi(\nu) d\mathcal{H}^{n-1} = F_{\text{hom}}(\rho\chi_V, \mathbb{R}^n) < +\infty \quad (91)$$

since otherwise there is nothing to prove. By the isoperimetric inequality there exists $C > 0$ such that

$$\min\{|V|, |\mathbb{R}^n \setminus V|\}^{\frac{n-1}{n}} \leq C\text{Per}(V, \mathbb{R}^n).$$

Without loss of generality we assume that $|V| < +\infty$. By the Fleming-Rishel formula we can find $\{R_k\}_k$ such that $R_k \rightarrow +\infty$ and

$$(i) \quad \mathcal{H}^{n-1}(V \cap \partial B_{R_k}) \leq \frac{1}{k} \quad \text{and} \quad (ii) \quad |V \cap B_{R_k}^c| \leq \frac{1}{k}. \quad (92)$$

We define $V_k = V \cap B_{R_k}$. Then, thanks to (92)(ii), $\chi_{V_k} \rightarrow \chi_V$ in $L^1(\mathbb{R}^n)$ and thus also $\rho_{\chi_{V_k}} \rightarrow \rho_{\chi_V}$ weakly* in $\mathcal{M}_+(\mathbb{R}^n)$. Furthermore, (92)(i) implies that

$$\lim_{k \rightarrow +\infty} F_{\text{hom}}(\rho_{\chi_{V_k}}) = F_{\text{hom}}(\rho_{\chi_V}).$$

It therefore suffices to construct the recovery sequence for $\rho_{\chi_{V_k}}$. Let $S_k = R_k + 2R$ and let $\mu_\varepsilon \rightarrow \rho_{V_k}$ in $L^1(B_{S_k})$ such that

$$\limsup_{\varepsilon \rightarrow 0} F_\varepsilon(\mu_\varepsilon, B_{S_k}) \leq F_{\text{hom}}(\rho_{\chi_V}, B_{S_k}). \quad (93)$$

We modify μ_ε such that

$$F_\varepsilon(\mu_\varepsilon) \leq F_\varepsilon(\mu_\varepsilon, B_{S_k}) + o(1), \quad (94)$$

and this, by (93), proves the statement. By Lemma 5.11, there exists $\eta_\varepsilon \rightarrow 0$ such that $|V_\varepsilon \cap A_{R_k, S}| \leq \eta_\varepsilon$ with V_ε defined in (75). Now, let us take $k \in \{[\frac{R_k}{\varepsilon} + R], \dots, [\frac{S}{\varepsilon} - 3R]\} =: \mathcal{K}_\varepsilon$. Noting that $\#\mathcal{K}_\varepsilon \approx \varepsilon^{-1}$, Lemma 5.9(i) implies that there exists $k_\varepsilon \in \mathcal{K}_\varepsilon$ such that

$$\begin{aligned} \omega_n \left(\frac{r}{2}\right)^n \varepsilon^{n-1} \#(X_\varepsilon \cap A_{k_\varepsilon \varepsilon, (k_\varepsilon + 3R)\varepsilon}) &\leq \omega_n \left(\frac{r}{2}\right)^n \varepsilon^n \sum_{k \in \mathcal{K}_\varepsilon} \#(X_\varepsilon \cap A_{k\varepsilon, (k+3R)\varepsilon}) \\ &\leq C |V_\varepsilon \cap A_{R_k, S}| \leq C \eta_\varepsilon. \end{aligned}$$

Here, we have used that for $k \in \mathcal{K}_\varepsilon$ fixed

$$A_{k\varepsilon, (k+3R)\varepsilon} \cap A_{j\varepsilon, (j+3R)\varepsilon} \neq \emptyset$$

for finitely many indices j independent of ε (clearly for $j > k + 3R$ or $j < k - 3R$ the intersection is empty). We can therefore define $\hat{X}_\varepsilon = \mathcal{L} \cap B_{k_\varepsilon}$ and employ Lemma 5.9(ii) to obtain

$$\begin{aligned} F_\varepsilon(\hat{X}_\varepsilon) &= F_\varepsilon(\hat{X}_\varepsilon, B_{k_\varepsilon}) \leq F_\varepsilon(X_\varepsilon, B_{k_\varepsilon}) + \bar{c} \max_{x \in \mathcal{L}} \#\mathcal{N}(x) \varepsilon^{n-1} \#(X_\varepsilon \cap A_{k_\varepsilon \varepsilon, (k_\varepsilon + 3R)\varepsilon}) \\ &\leq F_\varepsilon(X_\varepsilon, B_S) + C \eta_\varepsilon, \end{aligned}$$

which proves (94) and with it the claim. \square

In the following we show that for minimizing sequences we can improve Proposition 5.10 to obtain strong $L^1(\mathbb{R}^n)$ compactness. This implies that the rescaled empirical measures of such sequences converge to a suitably normalized Wulff shape for the limiting perimeter energy.

Lemma 5.15. (*Nucleation Lemma*) *For every $(p_0, v_0) \in (0, +\infty)^2$ there exists $m = m(p_0, v_0) > 0$ such that if $V \subset \mathbb{R}^n$, $\text{Per}(V) \leq p_0$, $|V| \geq v_0$, then there exists $x \in \mathbb{R}^n$ such that*

$$|V \cap B_1(x)| \geq m.$$

Proof. Assume that $|V \cap B_1(x)| < \delta \leq \omega_n/2$ for all $x \in \mathbb{R}^n$. Then, noting that $\chi_{B_1(x)}(y) = \chi_{B_1(y)}(x)$ for all $x, y \in \mathbb{R}^n$, we obtain

$$\omega_n |V| = \int_{x \in \mathbb{R}^n} \int_{y \in \mathbb{R}^n} \chi_V(x) \chi_{B_1(x)}(y) \, dy \, dx = \int_{x \in \mathbb{R}^n} \int_{y \in \mathbb{R}^n} \chi_{V \cap B_1(y)}(x) \, dy \, dx. \quad (95)$$

Due to the assumption we have $|V \cap B_1(x)| < \delta$ and therefore due to the relative isoperimetric inequality and Fubini's Theorem, we obtain

$$\begin{aligned} \int_{x \in \mathbb{R}^n} \int_{y \in \mathbb{R}^n} \chi_{V \cap B_1(y)}(x) \, dy \, dx &= \int_{y \in \mathbb{R}^n} |V \cap B_1(y)| \, dy \leq \delta^{1/n} \int_{y \in \mathbb{R}^n} |V \cap B_1(x)|^{1-\frac{1}{n}} \, dy \\ &\leq \frac{\delta^{1/n}}{C_B} \int_{y \in \mathbb{R}^n} \mathcal{H}^{n-1}(\partial^* V \cap B_1(y)) \, dy, \end{aligned} \quad (96)$$

where $C_B > 0$ denotes the relative isoperimetric constant of B_1 . Now, again due to Fubini's Theorem, we obtain

$$\begin{aligned} \int_{y \in \mathbb{R}^n} \mathcal{H}^{n-1}(\partial^* V \cap B_1(y)) \, dy &= \int_{y \in \mathbb{R}^n} \int_{x \in \mathbb{R}^n} \chi_{B_1(y)}(x) \, d\mathcal{H}^{n-1} \lfloor_{\partial^* V}(x) \, dy \\ &= \int_{x \in \mathbb{R}^n} \int_{y \in \mathbb{R}^n} \chi_{B_1(x)}(y) \, dy \, d\mathcal{H}^{n-1} \lfloor_{\partial^* V}(x) = \omega_n \mathcal{H}^{n-1}(\partial^* V) \end{aligned}$$

This together with (95) and (96) leads to

$$|V| \leq \frac{\delta^{1/n}}{C_B} \text{Per}(V).$$

Then, we conclude that $\delta \geq (C_B |V| / \text{Per}(V))^n$ and thus for any $m < \min\{\omega_n/2, (C_B v/p_0)^n\}$ the thesis holds. \square

Lemma 5.16. (*Concentration Lemma*) *Let $\{X_\varepsilon\}_\varepsilon$ and $\{n_\varepsilon\}_\varepsilon \subset \mathbb{N}$ be such that for all $\varepsilon > 0$ $X_\varepsilon \subset \varepsilon \mathcal{L}$, $\#X_\varepsilon = n_\varepsilon$, $n_\varepsilon \varepsilon^n \rightarrow \rho v$ as $\varepsilon \rightarrow 0$, and*

$$F_\varepsilon(X_\varepsilon) = \min_{\#X=n_\varepsilon} F_\varepsilon(X).$$

Then, there exists $\{\tau_\varepsilon\}_\varepsilon \subset \mathbb{R}^n$ such that $\tau_\varepsilon \in \varepsilon \text{span}_{\mathbb{Z}}\{e_1, \dots, e_n\}$, and $\mu_\varepsilon(\cdot - \tau_\varepsilon) \xrightarrow{} \rho \chi_W$, where W is the Wulff shape of φ_{hom} defined in Theorem 5.14 and such that $|W| = v$.*

Proof. Let $\{X_\varepsilon\}_\varepsilon$ be as in the assumptions of the Lemma.

Step 1. (Energy bound) We first show that

$$\sup_{\varepsilon > 0} F_\varepsilon(X_\varepsilon) < +\infty. \quad (97)$$

To this aim, for each $\varepsilon > 0$ we construct a competitor Y_ε such that $F_\varepsilon(Y_\varepsilon) \leq C$ for some constant $C > 0$ independent of ε . To this end let $r_\varepsilon > 0$ be the maximal $s > 0$ such that

$$\#(B_s \cap \varepsilon \mathcal{L}) \leq n_\varepsilon.$$

Due to Lemma 5.9(i), we have that $r_\varepsilon \leq r_0 < +\infty$ and

$$\omega_n \left(\frac{r}{2}\right)^n \varepsilon^n \#(A_{r_\varepsilon, r_\varepsilon+2\varepsilon R} \cap \varepsilon \mathcal{L}) \leq |A_{r_\varepsilon - \varepsilon r/2, r_\varepsilon+3\varepsilon R}| \leq C \varepsilon r_\varepsilon^{n-1} \leq C \varepsilon. \quad (98)$$

By the maximality in the choice of r_ε we have $\#(B_{r_\varepsilon+\varepsilon r} \cap \varepsilon \mathcal{L}) > n_\varepsilon$ and thus

$$n_\varepsilon - C \varepsilon^{1-n} \leq \#(B_{r_\varepsilon} \cap \varepsilon \mathcal{L}) \leq n_\varepsilon.$$

We set $Y_\varepsilon = (B_{r_\varepsilon} \cap \varepsilon \mathcal{L}) \cup Y_\varepsilon^0$, where $Y_\varepsilon^0 \subset \varepsilon \mathcal{L} \setminus B_{r_\varepsilon}$ such that $\#Y_\varepsilon = n_\varepsilon$. Due to Lemma 5.9 (ii), (iii), the minimality of X_ε , and (98), we have

$$\begin{aligned} F_\varepsilon(X_\varepsilon) &\leq F_\varepsilon(Y_\varepsilon) \\ &\leq \sum_{x \in \mathcal{L}_\varepsilon \cap B_{r_\varepsilon}} \varepsilon^{n-1} \#(\mathcal{N}_{\mathcal{L}_\varepsilon}(x) \setminus X) + C \varepsilon^{n-1} \#Y_\varepsilon^0 \leq C \#(A_{r_\varepsilon, r_\varepsilon+2\varepsilon R} \cap \mathcal{L}) \leq C. \end{aligned}$$

This shows (97) and concludes Step 1.

Step 2.(Nucleation) We show that if (97) holds, then there exist $v_1 > 0$ and $\{\tau_\varepsilon^1\}_\varepsilon \subset \mathbb{R}^n$ such that $\tau_\varepsilon^1 \in \varepsilon \text{span}_{\mathbb{Z}}\{e_1, \dots, e_n\}$ and, up to subsequences,

$$\mu_\varepsilon(\cdot - \tau_\varepsilon^1) \xrightarrow{*} \rho\chi_{V_1} \text{ in the sense of measures,} \quad (99)$$

where $|V_1| := v_1 \in (m, v]$, with $m = m(p_0, v_0) > 0$ given by Lemma 5.15. In fact, due to (97) and Lemma 5.9(iv) there exists $p_0 > 0$ such that $\text{Per}(V_\varepsilon) \leq p_0$ for all $\varepsilon > 0$. Additionally, due to Lemma 5.9(i) and the fact that $n_\varepsilon \varepsilon^n \rightarrow v > 0$, we have that $|V_\varepsilon| \geq v_1 > 0$ for all $\varepsilon > 0$ small enough. Thus, by Lemma 5.15 there exists $m = m(p_0, v_1)$ and $\{\tau_\varepsilon^1\}_\varepsilon \subset \mathbb{R}^d$ such that

$$|V_\varepsilon \cap B_1(\tau_\varepsilon^1)| \geq m. \quad (100)$$

Actually, by lowering m a bit we can assume without loss of generality that $\tau_\varepsilon^1 \in \varepsilon \text{span}_{\mathbb{Z}}\{e_1, \dots, e_n\}$.

By (100), (97), Proposition 5.10, and Lemma 5.11, we get (99) up to passing to a subsequence.

Step 3.(Splitting of the energy) We show that for any $0 < \delta < |V_1|$ there exists $S_\delta = S_\delta(V_1) > 0$ big enough and $S_\varepsilon \in (S_\delta, S_\delta + 1)$ such that, for $\varepsilon > 0$ small enough, there holds

$$\#(X_\varepsilon \cap B_{S_\delta}) \geq \omega_n^{-1} R^{-n} (|V_1| - \delta) \varepsilon^{-n} \quad \text{and} \quad \#(X_\varepsilon \cap A_{S_\varepsilon, S_\varepsilon + 3\varepsilon R}) \leq \delta \varepsilon^{1-n}. \quad (101)$$

First of all, we find S_δ such that for $\varepsilon > 0$ small enough

$$|V_1 \cap B_{S_\delta - R\varepsilon}| \geq |V_1 \cap B_{S_\delta - 1}| \geq |V_1| - \delta/2 \quad \text{and} \quad |V_1 \cap A_{S_\delta, S_\delta + 1}| \leq |V_1 \setminus B_{S_\delta - R\varepsilon}| \leq \delta/2.$$

Note that, due to Step 1 and Step 2, $\chi_{V_\varepsilon(\cdot - \tau_\varepsilon^1)} \rightarrow \chi_{V_1}$ in $L^1(B_{S_\delta + 1})$. Due to Lemma 5.9(i), we have for $\varepsilon > 0$ small enough

$$\omega_n \varepsilon^n R^n \#(X_\varepsilon \cap B_{S_\delta}) \geq |V_\varepsilon \cap B_{S_\delta - R\varepsilon}| \geq |V_1| - \delta; \quad (102)$$

$$\omega_n \varepsilon^n \left(\frac{r}{2}\right)^n \#(X_\varepsilon \cap A_{S_\delta + R\varepsilon, S_\delta + 1 - R\varepsilon}) \leq |V_\varepsilon \cap A_{S_\delta, S_\delta + 1}| \leq \delta. \quad (103)$$

Next, we find S_ε by averaging: We choose $k_\varepsilon \in \{[\frac{S}{\varepsilon} + R], \dots, [\frac{S+1}{\varepsilon} - 4R]\} =: \mathcal{K}_\varepsilon$ (note that $\mathcal{K}_\varepsilon \neq \emptyset$ and actually $\#\mathcal{K}_\varepsilon \approx \frac{S}{\varepsilon}$ for $\varepsilon > 0$ small enough) such that for some $C > 1$

$$\begin{aligned} \varepsilon^{n-1} \#(X_\varepsilon \cap A_{k_\varepsilon \varepsilon, (k_\varepsilon + 3R)\varepsilon}) &\leq C \sum_{k \in \mathcal{K}_\varepsilon} \varepsilon^n \#(X_\varepsilon \cap A_{k\varepsilon, (k+3R)\varepsilon}) \\ &\leq C \varepsilon^n \#(X_\varepsilon \cap A_{S - R\varepsilon, S + 1 - R\varepsilon}) \leq C\delta. \end{aligned} \quad (104)$$

In the latter estimate, we have used (103) and that, for $k \in \mathcal{K}_\varepsilon$ fixed, we have

$$A_{k\varepsilon, (k+3R)\varepsilon} \cap A_{j\varepsilon, (j+3R)\varepsilon} \neq \emptyset$$

for at most a finite number of indices j independent of ε (clearly for $j > k + 3R$ or $j < k - 3R$ the intersection is empty). Up to replacing δ by $C^{-1}\delta$, (102) and (104) give (101) and this concludes Step 3.

In the next steps we use the following notation. For any $\lambda > 0$ let $X_\varepsilon^\lambda \subset (W_\lambda)_1$ be such that, setting $V_\varepsilon^\lambda = V_\varepsilon(X_\varepsilon^\lambda)$, we have $\chi_{V_\varepsilon^\lambda} \rightarrow \chi_{W_\lambda}$ in $L^1(\mathbb{R}^n)$, and

$$\lim_{\varepsilon \rightarrow 0} F_\varepsilon(X_\varepsilon^\lambda) = \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(X_\varepsilon^\lambda) = \limsup_{\varepsilon \rightarrow 0} F_\varepsilon(X_\varepsilon^\lambda, (W_\lambda)_1) = F_{\text{hom}}(\rho\chi_{W_\lambda}). \quad (105)$$

Step 4.(Identification of V_1) We claim that $V_1 = W_{v_1}$ up to null sets, where W_{v_1} is the Wulff Shape of F_{hom} such that $|W_{v_1}| = v_1$. Assume by contradiction that this were not the case. By the anisotropic isoperimetric inequality we have that

$$0 < \frac{1}{2} (F_{\text{hom}}(\rho\chi_{V_1}) - F_{\text{hom}}(\rho\chi_{W_{v_1}})) =: \eta.$$

Up to translating, we may assume that $\tau_\varepsilon^1 = 0$. Then, there exists $S_0 > 0$ big enough such that for all $S \geq S_0$ we have $W_{v_1} \subset\subset B_S$ and

$$F_{\text{hom}}(\rho\chi_{V_1}, B_S) \geq F_{\text{hom}}(\rho\chi_{V_1}) - \eta.$$

By Theorem 5.14, we have that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(X_\varepsilon, B_S) - F_{\text{hom}}(\rho\chi_{W_{v_1}}, B_S) &\geq F_{\text{hom}}(\rho\chi_{V_1}, B_S) - F_{\text{hom}}(\rho\chi_{W_{v_1}}, B_S) \\ &\geq F_{\text{hom}}(\rho\chi_{V_1}) - \eta - F_{\text{hom}}(\rho\chi_{W_{v_1}}) \geq \eta > 0. \end{aligned} \quad (106)$$

We now construct a competitor \hat{X}_ε for $\varepsilon > 0$ small enough such that $\#X_\varepsilon = n_\varepsilon$ and

$$F_\varepsilon(\hat{X}_\varepsilon) < F_\varepsilon(X_\varepsilon). \quad (107)$$

This contradicts the assumptions of the Lemma on the minimality of X_ε and therefore $V_1 = W_{v_1}$ up to null sets. Let us take $\delta := \delta_\eta = (2C \max_{x \in \mathcal{L}} \#\mathcal{NN}(x))^{-1}\eta$ with C as in (73) and let $S_\delta, S_\varepsilon \in (S_\delta, S_\delta + 1)$ be as in Step 3 with $S_\delta \geq \text{diam } W_{v_1} + 1$. We define

$$\hat{X}_\varepsilon := X_\varepsilon^{v_1} \cup (X_\varepsilon \setminus B_{S_\varepsilon}) \cup Z_\varepsilon, \quad (108)$$

where $\#\hat{X}_\varepsilon = \#X_\varepsilon$. In what follows we assume for simplicity that $Z_\varepsilon = \emptyset$. Otherwise, one choose Z_ε such that its contributions to both energy and volume are negligible as $\varepsilon \rightarrow 0$. First of all note

$$F_\varepsilon(\hat{X}_\varepsilon) = F_\varepsilon(\hat{X}_\varepsilon, B_{S_\varepsilon}) + F_\varepsilon(\hat{X}_\varepsilon, A_{S_\varepsilon+3\varepsilon R}) + F_\varepsilon(\hat{X}_\varepsilon, B_{S_\varepsilon+3\varepsilon R}^c). \quad (109)$$

Then, by noting first that $X_\varepsilon = \hat{X}_\varepsilon$ on $(B_{S_\varepsilon+3\varepsilon R}^c)_{\varepsilon R}$, we have

$$F_\varepsilon(X_\varepsilon, B_{S_\varepsilon+3\varepsilon R}^c) = F_\varepsilon(\hat{X}_\varepsilon, B_{S_\varepsilon+3\varepsilon R}^c). \quad (110)$$

Furthermore,

$$F_\varepsilon(\hat{X}_\varepsilon, B_{S_\varepsilon}) = F_\varepsilon(X_\varepsilon^{v_1}, B_{S_\varepsilon}) = F_\varepsilon(X_\varepsilon^{v_1}). \quad (111)$$

Lastly, due to Lemma 5.9(iii) and the choice of δ , and (101), we have

$$F_\varepsilon(\hat{X}_\varepsilon, A_{S_\varepsilon, S_\varepsilon+3\varepsilon R}) \leq C \max_{x \in \mathcal{L}} \#\mathcal{NN}(x) \varepsilon^{n-1} \#(X_\varepsilon \cap A_{S_\varepsilon, S_\varepsilon+3\varepsilon R}) \leq \eta/2. \quad (112)$$

Comparing this to the energy of X_ε , also noting that $S_\delta \geq \text{diam } W_{v_1} + 1$ and, using (109)–(112), we obtain

$$\begin{aligned} F_\varepsilon(X_\varepsilon) &= F_\varepsilon(X_\varepsilon, B_{S_\varepsilon}) + F_\varepsilon(X_\varepsilon, A_{S_\varepsilon+3\varepsilon R}) + F_\varepsilon(X_\varepsilon, B_{S_\varepsilon+3\varepsilon R}^c) \\ &\geq F_\varepsilon(X_\varepsilon, B_{S_\delta}) + F_\varepsilon(X_\varepsilon, B_{S_\varepsilon+3\varepsilon R}^c) \geq F_\varepsilon(\hat{X}_\varepsilon) - \eta/2 + F_\varepsilon(X_\varepsilon, B_{S_\delta}) - F_\varepsilon(X_\varepsilon^{v_1}) \end{aligned}$$

Therefore, using (105) and (106), we obtain

$$\liminf_{\varepsilon \rightarrow 0} (F_\varepsilon(X_\varepsilon) - F_\varepsilon(\hat{X}_\varepsilon)) \geq \eta/2 > 0.$$

This yields (107) for $\varepsilon > 0$ small enough.

Step 5. ($v_1 = v$) Assume by contradiction that $v_1 < v$. We repeat Step 2 and Step 3 for $X_\varepsilon \setminus (W_{v_1}(\tau_\varepsilon^1))_1$ to find τ_ε^2 such that

$$\mu_\varepsilon(\cdot - \tau_\varepsilon^2) \xrightarrow{*} \rho\chi_{V_2} \text{ in the sense of measures,} \quad (113)$$

where $|V_2| = v_2 > 0$. Note that we can assume $|\tau_\varepsilon^1 - \tau_\varepsilon^2| \rightarrow +\infty$ since this would otherwise contradict $\mu_\varepsilon(\cdot - \tau_\varepsilon^1) \xrightarrow{*} \rho\chi_V$. By Step 4, we observe that $V_2 = W_{v_2}$. Note that $v \mapsto F_{\text{hom}}(W_v)$ is strictly concave in v . In fact there holds

$$F_{\text{hom}}(W_v) = v^{\frac{n-1}{n}} F_{\text{hom}}(W_1). \quad (114)$$

Therefore, given $v_1, v_2 > 0$, for $v = \lambda v_1 + (1 - \lambda)v_2$ for $\lambda \in (0, 1)$, using (114) and the strict concavity of the function $v \mapsto v^{\frac{n-1}{n}}$, we have

$$\begin{aligned} F_{\text{hom}}(W_v) &= F_{\text{hom}}(W_{\lambda v_1 + (1-\lambda)v_2}) = (\lambda v_1 + (1 - \lambda)v_2)^{\frac{n-1}{n}} F_{\text{hom}}(W_1) \\ &> \left(\lambda v_1^{\frac{n-1}{n}} + (1 - \lambda)v_2^{\frac{n-1}{n}} \right) F_{\text{hom}}(W_1) = \lambda F_{\text{hom}}(W_{v_1}) + (1 - \lambda)F_{\text{hom}}(W_{v_2}). \end{aligned}$$

Since $F_{\text{hom}}(W_0) = 0$, this in particular implies that $v \mapsto F_{\text{hom}}(W_v)$ is strictly subadditive, i.e.,

$$F_{\text{hom}}(W_{v_1+v_2}) < F_{\text{hom}}(W_{v_1}) + F_{\text{hom}}(W_{v_2})$$

for all $v_1, v_2 > 0$. Set

$$\eta := F_{\text{hom}}(W_{v_1}) + F_{\text{hom}}(W_{v_2}) - F_{\text{hom}}(W_{v_1+v_2}) > 0.$$

Due to Step 3 applied with $\delta = \delta_\eta := (3C \max_{x \in \mathcal{L}} \#\mathcal{NN}(x))^{-1} \eta$ and Step 4 we have that

$$\mu_\varepsilon(\cdot - \tau_\varepsilon^1) \xrightarrow{*} \rho\chi_{V_1} \quad \text{and} \quad \mu_\varepsilon(\cdot - \tau_\varepsilon^2) \xrightarrow{*} \rho\chi_{V_2}$$

and

$$\#(X_\varepsilon \cap A_{S_\varepsilon, S_\varepsilon + 3\varepsilon R}(\tau_\varepsilon^1)) \leq \delta_\eta \varepsilon^{1-n} \quad \text{and} \quad \#(X_\varepsilon \cap A_{\tilde{S}_\varepsilon, \tilde{S}_\varepsilon + 3\varepsilon R}(\tau_\varepsilon^2)) \leq \delta_\eta \varepsilon^{1-n},$$

where \tilde{S}_ε is associated with τ_ε^2 by following the same procedure to $X_\varepsilon \setminus B_{S_\varepsilon}(\tau_\varepsilon^1)$ as in Step 3. We can therefore define

$$\tilde{X}_\varepsilon = X_\varepsilon^{v_1+v_2} \cup (X_\varepsilon \setminus (B_{S_\varepsilon}(\tau_\varepsilon^1) \cup B_{\tilde{S}_\varepsilon}(\tau_\varepsilon^2))),$$

and without loss of generality (see Step 4) we can directly assume $\#\tilde{X}_\varepsilon = \#X_\varepsilon$. Now the argument follows very much in the spirit of Step 4. We first observe that

$$\begin{aligned} F_\varepsilon(\tilde{X}_\varepsilon) &= F_\varepsilon(\tilde{X}_\varepsilon, B_{S_\varepsilon}(\tau_\varepsilon^1)) + F_\varepsilon(\tilde{X}_\varepsilon, B_{\tilde{S}_\varepsilon}(\tau_\varepsilon^2)) + F_\varepsilon(\tilde{X}_\varepsilon, A_{S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1)) \\ &\quad + F_\varepsilon(\tilde{X}_\varepsilon, A_{S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2)) + F_\varepsilon(\tilde{X}_\varepsilon, (B_{S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1) \cup B_{\tilde{S}_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2))^c). \end{aligned} \quad (115)$$

Then, by noting first that $X_\varepsilon = \tilde{X}_\varepsilon$ on $((B_{S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1) \cup B_{\tilde{S}_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2))^c)_{\varepsilon R}$, we have

$$F_\varepsilon(X_\varepsilon, ((B_{S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1) \cup B_{\tilde{S}_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2))^c)_{\varepsilon R}) = F_\varepsilon(\tilde{X}_\varepsilon, ((B_{S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1) \cup B_{\tilde{S}_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2))^c)_{\varepsilon R}). \quad (116)$$

Furthermore,

$$F_\varepsilon(\tilde{X}_\varepsilon, B_{S_\varepsilon}(\tau_\varepsilon^1)) = F_\varepsilon(X_\varepsilon^{v_1}) \quad \text{and} \quad F_\varepsilon(\tilde{X}_\varepsilon, B_{\tilde{S}_\varepsilon}(\tau_\varepsilon^2)) = F_\varepsilon(X_\varepsilon^{v_2}). \quad (117)$$

Lastly, due to Lemma 5.9(iii) and the choice of δ , we have

$$F_\varepsilon(\tilde{X}_\varepsilon, A_{S_\varepsilon, S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1)) \leq C \max_{x \in \mathcal{L}} \#\mathcal{NN}(x) \varepsilon^{n-1} \#(X_\varepsilon \cap A_{S_\varepsilon, S_\varepsilon+3\varepsilon R}(\tau_\varepsilon^1)) \leq \eta/3 \quad (118)$$

and

$$F_\varepsilon(\tilde{X}_\varepsilon, A_{\tilde{S}_\varepsilon, \tilde{S}_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2)) \leq C \max_{x \in \mathcal{L}} \#\mathcal{NN}(x) \varepsilon^{n-1} \#(X_\varepsilon \cap A_{\tilde{S}_\varepsilon, \tilde{S}_\varepsilon+3\varepsilon R}(\tau_\varepsilon^2)) \leq \eta/3.$$

Comparing this to the energy of X_ε we obtain, using (115)–(118), as in Step 5,

$$\liminf_{\varepsilon \rightarrow 0} (F_\varepsilon(X_\varepsilon) - F_\varepsilon(\tilde{X}_\varepsilon)) \geq \eta/3 > 0.$$

This is a contradiction and therefore $v_1 = v$. Setting $\tau_\varepsilon := \tau_\varepsilon^1$ this concludes the proof. \square

Remark 5.17. We want to observe that Lemma 5.16 can be extended to the setting of [3] in which the functional F_ε also accounts for long-range interactions. In order to adapt the proof to the general case, the annulus $A_{r_\varepsilon, r_\varepsilon+3R\varepsilon}$ must be replaced by $A_{r_\varepsilon, r_\varepsilon+s_\varepsilon}$ where $s_\varepsilon = k_\varepsilon\varepsilon$ (here k_ε is such that $k_\varepsilon \rightarrow +\infty$ and $k_\varepsilon\varepsilon \rightarrow 0$). This choice ensures that $\hat{X}_\varepsilon \cap B_{r_\varepsilon}$ and $\hat{X}_\varepsilon \setminus B_{r_\varepsilon+s_\varepsilon}$ (resp. $X_\varepsilon \cap B_{r_\varepsilon}$ and $X_\varepsilon \setminus B_{r_\varepsilon+s_\varepsilon}$) are sufficiently distant such that the energy contribution that accounts for the interactions crossing the annulus are negligible as $\varepsilon \rightarrow 0$.

ACKNOWLEDGEMENTS

This work was supported by the DFG project FR 4083/1-1 and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2044 -390685587, Mathematics Münster: Dynamics–Geometry–Structure. The work of M. Cicalese was supported by the DFG Collaborative Research Center TRR 109, "Discretization in Geometry and Dynamics". The first and second author gratefully acknowledge the GNAMPA visiting professor program 2022 and the CIRM research in pairs program, Trento 2020, during which parts of the project have been carried out.

REFERENCES

- [1] N.L. ALLINGER. *Molecular structure: understanding steric and electronic effects from molecular mechanics*. John Wiley & Sons (2010).
- [2] R. ALICANDRO, A. BRAIDES, M. CICALESE, M. SOLCI. Discrete Variational Problems with Interfaces, book in preparation.
- [3] R. ALICANDRO, M. CICALESE, M. RUF. Domain formation in magnetic polymer composites: an approach via stochastic homogenization. *Arch. Ration. Mech. Anal.* **218** (2015), 945–984.
- [4] L. AMBROSIO, N. FUSCO, D. PALLARA. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 2000.
- [5] R. ALICANDRO, M.S. GELLI. *Local and non local continuum limits of Ising-type energies for spin systems*. *SIAM J. Math. Anal.* **48** (2016), 895–931.
- [6] Y. AU YEUNG. *Crystalline Order, Surface Energy Densities and Wulff Shapes: Emergence from Atomistic Models*. PhD thesis (2013), Technische Universität München.
- [7] Y. AU YEUNG, G. FRIESECKE, B. SCHMIDT. *Minimizing atomic configurations of short range pair potentials in two dimensions: crystallization in the Wulff shape*. *Calc. Var. Partial Differential Equations* **44** (2012) 81–100.
- [8] X. BLANC, M. LEWIN. *The crystallization conjecture: a review*. *EMS Surv. Math. Sci.* **2** (2015), 255–306.
- [9] A. BRAIDES. *Γ -convergence for Beginners*. Oxford University Press, Oxford 2002.
- [10] A. BRAIDES, L. TRUSKINOVSKY. *Asymptotic expansions by Γ -convergence*. *Contin. Mech. Thermodyn.* **20** (2008) 21–62.
- [11] A. CHAMBOLLE, L. KREUTZ. *Crystallinity of the homogenized surface energy density of periodic lattice systems*. Preprint (2021) ArXiv:2106.08111.
- [12] M. CICALESE, G.P. LEONARDI. *Maximal fluctuations on periodic lattices: an approach via quantitative Wulff inequalities*. *Comm. math. phys.* (2019) 1–14.
- [13] G. DAL MASO. *An introduction to Γ -convergence*. Springer Science & Business Media, **8** (2012).
- [14] E. DAVOLI, P. PIOVANO, U. STEFANELLI, *Wulff shape emergence in graphene*. *Math. Mod. and Meth. in Applied Sciences* **26** (2016) 2277–2310.
- [15] E. DAVOLI, P. PIOVANO, U. STEFANELLI, *Sharp $N^{3/4}$ Law for the Minimizers of the Edge-Isoperimetric Problem on the Triangular Lattice*. *Journal of Nonlinear Science* **27** (2017) 627–660.
- [16] L. DE LUCA, G. FRIESECKE. *Crystallization in two dimensions and a discrete Gauss-Bonnet Theorem*. *J. Nonlinear Sci.* **28** (2017), 69–90.
- [17] De Luca, L., Friesecke, G. *Classification of particle numbers with unique Heitmann-Radin minimizer*, *J. Stat. Phys.* **167** (2017), 1586–1592.
- [18] W. E, D. LI. *On the crystallization of 2D hexagonal lattices*. *Comm. Math. Phys.* **286** (2009), 1099–1140.
- [19] I. FONSECA, S. MÜLLER. *A uniqueness proof for the Wulff theorem*. *Proc. Roy. Soc. Edinburgh Sect.* **119** (1991) 125–136.

- [20] L. C. FLATLEY, F. THEIL. *Face-centered cubic crystallization of atomistic configurations*. Arch. Ration. Mech. Anal. **218** (2015), 363–416.
- [21] M. FRIEDRICH, L. KREUTZ. *Finite crystallization and Wulff shape emergence for ionic compounds in the square lattice*. Nonlinearity **33** (2020), 1240–1296.
- [22] M. FRIEDRICH, L. KREUTZ. *Crystallization in the hexagonal lattice for ionic dimers*. Math. Models Methods Appl. Sci. **29** (2019), 1853–1900.
- [23] M. FRIEDRICH, L. KREUTZ, B. SCHMIDT. *Emergence of rigid Polycrystals from atomistic Systems with Heitmann-Radin sticky disk energy*. Arch. Ration. Mech. Anal. **240** (2021), 627–698.
- [24] G. FRIESECKE, F. THEIL. *Molecular geometry optimization, models*. In the Encyclopedia of Applied and Computational Mathematics, B. Engquist (Ed.), Springer, 2015.
- [25] M. GARCÍA-BRAVO, T. RAJALA. *Strong BV-extension and $W^{1,1}$ -extension domains*. arXiv preprint arXiv:2105.05467 (2021).
- [26] C. S. GARDNER, C. RADIN. *The infinite-volume ground state of the Lennard-Jones potential*. J. Stat. Phys. **20** (1979), 719–724.
- [27] T.C. HALES. *A proof of the Kepler conjecture*. Ann. Math. **162** (2005), 1065–1185.
- [28] H. HARBORTH. *Lösung zu Problem 664 a*. Elem. Math. **29** (1974), 14–15.
- [29] L.H. HARPER. *Global methods for combinatorial isoperimetric problems*. Cambridge University Press **90** (2004).
- [30] R. HEITMANN, C. RADIN. *The ground state for sticky disks*. J. Stat. Phys. **22** (1980) 281–287.
- [31] E. G. LEWARS. *Computational Chemistry*. 2nd edition, Springer, 2011.
- [32] E. MAININI, P. PIOVANO, B. SCHMIDT, U. STEFANELLI. *$N^{3/4}$ Law in the Cubic Lattice*. Journal of statistical physics **176** (2019), 1480–1499.
- [33] E. MAININI, P. PIOVANO, U. STEFANELLI. *Finite crystallization in the square lattice*. Nonlinearity **27** (2014), 717–737.
- [34] E. MAININI, B. SCHMIDT. *Maximal fluctuations around the Wulff shape for edge-isoperimetric sets in \mathbb{Z}^d : a sharp scaling law..* Communications in Mathematical Physics **380** (2020), 947–971.
- [35] E. MAININI, U. STEFANELLI. *Crystallization in carbon nanostructures*. Comm. Math. Phys. **328** (2014), 545–571.
- [36] C. RADIN. *The ground state for soft disks*. J. Stat. Phys. **26** (1981), 365–373.
- [37] K. SCHÜTTE, B.L. VAN DER WAERDEN. *Das Problem der dreizehn Kugeln*. Math. Ann. **125** (1952), 324–334.
- [38] B. SCHMIDT. *Ground states of the 2D sticky disc model: fine properties and $N^{3/4}$ law for the deviation from the asymptotic Wulff shape*. J. Stat. Phys. **153(4)** (2013), 727–738.
- [39] F. THEIL. *A proof of crystallization in two dimensions*. Comm. Math. Phys. **262** (2006), 209–236.
- [40] G. WULFF. *Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Kristallflächen*. Z. Krist. **34** (1901), 449–530.

(Marco Cicalese) ZENTRUM MATHEMATIK - M7, TECHNISCHE UNIVERSITÄT MÜNCHEN, GARCHING, GERMANY
E-mail address: cicalese@ma.tum.de

(Leonard Kreutz) APPLIED MATHEMATICS MÜNSTER, WWU MÜNSTER, 48149 MÜNSTER, GERMANY
E-mail address: lkreutz@uni-muenster.de

(Gian Paolo Leonardi) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, 38050 POVO (TN), ITALY
E-mail address: gianpaolo.leonardi@unitn.it