

Nonlinear approximation of 3D smectic liquid crystals: sharp lower bound and compactness

Michael Novack*

Xiaodong Yan[†]

May 11, 2022

Abstract

We consider the 3D smectic energy

$$\mathcal{E}_\varepsilon(u) = \frac{1}{2} \int_\Omega \frac{1}{\varepsilon} \left(\partial_z u - \frac{(\partial_x u)^2 + (\partial_y u)^2}{2} \right)^2 + \varepsilon (\partial_x^2 u + \partial_y^2 u)^2 dx dy dz.$$

The model contains as a special case the well-known 2D Aviles-Giga model. We prove a sharp lower bound on \mathcal{E}_ε as $\varepsilon \rightarrow 0$ by introducing 3D analogues of the Jin-Kohn entropies [31]. The sharp bound corresponds to an equipartition of energy between the bending and compression strains and was previously demonstrated in the physics literature only when the approximate Gaussian curvature of each smectic layer vanishes. Also, for $\varepsilon_n \rightarrow 0$ and an energy-bounded sequence $\{u_n\}$ with $\|\nabla u_n\|_{L^p(\Omega)}, \|\nabla u_n\|_{L^2(\partial\Omega)} \leq C$ for some $p > 6$, we obtain compactness of ∇u_n in L^2 assuming that $\Delta_{xy} u_n$ has constant sign for each n .

1 Introduction

In this article, we analyze the energies

$$\mathcal{E}_\varepsilon(u) = \frac{1}{2} \int_\Omega \left[\frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right)^2 + \varepsilon (\Delta_\perp u)^2 \right] dx, \quad (1.1)$$

which represent the free energy of a smectic-A liquid crystal; see Section 2.1 for a discussion of the relevant physics literature regarding smectics. Here $\Omega \subset \mathbb{R}^3$ is a bounded domain with Lipschitz boundary, and $\mathbf{x} = (x, y, z)$. The subscript “ \perp ” denotes the restriction to the x, y variables of a differential operator or the projection from \mathbb{R}^3 to \mathbb{R}^2 , so

$$\nabla_\perp u = (\partial_x u, \partial_y u) \quad \text{and} \quad \Delta_\perp u = \partial_x^2 u + \partial_y^2 u.$$

and if $m = (m_1, m_2, m_3) \in \mathbb{R}^3$,

$$m_\perp = m_1 \hat{x} + m_2 \hat{y} \quad \text{and} \quad m = (m_\perp, m_3) \in \mathbb{R}^3.$$

Our main interest is the asymptotic behavior of energies (1.1) as $\varepsilon \rightarrow 0$, which corresponds to the regime in which the intrinsic length scale ε , cf. (2.5), is vanishingly small compared to Ω .

We prove the following main results when $\varepsilon \rightarrow 0$:

*Department of Mathematics, The University of Texas at Austin, Austin, TX, USA

[†]Department of Mathematics, The University of Connecticut, Storrs, CT, USA

- a lower bound, sharp when $\nabla u \in (BV \cap L^\infty)(\Omega; \mathbb{R}^3)$, on \mathcal{E}_ε when $\varepsilon \rightarrow 0$ (Theorem 3.6, Theorem 4.1), and
- a compactness theorem for the gradients of a sequence with bounded energies (Theorem 5.1) satisfying some additional technical assumptions.

These results generalize the authors' previous work [47] on the 2D model

$$\mathcal{J}_\varepsilon(u) = \frac{1}{2} \int_\Omega \left[\frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} (\partial_x u)^2 \right)^2 + \varepsilon (\partial_x^2 u)^2 \right] dx dz, \quad (1.2)$$

to the 3D energies (1.1). For $\varepsilon_n \rightarrow 0$ and a sequence $\{u_n\}$ with bounded energies $\mathcal{J}_{\varepsilon_n}(u_n)$, we proved compactness of ∇u_n in L^q for $1 \leq q < p$ under the additional assumption $\|\nabla u_n\|_{L^p} \leq C$ for some $p > 6$. Moreover, we obtained a lower bound on \mathcal{J}_ε and constructed a matching upper bound using on a 1D ansatz.

The sharp lower bound for the 3D energies, which was not previously shown in the physics literature, relies on a calibration argument which briefly works as follows. Letting

$$\Sigma(\nabla u) = (\Sigma_1, \Sigma_2, \Sigma_3) \quad (1.3)$$

where

$$\begin{aligned} \Sigma_1 &= \partial_z u \partial_x u - \frac{1}{2} \partial_x u (\partial_y u)^2 - \frac{1}{6} (\partial_x u)^3, \\ \Sigma_2 &= -\partial_x u \partial_y u + \frac{1}{2} \partial_y u (\partial_x u)^2 + \frac{1}{6} (\partial_y u)^3, \\ \Sigma_3 &= \frac{1}{2} (\partial_y u)^2 - \frac{1}{2} (\partial_x u)^2, \end{aligned}$$

direct calculation shows that for $u \in H^2(\Omega)$,

$$\operatorname{div} \Sigma(\nabla u) = \left(\partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right) (\partial_x^2 u - \partial_y^2 u). \quad (1.4)$$

Thus by the arithmetic mean-geometric mean inequality and the divergence theorem, cf. (2.24), $\mathcal{E}_\varepsilon(u)$ can be bounded below by

$$\mathcal{E}_\varepsilon(u) \geq \int_\Omega \operatorname{div} \Sigma(\nabla u) \, d\mathbf{x} - \varepsilon \|\nabla_\perp u\|_{H^{1/2}(\partial\Omega)}^2, \quad (1.5)$$

with approximate equality when

$$\frac{1}{\varepsilon} \int_\Omega \left(\partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right)^2 \approx \varepsilon \int_\Omega (\Delta_\perp u)^2. \quad (1.6)$$

By the rotational symmetry in the xy -plane of the energies (1.1), the same calculation holds for the rotations $\Sigma_{\xi, \eta}$ of Σ (see (3.1)) obtained by replacing $\{\hat{x}, \hat{y}\}$ with another orthonormal basis $\{\xi, \eta\}$ of \mathbb{R}^2 . Thus for a sequence $\varepsilon_n \rightarrow 0$ and $\{u_n\}$ converging to a limiting function u in a suitable space, we may bound $\liminf \mathcal{E}_{\varepsilon_n}(u_n)$ from below by taking the supremum of the divergences of $\Sigma_{\xi, \eta}$ over all $\{\xi, \eta\}$ as in [3, 6] for the Aviles-Giga problem. In fact, the energy (1.1) contains the 2D Aviles-Giga energy as a special case, which we explain in Section 2.2. If $\nabla u \in (BV \cap L^\infty)(\Omega; \mathbb{R}^3)$, this lower bound is given by

$$\int_{J_{\nabla u}} \frac{|\nabla_\perp^+ u - \nabla_\perp^- u|^4}{12 |\nabla^+ u - \nabla^- u|} \, d\mathcal{H}^2, \quad (1.7)$$

where $J_{\nabla u}$ is the jump set in the sense of BV . For the matching upper bound when $\nabla u \in (BV \cap L^\infty)(\Omega; \mathbb{R}^3)$, (1.5) implies the lower bound is optimal for $\mathcal{E}_\varepsilon(u)$ if (1.6) holds. When Ω is a cube, we show this is possible by using a 1D ansatz in which ∇u varies transverse to the defect set. The ansatz is chosen so that (1.6) holds. The upper bound for general Ω can then be shown by appealing to a result of Poliakovsky [49].

Physically, the 1D ansatz can be interpreted as equating the compression strain $(\partial_z u - \frac{1}{2} |\nabla_\perp u|^2)^2$ with the bending strain $\varepsilon^2 (\Delta_\perp u)^2$; see section Section 2.1 for physical background of model (1.1). Thus our analysis shows that the frustration coming from the competition between the compression and bending terms is resolved by an equipartition of energy between the two. Moreover, unlike many other problems from materials science where microstructure develops [35], microstructure does not appear in this smectics model. We remark that the same has been observed in the 2D problem [47].

The compactness of ∇u_n in L^2 is proved under the additional assumptions $\|\nabla u_n\|_{L^p(\Omega)} \leq C$ for some $p > 6$ and $\|\nabla u_n\|_{L^2(\partial\Omega)} \leq C$. We emphasize that these are physically justifiable, as the model (1.2) is only valid in the small strain regime [10, 55], cf. Section 2.1. Our compactness proof relies on a compensated compactness argument based on the work of Tartar [60, 61, 62] and Murat [44, 45]. The main challenge is to find the suitable entropies to apply Tartar and Murat's div-curl lemma. Assuming further that $\Delta_\perp u_n \geq 0$ a.e. in Ω , we show that $\text{curl } E_n$ and $\text{div } B_n$ are compact in $H^{-1}(\Omega)$, where

$$E_n = \left(\nabla_\perp u_n, \frac{1}{2} |\nabla_\perp u_n|^2 \right) \text{ and } B_n = \left(-\frac{\nabla_\perp u_n}{2} |\nabla_\perp u_n|^2, \frac{1}{2} |\nabla_\perp u_n|^2 \right).$$

Thus (E_n, B_n) satisfy the assumptions of the div-curl lemma. Applying the div-curl lemma to $E_n \cdot B_n$ yields strong convergence of $\nabla_\perp u_n$ in L^2 , and compactness of ∇u_n in L^2 follows from the fact $\partial_z u_n - \frac{1}{2} |\nabla_\perp u_n|^2 \rightarrow 0$ in L^2 .

The paper is organized as follows. Section 2 recalls the physical background of our model (1.1), summarizes the pertinent mathematical literature, and presents the main calculation behind the lower bound in a simplified setting. Also included in Section 2 are some preliminaries on functions of bounded variation. Section 3 is devoted to the lower bound. In Section 4 we construct a sequence which matches the lower bound from Section 3 when $\varepsilon \rightarrow 0$, and in Section 5 we prove compactness.

2 Background and Preliminaries

2.1 Physical background: Smectic A liquid crystals

Smectic liquid crystals are formed by elongated molecules that are aligned and arranged in fluid-like layers. They are a remarkable example of a geometrically frustrated, multi-layer, soft-matter system. Ground states of smectic liquid crystals are characterized by flat, equally spaced, parallel layers. Due to spontaneously broken translational and rotational symmetry, singularities form in regions where the smectic order breaks down. When defects are present, the layers must bend and the resulting curvature is, in general, incompatible with equal spacing between them. The subtle interplay between the geometry of the layers and equal spacing imposes theoretical complications, and understanding the layer structure of a smectic liquid crystal is a challenging task.

Smectics can be represented by the density modulation $\Delta\rho \propto \cos\left[\frac{2\pi}{a}\phi(\mathbf{x})\right]$, where $\mathbf{x} = (x, y, z) \in \mathbb{R}^3$, a is the layer spacing, and ϕ is the phase field of the order parameter [11, 20]. The peaks of the density wave where $\phi(\mathbf{x}) \in a\mathbb{Z}$ correspond to the smectic layers. We are interested in the smectic A phase, in which the nematic director coincides with the normal to the smectic layers $\mathbf{N} = \frac{\nabla\phi}{|\nabla\phi|}$. In terms of ϕ , the free energy of a smectic liquid crystal on a region Ω [56] is the sum of

the compression and bending energies

$$F = \frac{1}{2} \int_{\Omega} B \left[(1 - |\nabla\phi|)^2 + K_1 \left(\nabla \cdot \frac{\nabla\phi}{|\nabla\phi|} \right)^2 \right] d\mathbf{x}, \quad (2.1)$$

where B is the compression modulus and K_1 the bend modulus. The constant $\varepsilon = \sqrt{\frac{K_1}{B}}$ is the penetration length. In the presence of boundaries, there is also the saddle-splay term

$$F_K = \tilde{K} \int_{\Omega} \nabla \cdot [(\nabla \cdot \mathbf{N}) \mathbf{N} - (\mathbf{N} \cdot \nabla) \mathbf{N}] d\mathbf{x}.$$

The contribution of this term depends only on the boundary conditions and is often excluded from the energy. For configurations with topological defects, this term can contribute to the core energy of a defect.

The global minimizer of (2.1) is the zero-energy state $\phi(\mathbf{x}) = \mathbf{n} \cdot \mathbf{x} + \phi_0$ with $\mathbf{n} \in \mathbb{S}^2$ and $\phi_0 \in \mathbb{R}$. However, rarely do both terms in the free energy vanish. To understand the frustration of the problem in general, the following example of a smooth surface $\mathbf{x}_0 = \mathbf{x}_0(u, v)$ and its parallel surfaces $\mathbf{x}_n(u, v) := \mathbf{x}_0(u, v) + n\mathbf{a}\mathbf{N}(u, v)$, $n \in \mathbb{Z}$ is illuminating [33]. Let H_n and K_n denote the mean and Gaussian curvatures of \mathbf{x}_n . A standard calculation [19, Sec. 3-5 Exercise 11] yields the formulas

$$H_n = \frac{H_0 - naK_0}{1 - 2naH_0 + n^2a^2K_0}, \quad K_n = \frac{K_0}{1 - 2naH_0 + n^2a^2K_0}.$$

Since mean curvature can be expressed in terms of the surface normal by $H = -\frac{1}{2}\nabla \cdot \mathbf{N}$, the bending term is proportional to H^2 . Therefore, for evenly spaced layers, the only way for the bending to vanish (so $H_n = 0$) for all layers is for the Gaussian curvature K_0 of \mathbf{x}_0 to be zero, which in turn implies $K_n = 0$, so that the product of the principal curvatures is zero. On the other hand, $H_n = 0$ implies that the principal curvatures sum to 0 as well, so that they both vanish everywhere. Thus unless all the layers are flat, vanishing curvature is incompatible with the uniform layer spacing. The interaction between the layer spacing, the Gaussian curvature, and the mean curvature presents a major obstacle to finding minimal configurations for the energy (2.1). Throughout the physics literature there are numerous works on the search for exact or approximate solutions of deformations in smectics [8, 10, 17, 18, 30, 32, 33, 42, 54, 55, 56, 57].

In the study of smectic layers, it is typical to consider the deviation u from a fixed ground state $\bar{\phi}$, so $\phi = \bar{\phi} - u$. If we fix $\bar{\phi}(\mathbf{x}) = z$, then $u(\mathbf{x}) = z - \phi(\mathbf{x})$. Expressing the compression strain in powers of $\nabla u = \hat{z} - \nabla\phi$ and setting $\nabla_{\perp} = \partial_x \hat{x} + \partial_y \hat{y}$, $\Delta_{\perp} = \partial_x^2 + \partial_y^2$, we can expand the compression as

$$1 - |\nabla\phi| \approx \partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 + \mathcal{O}(u^3), \quad (2.2)$$

and bending strain as

$$\nabla \cdot \frac{\nabla\phi}{|\nabla\phi|} \approx -\Delta_{\perp} u + \mathcal{O}(u^3). \quad (2.3)$$

Keeping only the linear terms in the expansions (2.2) and (2.3) in the limit of small elastic strains $|\nabla u| \ll 1$ results in a linear theory [20, 34] of (2.1) which has been widely used in the study of strain fields and energetics of dislocations in smectics-A. On the other hand, the well-known example of the dilatative Helfrich-Hurault effect [23, 24], in which the layers wrinkle upon stretching, indicates that nonlinear effects can be important even for small strains. Observe that the linear model is only valid when $\frac{|\nabla_{\perp} u|^2}{\partial_z u} \ll 1$, and the nonlinear term is not negligible when $\partial_z u \sim |\nabla_{\perp} u|^2$. The first example of strain field taking into account the nonlinear effect was constructed by Brener

and Marchenko [10]. They considered the strain field for a single edge dislocation in the regime $\partial_z u \sim (\partial_x u)^2 \ll 1$ and found an exact solution to the Euler-Lagrange equation for the 2D nonlinear approximation of (2.1)

$$F = \frac{B}{2} \int_{\Omega} \left[\left(\partial_z u - \frac{1}{2} (\partial_x u)^2 \right)^2 + \varepsilon^2 (\partial_x^2 u)^2 \right] dx dz, \quad (2.4)$$

where

$$\varepsilon = \sqrt{K_1/B}. \quad (2.5)$$

Their construction deviates markedly from the strain field predicted by linear theory even away from the defects where the elastic strain and curvature are small. The solution was verified experimentally in a cholesteric finger texture by Ishikawa and Lavrentovich [30] and by Smalyukh and Lavrentovich [58] using confocal microscopy.

Brener and Marchenko found their solution by solving the fourth order Euler-Lagrange equation of (2.4) directly. Later, Santangelo and Kamien [55] approached the problem from a different perspective and discovered a large class of exact minima for the nonlinear approximations of (2.1). They studied the 3D version of (2.4)

$$F = \frac{B}{2} \int_{\Omega} \left[\left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right)^2 + \varepsilon^2 (\Delta_{\perp} u)^2 \right] d\mathbf{x}. \quad (2.6)$$

Completing the square in (2.6) yields

$$\begin{aligned} \mathcal{E}_{\varepsilon}(u) &= \frac{1}{2} \int_{\Omega} \left[\frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right)^2 + \varepsilon (\Delta_{\perp} u)^2 \right] d\mathbf{x} \\ &= \frac{1}{2} \int_{\Omega} \frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \mp \varepsilon \Delta_{\perp} u \right)^2 d\mathbf{x} \pm \int_{\Omega} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right) \Delta_{\perp} u d\mathbf{x}. \end{aligned} \quad (2.7)$$

Observe that

$$\partial_z u \Delta_{\perp} u = \nabla_{\perp} \cdot (\partial_z u \nabla_{\perp} u) - \frac{1}{2} \partial_z (|\nabla_{\perp} u|^2) \quad (2.8)$$

and

$$3 |\nabla_{\perp} u|^2 \Delta_{\perp} u = -4 \bar{K} u + \nabla_{\perp} \cdot (\nabla_{\perp} u |\nabla_{\perp} u|^2) + 2 \nabla_{\perp} \cdot \left(u \Delta_{\perp} u \nabla_{\perp} u - \frac{1}{2} u \nabla_{\perp} |\nabla_{\perp} u|^2 \right), \quad (2.9)$$

where $\bar{K} = \partial_x^2 u \partial_y^2 u - (\partial_{xy} u)^2$ is the lowest order approximation of the Gaussian curvature. By the Bochner formula $\frac{1}{2} \Delta |\nabla u|^2 = \nabla u \cdot \Delta(\nabla u) + |\nabla^2 u|^2$ from flat geometry or by direct calculation, \bar{K} can also be expressed as

$$\bar{K} = \frac{1}{2} \nabla_{\perp} \cdot \left(\nabla_{\perp} u \Delta_{\perp} u - \frac{1}{2} \nabla_{\perp} |\nabla_{\perp} u|^2 \right), \quad (2.10)$$

which is the form found for example in [55]. Substituting (2.8) and (2.9) back into (2.7) yields

$$\begin{aligned} \mathcal{E}_{\varepsilon}(u) &= \frac{1}{2} \int_{\Omega} \frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \mp \varepsilon \Delta_{\perp} u \right)^2 d\mathbf{x} \pm \frac{2}{3} \int_{\Omega} \bar{K} u d\mathbf{x} \\ &\quad \pm \int_{\Omega} \operatorname{div} \Xi(u) d\mathbf{x}, \end{aligned} \quad (2.11)$$

where

$$\Xi(u) = \left[\left(\partial_z u - \frac{1}{6} |\nabla_\perp u|^2 - \frac{1}{3} u \Delta_\perp u \right) \nabla_\perp u + \frac{1}{6} u \nabla_\perp |\nabla_\perp u|^2, -\frac{1}{2} |\nabla_\perp u|^2 \right]. \quad (2.12)$$

A direct conclusion from this decomposition is that the free energy of deformations with $\bar{K} = 0$ is always bounded below by the contributions from the boundary integrals involving $\Xi(u)$, and the minimum is achieved when

$$\partial_z u - \frac{1}{2} |\nabla_\perp u|^2 = \pm \varepsilon \Delta_\perp u. \quad (2.13)$$

The nonlinear differential equation (2.13) is called the BPS equation, with solutions referred to as BPS solutions. This equation is of reduced order compared to the Euler-Lagrange equation of the free energy (2.6). As observed in [55], (2.13) has a simple but important physical interpretation: equating the bending and compression energies so as to minimize their sum and alleviate the geometric frustration described earlier. This type of technique, called the BPS decomposition, was introduced by Bogomol'nyi [9], Prasad and Sommerfield [51] in the study of field configurations of magnetic monopoles and solitons in field theory. BPS-type decompositions have also been utilized in the analysis of thermal fluctuations in 2D smectics [22] and shape changes in vesicles [7].

When u is a function of z and x only, so that $\bar{K} = 0$, the BPS equation simplifies to

$$\partial_z u - \frac{1}{2} (\partial_x u)^2 = \pm \varepsilon \partial_x^2 u. \quad (2.14)$$

Through the Hopf-Cole transformation $S_\pm = \exp[\pm u(x, \pm z)/(2\varepsilon)]$, (2.14) becomes the diffusion equation

$$\partial_z S_\pm = \varepsilon \partial_x^2 S_\pm. \quad (2.15)$$

Solving (2.15) with the boundary conditions $S_\pm \rightarrow 1$ as $x \rightarrow -\infty$ and $S_\pm \rightarrow e^{\pm \frac{b}{4\varepsilon}}$ as $x \rightarrow +\infty$, where b is the Burgers vector of the dislocation, yields

$$S_\pm = 1 + \left(e^{\pm \frac{b}{4\varepsilon}} - 1 \right) \pi^{-1/2} \int_{-\infty}^{\frac{x}{2\sqrt{\varepsilon z}}} e^{-t^2} dt.$$

After inverting the Hopf-Cole transformation, this solution recovers the edge dislocation deformation calculated in [10].

When \bar{K} is small, the BPS solutions are energetically preferable compared to solutions from the linear theory [55]. Santangelo and Kamien [56] generalized this idea to the full energy (2.1) and established a specific set of minima of (2.1) when the $\bar{K} = 0$. Their arguments demonstrated that the layer deformation in the partially nonlinear theory [10, 55] is near the profile from the full energy.

Given the rigidity of the assumption that the Gaussian curvature of the smectic layers is zero, one might conjecture that BPS solutions are no longer (approximate) minimizers upon relaxing that assumption. Interestingly however, this is not the case. Indeed, we quote from [55], where some numerical simulations were done to investigate this issue. ‘‘Further study is needed to understand the precise role of the $u\bar{K}$ in the failure of the BPS configurations to minimize the energy. It is often the case that ‘near-BPS’ solutions are remarkably good approximants and it appears to be true here as well.’’ We utilize techniques drawn from the mathematical literature for singular perturbation problems to obtain a sharp lower bound for (2.6) while only assuming that $\int \varepsilon \bar{K}$ is small, providing an explanation for this phenomenon. Our analysis is inspired by the simple fact that the 3D smectic energy (2.6) is a generalization of the well-studied 2D Aviles-Giga functional, which, nevertheless, has not been observed previously in the literature to the best of our knowledge.

2.2 Mathematical Background: The Aviles-Giga energy

To illustrate the link between 3D smectics and Aviles-Giga functional, fix $\Omega \subset \mathbb{R}^2$ and consider the smectic energy (2.6) on the three dimensional cylinder $\Omega \times (0, 1)$ subject to the constraint $\partial_z u = 1/2$. With the dependence of ∇u on z eliminated, the energy (2.6) becomes

$$\frac{B}{2} \int_{\Omega} \left[\frac{1}{4} \left(1 - |\nabla_{\perp} u|^2 \right)^2 + \varepsilon^2 (\Delta_{\perp} u)^2 \right] dx dy. \quad (2.16)$$

Ignoring the harmless factor of $1/4$ on the first term, this is the 2D instance of the Aviles-Giga energy, which we now recall.

Aviles and Giga [5] formulated the energy

$$\mathcal{F}_{\varepsilon} = \int_{\Omega} \left[\frac{1}{\varepsilon} (|\nabla u|^2 - 1)^2 + \varepsilon (\Delta u)^2 \right] d\mathbf{x}, \quad (2.17)$$

known as the Aviles-Giga functional, as a model for smectic liquid crystals. Here $\Omega \subset \mathbb{R}^n$ is a bounded domain. When ε approaches zero, Aviles and Giga conjectured that the optimal transition layers are one dimensional and $\mathcal{F}_{\varepsilon}$ converges (in the sense of Γ -convergence) to the limiting energy

$$\mathcal{F}_0 = 1/3 \int_{J_{\nabla u}} |\nabla u|^3 d\mathcal{H}^{n-1}.$$

Here the limiting function u satisfies eikonal equation $|\nabla u| = 1$ a.e., $J_{\nabla u}$ is the defect set, cf. Definition 2.3, and $[\nabla u]$ is the jump in ∇u across $J_{\nabla u}$.

The Aviles-Giga functional has been extensively studied in the case $n = 2$. After extracting a boundary term, (2.17) is equal to

$$\tilde{\mathcal{F}}_{\varepsilon} = \int_{\Omega} \left[\frac{1}{\varepsilon} (|\nabla u|^2 - 1)^2 + \varepsilon |\nabla^2 u|^2 \right] dx dy. \quad (2.18)$$

Jin and Kohn [31] noticed that the divergence of the ‘‘Jin-Kohn entropy’’

$$\left(\partial_x u \left(1 - (\partial_y u)^2 - \frac{1}{3} (\partial_x u)^2 \right), -\partial_y u \left(1 - (\partial_x u)^2 - \frac{1}{3} (\partial_y u)^2 \right) \right), \quad (2.19)$$

calculated directly as $(1 - |\nabla u|^2)(\partial_x^2 u - \partial_y^2 u)$, bounds $\tilde{\mathcal{F}}_{\varepsilon}$ from below and the lower bound is asymptotically optimal if

$$|1 - |\nabla u_{\varepsilon}|^2| \approx |\partial_x^2 u_{\varepsilon} - \partial_y^2 u_{\varepsilon}| \quad \text{and} \quad \varepsilon \int_{\Omega} (\partial_x^2 u_{\varepsilon} \partial_y^2 u_{\varepsilon} - (\partial_{xy} u_{\varepsilon})^2) dx dy \approx 0.$$

For the unit square and boundary conditions $u = 0$, $\frac{\partial u}{\partial n} = -1$, they proved that the lower bound can be achieved by the ‘‘1D’’ ansatz $u_{\varepsilon} = ax + f_{\varepsilon}(y)$ when the associated defect set of the limiting map $\lim_{\varepsilon \rightarrow 0} u_{\varepsilon}$ is parallel to the x axis, corroborating Aviles-Giga’s conjecture regarding the one-dimensionality of the transition region. Recently, Ignat and Monteil [29] proved that any minimizer of (2.18) on an infinite strip is one-dimensional. By considering the supremum of the divergences of all rotated versions of Σu , Aviles and Giga [6] derived a limiting functional $J : W^{1,3}(\Omega) \rightarrow [0, \infty)$ which is lower semicontinuous with respect to strong topology in $W^{1,3}(\Omega)$ and coincides with \mathcal{F}_0 for any u satisfying the eikonal equation with $\nabla u \in BV(\Omega)$. Moreover,

$$J(u) \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{F}}_{\varepsilon_n}(u_n)$$

for any sequence u_n converging strongly to u in $W^{1,3}(\Omega)$. A construction which achieves conjectured Γ -limit when $\nabla u \in BV(\Omega)$ was provided in [12, 49]. For the Γ -convergence theory, another important question is the compactness of sequences with bounded energy when ε goes to zero. Such compactness results in two dimensions have been proved by two different groups [3, 16] using different approaches. For the Aviles-Giga functional in dimensions three or higher, the state of art is less clear. De Lellis [13] constructed a counterexample, showing that \mathcal{F}_0 is not the limiting energy for \mathcal{F}_ε and 1D ansatz is not optimal. The compactness and form of the limiting energy, however, are still open. The Aviles-Giga model and related topics such as the eikonal equation and other line-energy models have continued to be active areas of research in the past two decades; see [1, 15, 21, 25, 26, 27, 28, 36, 14, 37, 38, 39, 40, 41, 52, 53] and the references therein.

2.3 Heuristic Proof of the Lower Bound

The admissible class for (1.1) is $H^2(\Omega)$. We recall some trace properties for this space when $\partial\Omega$ is Lipschitz that are useful for the lower bound. First, since $u \in H^2$, it follows that $\nabla u|_{\partial\Omega} \in H^{1/2}(\partial\Omega; \mathbb{R}^3)$. Furthermore, momentarily replacing $\partial_x, \partial_y,$ and ∂_z by $\partial_1, \partial_2,$ and ∂_3 , the tangential derivative operators

$$\partial_{\tau_{jk}} = \nu_j \partial_k - \nu_k \partial_j : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega) = (H^{1/2}(\partial\Omega))^* \quad (2.20)$$

are well-defined, linear, and bounded for any $1 \leq j, k, l \leq 3$ [43, Proposition 2.1]. In particular, denoting by \langle, \rangle the pairing between $H^{-1/2}(\partial\Omega)$ and $H^{1/2}(\partial\Omega)$, the quantities

$$\langle \partial_m u, \partial_{\tau_{jk}}(\partial_l u) \rangle \quad (2.21)$$

are well-defined for $u \in H^2(\Omega)$ and $1 \leq j, k, l, m \leq 3$.

We recall the map

$$\left(\partial_z u \partial_x u - \frac{1}{2} \partial_x u (\partial_y u)^2 - \frac{1}{6} (\partial_x u)^3, -\partial_x u \partial_y u + \frac{1}{2} \partial_y u (\partial_x u)^2 + \frac{1}{6} (\partial_y u)^3, \frac{1}{2} (\partial_y u)^2 - \frac{1}{2} (\partial_x u)^2 \right)$$

from (1.3). We point out that the first two components of $\Sigma(\nabla u)$ are one half times the Jin-Kohn entropy for the Aviles-Giga energy when $\partial_z u = \frac{1}{2}$. Direct calculation shows that for $u \in H^2(\Omega)$,

$$\begin{aligned} \operatorname{div} \Sigma(\nabla u) &= \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right) (\partial_x^2 u - \partial_y^2 u) \\ &\leq \frac{1}{2\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right)^2 + \frac{\varepsilon}{2} (\partial_x^2 u - \partial_y^2 u)^2. \end{aligned} \quad (2.22)$$

To handle the fact that the second term differs from $(\Delta_{\perp} u)^2$, we recall $\bar{K} = \partial_x^2 u \partial_y^2 u - (\partial_{xy} u)^2$, so that

$$\int_{\Omega} [(\Delta_{\perp} u)^2 - (\partial_x^2 u - \partial_y^2 u)^2 - 4\bar{K}] \, d\mathbf{x} = \int_{\Omega} 4(\partial_{xy} u)^2 \, d\mathbf{x} \geq 0.$$

Combining this with (2.22), we arrive at

$$\int_{\Omega} \operatorname{div} \Sigma(\nabla u) \, d\mathbf{x} + \varepsilon \int_{\Omega} 2\bar{K} \, d\mathbf{x} \leq \mathcal{E}_{\varepsilon}(u). \quad (2.23)$$

Notice that $\overline{K} = \operatorname{div}_\perp(\partial_x u \partial_y^2 u, -\partial_x u \partial_{xy} u)$, so that

$$\varepsilon \int_\Omega 2\overline{K} \, d\mathbf{x} = 2\varepsilon \int_{\partial\Omega} \partial_x u \partial_{\tau_{12}} \partial_y u \, d\mathcal{H}^2 \quad (2.24)$$

and both terms on the left hand side of (2.23) depend only on boundary values. By the continuity of the tangential derivative operators from $H^{1/2}(\partial\Omega)$ to $H^{-1/2}(\partial\Omega)$, we can estimate

$$\left| \varepsilon \int_\Omega 2\overline{K} \, d\mathbf{x} \right| \leq 2\varepsilon \|\partial_x u\|_{H^{1/2}(\partial\Omega)} \|\partial_{\tau_{12}} \partial_y u\|_{H^{-1/2}(\partial\Omega)} \leq C(\Omega) \varepsilon \|\nabla_\perp u\|_{H^{1/2}(\partial\Omega)}^2. \quad (2.25)$$

The general lower bound as $\varepsilon \rightarrow 0$ is then derived by taking supremum over all the rotations of Σ as in (3.1). We also refer the reader to the end of Section 4 for a discussion of the implications of this analysis on the BPS solutions.

Remark 2.1. *There are few 3D examples where an explicit calibration can be found and, to our knowledge, there is no systematic approach to find such a calibration for an arbitrary energy. Our choice of the calibration (1.3) uses only ∇u while the calibration (2.12) from BPS decomposition involves second derivatives of u . For u smooth, since*

$$\begin{aligned} (\Delta_\perp u)^2 - (\partial_x^2 u - \partial_y^2 u)^2 &= 4\overline{K}, \\ (\partial_x u - \frac{1}{2}|\nabla_\perp u|^2) \cdot \Delta_\perp u &= \operatorname{div} \Xi(u) + \frac{2}{3}\overline{K}u, \\ (\partial_x u - \frac{1}{2}|\nabla_\perp u|^2) \cdot (\partial_x^2 u - \partial_y^2 u) &= \operatorname{div} \Sigma(\nabla u), \end{aligned}$$

the two calibrations can be linked by the following identity when $\overline{K} = 0$:

$$|\operatorname{div} \Xi| = |\operatorname{div} \Sigma(\nabla u)|.$$

Remark 2.2. *Comparing the BPS decomposition to (2.23), we see that the remainder term $\varepsilon \int_\Omega 2\overline{K}$ in the latter can be handled more easily than the term $\int_\Omega u \overline{K}$. This is the reason that we are able to obtain a lower bound even in the presence of non-vanishing Gaussian curvature. One might guess that since the BPS decomposition is also predicated on equipartition of energy between the bending and compression terms, the two arguments give the same lower bound*

$$\int_{J_{\nabla u}} \frac{|\nabla_\perp^+ u - \nabla_\perp^- u|^4}{12|\nabla^+ u - \nabla^- u|} \, d\mathcal{H}^2$$

in the limit $\varepsilon \rightarrow 0$ when $\overline{K} = 0$ for each u_ε and $\nabla u \in BV(\Omega; \mathbb{R}^3)$. This is indeed the case, although passing to the limit as $u_\varepsilon \rightarrow u$ in the term $\int_\Omega \operatorname{div} \Xi(u_\varepsilon)$ is non-trivial since $\Xi(u_\varepsilon)$ contains second order derivatives of u_ε . This can be accomplished via a blowup argument which then allows for a careful analysis of those higher order terms in the simplified setting of a flat jump set with limiting constant states ∇u^\pm on either side.

2.4 Properties of Functions of Bounded Variation

Our discussion draws from the relevant sections of [4, Chapter 3]. For the sake of generality and because the dimensions of the ambient/target spaces do not matter for these results, in this subsection we will consider functions defined on $\Omega \subset \mathbb{R}^N$ and taking values in \mathbb{R}^M .

Definition 2.1. [4, Def. 3.1] An element $m \in L^1(\Omega; \mathbb{R}^M)$ belongs to the space $BV(\Omega; \mathbb{R}^M)$ if the distributional derivative $Dm = (D_j m_i)$ of m is a finite $\mathbb{R}^{M \times N}$ -valued Radon measure.

Definition 2.2. [4, Def. 3.63] If $m \in L^1(\Omega; \mathbb{R}^M)$ we say that m has approximate limit $z = \text{ap lim}_{y \rightarrow x} m(y)$ at $x \in \Omega$ if

$$\lim_{r \rightarrow 0} \int_{B_r(x)} |m(y) - z| dy = 0.$$

If this property fails to hold at x for every $z \in \mathbb{R}^M$, then x belongs to S_m , the approximate discontinuity set.

To refer to solid half-balls in Ω , we define

$$B_r^+(x, \nu) := \{y \in B_r(x) : (y - x) \cdot \nu > 0\}, B_r^-(x, \nu) := \{y \in B_r(x) : (y - x) \cdot \nu < 0\}.$$

Definition 2.3. [4, Def. 3.67] For $m \in BV(\Omega; \mathbb{R}^M)$, we say that $x \in \Omega$ belongs to J_m , the set of approximate jump points, if there exist $m^+(x) \neq m^-(x) \in \mathbb{R}^M$ and $\nu_m(x) \in \mathbb{S}^{N-1}$ such that

$$\lim_{r \rightarrow 0} \int_{B_r^+(x, \nu_m)} |m(y) - m^+(x)| dy = 0 \text{ and } \lim_{r \rightarrow 0} \int_{B_r^-(x, \nu_m)} |m(y) - m^-(x)| dy = 0.$$

The vectors m^+ , m^- , and ν_m are uniquely determined up to permuting m^+ , m^- and exchanging $\nu_m(x)$ for $-\nu_m(x)$. Also, J_m is countably \mathcal{H}^{N-1} -rectifiable.

Definition 2.4. [4, Cor. 3.80] The precise representative of $m \in BV(\Omega; \mathbb{R}^M)$ is the function

$$\tilde{m}(x) := \begin{cases} \text{ap lim}_{y \rightarrow x} m(y) & \text{if } x \notin S_m, \\ \frac{m^+(x) + m^-(x)}{2} & \text{if } x \in J_m. \end{cases}$$

Theorem 2.3. [4, Thm. 3.78] If $m \in BV(\Omega; \mathbb{R}^M)$, then S_m is countably \mathcal{H}^{N-1} -rectifiable and $\mathcal{H}^{N-1}(S_m \setminus J_m) = 0$.

Next, we recall the BV Structure Theorem.

Theorem 2.4. [4, Section 3.9] For $m \in BV(\Omega; \mathbb{R}^M)$, the Radon measure Dm can be decomposed into three mutually singular measures

$$Dm = D^a m + D^j m + D^c m. \tag{2.26}$$

The first component is absolutely continuous with respect to the Lebesgue measure and is given by

$$D^a m = \nabla m \mathcal{L}^N,$$

where ∇m is the matrix of approximate partial derivatives defined \mathcal{L}^n -a.e. The component of Dm that is singular with respect to \mathcal{L}^N is $D^s m$. It can be written as $D^s m = D^j m + D^c m$, where

$$D^j m = (m^+ - m^-) \otimes \nu_m \mathcal{H}^{N-1} \llcorner J_m \tag{2.27}$$

and $D^c m$ is the Cantor part of Dm , which vanishes on sets that are \mathcal{H}^{N-1} σ -finite.

Remark 2.5. Since $D^a m$ and $D^c m$ both vanish on sets that are \mathcal{H}^{N-1} σ -finite, they vanish on S_m . Therefore, $\text{ap lim}_{y \rightarrow x} m(y)$, which is defined off of S_m , exists and is equal to \tilde{m} except on a set of $|D^a m|$ - and $|D^c m|$ -measure zero.

Lemma 2.6. *If $m \in BV(\Omega; \mathbb{R}^N)$ is equal to ∇u for some $u \in W^{1,1}(\Omega)$, then*

$$\nabla u^+ - \nabla u^- \ll \nu_{\nabla u}. \quad (2.28)$$

Proof. Since the Radon measure Dm is equal to $\nabla^2 u$, it is symmetric and can be decomposed into three mutually singular measures, cf. (2.26). Thus the jump part $(\nabla u^+ - \nabla u^-) \otimes \nu_{\nabla u} \mathcal{H}^2 \llcorner J_{\nabla u}$ is symmetric as well. But $(\nabla u^+ - \nabla u^-) \otimes \nu_{\nabla u}$ is symmetric if and only if $\nabla u^+ - \nabla u^- \ll \nu_{\nabla u}$, which is (2.28). \square

We state the BV Chain Rule [63, 2].

Theorem 2.7. [4, Thm 3.96] *Let $m \in BV(\Omega; \mathbb{R}^M)$ and $F : \mathbb{R}^M \rightarrow \mathbb{R}^P$ be C^1 with bounded gradient and $F(0) = 0$ if $\mathcal{L}^N(\Omega) = \infty$. Then $F \circ m \in BV(\Omega; \mathbb{R}^P)$ and*

$$D(F \circ m) = \nabla F(m) \nabla m \mathcal{L}^N + \nabla F(\tilde{m}) D^c m + (F(m^+) - F(m^-)) \otimes \nu_m \mathcal{H}^{N-1} \llcorner J_m. \quad (2.29)$$

When $M = P$, taking the trace on both sides of (2.29) yields

$$\operatorname{div}(F \circ m) = \operatorname{tr}(\nabla F(m) \nabla m) \mathcal{L}^2 + \operatorname{tr}(\nabla F(\tilde{m}) D^c m) + (F(m^+) - F(m^-)) \cdot \nu_m \mathcal{H}^{N-1} \llcorner J_m. \quad (2.30)$$

Remark 2.8. *As a consequence of Theorem 2.7, if F does not have bounded gradient, one must assume that m instead is bounded in order to apply the chain rule above.*

3 The 3D Aviles-Giga Space and the Lower Bound

In this section we prove the lower bound. Due to the connection between the 3D model (2.6) and 2D Aviles-Giga, many of the arguments leading to the lower bound are the natural 3D analogues of results from [3, 6]. The proof is centered around the 3D version of the 2D Aviles-Giga space considered in [6] and explicitly defined in [3]. Although alternate proofs are available when $\nabla u \in BV(\Omega; \mathbb{R}^3)$, for example via blowup or covering arguments, we follow the structure of [3] which gives the most general version of the lower bound without this assumption on ∇u .

Let $\{\xi, \eta\}$ be an orthonormal basis of \mathbb{R}^2 . For any vector $m = (m_\perp, m_3)$ with $m_\perp \in \mathbb{R}^2$, denote

$$m_\xi = m_\perp \cdot \xi, \quad m_\eta = m_\perp \cdot \eta$$

and set

$$\Sigma_{\xi\eta}(m) = \left(m_3 m_\xi - \frac{m_\xi m_\eta^2}{2} - \frac{m_\xi^3}{6} \right) \xi + \left(-m_3 m_\eta + \frac{m_\eta m_\xi^2}{2} + \frac{m_\eta^3}{6} \right) \eta + \left(-\frac{m_\xi^2}{2} + \frac{m_\eta^2}{2} \right) \hat{z}. \quad (3.1)$$

Note that for $m : \Omega \rightarrow \mathbb{R}^3$, $\Sigma_{\xi\eta}(m) \in L^1_{\operatorname{loc}}(\Omega; \mathbb{R}^3)$ if $m_\perp \in L^3_{\operatorname{loc}}(\Omega)$ and $m_3 \in L^{\frac{3}{2}}_{\operatorname{loc}}(\Omega)$. This and the fundamental equation (2.23) motivate the following definition.

Definition 3.1. *Let $u \in W^{1, \frac{3}{2}}_{\operatorname{loc}}(\Omega)$ be such that $\nabla_\perp u \in L^3_{\operatorname{loc}}(\Omega; \mathbb{R}^2)$. We say that $u \in AG^{3D}(\Omega)$ if $\operatorname{div} \Sigma_{\xi\eta}(\nabla u)$ is a finite Radon measure in Ω for all orthonormal bases $\{\xi, \eta\}$ of \mathbb{R}^2 .*

Definition 3.2. *If $u \in AG^{3D}(\Omega)$ and $\partial_z u = \frac{1}{2} |\nabla_\perp u|^2$, we say $u \in AG_0^{3D}(\Omega)$.*

Let $\{e_1, e_2\}$ be the standard basis of \mathbb{R}^2 , and let $\{\varepsilon_1, \varepsilon_2\}$ be given by

$$\varepsilon_1 = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right), \quad \varepsilon_2 = \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right). \quad (3.2)$$

A routine calculation, yields a formula for $\Sigma_{\xi\eta}$ in terms of $\Sigma_{e_1e_2}$ and $\Sigma_{\varepsilon_1\varepsilon_2}$. The \mathbb{R}^2 -valued version of this formula for the Jin-Kohn entropies was first derived in [3], and we do not include the proof.

Lemma 3.1. *For any $m \in L^{\frac{3}{2}}_{\text{loc}}(\Omega; \mathbb{R}^3)$ such that $m_{\perp} \in L^3_{\text{loc}}(\Omega; \mathbb{R}^2)$, if $\xi = (\cos \theta, \sin \theta)$ and $\eta = (-\sin \theta, \cos \theta)$, then*

$$\Sigma_{\xi\eta}(m) = \cos 2\theta \Sigma_{e_1e_2}(m) + \sin 2\theta \Sigma_{\varepsilon_1\varepsilon_2}(m). \quad (3.3)$$

Definition 3.3. *For any $u \in AG^{3D}(\Omega)$, let Iu be the finite vector-valued Radon measure*

$$Iu = (\text{div } \Sigma_{e_1e_2}(\nabla u), \text{div } \Sigma_{\varepsilon_1\varepsilon_2}(\nabla u)). \quad (3.4)$$

By (3.3), $u \in AG^{3D}(\Omega)$ if and only if $\text{div } \Sigma_{e_1e_2}(\nabla u)$, $\text{div } \Sigma_{\varepsilon_1\varepsilon_2}(\nabla u)$ are finite Radon measures. Let \mathcal{B}_+ denote the sets of all orthonormal bases of \mathbb{R}^2 with the same orientation as $\{e_1, e_2\}$. We decompose the domain and on each piece we consider the rotated measure $\text{div } \Sigma_{\xi\eta}(\nabla u)$. The following result shows that $|Iu|$ is the supremum of the measures $|\text{div } \Sigma_{\xi\eta}(\nabla u)|$ over $\{\xi, \eta\} \in \mathcal{B}_+$.

Theorem 3.2. (i) *For any $u \in AG^{3D}(\Omega)$ and Borel subset B of Ω ,*

$$\begin{aligned} |Iu|(B) &= \left(\bigvee_{\{\xi, \eta\} \in \mathcal{B}_+} |\text{div } \Sigma_{\xi\eta}(\nabla u)| \right) (B) \\ &= \sup \left\{ \sum_{j=1}^J |\text{div } \Sigma_{\xi_j \eta_j}(\nabla u)| (B_j) : J \in \mathbb{N}, \{B_j\}_{j=1}^J \text{ is a Borel partition of } B, \{\xi_j, \eta_j\} \in \mathcal{B}_+ \right\} \end{aligned} \quad (3.5)$$

(ii) *For an open set A ,*

$$|Iu|(A) = \sup \left\{ \sum_{j=1}^J |\text{div } \Sigma_{\xi_j \eta_j}(\nabla u)| (A_j) : J \in \mathbb{N}, A_j \text{ are open, disjoint, } A_j \subset\subset A, \{\xi_j, \eta_j\} \in \mathcal{B}_+ \right\}. \quad (3.6)$$

(iii) *If $\{u_n\} \subset AG^{3D}(\Omega)$ are such that*

$$\nabla_{\perp} u_n \xrightarrow{L^3} \nabla_{\perp} u, \quad \partial_z u_n \xrightarrow{L^{3/2}} \partial_z u,$$

and

$$|Iu_n|(\Omega) \leq C < \infty \quad \text{for all } n, \quad (3.7)$$

then $u \in AG^{3D}(\Omega)$ and

$$Iu_n \xrightarrow{*} Iu.$$

Thus for every open set $A \subset \Omega$,

$$|Iu|(A) \leq \liminf_{n \rightarrow \infty} |Iu_n|(A). \quad (3.8)$$

Proof. By Riesz's Theorem and (3.3), we have the equality of measures

$$\operatorname{div} \Sigma_{\xi\eta}(\nabla u) = (\cos 2\theta, \sin 2\theta) \cdot Iu = (\cos 2\theta, \sin 2\theta) \cdot g|Iu|$$

for some g which is unit-valued $|Iu|$ -a.e. One then has for any Borel set B and Borel partition $\{B_j\}_{j=1}^J$

$$\sum_{j=1}^J |\operatorname{div} \Sigma_{\xi_j\eta_j}(\nabla u)|(B_j) = \sum_{j=1}^J \int_{B_j} |(\cos 2\theta_j, \sin 2\theta_j) \cdot g| d|Iu| \leq \sum_{j=1}^J |Iu|(B_j). \quad (3.9)$$

Approximating g by functions that take finitely many values in \mathbb{S}^1 yields (3.5). The representation (3.6) when A is open is a consequence of the usual approximation theorems for Radon measures. To see this, note that for any partition $\{B_j\}_{j=1}^J$ of A , we can approximate B_j from inside by disjoint compact sets $\{K_j\}_{j=1}^J$ and then K_j by the desired open sets.

For (iii), by Hölder's inequality and the convergence of ∇u_n to ∇u , we have

$$\Sigma_{e_1 e_2}(\nabla u_n) \rightarrow \Sigma_{e_1 e_2}(\nabla u), \quad \Sigma_{\varepsilon_1 \varepsilon_2}(\nabla u_n) \rightarrow \Sigma_{\varepsilon_1 \varepsilon_2}(\nabla u) \quad \text{in } L^1(\Omega; \mathbb{R}^3),$$

so that the divergences converge in the sense of distributions. Thus (3.7) implies that Iu is Radon and $Iu_n \xrightarrow{*} Iu$, and (3.8) is a consequence of the weak-* convergence. \square

Proposition 3.3. *If $u \in W^{2,9/5}(\Omega)$, then*

$$\operatorname{div} \Sigma_{\xi\eta}(\nabla u) = \left(\partial_z u - \frac{1}{2} |\nabla u|^2 \right) (\partial_\xi^2 u - \partial_\eta^2 u) \quad (3.10)$$

and

$$|Iu| = \left| \partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right| |\lambda_1 - \lambda_2| \mathcal{L}^3 \llcorner \Omega \quad (3.11)$$

where λ_1, λ_2 are the eigenvalues of $\nabla_\perp^2 u$.

Proof. If u is smooth, we can calculate

$$\begin{aligned} \operatorname{div} \Sigma_{\xi\eta}(\nabla u) &= \partial_\xi (\Sigma_{\xi\eta}(\nabla u) \cdot \xi) + \partial_\eta (\Sigma_{\xi\eta}(\nabla u) \cdot \eta) + \partial_z (\Sigma_{\xi\eta}(\nabla u) \cdot \hat{z}) \\ &= \partial_\xi \partial_z u \partial_\xi u + \partial_z u \partial_\xi^2 u - \partial_\xi u \partial_\eta u \partial_\xi \partial_\eta u - \frac{\partial_\xi^2 u (\partial_\eta u)^2}{2} - \frac{(\partial_\xi u)^2 \partial_\xi^2 u}{2} \\ &\quad - \partial_\eta \partial_z u \partial_\eta u - \partial_z u \partial_\eta^2 u + \partial_\xi u \partial_\eta u \partial_\eta \partial_\xi u + \frac{\partial_\eta^2 u (\partial_\xi u)^2}{2} + \frac{(\partial_\eta u)^2 \partial_\eta^2 u}{2} \\ &\quad - \partial_\xi u \partial_z \partial_\xi u + \partial_\eta u \partial_z \partial_\eta u \\ &= \left(\partial_z u - \frac{|\nabla_\perp u|^2}{2} \right) (\partial_\xi^2 u - \partial_\eta^2 u). \end{aligned} \quad (3.12)$$

If $u \in W^{2,9/5}(\Omega)$, then by the Sobolev embedding, $\nabla u \in L^{9/2}(\Omega)$. For u_n smooth and converging to u in $W^{2,9/5}(\Omega)$, Hölder's inequality yields for any test function $\varphi \in C_0^\infty(\Omega)$,

$$\begin{aligned} \int_\Omega -\Sigma_{\xi\eta}(\nabla u) \cdot \nabla \varphi \, d\mathbf{x} &= \lim_{n \rightarrow \infty} \int_\Omega -\Sigma_{\xi\eta}(\nabla u_n) \cdot \nabla \varphi \, d\mathbf{x} \\ &= \lim_{n \rightarrow \infty} \int_\Omega \left(\partial_z u_n - \frac{|\nabla_\perp u_n|^2}{2} \right) (\partial_\xi^2 u_n - \partial_\eta^2 u_n) \varphi \, d\mathbf{x} \\ &= \int_\Omega \left(\partial_z u - \frac{|\nabla_\perp u|^2}{2} \right) (\partial_\xi^2 u - \partial_\eta^2 u) \varphi \, d\mathbf{x}, \end{aligned}$$

so that (3.10) is proved.

For (3.11), if $v = v(\mathbf{x})$ and $w = w(\mathbf{x})$ are the orthonormal eigenvectors of $\nabla_{\perp}^2 u(\mathbf{x})$ with corresponding eigenvalues $\lambda_1(\mathbf{x}) = \partial_v^2 u(\mathbf{x})$ and $\lambda_2(\mathbf{x}) = \partial_w^2 u(\mathbf{x})$, then direct calculation gives

$$|\partial_{\xi}^2 u - \partial_{\eta}^2 u| = |\partial_v^2 u - \partial_w^2 u| |(\xi \cdot v)^2 - (\eta \cdot v)^2| = |\partial_v^2 u - \partial_w^2 u| |(\xi \cdot w)^2 - (\eta \cdot w)^2|.$$

From this we may conclude that

$$\sup_{\{\xi, \eta\} \in \mathcal{B}_+} |\partial_{\xi}^2 u - \partial_{\eta}^2 u| = |\lambda_1 - \lambda_2|,$$

and thus

$$|Iu| = \left| \partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right| |\lambda_1 - \lambda_2| \mathcal{L}^3 \llcorner \Omega.$$

□

Remark 3.4. *The condition that $u \in W^{2, \frac{9}{5}}(\Omega)$ is stronger than merely requiring that $u \in AG^{3D}(\Omega)$. However, by Hölder's inequality and the Sobolev embedding, $9/5$ is the optimal exponent for which the measure $\text{div } \Sigma(\nabla u)$ is absolutely continuous with respect to the Lebesgue measure and is thus represented by the integration of an L^1 -function; cf. [3, Proposition 3.4] for the corresponding result in two dimensions.*

The next proposition gives a formula for $|Iu|$ under certain regularity conditions on $u \in AG_0^{3D}(\Omega)$ necessary to apply the BV Chain rule, cf. Theorem 2.7 and Remark 2.8.

Proposition 3.5. *If $u \in AG_0^{3D}(\Omega) \cap W^{1, \infty}(\Omega)$ and $\nabla u \in BV(\Omega; \mathbb{R}^3)$, then*

$$|Iu| = \frac{|\nabla_{\perp} u^+ - \nabla_{\perp} u^-|^4}{12 |\nabla u^+ - \nabla u^-|} \mathcal{H}^2 \llcorner J_{\nabla u}.$$

Proof. For any $\{\xi, \eta\}$, first notice that due to the BV Chain rule,

$$|\text{div } \Sigma_{\xi\eta}(\nabla u)| \llcorner J_{\nabla u} = |(\Sigma_{\xi\eta}(\nabla u^+) - \Sigma_{\xi\eta}(\nabla u^-)) \cdot \nu_{\nabla u}| \mathcal{H}^2 \llcorner J_{\nabla u}.$$

We compute the right hand side and then optimize over choices of $\{\xi, \eta\} \in \mathcal{B}_+$ at each point in $J_{\nabla u}$. To simplify the notation in the calculation, set $\nabla u = m$. Now since $u \in AG_0^{3D}(\Omega)$, m^+ and m^- satisfy

$$m_{\pm}^{\pm} = \frac{1}{2} ((m_{\xi}^{\pm})^2 + (m_{\eta}^{\pm})^2) \quad (3.13)$$

on J_m , which can be directly verified from Definition 2.3. This gives

$$\begin{aligned} \Sigma_{\xi\eta}(m^{\pm})_{\perp} &= \left(m_{\pm}^{\pm} m_{\xi}^{\pm} - \frac{m_{\xi}^{\pm} (m_{\eta}^{\pm})^2}{2} - \frac{(m_{\xi}^{\pm})^3}{6} \right) \xi + \left(-m_{\pm}^{\pm} m_{\eta}^{\pm} + \frac{m_{\eta}^{\pm} (m_{\xi}^{\pm})^2}{2} + \frac{(m_{\eta}^{\pm})^3}{6} \right) \eta \\ &= \frac{(m_{\xi}^{\pm})^3}{3} \xi - \frac{(m_{\eta}^{\pm})^3}{3} \eta. \end{aligned} \quad (3.14)$$

Using (3.14) to rewrite $\Sigma_{\xi\eta}(m^\pm)$ and then (3.13) and $m^+ - m^- // \nu_m$ to replace ν_m , cf. Equation (2.28), we have

$$\begin{aligned} & |(\Sigma_{\xi\eta}(m^+) - \Sigma_{\xi\eta}(m^-)) \cdot \nu_m| \\ &= \left| \left(\frac{(m_\xi^+)^3}{3} - \frac{(m_\xi^-)^3}{3} \right) \nu_\xi - \left(\frac{(m_\eta^+)^3}{3} - \frac{(m_\eta^-)^3}{3} \right) \nu_\eta + \left(-\frac{(m_\xi^+)^2}{2} + \frac{(m_\xi^-)^2}{2} + \frac{(m_\eta^+)^2}{2} - \frac{(m_\eta^-)^2}{2} \right) \nu_z \right| \\ &= \frac{1}{|m^+ - m^-|} \left| \left(\frac{(m_\xi^+)^3}{3} - \frac{(m_\xi^-)^3}{3} \right) (m_\xi^+ - m_\xi^-) - \left(\frac{(m_\eta^+)^3}{3} - \frac{(m_\eta^-)^3}{3} \right) (m_\eta^+ - m_\eta^-) \right. \\ &\quad \left. + \left(-\frac{(m_\xi^+)^2}{2} + \frac{(m_\xi^-)^2}{2} + \frac{(m_\eta^+)^2}{2} - \frac{(m_\eta^-)^2}{2} \right) \left(\frac{|m_\perp^+|^2}{2} - \frac{|m_\perp^-|^2}{2} \right) \right|. \end{aligned}$$

Expanding out the right hand side of the previous equation and combining like terms gives

$$|(\Sigma_{\xi\eta}(m^+) - \Sigma_{\xi\eta}(m^-)) \cdot \nu_m| = \frac{|(m_\xi^+ - m_\xi^-)^4 - (m_\eta^+ - m_\eta^-)^4|}{12|m^+ - m^-|} \leq \frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|}. \quad (3.15)$$

Equality is achieved for $\{\xi, \eta\}$ such that $(m^+ - m^-)_\perp // \xi$ or $(m^+ - m^-)_\perp // \eta$. Taking the supremum over $\{\xi, \eta\} \in \mathcal{B}_+$, we find that

$$|Iu| \llcorner J_{\nabla u} = \frac{|\nabla u_\perp^+ - \nabla u_\perp^-|^4}{12|\nabla u^+ - \nabla u^-|} \mathcal{H}^2 \llcorner J_{\nabla u}. \quad (3.16)$$

To complete the proof of the proposition, we must show that

$$|Iu|(\Omega \setminus J_{\nabla u}) = 0.$$

Recalling the BV Chain rule, Theorem 2.7, notice that away from $J_{\nabla u}$, $\operatorname{div} \Sigma_{\xi\eta}(\nabla u)$ can be computed using the usual chain rule formula by substituting $D^a(\nabla u)$ and $D^c(\nabla u)$ for the classical second derivatives of u . Therefore, by the same manipulations as in (3.12), we have

$$\begin{aligned} & \operatorname{div} \Sigma_{\xi\eta}(\nabla u) \llcorner (\Omega \setminus J_{\nabla u}) \\ &= \left(\widetilde{\partial_z u} - \frac{|\widetilde{\nabla_\perp u}|^2}{2} \right) [\xi^T D^a(\nabla u) \xi + \xi^T D^c(\nabla u) \xi - \eta^T D^a(\nabla u) \eta - \eta^T D^c(\nabla u) \eta]. \end{aligned}$$

Since $\widetilde{\partial_z u} - \frac{|\widetilde{\nabla_\perp u}|^2}{2} = 0$ for \mathbf{x} where the approximate limit $\widetilde{\nabla u}$ exists, it is zero $|D^a(\nabla u)|$ - and $|D^c(\nabla u)|$ -a.e. by Remark 2.5. Thus $|Iu| \llcorner (\Omega \setminus J_{\nabla u})$ vanishes as well. \square

We are ready to prove the lower bound. The theorem is stated under the assumption that $\varepsilon_n^2 \int \overline{K}_n \rightarrow 0$, which can be enforced by mild control on the boundary data as in (2.25).

Theorem 3.6. *Let $\Omega \subset \mathbb{R}^3$ be an open set. Consider $\varepsilon_n \searrow 0$ and $\{u_n\} \subset H^2(\Omega)$ such that*

$$\nabla_\perp u_n \xrightarrow{L^3} \nabla_\perp u, \quad \partial_z u_n \xrightarrow{L^{3/2}} \partial_z u \quad (3.17)$$

for some $u \in W^{1, \frac{3}{2}}(\Omega)$ with $\nabla_\perp u \in L^3(\Omega; \mathbb{R}^2)$. If $\liminf_{n \rightarrow \infty} \mathcal{E}_{\varepsilon_n}(u_n)$ is finite and

$$\lim_{n \rightarrow \infty} \varepsilon_n^2 \int_\Omega \overline{K}_n \, dx = 0, \quad (3.18)$$

then $u \in AG_0^{3D}(\Omega)$ and

$$\liminf_{n \rightarrow \infty} \mathcal{E}_{\varepsilon_n}(u_n) \geq |Iu|(\Omega). \quad (3.19)$$

When $u \in AG_0^{3D}(\Omega) \cap W^{1,\infty}(\Omega)$ and $\nabla u \in BV(\Omega; \mathbb{R}^3)$, then by Proposition 3.5, the lower bound is given by

$$|Iu| = \frac{|\nabla_{\perp} u^+ - \nabla_{\perp} u^-|^4}{12|\nabla u^+ - \nabla u^-|} \mathcal{H}^2 \llcorner J_{\nabla u}.$$

Proof. For each $u_n \in H^2(\Omega)$, the representation (3.11) of $|Iu_n|$ gives

$$\begin{aligned} |Iu_n| &= \left| \partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right| |\lambda_{1,n} - \lambda_{2,n}| \mathcal{L}^3 \llcorner \Omega \\ &\leq \left[\frac{1}{2\varepsilon_n} \left(\partial_z u_n - \frac{|\nabla_{\perp} u_n|^2}{2} \right)^2 + \frac{\varepsilon_n}{2} (\lambda_{1,n} - \lambda_{2,n})^2 \right] \mathcal{L}^3 \llcorner \Omega. \end{aligned} \quad (3.20)$$

Next, fix an open set $A \subset\subset \Omega$ and a test function $\varphi \in C_c^\infty(\Omega; [0, 1])$ such that $\varphi = 1$ on A . We estimate

$$\begin{aligned} \frac{\varepsilon_n}{2} \int_{\Omega} (\Delta_{\perp} u_n)^2 d\mathbf{x} &\geq \frac{\varepsilon_n}{2} \int_{\Omega} [\lambda_{1,n}^2 + \lambda_{2,n}^2 + 2\lambda_{1,n}\lambda_{2,n}] \varphi d\mathbf{x} \\ &= \frac{\varepsilon_n}{2} \int_{\Omega} [(\lambda_{1,n} - \lambda_{2,n})^2 + 4 \det(\nabla_{\perp}^2 u_n)] \varphi d\mathbf{x} \\ &= \frac{\varepsilon_n}{2} \int_{\Omega} (\lambda_{1,n} - \lambda_{2,n})^2 \varphi d\mathbf{x} - 2\varepsilon_n \int_{\Omega} \partial_x u_n (\partial_y^2 u_n \partial_x \varphi - \partial_{xy} u_n \partial_y \varphi) d\mathbf{x} \\ &\geq \frac{\varepsilon_n}{2} \int_A (\lambda_{1,n} - \lambda_{2,n})^2 d\mathbf{x} - 2\varepsilon_n \|\nabla \varphi\|_{L^\infty} \|\nabla_{\perp} u_n\|_{L^2} \|\nabla_{\perp}^2 u_n\|_{L^2}. \end{aligned} \quad (3.21)$$

Now since $\nabla_{\perp} u_n$ are bounded in L^3 , the square of the remainder in (3.21) can be estimated by

$$\begin{aligned} \varepsilon_n^2 \|\nabla_{\perp} u_n\|_{L^2}^2 \|\nabla_{\perp}^2 u_n\|_{L^2}^2 &= \varepsilon_n^2 \|\nabla_{\perp} u_n\|_{L^2}^2 \int_{\Omega} [\lambda_{1,n}^2 + \lambda_{2,n}^2 + 2\lambda_{1,n}\lambda_{2,n} - 2\lambda_{1,n}\lambda_{2,n}] d\mathbf{x} \\ &\leq C\varepsilon_n^2 \int_{\Omega} (\Delta_{\perp} u_n)^2 d\mathbf{x} + C\varepsilon_n^2 \left| 2 \int_{\Omega} \overline{K}_n d\mathbf{x} \right| \\ &\rightarrow 0. \end{aligned} \quad (3.22)$$

Combining (3.20)-(3.22), we conclude that

$$\begin{aligned} \mathcal{E}_{\varepsilon_n}(u_n) &\geq \frac{1}{2} \int_A \left[\frac{1}{\varepsilon_n} \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right)^2 + \varepsilon_n (\lambda_{1,n} - \lambda_{2,n})^2 \right] d\mathbf{x} - 2\varepsilon_n \|\nabla \varphi\|_{L^\infty} \|\nabla_{\perp} u_n\|_{L^2} \|\nabla_{\perp}^2 u_n\|_{L^2} \\ &\geq |Iu_n|(A) - o(1). \end{aligned} \quad (3.23)$$

Since the limit inferior of the energies is finite, we can appeal to Theorem 3.2.(iii) to find that $u \in AG^{3D}(A)$ and

$$\liminf_{n \rightarrow \infty} \mathcal{E}_{\varepsilon_n}(u_n) \geq \liminf_{n \rightarrow \infty} |Iu_n|(A) \geq |Iu|(A).$$

An exhaustion argument gives $u \in AG^{3D}(\Omega)$ and (3.19). The fact that $u \in AG_0^{3D}(\Omega)$ follows from

$$\int_{\Omega} \left(\partial_z u_n - \frac{|\nabla_{\perp} u_n|^2}{2} \right)^2 d\mathbf{x} \leq \varepsilon_n \mathcal{E}_{\varepsilon_n}(u_n) \rightarrow 0. \quad (3.24)$$

□

4 The Upper Bound

In this section we show that the lower bound Theorem 3.6 is sharp when $u \in AG_0^{3D}(\Omega)$ and $\nabla u \in (BV \cap L^\infty)(\Omega)$ by means of a construction, so that we have matching upper and lower bounds. Combined with the lower bound, this allows us to conclude that under reasonable assumptions, equipartition of energy in (2.6) is optimal.

Theorem 4.1. *Let $u \in AG_0^{3D}(\Omega) \cap W^{1,\infty}(\Omega)$ and $\nabla u \in BV(\Omega; \mathbb{R}^3)$. Then there exists a sequence $\{u_\varepsilon\} \subset C^2(\Omega)$ such that*

$$u_\varepsilon \rightarrow u \text{ in } W^{1,p}(\Omega) \text{ for all } 1 \leq p < \infty \quad (4.1)$$

and

$$\mathcal{E}_\varepsilon(u_\varepsilon) \rightarrow \int_{J_{\nabla u}} \frac{|\nabla_\perp u^+ - \nabla_\perp u^-|^4}{12|\nabla u^+ - \nabla u^-|} d\mathcal{H}^2. \quad (4.2)$$

The proof of Theorem 4.1 consists of two steps. In Proposition 4.2, we show that on a cube with jump set parallel to one of the faces, the sequence of one-dimensional competitors with constant gradient in the direction parallel to the jump set is asymptotically minimizing. Second, the cube construction can be leveraged to obtain the full upper bound Theorem 4.1 by using the results of [50].

To formulate the problem on a cube, let us fix an orthonormal basis $\{\zeta_1, \zeta_2, \nu\}$ of \mathbb{R}^3 and the set

$$C = \{\mathbf{x} \in \mathbb{R}^3 : |\mathbf{x} \cdot \nu| \leq 1/2, |\mathbf{x} \cdot \zeta_i| \leq 1/2 \text{ for } i = 1, 2\}.$$

Next, we choose boundary data that will be compatible with a limiting jump set $\{\mathbf{x} \in C : \mathbf{x} \cdot \nu = 0\}$. Let $m^+ \neq m^-$ be such that

$$m_3^\pm = \frac{1}{2}|m_\perp^\pm|^2 \quad \text{and} \quad \nu // (m^+ - m^-), \quad (4.3)$$

and consider the class

$$\mathcal{A}_C := \{u \in H^2 : \nabla u = m^\pm \text{ when } \mathbf{x} \cdot \nu = \pm 1/2 \text{ and } \nabla u \text{ is 1-periodic in the } \zeta_1, \zeta_2 \text{ directions}\}.$$

Note that since $m^+ \neq m^-$, the first equation in (4.3) enforces

$$\nu_\perp \neq 0,$$

so that we can define the planar unit vectors

$$\xi = \frac{\nu_\perp}{|\nu_\perp|} \quad \text{and} \quad \eta = (-\xi_2, \xi_1, 0).$$

The smaller set of 1D competitors is defined by

$$\mathcal{A}_C^{1D} := \{u \in \mathcal{A}_C : \nabla u \cdot \zeta_i = m^+ \cdot \zeta_i = m^- \cdot \zeta_i \text{ for } i = 1, 2\}.$$

We remark that due to the boundary conditions imposed on the class \mathcal{A}_C and the identity $\overline{K} = \det \nabla_\perp^2 u = \nabla_\perp (\partial_x u \partial_y^2 u, -\partial_x u \partial_{xy} u)$,

$$\begin{aligned} \int_C (\Delta_\perp u)^2 d\mathbf{x} - \int_C |\nabla_\perp^2 u|^2 d\mathbf{x} \\ = 2 \int_C \partial_x^2 u \partial_y^2 u - (\partial_{xy} u)^2 d\mathbf{x} \end{aligned}$$

$$\begin{aligned}
&= 2 \int_{\partial C \cap \{\mathbf{x} \cdot \nu = 1/2\}} \partial_x u \partial_{\tau_{1,2}}(\partial_y u) d\mathcal{H}^2 + 2 \sum_{i=1}^2 \int_{\partial C \cap \{|\mathbf{x} \cdot \zeta_i| = 1/2\}} \partial_x u \partial_{\tau_{1,2}}(\partial_y u) d\mathcal{H}^2 \\
&= 0.
\end{aligned} \tag{4.4}$$

We set

$$r_\varepsilon = \inf_{\mathcal{A}_C} \mathcal{E}_\varepsilon \quad \text{and} \quad r_\varepsilon^{1D} = \inf_{\mathcal{A}_C^{1D}} \mathcal{E}_\varepsilon.$$

Proposition 4.2. *For any $\varepsilon > 0$,*

$$\frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|} \leq r_\varepsilon \leq r_\varepsilon^{1D} \leq \frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|} + c_1 e^{-c_2 \varepsilon}. \tag{4.5}$$

The constants c_1 and c_2 depend only on m^+ and m^- .

Proof. The inequality $r_\varepsilon \leq r_\varepsilon^{1D}$ is immediate, since $\mathcal{A}_C^{1D} \subset \mathcal{A}_C$. Also, the inequality

$$\frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|} \leq r_\varepsilon$$

follows from (3.12), (3.15), and the boundary conditions for $u \in \mathcal{A}_C$. Indeed, since

$$\nu_\perp // (m^+ - m^-)_\perp // \xi,$$

we have

$$\begin{aligned}
\frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|} &\stackrel{(3.15)}{=} \left| \int_{C \cap \{\mathbf{x} \cdot \nu = 1/2\}} \Sigma_{\xi\eta}(m^+) \cdot \nu d\mathcal{H}^2 - \int_{C \cap \{\mathbf{x} \cdot \nu = -1/2\}} \Sigma_{\xi\eta}(m^-) \cdot \nu d\mathcal{H}^2 \right| \\
&= \left| \int_C \operatorname{div} \Sigma_{\xi\eta}(\nabla u) d\mathbf{x} \right| \\
&\stackrel{(3.12)}{\leq} \int_C \frac{1}{2\varepsilon} \left(\partial_z u - \frac{|\nabla_\perp u|^2}{2} \right)^2 + \frac{\varepsilon}{2} (\partial_\xi^2 u - \partial_\eta^2 u)^2 d\mathbf{x} \\
&\leq \int_C \frac{1}{2\varepsilon} \left(\partial_z u - \frac{|\nabla_\perp u|^2}{2} \right)^2 + \frac{\varepsilon}{2} \left((\partial_\xi^2 u)^2 + 2(\partial_\eta \partial_\xi u)^2 + (\partial_\eta^2 u)^2 \right) d\mathbf{x} \\
&\quad - \varepsilon \int_C (\partial_\xi^2 u \partial_\eta^2 u - (\partial_\eta \partial_\xi u)^2) d\mathbf{x} \\
&= \frac{1}{2} \int_C \frac{1}{\varepsilon} \left(\partial_z u - \frac{|\nabla_\perp u|^2}{2} \right)^2 + \varepsilon |\nabla_\perp^2 u|^2 d\mathbf{x} - \varepsilon \int_C \det(\nabla_\perp^2 u) d\mathbf{x} \\
&\stackrel{(4.4)}{=} \mathcal{E}_\varepsilon(u).
\end{aligned}$$

Finally, showing that

$$r_\varepsilon^{1D} \leq \frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|} + c_1 e^{-c_2 \varepsilon} \tag{4.6}$$

entails constructing a sequence $\{\nabla u_\varepsilon\}$ such that each ∇u_ε is a function of $\mathbf{x} \cdot \nu$ and

$$\mathcal{E}_\varepsilon(u_\varepsilon) \leq \frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|} + c_1 e^{-c_2 \varepsilon}.$$

Since the steps of such a construction are standard in the calculus of variations, we outline the procedure and refer to [47, Proposition 5.2], which contains a full proof in the 2D case, for some of the estimates.

Let

$$p = m^+ - m^-$$

and g be the solution to the initial value problem

$$\begin{cases} g'(t) = \frac{|gp_3 + m_3^- - (gp_1 + m_1^-)^2/2 - (gp_2 + m_2^-)^2/2|}{|p_\perp \cdot \nu_\perp|}, \\ g(0) = 1/2. \end{cases} \quad (4.7)$$

Note that the denominator $|p_\perp \cdot \nu_\perp| \neq 0$ since $m_3^\pm = \frac{1}{2}|m_\perp^\pm|^2$, $m^+ \neq m^-$ imply that $m_\perp^+ \neq m_\perp^-$. One can check that g exists for all time and approaches 1 and 0 exponentially as $t \rightarrow \pm\infty$ (see for example [59, Equation (1.21)]). Consider the family of functions

$$g\left(\frac{\mathbf{x} \cdot \nu}{\varepsilon}\right) p + m^- = \nabla \left[\varepsilon |p| G\left(\frac{\mathbf{x} \cdot \nu}{\varepsilon}\right) + m^- \cdot \mathbf{x} \right] =: \nabla w_\varepsilon(\mathbf{x}),$$

where G is an antiderivative of g , on the infinite strip $\{|\mathbf{x} \cdot \tau_i| \leq 1/2 : i = 1, 2\}$. Let ξ be the unit vector $\nu_\perp/|\nu_\perp|$. By direct calculation, we have for any $\varepsilon > 0$

$$\begin{aligned} & \mathcal{E}_\varepsilon(\nabla w_\varepsilon) \\ &= \int_{\{|\mathbf{x} \cdot \tau_i| \leq 1/2; i=1,2\}} \frac{1}{\varepsilon} \left(\partial_z w_\varepsilon - \frac{1}{2} |\nabla_\perp w_\varepsilon|^2 \right)^2 + \varepsilon (\Delta_\perp w_\varepsilon)^2 \, d\mathbf{x} \\ &= \int_{-\infty}^{\infty} \left[\frac{1}{\varepsilon} \left(g\left(\frac{t}{\varepsilon}\right) p_3 + m_3^- - \frac{(g\left(\frac{t}{\varepsilon}\right) p_1 + m_1^-)^2}{2} - \frac{(g\left(\frac{t}{\varepsilon}\right) p_2 + m_2^-)^2}{2} \right)^2 + \varepsilon g'\left(\frac{t}{\varepsilon}\right)^2 \frac{(p_\perp \cdot \nu_\perp)^2}{\varepsilon^2} \right] dt \\ &\stackrel{(4.7)}{=} \left| \int_{-\infty}^{\infty} \left(g\left(\frac{t}{\varepsilon}\right) p_3 + m_3^- - \frac{(g\left(\frac{t}{\varepsilon}\right) p_1 + m_1^-)^2}{2} - \frac{(g\left(\frac{t}{\varepsilon}\right) p_2 + m_2^-)^2}{2} \right) g'\left(\frac{t}{\varepsilon}\right) \frac{(p_\perp \cdot \nu_\perp)}{\varepsilon} dt \right| \\ &= \left| \int_{\{|\mathbf{x} \cdot \tau_i| \leq 1/2; i=1,2\}} \left(\partial_z w_\varepsilon - \frac{|\nabla_\perp w_\varepsilon|^2}{2} \right) \partial_\xi^2 w_\varepsilon \, d\mathbf{x} \right| \\ &= \left| \int_{\{|\mathbf{x} \cdot \tau_i| \leq 1/2; i=1,2\}} \left(\partial_z w_\varepsilon - \frac{|\nabla_\perp w_\varepsilon|^2}{2} \right) (\partial_\xi^2 w_\varepsilon - \partial_\eta^2 w_\varepsilon) \, d\mathbf{x} \right| \\ &\stackrel{(3.12)}{=} \left| \int_{\{|\mathbf{x} \cdot \tau_i| \leq 1/2; i=1,2\}} \operatorname{div} \Sigma_{\xi\eta}(\nabla w_\varepsilon) \, d\mathbf{x} \right| \\ &= |(\Sigma_{\xi\eta}(m^+) - \Sigma_{\xi\eta}(m^-)) \cdot \nu| \end{aligned}$$

In (3.16) in the lower bound, we saw that this was equal to

$$\frac{|m_\perp^+ - m_\perp^-|^4}{12|m^+ - m^-|}. \quad (4.8)$$

The sequence $\{\nabla u_\varepsilon\}$ is constructed by suitably truncating ∇w_ε so that it is in the class \mathcal{A}_C^{1D} . The estimate (4.6) follows from the exponential approach of g to 0 and 1 combined with (4.8); see [47, Proposition 5.2] for full details. \square

To prove Theorem 4.1, we appeal to a general theorem from [49]. The version applicable to this problem reads as follows.

Theorem 4.3. ([49, Theorem 1.2]) *Let $\Omega \subset \mathbb{R}^3$ be a bounded C^2 -domain and let*

$$F(a, b) : \mathbb{R}^{3 \times 3} \times \mathbb{R}^3 \rightarrow \mathbb{R}$$

be a C^1 function satisfying $F \geq 0$. Let $u \in W^{1,\infty}(\Omega)$ be such that $\nabla u \in BV(\Omega; \mathbb{R}^3)$ and $F(0, \nabla u(x)) = 0$ a.e. in Ω . Then there exists a family of functions $\{u_\varepsilon\} \subset C^2(\mathbb{R}^3)$ satisfying

$$u_\varepsilon \rightarrow u \text{ in } W^{1,p}(\Omega) \text{ for } 1 \leq p < \infty$$

and

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_{\Omega} F(\varepsilon \nabla^2 u_\varepsilon, \nabla u_\varepsilon) dx dz \\ &= \int_{J_{\nabla u}} \inf_{r \in \mathcal{R}_{\chi(x,z),0}} \left\{ \int_{-\infty}^{\infty} F(-r'(t)\nu(x,z) \otimes \nu(x,z), r(t)\nu(x,z) + \nabla u^-(x,z)) dt \right\} d\mathcal{H}^2. \end{aligned}$$

Here $\chi(x, z)$ is given by

$$\chi(x, z)\nu(x, z) = \nabla u^+(x, z) - \nabla u^-(x, z),$$

and

$$\mathcal{R}_{\chi(x,z),0} := \{r(t) \in C^1(\mathbb{R}) : \exists L > 0 \text{ s.t. } r(t) = \chi(x, z) \text{ for } t \leq -L, r(t) = 0 \text{ for } t \geq L\}.$$

Proof of Theorem 4.1. If we set

$$F(a, b) = \frac{1}{2} \left(b_3 - \frac{|b_\perp|^2}{2} \right)^2 + \frac{1}{2} \sum_{i=1}^2 a_{ii}^2,$$

then

$$\frac{1}{\varepsilon} \int_{\Omega} F(\varepsilon \nabla^2 u_\varepsilon, \nabla u_\varepsilon) d\mathbf{x} = \mathcal{E}_\varepsilon(u_\varepsilon).$$

To evaluate the infimum in Theorem 4.3, we can rescale and use Proposition 4.2 to see that it is

$$\frac{|\nabla u_\perp^+ - \nabla u_\perp^-|^4}{12|\nabla u^+ - \nabla u^-|}.$$

This finishes the proof. \square

Remark 4.4. *A recovery sequence with specified boundary data for u and ∇u could be constructed as in [12, Section 6] or [48, Theorem 1.1].*

Finally, let us rephrase the theorems of the last two sections in terms of the original problem involving smectics: roughly speaking, equipartition of energy is optimal when the Gaussian curvature induced by the boundary conditions is not prohibitively large. More precisely, denoting by λ_i the eigenvalues of $\nabla_\perp^2 u$, we have

$$\begin{aligned} \mathcal{E}_\varepsilon(u) &= \frac{1}{2} \int_{\Omega} \left[\varepsilon(\lambda_1 + \lambda_2)^2 + \frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right)^2 \right] d\mathbf{x} \\ &= \frac{1}{2} \int_{\Omega} \left[\varepsilon(\lambda_1 - \lambda_2)^2 + \frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right)^2 \right] d\mathbf{x} + 2\varepsilon \int_{\Omega} \lambda_1 \lambda_2 d\mathbf{x} \\ &= |Iu|(\Omega) + \frac{1}{2} \int_{\Omega} \left(\varepsilon^{1/2} |\lambda_1 - \lambda_2| - \frac{1}{\varepsilon^{1/2}} \left| \partial_z u - \frac{1}{2} |\nabla_\perp u|^2 \right| \right)^2 d\mathbf{x} + 2\varepsilon \int_{\Omega} \bar{K} d\mathbf{x}. \end{aligned} \quad (4.9)$$

If $\varepsilon \int_{\Omega} \bar{K} \, d\mathbf{x}$ is small compared to the energy $\mathcal{E}_{\varepsilon}(u)$, which by (2.24) can be enforced by choosing boundary conditions such that $\varepsilon \|\nabla_{\perp} u\|_{H^{1/2}(\partial\Omega)}^2$ is small, then contribution of the curvature term is negligible. Thus the energy $\mathcal{E}_{\varepsilon}$ is minimized by minimizing $|Iu|$ among competitors saturating the perfect square. The matching upper bound demonstrates that this procedure is optimal in a reasonable range of situations. Furthermore, saturation of the perfect square entails

$$\int_{\Omega} \varepsilon(\lambda_1 - \lambda_2)^2 \, d\mathbf{x} \approx \int_{\Omega} \frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right)^2 \, d\mathbf{x}. \quad (4.10)$$

Since $\varepsilon(\Delta_{\perp} u)^2$ and $\varepsilon(\lambda_1 - \lambda_2)^2$ differ by $4\varepsilon\bar{K}$, the assumption that the integral of the curvature is small and (4.10) imply that

$$\int_{\Omega} \varepsilon(\Delta_{\perp} u)^2 \, d\mathbf{x} \approx \int_{\Omega} \frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right)^2 \, d\mathbf{x}, \quad (4.11)$$

which is precisely the BPS equation squared and integrated over Ω .

Remark 4.5. *Our 1D ansatz satisfies BPS equation (2.13). Also, the condition that $\varepsilon \int_{\Omega} \bar{K}$ must be small for equipartition to be optimal coincides with the observation from [55] that BPS solutions are not competitive when the curvature is very large, so that the result is qualitatively sharp in some sense.*

Remark 4.6. *Both the arguments for the lower and upper bound hold for the sequence of energies*

$$\tilde{\mathcal{E}}_{\varepsilon}(u) = \frac{1}{2} \int_{\Omega} \left[\frac{1}{\varepsilon} \left(\partial_z u - \frac{1}{2} |\nabla_{\perp} u|^2 \right)^2 + \varepsilon |\nabla_{\perp}^2 u|^2 \right] \, d\mathbf{x}$$

with trivial modifications. For lower bound, we only need to assume $\liminf \tilde{\mathcal{E}}_{\varepsilon_n}(u_n)$ is finite.

5 Compactness

The main result in this section is the compactness theorem.

Theorem 5.1. *Let $\Omega \subset \mathbb{R}^3$ be a bounded domain with C^1 boundary, $\varepsilon_n \rightarrow 0$, and $\{u_n\} \subset H^2(\Omega)$ be a sequence of functions with uniformly bounded energies $\mathcal{E}_{\varepsilon_n}(u_n)$ such that $\|\nabla u_n\|_{L^p(\Omega)} \leq C$ for some $p > 6$ and $\|\nabla u_n\|_{L^2(\partial\Omega)} \leq C$. Assume also that $\Delta_{\perp} u_n \geq 0$ or $\Delta_{\perp} u_n \leq 0$ a.e. in Ω . Then ∇u_n is precompact in $L^q(\Omega)$ for any $1 \leq q < p$.*

Theorem 5.1 is a direct corollary of the following stronger proposition.

Proposition 5.2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded domain with C^1 boundary and $\{u_n\} \subset H^2(\Omega)$ be a sequence of functions satisfying*

$$\|\nabla u_n\|_{L^p(\Omega)} \leq C, \text{ for some } p > 6, \quad (5.1)$$

$$\|\nabla u_n\|_{L^2(\partial\Omega)} \leq C, \quad (5.2)$$

$$\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \rightarrow 0 \text{ strongly in } L^2(\Omega), \quad (5.3)$$

and

$$\left| \partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right| |\Delta_{\perp} u_n| \text{ is bounded in } L^1(\Omega). \quad (5.4)$$

If in addition

$$\Delta_{\perp} u_n \geq 0 \text{ a.e. in } \Omega \text{ or } \Delta_{\perp} u_n \leq 0 \text{ a.e. in } \Omega, \quad (5.5)$$

then $(\nabla_{\perp} u_n)$ is precompact in $L^q(\Omega)$ for $1 \leq q < p$.

We first prove a lemma used in the proof of Proposition 5.2.

Lemma 5.3. *Under the assumptions (5.1) – (5.5), $\operatorname{div} B_n$ is relatively compact in $H^{-1}(\Omega)$, where*

$$B_n = \left(-\frac{\nabla_{\perp} u_n}{2} |\nabla_{\perp} u_n|^2, \frac{1}{2} |\nabla_{\perp} u_n|^2 \right).$$

Proof. We prove the Lemma when u_n is smooth and general case follows by approximating. By (5.3),

$$\begin{aligned} & \partial_z (\partial_x u_n) - \partial_x \left(\frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \\ &= \partial_x \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \rightarrow 0 \text{ strongly in } H^{-1}(\Omega), \quad \text{and} \end{aligned} \quad (5.6)$$

$$\begin{aligned} & \partial_z (\partial_y u_n) - \partial_y \left(\frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \\ &= \partial_y \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \rightarrow 0 \text{ strongly in } H^{-1}(\Omega). \end{aligned} \quad (5.7)$$

Multiplying (5.6) by $\partial_x u_n$ and (5.7) by $\partial_y u_n$ then summing, we have

$$\begin{aligned} & \partial_z \left(\frac{1}{2} |\nabla_{\perp} u_n|^2 \right) - \partial_x \left(\frac{\partial_x u_n}{2} |\nabla_{\perp} u_n|^2 \right) - \partial_y \left(\frac{\partial_y u_n}{2} |\nabla_{\perp} u_n|^2 \right) \\ &= \partial_x \left(\partial_x u_n \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \right) + \partial_y \left(\partial_y u_n \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \right) \\ & \quad - \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \Delta_{\perp} u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \Delta_{\perp} u_n \\ &= I + II + III. \end{aligned} \quad (5.8)$$

Here

$$I = \partial_x \left(\partial_x u_n \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \right) + \partial_y \left(\partial_y u_n \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \right) \rightarrow 0$$

in $W^{-1, \frac{2p}{p+2}}(\Omega)$ up to a subsequence,

$$II = - \left(\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \right) \Delta_{\perp} u_n$$

is bounded in $L^1(\Omega)$, and

$$III = -\frac{1}{2} |\nabla_{\perp} u_n|^2 \Delta_{\perp} u_n.$$

If III is bounded in $\mathcal{M}(\Omega)$, the space of measures, then the right hand side of (5.8) is the sum of a term relatively compact in $W^{-1, \frac{2p}{p+2}}(\Omega)$ and a term bounded in $\mathcal{M}(\Omega)$, so that by the embedding theorem, the right hand side of (5.8) is relatively compact in $W^{-1, r}(\Omega)$ for some $1 \leq r < 2$. On the other hand, assumption (5.1) implies the left hand side of (5.8) is bounded in $W^{-1, \frac{p}{3}}(\Omega)$. Relative compactness of $\operatorname{div} B_n$ in $H^{-1}(\Omega)$ follows from interpolation.

To finish the proof, we show III is bounded in $\mathcal{M}(\Omega)$ under the additional assumption (5.5). Rewrite (5.8) as

$$\partial_z(\eta(u_n)) - \operatorname{div}_\perp(F(u_n)) + G(u_n) = \mu_n, \quad (5.9)$$

where

$$\begin{aligned} \eta(u_n) &= \frac{1}{2} |\nabla_\perp u_n|^2, \quad F(u_n) = (\partial_x u_n \partial_z u_n, \partial_y u_n \partial_z u_n), \\ G(u_n) &= \left(\partial_z u_n - \frac{1}{2} |\nabla_\perp u_n|^2 \right) \Delta_\perp u_n, \\ \mu_n &= -\frac{1}{2} |\nabla_\perp u_n|^2 \Delta_\perp u_n. \end{aligned}$$

Let $\Omega^- = \{\mathbf{x} \in \Omega : \mu_n(\mathbf{x}) \leq 0\}$ and $\Omega^+ = \{\mathbf{x} \in \Omega : \mu_n(\mathbf{x}) \geq 0\}$. Since (5.5) holds, then $\Omega = \Omega^-$ or $\Omega = \Omega^+$. Assume $\Omega = \Omega^-$ (the other case can be proved similarly). Integrating (5.9) over Ω and using the divergence theorem on the first two terms yields

$$\begin{aligned} \int_\Omega -\mu_n dx dy dz &= \int_{\partial\Omega} -\eta(u_n) \nu_3 d\mathcal{H}^2 + \int_{\partial\Omega} F(u_n) \cdot \nu_\perp d\mathcal{H}^2 - \int_\Omega G(u_n) d\mathbf{x} \\ &\leq \|\nabla_\perp u\|_{L^2(\partial\Omega)}^2 + \|\partial_z u\|_{L^2(\partial\Omega)} \|\nabla_\perp u\|_{L^2(\partial\Omega)} + \mathcal{E}_{\varepsilon_n}(u_n) \\ &\leq C \end{aligned}$$

for some constant C depending on the energy bound and $\|\nabla u_n\|_{L^2(\partial\Omega)}$. Therefore $-\mu_n \mathcal{L}^3 \llcorner \Omega$ is bounded in $\mathcal{M}(\Omega)$. \square

Remark 5.4.

- A special case satisfying (5.1) is $\|\nabla u_n\|_{L^\infty(\Omega)} \leq C$.
- In the proof of Lemma 5.3, the convergence result in (5.6) and (5.7) has not been used; however, it is used below in the proof of Proposition 5.2.

Proof of Proposition 5.2. Set

$$E_n = \left(\nabla_\perp u_n, \frac{1}{2} |\nabla_\perp u_n|^2 \right) \text{ and } B_n = \left(-\frac{\nabla_\perp u_n}{2} |\nabla_\perp u_n|^2, \frac{1}{2} |\nabla_\perp u_n|^2 \right).$$

Lemma 5.3 together with (5.6) and (5.7) implies

$$\operatorname{curl} E_n \text{ and } \operatorname{div} B_n \text{ are relatively compact in } H^{-1}(\Omega).$$

If $E_n \rightharpoonup E_\infty$, $B_n \rightharpoonup B_\infty$ in $L^2(\Omega)$, then Tartar-Murat's div-curl Lemma applied to E_n and B_n yields

$$E_n \cdot B_n \rightharpoonup E_\infty \cdot B_\infty \text{ in } \mathcal{D}'(\Omega). \quad (5.10)$$

We introduce the following notations for the weak limits in $L^r(\Omega)$ for $r > 1$:

$$\begin{aligned} (\partial_x u_n)^4 &\rightharpoonup U_4, \quad (\partial_y u_n)^4 \rightharpoonup V_4, \quad (\partial_x u_n)^3 \rightharpoonup U_3, \quad (\partial_y u_n)^3 \rightharpoonup V_3, \\ (\partial_x u_n)^2 \partial_y u_n &\rightharpoonup U_{21}, \quad \partial_x u_n (\partial_y u_n)^2 \rightharpoonup U_{12}, \quad (\partial_x u_n)^2 \rightharpoonup U_2, \quad (\partial_y u_n)^2 \rightharpoonup V_2, \end{aligned}$$

and

$$(\partial_x u_n)^2 (\partial_y u_n)^2 \rightharpoonup U_{22}, \quad \partial_x u_n \partial_y u_n \rightharpoonup U_{11}, \quad \partial_x u_n \rightharpoonup U_1, \quad \partial_y u_n \rightharpoonup V_1.$$

Here r depends on p and the term in question but is greater than 1 for each. Under these notations, (5.10) can be written as

$$-\frac{1}{4} |\nabla_{\perp} u_n|^4 \rightharpoonup \left(U_1, V_1, \frac{1}{2} (U_2 + V_2) \right) \cdot \left(-\frac{1}{2} U_3 - \frac{1}{2} U_{12}, -\frac{1}{2} U_{21} - \frac{1}{2} V_3, \frac{1}{2} (U_2 + V_2) \right).$$

From this it follows that

$$-\frac{1}{4} (U_4 + 2U_{22} + V_4) = -\frac{1}{2} U_1 U_3 - \frac{1}{2} U_1 U_{12} - \frac{1}{2} V_1 U_{21} - \frac{1}{2} V_1 V_3 + \frac{1}{4} (U_2 + V_2)^2,$$

or, equivalently,

$$0 = U_4 + 2U_{22} + V_4 - 2U_1 U_3 - 2U_1 U_{12} - 2V_1 U_{21} - 2V_1 V_3 + (U_2 + V_2)^2. \quad (5.11)$$

Next, we consider

$$\begin{aligned} &\left((\partial_x u_n)^2 - \partial_x u_n U_1 + (\partial_y u_n)^2 - \partial_y u_n V_1 \right)^2 \\ &= (\partial_x u_n)^2 (\partial_x u_n - U_1)^2 + 2\partial_x u_n \partial_y u_n (\partial_x u_n - U_1) (\partial_y u_n - V_1) + (\partial_y u_n)^2 (\partial_y u_n - V_1)^2 \\ &\rightharpoonup U_4 - 2U_1 U_3 + U_2 U_1^2 + 2(U_{22} - U_{21} V_1 - U_1 U_{12} + U_{11} U_1 V_1) + V_4 - 2V_1 V_3 + V_2 V_1^2 \\ &\stackrel{(5.11)}{=} -(U_2 + V_2)^2 + U_2 U_1^2 + V_2 V_1^2 + 2U_{11} U_1 V_1. \end{aligned}$$

Observe that

$$\begin{aligned} (\partial_x u_n U_1 + \partial_y u_n V_1 - U_2 - V_2)^2 &= (U_2 + V_2)^2 - 2(\partial_x u_n U_1 + \partial_y u_n V_1)(U_2 + V_2) + (\partial_x u_n U_1 + \partial_y u_n V_1)^2 \\ &\rightharpoonup (U_2 + V_2)^2 - 2(U_2 + V_2)(U_1^2 + V_1^2) + U_2 U_1^2 + V_2 V_1^2 + 2U_{11} U_1 V_1 \\ &\leq (U_2 + V_2)^2 - U_2 U_1^2 - V_2 V_1^2 - 2U_{11} U_1 V_1, \end{aligned} \quad (5.12)$$

where the last inequality follows from

$$U_2 V_1^2 + V_2 U_1^2 \geq 2U_{11} U_1 V_1,$$

which is a direct conclusion from weak limit

$$(\partial_x u_n V_1 - \partial_y u_n U_1)^2 \rightharpoonup U_2 V_1^2 + V_2 U_1^2 - 2U_{11} U_1 V_1.$$

Thus

$$\left((\partial_x u_n)^2 - \partial_x u_n U_1 + (\partial_y u_n)^2 - \partial_y u_n V_1 \right)^2 \rightharpoonup -(U_2 + V_2)^2 + U_2 U_1^2 + V_2 V_1^2 + 2U_{11} U_1 V_1 \stackrel{(5.12)}{\leq} 0.$$

From this we conclude that

$$(\partial_x u_n)^2 - \partial_x u_n U_1 + (\partial_y u_n)^2 - \partial_y u_n V_1 \rightarrow 0,$$

so that passing to the limit on the left side, we have

$$U_2 - U_1^2 + V_2 - V_1^2 = 0.$$

Since $F(s) = s^2$ is a convex function, by Lemma 2 in [60], we have $U_2 \geq U_1^2$, $V_2 \geq V_1^2$. Thus it must be the case that

$$U_2 = U_1^2, \quad V_2 = V_1^2;$$

in other words, $\lim_{n \rightarrow \infty} \|\nabla_{\perp} u_n\|_{L^2} = \|\lim_n \nabla_{\perp} u_n\|_{L^2}$, together with the weak convergence of $\nabla_{\perp} u_n$, the strong convergence of $\nabla_{\perp} u_n$ in L^2 follows. \square

Proof of Theorem 5.1. Boundedness of $\mathcal{E}_{\varepsilon_n}(u_n)$ implies (5.3) and (5.4). By Proposition 5.2, $\nabla_{\perp} u_n$ is precompact in $L^2(\Omega)$, and hence in $L^q(\Omega)$ for $1 \leq q < p$ by the uniform L^p bound. Compactness of $\partial_z u_n$ follows from uniform boundedness of $\|\nabla_{\perp} u_n\|_{L^p(\Omega)}$ for $p > 6$ and the fact $\partial_z u_n - \frac{1}{2} |\nabla_{\perp} u_n|^2 \rightarrow 0$ in L^2 . \square

Remark 5.5. *The additional assumptions (5.1), (5.2), (5.5) in Theorem 5.1 and Proposition 5.2 are used in the proof of relative compactness of $\operatorname{div} B_n$ in Lemma 5.3. We comment that the assumption (5.1) is physically justifiable since the model is only valid in the limit of small strains [55]. Assumption (5.2) is less restrictive than $\varepsilon_n \int_{\Omega} \bar{K} \rightarrow 0$. We would like to remove the technical assumption (5.5) in future work. An alternative approach to handling compactness is to rewrite the problem in terms of the geometric formulation of Tartar's conjecture [61]. Recall that the general question regarding upgrading weak convergence to strong convergence can be stated as follows: given a weakly convergent sequence of functions $z^\varepsilon : \mathbb{R}^m \rightarrow \mathbb{R}^N$ subject to linear differential constraints of the form*

$$\sum_{j=0}^{m-1} A_j \partial_j z^\varepsilon = \varphi^\varepsilon, \quad A_j \text{ a } s \times N \text{ constant matrix}, \quad (5.13)$$

and nonlinear algebraic constraints

$$\{z^\varepsilon(y)\} \subset M \quad \text{for a.e. } y \in \mathbb{R}^m, \quad (5.14)$$

where $M \subset \mathbb{R}^N$ is a subset, usually a manifold, what kind of structure on A_j and M would suppress oscillations in $\{z^\varepsilon\}$, so that $\{z^\varepsilon\}$ contains a strongly convergent subsequence? Tartar's conjecture can be expressed in terms of a geometric condition. We introduce the oscillation variety

$$V = \left\{ (\xi, \lambda) : \sum_{j=0}^{m-1} \xi_j A_j = 0, \quad \xi \neq 0 \right\} \subset \mathbb{R}^m \times \mathbb{R}^N,$$

and the wave cone, which is the projection of V to \mathbb{R}^N :

$$\Lambda = P_N V = \{\lambda : \exists \xi \neq 0, \text{ such that } (\xi, \lambda) \in V\}.$$

Given any a , let

$$\Lambda_a = a + \Lambda = \{a + \lambda, \lambda \in \Lambda\}$$

be the translated cone. Tartar conjectured:

Conjecture 1. *If the translated wave cone is separated from M in the sense that*

$$\Lambda_a \cap M = \{a\}$$

for all a , then the Young measure ν_x is a Dirac mass for almost every x , which implies the relative compactness in L^p

For a sequence with bounded energy (1.1), one may form a wave cone from (5.6), (5.7), and $\operatorname{curl} \nabla_{\perp} u = 0$ and construct a constitutive manifold from suitable entropy conditions. Our initial observation [46] shows the translated wave cone constructed this way is separated from the constitutive manifold. The final conclusion regarding compactness is still under investigation.

ACKNOWLEDGEMENTS We thank Robert V. Kohn for bringing this problem into our interest. M.N.'s research is supported by the NSF grant RTG-DMS 1840314. X.Y's research is supported by a Research Excellence Grant and a CLAS Dean's summer research grant from University of Connecticut. X.Y also thanks Fengbo Hang for helpful discussions on Tartar's compensated compactness results. Both authors thank the anonymous referee for the thorough review and helpful comments.

DATA AVAILABILITY STATEMENT Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- [1] F. ALOUGES, T. RIVIÈRE, AND S. SERFATY, *Néel and cross-tie wall energies for planar micromagnetic configurations*, vol. 8, 2002, pp. 31–68. A tribute to J. L. Lions.
- [2] L. AMBROSIO AND G. DAL MASO, *A general chain rule for distributional derivatives*, Proc. Amer. Math. Soc., 108 (1990), pp. 691–702.
- [3] L. AMBROSIO, C. DE LELLIS, AND C. MANTEGAZZA, *Line energies for gradient vector fields in the plane*, Calc. Var. Partial Differential Equations, 9 (1999), pp. 327–255.
- [4] L. AMBROSIO, N. FUSCO, AND D. PALLARA, *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 2000.
- [5] P. AVILES AND Y. GIGA, *A mathematical problem related to the physical theory of liquid crystal configurations*, in Miniconference on geometry and partial differential equations, 2 (Canberra, 1986), vol. 12 of Proc. Centre Math. Anal. Austral. Nat. Univ., Austral. Nat. Univ., Canberra, 1987, pp. 1–16.
- [6] ———, *On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields*, Proc. Roy. Soc. Edinburgh Sect. A, 129 (1999), pp. 1–17.
- [7] J. BENOIT, A. SAXENA, AND T. LOOKMAN, *Bogomol'nyi decomposition for vesicles of arbitrary genus*, Journal of Physics A: Mathematical and General, 34 (2001), pp. 9417–9423.
- [8] I. BLUESTEIN AND R. D. KAMIEN, *Nonlinear effects in the TGB a phase*, Europhysics Letters (EPL), 59 (2002), pp. 68–74.
- [9] E. B. BOGOMOL'NYI, *The stability of classical solutions*, Sov. J. Nucl. Phys. (Engl. Transl.); (United States), 24 (1976).

- [10] E. A. BRENER AND V. I. MARCHENKO, *Nonlinear theory of dislocations in smectic crystals: An exact solution*, Phys. Rev. E, 59 (1999), pp. R4752–R4753.
- [11] P. M. CHAIKIN AND T. C. LUBENSKY, *Principles of Condensed Matter Physics*, Cambridge University Press, 1995.
- [12] S. CONTI AND C. DE LELLIS, *Sharp upper bounds for a variational problem with singular perturbation*, Math. Ann., 338 (2007), pp. 119–146.
- [13] C. DE LELLIS, *An example in the gradient theory of phase transitions*, ESAIM: Control, Optimisation and Calculus of Variations, 7 (2002), pp. 285–289.
- [14] C. DE LELLIS AND R. IGNAT, *A regularizing property of the 2d-eikonal equation*, Communications in Partial Differential Equations, 40 (2015), pp. 1543–1557.
- [15] C. DE LELLIS AND F. OTTO, *Structure of entropy solutions to the eikonal equation*, J. Eur. Math. Soc. (JEMS), 5 (2003), pp. 107–145.
- [16] A. DESIMONE, S. MÜLLER, R. V. KOHN, AND F. OTTO, *A compactness result in the gradient theory of phase transitions*, Proceedings of the Royal Society of Edinburgh: Section A Mathematics, 131 (2001), p. 833–844.
- [17] B. A. DIDONNA AND R. D. KAMIEN, *Smectic phases with cubic symmetry: The splay analog of the blue phase*, Phys. Rev. Lett., 89 (2002), p. 215504.
- [18] ———, *Smectic blue phases: layered systems with high intrinsic curvature*, Phys. Rev. E (3), 68 (2003), pp. 041703, 11.
- [19] M. P. DO CARMO, *Differential geometry of curves and surfaces*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1976. Translated from the Portuguese.
- [20] P. D. GENNES AND J. PROST, *The physics of liquid crystals*, International series of monographs on physics (Oxford, England); 83, Clarendon Press ; Oxford University Press, Oxford : New York, 2nd ed.. ed., 1993.
- [21] F. GHIRALDIN AND X. LAMY, *Optimal Besov differentiability for entropy solutions of the eikonal equation*, Communications on Pure and Applied Mathematics, 73 (2020), pp. 317–349.
- [22] L. GOLUBOVIĆ AND Z.-G. WANG, *Kardar-Parisi-Zhang model and anomalous elasticity of two- and three-dimensional smectic-a liquid crystals*, Phys. Rev. E, 49 (1994), pp. 2567–2578.
- [23] W. HELFRICH, *Electrohydrodynamic and dielectric instabilities of cholesteric liquid crystals*, The Journal of Chemical Physics, 55 (1971), pp. 839–842.
- [24] J. P. HURAUULT, *Static distortions of a cholesteric planar structure induced by magnetic or ac electric fields*, The Journal of Chemical Physics, 59 (1973), pp. 2068–2075.
- [25] R. IGNAT, *Singularities of divergence-free vector fields with values into \mathbb{S}^1 or \mathbb{S}^2 : Applications to micromagnetics*, Confluentes Mathematici, 04 (2012), p. 1230001.
- [26] R. IGNAT, *Two-dimensional unit-length vector fields of vanishing divergence*, Journal of Functional Analysis, 262 (2012), pp. 3465–3494.

- [27] R. IGNAT AND B. MERLET, *Lower bound for the energy of Bloch walls in micromagnetics*, Arch. Ration. Mech. Anal., 199 (2011), pp. 369–406.
- [28] ———, *Entropy method for line-energies*, Calc. Var. Partial Differential Equations, 44 (2012), pp. 375–418.
- [29] R. IGNAT AND A. MONTEIL, *A De Giorgi-type conjecture for minimal solutions to a nonlinear stokes equation*, Communications on Pure and Applied Mathematics, 73 (2020), pp. 771–854.
- [30] T. ISHIKAWA AND O. D. LAVRENTOVICH, *Dislocation profile in cholesteric finger texture*, Phys. Rev. E, 60 (1999), pp. R5037–R5039.
- [31] W. JIN AND R. V. KOHN, *Singular perturbation and the energy of folds*, J. Nonlinear Sci., 10 (2000), pp. 355–390.
- [32] R. D. KAMIEN AND T. C. LUBENSKY, *Minimal surfaces, screw dislocations, and twist grain boundaries*, Phys. Rev. Lett., 82 (1999), pp. 2892–2895.
- [33] R. D. KAMIEN AND C. D. SANTANGELO, *Smectic liquid crystals: materials with one-dimensional, periodic order*, Geom. Dedicata, 120 (2006), pp. 229–240.
- [34] M. KLÉMAN, *Points, lines and walls in liquid crystals, magnetic systems and various ordered media*, Oxford Mathematical Monographs, John Wiley and Sons, New York, 1983.
- [35] R. V. KOHN, *Energy-driven pattern formation*, in International Congress of Mathematicians. Vol. I, Eur. Math. Soc., Zürich, 2007, pp. 359–383.
- [36] X. LAMY, A. LORENT, AND G. PENG, *Rigidity of a non-elliptic differential inclusion related to the Aviles-Giga conjecture*, Arch. Ration. Mech. Anal., 238 (2020), pp. 383–413.
- [37] A. LORENT, *A simple proof of the characterization of functions of low Aviles Giga energy on a ball via regularity*, ESAIM Control Optim. Calc. Var., 18 (2012), pp. 383–400.
- [38] ———, *A quantitative characterisation of functions with low Aviles Giga energy on convex domains*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 13 (2014), pp. 1–66.
- [39] A. LORENT AND G. PENG, *Regularity of the eikonal equation with two vanishing entropies*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 35 (2018), pp. 481–516.
- [40] ———, *Factorization for entropy production of the eikonal equation and regularity*, arXiv e-prints, (2021), p. arXiv:2104.01467.
- [41] E. MARCONI, *Characterization of minimizers of Aviles-Giga functionals in special domains*, arXiv e-prints, (2021), p. arXiv:2104.07125.
- [42] E. A. MATSUMOTO, C. D. SANTANGELO, AND R. D. KAMIEN, *Smectic pores and defect cores*, J. Interface Focus, 2 (2012), pp. 617–622.
- [43] I. MITREA, M. MITREA, AND M. WRIGHT, *Optimal estimates for the inhomogeneous problem for the bi-Laplacian in three-dimensional Lipschitz domains*, vol. 172, 2011, pp. 24–134. Problems in mathematical analysis. No. 51.
- [44] F. MURAT, *Compacité par compensation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 5 (1978), pp. 489–507.

- [45] ———, *Compacité par compensation: condition nécessaire et suffisante de continuité faible sous une hypothèse de rang constant*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 8 (1981), pp. 69–102.
- [46] M. NOVACK AND X. YAN, *Work in progress*.
- [47] ———, *Compactness and sharp lower bound for a 2d smectics model*, to appear in J. Non. Sci., (2021).
- [48] A. POLIAKOVSKY, *Upper bounds for singular perturbation problems involving gradient fields*, J. Eur. Math. Soc. (JEMS), 9 (2007), pp. 1–43.
- [49] ———, *A general technique to prove upper bounds for singular perturbation problems*, J. Anal. Math., 104 (2008), pp. 247–290.
- [50] ———, *On the Γ -limit of singular perturbation problems with optimal profiles which are not one-dimensional. Part I: The upper bound*, Differential Integral Equations, 26 (2013), pp. 1179–1234.
- [51] M. PRASAD AND C. M. SOMMERFIELD., *Solutions of classical gauge field theories with spin and internal symmetry*, Nuclear Physics B, 110 (1976), pp. 153 – 172.
- [52] T. RIVIÈRE AND S. SERFATY, *Limiting domain wall energy for a problem related to micromagnetics*, Comm. Pure Appl. Math., 54 (2001), pp. 294–338.
- [53] ———, *Compactness, kinetic formulation, and entropies for a problem related to micromagnetics*, Comm. Partial Differential Equations, 28 (2003), pp. 249–269.
- [54] C. D. SANTANGELO, *Geometry and the nonlinear elasticity of defects in smectic liquid crystals*, Liquid Crystals Today, 15 (2006), pp. 11–18.
- [55] C. D. SANTANGELO AND R. D. KAMIEN, *Bogomol’nyi, Prasad, and Sommerfield configurations in smectics*, Phys. Rev. Lett., 91 (2003), p. 045506.
- [56] C. D. SANTANGELO AND R. D. KAMIEN, *Curvature and topology in smectic-A liquid crystals*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci., 461 (2005), pp. 2911–2921.
- [57] ———, *Triply periodic smectic liquid crystals*, Phys. Rev. E (3), 75 (2007), pp. 011702, 12.
- [58] I. I. SMALYUKH AND O. D. LAVRENTOVICH, *Anchoring-mediated interaction of edge dislocations with bounding surfaces in confined cholesteric liquid crystals*, Phys. Rev. Lett., 90 (2003), p. 085503.
- [59] P. STERNBERG, *The effect of a singular perturbation on nonconvex variational problems*, Arch. Rational Mech. Anal., 101 (1988), pp. 209–260.
- [60] L. TARTAR, *Compensated compactness and applications to partial differential equations*, in Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV, vol. 39 of Res. Notes in Math., Pitman, Boston, Mass.-London, 1979, pp. 136–212.
- [61] L. TARTAR, *The compensated compactness method applied to systems of conservation laws*, in Systems of nonlinear partial differential equations (Oxford, 1982), vol. 111 of NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., Reidel, Dordrecht, 1983, pp. 263–285.

- [62] ———, *Compensation effects in partial differential equations*, Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5), 29 (2005), pp. 395–453.
- [63] A. I. VOL'PERT, *Spaces BV and quasilinear equations*, Mat. Sb. (N.S.), 73 (115) (1967), pp. 255–302.