

STABILITY FOR THE SURFACE DIFFUSION FLOW

ANTONIA DIANA, NICOLA FUSCO, AND CARLO MANTEGAZZA

ABSTRACT. We study the global existence and stability of surface diffusion flow (the normal velocity is given by the Laplacian of the mean curvature) of smooth boundaries of subsets of the n -dimensional flat torus. More precisely, we show that if a smooth set is “close enough” to a strictly stable critical set for the Area functional under a volume constraint, then the surface diffusion flow of its boundary hypersurface exists for all time and asymptotically converges to the boundary of a “translated” of the critical set. This result was obtained in dimension $n = 3$ by Acerbi, Fusco, Julin and Morini in [1] (extending previous results for spheres of Escher, Mayer and Simonett [15], Wheeler [37, 38] and Elliott and Garcke [14]). Our work generalizes such conclusion to any dimension $n \in \mathbb{N}$. For sake of clarity, we show all the details in dimension $n = 4$ and we list the necessary modifications to the quantities involved in the proof in the general n -dimensional case, in the last section.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. The surface diffusion flow and strictly stable critical sets	7
4. Evolution of geometric quantities and basic estimates	10
5. Global existence and stability	23
6. General dimensions	32
References	35

1. INTRODUCTION

Given a smooth immersed hypersurface in an n -dimensional flat torus $\varphi = \varphi_0 : M \rightarrow \mathbb{T}^n$ (or in \mathbb{R}^n), we say that a smooth family of smooth embeddings $\varphi_t : M \rightarrow \mathbb{T}^n$, for $t \in [0, T)$, is a *surface diffusion flow* for φ_0 if

$$\frac{\partial \varphi_t}{\partial t} = (\Delta H)\nu, \quad (1.1)$$

that is, the outer normal velocity (here ν is the outer normal) of the moving hypersurfaces is given by the Laplacian (in the induced metric) ΔH of the mean curvature, at every point and time. Such flow was first proposed by Mullins in [33] to study thermal grooving in material sciences and first analyzed mathematically more in detail in [15].

Date: October 23, 2023.

In particular, in the physically relevant case of three-dimensional space, it describes the evolution of interfaces between solid phases of a system, driven by surface diffusion of atoms under the action of a chemical potential (see for instance [25]).

Studying the flow in a flat torus \mathbb{T}^n , described as the quotient of \mathbb{R}^n by a discrete group of translations generated by some n linearly independent vectors, is equivalent to consider the flow of “periodic” hypersurfaces, invariant by such group of translations. Then, it is clear that our analysis also applies to compact hypersurfaces in \mathbb{R}^n or, more in general, in any (generalized) “cylinder” $\mathbb{S}^1 \times \dots \times \mathbb{S}^1 \times \mathbb{R} \times \dots \times \mathbb{R}$ of dimension n , with a flat metric.

Notice that, by the general equality $\Delta\varphi = -H\nu$ (see equation (2.2) below), the system (1.1) can be rewritten as

$$\frac{\partial\varphi_t}{\partial t} = -\Delta\Delta\varphi_t + \text{lower order terms} \quad (1.2)$$

hence, it is a fourth order, *quasilinear* and *degenerate*, parabolic system of PDEs. Indeed, it is quasilinear, as the coefficients (as second order partial differential operator) of the Laplacian associated to the induced metrics on the evolving hypersurfaces, depend on the first order derivatives of φ_t (and the coefficient of $\Delta\Delta$ on the third order derivatives) and the operator at the right hand side of system (1.1) is degenerate, as its symbol (the symbol of the linearized operator) admits zero eigenvalues, due to the invariance of the Laplacian by diffeomorphisms. The lack of maximum principle, as the flow is of fourth order, implies that it does not preserve convexity (see [27]), nor the embeddedness (see [20]), indeed it also does not have a “comparison principle”, while it is invariant by isometries of \mathbb{T}^n , reparametrizations and tangential perturbations of the velocity of the motion. Moreover, when it is restricted to closed embedded hypersurfaces which are boundaries of sets, the enclosed volume is preserved and actually it can be regarded as the \tilde{H}^{-1} -gradient flow of the *volume-constrained Area functional* (see [10] or [19], for instance).

There holds the following short time existence and uniqueness theorem (and also of dependence on the initial data) for the surface diffusion flow starting from a smooth hypersurface, proved by Escher, Mayer and Simonett in [15], which should be expected by the parabolic nature of the system (1.1), as shown by formula (1.2). The original result deals with the evolution in the whole space \mathbb{R}^n of a generic hypersurface, even only immersed, hence possibly with self-intersections. It is anyway straightforward to adapt the same arguments to our case, when the ambient is a flat torus \mathbb{T}^n (and the hypersurfaces are boundaries of sets).

Theorem 1.1. *Let $\varphi_0 : M \rightarrow \mathbb{R}^n$ be a smooth and compact, immersed hypersurface. Then, there exists a unique smooth $\varphi : [0, T) \times M \rightarrow \mathbb{R}^n$ such that $\varphi_t = \varphi(t, \cdot)$ is the surface diffusion flow of φ_0 , that is a solution of equation (1.1), for some maximal time of existence $T > 0$. Moreover, such flow and the maximal time of existence depend continuously on the $C^{2,\alpha}$ -norm of the initial hypersurface φ_0 .*

Actually, it is very likely, even if an explicit example is not present in literature (up to our knowledge), that this flow could develop singularities in finite time (as the mean curvature flow). Anyway, in the same mentioned paper [15] by Escher, Mayer and Simonett, the authors also showed that if the initial hypersurface is $C^{2,\alpha}$ -close enough to a sphere with the same enclosed volume, then the flow exists for every time and smoothly converges to a translate of such sphere. The analogous result was obtained by Wheeler in [37], for surfaces and in [38], for closed plane curves (see also the work of Elliott and Garcke [14] for curves) with a weaker initial $W^{2,2}$ -closedness condition and by Escher and Mucha in a previous work [16] with a Besov-type condition. Furthermore, in [39] Wheeler shows that any surface diffusion flow of curves that exists for all time, must converge smoothly, exponentially fast to a multiply-covered circle. We also mention a work by Miura e Okabe [32] where the authors prove a global existence result provided that an initial curve is $W^{2,2}$ -close to a multiply covered circle and sufficiently rotationally symmetric. Later on, Acerbi, Fusco, Julin and Morini in [1] extended these results, in dimensions two and three, to hypersurfaces close to boundaries of *strictly stable critical sets* (that we are going to define in a while) for the volume constrained Area functional (as it is every ball). Our aim in this work is to generalize such stability conclusion to any dimension $n \in \mathbb{N}$. Due to some necessary heavy analytic/algebraic computations that follows, we are going to show all the details of the proof in dimension $n = 4$ and we list the appropriate modifications for the general n -dimensional case, in the last section. The explicit and detailed computations and proofs in such general case will appear in the PhD thesis of Antonia Diana [13].

2. PRELIMINARIES

We introduce the basic notations and facts about hypersurfaces that we need in the paper, possible references are [18] or the first part of [34].

We will consider closed smooth hypersurfaces in the n -dimensional torus $\mathbb{T}^n \approx \mathbb{R}^n/\mathbb{Z}^n$ or in \mathbb{R}^n , given by smooth immersions $\varphi : M \rightarrow \mathbb{T}^n$ of a smooth, $(n - 1)$ -dimensional, compact manifold M , representing a hypersurface $\varphi(M)$ of \mathbb{T}^n . Taking local coordinates around any $p \in M$, we have local bases of the tangent space $T_p M$, which can be identified with the $(n - 1)$ -dimensional hyperplane $d\varphi_p(T_p M)$ of $\mathbb{R}^n \approx T_{\varphi(p)}\mathbb{T}^n$ which is tangent to $\varphi(M)$ at $\varphi(p)$ and of the cotangent space $T_p^* M$, respectively given by vectors $\{\frac{\partial}{\partial x_i}\}$ and 1-forms $\{dx_j\}$. So, we will denote vectors on M by $X = X^i$, which means $X = X^i \frac{\partial}{\partial x_i}$, covectors by $Y = Y_j$, that is, $Y = Y_j dx_j$ and a general mixed tensor with $T = T_{j_1 \dots j_l}^{i_1 \dots i_k}$.

In the whole paper the convention to sum over repeated indices will be adopted.

Sometimes we will need also to consider tensors along M , viewing it as a submanifold of \mathbb{T}^n or \mathbb{R}^n via the map φ , in that case we will use the Greek indices to denote the components of such tensors in the canonical basis $\{e_\alpha\}$ of \mathbb{R}^n , for instance, given a vector field X along M , not necessarily tangent, we will have $X = X^\alpha e_\alpha$.

The manifold M gets in a natural way a metric tensor g , pull-back via the map φ of the metric tensor of \mathbb{T}^n , coming from the standard scalar product $\langle \cdot | \cdot \rangle$ of \mathbb{R}^n , hence, turning it into a Riemannian manifold (M, g) . Then, the components of g in a local chart are

$$g_{ij} = \left\langle \frac{\partial \varphi}{\partial x_i} \middle| \frac{\partial \varphi}{\partial x_j} \right\rangle$$

and the ‘‘canonical’’ measure μ , induced on M by the metric g is then locally described by $\mu = \sqrt{\det g_{ij}} \mathcal{L}^{n-1}$, where \mathcal{L}^{n-1} is the standard Lebesgue measure on \mathbb{R}^{n-1} . The inner product on M , extended to tensors, is given by

$$g(T, S) = g_{i_1 s_1} \cdots g_{i_k s_k} g^{j_1 z_1} \cdots g^{j_l z_l} T_{j_1 \dots j_l}^{i_1 \dots i_k} S_{z_1 \dots z_l}^{s_1 \dots s_k}$$

where g_{ij} is the matrix of the coefficients of the metric tensor in the local coordinates and g^{ij} is its inverse. Clearly, the norm of a tensor is then

$$|T| = \sqrt{g(T, T)}.$$

The induced Levi-Civita covariant derivative on (M, g) of a vector field X and of a 1-form ω are respectively given by

$$\nabla_j X^i = \frac{\partial X^i}{\partial x_j} + \Gamma_{jk}^i X^k, \quad \nabla_j \omega_i = \frac{\partial \omega_i}{\partial x_j} - \Gamma_{ji}^k \omega_k,$$

where Γ_{jk}^i are the Christoffel symbols of the connection ∇ , expressed by the formula

$$\Gamma_{jk}^i = \frac{1}{2} g^{il} \left(\frac{\partial}{\partial x_j} g_{kl} + \frac{\partial}{\partial x_k} g_{jl} - \frac{\partial}{\partial x_l} g_{jk} \right).$$

The covariant derivative ∇T of a tensor $T = T_{j_1 \dots j_l}^{i_1 \dots i_k}$ will be denoted by $\nabla_s T_{j_1 \dots j_l}^{i_1 \dots i_k} = (\nabla T)_{s j_1 \dots j_l}^{i_1 \dots i_k}$ and with $\nabla^m T$ we will mean the m -th iterated covariant derivative of a tensor T .

The gradient ∇f of a function, the divergence $\operatorname{div} X$ of a tangent vector field and the Laplacian Δf at a point $p \in M$, are defined respectively by

$$g(\nabla f(p), v) = df_p(v) \quad \forall v \in T_p M,$$

$$\operatorname{div} X = \operatorname{tr} \nabla X = \nabla_i X^i = \frac{\partial X^i}{\partial x_i} + \Gamma_{ik}^i X^k$$

(in a local chart) and $\Delta f = \operatorname{div} \nabla f$. The Laplacian ΔT of a tensor T is $\Delta T = g^{ij} \nabla_i \nabla_j T$. We then recall that by the *divergence theorem* for compact manifolds (without boundary), there holds

$$\int_M \operatorname{div} X \, d\mu = 0,$$

for every tangent vector field X on M , which in particular implies

$$\int_M \Delta f \, d\mu = 0,$$

for every smooth function $f : M \rightarrow \mathbb{R}$.

Assuming that we have a globally defined unit normal vector field $\nu : M \rightarrow \mathbb{R}^n$ to $\varphi(M)$ (this will hold in our situation where the hypersurfaces are boundaries of sets $E \subseteq \mathbb{T}^n$, hence we will always consider ν to be the *outer unit normal vector* at every point of ∂E), we define the *second fundamental form* B which is a symmetric bilinear form given, in a local charts, by its components

$$h_{ij} = - \left\langle \frac{\partial^2 \varphi}{\partial x_i \partial x_j} \middle| \nu \right\rangle$$

and whose trace is the *mean curvature* $H = g^{ij} h_{ij}$ of the hypersurface (with these choices, the standard sphere of \mathbb{R}^n has positive mean curvature).

Remark 2.1. If the hypersurface $M \subseteq \mathbb{T}^n$ is the graph of a function $f : U \rightarrow \mathbb{R}$ with U an open subset of \mathbb{R}^{n-1} , that is, $\varphi(x) = (x, f(x))$, then we have

$$g_{ij} = \delta_{ij} + \frac{\partial f}{\partial x_i} \frac{\partial f}{\partial x_j}, \quad \nu = - \frac{(\nabla f, -1)}{\sqrt{1 + |\nabla f|^2}}$$

$$h_{ij} = - \frac{\text{Hess}_{ij} f}{\sqrt{1 + |\nabla f|^2}}$$

$$H = - \frac{\Delta f}{\sqrt{1 + |\nabla f|^2}} + \frac{\text{Hess} f(\nabla f, \nabla f)}{(\sqrt{1 + |\nabla f|^2})^3} = - \text{div} \left(\frac{\nabla f}{\sqrt{1 + |\nabla f|^2}} \right)$$

where $\text{Hess} f$ is the Hessian of the function f .

Then, the following *Gauss–Weingarten relations* hold,

$$\frac{\partial^2 \varphi}{\partial x_i \partial x_j} = \Gamma_{ij}^k \frac{\partial \varphi}{\partial x_k} - h_{ij} \nu \quad \frac{\partial \nu}{\partial x_j} = h_{jl} g^{ls} \frac{\partial \varphi}{\partial x_s}, \quad (2.1)$$

which easily imply $|\nabla \nu| = |B|$ and the identity

$$\Delta \varphi = g^{ij} \left(\frac{\partial^2 \varphi}{\partial x_i \partial x_j} - \Gamma_{ij}^k \frac{\partial \varphi}{\partial x_k} \right) = -g^{ij} h_{ij} \nu = -H \nu. \quad (2.2)$$

The Riemann tensor is expressed via the second fundamental form as follows (*Gauss equations*),

$$R_{ijkl} = h_{ik} h_{jl} - h_{il} h_{jk} \quad (2.3)$$

hence, the formulas for the interchange of covariant derivatives, which involve the Riemann tensor, become

$$\begin{aligned} \nabla_i \nabla_j X^s - \nabla_j \nabla_i X^s &= R_{ijkl} g^{ks} X^l = R_{ijl}^s X^l = (h_{ik} h_{jl} - h_{il} h_{jk}) g^{ks} X^l \\ \nabla_i \nabla_j \omega_k - \nabla_j \nabla_i \omega_k &= R_{ijkl} g^{ls} \omega_s = R_{ijk}^s \omega_s = (h_{ik} h_{jl} - h_{il} h_{jk}) g^{ls} \omega_s \end{aligned}$$

for every vector field X and 1-form ω .

We say that a set $E \subseteq \mathbb{T}^n$ is a *smooth set* if it is the closure of an open subset of \mathbb{T}^n and its boundary ∂E is a smooth embedded hypersurface (unless otherwise stated all the

sets we are going to consider will be smooth). Then, for a smooth set $E \subseteq \mathbb{T}^n$ and $\varepsilon > 0$ small enough, we let (d is the “Euclidean” distance on \mathbb{T}^n)

$$N_\varepsilon = \{x \in \mathbb{T}^n : d(x, \partial E) < \varepsilon\} \quad (2.4)$$

to be a *tubular neighborhood* of ∂E such that the *orthogonal projection map* $\pi_E : N_\varepsilon \rightarrow \partial E$ giving the (unique) closest point on ∂E and the *signed distance function* $d_E : N_\varepsilon \rightarrow \mathbb{R}$ from ∂E

$$d_E(x) = \begin{cases} d(x, \partial E) & \text{if } x \notin E \\ -d(x, \partial E) & \text{if } x \in E \end{cases} \quad (2.5)$$

are well defined and smooth in N_ε (for a proof of the existence of such tubular neighborhood and of all the subsequent properties, see [31] for instance). Moreover, for every $x \in N_\varepsilon$, the projection map is given explicitly by

$$\pi_E(x) = x - \nabla d_E^2(x)/2 = x - d_E(x)\nabla d_E(x)$$

and the unit vector $\nabla d_E(x)$ is orthogonal to ∂E at the point $\pi_E(x) \in \partial E$, indeed actually

$$\nabla d_E(x) = \nabla d_E(\pi_E(x)) = \nu(\pi_E(x)),$$

which means that the integral curves of the vector field ∇d_E are straight segments orthogonal to ∂E .

This clearly implies that the map

$$\partial E \times (-\varepsilon, \varepsilon) \ni (y, t) \mapsto L(y, t) = y + t\nabla d_E(y) = y + t\nu(y) \in N_\varepsilon \quad (2.6)$$

is a smooth diffeomorphism with inverse

$$N_\varepsilon \ni x \mapsto L^{-1}(x) = (\pi_E(x), d_E(x)) \in \partial E \times (-\varepsilon, \varepsilon).$$

Moreover, denoting with JL its Jacobian (relative to the hypersurface ∂E), there holds

$$0 < C_1 \leq JL(y, t) \leq C_2$$

on $\partial E \times (-\varepsilon, \varepsilon)$, for a couple of constants C_1, C_2 , depending on E and ε .

From now on, in all the rest of the work, with N_ε we will always denote a suitable tubular neighborhood of a smooth set, with the above properties.

By means of such tubular neighborhoods of smooth sets $E \subseteq \mathbb{T}^n$, we can speak of “ $W^{k,p}$ -closedness” (or of “ C^k -closedness”) of sets. Indeed, fixed a smooth set E , we say that $F, F' \subseteq \mathbb{T}^n$ are δ -close in $W^{k,p}$ (or in C^k), for some $\delta > 0$ “small enough”, if we have $\text{Vol}(F \Delta F') < \delta$ and that $\partial F, \partial F'$ are contained in a tubular neighborhood N_ε of E as above, described by

$$\partial F = \{y + \psi(y)\nu_E(y) : y \in \partial E\} \quad \text{and} \quad \partial F' = \{y + \psi'(y)\nu_E(y) : y \in \partial E\},$$

for two functions $\psi : \partial E \rightarrow \mathbb{R}$ with $\|\psi - \psi'\|_{W^{k,p}(\partial E)} < \delta$ (respectively, $\|\psi - \psi'\|_{C^k(\partial E)} < \delta$). That is, we are asking that the two sets F and F' differ by a set of small Lebesgue measure and that their boundaries are “close” in $W^{k,p}$ (or C^k) as graphs on ∂E .

Definition 2.2. Given a smooth set $E \subseteq \mathbb{T}^n$ and a tubular neighborhood N_ε of ∂E , as in formula (2.4), for any $M < \varepsilon$, we denote by $\mathfrak{C}_M^1(E)$, the class of all sets $F \subseteq E \cup N_\varepsilon$ such that $\text{Vol}(F \triangle E) \leq M$ and

$$\partial F = \{y + \psi_F(y)\nu_E(y) : y \in \partial E\},$$

for some function $\psi_F \in C^1(\partial E)$, with $\|\psi_F\|_{C^1(\partial E)} \leq M$ (hence, $\partial F \subseteq N_\varepsilon$).

Definition 2.3. Given a sequence of smooth sets $F_i \in \mathfrak{C}_M^1(E)$, for some smooth set $E \subseteq \mathbb{T}^n$, we will write $F_i \rightarrow F$ in $W^{k,p}$ if there exists $F \in \mathfrak{C}_M^1(E)$ such that for every $\delta > 0$, if $i \in \mathbb{N}$ is large enough there holds $\text{Vol}(F_i \triangle F) < \delta$ and, describing the boundaries of F_i, F as

$$\partial F_i = \{y + \psi_i(y)\nu_E(y) : y \in \partial E\} \quad \text{and} \quad \partial F = \{y + \psi(y)\nu_E(y) : y \in \partial E\},$$

for some smooth function $\psi_i, \psi : \partial E \rightarrow \mathbb{R}$, we have $\|\psi_i - \psi\|_{W^{k,p}(\partial E)} < \delta$.

3. THE SURFACE DIFFUSION FLOW AND STRICTLY STABLE CRITICAL SETS

In all the following $\mathbb{T}^n = \mathbb{R}^n / \sim$ is a flat n -dimensional torus, quotient of \mathbb{R}^n by a discrete group of translations generated by some n linearly independent vectors. Since we want to deal with the surface diffusion flow of embedded smooth hypersurfaces which are boundaries of smooth sets (recall that any of them is the closure of an open subset of \mathbb{T}^n), we give the following definition.

Definition 3.1. Let $E \subseteq \mathbb{T}^n$ be a smooth set. We say that the family $E_t \subseteq \mathbb{T}^n$, for $t \in [0, T)$ with $E_0 = E$, is a *surface diffusion flow* starting from E if the map $t \mapsto \chi_{E_t}$ is continuous from $[0, T)$ to $L^1(\mathbb{T}^n)$ and the hypersurfaces ∂E_t move by surface diffusion, that is, there exists a smooth family of embeddings $\varphi_t : \partial E \rightarrow \mathbb{T}^n$, for $t \in [0, T)$, with $\varphi_0 = \text{Id}$ and $\varphi_t(\partial E) = \partial E_t$, such that

$$\frac{\partial \varphi_t}{\partial t} = (\Delta H)\nu, \quad (3.1)$$

where, at every point and time, H and Δ are respectively the mean curvature and the Laplacian (with the Riemannian metric induced by \mathbb{T}^n , that is, by \mathbb{R}^n) of the moving hypersurface ∂E_t , while ν is the “outer” normal to the smooth set E_t .

Remark 3.2. An alternative way to describe the flow is to speak of the sets “enclosed” by the boundary hypersurfaces moving by surface diffusion. This anyway would introduce an ambiguity, since every hypersurface ∂E_t clearly “separate” \mathbb{T}^n in components and one should indicate which ones are actually the sets E_t at every time t . The use of the continuity of the map $t \mapsto \chi_{E_t}$ is a way to avoid such ambiguity. Moreover, it follows easily that being the solution of the PDE system (3.1) unique, by Theorem 1.1, the sets E_t are uniquely determined (being a “geometric flow”, actually the same “geometric” uniqueness also holds for the hypersurfaces ∂E_t , like for the mean curvature flow, see [30, Section 1.3]).

Then, we have the following proposition, which is actually Theorem 1.1 “adapted” to the above definition.

Proposition 3.3. *Given a smooth set $E \subseteq \mathbb{T}^n$ and a tubular neighborhood N_ε of ∂E , as in formula (2.4) and $M < \varepsilon$, for every $E_0 \subseteq \mathbb{T}^n$ smooth set in $\mathfrak{C}_M^1(E)$, whose boundary ∂E_0 is represented by*

$$\partial E_0 = \{y + \psi_0(y)\nu_E(y) : y \in \partial E\}$$

for a smooth function $\psi_0 : \partial E \rightarrow \mathbb{R}$, there exists a unique surface diffusion flow E_t , starting from E_0 , determined by

$$\partial E_t = \{y + \psi_t(y)\nu_E(y) : y \in \partial E\}$$

with smooth functions $\psi_t : \partial E \rightarrow \mathbb{R}$, for t in some maximal positive interval of time $[0, T(E_0))$, with $T(E_0)$ depending on the $C^{2,\alpha}$ -norm of ψ_0 .

We now define the *strictly stable critical sets* and we state our main theorem, we refer to [10, Section 2] for the following facts.

We consider the *Area functional* $\mathcal{A}(\partial E)$ on the family of smooth sets $E \subseteq \mathbb{T}^n$, giving the $(n-1)$ -dimensional “area” of the boundary of E , with a constraint on the $(n$ -dimensional) volume $\text{Vol}(E)$. It is then well known that a set E is a *critical set* (that is, with zero *constrained first variation*) if and only if its boundary satisfy $H = \lambda$, for some constant $\lambda \in \mathbb{R}$. This is obtained, by “testing” the first variation of the Area functional with all the *volume-preserving variations* of E , which turns out to have as infinitesimal generators, vector fields X on ∂E satisfying $X = \psi\nu_E$ with $\psi \in C^\infty(\partial E)$ such that $\int_{\partial E} \psi d\mu = 0$.

Then, at a critical set E , the second variation of the volume-constrained Area functional along such vector fields $X = \psi\nu_E$ on ∂E is given by

$$\left. \frac{d^2}{dt^2} \mathcal{A}(\partial E_t) \right|_{t=0} = \int_{\partial E} (|\nabla \psi|^2 - \psi^2 |B|^2) d\mu.$$

This motivates the following definition.

Definition 3.4. Given any smooth open set $E \subseteq \mathbb{T}^n$, we define the quadratic form

$$\Pi_E(\psi) = \int_{\partial E} (|\nabla \psi|^2 - \psi^2 |B|^2) d\mu, \quad (3.2)$$

for all ψ in the space of Sobolev functions

$$\tilde{H}^1(\partial E) = \left\{ \psi \in H^1(\partial E) : \int_{\partial E} \psi d\mu = 0 \right\}.$$

From now on we will extensively use Sobolev spaces on smooth hypersurfaces. Most of their properties hold as in \mathbb{R}^n , standard references are [2] in the Euclidean space and [3] on manifolds.

We notice that for every $\eta \in \mathbb{R}^n$, we have $\langle \eta | \nu_E \rangle \in \tilde{H}^1(\partial E)$, as in general, for every smooth set $E \subseteq \mathbb{T}^n$, there holds

$$\int_{\partial E} \langle \eta | \nu_E \rangle d\mu = \int_E \text{div } \eta dx = 0, \quad \text{hence also} \quad \int_{\partial E} \nu_E d\mu = 0. \quad (3.3)$$

Then, setting $E_t = E + t\eta$, by the translation invariance of the area functional \mathcal{A} , we have $\mathcal{A}(\partial E_t) = \mathcal{A}(\partial E)$, thus

$$0 = \frac{d^2}{dt^2} \mathcal{A}(\partial E_t) \Big|_{t=0} = \Pi_E(\langle \eta | \nu_E \rangle),$$

that is, the form Π_E is zero on the vector subspace

$$T(\partial E) = \{ \langle \eta | \nu_E \rangle : \eta \in \mathbb{R}^n \} \subseteq \tilde{H}^1(\partial E)$$

of dimension clearly less than or equal to n (and at least one). Then, we split

$$\tilde{H}^1(\partial E) = T(\partial E) \oplus T^\perp(\partial E),$$

where $T^\perp(\partial E) \subseteq \tilde{H}^1(\partial E)$ is the vector subspace L^2 -orthogonal to $T(\partial E)$ (with respect to the measure μ on ∂E), that is,

$$\begin{aligned} T^\perp(\partial E) &= \left\{ \psi \in \tilde{H}^1(\partial E) : \int_{\partial E} \psi \nu_E d\mu = 0 \right\} \\ &= \left\{ \psi \in H^1(\partial E) : \int_{\partial E} \psi d\mu = 0 \text{ and } \int_{\partial E} \psi \nu_E d\mu = 0 \right\} \end{aligned}$$

and we give the following ‘‘stability’’ conditions.

Definition 3.5. We say that a critical set $E \subseteq \mathbb{T}^n$ for \mathcal{A} under a volume constraint is *stable* if

$$\Pi_E(\psi) \geq 0 \quad \text{for all } \psi \in \tilde{H}^1(\partial E)$$

and *strictly stable* if actually

$$\Pi_E(\psi) > 0 \quad \text{for all } \psi \in T^\perp(\partial E) \setminus \{0\}.$$

As one can easily guess, these stability notions are related to the (sufficient and necessary) local minimality properties of a set for the volume-constrained Area functional, as it is shown in [1] (see also the discussion in [10, Section 2.2]).

We observe that there exists an orthonormal frame $\{e_1, \dots, e_n\}$ of \mathbb{R}^n such that

$$\int_{\partial E} \langle \nu_E | e_i \rangle \langle \nu_E | e_j \rangle d\mu = 0, \tag{3.4}$$

for all $i \neq j$, indeed, considering the symmetric $n \times n$ -matrix $A = (a_{ij})$ with components $a_{ij} = \int_{\partial E} \nu_E^i \nu_E^j d\mu$, where $\nu_E^i = \langle \nu_E | \varepsilon_i \rangle$ for some basis $\{\varepsilon_1, \dots, \varepsilon_n\}$ of \mathbb{R}^n , we have

$$\int_{\partial E} (O\nu_E)_i (O\nu_E)_j d\mu = (OAO^{-1})_{ij},$$

for every $O \in SO(n)$. Choosing O such that OAO^{-1} is diagonal and setting $e_i = O^{-1}\varepsilon_i$, relations (3.4) are clearly satisfied. Hence, considering such basis, the functions $\langle \nu_E | e_i \rangle$

which are not identically zero are an orthogonal basis of $T(\partial E)$ and we set

$$I_E = \{i \in \{1, \dots, n\} : \langle \nu_E | e_i \rangle \text{ is not identically zero}\}, \quad O_E = \text{Span}\{e_i : i \in I_E\}. \quad (3.5)$$

We observe that it is easy to see (by a dilation/contraction argument) that any strictly stable smooth critical set must be connected, but actually, Theorem 5.1 can be clearly applied also to finite unions of boundaries of strictly stable critical sets. Moreover, by the very definition above, if ∂E in \mathbb{T}^n is composed by flat pieces, hence its second fundamental form B is identically zero, the set E is critical and stable and with a little effort, actually strictly stable. It is a little more difficult to show that any ball in any dimension $n \in \mathbb{N}$ is strictly stable (it is obviously a critical set), which is connected to the study of the eigenvalues of the Laplacian on the sphere \mathbb{S}^{n-1} , see [22, Theorem 5.4.1], for instance. Then, the same holds for all the ‘‘cylinders’’ $\mathbb{R}^k \times \mathbb{S}^{n-k-1} \subseteq \mathbb{R}^n$, bounding $E \subseteq \mathbb{T}^n$ after taking their quotient by the same equivalence relation defining \mathbb{T}^n , determined by the standard integer lattice of \mathbb{R}^n .

If $n = 2$, it follows that the only bounded strictly stable critical sets of the *Length* functional (that is, the Area functional in 2-dimensional ambient space) in the plane are the disks and in \mathbb{T}^2 they are the disks and the ‘‘strips’’ with straight borders. In the three-dimensional case, a first classification of the smooth stable ‘‘periodic’’ critical sets for the volume-constrained Area functional, was given by Ros in [35], where it is shown that in the flat torus \mathbb{T}^3 , they are *balls*, *2-tori*, *gyroids* or *lamellae*, as in the following figure.




FIGURE 1. From left to right: balls, 2-tori, gyroids and lamellae.

Lamellae, 2-tori and balls are actually strictly stable, while in [23, 24, 36] the authors established the strict stability of gyroids only in some cases.

We are going to prove in Theorem 5.1 that if a smooth set is ‘‘close’’ enough to a strictly stable critical set for the volume constrained Area functional, then its surface diffusion flow exists for every time and smoothly converges to a ‘‘translate’’ of such set.

4. EVOLUTION OF GEOMETRIC QUANTITIES AND BASIC ESTIMATES

Along any surface diffusion flow $\varphi_t : M \rightarrow \mathbb{T}^n$ (or when the ambient is a general flat space) we have the following evolution equations (computed in detail in [29, Proposition 3.4] for a general geometric flow of hypersurfaces),

$$\frac{\partial}{\partial t} g_{ij} = 2\Delta H h_{ij}, \quad \frac{\partial}{\partial t} g^{ij} = -2\Delta H h^{ij}, \quad \frac{\partial}{\partial t} \mu = H\Delta H \mu \quad (4.1)$$

and

$$\frac{\partial}{\partial t} \Gamma_{jk}^i = \nabla B * \Delta H + B * \nabla \Delta H \quad (4.2)$$

where $T * S$ (following Hamilton [26]) denotes a tensor formed by a sum of terms each one of them obtained by the product of a real constant with the contraction on some indices of the pair T, S with the metric g_{ij} and/or its inverse g^{ij} . A very useful property of such $*$ -product is that $|T * S| \leq C|T||S|$ where the constant C depends only on the “algebraic structure” of $T * S$, moreover, it clearly holds $\nabla T * S = \nabla T * S + T * \nabla S$.

Then, arguing as in [30, Proposition 2.3.1], we get the following evolution equation for the mean curvature

$$\frac{\partial}{\partial t} H = -\Delta \Delta H - \Delta H |B|^2.$$

We now introduce some notation which will be useful for the computations that follow (see [29]). If T_1, \dots, T_l is a finite family of tensors (here l is not an index of the tensor T), with the symbol

$$\bigotimes_{i=1}^l T_i$$

we will mean $T_1 * T_2 * \dots * T_l$.

With the symbol $\mathfrak{p}_s(\nabla^\alpha T, \nabla^\beta S, \dots, \nabla^\gamma R)$ we will denote a polynomial in the tensors T, S, \dots, R and their iterated covariant derivatives with the $*$ product as

$$\mathfrak{p}_s(\nabla^\alpha T, \nabla^\beta S, \dots, \nabla^\gamma R) = \sum_{i+j+\dots+k=s} c_{ij\dots k} \nabla^i T * \nabla^j S * \dots * \nabla^k R$$

where the $c_{i_1\dots i_l}$ are some real constants and $i \leq \alpha, j \leq \beta, \dots, k \leq \gamma$. Moreover, we set $\mathfrak{p}_0(\cdot) = 0$. Notice that every tensor must be present in every additive term of $\mathfrak{p}_s(\nabla^\alpha T, \nabla^\beta S, \dots, \nabla^\gamma R)$ and there are no repetitions.

We will use instead the symbol $\mathfrak{q}^s(\nabla^\alpha B, \nabla^\beta H)$ for a completely contracted polynomial (hence a function) of the iterated covariant derivatives of B and H , respectively up to α and β (repetitions are allowed), where in every additive term both B and H must be present and H without derivatives is considered as a contracted B -factor. That is,

$$\mathfrak{q}^s(\nabla^\alpha B, \nabla^\beta H) = \sum_{k=1}^p \bigotimes_{k=1}^p \nabla^{i_k} B \bigotimes_{l=1}^q \nabla^{j_l} H$$

with $p, q \geq 1, i_1, \dots, i_p \leq \alpha$ and $1 \leq j_1, \dots, j_q \leq \beta$, then the coefficient s denotes the sum

$$s = \sum_{k=1}^p (i_k + 1) + \sum_{l=1}^q (j_l + 1). \quad (4.3)$$

We advise the reader that in the following the polynomials \mathfrak{p}_s and \mathfrak{q}^s could vary from a line to another in a computation, by addition of “similar” terms.

With this notation, we have the following “computation” lemmas.

Lemma 4.1. *For every tensor T and function f on M , we have*

$$\frac{\partial}{\partial t} \nabla^s T = \nabla^s \frac{\partial T}{\partial t} + \mathfrak{p}_s(\nabla^{s-1} T, \nabla^s B, \nabla^s \Delta H) \quad \text{for every } s \geq 1 \quad (4.4)$$

$$\frac{\partial}{\partial t} df = d \frac{\partial f}{\partial t} \quad \text{and} \quad \frac{\partial}{\partial t} \nabla^s f = \nabla^s \frac{\partial f}{\partial t} + \mathfrak{p}_{s-1}(\nabla^{s-2}(\nabla f), \nabla^{s-1} B, \nabla^{s-1} \Delta H) \quad (4.5)$$

for every $s \geq 2$.

Proof. We show the first equation by induction on $s \in \mathbb{N}$. If $s = 1$, we have

$$\begin{aligned} \frac{\partial}{\partial t} \nabla T &= \frac{\partial}{\partial t} (\partial T + T\Gamma) = \frac{\partial}{\partial t} \partial T + \frac{\partial}{\partial t} (T\Gamma) = \partial \frac{\partial T}{\partial t} + \frac{\partial T}{\partial t} \Gamma + T \frac{\partial \Gamma}{\partial t} \\ &= \nabla \frac{\partial T}{\partial t} + T * \nabla B * \Delta H + T * B * \nabla \Delta H = \nabla \frac{\partial T}{\partial t} + \mathfrak{p}_1(T, \nabla B, \nabla \Delta H), \end{aligned}$$

where we computed “schematically”, denoting with ∂ the standard derivative in coordinates (with commute with $\frac{\partial}{\partial t}$) and with Γ the Christoffel symbols, moreover, we used formula (4.2).

Now, assuming that formula (4.4) holds up to $s - 1$, we apply it to the tensor $S = \nabla T$

$$\begin{aligned} \frac{\partial}{\partial t} \nabla^s T &= \frac{\partial}{\partial t} \nabla^{s-1} S = \nabla^{s-1} \frac{\partial S}{\partial t} + \mathfrak{p}_{s-1}(\nabla^{s-2} S, \nabla^{s-1} B, \nabla^{s-1} \Delta H) \\ &= \nabla^{s-1} \frac{\partial}{\partial t} \nabla T + \mathfrak{p}_s(\nabla^{s-1} T, \nabla^{s-1} B, \nabla^{s-1} \Delta H) \\ &= \nabla^{s-1} \left(\nabla \frac{\partial T}{\partial t} + \mathfrak{p}_1(T, \nabla B, \nabla \Delta H) \right) + \mathfrak{p}_s(\nabla^{s-1} T, \nabla^{s-1} B, \nabla^{s-1} \Delta H) \\ &= \nabla^s \frac{\partial T}{\partial t} + \mathfrak{p}_s(\nabla^{s-1} T, \nabla^s B, \nabla^s \Delta H) \end{aligned}$$

by the properties of the $*$ -product. Hence, formula (4.4) is proved.

To get equation (4.5), we apply the previous formula to $T = \nabla f$ as follows

$$\begin{aligned} \frac{\partial}{\partial t} \nabla^s f &= \frac{\partial}{\partial t} \nabla^{s-1} \nabla f = \nabla^{s-1} \frac{\partial}{\partial t} \nabla f + \mathfrak{p}_{s-1}(\nabla^{s-2}(\nabla f), \nabla^{s-1} B, \nabla^{s-1} \Delta H) \\ &= \nabla^s \frac{\partial f}{\partial t} + \mathfrak{p}_{s-1}(\nabla^{s-2}(\nabla f), \nabla^{s-1} B, \nabla^{s-1} \Delta H) \end{aligned}$$

and we are done. □

We are now ready to compute the evolution equations and then estimate the key quantities for our flow in dimension $n = 4$.

Proposition 4.2. *Let $E_t \subseteq \mathbb{T}^4$ be a surface diffusion flow. Then, the following equations hold*

$$\frac{d}{dt} \int_{\partial E_t} |\nabla H|^2 d\mu_t = -2\Pi_{E_t}(\Delta H) + \int_{\partial E_t} H \Delta H |\nabla H|^2 d\mu_t \quad (4.6)$$

$$\begin{aligned} & - \int_{\partial E_t} 2B(\nabla H, \nabla H) \Delta H d\mu_t \\ \frac{d}{dt} \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t &= -2 \int_{\partial E_t} |\nabla^4 H|^2 d\mu_t + \int_{\partial E_t} \mathfrak{q}^{10}(B, \nabla^3 H) d\mu_t \quad (4.7) \\ & + \int_{\partial E_t} \mathfrak{q}^{10}(\nabla(B^2), \nabla^4 H) d\mu_t \end{aligned}$$

where Π_{E_t} is the quadratic form defined in formula (3.2) and where

- every “monomial” of $\mathfrak{q}^{10}(B, \nabla^3 H)$ has 4 factors in $B, \nabla H$ and their covariant derivatives, the factor B (or H , without derivatives) is present exactly one time and the other three factors are derivatives of ∇H up to $\nabla^3 H$,
- every “monomial” of $\mathfrak{q}^{10}(\nabla(B^2), \nabla^4 H)$ has 3 factors in $B^2, \nabla(B^2)$ and covariant derivatives of H , the factor B^2 or $\nabla(B^2)$ is present exactly one time, the other two factors are derivatives of ∇H up to $\nabla^4 H$.

Finally, the coefficients of these polynomials are algebraic, that is, they are the result of formal manipulations, in particular, they are independent of E_t .

Proof. Taking into account the above evolution equations, we compute

$$\begin{aligned} \frac{d}{dt} \int_{\partial E_t} |\nabla H|^2 d\mu_t &= \int_{\partial E_t} H |\nabla H|^2 \Delta H d\mu_t - \int_{\partial E_t} 2h^{ij} \nabla_i H \nabla_j H \Delta H d\mu_t \\ & - \int_{\partial E_t} 2g^{ij} \nabla_i H \nabla_j (|B|^2 \Delta H + \Delta \Delta H) d\mu_t \\ &= \int_{\partial E_t} H |\nabla H|^2 \Delta H d\mu_t - \int_{\partial E_t} 2B(\nabla H, \nabla H) \Delta H d\mu_t \\ & + \int_{\partial E_t} 2|B|^2 (\Delta H)^2 d\mu_t + \int_{\partial E_t} 2\Delta H \Delta \Delta H d\mu_t \\ &= \int_{\partial E_t} H |\nabla H|^2 \Delta H d\mu_t - \int_{\partial E_t} 2B(\nabla H, \nabla H) \Delta H d\mu_t \\ & + \int_{\partial E_t} 2|B|^2 (\Delta H)^2 d\mu_t - \int_{\partial E_t} 2|\nabla \Delta H|^2 d\mu_t, \end{aligned}$$

where the first term on the right comes from the area variation and the second from the evolution equation of the inverse of the metric (see formulas (4.1)). Then, we have formula (4.6), recalling the definition of Π_{E_t} .

To get equation (4.7) we compute analogously

$$\begin{aligned}
\frac{d}{dt} \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t &= \int_{\partial E_t} |\nabla^2 H|^2 H \Delta H d\mu_t + 2 \int_{\partial E_t} g\left(\nabla^2 H, \frac{\partial}{\partial t} \nabla^2 H\right) d\mu_t \\
&\quad - \int_{\partial E_t} 2h^{ij} g^{kl} \Delta H \nabla_{ik}^2 H \nabla_{jl}^2 H d\mu_t \\
&\quad - \int_{\partial E_t} 2h^{kl} g^{ij} \Delta H \nabla_{ik}^2 H \nabla_{jl}^2 H d\mu_t
\end{aligned} \tag{4.8}$$

where, as before, the first term comes from the area variation, while the last two from the evolution equation of the inverse of the metric.

Then we focus on the second integral, noticing that we can collect all the other terms inside the integrals in a “polynomial” $q^{10}(B, \nabla^1(\nabla H))$ such that every “monomial” has 4 factors in B , ∇H and their covariant derivatives (remember that we consider H as a contracted B -factor, in the first term – we will always do the same also in the following) and at least three of them are derivatives of ∇H .

By formula (4.5) in Lemma 4.1 with $f = H$ and $s = 2$, we have

$$\begin{aligned}
\frac{\partial}{\partial t} \nabla^2 H &= \nabla^2 \frac{\partial}{\partial t} H + \mathfrak{p}_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H) \\
&= \nabla^2(-\Delta \Delta H - \Delta H |B|^2) + \mathfrak{p}_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H)
\end{aligned}$$

hence, integrating by parts we get

$$\begin{aligned}
\int_{\partial E_t} g\left(\nabla^2 H, \frac{\partial}{\partial t} \nabla^2 H\right) d\mu_t &= \int_{\partial E_t} g\left(\nabla^2 H, \nabla^2(-\Delta \Delta H - \Delta H |B|^2)\right) d\mu_t \\
&\quad + \int_{\partial E_t} g\left(\nabla^2 H, \mathfrak{p}_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H)\right) d\mu_t \\
&= - \int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H (\Delta \Delta H + \Delta H |B|^2) d\mu_t \\
&\quad + \int_{\partial E_t} g\left(\nabla^2 H, \mathfrak{p}_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H)\right) d\mu_t \\
&= - \int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta \Delta H d\mu_t \\
&\quad - \int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta H |B|^2 d\mu_t \\
&\quad + \int_{\partial E_t} g\left(\nabla^2 H, \mathfrak{p}_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H)\right) d\mu_t.
\end{aligned} \tag{4.9}$$

Then, recalling the properties of $\mathfrak{p}_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H)$, integrating by parts in the last integral, we can “take away” the derivative from B (in the monomials containing it) and “move” it on the other three factors, which are derivatives of H . Hence, such integral becomes a term of kind $\int_{\partial E_t} q^{10}(B, \nabla^3 H) d\mu_t$, noticing that $\nabla^4 H$ cannot appear, as by

the properties of $p_1(\nabla^0(\nabla H), \nabla^1 B, \nabla^1 \Delta H)$ either it contains ∇B or $\nabla \Delta H$, but not both together in any of its monomials. Then, recalling equation (4.8) (and the subsequent discussion), we can write

$$\begin{aligned} \frac{d}{dt} \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t &= \int_{\partial E_t} |\nabla^2 H|^2 H \Delta H d\mu_t - 2 \int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta \Delta H d\mu_t \\ &\quad - 2 \int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta H |B|^2 d\mu_t + \int_{\partial E_t} q^{10}(B, \nabla^3 H) d\mu_t \end{aligned} \quad (4.10)$$

where every ‘‘monomial’’ of $q^{10}(B, \nabla^3 H)$ has 4 factors in B , ∇H and their covariant derivatives, moreover

- the factor B (or H , without derivatives) is present exactly one time,
- the other three factors are derivatives of ∇H up to $\nabla^3 H$.

Now we deal with the second term in the right hand side of this equation. We want to show that

$$\int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta \Delta H d\mu_t = \int_{\partial E_t} |\nabla^4 H|^2 d\mu_t + \int_{\partial E_t} q^{10}(\nabla(B^2), \nabla^4 H) d\mu_t \quad (4.11)$$

where every ‘‘monomial’’ of $q^{10}(\nabla(B^2), \nabla^4 H)$ has 3 factors in B^2 , $\nabla(B^2)$ and covariant derivatives of H , moreover

- the factor B^2 or $\nabla(B^2)$ is present exactly one time,
- the other two factors are derivatives of ∇H up to $\nabla^4 H$.

We recall that

$$\int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta \Delta H d\mu_t = \int_{\partial E_t} g^{ij} g^{kl} g^{ms} g^{pq} \nabla_i \nabla_k \nabla_j \nabla_l H \nabla_m \nabla_s \nabla_p \nabla_q H d\mu_t.$$

Then, since every interchange in a pair of subsequent covariant derivatives of ∇H and $\nabla^2 H$ produces an extra ‘‘error term’’ respectively of the form $\text{Riem} * \nabla H = B^2 * \nabla H$ or $\text{Riem} * \nabla^2 H = B^2 * \nabla^2 H$, by the Gauss equations (2.3) (while instead we can switch freely two derivatives on H , being the Hessian symmetric), after some of such interchanges and integrations by parts in order ‘‘to carry’’ derivatives from one of the two factors to the other, we obtain

$$\begin{aligned} \int_{\partial E_t} g^{ij} g^{kl} \nabla_{ikjl}^4 H \Delta \Delta H d\mu_t &= \int_{\partial E_t} g^{ij} g^{kl} g^{ms} g^{pq} \nabla_{qsjl}^4 H \nabla_{ikmp}^4 H d\mu_t \\ &\quad + \int_{\partial E_t} q^{10}(\nabla(B^2), \nabla^4 H) d\mu_t \\ &= \int_{\partial E_t} |\nabla^4 H|^2 d\mu_t + \int_{\partial E_t} q^{10}(\nabla(B^2), \nabla^4 H) d\mu_t. \end{aligned}$$

where the term $\int_{\partial E_t} q^{10}(\nabla(B^2), \nabla^4 H) d\mu_t$ collects all the ‘‘error terms’’ in such manipulation.

Finally, noticing that also the remaining term in formula (4.10) is of the form

$$\int_{\partial E_t} \mathfrak{q}^{10}(\nabla(B^2), \nabla^4 H) d\mu_t.$$

putting together equations (4.10) and (4.11), we get the second formula of the proposition. \square

In all the following, we will be interested in having uniform estimates for the families of sets in $\mathfrak{C}_{M_E}^1(E)$, given a smooth set $E \subseteq \mathbb{T}^n$ and a tubular neighborhood N_ε of ∂E as in formula (2.4), for $M_E < \varepsilon$. To this aim, we need that the constants in the Sobolev, Poincaré, Gagliardo–Nirenberg interpolation and Calderón–Zygmund inequalities relative to all the hypersurfaces ∂F boundaries of the sets $F \in \mathfrak{C}_{M_E}^1(E)$, are uniform. This is proved in detail in [11] (for the Calderón–Zygmund inequalities, we actually need that $F \in \mathfrak{C}_{M_E}^1(E)$, with $M_E > 0$ small enough), hence, from now on we will use the adjective “uniform” in order to underline such fact. We also highlight that in all the following we will denote with C a constant which may vary from a line to another and depends only on E and M_E .

Proposition 4.3 (Gagliardo–Nirenberg interpolation inequalities). *Let $E \subseteq \mathbb{T}^n$ be a smooth set, j, m be integer such that $0 \leq j < m$ and $0 < r, q \leq +\infty$. Then, for every $F \in \mathfrak{C}_{M_E}^1(E)$ and every covariant tensor $T = T_{i_1 \dots i_j}$ the following “uniform” interpolation inequalities hold*

$$\|\nabla^j T\|_{L^p(\partial F)} \leq C \left(\|\nabla^m T\|_{L^r(\partial F)} + \|T\|_{L^r(\partial F)} \right)^\theta \|T\|_{L^q(\partial F)}^{1-\theta}, \quad (4.12)$$

with the compatibility condition

$$\frac{1}{p} = \frac{j}{n-1} + \theta \left(\frac{1}{r} - \frac{m}{n-1} \right) + \frac{1-\theta}{q},$$

for all $\theta \in [j/m, 1)$ for which $p \in [1, +\infty)$ is nonnegative and where C is a constant depending only on n, j, m, p, q, r and E, M_E . Moreover, if $f : \partial F \rightarrow \mathbb{R}$ is a smooth function with $\int_{\partial F} f d\mu = 0$, inequality (4.12) becomes

$$\|\nabla^j f\|_{L^p(\partial F)} \leq C \|\nabla^m f\|_{L^r(\partial F)}^\theta \|f\|_{L^q(\partial F)}^{1-\theta}. \quad (4.13)$$

Finally, if $f : \partial F \rightarrow \mathbb{R}$ is any smooth function, there holds

$$\|\nabla^j f\|_{L^p(\partial F)} \leq C \|\nabla^m f\|_{L^r(\partial F)}^\theta \|\nabla f\|_{L^q(\partial F)}^{1-\theta} \quad (4.14)$$

with the compatibility condition

$$\frac{1}{p} = \frac{j-1}{n-1} + \theta \left(\frac{1}{r} - \frac{m-1}{n-1} \right) + \frac{1-\theta}{q}, \quad (4.15)$$

for all $\theta \in [j/m, 1)$ for which $p \in [1, +\infty)$ is nonnegative and where the constant C is like above.

By density, these inequalities clearly extend to functions and tensors in the appropriate Sobolev spaces.

Proof – Sketch. For a single fixed regular hypersurface ∂F , inequality (4.13) is given by Theorem 3.70 in [3], while inequality (4.12) for T equal to a function $f : \partial F \rightarrow \mathbb{R}$ can be obtained by repeating step by step the proof of such theorem, once established the standard Sobolev inequality for hypersurfaces without boundary,

$$\|f\|_{L^{p^*}(\partial F)} \leq C (\|\nabla f\|_{L^p(\partial F)} + \|f\|_{L^p(\partial F)}), \quad (4.16)$$

for every $p \in [1, +\infty)$ (an example of such argument can be found in [29, Section 6]). The extension of inequality (4.12) to tensors can be obtained as in [29, Sections 5 and 6], by means of the estimate (see [3], Proposition 2.11 and also [7, 8]),

$$\left| \nabla \sqrt{|T|^2 + \varepsilon^2} \right| = \left| \frac{\langle \nabla T, T \rangle}{\sqrt{|T|^2 + \varepsilon^2}} \right| \leq \frac{|T|}{\sqrt{|T|^2 + \varepsilon^2}} |\nabla T| \leq |\nabla T|$$

clearly leading to the previous Sobolev inequality for tensors, as $\sqrt{|T|^2 + \varepsilon^2}$ converges to $|T|$ when $\varepsilon \rightarrow 0$ (this argument is necessary as $|T|$ is not necessarily smooth).

In order to get inequality (4.14), we apply estimate (4.13) to the function $f - \bar{f}$, with $\bar{f} = \int_{\partial F} f d\mu$, and we get

$$\|\nabla^j f\|_{L^p(\partial F)} \leq C \|\nabla^m f\|_{L^r(\partial F)}^\theta \|f - \bar{f}\|_{L^{q^*}(\partial F)}^{1-\theta}$$

where

$$\frac{1}{p} = \frac{j}{n-1} + \theta \left(\frac{1}{r} - \frac{m}{n-1} \right) + \frac{1-\theta}{q^*},$$

with $q^* = \frac{(n-1)q}{n-1-q}$. Then, by means of the Sobolev–Poincaré inequality

$$\|f - \bar{f}\|_{L^{q^*}(\partial F)} \leq C \|\nabla f\|_{L^q(\partial F)}, \quad (4.17)$$

that also can be easily deduced by estimate (4.13), we obtain

$$\|\nabla^j f\|_{L^p(\partial F)} \leq C \|\nabla^m f\|_{L^r(\partial F)}^\theta \|\nabla f\|_{L^q(\partial F)}^{1-\theta}$$

and it is a straightforward check that relation (4.15) holds.

Finally, the “uniformity” in the constants of the inequalities, independently of $F \in \mathfrak{C}_{M_E}^1(E)$, follows by the same independence in the Sobolev inequalities (by checking the proof of Theorem 3.70 in [3]). This is shown and discussed in detail in [11]. \square

Remark 4.4. Notice that in the same hypotheses of this proposition, by inequality (4.17), we also have the following “uniform” Poincaré inequalities

$$\|f - \bar{f}\|_{L^p(\partial F)} \leq C \|\nabla f\|_{L^p(\partial F)}, \quad (4.18)$$

for every $p \in [1, +\infty)$.

Remark 4.5. Very similar “uniform” interpolation inequalities are worked out in [29], for any family of smooth, n -dimensional, regular hypersurfaces $N \subseteq \mathbb{R}^{n+1}$ satisfying $\text{Vol}(N) + \|H\|_{L^{n+\delta}(N)} \leq C$, for some $\delta > 0$, instead of being boundaries of sets belonging to $\mathfrak{C}_{M_E}^1(E)$.

From now on, we consider as ambient space the four-dimensional flat torus \mathbb{T}^4 .

Proposition 4.6. *Let $E_t \subseteq \mathbb{T}^4$ be a surface diffusion flow such that $E_t \in \mathfrak{C}_{M_E}^1(E)$, for some smooth set E . Then,*

$$\begin{aligned} \frac{d}{dt} \int_{\partial E_t} |\nabla H|^2 d\mu_t &\leq -2\Pi_{E_t}(\Delta H) + C_1 \int_{\partial E_t} |B| |\nabla H|^2 |\Delta H| d\mu_t \\ \frac{d}{dt} \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t &\leq -\|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C_2(1 + \|\nabla H\|_{L^2(\partial E_t)}^\tau) \|\nabla H\|_{L^2(\partial E_t)}^2 \end{aligned} \quad (4.19)$$

for some exponent $\tau > 0$ and constant C_1, C_2 such that C_1 depends only on E and M_E , while C_2 depends also on $\|B\|_{L^\infty(\partial E_t)}$ and $\|\nabla B\|_{L^6(\partial E_t)}$.

Proof. The first inequality follows immediately from formula (4.6).

About the second one, recalling formula (4.7), we start dealing with the integral $\int_{\partial E_t} \mathfrak{q}^{10}(B, \nabla^3 H) d\mu_t$, which is a sum of integrals each one like

$$\int_{\partial E_t} B \bigotimes_{l=1}^3 \nabla^{j_l} H d\mu_t,$$

with $0 \leq j_l \leq 3$, moreover, it must hold

$$10 = 1 + \sum_{l=1}^3 (j_l + 1) \quad \text{that is,} \quad \sum_{l=1}^3 j_l = 6,$$

by formula (4.3). Hence,

$$\int_{\partial E_t} |\mathfrak{q}^{10}(B, \nabla^3 H)| d\mu_t \leq C \sum \int_{\partial E_t} \left| B \bigotimes_{l=1}^3 \nabla^{j_l} H \right| d\mu_t \leq C \sum \int_{\partial E_t} |B| \prod_{l=1}^3 |\nabla^{j_l} H| d\mu_t.$$

We can estimate each integral of the last sum by means of Hölder inequality, as follows

$$\int_{\partial E_t} |B| \prod_{l=1}^3 |\nabla^{j_l} H| d\mu_t \leq \|B\|_{L^\infty(\partial E_t)} \prod_{l=1}^3 \|\nabla^{j_l} H\|_{L^{\beta_l}(\partial E_t)}$$

where $\beta_l = \frac{9}{j_l+1} > 2$, which clearly satisfy

$$\sum_{l=1}^3 \frac{1}{\beta_l} = \sum_{l=1}^3 \frac{j_l + 1}{9} = 1.$$

Then, we now estimate any of such products as follows: applying the uniform interpolation inequalities (4.14) to H , we get

$$\|\nabla^{j_l} H\|_{L^{\beta_l}(\partial E_t)} \leq C \|\nabla^4 H\|_{L^2(\partial E_t)}^{\theta_l} \|\nabla H\|_{L^2(\partial E_t)}^{1-\theta_l}, \quad (4.20)$$

for some constants C depending on β_l and coefficients θ_l which are given by

$$\theta_l = \frac{1}{3} \left(j_l - \frac{3}{\beta_l} + \frac{1}{2} \right) \in \left(\frac{j_l - 1}{3}, 1 \right).$$

Then, noticing that

$$\Theta = \sum_{l=1}^3 \theta_l = \frac{1}{3} \sum_{l=1}^3 j_l - \sum_{l=1}^3 \frac{1}{\beta_l} + \frac{1}{2} = 1 + \frac{1}{2} < 2,$$

multiplying together inequalities (4.20), applying the Young inequality, we have

$$\begin{aligned} \int_{\partial E_t} |B| \prod_{l=1}^3 |\nabla^{j_l} H| d\mu_t &\leq \|B\|_{L^\infty(\partial E_t)} \|\nabla^4 H\|_{L^2(\partial E_t)}^\Theta \|\nabla H\|_{L^2(\partial E_t)}^{3-\Theta} \\ &\leq \varepsilon \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \|\nabla H\|_{L^2(\partial E_t)}^\tau \end{aligned}$$

for any $\varepsilon > 0$, with $C = C(\varepsilon, \|B\|_{L^\infty(\partial E_t)})$ and $\tau > 2$.

Hence, we conclude (by choosing appropriately $\varepsilon > 0$ for each summand in $\mathfrak{q}^{10}(B, \nabla^3 H)$) that

$$\int_{\partial E_t} \mathfrak{q}^{10}(B, \nabla^3 H) d\mu_t \leq \frac{1}{2} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \|\nabla H\|_{L^2(\partial E_t)}^\tau \quad (4.21)$$

with $C = C(\|B\|_{L^\infty(\partial E_t)})$ and $\tau > 2$.

Now we deal with the last integral in formula (4.7), that is

$$\int_{\partial E_t} \mathfrak{q}^{10}(\nabla(B^2), \nabla^4 H) d\mu_t = \int_{\partial E_t} \nabla^i(B^2) * \nabla^j H * \nabla^4 H d\mu_t,$$

with $0 \leq i \leq 1$ and $0 \leq j \leq 3$, moreover, it must hold

$$10 = i + 2 + j + 1 + 5 \quad \text{that is,} \quad i + j = 2,$$

by formula (4.3), which actually implies that we have only two types of terms,

$$\int_{\partial E_t} B^2 * \nabla^2 H * \nabla^4 H d\mu_t \quad \text{and} \quad \int_{\partial E_t} B * \nabla B * \nabla H * \nabla^4 H d\mu_t.$$

We now respectively estimate their modulus as follows (after ‘‘carrying’’ the modulus inside the integrals and using the properties of the $*$ -product):

$$\begin{aligned} \int_{\partial E_t} |B^2| |\nabla^2 H| |\nabla^4 H| d\mu_t &\leq \varepsilon \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t, \\ \int_{\partial E_t} |B| |\nabla B| |\nabla H| |\nabla^4 H| d\mu_t &\leq \varepsilon \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \int_{\partial E_t} |\nabla B|^2 |\nabla H|^2 d\mu_t \end{aligned}$$

for any $\varepsilon > 0$, with $C = C(\varepsilon, \|B\|_{L^\infty(\partial E_t)})$.

By the uniform inequalities (4.14) and Young inequality, we have

$$\|\nabla^2 H\|_{L^2(\partial E_t)}^2 \leq C \|\nabla^4 H\|_{L^2(\partial E_t)}^{2/3} \|\nabla H\|_{L^2(\partial E_t)}^{4/3} \leq \varepsilon \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \int_{\partial E_t} |\nabla H|^2 d\mu_t$$

hence,

$$\int_{\partial E_t} |B^2| |\nabla^2 H| |\nabla^4 H| d\mu_t \leq \varepsilon \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \int_{\partial E_t} |\nabla H|^2 d\mu_t \quad (4.22)$$

for some constant $C = C(\varepsilon, \|B\|_{L^\infty(\partial E_t)})$.

Then, by Hölder inequality, we have

$$\begin{aligned} \int_{\partial E_t} |\nabla B|^2 |\nabla H|^2 d\mu_t &\leq \|\nabla B\|_{L^6(\partial E_t)}^2 \|\nabla H\|_{L^3(\partial E_t)}^2 \\ &\leq C \|\nabla B\|_{L^6(\partial E_t)}^2 \|\nabla^4 H\|_{L^2(\partial E_t)}^{1/3} \|\nabla H\|_{L^2(\partial E_t)}^{5/3}, \end{aligned}$$

where we used again the uniform interpolation inequalities (4.14) as follows,

$$\|\nabla H\|_{L^3(\partial E_t)}^2 \leq C \|\nabla^4 H\|_{L^2(\partial E_t)}^{1/3} \|\nabla H\|_{L^2(\partial E_t)}^{5/3}.$$

Hence, by means of Young inequality, we finally get

$$\int_{\partial E_t} |\nabla B|^2 |\nabla H|^2 d\mu_t \leq \varepsilon \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \int_{\partial E_t} |\nabla H|^2 d\mu_t \quad (4.23)$$

for some constant $C = C(\varepsilon, \|B\|_{L^\infty(\partial E_t)}, \|\nabla B\|_{L^6(\partial E_t)})$.

Hence, putting together estimates (4.22) and (4.23), we conclude (by choosing appropriately $\varepsilon > 0$ for each summand in $q^{10}(\nabla(B^2), \nabla^4 H)$ as before, that

$$\int_{\partial E_t} q^{10}(\nabla(B^2), \nabla^4 H) d\mu_t \leq \frac{1}{2} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \|\nabla H\|_{L^2(\partial E_t)}^2 \quad (4.24)$$

with $C = C(\|B\|_{L^\infty(\partial E_t)}, \|\nabla B\|_{L^6(\partial E_t)})$.

Estimating the two last terms in formula (4.7) by means of inequalities (4.21) and (4.24), we obtain inequality (4.19). \square

Remark 4.7. Recalling Remark 4.5, in the proof of this proposition we could alternatively uniformly control the constants in the interpolation inequalities by a function of $\text{Vol}(\partial E_t) + \|H\|_{L^4(\partial E_t)}$ (of $\text{Vol}(\partial E_t) + \|H\|_{L^n(\partial E_t)}$ in the n -dimensional case), instead of using Proposition 4.3, as it is done in [29]. It follows that this proposition holds also for only immersed (not boundaries of sets) smooth hypersurfaces moving by the surface diffusion flow with equibounded volumes.

Lemma 4.8. *Let $E \subseteq \mathbb{T}^4$ be a smooth set and N_ε be a tubular neighborhood of ∂E , as in formula (2.4). For M_E small enough and $\delta > 0$, there exists a constant $C = C(E, M_E, \delta)$ such that if $F \in \mathfrak{C}_{M_E}^1(E)$ with*

$$\partial F = \{y + \psi_F(y)\nu_E(y) : y \in \partial E\}$$

for a smooth function $\psi_F : \partial E \rightarrow \mathbb{R}$ and

$$\int_{\partial F} |\nabla^2 H|^2 d\mu + \int_{\partial F} |\nabla H|^2 d\mu \leq \delta,$$

there hold

$$\|B\|_{L^\infty(\partial F)} + \|\nabla B\|_{L^6(\partial F)} \leq C \quad \text{and} \quad \|\psi_F\|_{W^{4,2}(\partial E)} \leq C.$$

Moreover, for every $1 \leq p < 6$, there exists a monotone nondecreasing function $\omega : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, depending only on E and M_E , with $\lim_{\delta \rightarrow 0^+} \omega(\delta) = 0$ and such that if F satisfies the further condition

$$\text{Vol}(F \Delta E) \leq \delta,$$

then $\|\psi_F\|_{W^{3,p}(\partial E)} \leq \omega(\delta)$.

As a consequence, if $E_i \subseteq \mathfrak{C}_{M_E}^1(E)$ is a sequence of smooth sets such that

$$\sup_{i \in \mathbb{N}} \int_{\partial E_i} |\nabla^2 H|^2 d\mu_i + \int_{\partial E_i} |\nabla H|^2 d\mu_i < +\infty,$$

then there exists a (non necessarily smooth) set $E' \in \mathfrak{C}_{M_E}^1(E)$ such that, up to a (non relabeled) subsequence, $E_i \rightarrow E'$ in $W^{3,p}$ as $i \rightarrow \infty$, for all $1 \leq p < 6$. Moreover, if

$$\int_{\partial E_i} |\nabla^2 H|^2 d\mu_i + \int_{\partial E_i} |\nabla H|^2 d\mu_i \rightarrow 0,$$

as $i \rightarrow \infty$, the set E' is critical for the volume-constrained Area functional, that is, its mean curvature is constant.

Proof. Let $F \in \mathfrak{C}_{M_E}^1(E)$ with an associate function $\psi_F : \partial E \rightarrow \mathbb{R}$ as in the statement. We start by observing that, by the uniform Sobolev inequality (4.16) (extended to tensors, as for the other inequalities in Proposition 4.3) applied to ∇H , we have

$$\|\nabla H\|_{L^6(\partial F)} \leq C (\|\nabla^2 H\|_{L^2(\partial F)} + \|\nabla H\|_{L^2(\partial F)}) \leq C\sqrt{\delta}$$

then, by an easy slight variation of inequality (4.16) and by inequality (4.17) (both extended to p larger than the dimension $n - 1 = 3$), we get

$$\|H\|_{L^\infty(\partial F)} \leq C (\|\nabla H\|_{L^6(\partial F)} + \|H\|_{L^2(\partial F)}) \leq C\sqrt{\delta} + C\|H\|_{L^2(\partial F)}$$

$$\|H - \bar{H}\|_{L^\infty(\partial F)} \leq C \|\nabla H\|_{L^6(\partial F)} \leq C\sqrt{\delta} \quad (4.25)$$

where $\bar{H} = \int_{\partial F} H d\mu$ and all the constants depends only on E and M_E .

By the uniform C^1 -bounds on ∂F , we may find a finite family (only depending on E and M_E) of “solid” cylinders of the form $\mathcal{C}_k = D_k + \nu_E(x_k)\mathbb{R}$, with $D_k \subseteq T_{x_k}E$ a closed disk of fixed radius $R > 0$ centered at the origin, for a finite family of points $x_k \in E$, such that $\partial F \cap \mathcal{C}_k$ is the graph on D_k of a smooth function $f_k : D_k \rightarrow \mathbb{R}$, with

$$\|f_k\|_{C^1(D_k)} \leq M_E \quad (4.26)$$

for every k and $\partial F = \bigcup \partial F \cap \mathcal{C}_k$.

Since we want to estimate $\int_{\partial F \cap \mathcal{C}_k} H d\mu$, which is a “geometric” quantity, we can assume (by means of an isometry) that $T_{x_k}E = \langle e_1, e_2, e_3 \rangle$, hence $\nu_E(x_k) = e_4$, in the canonical orthonormal basis of \mathbb{R}^4 and

$$\partial F \cap \mathcal{C}_k = \{(x, f_k(x)) : x \in D_k\}.$$

Then, by formulas in Remark 2.1 we have

$$H = -\operatorname{div}\left(\frac{\nabla f_k}{\sqrt{1+|\nabla f_k|^2}}\right),$$

hence,

$$\begin{aligned}\int_{D_k} H dx &= -\int_{D_k} \operatorname{div}\left(\frac{\nabla f_k}{\sqrt{1+|\nabla f_k|^2}}\right) dx = -\int_{\partial D_k} \left\langle \frac{\nabla f_k}{\sqrt{1+|\nabla f_k|^2}} \middle| \frac{x}{|x|} \right\rangle d\sigma \\ &= \int_{\partial D_k} \left\langle \nu_F \middle| \frac{x}{|x|} \right\rangle d\sigma\end{aligned}$$

where σ is the canonical (standard) 2-dimensional measure on the sphere ∂D_k . Thus, being the last term at most equal to the area of the sphere ∂D_k , we get

$$\bar{H} \operatorname{Vol}(D_k) = \int_{D_k} (\bar{H} - H) dx + \int_{D_k} H dx \leq \int_{D_k} |H - \bar{H}| dx + C \leq C \int_{\partial F \cap C_k} |H - \bar{H}| dx + C$$

where in the last inequality we kept into account estimate (4.26) in changing the domain (and variables) of integration. Hence, controlling the last term of this inequality by estimate (4.25), it follows that \bar{H} is bounded by a constant depending on E, M_E, δ and the same then holds also for H . In particular, recalling that the volume of ∂F is uniformly bounded (as $F \in \mathfrak{C}_{M_E}^1(E)$), we have that $H \in L^q(\partial F)$ for every $q \in [1, +\infty)$. Then, choosing M_E small enough, Theorem 3.1 in [11] holds, saying that we have an analogous uniform estimate on B in $L^q(\partial F)$, for every $q \in [1, +\infty)$.

Once we have a control on $\|B\|_{L^q(\partial F)}$, for some exponent q larger than the dimension of the hypersurfaces, again if M_E is small enough, we have the following uniform higher order Calderón–Zygmund–type inequalities discussed in [11, Section 3.1], holding in any dimension,

$$\|\nabla^k B\|_{L^2(\partial F)} \leq C_k (1 + \|\nabla^k H\|_{L^2(\partial F)})$$

for every $k \in \mathbb{N}$, where the constants C_k depend on E, M_E and $\|B\|_{L^q(\partial F)}$ (and the dimension), hence in our situation they depend on E and M_E .

It then follows

$$\|\nabla^2 B\|_{L^2(\partial F)} \leq C(E, M_E) \tag{4.27}$$

and, by the uniform Sobolev embeddings in dimension 3, we conclude

$$\|B\|_{L^q(\partial F)} + \|\nabla B\|_{L^6(\partial F)} \leq C(E, M_E, \delta) \tag{4.28}$$

for every $q \in [1, +\infty)$.

These geometric estimates on B and their derivatives, can be “transferred” to estimates on the function $\psi_F : \partial E \rightarrow \mathbb{R}$, by means of the technique of localization/representation for any “graphical” hypersurface on ∂E introduced by Langer in [28] for surfaces, generalized to any dimension by Delladio [12] and fully developed in details by Breuning in the papers [4, 5, 6] (such technique is similar to the one we used to estimate \bar{H} above). In particular, by the results in [6], under a uniform control on $\|B\|_{L^q(\partial F)}$ with q larger than

the dimension of the hypersurface, we have that an estimate on $\|B\|_{W^{k,p}(\partial F)}$ implies a uniform estimate on $\|\psi_F\|_{W^{k+2,p}(\partial F)}$ and viceversa, for every set $F \in \mathfrak{C}_{M_E}^1(E)$. Hence, by the previous estimates (4.27) and (4.28) on B and its derivatives, we conclude

$$\|\psi_F\|_{W^{4,2}(\partial E)} \leq C(E, M_E, \delta).$$

Then, we notice that, by uniform Sobolev embeddings, we have

$$\|\nabla^2 \psi_F\|_{L^\infty(\partial E)} \leq C(E, M_E, \delta)$$

which in turn implies $\|B\|_{L^\infty(\partial F)} \leq C(E, M_E, \delta)$, by what we said above. Now, in the hypotheses of the lemma on a sequence of sets E_i , writing

$$\partial E_i = \{y + \psi_i(y)\nu_E(y) : y \in \partial E\},$$

by the previous estimates and the uniform Sobolev compact embeddings

$$W^{4,2}(\partial E) \hookrightarrow W^{3,p}(\partial E) \hookrightarrow C^1(\partial E)$$

for all $1 \leq p < 6$, up to a (not relabeled) subsequence there exists a set $E' \in \mathfrak{C}_{M_E}^1(E)$ such that $\psi_i \rightarrow \psi_{E'}$ in $W^{3,p}(\partial E)$ (and in $C^1(\partial E)$) where

$$\partial E' = \{y + \psi_{E'}(y)\nu_E(y) : y \in \partial E\},$$

for all $1 \leq p < 6$.

If actually

$$\int_{\partial E_i} |\nabla^2 H|^2 d\mu_i + \int_{\partial E_i} |\nabla H|^2 d\mu_i \rightarrow 0,$$

clearly for the limit set E' the mean curvature must be constant.

The fact that $\|\psi_F\|_{W^{3,p}(\partial E)}$ goes uniformly to zero as $\delta \rightarrow 0$, hence we have a function ω as in the statement, follows by the fact that, assuming $F_i \in \mathfrak{C}_{M_E}^1(E)$ and

$$\text{Vol}(F_i \triangle E) \leq \delta_i, \quad \int_{\partial F_i} |\nabla^2 H|^2 d\mu_i + \int_{\partial F_i} |\nabla H|^2 d\mu_i \leq \delta_i$$

with $\delta_i \rightarrow 0$, as $i \rightarrow \infty$, by the previous argument we have that $\psi_{F_i} : \partial E \rightarrow \mathbb{R}$ converges to some $\psi : \partial E \rightarrow \mathbb{R}$ in $W^{3,p}(\partial E)$, hence in $L^1(\partial E)$, while the limit $\text{Vol}(F_i \triangle E) \rightarrow 0$ implies that $\|\psi_{F_i}\|_{L^1(\partial E)} \rightarrow 0$, then we conclude that ψ must be zero and we have the thesis. \square

5. GLOBAL EXISTENCE AND STABILITY

Theorem 5.1. *Let $E \subseteq \mathbb{T}^4$ be a strictly stable critical set for the Area functional under a volume constraint and let N_ε be a tubular neighborhood of ∂E , as in formula (2.4). For $M_E < \varepsilon/2$ small enough, there exists $\delta > 0$ such that, if E_0 is a smooth set in $\mathfrak{C}_{M_E}^1(E)$ satisfying $\text{Vol}(E_0) = \text{Vol}(E)$ and*

$$\text{Vol}(E_0 \triangle E) \leq \delta \quad \text{and} \quad \int_{\partial E_0} |\nabla^2 H|^2 d\mu_0 + \int_{\partial E_0} |\nabla H|^2 d\mu_0 \leq \delta, \quad (5.1)$$

then, the unique smooth surface diffusion flow E_t starting from E_0 , given by Proposition 3.3, is defined for all $t \geq 0$. Moreover, E_t converges smoothly to $E' = E + \eta$ exponentially fast as

$t \rightarrow +\infty$, for some $\eta \in \mathbb{R}^4$, with the meaning that the sequence of smooth functions $\psi_t : \partial E \rightarrow \mathbb{R}$ representing ∂E_t as “normal graphs” on ∂E , that is,

$$\partial E_t = \{y + \psi_t(y)\nu_E(y) : y \in \partial E\},$$

satisfy, for every $k \in \mathbb{N}$,

$$\|\psi_t - \psi\|_{C^k(\partial E)} \leq C_k e^{-\beta_k t},$$

for every $t \in [0, +\infty)$, for some positive constants C_k and β_k , where $\psi : \partial E \rightarrow \mathbb{R}$ represents $\partial E' = \partial E + \eta$ as a “normal graph” on ∂E .

Remark 5.2. The request that E_0 belongs to $\mathfrak{C}_{M_E}^1(E)$ with M_E small enough, is necessary only in order to be able to represent its boundary as a graph of a function with bounded gradient on ∂E and to have uniform Sobolev, interpolation and Calderón–Zygmund inequalities, as proved in [11, Section 3], while the first condition (5.1) is a “closedness” assumption in L^1 for E_0 and E (that is, on ψ_0). The second “small energy” condition (5.1) in the theorem implies (see the last part of Lemma 4.8 and its proof) that the mean curvature of ∂E_0 is “close” to be constant, as it is for the strictly stable set E (actually for any critical set). Notice that this latter is a fourth order condition for the boundary of E_0 and that all these assumptions are clearly implied by an appropriate $W^{4,2}$ -closedness of E_0 to E , arguing as in Lemma 4.8.

Proof of Theorem 5.1. By choosing M_E small enough, we assume that for every set $F \in \mathfrak{C}_{2M_E}^1(E)$, all the constants in the inequalities we are going to consider for functions on ∂F are uniform, depending on E and M_E , by [11].

After choosing some small $\delta_0 > 0$, we consider the surface diffusion flow E_t starting from $E_0 \in \mathfrak{C}_{M_E}^1(E)$ satisfying

$$\text{Vol}(E_0 \triangle E) \leq \delta \quad \text{and} \quad \int_{\partial E_0} |\nabla^2 H|^2 d\mu_0 + \int_{\partial E_0} |\nabla H|^2 d\mu_0 \leq \delta,$$

for $\delta < \delta_0/2$ and we let $T(E_0) \in (0, +\infty]$ be the maximal time such that the flow is defined for t in the interval $[0, T(E_0))$, $E_t \in \mathfrak{C}_{2M_E}^1(E)$,

$$\text{Vol}(E_t \triangle E) \leq \delta_0 \quad \text{and} \quad \mathcal{F}(t) = \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t + \int_{\partial E_t} |\nabla H|^2 d\mu_t \leq \delta_0.$$

All the moving boundaries ∂E_t can be represented as normal graphs on ∂E as

$$\partial E_t = \{y + \psi_t(y)\nu_E(y) : y \in \partial E\}$$

for some smooth functions $\psi_t : \partial E \rightarrow \mathbb{R}$. Moreover, if $T(E_0) < +\infty$, then at least one of the three following conditions must hold:

- $\limsup_{t \rightarrow T(E_0)} \|\psi_t\|_{C^1(\partial E)} = 2M_E$
- $\limsup_{t \rightarrow T(E_0)} \mathcal{F}(t) = \delta_0$
- $\limsup_{t \rightarrow T(E_0)} \text{Vol}(E_t \triangle E) = \delta_0$

otherwise, restarting the flow from a time \bar{t} close enough to $T(E_0)$ by means of Proposition 3.3, we have the contradiction that $T(E_0)$ cannot be the maximal time defined

above. Indeed, the time interval of smooth existence of the flow given by such proposition is bounded below by a constant depending on the $C^{2,\alpha}$ -norm of $\psi_{\bar{t}}$ and this latter by a constant depending on δ_0 , by the first point of Lemma 4.8 and Sobolev (uniform) embeddings.

We are going to show that if δ_0 was chosen small enough, there exists $\delta > 0$ such that none of these conditions can occur, hence $T(E_0) = +\infty$, that is, the surface diffusion flow of E_0 exists for all time.

Let us define, for $K > 2$, the following “energy” function

$$\mathcal{E}(t) = \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t + K \int_{\partial E_t} |\nabla H|^2 d\mu_t \geq \mathcal{F}(t)$$

(notice that also holds $\mathcal{E}(t) \leq K\mathcal{F}(t)$). From Lemma 4.8 we easily have

$$\|B\|_{L^\infty(\partial E_t)} + \|\nabla B\|_{L^6(\partial E_t)} \leq S_0(\mathcal{F}(t)) \leq S_0(\mathcal{E}(t)), \quad (5.2)$$

for $t \in [0, T(E_0))$, where the function $S_0 : [0, +\infty) \rightarrow \mathbb{R}^+$ is continuous and monotone nondecreasing and it is determined by E and M_E .

We now split the rest of the proof into steps. Our first goal will be showing that the function \mathcal{E} decreases in time if δ is small enough, for an appropriate constant K .

Step 1 (Monotonicity of \mathcal{E}).

By Proposition 4.6, for any $t \in [0, T(E_0))$, we have

$$\begin{aligned} \frac{d}{dt}\mathcal{E}(t) &\leq -2K\Pi_{E_t}(\Delta H) + KC_1 \int_{\partial E_t} |B|\nabla H|^2|\Delta H| d\mu_t \\ &\quad - \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + S_1(\mathcal{E}(t))(1 + \|\nabla H\|_{L^2(\partial E_t)}^\tau) \|\nabla H\|_{L^2(\partial E_t)}^2, \end{aligned} \quad (5.3)$$

for some exponent $\tau > 0$ and a continuous, monotone nondecreasing function $S_1 : [0, +\infty) \rightarrow \mathbb{R}^+$ (as the function C above), depending on E , M_E , by inequality (5.2). Moreover, estimating the first integral at the right hand side as in the first part of Proposition 4.6, we get

$$C_1 \int_{\partial E_t} |B|\nabla H|^2|\Delta H| d\mu_t \leq \frac{1}{2K} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + S_2(\mathcal{E}(t))K \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau'},$$

with $\tau' > 0$ and the function S_2 is like S_0 and S_1 .

Hence, from inequality (5.3), we obtain

$$\begin{aligned} \frac{d}{dt}\mathcal{E}(t) &\leq -2K\Pi_{E_t}(\Delta H) + \frac{1}{2} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + S_2(\mathcal{E}(t))K^2 \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau'} \\ &\quad - \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + S_1(\mathcal{E}(t))(1 + \|\nabla H\|_{L^2(\partial E_t)}^\tau) \|\nabla H\|_{L^2(\partial E_t)}^2 \\ &\leq -2K\Pi_{E_t}(\Delta H) - \frac{1}{2} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 \\ &\quad + S_1(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^2 + S_1(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau} + S_2(\mathcal{E}(t))K^2 \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau'} \end{aligned} \quad (5.4)$$

with $\tau, \tau' > 0$. Then, by the “uniform” Poincaré inequality (4.18) with $f = H$, that is

$$\|H - \bar{H}\|_{L^2(\partial E_t)} \leq C \|\nabla H\|_{L^2(\partial E_t)} \quad (5.5)$$

and by interpolation, we have

$$\|\nabla^2 H\|_{L^2(\partial E_t)}^2 \leq C \|\nabla^4 H\|_{L^2(\partial E_t)} \|\mathbb{H} - \bar{\mathbb{H}}\|_{L^2(\partial E_t)} \leq C \|\nabla^4 H\|_{L^2(\partial E_t)} \|\nabla H\|_{L^2(\partial E_t)}.$$

Hence, using Young inequality again, we get

$$\|\nabla^2 H\|_{L^2(\partial E_t)}^2 \leq \frac{1}{2} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 + C \|\nabla H\|_{L^2(\partial E_t)}^2,$$

that is,

$$-\frac{1}{2} \|\nabla^4 H\|_{L^2(\partial E_t)}^2 \leq -\|\nabla^2 H\|_{L^2(\partial E_t)}^2 + C \|\nabla H\|_{L^2(\partial E_t)}^2.$$

Substituting into inequality (5.4), we conclude

$$\begin{aligned} \frac{d}{dt} \mathcal{E}(t) &\leq -2K \Pi_{E_t}(\Delta H) - \|\nabla^2 H\|_{L^2(\partial E_t)}^2 + S(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^2 \\ &\quad + S(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau} + S(\mathcal{E}(t)) K^2 \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau'}, \end{aligned}$$

with $S = \max\{S_1 + C, S_2\} : [0, +\infty) \rightarrow \mathbb{R}^+$ continuous, monotone nondecreasing and depending on E and M_E .

If we now assume that, for every $t \in [0, T(E_0))$,

$$\Pi_{E_t}(\Delta H) \geq \sigma \|\nabla H\|_{L^2(\partial E_t)}^2 \tag{5.6}$$

for some constant $\sigma > 0$, then there holds (recalling that $K > 2$)

$$\begin{aligned} \frac{d}{dt} \mathcal{E}(t) &\leq -[2K\sigma - S(\mathcal{E}(t)) - 2] \|\nabla H\|_{L^2(\partial E_t)}^2 - 2 \|\nabla H\|_{L^2(\partial E_t)}^2 - \|\nabla^2 H\|_{L^2(\partial E_t)}^2 \\ &\quad + S(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau} + S(\mathcal{E}(t)) K^2 \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau'} \\ &\leq -[2K\sigma - S(\mathcal{E}(t)) - 2] \|\nabla H\|_{L^2(\partial E_t)}^2 - 2(\|\nabla H\|_{L^2(\partial E_t)}^2 + \|\nabla^2 H\|_{L^2(\partial E_t)}^2 / K) \\ &\quad + S(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau} + S(\mathcal{E}(t)) K^2 \|\nabla H\|_{L^2(\partial E_t)}^{2+\tau'} \\ &= -P(\mathcal{E}(t)) \|\nabla H\|_{L^2(\partial E_t)}^2 - 2\mathcal{E}(t)/K \\ &\quad + S(\mathcal{E}(t)) \mathcal{E}(t)^{1+\tau/2} + S(\mathcal{E}(t)) K^2 \mathcal{E}(t)^{1+\tau'/2} \end{aligned}$$

where $P = [2K\sigma - S - 2] : [0, +\infty) \rightarrow \mathbb{R}$ is continuous and monotone nonincreasing, determined by E and M_E and $\tau, \tau' > 0$.

It is then an exercise of qualitative analysis of ordinary differential inequalities, to conclude that if $P(0)$ is positive and $\mathcal{E}(0)$ is small enough (such that $-\mathcal{E}(0)/K + S(\mathcal{E}(0))\mathcal{E}(0)^{1+\tau/2} + S(\mathcal{E}(0))K^2\mathcal{E}(0)^{1+\tau'/2} < 0$, which can be always achieved, once K is fixed), the first term starts and stays negative and the “energy” \mathcal{E} satisfies

$$\frac{d}{dt} \mathcal{E}(t) \leq -\mathcal{E}(t)/K \tag{5.7}$$

for every $t \in [0, T(E_0))$, that is, the function \mathcal{E} is never increasing, so it remains bounded by $\mathcal{E}(0)$ (moreover, it decreases exponentially and converges to zero, as $t \rightarrow +\infty$, if the flow is “eternal”). Thus, choosing an appropriate constant K , by the definition of the function S , it is easy to see that we can make $P(0) > 0$, hence if $\delta > 0$ is small,

since $\mathcal{E}(0) \leq K\mathcal{F}(0) \leq \delta K$, it follows that $\mathcal{E}(0)$ is small enough and we have the above conclusion.

Step 2 (Proof of estimate (5.6)).

We need the following key lemma which is implied by Proposition 2.35 in [10], that actually simply generalizes to any dimension Lemma 2.6 in [1].

Lemma 5.3. *Let $E \subseteq \mathbb{T}^4$ be a strictly stable critical set. For every $\theta \in (0, 1]$ there exist a constant $\sigma_\theta > 0$ such that if $F \in \mathfrak{C}_{2M_E}^1(E)$ satisfies*

$$\text{Vol}(F \triangle E) \leq \delta_0 \quad \text{and} \quad \int_{\partial F} |\nabla H|^2 d\mu \leq \delta_0, \quad (5.8)$$

for $\delta_0 > 0$ small enough, there holds

$$\Pi_F(\varphi) \geq \sigma_\theta \|\varphi\|_{L^2(\partial F)}^2,$$

for all $\varphi \in \tilde{H}^1(\partial F)$ satisfying

$$\min_{\eta \in O_E} \|\varphi - \langle \eta | \nu_F \rangle\|_{L^2(\partial F)} \geq \theta \|\varphi\|_{L^2(\partial F)}$$

where the vector subspace $O_E \subseteq \mathbb{R}^4$ was defined in formula (3.5).

Proof. Representing the boundary of $F \in \mathfrak{C}_{2M_E}^1(E)$ as $\partial F = \{y + \psi_F(y)\nu_E(y) : y \in \partial E\}$ for a smooth function $\psi_F : \partial E \rightarrow \mathbb{R}$, according to Proposition 2.35 in [10], fixed some $p > 3$, there exists a positive constant $C = C(\theta, p)$ such that the conclusion follows if $\|\psi_F\|_{W^{2,p}(\partial E)} \leq C$. This inequality follows if conditions (5.8) hold with δ_0 small enough, by the properties of the function ω stated in Lemma 4.8 (and Sobolev embeddings). \square

We now want to apply this lemma with $F = E_t$ and $\varphi = \Delta H$, for all $t \in [0, T(E_0))$, hence, we need to show that there exists a small constant $\theta > 0$ such that

$$\min_{\eta \in O_E} \|\Delta H - \langle \eta | \nu_t \rangle\|_{L^2(\partial E_t)} \geq \theta \|\Delta H\|_{L^2(\partial E_t)} \quad \text{for all } t \in [0, T(E_0)). \quad (5.9)$$

Considering the special basis $\{e_i\}$ of \mathbb{R}^4 and the associated set $i \in I_E$ in the discussion just after Definition 3.5, by the properties of the function ω stated in Lemma 4.8, if δ_0 is small enough we have that for every $t \in [0, T(E_0))$ the norm $\|\psi_F\|_{W^{4,2}(\partial E)}$ is small, hence the same holds for $\|\psi_F\|_{C^1(\partial E)}$. Then, it follows that there exists a constant $C_0 = C_0(E, M_E) > 0$ such that, for every $i \in I_E$, we have $\|\langle e_i | \nu_t \rangle\|_{L^2(\partial E_t)} \geq C_0 > 0$, holding $\|\langle e_i | \nu_E \rangle\|_{L^2(\partial E)} > 0$ (notice that this argument also shows that, with an appropriate choice of small δ_0 and δ , the condition $\limsup_{t \rightarrow T(E_0)} \|\psi_t\|_{C^1(\partial E)} = 2M_E$ cannot occur). It is then easy to see that the vector $\eta_t \in O_E$ realizing the above minimum for E_t is unique and satisfies

$$\Delta H = \langle \eta_t | \nu_t \rangle + g, \quad (5.10)$$

where $g \in L^2(\partial E_t)$ is a function L^2 -orthogonal (with respect to the measure μ_t on ∂E_t) to the vector subspace of $L^2(\partial E_t)$ spanned by the functions $\langle e_i | \nu_t \rangle$. Moreover, letting

$\eta_t = \eta_t^i e_i$, from relation (5.9) we have

$$\|\Delta H\|_{L^2(\partial E_t)}^2 \geq \|\langle \eta_t | \nu_t \rangle\|_{L^2(\partial E_t)}^2 = \int_{\partial E_t} |\eta_t^i \langle e_i | \nu_t \rangle|^2 d\mu_t \geq C_0^2 |\eta_t^i|^2 = C |\eta_t|^2, \quad (5.11)$$

where C is a constant depending only on E and M_E .

We now argue by contradiction, assuming $\|g\|_{L^2(\partial E_t)} < \theta \|\Delta H\|_{L^2(\partial E_t)}$.

We recall that, thanks to inequality (5.5), we have

$$\int_{\partial E_t} |H - \bar{H}|^2 d\mu_t \leq C \int_{\partial E_t} |\nabla H|^2 d\mu_t \leq C \|\Delta H\|_{L^2(\partial E_t)}^2 \quad (5.12)$$

where the second estimate can be obtained integrating by parts and using the Cauchy–Schwarz inequality.

Hence, by multiplying relation (5.10) by $H - \bar{H}$ and integrating over ∂E_t , we get

$$\begin{aligned} \left| \int_{\partial E_t} (H - \bar{H}) \Delta H d\mu_t \right| &= \left| \int_{\partial E_t} (H - \bar{H}) g d\mu_t \right| \\ &< \theta \|H - \bar{H}\|_{L^2(\partial E_t)} \|\Delta H\|_{L^2(\partial E_t)} \\ &\leq C \theta \|\Delta H\|_{L^2(\partial E_t)}^2, \end{aligned} \quad (5.13)$$

where the equality follows from the identities

$$\int_{\partial E_t} H \nu_t d\mu_t = 0 \quad \text{and} \quad \int_{\partial E_t} \nu_t d\mu_t = 0$$

holding for every embedded hypersurface. We notice that the first identity is a consequence of relation (2.2) and the divergence theorem, while the second is equality (3.3). Then, recalling estimate (5.11) and the fact that g is L^2 -orthogonal to $\langle \eta_t | \nu_t \rangle$, we have

$$\begin{aligned} \|\langle \eta_t | \nu_t \rangle\|_{L^2(\partial E_t)}^2 &= \int_{\partial E_t} \Delta H \langle \eta_t | \nu_t \rangle d\mu_t \\ &= - \int_{\partial E_t} \langle \nabla H | \nabla \langle \eta_t | \nu_t \rangle \rangle d\mu_t \\ &\leq |\eta_t| \|\nabla \nu_t\|_{L^2(\partial E_t)} \|\nabla H\|_{L^2(\partial E_t)} \\ &\leq C \|\Delta H\|_{L^2(\partial E_t)} \|\nabla \nu_t\|_{L^2(\partial E_t)} \left| \int_{\partial E_t} (H - \bar{H}) \Delta H d\mu_t \right|^{1/2} \\ &\leq C \sqrt{\theta} \|\Delta H\|_{L^2(\partial E_t)}^2, \end{aligned}$$

where in the last inequality we used relation (5.13) and we estimated $\|\nabla \nu_t\|_{L^2(\partial E_t)}$ by inequality (5.2) and the fact that $\mathcal{F}(t) \leq \delta_0$, as $\nabla \nu_t = B$ by the Gauss–Weingarten relations (2.1).

If then $\theta > 0$ is chosen so small that $C\sqrt{\theta} < 1 - \theta^2$ in the last inequality, we have a contradiction since equality (5.10) and the fact that $\|g\|_{L^2(\partial E_t)} < \theta \|\Delta H\|_{L^2(\partial E_t)}$ imply (by

L^2 -orthogonality) that

$$\|\langle \eta_t | \nu_t \rangle\|_{L^2(\partial E_t)}^2 > (1 - \theta^2) \|\Delta H\|_{L^2(\partial E_t)}^2.$$

All this argument shows that with such a choice of θ , condition (5.9) holds, hence by Lemma 5.3 we conclude

$$\Pi_{E_t}(\Delta H) \geq \sigma_\theta \|\Delta H\|_{L^2(\partial E_t)}^2 \quad \text{for all } t \in [0, T(E_0)).$$

Then, the second estimate (5.12) clearly proves assumption (5.6) and the proof of monotonicity of \mathcal{E} in Step 1 is concluded. Hence, if δ is small enough, $\mathcal{E}(t)$ remains bounded by δ during the flow, up to the time $t = T(E_0)$, thus the same clearly holds for $\mathcal{F}(t)$.

Step 3 (*Global existence of the flow*).

We have seen at Step 1 that choosing an appropriate constant K , if δ is small enough, then the “energy” $\mathcal{E}(t)$ is uniformly bounded and decreasing. More precisely, integrating the differential inequality (5.7), there holds

$$\mathcal{E}(t) \leq \mathcal{E}(0)e^{-t/K} \leq \delta e^{-t/K} \leq \delta \quad (5.14)$$

hence, we also have $\mathcal{F}(t) \leq \delta e^{-t/K} \leq \delta$, for every $t \in [0, T(E_0))$.

Moreover, at Step 2 we already saw that if δ_0 is chosen small enough,

$$\limsup_{t \rightarrow T(E_0)} \|\psi_t\|_{C^1(\partial E)} = 2M_E$$

is not possible. Hence, in order to obtain the global existence of the flow, we only have to show that also

$$\limsup_{t \rightarrow T(E_0)} \text{Vol}(E_t \Delta E) = \delta_0 \quad (5.15)$$

cannot occur.

We define the following quantity

$$D(t) = \int_{E_t \Delta E} d(x, \partial E) dx = \int_{E_t} d_E(x) dx - \int_E d_E(x) dx, \quad (5.16)$$

where $d_E : N_\varepsilon \rightarrow \mathbb{R}$ is the signed distance function defined in formula (2.5). We observe that,

$$\text{Vol}(E_t \Delta E) \leq C \|\psi_t\|_{L^1(\partial E)} \leq C \|\psi_t\|_{L^2(\partial E)}$$

and

$$\begin{aligned} \|\psi_t\|_{L^2(\partial E)}^2 &= 2 \int_{\partial E} \int_0^{|\psi_t(y)|} t dt d\mu(y) \\ &= 2 \int_{\partial E} \int_0^{|\psi_t(y)|} d(L(y, t), \partial E) dt d\mu(y) \\ &= 2 \int_{E_t \Delta E} d(x, \partial E) JL^{-1}(x) dx \\ &\leq CD(t), \end{aligned}$$

where the constants depend on E and M_E , $L : \partial E \times (-\varepsilon, \varepsilon) \rightarrow N_\varepsilon$ is the smooth diffeomorphism defined in formula (2.6) and JL is its Jacobian. It clearly follows

$$\text{Vol}(E_t \triangle E) \leq C \|\psi_t\|_{L^1(\partial E)} \leq C \|\psi_t\|_{L^2(\partial E)} \leq C \sqrt{D(t)}, \quad (5.17)$$

and

$$D(t) \leq \int_{E_t \triangle E} 2M_E dx = 2M_E \text{Vol}(E_t \triangle E). \quad (5.18)$$

Then, recalling formula (5.16), we compute

$$\frac{d}{dt} D(t) = \frac{d}{dt} \int_{E_t \triangle E} d(x, \partial E) dx = \int_{\partial E_t} d_E \Delta H d\mu_t \leq C \|\Delta H\|_{L^2(\partial E_t)} \leq C \sqrt{\delta} e^{-t/2K},$$

for all $t \leq T(E_0)$, where the last inequality clearly follows from the above estimate (5.14) for $\mathcal{E}(t)$.

By integrating this differential inequality on $[0, \bar{t}]$ with $\bar{t} \in [0, T(E_0))$ and taking into account estimate (5.17), we get

$$\text{Vol}(E_{\bar{t}} \triangle E) \leq C \|\psi_{\bar{t}}\|_{L^2(\partial E)} \leq C \sqrt{D(0) + 2KC\sqrt{\delta}} \leq C \sqrt[4]{\delta},$$

as $D(0) \leq C \text{Vol}(E_0 \triangle E) \leq C\delta$, by inequality (5.18) with $t = 0$. Hence, if $\delta > 0$ is small enough such that $C \sqrt[4]{\delta} < \delta_0$, we have that also condition (5.15) cannot happen.

We conclude that the surface diffusion flow of E_0 exists smooth for every time, moreover $E_t \in \mathfrak{C}_{2M_E}^1(E)$ and

$$\text{Vol}(E_t \triangle E) \leq C \sqrt[4]{\delta}, \quad \int_{\partial E_t} |\nabla^2 H|^2 d\mu_t + \int_{\partial E_t} |\nabla H|^2 d\mu_t \leq \delta e^{-t/K}, \quad (5.19)$$

for every $t \in [0, +\infty)$.

Step 4 (Convergence, up to a subsequence, to a translate of E).

Let $t_i \rightarrow +\infty$, then by estimates (5.19), the sets E_{t_i} satisfy the hypotheses of the last point of Lemma 4.8, hence, up to a (not relabeled) subsequence, we have that there exists a critical set $E' \in \mathfrak{C}_{2M_E}^1(E)$ such that $E_{t_i} \rightarrow E'$ in $W^{3,p}$, that is $\|\psi_{t_i} - \psi\|_{W^{3,p}(\partial E)} \rightarrow 0$ for some $\psi : \partial E \rightarrow \mathbb{R}$ representing $\partial E'$ as a “normal graph” on ∂E . As $\partial E'$ has constant mean curvature and it is a graph over ∂E of a C^1 function (by Sobolev embeddings), it follows by standard regularity theory for quasilinear equations that it is smooth (see [21] for instance), then by Proposition 2.7 in [1] (see also [10, Proposition 2.36]), we have that $E' = E + \eta$ for some (small) $\eta \in \mathbb{R}^4$. Such proposition actually states that E is a strict local minimum for the volume–constrained Area functional, up to translations and that a smooth set “close enough” to E (as E' in our situation) can be a critical set if and only if it is a translate of E .

Step 5 (Smooth exponential convergence of the full sequence).

Arguing similarly as above, we consider the function

$$\bar{D}(t) = \int_{E_t \triangle E'} d(x, \partial E) dx$$

with derivative

$$\frac{d}{dt}\bar{D}(t) = \frac{d}{dt} \int_{E_t \Delta E'} d(x, \partial E) dx = \int_{\partial E_t} \operatorname{sgn}(\psi_t - \psi) d_{\partial E} \Delta H d\mu_t, \quad (5.20)$$

where sgn is the “sign function”. By the exponential second estimate (5.19) and the fact that $E_t \in \mathfrak{C}_{2M_E}^1(E)$, we have

$$\left| \frac{d}{dt}\bar{D}(t) \right| \leq C \|\Delta H\|_{L^2(\partial E_t)} \leq C\sqrt{\delta} e^{-t/2K}$$

for all $t \geq 0$, moreover,

$$\bar{D}(t) \leq \int_{E_t \Delta E'} 2M_E dx = 2M_E \operatorname{Vol}(E_t \Delta E') \leq C \|\psi_t - \psi\|_{L^1(\partial E)} \leq C \|\psi_t - \psi\|_{L^2(\partial E)}$$

which implies $\bar{D}(t_i) \rightarrow 0$, as $i \rightarrow \infty$, by the previous step.

Integrating the differential inequality (5.20), we get

$$\begin{aligned} \bar{D}(t) - \bar{D}(t_i) &= - \int_{t_i}^t \frac{d}{ds} \bar{D}(s) ds \leq \int_t^{+\infty} \left| \frac{d}{ds} \bar{D}(s) \right| ds \leq \int_t^{+\infty} C\sqrt{\delta} e^{-s/2K} ds \\ &\leq 2CK\sqrt{\delta} e^{-t/2K}, \end{aligned}$$

hence, passing to the limit as $i \rightarrow \infty$, we conclude

$$\bar{D}(t) \leq C e^{-t/2K}$$

for every $t \geq 0$, thus $\lim_{t \rightarrow +\infty} \bar{D}(t) = 0$. Then, we have

$$\begin{aligned} \|\psi_t - \psi\|_{L^2(\partial E)}^2 &= 2 \int_{\partial E} \left| \int_{\psi(y)}^{\psi_t(y)} s ds \right| d\mu(y) \\ &= 2 \int_{\partial E} \left| \int_{\psi(y)}^{\psi_t(y)} d(L(y, s), \partial E) ds \right| d\mu(y) \\ &= 2 \int_{E_t \Delta E'} d(x, \partial E) JL^{-1}(x) dx \\ &\leq C\bar{D}(t) \\ &\leq C e^{-t/2K}, \end{aligned}$$

where $L : \partial E \times (-\varepsilon, \varepsilon) \rightarrow N_\varepsilon$ is, as before, the smooth diffeomorphism defined in formula (2.6) with Jacobian JL . By this exponential decay and the uniform bound on $\|\psi_t - \psi\|_{W^{4,2}(\partial E)}$ following from estimates (5.19) by means of Lemma 4.8, we obtain the convergence of the full sequence E_t to E' in $W^{3,p}$.

Finally, we have that the convergence of $E_t \rightarrow E + \eta$ is actually exponentially smooth, by arguing as in the proof of Theorem 5.1 in [17] (see also [9]), that is, via standard parabolic estimates and uniform interpolation inequalities (and Sobolev embeddings), holding the exponential convergence in $W^{3,p}$. \square

6. GENERAL DIMENSIONS

In general dimension $n \in \mathbb{N}$, the main global existence and stability Theorem 5.1 takes the following form.

Theorem 6.1. *Let $E \subseteq \mathbb{T}^n$, for $n \geq 3$, be a strictly stable critical set for the Area functional under a volume constraint and let N_ε be a tubular neighborhood of ∂E , as in formula (2.4). For $M_E < \varepsilon/2$ small enough, there exists $\delta > 0$ such that, if E_0 is a smooth set in $\mathfrak{C}_{M_E}^1(E)$ satisfying $\text{Vol}(E_0) = \text{Vol}(E)$ and*

$$\text{Vol}(E_0 \triangle E) \leq \delta \quad \text{and} \quad \int_{\partial E_0} |\nabla^{n-2} \mathbf{H}|^2 d\mu_0 + \int_{\partial E_0} |\nabla \mathbf{H}|^2 d\mu_0 \leq \delta,$$

then, the unique smooth surface diffusion flow E_t starting from E_0 , given by Proposition 3.3, is defined for all $t \geq 0$. Moreover, E_t converges smoothly to $E' = E + \eta$ exponentially fast as $t \rightarrow +\infty$, for some $\eta \in \mathbb{R}^n$, with the meaning that the sequence of smooth functions $\psi_t : \partial E \rightarrow \mathbb{R}$ representing ∂E_t as “normal graphs” on ∂E , that is,

$$\partial E_t = \{y + \psi_t(y)\nu_E(y) : y \in \partial E\},$$

satisfy, for every $k \in \mathbb{N}$,

$$\|\psi_t - \psi\|_{C^k(\partial E)} \leq C_k e^{-\beta_k t},$$

for every $t \in [0, +\infty)$, for some positive constants C_k and β_k , where $\psi : \partial E \rightarrow \mathbb{R}$ represents $\partial E' = \partial E + \eta$ as a “normal graph” on ∂E .

The case $n = 2$, where the boundary hypersurfaces of strictly stable critical sets are circles or straight curves in \mathbb{T}^2 , was analyzed by Elliott and Garcke in [14]. In the three-dimensional case, this theorem is Theorem 4.3 in the paper [1] by Acerbi, Fusco, Julin and Morini. We are going to list now the appropriate modification to the line of proof in dimension $n = 4$ in order to deal with the general dimensional case, noticing that actually up to Proposition 4.2 all the statements are n -dimensional. See the PhD thesis of Antonia Diana [13] for the full details.

□ Proposition 4.2 in dimension $n \in \mathbb{N}$.

Equation (4.7) must be substituted by

$$\begin{aligned} \frac{d}{dt} \int_{\partial E_t} |\nabla^{n-2} \mathbf{H}|^2 d\mu_t &= -2 \int_{\partial E_t} |\nabla^n \mathbf{H}|^2 d\mu_t + \int_{\partial E_t} \mathfrak{q}^{2n+2}(\nabla^{n-4} \mathbf{B}, \nabla^{n-1} \mathbf{H}) d\mu_t \\ &\quad + \int_{\partial E_t} \mathfrak{q}^{2n+2}(\nabla^{n-3}(\mathbf{B}^2), \nabla^n \mathbf{H}) d\mu_t \end{aligned} \quad (6.1)$$

where

- every “monomial” of $\mathfrak{q}^{2n+2}(\nabla^{n-4} \mathbf{B}, \nabla^{n-1} \mathbf{H})$ has 4 factors in \mathbf{B} , $\nabla \mathbf{H}$ and their covariant derivatives, the factor \mathbf{B} (or \mathbf{H} without derivatives) or its covariant derivative up to $\nabla^{n-4} \mathbf{B}$ is present exactly one time and the other three factors are derivatives of $\nabla \mathbf{H}$ up to $\nabla^{n-1} \mathbf{H}$,
- every “monomial” of $\mathfrak{q}^{2n+2}(\nabla^{n-3}(\mathbf{B}^2), \nabla^n \mathbf{H})$ has 3 factors in \mathbf{B}^2 , $\nabla \mathbf{H}$ and their covariant derivatives, the factor \mathbf{B}^2 or its covariant derivative up to $\nabla^{n-3}(\mathbf{B}^2)$ is present exactly one time, the other two factors are derivatives of $\nabla \mathbf{H}$ up to $\nabla^n \mathbf{H}$.

The proof of this formula can be obtained by following step by step the proof of Proposition 4.2, with the appropriate modifications due to the dimension n . The only term that needs a slightly different treatment is the analogue of the first one in the second line of formula (4.10), coming from the term

$$- \int_{\partial E_t} g(\nabla^{n-2}H, \nabla^{n-2}(\Delta H|B|^2)) d\mu_t$$

which would appear in the right hand side of the first line of the n -dimensional version of computation (4.9). Integrating by parts two times, the integrand becomes a contraction of $\nabla^n H$ with $\nabla^{n-4}(\Delta H|B|^2)$, which clearly is a polynomial of the form $q^{2n+2}(\nabla^{n-3}(B^2), \nabla^n H)$ and we are done.

□ Proposition 4.6 in dimension $n \in \mathbb{N}$.

Inequality (4.19) must be substituted by

$$\frac{d}{dt} \int_{\partial E_t} |\nabla^{n-2}H|^2 d\mu_t \leq -\|\nabla^n H\|_{L^2(\partial E_t)}^2 + C_2(1 + \|\nabla H\|_{L^2(\partial E_t)}^\tau) \|\nabla H\|_{L^2(\partial E_t)}^2 \quad (6.2)$$

with $\tau > 0$ and a constant C_2 depending on E , M_E and, for any $j \leq n-3$, on $\|\nabla^j B\|_{L^{\frac{2n-2}{2j-n+3}}(\partial E_t)}$ if $2j > n-3$, on $\|\nabla^j B\|_{L^6(\partial E_t)}$ if $2j = n-3$ and on $\|\nabla^j B\|_{L^\infty(\partial E_t)}$ if $2j < n-3$ (in particular, on $\|B\|_{L^\infty(\partial E_t)}$, when $n \geq 4$).

The proof of this inequality goes like the proof of Proposition 4.6, by choosing suitable exponents in the interpolation inequalities in dealing with the two polynomial terms

$$\int_{\partial E_t} q^{2n+2}(\nabla^{n-4}B, \nabla^{n-1}H) d\mu_t \quad \text{and} \quad \int_{\partial E_t} q^{2n+2}(\nabla^{n-3}(B^2), \nabla^n H) d\mu_t$$

in equality (6.1). In particular, expanding the iterated derivatives of B^2 in the second one, one gets factors like $\nabla^i B * \nabla^j B$ with $i+j \leq n-3$ that are estimated by the constant C_2 .

□ Lemma 4.8 in dimension $n \in \mathbb{N}$.

Lemma 6.2. Let $E \subseteq \mathbb{T}^n$ be a smooth set and N_ε be a tubular neighborhood of ∂E , as in formula (2.4). For M_E small enough and $\delta > 0$, there exists a constant $C = C(E, M_E, \delta)$ such that if $F \in \mathfrak{C}_{M_E}^1(E)$ with

$$\partial F = \{y + \psi_F(y)\nu_E(y) : y \in \partial E\}$$

for a smooth function $\psi_F : \partial E \rightarrow \mathbb{R}$ and

$$\int_{\partial F} |\nabla^{n-2}H|^2 d\mu + \int_{\partial F} |\nabla H|^2 d\mu \leq \Lambda,$$

there hold

$$\|B\|_{L^\infty(\partial F)} + \|\nabla^{n-3}B\|_{L^{\frac{2n-2}{n-3}}(\partial F)} \leq C \quad \text{and} \quad \|\psi_F\|_{W^{n,2}(\partial E)} \leq C.$$

Moreover, for every $1 \leq p < 1 \leq p < \frac{2n-2}{n-3}$, there exists a monotone nondecreasing function $\omega : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, depending only on E and M_E , with $\lim_{\delta \rightarrow 0^+} \omega(\delta) = 0$ and such that if F satisfies the further condition

$$\text{Vol}(F \Delta E) \leq \delta,$$

then $\|\psi_F\|_{W^{n-1,p}(\partial E)} \leq \omega(\delta)$.

As a consequence, if $E_i \subseteq \mathfrak{C}_{M_E}^1(E)$ is a sequence of smooth sets such that

$$\sup_{i \in \mathbb{N}} \int_{\partial E_i} |\nabla^2 H|^{n-2} d\mu_i + \int_{\partial E_i} |\nabla H|^2 d\mu_i < +\infty,$$

then there exists a (non necessarily smooth) set $E' \in \mathfrak{C}_{M_E}^1(E)$ such that, up to a (non relabeled) subsequence, $E_i \rightarrow E'$ in $W^{n-1,p}$ as $i \rightarrow \infty$, for all $1 \leq p < \frac{2n-2}{n-3}$. Moreover, if

$$\int_{\partial E_i} |\nabla^{n-2} H|^2 d\mu_i + \int_{\partial E_i} |\nabla H|^2 d\mu_i \rightarrow 0,$$

as $i \rightarrow \infty$, the set E' is critical for the volume-constrained Area functional, that is, its mean curvature is constant.

Remark 6.3. We notice that once we control $\|\nabla^{n-3} B\|_{L^{\frac{2n-2}{n-3}}(\partial F)}$ and $\|B\|_{L^\infty(\partial F)}$, by means of the uniform interpolation inequalities, we get an estimate on all the norms of the derivatives of B on which the constant C_2 in inequality (6.2), at the previous point, depends. Hence, C_2 is uniformly bounded for all the sets F like above.

Proof of Theorem 6.1. As in the proof of Theorem 5.1, by choosing M_E small enough in order that all the constants in the inequalities are uniform and after choosing some small δ_0 , we consider the surface diffusion flow E_t starting from $E_0 \in \mathfrak{C}_{M_E}^1(E)$ as in the hypotheses and the maximal time $T(E_0)$ such that $E_t \in C_{2M_E}^1(E)$,

$$\text{Vol}(E_t \Delta E) \leq \delta_0 \quad \text{and} \quad \mathcal{F}_n(t) = \int_{\partial E_t} |\nabla^{n-2} H|^2 d\mu_t + \int_{\partial E_t} |\nabla H|^2 d\mu_t \leq \delta_0,$$

for every $t \in [0, T(E_0))$.

Then, we aim to show that the smooth function ψ_t , which represents the moving boundary ∂E_t as a normal graph on ∂E , does not satisfy the following conditions:

- $\limsup_{t \rightarrow T(E_0)} \|\psi_t\|_{C^1(\partial E)} = 2M_E$
- $\limsup_{t \rightarrow T(E_0)} \mathcal{F}_n(t) = \delta_0$
- $\limsup_{t \rightarrow T(E_0)} \text{Vol}(E_t \Delta E) = \delta_0$

that is, the maximal time $T(E_0) = +\infty$, hence the flow exists for all times.

So, we define for a suitable $K > 2$ the following “energy” function

$$\mathcal{E}_n(t) = \int_{\partial E_t} |\nabla^{n-2} H|^2 d\mu_t + K \int_{\partial E_t} |\nabla H|^2 d\mu_t \geq \mathcal{F}_n(t).$$

Recalling Proposition 4.6 and estimate (6.2) and noticing that Lemma 5.3 holds in any dimension (as it is shown in [10]), following Step 1 and Step 2 in the proof of Theorem 5.1, we have

$$\frac{d}{dt}\mathcal{E}_n(t) \leq -\mathcal{E}_n(t)/K$$

for every $t \in [0, T(E_0))$, that is, the energy \mathcal{E}_n (hence \mathcal{F}_n) is uniformly bounded from above and decreasing. Then, the rest of the proof proceeds as the one of Theorem 5.1. \square

REFERENCES

1. E. Acerbi, N. Fusco, V. Julin, and M. Morini, *Nonlinear stability results for the modified Mullins–Sekerka flow and the surface diffusion flow*, J. Diff. Geom. **113** (2019), 1–53.
2. R. A. Adams and J. F. Fournier, *Sobolev spaces (second edition)*, Pure and Appl. Math., vol. 140, Elsevier/Academic Press, Amsterdam, 2013.
3. T. Aubin, *Some nonlinear problems in Riemannian geometry*, Springer–Verlag, 1998.
4. P. Breuning, *Compactness of immersions with local Lipschitz representation*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **29** (2012), no. 4, 545–572.
5. ———, *C^1 -regularity for local graph representations of immersions*, Trans. Amer. Math. Soc. **365** (2013), no. 12, 6185–6198.
6. ———, *Immersions with bounded second fundamental form*, J. Geom. Anal. **25** (2015), no. 2, 1344–1386.
7. M. Cantor, *Sobolev inequalities for Riemannian bundles*, Bull. Amer. Math. Soc. **80** (1974), 239–243.
8. ———, *Sobolev inequalities for Riemannian bundles*, Differential Geometry, Proc. Sympos. Pure Math., vol. 27, Amer. Math. Soc., 1975, pp. 171–184.
9. D. De Gennaro, A. Diana, A. Kubin, and A. Kubin, *Stability of volume-preserving geometric flows in the flat torus*, ArXiv Preprint Server – <http://arxiv.org>, 2023.
10. S. Della Corte, A. Diana, and C. Mantegazza, *Global existence and stability for the modified Mullins–Sekerka flow and surface diffusion flow*, Math. Engineering **4** (2022), Paper n.054, 104 pp.
11. ———, *Uniform Sobolev, interpolation and Calderón–Zygmund inequalities for graph hypersurfaces*, ArXiv Preprint Server – <http://arxiv.org>, 2023.
12. S. Delladio, *On hypersurfaces in \mathbb{R}^{n+1} with integral bounds on curvature*, J. Geom. Anal **11** (2001), no. 1, 17–42.
13. A. Diana, *in preparation*, Ph.D. thesis, Scuola Superiore Meridionale, Napoli.
14. C. M. Elliott and H. Garcke, *Existence results for diffusive surface motion laws*, Adv. Math. Sci. Appl. **7** (1997), no. 1, 467–490.
15. J. Escher, U. F. Mayer, and G. Simonett, *The surface diffusion flow for immersed hypersurfaces*, SIAM J. Math. Anal. **29** (1998), 1419–1433.
16. J. Escher and P. B. Mucha, *The surface diffusion flow on rough phase spaces*, Discrete Contin. Dyn. Syst. **26** (2010), 431–453.
17. N. Fusco, V. Julin, and M. Morini, *The surface diffusion flow with elasticity in three dimensions*, Arch. Rat. Mech. Anal. **237** (2020), 1325–1382.
18. S. Gallot, D. Hulin, and J. Lafontaine, *Riemannian geometry*, Springer–Verlag, 1990.
19. H. Garcke, *Curvature driven interface evolution*, Jahresber. Dtsch. Math.–Ver. **115** (2013), 63–100.
20. Y. Giga and K. Ito, *On pinching of curves moved by surface diffusion*, Comm. Appl. Anal. **2** (1998), no. 3, 393–405.
21. D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Springer–Verlag, 1983.
22. H. Groemer, *Geometric applications of Fourier series and spherical harmonics*, Encyclopedia of Mathematics and its Applications, vol. 61, Cambridge University Press, Cambridge, 1996.

23. K. Grosse-Brauckmann, *Gyroids of constant mean curvature*, Experiment. Math. **6** (1997), 33–50.
24. K. Grosse-Brauckmann and M. Wohlgemuth, *The gyroid is embedded and has constant mean curvature companions*, Calc. Var. PDE **4** (1996), 499–523.
25. M. E. Gurtin and M. E. Jabbour, *Interface evolution in three dimensions with curvature-dependent energy and surface diffusion: Interface-controlled evolution, phase transitions, epitaxial growth of elastic films*, Arch. Rat. Mech. Anal. **163** (2002), 171–208.
26. R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Diff. Geom. **17** (1982), no. 2, 255–306.
27. K. Ito, *The surface diffusion flow equation does not preserve the convexity*, RIMS Kokyuroku **1105** (1999), 10–21.
28. J. Langer, *A compactness theorem for surfaces with L_p -bounded second fundamental form*, Math. Ann. **270** (1985), 223–234.
29. C. Mantegazza, *Smooth geometric evolutions of hypersurfaces*, Geom. Funct. Anal. **12** (2002), no. 1, 138–182.
30. ———, *Lecture notes on mean curvature flow*, Progress in Mathematics, vol. 290, Birkhäuser/Springer Basel AG, Basel, 2011.
31. C. Mantegazza and A. C. Mennucci, *Hamilton–Jacobi equations and distance functions on Riemannian manifolds*, Appl. Math. Opt. **47** (2003), no. 1, 1–25.
32. T. Miura and S. Okabe, *On the isoperimetric inequality and surface diffusion flow for multiply winding curves*, Arch. Ration. Mech. Anal. **239** (2011), 1111–1129.
33. W. M. Mullins, *Two-dimensional motion of idealized grain boundaries*, J. Appl. Phys. **27** (1956), 900–904.
34. P. Petersen, *Riemannian geometry*, Springer-Verlag, 1998.
35. A. Ros, *Stable periodic constant mean curvature surfaces and mesoscopic phase separation*, Interf. Free Bound. **9** (2007), 355–365.
36. M. Ross, *Schwarz’ P and D surfaces are stable*, Diff. Geom. Appl. **2** (1992), 179–195.
37. G. Wheeler, *Surface diffusion flow near spheres*, Calc. Var. PDE **44** (2012), 131–151.
38. ———, *On the curve diffusion flow of closed plane curves*, Ann. Mat. Pura Appl. (2013), no. 5, 931–950.
39. ———, *Convergence for global curve diffusion flows*, Math. Engineering **4** (2022), no. 1, 1–13.

A. DIANA, SCUOLA SUPERIORE MERIDIONALE, ITALY

Email address: antonia.diana@unina.it

N. FUSCO, UNIVERSITÀ DI NAPOLI FEDERICO II & SCUOLA SUPERIORE MERIDIONALE, ITALY

Email address: n.fusco@unina.it

C. MANTEGAZZA, UNIVERSITÀ DI NAPOLI FEDERICO II & SCUOLA SUPERIORE MERIDIONALE, ITALY

Email address: carlo.mantegazza@unina.it