

ON THE SPECTRUM OF SETS MADE OF CORES AND TUBES

FRANCESCA BIANCHI, LORENZO BRASCO, AND ROBERTO OGNIBENE

Dedicated to Giuseppe Buttazzo, a master of Calculus of Variations

ABSTRACT. We analyze the spectral properties of a particular class of unbounded open sets. These are made of a central bounded “core”, with finitely many unbounded tubes attached to it. We adopt an elementary and purely variational point of view, studying the compactness (or the defect of compactness) of level sets of the relevant constrained Dirichlet integral. As a byproduct of our argument, we also get exponential decay at infinity of variational eigenfunctions. Our analysis includes as a particular case a planar set (sometimes called “bookcover”), already encountered in the literature on curved quantum waveguides. J. Hersch suggested that this set could provide the sharp constant in the *Makai-Hayman inequality* for the bottom of the spectrum of the Dirichlet-Laplacian of planar simply connected sets. We disprove this fact, by means of a singular perturbation technique.

CONTENTS

1. Introduction	2
1.1. Constrained critical points of Dirichlet integrals	2
1.2. Loss of compactness	3
1.3. A class of unbounded sets	4
1.4. A detour: the Makai-Hayman inequality	6
1.5. An example by Hersch	7
1.6. Plan of the paper	9
2. Preliminaries	9
3. The first eigenvalue	12
3.1. Set-up	12
3.2. Existence of a first eigenfunction	13
3.3. Examples	16
4. Higher eigenvalues	18
5. Some estimates for eigenfunctions	21
6. Defect of compactness along the tubes	26
7. Singular perturbation of Hersch’s pipe	29
Appendix A. The thin torsional rigidity	36
References	41

Date: December 30, 2023.

2010 Mathematics Subject Classification. 35P15, 35B38, 58E05.

Key words and phrases. Eigenvalue estimates, Poincaré inequality, inradius, Makai-Hayman inequality, Palais-Smale sequence, curved waveguide.

1. INTRODUCTION

1.1. Constrained critical points of Dirichlet integrals. For an open set $\Omega \subseteq \mathbb{R}^N$, we indicate by $\mathcal{S}_2(\Omega)$ the sphere

$$\mathcal{S}_2(\Omega) = \left\{ \varphi \in W_0^{1,2}(\Omega) : \|\varphi\|_{L^2(\Omega)} = 1 \right\}.$$

Here, by $W_0^{1,2}(\Omega)$ we mean the closure of $C_0^\infty(\Omega)$ in the standard Sobolev space $W^{1,2}(\Omega)$. Then, we define

$$\lambda_1(\Omega) := \inf_{\varphi \in \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla \varphi|^2 dx.$$

This is nothing but the sharp constant in the Poincaré inequality for functions in $W_0^{1,2}(\Omega)$. Of course, we have $\lambda_1(\Omega) = 0$ each time that Ω does not support such an inequality. Necessary and sufficient conditions on Ω assuring that $\lambda_1(\Omega) > 0$ can be found for example in [34, Chapter 15, Section 4].

Even when Ω supports the Poincaré inequality, the value $\lambda_1(\Omega)$ is not necessarily attained, i.e. it may be just an infimum. A sufficient condition for $\lambda_1(\Omega)$ to be a minimum is the compactness of the embedding

$$(1.1) \quad W_0^{1,2}(\Omega) \hookrightarrow L^2(\Omega).$$

In this case, there exists a minimizer $u_1 \in \mathcal{S}_2(\Omega)$ and by optimality it solves in weak sense

$$(1.2) \quad -\Delta u_1 = \lambda u_1, \quad \text{in } \Omega,$$

with $\lambda = \lambda_1(\Omega)$, i.e. $\lambda_1(\Omega)$ is an eigenvalue of the Dirichlet-Laplacian on Ω and u_1 is an associated eigenfunction. Observe that we can always suppose that u_1 is non-negative, since both the constraint and the functional are invariant by the change $\varphi \mapsto |\varphi|$.

However, in this situation, much more is true: the compactness of the embedding (1.1) entails that we can construct recursively a diverging sequence $\{\lambda_j(\Omega)\}_{j \in \mathbb{N} \setminus \{0\}}$ of positive numbers, each one being an eigenvalue of the Dirichlet-Laplacian on Ω . Moreover, if we define¹

$$\text{Spec}(\Omega) = \left\{ \lambda : (1.2) \text{ admits a weak solution } u \in \mathcal{S}_2(\Omega) \right\},$$

we have that

$$\text{Spec}(\Omega) = \left\{ \lambda_j(\Omega) \right\}_{j \in \mathbb{N} \setminus \{0\}}.$$

Finally, each λ_j has the following variational characterization

$$(1.3) \quad \lambda_j(\Omega) = \inf \left\{ \max_{u \in F \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla u|^2 dx : F \subseteq W_0^{1,2}(\Omega) \text{ subspace with } \dim F = j \right\}.$$

We refer for example to [13, Chapter VI] for these facts.

It is noteworthy that, according to Lagrange's multipliers rule, each element of $\text{Spec}(\Omega)$ can be understood as a critical point of the Dirichlet integral

$$\varphi \mapsto \int_{\Omega} |\nabla \varphi|^2 dx,$$

constrained to the “manifold” $\mathcal{S}_2(\Omega)$. In this interpretation, the associated eigenfunctions are the relevant critical points. The first eigenvalue $\lambda_1(\Omega)$ and an associated eigenfunction u_1 correspond to the global constrained minimum and a global minimizer, respectively.

¹The constraint $u \in \mathcal{S}_2(\Omega)$ has no real bearing, since the equation (1.2) is linear. This is just a cheap way of saying that (1.2) admits a non-trivial solution.

Thus, the discussion above entails that when the embedding (1.1) is compact, the Critical Point Theory of this constrained functional is completely clear: there is only a discrete sequence of critical values, which coincides with $\{\lambda_j(\Omega)\}_{j \in \mathbb{N} \setminus \{0\}}$. Actually, a much more sophisticated conclusion can be drawn in this “compact” situation. This is better elucidated by recalling the following definition, which is well-known in Critical Point Theory (see for example [41, Chapter II, Section 2]).

Definition 1.1. Let $\lambda \geq 0$, we say that $\{\varphi_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,2}(\Omega)$ is a *constrained Palais-Smale sequence at the level λ* if the following three properties hold:

- (1) $\varphi_n \in \mathcal{S}_2(\Omega)$, for every $n \in \mathbb{N}$;
- (2) $\lim_{n \rightarrow \infty} \|\nabla \varphi_n\|_{L^2(\Omega)}^2 = \lambda$;
- (3) we have²

$$\lim_{n \rightarrow \infty} \| -\Delta \varphi_n - \lambda \varphi_n \|_{W^{-1,2}(\Omega)} = 0.$$

Then, as a consequence of the previous discussion, the only values λ admitting a constrained Palais-Smale sequence are given by $\{\lambda_j(\Omega)\}_{j \in \mathbb{N} \setminus \{0\}}$. This is due to fact that if (1.1) is compact, then automatically a constrained Palais-Smale sequence would converge in $L^2(\Omega)$ (up to a subsequence). By condition (3), the limit function $u \in \mathcal{S}_2(\Omega)$ would be an eigenfunction and λ an eigenvalue.

1.2. Loss of compactness. What happens when the embedding (1.1) is no more compact? What can be said about the structure of critical values for the constrained Dirichlet integral, in this case? Is it possible to relate the behaviour of critical values to the “defect of compactness” of the embedding (1.1)?

In this paper, we will try to (partially) answer these questions, in a particular class of open sets where the loss of compactness is controlled, in a suitable sense. Before presenting the class of open sets we want to deal with, we wish to make a couple of further observations.

At first, we go back to the global infimum $\lambda_1(\Omega)$. Then we observe that the compactness of the embedding is certainly a too strong condition for this to be a minimum. Indeed, it would be sufficient to know that a certain constrained sublevel set

$$E_\Omega(t) := \left\{ \varphi \in \mathcal{S}_2(\Omega) : \int_\Omega |\nabla \varphi|^2 dx \leq t \right\}, \quad \text{with } t > \lambda_1(\Omega),$$

is relatively compact in $L^2(\Omega)$, to infer the existence of a global minimizer u_1 (and thus of a first eigenfunction). In other words, if compactness of the embedding (1.1) is lost only for “large energy levels”, we can still infer compactness of minimizing sequences.

More generally, one can prove that if we define $\lambda_j(\Omega)$ by (1.3) and compactness still holds for some $E_\Omega(t)$ with $t > \lambda_j(\Omega)$, then we still have room to prove that $\lambda_j(\Omega)$ actually defines a critical value (and thus an eigenvalue of the Dirichlet-Laplacian).

However, even these weaker conditions are in a certain sense too strong. By appealing to Definition 1.1, it would be sufficient to know that $\lambda_j(\Omega)$ defined by (1.3) admits a constrained Palais-Smale sequence (see above), enjoying a suitable form of compactness. More precisely, it would suffice that such a sequence admits a *weakly convergent subsequence* in $L^2(\Omega)$ to a limit function $u \neq 0$. Indeed,

²For every $u \in W_0^{1,2}(\Omega)$, we set

$$\| -\Delta u - \lambda u \|_{W^{-1,2}(\Omega)} = \sup_{\varphi \in C_0^\infty(\Omega)} \left\{ \left| \int_\Omega \langle \nabla u, \nabla \varphi \rangle dx - \lambda \int_\Omega u \varphi dx \right| : \|\varphi\|_{W^{1,2}(\Omega)} = 1 \right\}.$$




FIGURE 1. An example of set verifying Assumptions 1, in dimension $N = 2$ and with $k = 2$ disjoint tubes \mathcal{C}_1 and \mathcal{C}_2 .

even if this is not enough to assure that u belongs to the constraint $\mathcal{S}_2(\Omega)$, nevertheless by linearity of $\varphi \mapsto -\Delta\varphi - \lambda\varphi$ we would get that u is a non-trivial weak solution of (1.2). Accordingly, we could conclude again that $\lambda_j(\Omega)$ is an eigenvalue of the Dirichlet-Laplacian on Ω .

Ca va sans dire, this compactness requirement on constrained Palais-Smale sequences is much weaker than the compactness of the sublevel sets $E_\Omega(t)$.

1.3. A class of unbounded sets. We are interested in open sets made of a “central core” Ω_0 , to which we attach a finite number of infinite cylindrical sets $\mathcal{C}_1, \dots, \mathcal{C}_k$. We admit that these cylindrical sets may have a non-empty pairwise intersection, provided the latter is bounded, while we allow their boundaries to arbitrarily intersect. This class of sets is a generalization of those considered in [10], among others. We also cite [32], where sets of this type have been considered, together with their spectral properties.

More precisely, throughout the whole paper we will make the following

Assumptions 1. For $N \geq 2$, let $\Omega \subseteq \mathbb{R}^N$ be an open set such that

$$\Omega = \Omega_0 \cup \bigcup_{i=1}^k \mathcal{C}_i,$$

where $k \in \mathbb{N} \setminus \{0\}$ and

(A1) $\Omega_0 \subseteq \mathbb{R}^N$ is an open bounded set (it could be empty, see Example 3.2);

(A2) for $i = 1, \dots, k$, the open set \mathcal{C}_i is a cylindrical set given by

$$\mathcal{C}_i := \mathbf{R}_i(E_i \times (0, +\infty)) + \mathbf{t}_i,$$

where:

- $E_i \subseteq \mathbb{R}^{N-1}$ is an open bounded connected set;
- \mathbf{R}_i is a linear isometry represented by an $N \times N$ orthogonal matrix;
- $\mathbf{t}_i \in \mathbb{R}^N$ is a given point;

(A3) there exists $\varrho_0 > 0$ such that

$$\left(\mathcal{C}_i \cap \mathcal{C}_j \right) \setminus B_{\varrho_0}(0) = \emptyset, \quad \text{for every } i \neq j.$$

If we set $\mathbf{e}_N = (0, \dots, 0, 1)$, we also denote

$$\omega_i := \mathbf{R}_i(\mathbf{e}_N), \quad \text{for } i = 1, \dots, k,$$

which coincides with the direction of the axis of the cylindrical set \mathcal{C}_i (see Figure 1).

For these sets, we will prove (see Theorem 3.3) that if

$$\lambda_1(\Omega) < \mathcal{E}(\Omega) := \min \left\{ \lambda_1(E_1), \dots, \lambda_1(E_2) \right\},$$

then there exists a minimizing sequence

$$\{u_n\}_{n \in \mathbb{N}} \subseteq \mathcal{S}_2(\Omega), \quad \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^2 = \lambda_1(\Omega),$$

which is compact in $L^2(\Omega)$. This assures that $\lambda_1(\Omega)$ is attained and there exists a first eigenfunction.

In order to prove this, we will proceed as follows: we “truncate” Ω along the tubes, so to build a sequence of bounded sets $\{\Omega_n\}_{n \in \mathbb{N}}$ exhausting Ω . Then, as a minimizing sequence we will take the sequence made of the first eigenfunctions of these sets, let us call it $\{u_1^n\}_{n \in \mathbb{N}}$. To get compactness of this sequence, we will rely on the equation satisfied by u_1^n . Indeed, this permits to get a Caccioppoli-type inequality “localized at infinity” along the tubes. We then join this estimate with Poincaré’s inequality along the tubes \mathcal{C}_i , for which the relevant constant is given by³ $\lambda_1(E_i)$. By taking into account that $\lambda_1(\Omega) < \mathcal{E}(\Omega)$, we will get that $\{u_1^n\}_{n \in \mathbb{N}}$ must have a uniformly small L^2 norm at infinity. This finally gives the desired compactness, by means of the classical Riesz-Fréchet-Kolmogorov theorem.

In a nutshell: since the global infimum $\lambda_1(\Omega)$ is strictly less than the “energy” of all the tubes, minimizing sequences does not want to “occupy too much” a tube. Indeed, this would raise too much the value of the Dirichlet integral. Rather, they try to concentrate as much as possible towards the “core” Ω_0 . This gives a gain of compactness. It is likely that a similar result could be obtained by using some sophisticated *concentration-compactness* methods *à la* Lions (see for example [40, Theorem 3.1]). On the contrary, the method of proof presented above is simple and elementary, thus we believe it to be of independent interest.

We point out that, as a direct byproduct of this proof, we get that the L^2 norm of a first eigenfunction for Ω must decay exponentially to 0, at infinity. In turn, by means of a standard $L^\infty - L^2$ estimate “localized at infinity” (obtained for example by a classical Moser iteration), we can upgrade this exponential decay to a uniform one (see Theorem 5.1).

Actually, there is nothing specific to λ_1 in the previous argument. More generally, if for every $j \in \mathbb{N} \setminus \{0\}$ we define $\lambda_j(\Omega)$ by (1.3), we can prove that whenever we have

$$\lambda_j(\Omega) < \mathcal{E}(\Omega),$$

then $\lambda_j(\Omega)$ is an eigenvalue (see Theorem 4.1). The argument is essentially the same as above: we rely again on the equations for the first j -eigenfunctions of the truncated sets Ω_n .

From this argument, we see that $\mathcal{E}(\Omega)$ plays an important role: it can be understood as the energy threshold under which some compactness survives and above which compactness *may completely*

³We recall that λ_1 is invariant by orthogonal transformations and translations, thus we have

$$\lambda_1(\mathcal{C}_i) = \lambda_1(E_1 \times (0, +\infty)) = \lambda_1(E_1).$$

fail. We will make this precise and complement our analysis by showing in Proposition 6.1 that *for every*

$$\lambda \geq \mathcal{E}(\Omega),$$

there exists a sequence of “almost critical points” for the Dirichlet integral constrained to the sphere $\mathcal{S}_2(\Omega)$, for which compactness is completely lost, i.e. it weakly converges in $L^2(\Omega)$ to 0. Thus, it is impossible to retrieve a critical point associated to λ from such a sequence. More precisely, by using the definitions recalled above, we will show that for every such λ there exists a constrained Palais-Smale sequence weakly converging to 0. We will construct this sequence “by hands”, exploiting the geometry of our sets.

We can now build a bridge between Spectral Theory and Critical Point Theory: indeed, these particular sequences are nothing but a variational reformulation of *singular Weyl sequences*, appearing in Spectral Theory, see for example [8, Chapter 9, Section 2] and [45, Chapter 6, Section 4]. These sequences are important since they permit to characterize the *essential spectrum* of a self-adjoint operator, see for example [8, Theorem 9.2.2]. In this way, we retrieve a variational characterization of the essential spectrum of the Dirichlet-Laplacian on our sets.

1.4. A detour: the Makai-Hayman inequality. In order to neatly justify our original interest towards this class of sets, it is necessary to take a small *detour*. Thus, let us now briefly turn our attention to an apparently unrelated problem: the so-called *Makai-Hayman inequality* in Spectral Geometry. At this aim, we need to recall the definition of *inradius* r_Ω of an open set $\Omega \subseteq \mathbb{R}^N$. This is the quantity given by

$$r_\Omega := \sup \left\{ r > 0 : \text{there exists } x \in \Omega \text{ such that } B_r(x) \subseteq \Omega \right\}.$$

A remarkable result due to Makai (see [33]) asserts that there exists a universal constant $C > 0$ such that

$$(1.4) \quad \lambda_1(\Omega) \geq \frac{C}{r_\Omega^2},$$

for every $\Omega \subseteq \mathbb{R}^2$ open *simply connected set* with finite inradius. The same result was then rediscovered independently by Hayman in [24], by means of a different proof. For this reason, we will call (1.4) the Makai-Hayman inequality.

For completeness, we mention that this result can be extended to more general planar open sets, having nontrivial topology. These generalizations are contained in a series of papers appeared shortly after Hayman’s one. We cite for example Croke [14], Graversen and Rao [23], Osserman [36] and Taylor [43]. More recently, some of these results have been generalized to the p -Laplacian by Poliquin (see [37]) and to the fractional Laplacian by the first two authors (see [6, 7]).

The validity of (1.4) immediately leads to a natural question: what is the sharp constant in such an inequality? In other words, if for every $\Omega \subseteq \mathbb{R}^2$ open simply connected set with finite inradius, we define the scale invariant quantity

$$(1.5) \quad \rho(\Omega) := r_\Omega^2 \lambda_1(\Omega),$$

we want to determine the value of the following spectral optimization problem

$$C_{\text{MH}} = \inf \left\{ \rho(\Omega) : \Omega \subseteq \mathbb{R}^2 \text{ open simply connected with } r_\Omega < +\infty \right\}.$$




FIGURE 2. The set \mathbf{H} : it can be built by taking the disk $\{(x_1, x_2) : x_1^2 + x_2^2 < 1\}$, removing the segment $[0, 1] \times \{0\}$ and then adding two parallel half-strips.

In his paper [33], Makai showed that

$$\frac{1}{4} \leq C_{\text{MH}} < \frac{\pi^2}{4}.$$

It is noteworthy that

$$\frac{\pi^2}{4} = \inf \left\{ \rho(\Omega) : \Omega \subseteq \mathbb{R}^2 \text{ open convex set with } r_\Omega < +\infty \right\},$$

i.e. this is the sharp constant in (1.4), in the restricted class of convex sets. Such a value is attained for example by infinite strips. We recall that the determination of the sharp constant for convex sets is a celebrated result due to Hersch, see [27, Théorème 8.1].

In particular, we see that relaxing “convexity” to “simple connectedness” certainly lower the value of the optimal constant. However, the exact determination of C_{MH} is still an open problem: at present, the best known result is that

$$0.6197 < C_{\text{MH}} < 2.095.$$

The lower bound is due to Bañuelos and Carroll (see [5, Corollary 6.1]), while the upper bound has been proved by Brown (see [9, Section 5]), by slightly improving a method used in [5]. After these results, essentially no progress has been made on the problem. We refer the reader to [11] and [25] for a comprehensive overview of results in spectral optimization.

1.5. An example by Hersch. We can now explain our interest towards sets having the structure encoded by Assumptions 1. Indeed, in his review of Makai’s paper (see [26]), Hersch suggested that an optimal set providing the sharp value C_{MH} could be the following

$$(1.6) \quad \mathbf{H} := \left\{ (x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 < 1, x < 0 \right\} \cup \left\{ (x_1, x_2) \in \mathbb{R}^2 : 0 < |x_2| < 1, x_1 \geq 0 \right\},$$

see Figure 2. We will call it *Hersch’s pipe*. This is a slit disk, to which two infinite tubes of constant width are attached. Thus, it is not difficult to see that \mathbf{H} is just a particular instance of the sets studied in this paper. More precisely, we observe that \mathbf{H} satisfies Assumptions 1, by taking $N = 2$, $k = 2$ and

$$\begin{aligned} \Omega_0 &= \left\{ (x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 < 1 \right\} \setminus \left([0, 1] \times \{0\} \right) && (\text{slit disk}), \\ \mathcal{C}_1 &= (0, +\infty) \times (0, 1), && \mathcal{C}_2 = (0, +\infty) \times (-1, 0). \end{aligned}$$

It is not difficult to see that

$$\lambda_1(\Omega_0) = \pi^2,$$

with a first eigenfunction given by (in polar coordinates)

$$u(\varrho, \vartheta) = \frac{\sin(\pi \varrho)}{\sqrt{\pi \varrho}} \sin\left(\frac{\vartheta}{2}\right), \quad \text{for } 0 < \varrho < 1, 0 < \vartheta < 2\pi,$$

see for example [44, Exercise 6.8.9]. By strict monotonicity with respect to set inclusion, we thus have $\lambda_1(\mathbf{H}) < \lambda_1(\Omega_0) = \pi^2$, while $r_{\mathbf{H}} = r_{\Omega_0} = 1/2$. Thus, by recalling the notation (1.5), we have

$$\rho(\mathbf{H}) < \rho(\Omega_0) = \left(\frac{1}{2}\right)^2 \pi^2 = \frac{\pi^2}{4}.$$

However, we will show in Section 7 that actually \mathbf{H} is *not an optimal shape* for C_{MH} . We will prove this fact by a *singular perturbation technique*. Namely, we will see that adding a suitable (finite) number of thin tubes on the flat part of $\partial\mathbf{H}$ will decrease the quantity $\rho(\mathbf{H})$. In this part, we will greatly rely on the technique recently studied (in greater generality) in [1] by L. Abatangelo and the third author (see also [20]). We point out that, in order to apply the results of [1], we will need to know that $\lambda_1(\mathbf{H})$ is attained, i.e. it admits a first eigenfunction u_1 . It is precisely here that the results of the first part of the paper (i.e. Theorem 3.3) will be useful.

Let us briefly explain which is the crucial point permitting to lower the value ρ by adding tubes. We first observe that if we add to \mathbf{H} a small tube of width $\varepsilon \ll 1$, by calling \mathbf{H}_ε the new set we would get

$$r_{\mathbf{H}_\varepsilon}^2 - r_{\mathbf{H}}^2 \sim C_1 \varepsilon^2 \quad \text{and} \quad \lambda_1(\mathbf{H}) - \lambda_1(\mathbf{H}_\varepsilon) \sim C_2 \varepsilon^2.$$

That is, both the variation of inradius and that of λ_1 are of order ε^2 . As one may expect, the two variations compete: the inradius *increases*, while λ_1 *decreases*. Thus, in order to decide whether the product $r_{\mathbf{H}_\varepsilon}^2 \lambda_1(\mathbf{H}_\varepsilon)$ decreased or not, a very precise asymptotics for $\lambda_1(\mathbf{H}_\varepsilon)$ would be needed. This seems out of reach. On the other hand, one can observe that if we keep on adding thin tubes of the same width ε , the inradius is *insensitive* to the number n_0 of tubes attached. While, on the contrary, λ_1 is very much affected by these perturbations. Indeed, we can prove that each tube gives a fixed contribution of order ε^2 , proportional to the square of the value of the normal derivative of u_1 at the “junction point” where the tube is attached. It should be noticed that, since u_1 has an exponential decay along the tubes, this contribution tends to be weaker and weaker, as the tube is attached further and further away from the origin.

Incidentally, we notice that \mathbf{H} can also be seen as a particular *curved waveguide*, i.e. it can be regarded as the tubular neighborhood (having width $1/2$) of the $C^{1,1}$ curve

$$(1.7) \quad \gamma(s) = \begin{cases} \left(s - 1, \frac{1}{2}\right), & \text{for } s \geq 1, \\ \left(-\frac{1}{2} \cos\left(\frac{\pi}{2}s\right), \frac{1}{2} \sin\left(\frac{\pi}{2}s\right)\right), & \text{for } -1 < s < 1, \\ \left(-s - 1, -\frac{1}{2}\right), & \text{for } s \leq -1, \end{cases}$$

see Figure 3. The systematic study of the spectral theory of curved waveguides has been initiated in the landmark paper [18] by Exner and Šeba. In [18, Example 4.3] this specific example is called “bookcover”. Without any attempt of completeness, we refer to [3, 19, 22] and [28] for some thorough studies on the spectral properties of these sets.




FIGURE 3. In dashed line, the image of the curve γ defined by (1.7). By taking its tubular neighborhood of width $1/2$, we get the set \mathbf{H} .

1.6. Plan of the paper. In Section 2 we recall some technical facts, which will be useful along the paper. Then in Section 3 we show that if $\lambda_1(\Omega)$ is below the critical threshold $\mathcal{E}(\Omega)$, it admits a first eigenfunction. This section contains also some examples of sets to which our existence result applies. Section 4 generalizes the existence result to higher eigenvalues. In the subsequent Section 5 we proceed to prove some properties of the eigenfunctions: the main result here is the exponential decay to 0 at infinity. To complete the analysis, we discuss in Section 6 the loss of compactness along the tubes, i.e. we prove that every $\lambda \geq \mathcal{E}(\Omega)$ admits a constrained Palais-Smale sequence which “disappears” along one of the tubes \mathcal{C}_i . Finally, in Section 7 we consider Hersch’s pipe and disprove its optimality for the problem of determining the sharp Makai-Hayman inequality. The paper is complemented by an appendix, which contains some technical facts taken from [1].

Acknowledgments. We owe the knowledge of reference [31] to the kind courtesy of Giorgio Talenti, we wish to thank him. We thank Francesca Prinari for bringing our attention to [40]. Part of this research has been done during a visit of R.O. to the University of Ferrara in March 2023, as well as during the conference “*NonPUB23 – Nonlocal and Nonlinear Partial Differential Equations at the University of Bologna*”, held in Bologna in June 2023. Organizers and hosting institutions are gratefully acknowledged.

R.O. has been financially supported by the project *ERC VAREG – Variational approach to the regularity of the free boundaries* (grant agreement No. 853404) and by the INdAM-GNAMPA Project 2022 CUP_E55F22000270001.

F.B. and L.B. have been financially supported by the *Fondo di Ateneo per la Ricerca FAR 2020* and *FAR 2021* of the University of Ferrara.

F.B. and R.O. are members of the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). They are both supported by the INdAM-GNAMPA Project 2023 CUP_E53C22001930001.

2. PRELIMINARIES

The following technical result is quite classical, it will be useful in the paper. We enclose a proof, for completeness.

Lemma 2.1. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set. Let $U \in W_0^{1,2}(\Omega)$ be such that*

$$(2.1) \quad \int_{\Omega} \langle \nabla U, \nabla \varphi \rangle dx = \lambda \int_{\Omega} U \varphi dx, \quad \text{for every } \varphi \in W_0^{1,2}(\Omega),$$

for some $\lambda \geq 0$. Then

$$(2.2) \quad \int_{\Omega} \langle \nabla |U|, \nabla \varphi \rangle dx \leq \lambda \int_{\Omega} |U| \varphi dx, \quad \text{for every } \varphi \in W_0^{1,2}(\Omega) \text{ such that } \varphi \geq 0.$$

Proof. For every $\varepsilon > 0$, we consider the C^2 convex function

$$f_{\varepsilon}(t) = (\varepsilon^2 + t^2)^{\frac{1}{2}} - \varepsilon, \quad \text{for } t \in \mathbb{R}.$$

By the Chain Rule, we have $f_{\varepsilon}(U) \in W_0^{1,2}(\Omega)$ and $\nabla f_{\varepsilon}(U) = f'_{\varepsilon}(U) \nabla U$. We take a non-negative $\eta \in C_0^{\infty}(\Omega)$ and use the test function

$$\varphi = \eta f'_{\varepsilon}(U),$$

in (2.1). This gives

$$\int_{\Omega} \langle \nabla f_{\varepsilon} \circ U, \nabla \eta \rangle dx + \int_{\Omega} f''_{\varepsilon}(U) |\nabla U|^2 \eta dx = \lambda \int_{\Omega} U \eta f'_{\varepsilon}(U) dx.$$

By using the convexity of f_{ε} and the fact that

$$t f'_{\varepsilon}(t) \leq |t|, \quad \text{for every } t \in \mathbb{R},$$

we obtain

$$(2.3) \quad \int_{\Omega} \langle \nabla f_{\varepsilon} \circ U, \nabla \eta \rangle dx \leq \lambda \int_{\Omega} |U| \eta dx.$$

By using the Dominated Convergence Theorem and the form of f_{ε} , it is easily seen that

$$\lim_{\varepsilon \rightarrow 0^+} \|f_{\varepsilon} \circ U - |U|\|_{W^{1,2}(\Omega)} = 0.$$

Thus we can pass to the limit in (2.3). Finally, by a density argument, we can enlarge the class of test functions to $\eta \in W_0^{1,2}(\Omega)$, with $\eta \geq 0$, and conclude the proof. \square

We recall that for a general open set $\Omega \subseteq \mathbb{R}^N$, we have defined for every $j \in \mathbb{N} \setminus \{0\}$

$$\lambda_j(\Omega) := \inf \left\{ \max_{\varphi \in F \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla \varphi|^2 dx : F \subseteq W_0^{1,2}(\Omega) \text{ subspace with } \dim F = j \right\},$$

where

$$\mathcal{S}_2(\Omega) = \left\{ \varphi \in W_0^{1,2}(\Omega) : \|\varphi\|_{L^2(\Omega)} = 1 \right\}.$$

Thanks to its definition, it is easily seen that for every pair of open sets $\Omega', \Omega \subseteq \mathbb{R}^N$, we have

$$(2.4) \quad \Omega' \subseteq \Omega \quad \implies \quad \lambda_j(\Omega) \leq \lambda_j(\Omega'), \quad \text{for every } j \in \mathbb{N} \setminus \{0\}.$$

The next two results are apparently well-known, but we draw the reader's attention to the fact that we do not take any assumption on the open set. In particular, λ_j is not necessarily an eigenvalue.

Lemma 2.2. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set and let $\{\Omega_n\}_{n \in \mathbb{N}}$ be a sequence of open sets such that*

$$\Omega_n \subseteq \Omega_{n+1} \subseteq \Omega \quad \text{for every } n \in \mathbb{N} \quad \text{and} \quad \bigcup_{n \in \mathbb{N}} \Omega_n = \Omega.$$

Then we have

$$\lim_{n \rightarrow \infty} \lambda_j(\Omega_n) = \lambda_j(\Omega), \quad \text{for every } j \in \mathbb{N} \setminus \{0\}.$$

Proof. By (2.4), we already know that the limit of $\lambda_j(\Omega)$ exists and is such that

$$\lim_{n \rightarrow \infty} \lambda_j(\Omega_n) \geq \lambda_j(\Omega).$$

In order to prove the reverse inequality, we take $F \subseteq W_0^{1,2}(\Omega)$ a vector subspace with dimension j . By definition, this means that there exists j linearly independent functions $u_1, \dots, u_j \in W_0^{1,2}(\Omega)$ such that

$$F = \left\{ \sum_{i=1}^j \alpha_i u_i : \alpha_1, \dots, \alpha_j \in \mathbb{R} \right\}.$$

We can suppose that $\{u_1, \dots, u_j\}$ is an orthonormal set in $L^2(\Omega)$, by the Gram-Schmidt orthonormalization process. For every u_i with $i = 1, \dots, j$, there exists a sequence $\{u_i^m\}_{m \in \mathbb{N}} \subseteq C_0^\infty(\Omega)$ such that

$$\lim_{m \rightarrow \infty} \|u_i^m - u_i\|_{W^{1,2}(\Omega)} = 0.$$

We can choose such sequences such that $\{u_1^m, \dots, u_j^m\}$ is an orthonormal set in $L^2(\Omega)$, for every $m \in \mathbb{N}$. We then define

$$F_m = \left\{ \sum_{i=1}^j \alpha_i u_i^m : \alpha_1, \dots, \alpha_j \in \mathbb{R} \right\},$$

and observe that this is a j -dimensional vector subspace of $C_0^\infty(\Omega)$. Since the sequence $\{\Omega_n\}_{n \in \mathbb{N}}$ is exhausting Ω , we have that F_m is actually a j -dimensional vector subspace of $C_0^\infty(\Omega_n)$, for n large enough (depending on m). Thus we get

$$\lim_{n \rightarrow \infty} \lambda_j(\Omega_n) \leq \max_{u \in F_m \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla u|^2 dx.$$

By using the construction of F_m , it is easily seen that

$$\lim_{m \rightarrow \infty} \max_{u \in F_m \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla u|^2 dx = \max_{u \in F \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla u|^2 dx.$$

Finally, by arbitrariness of F , the last two equations in display and the definition of $\lambda_j(\Omega)$ give the desired conclusion. \square

Lemma 2.3. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set, then we have*

$$\lambda_j(\Omega) \leq \lambda_{j+1}(\Omega), \quad \text{for every } j \in \mathbb{N} \setminus \{0\}.$$

Proof. Let $F \subseteq W_0^{1,2}(\Omega)$ be a $(j+1)$ -dimensional vector subspace and let $\{u_1, \dots, u_j, u_{j+1}\}$ be a basis of F . We define \tilde{F} the j -dimensional subspace spanned by $\{u_1, \dots, u_j\}$. Then, $\tilde{F} \subseteq F$ and we have that

$$\lambda_j(\Omega) \leq \max_{u \in \tilde{F} \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla u|^2 dx \leq \max_{u \in F \cap \mathcal{S}_2(\Omega)} \int_{\Omega} |\nabla u|^2 dx.$$

By taking the infimum with respect to $(j+1)$ -dimensional subspaces F , we conclude. \square

3. THE FIRST EIGENVALUE

3.1. **Set-up.** We will use the same notations of Assumptions 1. For any $r > 0$, we set

$$Q_r^i := \mathbf{R}_i((-r, r)^N) + \mathbf{t}_i, \quad \text{for } i = 1, \dots, k,$$

and

$$\Omega_r^i := \Omega_0 \cup (Q_r^i \cap \mathcal{C}_i), \quad \text{for } i = 1, \dots, k.$$

Accordingly, we also set

$$(3.1) \quad \Omega_r = \bigcup_{i=1}^k \Omega_r^i.$$

We then take $r_0 = r_0(\Omega) > 0$ to be large enough such that $\Omega \setminus \Omega_r$ is a disjoint union of k cylindrical sets, for any $r \geq r_0$. More precisely, we have

$$(3.2) \quad \Omega \setminus \Omega_r = \bigcup_{i=1}^k (\mathcal{C}_i \setminus Q_r^i), \quad \text{for } r \geq r_0.$$

The existence of such r_0 is guaranteed by the construction of Ω , see Assumptions 1.

This crucial property entails that a Poincaré inequality holds for functions of $W_0^{1,2}(\Omega)$, even when restricted to $\mathcal{C}_i \setminus Q_r^i$. Namely, we have the following

Lemma 3.1 (Poincaré inequality at infinity). *Let $\Omega \subseteq \mathbb{R}^N$ be an open set satisfying Assumptions 1. With the notations above, we have*

$$\lambda_1(E_i) \int_{\mathcal{C}_i \setminus Q_r^i} |\varphi|^2 dx \leq \int_{\mathcal{C}_i \setminus Q_r^i} |\nabla \varphi|^2 dx,$$

for every $i = 1, \dots, k$, $r \geq r_0$ and every $\varphi \in W_0^{1,2}(\Omega)$.

Proof. We will use the notation $x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R}$. By a simple change of variable, we have that, for any $\varphi \in C_0^\infty(\Omega)$ and any $i = 1, \dots, k$, there holds

$$\int_{\mathcal{C}_i \setminus Q_r^i} |\varphi|^2 dx = \int_r^{+\infty} \int_{E_i} |\varphi(\mathbf{R}_i(x', x_N) + \mathbf{t}_i)|^2 dx' dx_N.$$

For every fixed x_N , we now use the $(N-1)$ -dimensional Poincaré inequality for the function

$$x' \mapsto \varphi(\mathbf{R}_i(x', x_N) + \mathbf{t}_i),$$

which is compactly supported in E_i . This yields

$$\int_{\mathcal{C}_i \setminus Q_r^i} |\varphi|^2 dx \leq \frac{1}{\lambda_1(E_i)} \int_r^{+\infty} \int_{E_i} |\nabla' \varphi(\mathbf{R}_i(x', x_N) + \mathbf{t}_i)|^2 dx' dx_N,$$

where we denoted by ∇' the gradient with respect to x' . By observing that

$$\int_r^{+\infty} \int_{E_i} |\nabla' \varphi(\mathbf{R}_i(x', x_N) + \mathbf{t}_i)|^2 dx' dx_N \leq \int_{\mathcal{C}_i \setminus Q_r^i} |\nabla \varphi|^2 dx,$$

we now easily get the desired Poincaré inequality, for functions in $C_0^\infty(\Omega)$. Finally, by density of $C_0^\infty(\Omega)$ in $W_0^{1,2}(\Omega)$, we conclude the proof. \square

Definition 3.2 (Tubular cut-off functions). We denote by $\eta : \mathbb{R} \rightarrow [0, +\infty)$ a C^∞ function such that

$$0 \leq \eta \leq 1, \quad |\eta'| \leq 2, \quad \eta(t) = \begin{cases} 0, & \text{for } t \leq 0, \\ 1, & \text{for } t \geq 1. \end{cases}$$

For every $i = 1, \dots, k$, we set

$$\eta_{E_i}(x', x_N) = \begin{cases} \eta(x_N), & \text{if } x' \in \overline{E_i}, \\ 0, & \text{otherwise,} \end{cases}$$

and for a pair $0 < r < R$, we also define

$$\eta_{r,R}^{(i)}(x) = \eta_{E_i}(D_{r,R}(\mathbf{R}_i^{-1}(x - \mathbf{t}_i) - r \mathbf{e}_N)), \quad \text{with } D_{r,R} = \begin{bmatrix} & & & 0 \\ & & & \vdots \\ & \text{Id}_{N-1} & & 0 \\ 0 & \dots & 0 & (R-r)^{-1} \end{bmatrix}.$$

Here \mathbf{R}_i and \mathbf{t}_i are still as in Assumptions 1. It is crucial to observe that, by construction, we have

$$\eta_{r,R}^{(i)}(x) \equiv 0, \quad \text{in } \mathcal{C}_i \cap Q_r^i, \quad \eta_{r,R}^{(i)}(x) \equiv 1, \quad \text{in } \mathcal{C}_i \setminus Q_R^i, \quad \left| \nabla \eta_{r,R}^{(i)}(x) \right| \leq \frac{2}{R-r}, \quad \text{in } \mathcal{C}_i.$$

In particular, this acts as a ‘‘cut-off at infinity’’ along the cylindrical set \mathcal{C}_i . In the particular case $R = r + 1$, we will simply use the notation

$$\eta_r^{(i)}(x) := \eta_{E_i}(\mathbf{R}_i^{-1}(x - \mathbf{t}_i) - r \mathbf{e}_N).$$

3.2. Existence of a first eigenfunction.

Theorem 3.3. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set satisfying Assumptions 1. If*

$$(3.3) \quad \lambda_1(\Omega) < \mathcal{E}(\Omega) := \min \left\{ \lambda_1(E_1), \dots, \lambda_1(E_k) \right\}.$$

then $\lambda_1(\Omega)$ is actually a minimum, i.e. there exists a first eigenfunction $u_1 \in W_0^{1,2}(\Omega)$ and $u_1 \geq 0$. Moreover, if Ω is connected, then every other first eigenfunction is proportional to u_1 .

Proof. Let $n \geq r_0$. With the notation of (3.1), we consider $\Omega_n \subseteq \mathbb{R}^N$. Since the latter is an open bounded set, the embedding $W_0^{1,2}(\Omega_n) \hookrightarrow L^2(\Omega_n)$ is compact and the quantity

$$\lambda_1(\Omega_n) = \inf_{\varphi \in \mathcal{S}_2(\Omega_n)} \int_{\Omega_n} |\nabla \varphi|^2 dx,$$

admits a nonnegative minimizer $u_1^n \in \mathcal{S}_2(\Omega_n)$, as recalled in the Introduction. We also observe that

$$(3.4) \quad \lim_{n \rightarrow \infty} \lambda_1(\Omega_n) = \lambda_1(\Omega),$$

thanks to Lemma 2.2. We will prove existence of a first eigenfunction for Ω by using the Direct Method in the Calculus of Variations: as a minimizing sequence, we will take $\{u_1^n\}_{n \in \mathbb{N}}$, with the previous notation. Indeed, in light of (3.4), this is a minimizing sequence.

In what follows, each function u_1^n is considered to be extended by 0 outside Ω_n . We first observe that this sequence is bounded in $W_0^{1,2}(\Omega)$. Thus, there exists $u_1 \in W_0^{1,2}(\Omega)$ such that u_1^n converges weakly to u_1 in $W_0^{1,2}(\Omega)$, up to a subsequence. Observe that the limit function still belongs to $W_0^{1,2}(\Omega)$, because the latter is a weakly closed space. Moreover, by lower semicontinuity, we have

$$\int_{\Omega} |\nabla u_1|^2 dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |\nabla u_1^n|^2 dx = \lambda_1(\Omega).$$

In order to conclude, it is sufficient to upgrade this convergence to the strong one in $L^2(\mathbb{R}^N)$, so to assure that $u_1 \in \mathcal{S}_2(\Omega)$. To this aim, we appeal to the classical Riesz-Fréchet-Kolmogorov Theorem. Since we work with a sequence of functions in $W_0^{1,2}(\Omega)$, which is bounded in the norm of $W^{1,2}(\mathbb{R}^N)$, we just need to verify that the sequence “does not lose mass at infinity”. Namely, we need to prove that there exists $n_0 \in \mathbb{N}$ such that, for every $\varepsilon > 0$, there exists $R_\varepsilon > 0$ such that

$$(3.5) \quad \int_{\mathbb{R}^N \setminus \Omega_{R_\varepsilon}} |u_1^n|^2 dx < \varepsilon, \quad \text{for all } n \geq n_0.$$

By minimality, we have that

$$(3.6) \quad \int_{\mathbb{R}^N} \langle \nabla u_1^n, \nabla \varphi \rangle dx = \lambda_1(\Omega_n) \int_{\mathbb{R}^N} u_1^n \varphi dx, \quad \text{for every } \varphi \in W_0^{1,2}(\Omega_n).$$

We now fix $i = 1, \dots, k$ and test (3.6) with $u_1^n (\eta_r^{(i)})^2 \in W_0^{1,2}(\Omega_n)$, where $\eta_r^{(i)}$ is as⁴ in Definition 3.2 and $r \geq r_0$. This gives

$$(3.7) \quad \int_{\mathbb{R}^N} |\nabla u_1^n|^2 |\eta_r^{(i)}|^2 dx + 2 \int_{\mathbb{R}^N} u_1^n \eta_r^{(i)} \langle \nabla u_1^n, \nabla \eta_r^{(i)} \rangle dx = \lambda_1(\Omega_n) \int_{\mathbb{R}^N} |u_1^n|^2 |\eta_r^{(i)}|^2 dx.$$

By Young's inequality and taking into account the properties of $\eta_r^{(i)}$, for $0 < \delta_n < 1$ we have that

$$\begin{aligned} \int_{\mathbb{R}^N} |\nabla u_1^n|^2 |\eta_r^{(i)}|^2 dx + 2 \int_{\mathbb{R}^N} u_1^n \eta_r^{(i)} \langle \nabla u_1^n, \nabla \eta_r^{(i)} \rangle dx \\ \geq \int_{\mathbb{R}^N} |\nabla u_1^n|^2 |\eta_r^{(i)}|^2 dx - \delta_n \int_{\mathbb{R}^N} |\nabla u_1^n|^2 |\eta_r^{(i)}|^2 dx - \frac{1}{\delta_n} \int_{\mathbb{R}^N} |u_1^n|^2 |\nabla \eta_r^{(i)}|^2 dx \\ \geq (1 - \delta_n) \int_{\mathcal{C}_i \setminus \Omega_{r+1}^i} |\nabla u_1^n|^2 dx - \frac{4}{\delta_n} \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_1^n|^2 dx. \end{aligned}$$

By spending this information in (3.7) and using Lemma 3.1, we obtain that

$$\mathcal{E}(\Omega) (1 - \delta_n) \int_{\mathcal{C}_i \setminus \Omega_{r+1}^i} |u_1^n|^2 dx \leq \frac{4}{\delta_n} \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_1^n|^2 dx + \lambda_1(\Omega_n) \int_{\mathcal{C}_i \setminus \Omega_r^i} |u_1^n|^2 dx.$$

We can decompose the last term as follows

$$\lambda_1(\Omega_n) \int_{\mathcal{C}_i \setminus \Omega_r^i} |u_1^n|^2 dx = \lambda_1(\Omega_n) \int_{\mathcal{C}_i \setminus \Omega_{r+1}^i} |u_1^n|^2 dx + \lambda_1(\Omega_n) \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_1^n|^2 dx.$$

This in turn gives

$$(3.8) \quad \left((1 - \delta_n) \mathcal{E}(\Omega) - \lambda_1(\Omega_n) \right) \int_{\mathcal{C}_i \setminus \Omega_{r+1}^i} |u_1^n|^2 dx \leq \left(\frac{4}{\delta_n} + \lambda_1(\Omega_n) \right) \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_1^n|^2 dx.$$

In light of (3.4) and of the assumption $\lambda_1(\Omega) < \mathcal{E}(\Omega)$, there exists $n_0 \geq r_0$ such that

$$\lambda_1(\Omega_n) \leq \frac{\mathcal{E}(\Omega) + \lambda_1(\Omega)}{2}, \quad \text{for every } n \geq n_0.$$

Thus, we also get

$$\mathcal{E}(\Omega) - \lambda_1(\Omega_n) \geq \frac{\mathcal{E}(\Omega) - \lambda_1(\Omega)}{2} > 0, \quad \text{for every } n \geq n_0$$

⁴This is a feasible test function, since for every $\varphi \in W_0^{1,2}(\Omega_n)$ and $\eta \in C^1(\overline{\Omega_n})$, we have $\varphi \eta \in W_0^{1,2}(\Omega_n)$, as well.

This entails that for every $n \geq n_0$ the following choice is feasible

$$\delta_n = \frac{\mathcal{E}(\Omega) - \lambda_1(\Omega_n)}{2 \mathcal{E}(\Omega)}.$$

From (3.8) we then obtain

$$(3.9) \quad \frac{\mathcal{E}(\Omega) - \lambda_1(\Omega_n)}{2} \int_{C_i \setminus \Omega_{r+1}^i} |u_1^n|^2 dx \leq \left(\frac{8 \mathcal{E}(\Omega)}{\mathcal{E}(\Omega) - \lambda_1(\Omega_n)} + \lambda_1(\Omega_n) \right) \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_1^n|^2 dx.$$

for every $n \geq n_0$. Observe that for $n \geq n_0$ we also have

$$\lambda_1(\Omega_n) \leq \frac{\mathcal{E}(\Omega) + \lambda_1(\Omega)}{2} < \mathcal{E}(\Omega),$$

thus from (3.9) we can get

$$\int_{C_i \setminus \Omega_{r+1}^i} |u_1^n|^2 dx \leq C_\Omega \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_1^n|^2 dx, \quad \text{for } n \geq n_0 \text{ and } i = 1, \dots, k,$$

where $C_\Omega > 0$ is the constant given by

$$C_\Omega := \frac{4}{\mathcal{E}(\Omega) - \lambda_1(\Omega)} \left(\frac{16 \mathcal{E}(\Omega)}{\mathcal{E}(\Omega) - \lambda_1(\Omega)} + \mathcal{E}(\Omega) \right).$$

We can sum over i such an estimate, so to get

$$(3.10) \quad \int_{\mathbb{R}^N \setminus \Omega_{r+1}} |u_1^n|^2 dx = \sum_{i=1}^k \int_{C_i \setminus \Omega_{r+1}^i} |u_1^n|^2 dx \leq C_\Omega \int_{\Omega_{r+1} \setminus \Omega_r} |u_1^n|^2 dx,$$

which holds for $r \geq r_0$ and $n \geq n_0$. It is not difficult to see that this estimate gives the desired uniform decay at infinity: indeed, if we set

$$A_n(r) = \int_{\mathbb{R}^N \setminus \Omega_r} |u_1^n|^2 dx,$$

the previous estimate can be recast into

$$A_n(r+1) \leq C_\Omega (A_n(r) - A_n(r+1)) \quad \text{that is} \quad A_n(r+1) \leq \frac{C_\Omega}{C_\Omega + 1} A_n(r).$$

In particular, by using that $r \mapsto A_n(r)$ is non-increasing, for every $R \geq r_0$ we can obtain⁵

$$A_n(R+1) \leq A_n(r_0 + \lfloor R - r_0 \rfloor + 1) \leq \left(\frac{C_\Omega}{C_\Omega + 1} \right)^{\lfloor R - r_0 \rfloor + 1} A_n(r_0) \leq \left(\frac{C_\Omega}{C_\Omega + 1} \right)^{R - r_0} A_n(r_0).$$

By using that

$$A_n(r_0) \leq \int_{\mathbb{R}^N} |u_1^n|^2 dx = 1,$$

we finally get the desired decay property (3.5). Therefore, up to a subsequence, we have that u_1^n strongly converges to u_1 in $L^2(\Omega)$, as n goes to ∞ . Moreover, $u_1 \geq 0$ since $u_1^n \geq 0$ for all $n \in \mathbb{N}$.

The last part of the statement is a classical fact. This completes the proof. \square

⁵For every $t \in \mathbb{R}$, we denote its *integer part* by

$$\lfloor t \rfloor = \max \{ n \in \mathbb{Z} : t \geq n \}.$$

Remark 3.4. If we admit the cylindrical sets \mathcal{C}_i to have an overlapping with infinite volume, i.e. if condition (A3) is violated, the previous result may fail to be true. As a simple example, take $N = 2$, $k = 2$ and

$$\Omega_0 = (0, 2) \times (0, 2), \quad \mathcal{C}_1 = (0, 2) \times (0, +\infty), \quad \mathcal{C}_2 = (1, 3) \times (0, +\infty).$$

In this case, we have

$$\Omega = \Omega_0 \cup \mathcal{C}_1 \cup \mathcal{C}_2 = (0, 3) \times (0, +\infty),$$

and

$$\lambda_1(\Omega) = \frac{\pi^2}{9} < \min \left\{ \lambda_1(\mathcal{C}_1), \lambda_1(\mathcal{C}_2) \right\} = \frac{\pi^2}{4}.$$

However, it is well-know that $\lambda_1(\Omega)$ is not attained, in this case.

3.3. Examples.

Example 3.1 (Massive core). Let us suppose that $\Omega \subseteq \mathbb{R}^N$ satisfies Assumptions 1, with its core Ω_0 having the following property:

$$\text{if } \mathcal{E}(\Omega) = \lambda_1(E_i), \text{ then } \Omega_0 \cap \mathcal{C}_i \neq \emptyset \quad \text{and} \quad \lambda_1(\Omega_0) \leq \lambda_1(\mathcal{C}_i).$$

In particular, we have $\Omega_0 \subseteq \Omega_0 \cup \mathcal{C}_i \subseteq \Omega$ and $\Omega_0 \cup \mathcal{C}_i$ is a connected open set such that

$$|(\Omega_0 \cup \mathcal{C}_i) \setminus \Omega_0| > 0.$$

Moreover, Ω_0 is bounded and thus $\lambda_1(\Omega_0)$ is actually an eigenvalue. These facts imply that

$$\lambda_1(\Omega_0 \cup \mathcal{C}_i) < \lambda_1(\Omega_0).$$

This in turn gives

$$\lambda_1(\Omega) < \lambda_1(\Omega_0) \leq \lambda_1(\mathcal{C}_i) = \mathcal{E}(\Omega).$$

Thus, we can apply Theorem 3.3.

Example 3.2 (Infinite cross). We take the open set

$$\Omega = \left(\mathbb{R} \times (-1, 1) \right) \times \left((-1, 1) \times \mathbb{R} \right).$$

Observe that such a set satisfies Assumptions 1: indeed, it can be also written as follows

$$\Omega = Q \cup \bigcup_{i=1}^4 \mathcal{C}_i,$$

where Q is the open square with vertices $(0, \pm 2)$ and $(\pm 2, 0)$ and

$$\begin{aligned} \mathcal{C}_1 &= (-1, +\infty) \times (-1, 1), & \mathcal{C}_2 &= (-\infty, 1) \times (-1, 1), \\ \mathcal{C}_3 &= (-1, 1) \times (1, +\infty), & \mathcal{C}_4 &= (-1, 1) \times (-\infty, 1). \end{aligned}$$

We observe that

$$\mathcal{E}(\Omega) = \lambda_1(\mathcal{C}_1) = \dots = \lambda_1(\mathcal{C}_4) = \frac{\pi^2}{4},$$

and that (see [25, Chapter 1])

$$\lambda_1(Q) = \frac{\pi^2}{(2\sqrt{2})^2} + \frac{\pi^2}{(2\sqrt{2})^2} = \mathcal{E}(\Omega).$$

Thus, this set is a particular case of Example 3.1 and we can apply Theorem 3.3. A study of the spectral properties of this set can be also found in [4, 35] and [38].




FIGURE 4. The set of Example 3.2. In dashed line, the square Q such that $\lambda_1(Q) \leq \mathcal{E}(\Omega)$.




FIGURE 5. The set of Example 3.3. In dashed line, the triangle T such that $\lambda_1(T) \leq \mathcal{E}(\Omega)$.

We point out that this set could also be realized simply as the union of the four cylindrical sets $\mathcal{C}_1, \dots, \mathcal{C}_4$, i.e. we could think that $\Omega_0 = \emptyset$. This shows that Theorem 3.3 may cover cases where the core is empty and the “compactness” is created by some non-trivial intersections of the “tubes”.

Example 3.3 (The broken strip). We fix $0 < \vartheta < \pi/2$ and take the open set Ω defined as follows

$$\Omega = \left\{ (x_1, x_2) \in \mathbb{R}^2 : x_1 \tan \vartheta < |x_2| < x_1 \tan \vartheta + \frac{1}{\cos \vartheta} \right\},$$

see Figure 5. This set satisfies Assumptions 1, by taking for example

$$\Omega_0 = T := \left\{ (x_1, x_2) \in \Omega : x_1 < 0 \right\} \quad (\text{isosceles triangle}),$$

and the two half-strips

$$\mathcal{C}_1 = \left\{ (x_1, x_2) \in \Omega : x_1 \cos \vartheta + x_2 \sin \vartheta > 0 \text{ and } x_2 > 0 \right\},$$

$$\mathcal{C}_2 = \left\{ (x_1, x_2) \in \Omega : x_1 \cos \vartheta - x_2 \sin \vartheta > 0 \text{ and } x_2 < 0 \right\}.$$

By [39, Theorem 1.1], we have the following Pólya-type upper bound for triangles

$$\lambda_1(T) \leq \left(\frac{\pi}{3} \right)^2 \left(\frac{L}{A} \right)^2,$$

where L and A are the perimeter and the area of T , respectively. It is easily seen that

$$L = \frac{2}{\sin \vartheta \cos \vartheta} + \frac{2}{\cos \vartheta} \quad \text{and} \quad A = \frac{1}{\sin \vartheta \cos \vartheta},$$

thus we have

$$\lambda_1(T) \leq \left(\frac{\pi}{3}\right)^2 (2 + 2 \sin \vartheta)^2.$$

In particular, for every $0 < \vartheta \leq \pi/6$ we have

$$\lambda_1(T) \leq \left(\frac{\pi}{3}\right)^2 (2 + 2 \sin \vartheta)^2 \leq \pi^2.$$

This entails that for all these angles we have

$$\lambda_1(T) \leq \pi^2 = \lambda_1(\mathcal{C}_1) = \lambda_1(\mathcal{C}_2) = \mathcal{E}(\Omega).$$

Thus, for every $0 < \vartheta \leq \pi/6$, also this set is a particular case of Example 3.1. Thus, we can apply Theorem 3.3 and get existence of a first eigenfunction. We refer to [4] and [15] for further studies on this set.

Example 3.4 (Hersch's pipe). We take the set \mathbf{H} introduced in (1.6). We have already observed in the introduction that \mathbf{H} satisfies Assumptions 1, by taking $N = 2$, $k = 2$ and

$$\Omega_0 = \left\{ (x, y) \in \mathbb{R}^2 : x_1^2 + x_2^2 < 1 \right\} \setminus \left([0, 1] \times \{0\} \right) \quad (\text{slit disk}),$$

$$\mathcal{C}_1 = (0, +\infty) \times (0, 1), \quad \mathcal{C}_2 = (0, +\infty) \times (-1, 0).$$

Moreover, we have already remarked that

$$\lambda_1(\Omega_0) = \pi^2.$$

By observing that $\mathcal{E}(\mathbf{H}) = \lambda_1(\mathcal{C}_1) = \lambda_1(\mathcal{C}_2) = \pi^2$ and that $\Omega_0 \cap \mathcal{C}_i \neq \emptyset$ for both $i = 1, 2$, we see that \mathbf{H} is actually a particular case of Example 3.1. Thus, we can apply Theorem 3.3, here as well.

4. HIGHER EIGENVALUES

In the next result, we still denote by $\mathcal{E}(\Omega)$ the threshold energy defined in (3.3) and we let $\lambda_j(\Omega)$ be as in (1.3).

Theorem 4.1. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set satisfying Assumptions 1. If $\ell \in \mathbb{N} \setminus \{0, 1\}$ is such that*

$$\lambda_\ell(\Omega) < \mathcal{E}(\Omega),$$

then for every $j = 1, \dots, \ell$ we have:

- (1) $\lambda_j(\Omega)$ is an eigenvalue, with associated eigenfunction $u_j \in \mathcal{S}_2(\Omega)$. Moreover, $\{u_1, \dots, u_j\}$ can be chosen to form an orthonormal set in $L^2(\Omega)$ and we have⁶

$$\lambda_j(\Omega) = \min_{u \in \mathcal{S}_2(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^2 dx : \int_{\Omega} \varphi u_m dx = 0, \text{ for } m = 1, \dots, j-1 \right\},$$

with the minimum attained by u_j ;

- (2) *if $\{\Omega_n\}_{n \in \mathbb{N}}$ is the exhausting sequence of sets defined in (3.1), for every $n \in \mathbb{N}$ there exists $\{u_1^n, \dots, u_j^n\} \subseteq \mathcal{S}_2(\Omega_n)$ eigenfunctions of Ω_n associated to $\lambda_1(\Omega_n), \dots, \lambda_j(\Omega_n)$ which form an orthonormal set in $L^2(\Omega_n)$ and such that (up to a subsequence)*

$$\lim_{n \rightarrow \infty} \|u_m^n - u_m\|_{L^2(\Omega)} = 0, \quad \text{for } m = 1, \dots, j.$$

⁶For $j = 1$, it is intended that the orthogonality condition is void.

Proof. We will prove the result by finite induction on $j \in \{1, \dots, \ell\}$. By observing that (see Lemma 2.3)

$$\lambda_1(\Omega) \leq \lambda_\ell(\Omega) < \mathcal{E}(\Omega),$$

we obtain that the case $j = 1$ has been proved in Theorem 3.3.

We now assume that the statement holds for every $j = 1, \dots, \ell - 1$. We need to show that this entails the validity of the case $j = \ell$, as well. In order to do this, we will adapt the same idea of Theorem 3.3, by taking into account the additional difficulties connected with the orthogonality relations.

The inductive assumptions implies that $\lambda_1(\Omega), \dots, \lambda_{\ell-1}(\Omega)$ are eigenvalues of Ω , with an orthonormal set of associated eigenfunctions $\{u_1, \dots, u_{\ell-1}\} \subseteq \mathcal{S}_2(\Omega)$. Moreover, we know that these eigenfunctions can be approximated, strongly in $L^2(\Omega)$, by an orthonormal set $\{u_1^n, \dots, u_{\ell-1}^n\} \subseteq \mathcal{S}_2(\Omega_n)$ of eigenfunctions of Ω_n . We then observe that (see [13, Chapter VI])

$$\begin{aligned} \lambda_\ell(\Omega_n) &= \inf \left\{ \max_{\varphi \in F \cap \mathcal{S}_2(\Omega_n)} \int_{\Omega_n} |\nabla \varphi|^2 dx : F \subseteq W_0^{1,2}(\Omega_n) \text{ subspace with } \dim F = \ell \right\} \\ &= \inf_{\varphi \in \mathcal{S}_2(\Omega_n)} \left\{ \int_{\Omega_n} |\nabla \varphi|^2 dx : \int_{\Omega} \varphi u_j^n = 0, \text{ for } j = 1, \dots, \ell - 1 \right\}. \end{aligned}$$

We take u_ℓ^n to be a minimizer of the last problem, thus by construction $\{u_1^n, \dots, u_{\ell-1}^n, u_\ell^n\}$ is an orthonormal set. Existence of u_ℓ^n is a plain consequence of the compactness of the embedding $W_0^{1,2}(\Omega_n) \hookrightarrow L^2(\Omega_n)$. By Lemma 2.2, we have that

$$\lim_{n \rightarrow \infty} \int_{\Omega_n} |\nabla u_\ell^n|^2 dx = \lim_{n \rightarrow \infty} \lambda_\ell(\Omega_n) = \lambda_\ell(\Omega),$$

thus the sequence $\{u_\ell^n\}_{n \geq r_0} \subseteq \mathcal{S}_2(\Omega_n)$ is bounded in $W_0^{1,2}(\Omega)$. This implies that, up to a subsequence, it weakly converges in $W^{1,2}(\Omega)$ to a function $u_\ell \in W_0^{1,2}(\Omega)$. We claim that u_ℓ is an eigenfunction associated to $\lambda_\ell(\Omega)$, such that

$$(4.1) \quad \lim_{n \rightarrow \infty} \|u_\ell - u_\ell^n\|_{L^2(\Omega)} = 0,$$

$$(4.2) \quad \int_{\Omega} u_\ell u_j = 0, \text{ for all } j = 1, \dots, \ell - 1 \quad \text{and} \quad \int_{\Omega} |u_\ell|^2 dx = 1,$$

and

$$(4.3) \quad \int_{\Omega} |\nabla u_\ell|^2 dx = \min_{\varphi \in \mathcal{S}_2(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^2 dx : \int_{\Omega} \varphi u_j = 0, \text{ for } m = 1, \dots, \ell - 1 \right\},$$

These facts would be sufficient to conclude the proof.

We start from (4.1): this strong convergence can be inferred by repeating verbatim the compactness argument in the proof of Theorem 3.3. We need to rely this time on the equation for u_ℓ^n . Observe that we already know that the sequence weakly converges to u_ℓ , thus once the strong compactness is obtained, the strong limit must be the same. Thus (4.1) is established.

This in particular implies that the second property in (4.2) holds true. As for the orthogonality conditions in (4.2), for $j = 1, \dots, \ell - 1$ we have

$$\left| \int_{\Omega} u_\ell u_j dx \right| = \lim_{n \rightarrow \infty} \left| \int_{\Omega} u_\ell^n u_j dx \right| = \lim_{n \rightarrow \infty} \left| \int_{\Omega} u_\ell^n (u_j - u_j^n) dx \right| \leq \lim_{n \rightarrow \infty} \|u_j - u_j^n\|_{L^2(\Omega)} = 0.$$

Observe that in the second equality we used that u_ℓ^n is orthogonal to u_j^n (see above).

In order to show that u_ℓ is an eigenfunction associated to $\lambda_\ell(\Omega)$, we first observe that u_ℓ is non-trivial, thanks to the normalization condition on the $L^2(\Omega)$ norm. It is then sufficient to pass to the limit in the equation for u_ℓ^n . Indeed, $\{\Omega_n\}_{n \geq r_0}$ is an exhausting sequence for Ω . Thus, for every $\varphi \in C_0^\infty(\Omega)$ we have that this is compactly supported in Ω_n , as well, for n large enough (depending on φ). Hence, from the weak convergence of $\{u_\ell^n\}_{n \in \mathbb{N}}$ and Lemma 2.2, for every fixed $\varphi \in C_0^\infty(\Omega)$ we get

$$0 = \lim_{n \rightarrow \infty} \left(\int_{\Omega_n} \langle \nabla u_\ell^n, \nabla \varphi \rangle dx - \lambda_\ell(\Omega_n) \int_{\Omega_n} u_\ell^n \varphi dx \right) = \int_{\Omega} \langle \nabla u_\ell, \nabla \varphi \rangle dx - \lambda_\ell(\Omega) \int_{\Omega} u_\ell \varphi dx.$$

This is valid for every $\varphi \in C_0^\infty(\Omega)$, thus by density we get that u_ℓ is an eigenfunction, as claimed.

We are only left with proving (4.3). From (4.2) and the fact that u_ℓ is an eigenfunction, we already know that

$$\lambda_\ell(\Omega) = \int_{\Omega} |\nabla u_\ell|^2 dx \geq \inf_{\varphi \in \mathcal{S}_2(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^2 dx : \int_{\Omega} \varphi u_j = 0, \text{ for } m = 1, \dots, \ell - 1 \right\}.$$

On the other hand, for every $u \in \mathcal{S}_2(\Omega)$ which is orthogonal to $u_1, \dots, u_{\ell-1}$ in $L^2(\Omega)$, we can consider the ℓ -dimensional vector subspace of $W_0^{1,2}(\Omega)$ generated by $\{u_1, \dots, u_{\ell-1}, u\}$. Then by definition we have

$$\begin{aligned} \lambda_\ell(\Omega) &\leq \max_{a \in \mathbb{S}^{\ell-1}} \int_{\Omega} \left| \sum_{j=1}^{\ell-1} a_j \nabla u_j + a_\ell \nabla u \right|^2 dx = \max_{a \in \mathbb{S}^{\ell-1}} \left(\sum_{j=1}^{\ell-1} a_j^2 \int_{\Omega} |\nabla u_j|^2 dx + a_\ell^2 \int_{\Omega} |\nabla u|^2 dx \right) \\ &= \max_{a \in \mathbb{S}^{\ell-1}} \left(\sum_{j=1}^{\ell-1} a_j^2 \lambda_j(\Omega) + a_\ell^2 \int_{\Omega} |\nabla u|^2 dx \right). \end{aligned}$$

Here we have used the orthogonality conditions and the fact that

$$\int_{\Omega} \langle \nabla u_j, \nabla u \rangle dx = \lambda_j(\Omega) \int_{\Omega} u_j u dx, \quad \text{for } j = 1, \dots, \ell - 1.$$

We now use the inductive assumption to assure that for $j = 1, \dots, \ell - 1$

$$\lambda_j(\Omega) = \inf_{\varphi \in \mathcal{S}_2(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^2 dx : \int_{\Omega} \varphi u_m = 0, \text{ for } m = 1, \dots, j - 1 \right\} \leq \int_{\Omega} |\nabla u|^2 dx.$$

Thus, from the estimate above we get

$$\lambda_\ell(\Omega) \leq \max_{a \in \mathbb{S}^{\ell-1}} \left(\sum_{j=1}^{\ell-1} a_j^2 \lambda_j(\Omega) + a_\ell^2 \int_{\Omega} |\nabla u|^2 dx \right) \leq \int_{\Omega} |\nabla u|^2 dx.$$

By recalling that u is an arbitrary trial functions for the minimization problem in the right-hand side of (4.3), we get that

$$\lambda_\ell(\Omega) \leq \inf_{u \in \mathcal{S}_2(\Omega)} \left\{ \int_{\Omega} |\nabla u|^2 dx : \int_{\Omega} u u_j = 0, \text{ for } m = 1, \dots, \ell - 1 \right\},$$

as well. Thus (4.3) holds true: observe that we also obtained that the infimum in (4.3) is attained by u_ℓ .

The proof is now over. \square

5. SOME ESTIMATES FOR EIGENFUNCTIONS

In this section, we prove that the eigenfunctions obtained in Sections 3 and 4 satisfy suitable decay estimates at infinity. A similar result is contained in [17, Section 4.2], but here we rely on genuinely nonlinear techniques.

Theorem 5.1. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set satisfying Assumptions 1. We suppose that*

$$\lambda_\ell(\Omega) < \mathcal{E}(\Omega),$$

for some $\ell \in \mathbb{N} \setminus \{0\}$. For every $j = 1, \dots, \ell$, there exists $0 < \beta_j < 1$ and two constants $C_{1,j}, C_{2,j} > 0$ such that for every eigenfunction u_j associated to $\lambda_j(\Omega)$ we have

$$(5.1) \quad \|u_j\|_{L^2(\Omega \setminus \Omega_R)} \leq C_{1,j} \|u_j\|_{L^2(\Omega)} \beta_j^R, \quad \text{for every } R > 0,$$

and

$$(5.2) \quad |u_j(x)| \leq C_{2,j} \|u_j\|_{L^2(\Omega)} \beta_j^R, \quad \text{for every } R > 0 \text{ and for a. e. } |x| > R.$$

Moreover, we also have

$$(5.3) \quad \|u_j\|_{L^\infty(\Omega)} \leq C_3 \left(\lambda_j(\Omega) \right)^{\frac{N}{4}} \|u_j\|_{L^2(\Omega)},$$

for a constant $C_3 = C_3(N) > 0$.

Proof. We first observe that $|u_j| \in W_0^{1,2}(\Omega)$ is a non-negative function such that

$$(5.4) \quad \int_{\Omega} \langle \nabla |u_j|, \nabla \varphi \rangle dx \leq \lambda \int_{\Omega} |u_j| \varphi dx, \quad \text{for every } \varphi \in W_0^{1,2}(\Omega) \text{ such that } \varphi \geq 0,$$

thanks to Lemma 2.1. We divide the proof in three parts, according to the property of u_j we need to prove.

Decay in L^2 . In order to prove (5.1), we first observe that u_j satisfies the following estimate

$$\left((1 - \delta) \mathcal{E}(\Omega) - \lambda_j(\Omega) \right) \int_{\mathcal{C}_i \setminus \Omega_{r+1}^i} |u_j|^2 dx \leq \left(\frac{4}{\delta} + \lambda_j(\Omega) \right) \int_{\Omega_{r+1}^i \setminus \Omega_r^i} |u_j|^2 dx.$$

Indeed, it is sufficient to start from (5.4) and reproduce verbatim the proof of (3.8). We then choose

$$\delta = \frac{\mathcal{E}(\Omega) - \lambda_j(\Omega)}{2 \mathcal{E}(\Omega)}.$$

With simple manipulations, along the lines we used to get (3.10), we now obtain

$$\int_{\mathbb{R}^N \setminus \Omega_{r+1}} |u_j|^2 dx = \sum_{i=1}^k \int_{\mathcal{C}_i \setminus \Omega_{r+1}^i} |u_j|^2 dx \leq C_{\Omega,j} \int_{\Omega_{r+1} \setminus \Omega_r} |u_j|^2 dx.$$

The constant $C_{\Omega,j} > 0$ is given by

$$C_{\Omega,j} := \frac{2}{\mathcal{E}(\Omega) - \lambda_j(\Omega)} \left(\frac{8 \mathcal{E}(\Omega)}{\mathcal{E}(\Omega) - \lambda_j(\Omega)} + \mathcal{E}(\Omega) \right).$$

With same argument as in the proof of Theorem 3.3, the previous estimate permits to infer that

$$\int_{\mathbb{R}^N \setminus \Omega_{R+1}} |u_j|^2 dx \leq \left(\frac{C_{\Omega,j}}{C_{\Omega,j} + 1} \right)^{R-r_0} \int_{\Omega} |u_j|^2 dx, \quad \text{for every } R \geq r_0.$$

By setting

$$\beta_j = \sqrt{\frac{C_{\Omega,j}}{C_{\Omega,j} + 1}} \quad \text{and} \quad C_{1,j} = \left(\frac{C_{\Omega,j}}{C_{\Omega,j} + 1} \right)^{-\frac{r_0}{2} - 1},$$

we get (5.1) for $R \geq r_0 + 1$. On the other hand, for $0 < R < r_0 + 1$ it is sufficient to notice that

$$\|u_j\|_{L^2(\Omega \setminus \Omega_R)} \leq \|u_j\|_{L^2(\Omega)} \leq \|u_j\|_{L^2(\Omega)} \left(\frac{1}{\beta_j^{r_0+1}} \right) \beta_j^R,$$

with the same choice of $0 < \beta_j < 1$ as above.

Global boundedness. We now prove that $u_j \in L^\infty(\Omega)$. In what follows, for simplicity we will simply write u in place of $|u_j|$. For every $\gamma \geq 1$ and $M > 0$, we insert in (5.4) the test function

$$\varphi = u_M^\gamma := (\min\{u, M\})^\gamma.$$

Thanks to the Chain Rule in $W_0^{1,2}(\Omega)$, this is a feasible test function. With standard computations, we obtain

$$(5.5) \quad \frac{4\gamma}{(\gamma+1)^2} \int_{\Omega} \left| \nabla u_M^{\frac{\gamma+1}{2}} \right|^2 dx = \lambda_j(\Omega) \int_{\Omega} u u_M^\gamma.$$

In order to estimate from below the left-hand side, we will use the following two functional inequalities: for $N \geq 3$, the *Sobolev inequality* (see [42])

$$T_N \left(\int_{\mathbb{R}^N} |\varphi|^{2^*} dx \right)^{\frac{2}{2^*}} \leq \int_{\mathbb{R}^N} |\nabla \varphi|^2, \quad \text{for } T_N = N(N-2)\pi \left(\frac{\Gamma(N/2)}{\Gamma(N)} \right)^{\frac{2}{N}},$$

while for $N = 2$ we will use the *Ladyzhenskaya inequality* (see [31, equation (1.11)])

$$(5.6) \quad \pi \left(\int_{\mathbb{R}^2} |\varphi|^4 dx \right) \leq \left(\int_{\mathbb{R}^2} |\nabla \varphi|^2 dx \right) \left(\int_{\mathbb{R}^2} |\varphi|^2 dx \right),$$

both holding for every $\varphi \in W^{1,2}(\mathbb{R}^N)$. By sticking for the moment to the case $N \geq 3$, we then obtain

$$\frac{4\gamma}{(\gamma+1)^2} T_N \left(\int_{\Omega} \left| u_M^{\frac{\gamma+1}{2}} \right|^{2^*} dx \right)^{\frac{2}{2^*}} \leq \lambda_j(\Omega) \int_{\Omega} u u_M^\gamma.$$

We now introduce the parameter

$$\vartheta = \frac{\gamma+1}{2} \geq 1,$$

and rewrite the previous estimate as

$$\left(\int_{\Omega} u_M^{2^*\vartheta} dx \right)^{\frac{2}{2^*}} \leq \frac{\vartheta^2}{2\vartheta-1} \frac{\lambda_j(\Omega)}{T_N} \int_{\Omega} u u_M^{2\vartheta-1} \leq \vartheta \frac{\lambda_j(\Omega)}{T_N} \int_{\Omega} u u_M^{2\vartheta-1}.$$

We then define the recursive sequence of exponents

$$(5.7) \quad \vartheta_0 = 1, \quad \vartheta_{m+1} = \frac{2^*}{2} \vartheta_m = \left(\frac{N}{N-2} \right)^{m+1}.$$

Moreover, with simple manipulations we also get

$$(5.8) \quad \left(\int_{\Omega} u_M^{2\vartheta_{m+1}} dx \right)^{\frac{1}{2\vartheta_{m+1}}} \leq \vartheta_i^{\frac{1}{2\vartheta_m}} \left(\frac{\lambda_j(\Omega)}{T_N} \right)^{\frac{1}{2\vartheta_m}} \left(\int_{\Omega} u u_M^{2\vartheta_m-1} \right)^{\frac{1}{2\vartheta_m}}.$$

We claim at first that (5.8) shows that $u \in L^{2^{\vartheta_m}}(\Omega)$, for every $m \in \mathbb{N}$. We prove this fact by induction: for $m = 0$, we have $2\vartheta_0 = 2$ and $u \in L^2(\Omega)$ holds true by assumption. Let us now assume that $u \in L^{2^{\vartheta_m}}(\Omega)$: this assumption, (5.8) and $u_M \leq u$ imply that

$$\left(\int_{\Omega} u_M^{2^{\vartheta_{m+1}}} dx \right)^{\frac{1}{2^{\vartheta_{m+1}}}} \leq \vartheta_i^{\frac{1}{2^{\vartheta_m}}} \left(\frac{\lambda_j(\Omega)}{T_N} \right)^{\frac{1}{2^{\vartheta_m}}} \left(\int_{\Omega} u^{2^{\vartheta_m}} \right)^{\frac{1}{2^{\vartheta_m}}}.$$

The right-hand side is finite and independent of $M > 0$: by taking the limit as M goes to ∞ and using Fatou's Lemma, we then obtain

$$(5.9) \quad \left(\int_{\Omega} u^{2^{\vartheta_{m+1}}} dx \right)^{\frac{1}{2^{\vartheta_{m+1}}}} \leq \vartheta_m^{\frac{1}{2^{\vartheta_m}}} \left(\frac{\lambda_j(\Omega)}{T_N} \right)^{\frac{1}{2^{\vartheta_m}}} \left(\int_{\Omega} u^{2^{\vartheta_m}} \right)^{\frac{1}{2^{\vartheta_m}}}.$$

Thus $u \in L^{2^{\vartheta_{m+1}}}(\Omega)$, as well.

By observing that $\{\vartheta_m\}_{i \in \mathbb{N}}$ is an increasingly diverging sequence, we then obtain in particular that $u \in L^{\gamma}(\Omega)$, for every $2 \leq \gamma < +\infty$. In order to obtain the boundedness of u , it is now sufficient to use recursively the scheme of reverse Hölder inequalities (5.9): after n iterations, we get

$$(5.10) \quad \left(\int_{\Omega} u^{2^{\vartheta_n}} dx \right)^{\frac{1}{2^{\vartheta_n}}} \leq \left(\prod_{m=0}^{n-1} \vartheta_m^{\frac{1}{2^{\vartheta_m}}} \right) \left(\frac{\lambda_j(\Omega)}{T_N} \right)^{\sum_{m=0}^{n-1} \frac{1}{2^{\vartheta_m}}} \left(\int_{\Omega} u^2 dx \right)^{\frac{1}{2}}.$$

By observing that

$$(5.11) \quad \lim_{n \rightarrow \infty} \sum_{m=0}^{n-1} \frac{1}{2^{\vartheta_m}} = \frac{1}{2} \sum_{m=0}^{\infty} \left(\frac{N-2}{N} \right)^m = \frac{N}{4},$$

and that

$$\lim_{n \rightarrow \infty} \left(\prod_{m=0}^{n-1} \vartheta_m^{\frac{1}{2^{\vartheta_m}}} \right) =: C_N < +\infty,$$

and passing to the limit as n goes to ∞ in (5.10), we finally get

$$\|u\|_{L^{\infty}(\Omega)} \leq C_N \left(\frac{\lambda_j(\Omega)}{T_N} \right)^{\frac{N}{4}} \|u\|_{L^2(\Omega)}.$$

This gives the boundedness of u , in the case $N \geq 3$.

We briefly describe how to adapt the proof to the case $N = 2$: we go back to (5.5) and multiply both sides by the L^2 norm of $u_M^{(\gamma+1)/2}$. This gives

$$\frac{4\gamma}{(\gamma+1)^2} \left(\int_{\Omega} \left| \nabla u_M^{\frac{\gamma+1}{2}} \right|^2 dx \right) \left(\int_{\Omega} u_M^{\gamma+1} dx \right) = \lambda_j(\Omega) \left(\int_{\Omega} u u_M^{\gamma} dx \right) \left(\int_{\Omega} u_M^{\gamma+1} dx \right).$$

We use the Ladyzhenskaya inequality on the left-hand side and use again the parameter $\vartheta = (\gamma+1)/2$. We now get

$$\int_{\Omega} u_M^{4\vartheta} dx \leq \vartheta \frac{\lambda_j(\Omega)}{\pi} \left(\int_{\Omega} u u_M^{2\vartheta-1} dx \right) \left(\int_{\Omega} u_M^{2\vartheta} dx \right).$$

This time we define the recursive sequence of exponents

$$\vartheta_0 = 1, \quad \vartheta_{m+1} = 2\vartheta_m = 2^{m+1},$$

thus, in place of (5.8), we now get

$$\left(\int_{\Omega} u_M^{2\vartheta_{m+1}} dx \right)^{\frac{1}{2\vartheta_{m+1}}} \leq \vartheta_m^{\frac{1}{4\vartheta_m}} \left(\frac{\lambda_j(\Omega)}{\pi} \right)^{\frac{1}{4\vartheta_m}} \left(\int_{\Omega} u u_M^{2\vartheta_{m-1}} \right)^{\frac{1}{4\vartheta_m}} \left(\int_{\Omega} u_M^{2\vartheta_m} dx \right)^{\frac{1}{4\vartheta_m}}.$$

We can now proceed as above, to prove at first that $u \in L^\gamma(\Omega)$ for every $2 \leq \gamma < +\infty$ and then obtain the estimate

$$\|u\|_{L^\infty(\Omega)} \leq C \sqrt{\frac{\lambda_j(\Omega)}{\pi}} \|u\|_{L^2(\Omega)},$$

where $C > 0$ is a universal constant.

Decay in L^∞ . In order to upgrade the decay estimate to the L^∞ norm, we will use again a suitable Moser's iteration, this time "localized at infinity". We keep on using the notation u in place of $|u_j|$ and we recall that r_0 is as in (3.2). We fix $i = 1, \dots, k$ and take $r_0 + 1 \leq r < R$, then in the equation (5.4) we use the test function

$$\varphi = u^\gamma \left(\eta_{r,R}^{(i)} \right)^2,$$

where $\eta_{r,R}^{(i)}$ is the "cut-off at infinity" defined in Definition 3.2 and $\gamma \geq 1$. Observe that this is a feasible test function, thanks to the fact that $u \in L^\infty(\Omega)$, from the previous step of the proof. Thus, in particular $u^\gamma \in W_0^{1,2}(\Omega)$. From the equation, we get

$$(5.12) \quad \frac{4\gamma}{(\gamma+1)^2} \int_{\mathbb{R}^N} \left| \nabla u^{\frac{\gamma+1}{2}} \right|^2 \left(\eta_{r,R}^{(i)} \right)^2 dx + 2 \int_{\mathbb{R}^N} \left\langle \nabla u, \nabla \eta_{r,R}^{(i)} \right\rangle u^\gamma \eta_{r,R}^{(i)} dx \\ \leq \lambda_j(\Omega) \int_{\mathbb{R}^N} u^{\gamma+1} \left(\eta_{r,R}^{(i)} \right)^2 dx.$$

By using the Cauchy-Schwarz and Young inequalities, we have for $\delta > 0$

$$2 \int_{\mathbb{R}^N} \left\langle \nabla u, \nabla \eta_{r,R}^{(i)} \right\rangle u^\gamma \eta_{r,R}^{(i)} dx \geq -\delta \int_{\mathbb{R}^N} |\nabla u|^2 u^{\gamma-1} \left(\eta_{r,R}^{(i)} \right)^2 dx - \frac{1}{\delta} \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx \\ = -\delta \frac{4}{(\gamma+1)^2} \int_{\mathbb{R}^N} \left| \nabla u^{\frac{\gamma+1}{2}} \right|^2 \left(\eta_{r,R}^{(i)} \right)^2 dx \\ - \frac{1}{\delta} \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx.$$

We choose $\delta = \gamma/2$ and insert this estimate in (5.12), this gives⁷

$$\frac{2\gamma}{(\gamma+1)^2} \int_{\mathbb{R}^N} \left| \nabla u^{\frac{\gamma+1}{2}} \right|^2 \left(\eta_{r,R}^{(i)} \right)^2 dx \leq \frac{2}{\gamma} \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx \\ + \lambda_j(\Omega) \int_{\mathbb{R}^N} u^{\gamma+1} \left(\eta_{r,R}^{(i)} \right)^2 dx.$$

We sum on both sides the quantity

$$\frac{2\gamma}{(\gamma+1)^2} \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx,$$

⁷Observe that both integrals on the right-hand side are finite for every $\gamma \geq 1$, thanks to the fact that $u \in L^2(\Omega) \cap L^\infty(\Omega)$.

and use that

$$\int_{\mathbb{R}^N} \left| \nabla u^{\frac{\gamma+1}{2}} \right|^2 \left(\eta_{r,R}^{(i)} \right)^2 dx + \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx \geq \frac{1}{2} \int_{\mathbb{R}^N} \left| \nabla \left(u^{\frac{\gamma+1}{2}} \eta_{r,R}^{(i)} \right) \right|^2 dx.$$

We thus obtain

$$\begin{aligned} \frac{\gamma}{(\gamma+1)^2} \int_{\mathbb{R}^N} \left| \nabla \left(u^{\frac{\gamma+1}{2}} \eta_{r,R}^{(i)} \right) \right|^2 dx &\leq \left[\frac{2}{\gamma} + \frac{2\gamma}{(\gamma+1)^2} \right] \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx \\ &\quad + \lambda_j(\Omega) \int_{\mathbb{R}^N} u^{\gamma+1} \left(\eta_{r,R}^{(i)} \right)^2 dx. \end{aligned}$$

We now polish a little bit this estimate. We multiply both sides by $(\gamma+1)^2/\gamma$ and use that

$$\frac{(\gamma+1)^2}{\gamma^2} \leq 4 \quad \text{and} \quad \frac{(\gamma+1)^2}{\gamma} \leq 2(\gamma+1), \quad \text{for } \gamma \geq 1,$$

so to get

$$\int_{\mathbb{R}^N} \left| \nabla \left(u^{\frac{\gamma+1}{2}} \eta_{r,R}^{(i)} \right) \right|^2 dx \leq 10 \int_{\mathbb{R}^N} \left| \nabla \eta_{r,R}^{(i)} \right|^2 u^{\gamma+1} dx + 2(\gamma+1) \lambda_j(\Omega) \int_{\mathbb{R}^N} u^{\gamma+1} \left(\eta_{r,R}^{(i)} \right)^2 dx.$$

Finally, from this we can obtain

$$(5.13) \quad \int_{\mathbb{R}^N} \left| \nabla \left(u^{\frac{\gamma+1}{2}} \eta_{r,R}^{(i)} \right) \right|^2 dx \leq 2(\gamma+1) \left(3 + \lambda_j(\Omega) \right) \int_{\mathbb{R}^N} \left[\left| \nabla \eta_{r,R}^{(i)} \right|^2 + \left(\eta_{r,R}^{(i)} \right)^2 \right] u^{\gamma+1} dx.$$

We restrict for simplicity to the case $N \geq 3$. By using Sobolev's inequality in the left-hand side of (5.13), we obtain

$$(5.14) \quad \left(\int_{\mathbb{R}^N} \left(u^{\frac{\gamma+1}{2}} \eta_{r,R}^{(i)} \right)^{2^*} dx \right)^{\frac{2}{2^*}} \leq \frac{2}{T_N} (\gamma+1) \left(3 + \lambda_j(\Omega) \right) \int_{\mathbb{R}^N} \left[\left| \nabla \eta_{r,R}^{(i)} \right|^2 + \left(\eta_{r,R}^{(i)} \right)^2 \right] u^{\gamma+1} dx.$$

We now use the properties of the cut-off functions, encoded by Definition 3.2. From (5.14), we get for $i = 1, \dots, k$ and $r_0 \leq r < R$

$$(5.15) \quad \left(\int_{\mathcal{C}_i \setminus Q_R^i} \left(u^{\frac{\gamma+1}{2}} \right)^{2^*} dx \right)^{\frac{2}{2^*}} \leq \frac{2}{T_N} (\gamma+1) \left(3 + \lambda_j(\Omega) \right) \left[\frac{4}{(R-r)^2} + 1 \right] \int_{\mathcal{C}_i \setminus Q_r^i} u^{\gamma+1} dx.$$

As in the previous step, we introduce the parameter $\vartheta = (\gamma+1)/2$ and rewrite (5.15) as follows

$$\left(\int_{\mathcal{C}_i \setminus Q_R^i} u^{2^* \vartheta} dx \right)^{\frac{1}{2^* \vartheta}} \leq \left(\frac{4}{T_N} \left(3 + \lambda_j(\Omega) \right) \right)^{\frac{1}{2\vartheta}} \vartheta^{\frac{1}{2\vartheta}} \left[\frac{4}{(R-r)^2} + 1 \right]^{\frac{1}{2\vartheta}} \left(\int_{\mathcal{C}_i \setminus Q_r^i} u^{2\vartheta} dx \right)^{\frac{1}{2\vartheta}}.$$

We take $R_0 \geq r_0$ and use the previous estimate with the choices

$$r = r_m := (R_0 + 2) - \frac{1}{2^m}, \quad R = r_{m+1}, \quad \text{for } m \in \mathbb{N},$$

and $\vartheta = \vartheta_m$, where the latter is again the sequence defined in (5.7). We thus get

$$\begin{aligned} \left(\int_{\mathcal{C}_i \setminus Q_{r_{m+1}}^i} u^{2^* \vartheta_{m+1}} dx \right)^{\frac{1}{2^* \vartheta_{m+1}}} &\leq \left(\frac{4}{T_N} \left(3 + \lambda_j(\Omega) \right) \right)^{\frac{1}{2\vartheta_m}} \vartheta_m^{\frac{1}{2\vartheta_m}} (16 \cdot 4^m + 1)^{\frac{1}{2\vartheta_m}} \\ &\quad \times \left(\int_{\mathcal{C}_i \setminus Q_{r_m}^i} u^{2\vartheta_m} dx \right)^{\frac{1}{2\vartheta_m}}. \end{aligned}$$

We iterate this estimate, starting from $m = 0$: after n steps, we get

$$(5.16) \quad \left(\int_{\mathcal{C}_i \setminus Q_{r_n}^i} u^{2\vartheta_n} dx \right)^{\frac{1}{2\vartheta_n}} \leq \left(\frac{4}{T_N} (3 + \lambda_j(\Omega)) \right)^{\frac{1}{2} \sum_{m=0}^{n-1} \frac{1}{\vartheta_m}} \prod_{m=0}^{n-1} \left\{ \frac{1}{\vartheta_m} (16 \cdot 4^m + 1) \right\}^{\frac{1}{2\vartheta_m}} \\ \times \left(\int_{\mathcal{C}_i \setminus Q_{R_0+1}^i} u^2 dx \right)^{\frac{1}{2}}.$$

We now pass to the limit as n goes to ∞ in (5.16). By recalling (5.11) and observing that

$$\lim_{n \rightarrow \infty} \prod_{m=0}^{n-1} \left\{ \frac{1}{\vartheta_m} (16 \cdot 4^m + 1) \right\}^{\frac{1}{2\vartheta_m}} =: C_N < +\infty, \quad \lim_{n \rightarrow \infty} r_n = R_0 + 2,$$

so to get

$$\|u\|_{L^\infty(\mathcal{C}_i \setminus Q_{R_0+2}^i)} \leq C \left(\frac{3 + \lambda_j(\Omega)}{T_N} \right)^{\frac{N}{4}} \left(\int_{\mathcal{C}_i \setminus Q_{R_0+1}^i} u^2 dx \right)^{\frac{1}{2}} \\ \leq C \left(\frac{3 + \lambda_j(\Omega)}{T_N} \right)^{\frac{N}{4}} \left(\int_{\Omega \setminus \Omega_{R_0+1}} u^2 dx \right)^{\frac{1}{2}}.$$

Since this holds for every $i = 1, \dots, k$, we finally get for every $R_0 \geq r_0$

$$\|u\|_{L^\infty(\Omega \setminus \Omega_{R_0+2})} \leq C \left(\frac{3 + \lambda_j(\Omega)}{T_N} \right)^{\frac{N}{4}} \left(\int_{\Omega \setminus \Omega_{R_0+1}} u^2 dx \right)^{\frac{1}{2}}.$$

This is now enough to conclude the proof for $N \geq 3$. For $N = 2$ we may conclude, analogously to the previous step, by using Ladyzhenskaya's inequality (5.6) in place of the Sobolev inequality. \square

Remark 5.2 (Quality of the constants). By inspecting the proof of the previous result, we easily see that both the base β_j and the constant $C_{1,j}$ depend on $\lambda_j(\Omega)$ and on the crucial gap

$$\mathcal{E}(\Omega) - \lambda_j(\Omega) > 0.$$


In particular, we have

$$\beta_j \nearrow 1 \quad \text{and} \quad C_{1,j} \nearrow 1, \quad \text{as} \quad \mathcal{E}(\Omega) - \lambda_j(\Omega) \searrow 0.$$

The estimate (5.3) is classical, here the constant C_3 only depends on the dimension N , through the sharp constant in the Sobolev inequality (see for example [21, Theorem 8.15]). See also [16, Example 2.1.8] for the same estimate, with a dimension-free constant, and [12, Theorem 2] for a sharp version of this estimate. Finally, the constant $C_{2,j}$ depends on N , the eigenvalue $\lambda_j(\Omega)$ and the constant $C_{1,j}$.

6. DEFECT OF COMPACTNESS ALONG THE TUBES

In this section, we analyze what happens for energy levels above the threshold $\mathcal{E}(\Omega)$. We will see that compactness is completely lost, in a suitable sense. We will need the definition of *constrained Palais-Smale sequence*, given in Definition 1.1. The main result of this section is the following

FIGURE 6. The construction of the sequence φ_n in the proof of Proposition 6.1.

Proposition 6.1. *Let $\Omega \subseteq \mathbb{R}^N$ be an open set satisfying Assumptions 1. Then every λ such that*

$$\lambda \geq \mathcal{E}(\Omega) = \min \left\{ \lambda_1(E_1), \dots, \lambda_1(E_k) \right\},$$

admits a constrained Palais-Smale sequence which is weakly converging to 0.

Proof. We give an explicit construction of such a sequence. Without loss of generality, we can assume that

$$\mathcal{E}(\Omega) = \lambda_1(E_1).$$

Let us indicate by $\psi_1 \in W_0^{1,2}(E_1)$ a first positive eigenfunction of E_1 . For simplicity, we take it normalized, i.e. with unit L^2 norm. Up to a rigid movement, we can suppose that

$$E_1 \subseteq \mathbb{R}^{N-1} \times \{0\} \quad \text{and} \quad \mathcal{C}_1 = E_1 \times (0, +\infty).$$

By using the notation $x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R}$, for $n \in \mathbb{N} \setminus \{0\}$ we set $\lambda_n = \lambda + 1/n$ and we take the function

$$\varphi_n(x', x_N) = \psi_1(x') \zeta_n(x_N),$$

where

$$\zeta_n(t) = \begin{cases} \sin \left(\sqrt{\lambda_n - \lambda_1(E_1)} (t - r_0) \right), & \text{if } r_0 < t < r_0 + R_n := r_0 + \frac{2n\pi}{\sqrt{\lambda_n - \lambda_1(E_1)}}, \\ 0, & \text{otherwise,} \end{cases}$$

and $r_0 = r_0(\Omega)$ is the same as in Subsection 3.1. The function φ_n is then extended by 0 to the remainder of Ω , i.e. it is supported in the cylindrical set \mathcal{C}_1 . By construction, we have that this weakly solves

$$-\Delta \varphi_n = \lambda_n \varphi_n, \quad \text{in } E_1 \times (r_0, r_0 + R_n).$$

Its L^2 norm is given by

$$\int_{\Omega} |\varphi_n|^2 dx = \left(\int_{E_1} |\psi_1|^2 dx' \right) \left(\int_{r_0}^{r_0+R_n} |\zeta_n|^2 dx_N \right) = \left(\int_0^{R_n} \left(\sin \left(\sqrt{\lambda_n - \lambda_1(E_1)} \tau \right) \right)^2 d\tau \right) = \frac{R_n}{2}.$$

We then set

$$U_n(x) = \frac{\varphi_n(x)}{\|\varphi_n\|_{L^2(\Omega)}} = \sqrt{\frac{2}{R_n}} \psi_1(x') \zeta_n(x_N).$$

By construction, it is easily seen that it converges to zero weakly in $L^2(\Omega)$, as well as in $W_0^{1,2}(\Omega)$.

We need to verify that this is a Palais-Smale sequence at the level λ . Its Dirichlet integral is given by

$$\begin{aligned} \int_{\Omega} |\nabla U_n|^2 dx &= \frac{2}{R_n} \left(\int_{E_1} |\nabla' \psi_1|^2 dx' \right) \left(\int_{r_0}^{r_0+R_n} |\zeta_n|^2 dx_N \right) \\ &\quad + \frac{2}{R_n} \left(\int_{E_1} |\psi_1|^2 dx' \right) \left(\int_{r_0}^{r_0+R_n} |\zeta_n'|^2 dx_N \right) \\ &= \lambda_1(E_1) + \frac{2}{R_n} (\lambda_n - \lambda_1(E_1)) \int_0^{R_n} \left(\cos \left(\sqrt{\lambda_n - \lambda_1(E_1)} \tau \right) \right)^2 d\tau = \lambda_n = \lambda + \frac{1}{n}, \end{aligned}$$

where we used that

$$\int_0^{R_n} \left(\cos \left(\sqrt{\lambda_n - \lambda_1(E_1)} \tau \right) \right)^2 d\tau = \int_0^{R_n} \left(\sin \left(\sqrt{\lambda_n - \lambda_1(E_1)} \tau \right) \right)^2 d\tau = \frac{R_n}{2}.$$

Thus, we obtained that $\{U_n\}_{n \in \mathbb{N} \setminus \{0\}}$ satisfies properties (1) and (2) in Definition 1.1. In order to verify point (3), we take $\varphi \in C_0^\infty(\Omega)$. We first observe that by standard Elliptic regularity, we have $\psi_1 \in C^\infty(E_1)$ (see for example [21, Corollary 8.11]). Thus, in particular, we get that

$$U_n \in C^2(\overline{\mathcal{O}} \times [r_0, r_0 + R_n]),$$

for every \mathcal{O} compactly contained in E_1 . We have enough regularity to justify the following identities: by using the equation satisfied by U_n and the Divergence Theorem, we get

$$\begin{aligned} \int_{\Omega} \langle \nabla U_n, \nabla \varphi \rangle dx &= \int_{E_1} \int_{r_0}^{r_0+R_n} \langle \nabla U_n, \nabla \varphi \rangle dx' dx_N \\ &= \int_{E_1} \int_{r_0}^{r_0+R_n} \operatorname{div}(\nabla U_n \varphi) dx' dx_N - \int_{E_1} \int_{r_0}^{r_0+R_n} \Delta U_n \varphi dx' dx_N \\ &= \int_{E_1} \frac{\partial U_n}{\partial x_N}(x', r_0 + R_n) \varphi(x', r_0 + R_n) dx' \\ &\quad - \int_{E_1} \frac{\partial U_n}{\partial x_N}(x', r_0) \varphi(x', r_0) dx' + \lambda_n \int_{\Omega} U_n \varphi dx. \end{aligned}$$

With simple manipulations and by recalling that $\lambda_n = \lambda + 1/n$, we get

$$\begin{aligned} \left| \int_{\Omega} \langle \nabla U_n, \nabla \varphi \rangle dx - \lambda \int_{\Omega} U_n \varphi dx \right| &\leq \frac{1}{n} \|U_n\|_{L^2(\Omega)} \|\varphi\|_{L^2(\Omega)} \\ &\quad + \left\| \frac{\partial U_n}{\partial x_N} \right\|_{L^2(E_1 \times \{r_0\})} \|\varphi\|_{L^2(E_1 \times \{r_0\})} \\ &\quad + \left\| \frac{\partial U_n}{\partial x_N} \right\|_{L^2(E_1 \times \{r_0+R_n\})} \|\varphi\|_{L^2(E_1 \times \{r_0+R_n\})}. \end{aligned}$$

By using the properties⁸ of U_n and the trace inequality for the trial function φ , we then obtain

$$\left| \int_{\Omega} \langle \nabla U_n, \nabla \varphi \rangle dx - \lambda \int_{\Omega} U_n \varphi dx \right| \leq \frac{C}{\sqrt{n}} \|\varphi\|_{W^{1,2}(\Omega)},$$

for some C independent of both n and φ . This finally shows that $\{U_n\}_{n \in \mathbb{N} \setminus \{0\}}$ is a constrained P.-S. sequence at the level λ . \square

Remark 6.2. As recalled in the Introduction, a constrained Palais-Smale sequence weakly converging to 0 can be seen as a variational reformulation of the concept of *singular Weyl sequence*, appearing in the Spectral Theory of self-adjoint operators, see for example [8, Chapter 9, Section 2] and [45, Chapter 6, Section 4]. With this in mind, according to [8, Theorem 9.2.2], the previous result shows that every $\lambda \geq \mathcal{E}(\Omega)$ belongs to the *essential spectrum* of the Dirichlet-Laplacian on Ω .

Remark 6.3. Under the assumptions of the previous result, in general it is not true that *every* constrained Palais-Smale sequence at a level $\lambda \geq \mathcal{E}(\Omega)$ weakly converges to 0. For example, by taking

$$\Omega = \left((-1, 1) \times (-1, 1) \right) \cup \left((2, +\infty) \times (-1, 1) \right),$$

we see that this set admits an eigenfunction $u \in W_0^{1,2}(\Omega)$, given by the first eigenfunction of $(-1, 1) \times (-1, 1)$, i.e. associated to the eigenvalue

$$\lambda = \lambda_1((-1, 1) \times (-1, 1)) = \frac{\pi^2}{2} > \frac{\pi^2}{4} = \lambda_1((2, +\infty) \times (-1, 1)) = \mathcal{E}(\Omega).$$

Thus the constant sequence $u_1^n = u$ is a Palais-Smale sequence at a level λ larger than $\mathcal{E}(\Omega)$, but of course it is not weakly converging to 0.

7. SINGULAR PERTURBATION OF HERSCH'S PIPE

We indicate by \mathbf{H} the set of Example 3.4. We define $I := (-1, 1)$ and, for $n \in \mathbb{N} \setminus \{0\}$, $0 < \varepsilon < 1/(2n)$ and $i = 0, \dots, n$, we denote

$$\Sigma_i^\varepsilon := 2 + \frac{i}{n} + \varepsilon I = \left(2 + \frac{i}{n} - \varepsilon, 2 + \frac{i}{n} + \varepsilon \right) \subseteq \mathbb{R} \quad \text{and} \quad T_i^\varepsilon := \Sigma_i^\varepsilon \times [1, 2).$$

We also denote

$$\Gamma_n^\varepsilon := \bigcup_{i=0}^n (\Sigma_i^\varepsilon \times \{1\}) \quad \text{and} \quad \mathbf{H}_n^\varepsilon := \mathbf{H} \cup \bigcup_{i=0}^n T_i^\varepsilon.$$

We can observe that the pair $(\mathbf{H}, \Gamma_n^\varepsilon)$ satisfies Assumptions 2 in Appendix A with $\Omega = \mathbf{H}$ and $\Sigma = \Gamma_n^\varepsilon$. Accordingly, with the notations of Appendix A we have $\Omega_\Sigma = \mathbf{H}_n^\varepsilon$.

The following simple regularity result is not optimal, but it will be largely sufficient for our needs. It asserts that the first eigenfunctions of \mathbf{H} is C^2 up to the boundary, in the flat upper part.

⁸Observe that by construction we have

$$\frac{\partial U_n}{\partial x_N}(x', r_0) = \sqrt{\frac{2}{R_n}} \psi_1(x') \zeta'_n(r_0) = \sqrt{\lambda_n - \lambda_1(E_1)} \sqrt{\frac{2}{R_n}} \psi_1(x'),$$

and

$$\frac{\partial U_n}{\partial x_N}(x', r_0 + R_n) = \sqrt{\frac{2}{R_n}} \psi_1(x') \zeta'_n(r_0 + R_n) = \sqrt{\lambda_n - \lambda_1(E_1)} \sqrt{\frac{2}{R_n}} \psi_1(x').$$




FIGURE 7. The construction of \mathbf{H}_n^ε , with $n = 3$: we add to \mathbf{H} an array of $n+1 = 4$ equally spaced vertical tubes. Each tube has height 1 and width 2ε , with $\varepsilon \ll 1$.

Lemma 7.1. *Let $u_1 \in \mathcal{S}_2(\mathbf{H})$ be the first positive eigenfunction of \mathbf{H} . Then for every $0 < \tau < L$ we have*

$$u_1 \in C^2\left([\tau, L] \times [\tau, 1]\right).$$

In particular, the function $\partial u_1 / \partial x_2$ has a C^1 trace on $(0, +\infty) \times \{1\}$. Moreover, for every $U \in W_0^{1,2}(\mathbf{H}_n^\varepsilon)$ the following identity holds

$$\int_{\mathbf{H}} \langle \nabla u_1, \nabla U \rangle dx = \lambda_1(\mathbf{H}) \int_{\mathbf{H}} u_1 U dx + \int_{\Gamma_\varepsilon} \frac{\partial u_1}{\partial x_2} U d\mathcal{H}^1.$$

Proof. For $\tau > 0$, we set $A_\tau = (\tau/2, +\infty) \times (0, 1)$ and extend u_1 to $A'_\tau = (\tau/2, +\infty) \times (0, 2)$ by odd reflection with respect to the second variable. By construction, we have that this is a weak solution of

$$-\Delta u = \lambda_1(\mathbf{H}) u, \quad \text{in } A'_\tau.$$

Let us still denote it by u_1 , for simplicity. By standard Elliptic Regularity (see [21, Corollary 8.11]), we have that u_1 is actually $C^\infty(E)$ for every open set E compactly contained in A'_τ . This shows in particular that u_1 is C^2 on $[\tau, L] \times [\tau, 1]$.

Let us now take $U \in C_0^\infty(\mathbf{H}_n^\varepsilon)$, we observe that this is not a feasible test function for the equation of u_1 . Since U is compactly supported, there exists $0 < \delta \ll 1$ such that its support is contained in

$$\mathbf{H}_n^\varepsilon(\delta) := \left\{ x \in \mathbf{H} : \text{dist}(x, \partial \mathbf{H}_n^\varepsilon) \geq \delta, x_1 \leq \frac{1}{\delta} \right\}.$$

By the previous point and the fact that $u_1 \in C^\infty(\mathbf{H})$, we know that

$$u_1 \in C^2\left(\mathbf{H}_n^\varepsilon(\delta) \cap \{(x_1, x_2) : x_2 \leq 1\}\right).$$

In particular u_1 solves in classical sense the equation, on this set. By the Divergence Theorem, we thus have

$$\begin{aligned} \lambda_1(\mathbf{H}) \int_{\mathbf{H}} u_1 U \, dx &= \lambda_1(\mathbf{H}) \int_{\mathbf{H}_n^\varepsilon(\delta) \cap \{x_2 \leq 1\}} u_1 U \, dx = - \int_{\mathbf{H}_n^\varepsilon(\delta) \cap \{x_2 \leq 1\}} \Delta u_1 U \, dx \\ &= - \int_{\mathbf{H}_n^\varepsilon(\delta) \cap \{x_2 \leq 1\}} \operatorname{div}(\nabla u_1 U) \, dx \\ &\quad + \int_{\mathbf{H}_n^\varepsilon(\delta) \cap \{x_2 \leq 1\}} \langle \nabla u_1, \nabla U \rangle \, dx \\ &= - \int_{\Gamma_n^\varepsilon} \frac{\partial u}{\partial x_2} U \, d\mathcal{H}^1 + \int_{\mathbf{H}} \langle \nabla u_1, \nabla U \rangle \, dx. \end{aligned}$$

This is the claimed identity, for a test function $U \in C_0^\infty(\mathbf{H}_n^\varepsilon)$. By a density argument (and using the continuity of the trace operator), we can then conclude that the identity holds for test functions in $W_0^{1,2}(\mathbf{H}_n^\varepsilon)$, as well. \square

We now present some results, aimed at giving an asymptotic estimate for $\lambda_1(\mathbf{H}_n^\varepsilon)$, as ε goes to 0. We will crucially exploit the concept of *thin torsional rigidity*, see Appendix A below. In particular, we will need the following quantity

$$\mathbf{T}_n^\varepsilon := \sup_{\varphi \in C_0^\infty(\mathbf{H}_n^\varepsilon)} \left\{ 2 \int_{\Gamma_n^\varepsilon} \frac{\partial u_1}{\partial x_2} \varphi \, d\mathcal{H}^1 - \int_{\mathbf{H}_n^\varepsilon} |\nabla \varphi|^2 \, dx \right\},$$

which is well-defined, in light of the construction and Lemma 7.1. We then observe that, with the notation of Definition A.1, we have

$$(7.1) \quad \mathbf{T}_n^\varepsilon = \mathcal{T}_{\Omega_\Sigma}(\Sigma; f), \quad \text{with } \Omega = \mathbf{H}, \Sigma = \Gamma_n^\varepsilon \text{ and } f = \frac{\partial u_1}{\partial x_2}.$$

The following two results will permit to quantify the decay rate to 0 of this quantity, as ε goes to 0.

Lemma 7.2 (Upper bound). *With the notation above, we have*

$$\mathbf{T}_n^\varepsilon \leq 2(n+1)\varepsilon^2 \left\| \frac{\partial u_1}{\partial x_2} \right\|_{L^\infty([1,4] \times \{1\})}^2, \quad \text{for every } n \in \mathbb{N} \setminus \{0\} \text{ and } 0 < \varepsilon \leq \frac{1}{4n}.$$

Proof. First of all, by construction we have $\Gamma_n^\varepsilon \subseteq [1,4] \times \{1\}$. Thus we get

$$\left\| \frac{\partial u_1}{\partial x_2} \right\|_{L^\infty(\Gamma_n^\varepsilon)} \leq \left\| \frac{\partial u_1}{\partial x_2} \right\|_{L^\infty([1,4] \times \{1\})}.$$

The last term is finite thanks to Lemma 7.1. By (7.1), we can thus apply Lemma A.4. This entails that

$$\mathbf{T}_n^\varepsilon \leq \gamma(\mathbf{H}_n^\varepsilon) \left\| \frac{\partial u_1}{\partial x_2} \right\|_{L^\infty([1,4] \times \{1\})}^2 \mathcal{H}^1(\Gamma_n^\varepsilon) = 2\varepsilon(n+1)\gamma(\mathbf{H}_n^\varepsilon) \left\| \frac{\partial u_1}{\partial x_2} \right\|_{L^\infty([1,4] \times \{1\})}^2,$$

where $\gamma(\mathbf{H}_n^\varepsilon)$ is the sharp trace-type constant defined in Lemma A.4. We will show that $\gamma(\mathbf{H}_n^\varepsilon) \leq \varepsilon$. By definition, we have

$$\begin{aligned} \gamma(\mathbf{H}_n^\varepsilon) &= \sup_{\varphi \in W_0^{1,2}(\mathbf{H}_n^\varepsilon) \setminus \{0\}} \frac{\int_{\Gamma_n^\varepsilon} |\varphi|^2 d\mathcal{H}^1}{\int_{\mathbf{H}_n^\varepsilon} |\nabla \varphi|^2 dx} = \sup_{\varphi \in C_0^\infty(\mathbf{H}_n^\varepsilon) \setminus \{0\}} \frac{\int_{\Gamma_n^\varepsilon} |\varphi|^2 d\mathcal{H}^1}{\int_{\mathbf{H}_n^\varepsilon} |\nabla \varphi|^2 dx} \\ &= \sup_{\varphi \in C_0^\infty(\mathbf{H}_n^\varepsilon) \setminus \{0\}} \frac{\sum_{i=0}^n \int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{\mathbf{H}_n^\varepsilon} |\nabla \varphi|^2 dx} \end{aligned}$$

We take $\varphi \in C_0^\infty(\mathbf{H}_n^\varepsilon) \setminus \{0\}$ and observe that

$$(7.2) \quad \text{if } \int_{T_i^\varepsilon} |\nabla \varphi|^2 dx = 0, \quad \text{then } \int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1 = 0.$$

Indeed, the first condition implies that φ is constant in the tube T_i^ε . Since φ is compactly supported in \mathbf{H}_n^ε and $\partial T_i^\varepsilon \cap \partial \mathbf{H}_n^\varepsilon \neq \emptyset$, this shows that φ must identically vanish on the tube $\overline{T_i^\varepsilon}$. Finally, since $\Sigma_i^\varepsilon \times \{1\} \subseteq \partial T_i^\varepsilon$, we get that φ identically vanishes on $\Sigma_i^\varepsilon \times \{1\}$, as well.

Thanks to (7.2), we can write

$$\gamma(\mathbf{H}_n^\varepsilon) = \sup_{\varphi \in C_0^\infty(\mathbf{H}_n^\varepsilon) \setminus \{0\}} \left\{ \frac{\sum_{i=0}^n \int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{\mathbf{H}_n^\varepsilon} |\nabla \varphi|^2 dx} : \exists i_0 \in \{0, \dots, n\} \text{ such that } \int_{T_{i_0}^\varepsilon} |\nabla \varphi|^2 dx \neq 0 \right\}.$$

Let us take $\varphi \in C_0^\infty(\mathbf{H}_n^\varepsilon) \setminus \{0\}$ which is admissible for the maximization problem on the right-hand side. Let $J_\varphi \subseteq \{0, \dots, n\}$ be the set of indices such that

$$\int_{T_i^\varepsilon} |\nabla \varphi|^2 dx \neq 0, \quad \text{for } i \in J_\varphi.$$

In light of (7.2) and the definition of J_φ , we then have⁹

$$\begin{aligned} \frac{\sum_{i=0}^n \int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{\mathbf{H}_n^\varepsilon} |\nabla \varphi|^2 dx} &\leq \frac{\sum_{i \in J_\varphi} \int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\sum_{i \in J_\varphi} \int_{T_i^\varepsilon} |\nabla \varphi|^2 dx} \leq \max_{i \in J_\varphi} \frac{\int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{T_i^\varepsilon} |\nabla \varphi|^2 dx} \\ &\leq \max_{i=1, \dots, n} \sup_{\varphi \in C_0^\infty(T_i^\varepsilon) \setminus \{0\}} \frac{\int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{T_i^\varepsilon} |\nabla \varphi|^2 dx}. \end{aligned}$$

Finally, by scaling and translating, we see that

$$\sup_{\varphi \in C_0^\infty(T_i^\varepsilon) \setminus \{0\}} \frac{\int_{\Sigma_i^\varepsilon \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{T_i^\varepsilon} |\nabla \varphi|^2 dx} = \varepsilon \sup_{\varphi \in C_0^\infty(\mathbb{I} \times [1, 2]) \setminus \{0\}} \frac{\int_{(-1, 1) \times \{1\}} |\varphi|^2 d\mathcal{H}^1}{\int_{(-1, 1) \times [1, 2]} |\nabla \varphi|^2 dx} \leq \varepsilon.$$

In the last estimate we used the following trace inequality

$$\int_{\mathbb{I} \times [1, 2]} |\nabla \varphi|^2 dx \geq \int_{\mathbb{I}} \left(\int_1^2 \left| \frac{\partial \varphi}{\partial x_2} \right|^2 dx_2 \right) dx_1 \geq \int_{\mathbb{I}} |\varphi(x_1, 1)|^2 dx_1 = \int_{\mathbb{I} \times \{1\}} |\varphi|^2 d\mathcal{H}^1,$$

which holds for every $\varphi \in C_0^\infty(\mathbb{I} \times [1, 2])$. This discussion proves that $\gamma(\mathbf{H}_n^\varepsilon) \leq \varepsilon$, as claimed. \square

Lemma 7.3 (Asymptotical lower bound). *With the notation above, there exists a universal constant $\alpha > 0$ such that for every $n \in \mathbb{N} \setminus \{0\}$ we have*

$$\mathbf{T}_n^\varepsilon \geq \alpha \varepsilon^2 \sum_{i=0}^n \left(\frac{\partial u_1}{\partial x_2} \left(2 + \frac{i}{n}, 1 \right) \right)^2 + o(\varepsilon^2), \quad \text{as } \varepsilon \searrow 0.$$

Proof. By using (7.1) and the super-additivity of the thin torsional rigidity (see Lemma A.3), we get

$$\mathbf{T}_n^\varepsilon \geq \sum_{i=0}^n \mathcal{T}_{\mathbf{H} \cup T_i^\varepsilon} \left(\Sigma_i^\varepsilon \times \{1\}; \frac{\partial u_1}{\partial x_2} \right).$$

Observe that we used that $\partial u_1 / \partial x_2$ has constant sign on Γ_n^ε . On the other hand, as a consequence of Theorem A.8 applied with

$$\mathbf{p}_i = \left(2 + \frac{i}{n}, 1 \right), \quad \Sigma_i = (\mathbb{I} \times \{1\}) + \mathbf{p}_i, \quad f_i = \frac{\partial u_1}{\partial x_2},$$

⁹We use the elementary inequality

$$\frac{\sum_{i=1}^n a_i}{\sum_{i=1}^n b_i} \leq \max \left\{ \frac{a_i}{b_i} : i = 1, \dots, n \right\}, \quad \text{for all } a_i \geq 0, b_i > 0.$$

we know that for every $i = 0, \dots, n$ we have

$$\mathcal{T}_{\mathbf{H} \cup T_i^\varepsilon} \left(\Sigma_i^\varepsilon \times \{1\}; \frac{\partial u_1}{\partial x_2} \right) = \alpha_i \left(\frac{\partial u_1}{\partial x_2} \left(2 + \frac{i}{n}, 1 \right) \right)^2 \varepsilon^2 + o(\varepsilon^2), \quad \text{as } \varepsilon \searrow 0,$$

for a universal constant $\alpha_i > 0$. More precisely, with the notations of Definition A.6, the latter is given by

$$\alpha_i = \mathfrak{T}_{\Pi_{\Sigma_i \times \{1\}}}(\Sigma_i \times \{1\}) = \mathfrak{T}_{\Pi_{\mathbf{I} \times \{0\}}}(\mathbf{I} \times \{0\}) > 0.$$

Observe in particular that α_i does not depend i , thanks to the fact that the quantity \mathfrak{T} is invariant by rigid movements. This is enough to conclude. \square

The relevance of \mathbf{T}_n^ε in our problem is encoded by the following estimate.

Proposition 7.4 (Eigenvalue estimate). *With the notation above, for every $n \in \mathbb{N} \setminus \{0\}$ we have that*

$$(7.3) \quad \lambda_1(\mathbf{H}_n^\varepsilon) \leq \lambda_1(\mathbf{H}) - \mathbf{T}_n^\varepsilon + o(\mathbf{T}_n^\varepsilon), \quad \text{as } \varepsilon \searrow 0,$$

Proof. We first observe that $\lambda_1(\mathbf{H}_n^\varepsilon) > 0$, since \mathbf{H}_n^ε is bounded in the x_2 direction. By appealing again to (7.1), we can apply Proposition A.2 and infer existence of $U_{\Gamma_n^\varepsilon} \in W_0^{1,2}(\mathbf{H}_n^\varepsilon)$ which attains the maximum in the definition of \mathbf{T}_n^ε .

We use the trial function $u_1 - U_{\Gamma_n^\varepsilon}$ in the variational problem which defines $\lambda_1(\mathbf{H}_n^\varepsilon)$. This gives

$$(7.4) \quad \lambda_1(\mathbf{H}_n^\varepsilon) \leq \frac{\int_{\mathbf{H}_n^\varepsilon} |\nabla(u_1 - U_{\Gamma_n^\varepsilon})|^2 dx}{\int_{\mathbf{H}_n^\varepsilon} (u_1 - U_{\Gamma_n^\varepsilon})^2 dx} = \frac{\lambda_1(\mathbf{H}) + \int_{\mathbf{H}_n^\varepsilon} |\nabla U_{\Gamma_n^\varepsilon}|^2 dx - 2 \int_{\mathbf{H}_n^\varepsilon} \langle \nabla u_1, \nabla U_{\Gamma_n^\varepsilon} \rangle dx}{1 + \int_{\mathbf{H}_n^\varepsilon} U_{\Gamma_n^\varepsilon}^2 dx - 2 \int_{\mathbf{H}_n^\varepsilon} u_1 U_{\Gamma_n^\varepsilon} dx} \\ \leq \frac{\lambda_1(\mathbf{H}) + \int_{\mathbf{H}_n^\varepsilon} |\nabla U_{\Gamma_n^\varepsilon}|^2 dx - 2 \int_{\mathbf{H}_n^\varepsilon} \langle \nabla u_1, \nabla U_{\Gamma_n^\varepsilon} \rangle dx}{1 - 2 \int_{\mathbf{H}_n^\varepsilon} u_1 U_{\Gamma_n^\varepsilon} dx}.$$

By using u_1 as a test function for the equation of $U_{\Gamma_n^\varepsilon}$ (see Proposition A.2) and observing that u_1 has a null trace on Γ_n^ε , we get

$$(7.5) \quad 0 = \int_{\mathbf{H}_n^\varepsilon} \langle \nabla u_1, \nabla U_{\Gamma_n^\varepsilon} \rangle dx = \int_{\mathbf{H}} \langle \nabla u_1, \nabla U_{\Gamma_n^\varepsilon} \rangle dx.$$

On the other hand, by using $U_{\Gamma_n^\varepsilon}$ as a test function for the equation of u_1 , one obtains from Lemma 7.1 that

$$(7.6) \quad \int_{\mathbf{H}_n^\varepsilon} u_1 U_{\Gamma_n^\varepsilon} dx = \int_{\mathbf{H}} u_1 U_{\Gamma_n^\varepsilon} dx \\ = \frac{1}{\lambda_1(\mathbf{H})} \int_{\mathbf{H}} \langle \nabla u_1, \nabla U_{\Gamma_n^\varepsilon} \rangle dx - \frac{1}{\lambda_1(\mathbf{H})} \int_{\Gamma_n^\varepsilon} U_{\Gamma_n^\varepsilon} \frac{\partial u_1}{\partial x_2} d\mathcal{H}^1 = -\frac{1}{\lambda_1(\mathbf{H})} \mathbf{T}_n^\varepsilon.$$

In the last identity, we used (A.1). Hence, from (7.4), (7.5) and (7.6) we get that

$$\lambda_1(\mathbf{H}_n^\varepsilon) \leq (\lambda_1(\mathbf{H}) + \mathbf{T}_n^\varepsilon) \left(1 + \frac{2}{\lambda_1(\mathbf{H})} \mathbf{T}_n^\varepsilon \right)^{-1}.$$

For every fixed $n \in \mathbb{N} \setminus \{0\}$, the quantity \mathbf{T}_n^ε is infinitesimal, as ε goes to 0 (thanks to Lemma 7.2). Thus we obtain that

$$\lambda_1(\mathbf{H}_n^\varepsilon) \leq (\lambda_1(\mathbf{H}) + \mathbf{T}_n^\varepsilon) \left(1 - \frac{2}{\lambda_1(\mathbf{H})} \mathbf{T}_n^\varepsilon + o(\mathbf{T}_n^\varepsilon) \right), \quad \text{as } \varepsilon \searrow 0.$$

This proves the claimed asymptotical estimate (7.3). \square

The main outcome of the previous discussion is contained in the following result. This simply follows by combining Proposition 7.4, Lemma 7.2 and Lemma 7.3.

Corollary 7.5. *For any fixed $n \in \mathbb{N} \setminus \{0\}$, there holds*

$$\lambda_1(\mathbf{H}_n^\varepsilon) \leq \lambda_1(\mathbf{H}) - \varepsilon^2 \alpha \sum_{i=0}^n \left(\frac{\partial u_1}{\partial x_2} \left(2 + \frac{i}{n}, 1 \right) \right)^2 + o(\varepsilon^2), \quad \text{as } \varepsilon \searrow 0.$$

Finally, by recalling the notation (1.5), we can prove the following

Theorem 7.6. *With the notation above, there exist $n_0 \in \mathbb{N} \setminus \{0\}$ such that*

$$\rho(\mathbf{H}_{n_0}^\varepsilon) \leq (1 - \varepsilon^2) \rho(\mathbf{H}) + o(\varepsilon^2), \quad \text{as } \varepsilon \searrow 0.$$

In particular, \mathbf{H} does not provide the sharp Makai-Hayman constant.

Proof. We first decide the number n_0 of tubes to be attached. At this aim, we observe that the following constant

$$\mathfrak{m} := \min_{x_1 \in [1,4]} \left(\frac{\partial u_1}{\partial x_2}(x_1, 1) \right)^2,$$

is positive, thanks to the Hopf Boundary Lemma. Accordingly, we set

$$n_0 := \min \left\{ n \in \mathbb{N} \setminus \{0\} : n \geq \frac{\lambda_1(\mathbf{H}) (1 + 2r_{\mathbf{H}}^2)}{2\alpha \mathfrak{m} r_{\mathbf{H}}^2} \right\},$$

and observe that such a choice is universal. Thanks to this definition, we have

$$(7.7) \quad \alpha \sum_{i=0}^{n_0} \left(\frac{\partial u_1}{\partial x_2} \left(2 + \frac{i}{n_0}, 1 \right) \right)^2 \geq \frac{\lambda_1(\mathbf{H}) (1 + 2r_{\mathbf{H}}^2)}{2r_{\mathbf{H}}^2}.$$

In order to estimate the quantity $\rho(\mathbf{H}_{n_0}^\varepsilon)$, one can easily check that

$$r_{\mathbf{H}_{n_0}^\varepsilon} = \frac{1}{2} + \frac{\varepsilon^2}{2} = r_{\mathbf{H}} + \frac{\varepsilon^2}{4r_{\mathbf{H}}},$$

which implies that

$$r_{\mathbf{H}_{n_0}^\varepsilon}^2 = r_{\mathbf{H}}^2 + \frac{\varepsilon^2}{2} + o(\varepsilon^2), \quad \text{as } \varepsilon \searrow 0.$$

Observe that the variation on the inradius does not depend on n_0 . This is the crucial point. Thanks to this fact and to Corollary 7.5, we have that, as ε goes to 0,

$$\begin{aligned} \rho(\mathbf{H}_{n_0}^\varepsilon) &= r_{\mathbf{H}_{n_0}^\varepsilon}^2 \lambda_1(\mathbf{H}_{n_0}^\varepsilon) \leq \left(r_{\mathbf{H}}^2 + \frac{\varepsilon^2}{2} + o(\varepsilon^2) \right) \left(\lambda_1(\mathbf{H}) - \varepsilon^2 \alpha \sum_{i=0}^{n_0} \left(\frac{\partial u_1}{\partial x_2} \left(2 + \frac{i}{n_0}, 1 \right) \right)^2 + o(\varepsilon^2) \right) \\ &= \rho(\mathbf{H}) + \varepsilon^2 \left(\frac{\lambda_1(\mathbf{H})}{2} - r_{\mathbf{H}}^2 \alpha \sum_{i=0}^{n_0} \left(\frac{\partial u_1}{\partial x_2} \left(2 + \frac{i}{n_0}, 1 \right) \right)^2 \right) + o(\varepsilon^2). \end{aligned}$$

By recalling (7.7), we get the desired conclusion. \square

APPENDIX A. THE THIN TORSIONAL RIGIDITY

In this section, we briefly recall some facts about the *thin torsional rigidity* used in Section 7, essentially taken from [1]. The main difference with [1] consists in the fact that we want to allow many tubes to be attached, rather than just one.

We start by describing the class of sets which will be singularly perturbed by thin tubes.

Assumptions 2. For $N \geq 2$, let $\Omega \subseteq \mathbb{R}^N$ be an open set with the following property. For some $M \in \mathbb{N} \setminus \{0\}$, there exist:

$$\mathbf{p}_1, \dots, \mathbf{p}_M \in \partial\Omega, \quad \nu_1, \dots, \nu_M \in \mathbb{S}^{N-1}, \quad r_1, \dots, r_M > 0,$$

such that, for all $i = 1, \dots, M$, there holds:

- i. $\partial\Omega \cap B_{r_i}(\mathbf{p}_i) = \partial\Omega \cap \{x \in B_{r_i}(\mathbf{p}_i) : \langle x - \mathbf{p}_i, \nu_i \rangle = 0\}$;
- ii. $(\mathbb{R}^N \setminus \bar{\Omega}) \cap B_{r_i}(\mathbf{p}_i) = (\mathbb{R}^N \setminus \bar{\Omega}) \cap \{x \in B_{r_i}(\mathbf{p}_i) : \langle x - \mathbf{p}_i, \nu_i \rangle > 0\}$.

In particular, the boundary of Ω is flat around each point \mathbf{p}_i , with ν_i representing the unit outer normal vector of $\partial\Omega$ around this point.

Moreover, for every $i = 1, \dots, M$ we take a relatively open connected subset $\Sigma_i \subseteq \partial\Omega \cap B_{r_i}(\mathbf{p}_i)$, such that $\mathbf{p}_i \in \Sigma_i$ and $\Sigma_i \cap \Sigma_j = \emptyset$ for $i \neq j$. Then we define

$$\Sigma := \bigcup_{i=1}^M \Sigma_i.$$

For a pair (Ω, Σ) as above, we define

$$T_{\Sigma_i} := \{x + t\nu_i : x \in \Sigma_i, t \in [0, 1)\} \quad \text{and} \quad \Omega_\Sigma := \Omega \cup \bigcup_{i=1}^M T_{\Sigma_i},$$

and we also assume that

$$T_{\Sigma_i} \cap \Omega = \emptyset \quad \text{and} \quad T_{\Sigma_i} \cap T_{\Sigma_j} = \emptyset,$$

for all $i, j = 1, \dots, M$ such that $i \neq j$.

Definition A.1. Let $\Omega \subseteq \mathbb{R}^N$ and $\Sigma \subseteq \partial\Omega$ be a pair satisfying Assumptions 2. For any $f \in L^2(\Sigma)$, we define the *thin f -torsional rigidity* of Σ relative to Ω_Σ as follows

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; f) := \sup_{\varphi \in C_0^\infty(\Omega_\Sigma)} \left\{ 2 \int_\Sigma f \varphi \, d\mathcal{H}^{N-1} - \int_{\Omega_\Sigma} |\nabla \varphi|^2 \, dx \right\}.$$

Of course, the supremum is unchanged if settled on $W_0^{1,2}(\Omega_\Sigma)$. In the particular case $f \equiv 1$, we will simply write

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma) := \mathcal{T}_{\Omega_\Sigma}(\Sigma; 1),$$

and we call it the *thin torsional rigidity* of Σ relative to Ω_Σ .

Proposition A.2. Let $\Omega \subseteq \mathbb{R}^N$ and $\Sigma \subseteq \partial\Omega$ be a pair satisfying Assumptions 2. Let us suppose that $\lambda_1(\Omega_\Sigma) > 0$. For any $f \in L^2(\Sigma)$, we have

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; f) = \max_{\varphi \in W_0^{1,2}(\Omega_\Sigma)} \left\{ 2 \int_\Sigma f \varphi \, d\mathcal{H}^{N-1} - \int_{\Omega_\Sigma} |\nabla \varphi|^2 \, dx \right\},$$

and this is a positive quantity. Moreover, such a maximum is (uniquely) attained by a function $U_{\Sigma,f} \in W_0^{1,2}(\Omega_\Sigma)$, which weakly satisfies

$$\begin{cases} -\Delta U_{\Sigma,f} = 0, & \text{in } \Omega_\Sigma \setminus \Sigma, \\ \frac{\partial U_{\Sigma,f}}{\partial \nu_i^+} + \frac{\partial U_{\Sigma,f}}{\partial \nu_i^-} = f, & \text{on } \Sigma_i, \text{ for } i = 1, \dots, M, \end{cases}$$

where

$$\frac{\partial U_{\Sigma,f}}{\partial \nu_i^\pm}(x) := \lim_{h \rightarrow 0^-} \frac{U_{\Sigma,f}(x \pm h \nu_i) - U_{\Sigma,f}(x)}{h}, \quad \text{for } x \in \Sigma_i.$$

More precisely, there holds

$$\int_{\Omega_\Sigma} \langle \nabla U_{\Sigma,f}, \nabla \varphi \rangle dx = \int_{\Sigma} \varphi f d\mathcal{H}^{N-1}, \quad \text{for every } \varphi \in W_0^{1,2}(\Omega_\Sigma).$$

Finally, we have

$$(A.1) \quad \mathcal{T}_{\Omega_\Sigma}(\Sigma; f) = \int_{\Sigma} f U_{\Sigma,f} d\mathcal{H}^{N-1}.$$

Proof. The assumption $\lambda_1(\Omega_\Sigma) > 0$ entails that

$$\varphi \mapsto \left(\int_{\Omega_\Sigma} |\nabla \varphi|^2 dx \right)^{\frac{1}{2}},$$

is an equivalent norm on $W_0^{1,2}(\Omega_\Sigma)$. Moreover, we have the continuity of the trace operator

$$\text{Tr} : W_0^{1,2}(\Omega_\Sigma) \rightarrow L^2 \left(\bigcup_{i=1}^M (\partial\Omega \cap B_{r_i}(\mathbf{p}_i)) \right),$$

see for example [30, Theorem 18.40]. Then the proof of this result is standard, it is sufficient to use the Direct Method in the Calculus of Variations. \square

Lemma A.3 (Superadditivity). *Let $\Omega \subseteq \mathbb{R}^N$ and $\Sigma \subseteq \partial\Omega$ be a pair satisfying Assumptions 2. Then for every $f \in L^2(\Sigma)$ we have*

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; |f|) = \mathcal{T}_{\Omega_\Sigma}(\Sigma; -|f|),$$

and

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; |f|) \geq \sum_{i=1}^M \mathcal{T}_{\Omega \cup T_{\Sigma_i}}(\Sigma_i; |f|).$$

Proof. The first fact simply follows by observing that

$$2 \int_{\Sigma} |f| \varphi d\mathcal{H}^{N-1} - \int_{\Omega_\Sigma} |\nabla \varphi|^2 dx = 2 \int_{\Sigma} (-|f|) (-\varphi) d\mathcal{H}^{N-1} - \int_{\Omega_\Sigma} |\nabla(-\varphi)|^2 dx.$$

For every $i = 1, \dots, M$, let $U_i \in W_0^{1,2}(\Omega \cup T_{\Sigma_i})$ be a function admissible for the optimization problem defining $\mathcal{T}_{\Omega \cup T_{\Sigma_i}}(\Sigma_i)$. We extend each U_i to the whole Ω_Σ , by defining it to be identically 0 in T_{Σ_j} , with $j \neq i$. We then observe that the trial function

$$U = \max \{ |U_1|, \dots, |U_M| \},$$

belongs to $W_0^{1,2}(\Omega_\Sigma)$. Thus, we get

$$\begin{aligned} \mathcal{T}_{\Omega_\Sigma}(\Sigma; |f|) &\geq 2 \int_\Sigma |f| U d\mathcal{H}^{N-1} - \int_{\Omega_\Sigma} |\nabla U|^2 dx \\ &\geq 2 \sum_{i=1}^M \int_{\Sigma_i} |f| U d\mathcal{H}^{N-1} - \sum_{i=1}^M \int_{\Omega_\Sigma} |\nabla U_i|^2 dx, \end{aligned}$$

thanks to the fact that

$$\int_{\Omega_\Sigma} |\nabla U|^2 dx \leq \sum_{i=1}^M \int_{\Omega_\Sigma} |\nabla |U_i||^2 dx = \sum_{i=1}^M \int_{\Omega_\Sigma} |\nabla U_i|^2 dx.$$

Moreover, by construction we have $U = |U_i| \geq U_i$ on Σ_i , which permits to infer that

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; |f|) \geq \sum_{i=1}^M \left[2 \int_{\Sigma_i} |f| U_i d\mathcal{H}^{N-1} - \int_{\Omega_\Sigma} |\nabla U_i|^2 dx \right],$$

By arbitrariness of the functions U_i , this gives the desired result. \square

Lemma A.4. *Let $\Omega \subseteq \mathbb{R}^N$ and $\Sigma \subseteq \partial\Omega$ be a pair satisfying Assumptions 2. Then, if $f \in L^\infty(\Sigma)$, there holds*

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; f) \leq \gamma(\Omega_\Sigma) \|f\|_{L^\infty(\Sigma)}^2 \mathcal{H}^{N-1}(\Sigma),$$

where

$$\gamma(\Omega_\Sigma) := \sup_{\substack{\varphi \in W_0^{1,2}(\Omega_\Sigma) \\ \varphi \neq 0}} \frac{\int_\Sigma |\varphi|^2 d\mathcal{H}^{N-1}}{\int_{\Omega_\Sigma} |\nabla \varphi|^2 dx}.$$

Proof. It is not difficult to see that $\mathcal{T}_{\Omega_\Sigma}(\Sigma; f)$ can be equivalently defined as

$$\mathcal{T}_{\Omega_\Sigma}(\Sigma; f) = \sup_{\substack{\varphi \in W_0^{1,2}(\Omega_\Sigma) \\ \varphi \neq 0}} \frac{\left(\int_\Sigma \varphi f d\mathcal{H}^{N-1} \right)^2}{\int_{\Omega_\Sigma} |\nabla \varphi|^2 dx},$$

see [1, Lemma 2.1]. We conclude by applying the Cauchy-Schwarz inequality to the numerator and using that $f \in L^\infty(\Sigma)$. \square

For any open set $E \subseteq \mathbb{R}^N$, we define the space $\mathcal{D}_0^{1,2}(E)$ as the completion of $C_0^\infty(E)$ with respect to the norm

$$(A.2) \quad \|\varphi\|_{\mathcal{D}_0^{1,2}(E)} := \left(\int_E |\nabla \varphi|^2 dx \right)^{\frac{1}{2}}, \quad \text{for every } \varphi \in C_0^\infty(E).$$

Remark A.5. We recall that for $N \geq 3$, the space $\mathcal{D}_0^{1,2}(E)$ can be identified with the closure of $C_0^\infty(E)$ in the Banach space

$$X^{2^*,2}(E) := \left\{ \varphi \in L^{2^*}(E) : \nabla \varphi \in L^2(E) \right\}, \quad \text{with } 2^* = \frac{2N}{N-2},$$

endowed with the natural norm

$$\|\varphi\|_{X^{2^*,2}(E)} := \|\varphi\|_{L^{2^*}(E)} + \|\nabla \varphi\|_{L^2(E)}.$$

This is possible thanks to Sobolev's inequality, which makes the two norms $\|\cdot\|_{X^{2^*,2}(E)}$ and $\|\cdot\|_{\mathcal{D}_0^{1,2}(E)}$ equivalent on $C_0^\infty(E)$.

The case $N = 2$ is more delicate, since we can not appeal to Sobolev's inequality. If $E \subsetneq \mathbb{R}^2$ is *simply connected*, we have at our disposal the following Hardy inequality

$$\frac{1}{16} \int_E \left| \frac{\varphi}{d_E} \right|^2 dx \leq \int_E |\nabla \varphi|^2 dx, \quad \text{for every } \varphi \in C_0^\infty(E),$$

see [2, page 278] and also [29]. Here $d_E : E \rightarrow \mathbb{R}$ is the distance from the boundary ∂E . In light of this result, $\mathcal{D}_0^{1,2}(E)$ can be identified with the closure of $C_0^\infty(E)$ in the Hilbert space

$$\dot{W}^{1,2}(E; d_E) = \left\{ \varphi \in L_{\text{loc}}^2(E) : \frac{\varphi}{d_E} \in L^2(E), \nabla \varphi \in L^2(E) \right\},$$

endowed with the norm

$$\|\varphi\|_{\dot{W}^{1,2}(E; d_E)} = \left(\left\| \frac{\varphi}{d_E} \right\|_{L^2(E)}^2 + \|\nabla \varphi\|_{L^2(E)}^2 \right)^{\frac{1}{2}}.$$

Definition A.6. Let $\mathbf{p} \in \mathbb{R}^N$ and $\nu \in \mathbb{S}^{N-1}$ be fixed. For any $\Sigma \subseteq \{x \in \mathbb{R}^N : \langle x - \mathbf{p}, \nu \rangle = 0\}$ relatively open bounded set, we let

$$T_\Sigma^\infty := \{x + t\nu : x \in \Sigma, t \in (-\infty, 0]\} \quad \text{and} \quad \Pi_\Sigma := \{x \in \mathbb{R}^N : \langle x - \mathbf{p}, \nu \rangle > 0\} \cup T_\Sigma^\infty.$$

For any $f \in L^2(\Sigma)$, we denote

$$\mathfrak{I}_{\Pi_\Sigma}(\Sigma; f) := \sup_{\varphi \in C_0^\infty(\Pi_\Sigma)} \left\{ 2 \int_\Sigma f \varphi d\mathcal{H}^{N-1} - \int_{\Pi_\Sigma} |\nabla \varphi|^2 dx \right\}.$$

Moreover, we denote

$$\mathfrak{I}_{\Pi_\Sigma}(\Sigma) := \mathfrak{I}_{\Pi_\Sigma}(\Sigma; 1).$$

We observe in particular that this quantity is invariant by rigid movements.

Proposition A.7. *For any $\Sigma \subseteq \{x \in \mathbb{R}^N : \langle x - \mathbf{p}, \nu \rangle = 0\}$ relatively open bounded set and any $f \in L^2(\Sigma)$, we have*

$$\mathfrak{I}_{\Pi_\Sigma}(\Sigma; f) = \max_{\varphi \in \mathcal{D}_0^{1,2}(\Pi_\Sigma)} \left\{ 2 \int_\Sigma f \varphi d\mathcal{H}^{N-1} - \int_{\Pi_\Sigma} |\nabla \varphi|^2 dx \right\},$$

and this is a positive quantity. Moreover, such a maximum is (uniquely) attained by a function $\tilde{U}_{\Sigma, f} \in \mathcal{D}_0^{1,2}(\Pi_\Sigma)$, which weakly satisfies

$$\begin{cases} -\Delta \tilde{U}_{\Sigma, f} = 0, & \text{in } \Pi_\Sigma \setminus \Sigma, \\ \frac{\partial \tilde{U}_{\Sigma, f}}{\partial \nu^+} + \frac{\partial \tilde{U}_{\Sigma, f}}{\partial \nu^-} = f, & \text{on } \Sigma, \end{cases}$$

where

$$\frac{\partial \tilde{U}_{\Sigma, f}}{\partial \nu^\pm}(x) := \lim_{h \rightarrow 0^-} \frac{\tilde{U}_{\Sigma, f}(x \pm h\nu) - \tilde{U}_{\Sigma, f}(x)}{h}, \quad \text{for } x \in \Sigma.$$

More precisely, there holds

$$\int_{\Pi_\Sigma} \langle \nabla \tilde{U}_{\Sigma, f}, \nabla \varphi \rangle dx = \int_\Sigma \varphi f d\mathcal{H}^{N-1}, \quad \text{for every } \varphi \in \mathcal{D}_0^{1,2}(\Pi_\Sigma).$$

Proof. Without loss of generality, we can suppose that $\mathbf{p} = 0$ and $\nu = \mathbf{e}_N$. In this case, $\Sigma \subseteq \mathbb{R}^{N-1}$ and

$$T_\Sigma^\infty := \Sigma \times (-\infty, 0] \quad \text{and} \quad \Pi_\Sigma := \{x = (x', x_N) \in \mathbb{R}^N : x_N > 0\} \cup T_\Sigma^\infty.$$

We need at first to show the existence of a continuous trace operator

$$\text{Tr} : \mathcal{D}_0^{1,2}(\Pi_\Sigma) \rightarrow L^2(\Sigma).$$

We take $\varphi \in C_0^\infty(\Pi_\Sigma)$ and write

$$\varphi(x', 0) = \varphi(x', x_N) + \int_0^{x_N} \frac{\partial \varphi}{\partial x_N}(x', \tau) d\tau, \quad \text{for every } x' \in \Sigma \text{ and } x_N \in (0, 1).$$

In particular, by raising to the power 2 and using Jensen's inequality, we get

$$\begin{aligned} |\varphi(x', 0)|^2 &\leq 2|\varphi(x', x_N)|^2 + 2x_N \int_0^{x_N} \left| \frac{\partial \varphi}{\partial x_N}(x', \tau) \right|^2 d\tau \\ &\leq 2|\varphi(x', x_N)|^2 + 2x_N \int_0^1 \left| \frac{\partial \varphi}{\partial x_N}(x', \tau) \right|^2 d\tau. \end{aligned}$$

We now integrate this estimate on $\Sigma \times (0, 1)$, so to get

$$(A.3) \quad \int_\Sigma |\varphi(x', 0)|^2 dx' \leq 2 \int_{\Sigma \times (0,1)} |\varphi|^2 dx + 2 \int_{\Sigma \times (0,1)} \left| \frac{\partial \varphi}{\partial x_N} \right|^2 dx.$$

If $N \geq 3$, we can estimate

$$\int_{\Sigma \times (0,1)} |\varphi|^2 dx \leq |\Sigma \times (0, 1)|^{1-\frac{2}{2^*}} \left(\int_{\Sigma \times (0,1)} |\varphi|^{2^*} dx \right)^{\frac{2}{2^*}} \leq \frac{|\Sigma \times (0, 1)|^{1-\frac{2}{2^*}}}{T_N} \int_{\Pi_\Sigma} |\nabla \varphi|^2 dx,$$

by Sobolev's inequality. If $N = 2$, we observe that Π_Σ is a proper simply connected subset of \mathbb{R}^2 and use the Hardy inequality recalled in Remark A.5 above. This permits to infer that

$$\int_{\Sigma \times (0,1)} |\varphi|^2 dx = \int_{\Sigma \times (0,1)} \left| \frac{\varphi}{d_{\Pi_\Sigma}} \right|^2 d_{\Pi_\Sigma}^2 dx \leq C_\Sigma^2 \int_{\Sigma \times (0,1)} \left| \frac{\varphi}{d_{\Pi_\Sigma}} \right|^2 \leq 16 C_\Sigma^2 \int_{\Pi_\Sigma} |\nabla \varphi|^2 dx.$$

We used that d_{Π_Σ} is bounded by a constant C_Σ on $\Sigma \times (0, 1)$. Thus, both for $N \geq 3$ and $N = 2$, from (A.3) we get

$$\left(\int_\Sigma |\varphi(x', 0)|^2 dx' \right)^{\frac{1}{2}} \leq C \left(\int_{\Pi_\Sigma} |\nabla \varphi|^2 dx \right)^{\frac{1}{2}}, \quad \text{for every } \varphi \in C_0^\infty(\Pi_\Sigma),$$

for some $C = C(N, \Sigma) > 0$. This estimate shows in particular that for every Cauchy sequence $\{\varphi_n\}_{n \in \mathbb{N}} \subseteq C_0^\infty(\Pi_\Sigma)$ with respect to the norm (A.2), we have that $\{\varphi(\cdot, 0)\}_{n \in \mathbb{N}}$ is a Cauchy sequence in the Hilbert space $L^2(\Sigma)$, as well. This permits to define the trace operator in a standard way, which is thus continuous.

The existence of a maximizer for $\mathfrak{T}_{\Pi_\Sigma}(\Sigma; f)$ can now be proved by means of the Direct Method in the Calculus of Variations. We leave the details to the reader. \square

Finally, we recall [1, Theorem 2.8], which contains a blow-up analysis for the thin torsional rigidity of shrinking sets, here suitably stated in our framework. We point out that [1, Theorem 2.8] has been proved only in dimension $N \geq 3$, but the very same argument can be repeated in dimension 2. It is sufficient to use the characterization of $\mathcal{D}_0^{1,2}$ on simply connected proper subsets of \mathbb{R}^2 , contained in Remark A.5.

In order to state it clearly for our needs, we need some further notation: let $\Omega \subseteq \mathbb{R}^N$ and $\Sigma \subseteq \partial\Omega$ be a pair satisfying Assumptions 2. For every $i = 1, \dots, M$ and for every $0 < \varepsilon < 1$, we set

$$\Sigma_i^\varepsilon := \varepsilon(\Sigma_i - \mathbf{p}_i) + \mathbf{p}_i.$$

Recall that $\mathbf{p}_i \in \Sigma_i$, thus Σ_i^ε is shrinking to this point, as ε goes to 0. Then we set

$$T_i^\varepsilon := T_{\Sigma_i^\varepsilon} = \{x + t\nu_i : x \in \Sigma_i^\varepsilon, t \in [0, 1)\} \quad \text{and} \quad \Omega_{\Sigma_i^\varepsilon} := \Omega \cup T_i^\varepsilon.$$

Theorem A.8. *With the notations above, let us suppose that*

$$\Omega \subseteq \{x \in \mathbb{R}^N : \langle x - \mathbf{p}_i, \nu_i \rangle \leq 0\}.$$

Let $f_i \in C^0(\partial\Omega \cap B_{r_i}(\mathbf{p}_i))$ be such that $f_i(\mathbf{p}_i) \neq 0$. Then, we have

$$\mathcal{T}_{\Omega \cup T_i^\varepsilon}(\Sigma_i^\varepsilon; f_i) = \varepsilon^2 \left(f_i(\mathbf{p}_i) \right)^2 \mathfrak{T}_{\Pi_{\Sigma_i}}(\Sigma_i) + o(\varepsilon^2), \quad \text{as } \varepsilon \searrow 0.$$

REFERENCES

- [1] L. Abatangelo, R. Ognibene, Sharp behavior of Dirichlet–Laplacian eigenvalues for a class of singularly perturbed problems, preprint (2023), available at <https://arxiv.org/abs/2301.11729> [8](#), [9](#), [36](#), [38](#), [40](#)
- [2] A. Ancona, On strong barriers and inequality of Hardy for domains in \mathbb{R}^n , *J. London Math. Soc.*, **34** (1986), 274–290. [39](#)
- [3] M. S. Ashbaugh, P. Exner, Lower bounds to bound state energies in bent tubes, *Phys. Lett. A*, **150** (1990), 183–186. [8](#)
- [4] Y. Avishai, D. Bessis, B. G. Giraud, G. Mantica, Quantum bound states in open geometries, *Phys. Rev. B*, **44** (1991), 8028–8034. [16](#), [18](#)
- [5] R. Bañuelos, T. Carroll, Brownian motion and the fundamental frequency of a drum, *Duke Math. J.*, **75** (1994), 575–602. [7](#)
- [6] F. Bianchi, L. Brasco, An optimal lower bound in fractional spectral geometry for planar sets with topological constraints, to appear on *J. London Math. Soc.* (2023), [doi:10.1112/jlms.12814](https://doi.org/10.1112/jlms.12814) [6](#)
- [7] F. Bianchi, L. Brasco, The fractional Makai–Hayman inequality, *Ann. Mat. Pura Appl.* (4), **201** (2022), 2471–2504. [6](#)
- [8] M. Sh. Birman, M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert space*. Translated from the 1980 Russian original by S. Khrushchev and V. Peller. Mathematics and its Applications (Soviet Series). D. Reidel Publishing Co., Dordrecht, 1987. [6](#), [29](#)
- [9] P. R. Brown, Constructing mappings onto radial slit domains, *Rocky Mountain J. Math.*, **37** (2007), 1791–1812. [7](#)
- [10] W. Bulla, F. Gesztesy, W. Renger, B. Simon, Weakly coupled bound states in quantum waveguides, *Proc. Amer. Math. Soc.*, **125** (1997), 1487–1495. [4](#)
- [11] G. Buttazzo, Spectral optimization problems, *Rev. Mat. Complut.*, **24** (2011), 277–322. [7](#)
- [12] G. Chiti, An isoperimetric inequality for the eigenfunctions of linear second order elliptic operators, *Boll. Un. Mat. Ital. A* (6), **1** (1982), 145–151. [26](#)
- [13] R. Courant, D. Hilbert, *Methods of mathematical physics. Vol. I*. Interscience Publishers, Inc., New York, N.Y., 1953. [2](#), [19](#)
- [14] C. B. Croke, The first eigenvalue of the Laplacian for plane domains, *Proc. Amer. Math. Soc.*, **81** (1981), 304–305. [6](#)
- [15] M. Dauge, N. Raymond, Plane waveguides with corners in the small angle limit, *J. Math. Phys.*, **53** (2012), 123529, 34 pp. [18](#)
- [16] E. B. Davies, *Heat kernels and spectral theory*. Cambridge Tracts in Math., **92**. Cambridge University Press, Cambridge, 1989. [26](#)
- [17] A. L. Delitsyn, B. T. Nguyen, D. S. Grebenkov, Exponential decay of Laplacian eigenfunctions in domains with branches of variable cross-sectional profiles, *Eur. Phys. J. B*, **85** (2012), 371. [21](#)
- [18] P. Exner, P. Šeba, Bound states in curved quantum waveguides, *J. Math. Phys.*, **30** (1989), 2574–2580. [8](#)
- [19] P. Exner, H. Kovařík, *Quantum waveguides*. Theoretical and Mathematical Physics. Springer, Cham, 2015. [8](#)

- [20] V. Felli, R. Ognibene, Sharp convergence rate of eigenvalues in a domain with a shrinking tube, *J. Differential Equations*, **269** (2020), 713–763. [8](#)
- [21] D. Gilbarg, N. S. Trudinger, *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics Math. Springer-Verlag, Berlin, 2001. [26](#), [28](#), [30](#)
- [22] J. Goldstone, R. L. Jaffe, Bound states in twisting tubes, *Phys. Rev. B*, **45** (1992), 14100–14107. [8](#)
- [23] S. E. Graversen, M. Rao, Brownian Motion and Eigenvalues for the Dirichlet Laplacian, *Math. Z.*, **203** (1990), 699–708. [6](#)
- [24] W. K. Hayman, Some bounds for principal frequency, *Applicable Anal.*, **7** (1977/78), 247–254. [6](#)
- [25] A. Henrot, *Extremum problems for eigenvalues of elliptic operators*. Frontiers in Mathematics. Birkhauser Verlag, Basel, 2006. [7](#), [16](#)
- [26] J. Hersch, Review of the article “A lower estimation of the principal frequencies of simply connected membranes” by E. Makai, *Mathematical Review MR0185263*. [7](#)
- [27] J. Hersch, Sur la fréquence fondamentale d’une membrane vibrante: évaluations par défaut et principe de maximum, *Z. Angew. Math. Phys.*, **11** (1960), 387–413. [7](#)
- [28] D. Krejčířík, J. Kríž, On the spectrum of curved planar waveguides, *Publ. Res. Inst. Math. Sci.*, **41** (2005), 757–791. [8](#)
- [29] A. Laptev, A. V. Sobolev, Hardy inequalities for simply connected planar domains. *Spectral theory of differential operators*, 133–140, Amer. Math. Soc. Transl. Ser. 2, 225, Adv. Math. Sci., 62, Amer. Math. Soc., Providence, RI, 2008. [39](#)
- [30] G. Leoni, *A first course in Sobolev spaces. Second edition*. Graduate Studies in Mathematics, **181**. American Mathematical Society, Providence, RI, 2017. [37](#)
- [31] H. A. Levine, An estimate for the best constant in a Sobolev inequality involving three integral norms, *Ann. Mat. Pura Appl.* (4), **124** (1980), 181–197. [9](#), [22](#)
- [32] M. Levitin, M. Marletta, A simple method of calculating eigenvalues and resonances in domains with infinite regular ends, *Proc. Roy. Soc. Edinburgh Sect. A*, **138** (2008), 1043–1065. [4](#)
- [33] E. Makai, A lower estimation of the principal frequencies of simply connected membranes, *Acta Math. Acad. Sci. Hungar.*, **16** (1965), 319–323. [6](#), [7](#)
- [34] V. Maz’ya, *Sobolev spaces with applications to elliptic partial differential equations*. Second, revised and augmented edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **342**. Springer, Heidelberg, 2011. [2](#)
- [35] S. A. Nazarov, Discrete spectrum of cross-shaped quantum waveguides, *J. Math. Sci. (N.Y.)*, **196** (2014), 346–376. [16](#)
- [36] R. Osserman, A note on Hayman’s theorem on the bass note of a drum, *Comment. Math. Helvetici*, **52** (1977), 545–555. [6](#)
- [37] G. Poliquin, Principal frequency of the p -Laplacian and the inradius of Euclidean domains, *J. Topol. Anal.*, **7** (2015), 505–511. [6](#)
- [38] R. L. Schult, D. G. Ravenhall, H. W. Wyld, Quantum bound states in a classically unbounded system of crossed wires, *Phys. Review B*, **39** (1989), 5476–5479. [16](#)
- [39] B. Siudeja, Sharp bounds for eigenvalues of triangles, *Michigan Math. J.*, **55** (2007), 243–254. [17](#)
- [40] D. Smets, A concentration-compactness lemma with applications to singular eigenvalue problems, *J. Funct. Anal.*, **167** (1999), 463–480. [5](#), [9](#)
- [41] M. Struwe, *Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems. Fourth edition*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, **34**. Springer-Verlag, Berlin, 2008. [3](#)
- [42] G. Talenti, Best constant in Sobolev inequality, *Ann. Mat. Pura Appl.* (4), **110** (1976), 353–372. [22](#)
- [43] M. E. Taylor, Estimate on the fundamental frequency of a drum, *Duke Math. J.* **46** (1979), 447–453. [6](#)
- [44] G. Talenti, A. Colesanti, P. Salani, *Un’introduzione al Calcolo delle Variazioni: teoria ed esercizi*. Unione Matematica Italiana, 2016. [8](#)
- [45] G. Teschl, *Mathematical methods in quantum mechanics. With applications to Schrödinger operators*. Second edition. Graduate Studies in Mathematics, **157**. American Mathematical Society, Providence, RI, 2014. [6](#), [29](#)

(F. Bianchi) DIPARTIMENTO DI SCIENZE MATEMATICHE, FISICHE E INFORMATICHE
UNIVERSITÀ DI PARMA
PARCO AREA DELLE SCIENZE 53/A, CAMPUS, 43124 PARMA, ITALY
E-mail address: francesca.bianchi@unipr.it

(L. Brasco) DIPARTIMENTO DI MATEMATICA E INFORMATICA
UNIVERSITÀ DEGLI STUDI DI FERRARA
VIA MACHIAVELLI 35, 44121 FERRARA, ITALY
E-mail address: lorenzo.brasco@unife.it

(R. Ognibene) DIPARTIMENTO DI MATEMATICA
UNIVERSITÀ DI PISA
LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY
E-mail address: roberto.ognibene@dm.unipi.it