

CONVERGENCE OF CRITICAL POINTS FOR A PHASE-FIELD APPROXIMATION OF 1D COHESIVE FRACTURE ENERGIES

MARCO BONACINI AND FLAVIANA IURLANO

ABSTRACT. Variational models for cohesive fracture are based on the idea that the fracture energy is released gradually as the crack opening grows. Recently, [21] proposed a variational approximation via Γ -convergence of a class of cohesive fracture energies by phase-field energies of Ambrosio-Tortorelli type, which may be also used as regularization for numerical simulations. In this paper we address the question of the asymptotic behaviour of critical points of the phase-field energies in the one-dimensional setting: we show that they converge to a selected class of critical points of the limit functional. Conversely, each critical point in this class can be approximated by a family of critical points of the phase-field functionals.

1. INTRODUCTION

Fracture models describe the evolution of surface cracks in elastic materials subjected to external loads or boundary conditions. The literature distinguishes between brittle models and cohesive models (also known as Griffith and Barenblatt models respectively). The former treat fracture as an instantaneous phenomenon: the body deforms elastically until a crack surface appears; the crack energy is instantaneously released and there is no transmission of force across the crack surface. The latter treat fracture as a gradual phenomenon: the bonds between the lips progressively weaken; the crack energy is released with the growth of the crack opening and the force transmitted across the crack surface gradually reduces to zero. Thanks to these features, cohesive models are better suited than brittle models for describing crack nucleation. We refer the interested reader to the book [16] and references therein for a comprehensive comparison between brittle and cohesive, and we work from now on in the cohesive setting.

The variational study of cohesive fracture started in the late 1990s and has been earning interest ever since [4, 8, 12, 13, 17–20, 23, 24, 26–29, 36, 41, 42, 44, 46–48]. The appropriate variational setting to model a cohesive fracture process was shown to be the space of functions of bounded variation or bounded deformation, allowing to describe the crack as the jump set of a discontinuous displacement, and the total energy as a competition between bulk and surface contributions [31]. The presence of free discontinuities, making the numerical treatment highly complex, lead to the development of regularized phase-field theories [2, 3, 12, 21, 22, 25], in the spirit of the classical Allen-Cahn (or Modica-Mortola) approximation for phase transitions [40], and the Ambrosio-Tortorelli approximation of the Mumford-Shah functional for image segmentation or brittle fracture [6, 7, 33]. The general approach of these works is to construct sequences of purely bulk energies, whose variables are forced to engage transitions in thin concentration sets, and to show the convergence of corresponding global minimizers to a global minimizer of the given energy as the thickness of the concentration sets vanishes.

Date: September 29, 2023.

Although this kind of results usually marks decisive enhancements in the mathematical comprehension of the corresponding phenomena, its energy-based formulation and its global minimization focus may be not completely satisfactory from the mechanical point of view. Indeed, fracture evolution might realistically occur along critical states rather than following global minimizers. In addition, numerical schemes based on alternate minimization for the regularized energies typically converge to critical points of the limit energy; hence, the sole convergence of global minimizers does not provide a complete theoretical justification of the adoption of the phase-field models for numerical simulations, see for example [1, 15, 16, 32, 35, 51, 52].

This motivated the investigation of better converging properties of the proposed regular functionals. On the one side, it lead to the study of the convergence of the corresponding gradient flows, see [5, 9] for brittle fracture. On the other side, it lead to the study of the convergence of critical points, see in particular [34, 38, 39, 43, 45, 49, 50] for the Allen-Cahn functionals, and [10, 11, 30, 37] for the Ambrosio-Tortorelli functionals.

In this paper we address the latter question in the context of one-dimensional cohesive fracture: we study the asymptotic behaviour of the critical points of the regularized functionals proposed in [21]. Thanks to its nature, the cohesive case allows for a deeper and more complete analysis with respect to the brittle case. Our approach heavily relies on one-dimensional arguments and the analysis is at the moment limited to this setting; its possible extension to the higher-dimensional case, in the spirit of the recent work [11] in the context of brittle fracture, poses significant challenges.

The rest of this Introduction is organized as follows: in Section 1.1 we provide notation and properties of the sharp cohesive model and of its critical points; in Section 1.2 we introduce our regularized models, which are slight modifications of those proposed in [12, 21]. A few additional technical assumptions will be needed. Our main results will be stated in Section 1.3.

1.1. The cohesive fracture energy and its critical points. We first introduce a one-dimensional cohesive fracture energy for a bar of total length $L > 0$ (at rest), and total elongation $a > 0$.

The deformation of the bar is described by a function of bounded variation $u \in \text{BV}(0, L)$, whose distributional derivative is a bounded Radon measure on $(0, L)$ that can be written as

$$Du = u' dx + D^c u + \sum_{x \in J_u} [u](x) \delta_x,$$

where $u' \in L^1(0, L)$ denotes the density of the absolutely continuous part (with respect to the Lebesgue measure), $D^c u$ is the Cantor part, $[u](x) := u(x^+) - u(x^-)$, where $u(x^+)$ and $u(x^-)$ are the approximate limits from the right and from the left of u at x respectively, and $J_u := \{x \in (0, L) : [u](x) \neq 0\}$ is the set of essential discontinuities. Since we want to include in the energy the boundary conditions, we set $u(0^-) = 0$, $u(L^+) = a$, we extend the definition of $[u](x) := u(x^+) - u(x^-)$ also at the endpoints $x = 0$, $x = L$, and we let $J_u^a := \{x \in [0, L] : [u](x) \neq 0\}$.

The *cohesive energy* of the bar is defined as

$$\Phi(u) := \int_0^L \phi(u') dx + \sum_{x \in J_u^a} g(|[u](x)|) + \sigma_c |D^c u|(0, L) \quad \text{for } u \in \text{BV}(0, L). \quad (1.1)$$

Here $\sigma_c \in (0, +\infty)$ and the *elastic energy density* $\phi : \mathbb{R} \rightarrow [0, +\infty)$ is given by

$$\phi(\xi) := \begin{cases} \xi^2 & \text{if } |\xi| \leq \frac{\sigma_c}{2}, \\ \sigma_c |\xi| - \frac{\sigma_c^2}{4} & \text{if } |\xi| > \frac{\sigma_c}{2}. \end{cases} \quad (1.2)$$

For the *cohesive energy density* we assume that $g : [0, +\infty) \rightarrow [0, +\infty)$ is a nondecreasing function of class C^1 with $g(0) = 0$, $g'(0) = \sigma_c$. We further assume that $\{g' > 0\} = [0, s_{\text{frac}})$ for some $s_{\text{frac}} \in (0, +\infty]$, and that g' is strictly decreasing in $[0, s_{\text{frac}})$.

It is convenient to include in the energy also the non-interpenetration constraint that the singular part $D^s u := D^c u + \sum_{x \in J_u^a} [u](x) \delta_x$ of Du is a nonnegative measure: we therefore define $\tilde{\Phi}(u) := \Phi(u)$ if $u \in \text{BV}(0, L)$ with $D^s u \geq 0$ (so that in particular $u(0^+) \geq 0$, $u(L^-) \leq a$), and $\tilde{\Phi}(u) = +\infty$ otherwise.

Critical points of the functional $\tilde{\Phi}$ are functions u such that $\tilde{\Phi}(u) < +\infty$ and

$$\liminf_{t \rightarrow 0^+} \frac{\tilde{\Phi}(u + tv) - \tilde{\Phi}(u)}{t} \geq 0 \quad \text{for all } v \in \text{BV}(0, L).$$

These are studied in details in [17] (see also [28]). We stress that nonnegativity of the unilateral lower limit as $t \rightarrow 0^+$ is required in place of the usual vanishing of the first variation. This is a standard way to give a meaningful notion of critical point in presence of a noninterpenetration constraint. Mechanically, such condition provides a critical stress to nucleation, in the sense that nucleation of a crack point is only possible when the stress reaches the critical value.

By [17, Theorem 6.3] one has that a function $u \in \text{BV}(0, L)$ with $D^s u \geq 0$ is a critical point of $\tilde{\Phi}$ if and only if there exists a constant $\sigma \in \mathbb{R}$ such that

- (i) $\sigma \leq \sigma_c$,
- (ii) $\phi'(u') = \sigma$ a.e. in $(0, L)$,
- (iii) $g'([u]) = \sigma$ on J_u^a ,
- (iv) $D^c u = 0$ if $\sigma < \sigma_c$.

As observed in [17, Remark 6.6], in the model described by the energy $\tilde{\Phi}$ the quantity $\phi'(u')$ represents the stress in the elastic part of the bar due to the deformation gradient u' ; $g'([u](x))$ represents the stress transmitted through the points of J_u^a of the reference configuration (where there is concentration of the strain); $D^c u$ can be seen as a singular, not concentrated strain which transmits a stress σ_c and can be present only if $\sigma = \sigma_c$, whereas if $\sigma < \sigma_c$ a critical point is necessarily in $\text{SBV}(0, L)$. The constant σ_c is the maximum possible stress for an equilibrium configuration.

In view of the previous conditions, we can give an explicit description of all critical points belonging to $\text{SBV}(0, L)$ corresponding to a prescribed elongation $a > 0$ (see Figure 1):

- (a) (*elastic states*) $u(x) = \frac{a}{L}x$ in $(0, L)$;
- (b) (*pre-fractured states*) $\sigma \in (0, \sigma_c)$, $\#J_u^a = k$, $1 \leq k < +\infty$, $u' = \frac{\sigma}{2}$ a.e. in $(0, L)$, $[u](x) = s_0$ for all $x \in J_u^a$ where $s_0 \in (0, s_{\text{frac}})$ obeys $g'(s_0) = \sigma$, and $\frac{\sigma}{2}L + ks_0 = a$;
- (c) (*fractured states*) $\sigma = 0$, $\#J_u^a = k$, $1 \leq k < +\infty$, $u' = 0$ a.e. in $(0, L)$, and $[u](x) \geq s_{\text{frac}}$ for all $x \in J_u^a$, with $\sum_{x \in J_u^a} [u](x) = a$.

Notice that in the first case $\sigma = \frac{2a}{L} \wedge \sigma_c \in [0, \sigma_c]$ and all the possible slopes $u' = \frac{a}{L}$ are allowed; if $\frac{2a}{L} \geq \sigma_c$ then $\sigma = \sigma_c$. In the second case u is piecewise affine with constant slope $\frac{\sigma}{2}$ and with a finite number of jumps of the same amplitude. In the third case $\sigma = 0$, u is piecewise constant with a finite number of jumps with amplitude larger than s_{frac} , and this case can only occur if $s_{\text{frac}} < +\infty$ and $a \geq s_{\text{frac}}$.




FIGURE 1. Critical points in $SBV(0, L)$ of the cohesive energy (1.1). Left: elastic state $u(x) = \frac{a}{L}x$. Center: pre-fractured state with $\sigma \in (0, \sigma_c)$. Right: fractured state with $\sigma = 0$.

Remark 1.1. By the results in [14] the functional $\tilde{\Phi}$ is the lower semicontinuous envelope with respect to the L^1 -convergence of the energy

$$u \mapsto \int_0^L |u'|^2 dx + \sum_{x \in J_u^a} g(|[u](x)|) \quad \text{for } u \in SBV(0, L) \text{ with } [u] \geq 0. \quad (1.3)$$

As it is observed in [17, Remark 6.5], every critical point of this functional is also a critical point of $\tilde{\Phi}$, and every critical point of $\tilde{\Phi}$ with $\sigma < \sigma_c$ is a critical point of (1.3). For $\sigma = \sigma_c$, there are critical points of $\tilde{\Phi}$ which are not critical for (1.3): in particular, elastic states $u(x) = \frac{a}{L}x$ with $\frac{2a}{L} > \sigma_c$, and critical points with $D^c u \neq 0$. Such configurations, however, are not *local minimizers* of $\tilde{\Phi}$: in particular, $\tilde{\Phi}$ and (1.3) have the same (local) minimizers, see [17, Theorem 7.2].

Remark 1.2. It is instructive to consider a quasistatic evolution for the cohesive model (1.3), corresponding to a time-dependent elongation $a \geq 0$, monotone increasing in time: at each time we assume that the deformation u_a of the bar is a critical state satisfying the boundary conditions $u_a(0) = 0$, $u_a(L) = a$ (in the weak sense, as specified above). Initially, for small values of a , the response of the bar is purely elastic and the evolution follows the elastic critical points $u_a(x) = \frac{a}{L}x$, until the critical stress is reached ($\frac{a}{L} = \frac{\sigma_c}{2}$). At this point, that is for $a = \frac{1}{2}\sigma_c L$, it is expected that the state of the bar switches to a pre-fractured state, a pre-fracture point appears, and the amplitude of the crack continuously increases from 0 to s_{frac} as a increases from $\frac{1}{2}\sigma_c L$ to s_{frac} . In this case the displacement u_a has constant slope $\frac{\sigma}{2}$, with $\sigma \in (0, \sigma_c)$, and jump amplitude $s_0 \in (0, s_{\text{frac}})$, which should satisfy the compatibility conditions

$$g'(s_0) = \sigma, \quad \frac{\sigma}{2}L + s_0 = a. \quad (1.4)$$

The limit case $a = s_{\text{frac}}$ corresponds to the complete fracture state, characterized by jump amplitude exactly equal to the boundary datum $s_0 = s_{\text{frac}} = a$ and vanishing stress $\sigma = 0$.

Concerning the compatibility conditions above, we remark that the existence of a solution $(\sigma, s_0) \in (0, \sigma_c) \times (0, s_{\text{frac}})$ to (1.4) for values $a > \frac{1}{2}\sigma_c L$, with the property that $s_0 \rightarrow 0^+$ and $\sigma \rightarrow \sigma_c^-$ as $a \rightarrow (\frac{1}{2}\sigma_c L)^+$, is not a priori guaranteed. In case of nonexistence, then, once the critical stress is reached, the evolution creates instantaneously a jump of strictly positive amplitude, possibly even brittle without cohesive effects (see [19, Section 9] for an explicit example in two dimensions).

The behaviour of the function g at the origin determines whether for elongations $a \rightarrow (\frac{1}{2}\sigma_c L)^+$ there are solutions of (1.4) such that $s_0 \rightarrow 0^+$. Assuming that g satisfies the expansion

$$g(s) = \sigma_c s - \tilde{\ell} s^p + o(s^p) \quad \text{as } s \rightarrow 0$$

for some $\tilde{\ell} > 0$ and $p > 1$, then it can be checked by elementary arguments that existence of solutions as above is guaranteed if $p > 2$, whereas it fails if $p < 2$. The case $p = 2$ is critical, the existence of solutions depends on the length L of the bar and fails for sufficiently large L ; this is an instance of the well-known size effects in fracture. A suitable choice of our regularized models will produce in the limit a density g with the previous asymptotic expansion, see Proposition 3.9. We refer to [27] for a more detailed discussion of the content of this remark.

1.2. Phase-field approximation. Following [21], we now introduce a family of functionals $\mathcal{F}_\varepsilon : L^1(0, L) \times L^1(0, L) \rightarrow [0, +\infty]$, depending on a real parameter $\varepsilon > 0$, which approximate the cohesive energy density (1.1) in the sense of Γ -convergence. We let f satisfy the following conditions, that will be assumed throughout the paper unless explicitly stated:

$$(f1) \quad f \in C^2([0, 1]; [0, +\infty)), \quad f^{-1}(0) = \{0\};$$

$$(f2) \quad \lim_{s \rightarrow 1^-} (1-s)f(s) = \sigma_c, \quad \text{with } \sigma_c \in (0, +\infty);$$

$$(f3) \quad \frac{d}{ds} [(1-s)f(s)] > 0 \quad \text{for all } s \in (0, 1);$$

$$(f4) \quad \frac{d}{ds} \left[\frac{(1-s)f'(s)}{f(s)} \right] < 0 \quad \text{for all } s \in (0, 1);$$

$$(f5) \quad \lim_{s \rightarrow 1^-} \frac{1}{(1-s)^3} \frac{d}{ds} [(1-s)f(s)] = +\infty;$$

$$(f6) \quad \text{the map } s \mapsto \sqrt{s}f(1-\sqrt{s}) \text{ is convex.}$$

The previous conditions look very technical and a few comments are required: firstly, they are satisfied by a large class of functions, as Remark 1.3 below shows. Secondly, the Γ -convergence result in [21] holds under weaker assumptions, but the analysis of the model was further improved in [12] where conditions (f1), (f2), (f3), (f6) were assumed in order to obtain more detailed properties of the limit functional, that we will recall in Section 3. Here, we also include the conditions (f4)–(f5) whose role is crucial for the analysis in Section 2. We believe that the optimal set of assumptions is (f1)–(f5), but dropping the convexity condition (f6) would require a much finer analysis that goes beyond the scopes of this paper (see Remark 3.5).

Remark 1.3. Prototype functions with the properties above are the maps

$$f_q(s) := \frac{\sigma_c [1 - (1-s)^q]}{1-s} \quad \text{and} \quad f^p(s) := \frac{(\sigma_c + p(1-s))s^2}{1-s} \quad (1.5)$$

for $\sigma_c > 0$, $q \in (0, 2]$, and $p \in (-\sigma_c, 2\sigma_c)$. (Notice that for $q \in (2, 4)$ the function f_q satisfies (f1)–(f5) but not (f6).)

We next need to truncate the function f in a neighbourhood of 1 in a smooth way. To this aim we fix points

$$s_\varepsilon \in (0, 1) \quad \text{such that} \quad s_\varepsilon \uparrow 1 \quad \text{and} \quad \sqrt{\varepsilon}f(s_\varepsilon) \rightarrow 1 \quad (1.6)$$

as $\varepsilon \rightarrow 0$. By (f2) and $\sqrt{\varepsilon}f(s_\varepsilon) \rightarrow 1$ we easily deduce that

$$\frac{1 - s_\varepsilon}{\sqrt{\varepsilon}} \rightarrow \sigma_c \quad \text{as } \varepsilon \rightarrow 0. \quad (1.7)$$

We then define the function $f_\varepsilon : [0, 1] \rightarrow \mathbb{R}$ by

$$f_\varepsilon(s) := \begin{cases} \sqrt{\varepsilon}f(s) & \text{if } 0 \leq s \leq s_\varepsilon, \\ \psi_\varepsilon(s) & \text{if } s_\varepsilon < s \leq 1, \end{cases} \quad (1.8)$$

where $\psi_\varepsilon : [0, 1] \rightarrow [0, 1]$ is any function satisfying the following conditions:

(ψ 1) $\psi_\varepsilon \in C^2([0, 1])$ is monotone nondecreasing, $\psi_\varepsilon(1) = 1$;

(ψ 2) $\psi_\varepsilon(s_\varepsilon) = \sqrt{\varepsilon}f(s_\varepsilon)$, $\psi'_\varepsilon(s_\varepsilon) = \sqrt{\varepsilon}f'(s_\varepsilon)$;

(ψ 3) the map $s \mapsto \frac{\psi'_\varepsilon(s)}{1-s}$ is monotone nonincreasing.

Notice that the condition (ψ 3) forces $\psi'_\varepsilon(1) = 0$. The function f_ε in (1.8) is of class $C^1(0, 1)$, and can be easily extended to a globally C^1 -function on \mathbb{R} by setting $f_\varepsilon(s) = \sqrt{\varepsilon}f'(0)s$ for $s < 0$, $f_\varepsilon(s) = 1$ for $s > 1$.

Remark 1.4. An explicit family of functions with the properties above are the exponentials

$$\psi_\varepsilon(s) := 1 - \alpha_\varepsilon e^{-\frac{\beta_\varepsilon}{1-s}}, \quad \alpha_\varepsilon := (1 - \sqrt{\varepsilon}f(s_\varepsilon))e^{\frac{\beta_\varepsilon}{1-s_\varepsilon}}, \quad \beta_\varepsilon := \frac{\sqrt{\varepsilon}f'(s_\varepsilon)(1-s_\varepsilon)^2}{1 - \sqrt{\varepsilon}f(s_\varepsilon)}.$$

The choice of the coefficients α_ε and β_ε guarantees that (ψ 2) holds. Moreover, it can be checked by an elementary computation that also (ψ 3) holds for ε small. See Figure 2 for a numerical plot of the resulting function f_ε .




FIGURE 2. Numerical plot of the function f_ε defined in (1.8), obtained by choosing the prototype function f_q (with $q = 1$) as in Remark 1.3 and the exponential junction ψ_ε as in Remark 1.4.

We next introduce a cohesive energy density $g : [0, +\infty) \rightarrow [0, +\infty)$, depending on f , by solving the following auxiliary optimal profile problem:

$$g(s) := \inf_{(\alpha, \beta) \in \mathcal{U}_s} \mathcal{G}(\alpha, \beta), \quad (1.9)$$

where

$$\mathcal{G}(\alpha, \beta) := \int_{-\infty}^{+\infty} \left(f^2(\beta) |\alpha'|^2 + \frac{(1-\beta)^2}{4} + |\beta'|^2 \right) dt, \quad (1.10)$$

$$\mathcal{U}_s := \left\{ (\alpha, \beta) \in H_{\text{loc}}^1(\mathbb{R}) \times H_{\text{loc}}^1(\mathbb{R}) : \alpha' \in L^1(\mathbb{R}), \left| \int_{-\infty}^{+\infty} \alpha'(t) dt \right| = s, \right. \\ \left. 0 \leq \beta \leq 1, \lim_{|t| \rightarrow +\infty} \beta(t) = 1 \right\}. \quad (1.11)$$

The properties of the function g are discussed in Section 3. Here we just note that g satisfies the conditions we required in Subsection 1.1 for a cohesive surface energy density, see in particular Proposition 3.1. A more operational formula for g is provided in Proposition 3.3. This may be used to obtain the explicit expression of g in the spirit of Remark 3.4, which treats the prototype case $f(t) = t/(1-t)$, $t \in [0, 1)$. The threshold of complete fracture s_{frac} (which is possibly infinite) is explicitly determined in terms of f , see (3.1) and Proposition 3.7.

With the previous definitions, we are now ready to introduce the approximating functionals and to state the main Γ -convergence result from [21]. We let $\mathcal{F}_\varepsilon : L^1(0, L) \times L^1(0, L) \rightarrow [0, +\infty]$ be defined by

$$\mathcal{F}_\varepsilon(u, v) := \begin{cases} \int_0^L \left(f_\varepsilon^2(v) |u'|^2 + \frac{(1-v)^2}{4\varepsilon} + \varepsilon |v'|^2 \right) dx & \text{if } u, v \in H^1(0, L), \\ +\infty & \text{if } 0 \leq v \leq 1 \text{ } \mathcal{L}^1\text{-a.e. in } (0, L), \\ & \text{otherwise.} \end{cases} \quad (1.12)$$

The following Γ -convergence result is proved in [21], see in particular Theorem 3.1, Remark 3.2 and Theorem 3.3.1. Although we state the theorem in dimension one, the result in [21] is actually proved in any dimension and under more general assumptions on f .

Theorem 1.5 (Conti-Focardi-Iurlano). *The functionals \mathcal{F}_ε defined in (1.12) Γ -converge as $\varepsilon \rightarrow 0^+$ in $L^1(0, L) \times L^1(0, L)$ to the functional*

$$\mathcal{F}(u, v) := \begin{cases} \Phi(u) & \text{if } u \in \text{BV}(0, L), v = 1 \text{ } \mathcal{L}^1\text{-a.e. in } (0, L), \\ +\infty & \text{otherwise,} \end{cases} \quad (1.13)$$

where Φ is the cohesive fracture energy given by (1.1) (with elastic energy density ϕ as in (1.2) and surface energy density g as in (1.9)). Moreover, if $(u_\varepsilon, v_\varepsilon)$ satisfies the uniform bound

$$\sup_\varepsilon \left(\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) + \|u_\varepsilon\|_{L^1(0, L)} \right) < +\infty, \quad (1.14)$$

then there exists a subsequence $(u_{\varepsilon_k}, v_{\varepsilon_k})_{k \in \mathbb{N}}$ and a function $u \in \text{BV}(0, L)$ such that $u_{\varepsilon_k} \rightarrow u$ almost everywhere in $(0, L)$ and $v_{\varepsilon_k} \rightarrow 1$ in $L^1(0, L)$.

Remark 1.6. We collect here for later use a few properties that follow immediately from the assumptions (f1)-(f5). Firstly, we observe that the function f is strictly increasing with $f'(s) > 0$ for $s \in (0, 1)$, by (f3). By de l'Hôpital theorem, using (f2) and (f4), one also has that

$$\lim_{s \rightarrow 1^-} (1-s)^2 f'(s) = \sigma_c. \quad (1.15)$$

Finally, by the monotonicity property (f4) one has that the function $\frac{f'(s)}{f(s)}$ has a limit as $s \rightarrow 0^+$, and since $f(0) = 0$ it is easily seen that it must be

$$\lim_{s \rightarrow 0^+} \frac{f'(s)}{f(s)} = +\infty. \quad (1.16)$$

1.3. Main results. We consider a family of *critical points* $(u_\varepsilon, v_\varepsilon)$ of the approximating functionals (1.12), i.e. $u_\varepsilon, v_\varepsilon \in H^1(0, L)$ satisfy the Euler-Lagrange equations (in the weak sense)

$$\begin{cases} -\varepsilon v_\varepsilon'' + f_\varepsilon(v_\varepsilon) f_\varepsilon'(v_\varepsilon) (u_\varepsilon')^2 + \frac{v_\varepsilon - 1}{4\varepsilon} = 0 & \text{in } (0, L), & (1.17a) \\ (f_\varepsilon^2(v_\varepsilon) u_\varepsilon')' = 0 & \text{in } (0, L), & (1.17b) \\ u_\varepsilon(0) = 0, u_\varepsilon(L) = a_\varepsilon, & & (1.17c) \\ v_\varepsilon(0) = v_\varepsilon(L) = 1, & & (1.17d) \end{cases}$$

where for the Dirichlet boundary condition for u_ε we also require that

$$a_\varepsilon \rightarrow a > 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (1.18)$$

In our first main result we show that any such a family of critical points with equibounded energies is precompact and that any limit point is necessarily a critical point of the cohesive energy (1.1). Moreover the Γ -convergence acts as a selection criterion, since the limit critical point has at most one crack point, located at the midpoint of the bar.

Theorem 1.7. *Assume that $(u_\varepsilon, v_\varepsilon)$ are critical points of the functionals \mathcal{F}_ε with*

$$\sup_\varepsilon \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) < +\infty. \quad (1.19)$$

Then there exists a subsequence $\varepsilon_k \rightarrow 0$ such that $(u_{\varepsilon_k}, v_{\varepsilon_k}) \rightarrow (u, 1)$ in $(L^1(0, L))^2$, for some $u \in \text{SBV}(0, L)$ such that $|Du|(0, L) = a$. Moreover, letting $m_0 := \lim_{k \rightarrow \infty} \min_{[0, L]} v_{\varepsilon_k}$, we have that:

- (i) *If $m_0 = 1$, then $u(x) = \frac{a}{L}x$.*
- (ii) *If $m_0 \in (0, 1)$, then $u(x) = c_0x + (a - c_0L)\chi_{(\frac{L}{2}, L)}(x)$ with $c_0 = \frac{1}{2}f(m_0)(1 - m_0) \in (0, \frac{\sigma_c}{2})$, $a - c_0L = [u](\frac{L}{2}) \in (0, s_{\text{frac}})$, and $g'([u](\frac{L}{2})) = 2c_0$.*
- (iii) *If $m_0 = 0$, then $s_{\text{frac}} \in \mathbb{R}$ and $u(x) = a\chi_{(\frac{L}{2}, L)}(x)$ with $a = [u](\frac{L}{2}) = s_{\text{frac}}$.*

Finally if $m_0 < 1$, or if $m_0 = 1$ and $\frac{a}{L} \leq \frac{\sigma_c}{2}$, the convergence of the energies holds:

$$\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \rightarrow \mathcal{F}(u, 1) \quad \text{as } \varepsilon \rightarrow 0. \quad (1.20)$$

If instead $m_0 = 1$ and $\frac{a}{L} > \frac{\sigma_c}{2}$, the convergence of the energies (1.20) does not hold.

Our second result is an existence counterpart of Theorem 1.7: we show that, for any critical point \bar{u} of the cohesive energy (1.1) that might appear as limit of critical points of the functionals \mathcal{F}_ε (that is, those that can be obtained as limits in Theorem 1.7), we can actually *construct* a family $(u_\varepsilon, v_\varepsilon)$ of critical points of \mathcal{F}_ε , with equibounded energy, approximating \bar{u} as $\varepsilon \rightarrow 0$.

Theorem 1.8. *Let $\bar{u} \in \text{SBV}(0, L)$ be either:*

- (i) *$\bar{u}(x) = \frac{a}{L}x$ for some $a > 0$, or*
- (ii) *$\bar{u}(x) = c_0x + (a - c_0L)\chi_{(\frac{L}{2}, L)}(x)$ with $c_0 \in (0, \frac{\sigma_c}{2})$ and $g'(a - c_0L) = 2c_0$, or*
- (iii) *$\bar{u}(x) = a\chi_{(\frac{L}{2}, L)}(x)$ with $a = s_{\text{frac}}$ (if s_{frac} is finite).*

Then for every $\varepsilon > 0$ sufficiently small there exists a critical point $(u_\varepsilon, v_\varepsilon)$ solution to (1.17a)–(1.17d) with $a_\varepsilon \rightarrow a$ and uniformly bounded energy, such that $u_\varepsilon \rightarrow \bar{u}$ in $L^1(0, L)$ as $\varepsilon \rightarrow 0$.

Let us discuss briefly our strategy of proof. Compared with the brittle case [10, 11, 30, 37], the main difficulty in our problem is that the behaviours of u_ε and v_ε are deeply related, meaning that their transitions happen in the same regions with the same scale. With this

idea in mind, we start the proof of Theorem 1.7 and the study of system (1.17a)-(1.17d) by computing u'_ε from (1.17b) and inserting it into (1.17a), so obtaining a second order ODE for v_ε (equation (4.6)). Analysis of the ODE (4.6) (performed in Section 2) shows that v_ε is a symmetric well with minimum $m_\varepsilon = v_\varepsilon(\frac{L}{2}) \in (0, 1]$ and that the interval $\{f_\varepsilon(v_\varepsilon) = \varepsilon^{1/2}f(v_\varepsilon)\}$ shrinks to 0 as $\varepsilon \rightarrow 0$ (Section 4). Also, by definition of such interval, we find $v_\varepsilon \rightarrow 1$ uniformly on compact sets not containing $\frac{L}{2}$. Now, we argue differently depending on the value of $m_0 := \lim_{\varepsilon \rightarrow 0} m_\varepsilon$, that is, on the fact we are in the elastic ($m_0 = 1$), pre-fractured ($m_0 \in (0, 1)$) or fractured ($m_0 = 0$) regime. The richest regime is the pre-fractured one, addressed in Section 5.1. Defining $c_\varepsilon := f_\varepsilon^2(v_\varepsilon)u'_\varepsilon$ and $c_0 := \lim_{\varepsilon \rightarrow 0} c_\varepsilon$, we get $u'_\varepsilon \rightarrow c_0$ uniformly on compact sets not containing $\frac{L}{2}$. This in particular implies $u \in \text{SBV}(0, L)$, $J_u \subset \{\frac{L}{2}\}$, $u' = c_0$ a.e.. We consider the blow-up of u_ε and v_ε around the minimum point $\frac{L}{2}$, setting $\tilde{u}_\varepsilon(t) = u_\varepsilon(\frac{L}{2} + \varepsilon t)$ and $\tilde{v}_\varepsilon(t) = v_\varepsilon(\frac{L}{2} + \varepsilon t)$. Passing to the limit, we get $(\tilde{u}_\varepsilon, \tilde{v}_\varepsilon) \rightarrow (\alpha_{\bar{s}}, \beta_{\bar{s}})$ optimal profile for $g(\bar{s})$, for some $\bar{s} \geq 0$. The most delicate point is to prove that $[u](\frac{L}{2}) = \bar{s}$. This is obtained by studying the bijective continuous map $s \in (0, s_{\text{frac}}) \mapsto m_s \in (0, 1)$, where m_s is the minimum of β_s with (α_s, β_s) optimal profile for $g(s)$. Its inverse can be written in integral form, see (3.7) in Proposition 3.3. Proofs in the elastic and fractured regimes, respectively performed in Sections 5.3 and 5.2, are not based on blow-up arguments, but rather on energy estimates. However, they also require fine ad hoc computations involving formula (3.7): indeed, in the elastic case we need to show that $[u](\frac{L}{2}) = 0$ and in the fractured case that $[u](\frac{L}{2}) = s_{\text{frac}}$.

The proof of the second main Theorem 1.8 is addressed in Section 6. The elastic case is trivial, since we can define $u_\varepsilon(x) = \frac{\alpha}{L}x$, $v_\varepsilon(x) = 1$ in $(0, L)$. The pre-fractured and fractured cases are again ODE-based. Take $c_\varepsilon \in (0, \frac{\sigma_c}{2})$ and set $c_0 := \lim_{\varepsilon \rightarrow 0} c_\varepsilon$. Then, we show that for all $\varepsilon > 0$ we can choose $m_\varepsilon > 0$ such that the unique solution v_ε to the second order ODE (4.6) with initial conditions $v_\varepsilon(\frac{L}{2}) = m_\varepsilon$, $v'_\varepsilon(\frac{L}{2}) = 0$, satisfies in addition $v_\varepsilon(0) = v_\varepsilon(L) = 1$. This strongly uses the analysis of the ODE in Section 2 and a continuous dependence argument on the initial value. Finally, u_ε is easily computed from (1.17b).

Structure of the paper. Section 2 contains a detailed analysis of the ODE that is solved by a critical profile v_ε . In Section 3 we discuss the properties of the cohesive energy density g appearing in the Γ -limit of the functionals \mathcal{F}_ε . In Sections 4 and 5 we give the proof of Theorem 1.7, whereas Section 6 contains the proof of Theorem 1.8.

2. ANALYSIS OF THE ODE SATISFIED BY CRITICAL POINTS

In this section we discuss the properties of the solution to the Cauchy problem

$$\begin{cases} y'' = \frac{1-y}{4} \left[\frac{(2\alpha)^2 f'(y)}{(1-y)f^3(y)} - 1 \right] \end{cases} \quad (2.1a)$$

$$\begin{cases} y(0) = m \end{cases} \quad (2.1b)$$

$$\begin{cases} y'(0) = 0 \end{cases} \quad (2.1c)$$

for fixed parameters $\alpha \in (0, \frac{\sigma_c}{2})$ and $m \in (0, 1)$. As it will be clear in the following of the paper, this equation is satisfied by an optimal profile β_s for the minimum problem defining g (see (1.9) and Section 3) and also corresponds to the blow-up of the critical points equation (1.17a) around a stationary point of v_ε .

In the whole of this section we always assume that the function f appearing in (2.1a) satisfies the conditions (f1)–(f5); in particular, the analysis of the Cauchy problem does not make use of (f6).

It is convenient to introduce the functions

$$\bar{f}(s) := \frac{f'(s)}{(1-s)f^3(s)} \quad (2.2)$$

and

$$h(s) := \frac{1-s}{4} \left[\frac{(2\alpha)^2 f'(s)}{(1-s)f^3(s)} - 1 \right] = \frac{1-s}{4} \left[(2\alpha)^2 \bar{f}(s) - 1 \right] \quad \text{for } s \in (0, 1), \quad (2.3)$$

extended by continuity by setting $h(1) = 0$. With this notation, the equation (2.1a) takes the form $y'' = h(y)$. The constant function $\bar{y} \equiv 1$ is always a stationary solution of the equation (2.1a); in the next proposition we show in particular that the function h has a unique zero $z_\alpha \in (0, 1)$, which corresponds to a second stationary solution $\bar{y} \equiv z_\alpha$.

Proposition 2.1 (Properties of \bar{f}). *The function \bar{f} defined in (2.2) is of class $C^1(0, 1)$ and strictly decreasing, with $\bar{f}'(s) < 0$ for all $s \in (0, 1)$. Moreover*

$$\lim_{s \rightarrow 0^+} \bar{f}(s) = +\infty, \quad \lim_{s \rightarrow 1^-} \bar{f}(s) = \frac{1}{\sigma_c^2}, \quad \lim_{s \rightarrow 0^+} \frac{|\bar{f}'(s)|}{\bar{f}(s)} = +\infty, \quad \lim_{s \rightarrow 1^-} \frac{|\bar{f}'(s)|}{(1-s)^3} = +\infty. \quad (2.4)$$

In particular, it follows that for every $\alpha \in (0, \frac{\sigma_c}{2})$ there exists a unique $z_\alpha \in (0, 1)$ such that

$$\bar{f}(z_\alpha) = \frac{1}{(2\alpha)^2}, \quad (2.5)$$

and moreover $(1 - z_\alpha)f(z_\alpha) > 2\alpha$.

Proof. The regularity of \bar{f} follows by (f1). By computing explicitly the derivative appearing in assumption (f4), one easily obtains the inequality

$$(1-s)f(s)f''(s) - f(s)f'(s) - (1-s)(f'(s))^2 < 0.$$

Then using this inequality we get for all $s \in (0, 1)$

$$\begin{aligned} \bar{f}'(s) &= \frac{(1-s)f(s)f''(s) + f(s)f'(s) - 3(1-s)(f'(s))^2}{(1-s)^2 f^4(s)} \\ &< \frac{2f'(s)[f(s) - (1-s)f'(s)]}{(1-s)^2 f^4(s)} = -\frac{2f'(s)[(1-s)f(s)]'}{(1-s)^2 f^4(s)} < 0, \end{aligned}$$

where the last inequality follows by (f3) (see also Remark 1.6); this proves the strict monotonicity of \bar{f} . The first limit in (2.4) follows from (1.16); the second limit in (2.4) is a consequence of (f2) and (1.15); for the third limit, one has, arguing as before and using (1.16),

$$\frac{|\bar{f}'(s)|}{\bar{f}(s)} > \frac{2f'(s)[(1-s)f'(s) - f(s)]}{(1-s)^2 f^4(s)} \cdot \frac{(1-s)f^3(s)}{f'(s)} = \frac{2f'(s)}{f(s)} - \frac{2}{1-s} \rightarrow +\infty \quad \text{as } s \rightarrow 0^+.$$

Similarly for the fourth limit in (2.4) we use assumption (f5), together with (f2) and (1.15):

$$\frac{|\bar{f}'(s)|}{(1-s)^3} > \frac{2f'(s)[(1-s)f(s)]'}{(1-s)^5 f^4(s)} = \frac{[(1-s)f(s)]'}{(1-s)^3} \cdot \frac{2(1-s)^2 f'(s)}{(1-s)^4 f^4(s)} \rightarrow +\infty \quad \text{as } s \rightarrow 1^-.$$

Existence and uniqueness of z_α are immediate consequences of the strict monotonicity of \bar{f} and of (2.4). To show that $(1 - z_\alpha)f(z_\alpha) > 2\alpha$, we observe that (2.5) and the monotonicity




FIGURE 3. Left: numerical plots of the solution to the Cauchy problem (2.1a)–(2.1c) for different initial values m , for the prototype function $f(s) = \frac{s}{1-s}$ (see (1.5) with $\sigma_c = q = 1$) and $\alpha = 0.2$. The two dashed lines correspond to the stationary solutions $\bar{y} \equiv 1$ and $\bar{y} \equiv z_\alpha$. The green, red and blue curves correspond to cases (i), (ii) and (iii) in Theorem 2.2 respectively. Right: phase diagram of the solutions to (2.1a)–(2.1c) in the plane (y, y') .

assumption (f4) give

$$(1 - z_\alpha)^2 f^2(z_\alpha) = \frac{(1 - z_\alpha) f'(z_\alpha)}{f(z_\alpha)} \cdot (2\alpha)^2 > \lim_{s \rightarrow 1^-} \frac{(1 - s) f'(s)}{f(s)} \cdot (2\alpha)^2 = (2\alpha)^2,$$

where we used (f2) and (1.15) in the last inequality. \square

By Cauchy-Lipschitz theorem, for every value of the initial datum $m \in (0, 1)$ the Cauchy problem (2.1a)–(2.1c) has a unique solution \bar{y} of class C^2 , which can be continued as long as $y(t) \in (0, 1)$ and is defined in a maximal interval $(-T, T)$ for some $T \in (0, +\infty]$. By uniqueness the solution is symmetric with respect to the origin, that is $\bar{y}(t) = \bar{y}(-t)$, and therefore we study the equation only on the positive real axis. In the following theorem we characterize the behaviour of the solution to (2.1a)–(2.1c) in terms on the relation between the parameters m and α . We focus on the case $m \in (0, z_\alpha)$; for the case $m \in (z_\alpha, 1)$, see Remark 2.4.

Theorem 2.2. *Let $\alpha \in (0, \frac{\sigma_c}{2})$ and $m \in (0, z_\alpha)$ be given, let f satisfy (f1)–(f5), and let \bar{y} be the solution to the Cauchy problem (2.1a)–(2.1c). Then there exists $t_0 > 0$ such that $\bar{y}(t_0) = z_\alpha$ and $m < \bar{y}(t) < z_\alpha$ for $t \in (0, t_0)$. Moreover:*

- (i) *if $(1 - m)f(m) < 2\alpha$, then there exists $t_1 \in (t_0, +\infty)$ such that $\bar{y}(t_1) = 1$, and \bar{y} is strictly increasing in $(0, t_1)$;*
- (ii) *if $(1 - m)f(m) = 2\alpha$, then $\bar{y}(t) < 1$ for all $t \in (0, +\infty)$, \bar{y} is strictly increasing, and $\bar{y}(t) \rightarrow 1$ as $t \rightarrow +\infty$;*
- (iii) *if $(1 - m)f(m) > 2\alpha$, then \bar{y} oscillates periodically between its minimum m and a maximum $M \in (z_\alpha, 1)$: more precisely, there exists $t_2 \in (t_0, +\infty)$ such that $\bar{y}(t_2) = M = \max \bar{y}$ and \bar{y} is strictly increasing in $(0, t_2)$. For $t \in (t_2, 2t_2)$ one has $\bar{y}(t) = \bar{y}(2t_2 - t)$, and \bar{y} is periodic with period $2t_2$.*

The values of t_0, t_1, t_2, M in the previous statements depend on m and α .

Proof. Notice that, since $(1 - z_\alpha)f(z_\alpha) > 2\alpha$ by Proposition 2.1 and the map $s \mapsto (1 - s)f(s)$ is strictly increasing by assumption (f3), for $m \in (0, z_\alpha)$ all the three cases may occur. We

also observe that, in view of Proposition 2.1, one has that $h(s) > 0$ for $s \in (0, z_\alpha)$ and $h(s) < 0$ for $s \in (z_\alpha, 1)$.

The solution \bar{y} satisfies $\bar{y}'(0) = 0$ and $\bar{y}''(0) = h(m) > 0$, therefore $\bar{y}' > 0$ and \bar{y} is strictly increasing and convex in a right neighbourhood of the origin. Since \bar{y} remains convex as long as $\bar{y}(t) < z_\alpha$, there exists $t_0 > 0$ such that $\bar{y}(t_0) = z_\alpha$, and \bar{y} is convex and strictly increasing in $(0, t_0)$.

At the point t_0 we have $\bar{y}'(t_0) > 0$ (otherwise $\bar{y} \equiv z_\alpha$) and $\bar{y}''(t_0) = 0$, and therefore $\bar{y}(t) > z_\alpha$ and $\bar{y}''(t) = h(\bar{y}(t)) < 0$ for t in a right neighbourhood of t_0 , so that \bar{y} becomes concave after t_0 . We let

$$t_1 := \sup\{t > 0 : \bar{y}(s) < 1 \text{ for all } s \in (0, t)\}$$

and we observe that $t_1 > t_0$. We can then identify three possible types of solutions.

Case I. If $t_1 \in \mathbb{R}$, then \bar{y} reaches the value 1 in finite time: indeed it must be $\bar{y}(t_1) = 1$, and $\bar{y}'(t_1) > 0$ (it cannot be $\bar{y}'(t_1) = 0$, or else the solution would coincide with the stationary solution constantly equal to 1).

If $t_1 = +\infty$, then the solution is defined in the whole positive real line and $\bar{y}(t) < 1$ for all $t > 0$. We let

$$t_2 := \sup\{t > 0 : \bar{y}'(s) > 0 \text{ for all } s \in (0, t)\}$$

and we observe that $t_2 > t_0$, as $\bar{y}'(t) > 0$ for all $t \in (0, t_0)$.

Case II. If $t_2 = +\infty$, then $\bar{y}'(t) > 0$ for all $t > 0$. Then the solution is strictly increasing in $(0, +\infty)$, is convex in $(0, t_0)$, concave in (t_0, ∞) , and $\bar{y}(t) \rightarrow 1$ as $t \rightarrow +\infty$.

Case III. If $t_2 \in \mathbb{R}$, then $\bar{y}'(t_2) = 0$, $M := \bar{y}(t_2) \in (z_\alpha, 1)$ and $\bar{y}''(t_2) = h(M) < 0$. Then t_2 is a local maximum of \bar{y} . The function $z(t) := \bar{y}(2t_2 - t)$, for $t \in (t_2, 2t_2)$, solves the equation (2.1a) in $(t_2, 2t_2)$, with $z(t_2) = M = \bar{y}(t_2)$ and $z'(t_2) = 0 = \bar{y}'(t_2)$, and therefore by uniqueness of the solution of the Cauchy problem it must be $\bar{y}(t) = z(t) = \bar{y}(2t_2 - t)$, for $t \in (t_2, 2t_2)$. At the point $2t_2$ we have $\bar{y}(2t_2) = m$, $\bar{y}'(2t_2) = 0$, and again by uniqueness we can conclude that $\bar{y}(t) = \bar{y}(t - 2t_2)$ for all $t > 2t_2$, that is, \bar{y} is periodic with period $2t_2$.

These are the only three possible behaviours of solutions to (2.1a)–(2.1c), for $m \in (0, z_\alpha)$. See Figure 3 for numerical plots of the three types of solutions. To conclude the proof, we only need to determine the form of the solution in terms of the value of the initial datum. We let

$$\tilde{t} := \begin{cases} t_1 & \text{in Case I,} \\ +\infty & \text{in Case II,} \\ t_2 & \text{in Case III.} \end{cases}$$

Since $\bar{y}' > 0$ in $(0, \tilde{t})$ in all cases, we can multiply the equation (2.1a) by \bar{y}' and integrate in $(0, t)$, for $t < \tilde{t}$: we obtain by a change of variables

$$\int_0^t \bar{y}''(\tau) \bar{y}'(\tau) \, d\tau = \int_0^t h(\bar{y}(\tau)) \bar{y}'(\tau) \, d\tau = \int_m^{\bar{y}(t)} h(s) \, ds,$$

and, recalling that $\bar{y}'(0) = 0$,

$$\begin{aligned} \frac{1}{2}(\bar{y}'(t))^2 &= \int_m^{\bar{y}(t)} h(s) \, ds = \int_m^{\bar{y}(t)} \frac{1-s}{4} \left[\frac{(2\alpha)^2 f'(s)}{(1-s)f^3(s)} - 1 \right] \, ds \\ &= \frac{1}{4} \int_m^{\bar{y}(t)} \left((2\alpha)^2 \frac{f'(s)}{f^3(s)} - 1 + s \right) \, ds \\ &= \frac{1}{4} \left(-\frac{(2\alpha)^2}{2f^2(\bar{y}(t))} + \frac{(2\alpha)^2}{2f^2(m)} - \bar{y}(t) + \frac{1}{2}\bar{y}^2(t) + m - \frac{1}{2}m^2 \right). \end{aligned}$$

Therefore for all $t \in (0, \tilde{t})$

$$(\bar{y}')^2 = \frac{1}{4} \left[(2\alpha)^2 \left(\frac{1}{f^2(m)} - \frac{1}{f^2(\bar{y})} \right) - (1-m)^2 + (1-\bar{y})^2 \right]. \quad (2.6)$$

If we are in Case I, then as $t \rightarrow \tilde{t} = t_1$ we have $\bar{y}(t) \rightarrow 1$ and $\bar{y}'(t) \rightarrow \bar{y}'(t_1) > 0$, therefore by letting $t \rightarrow t_1$ in (2.6) we obtain

$$0 < (\bar{y}'(t_1))^2 = \frac{1}{4} \left[\frac{(2\alpha)^2}{f^2(m)} - (1-m)^2 \right] = \frac{(1-m)^2}{4} \left[\left(\frac{2\alpha}{(1-m)f(m)} \right)^2 - 1 \right]$$

and hence $(1-m)f(m) < 2\alpha$.

If we are in Case II, by letting $t \rightarrow +\infty$ in (2.6) we obtain, since $\bar{y}(t) \rightarrow 1$ and $\bar{y}'(t) \rightarrow 0$,

$$0 = \lim_{t \rightarrow +\infty} (\bar{y}'(t))^2 = \frac{(1-m)^2}{4} \left[\left(\frac{2\alpha}{(1-m)f(m)} \right)^2 - 1 \right]$$

and hence $(1-m)f(m) = 2\alpha$.

Finally, if we are in Case III, by letting $t \rightarrow \tilde{t} = t_2$ in (2.6) we have, since $\bar{y}'(t_2) = 0$ and $\bar{y}(t_2) = M \in (z_\alpha, 1)$,

$$0 = \frac{1}{4} \left[(2\alpha)^2 \left(\frac{1}{f^2(m)} - \frac{1}{f^2(M)} \right) - (1-m)^2 + (1-M)^2 \right],$$

from which we get

$$\begin{aligned} (2\alpha)^2 &= \left(\frac{(1-m)^2 - (1-M)^2}{f^2(M) - f^2(m)} \right) f^2(m) f^2(M) \\ &= \left(\frac{f^2(M)(1-m)^2 - f^2(m)(1-M)^2}{(1-m)^2(f^2(M) - f^2(m))} \right) (f(m)(1-m))^2 \\ &< \left(\frac{f^2(M)(1-m)^2 - f^2(m)(1-m)^2}{(1-m)^2(f^2(M) - f^2(m))} \right) (f(m)(1-m))^2 = (f(m)(1-m))^2, \end{aligned}$$

where we used the fact that $f^2(M)(1-M)^2 > f^2(m)(1-m)^2$ since $M > m$ and the function $s \mapsto f(s)(1-s)$ is strictly increasing by (f3). Therefore $(1-m)f(m) > 2\alpha$.

Since these are the only possible cases, the characterization of \bar{y} in the statement holds. \square

Remark 2.3. Let $m_\alpha \in (0, z_\alpha)$ be the unique value such that $(1-m_\alpha)f(m_\alpha) = 2\alpha$. The three cases (i), (ii) and (iii) in Theorem 2.2 correspond to $m \in (0, m_\alpha)$, $m = m_\alpha$, and $m \in (m_\alpha, z_\alpha)$ respectively. If $m \in (m_\alpha, z_\alpha)$ (case (iii)) the solution has a maximum value $M \in (z_\alpha, 1)$ which is uniquely determined by α and m by the equation

$$(1-M)^2 - \frac{(2\alpha)^2}{f^2(M)} = (1-m)^2 - \frac{(2\alpha)^2}{f^2(m)}, \quad (2.7)$$

which is obtained by evaluating (2.6) at the maximum point t_2 . We observe that the function

$$x \mapsto (1-x)^2 - \frac{(2\alpha)^2}{f^2(x)}$$

vanishes at $x = m_\alpha$ and at $x = 1$, has a maximum point at $x = z_\alpha$, is increasing in (m_α, z_α) and decreasing in $(z_\alpha, 1)$, as can be easily checked by noting that its derivative is given by $2(1-x)((2\alpha)^2 \bar{f}(x) - 1)$.

It follows that M takes all the values in the interval $(z_\alpha, 1)$ when $m \in (m_\alpha, z_\alpha)$, with $M \rightarrow 1$ as $m \rightarrow m_\alpha$, and $M \rightarrow z_\alpha$ as $m \rightarrow z_\alpha$.

Remark 2.4. In the case $m \in (z_\alpha, 1)$, the solution \bar{y} is a translation of one of the periodic orbits obtained in Theorem 2.2(iii). Indeed, by Remark 2.3 every value of $m \in (z_\alpha, 1)$ corresponds to the maximum value of one of the solutions with initial datum in (m_α, z_α) , so that by uniqueness \bar{y} is a translation of that solution.

In the next proposition we obtain uniform estimates on the point t_η at which the solution to the Cauchy problem (2.1a)–(2.1c) reaches a value η arbitrarily close to 1 (or the maximum M for the periodic solutions). In the following, we will denote by $\omega(\cdot)$ a generic modulus of continuity (that is, $\omega : [0, 1] \rightarrow \mathbb{R}$ is a bounded, monotone increasing function such that $\omega(t) \rightarrow 0$ as $t \rightarrow 0^+$), which is independent of $\alpha \in (0, \frac{\sigma_c}{2})$ and $m \in (0, z_\alpha)$, but depends ultimately only on f . The function ω might change from line to line.

Proposition 2.5. *Under the assumptions of Theorem 2.2, let y be the solution to (2.1a)–(2.1c). Let $\eta \in (z_\alpha, 1)$ be fixed; in the case $(1-m)f(m) > 2\alpha$, assume further that $\eta \in (z_\alpha, M)$, where $M \in (z_\alpha, 1)$ is the maximum of y . Denote by $t_\eta > 0$ the first point such that $y(t_\eta) = \eta$.*

Then there exists a constant $C_\alpha > 0$, depending only on $\alpha \in (0, \frac{\sigma_c}{2})$, and a modulus of continuity ω independent of α and m , such that the following estimates hold:

$$t_\eta \leq \frac{C_\alpha}{\sqrt{1-\eta}} \quad \text{with} \quad \sup_{\alpha \in (0, \frac{\sigma_c}{2} - \delta)} C_\alpha < +\infty \quad \text{for all } \delta > 0, \quad (2.8)$$

$$t_\eta \leq \frac{\omega(1-m)}{(1-\eta)^2}. \quad (2.9)$$

Furthermore, we have the estimate from below

$$t_\eta \geq \log \left(\frac{1 - z_\alpha + C_{\alpha,m}}{1 - \eta + C_{\alpha,m}} \right), \quad \text{where} \quad C_{\alpha,m} := \left| 1 - \left(\frac{2\alpha}{(1-m)f(m)} \right)^2 \right|^{\frac{1}{2}}. \quad (2.10)$$

Proof. We first obtain a uniform estimate of the point t_0 where there is a change of convexity of the solution y , as in the statement of Theorem 2.2: recall that for $t \in (0, t_0)$ the solution is strictly increasing with $m < y(t) < z_\alpha$.

From the proof of Theorem 2.2, see in particular (2.6), we have that

$$(y')^2 = \frac{(1-y)^2}{4} \left[1 - \left(\frac{2\alpha}{(1-y)f(y)} \right)^2 \right] - \frac{(1-m)^2}{4} \left[1 - \left(\frac{2\alpha}{(1-m)f(m)} \right)^2 \right] =: \Psi(y). \quad (2.11)$$

By a straightforward computation and recalling the definition (2.2) of \bar{f} and (2.5) we find

$$\Psi'(s) = \frac{1}{2}(2\alpha)^2(1-s)(\bar{f}(s) - \bar{f}(z_\alpha)) \quad (2.12)$$

and in particular $\Psi'(s) > 0$ for $s \in (m, z_\alpha)$ since \bar{f} is strictly decreasing (see Proposition 2.1). By repeatedly applying the mean value theorem we have for all $s \in (m, z_\alpha)$ (using also $\Psi(m) = 0$)

$$\begin{aligned} \Psi(s) &= \Psi(m) + \Psi'(\xi)(s - m) && \text{for some } \xi \in (m, s) \\ &= \frac{1}{2}(2\alpha)^2(1 - \xi)(\bar{f}(\xi) - \bar{f}(z_\alpha))(s - m) \\ &= \frac{1}{2\bar{f}(z_\alpha)}(1 - \xi)(-\bar{f}'(\zeta))(z_\alpha - \xi)(s - m) && \text{for some } \zeta \in (\xi, z_\alpha) \\ &\geq \frac{1}{2}\left(-\frac{\bar{f}'(\zeta)}{\bar{f}(\zeta)}\right)(1 - z_\alpha)(z_\alpha - s)(s - m), \end{aligned} \quad (2.13)$$

where we used in particular (2.5) in the third equality, and the monotonicity of the function \bar{f} in the last inequality. Observe now that, being $\bar{f}' < 0$ and continuous in $(0, 1)$, and by (2.4), the ratio $|\bar{f}'|/\bar{f}$ is uniformly bounded from below by a positive constant in $(0, \frac{1}{2})$, whereas if $\zeta \in (\frac{1}{2}, 1)$ one has, by the fourth limit in (2.4), that $|\bar{f}'(\zeta)|/\bar{f}(\zeta) \geq |\bar{f}'(\zeta)|/\bar{f}(\frac{1}{2}) \geq (1 - \zeta)^3/\omega(1 - m)$, for a uniform modulus of continuity ω . Therefore we can write

$$\inf_{\zeta \in (m, z_\alpha)} \frac{|\bar{f}'(\zeta)|}{\bar{f}(\zeta)} \geq \frac{(1 - z_\alpha)^3}{\omega(1 - m)}. \quad (2.14)$$

Hence combining (2.11), (2.13), and (2.14) we find

$$t_0 = \int_0^{t_0} \frac{y'(t)}{\sqrt{\Psi(y(t))}} dt = \int_m^{z_\alpha} \frac{ds}{\sqrt{\Psi(s)}} \leq \frac{\omega(1 - m)}{(1 - z_\alpha)^2} \int_m^{z_\alpha} \frac{ds}{\sqrt{z_\alpha - s}\sqrt{s - m}} \quad (2.15)$$

which eventually yields

$$t_0 \leq \pi \frac{\omega(1 - m)}{(1 - z_\alpha)^2}. \quad (2.16)$$

We next fix $\eta > z_\alpha$ as in the statement. By the properties of the solution, there exists a point $t_\eta > t_0$ such that $y(t_\eta) = \eta$; in the interval (t_0, t_η) the solution satisfies $z_\alpha < y(t) < \eta$, $y'(t) > 0$. As before we have from (2.11)

$$t_\eta - t_0 = \int_{t_0}^{t_\eta} \frac{y'(t)}{\sqrt{\Psi(y(t))}} dt = \int_{z_\alpha}^\eta \frac{ds}{\sqrt{\Psi(s)}}, \quad (2.17)$$

and therefore we need to estimate from below $\Psi(s)$ for $s \in (z_\alpha, \eta)$. Notice that by (2.12) and monotonicity of \bar{f} , Ψ is decreasing in $(z_\alpha, 1)$. For all $s \in (z_\alpha, \eta)$ we have by the mean value theorem, using also $\Psi(\eta) \geq 0$, (2.5), and (2.12),

$$\begin{aligned} \Psi(s) &= \Psi(\eta) + \Psi'(\xi)(s - \eta) \\ &\geq \frac{1}{2}(2\alpha)^2(1 - \xi)(\bar{f}(z_\alpha) - \bar{f}(\xi))(\eta - s) \\ &\geq \frac{1}{2}(1 - \eta)(\eta - s) \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\bar{f}(z_\alpha)} \end{aligned} \quad (2.18)$$

for some $\xi \in (s, \eta)$. We next bound from below the quotient on the right-hand side of (2.18), and we consider first the case α uniformly bounded away from $\frac{\sigma_c}{2}$ (which will prove (2.8)), and then the case α in a small neighbourhood of $\frac{\sigma_c}{2}$ (which will prove (2.9)). It is important to recall from Proposition 2.1 that z_α depends continuously and monotonically on α , and that $z_\alpha \rightarrow 1$ as $\alpha \rightarrow \frac{\sigma_c}{2}$.

Notice first that

$$\text{if } \frac{\bar{f}(\xi)}{\bar{f}(z_\alpha)} \leq \frac{1}{2} \quad \text{then} \quad \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\bar{f}(z_\alpha)} = 1 - \frac{\bar{f}(\xi)}{\bar{f}(z_\alpha)} \geq \frac{1}{2} \geq \frac{1}{2}(\xi - z_\alpha). \quad (2.19)$$

Let $\delta > 0$ be such that $\alpha < \frac{\sigma_c}{2} - \delta$. Then $z_\alpha < 1 - c_\delta$ for some $c_\delta > 0$ depending only on δ . We have that

$$\text{if } \xi \geq 1 - \frac{c_\delta}{2} \quad \text{then} \quad \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\bar{f}(z_\alpha)} \geq 1 - \frac{\bar{f}(1 - \frac{c_\delta}{2})}{\bar{f}(1 - c_\delta)} =: C_\delta^1 \geq C_\delta^1(\xi - z_\alpha), \quad (2.20)$$

with $C_\delta^1 > 0$ depending only on δ , by the properties of \bar{f} . If instead $\frac{\bar{f}(\xi)}{\bar{f}(z_\alpha)} \geq \frac{1}{2}$ and $\xi \leq 1 - \frac{c_\delta}{2}$ we obtain

$$\begin{aligned} \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\bar{f}(z_\alpha)} &\geq \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\xi - z_\alpha} \cdot \frac{\xi - z_\alpha}{2\bar{f}(\xi)} = -\bar{f}'(\zeta) \cdot \frac{\xi - z_\alpha}{2\bar{f}(\xi)} && \text{for some } \zeta \in (z_\alpha, \xi) \\ &\geq \frac{|\bar{f}'(\zeta)|}{2\bar{f}(\zeta)} \cdot (\xi - z_\alpha) \geq \left(\inf_{(0, 1 - \frac{c_\delta}{2})} \frac{|\bar{f}'|}{2\bar{f}} \right) (\xi - z_\alpha) =: C_\delta^2(\xi - z_\alpha) \end{aligned}$$

where again $C_\delta^2 > 0$ by the properties of \bar{f} in Proposition 2.1. Hence

$$\text{if } \frac{\bar{f}(\xi)}{\bar{f}(z_\alpha)} \geq \frac{1}{2} \quad \text{and} \quad \xi \leq 1 - \frac{c_\delta}{2} \quad \text{then} \quad \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\bar{f}(z_\alpha)} \geq C_\delta^2(\xi - z_\alpha). \quad (2.21)$$

Therefore by inserting (2.19), (2.20), and (2.21) into (2.18) we find that for all $\alpha < \frac{\sigma_c}{2} - \delta$

$$\Psi(s) \geq C_\delta(1 - \eta)(\eta - s)(\xi - z_\alpha) \geq C_\delta(1 - \eta)(\eta - s)(s - z_\alpha) \quad \text{for all } s \in (z_\alpha, \eta)$$

for a uniform constant $C_\delta > 0$ depending only on δ . In turn by (2.17) we conclude that

$$t_\eta - t_0 \leq \frac{1}{\sqrt{C_\delta}\sqrt{1 - \eta}} \int_{z_\alpha}^\eta \frac{ds}{\sqrt{\eta - s}\sqrt{s - z_\alpha}} = \frac{\pi}{\sqrt{C_\delta}\sqrt{1 - \eta}}$$

which combined with (2.16) proves the estimate (2.8) in the statement.

We next show (2.9). If $\frac{\bar{f}(\xi)}{\bar{f}(z_\alpha)} \geq \frac{1}{2}$ then by using the last condition in (2.4) and arguing similarly to (2.14) we have

$$\begin{aligned} \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\bar{f}(z_\alpha)} &\geq \frac{\bar{f}(z_\alpha) - \bar{f}(\xi)}{\xi - z_\alpha} \cdot \frac{\xi - z_\alpha}{2\bar{f}(\xi)} = -\bar{f}'(\zeta) \cdot \frac{\xi - z_\alpha}{2\bar{f}(\xi)} && \text{for some } \zeta \in (z_\alpha, \xi) \\ &\geq \frac{|\bar{f}'(\zeta)|}{2\bar{f}(\zeta)} \cdot (\xi - z_\alpha) \geq \frac{(1 - \eta)^3}{\omega(1 - z_\alpha)} \cdot (\xi - z_\alpha) \end{aligned}$$

for a uniform modulus of continuity ω . Therefore combining this estimate with (2.19) and inserting them into (2.18) we can write

$$\Psi(s) \geq \frac{(1 - \eta)^4}{\omega(1 - z_\alpha)}(\eta - s)(\xi - z_\alpha) \geq \frac{(1 - \eta)^4}{\omega(1 - z_\alpha)}(\eta - s)(s - z_\alpha) \quad \text{for all } s \in (z_\alpha, \eta)$$

and in turn by (2.17) we conclude that

$$t_\eta - t_0 \leq \frac{\omega(1 - z_\alpha)}{(1 - \eta)^2} \int_{z_\alpha}^\eta \frac{ds}{\sqrt{\eta - s}\sqrt{s - z_\alpha}} = \frac{\pi\omega(1 - z_\alpha)}{(1 - \eta)^2}.$$

By combining this estimate with (2.16) we obtain (2.9).

We eventually prove the estimate from below (2.10). By the assumptions (f2)–(f3) we have that $(1-s)f(s) \leq \sigma_c$ for all $s \in (0, 1)$, so that we can bound the function Ψ in (2.11) by

$$\Psi(s) \leq \frac{(1-s)^2}{4} \left[1 - \left(\frac{2\alpha}{\sigma_c} \right)^2 \right] + \frac{(1-m)^2}{4} \left| 1 - \left(\frac{2\alpha}{(1-m)f(m)} \right)^2 \right|$$

and in turn we obtain the rough estimate $\sqrt{\Psi(s)} \leq (1-s) + C_{\alpha,m}$, where $C_{\alpha,m}$ is the constant defined in (2.10). By inserting this inequality into (2.17) and integrating we deduce (2.10) by an elementary computation. \square

We conclude this section by discussing the behaviour of solutions to (2.1a)–(2.1c) in the case $\alpha \geq \frac{\sigma_c}{2}$.

Proposition 2.6. *Let $\alpha \geq \frac{\sigma_c}{2}$ and $m \in (0, 1)$ be given, let f satisfy (f1)–(f5), and let \bar{y} be the solution to the Cauchy problem (2.1a)–(2.1c). Then there exists $t_1 > 0$ such that $\bar{y}(t_1) = 1$ and $\bar{y}(t)$ is monotone increasing in $(0, t_1)$. Moreover, denoting for $\eta \in (m, 1)$ by $t_\eta \in (0, t_1)$ the point such that $\bar{y}(t_\eta) = \eta$, the estimate (2.9) holds.*

Proof. By the properties of \bar{f} in Proposition 2.1 and since $\alpha \geq \frac{\sigma_c}{2}$, the function h in (2.3) is strictly positive in $(0, 1)$. Therefore, since initially we have $\bar{y}'(0) = 0$ and $\bar{y}''(0) = h(m) > 0$, the solution is strictly increasing and convex for $t > 0$ as long as $\bar{y} \in (0, 1)$. The existence of a point $t_1 > 0$ such that $\bar{y}(t_1) = 1$ follows easily.

Fix now $\tilde{\alpha} \in (0, \frac{\sigma_c}{2})$ such that $m \in (0, z_{\tilde{\alpha}})$ and $(1-m)f(m) < 2\tilde{\alpha}$, and let \tilde{y} be the solution to the Cauchy problem (2.1a)–(2.1c) with α replaced by $\tilde{\alpha}$. Since $\alpha \geq \frac{\sigma_c}{2} > \tilde{\alpha}$, by comparison we have $\tilde{y} \leq \bar{y}$.

We can apply to \tilde{y} the qualitative analysis in Theorem 2.2 and Proposition 2.5 to deduce, in particular, that for all $\eta \in (m, 1)$ there exists \tilde{t}_η such that $\tilde{y}(\tilde{t}_\eta) = \eta$, and \tilde{t}_η obeys the estimate (2.9). Then by comparison $t_\eta \leq \tilde{t}_\eta$ obeys the same estimate, as desired. \square

3. PROPERTIES OF THE COHESIVE ENERGY DENSITY, OLD AND NEW

We discuss in this section the main properties of the cohesive energy density g defined by (1.9), appearing in the Γ -limit of the phase-field functionals (1.12). Most of the properties of g have been studied in [12, 21], but we also include here some new results: in Proposition 3.6 we prove that $g \in C^1([0, +\infty))$ and a characterization of the derivative of g , and in Proposition 3.7 we determine explicitly the value of the threshold of complete fracture s_{frac} (see (3.1)). Eventually we determine the asymptotic expansion of g near the origin in terms of the behaviour of $f(s)$ as $s \rightarrow 1$, see Proposition 3.9: this is only relevant in connection with the existence of critical points of pre-fractured type, see Remark 1.2.

The following properties are proved in [21, Proposition 4.1].

Proposition 3.1. *Assume that f satisfies the assumptions (f1), (f2), (f3). The function g defined in (1.9) enjoys the following properties:*

- (i) $g(0) = 0$, and g is subadditive;
- (ii) g is nondecreasing, $g(s) \leq \min\{1, \sigma_c s\}$, and g is Lipschitz continuous with Lipschitz constant σ_c ;
- (iii) $\lim_{s \rightarrow +\infty} g(s) = 1$;
- (iv) $\lim_{s \rightarrow 0^+} \frac{g(s)}{s} = \sigma_c$.

We studied in [12] the existence and properties of optimal pairs for the minimum problem (1.9) which defines g , which we collect in the following two propositions. Notice that the first

part of the statement of Proposition 3.2 does not make use of the convexity assumption (f6), whereas this condition is crucial to deduce uniqueness of the optimal pair and the properties of the function \bar{m}_s in Proposition 3.3 (compare with Remark 3.5).

Proposition 3.2. *Assume that f satisfies the assumptions (f1), (f2), (f3). Let g be defined by (1.9) and let*

$$s_{\text{frac}} := \sup\{s : g(s) < 1\} \in (0, +\infty]. \quad (3.1)$$

Let $s \in (0, s_{\text{frac}})$, so that $g(s) < 1$. Then:

- (i) *There exists an optimal pair $(\alpha_s, \beta_s) \in \mathcal{U}_s$ such that $g(s) = \mathcal{G}(\alpha_s, \beta_s)$.*
- (ii) *If (α_s, β_s) is an optimal pair for $g(s)$, then there exist $-\infty \leq T_- < T_* < T_+ \leq +\infty$ such that $\{\beta_s < 1\} = (T_-, T_+)$, $\beta_s \in C^1(T_-, T_+)$, β_s is symmetric with respect to T_* and nonincreasing in $(-\infty, T_*)$, $\alpha_s \in C^1(\mathbb{R})$ is nondecreasing.*
- (iii) *Any optimal pair (α_s, β_s) solves the Euler-Lagrange equations*

$$\beta_s'' = f(\beta_s)f'(\beta_s)(\alpha_s')^2 - \frac{1 - \beta_s}{4} \quad \text{weakly in } \{\beta_s < 1\}, \quad (3.2)$$

$$f^2(\beta_s)\alpha_s' = \sigma_s \quad \text{pointwise on } \{\beta_s < 1\}, \text{ for a constant } \sigma_s \in \mathbb{R}. \quad (3.3)$$

- (iv) *Any optimal pair (α_s, β_s) satisfies pointwise on $\{\beta_s < 1\}$*

$$f^2(\beta_s)|\alpha_s'|^2 + |\beta_s'|^2 - \frac{(1 - \beta_s)^2}{4} = 0. \quad (3.4)$$

- (v) *Assuming in addition that f satisfies (f6), the optimal pair is unique up to translations, in the sense that if (α_s, β_s) and $(\hat{\alpha}_s, \hat{\beta}_s)$ are minimizers then there are $a_1, t_1 \in \mathbb{R}$ such that $\alpha_s(t) = a_1 + \hat{\alpha}_s(t - t_1)$, $\beta_s(t) = \hat{\beta}_s(t - t_1)$.*

If instead $s \geq s_{\text{frac}}$, one has $g(s) = 1$, and $g(s) = \mathcal{G}(0, \beta_s)$, where $\beta_s(t) = 1 - e^{-\frac{|t|}{2}}$.

Proof. See the proof of [12, Proposition 8.1], which only uses (f1)-(f3), and [12, Proposition 8.3] for the uniqueness part (v). \square

Proposition 3.3. *Assume that f satisfies the assumptions (f1)-(f6). For $s \in (0, s_{\text{frac}})$ let (α_s, β_s) be the optimal pair for $g(s)$ given by Proposition 3.2, translated so that β_s attains its minimum at the origin; for $s \geq s_{\text{frac}}$ let $\beta_s(t) = 1 - e^{-\frac{|t|}{2}}$. Define*

$$\bar{m}_s := \min_{t \in \mathbb{R}} \beta_s(t) = \beta_s(0) \quad (3.5)$$

(uniquely determined by s). The map $s \mapsto \bar{m}_s$ is continuous, strictly decreasing in $[0, s_{\text{frac}})$, with $\bar{m}_0 = 1$ and $\bar{m}_s = 0$ for $s \geq s_{\text{frac}}$.

Moreover, one has that $\{\beta_s < 1\} = \mathbb{R}$, and the constant σ_s in (3.3) is given by

$$\sigma_s = \frac{1}{2}(1 - \bar{m}_s)f(\bar{m}_s). \quad (3.6)$$

Denoting by $m \mapsto s(m)$ the inverse of the map $s \mapsto \bar{m}_s$, one has for all $m \in (0, 1)$

$$s(m) = 2(1 - m)f(m) \int_m^1 \frac{1}{f(t)((1 - t)^2 f^2(t) - (1 - m)^2 f^2(m))^{1/2}} dt, \quad (3.7)$$

$$g(s(m)) = 2 \int_m^1 \frac{(1 - t)^2 f(t)}{((1 - t)^2 f^2(t) - (1 - m)^2 f^2(m))^{1/2}} dt. \quad (3.8)$$

Proof. The first part of the statement concerning the properties of the map \bar{m}_s is proved in [12, Proposition 8.3] without the hypotheses (f4)-(f5). By substituting (3.3) into (3.4) and evaluating the resulting equation at the origin, recalling that $\beta_s(0) = \bar{m}_s$ and $\beta'_s(0) = 0$, we find (3.6). Inserting (3.3) into (3.2) we find that β_s solves in the set $\{\beta_s < 1\}$ the Cauchy problem (2.1a)–(2.1c) for the values of the parameters $\alpha = \sigma_s$ and $m = \bar{m}_s$. By (3.6) we can apply case (ii) in Theorem 2.2 to deduce that $\beta_s < 1$ in \mathbb{R} and $\beta'_s > 0$ in $(0, +\infty)$.

Finally, we prove (3.7) and (3.8). We have by a change of variable

$$\begin{aligned} s &= \int_{\mathbb{R}} \alpha'_s dt \stackrel{(3.3)}{=} 2 \int_0^{+\infty} \frac{\sigma_s}{f^2(\beta_s)} dt \stackrel{(3.4)}{=} 2\sigma_s \int_0^{+\infty} \frac{\beta'_s}{f^2(\beta_s)} \left(\frac{(1-\beta_s)^2}{4} - \frac{\sigma_s^2}{f^2(\beta_s)} \right)^{-\frac{1}{2}} dt \\ &= 2\sigma_s \int_{\bar{m}_s}^1 \frac{1}{f^2(t)} \left(\frac{(1-t)^2}{4} - \frac{\sigma_s^2}{f^2(t)} \right)^{-\frac{1}{2}} dt = 4\sigma_s \int_{\bar{m}_s}^1 \frac{dt}{f(t)[(1-t)^2 f^2(t) - 4\sigma_s^2]^{1/2}} \end{aligned}$$

from which we obtain (3.7) thanks to (3.6). Similarly

$$\begin{aligned} g(s) &= \mathcal{G}(\alpha_s, \beta_s) \stackrel{(3.4)}{=} 2 \int_{-\infty}^{+\infty} \frac{(1-\beta_s)^2}{4} dt = \int_0^{+\infty} (1-\beta_s)^2 dt \\ &\stackrel{(3.4)}{=} \int_0^{+\infty} (1-\beta_s)^2 \beta'_s \left(\frac{(1-\beta_s)^2}{4} - \frac{\sigma_s^2}{f^2(\beta_s)} \right)^{-\frac{1}{2}} dt \\ &= \int_{\bar{m}_s}^1 (1-t)^2 \left(\frac{(1-t)^2}{4} - \frac{\sigma_s^2}{f^2(t)} \right)^{-\frac{1}{2}} dt = \int_{\bar{m}_s}^1 \frac{2(1-t)^2 f(t)}{[(1-t)^2 f^2(t) - 4\sigma_s^2]^{1/2}} dt \end{aligned}$$

from which we obtain (3.8) thanks to (3.6). \square

Remark 3.4. In the prototype case $f(t) = t/(1-t)$, $t \in [0, 1)$, equations (3.7)–(3.8) give explicit formulas for the maps $m \in (0, 1) \mapsto s(m) \in (0, s_{\text{frac}})$ and $m \in (0, 1) \mapsto g(s(m)) \in (0, 1)$ (notice that in this case $s_{\text{frac}} = \pi$ by Proposition 3.9 below):

$$\begin{aligned} s(m) &= 2 \arctan \left(\frac{\sqrt{1-m^2}}{m} \right) - 2m \log \left(\frac{1+\sqrt{1-m^2}}{m} \right), \\ g(s(m)) &= m^2 \log \left(\frac{m}{\sqrt{1-m^2}+1} \right) + \sqrt{1-m^2}. \end{aligned}$$

See also Figure 4 for numeric plots obtained using these expressions.

Remark 3.5. If we drop the assumption (f6), there is numerical evidence (using the function f_q in (1.5) for $q > 2$) that the conclusions of Proposition 3.3 are no longer true in general. However, it seems that it is still possible to define a one-to-one correspondence $s \in (0, s_{\text{frac}}) \mapsto \bar{m}_s \in (m_{\text{frac}}, 1)$, for some $m_{\text{frac}} > 0$.

We next show that g is differentiable and characterize its derivative in terms of the minimum value \bar{m}_s (see (3.5)) of the optimal profile β_s .

Proposition 3.6. *Under the assumptions (f1)–(f6), the function g defined in (1.9) satisfies*

$$g'(s) = (1 - \bar{m}_s) f(\bar{m}_s) \quad \text{for all } s \geq 0. \quad (3.9)$$

Proof. We first observe that the identity (3.9) is valid for $s = 0$ since $g'(0) = \sigma_c$ by Proposition 3.1, $\bar{m}_0 = 1$ by Proposition 3.3, and $\lim_{t \rightarrow 1} (1-t)f(t) = \sigma_c$ by assumption (f2). It is also valid for $s > s_{\text{frac}}$, since $g(s) \equiv 1$ and $\bar{m}_s = 0$ for $s \in [s_{\text{frac}}, +\infty)$.




FIGURE 4. Left: plots of the functions $s(m)$ and $g(s(m))$ in (3.7)–(3.8), in the prototype case $f(t) = t/(1-t)$, see Remark 3.4. Right: plot of the function $g(s)$ obtained by using the expressions in Remark 3.4 and inverting numerically the function $s(m)$.

We then consider the case $s \in (0, s_{\text{frac}})$. Let $(\alpha_s, \beta_s) \in \mathcal{U}_s$ be an optimal pair for $g(s)$, according to Proposition 3.2. Let $s, s' \in (0, s_{\text{frac}})$ be two arbitrary points. Since $(\frac{s}{s'}\alpha_{s'}, \beta_{s'}) \in \mathcal{U}_s$ is an admissible competitor for the minimum problem (1.9) defining $g(s)$, we have

$$\begin{aligned}
 g(s) &\leq \mathcal{G}\left(\frac{s}{s'}\alpha_{s'}, \beta_{s'}\right) = \int_{-\infty}^{+\infty} \left(\left(\frac{s}{s'}\right)^2 f^2(\beta_{s'}) |\alpha'_{s'}|^2 + \frac{(1-\beta_{s'})^2}{4} + |\beta'_{s'}|^2 \right) dt \\
 &= \mathcal{G}(\alpha_{s'}, \beta_{s'}) + \left(\left(\frac{s}{s'}\right)^2 - 1 \right) \int_{-\infty}^{+\infty} f^2(\beta_{s'}) |\alpha'_{s'}|^2 dt \\
 &\stackrel{(3.3)}{=} g(s') + (s-s') \cdot \frac{s+s'}{(s')^2} \cdot \sigma_{s'} \int_{-\infty}^{+\infty} \alpha'_{s'} dt \\
 &\stackrel{(3.6)}{=} g(s') + (s-s') \cdot \frac{s+s'}{2s'} \cdot (1-\bar{m}_{s'}) f(\bar{m}_{s'})
 \end{aligned}$$

whence

$$g(s) \leq g(s') + (s-s') \cdot \frac{s+s'}{2s'} \cdot (1-\bar{m}_{s'}) f(\bar{m}_{s'}) \quad \text{for all } s, s' \in (0, s_{\text{frac}}). \quad (3.10)$$

If $s' < s$, by dividing both sides in (3.10) by $(s-s')$ we get

$$\frac{g(s) - g(s')}{s - s'} \leq \frac{s+s'}{2s'} \cdot (1-\bar{m}_{s'}) f(\bar{m}_{s'}),$$

and letting $s \rightarrow (s')^+$ or $s' \rightarrow (s)^-$, using the continuity of $s \mapsto \bar{m}_s$, we obtain the following inequalities for the right and left derivatives of g :

$$g'_+(s') \leq (1-\bar{m}_{s'}) f(\bar{m}_{s'}), \quad g'_-(s) \leq (1-\bar{m}_s) f(\bar{m}_s).$$

By arguing similarly for $s' > s$ in (3.10), we obtain the opposite inequalities, so that (3.9) follows for all $s \in (0, s_{\text{frac}})$.

To conclude the proof, it only remains to show that g is differentiable also at s_{frac} and $g'(s_{\text{frac}}) = 0$. This is easily obtained since $\lim_{s \rightarrow s_{\text{frac}}^-} g'(s) = 0$ by the first part of the proof and $g \equiv 1$ for $s > s_{\text{frac}}$. \square

A condition for the finiteness of the threshold s_{frac} is given in [12, Proposition 6.3]. In the following proposition we improve that result by determining explicitly the value of s_{frac} in terms of the derivative of f at the origin.

Proposition 3.7. *Under the assumptions (f1)–(f6), the threshold s_{frac} defined in (3.1) is given by*

$$s_{\text{frac}} = \frac{\pi}{f'(0)} \quad (3.11)$$

(with $s_{\text{frac}} = +\infty$ if $f'(0) = 0$).

Proof. For any fixed $\delta \in (0, 1)$ we have by (3.7)

$$\begin{aligned} s_{\text{frac}} &= \lim_{s \rightarrow s_{\text{frac}}^-} 2(1 - \bar{m}_s)f(\bar{m}_s) \int_{\bar{m}_s}^1 \frac{dt}{f(t)[(1-t)^2 f^2(t) - (1 - \bar{m}_s)^2 f^2(\bar{m}_s)]^{1/2}} \\ &= \lim_{s \rightarrow s_{\text{frac}}^-} 2(1 - \bar{m}_s)f(\bar{m}_s) \int_{\bar{m}_s}^{\delta} \frac{dt}{f(t)[(1-t)^2 f^2(t) - (1 - \bar{m}_s)^2 f^2(\bar{m}_s)]^{1/2}}, \end{aligned}$$

where in the last equality we have used that the denominator in the integrand is uniformly far from 0 in $(\delta, 1)$ and that $\bar{m}_s \rightarrow 0$. Denoting by $\phi(t) = (1-t)f(t)$, which is strictly increasing by assumption (f3), we can write the previous identity as

$$\begin{aligned} s_{\text{frac}} &= \lim_{s \rightarrow s_{\text{frac}}^-} 2\phi(\bar{m}_s) \int_{\bar{m}_s}^{\delta} \frac{\phi'(t)}{\phi(t)\sqrt{\phi^2(t) - \phi^2(\bar{m}_s)}} \cdot \frac{(1-t)}{\phi'(t)} dt \\ &\leq \lim_{s \rightarrow s_{\text{frac}}^-} 2\phi(\bar{m}_s) \left(\sup_{t \in (\bar{m}_s, \delta)} \frac{(1-t)}{\phi'(t)} \right) \int_{\bar{m}_s}^{\delta} \frac{\phi'(t)}{\phi(t)\sqrt{\phi^2(t) - \phi^2(\bar{m}_s)}} dt \\ &= \lim_{s \rightarrow s_{\text{frac}}^-} 2 \left(\sup_{t \in (\bar{m}_s, \delta)} \frac{(1-t)}{\phi'(t)} \right) \arctan \left(\frac{\sqrt{\phi^2(\delta) - \phi^2(\bar{m}_s)}}{\phi(\bar{m}_s)} \right) = \pi \sup_{t \in (0, \delta)} \frac{(1-t)}{\phi'(t)} \end{aligned}$$

since $\bar{m}_s \rightarrow 0$ as $s \rightarrow s_{\text{frac}}^-$. Arguing similarly we find

$$\pi \inf_{t \in (0, \delta)} \frac{(1-t)}{\phi'(t)} \leq s_{\text{frac}} \leq \pi \sup_{t \in (0, \delta)} \frac{(1-t)}{\phi'(t)}.$$

The conclusion follows by letting $\delta \rightarrow 0^+$. \square

Remark 3.8. For the prototype examples in Remark 1.3 one has $s_{\text{frac}} = \frac{\pi}{q\sigma_c} \in \mathbb{R}$ for the functions f_q , and $s_{\text{frac}} = +\infty$ for the functions f^p .

We conclude this section by determining the asymptotic expansion of the cohesive energy density g at the origin.

Proposition 3.9. *Assume that f satisfies the assumptions (f1)–(f6), and assume further that*

$$(1-s)f(s) = \sigma_c - \ell(1-s)^q + o((1-s)^q) \quad \text{as } s \rightarrow 1^- \quad (3.12)$$

for some $\ell > 0$ and $q \in (0, 2]$, where $o(t)$ denotes any quantity such that $\lim_{t \rightarrow 0} \frac{o(t)}{t} = 0$. Then the function g defined in (1.9) satisfies, for $p := \frac{4+q}{4-q}$ and for some $\tilde{\ell} > 0$,

$$g(s) = \sigma_c s - \tilde{\ell} s^p + o(s^p) \quad \text{as } s \rightarrow 0^+. \quad (3.13)$$

Proof. The proof is a straightforward adaptation of [12, Proposition 8.5], which deals with the case $q = 1$. The details are left to the reader. \square

Remark 3.10. The function f_q for $q \in (0, 4)$ in Remark 1.3 satisfies the condition (3.12).

4. PRELIMINARY PROPERTIES OF CRITICAL POINTS

We assume along this section that $(u_\varepsilon, v_\varepsilon) \in (H^1(0, L))^2$ is a family of critical points of the energies \mathcal{F}_ε , *i.e.* they are weak solutions to the system of equations (1.17a)–(1.17d). We also suppose that the Dirichlet boundary condition satisfies (1.18), and that the equiboundedness of the energy (1.19) holds. We first remark that by (1.17b) there exist constants $c_\varepsilon \in \mathbb{R}$ such that

$$f_\varepsilon^2(v_\varepsilon)u'_\varepsilon = c_\varepsilon \quad \text{a.e. in } (0, L). \quad (4.1)$$

Notice that from (4.1) it follows that u'_ε has constant sign a.e. in $(0, L)$ and therefore, in view of the boundary conditions, it must be $c_\varepsilon \geq 0$ and u_ε monotone nondecreasing in $(0, L)$. Moreover it cannot be $c_\varepsilon = 0$, or otherwise the second term in (1.17a) would vanish and v_ε would be a weak solution to $-\varepsilon v''_\varepsilon + \frac{v_\varepsilon - 1}{4\varepsilon} = 0$, with $v_\varepsilon(0) = v_\varepsilon(L) = 1$; however, this would imply that $v_\varepsilon \equiv 1$ and in turn, by (4.1), $u'_\varepsilon = 0$ almost everywhere in $(0, L)$, which is not possible in view of the boundary conditions (1.17c). Therefore $c_\varepsilon > 0$ for all ε .

We also have

$$\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq \int_0^L f_\varepsilon^2(v_\varepsilon)|u'_\varepsilon|^2 dx = c_\varepsilon \int_0^L u'_\varepsilon dx = c_\varepsilon a_\varepsilon,$$

so that by (1.18) and (1.19) we obtain that $\sup_\varepsilon c_\varepsilon < +\infty$ and, up to subsequences,

$$c_\varepsilon \rightarrow c_0 \geq 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (4.2)$$

Similarly to [30, Lemma 3.1] and to [10, Lemma 2.3], we show that v_ε obeys a maximum principle.

Lemma 4.1. *We have that $0 \leq v_\varepsilon \leq 1$ in $[0, L]$.*

Proof. By testing (1.17a) with the function $\varphi_\varepsilon := \max\{-v_\varepsilon, 0\}$, which is admissible since $\varphi_\varepsilon \in H_0^1(0, L)$ in view of the boundary conditions (1.17d), we have

$$-\int_0^L \varepsilon(\varphi'_\varepsilon)^2 dx + \int_0^L f_\varepsilon(v_\varepsilon)f'_\varepsilon(v_\varepsilon)(u'_\varepsilon)^2 \varphi_\varepsilon dx - \int_0^L \frac{\varphi_\varepsilon + 1}{4\varepsilon} \varphi_\varepsilon dx = 0,$$

and since all the terms are nonpositive (recall that $f_\varepsilon(s) := \sqrt{\varepsilon}f'(0)s$ for $s < 0$) we deduce that $\int_0^L \frac{\varphi_\varepsilon + 1}{4\varepsilon} \varphi_\varepsilon dx = 0$, and therefore $v_\varepsilon \geq 0$. Similarly, by testing the equation with the function $\varphi_\varepsilon := \max\{0, v_\varepsilon - 1\} \in H_0^1(0, L)$ we prove that $v_\varepsilon \leq 1$. \square

We next show that the solutions to the equations (1.17a)–(1.17d) satisfy a conservation law, which can also be seen as a consequence of the vanishing of the first variation of the functional \mathcal{F}_ε with respect to *inner variations*.

Proposition 4.2. *There exist constants $d_\varepsilon \in \mathbb{R}$ such that*

$$\frac{(1 - v_\varepsilon)^2}{4\varepsilon} - f_\varepsilon^2(v_\varepsilon)(u'_\varepsilon)^2 - \varepsilon(v'_\varepsilon)^2 = d_\varepsilon \quad \text{in } (0, L), \quad (4.3)$$

with $\sup_\varepsilon |d_\varepsilon| < +\infty$ and, up to subsequences, $d_\varepsilon \rightarrow d_0$ as $\varepsilon \rightarrow 0$.

Furthermore, $u_\varepsilon, v_\varepsilon \in C^2(0, L)$ with $v_\varepsilon > 0$ in $[0, L]$, and the equations (1.17a)–(1.17d) hold in the classical sense.

Proof. We first remark that, as v_ε is a weak solution to (1.17a), we have $v_\varepsilon'' \in L^1(0, L)$ and therefore $v_\varepsilon \in C^1([0, L])$. By (4.1) we have $u_\varepsilon' = \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)}$ almost everywhere in the open set $\{v_\varepsilon > 0\}$, that is, u_ε' is (almost everywhere equal to) a C^1 -function in $\{v_\varepsilon > 0\}$. In particular u_ε is of class C^2 in $\{v_\varepsilon > 0\}$. In turn, by (1.17a) the same holds for v_ε , and equation (1.17a) holds in the classical sense in $\{v_\varepsilon > 0\}$.

We can thus differentiate the left-hand side of (4.3) in $\{v_\varepsilon > 0\}$:

$$\begin{aligned} & \frac{1}{2} \left(\frac{(1-v_\varepsilon)^2}{4\varepsilon} - f_\varepsilon^2(v_\varepsilon)(u_\varepsilon')^2 - \varepsilon(v_\varepsilon')^2 \right)' \\ &= v_\varepsilon' \left(-\varepsilon v_\varepsilon'' - f_\varepsilon(v_\varepsilon) f_\varepsilon'(v_\varepsilon) (u_\varepsilon')^2 + \frac{v_\varepsilon - 1}{4\varepsilon} \right) - f_\varepsilon^2(v_\varepsilon) u_\varepsilon' u_\varepsilon'' \\ &\stackrel{(1.17a)}{=} -2v_\varepsilon' f_\varepsilon(v_\varepsilon) f_\varepsilon'(v_\varepsilon) (u_\varepsilon')^2 - f_\varepsilon^2(v_\varepsilon) u_\varepsilon' u_\varepsilon'' \\ &= -u_\varepsilon' \left(f_\varepsilon^2(v_\varepsilon) u_\varepsilon' \right)' \stackrel{(1.17b)}{=} 0, \end{aligned}$$

hence (4.3) holds in $\{v_\varepsilon > 0\}$, for a constant d_ε possibly changing with the connected components of $\{v_\varepsilon > 0\}$.

We next show that $v_\varepsilon > 0$ everywhere in $[0, L]$. Consider any connected component (a, b) of the open set $\{v_\varepsilon > 0\}$. By combining (4.3) and (4.1) we have

$$\frac{(1-v_\varepsilon)^2}{4\varepsilon} - \frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon)} - \varepsilon(v_\varepsilon')^2 = d_\varepsilon \quad \text{in } (a, b). \quad (4.4)$$

Assume by contradiction that v_ε vanishes at one of the endpoints, say $v_\varepsilon(a) = 0$. The point a must be in the interior $(0, L)$ by (1.17d); since a is a minimum point of v_ε by Lemma 4.1 and v_ε is of class $C^1([0, L])$, we have $v_\varepsilon'(a) = 0$. We can pass to the limit in (4.4) as $x \rightarrow a$ from the interior of (a, b) :

$$\frac{1}{4\varepsilon} - d_\varepsilon = \lim_{x \rightarrow a} \frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon(x))}$$

and since $f_\varepsilon(v_\varepsilon(x)) \rightarrow f_\varepsilon(0) = 0$, we conclude that it must be $c_\varepsilon = 0$. However, we already observed that $c_\varepsilon > 0$ (see the discussion after (4.1)), which is a contradiction proving that $(a, b) = (0, L)$, and since v_ε cannot vanish at the endpoints we obtain $\{v_\varepsilon > 0\} = [0, L]$.

Finally, by integrating (4.3) on $(0, L)$ we also have

$$|d_\varepsilon| = \frac{1}{L} \int_0^L \left| \frac{(1-v_\varepsilon)^2}{4\varepsilon} - f_\varepsilon^2(v_\varepsilon)(u_\varepsilon')^2 - \varepsilon(v_\varepsilon')^2 \right| dx \leq \frac{1}{L} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon),$$

and therefore $\sup_\varepsilon |d_\varepsilon| < +\infty$ by (1.19). \square

Notice that, by using (4.1), we can rewrite (4.3) in the form

$$\frac{(1-v_\varepsilon)^2}{4\varepsilon} - \frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon)} - \varepsilon(v_\varepsilon')^2 = d_\varepsilon \quad \text{in } (0, L). \quad (4.5)$$

Similarly, we can rewrite (1.17a) as an equation for the function v_ε alone:

$$-\varepsilon v_\varepsilon'' + \frac{c_\varepsilon^2 f_\varepsilon'(v_\varepsilon)}{f_\varepsilon^3(v_\varepsilon)} + \frac{v_\varepsilon - 1}{4\varepsilon} = 0 \quad \text{in } (0, L). \quad (4.6)$$

From this equation we can deduce the symmetry properties of the function v_ε , similarly to [30, Lemma 4.1 and Proposition 4.2] and to [10, Proposition 2.1].

Lemma 4.3. *The graph of v_ε in $[0, L]$ is a symmetric “well”: it is symmetric with respect to the point $\frac{L}{2}$, which is a global minimum, and v_ε is decreasing in $(0, \frac{L}{2})$.*

Proof. By (4.6), v_ε is a solution to an equation of the form $v_\varepsilon'' = h_\varepsilon(v_\varepsilon)$, where the function h_ε is Lipschitz continuous in $(0, +\infty)$ by definition (1.8) of f_ε and by assumption (f1). Notice that v_ε takes values in $(0, 1]$ by Lemma 4.1, and therefore Cauchy-Lipschitz theorem guarantees uniqueness.

If v_ε is not identically equal to 1, by Rolle’s Theorem there exists at least one critical point in $(0, L)$. Given any critical point $x_0 \in (0, L)$ of v_ε , we can symmetrize the graph of v_ε about the vertical line through x_0 : more precisely, if $x_0 \in (0, \frac{L}{2}]$ then we define $\tilde{v}_\varepsilon : (0, 2x_0) \rightarrow \mathbb{R}$ by $\tilde{v}_\varepsilon(x) = v_\varepsilon(x)$ for $x \in (0, x_0)$, $\tilde{v}_\varepsilon(x) = v_\varepsilon(2x_0 - x)$ for $x \in (x_0, 2x_0)$. Then \tilde{v}_ε is also a solution of $\tilde{v}_\varepsilon'' = h_\varepsilon(\tilde{v}_\varepsilon)$ in $(0, 2x_0)$, with $\tilde{v}_\varepsilon(x_0) = v_\varepsilon(x_0)$, $\tilde{v}_\varepsilon'(x_0) = v_\varepsilon'(x_0) = 0$, and Cauchy-Lipschitz theorem yields that $v_\varepsilon = \tilde{v}_\varepsilon$ in $(x_0, 2x_0)$. In particular the critical point x_0 is either a maximum or a minimum point. A symmetric argument can be repeated in the case of a critical point in the interval $(\frac{L}{2}, L)$.

Therefore the graph of v_ε is symmetric with respect to all the vertical lines passing through its critical points, which are either absolute maximum or absolute minimum points. If there is an interior maximum point at $x_0 \in (0, L)$, since $v_\varepsilon(0) = v_\varepsilon(L) = 1$ it must be $v_\varepsilon(x_0) = 1$, $v_\varepsilon'(x_0) = 0$. Then by uniqueness we conclude that $v_\varepsilon \equiv 1$, since the constant function 1 is also a solution of (4.6). Hence, if v_ε is not identically equal to 1, there are no interior maximum points and therefore there is a unique interior critical (minimum) point, located at $\frac{L}{2}$. The symmetric structure described in the statement follows. \square

Remark 4.4. The symmetry property of v_ε proved in the previous lemma is in accordance with the result in [10] for the Ambrosio-Tortorelli functional, where it is shown that, imposing the Dirichlet boundary conditions on v_ε , a much stronger symmetry property is obtained than in the Neumann case considered in [30], namely that v_ε has a unique critical point located at the midpoint $\frac{L}{2}$. We expect that, also in our setting, imposing Neumann conditions $v_\varepsilon'(0) = v_\varepsilon'(L) = 0$ we would obtain a weaker symmetry property, namely that there exists $n_\varepsilon \in \mathbb{N}$ such that the graph of v_ε in $(0, L)$ is made of n_ε repeated identical subgraphs, each of which is a symmetric “well” (with a unique interior critical point, which is a global minimum, and two maxima at the endpoints), or a symmetric “bell” (with a unique interior critical point, which is a global maximum, and two minima at the endpoints).

We conclude this section by collecting in the following lemma the compactness properties of the family $(u_\varepsilon, v_\varepsilon)$, together with a uniform bound of v_ε' .

Lemma 4.5. *We have that $v_\varepsilon \rightarrow 1$ in $L^1(0, L)$ and, up to extracting a subsequence $\varepsilon_k \rightarrow 0$, $u_\varepsilon \rightarrow u$ in $L^1(0, L)$ for some $u \in \text{BV}(0, L)$ with $|Du|(0, L) \leq a$. Moreover $\varepsilon \|v_\varepsilon'\|_\infty \leq \frac{1}{2}$ for all ε sufficiently small.*

Proof. In view of the bound (1.19) we have that $\sup_\varepsilon \frac{1}{\varepsilon} \int_0^L (1 - v_\varepsilon)^2 dx < +\infty$, which immediately implies the convergence of v_ε . Since u_ε is monotone increasing by (4.1), we have $|Du_\varepsilon|(0, L) = a_\varepsilon$, $\|u_\varepsilon\|_{L^\infty(0, L)} = a_\varepsilon$. Hence $(u_\varepsilon)_\varepsilon$ is bounded in BV and the second convergence follows by the compact embedding of BV into L^1 . By semicontinuity of the total variation $|Du|(0, L) \leq \liminf_\varepsilon |Du_\varepsilon|(0, L) = \liminf_\varepsilon a_\varepsilon = a$. Finally by (4.5)

$$\varepsilon^2 (v_\varepsilon')^2 \leq \frac{(1 - v_\varepsilon)^2}{4} + \varepsilon |d_\varepsilon| \leq \frac{1}{4} + \varepsilon \sup_\varepsilon |d_\varepsilon|,$$

with $\sup_\varepsilon |d_\varepsilon| < +\infty$ by Proposition 4.2. \square

As a consequence of Lemma 4.5 we remark for later use that the constant c_0 (see (4.2)) satisfies the bound

$$c_0 = \frac{1}{L} \int_0^L \lim_{\varepsilon \rightarrow 0} c_\varepsilon \, dx \stackrel{(4.1)}{\leq} \lim_{\varepsilon \rightarrow 0} \frac{1}{L} \int_0^L f_\varepsilon^2(v_\varepsilon) u'_\varepsilon \, dx \leq \lim_{\varepsilon \rightarrow 0} \frac{1}{L} \int_0^L u'_\varepsilon \, dx = \frac{a}{L}. \quad (4.7)$$

5. PROOF OF THE CONVERGENCE OF CRITICAL POINTS

This section is entirely devoted to the proof of Theorem 1.7. We assume along all this section that $(u_\varepsilon, v_\varepsilon)$ is a family of critical points of \mathcal{F}_ε satisfying the assumptions of Theorem 1.7. We recall that $(u_\varepsilon, v_\varepsilon)$ enjoys the regularity properties discussed in the previous section and that $v_\varepsilon(x) \in (0, 1]$, see in particular Lemma 4.1 and Proposition 4.2.

We denote by

$$m_\varepsilon := \min_{[0, L]} v_\varepsilon \in (0, 1]. \quad (5.1)$$

In view of the symmetry properties of the critical points observed in Lemma 4.3, we have that v_ε has a *single-well shape*, that is, its global minimum m_ε is achieved at the midpoint $\frac{L}{2}$, the graph of v_ε is symmetric with respect to $\frac{L}{2}$, and v_ε is decreasing in $(0, \frac{L}{2})$ and increasing in $(\frac{L}{2}, L)$, achieving its maximum at the endpoints $v_\varepsilon(0) = v_\varepsilon(L) = 1$.

We further assume that we have extracted a subsequence (not relabeled) such that $u_\varepsilon \rightarrow u$ and $v_\varepsilon \rightarrow 1$ in $L^1(0, L)$ as in Lemma 4.5, $c_\varepsilon \rightarrow c_0$ (see (4.2)), $d_\varepsilon \rightarrow d_0$ (see Proposition 4.2), and also

$$m_\varepsilon \rightarrow m_0 \in [0, 1] \quad \text{as } \varepsilon \rightarrow 0. \quad (5.2)$$

For later use it is convenient to introduce the discrepancy

$$\xi_\varepsilon(x) := \frac{(1 - v_\varepsilon(x))^2}{4\varepsilon} - \varepsilon(v'_\varepsilon(x))^2 \stackrel{(4.5)}{=} \frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon(x))} + d_\varepsilon. \quad (5.3)$$

By the second expression of ξ_ε in (5.3) and monotonicity of v_ε , the minimum of the function ξ_ε is attained at the maximum point of v_ε , that is $\min \xi_\varepsilon = \xi_\varepsilon(0) = -\varepsilon(v'_\varepsilon(0))^2 \leq 0$, and similarly the maximum of ξ_ε is attained at the midpoint, that is $\max \xi_\varepsilon = \xi_\varepsilon(\frac{L}{2}) = \frac{(1 - m_\varepsilon)^2}{4\varepsilon} \geq 0$. Hence there exists $y_\varepsilon \in [0, \frac{L}{2}]$ such that $\xi_\varepsilon(y_\varepsilon) = 0$, $\xi_\varepsilon \leq 0$ in $[0, y_\varepsilon]$ and $\xi_\varepsilon \geq 0$ in $[y_\varepsilon, \frac{L}{2}]$. Up to subsequences we can assume $y_\varepsilon \rightarrow y_0 \in [0, \frac{L}{2}]$. Notice that by evaluating (5.3) at the point y_ε we find

$$d_\varepsilon = -\frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon(y_\varepsilon))} < 0. \quad (5.4)$$

In the following lemma we show an explicit relation between the limit values c_0 and m_0 .

Lemma 5.1. *Assume that $f_\varepsilon(m_\varepsilon) \rightarrow 0$. Then*

$$c_0 = \frac{1}{2}(1 - m_0)f(m_0), \quad (5.5)$$

where c_0 and m_0 are the limits in (4.2) and (5.2) respectively, and the right-hand side of (5.5) must be interpreted as $\frac{\sigma_c}{2}$ if $m_0 = 1$, in view of (f2).

Proof. By evaluating (4.5) at a minimum point of v_ε we have

$$\frac{(1 - m_\varepsilon)^2}{4\varepsilon} - \frac{c_\varepsilon^2}{f_\varepsilon^2(m_\varepsilon)} = d_\varepsilon. \quad (5.6)$$

Since by assumption $f_\varepsilon(m_\varepsilon) \rightarrow 0$ and recalling the definition (1.8) of f_ε , it must be $f_\varepsilon(m_\varepsilon) = \sqrt{\varepsilon}f(m_\varepsilon)$ for ε small. Then

$$c_0^2 = \lim_{\varepsilon \rightarrow 0} c_\varepsilon^2 = \lim_{\varepsilon \rightarrow 0} \frac{1}{4\varepsilon} (1 - m_\varepsilon)^2 f_\varepsilon^2(m_\varepsilon) - d_\varepsilon f_\varepsilon^2(m_\varepsilon) = \frac{1}{4} (1 - m_0)^2 f^2(m_0),$$

where we used the uniform bound on d_ε in Proposition 4.2. \square

In the following lemma, which is a consequence of the qualitative study of the equation (4.6) for v_ε contained in Section 2, we deduce some general properties of the sequence $(u_\varepsilon, v_\varepsilon)$.

Lemma 5.2. *Assume that $m_\varepsilon < s_\varepsilon$, where s_ε is as in (1.8). Let*

$$A_\varepsilon := \{x \in (0, L) : v_\varepsilon(x) < s_\varepsilon\}.$$

Then $A_\varepsilon = (\frac{L}{2} - x_\varepsilon, \frac{L}{2} + x_\varepsilon)$ with $\lim_{\varepsilon \rightarrow 0} x_\varepsilon = 0$. If $m_0 < 1$, then $\lim_{\varepsilon \rightarrow 0} \frac{x_\varepsilon}{\varepsilon} = +\infty$.

Finally, $u \in \text{SBV}(0, L)$ with $J_u \subset \{\frac{L}{2}\}$ and $u' = c_0$ almost everywhere in $(0, L)$.

Proof. Since v_ε has a single-well shape, $v_\varepsilon(\frac{L}{2}) = m_\varepsilon < s_\varepsilon$, and $v_\varepsilon = 1$ at the endpoints of $(0, L)$, we have $A_\varepsilon = (\frac{L}{2} - x_\varepsilon, \frac{L}{2} + x_\varepsilon)$ for some $x_\varepsilon \in (0, \frac{L}{2})$. Notice that for $x \in A_\varepsilon$ we have $f_\varepsilon(v_\varepsilon(x)) = \sqrt{\varepsilon}f(v_\varepsilon(x))$. The rescaled function $\tilde{v}_\varepsilon(t) = v_\varepsilon(\frac{L}{2} + \varepsilon t)$ satisfies

$$\begin{aligned} \tilde{v}_\varepsilon'' &= \frac{1 - \tilde{v}_\varepsilon}{4} \left[\frac{(2c_\varepsilon)^2 f'(\tilde{v}_\varepsilon)}{(1 - \tilde{v}_\varepsilon) f^3(\tilde{v}_\varepsilon)} - 1 \right] & \text{for } t \in \left(-\frac{x_\varepsilon}{\varepsilon}, \frac{x_\varepsilon}{\varepsilon} \right) =: (-t_\varepsilon, t_\varepsilon), \\ \tilde{v}_\varepsilon(0) &= m_\varepsilon, \quad \tilde{v}_\varepsilon'(0) = 0, \end{aligned} \quad (5.7)$$

that is, \tilde{v}_ε is a solution in $(-t_\varepsilon, t_\varepsilon)$ of the Cauchy problem (2.1a)–(2.1c) studied in Section 2, for the values of the parameters $m := m_\varepsilon \in (0, 1)$ and $\alpha := c_\varepsilon > 0$.

We first consider the case $c_\varepsilon < \frac{\sigma_c}{2}$, which is the assumption in Theorem 2.2 and Proposition 2.5. Notice that necessarily $m_\varepsilon < z_{c_\varepsilon}$, where $z_{c_\varepsilon} \in (0, 1)$ is defined by the equation (2.5): indeed, if it were $m_\varepsilon > z_{c_\varepsilon}$ then by the qualitative analysis of the ODE (5.7) (see in particular Remark 2.4) the solution \tilde{v}_ε would have a local maximum at the origin, which is incompatible with the single-well structure; if $m_\varepsilon = z_{c_\varepsilon}$ then \tilde{v}_ε would be constant, which is again not possible.

Hence in the case $c_\varepsilon < \frac{\sigma_c}{2}$ we have $m_\varepsilon \in (0, z_{c_\varepsilon})$ and we are in position to apply Proposition 2.5 with $\eta := s_\varepsilon$ in order to estimate the time t_ε such that $\tilde{v}_\varepsilon(t_\varepsilon) = s_\varepsilon$. Notice that, in the case $(1 - m_\varepsilon)f(m_\varepsilon) > 2c_\varepsilon$, the additional assumption $\eta < M$ (where M is the maximum of the solution and then depends on ε) is certainly satisfied, or else the function \tilde{v}_ε would reach a maximum point before t_ε and then decrease, which is incompatible with its single-well shape. Hence t_ε obeys the bounds (2.8)–(2.9). In particular, if $m_\varepsilon \rightarrow m_0 \in [0, 1)$ then $c_\varepsilon \rightarrow c_0 \in [0, \frac{\sigma_c}{2})$ by Lemma 5.1 and therefore the estimate (2.8) holds with a constant which is uniformly bounded with respect to ε , namely

$$t_\varepsilon \leq \frac{C_0}{\sqrt{1 - s_\varepsilon}} \quad \text{if } m_0 \in [0, 1). \quad (5.8)$$

If, instead, $m_\varepsilon \rightarrow m_0 = 1$, then we can apply (2.9) to deduce

$$t_\varepsilon \leq \frac{\omega(1 - m_\varepsilon)}{(1 - s_\varepsilon)^2} \quad \text{if } m_0 = 1, \quad (5.9)$$

where $\omega(\cdot)$ is a modulus of continuity independent of ε . Hence if $m_0 \in [0, 1)$ we find

$$x_\varepsilon = \varepsilon t_\varepsilon \leq \frac{C_0 \varepsilon}{\sqrt{1-s_\varepsilon}} \stackrel{(1.7)}{\sim} \frac{C_0 \varepsilon}{\varepsilon^{\frac{1}{4}} \sqrt{\sigma_c}} \longrightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,$$

whereas if $m_0 \rightarrow 1$

$$x_\varepsilon = \varepsilon t_\varepsilon \leq \frac{\varepsilon \omega(1-m_\varepsilon)}{(1-s_\varepsilon)^2} \stackrel{(1.7)}{\sim} \frac{\omega(1-m_\varepsilon)}{\sigma_c^2} \longrightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,$$

proving that $x_\varepsilon \rightarrow 0$ in any case.

Consider now the case $c_\varepsilon \geq \frac{\sigma_c}{2}$. Notice that in this case it must be $m_\varepsilon \rightarrow 1$, or otherwise by Lemma 5.1 $c_\varepsilon \rightarrow c_0 = \frac{1}{2}(1-m_0)f(m_0) < \frac{\sigma_c}{2}$, which is not possible. We can then apply Proposition 2.6 to deduce that the estimate (5.9) continues to hold, and therefore $x_\varepsilon \rightarrow 0$, as before.

Assume now that $m_0 < 1$ and let us show that $\frac{x_\varepsilon}{\varepsilon} \rightarrow +\infty$. As already observed it must be $c_\varepsilon < \frac{\sigma_c}{2}$, and we can apply again Proposition 2.5 with $\eta = s_\varepsilon$ to deduce by (2.10) that

$$\frac{x_\varepsilon}{\varepsilon} = t_\varepsilon \geq \log\left(\frac{1-z_{c_\varepsilon}+k_\varepsilon}{1-s_\varepsilon+k_\varepsilon}\right) \quad k_\varepsilon := \left|1 - \left(\frac{2c_\varepsilon}{(1-m_\varepsilon)f(m_\varepsilon)}\right)^2\right|^{\frac{1}{2}}. \quad (5.10)$$

By (5.5) we have $k_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. Moreover $z_{c_\varepsilon} \rightarrow z_{c_0}$, where $c_0 = \frac{1}{2}(1-m_0)f(m_0)$ by (5.5). Since $m_0 < 1$, we then have $c_0 \in [0, \frac{\sigma_c}{2})$ and in turn $z_{c_0} \in [0, 1)$ by the properties of \bar{f} in Proposition 2.1. Therefore by passing to the limit as $\varepsilon \rightarrow 0$ in (5.10) we obtain $\frac{x_\varepsilon}{\varepsilon} \rightarrow +\infty$ as $\varepsilon \rightarrow 0$, which completes the proof of the first part of the statement.

For every fixed $\delta > 0$ it holds $A_\varepsilon \subset (\frac{L}{2} - \delta, \frac{L}{2} + \delta)$ for all ε sufficiently small. In particular for all $x \in (\frac{L}{2} - \delta, \frac{L}{2} + \delta)^c := (0, L) \setminus (\frac{L}{2} - \delta, \frac{L}{2} + \delta)$ we have $v_\varepsilon \geq s_\varepsilon$ and, in turn, $f_\varepsilon(v_\varepsilon(x)) \geq f_\varepsilon(s_\varepsilon) \rightarrow 1$, that is, $f_\varepsilon(v_\varepsilon)$ converges uniformly to 1 on compact sets not containing $\frac{L}{2}$. Hence by (4.1)

$$u'_\varepsilon = \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} \longrightarrow c_0 \quad \text{uniformly on } (\frac{L}{2} - \delta, \frac{L}{2} + \delta)^c, \text{ for all } \delta > 0. \quad (5.11)$$

We also have

$$\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq f_\varepsilon^2(s_\varepsilon) \int_{(\frac{L}{2}-\delta, \frac{L}{2}+\delta)^c} |u'_\varepsilon|^2 dx,$$

so that from the uniform bound on the energies (1.19) and the convergence $f_\varepsilon(s_\varepsilon) \rightarrow 1$, we have that u_ε is uniformly bounded in $H^1((\frac{L}{2} - \delta, \frac{L}{2} + \delta)^c)$ for all $\delta > 0$. We can conclude that $u \in H^1((\frac{L}{2} - \delta, \frac{L}{2} + \delta)^c)$ for all $\delta > 0$. The properties in the statement are then immediate consequences of the previous facts. \square

5.1. Case I: pre-fractured regime. We show that if $m_0 \in (0, 1)$ then the limit function u is a piecewise affine critical point of the cohesive energy (1.1) with a single jump at $\frac{L}{2}$ and constant slope. This is summarized in the following proposition, which is the main result of this subsection.

Proposition 5.3. *Assume that $m_0 \in (0, 1)$. Then $u \in \text{SBV}(0, L)$, $u' = c_0 \in (0, \frac{\sigma_c}{2})$ almost everywhere in $(0, L)$, $J_u = \{\frac{L}{2}\}$, $[u](\frac{L}{2}) \in (0, s_{\text{frac}})$, and*

$$g'([u](\frac{L}{2})) = 2c_0. \quad (5.12)$$

Moreover u attains the limit boundary conditions, that is, $|Du|(0, L) = c_0 L + [u](\frac{L}{2}) = a$. Finally, the convergence of the energies (1.20) holds.

Notice that, under the assumptions of the proposition, we can apply Lemma 5.2 since $m_\varepsilon < s_\varepsilon$ for ε small enough, as $m_0 \in (0, 1)$ and $s_\varepsilon \rightarrow 1$. We premise a lemma to the proof of the proposition.

Lemma 5.4. *Under the assumptions of Proposition 5.3, let A_ε be as in Lemma 5.2. Then we have as $\varepsilon \rightarrow 0$*

$$\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon; A_\varepsilon^c) := \int_{(0,L) \setminus A_\varepsilon} \left(f_\varepsilon^2(v_\varepsilon) |u'_\varepsilon|^2 + \frac{(1-v_\varepsilon)^2}{4\varepsilon} + \varepsilon |v'_\varepsilon|^2 \right) dx \rightarrow \int_0^L |u'|^2 dx.$$

Moreover $c_0^2 + d_0 = 0$.

Proof. By multiplying (4.6) by the function $v_\varepsilon - 1$ and integrating in $(0, \frac{L}{2} - x_\varepsilon)$ we have, after integration by parts,

$$\begin{aligned} \int_0^{\frac{L}{2}-x_\varepsilon} \left(\varepsilon |v'_\varepsilon|^2 + \frac{(1-v_\varepsilon)^2}{4\varepsilon} \right) dx &= \varepsilon (s_\varepsilon - 1) v'_\varepsilon \left(\frac{L}{2} - x_\varepsilon \right) + \int_0^{\frac{L}{2}-x_\varepsilon} \frac{c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{f_\varepsilon^3(v_\varepsilon)} (1-v_\varepsilon) dx \\ &\leq \varepsilon (1-s_\varepsilon) \|v'_\varepsilon\|_\infty + c_\varepsilon^2 \int_0^{\frac{L}{2}-x_\varepsilon} \frac{\psi'_\varepsilon(v_\varepsilon)}{(1-v_\varepsilon)\psi_\varepsilon^3(v_\varepsilon)} (1-v_\varepsilon)^2 dx \end{aligned}$$

where in the second equality we used the fact that $f_\varepsilon(v_\varepsilon) = \psi_\varepsilon(v_\varepsilon)$ in $(0, \frac{L}{2} - x_\varepsilon)$ since $v_\varepsilon \geq s_\varepsilon$ in this interval. In view of the monotonicity properties of ψ_ε in assumptions ($\psi 1$) and ($\psi 3$), the previous estimate yields

$$\int_0^{\frac{L}{2}-x_\varepsilon} \left(\varepsilon |v'_\varepsilon|^2 + \frac{(1-v_\varepsilon)^2}{4\varepsilon} \right) dx \leq \varepsilon (1-s_\varepsilon) \|v'_\varepsilon\|_\infty + \frac{c_\varepsilon^2 \psi'_\varepsilon(s_\varepsilon)}{(1-s_\varepsilon)\psi_\varepsilon^3(s_\varepsilon)} \int_0^{\frac{L}{2}-x_\varepsilon} (1-v_\varepsilon)^2 dx.$$

Now recalling (f2), (1.7), and that $c_0 < \frac{\sigma_c}{2}$ (by (5.5) and the assumption $m_0 \in (0, 1)$)

$$\lim_{\varepsilon \rightarrow 0} \frac{4\varepsilon c_\varepsilon^2 \psi'_\varepsilon(s_\varepsilon)}{(1-s_\varepsilon)\psi_\varepsilon^3(s_\varepsilon)} = \lim_{\varepsilon \rightarrow 0} \frac{4\varepsilon \sqrt{\varepsilon} c_\varepsilon^2}{(1-s_\varepsilon)^3} \cdot \frac{(1-s_\varepsilon)^2 f'(s_\varepsilon)}{f_\varepsilon^3(s_\varepsilon)} = \frac{4c_0^2}{\sigma_c^2} < 1,$$

hence we have that there exists a constant $C > 0$, independent of ε , such that for all ε sufficiently small

$$\frac{1}{4\varepsilon} - \frac{c_\varepsilon^2 \psi'_\varepsilon(s_\varepsilon)}{(1-s_\varepsilon)\psi_\varepsilon^3(s_\varepsilon)} \geq \frac{C}{4\varepsilon}.$$

In turn we find

$$\int_0^{\frac{L}{2}-x_\varepsilon} \left(\varepsilon |v'_\varepsilon|^2 + \frac{C(1-v_\varepsilon)^2}{4\varepsilon} \right) dx \leq \varepsilon (1-s_\varepsilon) \|v'_\varepsilon\|_\infty \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

in view of the bound $\varepsilon \|v'_\varepsilon\|_\infty \leq \frac{1}{2}$ for ε small in Lemma 4.5. By symmetry of v_ε with respect to the midpoint $\frac{L}{2}$ we can conclude that

$$\lim_{\varepsilon \rightarrow 0} \int_{(0,L) \setminus A_\varepsilon} \left(\varepsilon |v'_\varepsilon|^2 + \frac{(1-v_\varepsilon)^2}{4\varepsilon} \right) dx = 0. \quad (5.13)$$

In turn using (5.11)

$$\lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon; A_\varepsilon^c) = \lim_{\varepsilon \rightarrow 0} \int_{(0,L) \setminus A_\varepsilon} f_\varepsilon^2(v_\varepsilon) |u'_\varepsilon|^2 dx = c_0^2 L = \int_0^L |u'|^2 dx$$

proving the first part of the statement.

To show that $c_0^2 + d_0 = 0$, fix any $x_0 \in (0, \frac{L}{2})$ and notice that $(0, x_0) \subset A_\varepsilon^c$ for ε small, since $x_\varepsilon \rightarrow 0$ by Lemma 5.2. Then, using the uniform convergence of u'_ε to c_0 (see (5.11)) we find

$$|c_0^2 + d_0|x_0 = \lim_{\varepsilon \rightarrow 0} \int_0^{x_0} |c_\varepsilon u'_\varepsilon + d_\varepsilon| dx \stackrel{(4.3)}{=} \lim_{\varepsilon \rightarrow 0} \int_0^{x_0} \left| \frac{(1 - v_\varepsilon)^2}{4\varepsilon} - \varepsilon (v'_\varepsilon)^2 \right| dx \stackrel{(5.13)}{=} 0,$$

which completes the proof of the lemma. \square

We are now ready to give the proof of Proposition 5.3.

Proof of Proposition 5.3. We first prove that the limit function u satisfies $|Du|(0, L) = a$. Fix $\delta > 0$ such that $u_\varepsilon(\frac{L}{2} \pm \delta) \rightarrow u(\frac{L}{2} \pm \delta)$ and denote by $I_\delta := (0, L) \setminus (\frac{L}{2} - \delta, \frac{L}{2} + \delta)$. We have $I_\delta \subset A_\varepsilon^c$ for ε small enough, since $x_\varepsilon \rightarrow 0$ by Lemma 5.2; hence by (5.13)

$$\begin{aligned} 0 &= \lim_{\varepsilon \rightarrow 0} \int_{I_\delta} \left(\frac{(1 - v_\varepsilon)^2}{4\varepsilon} - \varepsilon |v'_\varepsilon|^2 \right) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{I_\delta} (c_\varepsilon u'_\varepsilon + d_\varepsilon) dx \\ &= c_0 \lim_{\varepsilon \rightarrow 0} (u_\varepsilon(L) - u_\varepsilon(\frac{L}{2} + \delta) + u_\varepsilon(\frac{L}{2} - \delta) - u_\varepsilon(0)) + d_0(L - 2\delta) \\ &= c_0(a - u(\frac{L}{2} + \delta) + u(\frac{L}{2} - \delta)) - c_0^2(L - 2\delta), \end{aligned}$$

where the second equality follows by (4.3) and (4.1), and the last one by Lemma 5.4. Hence by letting $\delta \rightarrow 0$ we find $a = c_0L + [u](\frac{L}{2}) = \int_0^L u' dx + [u](\frac{L}{2}) = |Du|(0, L)$.

To conclude the proof, it remains to show the criticality identity (5.12) and the convergence of the energies (1.20). We consider a blow-up of the functions u_ε and v_ε around the midpoint: let $\tilde{u}_\varepsilon(t) := u_\varepsilon(\frac{L}{2} + \varepsilon t)$, $\tilde{v}_\varepsilon(t) := v_\varepsilon(\frac{L}{2} + \varepsilon t)$ for $t \in (-\frac{L}{2\varepsilon}, \frac{L}{2\varepsilon})$. The idea of the proof is to show that the pair $(\tilde{u}_\varepsilon, \tilde{v}_\varepsilon)$ converges to an optimal pair $(\alpha_{s_0}, \beta_{s_0})$ for the minimum problem (1.9) which defines $g(s_0)$, with $s_0 = [u](\frac{L}{2})$. We refer to Section 3, and in particular to Proposition 3.2 and Proposition 3.3, for the existence and the main properties of optimal pairs for g .

We first remark that for all $T > 0$ and all ε sufficiently small

$$\int_{-T}^T \left(\frac{(1 - \tilde{v}_\varepsilon)^2}{4} + |\tilde{v}'_\varepsilon|^2 \right) dt = \int_{\frac{L}{2} - \varepsilon T}^{\frac{L}{2} + \varepsilon T} \left(\frac{(1 - v_\varepsilon)^2}{4\varepsilon} + \varepsilon |v'_\varepsilon|^2 \right) dx \leq C, \quad (5.14)$$

for a constant $C > 0$ independent of ε and T , by the uniform bound (1.19). Therefore, up to extracting a subsequence, we have that $\tilde{v}_\varepsilon \rightharpoonup \tilde{v}$ weakly in $H^1_{\text{loc}}(\mathbb{R})$, for some function \tilde{v} with $1 - \tilde{v} \in H^1(\mathbb{R})$, and the convergence is also uniform on compact sets.

Furthermore, the function \tilde{v}_ε solves the initial value problem (5.7) in $(-\frac{x_\varepsilon}{\varepsilon}, \frac{x_\varepsilon}{\varepsilon})$, where $t_\varepsilon := \frac{x_\varepsilon}{\varepsilon} \rightarrow +\infty$ as $\varepsilon \rightarrow 0$ by Lemma 5.2. For every $T > 0$ and every test function $\varphi \in C_c^\infty(-T, T)$, since $(-T, T) \subset (-\frac{x_\varepsilon}{\varepsilon}, \frac{x_\varepsilon}{\varepsilon})$ for all ε small enough we can pass to the limit in the weak formulation of (5.7):

$$0 = \int_{-T}^T \left(\tilde{v}'_\varepsilon \varphi' + \frac{c_\varepsilon^2 f'(\tilde{v}_\varepsilon)}{f^3(\tilde{v}_\varepsilon)} \varphi + \frac{\tilde{v}_\varepsilon - 1}{4} \varphi \right) dt \longrightarrow \int_{-T}^T \left(\tilde{v}' \varphi' + \frac{c_0^2 f'(\tilde{v})}{f^3(\tilde{v})} \varphi + \frac{\tilde{v} - 1}{4} \varphi \right) dt$$

where the convergence is justified since $\tilde{v}_\varepsilon \rightharpoonup \tilde{v}$ weakly in $H^1(-T, T)$ and uniformly on $[-T, T]$, and $\tilde{v} \geq \tilde{v}(0) = m_0 > 0$. In conclusion, we obtained that the limit function \tilde{v} is a weak solution

to the equation

$$\begin{aligned}\tilde{v}'' &= \frac{c_0^2 f'(\tilde{v})}{f^3(\tilde{v})} + \frac{\tilde{v} - 1}{4} && \text{in } \mathbb{R}, \\ \tilde{v}(0) &= m_0, \quad \tilde{v}'(0) = 0.\end{aligned}\tag{5.15}$$

Notice that the right-hand side of (5.15) is a continuous function, and therefore $\tilde{v} \in C^2(\mathbb{R})$ and the equation holds in the classical sense; moreover $\tilde{v}'(0) = 0$ since the origin is a minimum point of \tilde{v} .

Consider now the map $s \mapsto \bar{m}_s$ defined in Proposition 3.3, which associates to every $s \in [0, +\infty)$ the minimum value of the optimal profile β_s for $g(s)$. This map is a continuous bijection between $(0, s_{\text{frac}})$ and $(0, 1)$: in particular, since $m_0 \in (0, 1)$, we have that there exists $s_0 \in (0, s_{\text{frac}})$ such that $m_0 = \bar{m}_{s_0}$. By (3.2) and (3.3) the optimal profile β_{s_0} for $g(s_0)$ solves

$$\begin{aligned}\beta_{s_0}'' &= \frac{\sigma_{s_0}^2 f'(\beta_{s_0})}{f^3(\beta_{s_0})} + \frac{\beta_{s_0} - 1}{4} && \text{in } \mathbb{R}, \\ \beta_{s_0}(0) &= \bar{m}_{s_0}, \quad \beta_{s_0}'(0) = 0,\end{aligned}\tag{5.16}$$

where by (3.6) and (5.5) the constant σ_{s_0} is given by

$$\sigma_{s_0} = \frac{1}{2}(1 - \bar{m}_{s_0})f(\bar{m}_{s_0}) = \frac{1}{2}(1 - m_0)f(m_0) = c_0.\tag{5.17}$$

Therefore by comparing (5.15) and (5.16) we conclude, by uniqueness, that it must be

$$\tilde{v} = \beta_{s_0}.\tag{5.18}$$

In order to obtain the criticality condition (5.12), it is now sufficient to show that $s_0 = [u](\frac{L}{2})$: indeed in this case we would have by Proposition 3.6

$$g'([u](\frac{L}{2})) = g'(s_0) = (1 - \bar{m}_{s_0})f(\bar{m}_{s_0}) \stackrel{(5.17)}{=} 2c_0.$$

Therefore we now prove that $s_0 = [u](\frac{L}{2})$.

By using the properties of the optimal pair $(\alpha_{s_0}, \beta_{s_0})$ in Proposition 3.2, we have

$$\begin{aligned}s_0 &= \int_{\mathbb{R}} \alpha'_{s_0} dt \stackrel{(3.3)}{=} \int_{\mathbb{R}} \frac{\sigma_{s_0}}{f^2(\beta_{s_0})} dt \stackrel{(5.18)}{=} \int_{\mathbb{R}} \frac{c_0}{f^2(\tilde{v})} dt \\ &= \sup_{T>0} \int_{-T}^T \frac{c_0}{f^2(\tilde{v})} dt = \sup_{T>0} \lim_{\varepsilon \rightarrow 0} \int_{-T}^T \frac{c_\varepsilon}{f^2(\tilde{v}_\varepsilon)} dt = \sup_{T>0} \lim_{\varepsilon \rightarrow 0} \int_{\frac{L}{2}-\varepsilon T}^{\frac{L}{2}+\varepsilon T} \frac{c_\varepsilon}{\varepsilon f^2(v_\varepsilon)} dx \\ &\stackrel{(4.1)}{=} \sup_{T>0} \lim_{\varepsilon \rightarrow 0} \int_{\frac{L}{2}-\varepsilon T}^{\frac{L}{2}+\varepsilon T} u'_\varepsilon dx = \sup_{T>0} \lim_{\varepsilon \rightarrow 0} (u_\varepsilon(\frac{L}{2} + \varepsilon T) - u_\varepsilon(\frac{L}{2} - \varepsilon T)) \leq u(\frac{L}{2} + \delta) - u(\frac{L}{2} - \delta)\end{aligned}$$

for every $\delta > 0$ such that $u_\varepsilon(\frac{L}{2} \pm \delta) \rightarrow u(\frac{L}{2} \pm \delta)$, since u_ε is monotone nondecreasing. By letting $\delta \rightarrow 0$ we obtain $s_0 \leq [u](\frac{L}{2})$.

We next show the opposite inequality $s_0 \geq [u](\frac{L}{2})$. For $\delta > 0$ as above we have

$$\begin{aligned}
u(\frac{L}{2} + \delta) - u(\frac{L}{2} - \delta) &= \lim_{\varepsilon \rightarrow 0} (u_\varepsilon(\frac{L}{2} + \delta) - u_\varepsilon(\frac{L}{2} - \delta)) = \lim_{\varepsilon \rightarrow 0} \int_{\frac{L}{2} - \delta}^{\frac{L}{2} + \delta} u'_\varepsilon \, dx \\
&= \lim_{\varepsilon \rightarrow 0} \int_{\frac{L}{2} - \delta}^{\frac{L}{2} + \delta} \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} \, dx \leq \lim_{\varepsilon \rightarrow 0} \int_{\frac{L}{2} - x_\varepsilon}^{\frac{L}{2} + x_\varepsilon} \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} \, dx + 2\delta c_0 \\
&= \lim_{\varepsilon \rightarrow 0} 2 \int_0^{\frac{x_\varepsilon}{\varepsilon}} \frac{c_\varepsilon}{f^2(\tilde{v}_\varepsilon)} \, dt + 2\delta c_0 = \lim_{\varepsilon \rightarrow 0} 2c_0 \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{f^2(s)\sqrt{\Psi_\varepsilon(s)}} + 2\delta c_0 \\
&\leq 2c_0 \int_{m_0}^1 \frac{ds}{f^2(s)\sqrt{\Psi_0(s)}} + 2\delta c_0 \stackrel{(3.7)}{=} s_0 + 2\delta c_0, \tag{5.19}
\end{aligned}$$

where in the second line we have used (4.1) and $f_\varepsilon(v_\varepsilon) \geq f_\varepsilon(s_\varepsilon)$ in $(\frac{L}{2} - x_\varepsilon, \frac{L}{2} + x_\varepsilon)^c$, with $f_\varepsilon(s_\varepsilon) \rightarrow 1$ and $x_\varepsilon \rightarrow 0$ by Lemma 5.2; in the third line we have used

$$(\tilde{v}'_\varepsilon)^2 = \Psi_\varepsilon(\tilde{v}_\varepsilon)$$

with

$$\Psi_\varepsilon(s) := \frac{1}{4f^2(s)} \left((1-s)^2 f^2(s) - (1-m_\varepsilon)^2 f^2(m_\varepsilon) + 4\varepsilon |d_\varepsilon| (f^2(s) - f^2(m_\varepsilon)) \right), \quad s \in (m_0, 1),$$

which comes from (4.5), (5.6), and (5.4); and in the fourth line we have set

$$\Psi_0(s) := \frac{1}{4f^2(s)} \left((1-s)^2 f^2(s) - (1-m_0)^2 f^2(m_0) \right), \quad s \in (m_0, 1).$$

As $\delta \rightarrow 0$, we get $[u](\frac{L}{2}) \leq s_0$.

Hence $s_0 = [u](\frac{L}{2})$ which in turn yields, as we have seen before, that (5.12) holds. The only missing point to complete the proof of Proposition 5.3 is the convergence of the energies (1.20). However, this follows immediately by combining Lemma 5.4 with the computation below, which is based on the same arguments used in (5.19):

$$\begin{aligned}
&\lim_{\varepsilon \rightarrow 0} \int_{A_\varepsilon} \left(f_\varepsilon^2(v_\varepsilon) |u'_\varepsilon|^2 + \frac{(1-v_\varepsilon)^2}{4\varepsilon} + \varepsilon |v'_\varepsilon|^2 \right) dx \stackrel{(4.3)}{=} \lim_{\varepsilon \rightarrow 0} \left(2 \int_{A_\varepsilon} \frac{(1-v_\varepsilon)^2}{4\varepsilon} \, dx - \int_{A_\varepsilon} d_\varepsilon \, dx \right) \\
&= \lim_{\varepsilon \rightarrow 0} 2 \int_{-\frac{x_\varepsilon}{\varepsilon}}^{\frac{x_\varepsilon}{\varepsilon}} \frac{(1-\tilde{v}_\varepsilon)^2}{4} \, dt = \lim_{\varepsilon \rightarrow 0} \int_{m_\varepsilon}^{s_\varepsilon} \frac{(1-s)^2}{\sqrt{\Psi_\varepsilon(s)}} \, ds = \int_{m_0}^1 \frac{(1-s)^2}{\sqrt{\Psi_0(s)}} \, ds \stackrel{(3.8)}{=} g(s_0) = g([u](\frac{L}{2})),
\end{aligned}$$

where in the fourth equality we have used dominated convergence. This is allowed since, setting $\tilde{f}(s) := (1-s)f(s)$, which is monotone increasing by assumption (f3), we have

$$\frac{(1-s)^2}{\sqrt{\Psi_\varepsilon(s)}} \leq \frac{2\tilde{f}(s)(1-s)}{\sqrt{\tilde{f}^2(s) - \tilde{f}^2(m_\varepsilon)}} \leq \frac{2\tilde{f}(s)(1-s)}{\sqrt{2\tilde{f}(m_\varepsilon)\sqrt{\tilde{f}'(\zeta_\varepsilon(s))(s-m_\varepsilon)}}},$$

for all $s \in (m_\varepsilon, s_\varepsilon)$ and some $\zeta_\varepsilon(s) \in (m_\varepsilon, s)$. Since $\zeta_\varepsilon(s) \geq m_\varepsilon$ and $\inf_\varepsilon m_\varepsilon > 0$, in view of assumptions (f3) and (f5) we have that $\tilde{f}'(\zeta_\varepsilon(s)) \geq C(1-\zeta_\varepsilon(s))^3 \geq C(1-s)^3$ for all $s \in (m_\varepsilon, s_\varepsilon)$ and for a constant $C > 0$ independent of ε . Therefore we find for another constant $C_1 > 0$ independent of ε

$$\frac{(1-s)^2}{\sqrt{\Psi_\varepsilon(s)}} \leq \frac{C}{\sqrt{1-s}\sqrt{s-m_\varepsilon}}$$

which allows to apply the (generalized) dominated convergence theorem, as required. This concludes the proof. \square

5.2. Case II: complete fracture. We next show that if $m_0 = 0$ then the limit function u is a critical point of the cohesive energy (1.1) describing a completely fractured state, namely u has a single jump at $\frac{L}{2}$ and is constant elsewhere.

Proposition 5.5. *Assume that $m_0 = 0$. Then necessarily $s_{\text{frac}} \in \mathbb{R}$ and $a = s_{\text{frac}}$, and $u(x) = a\chi_{(\frac{L}{2}, L)}(x)$. Furthermore the convergence of the energies (1.20) holds.*

Proof. The proof is in part similar to that of Proposition 5.3. By Lemma 5.1 we have $c_0 = 0$. We can also apply Lemma 5.2 to deduce that $A_\varepsilon := \{v_\varepsilon < s_\varepsilon\} = (\frac{L}{2} - x_\varepsilon, \frac{L}{2} + x_\varepsilon)$ with $x_\varepsilon \rightarrow 0$ and $\frac{x_\varepsilon}{\varepsilon} \rightarrow +\infty$. Moreover, by the same lemma we have $u \in \text{SBV}(0, L)$ with $J_u \subset \{\frac{L}{2}\}$ and $u' = c_0 = 0$ almost everywhere in $(0, L)$, with uniform convergence $u'_\varepsilon \rightarrow 0$ on compact sets not containing $\frac{L}{2}$ by (5.11). We can also repeat word by word the proof of Lemma 5.4, so that

$$\lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon; A_\varepsilon^c) = 0, \quad c_0^2 + d_0 = 0 \quad (5.20)$$

(hence $d_0 = 0$).

For any $x \in (0, \frac{L}{2})$ we have $u_\varepsilon(x) = \int_0^x u'_\varepsilon dt \rightarrow c_0 x = 0$, whereas for $x \in (\frac{L}{2}, L)$ we have $u_\varepsilon(x) = u_\varepsilon(L) - \int_x^L u'_\varepsilon dt \rightarrow a - c_0(L - x) = a$. Therefore the limit function is $u(x) = a\chi_{(\frac{L}{2}, L)}(x)$ with a jump at the midpoint of amplitude $[u](\frac{L}{2}) = a$.

We now claim that

$$\lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) = 1. \quad (5.21)$$

We first observe that by (4.1) and the Dirichlet boundary condition (1.17c)

$$\lim_{\varepsilon \rightarrow 0} \int_0^L f_\varepsilon^2(v_\varepsilon)(u'_\varepsilon)^2 dx = \lim_{\varepsilon \rightarrow 0} c_\varepsilon \int_0^L u'_\varepsilon dx = \lim_{\varepsilon \rightarrow 0} c_\varepsilon a_\varepsilon = 0. \quad (5.22)$$

By (4.3) we have

$$\lim_{\varepsilon \rightarrow 0} \int_0^L \left| \frac{(1 - v_\varepsilon)^2}{4\varepsilon} - \varepsilon(v'_\varepsilon)^2 \right| dx = \lim_{\varepsilon \rightarrow 0} \int_0^L |c_\varepsilon u'_\varepsilon + d_\varepsilon| dx \leq \lim_{\varepsilon \rightarrow 0} (c_\varepsilon a_\varepsilon + |d_\varepsilon|L) = 0$$

(where we used in particular that $c_0 = 0$, $d_0 = 0$), and in turn it follows that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^L \left(\frac{(1 - v_\varepsilon)^2}{4\varepsilon} + \varepsilon(v'_\varepsilon)^2 - (1 - v_\varepsilon)|v'_\varepsilon| \right) dx &= \lim_{\varepsilon \rightarrow 0} \int_0^L \left(\frac{1 - v_\varepsilon}{2\sqrt{\varepsilon}} - \sqrt{\varepsilon}|v'_\varepsilon| \right)^2 dx \\ &\leq \lim_{\varepsilon \rightarrow 0} \int_0^L \left| \frac{(1 - v_\varepsilon)^2}{4\varepsilon} - \varepsilon(v'_\varepsilon)^2 \right| dx = 0. \end{aligned}$$

Therefore, combining this equation and (5.22), we find

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) &= \lim_{\varepsilon \rightarrow 0} \int_0^L f_\varepsilon^2(v_\varepsilon)(u'_\varepsilon)^2 dx + \int_0^L \left(\frac{(1 - v_\varepsilon)^2}{4\varepsilon} + \varepsilon(v'_\varepsilon)^2 \right) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^L (1 - v_\varepsilon)|v'_\varepsilon| dx = \lim_{\varepsilon \rightarrow 0} 2 \int_0^{\frac{L}{2}} (1 - v_\varepsilon)(-v'_\varepsilon) dx \\ &= \lim_{\varepsilon \rightarrow 0} (1 - v_\varepsilon(\frac{L}{2}))^2 = \lim_{\varepsilon \rightarrow 0} (1 - m_\varepsilon)^2 = 1, \end{aligned}$$

which proves the claim (5.21).

To conclude the proof, we need to show that

$$[u](\frac{L}{2}) = s_{\text{frac}} \quad (5.23)$$

(and, in particular, that s_{frac} is finite). Indeed, recalling that u is piecewise constant, in this case we have $\mathcal{F}(u, 1) = g([u](\frac{L}{2})) = 1$, and therefore (5.21) gives also the convergence of the energy. The rest of the proof is therefore devoted to showing (5.23).

Let $\tilde{u}_\varepsilon(t) := u_\varepsilon(\frac{L}{2} + \varepsilon t)$, $\tilde{v}_\varepsilon(t) := v_\varepsilon(\frac{L}{2} + \varepsilon t)$, for $t \in (-\frac{x_\varepsilon}{\varepsilon}, \frac{x_\varepsilon}{\varepsilon})$. We first check that $\tilde{v}_\varepsilon(t) \rightharpoonup 1 - e^{-|t|/2}$ weakly in $H^1_{\text{loc}}(\mathbb{R})$ and uniformly on compact sets. Indeed as in (5.14) we have that $(1 - \tilde{v}_\varepsilon)$ is uniformly bounded in $H^1(-T, T)$ for all fixed $T > 0$, so that \tilde{v}_ε converges weakly in $H^1_{\text{loc}}(\mathbb{R})$ and uniformly on compact sets to some function \tilde{v} as $\varepsilon \rightarrow 0$, with $1 - \tilde{v} \in H^1(\mathbb{R})$. Also, $\tilde{v}(0) = \lim_\varepsilon \tilde{v}_\varepsilon(0) = \lim_\varepsilon m_\varepsilon = 0$. We now check that $\{\tilde{v} = 0\} = \{0\}$. By the properties of v_ε , we have $\{\tilde{v} = 0\} = [-\tilde{x}, \tilde{x}]$, for some $\tilde{x} \geq 0$. Recalling (5.21) and that $\frac{x_\varepsilon}{\varepsilon} \rightarrow +\infty$ by Lemma 5.2, we have

$$1 = \lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} \int_{-\frac{x_\varepsilon}{\varepsilon}}^{\frac{x_\varepsilon}{\varepsilon}} \left(\frac{(1 - \tilde{v}_\varepsilon)^2}{4} + (\tilde{v}'_\varepsilon)^2 \right) dt \geq \int_{-T}^T \left(\frac{(1 - \tilde{v})^2}{4} + (\tilde{v}')^2 \right) dt$$

for all $T > 0$, so that

$$1 \geq \int_{\mathbb{R}} \left(\frac{(1 - \tilde{v})^2}{4} + (\tilde{v}')^2 \right) dt \geq \frac{\tilde{x}}{2} + \int_{\mathbb{R} \setminus (-\tilde{x}, \tilde{x})} \left(\frac{(1 - \tilde{v})^2}{4} + (\tilde{v}')^2 \right) dt \geq \frac{\tilde{x}}{2} + 1.$$

This implies $\tilde{x} = 0$ and in turn $\{\tilde{v} = 0\} = \{0\}$.

By writing in weak form the equation (5.7) satisfied by \tilde{v}_ε and passing to the limit as $\varepsilon \rightarrow 0$ we get for all $\varphi \in C_c^\infty(\mathbb{R} \setminus \{0\})$

$$0 = \int_{\text{supp}(\varphi)} \left(\tilde{v}'_\varepsilon \varphi' + \frac{c_\varepsilon^2 f'(\tilde{v}_\varepsilon)}{f^3(\tilde{v}_\varepsilon)} \varphi + \frac{\tilde{v}_\varepsilon - 1}{4} \varphi \right) dt \rightarrow \int_{\text{supp}(\varphi)} \left(\tilde{v}' \varphi' + \frac{\tilde{v} - 1}{4} \varphi \right) dt,$$

where we used the weak L^2 -convergence of \tilde{v}'_ε for the first term and the uniform convergence for the second and the third term. Together with $\tilde{v}(0) = 0$ and $\tilde{v}(+\infty) = 1$, this implies $\tilde{v}(t) = 1 - e^{-|t|/2}$.

We further recall that by changing variables in (4.5), using (5.6) and since $d_\varepsilon < 0$ by (5.4), we have in $(-\frac{x_\varepsilon}{\varepsilon}, \frac{x_\varepsilon}{\varepsilon})$

$$\begin{aligned} (\tilde{v}'_\varepsilon)^2 &= \frac{1}{4f^2(\tilde{v}_\varepsilon)} \left((1 - \tilde{v}_\varepsilon)^2 f^2(\tilde{v}_\varepsilon) - 4c_\varepsilon^2 - 4\varepsilon d_\varepsilon f^2(\tilde{v}_\varepsilon) \right) \\ &= \frac{1}{4f^2(\tilde{v}_\varepsilon)} \left((1 - \tilde{v}_\varepsilon)^2 f^2(\tilde{v}_\varepsilon) - (1 - m_\varepsilon)^2 f^2(m_\varepsilon) + 4\varepsilon |d_\varepsilon| (f^2(\tilde{v}_\varepsilon) - f^2(m_\varepsilon)) \right) =: \Psi_\varepsilon(\tilde{v}_\varepsilon). \end{aligned} \quad (5.24)$$

Let us now compute $[u](\frac{L}{2})$. We have

$$\begin{aligned} [u](\frac{L}{2}) &= a = \lim_{\varepsilon \rightarrow 0} \int_0^L u'_\varepsilon dx \stackrel{(4.1)}{=} \lim_{\varepsilon \rightarrow 0} 2 \int_{\frac{L}{2}}^L \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} dx = \lim_{\varepsilon \rightarrow 0} 2 \int_{\frac{L}{2}}^{\frac{L}{2} + x_\varepsilon} \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} dx \\ &= \lim_{\varepsilon \rightarrow 0} 2 \int_0^{\frac{x_\varepsilon}{\varepsilon}} \frac{c_\varepsilon}{f^2(\tilde{v}_\varepsilon)} dt \stackrel{(5.24)}{=} \lim_{\varepsilon \rightarrow 0} 2c_\varepsilon \int_0^{\frac{x_\varepsilon}{\varepsilon}} \frac{\tilde{v}'_\varepsilon}{f^2(\tilde{v}_\varepsilon) \sqrt{\Psi_\varepsilon(\tilde{v}_\varepsilon)}} dt \\ &= \lim_{\varepsilon \rightarrow 0} 2c_\varepsilon \int_{m_\varepsilon}^{s_\varepsilon} \frac{1}{f^2(s) \sqrt{\Psi_\varepsilon(s)}} ds, \end{aligned} \quad (5.25)$$

where in the fourth equality we have used that $f_\varepsilon(v_\varepsilon) \geq f_\varepsilon(v_\varepsilon(L/2 + x_\varepsilon)) \rightarrow 1$ in A_ε^c and that $c_\varepsilon \rightarrow 0$, and the last passage follows by a change of variables. Fix now any $\delta \in (0, 1)$. Since $m_\varepsilon < \delta < s_\varepsilon$ for ε small enough, we have by the definition of Ψ_ε that

$$\begin{aligned} 0 &\leq \lim_{\varepsilon \rightarrow 0} 2c_\varepsilon \int_\delta^{s_\varepsilon} \frac{ds}{f^2(s)\sqrt{\Psi_\varepsilon(s)}} \leq \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \int_\delta^{s_\varepsilon} \frac{ds}{f(s)\sqrt{(1-s)^2 f^2(s) - (1-m_\varepsilon)^2 f^2(m_\varepsilon)}} \\ &\leq \lim_{\varepsilon \rightarrow 0} \frac{4c_\varepsilon(s_\varepsilon - \delta)}{f(\delta)\sqrt{(1-\delta)^2 f^2(\delta) - (1-m_\varepsilon)^2 f^2(m_\varepsilon)}} = 0 \end{aligned}$$

where we used the monotonicity of the map $s \mapsto (1-s)f(s)$ given by assumption (f3). Therefore by (5.25) we see that for all $\delta \in (0, 1)$

$$[u](\frac{L}{2}) = a = \lim_{\varepsilon \rightarrow 0} 2c_\varepsilon \int_{m_\varepsilon}^\delta \frac{1}{f^2(s)\sqrt{\Psi_\varepsilon(s)}} ds. \quad (5.26)$$

By inserting the definition of Ψ_ε , we find

$$\begin{aligned} [u](\frac{L}{2}) &= \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \int_{m_\varepsilon}^\delta \frac{ds}{f(s)\sqrt{(1-s)^2 f^2(s) - (1-m_\varepsilon)^2 f^2(m_\varepsilon) + 4\varepsilon|d_\varepsilon|(f^2(s) - f^2(m_\varepsilon))}} \\ &\geq \lim_{\varepsilon \rightarrow 0} \frac{4c_\varepsilon}{\sqrt{(1-m_\varepsilon)^2 + 4\varepsilon|d_\varepsilon|}} \int_{m_\varepsilon}^\delta \frac{ds}{f(s)\sqrt{f^2(s) - f^2(m_\varepsilon)}} \\ &\geq \lim_{\varepsilon \rightarrow 0} 2f(m_\varepsilon) \inf_{s \in (m_\varepsilon, \delta)} \left(\frac{1}{f'(s)} \right) \int_{m_\varepsilon}^\delta \frac{f'(s)}{f(s)\sqrt{f^2(s) - f^2(m_\varepsilon)}} ds \\ &= \lim_{\varepsilon \rightarrow 0} 2f(m_\varepsilon) \inf_{s \in (m_\varepsilon, \delta)} \left(\frac{1}{f'(s)} \right) \int_{f(m_\varepsilon)}^{f(\delta)} \frac{dt}{t\sqrt{t^2 - f^2(m_\varepsilon)}} \\ &= \lim_{\varepsilon \rightarrow 0} 2 \inf_{s \in (m_\varepsilon, \delta)} \left(\frac{1}{f'(s)} \right) \arctan \left(\frac{\sqrt{f^2(\delta) - f^2(m_\varepsilon)}}{f(m_\varepsilon)} \right) = \pi \inf_{s \in (0, \delta)} \left(\frac{1}{f'(s)} \right). \end{aligned}$$

Similarly, again by (5.26) and using the definition of Ψ_ε , and denoting by $\tilde{f}(s) := (1-s)f(s)$, we have

$$\begin{aligned} [u](\frac{L}{2}) &= \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \int_{m_\varepsilon}^\delta \frac{ds}{f(s)\sqrt{(1-s)^2 f^2(s) - (1-m_\varepsilon)^2 f^2(m_\varepsilon) + 4\varepsilon|d_\varepsilon|(f^2(s) - f^2(m_\varepsilon))}} \\ &\leq \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \int_{m_\varepsilon}^\delta \frac{ds}{f(s)\sqrt{(1-s)^2 f^2(s) - (1-m_\varepsilon)^2 f^2(m_\varepsilon)}} \\ &= \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \int_{m_\varepsilon}^\delta \frac{\tilde{f}'(s)}{\tilde{f}(s)\sqrt{\tilde{f}^2(s) - \tilde{f}^2(m_\varepsilon)}} \cdot \frac{(1-s)}{\tilde{f}'(s)} ds \\ &\leq \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \sup_{s \in (m_\varepsilon, \delta)} \left(\frac{1-s}{\tilde{f}'(s)} \right) \int_{\tilde{f}(m_\varepsilon)}^{\tilde{f}(\delta)} \frac{dt}{t\sqrt{t^2 - \tilde{f}^2(m_\varepsilon)}} \\ &= \lim_{\varepsilon \rightarrow 0} 4c_\varepsilon \sup_{s \in (m_\varepsilon, \delta)} \left(\frac{1-s}{\tilde{f}'(s)} \right) \frac{1}{\tilde{f}(m_\varepsilon)} \arctan \left(\frac{\sqrt{\tilde{f}^2(\delta) - \tilde{f}^2(m_\varepsilon)}}{\tilde{f}(m_\varepsilon)} \right) = \pi \sup_{s \in (0, \delta)} \left(\frac{1-s}{\tilde{f}'(s)} \right) \end{aligned}$$

where we used the fact that $\frac{4c_\varepsilon}{\tilde{f}(m_\varepsilon)} \rightarrow 2$ by (5.6). By collecting the previous inequalities we conclude that for all $\delta \in (0, 1)$

$$\pi \inf_{s \in (0, \delta)} \left(\frac{1}{f'(s)} \right) \leq [u](\frac{L}{2}) \leq \pi \sup_{s \in (0, \delta)} \left(\frac{1-s}{\tilde{f}'(s)} \right)$$

so that by letting $\delta \rightarrow 0$ and recalling Proposition 3.7, we get $[u](\frac{L}{2}) = \frac{\pi}{f'(0)} = s_{\text{frac}}$. With a small abuse of notation, the previous computation says that if $f'(0) = 0$, then $[u](\frac{L}{2}) = +\infty$, which is a contradiction with $[u](\frac{L}{2}) = a < +\infty$. Hence necessarily s_{frac} is finite and (5.23) holds. \square

5.3. Case III: elastic regime. We eventually consider the case $m_0 = 1$. The limit behaviour of the family $(u_\varepsilon, v_\varepsilon)$ is an elastic critical point, as described by the following proposition.

Proposition 5.6. *Assume that $m_0 = 1$. Then $u(x) = \frac{a}{L}x$ and $c_0 = \frac{a}{L}$. Moreover the convergence of the energies (1.20) holds if and only if $\frac{a}{L} \leq \frac{\sigma_c}{2}$.*

Proof. We distinguish three cases, depending on whether the minimum value m_ε of v_ε is above or below the threshold s_ε (see (1.8)) and whether $f_\varepsilon(m_\varepsilon)$ converges to zero or not.

Step 1: $m_\varepsilon \geq s_\varepsilon$. In this case $f_\varepsilon(v_\varepsilon(x)) \geq f_\varepsilon(m_\varepsilon) = \psi_\varepsilon(m_\varepsilon) \rightarrow 1$, that is, $f_\varepsilon(v_\varepsilon)$ converges to 1 uniformly in $[0, L]$. In turn $u'_\varepsilon \rightarrow c_0$ uniformly in $[0, L]$ by (4.1) and $u' \equiv c_0$. In view of the boundary conditions (1.17c)

$$a = \lim_{\varepsilon \rightarrow 0} a_\varepsilon = \lim_{\varepsilon \rightarrow 0} \int_0^L u'_\varepsilon dx = c_0 L,$$

therefore $c_0 = \frac{a}{L}$ and $u(x) = \frac{a}{L}x$. To complete the proof in this case, it only remains to show that the convergence of the energy holds if and only if $c_0 \leq \frac{\sigma_c}{2}$.

If $c_0 > \frac{\sigma_c}{2}$, then

$$\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq \int_0^L f_\varepsilon^2(v_\varepsilon) |u'_\varepsilon|^2 dx \xrightarrow{\varepsilon \rightarrow 0} c_0^2 L > \left(\sigma_c c_0 - \frac{\sigma_c^2}{4} \right) L = \int_0^L \left(\sigma_c u' - \frac{\sigma_c^2}{4} \right) dx = \mathcal{F}(u, 1).$$

Conversely, assume that $c_0 \leq \frac{\sigma_c}{2}$. Recalling the definition of the discrepancy ξ_ε in (5.3) and evaluating it at the point y_ε where $\xi_\varepsilon(y_\varepsilon) = 0$, by the uniform convergence $f_\varepsilon(v_\varepsilon) \rightarrow 1$ we have

$$c_0^2 + d_0 = \lim_{\varepsilon \rightarrow 0} \left(\frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon(y_\varepsilon))} + d_\varepsilon \right) = \lim_{\varepsilon \rightarrow 0} \xi_\varepsilon(y_\varepsilon) = 0,$$

and therefore $c_0^2 + d_0 = 0$. Moreover, we compute

$$\begin{aligned} \int_0^L \left| \frac{(1-v_\varepsilon)^2}{4\varepsilon} - \varepsilon(v'_\varepsilon)^2 \right| dx &= 2 \int_0^{\frac{L}{2}} |\xi_\varepsilon(x)| dx = -2 \int_0^{y_\varepsilon} \xi_\varepsilon(x) dx + 2 \int_{y_\varepsilon}^{\frac{L}{2}} \xi_\varepsilon(x) dx \\ &= -2 \int_0^{y_\varepsilon} (c_\varepsilon u'_\varepsilon + d_\varepsilon) dx + 2 \int_{y_\varepsilon}^{\frac{L}{2}} (c_\varepsilon u'_\varepsilon + d_\varepsilon) dx \\ &= 2d_\varepsilon \left(\frac{L}{2} - 2y_\varepsilon \right) + 2c_\varepsilon \left(u_\varepsilon \left(\frac{L}{2} \right) - 2u_\varepsilon(y_\varepsilon) \right) \\ &= d_\varepsilon L + 2c_\varepsilon u_\varepsilon \left(\frac{L}{2} \right) - 4 \left(d_\varepsilon y_\varepsilon + c_\varepsilon \int_0^{y_\varepsilon} u'_\varepsilon dx \right), \end{aligned}$$

so that by passing to the limit, and assuming up to subsequences $y_\varepsilon \rightarrow y_0 \in [0, L]$,

$$\lim_{\varepsilon \rightarrow 0} \int_0^L \left| \frac{(1 - v_\varepsilon)^2}{4\varepsilon} - \varepsilon(v'_\varepsilon)^2 \right| dx = d_0L + c_0a - 4(d_0y_0 + c_0^2y_0) = 0, \quad (5.27)$$

where the last equality follows by the identities $c_0^2 + d_0 = 0$ and $c_0 = \frac{a}{L}$. We deduce that

$$\begin{aligned} \int_0^L \left(\frac{(1 - v_\varepsilon)^2}{4\varepsilon} + \varepsilon(v'_\varepsilon)^2 - (1 - v_\varepsilon)|v'_\varepsilon| \right) dx &= \int_0^L \left(\frac{1 - v_\varepsilon}{2\sqrt{\varepsilon}} - \sqrt{\varepsilon}|v'_\varepsilon| \right)^2 dx \\ &\leq \int_0^L \left| \frac{(1 - v_\varepsilon)^2}{4\varepsilon} - \varepsilon(v'_\varepsilon)^2 \right| dx \rightarrow 0, \end{aligned}$$

and in turn

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^L \left(\frac{(1 - v_\varepsilon)^2}{4\varepsilon} + \varepsilon(v'_\varepsilon)^2 \right) dx &= \lim_{\varepsilon \rightarrow 0} \int_0^L (1 - v_\varepsilon)|v'_\varepsilon| dx = \lim_{\varepsilon \rightarrow 0} -2 \int_0^{\frac{L}{2}} (1 - v_\varepsilon)v'_\varepsilon dx \\ &= \lim_{\varepsilon \rightarrow 0} (1 - v_\varepsilon(\frac{L}{2}))^2 = \lim_{\varepsilon \rightarrow 0} (1 - m_\varepsilon)^2 = 0. \end{aligned} \quad (5.28)$$

In view of (5.28), we conclude that the convergence of the energy holds:

$$\lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \stackrel{(5.28)}{=} \lim_{\varepsilon \rightarrow 0} \int_0^L f_\varepsilon^2(v_\varepsilon)|u'_\varepsilon|^2 dx = c_0^2L = \int_0^L (u')^2 dx = \mathcal{F}(u, 1),$$

where we used the uniform convergences $f_\varepsilon(v_\varepsilon) \rightarrow 1$ and $u'_\varepsilon \rightarrow c_0$, and $u' \equiv c_0 \leq \frac{\sigma_c}{2}$.

Step 2: $m_\varepsilon < s_\varepsilon$. In this case we can apply Lemma 5.2 to deduce that $|A_\varepsilon| \rightarrow 0$, where $A_\varepsilon = \{v_\varepsilon < s_\varepsilon\}$, and that $u'_\varepsilon \rightarrow c_0$ uniformly on compact subsets of $[0, L]$ not containing $\frac{L}{2}$, see in particular (5.11). For any $x \in (0, \frac{L}{2})$ we have $u_\varepsilon(x) = \int_0^x u'_\varepsilon dt \rightarrow c_0x$, whereas for $x \in (\frac{L}{2}, L)$ we have $u_\varepsilon(x) = u_\varepsilon(L) - \int_x^L u'_\varepsilon dt \rightarrow a - c_0(L - x)$. Therefore the limit function is

$$u(x) = \begin{cases} c_0x & \text{if } x \in (0, \frac{L}{2}), \\ c_0x + a - c_0L & \text{if } x \in (\frac{L}{2}, L), \end{cases} \quad (5.29)$$

with a possible jump at $\frac{L}{2}$ with amplitude $[u](\frac{L}{2}) = a - c_0L$. Notice that $c_0 \leq \frac{a}{L}$ by (4.7). We next distinguish two further subcases depending on the limit value of $f_\varepsilon(m_\varepsilon)$.

Step 2a: $f_\varepsilon(m_\varepsilon) \rightarrow 0$. In this case by Lemma 5.1 we have $c_0 = \frac{\sigma_c}{2}$. Let us first show that $[u](\frac{L}{2}) = 0$. We consider once more the blow-up $\tilde{v}_\varepsilon(t) := v_\varepsilon(\frac{L}{2} + \varepsilon t)$, which obeys the equation (5.24) in $(-\frac{x_\varepsilon}{\varepsilon}, \frac{x_\varepsilon}{\varepsilon})$. Denote by $\tilde{f}(s) := (1 - s)f(s)$ and recall that \tilde{f} is strictly increasing by assumption (f3). By monotonicity of f and \tilde{f} , the function Ψ_ε defined in (5.24) satisfies

$$\Psi_\varepsilon(s) \geq \frac{\tilde{f}^2(s) - \tilde{f}^2(m_\varepsilon)}{4f^2(s)} = \frac{\tilde{f}(s) - \tilde{f}(m_\varepsilon)}{4f^2(s)} \cdot (\tilde{f}(s) + \tilde{f}(m_\varepsilon)) \geq \frac{\tilde{f}(s) - \tilde{f}(m_\varepsilon)}{2f^2(s)} \cdot \tilde{f}(m_\varepsilon)$$

for $s \in (m_\varepsilon, 1)$. Then by arguing as in (5.25) we have, recalling that $c_\varepsilon \rightarrow \frac{\sigma_c}{2}$, $\tilde{v}_\varepsilon(\frac{x_\varepsilon}{\varepsilon}) = s_\varepsilon$, and $(\tilde{v}'_\varepsilon)^2 = \Psi_\varepsilon(\tilde{v}_\varepsilon)$ by (5.24),

$$\begin{aligned} [u](\frac{L}{2}) &= a - c_0L = \lim_{\varepsilon \rightarrow 0} 2 \int_{\frac{L}{2}}^L \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} dx - c_0L = \lim_{\varepsilon \rightarrow 0} 2c_\varepsilon \int_0^{\frac{x_\varepsilon}{\varepsilon}} \frac{dt}{f^2(\tilde{v}_\varepsilon)} + c_0L - c_0L \\ &= \lim_{\varepsilon \rightarrow 0} \sigma_c \int_0^{\frac{x_\varepsilon}{\varepsilon}} \frac{\tilde{v}'_\varepsilon}{f^2(\tilde{v}_\varepsilon) \sqrt{\Psi_\varepsilon(\tilde{v}_\varepsilon)}} dt = \lim_{\varepsilon \rightarrow 0} \sigma_c \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{f^2(s) \sqrt{\Psi_\varepsilon(s)}} \\ &\leq \lim_{\varepsilon \rightarrow 0} \frac{\sqrt{2}\sigma_c}{(\tilde{f}(m_\varepsilon))^{1/2}} \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{f(s)(\tilde{f}(s) - \tilde{f}(m_\varepsilon))^{1/2}} \\ &= \lim_{\varepsilon \rightarrow 0} \sqrt{2}\sigma_c \int_{m_\varepsilon}^{s_\varepsilon} \frac{(1-s) ds}{\tilde{f}(s)(\tilde{f}(s) - \tilde{f}(m_\varepsilon))^{1/2}} \leq \lim_{\varepsilon \rightarrow 0} \frac{\sqrt{2}\sigma_c}{\tilde{f}(m_\varepsilon)} \int_{m_\varepsilon}^{s_\varepsilon} \frac{(1-s) ds}{(\tilde{f}(s) - \tilde{f}(m_\varepsilon))^{1/2}} \\ &= \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{2}{\sigma_c}} \int_{m_\varepsilon}^{s_\varepsilon} \frac{(1-s) ds}{(\tilde{f}(s) - \tilde{f}(m_\varepsilon))^{1/2}}. \end{aligned}$$

We can now write, for $s \in (m_\varepsilon, s_\varepsilon)$, $\tilde{f}(s) - \tilde{f}(m_\varepsilon) = \tilde{f}'(\zeta_\varepsilon(s))(s - m_\varepsilon)$ for some point $\zeta_\varepsilon(s) \in (m_\varepsilon, s)$. Since $\zeta_\varepsilon(s) \geq m_\varepsilon \rightarrow 1$ and in view of assumption (f5), given any $M > 0$ we have that for all ε small enough

$$\tilde{f}'(\zeta_\varepsilon(s)) \geq M(1 - \zeta_\varepsilon(s))^3 \geq M(1 - s)^3 \quad \text{for all } s \in (m_\varepsilon, s_\varepsilon),$$

hence

$$\begin{aligned} [u](\frac{L}{2}) &\leq \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{2}{\sigma_c}} \cdot \frac{1}{M} \int_{m_\varepsilon}^{s_\varepsilon} \frac{(1-s) ds}{\sqrt{s - m_\varepsilon}(1-s)^{3/2}} \\ &\leq \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{2}{\sigma_c}} \cdot \frac{1}{M} \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{\sqrt{s - m_\varepsilon}\sqrt{s_\varepsilon - s}} = \sqrt{\frac{2}{\sigma_c}} \cdot \frac{\pi}{M}. \end{aligned}$$

Since M is arbitrarily large, we conclude that $[u](\frac{L}{2}) = 0$, as claimed. In turn $u(x) = \frac{a}{L}x$ by (5.29). In particular we also have $\frac{a}{L} = c_0 = \frac{\sigma_c}{2}$.

To conclude the proof in this case, we need to show that the convergence of the energy holds. We preliminary show that

$$\lim_{\varepsilon \rightarrow 0} \int_0^L \varepsilon (v'_\varepsilon(x))^2 dx = 0. \quad (5.30)$$

To this aim, we multiply (4.6) by the function $v_\varepsilon - 1$ and integrate in $(0, L)$: we have, after integration by parts,

$$\begin{aligned} \int_0^L \varepsilon |v'_\varepsilon|^2 dx &= \int_0^L \frac{c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{f_\varepsilon^3(v_\varepsilon)} (1 - v_\varepsilon) dx - \int_0^L \frac{(1 - v_\varepsilon)^2}{4\varepsilon} dx \\ &= \int_0^L \left(\frac{4\varepsilon c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{(1 - v_\varepsilon) f_\varepsilon^3(v_\varepsilon)} - 1 \right) \frac{(1 - v_\varepsilon)^2}{4\varepsilon} dx. \end{aligned}$$

Now, for $x \in A_\varepsilon$ we have $v_\varepsilon(x) \leq s_\varepsilon$ and by definition of f_ε (see (1.8)) and \bar{f} (see (2.2))

$$\frac{4\varepsilon c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{(1 - v_\varepsilon) f_\varepsilon^3(v_\varepsilon)} - 1 = (2c_\varepsilon)^2 \bar{f}(v_\varepsilon) - 1 \leq (2c_\varepsilon)^2 \bar{f}(m_\varepsilon) - 1,$$

where we used the monotonicity of \bar{f} (see Proposition 2.1). Similarly, if $x \in A_\varepsilon^c$ we have $v_\varepsilon(x) \geq s_\varepsilon$ and by the monotonicity properties of ψ_ε in assumptions $(\psi 1)$ and $(\psi 3)$

$$\frac{4\varepsilon c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{(1-v_\varepsilon)f_\varepsilon^3(v_\varepsilon)} - 1 = \frac{(2c_\varepsilon)^2 \varepsilon \psi'_\varepsilon(v_\varepsilon)}{(1-v_\varepsilon)\psi_\varepsilon^3(v_\varepsilon)} - 1 \leq \frac{(2c_\varepsilon)^2 \varepsilon \psi'_\varepsilon(s_\varepsilon)}{(1-s_\varepsilon)\psi_\varepsilon^3(s_\varepsilon)} - 1 = (2c_\varepsilon)^2 \bar{f}(s_\varepsilon) - 1.$$

Hence

$$\int_0^L \varepsilon |v'_\varepsilon|^2 dx \leq \left((2c_\varepsilon)^2 \bar{f}(m_\varepsilon) - 1 \right) \int_0^L \frac{(1-v_\varepsilon)^2}{4\varepsilon} dx \rightarrow 0$$

since the integral on the right-hand side is uniformly bounded by (1.19), and $(2c_\varepsilon)^2 \bar{f}(m_\varepsilon) \rightarrow (\frac{2c_0}{\sigma_c})^2 = 1$ again by Proposition 2.1. Hence (5.30) follows.

We next show $c_0^2 + d_0 = 0$. By evaluating (4.5) at $x = 0$ we have

$$c_0^2 + d_0 = \lim_{\varepsilon \rightarrow 0} (c_\varepsilon^2 + d_\varepsilon) = \lim_{\varepsilon \rightarrow 0} -\varepsilon (v'_\varepsilon(0))^2 \leq 0,$$

hence $c_0^2 + d_0 \leq 0$. On the other hand, for every fixed $x_0 \in (0, \frac{L}{2})$ we have that $f_\varepsilon(v_\varepsilon) \rightarrow 1$ uniformly in $(0, x_0)$, and therefore

$$\begin{aligned} (c_0^2 + d_0)x_0 &= \lim_{\varepsilon \rightarrow 0} \int_0^{x_0} \left(\frac{c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon)} + d_\varepsilon \right) dx \stackrel{(4.5)}{=} \lim_{\varepsilon \rightarrow 0} \int_0^{x_0} \left(\frac{(1-v_\varepsilon)^2}{4\varepsilon} - \varepsilon (v'_\varepsilon)^2 \right) dx \\ &\geq - \lim_{\varepsilon \rightarrow 0} \int_0^{x_0} \varepsilon (v'_\varepsilon)^2 dx \stackrel{(5.30)}{=} 0, \end{aligned}$$

which combined with the inequality obtained before gives $c_0^2 + d_0 = 0$, as desired.

By using (4.3), (4.1), (5.30), and the identities $c_0^2 + d_0 = 0$, $c_0 = \frac{a}{L}$, we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^L \frac{(1-v_\varepsilon)^2}{4\varepsilon} dx &= \lim_{\varepsilon \rightarrow 0} \int_0^L (\varepsilon (v'_\varepsilon)^2 + c_\varepsilon u'_\varepsilon) dx + d_0 L \\ &= c_0 a + d_0 L = c_0 (a - c_0 L) = 0, \end{aligned} \tag{5.31}$$

and similarly

$$\lim_{\varepsilon \rightarrow 0} \int_0^L \left(\frac{(1-v_\varepsilon)^2}{4\varepsilon} - f_\varepsilon^2(v_\varepsilon) (u'_\varepsilon)^2 \right) dx = \lim_{\varepsilon \rightarrow 0} \int_0^L \varepsilon (v'_\varepsilon)^2 dx + d_0 L = -c_0^2 L. \tag{5.32}$$

Therefore combining (5.30), (5.31), and (5.32) we find

$$\lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) = c_0^2 L + 2 \lim_{\varepsilon \rightarrow 0} \int_0^L \frac{(1-v_\varepsilon)^2}{4\varepsilon} dx = \int_0^L (u')^2 dx = \mathcal{F}(u, 1), \tag{5.33}$$

that is, the convergence of the energy holds.

Step 2b: $f_\varepsilon(m_\varepsilon) \rightarrow \alpha > 0$. By monotonicity of f_ε

$$\mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq \int_0^L f_\varepsilon^2(v_\varepsilon) |u'_\varepsilon|^2 dx \geq \frac{\alpha^2}{2} \int_0^L |u'_\varepsilon|^2 dx$$

for all ε sufficiently small; by the uniform bound (1.19) we then deduce that $\|u_\varepsilon\|_{H^1(0,L)}$ is uniformly bounded, and therefore that the limit u belongs to $H^1(0,L)$. In particular u cannot jump at $\frac{L}{2}$ and by (5.29) we conclude that $u(x) = \frac{a}{L}x$ and $c_0 = \frac{a}{L}$.

To conclude the proof, we have to show that also in this case the convergence of the energy holds if and only if $c_0 \leq \frac{\sigma_c}{2}$. Assume first that $c_0 > \frac{\sigma_c}{2}$: then for every $\delta > 0$, by the uniform convergences $u'_\varepsilon \rightarrow c_0$ and $f_\varepsilon(v_\varepsilon) \rightarrow 1$ in $(\frac{L}{2} - \delta, \frac{L}{2} + \delta)^c$ we find

$$\liminf_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} \int_{(\frac{L}{2} - \delta, \frac{L}{2} + \delta)^c} f_\varepsilon^2(v_\varepsilon) |u'_\varepsilon|^2 dx = c_0^2(L - 2\delta),$$

so that by letting $\delta \rightarrow 0$ we find

$$\liminf_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) \geq c_0^2 L > \left(\sigma_c c_0 - \frac{\sigma_c^2}{4} \right) L = \int_0^L \left(\sigma_c u' - \frac{\sigma_c^2}{4} \right) dx = \mathcal{F}(u, 1),$$

that is, the convergence of the energy does not hold. If, instead, $c_0 \leq \frac{\sigma_c}{2}$, then one can prove that the convergence of the energy holds just by repeating the argument in Step 2a leading to (5.33). \square

Proof of Theorem 1.7. The result follows by combining Proposition 5.3, Proposition 5.5 and Proposition 5.6. \square

6. PROOF OF THE APPROXIMATION OF CRITICAL POINTS

In this section we give the proof of Theorem 1.8. We premise a technical lemma to the proof, which shows that it is possible to construct a solution v_ε to the ODE (4.6) which attains the boundary conditions $v_\varepsilon(0) = v_\varepsilon(L) = 1$.

Lemma 6.1. *Let $c_\varepsilon \in (0, \frac{\sigma_c}{2})$ be such that $\sup_\varepsilon c_\varepsilon < \frac{\sigma_c}{2}$. Then there exists $\varepsilon_0 > 0$ with the following property: for every $\varepsilon \in (0, \varepsilon_0)$ there exists $m_\varepsilon \in (0, z_{c_\varepsilon})$, with $(1 - m_\varepsilon)f(m_\varepsilon) \leq 2c_\varepsilon$, such that the unique solution to the initial value problem*

$$\begin{cases} \varepsilon v_\varepsilon'' = \frac{c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{f_\varepsilon^3(v_\varepsilon)} + \frac{v_\varepsilon - 1}{4\varepsilon} \\ v_\varepsilon(\frac{L}{2}) = m_\varepsilon \\ v'_\varepsilon(\frac{L}{2}) = 0 \end{cases} \quad (6.1)$$

satisfies $v_\varepsilon(0) = v_\varepsilon(L) = 1$. Moreover:

- if $\inf_\varepsilon c_\varepsilon > 0$ then $\inf_\varepsilon m_\varepsilon > 0$;
- if $c_\varepsilon \rightarrow 0$ then $m_\varepsilon \rightarrow 0$ and $\frac{2c_\varepsilon}{f(m_\varepsilon)} \rightarrow 1$.

Proof. Recall that the value $z_{c_\varepsilon} \in (0, 1)$ appearing in the statement is defined by the relation (2.5). Since $\sup_\varepsilon c_\varepsilon < \frac{\sigma_c}{2}$ we have that $\sup_\varepsilon z_{c_\varepsilon} < 1$ and therefore by choosing ε_0 small enough we can guarantee that $z_{c_\varepsilon} < s_\varepsilon$ for all $\varepsilon \in (0, \varepsilon_0)$.

For all $m \in (0, z_{c_\varepsilon})$ we consider the solution $v_\varepsilon(\cdot; m)$ of the initial value problem (6.1) with $v_\varepsilon(0; m) = m$, which exists and is unique by Cauchy-Lipschitz Theorem. The solution is also symmetric with respect to the point $\frac{L}{2}$. The proof of the lemma amounts to show that we can choose a value m_ε such that $v_\varepsilon(L; m_\varepsilon) = 1$.

Fix any $m \in (0, z_{c_\varepsilon})$ such that $(1 - m)f(m) \leq 2c_\varepsilon$. We first observe that, in the region $\{v_\varepsilon < s_\varepsilon\}$ (which contains an interval centered at the point $\frac{L}{2}$, since $m < s_\varepsilon$ and the solution is symmetric), the rescaled function $\tilde{v}_\varepsilon(t) := v_\varepsilon(\frac{L}{2} + \varepsilon t; m)$ solves the initial value problem (2.1a)–(2.1c) studied in Section 2, for $\alpha = c_\varepsilon$. In view of Theorem 2.2, since we are assuming $(1 - m)f(m) \leq 2c_\varepsilon$, the solution reaches the value s_ε in finite time, namely

$$\forall m \in (0, z_{c_\varepsilon}) \text{ with } (1 - m)f(m) \leq 2c_\varepsilon \quad \exists x_1 = x_1(m, \varepsilon) > \frac{L}{2} \text{ such that } v_\varepsilon(x_1; m) = s_\varepsilon.$$

Furthermore, we can estimate x_1 by applying Proposition 2.5 with $\eta = s_\varepsilon$: after a rescaling we find

$$x_1(m, \varepsilon) \leq \frac{L}{2} + \frac{C\varepsilon}{\sqrt{1-s_\varepsilon}}$$

where the constant $C > 0$ is independent of ε and m , since $\sup_\varepsilon c_\varepsilon < \frac{\sigma_\varepsilon}{2}$ (see (2.8)). By (1.7), up to reducing the value of ε_0 if necessary, we can therefore guarantee that

$$\frac{L}{2} \leq x_1(m, \varepsilon) \leq \frac{3}{4}L \quad (6.2)$$

for all $\varepsilon \in (0, \varepsilon_0)$ and for all $m \in (0, z_{c_\varepsilon})$ such that $(1-m)f(m) \leq 2c_\varepsilon$.

In the following argument we work with a fixed $\varepsilon \in (0, \varepsilon_0)$ and we study the family of solutions $\{v_\varepsilon(\cdot; m)\}$ depending on the parameter m . By Theorem 2.2 it also follows that v_ε is strictly increasing in $(\frac{L}{2}, x_1)$ with $v'_\varepsilon(x_1; m) > 0$. We let

$$x_2 = x_2(m, \varepsilon) := \sup\{x > x_1(m, \varepsilon) : v_\varepsilon(\cdot; m) < 1 \text{ and } v'_\varepsilon(\cdot; m) > 0 \text{ in } (x_1, x)\}$$

so that v_ε is strictly increasing in $(\frac{L}{2}, x_2)$. By multiplying (6.1) by v'_ε and integrating in $(\frac{L}{2}, x)$, for $x < x_2$, we find with a change of variables

$$\begin{aligned} \frac{\varepsilon}{2}(v'_\varepsilon(x))^2 &= \int_{\frac{L}{2}}^x \varepsilon v'_\varepsilon v''_\varepsilon dt = \int_{\frac{L}{2}}^x \left(\frac{c_\varepsilon^2 f'_\varepsilon(v_\varepsilon)}{f_\varepsilon^3(v_\varepsilon)} + \frac{v_\varepsilon - 1}{4\varepsilon} \right) v'_\varepsilon dt \\ &= \int_m^{v_\varepsilon(x)} \left(\frac{c_\varepsilon^2 f'_\varepsilon(s)}{f_\varepsilon^3(s)} + \frac{s-1}{4\varepsilon} \right) ds \\ &= \frac{c_\varepsilon^2}{2} \left(\frac{1}{f_\varepsilon^2(m)} - \frac{1}{f_\varepsilon^2(v_\varepsilon(x))} \right) + \frac{1}{8\varepsilon} ((1-v_\varepsilon(x))^2 - (1-m)^2), \end{aligned}$$

whence

$$(v'_\varepsilon)^2 = \frac{1}{\varepsilon^2} \left[\left(\frac{(1-v_\varepsilon)^2}{4} - \frac{\varepsilon c_\varepsilon^2}{f_\varepsilon^2(v_\varepsilon)} \right) - \left(\frac{(1-m)^2}{4} - \frac{c_\varepsilon^2}{f_\varepsilon^2(m)} \right) \right] =: H(v_\varepsilon; m, \varepsilon) \quad (6.3)$$

in $(\frac{L}{2}, x_2(m, \varepsilon))$. Let

$$i(\varepsilon) := \inf_{s \in (s_\varepsilon, 1)} \left(\frac{(1-s)^2}{4} - \frac{\varepsilon c_\varepsilon^2}{f_\varepsilon^2(s)} \right)$$

and notice for later use that

$$-\frac{\varepsilon c_\varepsilon^2}{f_\varepsilon^2(s_\varepsilon)} \leq i(\varepsilon) \leq -\varepsilon c_\varepsilon^2 < 0. \quad (6.4)$$

The map $m \mapsto \frac{(1-m)^2}{4} - \frac{c_\varepsilon^2}{f_\varepsilon^2(m)}$ is strictly increasing for $m \in (0, z_{c_\varepsilon})$ (by Proposition 2.1), tends to $-\infty$ as $m \rightarrow 0^+$ and vanishes if $(1-m)f(m) = 2c_\varepsilon$. Hence there exists a unique $\hat{m} = \hat{m}(\varepsilon) \in (0, z_{c_\varepsilon})$ such that $(1-\hat{m})f(\hat{m}) < 2c_\varepsilon$ and

$$i(\varepsilon) - \left(\frac{(1-\hat{m}(\varepsilon))^2}{4} - \frac{c_\varepsilon^2}{f_\varepsilon^2(\hat{m}(\varepsilon))} \right) = 0.$$

By the definition (6.3) of the function H it follows that

$$\inf_{s \in (s_\varepsilon, 1)} H(s; \hat{m}(\varepsilon), \varepsilon) = 0, \quad \inf_{s \in (s_\varepsilon, 1)} H(s; m, \varepsilon) > 0 \quad \text{for all } m \in (0, \hat{m}(\varepsilon)).$$

In turn, since $v_\varepsilon(x; m) \in (s_\varepsilon, 1)$ for $x \in (x_1(m, \varepsilon), x_2(m, \varepsilon))$, by (6.3) we have that

$$\inf_{x \in (x_1(m, \varepsilon), x_2(m, \varepsilon))} v'_\varepsilon(x; m) = \inf_{s \in (s_\varepsilon, 1)} \sqrt{H(s; m, \varepsilon)} > 0 \quad \text{for all } m \in (0, \hat{m}(\varepsilon)). \quad (6.5)$$

Then for $m \in (0, \hat{m}(\varepsilon))$ the solution $v_\varepsilon(\cdot; m)$ reaches the value 1 at the finite point $x_2(m, \varepsilon) \in (x_1(m, \varepsilon), +\infty)$.

Summing up, we have proved so far that for all $m \in (0, \hat{m}(\varepsilon))$ there exist two points $x_1(m, \varepsilon) \in (\frac{L}{2}, \frac{3}{4}L)$ and $x_2(m, \varepsilon) \in (x_1(m, \varepsilon), +\infty)$ such that

$$v_\varepsilon(x_1(m, \varepsilon); m) = s_\varepsilon, \quad v_\varepsilon(x_2(m, \varepsilon); m) = 1.$$

The goal is now to show the existence of a value $m_\varepsilon \in (0, \hat{m}(\varepsilon))$ such that $x_2(m_\varepsilon, \varepsilon) = L$.

By the continuous dependence of the solution to (6.1) on the initial value m , the point x_2 is a continuous function of m . We can write by (6.3)

$$\begin{aligned} x_2(m, \varepsilon) &= x_1(m, \varepsilon) + \int_{x_1(m, \varepsilon)}^{x_2(m, \varepsilon)} \frac{v'_\varepsilon(x; m)}{\sqrt{H(v_\varepsilon(x; m); m, \varepsilon)}} dx \\ &= x_1(m, \varepsilon) + \int_{s_\varepsilon}^1 \frac{ds}{\sqrt{H(s; m, \varepsilon)}} \\ &\leq \frac{3}{4}L + (1 - s_\varepsilon) \left(\inf_{s \in (s_\varepsilon, 1)} H(s; m, \varepsilon) \right)^{-\frac{1}{2}} \end{aligned} \quad (6.6)$$

and since $\lim_{m \rightarrow 0^+} \inf_{s \in (s_\varepsilon, 1)} H(s; m, \varepsilon) = +\infty$, we see that $x_2(m, \varepsilon) < L$ for all m sufficiently small. On the other hand, for $m = \hat{m}(\varepsilon)$ we have

$$\inf_{x \in (x_1(\hat{m}(\varepsilon), \varepsilon), x_2(\hat{m}(\varepsilon), \varepsilon))} v'_\varepsilon(x; \hat{m}(\varepsilon)) = \inf_{s \in (s_\varepsilon, 1)} \sqrt{H(s; \hat{m}(\varepsilon), \varepsilon)} = 0$$

and therefore $x_2(\hat{m}(\varepsilon), \varepsilon) = +\infty$, which implies $\lim_{m \rightarrow \hat{m}(\varepsilon)^-} x_2(m, \varepsilon) = +\infty$. By continuity of $m \mapsto x_2(m, \varepsilon)$, we conclude that there exists $m_\varepsilon \in (0, \hat{m}(\varepsilon))$ such that $x_2(m_\varepsilon, \varepsilon) = L$ and therefore $v_\varepsilon(L; m_\varepsilon) = 1$, as claimed.

We eventually prove the second part of the statement. By (6.6) it also follows that

$$\frac{1}{\varepsilon^2} \left[i(\varepsilon) - \left(\frac{(1 - m_\varepsilon)^2}{4} - \frac{c_\varepsilon^2}{f^2(m_\varepsilon)} \right) \right] = \inf_{s \in (s_\varepsilon, 1)} H(s; m_\varepsilon, \varepsilon) \leq \frac{16(1 - s_\varepsilon)^2}{L^2}.$$

By elementary manipulations in the previous inequality, and recalling that by construction $(1 - m_\varepsilon)f(m_\varepsilon) < 2c_\varepsilon$, we find

$$\frac{(1 - m_\varepsilon)^2}{4} \leq \frac{c_\varepsilon^2}{f^2(m_\varepsilon)} \leq \frac{16\varepsilon^2(1 - s_\varepsilon)^2}{L^2} - i(\varepsilon) + \frac{(1 - m_\varepsilon)^2}{4}. \quad (6.7)$$

Suppose that $\inf_\varepsilon c_\varepsilon > 0$. If by contradiction $m_\varepsilon \rightarrow 0$, then by passing to the limit as $\varepsilon \rightarrow 0$ in (6.7) we would have that the middle term would tend to $+\infty$, whereas the right-hand side would tend to $\frac{1}{4}$ (since $i(\varepsilon) \rightarrow 0$ by (6.4)). This contradiction proves that if $\inf_\varepsilon c_\varepsilon > 0$ then $\inf_\varepsilon m_\varepsilon > 0$.

Similarly, if $c_\varepsilon \rightarrow 0$ then by $(1 - m_\varepsilon)f(m_\varepsilon) < 2c_\varepsilon$ we must have $m_\varepsilon \rightarrow 0$. Again by passing to the limit as $\varepsilon \rightarrow 0$ in (6.7) we easily deduce that $\frac{2c_\varepsilon}{f(m_\varepsilon)} \rightarrow 1$. \square

Proof of Theorem 1.8. We divide the proof into three cases according to the form of the critical point \bar{u} , as in the statement of the theorem.

Case (i). Assume $\bar{u}(x) = \frac{a}{L}x$ for some $a > 0$. In this case it is sufficient to take $u_\varepsilon(x) = \frac{a}{L}x$ and $v_\varepsilon(x) \equiv 1$. Since $f'_\varepsilon(1) = \psi'_\varepsilon(1) = 0$, it is immediately checked that the pair $(u_\varepsilon, v_\varepsilon)$ is indeed a solution to (1.17a)–(1.17d).

Case (ii). Assume $\bar{u}(x) = c_0x + (a - c_0L)\chi_{(\frac{L}{2}, L)}(x)$ with $c_0 \in (0, \frac{\sigma\varepsilon}{2})$ and $g'(a - c_0L) = 2c_0$.

We apply Lemma 6.1 with $c_\varepsilon = c_0$ for all ε , to find values m_ε and functions v_ε solving (6.1) such that $v_\varepsilon(0) = v_\varepsilon(L) = 1$. Notice also that $m_0 := \inf_\varepsilon m_\varepsilon > 0$. We define

$$u_\varepsilon(x) := \int_0^x \frac{c_0}{f_\varepsilon^2(v_\varepsilon(x))} dx$$

and we obtain that $(u_\varepsilon, v_\varepsilon)$ is a family of critical points for \mathcal{F}_ε , i.e. they solve the system of equations (1.17a)–(1.17d) for $a_\varepsilon := \int_0^L \frac{c_0}{f_\varepsilon^2(v_\varepsilon)} dx$. To conclude, we need to show that $a_\varepsilon \rightarrow a$ and that $u_\varepsilon \rightarrow \bar{u}$ in $L^1([0, L])$.

To this aim, we first show that the equiboundedness of the energy (1.19) holds for the family $(u_\varepsilon, v_\varepsilon)$. By the construction in Lemma 6.1 the function v_ε obeys the equation (6.3) (with $c_\varepsilon = c_0$ and $m = m_\varepsilon$). Denoting by $H_\varepsilon(s) := H(s; m_\varepsilon, \varepsilon)$ the function appearing in (6.3), we have for all $s \in (m_\varepsilon, 1)$ after some elementary manipulations

$$\begin{aligned} H_\varepsilon(s) &= \frac{1}{4\varepsilon^2 f_\varepsilon^2(s)} [(1-s)^2 f_\varepsilon^2(s) - (1-m_\varepsilon)^2 f_\varepsilon^2(m_\varepsilon)] + \frac{f_\varepsilon^2(s) - f_\varepsilon^2(m_\varepsilon)}{4\varepsilon^2 f_\varepsilon^2(s)} \left(\frac{4c_0^2}{f_\varepsilon^2(m_\varepsilon)} - (1-m_\varepsilon)^2 \right) \\ &\geq \frac{1}{4\varepsilon^2 f_\varepsilon^2(s)} \left((1-s)^2 f_\varepsilon^2(s) - (1-m_\varepsilon)^2 f_\varepsilon^2(s) \right) \end{aligned}$$

where we used the monotonicity of the map f_ε and the fact that $(1-m_\varepsilon)f(m_\varepsilon) \leq 2c_0$ by the construction in Lemma 6.1. Then, denoting by $\tilde{f}(s) := (1-s)f(s)$ (which is strictly increasing by assumption (f3)), we have

$$\begin{aligned} \int_0^L f_\varepsilon^2(v_\varepsilon)(u'_\varepsilon)^2 dx &= c_0^2 \int_0^L \frac{dx}{f_\varepsilon^2(v_\varepsilon(x))} \leq c_0^2 \int_{\{v_\varepsilon \leq s_\varepsilon\}} \frac{|v'_\varepsilon|}{f_\varepsilon^2(v_\varepsilon)\sqrt{H_\varepsilon(v_\varepsilon)}} dx + \frac{c_0^2 |\{v_\varepsilon > s_\varepsilon\}|}{f_\varepsilon^2(s_\varepsilon)} \\ &\leq 2c_0^2 \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{f_\varepsilon^2(s)\sqrt{H_\varepsilon(s)}} + \frac{Lc_0^2}{f_\varepsilon^2(s_\varepsilon)} \\ &\leq 4c_0^2 \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{f(s)(\tilde{f}^2(s) - \tilde{f}^2(m_\varepsilon))^{1/2}} + \frac{Lc_0^2}{f_\varepsilon^2(s_\varepsilon)}. \end{aligned} \tag{6.8}$$

For $s \in (m_\varepsilon, s_\varepsilon)$ we write $\tilde{f}(s) - \tilde{f}(m_\varepsilon) = \tilde{f}'(\zeta_\varepsilon(s))(s - m_\varepsilon)$ for some point $\zeta_\varepsilon(s) \in (m_\varepsilon, s)$. Since $\zeta_\varepsilon(s) \geq m_\varepsilon \geq m_0 > 0$, in view of assumptions (f3) and (f5) we have that $\tilde{f}'(\zeta_\varepsilon(s)) \geq C(1 - \zeta_\varepsilon(s))^3 \geq C(1-s)^3$ for all $s \in (m_\varepsilon, s_\varepsilon)$ and for a constant C independent of ε . Therefore

$$\begin{aligned} \int_0^L f_\varepsilon^2(v_\varepsilon)(u'_\varepsilon)^2 dx &\leq \frac{4c_0^2}{\sqrt{2C\tilde{f}(m_\varepsilon)}} \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{f(s)\sqrt{s - m_\varepsilon}(1-s)^{3/2}} + \frac{Lc_0^2}{f_\varepsilon^2(s_\varepsilon)} \\ &\leq \frac{4c_0^2}{\sqrt{2C\tilde{f}(m_\varepsilon)}} \frac{1}{\tilde{f}(m_\varepsilon)} \int_{m_\varepsilon}^{s_\varepsilon} \frac{ds}{\sqrt{s - m_\varepsilon}\sqrt{1-s}} + \frac{Lc_0^2}{f_\varepsilon^2(s_\varepsilon)} \leq C' \end{aligned} \tag{6.9}$$

for another constant C' uniform in ε . By multiplying (6.1) by $(v_\varepsilon - 1)$ and integrating, after integration by parts we have

$$\begin{aligned} \int_0^L \left(\varepsilon (v'_\varepsilon)^2 + \frac{(1 - v_\varepsilon)^2}{4\varepsilon} \right) dx &= c_0^2 \int_0^L \frac{(1 - v_\varepsilon) f'_\varepsilon(v_\varepsilon)}{f_\varepsilon^3(v_\varepsilon)} dx \\ &\leq \sup_{s \in (m_\varepsilon, 1)} \left(\frac{(1 - s) f'_\varepsilon(s)}{f_\varepsilon(s)} \right) \int_0^L f_\varepsilon^2(v_\varepsilon) (u'_\varepsilon)^2 dx \leq C'' \end{aligned} \quad (6.10)$$

for a constant C'' independent of ε , where the last estimate follows by (6.9) and from the assumptions (f4), (ψ 1), (ψ 2), (ψ 3) and by (1.7) (recalling that $m_\varepsilon \geq m_0 > 0$).

Combining (6.9) and (6.10) we obtain that $\sup_\varepsilon \mathcal{F}_\varepsilon(u_\varepsilon, v_\varepsilon) < +\infty$. The critical points $(u_\varepsilon, v_\varepsilon)$ then satisfy the assumption of Theorem 1.7. Since $\lim_{\varepsilon \rightarrow 0} m_\varepsilon \in (0, 1)$, we are in case (ii) and we can conclude that up to extraction of a subsequence $u_{\varepsilon_k} \rightarrow u_0$ in $L^1([0, L])$, where $u_0(x) = c_0 x + (a_0 - c_0 L) \chi_{(\frac{L}{2}, L)}(x)$, $a_0 = \lim_k a_{\varepsilon_k}$, and $g'(a_0 - c_0 L) = 2c_0$. Since we also have $g'(a - c_0 L) = 2c_0$ and g' is injective in $(0, s_{\text{frac}})$ by Proposition 3.6, we conclude that $a_0 = a$ and $u_0 = \bar{u}$. Hence, as the limit of any subsequence of u_ε converges to \bar{u} , we conclude that $u_\varepsilon \rightarrow \bar{u}$ in $L^1([0, L])$.

Case (iii). Assume that s_{frac} is finite (i.e. $f'(0) > 0$ by Proposition 3.7) and that $\bar{u}(x) = a \chi_{(\frac{L}{2}, L)}(x)$ with $a = s_{\text{frac}}$.

Take any sequence $c_\varepsilon \rightarrow 0$, with $c_\varepsilon > 0$, and apply Lemma 6.1 to find values m_ε and functions v_ε solving (6.1) such that $v_\varepsilon(0) = v_\varepsilon(L) = 1$. Notice also that $m_\varepsilon \rightarrow 0$ and $\frac{2c_\varepsilon}{f(m_\varepsilon)} \rightarrow 1$. We define

$$u_\varepsilon(x) := \int_0^x \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon(x))} dx$$

and we obtain that $(u_\varepsilon, v_\varepsilon)$ is a family of critical points for \mathcal{F}_ε , i.e. they solve the system of equations (1.17a)–(1.17d) for $a_\varepsilon := \int_0^L \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} dx$. To conclude, we need to show that $a_\varepsilon \rightarrow s_{\text{frac}}$ and that $u_\varepsilon \rightarrow \bar{u}$ in $L^1([0, L])$.

As in the previous step, we first show that the equiboundedness of the energy (1.19) holds for the family $(u_\varepsilon, v_\varepsilon)$. We indeed have, similarly to (6.8), for $\delta > 0$,

$$\begin{aligned} a_\varepsilon &= \int_0^L \frac{c_\varepsilon}{f_\varepsilon^2(v_\varepsilon)} dx \leq 4c_\varepsilon \int_{m_\varepsilon}^\delta \frac{ds}{f(s) (\tilde{f}^2(s) - \tilde{f}^2(m_\varepsilon))^{1/2}} + \frac{4c_\varepsilon(1 - \delta)}{f(\delta) (\tilde{f}^2(\delta) - \tilde{f}^2(m_\varepsilon))^{1/2}} + \frac{Lc_\varepsilon}{f_\varepsilon^2(s_\varepsilon)} \\ &\leq 4c_\varepsilon \sup_{s \in (m_\varepsilon, \delta)} \left(\frac{1 - s}{\tilde{f}'(s)} \right) \int_{m_\varepsilon}^\delta \frac{\tilde{f}'(s) ds}{\tilde{f}(s) (\tilde{f}^2(s) - \tilde{f}^2(m_\varepsilon))^{1/2}} + C_\delta c_\varepsilon \\ &= 4c_\varepsilon \sup_{s \in (m_\varepsilon, \delta)} \left(\frac{1 - s}{\tilde{f}'(s)} \right) \int_{\tilde{f}(m_\varepsilon)}^{\tilde{f}(\delta)} \frac{dt}{t (t^2 - \tilde{f}^2(m_\varepsilon))^{1/2}} + C_\delta c_\varepsilon \\ &= \frac{4c_\varepsilon}{\tilde{f}(m_\varepsilon)} \sup_{s \in (m_\varepsilon, \delta)} \left(\frac{1 - s}{\tilde{f}'(s)} \right) \arctan \left(\frac{(\tilde{f}^2(\delta) - \tilde{f}^2(m_\varepsilon))^{1/2}}{\tilde{f}(m_\varepsilon)} \right) + C_\delta c_\varepsilon, \end{aligned}$$

where C_δ is a constant depending on δ , for ε is small enough. Now, using the fact that $c_\varepsilon \rightarrow 0$, $m_\varepsilon \rightarrow 0$, $f'(0) > 0$ and that $\frac{2c_\varepsilon}{f(m_\varepsilon)} \rightarrow 1$, one can see that the right-hand side in the previous

chain of inequalities is uniformly bounded. Therefore

$$\sup_{\varepsilon} a_{\varepsilon} < +\infty, \quad \lim_{\varepsilon \rightarrow 0} \int_0^L f_{\varepsilon}^2(v_{\varepsilon})(u'_{\varepsilon})^2 dx = \lim_{\varepsilon \rightarrow 0} c_{\varepsilon} a_{\varepsilon} = 0. \quad (6.11)$$

Coming to the energy of v_{ε} , we fix $\delta \in (0, 1)$ and as in (6.10) we have

$$\begin{aligned} \int_0^L \left(\varepsilon(v'_{\varepsilon})^2 + \frac{(1-v_{\varepsilon})^2}{4\varepsilon} \right) dx &= c_{\varepsilon}^2 \int_0^L \frac{(1-v_{\varepsilon})f'_{\varepsilon}(v_{\varepsilon})}{f_{\varepsilon}^3(v_{\varepsilon})} dx \\ &\leq \frac{c_{\varepsilon}}{f(m_{\varepsilon})} \left(\sup_{s \in (m_{\varepsilon}, \delta)} f'(s) \right) \int_{\{v_{\varepsilon} \leq \delta\}} \frac{c_{\varepsilon}}{f_{\varepsilon}^2(v_{\varepsilon})} dx \\ &\quad + \sup_{s \in (\delta, 1)} \left(\frac{(1-s)f'_{\varepsilon}(s)}{f_{\varepsilon}(s)} \right) \int_{\{v_{\varepsilon} > \delta\}} f_{\varepsilon}^2(v_{\varepsilon})(u'_{\varepsilon})^2 dx. \end{aligned}$$

Again by (6.11), by $\frac{2c_{\varepsilon}}{f(m_{\varepsilon})} \rightarrow 1$, and by all the assumptions on f_{ε} , it is possible to check that the previous quantities are uniformly bounded with respect to ε . Hence $\sup_{\varepsilon} \mathcal{F}_{\varepsilon}(u_{\varepsilon}, v_{\varepsilon}) < +\infty$.

The critical points $(u_{\varepsilon}, v_{\varepsilon})$ then satisfy the assumption of Theorem 1.7. Since $\lim_{\varepsilon \rightarrow 0} m_{\varepsilon} = 0$, we are in case (iii) and, as at the end of the previous step, we can conclude that $u_{\varepsilon} \rightarrow \bar{u}$ in $L^1([0, L])$ and $a = s_{\text{frac}}$. \square

Acknowledgments. The authors are thankful to Cinzia Soresina for fruitful conversations. MB is member of the GNAMPA group of the Istituto Nazionale di Alta Matematica (INdAM).

REFERENCES

- [1] R. ALESSI, J.-J. MARIGO, AND S. VIDOLI, *Gradient damage models coupled with plasticity and nucleation of cohesive cracks*, Arch. Ration. Mech. Anal., 214 (2014), pp. 575–615.
- [2] R. ALICANDRO, A. BRAIDES, AND J. SHAH, *Free-discontinuity problems via functionals involving the L^1 -norm of the gradient and their approximations*, Interfaces Free Bound., 1 (1999), pp. 17–37.
- [3] R. ALICANDRO AND M. FOCARDI, *Variational approximation of free-discontinuity energies with linear growth*, Commun. Contemp. Math., 4 (2002), pp. 685–723.
- [4] S. ALMI, *Energy release rate and quasi-static evolution via vanishing viscosity in a fracture model depending on the crack opening*, ESAIM Control Optim. Calc. Var., 23 (2017), pp. 791–826.
- [5] S. ALMI, S. BELZ, AND M. NEGRI, *Convergence of discrete and continuous unilateral flows for Ambrosio-Tortorelli energies and application to mechanics*, ESAIM Math. Model. Numer. Anal., 53 (2019), pp. 659–699.
- [6] L. AMBROSIO AND V. M. TORTORELLI, *Approximation of functionals depending on jumps by elliptic functionals via Γ -convergence*, Comm. Pure Appl. Math., 43 (1990), pp. 999–1036.
- [7] L. AMBROSIO AND V. M. TORTORELLI, *On the approximation of free discontinuity problems*, Boll. Un. Mat. Ital. B (7), 6 (1992), pp. 105–123.
- [8] M. ARTINA, F. CAGNETTI, M. FORNASIER, AND F. SOLOMBRINO, *Linearly constrained evolutions of critical points and an application to cohesive fractures*, Math. Models Methods Appl. Sci., 27 (2017), pp. 231–290.
- [9] J.-F. BABADJIAN AND V. MILLOT, *Unilateral gradient flow of the Ambrosio-Tortorelli functional by minimizing movements*, Ann. Inst. H. Poincaré C Anal. Non Linéaire, 31 (2014), pp. 779–822.
- [10] J.-F. BABADJIAN, V. MILLOT, AND R. RODIAC, *A note on the one-dimensional critical points of the Ambrosio-Tortorelli functional*, Preprint, (2022).
- [11] ———, *On the convergence of critical points of the Ambrosio-Tortorelli functional*, Preprint, (2022).
- [12] M. BONACINI, S. CONTI, AND F. IURLANO, *Cohesive fracture in 1D: quasi-static evolution and derivation from static phase-field models*, Arch. Ration. Mech. Anal., 239 (2021), pp. 1501–1576.
- [13] E. BONETTI, C. CAVATERRA, F. FREDDI, AND F. RIVA, *On a phase-field model of damage for hybrid laminates with cohesive interface*, Math. Methods Appl. Sci., 45 (2022), pp. 3520–3553.

- [14] G. BOUCHITTÉ, A. BRAIDES, AND G. BUTTAZZO, *Relaxation results for some free discontinuity problems*, J. Reine Angew. Math., 458 (1995), pp. 1–18.
- [15] B. BOURDIN, *Image segmentation with a finite element method*, M2AN Math. Model. Numer. Anal., 33 (1999), pp. 229–244.
- [16] B. BOURDIN, G. A. FRANCFORT, AND J.-J. MARIGO, *The variational approach to fracture*, J. Elasticity, 91 (2008), pp. 5–148.
- [17] A. BRAIDES, G. DAL MASO, AND A. GARRONI, *Variational formulation of softening phenomena in fracture mechanics: the one-dimensional case*, Arch. Ration. Mech. Anal., 146 (1999), pp. 23–58.
- [18] L. CAFFARELLI, F. CAGNETTI, AND A. FIGALLI, *Optimal regularity and structure of the free boundary for minimizers in cohesive zone models*, Arch. Ration. Mech. Anal., 237 (2020), pp. 299–345.
- [19] F. CAGNETTI, *A vanishing viscosity approach to fracture growth in a cohesive zone model with prescribed crack path*, Math. Models Methods Appl. Sci., 18 (2008), pp. 1027–1071.
- [20] F. CAGNETTI AND R. TOADER, *Quasistatic crack evolution for a cohesive zone model with different response to loading and unloading: a Young measures approach*, ESAIM Control Optim. Calc. Var., 17 (2011), pp. 1–27.
- [21] S. CONTI, M. FOCARDI, AND F. IURLANO, *Phase field approximation of cohesive fracture models*, Ann. Inst. H. Poincaré C Anal. Non Linéaire, 33 (2016), pp. 1033–1067.
- [22] ———, *Phase-field approximation of a vectorial, geometrically nonlinear cohesive fracture energy*, Preprint arXiv:2205.06541, (2022).
- [23] V. CRISMALE, G. LAZZARONI, AND G. ORLANDO, *Cohesive fracture with irreversibility: quasistatic evolution for a model subject to fatigue*, Math. Models Methods Appl. Sci., 28 (2018), pp. 1371–1412.
- [24] G. DAL MASO AND A. GARRONI, *Gradient bounds for minimizers of free discontinuity problems related to cohesive zone models in fracture mechanics*, Calc. Var. Partial Differential Equations, 31 (2008), pp. 137–145.
- [25] G. DAL MASO, G. ORLANDO, AND R. TOADER, *Fracture models for elasto-plastic materials as limits of gradient damage models coupled with plasticity: the antiplane case*, Calc. Var. Partial Differential Equations, 55 (2016), pp. Art. 45, 39.
- [26] G. DAL MASO AND C. ZANINI, *Quasi-static crack growth for a cohesive zone model with prescribed crack path*, Proc. Roy. Soc. Edinburgh Sect. A, 137 (2007), pp. 253–279.
- [27] G. DEL PIERO, *A variational approach to fracture and other inelastic phenomena*, J. Elasticity, 112 (2013), pp. 3–77.
- [28] G. DEL PIERO AND L. TRUSKINOVSKY, *A one-dimensional model for localized and distributed failure*, J. Phys. IV France, 8 (1998).
- [29] ———, *Elastic bars with cohesive energy*, Contin. Mech. Thermodyn., 21 (2009), pp. 141–171.
- [30] G. A. FRANCFORT, N. Q. LE, AND S. SERFATY, *Critical points of Ambrosio-Tortorelli converge to critical points of Mumford-Shah in the one-dimensional Dirichlet case*, ESAIM Control Optim. Calc. Var., 15 (2009), pp. 576–598.
- [31] G. A. FRANCFORT AND J.-J. MARIGO, *Revisiting brittle fracture as an energy minimization problem*, J. Mech. Phys. Solids, 46 (1998), pp. 1319–1342.
- [32] F. FREDDI AND F. IURLANO, *Numerical insight of a variational smeared approach to cohesive fracture*, J. Mech. Phys. Solids, 98 (2017), pp. 156–171.
- [33] A. GIACOMINI, *Ambrosio-Tortorelli approximation of quasi-static evolution of brittle fractures*, Calc. Var. Partial Differential Equations, 22 (2005), pp. 129–172.
- [34] J. E. HUTCHINSON AND Y. TONEGAWA, *Convergence of phase interfaces in the van der Waals-Cahn-Hilliard theory*, Calc. Var. Partial Differential Equations, 10 (2000), pp. 49–84.
- [35] H. LAMMEN, S. CONTI, AND J. MOSLER, *A finite deformation phase field model suitable for cohesive fracture*, J. Mech. Phys. Solids, 178 (2023), pp. Paper No. 105349, 17.
- [36] C. J. LARSEN AND V. SLASTIKOV, *Dynamic cohesive fracture: models and analysis*, Math. Models Methods Appl. Sci., 24 (2014), pp. 1857–1875.
- [37] N. Q. LE, *Convergence results for critical points of the one-dimensional Ambrosio-Tortorelli functional with fidelity term*, Adv. Differential Equations, 15 (2010), pp. 255–282.
- [38] N. Q. LE AND P. J. STERNBERG, *Asymptotic behavior of Allen-Cahn-type energies and Neumann eigenvalues via inner variations*, Ann. Mat. Pura Appl. (4), 198 (2019), pp. 1257–1293.
- [39] S. LUCKHAUS AND L. MODICA, *The Gibbs-Thompson relation within the gradient theory of phase transitions*, Arch. Rational Mech. Anal., 107 (1989), pp. 71–83.

- [40] L. MODICA, *The gradient theory of phase transitions and the minimal interface criterion*, Arch. Rational Mech. Anal., 98 (1987), pp. 123–142.
- [41] M. NEGRI AND R. SCALA, *A quasi-static evolution generated by local energy minimizers for an elastic material with a cohesive interface*, Nonlinear Anal. Real World Appl., 38 (2017), pp. 271–305.
- [42] M. NEGRI AND R. SCALA, *Existence, energy identity, and higher time regularity of solutions to a dynamic viscoelastic cohesive interface model*, SIAM J. Math. Anal., 53 (2021), pp. 5682–5730.
- [43] P. PADILLA AND Y. TONEGAWA, *On the convergence of stable phase transitions*, Comm. Pure Appl. Math., 51 (1998), pp. 551–579.
- [44] F. RIVA, *Energetic evolutions for linearly elastic plates with cohesive slip*, Nonlinear Anal. Real World Appl., 74 (2023), pp. Paper No. 103934, 27.
- [45] M. RÖGER AND Y. TONEGAWA, *Convergence of phase-field approximations to the Gibbs-Thomson law*, Calc. Var. Partial Differential Equations, 32 (2008), pp. 111–136.
- [46] T. ROUBÍČEK, *A general thermodynamical model for adhesive frictional contacts between viscoelastic or poro-viscoelastic bodies at small strains*, Interfaces Free Bound., 21 (2019), pp. 169–198.
- [47] M. THOMAS, *A comparison of delamination models: modeling, properties, and applications*, in Mathematical analysis of continuum mechanics and industrial applications. II, vol. 30 of Math. Ind. (Tokyo), Springer, Singapore, 2018, pp. 27–38.
- [48] M. THOMAS AND C. ZANINI, *Cohesive zone-type delamination in visco-elasticity*, Discrete Contin. Dyn. Syst. Ser. S, 10 (2017), pp. 1487–1517.
- [49] Y. TONEGAWA, *Phase field model with a variable chemical potential*, Proc. Roy. Soc. Edinburgh Sect. A, 132 (2002), pp. 993–1019.
- [50] ———, *A diffused interface whose chemical potential lies in a Sobolev space*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 4 (2005), pp. 487–510.
- [51] J.-Y. WU, *A unified phase-field theory for the mechanics of damage and quasi-brittle failure*, J. Mech. Phys. Solids, 103 (2017), pp. 72–99.
- [52] J.-Y. WU AND V. P. NGUYEN, *A length scale insensitive phase-field damage model for brittle fracture*, J. Mech. Phys. Solids, 119 (2018), pp. 20–42.

(Marco Bonacini) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, ITALY
E-mail address: marco.bonacini@unitn.it

(Flaviana Iurlano) SORBONNE UNIVERSITÉ, CNRS, UNIVERSITÉ DE PARIS, LABORATOIRE JACQUES-LOUIS LIONS, 75005 PARIS, FRANCE
E-mail address: iurlano@ljl1.math.upmc.fr