

AN OPTIMAL DESIGN PROBLEM FOR A CHARGE QUBIT

DARIO MAZZOLENI, CYRILL B. MURATOV, AND BERARDO RUFFINI

ABSTRACT. In this paper we introduce a simple variational model describing the ground state of a superconducting charge qubit. The model gives rise to a shape optimization problem that aims at maximizing the number of qubit states at a given gating voltage. We show that for small values of the charge optimal shapes exist and are $C^{2,\alpha}$ -nearly spherical sets. In contrast, for large values of the charge the optimal shape does not exist, with the energy favoring disjoint collections of sets.

CONTENTS

1. Introduction	1
1.1. Main results and detailed strategy of their proof	3
2. Preliminaries	4
2.1. Minimization of $E_q(u, \Omega)$ in u .	5
2.2. Fraenkel asymmetry and quantitative Faber-Krahn inequality	7
2.3. Some facts about quasi-open sets	7
3. An existence result for an auxiliary problem	8
3.1. Removing the L^2 norm constraint in E_q	8
3.2. Another auxiliary problem: removing the volume constraint	9
3.3. Free boundary formulation	11
3.4. Equivalence between the minimizations of E_q and $E_{q,M,\eta}$	17
4. Optimality conditions and improvement of flatness	21
4.1. The structure of μ_u : blow up limits	22
5. The surgery result and the proof of Theorem 1.3	24
6. Non-existence of minimizers for q large: proof of Theorem 1.6	29
Appendix A. The physical model and non-dimensionalization	32
References	32

1. INTRODUCTION

This paper studies the shape optimization problem associated with the energy functional

$$(1) \quad E_q(\Omega) := \inf_{u \in H_0^1(\Omega)} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \frac{q}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x)|^2 |u(y)|^2}{|x-y|} dx dy : \int_{\Omega} |u(x)|^2 dx = 1 \right),$$

for a fixed parameter $q > 0$ referred to as “charge” from now on, among all measurable sets $\Omega \subset \mathbb{R}^3$ with prescribed volume. With $q = 0$, this problem is the classical shape optimization problem for the first Dirichlet eigenvalue of the Laplacian, whose solution is known to be a ball (see [24] for an overview). The Dirichlet energy, together with a volume constraint on Ω , acts as a cohesive term that forces the perimeter of the set to be minimized. At the same time the second term in the definition of the energy is the Coulombic repulsive energy for $q > 0$, favoring separation of charges and domain splitting. Thus the considered problem falls into a general class of geometric variational problems with competing attractive and repulsive interactions that received a significant attention recently (for a broad overview, see [14]). It is the competition of these two

D. Mazzoleni and B. Ruffini have been partially supported by the MUR via PRIN project P2022R537CS. D. Mazzoleni has been partially supported by the MUR via PRIN project P2020F3NCPX. B. Ruffini was also supported by the project ANR-18-CE40-0013 SHAPO financed by the French Agence Nationale de la Recherche. C. B. Muratov was partially supported by NSF via grant DMS-1908709 and MUR via PRIN 2022 PNRR project P2022WJW9H. C. B. Muratov also acknowledges the MUR Excellence Department Project awarded to the Department of Mathematics, University of Pisa, CUP I57G22000700001. The authors are members of INdAM-GNAMPA. Finally, the authors are grateful to A. Bernard-Mantel for multiple enlightening discussions on the underlying physics of the problem and warmly thank B. Velichkov for suggesting a strategy for the proof of Lemma 3.11.

interactions that makes the behavior of such problems highly non-trivial and interesting, and also makes their analysis challenging.

The choice of the particular energy functional in (1) is motivated by a basic model in quantum mechanics introduced by Hartree in 1927 [22]. We note that although this model was originally meant to describe the behavior of electrons in an atom, it suffers from a deficiency due to its lack of accounting for the fermionic nature of the electrons and in that context was superseded by a more appropriate Hartree-Fock model (see [31, 29] for a mathematical discussion of these models). Nevertheless, the so-called restricted Hartree equation

$$-\Delta u + Vu + \left(|u|^2 \star \frac{q}{|x|} \right) u = \varepsilon_q u \quad \text{in } H^1(\mathbb{R}^3; \mathbb{C}),$$

where V is the external potential, ε_q is the lowest energy level and “ \star ” denotes a convolution, together with its associated energy functional naturally re-emerge in the context of superconductors, in which the variable u stands for the self-consistent ground state wave function of the Cooper pairs in the Bose-Einstein condensate at zero temperature. Thus, we interpret the energy in (1) as a simple model for the ground state energy of the Cooper pairs confined to a nanoscale superconducting island Ω that is embedded into an insulator ($V = 0$ in Ω and $V = +\infty$ in Ω^c) [4]. The obtained energy is a single-orbital Hartree functional for bosons, with the total number of condensate particles proportional to $q > 0$. By the well-known property of the single-orbital Schrödinger operator, the function u may be chosen to be real-valued and nonnegative [30].

A Cooper pair box, or a charge qubit, is an example of a quantum bit device that uses the charge states of the Cooper pairs in a superconducting nanoscale island to represent quantum information [13, 6, 39, 26]. Ordinarily, a charge qubit is treated simply as a capacitor, with its quantum state corresponding to the discrete number of charges on the capacitor. Therefore, the shape of the island enters into the consideration solely through the value of the island’s capacitance. At the same time, as the island’s dimensions become smaller, as well as in the presence of a high dielectric constant matrix, an interplay between the kinetic and the potential energies of the Cooper pairs may become notable, making the dependence of the system’s characteristics on the island shape less straightforward. A natural question one may then ask is whether one could take advantage of the superconducting island’s shape to optimize some characteristics of the charge qubit. For example, one may ask what shape of the island at a given volume would maximize the number of stable charge states at a given gate voltage (i.e., for a given $V = V_0 < 0$ fixed in Ω), which is equivalent to minimizing the energy in (1) among all such domains. This is precisely the mathematical problem treated in the present paper.

As it is our interest to optimize $E_q(\Omega)$ with respect to Ω , we consider the optimal design problem

$$(2) \quad \inf_{\substack{\Omega \subset \mathbb{R}^3 \\ |\Omega| = |B_1|}} E_q(\Omega) = \inf_{\substack{\Omega \subset \mathbb{R}^3 \\ |\Omega| = |B_1|}} \inf_{u \in H_0^1(\Omega)} \left\{ E_q(u, \Omega) : \int_{\Omega} u^2 dx = 1 \right\},$$

where

$$(3) \quad E_q(u, \Omega) := \int_{\Omega} |\nabla u(x)|^2 dx + \frac{q}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x)|^2 |u(y)|^2}{|x - y|} dx dy.$$

Notice that due to the scaling properties of the energy functional in (3) any choice of the volume of Ω may be reduced to that of $|\Omega| = |B_1| = \frac{4}{3}\pi$ in (2) by redefining q . By the same reason all the physical constants in the problem may be absorbed in the value of the dimensionless constant q , which is thus the only non-trivial parameter of the model (see Appendix A for details).

Remark 1.1 (The shape optimization viewpoint). From a purely mathematical point of view, problem (2) is a shape optimization problem in which the functional to be optimized is a nonlocal, nonlinear perturbation of the first eigenvalue of the Dirichlet Laplacian. The nonlocal nature of the perturbation, even under some very strong regularity assumptions on the state variable u or the set Ω drastically prevents adapting standard shape optimization techniques. In particular, to our knowledge no second-order shape derivatives or symmetrization results are available in the literature for such functionals, implying that even local rigidity of critical points is in principle a highly nontrivial question.

Remark 1.2. Again, from a mathematical perspective it might be of some interest to consider the more general energy

$$E_q^{N,\alpha}(u, \Omega) := \int_{\Omega} |\nabla u|^2 dx + \frac{q}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x)|^2 |u(y)|^2}{|x - y|^{N-\alpha}} dx dy, \quad u \in H_0^1(\Omega),$$

where $u : \Omega \subset \mathbb{R}^N \rightarrow \mathbb{R}$ and $\alpha \in (0, N)$. For the sake of presentation we stick to the physically relevant case $N = 3$, $\alpha = 2$, but we stress that the strategy contained in this paper could be adapted in the more general case $N \geq 3$ and $\alpha \in (0, N)$.

1.1. Main results and detailed strategy of their proof. The first main result of the paper is the following:

Theorem 1.3. *For all $\varepsilon > 0$ there exists $q^* = q^*(\varepsilon) > 0$ and $\alpha \in (0, 1)$ such that, for all $0 < q \leq q^*$, there exists an optimal set for problem (2). Furthermore, every optimal set Ω is $C^{2,\alpha}$ -nearly spherical, namely, there is a function $\varphi_\varepsilon: \partial B_1 \rightarrow \mathbb{R}$ of class $C^{2,\alpha}$ such that $\|\varphi_\varepsilon\|_{C^{2,\alpha}} \leq \varepsilon$ and*

$$\partial\Omega = \left\{ (1 + \varphi_\varepsilon(x))x : x \in \partial B_1 \right\}.$$

Remark 1.4. It is natural to conjecture that for sufficiently small q problem (2) admits a unique minimizer: the ball. A natural strategy to prove that conjecture would be to develop a quantified second order shape derivative of the energy. To our knowledge, no second order shape derivative for such a nonlocal energy is available at present, and providing one appears to be quite challenging.

The strategy of the proof of Theorem 1.3 is quite involved, hence we offer an outline here. Its overall structure follows ideas developed in [28, 27, 18, 19, 20, 36, 37] for isoperimetric-type variational problems and developed later on in [35, 7] for spectral optimization problems (this list is not exhaustive). In our case it can be summarized, broadly speaking, in three main steps.

- (1) Prove existence of an optimal set among equibounded sets.
- (2) Prove regularity of the optimal shapes: any minimizer previously found is a nearly spherical set.
- (3) Remove the equiboundedness hypothesis.

Point (3) is almost independent and is dealt with in Section 5. It is in fact an improvement of spectral surgery techniques developed in [34, 35], allowing to reduce to the uniformly bounded setting. The technique nevertheless works on connected sets, hence its proof relies on some mild regularity that we have to show in the previous sections.

The core of the proof is contained in points (1) and (2). Their proof, performed in the (quite long) Section 3, is convoluted. Since its technical details may hide the ideas behind it, we describe it here.

- First we restrict ourselves to considering the problem for equibounded sets, that is, sets uniformly contained in a large enough ball B_R , $R \gg 1$. This greatly simplifies the ensuing compactness arguments, and does not lead to a loss of generality in view of point (3).
- Then we show the existence of a minimizer. In order to do that, we transform the energy E_q into a new equivalent energy $E_{q,M}$. The related ground state energy is such that the L^2 constraint on the density function u is replaced by a Lagrange multiplier¹, see formula (12).
- Next we eliminate the volume constraint on the admissible sets: following ideas from [1, 7, 35] we consider a new energy $\Omega \mapsto E_{q,M,\eta}(\Omega)$ where the volume constraint on Ω is replaced by a Lagrange multiplier (more precisely we need to use a penalizing piecewise linear function $\eta \mapsto f_\eta(|\Omega|)$, because of the different scalings of the addends in $E_{q,M}$). We are now able to show that a minimizer for $E_{q,M,\eta}$ exists, and it has finite perimeter. Such a property plays a crucial role later on in the proof.

A major complication arises now: at this stage we are not able to show that the problems of minimizing $E_{q,M}$ under volume constraint and the unconstrained minimization of $E_{q,M,\eta}$ are equivalent. To establish this, we hence need to show some deeper regularity results on the minimizers of the latter energy. To that end:

- We consider an equivalent *free boundary* formulation of the problem, where the minimization in Ω is replaced by the minimization of an energy functional $\mathcal{E}_{q,M,\eta}(u)$ for $u \in H_0^1(B_R)$. Every optimal Ω happens to be the support of an optimal function for $\mathcal{E}_{q,M,\eta}$.
- We then adapt techniques from the free boundary regularity theory to show that any minimizer of $\mathcal{E}_{q,M,\eta}$ is Lipschitz continuous and nondegenerate, thus its positivity set is open and satisfies density estimates from below and above. Note that here a major technical, but crucial, point is rendering all the implicit constants independent of q .
- With such a regularity at hand, by exploiting a quantitative version of the Faber–Krahn inequality we can prove that for q small minimizers of $E_{M,q,\eta}$ are close to a ball in the L^1 and in the Hausdorff distance. This is just enough to show the equivalence of the (minimization of) $E_{q,M}$ and $E_{q,M,\eta}$.
- Eventually, thanks to the fact that minimizers have finite perimeter, by means of a shape variation analysis and an improvement of flatness, we show that minimizers are in fact $C^{2,\alpha}$ -regular.

¹We will transform the energy two more times, for a total of four (equivalent) energies! This appears quite cumbersome, yet we are not able to reduce these complications.

Remark 1.5. It is quite natural to suppose that the closeness of minimizers to a ball should directly imply that the equiboundedness constraint might be removed. This would simplify the above general strategy. Nevertheless we are not able to do that as all the regularity estimates (in particular the crucial density estimates) do depend on R , with constants diverging as $R \rightarrow +\infty$.

The second main result of the paper is a counterpart of Theorem 1.3 when q is large. We show that in this case problem (2) has no minimizers.

Theorem 1.6. *There exists a universal $\bar{q} > 0$ such that if $q > \bar{q}$ then problem (2) has no solution.*

The proof of this latter result follows an elegant argument proposed in [32] to show non-existence of ground states of the Thomas-Fermi-Dirac-Von Weizsäcker energy. Failure of existence in this regime is due to the tendency of the minimizing sequences of $E_q(\Omega)$ to split into disjoint sets, which is common for variational problems with competing interactions [14]. Here we generalize the argument of [32] to our situation, in which there is no a priori L^∞ control on the minimizer u of $E_q(u, \Omega)$ for a given Ω , which is of independent interest. Thus some shape optimization arguments are also needed to conclude the proof.

Plan of the paper. In Section 2 we show some preliminary results on the nature of the energy $E_q(\cdot, \Omega)$ and in particular we study its Euler-Lagrange equation. In doing so we investigate the nonlinear and nonlocal eigenvalue problem associated to the Euler-Lagrange equation of the energy. Then we recall some notions about quasi-open sets and the quantitative Faber-Krahn inequality. Section 3 is devoted to the proof of Theorem 1.3 in the equibounded setting (namely Theorem 3.1). More precisely, we introduce a new functional without the measure constraint, then prove existence of minimizers and their mild regularity properties (using a free boundary formulation), and finally we show the equivalence with problem (2) in an equibounded setting. In Section 4, we first show what is the optimality condition at the free boundary and then prove that minimizers are $C^{2,\alpha}$ -nearly spherical, employing free boundary regularity techniques. Section 5 is devoted to a surgery argument which allows to conclude the proof of Theorem 1.3. Finally, in Section 6 we prove Theorem 1.6 concerning the case of nonexistence of minimizers.

2. PRELIMINARIES

We present here some basic properties of the functionals that will be used in the proofs. Throughout the paper, we adopt the following notations: for $\Omega \subset \mathbb{R}^3$ a bounded open set and $u \in H_0^1(\Omega)$ denoting with \star the usual convolution, we let

$$v_u(x) = u^2(x) \star \frac{1}{|x|} = \int_{\Omega} \frac{u^2(y)}{|x-y|} dy,$$

with $v_u \in W_{\text{loc}}^{2,3}(\mathbb{R}^3)$ by [17, Theorem 9.9] and Sobolev embedding. We begin by establishing a uniform bound on v_u .

Lemma 2.1. *Let $\bar{x} \in \mathbb{R}^3$, let $\Omega \subset \mathbb{R}^3$ be a bounded open set, and let $\varphi \in H_0^1(\Omega)$. Then*

$$\int_{\Omega} \frac{\varphi^2(x)}{|x-\bar{x}|} dx \leq 2\|\varphi\|_{L^2(\Omega)} \|\nabla\varphi\|_{L^2(\Omega)}.$$

Proof. Up to extending with the value $\varphi(x) = 0$ for $x \in \mathbb{R}^3 \setminus \Omega$, we can suppose $\varphi \in H_0^1(\mathbb{R}^3)$. By Hölder inequality we obtain

$$\int \frac{\varphi^2(x)}{|x-\bar{x}|} dx \leq \left\| \frac{\varphi(\cdot)}{|\cdot-\bar{x}|} \right\|_{L^2(\mathbb{R}^3)} \|\varphi\|_{L^2(\mathbb{R}^3)}$$

Then we can apply the classical Hardy-Sobolev inequality, see for instance [33, Corollary 2 of Section 2.1.7]

$$\left\| \frac{\varphi}{|\cdot-\bar{x}|} \right\|_{L^2(\mathbb{R}^3)} \leq 2\|\nabla\varphi\|_{L^2(\mathbb{R}^3)},$$

to obtain the result. \square

We recall now that for $\phi, \psi : \Omega \rightarrow \mathbb{R}$ the Coulomb energy denoted by

$$D(\phi, \psi) = \int_{\Omega} \int_{\Omega} \frac{\phi(x)\psi(y)}{|x-y|} dx dy$$

is a well defined positive definite bilinear form, provided $D(|\phi|, |\psi|) < \infty$ [30, Theorem 9.8]. We note that in the definition of $D(\cdot, \cdot)$ the dependence on Ω is implicit, as it is the domain of the functions ϕ, ψ .

2.1. Minimization of $E_q(u, \Omega)$ in u . Existence of u achieving the infimum in the definition of $E_q(\Omega)$ is an application of the Direct Method in the Calculus of Variations.

Lemma 2.2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded open set and let $q > 0$. Then the minimization problem*

$$(4) \quad E_q(\Omega) = \inf \left\{ E_q(v, \Omega) : v \in H_0^1(\Omega), \int_{\Omega} v^2 dx = 1 \right\}.$$

admits a solution with constant sign (nonnegative, without loss of generality).

Proof. Let $(u_n)_n$ be a minimizing sequence for the energy. As all the addends in the definition of $E_q(u, \Omega)$ are positive, we infer that $(u_n)_n$ is bounded in $H_0^1(\Omega)$.

Then up to passing to a subsequence, $(u_n)_n$ converges weakly in $H_0^1(\Omega)$ and strongly in $L^2(\Omega)$ to some function $u \in H_0^1(\Omega)$. In particular the L^2 -convergence implies that $\|u\|_{L^2(\Omega)} = 1$. By lower semicontinuity with respect to the weak convergence, we have that

$$\int |\nabla u|^2 dx \leq \liminf_{n \rightarrow +\infty} \int |\nabla u_n|^2 dx$$

and, by Fatou Lemma

$$D(u^2, u^2) = \iint_{\Omega \times \Omega} \frac{u^2(x)u^2(y)}{|x-y|} dx dy \leq \liminf_{n \rightarrow +\infty} D(u_n^2, u_n^2).$$

Hence, the energy is lower semicontinuous

$$E_q(u, \Omega) \leq \liminf_{n \rightarrow +\infty} E_q(u_n, \Omega),$$

and u is a minimizer. The fact that u can be chosen of constant sign follows since

$$E_q(|u|, \Omega) = E_q(u, \Omega),$$

and since $|u|$ is an admissible competitor. □

Remark 2.3. It is not difficult to check that the scale invariant functional

$$\tilde{E}_q(v, \Omega) := \frac{\int_{\Omega} |\nabla v|^2 dx}{\|v\|_{L^2}^2} + \frac{q \int_{\Omega} \int_{\Omega} \frac{v^2(x)v^2(y)}{|x-y|} dx dy}{\|v\|_{L^2}^4}, \quad v \in H_0^1(\Omega),$$

leads to an equivalent, yet unconstrained, minimization problem

$$\min \left\{ E_q(v, \Omega) : v \in H_0^1(\Omega), \int_{\Omega} v^2 dx = 1 \right\} = \min \left\{ \tilde{E}_q(v, \Omega) : v \in H_0^1(\Omega), v \neq 0 \right\}.$$

In particular we can always choose an optimal function $u \in H_0^1(\Omega)$ for the unconstrained problem which satisfies, a posteriori, the constraint $\|u\|_{L^2} = 1$.

Let us introduce the semilinear operator on $H_0^1(\Omega)$,

$$L_{\Omega, q}(u) = -\Delta u + q \left(u^2 \star \frac{1}{|\cdot|} \right) u.$$

One easily shows that $L_{\Omega, q}$ is a positive operator, that is, $(L_{\Omega, q}(u), u)_{L^2} \geq 0$. We say that a number $\lambda > 0$ is a (nonlinear) eigenvalue for $L_{\Omega, q}$ if there exists a non-null function v such that

$$(5) \quad L_{\Omega, q}(v) = \lambda v, \quad \int_{\Omega} v^2 dx = 1,$$

in distributional sense. Such a function is then called an eigenfunction corresponding to λ .

By means of a first variation, it is possible to check that (as it is observed in [31]) the optimal function u attaining $E_q(\Omega)$ is a (nonnegative) normalized eigenfunction for the operator $L_{\Omega, q}$ associated to the eigenvalue

$$\lambda_q = \int_{\Omega} |\nabla u|^2 dx + qD(u^2, u^2).$$

We note that for $q \in (0, 1]$ and $|\Omega| = |B_1|$ the quantity λ_q is uniformly bounded from below and above:

$$(6) \quad \lambda_0(B_1) \leq \lambda_0(\Omega) \leq \lambda_q \leq 2E_q(\Omega) \leq 2E_1(B_1),$$

where $\lambda_0(A)$ denotes the first eigenvalue of the Dirichlet Laplacian of an open set $A \subset \mathbb{R}^3$, and we used the Faber-Krahn inequality in the first inequality of (6). In other words, the Euler-Lagrange equation for $E_q(\Omega)$ is

$$(7) \quad L_{\Omega, q}(u) = \lambda_q u, \quad u \in H_0^1(\Omega).$$

By classical elliptic theory we deduce now a useful uniform bound on the L^∞ norm of solutions to (7). Precisely, we have the following result.

Lemma 2.4. *Let $\Omega \subset \mathbb{R}^3$ be a bounded open set, let $q \in (0, 1]$ and let u be a solution of*

$$(8) \quad L_{\Omega, q}(u) = \lambda_q u \quad \text{in } H_0^1(\Omega), \quad \int_{\Omega} u^2 dx = 1.$$

Then $u \in C^\infty(\Omega)$ and, hence, is a classical solution of (7). Furthermore, if $\lambda_q \leq M$ then

$$\|u\|_{L^\infty(\Omega)} \leq C,$$

for a constant $C = C(M, |\Omega|) > 0$ depending only on M and $|\Omega|$.

Proof. Thanks to Lemma 2.1, and by the Young inequality, we deduce that

$$\|v_u\|_{L^\infty(\Omega)} \leq 1 + \|\nabla u\|_{L^2(\Omega)}^2 \leq 1 + \lambda_q.$$

Hence the function

$$c(x) = \lambda_q - qv_u(x)$$

is equibounded in Ω by a constant depending only on λ_q . As u solves

$$\Delta u + c(x)u = 0, \quad \text{in } H_0^1(\Omega),$$

by [17, Theorem 8.8] we can assert that $u \in H_{\text{loc}}^2(\Omega)$ and by a simple bootstrap argument, $u \in C^\infty(\Omega)$ so that it is a classical solution of (7). Eventually, by [17, Theorem 8.15] we conclude that

$$\|u\|_{L^\infty(\Omega)} \leq C(M, |\Omega|)\|u\|_{L^2(\Omega)} = C(M, |\Omega|),$$

where $C(M, |\Omega|) > 0$ is a constant depending only on M and $|\Omega|$. \square

Corollary 2.5. *There exists $q_0 \in (0, 1]$ such that for every $q \in (0, q_0]$ and every bounded open set $\Omega \subset \mathbb{R}^3$ with $|\Omega| = |B_1|$ every nonnegative minimizer u of $E_q(u, \Omega)$ solves*

$$-\Delta u = c(x)u \quad \text{in } H_0^1(\Omega), \quad \int_{\Omega} u^2 dx = 1,$$

for some $c \in C^\infty(\Omega; \mathbb{R}^+)$. In particular, u is superharmonic.

Proof. We set $c(x) = \lambda_q - qv_u(x)$. The positivity of c for all q sufficiently small universal follows by the uniform L^∞ bound proven in Lemma 2.4 and by the uniform bounds on λ_q from (6). Since $u \geq 0$, then it is superharmonic for q below a certain threshold q_0 . \square

From now on $q_0 > 0$ always refers to the universal constant in Corollary 2.5.

Remark 2.6. We note that in general one cannot expect that a minimizer u of $E_q(u, \Omega)$ has the entire set Ω as its support. This can be already seen in the case $q = 0$ with Ω consisting of two disjoint sets whose first Dirichlet eigenvalues are distinct. It is not difficult to show that this situation persists to the case of $q > 0$ sufficiently small depending on Ω .

On the other hand, we can prove that the energy E_q is the same on Ω and on the positivity set of u and give some further characterizations.

Lemma 2.7. *Let $q > 0$, let Ω be a bounded open set and let $u \in H_0^1(\Omega)$ be a nonnegative minimizer of $E_q(u, \Omega)$. Then $E_q(\Omega) = E_q(\{u > 0\})$. Furthermore, if Ω_i is a connected component of Ω then either $u > 0$ or $u \equiv 0$ on Ω_i .*

Proof. It was already observed that we can choose $u \geq 0$. By the optimality of u , $E_q(\Omega) \geq E_q(\{u > 0\})$. On the other hand, $\{u > 0\} \subset \Omega$ and by monotonicity of the functional $E_q(\Omega)$ in Ω , we infer $E_q(\Omega) = E_q(\{u > 0\})$. The last part of the statement follows from Lemma 2.4 and [30, Theorem 9.10]. \square

Remark 2.8. We highlight again that if Ω is not connected, then we only know up to this point that the set $\{u > 0\}$ coincides with the union of some of the connected components of Ω and has the same energy as Ω .

We show now that $E_q(u, \Omega)$ has a unique (up to sign) minimizer u if Ω is a *connected* set (or, more generally, if $u > 0$ on all connected components of Ω). To do so, we follow an approach proposed in [5] (later revisited in [8]), based on a *hidden convexity property* of the functional.

Proposition 2.9. *Let $q > 0$ and let Ω be a connected bounded open set. Let $u, v \in H_0^1(\Omega)$ be such that $\|u\|_{L^2(\Omega)} = \|v\|_{L^2(\Omega)} = 1$. Let $\sigma_t = \sigma_t(u, v)$ be defined as*

$$\sigma_t = ((1-t)u^2 + tv^2)^{1/2}, \quad \text{for } t \in (0, 1).$$

Then the map $g : t \mapsto E_q(\sigma_t, \Omega)$ is strictly convex. In particular, the function u attaining the minimum of $E_q(u, \Omega)$ is unique.

Proof. Thanks to Lemma 2.7 and the connectedness of Ω , we deduce that the optimal function u for $E_q(\Omega)$ is strictly positive in Ω . Then the claim follows directly from [5, Lemma 4], which assures that the map $t \mapsto E_q(\sigma, \Omega)$ is strictly convex. Eventually, if u, v are two distinct (strictly positive) minimizers, since $\|\sigma_t(u, v)\|_{L^2(\Omega)} = 1$ by strict convexity it follows immediately that $E_q(\sigma_t, \Omega) < E_q(u, \Omega) = E_q(v, \Omega)$, a contradiction. \square

As a side result of Proposition 2.9, we have also the radially of u , when Ω is a ball.

Corollary 2.10. *Let $\Omega \subset \mathbb{R}^3$ be a ball. Then the unique solution to*

$$\min \left\{ E_q(u, \Omega) : u \in H_0^1(\Omega), \int_{\Omega} u^2 dx = 1 \right\},$$

is radial.

Proof. The uniqueness of the solution follows from Proposition 2.9. Concerning the radially, it follows from the rotational invariance of the energy $E_q(u, \Omega)$. \square

2.2. Fraenkel asymmetry and quantitative Faber-Krahn inequality. Here we recall the sharp quantitative version of the Faber–Krahn inequality. We first remind the notion of *Fraenkel asymmetry*: for a set $\Omega \subset \mathbb{R}^3$ with finite measure we define

$$(9) \quad \mathcal{A}(\Omega) = \inf_{x \in \mathbb{R}^3} \frac{|\Omega \Delta (B + x)|}{|\Omega|},$$

where B denotes the ball of measure $|\Omega|$ centered at the origin. We also recall that we denote by $\lambda_0(A)$ the first eigenvalue of the Dirichlet Laplacian of a set $A \subset \mathbb{R}^3$.

Theorem 2.11. [7] *There exists a universal positive constant $\hat{\sigma} > 0$ such that for all open sets $\Omega \subset \mathbb{R}^3$ with finite measure we have*

$$|\Omega|^{2/3} \lambda_0(\Omega) - |B_1|^{2/3} \lambda_0(B_1) \geq \hat{\sigma} \mathcal{A}(\Omega)^2,$$

where \mathcal{A} the Fraenkel asymmetry.

2.3. Some facts about quasi-open sets. Finally, we recall some definitions and facts about quasi-open sets which will be needed in the following.

Definition 2.12. A *quasi-open* set is a measurable set $\Omega \subset \mathbb{R}^3$ such that for all $\varepsilon > 0$ there exists K_ε compact such that its Newtonian capacity $\text{cap}(K_\varepsilon) < \varepsilon$ and $\Omega \setminus K_\varepsilon$ is open. Similarly, a function $u : \Omega \rightarrow \mathbb{R}$ is *quasi-continuous* if for all $\varepsilon > 0$ there exists a compact set K_ε such that $\text{cap}(K_\varepsilon) < \varepsilon$ and the restriction of u to $\Omega \setminus K_\varepsilon$ is continuous. Eventually, we say that a property holds *quasi-everywhere* on a set if it holds up to a set of null-capacity.

It is well-known that every $u \in H^1(B_R)$ admits a quasi-continuous representative \tilde{u} . Moreover, if \tilde{u} and \hat{u} are two quasi-continuous representatives of u , then they are equal quasi-everywhere. Therefore, in this paper, for every $u \in H^1(B_R)$ we identify it with its quasi-continuous representative. A quasi-open set is then simply a superlevel set of (the quasi-continuous representative of) a function $u \in H^1(B_R)$. For more details on quasi-open sets and quasi-continuous functions, and the definition of capacity we refer to [24, Chapter 2], [25, Chapter 3] or [23, Chapter 2 and 4].

Let us also stress that it is standard to define the Sobolev space H_0^1 on a quasi-open set $\Omega \subset \mathbb{R}^3$ as

$$H_0^1(\Omega) = \{u \in H^1(\mathbb{R}^3) : u = 0 \text{ quasi-everywhere in } \mathbb{R}^3 \setminus \Omega\}.$$

If Ω is an open set, this definition coincides with the usual one, see [25, Section 3.3.5] for more details.

Remark 2.13. If $\Omega \subset \mathbb{R}^3$ is a quasi-open set and $u \in H_0^1(\Omega)$ is a nonnegative solution to (8), then $u \in L^\infty(\Omega)$ and its L^∞ norm is bounded by a constant as in Lemma 2.4. This can be checked with an approximation argument directly from the definition of a quasi-open set.

3. AN EXISTENCE RESULT FOR AN AUXILIARY PROBLEM

In this section we begin the proof of the following result, which can be seen as an equibounded version of Theorem 1.3.

Theorem 3.1. *There exists $R_0 > 1$ such that, for all $R \geq R_0$ there exists $\bar{q} = \bar{q}(R) > 0$ such that for all $q \leq \bar{q}$ there exists a minimizer for the problem*

$$(10) \quad \min \left\{ E_q(\Omega) : \Omega \subset B_R, |\Omega| = |B_1| \right\}.$$

Moreover, for all $\varepsilon > 0$ there exists $q_\varepsilon = q_\varepsilon(R) > 0$ such that if $q \leq q_\varepsilon$ then any minimizer is an $(\varepsilon, C^{2,\alpha})$ -nearly spherical set, with $\alpha = \alpha(\varepsilon) \in (0, 1)$.

The proof of this result is carried out through several steps, divided in the next (sub)sections, as outlined in the introduction.

3.1. Removing the L^2 norm constraint in E_q .

Proposition 3.2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded quasi-open set of measure $|B_1|$ and let $q \leq q_0$. There exists $\bar{M} > 0$ (uniform for all $q \leq q_0$) such that for all $M \geq \bar{M}$, the minimizers of the problem (4) are the same as those of*

$$(11) \quad \min \left\{ E_{q,M}(v, \Omega) : v \in H_0^1(\Omega) \right\},$$

where

$$E_{q,M}(v, \Omega) = E_q(v, \Omega) + M \left| \int v^2 dx - 1 \right|.$$

Proof. Problem (11) admits a minimizer, as it can be seen as in the proof of Lemma 2.2. We set

$$0 < \alpha = \min \left\{ E_q(v, \Omega) : v \in H_0^1(\Omega), \int v^2 dx = 1 \right\} = E_q(u, \Omega).$$

Since $\int u^2 = 1$, clearly for all $M > 0$, $E_{q,M}(u, \Omega) = E_q(u, \Omega)$, hence

$$\alpha \geq \min \left\{ E_{q,M}(v, \Omega) : v \in H_0^1(\Omega) \right\}.$$

Let us assume by contradiction that there is a sequence $M_k \rightarrow +\infty$ such that

$$\min \left\{ E_{q,M_k}(v, \Omega) : v \in H_0^1(\Omega) \right\} = E_{q,M_k}(\hat{u}_k, \Omega) < \alpha.$$

We call $\sigma_k = \|\hat{u}_k\|_{L^2} - 1$ and define $u_k = \frac{\hat{u}_k}{1+\sigma_k}$, so that $\|u_k\|_{L^2} = 1$. This definition makes sense for all large enough k , since $\sigma_k \rightarrow 0$ as $k \rightarrow +\infty$. We can then compute

$$\begin{aligned} \alpha > E_{q,M_k}(\hat{u}_k, \Omega) &= (1 + \sigma_k)^2 \int |\nabla u_k|^2 dx + \frac{q}{2}(1 + \sigma_k)^4 D(u_k^2, u_k^2) + M_k \left| (1 + \sigma_k)^2 - 1 \right| \\ &= E_q(u_k, \Omega) + (2\sigma_k + o(\sigma_k)) \int |\nabla u_k|^2 dx + \frac{q}{2}(4\sigma_k + o(\sigma_k)) D(u_k^2, u_k^2) + 2M_k(|\sigma_k| + o(\sigma_k)). \end{aligned}$$

Noting that $\int |\nabla u_k|^2 dx$ and $D(u_k^2, u_k^2)$ are uniformly bounded by 2α and $q \leq q_0$, for $M_k \geq 10\alpha$ and k large enough, we conclude that

$$\alpha > E_{q,M_k}(\hat{u}_k, \Omega) \geq E_q(u_k, \Omega) \geq \alpha,$$

a contradiction. We note that the choice of M_k for reaching a contradiction is independent of q and Ω . \square

In accordance with the notation of the previous theorem, we set

$$(12) \quad E_{q,M}(\Omega) := \min \left\{ E_q(v, \Omega) + M \left| \int v^2 dx - 1 \right| : v \in H_0^1(\Omega) \right\}$$

and note that, as a consequence of Proposition 3.2, if u_Ω is a minimizer of $E_{q,M}(\Omega)$ for $M \geq \bar{M}$, then $\int u_\Omega^2 dx = 1$, thus, a posteriori, $E_q(\Omega) = E_{q,M}(\Omega)$.

Remark 3.3. From now on, we fix once and for all a constant $M > \bar{M}$, and we stress that this constant does not depend on $q \leq q_0$ and will not be changed later in the paper.

3.2. Another auxiliary problem: removing the volume constraint. In order to get rid of the measure constraint, we follow an approach first proposed by Aguilera, Alt and Caffarelli [1]. Let $\eta \in (0, 1)$ and consider the piecewise linear function

$$f_\eta: \mathbb{R}^+ \rightarrow \mathbb{R}, \quad f_\eta(s) = \begin{cases} \eta(s - |B_1|), & \text{if } s \leq |B_1|, \\ \frac{1}{\eta}(s - |B_1|), & \text{if } s \geq |B_1|. \end{cases}$$

It is easy to check that, for all $0 \leq s_2 \leq s_1$, there holds

$$(13) \quad \eta(s_1 - s_2) \leq f_\eta(s_1) - f_\eta(s_2) \leq \frac{1}{\eta}(s_1 - s_2).$$

We introduce then the functional (recalling that M is fixed, see Remark 3.3 and Proposition 3.2),

$$(14) \quad E_{q,M,\eta}(\Omega) := E_{q,M}(\Omega) + f_\eta(|\Omega|),$$

and, for $R > 1$, the minimization problem

$$(15) \quad \min \{E_{q,M,\eta}(\Omega) : \Omega \subset B_R, \Omega \text{ open}\}.$$

To prove the existence of a minimizer for problem (15), we need first to work in the setting of quasi-open sets (see Section 2.3) and then recover the regularity.

Thus, we first focus on the problem:

$$(16) \quad \min \{E_{q,M,\eta}(\Omega) : \Omega \subset B_R, \Omega \text{ quasi-open}\}.$$

We aim to prove that problem (16) is equivalent to the unconstrained problem (11) (and thus also to (10)), at least for q and η small enough. To do that we first have to prove existence and some mild regularity of minimizers of $E_{q,M,\eta}$. We begin by showing a lower bound for $E_{q,M,\eta}$ on equibounded sets.

Lemma 3.4. *Let $R > 1$, $q \in (0, q_0]$ and $\eta \in (0, 1)$. Then, for all quasi-open $\Omega \subset B_R$, we have*

$$E_{q,M,\eta}(\Omega) \geq \lambda_0(B_1)R^{-2} - |B_1|.$$

Proof. Let $v \in H_0^1(\Omega)$, with $\int_\Omega v^2 dx = 1$ be a function attaining the infimum in the definition of $E_{q,M}(\Omega) = E_q(\Omega)$, so that

$$E_{q,M,\eta}(\Omega) = \int |\nabla v|^2 dx + \frac{q}{2}D(v^2, v^2) + f_\eta(|\Omega|).$$

By the monotonicity of Dirichlet eigenvalues, the inclusion $\Omega \subset B_R$, the positivity of $D(\cdot, \cdot)$, the scaling properties of λ_0 , the isoperimetric inequality and the definition of f_η , we obtain

$$\int |\nabla v|^2 dx + \frac{q}{2}D(v^2, v^2) + f_\eta(|\Omega|) \geq \lambda_0(\Omega) - \eta \max\{|B_1| - |\Omega|, 0\} \geq \lambda_0(B_1)R^{-2} - |B_1|,$$

so the claim is proved. \square

The following existence result is mostly an adaptation to our situation of [7, Lemma 4.6] and [35, Lemma 3.2], which are in turn inspired by [10, Theorem 2.2 and Lemma 2.3].

Lemma 3.5. *Let $\eta \in (0, 1)$, $q \in (0, q_0]$ and let $R > 1$. There exists a minimizer for problem (16). Moreover all minimizers have perimeter uniformly bounded by a constant depending on R, η .*

Proof. Let $(\Omega_n)_n \subset B_R$ be such that

$$E_{q,M,\eta}(\Omega_n) \leq \inf \{E_{q,M,\eta}(\Omega) : \Omega \subset B_R, \text{ quasi-open}\} + \frac{1}{n}.$$

Let u_n be an optimal function with unit L^2 norm for the minimization of $E_q(\Omega_n)$ or equivalently $E_{q,M}(\Omega_n)$, so that by Lemma 2.7, either $\Omega_n = \{u_n > 0\}$, or $\{u_n > 0\}$ is a union of some connected components² of Ω_n , with $E_{q,M,\eta}(\{u_n > 0\}) \leq E_{q,M,\eta}(\Omega_n)$, so that $(\{u_n > 0\})_n$ still forms a minimizing sequence and we can replace Ω_n with $\{u_n > 0\}$. Let $t_n = 1/\sqrt{n}$. We define

$$\tilde{\Omega}_n := \{u_n > t_n\}.$$

We have

$$E_{q,M,\eta}(\Omega_n) \leq E_{q,M,\eta}(\tilde{\Omega}_n) + \frac{1}{n},$$

²Notice that as u_n are regular functions, there is no measure-theoretical issues in defining connected components.

and since we can take $\tilde{u}_n := (u_n - t_n)_+$ as a competitor in the minimization problem defining $E_M(\tilde{\Omega}_n)$, we obtain

$$(17) \quad \begin{aligned} & \int_{\{u_n > 0\}} |\nabla u_n|^2 dx + \frac{q}{2} D(u_n^2, u_n^2) + M \left| \int u_n^2 dx - 1 \right| + f_\eta(|\{u_n > 0\}|) \\ & \leq \int_{\{u_n > t_n\}} |\nabla u_n|^2 dx + \frac{q}{2} D(\tilde{u}_n^2, \tilde{u}_n^2) + M \left| \int (\tilde{u}_n)^2 dx - 1 \right| + f_\eta(|\{u_n > t_n\}|) + \frac{1}{n}. \end{aligned}$$

Using the uniform bound on the L^∞ norm of u_n , see Lemma 2.4, we observe that

$$\begin{aligned} \int_{\{u_n > t_n\}} u_n^2 - (u_n - t_n)^2 dx &= \int_{\{u_n > t_n\}} 2t_n u_n - t_n^2 dx \leq 2\|u_n\|_{L^\infty} t_n |\{u_n > t_n\}| \leq C(R) t_n |\{u_n > t_n\}|, \\ \text{and } \int_{\{0 \leq u_n \leq t_n\}} u_n^2 dx &\leq t_n^2 |\{0 < u_n \leq t_n\}|, \end{aligned}$$

and obtain the estimate (possibly increasing the value of $C(R)$)

$$(18) \quad \int_{\{u_n > 0\}} u_n^2 dx - \int_{\{u_n > t_n\}} (u_n - t_n)^2 dx \leq C(R) t_n |\{u_n > 0\}|.$$

Noting that $D(\tilde{u}_n^2, \tilde{u}_n^2) - D(u_n^2, u_n^2) \leq 0$, recalling the property (13) of f_η and (18), we can rewrite (17) as

$$(19) \quad \int_{\{0 < u_n < t_n\}} |\nabla u_n|^2 dx + \eta |\{0 < u_n < t_n\}| \leq C(R) M t_n |\{u_n > 0\}| + \frac{1}{n} \leq C(R, M) t_n + \frac{1}{n}.$$

On the other hand, since $\eta < 1$, using coarea formula, the arithmetic-geometric-mean inequality and (19), we obtain

$$\begin{aligned} 2\eta \int_0^{t_n} P(\{u_n > s\}) ds &= 2\eta \int_{\{0 < u_n < t_n\}} |\nabla u_n| dx \\ &\leq \eta \int_{\{0 < u_n < t_n\}} |\nabla u_n|^2 dx + \eta |\{0 < u_n < t_n\}| \leq C(R, M) t_n + \frac{1}{n}. \end{aligned}$$

As $t_n = 1/\sqrt{n}$ we can find a level $0 < s_n < 1/\sqrt{n}$ such that the sets $W_n := \{u_n > s_n\}$ satisfy

$$P(W_n) \leq \frac{2}{t_n} \int_0^{t_n} P(\{u_n > s\}) ds \leq \frac{C(R, M)}{\eta} + \frac{1}{\eta t_n n} \leq C(R, M, \eta) + \frac{1}{\eta \sqrt{n}}.$$

It is easy to check that $(W_n)_n$ is still a minimizing sequence for problem (16). In fact, with arguments similar to the ones used above, we obtain:

$$(20) \quad \begin{aligned} E_{q, M, \eta}(W_n) &= \int_{\{u_n > s_n\}} |\nabla u_n|^2 dx + \frac{q}{2} D((u_n - s_n)_+^2, (u_n - s_n)_+^2) \\ &\quad + M \left| \int_{\{u_n > s_n\}} (u_n - s_n)^2 dx - 1 \right| + f_\eta(|\{u_n > s_n\}|) \\ &\leq E_{q, M, \eta}(\Omega_n) + C(R, M) s_n + f_\eta(|\{u_n > s_n\}|) - f_\eta(|\{u_n > 0\}|) \\ &\leq E_{q, M, \eta}(\Omega_n) + \frac{C(R, M)}{\sqrt{n}} - \eta |\{0 < u_n < s_n\}| \leq E_{q, M, \eta}(\Omega_n) + \frac{C(R, M)}{\sqrt{n}}, \end{aligned}$$

where we used that $D((u_n - s_n)_+^2, (u_n - s_n)_+^2) - D(u_n^2, u_n^2) \leq 0$ and property (18) with s_n in place of t_n . Moreover, since the sets of the sequence $(W_n)_n$ have equibounded perimeter, there exists a Borel set W_∞ such that (up to passing to subsequences)

$$(21) \quad W_n \rightarrow W_\infty, \text{ in } L^1(B_R), \quad P(W_\infty) \leq C(R, M, \eta).$$

On the other hand, an optimal function (normalized in L^2) w_n attaining $E_{q, M, \eta}(W_n)$, is equibounded in $H^1(B_R)$. In fact, being $(W_n)_n$ a minimizing sequence for $E_{q, M, \eta}$, we have that

$$\int |\nabla w_n|^2 dx + \int w_n^2 dx \leq 1 + E_{q, M, \eta}(W_n) \leq 1 + C.$$

Hence, up to passing to subsequences, there is $w \in H_0^1(B_R)$ such that

$$(22) \quad w_n \rightarrow w \text{ strongly in } L^2(B_R), \text{ weakly in } H_0^1(B_R) \text{ and pointwise a.e.}$$

Let $W := \{w > 0\}$, and recall that we are identifying w with its quasi-continuous representative. Then, thanks to (21) and (22), we deduce

$$\chi_W(x) \leq \liminf_n \chi_{W_n}(x) = \chi_{W_\infty}(x), \quad \text{for a.e. } x \in B_R,$$

hence $|W \setminus W_\infty| = 0$, that is $W \subset W_\infty$ up to a negligible set. We now observe that $\Omega \mapsto f_\eta(|\Omega|)$ is continuous with respect to the L^1 convergence of sets, the Dirichlet energy is lower semicontinuous with respect to the weak H^1 convergence and the functional $D(\cdot, \cdot)$ is lower semicontinuous with respect to the strong L^2 convergence by Fatou lemma. We can therefore pass to the limit in (20) and obtain

$$(23) \quad \begin{aligned} \int_W |\nabla w|^2 dx + \frac{q}{2} D(w^2, w^2) + f_\eta(|W_\infty|) &\leq \liminf_n \int_{W_n} |\nabla w_n|^2 dx + \frac{q}{2} D(w_n^2, w_n^2) + f_\eta(|W_\infty|) \\ &\leq \liminf_n E_{q,M,\eta}(W_n) = \inf_{\Omega \subset B_R} E_{q,M,\eta}(\Omega) \leq E_q(W) + f_\eta(|W|). \end{aligned}$$

In conclusion, using also (23), we have

$$\eta |W_\infty \setminus W| = \eta(|W_\infty| - |W|) \leq f_\eta(|W_\infty|) - f_\eta(|W|) \leq 0,$$

thus $|W_\infty \setminus W| = 0$, which entails $W = W_\infty$ a.e. and this is the desired minimizer for problem (16). \square

3.3. Free boundary formulation. In the previous section we introduced the functional $E_{q,M,\eta}$, see (14), we proved existence and mild regularity properties of minimizers (namely: they are sets of finite perimeter). In this section we improve the regularity for such sets. This will be needed in the sequel but allows us also to show the equivalence between unconstrained minimizers of $E_{q,M,\eta}$ and volume constrained minimizers of E_q and $E_{q,M}$. The crucial remark is that one may consider, in place of the shape functional $E_{q,M,\eta}$, a functional defined on the larger space $H_0^1(B_R)$ and take a *free boundary* approach. Let us define, for $u \in H_0^1(B_R)$,

$$\mathcal{E}_{q,M,\eta}(u) = \int_{\{u>0\}} |\nabla u|^2 dx + \frac{q}{2} \int_{\{u>0\}} \int_{\{u>0\}} \frac{u^2(x)u^2(y)}{|x-y|} dx dy + M \left| \int_{\{u>0\}} u^2 dx - 1 \right| + f_\eta(|\{u > 0\}|),$$

and we note that one could have equivalently integrated over B_R in all the integrals above.

Lemma 3.6. *Let $\eta \in (0, 1)$, $q \in (0, q_0]$ and Ω be a minimizer for problem (16). Then every minimal function of $E_q(\Omega)$ is a minimizer of $\mathcal{E}_{q,M,\eta}$. Viceversa, if w minimizes $\mathcal{E}_{q,M,\eta}$, then $\Omega = \{w > 0\}$ is a minimizer of $E_{q,M,\eta}$. Furthermore, it is possible to select a minimizer Ω for $E_{q,M,\eta}$ which coincides with the support of an optimal function for $E_q(\Omega)$.*

Proof. Concerning the first claim, let Ω be an optimal set for $E_{q,M,\eta}$ and u an optimal function for $E_q(\Omega)$. We immediately note that $\{u > 0\} = \Omega$ up to sets of zero measure, otherwise $\{u > 0\} \subset \Omega$, thus $f_\eta(|\{u > 0\}|) < f_\eta(|\Omega|)$ so that $E_{q,M,\eta}(\Omega) > E_{q,M,\eta}(\{u > 0\})$, a contradiction with the optimality of Ω . As a consequence, $\mathcal{E}_{q,M,\eta}(u) = E_{q,M,\eta}(\{u > 0\}) = E_{q,M,\eta}(\Omega)$. Therefore, for all $v \in H_0^1(B_R)$, we have

$$\mathcal{E}_{q,M,\eta}(u) = E_{q,M,\eta}(\Omega) \leq E_{q,M,\eta}(\{v > 0\}) = \mathcal{E}_{q,M,\eta}(v),$$

namely u is a minimizer for $\mathcal{E}_{q,M,\eta}$.

Let us focus on the second claim: let u be an optimal function for $\mathcal{E}_{q,M,\eta}$, and we call $\Omega = \{u > 0\}$. For all $\tilde{\Omega} \subset B_R$, calling \tilde{u} any optimal function attaining $E_q(\tilde{\Omega})$, we have

$$E_{q,M,\eta}(\Omega) = \mathcal{E}_{q,M,\eta}(u) \leq \mathcal{E}_{q,M,\eta}(\tilde{u}) \leq E_{q,M,\eta}(\tilde{\Omega}),$$

as requested. Notice that the last inequality is not a priori an equality, as in principle $\{\tilde{u} > 0\} \subset \tilde{\Omega}$.

Concerning the last part of the statement, it is enough to notice that, given an optimal function u for $E_q(\Omega)$, $\{u > 0\}$ is always a minimizer for $E_{q,M,\eta}$. \square

In the rest of this section, we focus on the new (equivalent) formulation of problem (16)

$$(24) \quad \min \{ \mathcal{E}_{q,M,\eta}(u) : u \in H_0^1(B_R) \}.$$

As a consequence of Lemma 3.6 and Remark 2.8, working on problem (24) means selecting an optimal set $\{u > 0\}$ for the original problem (2) or for (16), namely the union of the connected components of Ω where u is nonzero.

In the next results, we work with a minimizer of the form $\Omega = \{u > 0\}$. This is not restrictive, as next lemma shows.

Lemma 3.7. *For all $R \geq 1$, $\eta \in (0, 1)$ and $q \leq q_0$, if Ω is an optimal set for problem (16) and u is an associated (nonnegative) optimal function attaining $E_q(\Omega)$, then $\{u > 0\} = \Omega$ (up to a negligible set). Moreover, Ω is connected.*

Proof. Let us argue for the sake of contradiction and assume that $\{u > 0\}$ is strictly contained in Ω . We already know from Remark 2.8 that the above assumptions entail that $u \equiv 0$ on a connected component ω , with positive measure. Therefore, $f_\eta(|\{u > 0\}|) < f_\eta(|\Omega|)$ and we have contradicted the minimality of Ω for $E_{q,M,\eta}$. Concerning the last part of the statement, if $\Omega = \{u > 0\}$ is the disjoint union of two components Ω_1 and Ω_2 , by increasing the distance between the two components we are strictly decreasing the Coulomb energy term, while the other terms of $E_{q,M,\eta}$ are unchanged. Thus we contradict again the optimality of Ω . \square

Remark 3.8. We stress that if we could prove that an optimal function for (24) is a quasi-minimizer for the functional

$$J(u) = \int_{\{u>0\}} |\nabla u|^2 dx + |\{u > 0\}|,$$

i.e. letting $J_{x,r}(u) = \int_{\{u>0\} \cap B_r(x)} |\nabla u|^2 + |\{u > 0\} \cap B_r(x)|$, we have, for some $\beta > 0$,

$$J_{x,r}(u) \leq (1 + Kr^\beta)J_{x,r}(v), \quad \text{for all admissible } v \text{ and for all } x, r,$$

this would strongly simplify the regularity proof, see for example [38]. Unfortunately, this does not seem to be the case in our setting, due to the presence of the nonlocal double integral term $D(u^2, u^2)$. Therefore we use a careful modification of the standard free boundary regularity techniques developed starting from [2].

Lemma 3.9. *Let $R > 1$, $\eta \in (0, 1)$, $q \in (0, q_0]$, let Ω be an optimal set for problem (16), and let $u \in H_0^1(\Omega)$ be any (nonnegative) function attaining $E_q(\Omega) = E_{q,M}(\Omega)$. Then for every $\kappa \in (0, 1)$ there are positive constants K_0, ρ_0 depending only on κ, η, R such that the following assertion holds: if $\rho \leq \rho_0$ and $x_0 \in B_R$, then*

$$(25) \quad \int_{\partial B_\rho(x_0) \cap B_R} u d\mathcal{H}^2 \leq K_0 \rho \implies u \equiv 0 \text{ in } B_{\kappa\rho}(x_0) \cap B_R.$$

Proof. By Lemma 3.7, we know that $\Omega = \{u > 0\}$, that u is optimal for problem (24) and has unitary L^2 norm. Without loss of generality, we fix $x_0 = 0$. We also extend u to zero outside B_R , so that by Lemma 2.4, u solves distributionally $-\Delta u \leq \gamma_1$ in \mathbb{R}^3 where (since $q \leq q_0$)

$$\gamma_1 := 2 \sup_x (\lambda_q - qv_u(x)) \|u\|_{L^\infty(\Omega)} > 0.$$

The positivity of γ_1 follows by the bound (uniform in q) on $\|u\|_{L^\infty(\Omega)}$ (see Lemma 2.4) and, consequently, on $\|v_u\|_{L^\infty(\Omega)}$, while $\lambda_q \geq \lambda_0(\Omega) \geq \lambda_0(B_1) > 0$. Then the function

$$x \mapsto u(x) + \gamma_1 \frac{|x|^2 - \rho^2}{6}$$

is subharmonic in B_ρ (recalling that we are in three dimensional setting). Thus, for every $\kappa \in (0, 1)$, there exists $c = c(\kappa)$ such that

$$(26) \quad \delta_\rho := \sup_{B_{\sqrt{\kappa}\rho}} u \leq c \left(\int_{\partial B_\rho \cap B_R} u d\mathcal{H}^2 + \gamma_1 \rho^2 \right) \leq c(K_0 \rho + \rho^2).$$

Let us show now that there exists $\rho > 0$ small enough so that there exists a positive solution w of

$$(27) \quad \begin{cases} -\Delta w = \frac{M}{2}(u + w), & \text{in } B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}, \\ w = \delta_\rho, & \text{on } \partial B_{\sqrt{\kappa}\rho}, \\ w = 0, & \text{on } B_{\kappa\rho}, \end{cases}$$

where M is fixed large enough so that the statement of Proposition 3.2 holds. To show existence of a solution, let $\rho > 0$ be such that

$$\alpha(\rho) := \lambda_0(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho})^{-1} \left(\frac{M}{2} \|u\|_{L^2(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho})} + \frac{M}{2} \right) \leq \frac{1}{4}.$$

This can be easily obtained as $\lambda_0(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \rightarrow +\infty$ as $\rho \rightarrow 0$ and since $u \in L^\infty(\Omega)$. Then any minimizing sequence for the energy

$$(28) \quad \varphi \mapsto \frac{1}{2} \int_{B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}} |\nabla \varphi|^2 dx - \frac{M}{2} \int_{B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}} \left(\frac{\varphi}{2} + u \right) \varphi dx$$

with boundary conditions as in (27) can be chosen, after standard computations, made of nonnegative functions (as passing to the modulus decreases the energy) and such that

$$\int_{B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}} |\varphi_n|^2 dx \leq \alpha(\rho) \int_{B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}} |\nabla \varphi_n|^2 dx \leq \frac{1}{4} \int_{B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}} |\nabla \varphi_n|^2 dx.$$

Hence the sequence $(\|\varphi_n\|_{H^1})$ is uniformly bounded and a positive minimizer for the energy (28) exists and solves its Euler-Lagrange equation, namely (27). By standard elliptic regularity, we obtain that the L^∞ norm of any positive solution w is bounded by a constant $\gamma_2 > 0$ depending only on κ and M . By definition, $w \geq u$ on $\partial B_{\sqrt{\kappa}\rho}$, therefore the function

$$v = \begin{cases} u, & \text{in } \mathbb{R}^3 \setminus B_{\sqrt{\kappa}\rho}, \\ \min\{u, w\}, & \text{in } B_{\sqrt{\kappa}\rho}, \end{cases}$$

satisfies

$$v \leq u, \quad \{v > 0\} \subset \{u > 0\}, \quad \{v > 0\} \setminus B_{\sqrt{\kappa}\rho} = \{u > 0\} \setminus B_{\sqrt{\kappa}\rho},$$

thus $D(v^2, v^2) \leq D(u^2, u^2)$ (and we can neglect these contributions in the following computations). Since $v \in H_0^1(B_R)$, inequality (24) gives

$$\begin{aligned} & \int_{B_{\sqrt{\kappa}\rho}} |\nabla u|^2 dx + f_\eta(|\{u > 0\}|) \\ & \leq \int_{B_{\sqrt{\kappa}\rho}} |\nabla v|^2 dx + M \left| \int_{B_{\sqrt{\kappa}\rho}} (v^2 - u^2) dx \right| + f_\eta(|\{v > 0\}|). \end{aligned}$$

As $v = 0$ in $B_{\kappa\rho}$, using also (13), we get

$$\begin{aligned} \eta|\{u > 0\} \cap B_{\kappa\rho}| & \leq \eta(|\{u > 0\} \setminus \{v > 0\}| \cap B_{\sqrt{\kappa}\rho}) \\ & \leq f_\eta(|\{u > 0\}|) - f_\eta(|\{v > 0\}|). \end{aligned}$$

On the other hand, we can rewrite the term involving the L^2 norm of the functions as

$$\left| \int_{B_{\sqrt{\kappa}\rho}} (v^2 - u^2) dx \right| = \int_{B_{\sqrt{\kappa}\rho}} (u^2 - v^2) dx = \int_{B_{\kappa\rho}} u^2 dx + \int_{(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \cap \{u > w\}} (u^2 - w^2) dx.$$

Thanks to the two inequalities above and the definition of v , we can infer

$$\begin{aligned} & \int_{B_{\kappa\rho}} |\nabla u|^2 dx + \eta|\{u > 0\} \cap B_{\kappa\rho}| \leq \int_{B_{\kappa\rho}} |\nabla u|^2 dx + f_\eta(|\{u > 0\}|) - f_\eta(|\{v > 0\}|) \\ (29) \quad & \leq \int_{B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}} (|\nabla v|^2 - |\nabla u|^2) dx + M \int_{B_{\kappa\rho}} u^2 dx + M \int_{(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \cap \{u > w\}} (u^2 - w^2) dx \\ & \leq 2 \int_{(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \cap \{u > w\}} (|\nabla w|^2 - \nabla u \cdot \nabla w) dx + M \int_{B_{\kappa\rho}} u^2 dx + M \int_{(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \cap \{u > w\}} (u^2 - w^2) dx. \end{aligned}$$

On the other hand testing (27) with $(u - w)_+$ and integrating over $B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}$, we obtain

$$(30) \quad \int_{(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \cap \{u > w\}} (|\nabla w|^2 - \nabla u \cdot \nabla w) dx + \frac{M}{2} \int_{(B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}) \cap \{u > w\}} (u^2 - w^2) dx = \int_{\partial B_{\kappa\rho}} \frac{\partial w}{\partial \nu} u d\mathcal{H}^2,$$

where ν denotes the outer unit normal exiting from $B_{\kappa\rho}$ and thanks to the fact that $w = 0$ on $\partial B_{\kappa\rho}$ and $w \geq u$ on $\partial B_{\sqrt{\kappa}\rho}$. Recalling that $\|u\|_{L^\infty} \leq \gamma_1$ and $\|w\|_{L^\infty} \leq \gamma_2$, we now fix $\gamma_3 = \frac{M}{2}(\gamma_1 + \gamma_2)$ and consider the solution to the problem

$$\begin{cases} -\Delta \tilde{w} = \gamma_3, & \text{in } B_{\sqrt{\kappa}\rho} \setminus B_{\kappa\rho}, \\ \tilde{w} = \delta_\rho, & \text{on } \partial B_{\sqrt{\kappa}\rho}, \\ \tilde{w} = 0, & \text{on } B_{\kappa\rho}, \end{cases}$$

since the torsion function on an annulus is explicit (see [7]), with a direct computation one obtains

$$\left| \frac{\partial \tilde{w}}{\partial \nu} \right| \leq \beta_1 \frac{\delta_\rho + \rho^2}{\rho}, \quad \text{on } \partial B_{\kappa\rho},$$

for some $\beta_1 = \beta_1(\kappa, M)$. By comparison, since $\tilde{w} - w$ is superharmonic (by (27)) it follows

$$\left| \frac{\partial w}{\partial \nu} \right| \leq \left| \frac{\partial \tilde{w}}{\partial \nu} \right| \leq \beta_1 \frac{\delta_\rho + \rho^2}{\rho}, \quad \text{on } \partial B_{\kappa\rho}.$$

We can now combine (29) and (30) to obtain

$$\int_{B_{\kappa\rho}} |\nabla u|^2 dx + \eta|\{u > 0\} \cap B_{\kappa\rho}| \leq \beta_1(\kappa) \frac{\delta_\rho + \rho^2}{\rho} \int_{\partial B_{\kappa\rho}} u d\mathcal{H}^2 + M \int_{B_{\kappa\rho}} u^2 dx.$$

By the uniform bound on the L^∞ norm of u , we have the estimate

$$M \int_{B_{\kappa\rho}} u^2 dx \leq \beta_2 \delta_\rho^2 |\{u > 0\} \cap B_{\kappa\rho}|,$$

for some $\beta_2(\kappa, M)$.

Then, using the definition of δ_ρ , the trace inequality in $W^{1,1}$ and the arithmetic geometric mean inequality we obtain

$$\begin{aligned} \int_{\partial B_{\kappa\rho}} u d\mathcal{H}^2 &\leq C(\kappa) \left(\frac{1}{\rho} \int_{B_{\kappa\rho}} u dx + \int_{B_{\kappa\rho}} |\nabla u| dx \right) \\ &\leq \beta_3 \left(\left(\frac{\delta_\rho}{\rho} + \frac{1}{2} \right) |\{u > 0\} \cap B_{\kappa\rho}| + \frac{1}{2} \int_{B_{\kappa\rho}} |\nabla u|^2 dx \right), \end{aligned}$$

for some $\beta_3 = \beta_3(\kappa) > 0$. By collecting the above estimates, recalling again (26) we have, for all $\rho \leq \rho_0$

$$\begin{aligned} &\eta \int_{B_{\kappa\rho}} |\nabla u|^2 dx + \eta |\{u > 0\} \cap B_{\kappa\rho}| \\ &\leq \beta_1 \frac{\delta_\rho + \rho^2}{\rho} \int_{\partial B_{\kappa\rho}} u d\mathcal{H}^2 + \delta_\rho (1 + \delta_\rho \beta_2) |\{u > 0\} \cap B_{\kappa\rho}| \\ &\leq \beta_1 (c(K_0 + \rho) + \rho) \int_{\partial B_{\kappa\rho}} u d\mathcal{H}^2 + c(K_0 \rho + \rho^2) (1 + c(K_0 \rho + \rho^2) \beta_3) |\{u > 0\} \cap B_{\kappa\rho}| \\ &\leq \beta_1 \beta_3 (c(K_0 + \rho) + \rho) \left[\left(\frac{\delta_\rho}{\rho} + \frac{1}{2} \right) |\{u > 0\} \cap B_{\kappa\rho}| + \frac{1}{2} \int_{B_{\kappa\rho}} |\nabla u|^2 dx \right] \\ &\quad + c(K_0 \rho + \rho^2) (1 + c(K_0 \rho + \rho^2) \beta_2) |\{u > 0\} \cap B_{\kappa\rho}| \\ &\leq \beta_1 \beta_3 (c(K_0 + \rho) + \rho) \left(2c(K_0 + \rho) + \frac{1}{2} \right) \left[\int_{B_{\kappa\rho}} |\nabla u|^2 + |\{u > 0\} \cap B_{\kappa\rho}| \right]. \end{aligned}$$

Eventually, by choosing $K_0, \rho_0 < \bar{\rho}$ small enough so that

$$\beta_1 \beta_3 (c(K_0 + \rho_0) + \rho_0) \left(2c(K_0 + \rho_0) + \frac{1}{2} \right) \leq \eta/4,$$

we conclude that $u \equiv 0$ in $B_{\kappa\rho}$, for all $\rho \leq \rho_0$. \square

Remark 3.10. The statement of Lemma 3.9 and in particular (25) can be also equivalently stated as

$$\|u\|_{L^\infty(B_\rho(x_0))} \leq K_0 \rho \implies u \equiv 0 \text{ in } B_{\kappa\rho}(x_0) \cap B_R,$$

see for example [35, Remark 4.3]. In other words, Lemma 3.9 implies that if $x_0 \in \bar{\Omega}$, then there is a constant $C = C(R, \eta) > 0$, which can be taken independent of x_0 , such that

$$\sup_{B_\rho(x_0) \cap B_R} u \geq C\rho, \quad \text{and} \quad \int_{\partial B_\rho(x_0) \cap B_R} u d\mathcal{H}^2 \geq C\rho.$$

Lemma 3.11. *Let R, η, q, Ω and u be as in Lemma 3.9. The function u can be extended to a Lipschitz continuous function defined in the whole B_R , with Lipschitz constant $L = L(R, \eta)$. In particular, $\Omega = \{u > 0\} \subset B_R$ is an open set.*

Proof. We follow the approach of [38, Section 3.2], first proposed in [9].

Step 1. We prove an estimate on the nonnegative Radon measure $|\Delta u|$, namely

$$|\Delta u|(B_r) \leq Cr^2, \quad \text{for all } B_r \subset B_R$$

for a universal constant $C > 0$. Let $\psi \in C_c^\infty(B_r)$ for some $B_r \subset B_R$, with $\|\psi\|_{L^\infty} \leq c$, and we test the optimality of u against $u + \psi$, obtaining:

$$\int_{\{u>0\}} |\nabla u|^2 dx + \frac{q}{2} D(u^2, u^2) + f_\eta(|\{u > 0\}|) \leq \int_{\{u>0\}} |\nabla(u+\psi)|^2 dx + \frac{q}{2} D((u+\psi)^2, (u+\psi)^2) + f_\eta(|\{u+\psi > 0\}|),$$

which implies

$$-2 \int_{\{u>0\} \cap B_r} \nabla u \cdot \nabla \psi dx \leq \int_{B_r} |\nabla \psi|^2 dx + C_\eta |\{u = 0\} \cap B_r| + \frac{q}{2} \int_{\{u>0\} \cup B_r} \int_{\{u>0\} \cup B_r} P(x, y) dx dy$$

where

$$P(x, y) = \frac{4u(x)\psi(x)u^2(y) + 4u(x)\psi(x)\psi^2(y) + 2u^2(x)\psi^2(y) + 4u(x)\psi(x)u(y)\psi(y) + \psi^2(x)\psi^2(y)}{|x - y|}.$$

Recalling that $\|\psi\|_{L^\infty} \leq c$, that we can assume $r < 1$, using Lemma 2.4 and [16, Lemma 2.4], we obtain

$$(31) \quad -2 \int_{\{u>0\} \cap B_r} \nabla u \cdot \nabla \psi \, dx \leq \int_{B_r} |\nabla \psi|^2 \, dx + C_\eta |\{u = 0\} \cap B_r| + qCr^3.$$

We now set, for all $\varphi \in C_c^\infty(B_r)$, $\psi = Cr^{3/2} \|\nabla \varphi\|_{L^2}^{-1} \varphi$ and from (31) we deduce, for some $\tilde{C} > 0$

$$\left| \int_{\{u>0\} \cap B_r} \nabla u \cdot \nabla \varphi \, dx \right| \leq \tilde{C} r^{3/2} \|\nabla \varphi\|_{L^2(B_r)}$$

It is then enough to choose $\varphi \in C_c^\infty(B_{2r})$ with $\varphi \geq 0$ and $\varphi = 1$ in B_r and with $\|\nabla \varphi\|_{L^\infty(B_{2r})} \leq \frac{2}{r}$ (notice that this is compatible with the requirement $\|\psi\|_{L^\infty} \leq c$) to obtain, for some constant $C > 0$:

$$(32) \quad |\Delta u|(B_r) \leq |\Delta u|(\varphi) = \left| \int_{\{u>0\} \cap B_r} \nabla u \cdot \nabla \varphi \, dx \right| \leq \left| \int_{\{u>0\} \cap B_r} \nabla u \cdot \nabla \varphi \, dx \right| \leq Cr^2.$$

Step 2. We prove that the Laplacian estimate (32) of Step 1 entails (recall that $\mathcal{H}^2(\partial B_r) = 4\pi r^2$)

$$(33) \quad \frac{1}{4\pi r^2} \int_{\partial B_r} u \, d\mathcal{H}^2 \leq Cr,$$

for some constant $C > 0$. This follows from [9, Lemma 3.6], which assures that, for all $x_0 \in B_R$, it holds

$$(34) \quad \frac{1}{4\pi r^2} \int_{\partial B_r(x_0)} u \, d\mathcal{H}^2 - u(x_0) = \int_0^r \frac{1}{4\pi s^2} \Delta u(B_s(x_0)) \, ds.$$

It is then enough to put together (34) and (32) to obtain (33). We note that applying (33) to a point $x_0 \in \partial\{u > 0\} \cap B_R$ and passing to the limit $r \rightarrow 0$, we deduce that $u(x_0) = 0$, thus $\Omega = \{u > 0\}$ is an open set. *Step 3.* We conclude, using Step 2, that u is Lipschitz continuous in B_R , as in [38, Lemma 3.5], see also [9, Theorem 3.1 and 4.1]. \square

An immediate and fundamental consequence of Lemmas 3.9 and 3.11 is the following density estimate on Ω .

Lemma 3.12. *Let $R > 1$ and $\eta \in (0, 1)$. There exists $q_1 \in (0, q_0]$ such that for all $q \in (0, q_1]$, calling Ω an optimal set for problem (15) and u a positive normalized function attaining $E_{q, M, \eta}(\Omega)$, there exist positive constants $\theta = \theta(R, \eta)$ and $\rho_0 = \rho_0(R, \eta) < 1$ such that for every $x_0 \in \partial\Omega$ and every $\rho \leq \rho_0$, we have*

$$\theta \leq \frac{|\Omega \cap B_\rho(x_0)|}{|B_\rho|} \leq (1 - \theta).$$

Proof. Let us start from the lower bound. We can assume that $x_0 = 0 \in \partial\Omega = \partial\{u > 0\}$. Thus, the nondegeneracy condition of Remark 3.10 implies that

$$\|u\|_{L^\infty(B_{\rho/2})} \geq C \frac{\rho}{2}.$$

Thus, there is a point $y \in B_{\rho/2}$ such that $u(y) \geq C \frac{\rho}{2}$. On the other hand, the Lipschitz continuity of u , with constant $L = L(R, \eta)$, implies that $u > 0$ on a ball with radius $\frac{\rho}{2} \min\{1, \frac{C}{L}\}$, and so we conclude.

The upper bound can be obtained as in [2], see also [38, Section 5.1], with a few modifications. Precisely, let $x_0 = 0 \in \partial\Omega$ and consider the function h which is equal to u outside B_ρ , and inside B_ρ it is the solution of

$$\begin{cases} -\Delta h = \gamma_1 & \text{in } B_\rho, \\ h = u & \text{on } \partial B_\rho, \end{cases}$$

where $\gamma_1 = \gamma_1(\rho) := 2 \sup_x (\lambda_q - qv_u(x)) \|u\|_{L^\infty(B_\rho)} > 0$. As a consequence, we obtain that $-\Delta(h - u) = \gamma_1 - (\lambda_q u - qv_u)u \geq 0$ in B_ρ . In particular, we have that $u \leq h$ and $\{u > 0\} \subset \{h > 0\}$ in B_ρ . Moreover, since h is the torsion function multiplied by γ_1 with boundary datum u , recalling that $u(0) = 0$ and that u is Lipschitz continuous with constant $L = L(R, \eta)$, see Lemma 3.11, we deduce by a simple comparison argument that

$$(35) \quad \|h\|_{L^\infty(B_\rho)} \leq C_h \rho,$$

with a constant C_h depending only on R, η . Thus, testing the optimality of u with h , using also (13) and an integration by parts, we have

$$\begin{aligned} \frac{1}{\eta}|B_\rho \cap \{u = 0\}| + \frac{q}{2} \int_{B_\rho} \int_{B_\rho} \frac{h^2(x)h^2(y)}{|x-y|} dx dy + M \left| \int_{B_\rho} (h^2 - u^2) dx \right| &\geq \int_{B_\rho} |\nabla u|^2 dx - \int_{B_\rho} |\nabla h|^2 dx \\ &= \int_{B_\rho} |\nabla(u-h)|^2 dx + 2 \int_{B_\rho} (\nabla h \cdot \nabla(u-h)) dx \\ &= \int_{B_\rho} |\nabla(u-h)|^2 dx + 2 \int_{B_\rho} (-\Delta h)(u-h) dx + 2 \int_{B_\rho} (u-h) \frac{\partial h}{\partial \nu} d\mathcal{H}^2 \\ &= \int_{B_\rho} |\nabla(u-h)|^2 dx - 2\gamma_1 \int_{B_\rho} (h-u) dx. \end{aligned}$$

Let us first treat the terms not involving the gradient: thanks to (35), we can bound the term (using also the scaling of the Riesz energy and the fact that $\rho \leq 1$)

$$(36) \quad \frac{q}{2} \int_{B_\rho} \int_{B_\rho} \frac{h^2(x)h^2(y)}{|x-y|} dx dy \leq C_h^4 \rho^4 \rho^5 D(\chi_{B_1}, \chi_{B_1}) \frac{q}{2} \leq C_a |B_\rho| q,$$

for a constant $C_a > 0$, depending only on R, η . On the other hand, using again the bound L^∞ on h and u , we have

$$(37) \quad \int_{B_\rho} (h^2 - u^2) dx + \int_{B_\rho} (h-u) dx \leq (C_h + 1) \int_{B_\rho} h dx \leq (C_h + 1) C_h \rho |B_\rho| = C'_h \rho |B_\rho|,$$

for a constant $C'_h > 0$ depending only on R, η .

Let us now focus on the gradient term. By the Poincaré and Cauchy-Schwarz inequalities, we have

$$\int_{B_\rho} |\nabla(u-h)|^2 dx \geq \frac{C_d}{|B_\rho|} \left(\frac{1}{\rho} \int_{B_\rho} (h-u) dx \right)^2,$$

so in order to prove the upper bound in the claim, we first need to show that $\frac{1}{\rho |B_\rho|} \int_{B_\rho} (h-u) dx$ is bounded from below by a positive constant. Notice that, by the non-degeneracy of u (see Remark 3.10), we have

$$C\rho \leq \sup_{B_{\rho/2}} u \leq \sup_{B_{\rho/2}} h.$$

On the other hand, since $h(x) + \frac{\gamma_1}{6}|x|^2$ is harmonic in B_ρ , the Harnack inequality in B_ρ implies

$$C\rho \leq \sup_{B_{\rho/2}} h \leq C_d (h(x) + \gamma_1 \rho^2) \quad \text{for every } x \in B_{\frac{\rho}{2}}.$$

Thus, by taking ρ_0 such that $2C_d \rho_0 \gamma_1 \leq C$, we obtain that $h \geq C_d C\rho = \bar{C}\rho$ in $B_{\frac{\rho}{2}}$. On the other hand, if $L = L(R, \eta)$ is the Lipschitz constant of u (by Lemma 3.11), then for any $\varepsilon \in (0, 1)$, $u \leq L\varepsilon\rho$ in $B_{\varepsilon\rho}$. Then

$$\int_{B_\rho} (h-u) dx \geq \int_{B_{\varepsilon\rho}} (h-u) dx \geq (\bar{C}\rho - L\varepsilon\rho) |B_{\varepsilon\rho}|,$$

which, after choosing $\varepsilon \leq \frac{1}{2}$ small enough, shows that

$$\frac{1}{\rho} \int_{B_\rho} (h-u) dx \geq C_0 |B_\rho|,$$

for some constant $C_0 > 0$ depending only on R, η and thus

$$\int_{B_\rho} |\nabla(u-h)|^2 dx \geq C_b |B_\rho|,$$

for some $C_b > 0$, depending only on R, η .

At this point, using also (37) and (36), we have

$$C_b |B_\rho| \leq \frac{1}{\eta} |B_\rho \cap \{u = 0\}| + C_a q |B_\rho| + (M + 2\gamma_1) \rho |B_\rho| \leq \frac{1}{\eta} |B_\rho \cap \{u = 0\}| + C_a q |B_\rho| + C_g \rho |B_\rho|,$$

for a universal constant $C_g > 0$, recalling that M is fixed (see Remark 3.3) and that γ_1 is uniformly bounded by a constant depending only on $\|u\|_{L^\infty(B_R)}$, which in turn is uniform in q (see Lemma 2.4). It is then enough to take

$$q_1 \leq \min \left\{ q_0, \frac{C_b}{4C_a} \right\}, \quad \rho_0 \leq \frac{C_b}{4C_g},$$

and we obtain that

$$\frac{C_b}{4} |B_\rho| \leq \frac{1}{\eta} |B_\rho \cap \{u = 0\}|,$$

which entails the density estimate from above, so the claim is proved. \square

Remark 3.13. Notice that the constants determining the Lipschitz regularity and the density estimates of the previous result do not depend on q for q small enough.

From now on, q_1 always refers to the constant defined in Lemma 3.12.

3.4. Equivalence between the minimizations of E_q and $E_{q,M,\eta}$. In this section we show that unconstrained minima of $E_{q,M,\eta}$ and volume constrained minima of E_q (or equivalently $E_{q,M}$) are actually the same. We begin by showing that for q small, the minimizers of $E_{q,M,\eta}$ in B_R are close to a ball in L^∞ . To do that, we first start with an estimate that assures the L^1 -proximity of an optimal set for problem (16) to a ball with radius not too large.

Lemma 3.14. *Let $R > 2$, $q \in (0, q_1]$ and $\eta \in (0, 1)$. Let $\Omega = \Omega_{q,M,\eta}$ be an optimal set for (16) such that $\Omega = \{u_\Omega > 0\}$, where $u_\Omega \in H_0^1(\Omega)$ is the (positive) function attaining $E_q(\Omega) = E_{q,M}(\Omega)$, and $B = B_{q,M,\eta}$ a ball of measure $|\Omega|$ attaining the Fraenkel asymmetry for Ω , namely such that*

$$\mathcal{A}(\Omega) = \frac{|\Omega \Delta B|}{|\Omega|}.$$

Then, setting $u_B \in H_0^1(B)$ the function attaining $E_q(B)$, normalized so that $\|u_B\|_{L^2(B)} = \|u_\Omega\|_{L^2(\Omega)} = 1$, we have

$$|\{u_B > 0\} \Delta \{u_\Omega > 0\}|^2 \leq Cq|\Omega|^{\frac{13}{3}}.$$

for some universal constant $C > 0$.

Proof. Let w_B be the normalized first eigenfunction of the Dirichlet Laplacian in B , and note that it is an admissible competitor for $E_q(B)$. Thanks to the quantitative Faber–Krahn inequality (Theorem 2.11), we have

$$\begin{aligned} |\Omega|^{2/3} \int_\Omega |\nabla u_\Omega|^2 dx &\geq |\Omega|^{2/3} \lambda_0(\Omega) \\ (38) \quad &\geq |\Omega|^{2/3} \lambda_0(B) + \hat{\sigma} \frac{|\{u_B > 0\} \Delta \{u_\Omega > 0\}|^2}{|\Omega|^2} \\ &= |\Omega|^{2/3} \int_B |\nabla w_B|^2 dx + \hat{\sigma} \frac{|\{u_B > 0\} \Delta \{u_\Omega > 0\}|^2}{|\Omega|^2}. \end{aligned}$$

From the optimality of Ω , we deduce,

$$\begin{aligned} \int_\Omega |\nabla u_\Omega|^2 dx + \frac{q}{2} \int_\Omega \int_\Omega \frac{u_\Omega^2(x) u_\Omega^2(y)}{|x-y|} dx dy + f_\eta(|\Omega|) &\leq E_q(B) + f_\eta(|B|) \\ &\leq \int_B |\nabla w_B|^2 dx + \frac{q}{2} \int_B \int_B \frac{w_B^2(x) w_B^2(y)}{|x-y|} dx dy + f_\eta(|B|), \end{aligned}$$

and using also (38) we obtain

$$\begin{aligned} \hat{\sigma} \frac{|\{u_B > 0\} \Delta \{u_\Omega > 0\}|^2}{|\Omega|^2} &\leq |\Omega|^{2/3} \int_\Omega |\nabla u_\Omega|^2 dx - |\Omega|^{2/3} \int_B |\nabla w_B|^2 dx \\ &\leq |\Omega|^{2/3} \frac{q}{2} \left(D(u_B^2, u_B^2) - D(u_\Omega^2, u_\Omega^2) \right) \leq |\Omega|^{2/3} \frac{q}{2} D(u_B^2, u_B^2). \end{aligned}$$

Now, we recall that we have an uniform L^∞ bound on u_B (see Lemma 2.4), thus, using also the scaling of the Riesz functional,

$$D(u_B^2, u_B^2) \leq C \int_B \int_B \frac{dx dy}{|x-y|} = C|\Omega|^{\frac{5}{3}},$$

for some constant $C > 0$ depending only on the L^∞ norm of u_B , which is uniformly bounded, see Lemma 2.4, thus C is universal.

The previous two estimates lead to

$$|\{u_B > 0\} \Delta \{u_\Omega > 0\}|^2 \leq \frac{C}{\hat{\sigma}} |\Omega|^{\frac{13}{3}} \frac{q}{2},$$

so the claim is proved. \square

A simple but important consequence of the previous result is the following lemma, stating that the measure of the ball $B = B_{q,M,\eta}$ (to which any optimal set $\Omega_{q,M,\eta}$ is L^1 -close) is not too large, as we show in the next result.

Lemma 3.15. *Let $R > 2$. There exists $\eta_1 \in (0, 1)$ such that for all $q \in (0, q_1]$ and $\eta \leq \eta_1$, we have that any optimal set $\Omega_{q,M,\eta}$ for problem (16), such that $\Omega_{q,M,\eta} = \{u > 0\}$, where $u \in H_0^1(\Omega)$ is any (positive) function attaining $E_q(\Omega) = E_{q,M}(\Omega)$, satisfies*

$$(39) \quad |\Omega_{q,M,\eta}| \leq |B_2|, \quad |\Omega_{q,M,\eta} \Delta B_{q,M,\eta}| \leq c_1 q,$$

for some universal constant $c_1 > 0$.

Proof. Let us suppose for the sake of contradiction that $|\Omega_{q,M,\eta}| > |B_2|$. We are then going to reach a contradiction as long as

$$1/\eta \geq E_{q_0}(B_1).$$

Since the functional

$$q \mapsto E_{q,M,\eta}(\Omega_{q,M,\eta}),$$

is nondecreasing, we obtain

$$\sup_{q \in (0, q_0)} E_{q,M,\eta}(\Omega_{q,M,\eta}) = E_{q_0,M,\eta}(\Omega_{q_0,M,\eta}) \leq E_{q_0,M,\eta}(B_1) = E_{q_0}(B_1),$$

recalling that the optimal function for $E_{q_0}(B_1)$ is with unit L^2 norm. On the other hand, using the Faber–Krahn inequality and the positivity of E , since $|\Omega_{q,M,\eta}| > |B_2|$ we have

$$E_{q_0}(B_1) \geq E_{q_0,M,\eta}(\Omega_{q_0,M,\eta}) \geq \frac{1}{\eta} (|\Omega_{q_0,M,\eta}| - |B_1|) \geq \frac{1}{\eta} (|B_2| - |B_1|).$$

By choosing η_1 such that $\eta_1 < 1$ and

$$\frac{(|B_2| - |B_1|)}{\eta_1} > E_{q_0}(B_1),$$

we reach the desired contradiction. The second part of the claim then follows from Lemma 3.14. \square

We note that in the above lemma, η_1 does not depend on R . Next we show that, for q small, the boundary of any optimizer $\Omega_{q,M,\eta}$ (such that $\Omega_{q,M,\eta} = \{u > 0\}$, where $u \in H_0^1(\Omega)$ is the (positive) function attaining $E_q(\Omega) = E_{q,M}(\Omega)$) is close to the one of the corresponding optimal ball of the same measure $B_{q,M,\eta}$ in the definition of Fraenkel asymmetry, with respect to the Hausdorff distance d_H (see [3, Definition 4.4.9] for more details about the Hausdorff distance).

Lemma 3.16. *Under the assumptions of Lemma 3.15, for all $\delta > 0$ there exists $q_\delta = q_\delta(R, \eta) \in (0, q_1]$ such that for all $q \leq q_\delta$, we have*

$$\text{dist}_H(\partial\Omega_{q,M,\eta}, \partial B_{q,M,\eta}) \leq \delta.$$

Proof. This follows exactly as in [35, Lemma 5.4], we report here the proof for the sake of completeness. We fix $\delta > 0$ and call $B_\delta(B_{q,M,\eta}) := B_{q,M,\eta} + B_\delta$ the δ -neighborhood of $B_{q,M,\eta}$. First of all, we aim to prove that $\Omega_{q,M,\eta} \subset B_\delta(B_{q,M,\eta})$. If $\Omega_{q,M,\eta} \setminus B_\delta(B_{q,M,\eta})$ is empty, then there is nothing to prove. Otherwise there exists $x \in \Omega_{q,M,\eta} \setminus B_\delta(B_{q,M,\eta})$ so that by Lemma 3.12 there exists $\rho_0(R, \eta)$ such that for $\rho \leq \rho_1 := \min\{\rho_0(R, \eta), \delta\}$ it holds

$$|B_1| \theta \rho^3 \leq |B_\rho(x) \cap \Omega_{q,M,\eta}| \leq |\Omega_{q,M,\eta} \setminus B_{q,M,\eta}| \leq c_1 q,$$

where the last estimate follows from Lemma 3.15 and precisely (39). Notice that the choice of $\rho_1 \leq \delta$ assures that $|B_\rho(x) \cap \Omega_{q,M,\eta}| \leq |\Omega_{q,M,\eta} \setminus B_{q,M,\eta}|$. In conclusion, choosing $\rho = \rho_1$, we have

$$|B_1| \theta \rho_1^3 \leq c_1 q,$$

which is not possible as soon as

$$q \leq q_\delta := \frac{|B_1| \theta}{c_1} \rho_1^3.$$

With the same argument, thanks to the outer density estimate from Lemma 3.12 and again to the L^1 proximity from Lemma 3.15, we can show also that $B_R \setminus \Omega_{q,M,\eta} \subset B_\delta(B_R \setminus B_{q,M,\eta})$, with the same notation as above. This concludes the proof. \square

Remark 3.17. It is worth noting that the constant q_δ in the lemma above depends also on R . This is one of the main obstacles while trying to get rid of the equiboundedness assumption of Theorem 3.1.

Remark 3.18. In view of the previous result, we fix $q_2(R) := \min\{q_1, q_\delta\}$, where q_δ is the constant from Lemma 3.16 with the choice of $\delta := 1/2$.

Therefore, if $q \in (0, q_2]$, in the proof of the next Theorem 3.19, we are allowed to inflate a set without touching the boundary of the geometric constraint $\Omega \subset B_R$.

We can now show the equivalence between the constrained and the unconstrained problems. We recall that the constant M has been already fixed, see Remark 3.3.

Theorem 3.19. *There exists a universal constant $R_0 \geq 10$ such that, for all $R \geq R_0$, there exists $q_3 = q_3(R) \leq q_2$ and $\eta_2 = \eta_2(R) \leq \eta_1$ such that, for all $\eta \leq \eta_2$ and $q \in (0, q_3]$, we have that*

$$\min \{E_{q,M,\eta}(\Omega) : \Omega \subset B_R\} = \inf \{E_q(\Omega) : \Omega \subset B_R, |\Omega| = |B_1|\}.$$

As a consequence, problems (10) and (16) are equivalent for these values of q and η .

Proof. It is easy to check that

$$\min \{E_{q,M,\eta}(\Omega) : \Omega \subset B_R\} \leq \inf \{E_q(\Omega) : \Omega \subset B_R, |\Omega| = |B_1|\} =: \mu(q, R),$$

as the two functionals coincide on sets of measure $|B_1|$, thanks to the definition of f_η . Then, if the reversed inequality holds, it follows that on the set of minimizers (of the first or of the second problem) the two functionals do coincide, that is, problems (10) and (16) are equivalent.

We prove the claim of the theorem by contradiction. Let

$$\Omega_{q,M,\eta} \subset B_R, \quad \sigma_{q,M,\eta} \in \mathbb{R}, \quad |\Omega_{q,M,\eta}| = |B_1| + \sigma_{q,M,\eta}, \quad E_{q,M,\eta}(\Omega_{q,M,\eta}) < \mu,$$

and we also note that, $\mu \leq E_q(B_1)$, by definition of infimum. We moreover assume, without loss of generality, that $\Omega_{q,M,\eta}$ are minimizers for problem (16). We treat separately the case $\sigma_{q,M,\eta} > 0$ and $\sigma_{q,M,\eta} < 0$.

Case $\sigma_{q,M,\eta} > 0$. We first observe that $\sigma_{q,M,\eta} \rightarrow 0$ as $\eta \rightarrow 0$. Indeed (recalling also that $E_q = E_{q,M}$ thanks to Proposition 3.2)

$$E_{q,M,\eta}(\Omega_{q,M,\eta}) = E_q(\Omega_{q,M,\eta}) + \frac{1}{\eta} \sigma_{q,M,\eta}$$

and so

$$0 \leq \frac{\sigma_{q,M,\eta}}{\eta} = E_{q,M,\eta}(\Omega_{q,M,\eta}) - E_q(\Omega_{q,M,\eta}) \leq E_q(B_1),$$

using the assumption $E_{q,M,\eta}(\Omega_{q,M,\eta}) \leq \mu \leq E_q(B_1)$ and the positivity of the energy. This implies that $\sigma_{q,M,\eta} \rightarrow 0$ as $\eta \rightarrow 0$.

Let now $\lambda_{q,M,\eta} < 1$ be such that $|\lambda_{q,M,\eta} \Omega_{q,M,\eta}| = |B_1|$, therefore

$$\lambda_{q,M,\eta} = 1 - \frac{\sigma_{q,M,\eta}}{3|B_1|} + C\sigma_{q,M,\eta}^2,$$

for some $C \in \mathbb{R}$.

We call $u = u_{q,M,\eta}$ an optimal normalized function attaining $E_q(\Omega_{q,M,\eta})$, thus the function

$$\tilde{u}(y) = \lambda_{q,M,\eta}^{-\frac{3}{2}} u\left(\frac{y}{\lambda_{q,M,\eta}}\right), \quad y \in \lambda_{q,M,\eta} \Omega_{q,M,\eta},$$

is an admissible competitor with unitary L^2 -norm for $E_q(\lambda_{q,M,\eta} \Omega_{q,M,\eta})$. We have the following scalings

$$\begin{aligned} \int_{\lambda_{q,M,\eta} \Omega_{q,M,\eta}} |\nabla \tilde{u}(y)|^2 dy &= \lambda_{q,M,\eta}^{-2} \int_{\Omega_{q,M,\eta}} |\nabla u(x)|^2 dx, \\ D(\tilde{u}^2, \tilde{u}^2) &= \int_{\lambda_{q,M,\eta} \Omega_{q,M,\eta}} \int_{\lambda_{q,M,\eta} \Omega_{q,M,\eta}} \frac{\tilde{u}^2(x) \tilde{u}^2(y)}{|x-y|} dx dy \\ &= \lambda_{q,M,\eta}^{-1} \int_{\Omega_{q,M,\eta}} \int_{\Omega_{q,M,\eta}} \frac{u^2(w) u^2(z)}{|w-z|} dw dz \\ &= D(u^2, u^2). \end{aligned}$$

Since the new set $\lambda_{q,M,\eta}\Omega_{q,M,\eta}$ is now admissible in the constrained minimization problem (10), using the above scaling we obtain

$$\begin{aligned}
E_{q,M,\eta}(\Omega_{q,M,\eta}) &= E_q(\Omega_{q,M,\eta}) + \frac{\sigma_{q,M,\eta}}{\eta} \\
&< \mu \\
&\leq E_q(\lambda_{q,M,\eta}\Omega_{q,M,\eta}) \\
&\leq \int_{\lambda_{q,M,\eta}\Omega_{q,M,\eta}} |\nabla \tilde{u}(y)|^2 dy + \frac{q}{2} D(\tilde{u}^2, \tilde{u}^2) \\
&= \lambda_{q,M,\eta}^{-2} \int_{\Omega_{q,M,\eta}} |\nabla u(x)|^2 dx + \lambda_{q,M,\eta}^{-1} \frac{q}{2} D(u^2, u^2) \\
&= \int_{\Omega_{q,M,\eta}} |\nabla u(x)|^2 dx \left(1 + \frac{2\sigma_{q,M,\eta}}{3|B_1|} + C\sigma_{q,M,\eta}^2 \right) \\
&\quad + \frac{q}{2} D(u^2, u^2) \left(1 + \frac{\sigma_{q,M,\eta}}{3|B_1|} + C\sigma_{q,M,\eta}^2 \right),
\end{aligned}$$

we deduce that (up to increasing C , recalling also that $E_q(\Omega_{q,M,\eta}) \leq E_q(B_1)$)

$$\begin{aligned}
\frac{\sigma_{q,M,\eta}}{\eta} &< \int_{\Omega_{q,M,\eta}} |\nabla u(x)|^2 dx \left(\frac{2\sigma_{q,M,\eta}}{3|B_1|} \right) + \frac{q}{2} D(u^2, u^2) \left(\frac{\sigma_{q,M,\eta}}{3|B_1|} \right) + 2E_q(B_1)C\sigma_{q,M,\eta}^2 \\
&\leq \frac{\sigma_{q,M,\eta}}{3|B_1|} 2E_q(\Omega_{q,M,\eta}) + C\sigma_{q,M,\eta}^2.
\end{aligned}$$

Thus, for some universal $C > 0$,

$$\frac{1}{\eta} \leq CE_q(\Omega_{q,M,\eta}) + C\sigma \leq CE_q(B_1),$$

which leads to a contradiction as soon as $\eta_2 < \frac{1}{C(E_q(B_1))}$.

Case $\sigma_{q,M,\eta} < 0$. For this case let us call

$$\rho_{q,M,\eta} := \left(1 + \frac{\sigma_{q,M,\eta}}{|B_1|} \right)^{-1/3},$$

so that $|\rho_{q,M,\eta}\Omega_{q,M,\eta}| = |B_1|$.

We recall from the previous sections that a minimizer $\Omega_{q,M,\eta}$ for $E_{q,M,\eta}$ exists, and by Lemma 3.16, up to taking $q_3 \leq q_2$ as in Remark 3.18, and $\eta_2 < \eta_1$ as in Lemma 3.15, the rescaled set $\rho_{q,M,\eta}\Omega_{q,M,\eta}$ is still contained in B_R , as soon as R_0 is big enough.

In fact, we show that $\sigma_{q,M,\eta} \geq -\frac{3}{4}|B_1|$, thus $\rho_{q,M,\eta} \leq 4^3$ (hence, recalling Remark 3.18 we can take any $R_0 > 2 \cdot 4^3$).

If, for the sake of contradiction, $\sigma_{q,M,\eta} < -\frac{3}{4}|B_1|$, then $|\Omega_{q,M,\eta}| \leq \frac{1}{4}|B_1|$ and $|2^{1/3}\Omega_{q,M,\eta}| \leq \frac{1}{2}|B_1|$. The optimality of $\Omega_{q,M,\eta}$ entails,

$$\begin{aligned}
E_{q,M,\eta}(\Omega_{q,M,\eta}) &= E_q(\Omega_{q,M,\eta}) + \eta\sigma_{q,M,\eta} \leq E_q(2^{1/3}\Omega_{q,M,\eta}) + \eta(2|\Omega_{q,M,\eta}| - |B_1|) \\
&= E_q(2^{1/3}\Omega_{q,M,\eta}) + \eta(|B_1| + 2\sigma_{q,M,\eta}),
\end{aligned}$$

which is equivalent to say

$$E_q(\Omega_{q,M,\eta}) - E_q(2^{1/3}\Omega_{q,M,\eta}) \leq \eta(|B_1| + \sigma_{q,M,\eta}).$$

Also in view of Remark 2.3, we note that

$$E_q(2^{1/3}\Omega_{q,M,\eta}) = 2^{-2/3}E_q(\Omega_{q,M,\eta}),$$

thus we have (using also the Faber-Krahn inequality)

$$E_q(\Omega_{q,M,\eta}) - E_q(2^{1/3}\Omega_{q,M,\eta}) \geq (1 - 2^{-2/3})\lambda_0(\Omega_{q,M,\eta}) \geq (1 - 2^{-2/3})\lambda_0(B_1).$$

Finally, $(|B_1| + \sigma_{q,M,\eta}) \leq |B_1|$, so we reach a contradiction as soon as $\eta \leq \eta_2$ and

$$\eta_2 < (1 - 2^{-\frac{2}{3}}) \frac{\lambda_0(B_1)}{|B_1|}.$$

Let us define the function, using the same notations for u, \tilde{u} as in the previous case,

$$g: [1, \rho_{q,M,\eta}] \rightarrow \mathbb{R}, \quad g(r) = \int_{r\Omega_{q,M,\eta}} |\nabla \tilde{u}|^2 dx + \frac{q}{2} \int_{r\Omega_{q,M,\eta}} \int_{r\Omega_{q,M,\eta}} \frac{\tilde{u}^2(x)\tilde{u}^2(y)}{|x-y|} dx dy + \eta(r^3|\Omega_{q,M,\eta}| - |B_1|).$$

We want to show that the minimum of the function g is attained at $r = \rho := \rho_{q,M,\eta}$. This is equivalent to show that for some η the inequality

$$g(r) \geq \int_{\rho\Omega_{q,M,\eta}} |\nabla \tilde{u}|^2 dx + \frac{q}{2} \int_{\rho\Omega_{q,M,\eta}} \int_{\rho\Omega_{q,M,\eta}} \frac{\tilde{u}^2(x)\tilde{u}^2(y)}{|x-y|} dx dy, \quad \text{for all } r \in [1, \rho],$$

holds true. Up to rearranging the terms, and by the rescaling of the involved integrals, such an inequality reads as

$$\eta \left(1 - \left(\frac{r}{\rho} \right)^3 \right) \leq \int_{\rho\Omega_{q,M,\eta}} |\nabla \tilde{u}|^2 \left(\left(\frac{r}{\rho} \right)^{-2} - 1 \right) + \frac{q}{2} D(\tilde{u}^2, \tilde{u}^2) \left(\left(\frac{r}{\rho} \right)^{-1} - 1 \right).$$

Setting $t := \frac{r}{\rho} < 1$, and observing that $r^3 |\Omega_{q,M,\eta}| = t^3$, the last inequality is equivalent to

$$\eta \leq \frac{\int_{\rho\Omega_{q,M,\eta}} |\nabla \tilde{u}|^2 dx (t^{-2} - 1) + \frac{q}{2} D(\tilde{u}^2, \tilde{u}^2) (t^{-1} - 1)}{1 - t^3}.$$

It is easy to check that the right hand side is bounded from below by the function

$$t \mapsto \lambda_0(B) \frac{t^{-2} - 1}{1 - t^3}, \quad t \in (0, 1),$$

which is a function strictly decreasing in its domain and with infimum given by

$$\lim_{t \rightarrow 1^-} \lambda_0(B) \frac{t^{-2} - 1}{1 - t^3} = \frac{2}{3} > 0.$$

Thus it is enough to take $\eta \leq \eta_2 \leq 2/3$ and we immediately deduce that g has minimum for $r = \rho$. This concludes the proof. \square

We highlight that if $\eta > 0$ is such that Theorem 3.19 holds true, we can freely assume the equivalence of the volume-constrained minimization of E_q and the unconstrained one for $E_{q,M,\eta}$ (since M has been already fixed, see Remark 3.3). On the other hand, we stress that this choice of η depends, in all our estimates, on R .

4. OPTIMALITY CONDITIONS AND IMPROVEMENT OF FLATNESS

We have now the following picture. We know that minimizers for the auxiliary functional $E_{q,M,\eta}$ exist and (with the choices we made for M, η) are the same of those of the volume constrained functional E_q . These minimizers satisfy density estimates which are uniform with respect to $q \leq q_3$ (see Lemma 3.12 for the density estimate and Theorem 3.19 for the constant q_3). Moreover in Lemma 3.16 we have shown that such minimizers are close in Hausdorff distance to a given ball (any ball achieving the minimum in the definition of Fraenkel asymmetry of Ω). We now improve such a regularity with the final scope of showing that optimal sets are uniform $C^{2,\alpha}$ parametrizations on the boundary of the ball. The results and the proofs in this section are borrowed (with nontrivial adjustments) from results in [2, Theorem 4.5 and Theorem 4.8], [1, Theorem 2]. We begin with the following theorem, in which we use the notation $\partial^* F$ for the reduced boundary of a set of finite perimeter F .

Theorem 4.1. *Let $q \in (0, q_3]$, let Ω be a minimizer of (2), and let u be an optimal function attaining $E_q(\Omega)$, thus also solution of (5). Then we have that:*

(i) *There is a Borel function $\mu_u: \partial\Omega \rightarrow \mathbb{R}$ such that, in the sense of the distributions, one has*

$$(40) \quad -\Delta u = (\lambda_q - qv_u)u - \mu_u \mathcal{H}^2 \llcorner \partial\Omega, \quad \text{in } B_R.$$

(ii) *There exist constants $0 < c < C < +\infty$, depending on R , such that $c \leq \mu_u \leq C$.*

(iii) *For all points $\bar{x} \in \partial^* \Omega = \partial^* \{u > 0\}$, the measure theoretic inner unit normal $\nu_u(\bar{x})$ is well defined and, as $\rho \rightarrow 0$,*

$$\frac{\Omega - \bar{x}}{\rho} \rightarrow \{x : x \cdot \nu_u(\bar{x}) \geq 0\}, \quad \text{in } L^1(B_R).$$

(iv) *For \mathcal{H}^2 almost all $\bar{x} \in \partial^* \{u > 0\}$ we have*

$$\frac{u(\bar{x} + \rho x)}{\rho} \rightarrow \mu_u(\bar{x}) (x \cdot \nu_u(\bar{x}))_+, \quad \text{in } W^{1,p}(B_R) \text{ for every } p \in [1, +\infty).$$

(v) $\mathcal{H}^2(\partial\Omega \setminus \partial^* \Omega) = 0$.

Proof. The proof is essentially identical to that in [2, Section 4]. We only have to check that our hypotheses match with those in [2]. First by Lemma 2.4 u satisfies

$$-\Delta u - Q(x) = 0 \quad \text{in } \mathcal{D}'(\Omega),$$

where $Q = (\lambda_q - qv_u)u \in L^\infty(\Omega)$ and $u \in H_0^1(\Omega)$. Hence, by repeating the proof of [2, Theorem 4.5] or by directly applying [12, Proposition 2.3] one obtains that there exists a positive Radon measure concentrated on $\partial\Omega$ that we denote $\mu_u \mathcal{H}^2 \llcorner \partial\Omega$. Moreover, thanks to the non-degeneracy, see Remark 3.10 and the Lipschitz continuity of u we have that there exist constant $C > c > 0$ depending on q and R such that

$$c \leq \frac{1}{r} \int_{\partial B_r} u \, d\mathcal{H}^2 \leq C.$$

Hence we can work under the hypotheses of [2, Theorem 4.5] so that μ_u is a density of a Radon measure on $\partial\Omega$ and, denoting still with μ_u the function defining it, μ_u satisfies (i) – (v). \square

4.1. The structure of μ_u : blow up limits. We show now the following result.

Proposition 4.2. *Let Ω be a minimizer of (10). Then function $\mu_u : \partial\Omega \rightarrow \mathbb{R}$ found in Theorem 4.1 is constant on $\partial^*\Omega$.*

Proof. The proof follows the path of [35, Theorem 6.5], in turn inspired by [1]. Due to the nonlocal term, we will have to perform some new and non-straightforward computations.

We reason by contradiction and we assume that there exists $x_0, x_1 \in \partial^*\Omega$ such that

$$\mu_u(x_0) < \mu_u(x_1).$$

Then we construct a family of volume preserving diffeomorphisms as follows: let $\kappa < 1$ and $\rho < 1$ and let $\varphi \in C_0^1(B_1(0))$ be a non-null, radially symmetric function supported in $B_1(0)$. We define

$$\tau_{\rho, \kappa}(x) = \tau(x) = x + \sum_{i \in \{0,1\}} (-1)^i \kappa \rho \varphi \left(\frac{|x - x_i|}{\rho} \right) \nu_{x_i} \chi_{B_\rho(x_i)},$$

where ν_{x_i} are the measure theoretic inner normals to $\partial^*\Omega$ at x_i , $i = 1, 2$.

It is easy to notice that τ is indeed a diffeomorphism for ρ and κ small enough and that $\tau(x) - x$ vanishes outside $B_\rho(x_0) \cup B_\rho(x_1)$. Moreover we have:

$$(41) \quad \nabla \tau(x) = Id + \sum_{i \in \{0,1\}} (-1)^i \kappa \varphi' \left(\frac{|x - x_i|}{\rho} \right) \frac{x - x_i}{|x - x_i|} \otimes \nu_{x_i} \chi_{B_\rho(x_i)},$$

so that³

$$\det(\nabla \tau(x)) = 1 + \sum_{i \in \{0,1\}} (-1)^i \kappa \varphi' \left(\frac{|x - x_i|}{\rho} \right) \frac{x - x_i}{|x - x_i|} \cdot \nu_{x_i} \chi_{B_\rho(x_i)} + o(\kappa).$$

We call $\Omega_\rho = \tau(\Omega)$. We aim to show that for κ, ρ small enough it holds $E_{q,M,\eta}(\Omega_\rho) < E_{q,M,\eta}(\Omega)$, hence contradicting the minimality of Ω . To do that, we deal with the first variation of each term of the sum defining $E_{q,M,\eta}$. We stress that the computations regarding the volume and the Dirichlet energy contributions are identical to those performed originally in [1] (see also [7] and [15], where the same idea is applied). Moreover, exactly as in the proof of [35, Theorem 6.5] one obtains that

$$(42) \quad f_\eta(\Omega_\rho) - f_\eta(\Omega) = o(\rho^3), \quad \text{as } \rho \rightarrow 0,$$

and that

$$(43) \quad \frac{1}{\rho^3} \left(\int_{\Omega_\rho} |\nabla u_\rho|^2 \, dx - \int_{\Omega} |\nabla u|^2 \, dx \right) \leq \kappa (\mu_u^2(x_0) - \mu_u^2(x_1)) C(\varphi) + o_\rho(1) + o(\kappa),$$

where u_ρ and u are the functions attaining $E_q(\Omega_\rho)$ and $E_q(\Omega)$ respectively, and

$$C(\varphi) = \int_{B_1(0) \cap \{y \cdot \nu = 0\}} \varphi(|y|) \, d\mathcal{H}^2(y) = - \int_{B_1(0) \cap \{y \cdot \nu > 0\}} \varphi'(|y|) \frac{y \cdot \nu}{|y|} \, dy,$$

with the last equality that follows from the Divergence Theorem, recalling that ν is a inner normal and

$$\operatorname{div}(\varphi(|y|)\nu) = \varphi'(|y|) \frac{y \cdot \nu}{|y|}.$$

Notice also that by the radial symmetry of φ the value of $C(\varphi)$ is not affected by the choice of ν .

³We are using the formula $\det(Id + \xi A) = 1 + \operatorname{trace}(A)\xi + o(\xi)$ for a matrix $A \in \mathbb{R}^{N \times N}$.

We are left to compute the variation of the nonlocal term $D(\cdot, \cdot)$. This is the major technical difference with respect to the proof of [35, Theorem 6.5]. We claim that (recalling that u_ρ and u are the functions attaining $E_q(\Omega_\rho)$ and $E_q(\Omega)$ respectively)

$$(44) \quad \frac{1}{\rho^3} (D(u_\rho^2, u_\rho^2) - D(u^2, u^2)) = o(\kappa) + o(\rho).$$

Once that (44) is proved, the conclusion then readily follows: by minimality of Ω and thanks to (42), (43) and (44) we have that

$$\begin{aligned} 0 &\leq E_{q,M,\eta}(\Omega_\rho) - E_{q,M,\eta}(\Omega) \\ &\leq \kappa \rho^3 C(\varphi) \left((\mu_u(x_0)^2 - \mu_u(x_1)^2) \right) + o(\rho^3) + \rho^3 o(\kappa). \end{aligned}$$

Since from the assumptions there holds $\mu_u(x_0)^2 - \mu_u(x_1)^2 < 0$ we get the desired contradiction by choosing ρ and κ small enough.

It remains to show the validity of (44). To do so, we set

$$\tilde{u}(x) = u(\tau^{-1}(x)) \quad \text{and} \quad \tilde{w}(x) = v_{\tilde{u}}(x) \tilde{u}(x)^2,$$

where $v_u(x) = \int_\Omega \frac{u^2(y)}{|x-y|} dy$. With such a notation in force we compute, using also formula (41),

$$(45) \quad \begin{aligned} \frac{1}{\rho^3} (D(u_\rho^2, u_\rho^2) - D(u^2, u^2)) &= \frac{1}{\rho^3} \left(\int_{\Omega_\rho} \tilde{w} dx - \int_\Omega w dx \right) \\ &= \frac{1}{\rho^3} \int_\Omega (\tilde{w}(\tau(x)) \det(\nabla \tau(x)) - w(x)) dx \\ &= \frac{1}{\rho^3} \int_\Omega (\tilde{w}(\tau(x)) - w(x)) dx \\ &\quad + \frac{1}{\rho^3} \int_\Omega \tilde{w}(\tau(x)) \sum_{i \in \{0,1\}} (-1)^i \kappa \varphi' \left(\frac{|x-x_i|}{\rho} \right) \frac{x-x_i}{|x-x_i|} \cdot \nu_{x_i} \chi_{B_\rho(x_i)} dx + o(\kappa). \end{aligned}$$

We observe that since $\tilde{w}(x) = v_{\tilde{u}}(x) \tilde{u}(x)^2$, with $v_{\tilde{u}}$ uniformly bounded and \tilde{u} Lipschitz continuous in Ω , then $|\tilde{w}(\tau(x))| \leq C \rho^2$ in $\Omega \cap B_\rho(x_i)$, whenever $\tilde{u}(x_i) = 0$. With this in mind, we can compute

$$(46) \quad \begin{aligned} &\left| \frac{1}{\rho^3} \int_\Omega \tilde{w}(\tau(x)) \sum_{i \in \{0,1\}} (-1)^i \kappa \varphi' \left(\frac{|x-x_i|}{\rho} \right) \frac{x-x_i}{|x-x_i|} \cdot \nu_{x_i} \chi_{B_\rho(x_i)} dx \right| \\ &= \left| \frac{1}{\rho^3} \sum_{i \in \{0,1\}} (-1)^i \kappa \int_{\Omega \cap B_\rho(x_i)} \tilde{w}(\tau(x)) \varphi' \left(\frac{|x-x_i|}{\rho} \right) \frac{x-x_i}{|x-x_i|} \cdot \nu_{x_i} dx \right| \\ &\leq \left| \frac{C}{\rho} \sum_{i \in \{0,1\}} (-1)^i \kappa \int_{\Omega \cap B_\rho(x_i)} \varphi' \left(\frac{|x-x_i|}{\rho} \right) \frac{x-x_i}{|x-x_i|} \cdot \nu_{x_i} dx \right| \\ &= C \kappa \rho^2 \left| \int_{\frac{\Omega-x_i}{\rho} \cap B_1} \varphi'(|y|) \frac{y}{|y|} \cdot \nu_{x_i} dx \right| = o(\rho). \end{aligned}$$

Moreover

$$\begin{aligned} \frac{1}{\rho^3} \int_\Omega (\tilde{w}(\tau(x)) - w(x)) dx &= \frac{1}{\rho^3} \int_\Omega u^2(x) \left(\int_{\Omega_\rho} \frac{\tilde{u}^2(y)}{|\tau(x)-y|} dy - \int_\Omega \frac{u^2(y)}{|x-y|} dy \right) dx \\ &= \frac{1}{\rho^3} \int_\Omega u^2(x) \left(\int_{\Omega_\rho} \frac{u^2(\tau^{-1}y)}{|\tau(x)-y|} dy - \int_\Omega \frac{u^2(y)}{|x-y|} dy \right) dx \\ &= \frac{1}{\rho^3} \int_\Omega u^2(x) \int_\Omega \left(\frac{u^2(y)}{|\tau(x)-\tau(y)|} \det(\nabla \tau(y)) - \frac{u^2(y)}{|x-y|} \right) dy dx \end{aligned}$$

and, with a computation similar to the one done in (46) we obtain that

$$\frac{1}{\rho^3} \int_\Omega (\tilde{w}(\tau(x)) - w(x)) dx = \frac{1}{\rho^3} \int_\Omega u^2(x) \int_{\Omega \cap (B_\rho(x_0) \cup B_\rho(x_1))} u^2(y) \left(\frac{1}{|x-y|} - \frac{1}{|\tau(x)-\tau(y)|} \right) dy dx.$$

Now, by Lemma 3.11, we know that u is Lipschitz so that $u^2 \lesssim \rho^2$ in $\Omega \cap B_\rho(x_i)$, for $i = 1, 2$, since $u(x_i) = 0$. Hence we obtain by the previous formula and an elementary computation that

$$(47) \quad \left| \frac{1}{\rho^3} \int_{\Omega} (\tilde{w}(\tau(x)) - w(x)) dx \right| \leq \frac{L}{\rho} \int_{\Omega} u^2(x) \int_{\Omega \cap (B_\rho(x_0) \cup B_\rho(x_1))} \left| \frac{1}{|x-y|} - \frac{1}{|\tau(x) - \tau(y)|} \right| dy dx \\ \leq c\rho^2 = o(\rho),$$

for some universal $c > 0$. By (45), (46) and (47) we deduce (44), and the proof is concluded. \square

We are now in position to show $C^{2,\alpha}$ -regularity of the boundary of a minimizer Ω . This can be done in two steps: first one shows that such a boundary is locally the graph of a $C^{2,\alpha}$ function defined on the boundary of a ball. To do that one exploits the improvement of flatness technique from [2, Section 7 and 8], readapted with minimal changes to our setting as in [21, Appendix]. Then, as we already know by the previous section that the boundary of Ω is close in Hausdorff distance to that of a ball, we obtain that the local parametrization is a global parametrization of class $C^{2,\alpha}$ on the boundary of the ball. We first need a definition (see [2, Definition 7.1]).

Definition 4.3. Let $\gamma_{\pm} \in (0, 1]$ and $k > 0$. A weak solution u of (40) is of class $F(\gamma_-, \gamma_+, k)$ in $B_\rho(x_0)$ with respect to direction $\nu \in \mathbb{S}^{N-1}$ if

(a) $x_0 \in \partial\{u > 0\}$ and

$$u = 0, \quad \text{for } (x - x_0) \cdot \nu \leq -\gamma_- \rho, \quad x \in B_\rho(x_0), \\ u(x) \geq \mu_u(x_0)[(x - x_0) \cdot \nu - \gamma_+ \rho], \quad \text{for } (x - x_0) \cdot \nu \geq \gamma_+ \rho, \quad x \in B_\rho(x_0).$$

(b) $|\nabla u(x_0)| \leq \mu_u(x_0)(1+k)$ in $B_\rho(x_0)$ and $\text{osc}_{B_\rho(x_0)} \mu_u \leq k\mu_u(x_0)$.

We note that when $k = +\infty$, then condition (b) is automatically satisfied. We can show the following result.

Theorem 4.4. Let $q \in (0, q_3]$, Ω be an optimal set for (2), and u a function attaining $E_q(\Omega)$ and a weak solution to (40) in B_R . Then there are constants $\bar{\gamma}$ and \bar{k} , depending only on R, μ_u , such that if u is of class $F(\gamma, 1, +\infty)$ in $B_{4\rho}(x_0)$ with respect to some direction $\nu \in \mathbb{S}^{N-1}$ with $\gamma \leq \bar{\gamma}$ and $\rho \leq \bar{k}\gamma^2$, then there exists a $C^{2,\alpha}$ function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ with $\|f\|_{C^{2,\alpha}} \leq C(R, \mu_u)$ such that, calling

$$\text{graph}_\nu f := \{x \in \mathbb{R}^3 : x \cdot \nu = f(x - (x \cdot \nu)\nu)\},$$

then

$$\partial\{u > 0\} \cap B_\rho(x_0) = (x_0 + \text{graph}_\nu(f)) \cap B_\rho(x_0).$$

Moreover for all $\varepsilon_0 > 0$ there exists $q_\varepsilon \in (0, q_3]$ such that if $q < q_\varepsilon$ then

$$\partial\{u > 0\} = \left\{ \left(r + \varphi\left(\frac{x}{|x|}\right) \right) \frac{x}{|x|} : x \in \partial B_r \right\}$$

where $\varphi: \partial B_1 \rightarrow \mathbb{R}$ is a function with the same regularity of f and $\|\varphi\|_{C^{2,\alpha}} \leq \varepsilon_0$.

We omit the proof which is identical to that in [35, Theorems 1.2 and 6.8] (which is in turn inspired by [2, Theorem 8.1] and [21, Theorem 2.17 and Appendix]). We note that in our setting μ_u is constant, thus the requirement to be $C^{1,\alpha}$ regular is trivially satisfied.

We are now in position to prove Theorem 3.1.

Proof of Theorem 3.1. The existence of a minimizer follows from Lemma 3.5 and Theorem 3.19. On the other hand, the fact that any optimal set is $C^{2,\alpha}$ nearly spherical follows from Theorem 4.4. \square

5. THE SURGERY RESULT AND THE PROOF OF THEOREM 1.3

In this section, we remove the equiboundedness assumption that was present in Theorem 3.1. The surgery strategy that we employ is very similar to the one proposed in [34] (see also [11]) and used for the spectral Gamow problem in [35]. We recall here, for the reader's sake, the main notations and the changes that are needed in our setting.

Lemma 5.1. *There exist universal constants $D, \bar{\delta} < 1$ and $\bar{q} \in (0, q_3]$ such that if $q \leq \bar{q}$ then for any open and connected set $\Omega \subset \mathbb{R}^3$ of measure $|B_1|$ satisfying $E_q(\Omega) - \lambda_0(B) \leq \bar{\delta}$ there exists an open, connected set $\hat{\Omega}$ of measure $|B_1|$ with diameter bounded by D and such that*

$$E_q(\hat{\Omega}) \leq E_q(\Omega).$$

Let us introduce some notation. Let Ω be a connected set of measure $|B_1|$ such that $\lambda_0(\Omega) - \lambda_0(B_1) \leq E_q(\Omega) - \lambda_0(B_1) \leq \bar{\delta}$, and we fix B_1 the ball attaining the minimum in the Fraenkel asymmetry for Ω (see (9)). We can clearly assume (up to a traslation of Ω) that B_1 is centered at the origin. Then, by the quantitative Faber–Krahn inequality (see Theorem 2.11 or [7]), we have

$$|\Omega \Delta B_1| = \mathcal{A}(\Omega) \leq |B_1|^{1/3} \left(\frac{\bar{\delta}}{\hat{\sigma}} \right)^{1/2},$$

where $\hat{\sigma}$ is the constant from Theorem 2.11. By defining

$$K := \lambda_0(B_1) + 1 \geq \lambda_0(B_1) + \bar{\delta}$$

we obtain immediately

$$E_q(\Omega) \leq K, \quad \text{and in particular,} \quad \int_{\Omega} |\nabla u|^2 dx \leq K,$$

where $u = u_{q,\Omega}$ from now on is the function attaining $E_q(\Omega)$. We then note that (since B_1 has unit radius)

$$|\Omega \setminus [-t, t]^3| \leq |\Omega \Delta B| = \mathcal{A}(\Omega), \quad \text{for all } t \geq 1.$$

Let $\hat{m} \in (0, 1/4)$ be such that

$$(48) \quad \frac{(4\hat{m})^{2/3}}{\lambda_0(B_1)|B_1|^{2/3}} K \leq \frac{1}{2}.$$

Moreover, we choose $\bar{\delta}$ small enough so that

$$|\Omega \setminus [-1, 1]^3| \leq \mathcal{A}(\Omega) \leq |B_1| \left(\frac{\bar{\delta}}{\hat{\sigma}} \right) \leq \frac{\hat{m}}{2^6}.$$

We first focus on the direction e_1 and detail the construction in this case. We shall denote $z = (x, y) \in \mathbb{R} \times \mathbb{R}^2$ and by z_i the i -th component of $z \in \mathbb{R}^3$. For any $t \in \mathbb{R}$, we define

$$\Omega_t := \left\{ y \in \mathbb{R}^2 : (t, y) \in \Omega \right\},$$

and given any set $\Omega \subseteq \mathbb{R}^3$, we define its 1-dimensional projections for $p \in \{1, 2, 3\}$ as

$$\pi_p(\Omega) := \left\{ t \in \mathbb{R} : \exists (z_1, z_2, z_3) \in \Omega, z_p = t \right\}.$$

For every $t \leq -1$ we call

$$\Omega^+(t) := \left\{ (x, y) \in \Omega : x > t \right\}, \quad \Omega^-(t) := \left\{ (x, y) \in \Omega : x < t \right\}, \quad \varepsilon(t) := \mathcal{H}^2(\Omega_t).$$

Observe that

$$m(t) := |\Omega^-(t)| = \int_{-\infty}^t \varepsilon(s) ds \leq 2\hat{m}.$$

We call u the optimizer for $E_q(\Omega)$ (we note that it is unique since $\Omega = \{u > 0\}$ is connected). We define then also, for every $t \leq -1$,

$$\delta(t) := \int_{\Omega_t} |\nabla u(t, y)|^2 d\mathcal{H}^2(y), \quad \mu(t) := \int_{\Omega_t} u(t, y)^2 d\mathcal{H}^2(y),$$

which makes sense since u is smooth inside Ω . Applying the Faber–Krahn inequality in \mathbb{R}^2 to the set Ω_t , and using the rescaling property of eigenvalues on \mathbb{R}^2 , we know that

$$\varepsilon(t)\lambda_0(\Omega_t) = \mathcal{H}^2(\Omega_t)\lambda_0(\Omega_t) \geq \lambda_0(B_{\mathbb{R}^2}),$$

calling $B_{\mathbb{R}^2}$ the ball of unit measure in \mathbb{R}^2 . As a trivial consequence, we can estimate μ in terms of ε and δ : in fact, noting that $u(t, \cdot) \in H_0^1(\Omega_t)$ and writing $\nabla u = (\nabla_1 u, \nabla_y u)$, we have

$$(49) \quad \mu(t) = \int_{\Omega_t} u(t, \cdot)^2 d\mathcal{H}^2 \leq \frac{1}{\lambda_0(\Omega_t)} \int_{\Omega_t} |\nabla_y u(t, \cdot)|^2 d\mathcal{H}^2 \leq C\varepsilon(t)\delta(t).$$

We can now present two estimates which assure that u and ∇u cannot be too big in $\Omega^-(t)$.

Lemma 5.2. *Let $\Omega \subseteq \mathbb{R}^3$ and u be as in Lemma 5.1. For every $t \leq -1$ the following inequalities hold:*

$$(50) \quad \int_{\Omega^-(t)} u^2 dx \leq C_1 \varepsilon(t)^{\frac{1}{2}} \delta(t), \quad \int_{\Omega^-(t)} |\nabla u|^2 dx \leq C_1 \varepsilon(t)^{\frac{1}{2}} \delta(t),$$

for some universal constant $C_1 > 0$.

The proof of the above Lemma follows, up to a few minor changes, as in [34, Lemma 2.3], by working on u (and recalling it solves the PDE (7)) instead of the first eigenfunction of the Dirichlet Laplacian in Ω . We reproduce it here for the sake of completeness.

Proof. Let us fix $t \leq -1$. Consider the set Ω_S^- obtained by the union of $\Omega^-(t)$ and its reflection with respect to the plane $\{x = t\}$, and call $u_S \in H_0^1(\Omega_S^-)$ the function obtained by reflecting u . Using the Faber-Krahn inequality, we find then

$$\frac{\lambda_0(B_1)|B_1|^{\frac{2}{3}}}{(2m(t))^{\frac{2}{3}}} = \frac{\lambda_0(B_1)|B_1|^{\frac{2}{3}}}{|\Omega_S^-|^{\frac{2}{3}}} \leq \lambda_0(\Omega_S^-) \leq \frac{\int_{\Omega_S^-} |\nabla u_S|^2 dx}{\int_{\Omega_S^-} u_S^2 dx} = \frac{\int_{\Omega^-(t)} |\nabla u|^2 dx}{\int_{\Omega^-(t)} u^2 dx} = \frac{\int_{\Omega^-(t)} |\nabla u|^2 dx}{\int_{\Omega^-(t)} u^2 dx},$$

by the symmetry of Ω_S^- , and using the scaling. This estimate gives

$$(51) \quad \int_{\Omega^-(t)} u^2 dx \leq \frac{(2m(t))^{\frac{2}{3}}}{\lambda_0(B_1)|B_1|^{\frac{2}{3}}} \int_{\Omega^-(t)} |\nabla u|^2 dx$$

which in particular, being $m(t) \leq 2\widehat{m}$ and recalling (48), implies

$$\int_{\Omega^-(t)} u^2 dx \leq \frac{1}{2}.$$

On the other hand, recalling that $-\Delta u \leq \lambda_q u$ in Ω , by Schwarz inequality and using (49) we have

$$(52) \quad \begin{aligned} \int_{\Omega^-(t)} |\nabla u|^2 dx &\leq \int_{\Omega^-(t)} \lambda_q u^2 dx + \int_{\Omega_t} u \frac{\partial u}{\partial \nu} d\mathcal{H}^2 \leq K \int_{\Omega^-(t)} u^2 dx + \sqrt{\int_{\Omega_t} u^2 d\mathcal{H}^2} \int_{\Omega_t} |\nabla u|^2 d\mathcal{H}^2 \\ &\leq K \int_{\Omega^-(t)} u^2 dx + C\varepsilon(t)^{\frac{1}{2}} \delta(t). \end{aligned}$$

It is now easy to obtain (50) combining (51) and (52). In fact, by inserting the latter into the first, we find

$$\int_{\Omega^-(t)} u^2 dx \leq \frac{(2m(t))^{\frac{2}{3}}}{\lambda_0(B_1)|B_1|^{\frac{2}{3}}} \left(K \int_{\Omega^-(t)} u^2 dx + C\varepsilon(t)^{\frac{1}{2}} \delta(t) \right),$$

which by (48) again yields

$$(53) \quad \frac{1}{2} \int_{\Omega^-(t)} u^2 dx \leq \frac{(2m(t))^{\frac{2}{3}}}{\lambda_0(B_1)|B_1|^{\frac{2}{3}}} C\varepsilon(t)^{\frac{1}{2}} \delta(t) \leq C\varepsilon(t)^{\frac{1}{2}} \delta(t).$$

The left estimate in (50) is then obtained. To get the right one, one has then just to insert (53) into (52). \square

Let us go further into the construction, giving some additional definitions. For any $t \leq -1$ and $\sigma(t) > 0$, we define the cylinder $Q(t)$ as

$$Q(t) := \left\{ (x, y) \in \mathbb{R}^3 : t - \sigma(t) < x < t, (t, y) \in \Omega \right\} = (t - \sigma(t), t) \times \Omega_t,$$

where for any $t \leq -1$ we set

$$\sigma(t) = \varepsilon(t)^{\frac{1}{2}}.$$

We let also $\widetilde{\Omega}(t) = \Omega^+(t) \cup Q(t)$, and we introduce $\widetilde{u} \in H_0^1(\widetilde{\Omega}(t))$ as

$$\widetilde{u}(x, y) := \begin{cases} u(x, y) & \text{if } (x, y) \in \Omega^+(t), \\ \frac{x - t + \sigma(t)}{\sigma(t)} u(t, y) & \text{if } (x, y) \in Q(t). \end{cases}$$

The fact that \tilde{u} vanishes on $\partial\tilde{\Omega}(t)$ is obvious; moreover, $\nabla u = \nabla\tilde{u}$ on $\Omega^+(t)$, while on $Q(t)$ one has

$$\nabla\tilde{u}(x, y) = \left(\frac{u(t, y)}{\sigma(t)}, \frac{x - t + \sigma(t)}{\sigma(t)} \nabla_y u(t, y) \right).$$

A simple calculation allows us to estimate the integrals of \tilde{u} and $\nabla\tilde{u}$ on $Q(t)$.

Lemma 5.3. *For every $t \leq -1$, one has*

$$\int_{Q(t)} |\nabla\tilde{u}|^2 dx \leq C_2 \varepsilon(t)^{\frac{1}{2}} \delta(t), \quad \int_{Q(t)} \tilde{u}^2 dx \leq C_2 \varepsilon(t)^{\frac{3}{2}} \delta(t),$$

for a universal constant $C_2 > 0$.

The proof of the above Lemma follows as [34, Lemma 2.4].

Another simple but useful estimate concerns the Rayleigh quotients of the functions \tilde{u} on the sets $\tilde{\Omega}(t)$: notice that, while u has unit L^2 norm, the modified function \tilde{u} in general is not normalized so we need to take care also of its norm.

Lemma 5.4. *There exists a universal constant $C_3 > 0$ such that for every $t \leq -1$, one has*

$$\int_{\tilde{\Omega}(t)} |\nabla\tilde{u}|^2 dx \leq \int_{\Omega} |\nabla u|^2 dx + C_3 \varepsilon(t)^{\frac{1}{2}} \delta(t), \quad \int_{\tilde{\Omega}(t)} \tilde{u}^2 dx \geq \int_{\Omega} u^2 dx - C_3 \varepsilon(t)^{\frac{1}{2}} \delta(t).$$

Proof. It is enough to note that, by definition of $\tilde{\Omega}(t)$ and using Lemma 5.2 and 5.3, we obtain for the gradient term

$$\begin{aligned} \int_{\tilde{\Omega}(t)} |\nabla\tilde{u}|^2 dx &= \int_{\Omega^+(t)} |\nabla u|^2 dx + \int_{Q(t)} |\nabla\tilde{u}|^2 dx \\ &= \int_{\Omega} |\nabla u|^2 dx + \int_{Q(t)} |\nabla\tilde{u}|^2 dx - \int_{\Omega^-(t)} |\nabla u|^2 dx \leq \int_{\Omega} |\nabla u|^2 dx + C_2 \varepsilon(t)^{\frac{1}{2}} \delta(t), \end{aligned}$$

while for the function, we have

$$\int_{\tilde{\Omega}(t)} \tilde{u}^2 dx = \int_{\Omega^+(t)} u^2 dx + \int_{Q(t)} \tilde{u}^2 dx = \int_{\Omega} u^2 dx + \int_{Q(t)} \tilde{u}^2 dx - \int_{\Omega^-(t)} u^2 dx \geq \int_{\Omega} u^2 dx - C_1 \varepsilon(t)^{\frac{1}{2}} \delta(t).$$

□

We can now enter in the central part of our construction. Basically, we aim to show that either Ω already has bounded left “tail” in direction e_1 , or some rescaling of $\tilde{\Omega}(t)$ has energy lower than that of Ω .

Lemma 5.5. *Let Ω be as in the assumptions of Lemma 5.2, and let $t \leq -1$. There exist universal $\bar{q} \in (0, q_3]$ and $C_4 > 2$ such that, for all $q \leq \bar{q}$ exactly one of the three following conditions hold:*

- (1) $\max\{\varepsilon(t), \delta(t)\} > 1$;
- (2) (1) does not hold and $m(t) \leq C_4(\varepsilon(t) + \delta(t))\varepsilon(t)^{\frac{1}{2}}$;
- (3) (1) and (2) do not hold and one has that

$$\frac{\int_{\hat{\Omega}(t)} |\nabla\hat{u}|^2 dx}{\int_{\hat{\Omega}(t)} \hat{u}^2 dx} \leq \int_{\Omega} |\nabla u|^2 dx, \quad \text{and} \quad E(\hat{\Omega}(t)) < E_q(\Omega),$$

where for $t \leq -1$ we set

$$\hat{\Omega}(t) := |B_1|^{\frac{1}{3}} |\tilde{\Omega}(t)|^{-\frac{1}{3}} \tilde{\Omega}(t), \quad \text{and} \quad \hat{u}(x) = \tilde{u}(|B_1|^{-\frac{1}{3}} |\tilde{\Omega}(t)|^{\frac{1}{3}} x), \quad \text{for } x \in \hat{\Omega}(t).$$

Proof. Assume (1) is false. Then it is possible to apply Lemma 5.4, to obtain

$$(54) \quad \int_{\tilde{\Omega}(t)} |\nabla\tilde{u}|^2 dx \leq \int_{\Omega} |\nabla u|^2 dx + C_3 \varepsilon(t)^{\frac{1}{2}} \delta(t), \quad \int_{\tilde{\Omega}(t)} \tilde{u}^2 dx \geq \int_{\Omega} u^2 dx - C_3 \varepsilon(t)^{\frac{1}{2}} \delta(t) = 1 - C_3 \varepsilon(t)^{\frac{1}{2}} \delta(t).$$

By the scaling properties of the eigenvalue and the fact that $|\hat{\Omega}(t)| = |B_1|$, we know that

$$\frac{\int_{\hat{\Omega}(t)} |\nabla\hat{u}|^2 dx}{\int_{\hat{\Omega}(t)} \hat{u}^2 dx} = \frac{|\tilde{\Omega}(t)|^{\frac{2}{3}} \int_{\tilde{\Omega}(t)} |\nabla\tilde{u}|^2 dx}{|B_1|^{\frac{2}{3}} \int_{\tilde{\Omega}(t)} \tilde{u}^2 dx}.$$

By construction,

$$|\tilde{\Omega}(t)| = |\Omega^+(t)| + |Q(t)| = |B_1| - m(t) + \varepsilon(t)^{\frac{3}{2}},$$

hence the above estimates, the scaling of the integrals due to the definition of \widehat{u} and (54) lead to

$$\begin{aligned} \frac{\int_{\widehat{\Omega}(t)} |\nabla \widehat{u}|^2 dx}{\int_{\widehat{\Omega}(t)} \widehat{u}^2 dx} &= \left(1 - \frac{m(t)}{|B_1|} + \frac{\varepsilon(t)^{\frac{3}{2}}}{|B_1|}\right)^{\frac{2}{3}} \frac{\int_{\widehat{\Omega}(t)} |\nabla \widetilde{u}|^2 dx}{\int_{\widehat{\Omega}(t)} \widetilde{u}^2 dx}, \\ &\leq \left(1 - \frac{2}{3|B_1|} m(t) + \frac{2}{3|B_1|} \varepsilon(t)^{\frac{3}{2}}\right) \left(1 + C_3 \varepsilon^{\frac{1}{2}}(t) \delta(t)\right) \left(\int_{\Omega} |\nabla u|^2 dx + C_3 \varepsilon(t)^{\frac{1}{2}} \delta(t)\right) \\ &\leq \left(\int_{\Omega} |\nabla u|^2 dx - \frac{2\lambda_0(B_1)}{3|B_1|} m(t) + \frac{2K}{3|B_1|} \varepsilon(t)^{\frac{3}{2}} + \left(2C_3 + KC_3 + \frac{2}{3|B_1|}\right) \varepsilon(t)^{\frac{1}{2}} \delta(t)\right). \end{aligned}$$

At this point, defining $C_4 := \max\{\frac{2(K+1)}{3|B_1|} + 2C_3 + KC_3, 2\}$, if

$$m(t) \leq C_4(\varepsilon(t) + \delta(t))\varepsilon(t)^{\frac{1}{2}},$$

then condition (2) holds true. Otherwise, we immediately have that

$$(55) \quad \frac{\int_{\widehat{\Omega}(t)} |\nabla \widehat{u}|^2 dx}{\int_{\widehat{\Omega}(t)} \widehat{u}^2 dx} \leq \left(\int_{\Omega} |\nabla u|^2 dx - \left(\frac{2\lambda_0(B_1)}{3|B_1|} - 1\right) m(t)\right) \leq \int_{\Omega} |\nabla u|^2 dx - C_5 m(t),$$

for a universal constant $C_5 > 0$, therefore the first part of the third claim is verified.

On the other hand, we note that, using the L^∞ bound of u , see Lemma 2.4, the fact that $\widetilde{u} \leq u$ by construction and also [16, Lemma 2.4],

$$D(\widetilde{u}^2, \widetilde{u}^2) = D(u^2, u^2) + 2 \int_{\Omega^+(t)} \int_{Q(t)} \frac{\widetilde{u}^2(x)\widetilde{u}^2(y)}{|x-y|} dx dy + \int_{Q(t)} \int_{Q(t)} \frac{\widetilde{u}^2(x)\widetilde{u}^2(y)}{|x-y|} dx dy \leq D(u^2, u^2) + C_{fp} \varepsilon^{\frac{3}{2}}(t).$$

Then we can estimate, using the appropriate scalings,

$$\begin{aligned} (56) \quad \frac{D(\widehat{u}^2, \widehat{u}^2)}{\left(\int_{\widehat{\Omega}(t)} \widehat{u}^2 dx\right)^2} &\leq \frac{D(\widetilde{u}^2, \widetilde{u}^2)}{\left(\int_{\widehat{\Omega}(t)} \widetilde{u}^2 dx\right)^2} \left(1 - \frac{m(t)}{|B_1|} + \frac{\varepsilon(t)^{\frac{3}{2}}}{|B_1|}\right)^{-\frac{2}{3}} \leq \left(1 + \frac{2}{3|B_1|} m(t)\right) \frac{D(\widetilde{u}^2, \widetilde{u}^2)}{\left(\int_{\widehat{\Omega}(t)} \widetilde{u}^2 dx\right)^2} \\ &\leq \left(1 + \frac{2}{3|B_1|} m(t)\right) \left(1 + C_3 \varepsilon^{\frac{1}{2}}(t) \delta(t)\right) \left(D(u^2, u^2) + C_{fp} \varepsilon^{\frac{3}{2}}(t)\right) \\ &\leq D(u^2, u^2) + C \|u\|_{L^\infty}^2 m(t) + C_{fp} \varepsilon^{\frac{3}{2}}(t) + C_3 \|u\|_{L^\infty}^2 \varepsilon^{\frac{1}{2}}(t) \delta(t) \\ &\leq D(u^2, u^2) + C \|u\|_{L^\infty}^2 m(t) + (C_{fp} + C_3 \|u\|_{L^\infty}^2) m(t) \\ &= D(u^2, u^2) + C_6 m(t). \end{aligned}$$

Then, putting together (55) and (56), recalling also Remark 2.3 for the equivalence of the scale invariant energy,

$$\begin{aligned} E_q(\widehat{\Omega}(t)) &\leq \frac{\int_{\widehat{\Omega}(t)} |\nabla \widehat{u}|^2 dx}{\int_{\widehat{\Omega}(t)} \widehat{u}^2 dx} + \frac{q}{2} \frac{D(\widehat{u}^2, \widehat{u}^2)}{\left(\int_{\widehat{\Omega}(t)} \widehat{u}^2 dx\right)^2} \\ &\leq \int_{\Omega} |\nabla u|^2 dx + \frac{q}{2} D(u^2, u^2) - (C_5 - \frac{q}{2} C_6) m(t) \\ &\leq \int_{\Omega} |\nabla u|^2 dx + \frac{q}{2} D(u^2, u^2) - \frac{C_5}{2} m(t), \end{aligned}$$

up to taking $q \leq \bar{q} < \sqrt{\frac{C_5}{2C_6}}$, so that in this case condition (3) holds and the proof is concluded. \square

Once we have Lemma 5.5, the rest of the proof follows as in [34] or [35] as we detail here below.

Proof of Lemma 5.1. It is enough to repeat the analogs of [35, Lemma 8.7, Lemma 8.8, Proposition 8.1 and Section 9.2], noting that it is only a geometric argument and having $\int_{\Omega} |\nabla u|^2 dx$ instead of $\lambda_0(\Omega)$ does not change anything. \square

Proof of Theorem 1.3. We aim to apply the surgery result Lemma 5.1 and then to employ Theorem 3.1. Precisely, first, as in Section 9.2 of [35] we select a minimizing sequence for problem (2) made of connected sets. Then, by Lemma 5.1 we select another minimizing sequence of equibounded sets. At this point we are in position to apply Theorem 3.1 and we conclude. \square

6. NON-EXISTENCE OF MINIMIZERS FOR q LARGE: PROOF OF THEOREM 1.6

In this section we show that problem (2) does not admit minimizers for q large enough. The proof of Theorem 1.6 is a nontrivial adaptation and refinement of the argument in [32] to show non-existence of ground states of the Thomas-Fermi-Dirac-Von Weizsäcker energy for large values of the L^2 norm of the wavefunction.

First of all, we recall the equivalence of problem (2) with the case in which the parameter is the L^2 norm of the optimal function, which is the setting of [32]. We already discussed it in Appendix A and in Section 1, but we stress it also here for the reader's convenience.

Remark 6.1. Notice that problem (2) with $q = m$ is equivalent to

$$(57) \quad \min \{ \mathcal{F}_m(\Omega) : \Omega \subset \mathbb{R}^3, \text{ open, with } |\Omega| = |B_1| \}$$

where

$$(58) \quad \mathcal{F}_m(\Omega) = \min_{\varphi \in H_0^1(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^2 dx + \frac{1}{2} D(\varphi^2, \varphi^2) : \int_{\Omega} \varphi^2 dx = m \right\},$$

Indeed, let us fix $\Omega \subset \mathbb{R}^3$ an open set of unit measure and let $\varphi \in H_0^1(\Omega)$ with $\int_{\Omega} \varphi^2 dx = m$ be the function attaining $\mathcal{F}_m(\Omega)$. Then, since $\int_{\Omega} (m^{-1/2} \varphi)^2 dx = 1$,

$$E_m(\Omega) = m^{-1} \mathcal{F}_m(\Omega).$$

Proof of Theorem 1.6. We set in this proof, with a slight abuse of notation,

$$\mathcal{F}(\varphi) = \int_{\Omega} |\nabla \varphi|^2 dx + \frac{1}{2} D(\varphi^2, \varphi^2).$$

We show that if a minimizer for problem (57) exists, then m is bounded from above by a universal constant. Let $\Omega = \Omega_m$ be a minimizer of (57) for $m > 0$ and $\varphi = \varphi_{\Omega} > 0$ be its ground state function, that is, the function achieving the minimum for $\mathcal{F}_m(\Omega)$. We extend φ by zero outside Ω and define, for $x \in \mathbb{R}^3$ and $r > 0$,

$$V(x, r) = \int_{B_r(x)} \varphi^2(y) dy.$$

Note that now $\Omega = \text{supp}(\varphi)$.

We divide the proof into three steps.

Step 1. We claim that for all $x \in \mathbb{R}^3$, for all $t, r > 0$ and for all $R > r + t$, the following discrete-ODE type inequality holds:

$$(59) \quad \frac{1}{t^2} \left(V(x, r + t) - V(x, r) \right) \geq \frac{C}{R + r} \left(V(x, R) - V(x, r + t) \right) V(x, r),$$

where $C > 0$ is a universal constant.

To prove (59), let $f_1, f_2 \in C^\infty(\mathbb{R}^3, [0, 1])$ be radial functions such that $f_1 = 1$ on $B_r(x)$, $f_1 = 0$ outside $B_{r+t}(x)$, $f_2 = \sqrt{1 - f_1^2}$ and $|\nabla f_1| + |\nabla f_2| \leq \frac{C}{t}$ for some universal $C > 0$. Let $\varphi_i = f_i \varphi$, for $i = 1, 2$ and define

$$\psi_L(z) = C_L \left(\varphi_1(z) + \varphi_2(z + L\nu) \right), \quad z \in \mathbb{R}^3,$$

where $\nu \in \mathbb{R}^3$ is any non-null vector (its choice is irrelevant in the proof). Here $C_L > 0$ is a constant chosen so that $\|\psi_L\|_{L^2(\mathbb{R}^3)}^2 = m$. Let us set eventually $\Omega_L = \{\psi_L > 0\}$. An elementary computation shows that $C_L \leq 1$ and $C_L \rightarrow 1$ as $L \rightarrow +\infty$. Notice also that

$$|\Omega_L| = |\Omega| - |(\{\varphi_1 > 0\} - L\nu) \cap \{\varphi_2 > 0\}| \leq |\Omega| = |B_1|,$$

so that, since $\Omega \mapsto \mathcal{F}_m(\Omega)$ (see (58)) is decreasing with respect to set inclusion (as the minimization is done under Dirichlet boundary condition), there holds, for any measurable set A_L of measure $|B_1|$ and containing $\{\psi_L > 0\}$, that

$$\begin{aligned} \mathcal{F}(\varphi) &= \mathcal{F}_m(\Omega) \\ &\leq \mathcal{F}_m(A_L), && \text{by minimality of } \Omega \\ &\leq \mathcal{F}_m(\{\psi_L > 0\}) && \text{as } \{\psi_L > 0\} \subset A_L \\ &\leq \mathcal{F}(\psi_L) && \text{as } \psi_L \in H_0^1(\{\psi_L > 0\}). \end{aligned}$$

One easily sees that

$$\mathcal{F}(\psi_L) \rightarrow \mathcal{F}(\varphi_1) + \mathcal{F}(\varphi_2), \quad \text{as } L \rightarrow +\infty,$$

so that

$$(60) \quad \mathcal{F}(\varphi_1) + \mathcal{F}(\varphi_2) - \mathcal{F}(\varphi) \geq 0.$$

At this point a direct computation entails that for some universal constant C

$$(61) \quad \begin{aligned} |\nabla\varphi_1|^2 + |\nabla\varphi_2|^2 - |\nabla\varphi|^2 &= \varphi^2(|\nabla f_1|^2 + |\nabla f_2|^2) + 2(f_1\partial_r f_1\varphi\partial_r\varphi + f_2\partial_r f_2\varphi\partial_r\varphi) \\ &= \varphi^2(|\nabla f_1|^2 + |\nabla f_2|^2) \leq \frac{C}{t^2}\varphi^2\chi_{B_{r+t}(x)\setminus B_r(x)}, \end{aligned}$$

where ∂_r denotes the radial derivative with respect to r , χ is a characteristic function and where we used that

$$f_1\partial_r f_1 + f_2\partial_r f_2 = \frac{1}{2}\partial_r(f_1^2 + f_2^2) = 0.$$

Thus integrating (61) over the whole \mathbb{R}^3 (recalling that φ is extended by zero outside Ω), we obtain

$$(62) \quad \int_{\mathbb{R}^3} |\nabla\varphi_1(z)|^2 + |\nabla\varphi_2(z)|^2 - |\nabla\varphi(z)|^2 dz \leq \frac{C}{t^2} \int_{B_{r+t}(x)\setminus B_r(x)} \varphi^2(z) dz.$$

On the other hand, concerning the Coulombic term, we have (as in [32, equation (3.6)]):

$$(63) \quad \begin{aligned} D(\varphi_1^2, \varphi_1^2) + D(\varphi_2^2, \varphi_2^2) - D(\varphi^2, \varphi^2) &= -2 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\varphi_1^2(z)\varphi_2^2(y)}{|z-y|} dz dy \\ &\leq -2 \int_{B_r(x)} \int_{B_R(x)\setminus B_{r+t}(x)} \frac{\varphi_1^2(z)\varphi_2^2(y)}{|z-y|} dz dy \\ &= -2 \int_{B_r(x)} \int_{B_R(x)\setminus B_{r+t}(x)} \frac{\varphi^2(z)\varphi^2(y)}{|z-y|} dz dy \\ &\leq -\frac{2}{R+r} \int_{B_r(x)} \int_{B_R(x)\setminus B_{r+t}(x)} \varphi^2(z)\varphi^2(y) dz dy. \end{aligned}$$

We can eventually deduce the claim (59) from (62), (63) and (60).

Step 2. We prove now the following: there exists a universal constant $C_0 > 0$ such that for every given $x \in \mathbb{R}^3$ and $R_0 > 0$ the following implication holds

$$V(x, R_0) \geq \frac{C_0}{R_0} \implies V(x, 2R_0) \geq \frac{m}{2}.$$

To prove that, let $R > 0$. By (59) with

$$r = \frac{R}{4}, \quad t = \frac{R}{4N},$$

where $N \in \mathbb{N}$, $N \geq 2$ will be fixed later, we obtain, for all $j = 1, \dots, N$,

$$V\left(x, \frac{R}{4}\left(1 + \frac{j}{N}\right)\right) - V\left(x, \frac{R}{4}\left(1 + \frac{j-1}{N}\right)\right) \geq \frac{CR}{4N^2} [V(x, R/4)(V(x, R) - V(x, R/2))].$$

Then summing up over $j = 1, \dots, N$, we obtain, for all $R > 0$,

$$(64) \quad V(x, R/2) - V(x, R/4) \geq C \frac{R}{4N} [V(x, R/4)(V(x, R) - V(x, R/2))].$$

We set now $N = 2$ and $C_1 := 2/C$, where C is the universal constant in (64). By the same argument as above, for all $k \geq 1$ we have

$$(65) \quad \begin{aligned} V(x, 2^{k-1}R) - V(x, 2^{k-2}R) &\geq \frac{RV(x, 2^{k-2}R)}{4C_1} [(V(x, 2^kR) - V(x, 2^{k-1}R))] \\ &\geq \frac{RV(x, R/4)}{4C_1} [(V(x, 2^kR) - V(x, 2^{k-1}R))], \end{aligned}$$

where in the last inequality we used the monotonicity of $V(x, \cdot)$. We fix now $C_0 = 2C_1$ and for all $k \geq 1$ we compute (iterating (65))

$$\begin{aligned} V(x, 2^kR) - V(x, R/2) &= \sum_{i=0}^k [V(x, 2^iR) - V(x, 2^{i-1}R)] \\ &\leq \sum_{i=0}^k \left(\frac{2C_0}{RV(x, R/4)}\right)^{i+1} [V(x, R/2) - V(x, R/4)] \leq [V(x, R/2) - V(x, R/4)], \end{aligned}$$

where in the last inequality we have used the assumption of the claim with $R_0 = R/4$, namely that $\frac{R}{4}V(x, R/4) \geq C_0$. Therefore we deduce that for all $k \geq 1$

$$2V(x, 2R_0) = 2V(x, R/2) \geq V(x, 2^k R) = \int_{B_{2^k R}} \varphi^2(z) dz,$$

and the claim follows by taking the supremum with respect to k .

Step 3. There exists a universal constant $\beta > 0$, such that

$$m \leq \beta.$$

Indeed, we set

$$R_0 = \inf \left\{ r > 0 : \exists x \in \mathbb{R}^3, V(x, r) \geq \frac{C_0}{r} \right\}.$$

where $C_0 > 0$ is the constant in Step 2. We check that $R_0 \in (0, +\infty)$. Otherwise, either we may find sequences $R_k \rightarrow 0$ and $(x_k) \subset \mathbb{R}^3$ such that

$$m \geq V(x_k, R_k) = \int_{B_{R_k}(x_k)} \varphi^2(z) dz \geq \frac{C_0}{R_k} \rightarrow +\infty,$$

leading to an immediate contradiction. Thus $R_0 > 0$. Or, since we are taking the infimum over a nonempty set, it is clear that $R_0 < +\infty$.

Now, by Step 2 (and definition of R_0), we have

$$\frac{m}{2} \leq V(x, 2R_0).$$

Using a standard covering argument, $B_{2R_0}(x)$ can be covered by a universally bounded number of balls of radius proportional to $2R_0$: in particular, $B_{2R_0}(x) \subset \cup_{i=1}^n B_{R_0/2}(x_i)$ for some universal $n \in \mathbb{N}$. By our choice of R_0 , it follows that

$$(66) \quad \frac{m}{2} \leq V(x, 2R_0) \leq \sum_{i=1}^n V(x_i, R_0/2) \leq \frac{nC_0}{R_0}.$$

To conclude, we need now to compare m and R_0 . First, it is easy to see, using the definition of R_0 that

$$(67) \quad \mathcal{F}(\varphi) \geq \frac{1}{2}D(\varphi^2, \varphi^2) \geq \frac{1}{2} \int_{B_{2R_0}(x)} \int_{B_{2R_0}(x)} \frac{\varphi^2(z)\varphi^2(y)}{|z-y|} dy dz \geq \frac{1}{8R_0} \cdot \frac{C_0^2}{4R_0^2} = \frac{C_0^2}{2^5 R_0^3}.$$

On the other hand, we can estimate the Coulomb energy from above on the ball of unit radius⁴, obtaining (here we call φ_B the first Dirichlet Laplacian eigenfunction of B_1 with $\int \varphi_B^2 = m$)

$$\mathcal{F}_m(B_1) \leq \int_{B_1} |\nabla \varphi_B|^2 + \frac{1}{2}D(\varphi_B^2, \varphi_B^2) \leq \lambda_0(B_1)m + \frac{1}{2}\sqrt{\lambda_0(B_1)}C_{univ}m^2,$$

for a universal constant $C_{univ} > 0$ (from Lemma 2.1). If $m \leq 1$ we have already finished, otherwise $m^2 \geq m$ and thus

$$\mathcal{F}_m(B_1) \leq C_1 m^2,$$

for a universal positive constant $C_1 = \lambda_0(B_1) + \frac{1}{2}\sqrt{\lambda_0(B_1)}C$.

All in all, from the above estimate, (66) and (67) we obtain

$$C_1 m^2 \geq \frac{C_0^2}{2^5 R_0^3} \geq \frac{m^3}{2^8 n^3 C_0}.$$

This estimate concludes the proof, with the choice of the universal constant $\beta = \max\{1, 2^8 n^3 C_0 C_1\}$. \square

⁴Actually we can obtain a slightly better estimate by comparing with a union of N balls at infinite distance, but this does not change our final result.

APPENDIX A. THE PHYSICAL MODEL AND NON-DIMENSIONALIZATION

In its dimensional form, the ground state bosonic Hartree energy for the Cooper pairs takes the form (in the SI units)

$$\mathcal{E}(\phi, \Omega) = \frac{N\hbar^2}{2m^*} \int_{\Omega} |\nabla\phi(x)|^2 d^3x + \frac{N(N-1)e^2}{2\pi\epsilon_0\epsilon} \int_{\Omega} \int_{\Omega} \frac{\phi^2(x)\phi^2(y)}{|x-y|} d^3x d^3y,$$

where m^* is the effective mass of a Cooper pair and $-2|e|$ is its charge, where $-|e|$ is the elementary charge, ϵ_0 is the vacuum permittivity, ϵ is the dielectric constant of the surrounding matrix (within a simplified local treatment of the dielectric), N is the number of Cooper pairs in the island and ϕ is a single-orbital wave function subject to the normalization

$$(68) \quad \int_{\Omega} \phi^2(x) d^3x = 1.$$

We now perform a rescaling

$$x \rightarrow Lx, \quad \phi \rightarrow L^{-3/2}\phi,$$

that keeps the normalization condition in (68) unchanged. After some simple algebra we arrive at

$$\mathcal{E}(L^{-3/2}u(\cdot/L), L\Omega) = \frac{N\hbar^2}{2m^*L^2} E_q(u, \Omega),$$

where

$$q = \frac{2e^2(N-1)m^*L}{\pi\hbar^2\epsilon_0\epsilon}.$$

With the choice of $L = \left(\frac{3V}{4\pi}\right)^{1/3} = \left(\frac{V}{|B_1|}\right)^{1/3}$ we then arrive at the shape optimization problem in (2).

REFERENCES

- [1] N. Aguilera, H. W. Alt, and L. A. Caffarelli. An optimization problem with volume constraint. *SIAM J. Control Optim.*, 24(2):191–198, 1986.
- [2] H. W. Alt and L. A. Caffarelli. Existence and regularity for a minimum problem with free boundary. *J. Reine Angew. Math.*, 325:105–144, 1981.
- [3] L. Ambrosio and P. Tilli. *Topics on analysis in metric spaces*, volume 25 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2004.
- [4] I. Beloborodov, A. Lopatin, V. Vinokur, and K. Efetov. Granular electronic systems. *Reviews of Modern Physics*, 79(2):469–518, 2007.
- [5] R. Benguria, H. Brézis, and E. H. Lieb. The Thomas-Fermi-von Weizsäcker theory of atoms and molecules. *Comm. Math. Phys.*, 79(2):167–180, 1981.
- [6] V. Bouchiat, D. Vion, P. Joyez, D. Esteve, and M. Devoret. Quantum coherence with a single cooper pair. volume 76, pages 165–170, 1998.
- [7] L. Brasco, G. De Philippis, and B. Velichkov. Faber-Krahn inequalities in sharp quantitative form. *Duke Math. J.*, 164(9):1777–1831, 2015.
- [8] L. Brasco and G. Franzina. A note on positive eigenfunctions and hidden convexity. *Arch. Math. (Basel)*, 99(4):367–374, 2012.
- [9] T. Briançon, M. Hayouni, and M. Pierre. Lipschitz continuity of state functions in some optimal shaping. *Calc. Var. Partial Differential Equations*, 23(1):13–32, 2005.
- [10] D. Bucur. Minimization of the k -th eigenvalue of the Dirichlet Laplacian. *Arch. Ration. Mech. Anal.*, 206(3):1073–1083, 2012.
- [11] D. Bucur and D. Mazzoleni. A surgery result for the spectrum of the Dirichlet Laplacian. *SIAM J. Math. Anal.*, 47(6):4451–4466, 2015.
- [12] D. Bucur, D. Mazzoleni, A. Pratelli, and B. Velichkov. Lipschitz regularity of the eigenfunctions on optimal domains. *Arch. Ration. Mech. Anal.*, 216(1):117–151, 2015.
- [13] M. Büttiker. Zero-current persistent potential drop across small-capacitance josephson junctions. *Physical Review B*, 36(7):3548–3555, 1987.
- [14] R. Choksi, C. B. Muratov, and I. Topaloglu. An old problem resurfaces nonlocally: Gamow’s liquid drops inspire today’s research and applications. *Notices Amer. Math. Soc.*, 64(11):1275–1283, 2017.
- [15] G. De Philippis, M. Marini, and E. Mukoseeva. The sharp quantitative isocapacitary inequality. *Rev. Mat. Iberoam.*, 37(6):2191–2228, 2021.
- [16] N. Fusco and A. Pratelli. Sharp stability for the Riesz potential. *ESAIM Control Optim. Calc. Var.*, 26:Paper No. 113, 24, 2020.
- [17] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*, volume 224 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1983.
- [18] M. Goldman, M. Novaga, and B. Ruffini. Existence and stability for a non-local isoperimetric model of charged liquid drops. *Arch. Ration. Mech. Anal.*, 217(1):1–36, 2015.
- [19] M. Goldman, M. Novaga, and B. Ruffini. On minimizers of an isoperimetric problem with long-range interactions under a convexity constraint. *Anal. PDE*, 11(5):1113–1142, 2018.

- [20] M. Goldman, M. Novaga, and B. Ruffini. Rigidity of the ball for an isoperimetric problem with strong capacitary repulsion. *JEMS*, In press. DOI 10.4171/JEMS/1451, 2024.
- [21] B. Gustafsson and H. Shahgholian. Existence and geometric properties of solutions of a free boundary problem in potential theory. *J. Reine Angew. Math.*, 473:137–179, 1996.
- [22] D. Hartree. The wave mechanics of an atom with a non-Coulomb central field. part I: Theory and methods. *Mathematical Proceedings of the Cambridge Philosophical Society*, 24(1):89 – 110, 1928.
- [23] J. Heinonen, T. Kilpeläinen, and O. Martio. *Nonlinear potential theory of degenerate elliptic equations*. Dover Publications, Inc., Mineola, NY, 2006. Unabridged republication of the 1993 original.
- [24] A. Henrot. *Extremum problems for eigenvalues of elliptic operators*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2006.
- [25] A. Henrot and M. Pierre. *Shape variation and optimization. A geometrical analysis*, volume 28 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2018.
- [26] M. Kjaergaard, M. E. Schwartz, J. Braumüller, P. Krantz, J. I.-J. Wang, S. Gustavsson, and W. D. Oliver. Superconducting qubits: Current state of play. *Annual Review of Condensed Matter Physics*, 11:369 – 395, 2020.
- [27] H. Knüpfer and C. B. Muratov. On an isoperimetric problem with a competing nonlocal term I: The planar case. *Comm. Pure Appl. Math.*, 66(7):1129–1162, 2013.
- [28] H. Knüpfer and C. B. Muratov. On an isoperimetric problem with a competing nonlocal term II: The general case. *Comm. Pure Appl. Math.*, 67(12):1974–1994, 2014.
- [29] C. Le Bris and P.-L. Lions. From atoms to crystals: a mathematical journey. *Bull. Amer. Math. Soc. (N.S.)*, 42(3):291–363, 2005.
- [30] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [31] P.-L. Lions. Solutions of Hartree-Fock equations for Coulomb systems. *Comm. Math. Phys.*, 109(1):33–97, 1987.
- [32] J. Lu and F. Otto. Nonexistence of a minimizer for Thomas-Fermi-Dirac-von Weizsäcker model. *Comm. Pure Appl. Math.*, 67(10):1605–1617, 2014.
- [33] V. G. Maz'ja. *Sobolev spaces*. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1985. Translated from the Russian by T. O. Shaposhnikova.
- [34] D. Mazzoleni and A. Pratelli. Existence of minimizers for spectral problems. *J. Math. Pures Appl. (9)*, 100(3):433–453, 2013.
- [35] D. Mazzoleni and B. Ruffini. A spectral shape optimization problem with a nonlocal competing term. *Calc. Var. Partial Differential Equations*, 60(3):Paper No. 114, 46, 2021.
- [36] C. B. Muratov, M. Novaga, and B. Ruffini. On equilibrium shape of charged flat drops. *Comm. Pure Appl. Math.*, 71(6):1049–1073, 2018.
- [37] C. B. Muratov, M. Novaga, and B. Ruffini. Conducting flat drops in a confining potential. *Arch. Ration. Mech. Anal.*, 243(3):1773–1810, 2022.
- [38] B. Velichkov. *Regularity of the one-phase free boundaries*, volume 28 of *Lecture Notes of the Unione Matematica Italiana*. Springer, 2023.
- [39] D. Vion. Josephson quantum bits based on a Cooper pair box. *Les Houches Summer School Proceedings*, 79(C):521–523, 2004.

DIPARTIMENTO DI MATEMATICA F. CASORATI, UNIVERSITÀ DI PAVIA, VIA FERRATA 5, 27100 PAVIA, ITALY
Email address: `dario.mazzoleni@unipv.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO B. PONTECORVO, 5, 56127 PISA, ITALY
Email address: `cyrill.muratov@unipi.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY
Email address: `berardo.ruffini@unibo.it`