

MEASURES IN THE DUAL OF BV : PERIMETER BOUNDS AND RELATIONS WITH DIVERGENCE-MEASURE FIELDS

GIOVANNI E. COMI AND GIAN PAOLO LEONARDI

ABSTRACT. We analyze some properties of the measures in the dual of the space BV , by considering (signed) Radon measures satisfying a perimeter bound condition, which means that the absolute value of the measure of a set is controlled by the perimeter of the set itself, and whose total variations also belong to the dual of BV . We exploit and refine the results of [25], in particular exploring the relation with divergence-measure fields and proving the stability of the perimeter bound from sets to BV functions under a suitable approximation of the given measure. As an important tool, we obtain a refinement of Anzellotti-Giaquinta approximation for BV functions, which is of separate interest in itself and, in the context of Anzellotti's pairing theory for divergence-measure fields, implies a new way of approximating λ -pairings, as well as new bounds for their total variation. These results are also relevant due to their application in the study of weak solutions to the non-parametric prescribed mean curvature equation with measure data, which is explored in a subsequent work.

CONTENTS

1. Introduction	1
2. Preliminaries	6
2.1. Functions of bounded variation	6
2.2. Divergence-measure fields and λ -pairings	11
2.3. Measures in the dual of BV	15
3. λ -approximation	16
4. Perimeter bounds and admissible measures	26
References	40

1. INTRODUCTION

Given the ubiquitous presence of the space of functions of bounded variation BV in the Calculus of Variation, it is natural to try and characterize its dual space. An integral representation of the elements of $BV(\Omega)^*$, for any open set $\Omega \subset \mathbb{R}^n$, was obtained in [17], under the Continuum

2020 *Mathematics Subject Classification*. Primary: 26B30. Secondary: 26B20, 46E27.

Key words and phrases. Functions of bounded variation, divergence-measure fields, Gauss–Green formulas, dual of BV .

This work has been financially supported by GNAMPA - INdAM. Part of this work was undertaken while the second author was visiting the University of Trento. He would like to thank this institution for the support and warm hospitality during the visits. The authors are particularly grateful to Lorenzo Brasco and Giorgio Saracco for their support and encouragement during the preparation of the paper. The authors are also grateful to Virginia De Cicco for her comments to a preliminary version of the paper.

Hypothesis (we refer to the introduction of [17] for a detailed account of the previous research related to BV^*). Of particular interest is the subspace of $BV(\Omega)^*$ of those linear continuous functionals \mathfrak{T}_μ whose action on $u \in BV(\Omega) \cap L^\infty(\Omega)$ can be represented as

$$\mathfrak{T}_\mu(u) = \int_\Omega u^* d\mu, \quad (1.1)$$

where u^* is the precise representative of u and μ is a suitable finite Radon measure: these functionals are often called Radon measures in the dual of BV [23–25]. Given that u^* is well defined up to a set with zero $(n-1)$ -dimensional Hausdorff measure \mathcal{H}^{n-1} , an immediate property of the measures μ in the dual of BV is that $|\mu|(B) = 0$ for every Borel set with $\mathcal{H}^{n-1}(B) = 0$. As explored in [24, 25], under mild assumptions on the regularity of Ω we have that a finite Radon measure μ belongs to $BV(\Omega)^*$ in this sense if and only if $\mu = \operatorname{div} F$ for some vector field $F \in L^\infty(\Omega; \mathbb{R}^n)$. This shows a natural connection between the Radon measures in the dual of the space BV and the divergence-measure fields.

Given $p \in [1, \infty)$, a p -summable divergence-measure field on an open set Ω is a vector field $F \in L^p(\Omega; \mathbb{R}^n)$ such that its distributional divergence $\operatorname{div} F$ is a finite Radon measure on Ω . We denote the space of such fields as $\mathcal{DM}^p(\Omega)$. Initially introduced by Anzellotti in [3], the divergence-measure fields have been extensively studied, among other motivations, because of their natural applications in relation with generalized Gauss–Green formulas and weak formulations of some families of PDEs (see for instance [1, 4–9, 11–14, 21, 25–27, 29, 30]). Under this respect, it is important to note that, when $p \in [1, \infty)$, these fields can be naturally paired with scalar functions $u \in W^{1,p'}(\Omega)$, where p' is the conjugate exponent of p , since the scalar product $F \cdot \nabla u$ belongs to $L^1(\Omega)$. On the other hand, the case $p = \infty$ is more complex and interesting to study, since $p' = 1$, which includes the case of $u \in BV(\Omega)$. Following the approach of [14], for a given Borel function $\lambda : \Omega \rightarrow [0, 1]$, we can define the λ -pairing distribution between a vector field $F \in \mathcal{DM}^\infty(\Omega)$ and a scalar function $u \in BV(\Omega)$ as follows:

$$(F, Du)_\lambda := \operatorname{div}(uF) - u^\lambda \operatorname{div} F,$$

assuming $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$, where u^λ is defined \mathcal{H}^{n-1} -almost everywhere as the convex combination of u^+ and u^- (the upper and lower approximate limits of u) using the coefficients λ and $1 - \lambda$, respectively. We point out that this expression is well posed, since sets with zero \mathcal{H}^{n-1} -measure are also $|\operatorname{div} F|$ -negligible. As proved in [14], the λ -pairing is a distribution of order zero, i.e., it is a (finite) Radon measure. Moreover, the classical pairing introduced by Anzellotti in [3] corresponds to $\lambda \equiv \frac{1}{2}$, and in this case u^λ coincides \mathcal{H}^{n-1} -almost everywhere with u^* , the *precise representative* of u . For this reason, we set $(F, Du)_* := (F, Du)_{\frac{1}{2}}$. It is also clear that $(F, Du)_* = (F, Du)_\lambda$ for all $u \in BV(\Omega)$ such that $u^* \in L^1(\Omega; |\operatorname{div} F|)$ as long as the Borel function $\lambda : \Omega \rightarrow [0, 1]$ satisfies $\lambda(x) = \frac{1}{2}$ for $|\operatorname{div} F|$ -a.e. $x \in \Omega$.

Interestingly, every λ -pairing enjoys the same absolute continuity property with respect to the weak gradient of u :

$$|(F, Du)_\lambda| \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} |Du| \quad \text{on } \Omega. \quad (1.2)$$

This implies that the mapping

$$BV(\Omega) \cap L^\infty(\Omega) \ni u \mapsto (F, Du)_\lambda(\Omega)$$

defines a functional which is continuous with respect to the BV -(semi)norm. As noted in [14, Remark 4.6], the λ -pairing is linear with respect to u if and only if it coincides with the classical

one $(F, Du)_*$; that is, if $\lambda \equiv \frac{1}{2}$ up to $|\operatorname{div} F|$ -negligible sets. Therefore, the mapping

$$BV(\Omega) \cap L^\infty(\Omega) \ni u \mapsto (F, Du)_*(\Omega) \quad (1.3)$$

is a linear functional, continuous with respect to the BV -norm. These observations suggest a relation between the pairings and the functionals in the dual of BV .

In addition, in [25] it was proved that, in the case Ω is an open bounded set with Lipschitz boundary, a finite Radon measure μ belongs to $BV(\Omega)^*$ – that is, the functional $\mathfrak{T}_\mu : BV(\Omega) \cap L^\infty(\Omega) \rightarrow \mathbb{R}$ is continuous with respect to the BV -norm – if and only if there exists $C > 0$ such that

$$|\mu(U \cap \Omega)| \leq C P(U) \quad \text{for all open bounded sets } U \subset \mathbb{R}^n \text{ with smooth boundary,} \quad (1.4)$$

where $P(U)$ is the perimeter of the set U , which coincides with the surface measure of its boundary, $\mathcal{H}^{n-1}(\partial U)$, for smooth sets. Therefore, $\mu \in BV(\Omega)^*$ if and only if the measure μ enjoys this bound for its absolute value on open smooth sets in terms of their perimeter.

Finally, given $\mu \in BV(\Omega)^*$ there is the non-trivial question of whether $\mathfrak{T}_\mu(u)$ can be represented as the integral in (1.1) even for $u \in BV(\Omega) \setminus L^\infty(\Omega)$. Indeed, it can happen that $u^* \notin L^1(\Omega; |\mu|)$ (as we show in Remark 4.12 below), so that the integral representation would not be well-posed.

In this paper, we explore all these known connections recalled so far, in particular integrating and refining some of the results of [25].

More precisely, we consider the natural generalization of the condition (1.4): we say that a finite Radon measure $\mu \in \mathcal{M}(\Omega)$ satisfies a perimeter bound condition if there exists $L > 0$ such that

$$|\mu(E^1 \cap \Omega)| \leq L P(E) \quad \text{for all measurable sets } E \subset \Omega, \quad (1.5)$$

where E^1 is the set of points where the Lebesgue density of E is 1, also called the measure theoretic interior of E . In this case, we write $\mu \in \mathcal{PB}_L(\Omega)$, and we also set $\mathcal{PB}(\Omega) := \bigcup_{L>0} \mathcal{PB}_L(\Omega)$. We notice that in this definition we are testing the perimeter bound condition on the entire family of sets with finite perimeter. We actually prove that, if Ω is weakly regular (that is, it is a bounded open set such that $\mathcal{H}^{n-1}(\partial\Omega) = P(\Omega) < +\infty$) then (1.4) and (1.5) are equivalent (see Lemma 4.4 and Proposition 4.10(2)). However, our apparently stronger definition is motivated by the fact that the divergence measure of any $F \in \mathcal{DM}^\infty(\Omega)$ satisfies (1.5) for $L = \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}$, due to the generalized Gauss–Green formula: if $E \subseteq \Omega$ is a set of finite perimeter in \mathbb{R}^n such that either $E \subset\subset \Omega$ or Ω is weakly regular, then

$$\operatorname{div} F(E^1 \cap \Omega) = - \int_{\partial^* E} \operatorname{Tr}^i(F, \partial^* E) d\mathcal{H}^{n-1}, \quad (1.6)$$

where $\operatorname{Tr}^i(F, \partial^* E)$ is the interior normal trace of F and satisfies

$$\|\operatorname{Tr}^i(F, \partial^* E)\|_{L^\infty(\partial^* E; \mathcal{H}^{n-1})} \leq \|F\|_{L^\infty(E; \mathbb{R}^n)}.$$

This and similar formulas have been widely studied in the literature [4, 6, 7, 9, 11, 14, 29, 30], but we present in Section 2.2 some refinements which are suitable for our purposes.

Overall, we obtain the following refinement of [25, Theorem 8.2]: if Ω is a bounded open set with Lipschitz boundary, then

$$\mu \in \mathcal{PB}(\Omega) \iff \mu \in BV(\Omega)^* \iff \mu = \operatorname{div} F \quad \text{for some } F \in \mathcal{DM}^\infty(\Omega).$$

Thanks to the Gauss-Green formula (1.6), it is indeed intuitively clear that, if $\mu = \operatorname{div} F$ for some $F \in \mathcal{DM}^\infty(\Omega)$, then $\mu \in \mathcal{PB}_L(\Omega)$ for $L = \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}$. However, it is interesting to ask whether some kind of opposite implication holds true; that is, if, given $\mu \in \mathcal{PB}_L(\Omega)$, we can find $F \in \mathcal{DM}^\infty(\Omega)$ such that $\mu = \operatorname{div} F$ and $\|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq L$. Such result would be the natural extension of [25, Lemma 7.3], which states that, if Ω is a bounded open set with Lipschitz boundary and $\mathfrak{T} \in W_0^{1,1}(\Omega)^*$, then there exists $F \in \mathcal{DM}^\infty(\Omega)$ such that

$$\mathfrak{T}(u) = \int_{\Omega} u^* d \operatorname{div} F \quad \text{for all } u \in W_0^{1,1}(\Omega) \cap L^\infty(\Omega) \quad (1.7)$$

and we have

$$\|\mathfrak{T}\|_{W_0^{1,1}(\Omega)^*} = \min \left\{ \|G\|_{L^\infty(\Omega; \mathbb{R}^n)} : G \in \mathcal{DM}^\infty(\Omega) \text{ satisfying (1.7)} \right\}.$$

Therefore, we need to show that, if $\mu \in \mathcal{PB}_L(\Omega)$, then the related functional \mathfrak{T}_μ is well defined on the whole $W_0^{1,1}(\Omega)$ and it satisfies a bound for Sobolev functions similar to the perimeter bound; that is,

$$\left| \int_{\Omega} u^* d\mu \right| \leq L \int_{\Omega} |\nabla u| dx \quad \text{for all } u \in W_0^{1,1}(\Omega),$$

so that we get

$$\|\mathfrak{T}_\mu\|_{W_0^{1,1}(\Omega)^*} = \sup \left\{ \int_{\Omega} u^* d\mu : u \in W_0^{1,1}(\Omega), \|\nabla u\|_{L^1(\Omega; \mathbb{R}^n)} \leq 1 \right\} \leq L.$$

However, as already noted, it is in general not true that, if $\mu \in \mathcal{PB}(\Omega)$, then $u^* \in L^1(\Omega; |\mu|)$ for unbounded BV or even Sobolev functions (we give an example of such case in Remark 4.12). Therefore, it turns out that we need a stronger assumption on the measure μ : we say that μ is *admissible*, if $|\mu| \in BV(\Omega)^*$, see Definition 4.6. While it might be challenging to verify the admissibility assumption in general, we notice that, if Ω is an open bounded set with Lipschitz boundary, a relevant subclass of admissible measures consists of those that can be written as $\mu = h\mathcal{L}^n + \gamma\mathcal{H}^{n-1} \llcorner \Gamma$, where $h \in L^q(\Omega)$ for some $q > n$, $\gamma \in L^\infty(\Gamma; \mathcal{H}^{n-1})$ and $\Gamma \subset\subset \Omega$ is a compact set with finite \mathcal{H}^{n-1} measure and suitable decay properties on balls (see Example 4.14). If $\mu \in \mathcal{M}(\Omega)$ is admissible, it is indeed true that $u^\lambda \in L^1(\Omega; |\mu|)$ for all $u \in BV(\Omega)$ and $\lambda : \Omega \rightarrow [0, 1]$ Borel, thus including also the case $\lambda \equiv \frac{1}{2}$. A key step in the proof is Theorem 3.2, which is a refined version of the Anzellotti-Giaquinta approximation for functions $u \in BV(\Omega)$, since it additionally guarantees pointwise \mathcal{H}^{n-1} -a.e. convergence to any given λ -representative u^λ . It is also worth mentioning that such a result is of interest in itself, particularly concerning the theory of λ -pairings mentioned earlier, since it provides a new way of deriving (1.2) and it allows obtaining a new bound for the pairings in terms of the area functional, see Theorem 3.4; moreover, it plays a fundamental role in several other proofs along the paper. All in all, we can show that, if Ω is an open bounded set with Lipschitz boundary, $L > 0$ and $\mu \in \mathcal{PB}_L(\Omega)$ is admissible, then $\mu = \operatorname{div} F$ for some $F \in \mathcal{DM}^\infty(\Omega)$ with $\|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq L$ and

$$\left| \int_{\Omega} u^\lambda d\mu \right| \leq L \left(|Du|(\Omega) + \int_{\partial\Omega} |\operatorname{Tr}_{\partial\Omega}(u)| d\mathcal{H}^{n-1} \right) \quad \text{for all } u \in BV(\Omega) \text{ and } \lambda : \Omega \rightarrow [0, 1] \text{ Borel,} \quad (1.8)$$

see Proposition 4.19 and Lemma 4.21. This means that the constant L remains unchanged from the perimeter bound for sets to the estimate involving functions of bounded variations, and in this

way it refines the results of [25]. It is also interesting to notice that, if $\mu \in \mathcal{M}(\Omega)$ is admissible, then, for any vector field F such that $\operatorname{div} F = \mu$, the λ -pairing $(F, Du)_\lambda$ is well-defined for all BV function u and all Borel functions λ with values in $[0, 1]$ (see Remark 4.8). In addition, if Ω is an open bounded set with Lipschitz boundary, the action of the functional represented by an admissible measure μ can be expressed in terms of the pairing between $F \in \mathcal{DM}^\infty(\Omega)$, such that $\mu = \operatorname{div} F$, and functions in $BV(\Omega)$; that is,

$$\mathfrak{F}_\mu(u) = \int_\Omega u^* d\mu = -(F, Du)_*(\Omega) - \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(u) \operatorname{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1} \quad \text{for all } u \in BV(\Omega)$$

where $\operatorname{Tr}_{\partial\Omega}(u) \in L^1(\partial\Omega; \mathcal{H}^{n-1})$ and $\operatorname{Tr}^i(F, \partial\Omega) \in L^\infty(\partial\Omega; \mathcal{H}^{n-1})$ are the interior trace of u and the interior normal trace of F on the boundary of Ω , respectively. This fact relates to the previous observations on the pairing functional (1.3), since, if $\mu = \operatorname{div} F$ for some $F \in \mathcal{DM}^\infty(\Omega)$, then

$$\mathfrak{F}_\mu(u) = -(F, Du)_*(\Omega) \quad \text{for all } u \in BV_0(\Omega);$$

that is, $u \in BV(\Omega)$ with $\operatorname{Tr}_{\partial\Omega}(u) = 0$.

Finally, we prove that the perimeter bound and the admissibility conditions combined ensure enough stability under a suitable type of smooth approximation procedure (Proposition 4.23). More precisely, if Ω is an open bounded set with Lipschitz boundary and $\mu \in \mathcal{PB}_L(\Omega)$ is an admissible measure, then $\mu = \operatorname{div} F$ for some $F \in \mathcal{DM}^\infty(\Omega)$ with $\|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq L$ and so, relying on the fact that the Anzellotti-Giaquinta-type regularization of a vector field almost preserves its L^∞ norm, we can approximate μ in the weak- $*$ sense by a sequence of absolutely continuous measures $\mu_j \mathcal{L}^n \in \mathcal{PB}_{L_j}(\Omega)$ with smooth density functions, where $L_j \rightarrow L$ as $j \rightarrow +\infty$.

All these facts are of great relevance in their application to the study of the prescribed mean curvature measure equation

$$\operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = \mu \quad \text{on } \Omega, \quad (1.9)$$

for an admissible measure $\mu \in \mathcal{PB}_L(\Omega)$ for some $L \in (0, 1)$, which is the core of our subsequent work [20]. Following the approach of [26], we consider the following weak formulation of the equation (1.9).

Definition 1.1. *We say that $u \in BV(\Omega)$ is a weak solution to the prescribed mean curvature measure equation if there exist $T \in L^\infty(\Omega; \mathbb{R}^n)$ and a Borel function $\lambda : \Omega \rightarrow [0, 1]$ such that*

$$\begin{aligned} \|T\|_{L^\infty(\Omega; \mathbb{R}^n)} &\leq 1, \\ \operatorname{div} T &= \mu \quad \text{on } \Omega, \\ (T, Du)_\lambda &= \sqrt{1 + |Du|^2} - \sqrt{1 - |T|^2} \mathcal{L}^n \quad \text{on } \Omega, \end{aligned}$$

where the last two identities involve scalar Radon measures in $\mathcal{M}(\Omega)$ and $\lambda = \lambda_\mu$ is the characteristic function of a Borel set satisfying $(1 - \lambda)\mu = \mu^+$, where μ^\pm are the positive and negative parts of μ .

The existence of weak solutions of (1.9) is obtained via the direct method of Calculus of Variations, which involves proving coercivity and lower semicontinuity for the functional

$$\mathfrak{J}_\mu[u] := \sqrt{1 + |Du|^2}(B) + \int_\Omega u^- d\mu^+ - \int_\Omega u^+ d\mu^-,$$

In particular, (1.8) directly implies that \mathfrak{J}_μ is coercive, and the stability of the admissibility and perimeter bound conditions are exploited in order to prove a Gamma-convergence result for a sequence of functionals \mathfrak{J}_{μ_j} (see [20, Theorem 6.2]).

2. PRELIMINARIES

Through the rest of the paper, we work in an open set $\Omega \subset \mathbb{R}^n$. We denote by \mathcal{L}^n the Lebesgue measure, and by \mathcal{H}^m the m -dimensional Hausdorff measure, for $m \in [0, n]$, although we shall focus on the case $m = n - 1$. We denote by $\mathcal{M}(\Omega)$ the space of finite Radon measures on Ω , and by $\mathcal{M}_{\mathcal{H}}(\Omega)$ the space of measures in $\mathcal{M}(\Omega)$ that are absolutely continuous with respect to \mathcal{H}^{n-1} ; that is,

$$\mathcal{M}_{\mathcal{H}}(\Omega) := \{\mu \in \mathcal{M}(\Omega) : |\mu|(B) = 0 \text{ for all Borel sets } B \subset \Omega \text{ such that } \mathcal{H}^{n-1}(B) = 0\}. \quad (2.1)$$

Given two measures $\mu_1, \mu_2 \in \mathcal{M}(\Omega)$, we say that $\mu_1 \leq \mu_2$ on Ω if $\mu_1(B) \leq \mu_2(B)$ for all Borel sets $B \subseteq \Omega$. If $\mu \in \mathcal{M}(\Omega)$ satisfies $\mu \geq 0$ on Ω , then we say that μ is nonnegative. Thanks to the Hahn's decomposition theorem, we know that for any $\mu \in \mathcal{M}(\Omega)$ there exist two nonnegative measures μ^+ and μ^- (the positive and negative part of μ , respectively), which satisfy $\mu = \mu^+ - \mu^-$ and are concentrated on mutually disjoint Borel sets Ω_+, Ω_- such that $\Omega = \Omega_+ \cup \Omega_-$. In particular, we have $\mu_+ = \mu \llcorner \Omega_+$, $\mu_- = -\mu \llcorner \Omega_-$, and $|\mu| = \mu^+ + \mu^-$. In addition, by Lebesgue-Besicovitch differentiation theorem, for $|\mu|$ -a.e. $x \in \Omega$ we have

$$\frac{d\mu^\pm}{d|\mu|}(x) = \begin{cases} 1 & \text{if } x \in \Omega^\pm \\ 0 & \text{if } x \in \Omega^\mp. \end{cases} \quad (2.2)$$

Given $\mu \in \mathcal{M}(\Omega)$, Radon-Nikodym theorem and Lebesgue's decomposition theorem ensure that we can decompose it as

$$\mu = \mu^{ac} + \mu^s$$

where $\mu^{ac} = g\mathcal{L}^n$, for some $g \in L^1(\Omega)$, is the absolutely continuous part and μ^s is the singular part.

2.1. Functions of bounded variation. We say that $u \in L^1(\Omega)$ is a function of bounded variation, and we write $u \in BV(\Omega)$, if its distributional gradient Du is a vector valued Radon measure whose total variation $|Du|$ is a finite measure on Ω . $BV(\Omega)$ is a Banach space once equipped with the norm

$$\|u\|_{BV(\Omega)} = \|u\|_{L^1(\Omega)} + |Du|(\Omega).$$

However, the convergence induced by this norm is too strong, and hence it is customary to consider a weaker type of convergence for sequences of BV functions, the *strict convergence*: given $u \in BV(\Omega)$, we say that a sequence $(u_j)_{j \in \mathbb{N}}$ converges to u in $BV(\Omega)$ -strict if

$$\lim_{j \rightarrow +\infty} \|u - u_j\|_{L^1(\Omega)} + \left| |Du|(\Omega) - |Du_j|(\Omega) \right| = 0.$$

Following the notation of [2, Section 3.6], we say that a function $u \in L^1_{\text{loc}}(\Omega)$ has approximate limit at $x \in \Omega$ if there exists $z \in \mathbb{R}$ such that

$$\lim_{r \rightarrow 0} \int_{B_r(x)} |u(y) - z| dy = 0, \quad (2.3)$$

and we denote by $\tilde{u}(x)$ the value, z , of the approximate limit of u at x . In this case, we say that x is a Lebesgue point of u . For a given measurable set $E \subset \mathbb{R}^n$, we define the *measure theoretic interior* of E as

$$E^1 := \left\{ x \in \mathbb{R}^n : \lim_{r \rightarrow 0} \frac{|E \cap B_r(x)|}{|B_r(x)|} = 1 \right\} = \{x \in \mathbb{R}^n : \tilde{\chi}_E(x) = 1\}.$$

As customary, the approximate discontinuity set S_u is defined as the set of points where the approximate limit does not exist. In addition, we say that x belongs to J_u (the set of approximate jump points of u) if there exists $a, b \in \mathbb{R}$, $a \neq b$, and $\nu \in \mathbb{S}^{n-1}$ such that

$$\lim_{r \rightarrow 0} \int_{B_r^+(x, \nu)} |u(y) - a| dy = 0 \quad \text{and} \quad \lim_{r \rightarrow 0} \int_{B_r^-(x, \nu)} |u(y) - b| dy = 0, \quad (2.4)$$

where $B_r^\pm(x, \nu) := \{y \in B_r(x) : \pm(y - x) \cdot \nu \geq 0\}$. The triplet (a, b, ν) is uniquely determined by (2.4) up to a permutation of (a, b) and a change of sign of ν , and we denote it by $(u^+(x), u^-(x), \nu_u(x))$. For the approximate traces $u^\pm(x)$, we adopt the convention of having $u^+(x) > u^-(x)$. We can actually extend the approximate traces also for $x \in \Omega \setminus S_u$, by setting $u^+(x) = u^-(x) = \tilde{u}(x)$.

Given $u \in BV_{\text{loc}}(\Omega)$, following [2, Corollary 3.80] we define its precise representative

$$u^* : \Omega \setminus (S_u \setminus J_u) \rightarrow \mathbb{R}$$

by setting

$$u^*(x) = \begin{cases} \tilde{u}(x) & \text{if } x \in \Omega \setminus S_u, \\ \frac{u^+(x) + u^-(x)}{2} & \text{if } x \in J_u. \end{cases}$$

Since $\mathcal{H}^{n-1}(S_u \setminus J_u) = 0$ by [2, Theorem 3.78], $u^*(x)$ is well defined for \mathcal{H}^{n-1} -a.e. $x \in \Omega$ and satisfies

$$u^*(x) = \lim_{r \rightarrow 0} \int_{B_r(x)} u(y) dy \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in \Omega. \quad (2.5)$$

In addition, if we consider the extensions of the approximate traces, we also get

$$u^*(x) = \frac{u^+(x) + u^-(x)}{2} \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in \Omega. \quad (2.6)$$

In this way, we can see u^* as the average between u^+ and u^- . Therefore, it seems natural to generalize u^* by taking any convex combination of the approximate traces. More precisely, for a fixed a Borel function $\lambda : \Omega \rightarrow [0, 1]$ we define the λ -representative of u as

$$u^\lambda = \lambda u^+ + (1 - \lambda)u^- \quad \text{on } \Omega \setminus (S_u \setminus J_u),$$

which is well-defined for \mathcal{H}^{n-1} -a.e. $x \in \Omega$. Clearly, such representative could be defined for any $\lambda \in \mathcal{B}_b(\Omega)$, that is, any bounded Borel function $\lambda : \Omega \rightarrow \mathbb{R}$, but only convex combinations of u^\pm have relevant properties in relation with generalized notions of pairings between divergence-measure fields and scalar functions of bounded variation.

Let $N > 0$ and T_N be the truncation operator, that is, the 1-Lipschitz map defined as

$$T_N(t) = \begin{cases} N & \text{if } t > N, \\ t & \text{if } |t| \leq N, \\ -N & \text{if } t < -N. \end{cases}$$

We recall the statement of [2, Proposition 3.69 (c)]: if $u \in L^1_{\text{loc}}(\Omega)$ and $x \in J_u$, then $x \in J_{T_N(u)}$ if and only if $T_N(u^+(x)) \neq T_N(u^-(x))$, and in this case we have $T_N(u)^\pm(x) = T_N(u^\pm(x))$. Otherwise, $x \notin S_{T_N(u)}$ and

$$\widetilde{T_N(u)}(x) = T_N(\tilde{u})(x) = T_N(u^+(x)) = T_N(u^-(x)).$$

Hence, if we extend the approximate traces of $T_N(u)$ as above, we get

$$T_N(u)^\pm(x) = T_N(u^\pm)(x) \text{ for all } x \in \Omega \setminus (S_u \setminus J_u).$$

By exploiting the fact that

$$\left| \frac{T_N(\alpha) + T_N(\beta)}{2} \right| \leq \left| T_N\left(\frac{\alpha + \beta}{2}\right) \right| \leq \left| \frac{\alpha + \beta}{2} \right| \text{ for all } \alpha, \beta \in \mathbb{R},$$

we prove that

$$|T_N(u)^*| \leq |u^*| \text{ on } \Omega \setminus (S_u \setminus J_u).$$

Given $u \in BV(\Omega)$, we have that $T_N(u) \in BV(\Omega)$ for all $N > 0$. In particular, we deduce the following well-known estimate on the precise representative of the truncation of u :

$$|T_N(u)^*(x)| \leq |u^*(x)| \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in \Omega. \quad (2.7)$$

In addition, the λ -representative of $T_N(u)$ is well-defined, and we can get convergence properties and bounds for $T_N(u)^\lambda$ that are uniform in N (see Proposition 2.1 below). However, we notice that we cannot obtain an analogous estimate $|T_N(u)^\lambda(x)| \leq |u^\lambda(x)|$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$, unless $\lambda \in \{0, \frac{1}{2}, 1\}$. As an example, let $N > 0$, $\lambda \equiv \frac{1}{3}$ and

$$u(x) = \begin{cases} 2N & \text{if } x \in B(0, 1), \\ -\frac{5}{4}N & \text{if } x \notin B(0, 1). \end{cases}$$

Then, for all $x \in \partial B(0, 1)$, we have $u^+(x) = 2N$ and $u^-(x) = -\frac{5}{4}N$, so that

$$u^\lambda(x) = \frac{1}{3}u^+(x) + \frac{2}{3}u^-(x) = \frac{2}{3}N \left(1 - \frac{5}{4}\right) = -\frac{N}{6} \text{ for all } x \in \partial B(0, 1).$$

On the other hand, notice that

$$T_N(u)^\lambda(x) = \frac{1}{3}T_N(u^+)(x) + \frac{2}{3}T_N(u^-)(x) = \frac{N}{3} - \frac{2N}{3} = -\frac{N}{3} \text{ for all } x \in \partial B(0, 1),$$

so that we do not have $|T_N(u)^\lambda| \leq |u^\lambda|$ on $\partial B(0, 1)$. Nevertheless, we are able to obtain the following result (which is a somewhat modified version of [14, Proposition 3.4]).

Proposition 2.1. *Let $u \in BV_{\text{loc}}(\Omega)$, $\lambda : \Omega \rightarrow [0, 1]$ be a Borel function and $N > 0$. For all $x \in \Omega \setminus (S_u \setminus J_u)$ we set*

$$M[u, \lambda](x) := \min \left\{ \lambda(x)|u^+(x)| + (1 - \lambda(x))|u^-(x)|, \right. \\ \left. |u^*(x)| + \left| \lambda(x) - \frac{1}{2} \right| (|u^+(x)| + |u^-(x)|) \right\}. \quad (2.8)$$

Then $M[u, \lambda](x)$ is well defined for \mathcal{H}^{n-1} -a.e. $x \in \Omega$, and it satisfies

$$|T_N(u)^\lambda(x)| \leq M[u, \lambda](x) \text{ and } |u^\lambda(x)| \leq M[u, \lambda](x) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in \Omega. \quad (2.9)$$

Moreover, we have

$$T_N(u)^\lambda(x) = u^\lambda(x) \quad (2.10)$$

for all $x \in \Omega$ such that $-N \leq u^-(x) \leq u^+(x) \leq N$, which in turn implies $T_N(u)^\lambda(x) \rightarrow u^\lambda(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$, as $N \rightarrow +\infty$.

Proof. Assuming $x \in \Omega$ is such that $-N \leq u^-(x) \leq u^+(x) \leq N$, we have $T_N(u)^\pm(x) = u^\pm(x)$. This immediately implies that

$$T_N(u)^\lambda(x) = \lambda(x)T_N(u)^+(x) + (1 - \lambda(x))T_N(u)^-(x) = \lambda(x)u^+(x) + (1 - \lambda(x))u^-(x) = u^\lambda(x)$$

and this shows (2.10). Since $u^+(x)$ and $u^-(x)$ are well-defined and finite for \mathcal{H}^{n-1} -a.e. $x \in \Omega$, we notice that (2.10) easily implies $T_N(u)^\lambda(x) \rightarrow u^\lambda(x)$ as $N \rightarrow +\infty$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$.

Given that $\mathcal{H}^{n-1}(S_u \setminus J_u) = 0$, it is clear that $M[u, \lambda](x)$ is well defined for \mathcal{H}^{n-1} -a.e. $x \in \Omega$. Now, in order to prove the first inequality in (2.9), we must show two separate inequalities. The first inequality directly follows from the definition of T_N :

$$\begin{aligned} |T_N(u)^\lambda(x)| &\leq \lambda(x)|T_N(u)^+(x)| + (1 - \lambda(x))|T_N(u)^-(x)| \\ &\leq \lambda(x)|u^+(x)| + (1 - \lambda(x))|u^-(x)|. \end{aligned}$$

The second inequality is obtained by adding and subtracting $T_N(u)^*$ and by using (2.7), that is, for \mathcal{H}^{n-1} -a.e. $x \in \Omega$ we have

$$\begin{aligned} |T_N(u)^\lambda(x)| &\leq |T_N(u)^*(x)| + |T_N(u)^\lambda(x) - T_N(u)^*(x)| \\ &\leq |u^*(x)| + |(\lambda(x) - 1/2)T_N(u)^+(x) + (1/2 - \lambda(x))T_N(u)^-(x)| \\ &\leq |u^*(x)| + |\lambda(x) - 1/2|(|u^+(x)| + |u^-(x)|). \end{aligned}$$

By combining the two previous estimates, we get (2.9) as wanted. Finally, we argue in a similar way with $|u^\lambda|$ to conclude. \square

Remark 2.2. We observe that (2.9) gives a finer upper bound for $T_N(u)^\lambda$ than the one given in [14, Proposition 3.4]. In addition, we notice that, if $\lambda \equiv \frac{1}{2}$, then (2.9) reduces to the standard estimate (2.7). If we consider instead the cases $\lambda \equiv 1$ and $\lambda \equiv 0$, then we have $M[u, 1] = |u^+|$ and $M[u, 0] = |u^-|$.

Given a function $u \in BV(\Omega)$ we define $\sqrt{1 + |Du|^2}$ as the distributional area factor of the graph $\{(x, t) : u(x) = t\} \subset \Omega \times \mathbb{R}$, as done in [19]. While the expression makes sense for \mathcal{L}^n -a.e. $x \in \Omega$ for a function $u \in W_{\text{loc}}^{1,1}(\Omega)$, when $u \in BV(\Omega)$ we more generally define

$$\int_U \sqrt{1 + |Du|^2} := \sup \left\{ \int_\Omega \eta + u \operatorname{div} \phi \, dx : (\phi, \eta) \in C_c^1(U; \mathbb{R}^n \times \mathbb{R}), |(\phi, \eta)| \leq 1 \right\},$$

for any open set $U \subset \Omega$. It is then easy to see that, if we denote the absolutely continuous part of Du by $\nabla u \mathcal{L}^n$, and the singular part by $D^s u$, we have

$$\sqrt{1 + |Du|^2} = \sqrt{1 + |\nabla u|^2} \mathcal{L}^n + |D^s u|. \quad (2.11)$$

Given a measurable set E , we say that E is a set of finite perimeter in Ω if $|D\chi_E|(\Omega) < +\infty$, and we set $P(E; \Omega) = |D\chi_E|(\Omega)$ to be the perimeter of E in Ω . When $\Omega = \mathbb{R}^n$ we simply write $P(E)$. We refer to [2, 22] for the definition and properties of $\partial^* E$, the *reduced boundary* of E , for which

one has $P(E; \Omega) = \mathcal{H}^{n-1}(\partial^* E \cap \Omega)$. In particular, we recall that $\chi_E^+ = \chi_{E^1 \cup \partial^* E}$ and $\chi_E^- = \chi_{E^1}$, so that we deduce the following explicit formulas for the precise and λ -representatives of χ_E :

$$\chi_E^* = \chi_{E^1} + \frac{1}{2}\chi_{\partial^* E} \quad \text{and} \quad \chi_E^\lambda = \chi_{E^1} + \lambda\chi_{\partial^* E} \quad \mathcal{H}^{n-1}\text{-a.e. in } \Omega. \quad (2.12)$$

A remarkable property of sets of finite perimeter E in \mathbb{R}^n is the isoperimetric inequality [2, Theorem 3.46]: there exists a constant $c_n > 0$ such that

$$\min\{|E|, |\mathbb{R}^n \setminus E|\}^{1-\frac{1}{n}} \leq c_n P(E).$$

As a consequence, as long as $|\Omega| < +\infty$, for all sets $E \subset \Omega$ of finite perimeter in \mathbb{R}^n we have

$$|E \cap \Omega| \leq |\Omega|^{\frac{1}{n}} |E|^{1-\frac{1}{n}} \leq c_n |\Omega|^{\frac{1}{n}} P(E),$$

which immediately implies

$$\|\chi_E\|_{BV(\Omega)} = |E \cap \Omega| + P(E, \Omega) \leq (c_n |\Omega|^{\frac{1}{n}} + 1) P(E). \quad (2.13)$$

If Ω has bounded Lipschitz boundary, we denote by $\text{Tr}_{\partial\Omega}(u)$ the trace of u over $\partial\Omega$ and recall that the trace operator $\text{Tr}_{\partial\Omega} : BV(\Omega) \rightarrow L^1(\partial\Omega; \mathcal{H}^{n-1})$ is linear, surjective and continuous with respect to the topology induced by the strict convergence (see for instance [2, Theorem 3.88]). For the ease of the reader, we prove a useful result concerning a Cavalieri-type inequality involving the trace operator for BV functions.

Lemma 2.3. *Let Ω be an open, bounded set with Lipschitz boundary. Let $f \in BV(\Omega)$ be a nonnegative function, and set $E_t = \{x : f(x) > t\}$ for $t > 0$. Then*

$$\int_0^{+\infty} \int_{\partial\Omega} \text{Tr}_{\partial\Omega}(\chi_{E_t}) d\mathcal{H}^{n-1} dt \leq \int_{\partial\Omega} \text{Tr}_{\partial\Omega}(f) d\mathcal{H}^{n-1}.$$

Proof. We have

$$\begin{aligned} \int_{\partial\Omega} \text{Tr}_{\partial\Omega}(f)(x) d\mathcal{H}^{n-1}(x) &= \int_{\partial\Omega} \lim_{r \rightarrow 0} \frac{1}{|\Omega \cap B_r(x)|} \int_{\Omega \cap B_r(x)} f(y) dy d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial\Omega} \lim_{r \rightarrow 0} \frac{1}{|\Omega \cap B_r(x)|} \int_{\Omega \cap B_r(x)} \int_0^{+\infty} \chi_{E_t}(y) dt dy d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial\Omega} \lim_{r \rightarrow 0} \int_0^{+\infty} \frac{1}{|\Omega \cap B_r(x)|} \int_{\Omega \cap B_r(x)} \chi_{E_t}(y) dy dt d\mathcal{H}^{n-1}(x) \\ &\geq \int_{\partial\Omega} \int_0^{+\infty} \liminf_{r \rightarrow 0} \frac{1}{|\Omega \cap B_r(x)|} \int_{\Omega \cap B_r(x)} \chi_{E_t}(y) dy dt d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial\Omega} \int_0^{+\infty} \text{Tr}_{\partial\Omega}(\chi_{E_t})(x) dt d\mathcal{H}^{n-1}(x) \\ &= \int_0^{+\infty} \int_{\partial\Omega} \text{Tr}_{\partial\Omega}(\chi_{E_t})(x) d\mathcal{H}^{n-1}(x) dt, \end{aligned}$$

where the inequality in the fourth line follows from Fatou's lemma, while the exchange of the integration orders is a consequence of Tonelli's theorem. \square

2.2. Divergence-measure fields and λ -pairings. We recall the notions of divergence-measure field and of pairing between a function of bounded variation and an essentially bounded divergence-measure field.

Definition 2.4. *A vector field $F \in L^\infty(\Omega; \mathbb{R}^n)$ is called an essentially bounded divergence-measure field, and we write $F \in \mathcal{DM}^\infty(\Omega)$, if $\operatorname{div} F \in \mathcal{M}(\Omega)$. A vector field $F \in L^\infty_{\text{loc}}(\Omega; \mathbb{R}^n)$ is called a locally essentially bounded divergence-measure field, and we write $F \in \mathcal{DM}^\infty_{\text{loc}}(\Omega)$, if $F \in \mathcal{DM}^\infty(U)$ for any open set $U \subset\subset \Omega$.*

It has been proved by Chen and Frid [5] (see also [18], for an improved proof), that, if $F \in \mathcal{DM}^\infty(\Omega)$ and $u \in BV(\Omega) \cap L^\infty(\Omega)$, then the product uF belongs to $\mathcal{DM}^\infty(\Omega)$; and an analogous result holds also locally. Furthermore, we recall the fact that, if $F \in \mathcal{DM}^\infty_{\text{loc}}(\Omega)$, then $|\operatorname{div} F| \ll \mathcal{H}^{n-1}$, for which we refer to [5] and [29]. Hence, if u^* is the precise representative of u , the measure $u^* \operatorname{div} F$ is well defined, u^* being defined \mathcal{H}^{n-1} -a.e. on Ω .

Actually, the L^∞ -assumption on u is not strictly necessary, as one can see by following the approach of [14] in defining the notion of λ -pairing.

Definition 2.5. *Given a vector field $F \in \mathcal{DM}^\infty_{\text{loc}}(\Omega)$, a scalar function $u \in BV_{\text{loc}}(\Omega)$ and a Borel function $\lambda : \Omega \rightarrow [0, 1]$ such that $u^\lambda \in L^1_{\text{loc}}(\Omega; |\operatorname{div} F|)$, we define the λ -pairing between F and Du as the distribution $(F, Du)_\lambda$ given by*

$$(F, Du)_\lambda := \operatorname{div}(uF) - u^\lambda \operatorname{div} F. \quad (2.14)$$

Roughly speaking, the notion of λ -pairing extends the classical dot product between F and Du . We also remark that in the case $\lambda \equiv \frac{1}{2}$ one recovers the classical pairing first considered by Anzellotti in [3], which we denote by $(\bar{F}, Du)_*$. As proved in [14], the λ -pairing is indeed a Radon measure. For the ease of the reader, we recall its main properties in the following statement, that summarizes [14, Proposition 4.4] (see also [7, Remark 3.6]).

Theorem 2.6. *Let $F \in \mathcal{DM}^\infty(\Omega)$, $u \in BV(\Omega)$ and $\lambda : \Omega \rightarrow [0, 1]$ be a Borel function. If $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$, then we have $\operatorname{div}(uF), (F, Du)_\lambda \in \mathcal{M}(\Omega)$, with*

$$\operatorname{div}(uF) = u^\lambda \operatorname{div} F + (F, Du)_\lambda \quad \text{on } \Omega, \quad (2.15)$$

and

$$|(F, Du)_\lambda| \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} |Du| \quad \text{on } \Omega. \quad (2.16)$$

The assumption $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$ in Theorem 2.6 is equivalent to the summability of any $\bar{\lambda}$ -representative of u with respect to the measure $|\operatorname{div} F|$, as showed in [14, Lemma 3.2]. In particular, it is also equivalent to the summability of the majorant function $M[u, \lambda]$ given by (2.8). We collect this results in the proposition below.

Proposition 2.7. *Let $F \in \mathcal{DM}^\infty(\Omega)$ and $u \in BV(\Omega)$. Let $\lambda_1, \lambda_2 : \Omega \rightarrow [0, 1]$ be Borel functions. Then we have $u^{\lambda_1} \in L^1(\Omega; |\operatorname{div} F|)$ if and only if $u^{\lambda_2} \in L^1(\Omega; |\operatorname{div} F|)$. In addition, if $\lambda : \Omega \rightarrow [0, 1]$ is a Borel function, then $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$ if and only if $M[u, \lambda] \in L^1(\Omega; |\operatorname{div} F|)$.*

Proof. The first equivalence is given by [14, Lemma 3.2]. Then, if we assume $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$ for some $\lambda : \Omega \rightarrow [0, 1]$, by the first part of the statement, we have $u^+, u^- \in L^1(\Omega; |\operatorname{div} F|)$, since they correspond to the choices $\bar{\lambda} \equiv 1$ and $\bar{\lambda} \equiv 0$, respectively. Hence, we see that $M[u, \lambda] \in L^1(\Omega; |\operatorname{div} F|)$, in the light of (2.8) and the fact that $|\operatorname{div} F| \ll \mathcal{H}^{n-1}$. Finally, the second estimate in (2.9) immediately gives the opposite implication. \square

One of the main applications of the theory of divergence-measure fields has consisted in generalizing integration by parts formulas for rough domains and weakly differentiable functions. For the ease of the reader we collect in the following statements some versions of these results, suitably refined for the purposes of this paper. The first one is simply [9, Lemma 3.1].

Lemma 2.8. *Let $F \in \mathcal{DM}^\infty(\Omega)$ be such that $\text{supp}(F) \subset\subset \Omega$. Then we have $\text{div } F(\Omega) = 0$.*

Now we deal with the integration on sets with finite perimeter. Such Gauss–Green formulas have been widely studied in the literature [4, 6, 7, 9, 11, 14, 29, 30]. However, we provide here a general result, allowing for sets which can touch the boundary of the definition domain Ω , under minimal regularity assumptions. To this purpose, we recall that an open set Ω is said to be *weakly regular* if it is a bounded set with finite perimeter in \mathbb{R}^n such that $\mathcal{H}^{n-1}(\partial\Omega) = \mathcal{H}^{n-1}(\partial^* \Omega)$, or, equivalently, $\mathcal{H}^{n-1}(\partial\Omega \setminus \partial^* \Omega) = 0$.

Theorem 2.9. *Let $F \in \mathcal{DM}^\infty(\Omega)$ and let $E \subseteq \Omega$ be of finite perimeter in \mathbb{R}^n . Assume that either $E \subset\subset \Omega$ or Ω is weakly regular. Then, there exist the interior and exterior normal traces of F on $\partial^* E$; that is, the functions*

$$\text{Tr}^i(F, \partial^* E), \text{Tr}^e(F, \partial^* E) \in L^\infty(\partial^* E; \mathcal{H}^{n-1}),$$

which satisfy

$$\begin{cases} \text{div } F(E^1 \cap \Omega) = - \int_{\partial^* E} \text{Tr}^i(F, \partial^* E) d\mathcal{H}^{n-1} \\ \|\text{Tr}^i(F, \partial^* E)\|_{L^\infty(\partial^* E; \mathcal{H}^{n-1})} \leq \|F\|_{L^\infty(E; \mathbb{R}^n)}, \end{cases} \quad (2.17)$$

and

$$\begin{cases} \text{div } F((E^1 \cup \partial^* E) \cap \Omega) = - \int_{\partial^* E} \text{Tr}^e(F, \partial^* E) d\mathcal{H}^{n-1} \\ \|\text{Tr}^e(F, \partial^* E)\|_{L^\infty(\partial^* E; \mathcal{H}^{n-1})} \leq \begin{cases} \|F\|_{L^\infty(\Omega \setminus E; \mathbb{R}^n)} & \text{if } E \subset\subset \Omega \\ \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} & \text{otherwise} \end{cases}. \end{cases} \quad (2.18)$$

In particular, if Ω is weakly regular, then there exists $\text{Tr}^i(F, \partial\Omega) \in L^\infty(\partial\Omega; \mathcal{H}^{n-1})$ satisfying

$$\text{div } F(\Omega) = - \int_{\partial\Omega} \text{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1} \quad \text{and} \quad \|\text{Tr}^i(F, \partial\Omega)\|_{L^\infty(\partial\Omega; \mathcal{H}^{n-1})} \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}. \quad (2.19)$$

Proof. For the proof of (2.17) and (2.18) in the case $E \subset\subset \Omega$ we refer to [9, Theorem 3.2 and Corollary 3.6]. Then, if Ω is weakly regular, by [9, Corollary 5.5 and Remark 5.6] we have that the zero extension of F to \mathbb{R}^n , defined as

$$\hat{F}(x) := \begin{cases} F(x) & \text{if } x \in \Omega \\ 0 & \text{if } x \in \mathbb{R}^n \setminus \Omega \end{cases},$$

satisfies $\hat{F} \in \mathcal{DM}^\infty(\mathbb{R}^n)$, and there exist $\text{Tr}^i(\hat{F}, \partial\Omega), \text{Tr}^e(\hat{F}, \partial\Omega) \in L^\infty(\partial\Omega; \mathcal{H}^{n-1})$ such that

$$\text{div } \hat{F} = \text{div } F \llcorner \Omega + \text{Tr}^i(\hat{F}, \partial\Omega) \mathcal{H}^{n-1} \llcorner \partial\Omega \quad \text{and} \quad \text{Tr}^e(\hat{F}, \partial\Omega) = 0. \quad (2.20)$$

Hence, if $E \subset \Omega$ is a set of finite perimeter in \mathbb{R}^n , we can apply (2.17) and (2.18) to \hat{F} and obtain

$$\begin{aligned} \operatorname{div} \hat{F}(E^1) &= - \int_{\partial^* E} \operatorname{Tr}^i(\hat{F}, \partial^* E) d\mathcal{H}^{n-1}, \\ \operatorname{div} \hat{F}(E^1 \cup \partial^* E) &= - \int_{\partial^* E} \operatorname{Tr}^e(\hat{F}, \partial^* E) d\mathcal{H}^{n-1}. \end{aligned} \quad (2.21)$$

Then, we notice that it must be $\mathcal{H}^{n-1}(E^1 \setminus \Omega) = 0$, since $\mathcal{H}^{n-1}(\Omega^1 \setminus \Omega) = 0$ (see [9, Corollary 5.5]) and $E^1 \subset \Omega^1$. Therefore, we have

$$\operatorname{div} \hat{F}(E^1) = \operatorname{div} \hat{F}(E^1 \cap \Omega) + \operatorname{div} \hat{F}(E^1 \setminus \Omega) = \operatorname{div} F(E^1 \cap \Omega),$$

since $|\operatorname{div} \hat{F}| \ll \mathcal{H}^{n-1}$. Hence, given that

$$\|\operatorname{Tr}^i(F, \partial^* E)\|_{L^\infty(\partial^* E; \mathcal{H}^{n-1})} \leq \|\hat{F}\|_{L^\infty(E; \mathbb{R}^n)} = \|F\|_{L^\infty(E; \mathbb{R}^n)},$$

we can set $\operatorname{Tr}^i(F, \partial^* E) := \operatorname{Tr}^i(\hat{F}, \partial^* E)$ to include the case $\mathcal{H}^{n-1}(\partial^* E \cap \partial\Omega) > 0$, with a little abuse of notation, thus obtaining (2.17). As for (2.18), we exploit (2.20) and the fact that $\mathcal{H}^{n-1}(E^1 \setminus \Omega) = 0$ to get

$$\operatorname{div} \hat{F}(E^1 \cup \partial^* E) = \operatorname{div} F((E^1 \cup \partial^* E) \cap \Omega) + \int_{\partial^* E \cap \partial\Omega} \operatorname{Tr}^i(\hat{F}, \partial\Omega) d\mathcal{H}^{n-1}.$$

In addition, we notice that $\nu_E(x) = \nu_\Omega(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \partial^* E \cap \partial\Omega$, since $E \subset \Omega$, so that, by [9, Proposition 4.10], we have

$$\operatorname{Tr}^i(\hat{F}, \partial^* E) = \operatorname{Tr}^i(\hat{F}, \partial\Omega) \quad \text{and} \quad \operatorname{Tr}^e(\hat{F}, \partial^* E) = \operatorname{Tr}^e(\hat{F}, \partial\Omega) = 0 \quad \mathcal{H}^{n-1}\text{-a.e. on } \partial^* E \cap \partial\Omega,$$

due to (2.21). Hence, exploiting (2.20) we get

$$\begin{aligned} \operatorname{div} F((E^1 \cup \partial^* E) \cap \Omega) &= - \int_{\partial^* E} \operatorname{Tr}^e(\hat{F}, \partial^* E) d\mathcal{H}^{n-1} - \int_{\partial^* E \cap \partial\Omega} \operatorname{Tr}^i(\hat{F}, \partial\Omega) d\mathcal{H}^{n-1} \\ &= - \int_{\partial^* E \setminus \partial\Omega} \operatorname{Tr}^e(\hat{F}, \partial^* E) d\mathcal{H}^{n-1} - \int_{\partial^* E \cap \partial\Omega} \operatorname{Tr}^i(\hat{F}, \partial^* E) d\mathcal{H}^{n-1}. \end{aligned}$$

Hence, we obtain (2.18) by setting

$$\operatorname{Tr}^e(F, \partial^* E) := \begin{cases} \operatorname{Tr}^e(\hat{F}, \partial^* E) & \text{on } \partial^* E \setminus \partial\Omega \\ \operatorname{Tr}^i(\hat{F}, \partial^* E) & \text{on } \partial^* E \cap \partial\Omega \end{cases},$$

which clearly satisfies

$$\begin{aligned} \|\operatorname{Tr}^e(F, \partial^* E)\|_{L^\infty(\partial^* E; \mathcal{H}^{n-1})} &\leq \max\{\|\hat{F}\|_{L^\infty(\mathbb{R}^n \setminus E; \mathbb{R}^n)}, \|\hat{F}\|_{L^\infty(E; \mathbb{R}^n)}\} = \|\hat{F}\|_{L^\infty(\mathbb{R}^n; \mathbb{R}^n)} \\ &= \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \end{aligned}$$

Finally, we obtain (2.19) if we choose $E = \Omega$ in (2.17). \square

Remark 2.10. We point out that in Theorem 2.9 we can equivalently assume that E has finite perimeter only in Ω . Indeed, if $E \subset\subset \Omega$, then $P(E) = P(E, \Omega)$. If instead Ω is weakly regular, then there exists a family of bounded open sets with smooth boundary $(\Omega_k)_{k \in \mathbb{N}}$ such that

$$\Omega_k \subset \Omega_{k+1}, \quad \bigcup_{k=0}^{+\infty} \Omega_k = \Omega \quad \text{and} \quad P(\Omega_k) \rightarrow P(\Omega)$$

(see [28, Theorem 1.1] and the subsequent discussion). Hence, given $E \subset \Omega$ with $P(E, \Omega) < +\infty$, it is clear that

$$P(E \cap \Omega_k) = P(E \cap \Omega_k, \Omega) \leq P(E, \Omega) + P(\Omega_k, \Omega) = P(E, \Omega) + P(\Omega_k)$$

Thus, by the lower semicontinuity of the perimeter we get

$$P(E) \leq \liminf_{k \rightarrow +\infty} P(E \cap \Omega_k) \leq P(E, \Omega) + \lim_{k \rightarrow +\infty} P(\Omega_k) = P(E, \Omega) + P(\Omega).$$

Since clearly $P(E) \geq P(E, \Omega)$, we conclude that $E \subset \Omega$ has finite perimeter in Ω if and only if it has finite perimeter in \mathbb{R}^n .

Exploiting the Leibniz rule given by Theorem 2.6, we are able to provide an integration by parts formula up to the boundary of a Lipschitz domain for the divergence of a product between a divergence-measure field and a suitable scalar function of bounded variation. Although similar results are already present in the literature (see for instance [3–5, 7, 9, 12, 18]), to the best of our knowledge it has never been formulated with this level of generality before.

Theorem 2.11. *Let Ω be an open bounded set with Lipschitz boundary. Let $F \in \mathcal{DM}^\infty(\Omega)$ and $u \in BV(\Omega)$ be such that $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$ for some Borel function $\lambda : \Omega \rightarrow [0, 1]$. Then we have*

$$\int_{\Omega} u^\lambda d \operatorname{div} F + (F, Du)_\lambda(\Omega) = \operatorname{div}(uF)(\Omega) = - \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(u) \operatorname{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1}. \quad (2.22)$$

Proof. We notice that Ω is a weakly regular set. Hence, arguing as in the proof of Theorem 2.9, we consider the zero extension \hat{F} of F to \mathbb{R}^n . In addition, by [2, Theorem 3.87] we know that $\operatorname{Tr}_{\partial\Omega}(u) \in L^1(\partial\Omega; \mathcal{H}^{n-1})$ and that the zero extension \hat{u} of u to \mathbb{R}^n satisfies $\hat{u} \in BV(\mathbb{R}^n)$ and $D\hat{u} = Du$ on Ω . For $N > 0$, we consider the truncation $T_N(\hat{u})$, which clearly satisfies $T_N(\hat{u}) = \widehat{T_N(u)}$. Thanks to [11, Proposition 3.1], we know that

$$\operatorname{Tr}^i(T_N(\hat{u})\hat{F}, \partial\Omega) = \operatorname{Tr}_{\partial\Omega}(T_N(u)) \operatorname{Tr}^i(\hat{F}, \partial\Omega) \quad \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega.$$

By applying (2.17) to the set $E = \Omega$ in the domain \mathbb{R}^n , we get

$$\operatorname{div}(\widehat{T_N(u)\hat{F}})(\Omega^1) = - \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(T_N(u)) \operatorname{Tr}^i(\hat{F}, \partial\Omega) d\mathcal{H}^{n-1}.$$

However, Ω is weakly regular, which implies that $\mathcal{H}^{n-1}(\Omega^1 \setminus \Omega) = 0$ (see [9, Corollary 5.5]), and so

$$\operatorname{div}(\widehat{T_N(u)\hat{F}})(\Omega^1) = \operatorname{div}(\widehat{T_N(u)\hat{F}})(\Omega) = \operatorname{div}(T_N(u)F)(\Omega),$$

since $|\operatorname{div}(\widehat{T_N(u)\hat{F}})| \ll \mathcal{H}^{n-1}$ and $\widehat{T_N(u)\hat{F}} = T_N(u)F$ on Ω . Hence, by (2.15), we get

$$\int_{\Omega} T_N(u)^\lambda d \operatorname{div} F + (F, DT_N(u))_\lambda(\Omega) = \operatorname{div}(T_N(u)F)(\Omega) = - \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(T_N(u)) \operatorname{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1}, \quad (2.23)$$

where we set $\operatorname{Tr}^i(F, \partial\Omega) := \operatorname{Tr}^i(\hat{F}, \partial\Omega)$, as in the proof of Theorem 2.9. Since $T_N(u) \rightarrow u$ in $BV(\Omega)$, and therefore in $BV(\Omega)$ -strict as $N \rightarrow +\infty$, [2, Theorem 3.88] implies that $\operatorname{Tr}_{\partial\Omega}(T_N(u)) \rightarrow \operatorname{Tr}_{\partial\Omega}(u)$ in $L^1(\partial\Omega; \mathcal{H}^{n-1})$ as $N \rightarrow +\infty$. As for the left hand side, combining Proposition 2.1 and Proposition 2.7, we exploit Lebesgue's dominated convergence theorem to obtain $T_N(u)^\lambda \rightarrow u^\lambda$ in $L^1(\Omega; |\operatorname{div} F|)$ as $N \rightarrow +\infty$. Then, we know that $(F, DT_N(u))_\lambda \rightarrow (F, Du)_\lambda$ in $\mathcal{M}(\Omega)$ by [14, Remark 4.5]. Let $\Omega_\delta = \{x \in \Omega : \operatorname{dist}(x, \partial\Omega) > \delta\}$ for $\delta > 0$ such

that $\Omega_\delta \neq \emptyset$. By (2.16), it is clear that the sequence of measures $(|(F, DT_N(u))_\lambda|)_N$ is uniformly bounded, so that there exists some measure $\gamma \geq |(F, Du)_\lambda|$ such that $|(F, DT_N(u))_\lambda| \rightarrow \gamma$ in $\mathcal{M}(\Omega)$, up to a subsequence. Due to the non-concentration property of Radon measures, we know that $\gamma(\partial\Omega_\delta) = 0$ for \mathcal{L}^1 -a.e. $\delta > 0$. We fix a sequence $(\delta_k)_{k \in \mathbb{N}}, \delta_k \rightarrow 0^+$, of such good values of $\delta > 0$. Thus, by [2, Proposition 1.62], possibly up to a subsequence we get

$$(F, DT_N(u))_\lambda(\Omega_{\delta_k}) \rightarrow (F, Du)_\lambda(\Omega_{\delta_k}) \quad \text{as } N \rightarrow +\infty. \quad (2.24)$$

On the other hand, $|DT_N(u)| \leq |Du|$ and (2.16) yield

$$|(F, DT_N(u))_\lambda|(\Omega \setminus \Omega_{\delta_k}) \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} |Du|(\Omega \setminus \Omega_{\delta_k}) \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

This means that, for all $\varepsilon > 0$, there exists $k_0 \in \mathbb{N}$ such that $|(F, DT_N(u))_\lambda|(\Omega \setminus \Omega_{\delta_k}) < \varepsilon$ for all $N > 0$ and $k \geq k_0$. Thus, thanks to (2.24) we obtain

$$\begin{aligned} \limsup_{N \rightarrow +\infty} |(F, DT_N(u))_\lambda(\Omega) - (F, Du)_\lambda(\Omega)| &\leq \lim_{N \rightarrow +\infty} |(F, DT_N(u))_\lambda(\Omega_{\delta_k}) - (F, Du)_\lambda(\Omega_{\delta_k})| \\ &\quad + \limsup_{N \rightarrow +\infty} |(F, DT_N(u))_\lambda|(\Omega \setminus \Omega_{\delta_k}) + |(F, Du)_\lambda|(\Omega \setminus \Omega_{\delta_k}) \\ &\leq 2\varepsilon, \end{aligned}$$

which, since ε is arbitrary, implies

$$\lim_{N \rightarrow +\infty} (F, DT_N(u))_\lambda(\Omega) = (F, Du)_\lambda(\Omega).$$

All in all, we pass to the limit as $N \rightarrow +\infty$ in (2.23), finally obtaining (2.22). \square

2.3. Measures in the dual of BV . We recall the definition of the dual of the space BV .

Definition 2.12. *We denote by $BV(\Omega)^*$ the dual of the space $BV(\Omega)$; that is, the space of linear functionals $\mathfrak{T} : BV(\Omega) \rightarrow \mathbb{R}$ for which there exists a constant $C > 0$ such that*

$$|\mathfrak{T}(u)| \leq C \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega).$$

It is well known that there are some elements in the dual of BV whose action can be represented as the integration of a suitable representative of the BV function against a Radon measure (see for instance [25]). Indeed, given $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$, the linear functional

$$\mathfrak{T}_\mu(u) = \int_{\Omega} u^* d\mu \quad \text{for } u \in BV(\Omega) \cap L^\infty(\Omega)$$

is well defined, although not necessarily continuous. We point out that the assumption $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is necessary, since otherwise $u^*(x)$ might not be well defined for $|\mu|$ -a.e. $x \in \Omega$, and that the choice of the precise representative u^* ensures the linearity of \mathfrak{T}_μ , due to (2.5). Therefore, we choose the following notation:

given $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$, we say that $\mu \in BV(\Omega)^$ if there exists $\mathfrak{T} \in BV(\Omega)^*$ such that*

$$\mathfrak{T}(u) = \mathfrak{T}_\mu(u) \quad \text{for all } u \in BV(\Omega) \cap L^\infty(\Omega).$$

Under the additional assumption that Ω is an open bounded set with Lipschitz boundary, it was proved in [25, Theorem 8.2] that \mathfrak{T}_μ is continuous on $BV(\Omega) \cap L^\infty(\Omega)$ with respect to the BV -norm if and only if there exists $C > 0$ such that

$$|\mu(U \cap \Omega)| \leq C P(U) \quad \text{for all open sets } U \subset \mathbb{R}^n \text{ with smooth boundary.}$$

We further explore this and similar results in Section 4.

Given a pair $(\mu, \lambda) \in \mathcal{M}_{\mathcal{H}}(\Omega) \times \mathcal{B}_b(\Omega)$, it is also possible to define

$$\mathfrak{T}_{\mu, \lambda}(u) = \int_{\Omega} u^{\lambda} d\mu \quad \text{for } u \in BV(\Omega) \cap L^{\infty}(\Omega). \quad (2.25)$$

However, unless $\lambda(x) \equiv \frac{1}{2}$ for $|\mu|$ -a.e. $x \in \Omega$ or $|\mu|(\Sigma) = 0$ for all \mathcal{H}^{n-1} -rectifiable sets Σ , this mapping is not linear. Indeed, arguing analogously as in [14, Remark 4.6], we notice that for any $u \in BV(\Omega) \cap L^{\infty}(\Omega)$ we have

$$\mathfrak{T}_{\mu, \lambda}(u) + \mathfrak{T}_{\mu, \lambda}(-u) = \int_{\Omega} (u^{\lambda} + (-u)^{\lambda}) d\mu = \int_{\Omega} (2\lambda - 1)(u^{+} - u^{-}) d\mu.$$

Nevertheless, it is worth remarking that for a relevant class of measures the functional $\mathfrak{T}_{\mu, \lambda}$ can be extended to a continuous functional defined on the whole space $BV(\Omega)$, see Definition 4.6 and Lemma 4.17.

3. λ -APPROXIMATION

It is natural to expect that many properties of the λ -pairing $(F, Du)_{\lambda}$ can be obtained as direct extensions of properties satisfied by the pairing when u is a smooth function. The goal of this section is to provide the required approximation result, Theorem 3.4, which in turn is based on Theorem 3.2, a finer version of Anzellotti-Giaquinta's approximation that is specifically designed to enforce also the \mathcal{H}^{n-1} -a.e. approximation of the λ -representative u^{λ} .

We start by stating a first refinement of Anzellotti-Giaquinta approximation theorem for BV functions involving also the \mathcal{H}^{n-1} -a.e. pointwise convergence and the strict convergence with respect to the area functional. As customary, we say that any radial function $\rho \in C_c^{\infty}(B_1(0))$ such that $\rho \geq 0$ and $\int_{B_1(0)} \rho dx = 1$ is a *standard mollifier*. For all $\varepsilon > 0$ we set $\rho_{\varepsilon}(x) = \varepsilon^{-n} \rho(\frac{x}{\varepsilon})$, and we recall that, for all $u \in BV(\Omega)$, by [2, Corollary 3.80] we have

$$\lim_{\varepsilon \rightarrow 0^+} (\rho_{\varepsilon} * u)(x) = u^{*}(x) \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in \Omega. \quad (3.1)$$

Theorem 3.1. *Let $u \in BV(\Omega)$. Then there exists $(u_{\varepsilon})_{\varepsilon > 0} \subset BV(\Omega) \cap C^{\infty}(\Omega)$ such that*

- (1) $u_{\varepsilon} \rightarrow u$ in $L^1(\Omega)$ as $\varepsilon \rightarrow 0$,
- (2) $|Du_{\varepsilon}|(\Omega) \leq |Du|(\Omega) + 4\varepsilon$ for all $\varepsilon > 0$, and so $\lim_{\varepsilon \rightarrow 0} |Du_{\varepsilon}|(\Omega) = |Du|(\Omega)$,
- (3) $\sqrt{1 + |Du_{\varepsilon}|^2}(\Omega) \leq \sqrt{1 + |Du|^2}(\Omega) + 4\varepsilon$ for all $\varepsilon > 0$, and so $\lim_{\varepsilon \rightarrow 0} \sqrt{1 + |Du_{\varepsilon}|^2}(\Omega) = \sqrt{1 + |Du|^2}(\Omega)$,
- (4) $u_{\varepsilon}(x) \rightarrow u^{*}(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$ as $\varepsilon \rightarrow 0$,
- (5) $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)} \leq (1 + \varepsilon)\|u\|_{L^{\infty}(\Omega)}$ for all $\varepsilon > 0$,
- (6) if Ω is an open set with bounded Lipschitz boundary, then $\text{Tr}_{\partial\Omega}(u_{\varepsilon}) = \text{Tr}_{\partial\Omega}(u)$ for all $\varepsilon > 0$.

Proof. The proof is based on a slight modification of the construction of the approximating sequence in Anzellotti-Giaquinta theorem, for which we refer for instance to [16, Theorem 3, Section 5.2.2].

Fix $\varepsilon > 0$. Given a positive integer m , we set $\Omega_0 = \emptyset$, define for each $k \in \mathbb{N}, k \geq 1$ the sets

$$\Omega_k = \left\{ x \in \Omega : \text{dist}(x, \partial\Omega) > \frac{1}{m+k} \right\} \cap B(0, k+m)$$

and then we choose m such that $|Du|(\Omega \setminus \Omega_1) < \varepsilon$.

We define now $\Sigma_k := \Omega_{k+1} \setminus \overline{\Omega_{k-1}}$. Since $(\Sigma_k)_{k \geq 1}$ is an open cover of Ω , then there exists a partition of unity subordinate to that open cover; that is, a sequence of functions $(\zeta_k)_{k \geq 1}$ such that:

- (1) $\zeta_k \in C_c^\infty(\Sigma_k)$;
- (2) $0 \leq \zeta_k \leq 1$;
- (3) $\sum_{k=1}^{+\infty} \zeta_k = 1$ on Ω .

Next, we take a standard mollifier ρ and for all $k \in \mathbb{N}, k \geq 1$ we choose $\delta_k = \delta_k(\varepsilon) > 0$ small enough such that the following conditions hold

$$\text{supp}(\rho_{\delta_k} * \zeta_k) \subset \Sigma_k, \quad (3.2)$$

$$\|\rho_{\delta_k} * (u\zeta_k) - u\zeta_k\|_{L^1(\Omega)} < \frac{\varepsilon}{2^k}, \quad (3.3)$$

$$\|\rho_{\delta_k} * (u\nabla\zeta_k) - u\nabla\zeta_k\|_{L^1(\Omega; \mathbb{R}^n)} < \frac{\varepsilon}{2^k}, \quad (3.4)$$

$$\|\rho_{\delta_k} * \zeta_k - \zeta_k\|_{L^\infty(\Omega)} < \frac{\varepsilon}{2^k}. \quad (3.5)$$

Then we define $u_\varepsilon := \sum_{k=1}^{+\infty} \rho_{\delta_k} * (u\zeta_k)$. We notice that, for any fixed $x \in \Omega$, there exists a unique $k = k(x) \geq 1$ such that $x \in \Sigma_k \cap \Sigma_{k+1}$, so that, by (3.2),

$$u_\varepsilon(x) = \sum_{j=0}^1 (\rho_{\delta_{k+j}} * (u\zeta_{k+j}))(x). \quad (3.6)$$

Hence, $u_\varepsilon \in C^\infty(\Omega)$, since locally there are at most two nonzero terms in the sum. The proof of points (1) and (2) follows in a standard way from (3.2), (3.3) and (3.4) (see for instance [16, Theorem 3, Section 5.2.2] and [19, Theorem 1.17]). Arguing in a similar way, for any $(\varphi, \eta) \in C_c^1(\Omega; \mathbb{R}^n \times \mathbb{R})$ such that $\|(\varphi, \eta)\|_{L^\infty(\Omega; \mathbb{R}^n \times \mathbb{R})} \leq 1$ we obtain

$$\begin{aligned} \int_{\Omega} \eta + u_\varepsilon \operatorname{div} \varphi \, dx &= \int_{\Omega} \eta + u \operatorname{div}(\zeta_1(\rho_{\delta_1} * \varphi)) \, dx + \sum_{k=2}^{+\infty} \int_{\Omega} u \operatorname{div}(\zeta_k(\rho_{\delta_k} * \varphi)) \, dx + \\ &\quad - \sum_{k=1}^{+\infty} \int_{\Omega} \varphi \cdot (\rho_{\delta_k} * (u\nabla\zeta_k) - u\nabla\zeta_k) \, dx \\ &\leq \sqrt{1 + |Du|^2}(\Omega) + \sum_{k=2}^{+\infty} |Du|(\Sigma_k) + \sum_{k=1}^{+\infty} \|\rho_{\delta_k} * (u\nabla\zeta_k) - u\nabla\zeta_k\|_{L^1(\Omega; \mathbb{R}^n)} \\ &\leq \sqrt{1 + |Du|^2}(\Omega) + 2|Du|(\Omega \setminus \Omega_1) + \sum_{k=1}^{+\infty} \frac{\varepsilon}{2^k} \\ &\leq \sqrt{1 + |Du|^2}(\Omega) + 3\varepsilon. \end{aligned}$$

Together with the lower semicontinuity of the area functional with respect to the L^1 convergence, this implies point (3). Then, since ζ_k is continuous for every k and $u \in BV(\Omega)$, by (3.1) we have

$$\lim_{\varepsilon \rightarrow 0^+} (\rho_{\delta_k} * (u\zeta_k))(x) = \zeta_k(x)u^*(x) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in \Omega,$$

where the \mathcal{H}^{n-1} -negligible set of points depends only on u . Hence, for \mathcal{H}^{n-1} -a.e. $x \in \Omega$, by (3.6) we get

$$\lim_{\varepsilon \rightarrow 0^+} u_\varepsilon(x) = \lim_{\varepsilon \rightarrow 0^+} \sum_{j=0}^1 (\rho_{\delta_{k+j}} * (u\zeta_{k+j}))(x) = u^*(x) \sum_{j=0}^1 \zeta_{k+j}(x) = u^*(x).$$

As for point (5), we can assume without loss of generality that $u \in L^\infty(\Omega)$. For any $x \in \Omega$ there exists a unique $k \geq 1$ such that $\sum_{j=0}^1 \zeta_{k+j}(x) = 1$ and (3.5) holds true. Therefore we apply (3.6) to get

$$\begin{aligned} |u_\varepsilon(x)| &\leq \|u\|_{L^\infty(\Omega)} \sum_{j=0}^1 (\rho_{\delta_{k+j}} * \zeta_{k+j})(x) \leq \|u\|_{L^\infty(\Omega)} \left(1 + \sum_{j=0}^1 |(\rho_{\delta_{k+j}} * \zeta_{k+j})(x) - \zeta_{k+j}(x)| \right) \\ &\leq (1 + \varepsilon) \|u\|_{L^\infty(\Omega)}. \end{aligned}$$

Finally, if Ω is an open set with bounded Lipschitz boundary, then the trace operator is well defined and, thanks to [2, Theorem 3.87], it satisfies

$$\lim_{\rho \rightarrow 0} \frac{1}{\rho^n} \int_{\Omega \cap B_\rho(x)} |u(y) - \text{Tr}_{\partial\Omega}(u)(x)| dy = 0 \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in \partial\Omega.$$

Hence, arguing as in [19, Remarks 1.18 and 2.12], we conclude that (6) holds true. \square

We exploit Theorem 3.1 in order to obtain the following finer λ -approximation theorem, which is then applied to the approximation of λ -pairings (Theorem 3.4 below) and is exploited at length in Section 4. In addition, it represents one of the key tools in the proofs of the main results of [20].

Theorem 3.2 (λ -approximation). *Let $\lambda : \Omega \rightarrow [0, 1]$ be a given Borel function. Then for every $u \in BV(\Omega)$ there exists a sequence $(u_k^\lambda)_{k \in \mathbb{N}} \subset C^\infty(\Omega) \cap BV(\Omega)$ such that we have the following:*

- (1) $u_k^\lambda \rightarrow u$ in $BV(\Omega)$ -strict as $k \rightarrow +\infty$,
- (2) $u_k^\lambda(x) \rightarrow u^\lambda(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$ as $k \rightarrow +\infty$,
- (3) $\lim_{k \rightarrow +\infty} \sqrt{1 + |Du_k^\lambda|^2(\Omega)} = \sqrt{1 + |Du|^2(\Omega)}$,
- (4) $\|u_k^\lambda\|_{L^\infty(\Omega)} \leq (1 + \frac{1}{k}) \|u\|_{L^\infty(\Omega)}$ for all $k \in \mathbb{N}$,
- (5) if Ω is an open set with bounded Lipschitz boundary, then $\text{Tr}_{\partial\Omega}(u_k^\lambda)(x) = \text{Tr}_{\partial\Omega}(u)(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$ and all $k \in \mathbb{N}$.

Proof. We first prove the theorem under the assumption $\|u\|_{L^\infty(\Omega)} < +\infty$. It is enough to construct a sequence $(u_k^\lambda)_k$ of smooth functions that converge to u in $BV(\Omega)$ -strict, to $u^* = \tilde{u} = u^\lambda$ \mathcal{H}^{n-1} -a.e. on $\Omega \setminus J_u$, and to u^λ locally in measure with respect to $\mu = \mathcal{H}^{n-1} \llcorner J_u$. Then, we conclude by extracting a suitable subsequence that converges μ -almost everywhere to u^λ , hence \mathcal{H}^{n-1} -a.e. on Ω .

Let $(u_\varepsilon)_{\varepsilon > 0}$ be the smooth approximation of u given by Theorem 3.1. We choose a sequence $\varepsilon = \frac{1}{k}$, and we set $u_k := u_{\frac{1}{k}}$ for simplicity. Since $u_k(x) \rightarrow u^*(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$, the idea

is to define u_k^λ as a suitable perturbation of u_k near the jump set J_u , and then show that u_k^λ satisfies the convergence in measure stated above. The proof will be split into some steps.

Step one: local construction and estimates.

We fix $\varepsilon > 0$ and a Borel set $A \subset\subset \Omega$, then we consider the set

$$S = S(u, A, \varepsilon) = \{x \in A \cap J_u : u^+(x) - u^-(x) > \varepsilon\}.$$

Notice that $\mathcal{H}^{n-1}(S) < +\infty$, hence $\mu(S) < +\infty$. Up to choosing the parameter m in the proof of Theorem 3.1 to be large enough, we can assume that $u_k = u * \rho_k$ on A , where $\rho_k(|x|) = k^n \rho(k|x|)$ is a standard mollifier with support in $B_{1/k}$, for k sufficiently large. For \mathcal{H}^{n-1} -a.e. $x \in S$, we define the *blow-up of u at x* as the step function $u_{x,\infty} : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$u_{x,\infty}(y) = \begin{cases} u^+(x) & \text{if } (y-x) \cdot \nu_u(x) \geq 0, \\ u^-(x) & \text{otherwise.} \end{cases}$$

Therefore we have

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{B_r(x)} |u(y) - u_{x,\infty}(y)| dy = 0.$$

For \mathcal{H}^{n-1} -a.e. $x \in S$ we define the Borel function $\tau = \tau_k(x) \in [-1, 1]$ as the unique implicit solution of

$$u_{x,\infty} * \rho_k(x + \tau \nu_u(x)/k) = u^\lambda(x). \quad (3.7)$$

Note that $\tau_k(x)$ can be written as the composition of a continuous function, depending only on the mollifier ρ_k , with the function $\lambda(x)$.

Fix $\delta > 0$ to be later chosen, and consider the following perturbation of u_k inside a ball $B = B_r(x)$ centered at $x \in S$:

$$u_k^{\tau,B}(y) := \phi_{r,\delta}(y-x) u_k(y + \tau \nu_u(x)/k) + (1 - \phi_{r,\delta}(y-x)) u_k(y), \quad (3.8)$$

where $\tau \in [-1, 1]$ and $\phi_{r,\delta}(z) \in [0, 1]$ is a radially symmetric cut-off function of class C^∞ , with compact support in $B_r(0)$ and such that $\phi_{r,\delta}(z) = 1$ if $|z| < (1-\delta)r$ and $\|\nabla \phi_{r,\delta}\|_\infty < \frac{2}{\delta r}$. Clearly, $u_k^{\tau,B}$ is a smooth function obtained by locally ‘‘gluing’’ a suitable translation of u_k with u_k itself, it coincides with u_k outside B , and satisfies

$$u_k^{\tau,B}(x) = u_k(x + \tau \nu_u(x)/k) = u * \rho_k(x + \tau \nu_u(x)/k). \quad (3.9)$$

Its gradient is given by

$$\nabla u_k^{\tau,B}(y) = \nabla u_k(y) + \nabla \phi_{r,\delta}(y-x) (u_k(y + v_k) - u_k(y)) + \phi_{r,\delta}(y-x) (\nabla u_k(y + v_k) - \nabla u_k(y)).$$

where we have set $v_k = \frac{\tau}{k} \nu_u(x)$. For the sake of simplicity, we also set $B_\delta = B_{(1-\delta)r}(x)$, and $C_\delta = B \setminus B_\delta$. We find that

$$\begin{aligned} \int_B |\nabla u_k^{\tau,B}| &\leq \int_{B_\delta} |\nabla u_k(y + v_k)| dy + \int_{C_\delta} |\nabla u_k| + \frac{2}{\delta r} \int_{C_\delta} |u_k(y + v_k) - u_k(y)| dy \\ &\quad + \int_{C_\delta} |\nabla u_k(y + v_k) - \nabla u_k(y)| dy. \end{aligned} \quad (3.10)$$

Then, owing to the fact that $|Du_k|(A) \leq |Du|(A + B_{1/k}(0))$, the first term in the right-hand side of (3.10) can be estimated as follows:

$$\int_{B_\delta} |\nabla u_k(y + v_k)| dy \leq |Du|(B_{\delta,k}), \quad (3.11)$$

where we have set

$$B_{\delta,k} = B_\delta + B_{2/k}(0) = B_{(1-\delta)r+2/k}(x).$$

On observing that $|v_k| \leq k^{-1}$, the third term in the right-hand side of (3.10) can be estimated as follows:

$$\frac{2}{\delta r} \int_{C_\delta} |u_k(y + v_k) - u_k(y)| dy \leq \frac{2}{k\delta r} |Du|(C_{\delta,k}), \quad (3.12)$$

where $C_{\delta,k} = C_\delta + B_{2/k}(0)$. Concerning the fourth term in the right-hand side of (3.10), we set $\rho_{k,x}(z) = \rho_k(z - v_k)$ and denote by $u_{k,z}$ the average of u on $B_{2/k}(z)$, so that we obtain

$$\begin{aligned} \int_{C_\delta} |\nabla u_k(y + v_k) - \nabla u_k(y)| dy &\leq \int_{C_\delta} \left| \int_{\Omega} u(y) \left(\nabla \rho_{k,x}(z - y) - \nabla \rho_k(z - y) \right) dy \right| dz \\ &= \int_{C_\delta} \left| \int_{B_{2/k}(z)} (u(y) - u_{k,z}) \left(\nabla \rho_{k,x}(z - y) - \nabla \rho_k(z - y) \right) dy \right| dz \\ &\leq \frac{\|\nabla^2 \rho_k\|_\infty}{k} \int_{C_\delta} \int_{B_{2/k}(z)} |u(y) - u_{k,z}| dy dz \\ &\leq Ck^{n+1} \int_{C_\delta} \int_{B_{2/k}(z)} |u(y) - u_{k,z}| dy dz \\ &\leq Ck^n \int_{C_\delta} |Du|(B_{2/k}(z)) dz, \end{aligned}$$

where we have used the fact that $\|\nabla^2 \rho_k\|_\infty \leq Ck^{n+2}$ and, in the last step, the Poincarè-Wirtinger inequality on $B_{2/k}(z)$ (note that in this last estimate, as well as in the next ones, we will denote by C a dimensional constant that can possibly change from one line to another). We can push further the estimate by noticing that

$$\begin{aligned} \int_{C_\delta} |Du|(B_{2/k}(z)) dz &= \int_{C_\delta} \int_{B_{2/k}(z)} d|Du|(y) dz \\ &= \int_{C_\delta + B_{2/k}} \int_{C_\delta \cap B_{2/k}(y)} dz d|Du|(y) \\ &= \int_{C_\delta + B_{2/k}} |C_\delta \cap B_{2/k}(y)| d|Du|(y) \\ &\leq \frac{C}{k^n} |Du|(C_\delta + B_{2/k}) = \frac{C}{k^n} |Du|(C_{\delta,k}). \end{aligned}$$

This leads us to

$$\int_{C_\delta} |\nabla u_k(y + v_k) - \nabla u_k(y)| dy \leq C |Du|(C_{\delta,k}). \quad (3.13)$$

Consequently, if we plug (3.11), (3.12), (3.13) into (3.10), we obtain

$$\int_B |\nabla u_k^{\tau, B}| \leq |Du|(B_{\delta, k}) + \left(C + \frac{2}{k\delta r}\right) |Du|(C_{\delta, k}), \quad (3.14)$$

where C is a constant only depending on the dimension n . Note that as soon as $k \geq (r\delta)^{-1}$ the inequality improves to

$$\int_B |\nabla u_k^{\tau, B}| \leq |Du|(B_{\delta, k}) + C|Du|(C_{\delta, k}),$$

where C is another dimensional constant.

Step two: from local to global.

Here we show how to use the local construction and the estimate (3.14) provided by Step one, to define a sequence of approximations that behaves well on a compact subset of S with large measure. To this aim we must first guarantee the continuity and the uniformity of some quantities that appear in Step one, and then apply the appropriate covering theorem.

By Lusin and Egoroff Theorems, for every $\eta \in (0, 1)$ we can find a compact set $S_\eta \subset S$ satisfying the following properties:

- (i) $\mathcal{H}^{n-1}(S_\eta) \geq (1 - \eta)\mathcal{H}^{n-1}(S)$;
- (ii) the functions λ, ν_u, u^+, u^- restricted to S_η are continuous;
- (iii) if we set

$$\Delta_\eta(r) := \sup_{x \in S_\eta} \frac{1}{r^n} \int_{B_r(x)} |u(y) - u_{x, \infty}(y)| dy,$$

we have $\lim_{r \rightarrow 0} \Delta_\eta(r) = 0$;

- (iv) there exists $0 < r_{\delta, \eta} < 1$ such that

$$|Du|(C_{\delta, k}(x, r)) \leq 6(n-1)\delta |Du|(B_r(x))$$

for all $0 < r < r_{\delta, \eta}$, $k > (\delta r)^{-1}$, and $x \in S_\eta$, where $C_{\delta, k}(x, r)$ is defined as in the previous step (here the dependence upon x and r is explicitly written for the sake of clarity).

We remark that showing (i), (ii), and (iii) is standard. As for (iv), we notice that the asymptotic behaviour of $|Du|(C_{\delta, k})$ for $k > (\delta r)^{-1}$ and as $r \rightarrow 0$ is

$$\begin{aligned} |Du|(C_{\delta, k}) &= |D^d u|(C_{\delta, k}) + |D^j u|(C_{\delta, k}) \\ &= (1 + o(1))\omega_{n-1}|u^+(x) - u^-(x)| \left[(r + 2/k)^{n-1} - (r(1 - \delta) - 2/k)^{n-1} \right] \\ &= (1 + o(1))\omega_{n-1}|u^+(x) - u^-(x)| r^{n-1} \left[(1 + 2/(kr))^{n-1} - ((1 - \delta) - 2/(kr))^{n-1} \right] \\ &= (1 + o(1))(n-1)\omega_{n-1}|u^+(x) - u^-(x)| r^{n-1} \delta [4/(k\delta r) + 1] \\ &\leq 6(n-1)\delta |Du|(B_r(x)), \end{aligned} \quad (3.15)$$

Which gives (iv) at once.

Fix now an open set U containing S_η , then consider the family \mathcal{F} of balls centered in S_η and contained in U , with radius r so small that $\Delta_\eta(r) < \varepsilon$. By Vitali-Besicovitch Covering Theorem, we can find a finite and mutually disjoint family of balls $\{B^i\}_{i=1}^N$, with $B^i = B_{r_i}(x_i) \in \mathcal{F}$,

$0 < r_i < r_{\delta, \eta}$, and N depending on δ and η , such that

$$\mathcal{H}^{n-1} \left(S_\eta \cap \bigcup_{i=1}^N B^i \right) \geq (1 - \eta) \mathcal{H}^{n-1}(S_\eta) \geq (1 - \eta)^2 \mathcal{H}^{n-1}(S),$$

where we have used property (i) in the last inequality.

Step three: total variation estimate.

We set $r_0 = \min\{r_i : i = 1, \dots, N\}$ and consider the sequence u_k^λ constructed by replacing u_k with $u_k^{\tau, B}$ inside each ball $B = B^i$, as obtained and described in the previous steps. Hereafter we show that the total variation of Du_k^λ is controlled by that of Du , up to an error that goes to zero as $\delta \rightarrow 0$ (hence as $k \rightarrow \infty$). Indeed, if we set $C_\delta^i = B^i \setminus B_\delta^i$, we can assume, up to a small perturbation of r_i , that $|Du|(\partial C_\delta^i) = 0$ for all i , hence thanks to the estimate (3.14) we obtain

$$\begin{aligned} |Du_k^\lambda|(\Omega) &\leq |Du_k| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right) + \sum_{i=1}^N |Du_k^\lambda|(B^i) \\ &\leq |Du_k| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right) + \sum_{i=1}^N \left(|Du|(B_{\delta, k}^i) + \left(C + \frac{2}{k\delta r_i} \right) |Du|(C_{\delta, k}^i) \right). \end{aligned} \quad (3.16)$$

Let now assume $k \geq (\delta r_0)^{-1}$. Thanks to (iv) and recalling that the balls B^i are mutually disjoint, we have

$$\sum_{i=1}^N \left(C + \frac{2}{k\delta r_i} \right) |Du|(C_{\delta, k}^i) \leq (C + 2) \sum_{i=1}^N |Du|(C_{\delta, k}^i) \leq \tilde{C}\delta \sum_{i=1}^N |Du|(B^i) \leq \tilde{C}\delta |Du|(\Omega). \quad (3.17)$$

Thanks to Theorem 3.1, we have the strict convergence of u_k to u on Ω , hence by the lower semicontinuity of $|Du_k|$ restricted to the open set $\bigcup_{i=1}^N B^i$ we obtain

$$\limsup_{k \rightarrow \infty} |Du_k| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right) = |Du|(\Omega) - \liminf_{k \rightarrow \infty} |Du_k| \left(\bigcup_{i=1}^N B^i \right) \leq |Du| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right),$$

hence by selecting k large enough we can enforce

$$|Du_k| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right) \leq |Du| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right) + \delta r_0. \quad (3.18)$$

By combining (3.16), (3.17), and (3.18) we finally get

$$\begin{aligned} |Du_k^\lambda|(\Omega) &\leq |Du| \left(\Omega \setminus \bigcup_{i=1}^N B^i \right) + \delta r_0 + \sum_{i=1}^N |Du|(B^i) + \tilde{C}\delta |Du|(\Omega) \\ &= (1 + \tilde{C}\delta) |Du|(\Omega). \end{aligned} \quad (3.19)$$

This shows that the total variation of Du_k^λ is arbitrarily close to the total variation of Du , up to choosing k large enough. This will eventually lead to the BV -strict approximation property (see Step five). Arguing in a similar way, we can obtain an analogous upper bound for the area functional. This is due to the coincidence of the singular parts of the area and the total

variation functionals (see (2.11)) hence the previous construction leads to an estimate like (3.19) with $\sqrt{1 + |Du_k^\lambda|^2}$ and $\sqrt{1 + |Du|^\lambda}$ replacing the total variation functionals.

Step four: pointwise closeness of u_k^λ to u^λ on S_η .

Let us fix $B^i = B_{r_i}(x_i)$ for $i = 1, \dots, N$, and choose $x \in B_\delta^i \cap S_\eta$, where $B_\delta^i = B_{(1-\delta)r_i}(x_i)$. We would like to prove that $u_k^\lambda(x)$ is close to $u^\lambda(x)$. We have

$$\begin{aligned} |u_k^\lambda(x) - u^\lambda(x)| &\leq |u_k^\lambda(x) - u^\lambda(x_i)| + |u^\lambda(x_i) - u^\lambda(x)| \leq |u_k^\lambda(x) - u_{x,\infty} * \rho_k(x + v_k(x))| + \\ &\quad + |u_{x,\infty} * \rho_k(x + v_k(x)) - u_{x_i,\infty} * \rho_k(x_i + v_k(x_i))| + \\ &\quad + |u_{x_i,\infty} * \rho_k(x_i + v_k(x_i)) - u^\lambda(x_i)| + |u^\lambda(x_i) - u^\lambda(x)| \\ &=: A_1 + A_2 + A_3 + A_4. \end{aligned}$$

First of all, by (3.9) and (iii) we obtain

$$\begin{aligned} A_1 &\leq \int_{B_{1/k}(x+v_k(x))} \rho_k(x + v_k(x) - y) |u(y) - u_{x,\infty}(y)| dy \\ &\leq k^n \|\rho\|_\infty \int_{B_{2/k}(x)} |u(y) - u_{x,\infty}(y)| dy \rightarrow 0 \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Then, $A_3 = 0$ by (3.7), recalling that $v_k(y) = \frac{\tau(y)}{k} \nu_u(y)$, while thanks to (ii) the terms A_2 and A_4 are close to 0 if x is close to x_i , which in turn depends on the fact that r_i is taken small enough. This shows that, by choosing r_i small and $k \geq (\delta r_0)^{-1}$ large enough, we can enforce the required pointwise closeness.

Step five: conclusion. Let us fix three positive and infinitesimal sequences $\varepsilon_j, \eta_j, \delta_j$, as well as a monotone sequence $A_j \subset\subset \Omega$ of Borel sets, such that $\bigcup_j A_j = \Omega$. Let U_j be a sequence of open sets as in Step two, satisfying the extra condition $|U_j| < 2^{-j}$ for all j . For any integer $j \geq 1$ we can apply the previous steps with $A = A_j$ (Step one) and $U = U_j$ (Step two), and select from the initial sequence $(u_k)_k$ a suitable subsequence, that we do not relabel, whose elements can be locally perturbed according to the procedure described in Step one. We shall perform this construction iteratively, so that the sequence that we extract at the $(j+1)$ -th stage is also a subsequence of the one obtained at the j -th stage. By diagonal selection we obtain a sequence relabeled as u_k^λ . Owing to Step four, u_k^λ converges to u^λ in \mathcal{H}^{n-1} -measure on the whole jump set J_u as $k \rightarrow +\infty$, while by the choice of U_j we have that u_k^λ converges to u in $L^1(\Omega)$. Hence, by Step three, u_k^λ converges to u in $BV(\Omega)$ -strict, which is point (1) of the statement. Similarly, we deduce point (3). Up to a further extraction of a subsequence, we obtain the pointwise convergence \mathcal{H}^{n-1} -a.e. on Ω . Then, the bound $|u_k^\lambda(x)| \leq (1 + \frac{1}{k}) \|u\|_{L^\infty(\Omega)}$ for all $x \in \Omega$ follows immediately from the definition of u_k^λ in terms of $u_k^{\tau, B}$ and (3.8), thus concluding the proof of the theorem under the assumption $\|u\|_{L^\infty(\Omega)} < +\infty$. To obtain the complete proof, we apply the previous steps to the sequence of truncations $T_m(u)$, and obtain $(T_m(u)_k^\lambda)_k$ for each $m \in \mathbb{N}$. The diagonal sequence $T_k(u)_k^\lambda$ can then be easily shown to satisfy all the required properties. Finally, point (5) can be proved as in Theorem 3.1, given that the sequence $(u_k^\lambda)_k$ is obtained by a local perturbation of the approximating sequence $(u_k)_k$ from Anzellotti-Giaquinta approximation theorem. \square

We state for later use a simple consequence of Theorem 3.2.

Corollary 3.3. *Let $\lambda : \Omega \rightarrow [0, 1]$ be a given Borel function. Let $u \in BV(\Omega)$ and $(u_k^\lambda)_{k \in \mathbb{N}} \subset C^\infty(\Omega)$ be the approximating sequence of Theorem 3.2. Then we have*

$$\sqrt{1 + |Du_k^\lambda|^2} \rightharpoonup \sqrt{1 + |Du|^2} \text{ in } \mathcal{M}(\Omega).$$

Proof. The lower semicontinuity on open sets is an immediate consequence of the fact that $u_k^\lambda \rightarrow u$ in $L^1(\Omega)$, thanks to point (1) of Theorem 3.2; while the upper semicontinuity on compact sets follows from the lower semicontinuity on the open sets and point (3) of Theorem 3.2. \square

We conclude this section with an approximation result for the λ -pairings, which, as a byproduct, allows us to derive a different proof for the estimate (2.16), see Remark 3.6 below. In addition, we provide a useful alternative bound for the λ -pairing in our framework, a particular case of which was proved in [26, Lemma 5.5].

Theorem 3.4. *Let $F \in \mathcal{DM}^\infty(\Omega)$, $u \in BV(\Omega)$ and $\lambda : \Omega \rightarrow [0, 1]$ be a Borel function. If $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$, then there exists a sequence $(u_j)_{j \in \mathbb{N}} \subset C^\infty(\Omega) \cap BV(\Omega) \cap L^\infty(\Omega)$ such that $u_j \rightarrow u$ in $BV(\Omega)$ -strict and*

$$(F \cdot \nabla u_j) \mathcal{L}^n \rightharpoonup (F, Du)_\lambda \text{ in } \mathcal{M}(\Omega). \quad (3.20)$$

If $u \in BV(\Omega) \cap L^\infty(\Omega)$, then $uF \in \mathcal{DM}^\infty(\Omega)$ and (3.20) holds true for the approximating sequence $(u_k^\lambda)_{k \in \mathbb{N}}$ given by Theorem 3.2.

Finally, for all $a \geq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}$ we have

$$|(F, Du)_\lambda| \leq a \sqrt{1 + |Du|^2} - \sqrt{a^2 - |F|^2} \mathcal{L}^n \text{ on } \Omega. \quad (3.21)$$

Proof. Thanks to Theorem 2.6, we know that $\operatorname{div}(uF), (F, Du)_\lambda \in \mathcal{M}(\Omega)$ and that the Leibniz rule (2.15) holds true. In addition, Proposition 2.7 implies that $M[u, \lambda] \in L^1(\Omega; |\operatorname{div} F|)$, given that $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$.

Now, we exploit Theorem 3.2 in order to construct the smooth approximation $T_N(u)_k^\lambda$ of the truncation $T_N(u)$, for all $N > 0$. Thanks to (2.15), for all $k \in \mathbb{N}$ and $N > 0$, we have

$$\operatorname{div}(T_N(u)_k^\lambda F) = T_N(u)_k^\lambda \operatorname{div} F + (F \cdot \nabla T_N(u)_k^\lambda) \mathcal{L}^n \text{ on } \Omega.$$

Hence, Theorem 3.2 implies that

$$\left| \operatorname{div}(T_N(u)_k^\lambda F) \right|(\Omega) \leq \|\varphi\|_{L^\infty(\Omega)} \left(\left(1 + \frac{1}{k}\right) N |\operatorname{div} F|(\Omega) + \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \sup_{k \in \mathbb{N}} \|\nabla T_N(u)_k^\lambda\|_{L^1(\Omega; \mathbb{R}^n)} \right).$$

Therefore, we conclude that $(\operatorname{div}(T_N(u)_k^\lambda F))_{k \in \mathbb{N}}$ is bounded sequence in $\mathcal{M}(\Omega)$ for each fixed $N > 0$, and, since $\operatorname{div}(T_N(u)_k^\lambda F)$ converges to $\operatorname{div}(T_N(u)F)$ in the sense of distributions, we deduce that it also weakly converges in the sense of Radon measures. In addition, Theorem 3.2 and the Lebesgue theorem with respect to the measure $|\operatorname{div} F|$ imply that

$$T_N(u)_k^\lambda \operatorname{div} F \rightharpoonup T_N(u)^\lambda \operatorname{div} F \text{ in } \mathcal{M}(\Omega).$$

All in all, we see that

$$\begin{aligned} (F \cdot \nabla T_N(u)_k^\lambda) \mathcal{L}^n &= \operatorname{div}(T_N(u)_k^\lambda F) - T_N(u)_k^\lambda \operatorname{div} F \\ &\rightharpoonup \operatorname{div}(T_N(u)F) - T_N(u)^\lambda \operatorname{div} F = (F, DT_N(u))_\lambda \text{ as } k \rightarrow +\infty. \end{aligned}$$

Then we notice that (2.15) holds true for $T_N(u)$ and F , so that we get

$$\int_{\Omega} \varphi d(F, DT_N(u))_{\lambda} = - \int_{\Omega} T_N(u) F \cdot \nabla \varphi dx - \int_{\Omega} \varphi (T_N(u))^{\lambda} d \operatorname{div} F \quad (3.22)$$

for all $\varphi \in C_c^1(\Omega)$. In particular, this implies that the family of measures $((F, DT_N(u))_{\lambda})_{N>0}$ is uniformly bounded in $N > 0$, so that, up to extracting a subsequence, we may pass to the limit as $N \rightarrow +\infty$. Thanks to Proposition 2.1, we see that the truncation $T_N(u)$ for $N > 0$ satisfies

$$(T_N(u))^{\lambda}(x) \rightarrow u^{\lambda}(x) \text{ as } N \rightarrow +\infty \text{ and } |(T_N(u))^{\lambda}(x)| \leq M[u, \lambda](x) \text{ for } |\operatorname{div} F| \text{-a.e. } x \in \Omega,$$

so that the right hand side of (3.22) clearly converges to

$$- \int_{\Omega} u F \cdot \nabla \varphi dx - \int_{\Omega} \varphi u^{\lambda} d \operatorname{div} F = \int_{\Omega} \varphi d(F, Du)_{\lambda},$$

thanks to Lebesgue's Dominated Convergence Theorem with respect to the measure $|\operatorname{div} F|$. Hence, we deduce that

$$(F, DT_N(u))_{\lambda} \rightharpoonup (F, Du)_{\lambda} \text{ in } \mathcal{M}(\Omega) \text{ as } N \rightarrow +\infty. \quad (3.23)$$

All in all, we see that there exists a sequence $(k_j)_{j \in \mathbb{N}} \subset \mathbb{N}$ such that $u_j = T_{k_j}(u)_{k_j}^{\lambda} \in C^{\infty}(\Omega) \cap BV(\Omega) \cap L^{\infty}(\Omega)$ satisfies (3.20). If in addition $u \in BV(\Omega) \cap L^{\infty}(\Omega)$, then clearly $uF \in \mathcal{DM}^{\infty}(\Omega)$ and it is not necessary to consider the truncation of u , and therefore the first part of the argument above still holds true for the sequence $(u_k^{\lambda})_{k \in \mathbb{N}}$ given by Theorem 3.2.

Finally, we deal with (3.21). Due to the homogeneity of the λ -pairing in the first component, without loss of generality we assume $\|F\|_{L^{\infty}(\Omega; \mathbb{R}^n)} \leq 1$ and $a = 1$. For all $\phi \in C_c(\Omega)$ with $\phi \geq 0$, thanks to (3.20) and Corollary 3.3 applied to $T_N(u)$ for $N > 0$, we have

$$\begin{aligned} \int_{\Omega} \phi d(F, DT_N(u))_{\lambda} + \int_{\Omega} \phi \sqrt{1 - |F|^2} dx &= \lim_{k \rightarrow +\infty} \int_{\Omega} \phi (F \cdot \nabla T_N(u)_k^{\lambda} + \sqrt{1 - |F|^2}) dx \\ &\leq \liminf_{k \rightarrow +\infty} \int_{\Omega} \phi \left| (F, \sqrt{1 - |F|^2}) \cdot (\nabla T_N(u)_k^{\lambda}, 1) \right| dx \\ &\leq \lim_{k \rightarrow +\infty} \int_{\Omega} \phi \sqrt{1 + |\nabla T_N(u)_k^{\lambda}|^2} dx \\ &= \int_{\Omega} \phi d\sqrt{1 + |DT_N(u)|^2} \leq \int_{\Omega} \phi d\sqrt{1 + |Du|^2}. \end{aligned}$$

We see that (3.23) yields

$$\int_{\Omega} \phi d(F, Du)_{\lambda} = \lim_{N \rightarrow +\infty} \int_{\Omega} \phi d(F, DT_N(u))_{\lambda} \leq \int_{\Omega} \phi d\sqrt{1 + |Du|^2} - \int_{\Omega} \phi \sqrt{1 - |F|^2} dx.$$

Due to the fact that $\phi \geq 0$, this implies that

$$(F, Du)_{\lambda}^+ \leq \sqrt{1 + |Du|^2} - \sqrt{1 - |F|^2} \mathcal{L}^n \text{ on } \Omega.$$

By an analogous computation, in which the term $\int_{\Omega} \phi \sqrt{1 - |F|^2} dx$ is subtracted instead of added, and ϕ is replaced with $-\phi$, we obtain

$$- \int_{\Omega} \phi d(F, Du)_{\lambda} + \int_{\Omega} \phi \sqrt{1 - |F|^2} dx \leq \int_{\Omega} \phi d\sqrt{1 + |Du|^2},$$

which implies

$$(F, Du)_\lambda^- \leq \sqrt{1 + |Du|^2} - \sqrt{1 - |F|^2} \mathcal{L}^n \quad \text{on } \Omega.$$

All in all, we obtain (3.21). \square

Remark 3.5. *The optimal value of a in (3.21) is not necessarily $a = \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}$. Indeed, let us consider $F(x) = (c, 0, \dots, 0)$, for some $c > 0$, and $u(x) = x_1$. Then (3.21) reduces to*

$$c \leq a\sqrt{2} - \sqrt{a^2 - c^2}$$

for all $a \geq c$. However, it is plain to see that the minimum of the function $g(a) = a\sqrt{2} - \sqrt{a^2 - c^2}$ on $[c, +\infty)$ is attained at $a = c\sqrt{2}$, and we have $g(c\sqrt{2}) = c$, while $g(c) = c\sqrt{2}$.

Remark 3.6. *We notice that we can also prove the estimate on the total variation of the λ -pairings (2.16) as a consequence of the λ -approximation (3.20). Indeed, it is not difficult to check that $u_j \rightarrow u$ in $BV(\Omega)$ -strict, thanks to Theorem 3.2 and the well-known properties of the truncation operator. Given that the strict convergence implies the weak convergence $|\nabla u_j| \mathcal{L}^n \rightharpoonup |Du|$, for all $\phi \in C_c(\Omega)$ we get*

$$\begin{aligned} \int_\Omega \phi d(F, Du)_\lambda &= \lim_{j \rightarrow +\infty} \int_\Omega \phi (F \cdot \nabla u_j) dx \\ &\leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \lim_{j \rightarrow +\infty} \int_\Omega |\phi| |\nabla u_j| dx = \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \int_\Omega |\phi| d|Du|. \end{aligned}$$

Then, this easily implies $|(F, Du)_\lambda| \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} |Du|$ on Ω .

4. PERIMETER BOUNDS AND ADMISSIBLE MEASURES

We give a simple characterization of measures which are the distributional divergence of an essentially bounded vector field.

In the following we shall use the notation $\mathcal{M}_{\mathcal{H}}(\Omega)$ introduced in (2.1).

Lemma 4.1. *Let $\mu \in \mathcal{M}(\Omega)$ and assume that there exists $F \in L^\infty(\Omega; \mathbb{R}^n)$ such that $\operatorname{div} F = \mu$. Then μ enjoys the following properties:*

- (1) $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$,
- (2) $\max\{|\mu(E^1)|, |\mu(E^1 \cup \partial^* E)|\} \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} P(E)$ for all sets $E \subset\subset \Omega$ of finite perimeter in \mathbb{R}^n ,
- (3) if in addition Ω is weakly regular, then we have

$$\max\{|\mu(E^1 \cap \Omega)|, |\mu((E^1 \cup \partial^* E) \cap \Omega)|\} \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} P(E)$$

for all sets $E \subset \Omega$ of finite perimeter in \mathbb{R}^n .

Proof. It is clear that $F \in \mathcal{DM}^\infty(\Omega)$, and so $\mu = \operatorname{div} F \in \mathcal{M}_{\mathcal{H}}(\Omega)$ by [29, Theorem 3.2]. Then, (2) and (3) are easy consequences of the Gauss–Green formulas, see Theorem 2.9. Indeed, by (2.17) and (2.18) we get

$$|\mu(E^1 \cap \Omega)| = |\operatorname{div} F(E^1 \cap \Omega)| \leq \|F\|_{L^\infty(E; \mathbb{R}^n)} P(E) \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} P(E),$$

and

$$|\mu((E^1 \cup \partial^* E) \cap \Omega)| = |\operatorname{div} F((E^1 \cup \partial^* E) \cap \Omega)| \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} P(E).$$

\square

In analogy with the bounds obtained in Lemma 4.1 for measure which are the divergence of a \mathcal{DM}^∞ vector field, we define a general class of measures satisfying similar bounds with respect to the perimeter.

Definition 4.2. *Given $\mu \in \mathcal{M}(\Omega)$ and $L > 0$, we say that μ belongs to $\mathcal{PB}_L(\Omega)$ if*

$$|\mu(E^1 \cap \Omega)| \leq LP(E) \quad \text{for all measurable sets } E \subset \Omega. \quad (4.1)$$

We also set $\mathcal{PB}(\Omega) := \bigcup_{L>0} \mathcal{PB}_L(\Omega)$ and, if $\mu \in \mathcal{PB}(\Omega)$, we say that μ satisfies a perimeter bound condition.

Remark 4.3. *If $\mu \in \mathcal{PB}(\Omega)$, then there exists $L > 0$ such that $|\mu(B_r(x))| \leq Ln\omega_n r^{n-1}$ for all $x \in \Omega$ and $r > 0$ small enough so that $B_r(x) \subset \Omega$. In particular, by [25, Theorem 4.2 and Corollary 4.3] we have $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$.*

It is interesting to notice that, under some mild regularity assumptions on Ω , the perimeter bound condition (4.1) needs only to be satisfied on open sets with smooth boundary compactly contained in Ω .

Lemma 4.4. *Let Ω be weakly regular, $\mu \in \mathcal{M}(\Omega)$ and $L > 0$. Then we have $\mu \in \mathcal{PB}_L(\Omega)$ if and only if*

$$|\mu(A)| \leq LP(A) \quad \text{for all open sets with smooth boundary } A \subset\subset \Omega. \quad (4.2)$$

Proof. Clearly, any $\mu \in \mathcal{PB}_L(\Omega)$ satisfies (4.2). As for the opposite implication, we start by noticing that, if μ satisfies (4.2), then $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$, thanks to [25, Theorem 4.2 and Corollary 4.3]. By [28, Theorem 1.1], for all $\varepsilon > 0$ there exists an increasing family of open sets with smooth boundary $(\Omega_\varepsilon)_{\varepsilon>0}$ such that $\Omega_\varepsilon \subset\subset \Omega$, $\bigcup_{\varepsilon>0} \Omega_\varepsilon = \Omega$ and $\lim_{\varepsilon \rightarrow 0^+} P(\Omega_\varepsilon) = P(\Omega)$. We consider now $E \subset \Omega$ be a set of finite perimeter, and we set $E_\varepsilon = E \cap \Omega_\varepsilon$. Due to the smoothness of Ω_ε , we obtain that $E_\varepsilon^1 = E^1 \cap \Omega_\varepsilon$. Since $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$, we can apply [10, Theorem 3.1] to conclude that there exists a smooth approximation $(E_{\varepsilon,k})_{k \in \mathbb{N}}$ of the set of finite perimeter E_ε such that

$$\lim_{k \rightarrow +\infty} \mu(E_{\varepsilon,k}) = \mu(E_\varepsilon^1) = \mu(E^1 \cap \Omega_\varepsilon) \quad \text{and} \quad \lim_{k \rightarrow +\infty} P(E_{\varepsilon,k}) = P(E \cap \Omega_\varepsilon).$$

In particular, the construction performed in [10] implies that $E_{\varepsilon,k} \subset\subset \Omega$, given that $E_\varepsilon \subset\subset \Omega$. Hence, we can apply (4.2) to $E_{\varepsilon,k}$, and pass to the limit as $k \rightarrow +\infty$ in order to get

$$|\mu(E^1 \cap \Omega_\varepsilon)| = \lim_{k \rightarrow +\infty} \mu(E_{\varepsilon,k}) \leq L \lim_{k \rightarrow +\infty} P(E_{\varepsilon,k}) = LP(E \cap \Omega_\varepsilon).$$

Now, we pass to the limit as $\varepsilon \rightarrow 0^+$ and we employ the submodularity of the perimeter [2, Proposition 3.38] to obtain

$$\begin{aligned} |\mu(E^1 \cap \Omega)| &= \lim_{\varepsilon \rightarrow 0^+} |\mu(E^1 \cap \Omega_\varepsilon)| \leq L \limsup_{\varepsilon \rightarrow 0^+} (P(E) + P(\Omega_\varepsilon) - P(E \cup \Omega_\varepsilon)) \\ &= L \left(P(E) + P(\Omega) - \liminf_{\varepsilon \rightarrow 0^+} P(E \cup \Omega_\varepsilon) \right) \leq L (P(E) + P(\Omega) - P(\Omega)) = LP(E). \end{aligned}$$

Thus, we prove (4.1). □

We state now a basic result concerning nonnegative measures in the dual of BV .

Lemma 4.5. *Let $\nu \in BV(\Omega)^*$ be nonnegative, and let $C > 0$ be such that*

$$\left| \int_{\Omega} u^* d\nu \right| \leq C \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega) \cap L^\infty(\Omega).$$

Then for any $u \in BV(\Omega)$ we have $u^\pm \in L^1(\Omega; \nu)$, as well as $u^, u^\lambda, M[u, \lambda] \in L^1(\Omega; \nu)$ for any Borel function $\lambda : \Omega \rightarrow [0, 1]$. In particular, we obtain*

$$\int_{\Omega} |u^\lambda| d\nu \leq C \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega). \quad (4.3)$$

Proof. By Theorem 3.2 there exist two sequences (u_k^0) and (u_k^1) (corresponding to the choices $\lambda \equiv 0$ and $\lambda \equiv 1$) of smooth functions in $BV(\Omega)$ converging to u in $BV(\Omega)$ -strict, and respectively to u^- and u^+ ν -almost everywhere on Ω . With a little abuse of notation, we shall write $u_k^- = u_k^0$ and $u_k^+ = u_k^1$, from this point onwards. For any fixed $N \in \mathbb{N}$, the truncation operator T_N is Lipschitz, so that the sequences of truncations $(T_N(|u_k^\pm|))$ converge to $T_N(|u^\pm|)$ ν -almost everywhere on Ω , as $k \rightarrow \infty$. By Lebesgue's Dominated Convergence Theorem we obtain

$$\begin{aligned} \int_{\Omega} T_N(|u^\pm|) d\nu &= \lim_{k \rightarrow \infty} \int_{\Omega} T_N(|u_k^\pm|) d\nu \leq C \limsup_{k \rightarrow \infty} \|T_N(|u_k|)\|_{BV(\Omega)} \\ &\leq C \limsup_{k \rightarrow \infty} \| |u_k| \|_{BV(\Omega)} \leq C \limsup_{k \rightarrow \infty} \|u_k\|_{BV(\Omega)} = C \|u\|_{BV(\Omega)}. \end{aligned}$$

By monotone convergence we can take the limit as $N \rightarrow \infty$ and conclude

$$\int_{\Omega} |u^\pm| d\nu \leq C \|u\|_{BV(\Omega)}. \quad (4.4)$$

This easily implies that $u^*, u^\lambda, M[u, \lambda] \in L^1(\Omega; \nu)$ for any Borel function $\lambda : \Omega \rightarrow [0, 1]$, and that (4.3) holds true. \square

We introduce a special class of measures related to the dual of BV , which we call admissible in view of their role in the subsequent paper [20].

Definition 4.6. *We say that $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is admissible if $|\mu| \in BV(\Omega)^*$.*

Lemma 4.7. *If $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is admissible, then, for all $u \in BV(\Omega)$ and all Borel functions $\lambda : \Omega \rightarrow [0, 1]$, we have $u^\lambda, M[u, \lambda] \in L^1(\Omega; |\mu|)$. In particular, there exists a constant $C > 0$ such that*

$$\int_{\Omega} |u^\lambda| d|\mu| \leq C \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega) \text{ and all Borel functions } \lambda : \Omega \rightarrow [0, 1]. \quad (4.5)$$

Proof. Since $|\mu| \in BV(\Omega)^*$, then Lemma 4.5 immediately yields the conclusion. \square

Remark 4.8. *We notice that, thanks to Lemma 4.7, if the measure $\operatorname{div} F$ is admissible in Theorems 2.6, 2.11 and 3.4, then we do not need to require $u^\lambda \in L^1(\Omega; |\operatorname{div} F|)$. In other words, if $\operatorname{div} F$ is admissible, then the Leibniz rule and the integration by parts formula hold true for all BV functions.*

Remark 4.9. *We notice that, if $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is admissible, then we have also $\mu \in BV(\Omega)^*$. Indeed, due to Lemma 4.7, we know that $u^* \in L^1(\Omega, |\mu|)$ for all $u \in BV(\Omega)$, and, by exploiting*

(4.5) for $\lambda = \frac{1}{2}$, we see that

$$\left| \int_{\Omega} u^* d\mu \right| \leq \int_{\Omega} |u^*| d|\mu| \leq C \|u\|_{BV(\Omega)}.$$

We underline that, in general, the converse is not true, as shown in Remark 4.12.

In the following proposition, we explore the relations between the divergence-measure fields, the perimeter bound, and the admissibility condition.

Proposition 4.10. *The following hold true:*

- (1) *If Ω is weakly regular and $F \in \mathcal{DM}^\infty(\Omega)$, then $\operatorname{div} F \in \mathcal{PB}_L(\Omega)$ for $L = \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}$; if in addition Ω has Lipschitz boundary, then $\operatorname{div} F \in BV(\Omega)^*$.*
- (2) *If Ω is a bounded open set with Lipschitz boundary and $\mu \in \mathcal{PB}(\Omega)$, then there exists $F \in \mathcal{DM}^\infty(\Omega)$ such that $\operatorname{div} F = \mu$ on Ω , there exists $\tilde{L} > 0$ such that*

$$|\mu(U \cap \Omega)| \leq \tilde{L} P(U) \quad \text{for all bounded open sets } U \subset \mathbb{R}^n \text{ with smooth boundary,} \quad (4.6)$$

we have $\mu \in BV(\Omega)^$, with its action given for all $u \in BV(\Omega) \cap L^\infty(\Omega)$ by*

$$\mathfrak{T}_\mu(u) = \int_{\Omega} u^* d\mu = -(F, Du)_*(\Omega) - \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(u) \operatorname{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1}, \quad (4.7)$$

and, if μ is admissible, formula (4.7) holds for all $u \in BV(\Omega)$. If $n = 1$, (4.6) holds for all finite unions of open intervals.

- (3) *If $|\Omega| < +\infty$ and $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is an admissible measure, then $\mu \in \mathcal{PB}(\Omega)$.*
- (4) *If $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is an admissible measure, then there exists $F, \bar{F} \in \mathcal{DM}^\infty(\Omega)$ such that $\mu = \operatorname{div} F$ and $|\mu| = \operatorname{div} \bar{F}$ on Ω .*
- (5) *If $n = 1$ and $\mu \in \mathcal{M}(\Omega)$, then μ is admissible and there exists $f, \bar{f} \in BV(\Omega)$ such that $\mu = Df$ and $|\mu| = D\bar{f}$ on Ω ; if in addition $|\mathbb{R} \setminus \Omega| > 0$, then $\mathcal{M}(\Omega) = \mathcal{PB}(\Omega)$.*

Proof. The first part of point (1) is an immediate consequence of Lemma 4.1. The second is a consequence of the integration by parts formula (2.22) for $\lambda \equiv \frac{1}{2}$: for all $u \in BV(\Omega) \cap L^\infty(\Omega)$ we exploit the continuity of the trace operator, (2.16) and (2.19) to get

$$\begin{aligned} \left| \int_{\Omega} u^* d \operatorname{div} F \right| &\leq |(F, Du)_*(\Omega)| + \left| \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(u) \operatorname{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1} \right| \\ &\leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} (|Du|(\Omega) + C \|u\|_{BV(\Omega)}). \end{aligned}$$

As for point (2), the existence of $F \in \mathcal{DM}^\infty(\Omega)$ such that $\operatorname{div} F = \mu$ on Ω follows from [25, Theorem 7.4]. Now, if $n = 1$, we notice that

$$\left| \int_{\Omega} u^\lambda d\mu \right| \leq |\mu|(\Omega) \|u\|_{L^\infty(\Omega)} \leq C \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega) \text{ and } \lambda : \Omega \rightarrow [0, 1] \text{ Borel,}$$

thanks to the embedding $BV(\Omega) \subset L^\infty(\Omega)$. Therefore, if $\lambda \equiv \frac{1}{2}$, we see that $\mu \in BV(\Omega)^*$. In addition, if $U \subset \mathbb{R}$ is a finite union of open intervals, and $\lambda \equiv 0$, so that $\chi_U^\lambda = \chi_U^- = \chi_U$, we obtain

$$|\mu(U \cap \Omega)| \leq |\mu|(\Omega) \|\chi_U\|_{L^\infty(\Omega)} \leq |\mu|(\Omega) P(U),$$

since $P(U) \geq 1$. If instead $n \geq 2$, we consider an open bounded set U with smooth boundary and apply (2.22) to the vector field F and to the function $\chi_U \in BV(\Omega) \cap L^\infty(\Omega)$, by choosing $\lambda \equiv 0$.

Given that $U = U^1$ due to the smoothness of the boundary, by (2.12) we get $\chi_U^\lambda = \chi_U^- = \chi_U$ and

$$\int_{\Omega} \chi_U d \operatorname{div} F = -(F, D\chi_U)_0(\Omega) - \int_{\partial\Omega} \operatorname{Tr}_{\partial\Omega}(\chi_U) \operatorname{Tr}^i(F, \partial\Omega) d\mathcal{H}^{n-1}.$$

Since $\operatorname{div} F = \mu$, we exploit the continuity of the trace operator, (2.16) and (2.13) to obtain

$$\begin{aligned} |\mu(U \cap \Omega)| &\leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} (|D\chi_U|(\Omega) + \|\operatorname{Tr}_{\partial\Omega}(\chi_U)\|_{L^1(\partial\Omega; \mathcal{H}^{n-1})}) \\ &\leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} (P(U, \Omega) + C\|\chi_U\|_{BV(\Omega)}) \\ &\leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \left(P(U, \Omega) + C(c_n|\Omega|^{\frac{1}{n}} + 1)P(U) \right). \end{aligned}$$

All in all, we obtain (4.6) with $\tilde{L} = \|F\|_{L^\infty(\Omega; \mathbb{R}^n)}(1 + C(c_n|\Omega|^{\frac{1}{n}} + 1))$. Thus, we exploit [25, Theorem 8.2]¹ to conclude that $\mu \in BV(\Omega)^*$. Finally, (4.7) is an immediate consequence of Theorem 2.11, combined with Lemma 4.7 and Remark 4.8 if μ is admissible.

Then, we notice that, if $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$, by (2.12) we have

$$\left| \int_{\Omega} \chi_E^* d|\mu| \right| = |\mu|(E^1 \cap \Omega) + \frac{1}{2}|\mu|(\partial^* E \cap \Omega) \geq |\mu|(E^1 \cup \Omega) \quad (4.8)$$

for any measurable set $E \subset \Omega$ of finite perimeter in \mathbb{R}^n .

Let now $|\Omega| < +\infty$, $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ be an admissible measure and $C > 0$ be such that (4.5) holds true. Hence, we choose $u = \chi_E$ for some measurable set $E \subset \Omega$ of finite perimeter in \mathbb{R}^n and we exploit (2.13) to obtain

$$\left| \int_{\Omega} \chi_E^* d|\mu| \right| \leq C\|\chi_E\|_{BV(\Omega)} \leq C(c_n|\Omega|^{\frac{1}{n}} + 1)P(E).$$

Thus, in the light of (4.8), we get $\mu \in \mathcal{PB}_L(\Omega)$ for $L = C(c_n|\Omega|^{\frac{1}{n}} + 1)$, and this proves point (3).

Point (4) is stated in [25, Lemma 7.3] in the case Ω is an open bounded set with Lipschitz boundary; however, given that the proof is completely analogous to the one in the case $\Omega = \mathbb{R}^n$ (see [25, Lemma 4.1]), we stress the fact that this representation result holds true without any additional assumption on the set Ω .

Finally, let $n = 1$. We notice that $\mathcal{M}_{\mathcal{H}}(\Omega) = \mathcal{M}(\Omega)$, and we exploit point (4) and the fact that $\mathcal{DM}^\infty(\Omega) = BV(\Omega)$. Then, if $|\mathbb{R} \setminus \Omega| > 0$, we have $P(E) \geq 1$ as long as $|E| > 0$, so that we get

$$\left| \int_{\Omega} \chi_E^* d|\mu| \right| \leq |\mu|(\Omega)P(E).$$

Combining this bound with (4.8) we obtain $\mu \in \mathcal{PB}_L(\Omega)$ for $L = |\mu|(\Omega)$, and so we conclude that $\mathcal{M}(\Omega) = \mathcal{PB}(\Omega)$ if $n = 1$. \square

As an immediate consequence of Proposition 4.10(2), we obtain the following refinement of [25, Theorem 8.2].

¹In this result, the authors extend μ to 0 on $\mathbb{R}^n \setminus \Omega$, which is equivalent to taking the intersection with Ω on the left hand side of (4.6). Also, by closely inspecting the proof, we notice that it is sufficient to take bounded open sets with smooth boundary.

Corollary 4.11. *Let Ω be a bounded open set with Lipschitz boundary and $\mu \in \mathcal{M}(\Omega)$. Then we have $\mu \in BV(\Omega)^*$ if and only if there exists $L > 0$ such that*

$$|\mu(U)| \leq LP(U) \quad \text{for all open sets with smooth boundary } U \subset\subset \Omega. \quad (4.9)$$

Proof. If $\mu \in BV(\Omega)^*$, then [25, Theorem 8.2] implies (4.9). On the other hand, combining Lemma 4.4 and (4.9), we get $\mu \in \mathcal{PB}_L(\Omega)$. Therefore, by Proposition 4.10(2) we conclude that $\mu \in BV(\Omega)^*$. \square

In the light of Proposition 4.10(5), we see that in the one dimensional case every Radon measure is admissible. This fails to be true if $n \geq 2$, even assuming the perimeter bound condition, as we show in the following remark.

Remark 4.12. *There exist measures in $\mathcal{PB}(\Omega)$ which are not admissible. To see this we can closely follow [25, Proposition 5.1], which shows the existence of a measure $\mu \in BV^*(\mathbb{R}^n)$ such that $|\mu| \notin BV^*(\mathbb{R}^n)$. We let $\mu_\varepsilon = f_\varepsilon \mathcal{L}^n$, where*

$$f_\varepsilon(x) = \varepsilon|x|^{-1-\varepsilon} \sin(|x|^{-\varepsilon}) + (n-1)|x|^{-1} \cos(|x|^{-\varepsilon}) \quad \text{for } x \neq 0$$

and some $\varepsilon \in (0, n-1)$. Then, we have $f_\varepsilon = \operatorname{div} F_\varepsilon$ for

$$F_\varepsilon(x) = \frac{x}{|x|} \cos(|x|^{-\varepsilon}).$$

Clearly, $F_\varepsilon \in \mathcal{DM}_{\text{loc}}^\infty(\mathbb{R}^n)$ and $\|F_\varepsilon\|_{L^\infty(\mathbb{R}^n; \mathbb{R}^n)} = 1$, and so, due to Proposition 4.10(1), we have $\mu_\varepsilon \in BV(\Omega)^* \cap \mathcal{PB}_1(\Omega)$ for any open bounded set Ω with Lipschitz boundary.

Then, we consider the function $u_\varepsilon(x) = |x|^{-n+1+\varepsilon}$, which satisfies $u_\varepsilon \in W_{\text{loc}}^{1,1}(\mathbb{R}^n)$. We notice that $u_\varepsilon \notin L^1(B_r; |\mu_\varepsilon|)$ for any $r > 0$. Indeed, while it is easy to check that

$$\int_{B_r} \frac{|\cos(|x|^{-\varepsilon})|}{|x|^{n-\varepsilon}} dx < +\infty,$$

we see that, if we set $c_n = \mathcal{H}^{n-1}(\partial B_1)$,

$$\int_{B_r} \frac{|\sin(|x|^{-\varepsilon})|}{|x|^n} dx = c_n \int_0^r \frac{|\sin(\rho^{-\varepsilon})|}{\rho} d\rho = [\rho^{-\varepsilon} = t] = c_n \int_{r^{-\frac{1}{\varepsilon}}}^{+\infty} \frac{1}{\varepsilon} \frac{|\sin(t)|}{t} dt = +\infty,$$

due to the well-known fact that the function $t \rightarrow \frac{\sin(t)}{t} \notin L^1((\delta, +\infty))$ for all $\delta > 0$. This proves that $u_\varepsilon f_\varepsilon \notin L^1(B_r)$, and therefore $u_\varepsilon \notin L^1(B_r; |\mu_\varepsilon|)$. More in general, for any open set Ω containing the origin we have $u_\varepsilon \notin L^1(\Omega; |\mu_\varepsilon|)$, and therefore, due to Lemma 4.7, we obtain $|\mu_\varepsilon| \notin BV(\Omega)^*$.

The previous remark shows that we cannot drop the admissibility condition on μ , if we want to ensure that $u^\lambda \in L^1(\Omega; |\mu|)$ for all $u \in BV(\Omega)$ and $\lambda : \Omega \rightarrow [0, 1]$ Borel, even in the case Ω is a bounded open set with Lipschitz boundary. However, even without the admissibility condition, it is enough to assume that at least one λ -representative of $u \in BV(\Omega)$ is $|\mu|$ -summable, in order to ensure this summability for all of them.

Remark 4.13. *If $\mu \in \mathcal{PB}(\Omega)$, Proposition 4.10(2) implies that $\mu = \operatorname{div} F$ for some $F \in \mathcal{DM}^\infty(\Omega)$, hence we can apply Proposition 2.7 to conclude that, given some Borel functions $\lambda_1, \lambda_2 : \Omega \rightarrow [0, 1]$, we have $u^{\lambda_1} \in L^1(\Omega; |\mu|)$ if and only if $u^{\lambda_2} \in L^1(\Omega; |\mu|)$. In addition, for any Borel function $\lambda : \Omega \rightarrow [0, 1]$, we have $u^\lambda \in L^1(\Omega; |\mu|)$ if and only if $M[u, \lambda] \in L^1(\Omega; |\mu|)$, again by Proposition 2.7.*

We provide a few basic examples of admissible measures, which, in the light of Proposition 4.10(3), also belong to in $\mathcal{PB}(\Omega)$, as long as $|\Omega| < +\infty$.

Example 4.14. Let Ω be a bounded open set with Lipschitz boundary. We consider $\mu \in \mathcal{M}(\Omega)$ defined as

$$\mu = h \mathcal{L}^n + \gamma \mathcal{H}^{n-1} \llcorner \Gamma$$

where $h \in L^q(\Omega)$ for some $q > n$, $\gamma \in L^\infty(\Omega; \mathcal{H}^{n-1} \llcorner \Gamma)$, and $\Gamma \subset \Omega$ is a compact set such that there exist $\Lambda, \bar{\rho} > 0$ for which

$$\mathcal{H}^{n-1}(\Gamma \cap B_\rho(x)) \leq \Lambda \rho^{n-1} \quad \forall 0 < \rho < \bar{\rho} \text{ and } \forall x \in \Gamma. \quad (4.10)$$

We claim that μ is admissible, and we prove it by showing that both the measures $h \mathcal{L}^n$ and $\gamma \mathcal{H}^{n-1} \llcorner \Gamma$ are admissible. In particular, this implies that all these measures belong to $\mathcal{PB}(\Omega)$, thanks to Proposition 4.10(3). We first point out that, thanks to the Sobolev embedding of $BV(\Omega)$ into $L^{\frac{n}{n-1}}(\Omega)$, we get

$$\left| \int_{\Omega} u |h| dx \right| \leq \|h\|_{L^n(\Omega)} \|u\|_{L^{\frac{n}{n-1}}(\Omega)} \leq C \|h\|_{L^n(\Omega)} \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega),$$

This shows that $h \mathcal{L}^n$ is an admissible measure.

Then, we notice that the reason for requiring (4.10) is that it ensures a crucial continuity property of the upper and lower trace operators from BV to $L^1(\Gamma; \mathcal{H}^{n-1})$. Indeed, there exists a constant $C > 0$ depending only on $\Omega, \mathcal{H}^{n-1}(\Gamma), \bar{\rho}$ and Λ such that

$$\int_{\Gamma} |u^\pm| d\mathcal{H}^{n-1} \leq C \|u\|_{BV(\Omega)} \quad \text{for all } u \in BV(\Omega). \quad (4.11)$$

In order to show (4.11), we consider the zero-extension of u to \mathbb{R}^n , which we denote by u_0 : we have $u_0 \in BV(\mathbb{R}^n)$ and

$$\|u_0\|_{BV(\mathbb{R}^n)} \leq c \|u\|_{BV(\Omega)}, \quad (4.12)$$

for some $c = c(\Omega) > 0$ (see for instance [31, Lemma 5.10.4]). Due to (4.10), we can apply [2, Theorem 3.86] to get

$$\int_{\Gamma} |u^\pm| d\mathcal{H}^{n-1} \leq C \|u_0\|_{BV(\mathbb{R}^n)} \quad (4.13)$$

where $C > 0$ is a constant depending only on $\mathcal{H}^{n-1}(\Gamma), \bar{\rho}$ and Λ . Combining (4.12) and (4.13), we obtain (4.11), which in particular implies $\mathcal{H}^{n-1} \llcorner \Gamma \in BV(\Omega)^*$ and that $u^\pm \in L^1(\Gamma; \mathcal{H}^{n-1})$. Hence, for all $u \in BV(\Omega)$ we obtain

$$\left| \int_{\Gamma} u^* |\gamma| d\mathcal{H}^{n-1} \right| \leq \int_{\Gamma} |u^*| |\gamma| d\mathcal{H}^{n-1} \leq C \|\gamma\|_{L^\infty(\Gamma; \mathcal{H}^{n-1})} \|u\|_{BV(\Omega)},$$

where $C > 0$ is a constant depending only on $\Omega, \mathcal{H}^{n-1}(\Gamma), \bar{\rho}$ and Λ . Therefore, the measure $\gamma \mathcal{H}^{n-1} \llcorner \Gamma$ is admissible. All in all, this proves that μ is an admissible measure.

Example 4.15. Let $n \geq 2$ and $\mu = h \mathcal{L}^n$, where

$$h(x) = \frac{(n-1)}{|x-x_0|} \quad \text{for } x \neq x_0,$$

for some $x_0 \in \mathbb{R}^n$. We notice that $h \in L_{\text{loc}}^q(\mathbb{R}^n)$ only for $q \in [1, n)$. However, it is easy to check that $\mu = \text{div } F$, where

$$F(x) = \frac{(x - x_0)}{|x - x_0|} \text{ for } x \neq x_0.$$

By Proposition 4.10(1), for any open bounded set Ω with Lipschitz boundary we have $\mu \in BV(\Omega)^*$, so that μ is indeed an admissible measure, given that $\mu \geq 0$. In addition, since $\|F\|_{L^\infty(\mathbb{R}^n; \mathbb{R}^n)} = 1$, we see that $\mu \in \mathcal{PB}_1(\Omega)$, by Lemma 4.1.

Given that the admissibility of a measure depends on its total variation, it seems natural to characterize it in terms of properties of the positive and negative parts of the given measure. In the following lemma we prove that any measure whose positive and negative parts belong to $\mathcal{PB}(\Omega)$ is indeed admissible.

Lemma 4.16. *Let Ω be an open bounded set with Lipschitz boundary and let $\mu \in \mathcal{M}(\Omega)$. Assume that its positive and negative parts, μ^+ and μ^- , belong to $\mathcal{PB}_{L_+}(\Omega)$ and $\mathcal{PB}_{L_-}(\Omega)$ for some $L_+, L_- > 0$, respectively. Then μ is admissible and belongs to $\mathcal{PB}_L(\Omega)$, where $L = \max\{L_+, L_-\}$. In particular, if $\mu \geq 0$ and $\mu \in \mathcal{PB}(\Omega)$, then it is admissible.*

Proof. Thanks to Proposition 4.10(2), we know that $\mu^+, \mu^- \in BV(\Omega)^*$. Hence, a simple application of the triangle inequality implies that $|\mu| \in BV(\Omega)^*$. Let now $L_+, L_- > 0$ be the constants for which μ^+ and μ^- satisfy (4.1), respectively. We notice that, for all measurable sets $E \subset \Omega$, we have

$$|\mu(E^1 \cap \Omega)| = |\mu^+(E^1 \cap \Omega) - \mu^-(E^1 \cap \Omega)| \leq \max\{\mu^+(E^1 \cap \Omega), \mu^-(E^1 \cap \Omega)\} \leq \max\{L_+, L_-\}P(E),$$

so that $\mu \in \mathcal{PB}_L(\Omega)$, for $L = \max\{L_+, L_-\}$. \square

A relevant consequence of the results explored so far is that, for admissible measures μ , the functional $\mathfrak{T}_{\mu, \lambda}$ defined in (2.25) can be naturally extended to all functions in $BV(\Omega)$.

Lemma 4.17. *Let $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ be admissible and $\lambda : \Omega \rightarrow [0, 1]$ be a Borel function. Then the functional $\mathfrak{T}_{\mu, \lambda} : BV(\Omega) \rightarrow \mathbb{R}$ given by (2.25) is well-defined. Then, given $u \in BV(\Omega)$, if we consider the truncation of u , $T_N(u)$ for $N > 0$, and its smooth approximation $T_N(u)_k^\lambda$ given by Theorem 3.2, we have*

$$\lim_{N \rightarrow +\infty} \int_{\Omega} T_N(u)^\lambda d\mu = \int_{\Omega} u^\lambda d\mu \quad (4.14)$$

and

$$\lim_{N \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{\Omega} T_N(u)_k^\lambda d\mu = \int_{\Omega} u^\lambda d\mu. \quad (4.15)$$

Finally, for any $u \in BV(\Omega)$ we can find a sequence $(u_j)_{j \in \mathbb{N}} \subset C^\infty(\Omega) \cap BV(\Omega) \cap L^\infty(\Omega)$ such that

$$\lim_{j \rightarrow +\infty} \mathfrak{T}_{\mu, \lambda}(u_j) = \mathfrak{T}_{\mu, \lambda}(u). \quad (4.16)$$

Proof. We recall that $\mathfrak{T}_{\mu, \lambda}$ is well-defined on $BV(\Omega) \cap L^\infty(\Omega)$. Thanks to Lemma 4.7, $|\mu| \in BV(\Omega)^*$ implies $u^\lambda, M[u, \lambda] \in L^1(\Omega; |\mu|)$. Hence, for any $u \in BV(\Omega)$ we get (4.14) by Lebesgue's Dominated Convergence Theorem with respect to the measure $|\mu|$, thanks to (2.9). Hence, the functional

$$\mathfrak{T}_{\mu, \lambda}(u) = \int_{\Omega} u^\lambda d\mu$$

is well-defined for any $u \in BV(\Omega)$. Let now $N > 0$ and $T_N(u)_k^\lambda$ be the smooth approximation of $T_N(u)$ given by Theorem 3.2. Then, we have

$$T_N(u)_k^\lambda(x) \rightarrow T_N(u)^\lambda(x) \text{ as } k \rightarrow +\infty \text{ for } |\mu|\text{-a.e. } x \in \Omega,$$

since $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$, and $|T_N(u)_k^\lambda(x)| \leq 2N$ for every $x \in \Omega$ and $k \geq 1$. Therefore we obtain

$$\lim_{k \rightarrow \infty} \int_{\Omega} T_N(u)_k^\lambda d\mu = \int_{\Omega} T_N(u)^\lambda d\mu$$

by Lebesgue's Dominated Convergence Theorem with respect to $|\mu|$, since the constant N is a summable majorant, given that $|\mu|(\Omega) < \infty$. Hence, combining this result with (4.14), we get

$$\lim_{N \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{\Omega} T_N(u)_k^\lambda d\mu = \lim_{N \rightarrow +\infty} \int_{\Omega} T_N(u)^\lambda d\mu = \int_{\Omega} u^\lambda d\mu.$$

This proves (4.16). Consequently, there exists a sequence $(k_j)_{j \in \mathbb{N}} \subset \mathbb{N}$ such that $u_j := T_{k_j}(u)_{k_j}^\lambda$ belongs to $C^\infty(\Omega) \cap BV(\Omega) \cap L^\infty(\Omega)$ and satisfies (4.15). \square

Remark 4.18. *It is easy to see that, if $\mu \in BV^*(\Omega)$ and $\mu \geq 0$, then μ is an admissible measure. Hence, if we choose $\lambda \equiv \frac{1}{2}$, Lemma 4.17 provides an extension of [25, Theorem 7.4 (iv) and Remark 8.3], with no additional assumptions on Ω .*

We exploit the previous result to find a sharp estimate of $|\mathfrak{T}_{\mu, \lambda}(u)|$ in terms of the total variation of u and its trace on the boundary of the domain.

Proposition 4.19. *Let Ω be an open bounded set with Lipschitz boundary. Let $L > 0$ and $\mu \in \mathcal{PB}_L(\Omega)$. Let $u \in BV(\Omega)$ and $\lambda : \Omega \rightarrow [0, 1]$ be a Borel function. If $u^\lambda \in L^1(\Omega; |\mu|)$, then*

$$\left| \int_{\Omega} u^\lambda d\mu \right| \leq L \left(|Du|(\Omega) + \int_{\partial\Omega} |\text{Tr}_{\partial\Omega}(u)| d\mathcal{H}^{n-1} \right). \quad (4.17)$$

In particular, (4.17) holds true for all $u \in BV(\Omega)$ as long as μ is admissible.

Proof. We start by noticing that, thanks to Remark 4.3, we know that $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$. In addition, by Remark 4.13, we deduce that $M[u, \lambda] \in L^1(\Omega; |\mu|)$, given that $u^\lambda \in L^1(\Omega; |\mu|)$. In particular, if μ is admissible, Lemma 4.7 ensures that $u^\lambda, M[u, \lambda] \in L^1(\Omega; |\mu|)$ for all $u \in BV(\Omega)$. Take now $N > 0$ and let $T_N(u)_k^\lambda$ be the smooth approximation of $T_N(u)$ given by Theorem 3.2. Arguing as in the proof of Lemma 4.17, we see that

$$\int_{\Omega} T_N(u)^\lambda d\mu = \lim_{k \rightarrow \infty} \int_{\Omega} T_N(u)_k^\lambda d\mu. \quad (4.18)$$

We notice now that $T_N(u)_k^\lambda \in C^\infty(\Omega)$, so that the superlevel and sublevel sets

$$\{T_N(u)_k^\lambda > t\} = \{x \in \Omega : T_N(u)_k^\lambda(x) > t\} \text{ and } \{T_N(u)_k^\lambda < t\} = \{x \in \Omega : T_N(u)_k^\lambda(x) < t\}$$

are open sets with smooth boundary for \mathcal{L}^1 -a.e. $t \in \mathbb{R}$, by Morse-Sard lemma [22, Lemma 13.15]. In particular, we deduce that, for \mathcal{L}^1 -a.e. $t \in \mathbb{R}$,

$$\{T_N(u)_k^\lambda > t\}^1 = \{T_N(u)_k^\lambda > t\} \text{ and } \{T_N(u)_k^\lambda < t\}^1 = \{T_N(u)_k^\lambda < t\}.$$

We exploit now Theorem 3.2 to notice that

$$|T_N(u)_k^\lambda| \leq 2N \text{ for all } k \geq 1, \quad (4.19)$$

$$T_N(u)_k^\lambda \rightarrow T_N(u) \text{ in } BV(\Omega)\text{-strict as } k \rightarrow +\infty, \quad (4.20)$$

and

$$\mathrm{Tr}_{\partial\Omega} \left(T_N(u)_k^\lambda \right) (x) = \mathrm{Tr}_{\partial\Omega}(T_N(u))(x) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in \partial\Omega. \quad (4.21)$$

Therefore, by using (4.18), the layer-cake representation together with (4.19), the fact that μ satisfies (4.1), the coarea formula, Lemma 2.3, the convergence (4.20) and the identity (4.21), we get

$$\begin{aligned} \left| \int_{\Omega} T_N(u)^\lambda d\mu \right| &= \lim_{k \rightarrow \infty} \left| \int_0^\infty \int_{\{T_N(u)_k^\lambda > t\}} d\mu dt - \int_{-\infty}^0 \int_{\{T_N(u)_k^\lambda < t\}} d\mu dt \right| \\ &\leq \limsup_{k \rightarrow \infty} \int_0^{2N} \left| \mu(\{T_N(u)_k^\lambda > t\}) \right| dt + \int_{-2N}^0 \left| \mu(\{T_N(u)_k^\lambda < t\}) \right| dt \\ &\leq L \limsup_{k \rightarrow \infty} \int_0^{2N} P(\{T_N(u)_k^\lambda > t\}) dt + \int_{-2N}^0 P(\{T_N(u)_k^\lambda < t\}) dt \\ &\leq L \limsup_{k \rightarrow \infty} \int_{-2N}^{2N} |D\chi_{\{T_N(u)_k^\lambda > t\}}|(\Omega) dt + \int_0^{2N} \int_{\partial\Omega} \mathrm{Tr}_{\partial\Omega}(\chi_{\{T_N(u)_k^\lambda > t\}}) d\mathcal{H}^{n-1} dt \\ &\leq L \limsup_{k \rightarrow \infty} \int_{-\infty}^\infty |D\chi_{\{T_N(u)_k^\lambda > t\}}|(\Omega) dt + \int_0^{+\infty} \int_{\partial\Omega} \mathrm{Tr}_{\partial\Omega}(\chi_{\{T_N(u)_k^\lambda > t\}}) d\mathcal{H}^{n-1} dt \\ &\leq L \lim_{k \rightarrow \infty} |DT_N(u)_k^\lambda|(\Omega) + \int_{\partial\Omega} |\mathrm{Tr}_{\partial\Omega}(T_N(u)_k^\lambda)| d\mathcal{H}^{n-1} \\ &= L \left(|DT_N(u)|(\Omega) + \int_{\partial\Omega} |\mathrm{Tr}_{\partial\Omega}(T_N(u))| d\mathcal{H}^{n-1} \right) \\ &\leq L \left(|Du|(\Omega) + \int_{\partial\Omega} |\mathrm{Tr}_{\partial\Omega}(T_N(u))| d\mathcal{H}^{n-1} \right), \end{aligned}$$

since $|DT_N(u)| \leq |Du|$. In addition, we notice that $T_N(u) \rightarrow u$ in $BV(\Omega)$ -strict as $N \rightarrow +\infty$, so that we obtain

$$\mathrm{Tr}_{\partial\Omega}(T_N(u)) \rightarrow \mathrm{Tr}_{\partial\Omega}(u) \text{ in } L^1(\partial\Omega; \mathcal{H}^{n-1}) \text{ as } N \rightarrow +\infty,$$

see for instance to [2, Theorem 3.88]. Thus, we obtain (4.17) by passing to the limit as $N \rightarrow +\infty$ also on the left hand side, again thanks to the Lebesgue's Dominated Convergence Theorem, employing (2.9), (2.10), and the fact that $M[u, \lambda] \in L^1(\Omega; |\mu|)$. \square

Remark 4.20. *We stress that we could not directly apply the Cavalieri representation formula to $\int_{\Omega} T_N(u)^\lambda d\mu$, since, a priori, we do not know whether*

$$\{T_N(u)^\lambda > t\} = \{T_N(u)^\lambda > t\}^1 \text{ or } \{T_N(u)^\lambda > t\} = \{T_N(u)^\lambda > t\}^1 \cup \partial^* \{T_N(u)^\lambda > t\}.$$

Therefore, we would not be able to apply the property (4.1) and proceed with the proof. Instead, we see here the usefulness of Theorem 3.2 in ensuring that, for \mathcal{L}^1 -a.e. $t \in \mathbb{R}$,

$$\begin{aligned} \{(T_N(u))_k^\lambda > t\}^1 &= \{(T_N(u))_k^\lambda > t\}, \quad \partial^* \{(T_N(u))_k^\lambda > t\} = \partial \{(T_N(u))_k^\lambda > t\}, \\ \{(T_N(u))_k^\lambda < t\}^1 &= \{(T_N(u))_k^\lambda < t\}, \quad \partial^* \{(T_N(u))_k^\lambda < t\} = \partial \{(T_N(u))_k^\lambda < t\}. \end{aligned}$$

Lemma 4.21. *Let Ω be a bounded open set with Lipschitz boundary. Let $L > 0$ and $\mu \in \mathcal{PB}_L(\Omega)$ be an admissible measure. Then there exists $F \in \mathcal{DM}^\infty(\Omega)$ such that $\operatorname{div} F = \mu$ on Ω and $\|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq L$.*

Proof. Thanks to Lemma 4.7, for all $u \in BV(\Omega)$ and Borel functions $\lambda : \Omega \rightarrow [0, 1]$ we have $u^\lambda \in L^1(\Omega; |\mu|)$. Hence, Proposition 4.19 implies that $\mu \in W_0^{1,1}(\Omega)^*$; that is, it defines a linear continuous functional on $W_0^{1,1}(\Omega)$, since

$$\left| \int_{\Omega} \tilde{u} d\mu \right| \leq L \|\nabla u\|_{L^1(\Omega; \mathbb{R}^n)},$$

by (4.17), given that $\operatorname{Tr}_{\partial\Omega}(u) = 0$ and $u^\lambda(x) = u^*(x) = \tilde{u}(x)$ for \mathcal{H}^{n-1} -a.e. $x \in \Omega$ and all Borel functions $\lambda : \Omega \rightarrow [0, 1]$. Hence, [25, Lemma 7.3] yields

$$\begin{aligned} \|\mu\|_{W_0^{1,1}(\Omega)^*} &= \sup \left\{ \int_{\Omega} \tilde{u} d\mu : u \in W_0^{1,1}(\Omega), \|\nabla u\|_{L^1(\Omega; \mathbb{R}^n)} \leq 1 \right\} \\ &= \min \left\{ \|G\|_{L^\infty(\Omega; \mathbb{R}^n)} : G \in L^\infty(\Omega; \mathbb{R}^n), \operatorname{div} G = \mu \text{ on } \Omega \right\}. \end{aligned}$$

Thus, this ends the proof. \square

Remark 4.22. *Under the same assumptions of Lemma 4.21, we notice that the measure μ actually satisfies*

$$|\mu((E^1 \cup \partial^* E) \cap \Omega)| \leq L P(E),$$

for all sets $E \subset \Omega$ of finite perimeter in \mathbb{R}^n , coherently with properties (2) and (3) of Lemma 4.1. Indeed, Ω is an open bounded set with Lipschitz boundary, and hence it is weakly regular. Therefore, we can choose a vector field F as in Lemma 4.21 and, using the Gauss-Green formula (2.18), we immediately obtain

$$\begin{aligned} |\mu((E^1 \cup \partial^* E) \cap \Omega)| &= |\operatorname{div} F((E^1 \cup \partial^* E) \cap \Omega)| = \left| \int_{\partial^* E} \operatorname{Tr}^e(F, \partial^* E) d\mathcal{H}^{n-1} \right| \\ &\leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} P(E) \leq L P(E) \end{aligned}$$

for all sets $E \subset \Omega$ of finite perimeter in \mathbb{R}^n .

In addition, we notice that, if $\mu \in \mathcal{M}_{\mathcal{H}}(\Omega)$ is an admissible measure, then Theorem 2.11 (taking into account Remark 4.8), (2.16) and (2.19) implies the estimate

$$\left| \int_{\Omega} u^\lambda d\mu \right| \leq \Lambda \left(|Du|(\Omega) + \int_{\partial\Omega} |\operatorname{Tr}_{\partial\Omega}(u)| d\mathcal{H}^{n-1} \right),$$

where

$$\Lambda = \inf \{ \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} : F \in \mathcal{DM}^\infty(\Omega) \text{ with } \operatorname{div} F = \mu \}.$$

However, assuming $\mu \in \mathcal{PB}_L(\Omega)$ for some $L > 0$, we cannot prove that $\Lambda \leq L$ without exploiting Proposition 4.19 itself, since in the proof of Lemma 4.21 it is essential to know that the perimeter bound for sets holds at least for functions in $W_0^{1,1}(\Omega)$.

As a final result, we show the stability of the admissibility and the perimeter bound conditions under a suitable type of smooth approximation procedure. More precisely, we prove that any admissible measure $\mu \in \mathcal{PB}_L(\Omega)$ can be approximated in the weak- $*$ sense by a sequence of absolutely continuous measures $\mu_j \mathcal{L}^n \in \mathcal{PB}_{L_j}(\Omega)$ with smooth density functions, where $L_j \rightarrow L$ as $j \rightarrow +\infty$. The key idea is to use the duality between admissible measures in $\mathcal{PB}(\Omega)$ and

essentially bounded divergence-measure fields. While it is not clear if a direct mollification of μ would in general still belong to $\mathcal{PB}_L(\Omega)$ (even assuming μ of compact support in Ω), we shall instead rely on the fact that the Anzellotti-Giaquinta-type regularization of a vector field almost preserves its L^∞ norm. Similar ideas can be found for instance in [15].

Proposition 4.23. *Let Ω be an open bounded set with Lipschitz boundary. Let $L > 0$ and $\mu \in \mathcal{PB}_L(\Omega)$ be an admissible measure. Then there exists a sequence of functions $(\mu_j)_j \subset C^\infty(\Omega)$ such that $\mu_j \mathcal{L}^n$ is an admissible measure and it belongs to $\mathcal{PB}_{L(1+1/j)}(\Omega)$ for all $j \in \mathbb{N}$, and $\mu_j \mathcal{L}^n \rightharpoonup \mu$ as $j \rightarrow +\infty$. Moreover, if $\mu = h \mathcal{L}^n + \mu^s$ with $h \in L^q(\Omega)$, for some $q \in [1, +\infty]$, and μ^s has compact support in Ω , then for any open set $\Omega' \subset\subset \Omega$ containing the support of μ^s we have the following properties:*

- *there exists $(h_j)_j$ such that $h_j \rightarrow h$ in $L^p(\Omega)$, for all $p \in [1, q)$ and $p = q$ if $q < +\infty$, and $\mu_j - h_j \rightarrow 0$ uniformly on $\Omega \setminus \Omega'$;*
- *$\mu_j = \rho_j * \mu^s + h_j$ on Ω' for j large enough and $\rho_j(x) = \delta_j^{-n} \rho(x/\delta_j)$, for some sequence $\delta_j \downarrow 0$ and some fixed mollifier $\rho \in C_c^\infty(B_1)$.*

Proof. The construction of the sequence μ_j starting from the measure μ can be done as follows. We recall that, thanks to Proposition 4.10(4) and Lemma 4.21, if $\mu \in \mathcal{PB}_L(\Omega)$ and it is admissible, then there exist vector fields $F, G \in \mathcal{DM}^\infty(\Omega)$ such that

- $\|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq L$ and $\operatorname{div} F = \mu$ on Ω ,
- $\operatorname{div} G = |\mu|$ on Ω .

Analogously as in the proof of Theorem 3.1, we set $\Omega_0 = \emptyset$,

$$\Omega_k = \left\{ x \in \Omega : \operatorname{dist}(x, \partial\Omega) > \frac{1}{k} \right\} \quad \text{for all } k \geq 1,$$

and then $\Sigma_k := \Omega_{k+1} \setminus \overline{\Omega_{k-1}}$. We let $(\zeta_k)_{k \geq 1}$ be a partition of unity subordinate to the open cover $(\Sigma_k)_{k \geq 1}$ of Ω . We choose a nonnegative sequence $(\delta_{j,k})_{j,k \geq 1}$ such that $\delta_{j,k} \rightarrow 0$ as $j \rightarrow +\infty$ for all $k \geq 1$ and

$$\operatorname{supp}(\zeta_k) + B_{\delta_{j,k}} \subset \Sigma_k. \quad (4.22)$$

Finally, we take a standard mollifier ρ and we require that the family of mollifiers $\rho_{j,k} = \rho_{\delta_{j,k}}$ satisfies the following set of conditions:

$$\|\rho_{\delta_{j,k}} * \zeta_k - \zeta_k\|_{L^\infty(\Omega)} \leq \frac{1}{j2^k}, \quad (4.23)$$

$$\|\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k\|_{L^r(\Omega)} \leq \frac{1}{j2^k} \quad (4.24)$$

for all $r \in [1, n]$,

$$\|\rho_{j,k} * (G \cdot \nabla \zeta_k) - G \cdot \nabla \zeta_k\|_{L^n(\Omega)} \leq \frac{1}{j2^k} \quad (4.25)$$

and, if $\mu = h \mathcal{L}^n + \mu^s$,

$$\|\rho_{j,k} * (\zeta_k h) - \zeta_k h\|_{L^p(\Omega)} \leq \frac{1}{j2^k}, \quad (4.26)$$

for $p \in [1, q)$ and $p = q$ if $q < +\infty$. Then we set

$$F_j = \sum_{k=1}^{+\infty} \rho_{j,k} * (\zeta_k F)$$

and, finally,

$$\mu_j = \operatorname{div} F_j. \quad (4.27)$$

With reference to the proof of Theorem 3.1, we fix $\varphi \in C_c^\infty(\Omega)$ and obtain

$$\begin{aligned} \int_{\Omega} \varphi \mu_j dx &= - \int_{\Omega} F_j \cdot \nabla \varphi dx \\ &= - \sum_{k=1}^{+\infty} \int_{\Omega} (\zeta_k F) * \rho_{j,k} \cdot \nabla \varphi dx \\ &= \sum_{k=1}^{+\infty} \int_{\Omega} \zeta_k (\rho_{j,k} * \varphi) d\mu + \sum_{k=1}^{+\infty} \int_{\Omega} \varphi \left(\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k \right) dx \\ &\rightarrow \int_{\Omega} \varphi d\mu, \quad \text{as } j \rightarrow \infty, \end{aligned}$$

where the final passage to the limit is justified because

$$\left| \int_{\Omega} \varphi \left(\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k \right) dx \right| \leq \|\varphi\|_{L^\infty(\Omega)} \|\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k\|_{L^1(\Omega)} \leq \frac{\|\varphi\|_{L^\infty(\Omega)}}{j2^k}$$

thanks to (4.24). As a byproduct of the previous calculations, we also get

$$\mu_j = \sum_{k=1}^{+\infty} \rho_{j,k} * (\zeta_k \mu) + \sum_{k=1}^{+\infty} \left(\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k \right). \quad (4.28)$$

In order to prove that $\mu_j \mathcal{L}^n$ is admissible, we consider separately the two sums on the right hand side. Given that $\operatorname{div} G = |\mu|$, it is easy to see that

$$|\mu_j| \leq \sum_{k=1}^{+\infty} \rho_{j,k} * (\zeta_k \operatorname{div} G) + \sum_{k=1}^{+\infty} |\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k|.$$

Due to (4.24) and the Sobolev embedding $BV(\Omega) \subset L^{\frac{n}{n-1}}(\Omega)^2$, for all $u \in BV(\Omega)$, we have

$$\begin{aligned} \left| \int_{\Omega} u \sum_{k=1}^{+\infty} |\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k| dx \right| &\leq \sum_{k=1}^{+\infty} \|u\|_{L^{\frac{n}{n-1}}(\Omega)} \|\rho_{j,k} * (F \cdot \nabla \zeta_k) - F \cdot \nabla \zeta_k\|_{L^n(\Omega)} \\ &\leq C \|u\|_{BV(\Omega)}. \end{aligned}$$

As for the first sum, given that $\operatorname{div} G$ is an admissible measure, in the light of Remark 4.8 we can exploit the integration by parts formula (2.22) to obtain

$$\begin{aligned} \int_{\Omega} u \sum_{k=1}^{+\infty} \rho_{j,k} * (\zeta_k \operatorname{div} G) dx &= \sum_{k=1}^{+\infty} \int_{\Omega} \zeta_k (\rho_{j,k} * u) d \operatorname{div} G \\ &= - \sum_{k=1}^{+\infty} \int_{\Omega} G \cdot \nabla (\zeta_k (\rho_{j,k} * u)) dx, \end{aligned}$$

²If $n = 1$, $BV(\Omega) \subset L^\infty(\Omega)$.

since $\text{supp}(\zeta_k) \subset\subset \Omega$. Now, we exploit the Leibniz rule to get

$$\begin{aligned} \sum_{k=1}^{+\infty} \int_{\Omega} G \cdot \nabla (\zeta_k(\rho_{j,k} * u)) \, dx &= \int_{\Omega} G \cdot \left(\sum_{k=1}^{+\infty} \zeta_k(\rho_{j,k} * Du) \right) \, dx + \sum_{k=1}^{+\infty} \int_{\Omega} u \nabla \zeta_k \cdot (\rho_{j,k} * G - G) \, dx \\ &\quad + \sum_{k=1}^{+\infty} \int_{\Omega} u \nabla \zeta_k \cdot G \, dx. \end{aligned}$$

We deal with the three terms separately. As for the first term, we exploit the fact that $\sum_{k=1}^{+\infty} \chi_{\Sigma_k} \leq 2$ and (4.23) in order to get

$$\begin{aligned} \left| \int_{\Omega} G \cdot \left(\sum_{k=1}^{+\infty} \zeta_k(\rho_{j,k} * Du) \right) \, dx \right| &\leq \|G\|_{L^\infty(\Omega; \mathbb{R}^n)} \sum_{k=1}^{+\infty} \int_{\text{supp}(\zeta_k)} \rho_{j,k} * |Du| \, dx \\ &\leq 2\|G\|_{L^\infty(\Omega; \mathbb{R}^n)} |Du|(\Omega). \end{aligned}$$

Then, by (4.25) we see that

$$\left| \sum_{k=1}^{+\infty} \int_{\Omega} u \nabla \zeta_k \cdot (\rho_{j,k} * G - G) \, dx \right| \leq \sum_{k=1}^{+\infty} \|u\|_{L^{\frac{n}{n-1}}(\Omega)} \|\rho_{j,k} * (G \cdot \nabla \zeta_k) - G \cdot \nabla \zeta_k\|_{L^n(\Omega)} \leq C\|u\|_{BV(\Omega)},$$

again thanks to the Sobolev embedding for BV functions. Finally, we exploit again (2.22) to obtain

$$\sum_{k=1}^{+\infty} \int_{\Omega} u \nabla \zeta_k \cdot G \, dx = - \sum_{k=1}^{+\infty} \int_{\Omega} \zeta_k \, d \text{div}(uG) = - \int_{\Omega} d \text{div}(uG) = \int_{\partial\Omega} \text{Tr}^i(G, \partial\Omega) \text{Tr}_{\partial\Omega}(u) \, d\mathcal{H}^{n-1},$$

so that, thanks to (2.19), we get

$$\left| \sum_{k=1}^{+\infty} \int_{\Omega} u \nabla \zeta_k \cdot G \, dx \right| \leq \|G\|_{L^\infty(\Omega; \mathbb{R}^n)} \|u\|_{BV(\Omega)}.$$

All in all, this shows that $|\mu_j|_{\mathcal{L}^n} \in BV(\Omega)^*$.

In addition, when μ^s has compact support in $\Omega' \subset\subset \Omega$, and for j large, the first sum in (4.28) coincides with $\rho_{j,1} * \mu^s + h_j$ if restricted to Ω' , where h_j is the Anzellotti-Giaquinta's regularization of h , and it coincides with h_j when restricted to $\Omega \setminus \Omega'$. At the same time, the second sum in (4.28) is uniformly small and tends to 0 as $j \rightarrow \infty$ by (4.24). Finally, $h_j \rightarrow h$ in $L^p(\Omega)$ with p as above, by (4.26). This shows the last part of the statement.

Finally, we prove that $\mu_j \in \mathcal{PB}_{L(1+1/j)}(\Omega)$. Given any set $E \subset \Omega$ of finite perimeter in \mathbb{R}^n , by Theorem 2.9 we have

$$\left| \int_{E \cap \Omega} \mu_j \, dx \right| = \left| \int_{\partial^* E} \text{Tr}^i(F_j, \partial^* E) \, d\mathcal{H}^{n-1} \right| \leq \|F_j\|_{L^\infty(\Omega; \mathbb{R}^n)} P(E).$$

Then, we notice that, for any $x \in \Omega$, there exists a unique $k_0 \geq 1$ such that $x \in \Sigma_{k_0} \cap \Sigma_{k_0+1}$, so that, arguing as in the proof of point (5) of Theorem 3.1, thanks to the assumptions on ζ_k and $\rho_{j,k}$, in particular (4.22) and (4.23), we obtain

$$|F_j(x)| \leq \|F\|_{L^\infty(\Omega; \mathbb{R}^n)} \left(1 + \sum_{k=k_0}^{k_0+1} |(\zeta_k * \rho_{j,k})(x) - \zeta_k(x)| \right) \leq L(1 + 1/j),$$

□

REFERENCES

- [1] Luigi Ambrosio, Gianluca Crippa, and Stefania Maniglia. Traces and fine properties of a BD class of vector fields and applications. In *Annales de la Faculté des sciences de Toulouse: Mathématiques*, volume 14, pages 527–561, 2005.
- [2] Luigi Ambrosio, Nicola Fusco, and Diego Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford University Press, 2000.
- [3] Gabriele Anzellotti. Pairings between measures and bounded functions and compensated compactness. *Annali di Matematica Pura ed Applicata*, 135(1):293–318, 1983.
- [4] Gui-Qiang Chen, Giovanni E. Comi, and Monica Torres. Cauchy fluxes and Gauss–Green formulas for divergence-measure fields over general open sets. *Archive for Rational Mechanics and Analysis*, 233: 87–166, 2019.
- [5] Gui-Qiang Chen and Hermano Frid. Divergence-measure fields and hyperbolic conservation laws. *Archive for rational mechanics and analysis*, 147(2):89–118, 1999.
- [6] Gui-Qiang Chen, Monica Torres, and William P. Ziemer. Gauss–Green theorem for weakly differentiable vector fields, sets of finite perimeter, and balance laws. *Communications on Pure and Applied Mathematics*, 62(2): 242–304, 2009.
- [7] Giovanni E. Comi, Graziano Crasta, Virginia De Cicco, and Annalisa Malusa. Representation formulas for pairings between divergence-measure fields and BV functions. *Journal of Functional Analysis* 286.1: 110192, 2024.
- [8] Giovanni E. Comi, Virginia De Cicco, and Giovanni Scilla. Beyond BV : new pairings and Gauss–Green formulas for measure fields with divergence measure. *Preprint*, 2023.
- [9] Giovanni E. Comi and Kevin R. Payne. On locally essentially bounded divergence measure fields and sets of locally finite perimeter. *Advances in Calculus of Variations*, 13.2: 179–217, 2020.
- [10] Giovanni E. Comi and Monica Torres. One-sided approximation of sets of finite perimeter. *Rendiconti Lincei*, 28.1: 181–190, 2017.
- [11] Graziano Crasta and Virginia De Cicco. Anzellotti’s pairing theory and the Gauss–Green theorem. *Advances in Mathematics*, 343:935–970, 2019.
- [12] Graziano Crasta and Virginia De Cicco. An extension of the pairing theory between divergence-measure fields and BV functions. *J. Funct. Anal.* 276 (8):2605–2635, 2019.
- [13] Graziano Crasta and Virginia De Cicco. On the variational nature of the Anzellotti pairing. *arXiv preprint arXiv:2207.06469*, 2022.
- [14] Graziano Crasta, Virginia De Cicco, and Annalisa Malusa. Pairings between bounded divergence-measure vector fields and BV functions. *Advances in Calculus of Variations*, 2020.
- [15] Gianni Dal Maso, François Murat, Luigi Orsina, and Alain Prignet. Renormalized solutions of elliptic equations with general measure data. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 28(4):741–808, 1999.
- [16] Lawrence C. Evans and Ronald F. Gariepy. *Measure theory and fine properties of functions*. CRC press, 2015.
- [17] Nicola Fusco and Daniel Spector. A remark on an integral characterization of the dual of BV . *Journal of Mathematical Analysis and Applications*, 457.2: 1370–1375 (2018).
- [18] Hermano Frid. Divergence-measure fields on domains with Lipschitz boundary. In *Hyperbolic Conservation Laws and Related Analysis with Applications*, pages 207–225. Springer, 2014.
- [19] Enrico Giusti. Minimal surfaces and functions of bounded variation. *Monogr. Math.*, 80, 1984.
- [20] Gian Paolo Leonardi and Giovanni E. Comi. The prescribed mean curvature measure equation in non-parametric form. *Preprint*, 2024.
- [21] Gian Paolo Leonardi and Giorgio Saracco. Rigidity and trace properties of divergence-measure vector fields. *Adv. Calc. Var.*, 15(1):133–149, 2022.
- [22] Francesco Maggi. *Sets of finite perimeter and geometric variational problems: an introduction to Geometric Measure Theory*, volume 135 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory.

- [23] Norman G. Meyers and William P. Ziemer. Integral inequalities of Poincaré and Wirtinger type for BV functions. *American Journal of Mathematics*: 1345-1360, 1977.
- [24] Nguyen Cong Phuc and Monica Torres. Characterizations of the existence and removable singularities of divergence-measure vector fields. *Indiana University mathematics journal*, pages 1573–1597, 2008.
- [25] Nguyen Cong Phuc and Monica Torres. Characterizations of signed measures in the dual of BV and related isometric isomorphisms. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 17(1):385–417, 2017.
- [26] Christoph Scheven and Thomas Schmidt. BV supersolutions to equations of 1-Laplace and minimal surface type. *Journal of Differential Equations*, 261(3):1904–1932, 2016.
- [27] Christoph Scheven and Thomas Schmidt. On the dual formulation of obstacle problems for the total variation and the area functional. In *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, volume 35, pages 1175–1207. Elsevier, 2018.
- [28] Thomas Schmidt. Strict interior approximation of sets of finite perimeter and functions of bounded variation. *Proc. Amer. Math. Soc.*, 143(5):2069–2084, 2015.
- [29] Miroslav Šilhavý. Divergence measure fields and Cauchy's stress theorem. *Rendiconti del Seminario Matematico della Università di Padova*, 113:15–45, 2005.
- [30] Miroslav Šilhavý. The Gauss-Green theorem for bounded vector fields with divergence measure on sets of finite perimeter. *Indiana Univ. Math. J.* 72 (1): 29–42, 2023.
- [31] William P. Ziemer. *Weakly differentiable functions*, volume 120 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1989. Sobolev spaces and functions of bounded variation.

(Giovanni E. Comi) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA (ITALY)

Email address: giovannieugenio.comi@unibo.it

(Gian Paolo Leonardi) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TRENTO, VIA SOMMARIVE 14, IT-38123 POVO - TRENTO (ITALY)

Email address: gianpaolo.leonardi@unitn.it