

MINIMAL PERIODIC FOAMS WITH FIXED INRADIUS

ANNALISA CESARONI AND MATTEO NOVAGA

ABSTRACT. In this note we show existence and regularity of periodic tilings of the Euclidean space into equal cells containing a ball of fixed radius, which minimize either the classical or the fractional perimeter. We also discuss some qualitative properties of minimizers in dimensions 3 and 4.

CONTENTS

| | |
|---|----|
| 1. Introduction | 1 |
| 2. Lattices | 2 |
| 3. Existence of minimizers | 5 |
| 4. Regularity | 7 |
| 5. Comparison with other tilings | 8 |
| 5.1. The rhombic dodecahedral honeycomb and the Kelvin foam in \mathbb{R}^3 | 8 |
| 5.2. The 24-cell honeycomb in \mathbb{R}^4 | 10 |
| References | 11 |

1. INTRODUCTION

An important question, which has been considered since ancient times, is finding a partition of \mathbb{R}^2 into cells of equal area and with minimal perimeter, see [27]. This problem was eventually solved by Hales in [17], who proved that the hexagonal honeycomb is the unique minimizer. The analogous problem in dimension $N \geq 3$ is commonly known as Kelvin problem; the name comes from a famous conjecture due to Lord Kelvin [33], who claimed that in a tiling of \mathbb{R}^3 with cells of unit volume, the region with minimal surface area is given by a slightly modified truncated octahedron, also called Kelvin cell or tetrakaidecahedron. This conjecture was recently disproved by Weaire and Phelan, see in [35] and the discussion in Section 5.1. The Kelvin problem remains open for every $N \geq 3$, even if one restricts to lattice tilings, that is, to periodic partitions where all the cells are equal and given by a fundamental domain of a lattice. The counterexample by Weaire and Phelan does not apply to this case, so that the Kelvin foam could be a minimizer in \mathbb{R}^3 .

We notice that the lattice periodic Kelvin problem and, more generally, the isoperimetric problem for fundamental domains of a closed manifold, has been considered in the literature, starting from the paper [12], see also [25]. The concentration compactness argument for infinite partitions has been exploited in [10, 29, 30], whereas general perimeter functionals of local and nonlocal type have been considered in [9–11], see also [28] for a variant without periodicity.

1991 *Mathematics Subject Classification.* 49Q05 58E12 35R11.

Key words and phrases. Periodic tiling, perimeter, Kelvin foam, rhombic dodecahedron.

In this note we consider an isoperimetric problem which is related to the lattice periodic Kelvin problem. In particular we fix a perimeter functional, which is the classical perimeter or the fractional perimeter, and we show existence of periodic partitions of \mathbb{R}^N in equal cells with minimal perimeter, under the constraint that each cell contains a ball of fixed radius. We prove that this problem admits always a solution, by combining a concentration compactness argument (see [1, 23]), and a compactness theorem for lattices due to Mahler, see [24]. We provide also some basic regularity property for the minimizers, based on the observation that the partition of the space generated by a minimal cell is locally (almost) minimizing the perimeter, up to compact sufficiently small perturbations, see Section 4.

The problem we study in this note has been considered in dimension $N = 3$, restricting the class of admissible cells to parallelehedra, that is, adding a convexity constraint, see [3, 4, 21]. In this direction, Hales proved in [17] that among all Voronoi cells of lattices in \mathbb{R}^3 containing a ball of fixed radius the one with minimal perimeter is the regular rhombic dodecahedron, see Section 5.1. Nonetheless this polyhedron cannot be a solution of our isoperimetric problem, since it does not satisfy the necessary Plateau's conditions for local minimality, see Proposition 5.2. So, apart from dimension $N = 2$, the problem of identifying the shape of the minimizer remains open, even if one may formulate some conjectures, at least in dimension $N = 3, 4$, see Sections 5.1, 5.2.

The paper is organized as follows. In Section 2 we recall the basic definitions and properties of lattices in \mathbb{R}^N and we provide some important examples. Section 3 contains the main result of the paper, that is the existence of the solution of the isoperimetric problem, whereas in Section 4 we provide some regularity properties of the solution. Finally Sections 5.1 and 5.2 are devoted to a discussion of the case of dimensions $N = 3$ and $N = 4$ respectively.

Acknowledgements. The first author was supported by the PRIN Project 2022W58BJ5; the second author was supported by the PRIN 2022 Project 2022E9CF89 and by the MUR Excellence Department Project awarded to the Department of Mathematics of the University of Pisa. The authors are members of INDAM-GNAMPA.

2. LATTICES

A lattice is a discrete subgroup G of $(\mathbb{R}^N, +)$ of rank N . The elements of G can be expressed as $\sum k_i v_i$, for a given basis (v_1, \dots, v_N) of \mathbb{R}^N , with coefficients $k_i \in \mathbb{Z}$. Any two bases for a lattice G are related by a matrix with integer coefficients and determinant equal to ± 1 . Equivalently, every lattice can be viewed as a discrete group of isometries of \mathbb{R}^N .

The absolute value of the determinant of the matrix of any set of generators v_i of a lattice is uniquely determined and it is equal to $d(G) \in (0, +\infty)$, which we call *volume of the lattice* G . Equivalently, if we interpret G as a discrete group of isometries, $d(G)$ coincides with the volume of the quotient torus \mathbb{R}^N/G .

We recall this characterization of lattices, see [9, 10].

Lemma 2.1. *A closed subgroup G of $(\mathbb{R}^N, +)$ is discrete if and only if it does not contain a line.*

We let $\lambda(G) > 0$ be the minimal norm of the nonzero elements of G . In particular, for every $p, q \in G$, there holds that $|p - q| \geq \lambda(G)$. Other important values associated to a lattice G

are its *inradius* ρ_G and its *covering radius* r_G defined as

$$\begin{aligned}\rho_G &:= \sup\{r : \forall x \neq y \in G, B_r(x) \cap B_r(y) = \emptyset\} = \frac{\lambda(G)}{2} \\ r_G &:= \inf\{r : G + B_r = \mathbb{R}^N\}.\end{aligned}$$

The inradius is related to the problem of optimal packing of spheres, which can be stated as follows: find the arrangement of non-overlapping identical spheres (of radius r) in \mathbb{R}^N which maximizes the packing density, that is the proportion of space filled by the spheres. A lattice arrangement is an arrangement of spheres centered at the points of the lattice and with radius given by the inradius. The covering radius is related to the dual problem of the optimal packing, that is finding the most economical way to cover the space \mathbb{R}^N with overlapping spheres (of radius r). For more details on such problems we refer to the monograph [14, Chapters 1,2]. We recall that in dimension 2 the arrangement of spheres which solves all these problems is the regular hexagonal lattice arrangement, see Remark 2.8.

A lattice G is symmetric if it has a basis (v_1, \dots, v_N) of \mathbb{R}^N of generators such that $|v_i| = \lambda(G)$. It is clear that there exist non symmetric lattices, nonetheless it is always possible to choose a reduced set of generators for lattices (see [24, Theorem 1]), as follows:

Lemma 2.2. *There exists a dimensional constant C_N such that every lattice G admits a set of generators v_1, \dots, v_n with $\prod_{i=1}^N |v_i| \leq C_N d(G)$.*

In the following we present some examples of well known lattices (which are all symmetric).

Remark 2.3 (Root lattice and dual). The root lattice A_N is a N dimensional lattice which lies in the hypersurface $\sum_{i=1}^{N+1} x_i = 0$ of \mathbb{R}^{N+1} . It is defined as

$$A_N = \{z \in \mathbb{Z}^{N+1}, \sum_i z_i = 0\}.$$

The minimal vectors are permutations of $(1, -1, 0, \dots, 0)$ and the norm is $\sqrt{2}$, so the inradius $\rho_{A_N} = \frac{\sqrt{2}}{2}$.

The dual of the root lattice is A_N^* defined as $A_N^* = \cup_{i=0}^N A_N + [i]$, where $[i] \in \mathbb{R}^{n+1}$ has the first $N+1-i$ components equal to $\frac{i}{N+1}$ and the last i components equal to $\frac{i}{N+1} - 1$. The inradius of this lattice is $\rho_{A_N^*} = \frac{1}{2} \sqrt{\frac{N}{N+1}}$.

Remark 2.4 (Checkboard lattice). The checkboard lattice D_N is given by $\{z \in \mathbb{Z}^N, \sum_i z_i \text{ is even}\}$.

Its inradius is $\rho_{D_N} = \frac{\sqrt{2}}{2}$. The covering radius is $r_{D_N} = \frac{\sqrt{N}}{2}$ for $N \geq 4$.

When $N = 8$ the covering radius equal the minimum distance between two points of the lattice, and it is possible to add another copy of the lattice, so for $N \geq 8$ we define

$$D_N^+ = D_N \cup \left(D_N + \left(\frac{1}{2}, \dots, \frac{1}{2} \right) \right).$$

For $N = 8$, this lattice is denoted by E_8 . The inradius is given by $\rho_{D_N^+} = \frac{\sqrt{2}}{2}$.

Remark 2.5 (Leech lattice). The Leech lattice Λ_{24} is a lattice in \mathbb{R}^{24} , see [14, Chapter 4] for the precise construction. Its inradius is 1. It is the lattice which optimizes the sphere packing problem.

We recall the definition of fundamental domain of a lattice

Definition 2.6 (Fundamental Domain). A fundamental domain for the action of G is a set which contains almost all representatives for the orbits of G and such that the points whose orbit has more than one representative has measure zero, i.e. a measurable set $D \subseteq \mathbb{R}^N$ such that $|D + g \cap D| = 0$ for every $g \in G$ with $g \neq id$, and $|\mathbb{R}^N \setminus (D + G)| = 0$.

We will denote by \mathcal{D}_G the set of all fundamental domains of G .

If we fix a group G and consider **convex fundamental domains** associated to G , we obtain the class of **parallelohedra**, which are centrally symmetric polyhedra tiling the space by translations (see [26]), with at most $2(2^N - 1)$ facets. A special parallelohedron associated to the lattice G is its Voronoi cell, which is defined as

$$V_G := \{x \in \mathbb{R}^N : |x| \leq |x - g| \quad \forall g \in G, g \neq 0\}.$$

The distance between the center and every facet of the Voronoi cell associated to a lattice is at least equal to ρ_G , so in particular the Voronoi cell contains a ball of radius ρ_G . In a symmetric lattice the facets of the Voronoi cell are all the same distance from the origin.

Observe that not every parallelohedron is a Voronoi cell. In dimension 2, two-dimensional parallelohedra (parallelograms and centrally symmetric hexagons) are Voronoi cells if and only if they are inscribed in a circle.

Remark 2.7. The permutahedron P_N in \mathbb{R}^{N+1} is a N -dimensional polytope given by the convex envelope of the $(N + 1)!$ points obtained by permutations of the coordinates of the point

$$\left(-\frac{N}{2}, -\frac{N}{2} + 1, \dots, \frac{N}{2}\right).$$

It is contained in the hypersurface $(1, \dots, 1)^\perp$, it is the Voronoi cell of the lattice A_N^* , and has exactly $2(2^N - 1)$ facets.

Remark 2.8 (Hexagonal lattice in \mathbb{R}^2). The lattice A_2 is equivalent to the hexagonal lattice, whose Voronoi cell is the regular hexagon. The hexagonal lattice is generated by the vectors $(1, 0)$ and $(-1/2, \sqrt{3}/2)$. In this form, the determinant is $\sqrt{3}/2$ and the norm is 1. So the inradius is $1/2$, whereas the covering radius is $1/\sqrt{3}$.

Remark 2.9 (FCC and BCC lattices in \mathbb{R}^3). The checkboard lattice D_3 is called FCC (face-centered cubic) lattice, a family of generators is $(-1, -1, 0), (1, -1, 0), (0, 1, -1)$. The volume is 2, the norm is $\sqrt{2}$, and so the inradius is $1/\sqrt{2}$ and the covering radius is 1. The Voronoi cell of the FCC lattice is the rhombic dodecahedron. If centered at the origin it has 6 vertices $(\pm 1, 0, 0)$ and 8 vertices $(\pm 1/2, \pm 1/2, \pm 1/2)$.

The dual of the root lattice A_3^* is the BCC (body-centered cubic) lattice, a family of generators is $(2, 0, 0), (0, 2, 0), (1, 1, 1)$. The volume is 4, the norm is $\sqrt{3}$ and so the inradius is $\sqrt{3}/2$ and the covering radius is $\sqrt{5}/2$. The Voronoi cell of the BCC lattice is the truncated octahedron. If centered at the origin it has 24 vertices $(\pm 1, \pm 1/2, 0)$.

We introduce a notion of convergence of lattices see [8, 24]. Note that, if G is a lattice, then for every compact subset $K \subset \mathbb{R}^N$, the set $G \cap K$ is finite.

Definition 2.10. A sequence of lattices G_k converges to G if there exists for all k a set of generators g_k^i of G_k such that $g_k^i \rightarrow g^i$, and g^i is a set of generators of the lattice G .

A sequence of lattice G_k converges in the Kuratowski sense to G_i , if

$$G = \{g \in \mathbb{R}^N \mid \limsup_{i \rightarrow +\infty} d(g, G_i) = 0\}.$$

Note that G is a closed subgroup of $(\mathbb{R}^N, +)$. Actually, the two notion of convergence are equivalent, see [8, Section V.3, Theorem 1].

We recall the following compactness theorem for lattices due to Mahler [24, Theorem 2] (see also [8, Chapter V]).

Theorem 2.11. *Let G_i , $i \in \mathbb{N}$, be a sequence of lattices and assume that there exist two constants $k, \delta > 0$ such that $\lambda(G_i) \geq \delta > 0$ for all i and $d(G_i) = |\mathbb{R}^N/G_i| \leq m$ for all i and for some $m \in (0, +\infty)$. Then there exists a subsequence G_{i_n} and a lattice G such that $G_{i_n} \rightarrow G$, and $\lambda(G) \geq \delta$, $d(G) \leq m$.*

As a byproduct on this compactness theorem in [10, Lemma 3.7], see also [9, Theorem 2.10], it has been proved the following result.

Theorem 2.12. *Let G_h be a sequence of lattices with $d(G_h) \leq m$ for every h and for some $m \in (0, +\infty)$. Then, up to subsequences, $G_h \rightarrow G$ in the Kuratowski sense, where G is either a lattice with $d(G) \leq m$ or a closed group which contains a line. In the second case, every sequence D_h of fundamental domains for G_h converges to \emptyset in $L^1_{loc}(\mathbb{R}^N)$.*

3. EXISTENCE OF MINIMIZERS

We fix a general class of perimeter functionals $\text{Per}(E)$ defined on measurable sets $E \subseteq \mathbb{R}^N$, which are of two possible types.

- Classical perimeter: for every measurable set $E \subseteq \mathbb{R}^N$, we define

$$(1) \quad \text{Per}(E) = \int_{\partial^* E} dH^{n-1}(x)$$

where $\partial^* E$ is the reduced boundary of E and $\nu(x)$ is the (measure theoretic) exterior normal at E in x (see [23]).

- Fractional perimeter: let $s \in (0, 1)$ and for every measurable set $E \subseteq \mathbb{R}^N$, we define

$$(2) \quad \text{Per}(E) = \int_E \int_{\mathbb{R}^N \setminus E} \frac{1}{|x-y|^{N+s}} dx dy.$$

We fix a constant $m > 0$ and we introduce the following class of lattices

$$(3) \quad \mathcal{G}_m = \{G \text{ lattice in } \mathbb{R}^N, \text{ such that } \rho_G = m\}$$

and moreover for a given lattice $G \in \mathcal{G}_m$ we consider the class of fundamental domains which contain a ball of radius m :

$$(4) \quad \mathcal{D}_G^m = \{D \in \mathcal{D}_G, \text{ such that } \exists x \in D, |B_m(x) \setminus D| = 0\}$$

Note that this class is always not empty, since the Voronoi cell associated to a lattice $G \in \mathcal{G}_m$ is in \mathcal{D}_G^m .

First of all we observe that, for every fixed group G , every fundamental domain which minimizes the classical perimeter in (1) is indecomposable.

Lemma 3.1. *Let G be a lattice and $E \in \mathcal{D}_G$ be a solution to the isoperimetric problem*

$$\inf_{D \in \mathcal{D}_G} \text{Per}(D),$$

where Per is the classical perimeter. Then E is indecomposable.

Proof. Let $E = \cup_i E^i$ where E^i are the indecomposable components of E (see [2]), and assume by contradiction that there exists i such that $|E^i| \neq 0$ and $|F| \neq 0$ where $F := \cup_{j \neq i} E^j$.

Then for every $g \in G$, $\mathcal{H}^{N-1}(\partial^* E^i \cap \partial^*(F+g)) = 0$, where ∂^* denotes the reduced boundary of a set (see [23]): if it were not the case for some $g \in G$, then $E^i \cup (F+g)$ would be a fundamental domain with $\text{Per}(E^i \cup (F+g)) < \text{Per}(E)$. Let us consider the measurable sets $E^i + G = \cup_{g \in G} E^i + g$ and $F + G = \cup_{g \in G} F + g$. We have that $|(E^i + G) \cap (F + G)| = 0$, $|\mathbb{R}^N \setminus ((E^i + G) \cup (F + G))| = 0$, $|E^i + G|, |F + G| > 0$ and $H^{N-1}(\partial^*(E^i + G) \cap \partial^*(F + G)) = 0$, which is not possible. \square

We now prove the following existence result.

Theorem 3.2. *Let Per be a perimeter functional as in (1), (2). Then there exists a solution of the isoperimetric problem*

$$(5) \quad \inf_{G \in \mathcal{G}_m} \inf_{D \in \mathcal{D}_G^m} \text{Per}(D).$$

Proof. The argument is divided in three steps.

Step 1: First of all we consider a fixed $G \in \mathcal{G}_m$, as defined in (3) and we show that there exists $E \in \mathcal{D}_G$ such that

$$\text{Per}(E) = \inf_{D \in \mathcal{D}_G^m} \text{Per}(D).$$

Observe that since $G \in \mathcal{G}_m$, $d(G) = d_m > 0$. The existence of a fundamental domain of minimal perimeter for a fixed lattice has been obtained in [10, Theorem 3.2], with a concentration compactness argument dating back to Almgren and by lower semicontinuity of the perimeter functional (see [1] and the monograph [23]).

In particular given a minimizing sequence of fundamental domains $D_k \in \mathcal{D}_G^m$, then there exist $g_k^i \in G$, for $i \in I \subseteq \mathbb{N}$, and $E^i \subseteq \mathbb{R}^N$ such that $|g_k^i - g_k^j| \rightarrow +\infty$ if $j \neq i$ as $k \rightarrow +\infty$, and $D_k + g_k^i \rightarrow E^i$ locally in L^1 as $k \rightarrow +\infty$. Moreover $\cup_i E^i \in \mathcal{D}_G$: so $|\cup_i E^i| = |D_k| = d_m$ and $\text{Per}(\cup_i E^i) = \sum_i \text{Per}(E^i) = \inf_{D \in \mathcal{D}_G^m} \text{Per}(D)$.

Step 2: We prove that $E \in \mathcal{D}_G^m$ (so it contains a ball of radius m).

If the perimeter functional is the fractional perimeter (2) then there exists a unique i such that $|E^i| \neq 0$. This is an immediate consequence of the fact that $\text{Per}(\cup_i E^i) < \sum_i \text{Per}(E^i)$ if there are at least two elements $|E^i|, |E^j| \neq 0$, and on the other hand by Step 1 we have that $\text{Per}(\cup_i E^i) = \sum_i \text{Per}(E^i)$ (see [10, Remark 3.6]). Since $D_k \rightarrow E^i$ locally in L^1 and $|D_k| = |E^i|$, we conclude that actually the convergence is in $L^1(\mathbb{R}^N)$, which implies that $E^i \in \mathcal{D}_G^m$.

If the perimeter functional is the local perimeter (1), by Lemma 3.1 E is indecomposable. So $D_k \rightarrow E$ in $L^1(\mathbb{R}^N)$, and we conclude as above.

Step 3: we minimize among all possible lattices $G \in \mathcal{G}_m$.

Let us fix a minimizing sequence, that is a sequence of lattices $G_n \in \mathcal{G}_m$ and a sequence of fundamental domains $D_n \in \mathcal{D}_{G_n}^m$. So $\text{Per}(D_n) \leq \inf_{G \in \mathcal{G}_m} \inf_{D \in \mathcal{D}_G^m} \text{Per}(D) + 1 := C$, and by the isoperimetric inequality this implies $|D_n| \leq K$, and so $d(G_n) \leq K$ for every n . Moreover by the constraint $|D_n| \geq |B_m| = m^N \omega_N$.

By Theorem 2.12, up to subsequences either $G_n \rightarrow G$ where $d(G) \leq K$, or $D_n \rightarrow \emptyset$ in L^1_{loc} .

By the same argument of concentration compactness recalled in Step 1, see [10, Lemma 3.4], since D_n are measurable sets with $|D_n| \geq m^N \omega_N$ and $\text{Per}(D_n) \leq C$, up to passing to a subsequence, there exist $z_h^i \in \mathbb{Z}^N$ ($i \in \mathbb{N}, h \in \mathbb{N}$), with $|z_h^i - z_h^j| \rightarrow +\infty$ as $h \rightarrow +\infty$ for $i \neq j$, and a family (D^i) of measurable sets in \mathbb{R}^N such that $D_h - z_h^i \rightarrow D^i$ in $L^1_{loc}(\mathbb{R}^N)$ and

$\sum_i |D^i| = m$. This implies that it is not possible that $D_n \rightarrow \emptyset$ locally in L^1 , and so, up to a subsequence, $G_n \rightarrow G$ where G is a lattice.

Then we proceed with a similar proof as in Step 1, see the details in [9, Lemma 3.1, Theorem 1.2]. \square

For a fundamental domain D associated to a lattice we can introduce the ratio

$$\mathcal{I}(D) = \frac{\text{Per}D}{\rho_D^2} \quad \text{where } \rho_D = \max\{r \geq 0 : \exists x \in D, |D \setminus B(x, r)| = 0\}.$$

Note that problem (5) is equivalent to the following minimum problem:

$$(6) \quad \min_{G \text{ lattice in } \mathbb{R}^3} \min_{D \in \mathcal{D}_G} \mathcal{I}(D).$$

Remark 3.3. The same result as Theorem 3.2 holds under a more general class of constraints.

Let us fix a nonnegative functional \mathcal{F} defined on measurable sets, a constant $m > 0$ and the following class

$$\mathcal{G}_m = \{G \text{ lattice in } \mathbb{R}^N, \text{ such that } \exists D \in \mathcal{D}_G, \mathcal{F}(D) = m\}.$$

The isoperimetric problem reads:

$$(7) \quad \inf_{G \in \mathcal{G}_m} \inf_{D \in \mathcal{D}_G, \mathcal{F}(D)=m} \text{Per}(D).$$

The same argument as in the proof of Theorem 3.2 gives the existence of a solution to (7) if we assume that \mathcal{F} satisfies the following conditions:

- \mathcal{F} is continuous in L^1 , that is if $E_i \rightarrow E$ in measure (that is $\chi_{E_i} \rightarrow \chi_E$ in L^1), then $\mathcal{F}(E_i) \rightarrow \mathcal{F}(E)$;
- \mathcal{F} is nondegenerate in the sense that for any $C > 0$ there exists a constant $k(C) = k > 0$ such that if $\mathcal{F}(E) \geq C$ for $E \in \mathcal{M}$, then $|E| \geq k$.

The case of the inradius is obtained by considering

$$\mathcal{F}(E) = \sup\{r > 0, \text{ such that } |B_r(x) \setminus E| = 0 \text{ for some } x \in E\}.$$

Another possible functional satisfying the previous assumptions is the Riesz energy (see [22]):

$$\mathcal{F}(E) := \int_E \int_E \frac{1}{|x - y|^{N-\alpha}} dx dy, \quad \text{for some } \alpha \in (0, N).$$

4. REGULARITY

In order to study the regularity of the minimal fundamental domain we introduce the notion of partitions.

Definition 4.1. A partition of \mathbb{R}^N is a collection of measurable subsets $\{E_k\}_{k \in \mathbb{I}}$, where \mathbb{I} is either a finite or a countable set of ordered indices, such that

- (1) $|E_k| > 0$ for all k ,
- (2) $|E_k \cap E_j| = 0$ for all $k \neq j$,
- (3) $|\mathbb{R}^N \setminus \cup_k E_k| = 0$.

A fundamental domain D of a lattice G naturally induces the G -periodic partition $\{E_g\}_{g \in G}$, where $E_g = gD$. Moreover if D is bounded, then the associated partition is locally finite.

Definition 4.2. We say that a partition $\{E_k\}_k$ is Λ -minimal in an open set $A \subset \mathbb{R}^N$, for some $\Lambda \geq 0$, if $\sum_k \text{Per}(E_k, A) < +\infty$ and

$$\sum_k \text{Per}(E_k, A) \leq \sum_k [\text{Per}(F_k, A) + \Lambda \mu(E_k \Delta F_k)],$$

for every partition $\{F_k\}_k$ of X such that $E_k \Delta F_k \Subset A$ for all k .

We say that the partition is (Λ, r) -minimal (see [23]) for some $\Lambda \geq 0$ and $r > 0$, if it is Λ -minimal in $B_r(x)$ for all $x \in \mathbb{R}^N$.

Theorem 4.3. *Let D be a solution to (5) with $\rho_G = 1$ and $B \subset D$ is the ball of radius 1. Then the partition $D + G$ is (Λ, r) -minimal in $\mathbb{R}^N \setminus (B + G)$ for every $r \leq r_0 < 1 = \rho_G$, with $\Lambda = 0$ in the case of the local perimeter, and $\Lambda = \frac{\omega_N}{s(2-2r_0)^s} = \int_{|h|>2-2r_0} |h|^{-s-N} dh$, in the case of the nonlocal perimeter.*

Then, there exists a closed singular set $\Sigma \neq \emptyset$ with $\mathcal{H}^{N-1}(\Sigma) = 0$, such that $\partial D \setminus \Sigma$ is a $C^{1, \frac{1+s}{2}}$ hypersurface, where $s \in (0, 1)$ is the order of the fractional perimeter in the case (2) and $s = 1$ in the case of local perimeter (1).

Proof. The first part of the statement follows the same argument of the proof of (Λ, r) -minimality in \mathbb{R}^N of partitions associated to fundamental domains with minimal perimeter (without assuming the constraint on the inradius), given in [10, Proposition 4.1].

As for the regularity of the boundary of D , we may refer to classical regularity results about (local) minimal surfaces with obstacles [6, 19], whereas for nonlocal minimal surfaces with obstacle we refer to [7, Theorem 1.2]. \square

5. COMPARISON WITH OTHER TILINGS

5.1. The rhombic dodecahedral honeycomb and the Kelvin foam in \mathbb{R}^3 . In this section we consider the isoperimetric problem (5) for the classical perimeter in dimension $N = 3$. A restricted version of this isoperimetric problem has been already discussed in the literature, by considering as admissible competitors only convex fundamental domains containing a ball of given radius. In particular in [4] (see also [21]) it has been posed the problem of finding the parallelohedron with minimal perimeter containing a ball of radius 1, stating the following conjecture:

Rhombic Dodecahedral Conjecture: The perimeter of any parallelohedron containing a ball of radius 1 is at least that of a regular rhombic dodecahedron.

If we further restrict the class of competitors, admitting only Voronoi cells containing a ball of radius 1, then the conjecture is true, and it has been proved by Hales [17] in a more general setting, admitting also non periodic configurations:

Theorem 5.1 (Strong Dodecahedral Theorem). *The perimeter of any Voronoi cell associated to a packing of spheres of radius 1 in \mathbb{R}^3 is at least that of a regular rhombic dodecahedron.*

Going back to our problem, the first observation is that the regular rhombic dodecahedron cannot be a solution.

Proposition 5.2. *The regular rhombic dodecahedron is not a solution to (5) in dimension $N = 3$ with the standard perimeter.*

Proof. Let D be a solution of the isoperimetric problem (5) for the standard perimeter and inradius 1 in dimension $N = 3$. Let us fix $x \in \mathbb{R}^N$ and $r < 1 = \rho_G$ and consider the partition $D + G$ inside a ball $B_r(x)$. By Theorem 4.3, this partition inside the ball is locally minimal for the perimeter and so either it is empty or a minimal cone in \mathbb{R}^3 . The complete classification of minimal singular cones is available in dimensions 3 (see [32]): three half-hyperplanes meeting at 120 degrees or the cone over the 1-skeleton of a tetrahedron. So every minimal cone either is an hyperplane or has to coincide with one of the previous singular cones: these necessary conditions for minimality are called Plateau's conditions.

Regular rhombic dodecahedron has two types of vertices which are threefold and fourfold. If we look at the partition generated by a regular rhombic dodecahedron inside a ball $B_r(x)$ with $r < 1$ and $x \in \mathbb{R}^N$ one of the fourfold vertices, we obtain a cone with more than 4 chambers, so it cannot coincide with any of the previous configurations. On the other hand if x is one of the threefold vertices, we obtain the cone over the 1-skeleton of a cube, which again is not minimal. So regular rhombic dodecahedron cannot be minimal. \square

Remark 5.3. The same argument as in Proposition 5.2 shows that the solution to (5) cannot be a parallelohedron, since parallelohedra never satisfy Plateau's conditions at vertices and edges.

Besides the regular rhombic dodecahedral honeycomb there is another important structure arising in the Kelvin problem, which is the variational problem corresponding to the surface minimizing partition into cells of equal volume [33].

The construction of Lord Kelvin in [33] produces a partition of the space which is a critical point for the Kelvin problem. As describes in detail in [35, Section 7], Lord Kelvin first considered the partition generated by the regular rhombic dodecahedron, which as shown in Proposition 5.2 cannot be a local minimizer. The idea was then to substitute the partition locally around any fourfold vertex with the Plateau minimal partition with boundary given by the 1-skeleton of a cube. This introduces a new flat facet with four curved edges, which may be directed along any of the three cubic axes (and one may choose the same cubic axis for all of them). So he was attaching many copies of Plateau's wire cube together, edge to edge and in the rhombohedral cell a square face is introduced at the top and bottom and new horizontal edges appear at the four fourfold vertices that lie on the equator. These are the edges of square faces belonging to adjacent cells. The resulting cell, that we may call **anisotropic Kelvin cell**, is a candidate structure which satisfies Plateau's conditions and is a fundamental domain for the FCC lattice, like the regular rhombic dodecahedron.

Since this cell is not isotropic, Lord Kelvin further modified it by changing the ratio between the axis in the Plateau's wire cube, taking as a boundary to construct the minimal surface a parallelepiped and not a cube. By varying this ratio from 1 to $1/\sqrt{2}$ he obtained an isotropic cell, that we call **Kelvin cell** \mathcal{T}_K . This cell actually is space filling when centered at the points of the BCC lattice, even if it is not the Voronoi cell of such lattice (which is the regular truncated octahedron): in particular it is a fundamental domain of the BCC lattice and it contains a ball of radius less or equal to the ball contained in the regular truncated octahedron of the same volume. Indeed it can be also obtained as a small deformation of the faces of the regular truncated octahedron, see [31] for a description of this construction, and it has six planar quadrilateral faces and eight nonplanar hexagons of zero curvature. If we denote with \mathcal{T} the truncated octahedron we get that

$$\rho_{\mathcal{T}_K} \leq \rho_{\mathcal{T}}$$

and so

$$(8) \quad \mathcal{I}(\mathcal{T}_K) \geq \mathcal{I}(\mathcal{T}) \frac{\text{Per}(\mathcal{T}_K)}{\text{Per}(\mathcal{T})} \sim 0.998 \mathcal{I}(\mathcal{T}),$$

by using the computation of the ratio between the perimeter of the Kelvin cell and of the regular truncated octahedron obtained in [31].

Proposition 5.4. *The Kelvin cell \mathcal{T}_K is not a solution to (5).*

Proof. In order to show the result, it is sufficient to show that $\mathcal{I}(\mathcal{T}_K) > \mathcal{I}(\mathcal{D})$, where \mathcal{D} is the regular rhombic dodecahedron.

We have that the perimeter of the regular rhombic dodecahedron \mathcal{D} of inradius 1 is given by $12\sqrt{2}$, see [3, 17], so in particular

$$\mathcal{I}(\mathcal{D}) = 12\sqrt{2} \sim 16.97.$$

On the other hand the inradius of the regular truncated octahedron \mathcal{T} of volume 1 is given by $\rho = \frac{\sqrt{3}}{2^{\frac{3}{4}}}$ (see [14, Chapter 2]) and its perimeter $\frac{6\sqrt{3}+3}{2^{\frac{3}{4}}}$ (see [31]). So

$$\mathcal{I}(\mathcal{T}) = \frac{6\sqrt{3} + 3 \cdot 2^{\frac{10}{3}}}{2^{\frac{4}{3}} \cdot 3} = 4(2\sqrt{3} + 1) = 17.856,$$

hence by (8) we conclude

$$\mathcal{I}(\mathcal{T}_K) \geq 0,998 \cdot 17.856 \sim 17.82 > \mathcal{I}(\mathcal{D}).$$

□

It remains open the problem of identify a solution to problem (5) in \mathbb{R}^3 . Since the Kelvin cell is not a solution, and in view of the Strong Dodecahedral Theorem proved by Hales, one might be led to state the following:

FCC Conjecture: the optimal lattice for problem (5) is FCC.

5.2. The 24-cell honeycomb in \mathbb{R}^4 . In dimension $N = 4$, there exists a well known regular tessellation, which is the 24-cell honeycomb. The 24-cell is the convex hull of its vertices which can be described as the 24 coordinate permutations of $(\pm 1, \pm 1, 0, 0)$. In this frame of reference the 24-cell has inradius $\frac{\sqrt{2}}{2}$. Up to a rescaling, the 24-cell is the Voronoi cell of the D_4 lattice.

If a sphere is inscribed in each 24 cell of the 24-cell honeycomb, the resulting arrangement is the densest known regular sphere packing in four dimensions, with kissing number 24, even if the sphere packing problem is still unsolved in dimension 4. This suggests that the 24-cell could be a solution to problem (5).

In this direction, we observe that the 24-cell honeycomb satisfies the following local minimality property, which is a necessary condition for being a solution to problem (5), as stated in Theorem 4.3.

Proposition 5.5. *The 24-cell honeycomb is a minimizer for the perimeter in every ball $B_r(x)$, with $x \in \mathbb{R}^4$ and $0 < r < \frac{\sqrt{2}}{2}$. In particular, it is a critical point of problem (5).*

Proof. The thesis follows from the fact that inside a ball $B_r(x)$ with r smaller than the inradius, the 24-cell honeycomb is either empty or coincides, up to a translation, with one of the following cones in \mathbb{R}^4 : a hyperplane, three half-hyperplanes meeting at 120 degrees, the cone over the 1-skeleton of a tetrahedron times \mathbb{R} , the cone over the 2-skeleton of a hypercube.

Thanks to the results by Taylor [32] and Brakke [5] it is known that such cones define partitions of \mathbb{R}^4 which are minimal for the perimeter under compact perturbations. \square

Due to the properties of the 24-cell, in analogy with the rhombic dodecahedron conjecture in \mathbb{R}^3 , we may state the following conjecture:

24-cell Conjecture: The perimeter of any fundamental domain of a lattice in \mathbb{R}^4 , containing a ball of radius 1, is at least that of a 24-cell of inradius 1.

REFERENCES

- [1] F. J. Almgren, *Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints*, Mem. Amer. Math. Soc. **4** (1976), no. 165.
- [2] L. Ambrosio, V. Caselles, S. Masnou, and J.-M. Morel, *Connected components of sets of finite perimeter and applications to image processing*, J. Eur. Math. Soc. (JEMS) **3** (2001), no. 1, 39–92.
- [3] K. Bezdek, *On rhombic dodecahedra*, Beiträge Algebra Geom. **41** (2000), no. 2, 411–416.
- [4] ———, *On a strong version of the Kepler conjecture*, Mathematika **59** (2013), no. 1, 23–30.
- [5] K. A. Brakke, *Minimal cones on hypercubes*, J. Geom. Anal. **1** (1991), no. 4, 329–338.
- [6] L. A. Caffarelli, *The obstacle problem revisited*, J. Fourier Anal. Appl. **4** (1998), no. 4–5, 383–402.
- [7] L. A. Caffarelli, D. De Silva, and O. Savin, *Obstacle-type problems for minimal surfaces*, Comm. Partial Differential Equations **41** (2016), no. 8, 1303–1323.
- [8] J. W. S. Cassels, *An introduction to the geometry of numbers*, Classics in Mathematics, Springer-Verlag, Berlin, 1997. Corrected reprint of the 1971 edition.
- [9] A. Cesaroni, I. Fragalà, and M. Novaga, *Lattice tilings minimizing nonlocal perimeters*, Commun. Contemp. Math. (to appear).
- [10] A. Cesaroni and M. Novaga, *Periodic partitions with minimal perimeter*, Nonlinear Anal. **243** (2024), Paper No. 113522, 16 pp.
- [11] ———, *Minimal periodic foams with equal cells*, <https://arxiv.org/abs/2302.07112> (2023).
- [12] J. Choe, *On the existence and regularity of fundamental domains with least boundary area*, J. Differential Geom. **29** (1989), no. 3, 623–663.
- [13] M. Colombo and F. Maggi, *Existence and almost everywhere regularity of isoperimetric clusters for fractional perimeters*, Nonlinear Anal. **153** (2017), 243–274.
- [14] J.H. Conway and N.J.A. Sloane, *Sphere packings, lattices and groups. Third edition*, Grundlehren der mathematischen Wissenschaften, vol. 290, Springer-Verlag, New York, 1999.
- [15] T.C. Hales, *The honeycomb conjecture*, Discrete Comput. Geom. **25** (2001), no. 1, 1–22.
- [16] T. C. Hales, *A proof of the Kepler conjecture*, Ann. of Math. (2) **162** (2005), no. 3, 1065–1185.
- [17] ———, *Dense sphere packings: a blueprint for formal proofs*, London Math. Soc. Lecture Note Series, vol. 400, Cambridge University Press, Cambridge, 2012.
- [18] Á. G. Horváth, *Extremal polygons with minimal perimeter*, Period. Math. Hungar. **34** (1997), no. 1–2, 83–92. 3rd Geometry Festival: an International Conference on Packings, Coverings and Tilings (Budapest, 1996).
- [19] D. Kinderlehrer, *How a minimal surface leaves an obstacle*, Bull. Amer. Math. Soc. **78** (1972), 969–970.
- [20] G. Kindler, A. Rao, R. O’Donnell, and A. Wigderson, *Spherical cubes: Optimal foams from computational hardness amplification*, Communications of the ACM **55** (2012), no. 10, 90–97.
- [21] Zsolt Lángi, *An isoperimetric problem for three-dimensional parallelotopes*, Pacific J. Math. **316** (2022), no. 1, 169–181.
- [22] Elliott H. Lieb and Michael Loss, *Analysis*, Graduate Studies in Mathematics, vol. 14, American Mathematical Society, Providence, RI, 2001.
- [23] F. Maggi, *Sets of finite perimeter and geometric variational problems*, Cambridge Studies in Advanced Mathematics, vol. 135, Cambridge University Press, Cambridge, 2012.
- [24] K. Mahler, *On lattice points in n -dimensional star bodies. I. Existence theorems*, Proc. Roy. Soc. London Ser. A **187** (1946), 151–187.
- [25] B. Martelli, M. Novaga, A. Pluda, and S. Riolo, *Spines of minimal length*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **17** (2017), no. 3, 1067–1090.
- [26] P. McMullen, *Convex bodies which tile space by translation*, Mathematika **27** (1980), no. 1, 113–121.
- [27] F. Morgan, *The hexagonal honeycomb conjecture*, Trans. Amer. Math. Soc. **351** (1999), no. 5, 1753–1763.

- [28] F. Nobili and M. Novaga, *Lattice tilings with minimal perimeter and unequal volumes*, arxiv preprint <https://arxiv.org/abs/2406.12461> (2024).
- [29] M. Novaga, E. Paolini, E. Stepanov, and V.M. Tortorelli, *Isoperimetric clusters in homogeneous spaces via concentration compactness*, *J. Geom. Anal.* **32** (2022), no. 11, Paper No. 263.
- [30] ———, *Isoperimetric planar clusters with infinitely many regions*, *Netw. Heterog. Media* **18** (2023), no. 3, 1226–1235.
- [31] H.M Princen and P Levinson, *The surface area of Kelvin's minimal tetrakaidecahedron: The ideal foam cell (?)*, *Journal of Colloid and Interface Science* **120** (1987), no. 1, 172-175.
- [32] J. E. Taylor, *The structure of singularities in solutions to ellipsoidal variational problems with constraints in \mathbb{R}^3* , *Ann. of Math. (2)* **103** (1976), no. 3, 541–546.
- [33] W. Thomson (Lord Kelvin), *On the division of space with minimum partitional area*, *Acta Math.* **11** (1887), no. 1-4, 121–134.
- [34] D. Weaire (ed.), *The Kelvin problem*, Taylor & Francis, London, 1996. Foam structures of minimal surface area.
- [35] ———, *Kelvin's foam structure: a commentary*, *Philosophical Magazine Letters* **88** (2008), no. 2, 91–102.

DIPARTIMENTO DI MATEMATICA TULLIO LEVI-CIVITA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35131 PADOVA, ITALY

Email address: `annalisa.cesaroni@unipd.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY

Email address: `matteo.novaga@unipi.it`