

***H*-COMPACTNESS FOR NONLOCAL LINEAR OPERATORS IN FRACTIONAL DIVERGENCE FORM**

MAICOL CAPONI, ALESSANDRO CARBOTTI, AND ALBERTO MAIONE

ABSTRACT. We study the *H*-convergence of nonlocal linear operators in fractional divergence form, where the oscillations of the matrices are prescribed outside the reference domain. Our compactness argument bypasses the failure of the classical localisation techniques that mismatch with the nonlocal nature of the operators involved. If symmetry is also assumed, we extend the equivalence between *H*-convergence of the operators and Γ -convergence of the associated energies. As a consequence, we establish the uniqueness of the *H*-limit in the symmetric case.

1. INTRODUCTION

At the beginning of the '70s the French school of J.-L. Lions set the stage for the mathematical theory of the *homogenisation of composite materials*, made by the superposition of sheets of different materials, or considering homogeneous materials with holes filled by another material. For a given source term f , this problem is mathematically modelled by a symmetric matrix-valued function $A(x)$, identifying the different materials at each point x , and expressed through partial differential equations in divergence form

$$(1.1) \quad -\operatorname{div}(A(x)\nabla u(x)) = f(x).$$

An equivalent statement of problem (1.1) considers the momentum $p(x) := A(x)\nabla u(x)$ and is formulated as

$$-\operatorname{div}(p(x)) = f(x).$$

Such equations appear in many applications in physics and engineering: in the *equation of the electrostatics*, u is the electric potential, p is the electric displacement, and A is the dielectric constant, while, in the *equation of magnetostatics*, u is the magnetic potential, p is the magnetic induction, and A is the magnetic permeability. Other examples are the *equation of the time-independent heat transfer*, where u is the temperature, p is the heat flux, and A represents the thermal conductivity and, in the vectorial case, the *equation of the linear elasticity for composite materials*, where u is the displacement field, p (usually denoted by σ) is the Cauchy stress tensor, and A represents the elasticity tensor.

From a numerical point of view, when the coefficients of the matrix A oscillate too fast, it can be very difficult to deal with the above equations, for example when the heterogeneous material has a periodic structure. A possible way to overcome this problem is to approximate the solutions by the (unique) solution of a limit problem, much easier to deal with numerically. This is the so-called *homogenised* problem, in which the matrix A is independent on x .

The first relevant contributions to this theory are by Sánchez-Palencia [29, 30, 31], through applications of the *asymptotic expansions* method. It consists in approximating solutions of (1.1) with a series depending on the layers of the material, and then estimating the resulting error.

A more general approach, introduced by Spagnolo in the pioneering works [37, 38, 39], is the so-called *G*-convergence of linear partial differential equations of the form

$$(1.2) \quad -\operatorname{div}(A_h(x)\nabla u(x)) = f(x), \quad h \in \mathbb{N}.$$

The *G*-convergence theory studies the asymptotic behaviour of the solutions of problems (1.2), assuming very mild hypotheses on the coefficients of any matrix A_h , and it is based on the distributional nature of the problems (1.2) and some standard tools of functional analysis.

2020 *Mathematics Subject Classification.* (Primary) 35B40, (Secondary) 35B27, 35R11, 49J45, 74S40.

Key words and phrases. *H*-convergence, Γ -convergence, fractional operators, Riesz potentials.

Nevertheless, at its early stage, it was not clear what properties the limit with respect to the G -convergence would have satisfied. A first characterisation of the limit operator (the so-called G -limit) was only later obtained by De Giorgi and Spagnolo [18], who show that the G -limit still acts as a divergence form operator depending on a limit homogenised matrix A_∞ (sometimes called *effective* matrix and denoted by A^{eff}). We underline that, whenever $A_h(x) = A(hx)$, we recover the former theory of the homogenisation of composite materials but, differently from it, the limit matrix of the G -convergence, $A_\infty = A_\infty(x)$, may in general still be x -dependent.

By its very nature, the G -convergence is sensible according to the class of operators considered and, without further requirements, it appears incomplete when dealing with not necessarily symmetric matrices, needed in applications involving e.g. porous media. Tartar observes in [42] that the divergence operator does not in fact recognise perturbations from skew-symmetric matrices leading, in this case, to the non-uniqueness of the matrix A_∞ .

This lack of uniqueness is bypassed by Tartar [41] and Murat [26, 27] at the end of the '70s, and this theory is nowadays called H -convergence, where H stands for *homogenisation*. The main difference with the former G -convergence naturally appears in the non-symmetric case: the uniqueness of the homogenised matrix A_∞ is now guaranteed, whenever the sequence of the momenta $p_h = A_h \nabla u_h$ converges (in the weak topology) to the momentum of the limit solution $p_\infty = A_\infty \nabla u_\infty$.

In the last decade there has been an increasing interest in extending these results to a *nonlocal* framework, motivated by the huge number of applications in fluid mechanics, image denoising, nonlinear elasticity, nonlocal minimal surfaces, anomalous diffusion, and stable Lévy processes (see e.g. [10, 11] and the references therein). First attempts to deal with nonlocal H -convergence appear in [4, 19], where the authors study scalar perturbations of the fractional p -Laplace operator, and in [9], regarding the H -convergence of fractional powers of elliptic operators in divergence form. We also mention [6, 20, 28] for the homogenisation of more general nonlocal energies.

All the aforementioned contributions in nonlocal H -convergence consider, however, only the case of scalar weights, in contrast to the classical results by Tartar [42]. On the other hand, the case of matrix weights is of great interest in applications, as it allows to study anisotropic heterogeneous materials. We observe that it is not immediately clear how to get a momentum operator in problems involving Gagliardo seminorms, since the latter are defined via integration of scalar energies.

We hence wonder whether it might be possible to formulate a nonlocal H -convergence-type problem à la Tartar. To this aim, we exploit a suitable notion of fractional-order divergence div^s and fractional-order gradient ∇^s (see Definition 2.2), in a way that the fractional divergence acts on the nonlocal momentum $p(x) = A(x) \nabla^s u(x)$. The construction of this class of fractional-order operators is explained in details in Section 2, and relies on the pioneering contributions by Shieh and Spector [34, 35], Šilhavý [45], and the subsequent works [8, 12, 13, 14].

This leads to the study of the H -convergence of the following sequence of elliptic problems

$$(P_h^f) \quad \begin{cases} -\text{div}^s(A_h \nabla^s u_h) = f & \text{in } \Omega, \\ u_h = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

$h \in \mathbb{N}$, on the same spirit of Tartar and Murat.

When dealing with the asymptotics of problems (P_h^f) , there are two main issues to be taken into account. The first one is to overcome the localisation techniques needed in the H -convergence (see e.g. [42] for a deep discussion on this topic), which is done by exploiting the distributional nature of (P_h^f) and using a suitable Leibniz rule (see Proposition 2.7). The second question concerns the behaviour of the matrices $A_h(x)$ outside the reference domain Ω . Indeed, differently from the local scenario, since the fractional gradient ∇^s must be defined on the whole space \mathbb{R}^n , then the fractional divergence div^s acts on vector fields globally defined over \mathbb{R}^n (see Definition 2.2). Hence, the matrix-valued functions $A_h(x)$ must be defined on the whole space \mathbb{R}^n . On the other hand, $-\text{div}^s(A_h \nabla^s u_h)$ is bounded only in Ω , and so we cannot hope to obtain compactness of $(A_h)_h$ outside the reference domain.

In order to tackle this lack of compactness, the values of any matrix $A_h(x)$ are prescribed outside Ω by a fixed matrix $A_0(x)$ which satisfies standard growth conditions.

In details, we study the limiting behaviour, in the sense of the H -convergence, of the class of matrix-valued functions

$$\mathcal{M}(\lambda, \Lambda, \Omega, A_0) := \{A \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n) : A = A_0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}$$

for positive constants $\lambda \leq \Lambda$ and $A_0 \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$, where $\mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$ denotes the class of matrix-valued measurable functions $A: \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$ satisfying the growth conditions

$$\begin{aligned} A(x)\xi \cdot \xi &\geq \lambda|\xi|^2 && \text{for all } \xi \in \mathbb{R}^n \text{ and for a.e. } x \in \mathbb{R}^n, \\ A(x)\xi \cdot \xi &\geq \Lambda^{-1}|A(x)\xi|^2 && \text{for all } \xi \in \mathbb{R}^n \text{ and for a.e. } x \in \mathbb{R}^n. \end{aligned}$$

Our first main result is Theorem 3.1, which is the H -compactness of the class $\mathcal{M}(\lambda, \Lambda, \Omega, A_0)$ with respect to a suitable notion of nonlocal H -convergence, see Definition 2.13. More precisely, we prove that any sequence of matrix-valued functions $(A_h)_h$ in $\mathcal{M}(\lambda, \Lambda, \Omega, A_0)$ admits, up to a subsequence, a limit matrix A_∞ such that $(A_h)_h$ H -converges to A_∞ . Moreover, A_∞ still belongs to $\mathcal{M}(\lambda, \Lambda, \Omega, A_0)$, leading to the compactness of the class. It would be of great interest to compare our results with the ones obtained in [43, 44], where a different notion of nonlocal H -convergence, based on a functional analytic approach, is introduced.

We point out that, within the reference domain, our H -limit A_∞ coincides with its local counterpart, showing a consistency between the local and nonlocal H -convergence. As a by-product, in Theorem 3.2 we show that also the subclass of symmetric matrices, here denoted by $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, is H -compact, i.e. if $(A_h)_h$ is in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, then A_∞ belongs to $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$.

In the second part of this paper, we focus exclusively on the symmetric case and prove Theorem 4.1, which is the Γ -compactness of the class of nonlocal energies associated with $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$. This is achieved by exploiting the classical local Γ -compactness result for quadratic functionals, see Proposition 2.19, and proving the equivalence between Γ -convergence of the nonlocal energies and Γ -convergence of the corresponding local ones, see Proposition 4.3. This equivalence relies on Lemma 4.2, where we show that the integral representation of the nonlocal energies in term of L^∞ -matrices is unique (as in the local case), and the properties of the Riesz potential, which allow us to pass from the nonlocal scenario to the local one.

A very non-exhaustive list of recent references about the Γ -convergence of fractional quadratic energies is given by [1, 7, 32, 36]. In particular, in [7], the Γ -compactness is obtained using the Beurling-Deny criteria for Dirichlet forms, which are unfortunately not available in our setting since it is not known whether quadratic forms of the fractional gradient are Dirichlet forms.

In the last part of the paper, we prove in Theorem 5.1 the equivalence between nonlocal H -convergence in the symmetric case and Γ -convergence of the associated energies. This result is obtained by adapting the following classical argument: in Lemma 5.2, we first compare the Γ -convergence of the nonlocal energies with a suitable notion of nonlocal G -convergence. Then, in Proposition 5.4, we show the convergence of the momenta through the Γ -convergence.

Finally, as a consequence of the uniqueness representation of Lemma 4.2 and the equivalence Theorem 5.1, we derive the uniqueness of the H -limit A_∞ in the symmetric case, see Corollary 5.5. Moreover, in view of the Γ -compactness Theorem 4.1, we show that the nonlocal H -compactness of $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, already proved in Theorem 3.2, can be equivalently obtained only by Γ -convergence arguments.

The paper is organized as follows. In Section 2, we provide and discuss in details all the preliminaries. Section 3 is dedicated to the proofs of both H -compactness results of Theorem 3.1 and Theorem 3.2. Then, in Section 4 and Section 5, we focus specifically on the symmetric case. We prove the Γ -compactness result in Theorem 4.1, establish the equivalence between Γ -convergence and H -convergence in Theorem 5.1, and gives an alternative proof of the H -compactness result in Theorem 3.2, based on Γ -convergence. Finally, in Section 6, we present a list of open problems and new research directions drawn from this work.

2. PRELIMINARIES

2.1. Notation. We assume that $s \in (0, 1)$, $n \geq 2$, and that Ω is a bounded open subset of \mathbb{R}^n . The space of $n \times n$ real matrices is denoted by $\mathbb{R}^{n \times n}$, while $\mathbb{R}_{\text{sym}}^{n \times n}$ denotes the subspace of symmetric matrices. We adopt standard notation for Lebesgue spaces on measurable subsets $E \subseteq \mathbb{R}^n$ and Sobolev spaces on open subsets $O \subseteq \mathbb{R}^n$. We denote by $\|\cdot\|_X$ the norm of a Banach space X , by X' the dual space of X , and by $\langle \cdot, \cdot \rangle_{X' \times X}$ the duality product between X' and X .

2.2. The functional setting. For any $\alpha \in (0, n)$, the α -Riesz potential of a measurable function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is defined as

$$I_\alpha f(x) := \frac{1}{\gamma_\alpha} \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}} dy, \quad \text{with } \gamma_\alpha := 2^\alpha \pi^{\frac{n}{2}} \frac{\Gamma(\frac{\alpha}{2})}{\Gamma(\frac{n-\alpha}{2})}.$$

Notice that I_α is a Fourier multiplier that acts in the Fourier space as

$$\mathcal{F}(I_\alpha f)(\xi) = |\xi|^{-\alpha} \mathcal{F}(f)(\xi),$$

being \mathcal{F} the Fourier transform. Moreover, as shown in [40, Theorem 1, pag. 119], we have the following result.

Proposition 2.1. *Let $\alpha \in (0, n)$ and $p \in (1, \frac{n}{\alpha})$. For any $f \in L^p(\mathbb{R}^n)$, the α -Riesz potential $I_\alpha f$ is well-defined and there exists a positive constant C , depending only on α , n , and p , such that*

$$\|I_\alpha f\|_{L^{p_\alpha^*}(\mathbb{R}^n)} \leq C \|f\|_{L^p(\mathbb{R}^n)},$$

where $p_\alpha^* := \frac{np}{n-\alpha p}$.

Let us recall the notions of fractional divergence and fractional gradient that will be used to introduce the H -convergence problem in our framework.

Definition 2.2. *Let $\psi \in C_c^\infty(\mathbb{R}^n)$ and $\phi \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$ be fixed. We define the s -fractional gradient $\nabla^s \psi$ and the s -fractional divergence $\text{div}^s \phi$, respectively, as*

$$\begin{aligned} \nabla^s \psi(x) &:= \frac{n-1+s}{\gamma_{1-s}} \int_{\mathbb{R}^n} \frac{(\psi(x) - \psi(y))(x-y)}{|x-y|^{n+s+1}} dy \quad \text{for all } x \in \mathbb{R}^n, \\ \text{div}^s \phi(x) &:= \frac{n-1+s}{\gamma_{1-s}} \int_{\mathbb{R}^n} \frac{(\phi(x) - \phi(y)) \cdot (x-y)}{|x-y|^{n+s+1}} dy \quad \text{for all } x \in \mathbb{R}^n. \end{aligned}$$

The next result links the notions of fractional gradient and fractional divergence with the classical ones. A proof can be found in [34, Theorem 1.2] and [12, Proposition 2.2].

Proposition 2.3. *Let $\psi \in C_c^\infty(\mathbb{R}^n)$ and $\phi \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$. It holds that*

$$\begin{aligned} \nabla^s \psi &= \nabla(I_{1-s}\psi) = I_{1-s}(\nabla\psi) \quad \text{in } \mathbb{R}^n, \\ \text{div}^s \phi &= I_{1-s}(\text{div} \phi) = \text{div}(I_{1-s}\phi) \quad \text{in } \mathbb{R}^n. \end{aligned}$$

We now introduce the functional framework for our problems in fractional divergence form.

Definition 2.4. *Let $O \subseteq \mathbb{R}^n$ be an open set. We denote by $H_0^s(O)$ the Hilbert space defined as the closure of $C_c^\infty(O)$ with respect to the following norm*

$$\|\cdot\|_{H^s(\mathbb{R}^n)} := \left(\|\cdot\|_{L^2(\mathbb{R}^n)}^2 + \|\nabla^s \cdot\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)}^2 \right)^{1/2}$$

and, by $H^{-s}(O)$, the dual space of $H_0^s(O)$. When $O = \mathbb{R}^n$, we simply write $H_0^s(\mathbb{R}^n) = H^s(\mathbb{R}^n)$.

As a consequence of Proposition 2.3, one can immediately show, through Fourier transform, that for all $\psi \in C_c^\infty(\mathbb{R}^n)$ and $s, \sigma \in (0, 1)$ we have

$$(2.1) \quad -\text{div}^s(\nabla^\sigma \psi) = (-\Delta)^{\frac{s+\sigma}{2}} \psi.$$

Moreover, by Fubini's Theorem, it holds that

$$(2.2) \quad \int_{\mathbb{R}^n} \nabla^s \psi(x) \cdot \phi(x) dx = - \int_{\mathbb{R}^n} \psi(x) \text{div}^s \phi(x) dx$$

for all $\phi \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$ and $\psi \in C_c^\infty(\mathbb{R}^n)$.

If we extend the operators ∇^s and div^s respectively to $H^s(\mathbb{R}^n)$ and $H^s(\mathbb{R}^n; \mathbb{R}^n)$, then (2.2) holds for all $\phi \in H^s(\mathbb{R}^n; \mathbb{R}^n)$ and $\psi \in H^s(\mathbb{R}^n)$. In particular, we can define the operator $\operatorname{div}^s: L^2(\mathbb{R}^n; \mathbb{R}^n) \rightarrow H^{-s}(\Omega)$ as

$$\langle \operatorname{div}^s u, v \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} := - \int_{\mathbb{R}^n} u(x) \cdot \nabla^s v(x) \, dx$$

for all $u \in L^2(\mathbb{R}^n; \mathbb{R}^n)$ and $v \in H_0^s(\Omega)$.

The following two Propositions provide a useful connection between the Sobolev spaces $H^1(\mathbb{R}^n)$ and $H^s(\mathbb{R}^n)$. For the proof of the next Proposition we refer the interested reader to [8, Lemma A.4].

Proposition 2.5. *Let $w \in H^1(\mathbb{R}^n)$ and set $u := (-\Delta)^{\frac{1-s}{2}} w$. It holds that*

$$u \in H^s(\mathbb{R}^n) \quad \text{and} \quad \nabla^s u = \nabla w \text{ a.e. in } \mathbb{R}^n.$$

Moreover, there exists a positive constant C , depending only on n and s , such that the following estimate holds true

$$(2.3) \quad \|u\|_{L^2(\mathbb{R}^n)} \leq C(n, s) \|w\|_{L^2(\mathbb{R}^n)}^s \|\nabla w\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)}^{1-s}.$$

Proposition 2.6. *Let $u \in H^s(\mathbb{R}^n)$ and set $w := I_{1-s} u$. It holds that*

$$w \in \left\{ v \in L^{2_{1-s}^*}(\mathbb{R}^n) : \nabla v \in L^2(\mathbb{R}^n) \right\} \quad \text{and} \quad \nabla w = \nabla^s u \text{ a.e. in } \mathbb{R}^n,$$

where $2_{1-s}^* = \frac{2n}{n-2+2s}$.

In particular, $w \in H_{\text{loc}}^1(\mathbb{R}^n)$ and, for every open set $O \Subset \mathbb{R}^n$, there exists a positive constant C , depending only on n , s and O , such that the following estimate holds true

$$(2.4) \quad \|w\|_{H^1(O)} \leq C(n, s) \|u\|_{H^s(\mathbb{R}^n)}.$$

Proof. If $u \in C_c^\infty(\mathbb{R}^n)$, the result is a consequence of Proposition 2.1 (with $p = 2$ and $\alpha = 1 - s$) and of Proposition 2.3.

Let now $(u_h)_h \subset C_c^\infty(\mathbb{R}^n)$ be such that $u_h \rightarrow u$ strongly in $H^s(\mathbb{R}^n)$. Then,

$$w_h := I_{1-s} u_h \rightarrow w := I_{1-s} u \quad \text{strongly in } L^{2_{1-s}^*}(\mathbb{R}^n) \text{ as } h \rightarrow \infty$$

and

$$\nabla w_h = \nabla^s u_h \rightarrow \nabla^s u \quad \text{strongly in } L^2(\mathbb{R}^n; \mathbb{R}^n) \text{ as } h \rightarrow \infty,$$

which gives the existence of $\nabla w = \nabla^s u \in L^2(\mathbb{R}^n; \mathbb{R}^n)$. Thus, the estimate (2.4) follows again in virtue of Proposition 2.1. \square

We conclude this subsection recalling some extensions of classical results to the framework of fractional calculus, that will be used throughout the paper. They are a Leibniz-type rule for the fractional gradient [13, Eq. (1.5), (1.6)] and [21, Eq. (2.11)], a Poincaré-type inequality [34, Theorem 3.3], and a Rellich-type Theorem [35, Theorem 2.2].

Proposition 2.7 (Leibniz rule). *Let $\varphi \in C_c^1(\Omega)$ and $u \in H^s(\mathbb{R}^n)$. Then, $\varphi u \in H_0^s(\Omega)$ and*

$$\nabla^s(\varphi u) = \varphi \nabla^s u + u \nabla^s \varphi + \nabla_{\text{NL}}^s(\varphi, u)$$

where, for every $x \in \mathbb{R}^n$, the remainder term $\nabla_{\text{NL}}^s(\varphi, u)(x)$ is

$$\nabla_{\text{NL}}^s(\varphi, u)(x) := \frac{n-1+s}{\gamma_{1-s}} \int_{\mathbb{R}^n} \frac{(\varphi(x) - \varphi(y))(u(x) - u(y))(x-y)}{|x-y|^{n+s+1}} \, dy.$$

Moreover, there exists a positive constant C , depending only on n and s , such that

$$\|\nabla_{\text{NL}}^s(\varphi, u)\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)} \leq C \|u\|_{L^2(\mathbb{R}^n)} \|\varphi\|_{L^\infty(\Omega)}^{1-s} \|\nabla \varphi\|_{L^\infty(\Omega; \mathbb{R}^n)}.$$

Proposition 2.8 (Poincaré inequality). *For every set $O \Subset \mathbb{R}^n$ there exists a positive constant C , depending only on n , s and O , such that*

$$\|u\|_{L^2(O)} \leq C \|\nabla^s u\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)} \quad \text{for all } u \in H^s(\mathbb{R}^n).$$

Proposition 2.9 (Rellich Theorem). *For every set $O \in \mathbb{R}^n$ the space $H_0^s(O)$ is compactly embedded into $L^2(O)$.*

2.3. The H -convergence problem. We now want to introduce our notion of H -convergence associated with the nonlocal operators introduced above. As in the local counterpart (see e.g. [42]), we begin by defining the class of matrices in which we are interested.

Definition 2.10. *Given $0 < \lambda \leq \Lambda < \infty$ and a measurable subset E of \mathbb{R}^n , we define $\mathcal{M}(\lambda, \Lambda, E)$ as the collection of all matrix-valued measurable functions $A: E \rightarrow \mathbb{R}^{n \times n}$ satisfying*

$$(2.5) \quad A(x)\xi \cdot \xi \geq \lambda|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n \text{ and for a.e. } x \in E,$$

$$(2.6) \quad A(x)\xi \cdot \xi \geq \Lambda^{-1}|A(x)\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n \text{ and for a.e. } x \in E.$$

We also set

$$\mathcal{M}^{\text{sym}}(\lambda, \Lambda, E) := \{A \in \mathcal{M}(\lambda, \Lambda, E) : A = A^T \text{ a.e. in } E\}.$$

The estimate (2.6) above is needed to obtain the compactness of the class $\mathcal{M}(\lambda, \Lambda, \Omega)$ with respect to the H -convergence topology, even in the local setting. More precisely, it is well-known that the H -limit of sequences of matrices satisfying the standard growth condition

$$(2.7) \quad \lambda|\xi|^2 \leq A(x)\xi \cdot \xi \leq \Lambda|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n \text{ and for a.e. } x \in \Omega,$$

instead of (2.5) and (2.6), may belong to a wider class $\mathcal{M}(\lambda, \Lambda', \Omega)$ with $\Lambda' \geq \Lambda$ (see e.g. the observations of Tartar in [42, Chapter 6, Pag. 81]).

We also point out that, in the symmetric case $A = A^T$, conditions (2.5)–(2.6) are actually equivalent to (2.7), as shown in the following result.

Lemma 2.11. *Given $0 < \lambda \leq \Lambda < \infty$, let $B \in \mathbb{R}^{n \times n}$ satisfy*

$$(2.8) \quad B\xi \cdot \xi \geq \lambda|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n.$$

Then, B is invertible and condition

$$(2.9) \quad B\xi \cdot \xi \geq \Lambda^{-1}|B\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n$$

is equivalent to

$$(2.10) \quad B^{-1}\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n.$$

In particular, B satisfies

$$(2.11) \quad B\xi \cdot \xi \leq \Lambda|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n.$$

Moreover, if B is symmetric ($B = B^T$), then conditions (2.9), (2.10) and (2.11) are all equivalent.

Proof. Let $B \in \mathbb{R}^{n \times n}$ satisfy (2.8). By Lax-Milgram Theorem, for all $\eta \in \mathbb{R}^n$ there exists a unique vector $\xi \in \mathbb{R}^n$ such that $B\xi = \eta$, i.e. B is invertible. Hence, (2.9) implies

$$\Lambda^{-1}|\xi|^2 = \Lambda^{-1}|BB^{-1}\xi|^2 \leq BB^{-1}\xi \cdot B^{-1}\xi = B^{-1}\xi \cdot \xi \quad \text{for all } \xi \in \mathbb{R}^n.$$

Conversely, by (2.10), we get

$$\Lambda^{-1}|B\xi|^2 \leq B^{-1}B\xi \cdot B\xi = B\xi \cdot \xi \quad \text{for all } \xi \in \mathbb{R}^n.$$

Assume now that B satisfies condition (2.9). By Cauchy-Schwartz inequality, we get

$$\Lambda^{-1}|B\xi|^2 \leq B\xi \cdot \xi \leq |B\xi||\xi| \quad \text{for all } \xi \in \mathbb{R}^n,$$

leading to

$$(2.12) \quad |B\xi| \leq \Lambda|\xi|.$$

Therefore, by applying again Cauchy-Schwartz inequality, we get

$$B\xi \cdot \xi \leq |B\xi||\xi| \leq \Lambda|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n$$

and so condition (2.9) implies (2.11).

Finally, assume that B is symmetric. It is enough to show that (2.11) implies (2.9). We start by observing that, by the symmetry of B , the bilinear form $(\xi, \eta) \mapsto (B\xi, \eta)$ is a scalar product in \mathbb{R}^n , in virtue of (2.11). Hence, by Cauchy-Schwartz inequality and (2.11), it holds that

$$|B\xi \cdot \eta|^2 \leq (B\xi \cdot \xi)(B\eta \cdot \eta) \leq (B\xi \cdot \xi)\Lambda|\eta|^2 \quad \text{for all } \xi, \eta \in \mathbb{R}^n.$$

In particular, for $\eta = B\xi$, we get

$$|B\xi|^4 = |B\xi \cdot B\xi|^2 \leq (B\xi \cdot \xi)\Lambda|B\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n,$$

which implies (2.9). \square

From now on, we fix $0 < \lambda \leq \Lambda < \infty$. Given a sequence of matrices $(A_h)_h \subset \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$, for all $f \in H^{-s}(\Omega)$ and $h \in \mathbb{N}$ we consider the following elliptic problems

$$(P_h^f) \quad \begin{cases} -\operatorname{div}^s(A_h \nabla^s u_h) = f & \text{in } \Omega, \\ u_h = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where div^s and ∇^s are the fractional differential operators introduced in Definition 2.2. The following result ensures that the problem (P_h^f) is well-defined.

Lemma 2.12. *Let $A \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$. For every $f \in H^{-s}(\Omega)$ there exists a unique (weak) solution $u \in H_0^s(\Omega)$ of the elliptic problem*

$$\begin{cases} -\operatorname{div}^s(A \nabla^s u) = f & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

i.e. satisfying

$$\int_{\mathbb{R}^n} A(x) \nabla^s u(x) \cdot \nabla^s v(x) \, dx = \langle f, v \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} \quad \text{for all } v \in H_0^s(\Omega).$$

Moreover, the solution u satisfies the following estimate

$$\|\nabla^s u\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)} \leq \lambda^{-1} \|f\|_{H^{-s}(\Omega)}.$$

Proof. The proof is a direct application of Lax-Milgram Theorem and Proposition 2.8. Indeed, the bilinear form $a : H_0^s(\Omega) \times H_0^s(\Omega) \rightarrow \mathbb{R}$, defined as

$$a(u, v) := \int_{\mathbb{R}^n} A(x) \nabla^s u(x) \cdot \nabla^s v(x) \, dx \quad \text{for all } u, v \in H_0^s(\Omega),$$

is continuous and coercive, being $A \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$. \square

The notion of nonlocal H -convergence of $(A_h)_h \subset \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$ we propose consists in finding a limit matrix $A_\infty \in \mathcal{M}(\lambda', \Lambda', \mathbb{R}^n)$, with $0 < \lambda' \leq \lambda \leq \Lambda \leq \Lambda' < \infty$, such that the sequence of problems $(P_h^f)_h$ is related to the limit problem

$$(P_\infty^f) \quad \begin{cases} -\operatorname{div}^s(A_\infty \nabla^s u_\infty) = f & \text{in } \Omega, \\ u_\infty = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

in the sense of the next definition. In what follows, we denote by $u_h = u_h(f) \in H_0^s(\Omega)$, $h \in \mathbb{N}$, and by $u_\infty = u_\infty(f) \in H_0^s(\Omega)$ the unique weak solutions of (P_h^f) and (P_∞^f) , respectively.

Definition 2.13 (Nonlocal H -convergence). *Let $0 < \lambda' \leq \lambda \leq \Lambda \leq \Lambda' < \infty$ and consider $(A_h)_h \subset \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$ and $A_\infty \in \mathcal{M}(\lambda', \Lambda', \mathbb{R}^n)$. We say that*

$$(A_h)_h \text{ H-converges to } A_\infty \text{ in } H_0^s(\Omega)$$

if for all $f \in H^{-s}(\Omega)$ the following convergences simultaneously hold as $h \rightarrow \infty$:

$$(2.13) \quad \textbf{convergence of solutions:} \quad u_h \rightarrow u_\infty \quad \text{weakly in } H_0^s(\Omega),$$

$$(2.14) \quad \textbf{convergence of momenta:} \quad A_h \nabla^s u_h \rightarrow A_\infty \nabla^s u_\infty \quad \text{weakly in } L^2(\mathbb{R}^n; \mathbb{R}^n).$$

Definition 2.13 is the natural counterpart in the nonlocal setting of the local H -convergence, see e.g. [42, Definition 6.4]. For the readers' convenience, we recall such a notion in what follows.

Let $(B_h)_h \subset \mathcal{M}(\lambda, \Lambda, \Omega)$ be fixed. For any $h \in \mathbb{N}$ and $g \in H^{-1}(\Omega)$, we consider the following sequence of elliptic problems

$$(Q_h^g) \quad \begin{cases} -\operatorname{div}(B_h \nabla w_h) = g & \text{in } \Omega, \\ w_h = 0 & \text{on } \partial\Omega. \end{cases}$$

Given $B_\infty \in \mathcal{M}(\lambda', \Lambda', \Omega)$, for some $0 < \lambda' \leq \lambda \leq \Lambda \leq \Lambda' < \infty$, we also consider the problem

$$(Q_\infty^g) \quad \begin{cases} -\operatorname{div}(B_\infty \nabla w_\infty) = g & \text{in } \Omega, \\ w_\infty = 0 & \text{on } \partial\Omega, \end{cases}$$

and denote by $w_h = w_h(g) \in H_0^1(\Omega)$, $h \in \mathbb{N}$, and by $w_\infty = w_\infty(g) \in H_0^1(\Omega)$ the unique weak solutions of (Q_h^g) and (Q_∞^g) , respectively.

Definition 2.14 (Local H -convergence). *Let $0 < \lambda' \leq \lambda \leq \Lambda \leq \Lambda' < \infty$, and consider $(B_h)_h \subset \mathcal{M}(\lambda, \Lambda, \Omega)$ and $B_\infty \in \mathcal{M}(\lambda', \Lambda', \Omega)$. We say that*

$$(B_h)_h \text{ } H\text{-converges to } B_\infty \text{ in } H_0^1(\Omega),$$

if for all $g \in H^{-1}(\Omega)$ the following convergences simultaneously hold as $h \rightarrow \infty$:

$$(2.15) \quad \textbf{convergence of solutions:} \quad w_h \rightarrow w_\infty \quad \text{weakly in } H_0^1(\Omega),$$

$$(2.16) \quad \textbf{convergence of momenta:} \quad B_h \nabla w_h \rightarrow B_\infty \nabla w_\infty \quad \text{weakly in } L^2(\Omega; \mathbb{R}^n).$$

We recall that the class $\mathcal{M}(\lambda, \Lambda, \Omega)$ is compact with respect to the local H -convergence in $H_0^1(\Omega)$. A proof of this classical result can be found e.g. in [42, Theorem 6.5].

Proposition 2.15 (Local H -compactness). *For any $(B_h)_h \subset \mathcal{M}(\lambda, \Lambda, \Omega)$, there exist a not relabeled subsequence and a matrix-valued function $B_\infty \in \mathcal{M}(\lambda, \Lambda, \Omega)$ such that*

$$(B_h)_h \text{ } H\text{-converges to } B_\infty \text{ in } H_0^1(\Omega).$$

We also recall that the local H -convergence is stable with respect to the transpose operation. More precisely, we have the following result, due to Tartar, whose proof can be found in [42, Lemma 10.2].

Proposition 2.16 (H -convergence of the transpose). *Let $B_h, B_\infty \in \mathcal{M}(\lambda, \Lambda, \Omega)$, $h \in \mathbb{N}$, and assume that*

$$(B_h)_h \text{ } H\text{-converges to } B_\infty \text{ in } H_0^1(\Omega).$$

Then,

$$(B_h^T)_h \text{ } H\text{-converges to } B_\infty^T \text{ in } H_0^1(\Omega).$$

The first goal of this paper is to show that suitable subclasses of $\mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$ are compact with respect to the *nonlocal* H -convergence. The classes of matrices we are interested in are defined as follows.

Definition 2.17. *For any $A_0 \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$, we define*

$$\mathcal{M}(\lambda, \Lambda, \Omega, A_0) := \{A \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n) : A = A_0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}.$$

For any $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$, we also set

$$\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0) := \{A \in \mathcal{M}(\lambda, \Lambda, \Omega, A_0) : A = A^T \text{ a.e. in } \mathbb{R}^n\}.$$

In the next section we show that any sequence $(A_h)_h$ in $\mathcal{M}(\lambda, \Lambda, \Omega, A_0)$ (respectively in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$), for a fixed matrix $A_0 \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$ (respectively in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$), admits a not relabeled subsequence and a limit matrix A_∞ in $\mathcal{M}(\lambda, \Lambda, \Omega, A_0)$ (respectively in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$) such that

$$(A_h)_h \text{ } H\text{-converges to } A_\infty \text{ in } H_0^s(\Omega)$$

in the sense of Definition 2.13. In particular, the H -limit A_∞ satisfies (2.5)–(2.6) with the same constants λ and Λ of the sequence $(A_h)_h$, leading to the compactness of both classes.

2.4. The Γ -convergence problem. Let us fix $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$ and a sequence $(A_h)_h$ in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$. For any $h \in \mathbb{N}$, we consider the nonlocal energies $F_h: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ associated with A_h , represented by

$$(2.17) \quad F_h(u) := \begin{cases} \frac{1}{2} \int_{\mathbb{R}^n} A_h(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx & \text{if } u \in H_0^s(\Omega), \\ \infty & \text{if } u \in L^2(\mathbb{R}^n) \setminus H_0^s(\Omega). \end{cases}$$

The second goal of this paper is to show the Γ -compactness in the strong topology of $L^2(\mathbb{R}^n)$ for the class of nonlocal energies (2.17). In particular, we prove the existence of a limit symmetric matrix-valued function $A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ such that, up to a not relabeled subsequence,

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

where the limit nonlocal energy $F_\infty: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ is represented by

$$(2.18) \quad F_\infty(u) := \begin{cases} \frac{1}{2} \int_{\mathbb{R}^n} A_\infty(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx & \text{if } u \in H_0^s(\Omega), \\ \infty & \text{if } u \in L^2(\mathbb{R}^n) \setminus H_0^s(\Omega). \end{cases}$$

We start recalling the notion of Γ -convergence for functionals. For a complete treatment of the topic, we refer the interested reader to the monographs [5, 16].

Definition 2.18. Let X be a Banach space and let $E_h, E_\infty: X \rightarrow [0, \infty]$, $h \in \mathbb{N}$. We say that

$$(E_h)_h \text{ } \Gamma\text{-converges to } E_\infty \text{ strongly in } X$$

if the following two conditions simultaneously hold:

- **Γ -liminf inequality:** for all $x \in X$ and for any sequence $(x_h)_h \subset X$ strongly converging to x in X one has

$$E_\infty(x) \leq \liminf_{h \rightarrow \infty} E_h(x_h);$$

- **Γ -limsup inequality:** for all $x \in X$ there exists a recovery sequence $(y_h)_h \subset X$ strongly converging to x in X and such that

$$E_\infty(x) \geq \limsup_{h \rightarrow \infty} E_h(y_h).$$

In analogy with Subsection 2.3, we remind that the class of local energies $G_h: L^2(\Omega) \rightarrow [0, \infty]$, $h \in \mathbb{N}$, associated with a sequence $(B_h)_h$ in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega)$ and represented by

$$(2.19) \quad G_h(v) := \begin{cases} \frac{1}{2} \int_{\Omega} B_h(x) \nabla v(x) \cdot \nabla v(x) \, dx & \text{if } v \in H^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H^1(\Omega), \end{cases}$$

is compact with respect to the Γ -convergence in the strong topology of $L^2(\Omega)$. A proof of the next result can be found in [33] and [16, Theorem 22.2].

Proposition 2.19 (Γ -compactness of local functionals). Let $(B_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega)$ and, for any $h \in \mathbb{N}$, let $G_h: L^2(\Omega) \rightarrow [0, \infty]$ be the local energy associated with B_h as in (2.19). Then, there exists $G_\infty: L^2(\Omega) \rightarrow [0, \infty]$ such that, up to a not relabeled subsequence,

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega).$$

Moreover, there exists a matrix-valued function $B_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega)$ such that the Γ -limit G_∞ has the following integral representation

$$(2.20) \quad G_\infty(v) := \begin{cases} \frac{1}{2} \int_{\Omega} B_\infty(x) \nabla v(x) \cdot \nabla v(x) \, dx & \text{if } v \in H^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H^1(\Omega). \end{cases}$$

Remark 2.20. In order to derive Proposition 2.19 from [33] and [16, Theorem 22.2], we recall that, according to Lemma 2.11, any symmetric matrix-valued function B belongs to the space $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega)$ if and only if the following growth condition is satisfied

$$(2.21) \quad \lambda |\xi|^2 \leq B(x) \xi \cdot \xi \leq \Lambda |\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^n \text{ and for a.e. } x \in \Omega.$$

2.5. The equivalence problem. The third goal of this paper is to show the equivalence between nonlocal H -convergence and Γ -convergence of the associated nonlocal energies.

Let us fix $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. For any $h \in \mathbb{N}$, let $A_h, A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and let $F_h, F_\infty: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ be the corresponding nonlocal energies, defined respectively in (2.17) and (2.18). We want to show that

$$(A_h)_h \text{ } H\text{-converges to } A_\infty \text{ in } H_0^s(\Omega)$$

if and only if

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n).$$

In order to prove the above equivalence, we need to introduce the following notion of *nonlocal G -convergence* of $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$.

Definition 2.21 (Nonlocal G -convergence). *Let $0 < \lambda' \leq \lambda \leq \Lambda \leq \Lambda' < \infty$ and consider $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$ and $A_\infty \in \mathcal{M}^{\text{sym}}(\lambda', \Lambda', \mathbb{R}^n)$. We say that*

$$(A_h)_h \text{ } G\text{-converges to } A_\infty \text{ in } H_0^s(\Omega)$$

if for all $f \in H^{-s}(\Omega)$ the following convergence holds as $h \rightarrow \infty$:

$$(2.22) \quad u_h \rightarrow u_\infty \quad \text{weakly in } H_0^s(\Omega),$$

where $u_h = u_h(f)$ and $u_\infty = u_\infty(f)$ are, respectively, the unique solutions of (P_h^f) and (P_∞^f) .

Definition 2.21 is the natural counterpart in the nonlocal setting of the classical G -convergence, introduced by Spagnolo [37]. Clearly nonlocal H -convergence implies nonlocal G -convergence, see Definition 2.13.

In Section 5, we show that the nonlocal G -convergence of $(A_h)_h$ is equivalent to the Γ -convergence of the associated nonlocal energies $(F_h)_h$. Moreover, we prove that for any sequence $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, the Γ -convergence of the associated nonlocal energies $(F_h)_h$ implies the convergence of the nonlocal momenta.

As a consequence, in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ the notions of nonlocal H -convergence, nonlocal G -convergence, and Γ -convergence of the nonlocal energies, are all equivalent.

3. H -COMPACTNESS OF NONLOCAL OPERATORS

This first subsection is dedicated to the following H -compactness result for nonlocal operators. The proof of the following Theorem 3.1 relies on Proposition 2.15 and Proposition 2.16.

Theorem 3.1. *Let $A_0 \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$. For any $(A_h)_h \subset \mathcal{M}(\lambda, \Lambda, \Omega, A_0)$, there exist a not relabeled subsequence and a matrix-valued function $A_\infty \in \mathcal{M}(\lambda, \Lambda, \Omega, A_0)$ such that*

$$(A_h)_h \text{ } H\text{-converges to } A_\infty \text{ in } H_0^s(\Omega).$$

Proof. For any $h \in \mathbb{N}$, we define

$$(3.1) \quad B_h := A_h|_\Omega \in \mathcal{M}(\lambda, \Lambda, \Omega).$$

By Proposition 2.15, there exists a limit matrix

$$(3.2) \quad B_\infty \in \mathcal{M}(\lambda, \Lambda, \Omega)$$

such that, up to subsequences,

$$(B_h)_h \text{ } H\text{-converges to } B_\infty \text{ in } H_0^1(\Omega).$$

For a.e. $x \in \mathbb{R}^n$ we define

$$(3.3) \quad A_\infty(x) := \begin{cases} B_\infty(x) & \text{if } x \in \Omega, \\ A_0(x) & \text{if } x \in \mathbb{R}^n \setminus \Omega. \end{cases}$$

By construction, $A_\infty \in \mathcal{M}(\lambda, \Lambda, \Omega, A_0)$. To conclude the proof, we show that A_∞ is the H -limit of $(A_h)_h$ in $H_0^s(\Omega)$, in the sense of Definition 2.13.

Fix $f \in H^{-s}(\Omega)$ and consider the sequence $(u_h)_h \subset H_0^s(\Omega)$ of unique solutions of $(P_h^f)_h$. Since $(u_h)_h$ and $(A_h \nabla^s u_h)_h$ are respectively bounded in $H_0^s(\Omega)$ and $L^2(\mathbb{R}^n; \mathbb{R}^n)$, then there exist $u^* \in H_0^s(\Omega)$ and $m \in L^2(\mathbb{R}^n; \mathbb{R}^n)$ such that, up to a not relabeled subsequence,

$$(3.4) \quad u_h \rightarrow u^* \quad \text{weakly in } H_0^s(\Omega) \quad \text{and} \quad A_h \nabla^s u_h \rightarrow m \quad \text{weakly in } L^2(\mathbb{R}^n; \mathbb{R}^n) \quad \text{as } h \rightarrow \infty.$$

Our goal is to show that

$$(3.5) \quad m = A_\infty \nabla^s u^* \quad \text{a.e. in } \mathbb{R}^n.$$

Indeed, once obtained (3.5), by passing to the limit (as $h \rightarrow \infty$) in the weak formulation of (P_h^f) , we obtain that u^* is a solution of (P_∞^f) . Thus, by uniqueness, $u^* = u_\infty$ and, by (3.4),

$$(A_h)_h \text{ H-converges to } A_\infty \text{ in } H_0^s(\Omega),$$

as desired.

First, notice that

$$(3.6) \quad m = A_0 \nabla^s u^* = A_\infty \nabla^s u^* \quad \text{a.e. in } \mathbb{R}^n \setminus \Omega.$$

In fact, by (3.4), for all $\Phi \in L^2(\mathbb{R}^n \setminus \Omega; \mathbb{R}^n)$ it holds that

$$\begin{aligned} \int_{\mathbb{R}^n \setminus \Omega} m(x) \cdot \Phi(x) \, dx &= \lim_{h \rightarrow \infty} \int_{\mathbb{R}^n \setminus \Omega} A_h(x) \nabla^s u_h(x) \cdot \Phi(x) \, dx \\ &= \lim_{h \rightarrow \infty} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u_h(x) \cdot \Phi(x) \, dx = \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u^*(x) \cdot \Phi(x) \, dx, \end{aligned}$$

which implies (3.6).

It remains to show that

$$(3.7) \quad m = A_\infty \nabla^s u^* \quad \text{a.e. in } \Omega.$$

Fix $g \in H^{-1}(\Omega)$ and consider $(w_h)_h \subset H_0^1(\Omega)$ and $w_\infty \in H_0^1(\Omega)$, respectively (unique) solutions of the transpose problems

$$(3.8) \quad \begin{cases} -\operatorname{div}(B_h^T \nabla w_h) = g & \text{in } \Omega, \\ w_h = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{and} \quad \begin{cases} -\operatorname{div}(B_\infty^T \nabla w_\infty) = g & \text{in } \Omega, \\ w_\infty = 0 & \text{on } \partial\Omega. \end{cases}$$

By Proposition 2.16,

$$(3.9) \quad w_h \rightarrow w_\infty \quad \text{weakly in } H_0^1(\Omega) \quad \text{and} \quad B_h^T \nabla w_h \rightarrow B_\infty^T \nabla w_\infty \quad \text{weakly in } L^2(\Omega; \mathbb{R}^n) \quad \text{as } h \rightarrow \infty.$$

Let $\varphi \in C_c^\infty(\Omega)$ and define

$$M_h := \int_{\mathbb{R}^n} \varphi(x) A_h(x) \nabla^s u_h(x) \cdot \nabla w_h(x) \, dx.$$

We claim that for all $\varphi \in C_c^\infty(\Omega)$

$$(3.10) \quad \lim_{h \rightarrow \infty} M_h = \int_{\Omega} \varphi(x) m(x) \cdot \nabla w_\infty(x) \, dx = \int_{\Omega} \varphi(x) A_\infty(x) \nabla^s u^*(x) \cdot \nabla w_\infty(x) \, dx.$$

Once the claim is proved, we obtain (3.7). Indeed, since the operator $-\operatorname{div}(B_\infty^T \nabla \cdot)$ defines a bijection between the spaces $H_0^1(\Omega)$ and $H^{-1}(\Omega)$ and, since $g \in H^{-1}(\Omega)$ can be arbitrarily taken in (3.8), then for all $\varphi \in C_c^\infty(\Omega)$ and $w \in H_0^1(\Omega)$ the following identity holds

$$\int_{\Omega} \varphi(x) m(x) \cdot \nabla w(x) \, dx = \int_{\Omega} \varphi(x) A_\infty(x) \nabla^s u^*(x) \cdot \nabla w(x) \, dx.$$

Hence,

$$(3.11) \quad m(x) \cdot \nabla w(x) = A_\infty(x) \nabla^s u^*(x) \cdot \nabla w(x) \quad \text{for a.e. } x \in \Omega \quad \text{and for all } w \in H_0^1(\Omega)$$

and the collections of points of Ω where (3.11) fails can be chosen independent of w by a density argument. Therefore, by fixing $\Omega' \Subset \Omega$ and $\phi \in C_c^1(\Omega)$, such that $\phi = 1$ on Ω' , and by defining

$$w(x) := \phi(x) \xi \cdot x \quad \text{for a.e. } x \in \Omega \quad \text{and for all } \xi \in \mathbb{R}^n,$$

by (3.11), we get that

$$m(x) \cdot \xi = A_\infty(x) \nabla^s u^*(x) \cdot \xi \quad \text{for a.e. } x \in \Omega' \text{ and for all } \xi \in \mathbb{R}^n,$$

which implies the validity of (3.7) in Ω' . Moreover, since this is true for every $\Omega' \Subset \Omega$, we get (3.7) in all Ω , which completes the proof.

We then conclude by showing the validity of the claim (3.10). Its proof is divided into two steps.

Step 1. We first show that

$$(3.12) \quad \lim_{h \rightarrow \infty} M_h = \int_{\Omega} \varphi(x) A_\infty(x) \nabla^s u^*(x) \cdot \nabla w_\infty(x) \, dx.$$

In virtue of Proposition 2.3, we have

$$(3.13) \quad \begin{aligned} M_h &= \int_{\Omega} \varphi(x) \nabla^s u_h(x) \cdot B_h^T(x) \nabla w_h(x) \, dx \\ &= \int_{\Omega} \varphi(x) \nabla(I_{1-s} u_h)(x) \cdot B_h^T(x) \nabla w_h(x) \, dx \\ &= \int_{\Omega} \nabla(\varphi I_{1-s} u_h)(x) \cdot B_h^T(x) \nabla w_h(x) \, dx - \int_{\Omega} I_{1-s} u_h(x) \nabla \varphi(x) \cdot B_h^T(x) \nabla w_h(x) \, dx. \end{aligned}$$

By (3.4) and Proposition 2.9,

$$(3.14) \quad u_h \rightarrow u^* \quad \text{strongly in } L^2(\mathbb{R}^n) \text{ as } h \rightarrow \infty,$$

and so, by Proposition 2.1, we get

$$I_{1-s} u_h \rightarrow I_{1-s} u^* \quad \text{strongly in } L^2(\Omega) \text{ as } h \rightarrow \infty.$$

This last convergence, coupled with (3.9), implies that

$$(3.15) \quad \lim_{h \rightarrow \infty} \int_{\Omega} I_{1-s} u_h(x) \nabla \varphi(x) \cdot B_h^T(x) \nabla w_h(x) \, dx = \int_{\Omega} I_{1-s} u^*(x) \nabla \varphi(x) \cdot B_\infty^T(x) \nabla w_\infty(x) \, dx.$$

We observe that $\varphi I_{1-s} u_h \in H_0^1(\Omega)$. Hence, by (3.4), (3.14), and Proposition 2.6,

$$\varphi I_{1-s} u_h \rightarrow \varphi I_{1-s} u^* \quad \text{weakly in } H_0^1(\Omega) \text{ as } h \rightarrow \infty.$$

Thus, since w_h solves problem (3.8), we get

$$(3.16) \quad \begin{aligned} \lim_{h \rightarrow \infty} \int_{\Omega} \nabla(\varphi I_{1-s} u_h)(x) \cdot B_h^T(x) \nabla w_h(x) \, dx &= \lim_{h \rightarrow \infty} \langle g, \varphi I_{1-s} u_h \rangle_{H^{-1}(\Omega) \times H_0^1(\Omega)} \\ &= \langle g, \varphi I_{1-s} u^* \rangle_{H^{-1}(\Omega) \times H_0^1(\Omega)} \\ &= \int_{\Omega} \nabla(\varphi I_{1-s} u^*)(x) \cdot B_\infty^T(x) \nabla w_\infty(x) \, dx. \end{aligned}$$

By combining (3.13), (3.15), and (3.16), we then obtain (3.12), being

$$\begin{aligned} \lim_{h \rightarrow \infty} M_h &= \int_{\Omega} \nabla(\varphi I_{1-s} u^*)(x) \cdot B_\infty^T(x) \nabla w_\infty(x) \, dx - \int_{\Omega} I_{1-s} u^*(x) \nabla \varphi(x) \cdot B_\infty^T(x) \nabla w_\infty(x) \, dx \\ &= \int_{\Omega} \varphi(x) \nabla(I_{1-s} u^*)(x) \cdot B_\infty^T(x) \nabla w_\infty(x) \, dx = \int_{\Omega} \varphi(x) A_\infty(x) \nabla^s u^*(x) \cdot \nabla w_\infty(x) \, dx. \end{aligned}$$

Step 2. We conclude by showing that

$$(3.17) \quad \lim_{h \rightarrow \infty} M_h = \int_{\Omega} \varphi(x) m(x) \cdot \nabla w_\infty(x) \, dx.$$

By Proposition 2.5 and Proposition 2.7, we have

$$\begin{aligned}
(3.18) \quad M_h &= \int_{\Omega} \varphi(x) A_h(x) \nabla^s u_h(x) \cdot \nabla w_h(x) \, dx \\
&= \int_{\mathbb{R}^n} \varphi(x) A_h(x) \nabla^s u_h(x) \cdot \nabla^s ((-\Delta)^{\frac{1-s}{2}} w_h)(x) \, dx \\
&= \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla^s (\varphi (-\Delta)^{\frac{1-s}{2}} w_h)(x) \, dx \\
&\quad - \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla^s \varphi(x) (-\Delta)^{\frac{1-s}{2}} w_h(x) \, dx \\
&\quad - \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla_{\text{NL}}^s (\varphi, (-\Delta)^{\frac{1-s}{2}} w_h)(x) \, dx.
\end{aligned}$$

For what concerns the second integral in (3.18), in view of (3.9) and Proposition 2.9, we have

$$w_h \rightarrow w_{\infty} \quad \text{strongly in } L^2(\mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

Moreover, the sequence $(w_h)_h \subset H_0^1(\Omega)$ is uniformly bounded. Hence, by Proposition 2.5,

$$(3.19) \quad (-\Delta)^{\frac{1-s}{2}} w_h \rightarrow (-\Delta)^{\frac{1-s}{2}} w_{\infty} \quad \text{strongly in } L^2(\mathbb{R}^n)$$

and, since $\nabla^s \varphi \in L^{\infty}(\mathbb{R}^n)$, by (3.4) we get

$$(3.20) \quad \lim_{h \rightarrow \infty} \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla^s \varphi(x) (-\Delta)^{\frac{1-s}{2}} w_h(x) \, dx = \int_{\mathbb{R}^n} m(x) \cdot \nabla^s \varphi(x) (-\Delta)^{\frac{1-s}{2}} w_{\infty}(x) \, dx.$$

Regarding the third integral in (3.18), by Proposition 2.7 and (3.19), since ∇_{NL}^s is a bilinear operator, we deduce that

$$\nabla_{\text{NL}}^s (\varphi, (-\Delta)^{\frac{1-s}{2}} w_h) \rightarrow \nabla_{\text{NL}}^s (\varphi, (-\Delta)^{\frac{1-s}{2}} w_{\infty}) \quad \text{strongly in } L^2(\mathbb{R}^n; \mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

Therefore, in virtue of (3.4), we have

$$(3.21) \quad \lim_{h \rightarrow \infty} \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla_{\text{NL}}^s (\varphi, (-\Delta)^{\frac{1-s}{2}} w_h)(x) \, dx$$

$$(3.22) \quad = \int_{\mathbb{R}^n} m(x) \cdot \nabla_{\text{NL}}^s (\varphi, (-\Delta)^{\frac{1-s}{2}} w_{\infty})(x) \, dx.$$

Finally, $(-\Delta)^{\frac{1-s}{2}} w_h \in H^s(\mathbb{R}^n)$ by Proposition 2.5, which implies that $\varphi (-\Delta)^{\frac{1-s}{2}} w_h \in H_0^s(\Omega)$. Then,

$$\varphi (-\Delta)^{\frac{1-s}{2}} w_h \rightarrow \varphi (-\Delta)^{\frac{1-s}{2}} w_{\infty} \quad \text{weakly in } H_0^s(\Omega) \text{ as } h \rightarrow \infty.$$

Therefore, regarding the first integrals in (3.18), since u_h is a solution of (P_h^f) then, by (3.4),

$$\begin{aligned}
(3.23) \quad & \lim_{h \rightarrow \infty} \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla^s (\varphi (-\Delta)^{\frac{1-s}{2}} w_h)(x) \, dx \\
&= \lim_{h \rightarrow \infty} \langle f, \varphi (-\Delta)^{\frac{1-s}{2}} w_h \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} \\
&= \langle f, \varphi (-\Delta)^{\frac{1-s}{2}} w_{\infty} \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} \\
&= \lim_{h \rightarrow \infty} \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla^s (\varphi (-\Delta)^{\frac{1-s}{2}} w_{\infty})(x) \, dx \\
&= \int_{\mathbb{R}^n} m(x) \cdot \nabla^s (\varphi (-\Delta)^{\frac{1-s}{2}} w_{\infty})(x) \, dx.
\end{aligned}$$

By combining (3.18), (3.20), (3.21), and (3.23), we finally get (3.17), being

$$\begin{aligned} \lim_{h \rightarrow \infty} M_h &= \int_{\mathbb{R}^n} m(x) \cdot \nabla^s (\varphi(-\Delta)^{\frac{1-s}{2}} w_\infty)(x) dx \\ &\quad - \int_{\mathbb{R}^n} m(x) \cdot \nabla^s \varphi(x) (-\Delta)^{\frac{1-s}{2}} w_\infty(x) dx - \int_{\mathbb{R}^n} m(x) \cdot \nabla_{\text{NL}}^s (\varphi, (-\Delta)^{\frac{1-s}{2}} w_\infty)(x) dx \\ &= \int_{\mathbb{R}^n} \varphi(x) m(x) \cdot \nabla^s ((-\Delta)^{\frac{1-s}{2}} w_\infty)(x) dx = \int_{\Omega} \varphi(x) m(x) \cdot \nabla w_\infty(x) dx. \end{aligned}$$

Hence, the claim (3.10) holds true and the proof of the Theorem is accomplished. \square

As a consequence of Proposition 2.16, the H -compactness Theorem 3.1 also applies to the subclass of symmetric matrices $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, for a fixed $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. More precisely, the H -limit A_∞ of sequences $(A_h)_h$ in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ still lies in $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$.

Theorem 3.2. *Let $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. For any $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, there exist a not relabeled subsequence and a matrix-valued function $A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ such that*

$$(A_h)_h \text{ } H\text{-converges to } A_\infty \text{ in } H_0^s(\Omega).$$

Proof. The proof of the Theorem in the symmetric case follows verbatim the construction already presented in the proof of Theorem 3.1 for the general case, combined with the following observations.

The limit matrix B_∞ in the local H -convergence of the sequence $(B_h)_h$, introduced in (3.1), whose existence is shown in (3.2), is now symmetric in view of Proposition 2.16 and by the uniqueness of the local H -limit. This implies that the matrix A_∞ , defined in (3.3), belongs to the class $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, and it is the H -limit of $(A_h)_h$, in virtue of Theorem 3.1. \square

4. Γ -COMPACTNESS OF NONLOCAL ENERGIES

The goal of this subsection is to show the following Γ -compactness Theorem.

Theorem 4.1. *Let $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. Let $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and let $(F_h)_h$ be the nonlocal energies introduced in (2.17). Then, there exist a not relabeled subsequence of $(A_h)_h$ and $A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ such that*

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

where $F_\infty: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ is the nonlocal energy associated with A_∞ , as in (2.18).

In order to prove Theorem 4.1, we begin with the following lemma, which guarantees the uniqueness of the integral representation of nonlocal energies by matrices.

Lemma 4.2. *Let $A, \widehat{A} \in L^\infty(\mathbb{R}^n; \mathbb{R}_{\text{sym}}^{n \times n})$ and let $\Omega \subseteq \mathbb{R}^n$ be an open set. Assume that*

$$A(x)\xi \cdot \xi \geq 0 \quad \text{and} \quad \widehat{A}(x)\xi \cdot \xi \geq 0 \quad \text{for a.e. } x \in \Omega \text{ and all } \xi \in \mathbb{R}^n,$$

and

$$(4.1) \quad \int_{\mathbb{R}^n} A(x) \nabla^s \psi(x) \cdot \nabla^s \psi(x) dx = \int_{\mathbb{R}^n} \widehat{A}(x) \nabla^s \psi(x) \cdot \nabla^s \psi(x) dx \quad \text{for all } \psi \in C_c^\infty(\Omega).$$

Then

$$(4.2) \quad A(x_0) = \widehat{A}(x_0) \quad \text{for a.e. } x_0 \in \Omega.$$

Proof. We define

$$C := A - \widehat{A} \in L^\infty(\mathbb{R}^n; \mathbb{R}_{\text{sym}}^{n \times n}).$$

Since $C \in L^\infty(\mathbb{R}^n; \mathbb{R}_{\text{sym}}^{n \times n})$, for a.e. $x_0 \in \mathbb{R}^n$ and for all $M \in (0, \infty)$ we have

$$\lim_{r \rightarrow \infty} \int_{B_M(0)} \left| C \left(\frac{y}{r} + x_0 \right) - C(x_0) \right| dy = 0.$$

In particular, there exists $(r_k)_k \subset (0, \infty)$ with $r_k \rightarrow \infty$ as $k \rightarrow \infty$ such that

$$(4.3) \quad \left| C\left(\frac{y}{r_k} + x_0\right) - C(x_0) \right| \rightarrow 0 \quad \text{as } k \rightarrow \infty \text{ for a.e. } y \in \mathbb{R}^n.$$

Step 1: Blow-up. We fix $\varphi \in C_c^\infty(\mathbb{R}^n)$. For all $x_0 \in \Omega$ and $r \in (0, \infty)$ we define

$$\varphi_{x_0, r}(x) := \varphi(r(x - x_0)) \quad \text{for all } x \in \mathbb{R}^n.$$

There exists $r_0 = r_0(x_0, \Omega) \in (0, \infty)$ such that

$$\varphi_{x_0, r} \in C_c^\infty(\Omega) \quad \text{for all } r \in (r_0, \infty).$$

We have that

$$\begin{aligned} \nabla^s \varphi_{x_0, r}(x) &= \frac{1}{\gamma_{1-s}} \int_{\mathbb{R}^n} \frac{\nabla \varphi_{x_0, r}(y)}{|y - x|^{n+s-1}} dy = \frac{r}{\gamma_{1-s}} \int_{\mathbb{R}^n} \frac{\nabla \varphi(r(y - x_0))}{|y - x|^{n+s-1}} dy \\ &= \frac{r^s}{\gamma_{1-s}} \int_{\mathbb{R}^n} \frac{\nabla \varphi(z)}{|z + r(x_0 - x)|^{n+s-1}} dz = r^s \nabla^s \varphi(r(x - x_0)). \end{aligned}$$

By (4.1), for all $x_0 \in \Omega$ and $r \in (r_0, \infty)$ we have

$$(4.4) \quad \begin{aligned} 0 &= r^{n-2s} \int_{\mathbb{R}^n} C(x) \nabla^s \varphi_{x_0, r}(x) \cdot \nabla^s \varphi_{x_0, r}(x) dx \\ &= r^n \int_{\mathbb{R}^n} C(x) \nabla^s \varphi(r(x - x_0)) \cdot \nabla^s \varphi(r(x - x_0)) dx \\ &= \int_{\mathbb{R}^n} C\left(\frac{y}{r} + x_0\right) \nabla^s \varphi(y) \cdot \nabla^s \varphi(y) dy. \end{aligned}$$

Moreover, by (4.3), we derive

$$C\left(\frac{y}{r_k} + x_0\right) \nabla^s \varphi(y) \cdot \nabla^s \varphi(y) \rightarrow C(x_0) \nabla^s \varphi(y) \cdot \nabla^s \varphi(y) \quad \text{for a.e. } y \in \mathbb{R}^n \text{ as } k \rightarrow \infty.$$

Since

$$\left| C\left(\frac{y}{r_k} + x_0\right) \nabla^s \varphi(y) \cdot \nabla^s \varphi(y) \right| \leq \|C\|_{L^\infty(\mathbb{R}^n; \mathbb{R}^{n \times n})} |\nabla^s \varphi(y)|^2 \quad \text{for a.e. } y \in \mathbb{R}^n \text{ and all } k \in \mathbb{N},$$

in virtue of the dominated convergence Theorem and (4.4), we conclude that

$$\int_{\mathbb{R}^n} C(x_0) \nabla^s \varphi(y) \cdot \nabla^s \varphi(y) dy = 0 \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^n).$$

Step 2: Reduction to the local case. By a density argument, we derive that

$$\int_{\mathbb{R}^n} C(x_0) \nabla^s u(y) \cdot \nabla^s u(y) dy = 0 \quad \text{for all } u \in H^s(\mathbb{R}^n).$$

Let $v \in H^1(\mathbb{R}^n)$. By Proposition 2.5, we have that $u := (-\Delta)^{\frac{1-s}{2}} v \in H^s(\mathbb{R}^n)$ and

$$\nabla^s u(x) = \nabla v(x) \quad \text{for a.e. } x \in \mathbb{R}^n.$$

Therefore, we derive that

$$\int_{\mathbb{R}^n} A(x_0) \nabla v(y) \cdot \nabla v(y) dy = \int_{\mathbb{R}^n} \widehat{A}(x_0) \nabla v(y) \cdot \nabla v(y) dy \quad \text{for all } v \in H^1(\mathbb{R}^n).$$

Hence, we can apply [16, Lemma 22.5] to obtain (4.2). \square

As a consequence of Lemma 4.2, we derive the following equivalence between Γ -convergence of the nonlocal energies $(F_h)_h$ and Γ -convergence of the local ones $(G_h)_h$. The proof is inspired by some recent ideas presented in [15, 21].

Proposition 4.3. *Let $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. For any $h \in \mathbb{N}$, let $A_h, A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and $F_h, F_\infty: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ be the nonlocal energies, respectively defined in (2.17) and (2.18). For any $h \in \mathbb{N}$, define*

$$B_h := A_h|_\Omega \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega), \quad B_\infty := A_\infty|_\Omega \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega),$$

and consider the local energies $G_h, G_\infty: L^2(\Omega) \rightarrow [0, \infty]$, respectively defined in (2.19) and (2.20). Then,

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n)$$

if and only if

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega).$$

Proof. Step 1: Γ -convergence of $(G_h)_h$ implies Γ -convergence of $(F_h)_h$. We assume that

$$(4.5) \quad (G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega),$$

and we want to show that

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n).$$

Γ -liminf inequality. Let $u_h, u \in L^2(\mathbb{R}^n)$, $h \in \mathbb{N}$, be such that $(u_h)_h$ strongly converges to u in $L^2(\mathbb{R}^n)$ as $h \rightarrow \infty$. We show that

$$F_\infty(u) \leq \liminf_{h \rightarrow \infty} F_h(u_h).$$

Without loss of generality, we assume that

$$\liminf_{h \rightarrow \infty} F_h(u_h) < \infty,$$

the conclusion being otherwise trivial, and that the limit is actually achieved up to a not relabeled subsequence, i.e.

$$\liminf_{h \rightarrow \infty} F_h(u_h) = \lim_{h \rightarrow \infty} F_h(u_h).$$

According to its own definition, $(F_h)_h$ is finite only on $H_0^s(\Omega)$, thus forcing the sequence $(u_h)_h$ to lie therein. Since $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, there exists a positive constant C such that

$$\sup_{h \in \mathbb{N}} \|\nabla^s u_h\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)} \leq C,$$

which yields that $(u_h)_h$ is uniformly bounded in $H_0^s(\Omega)$. Then, the limit u also lies on $H_0^s(\Omega)$ and

$$(4.6) \quad u_h \rightarrow u \quad \text{weakly in } H_0^s(\Omega) \text{ as } h \rightarrow \infty.$$

For any $h \in \mathbb{N}$, we define

$$v_h := I_{1-s} u_h \quad \text{and} \quad v := I_{1-s} u.$$

By Proposition 2.6, $v_h, v \in H^1(\Omega)$ for any $h \in \mathbb{N}$ and, by (4.6) and the continuity of the linear operator $I_{1-s}: H^s(\mathbb{R}^n) \rightarrow H^1(\Omega)$,

$$v_h \rightarrow v \quad \text{strongly in } L^2(\Omega) \text{ and weakly in } H^1(\Omega) \text{ as } h \rightarrow \infty.$$

By (4.5),

$$(4.7) \quad G_\infty(v) \leq \liminf_{h \rightarrow \infty} G_h(v_h),$$

in virtue of the Γ -liminf inequality. We also note that, by Proposition 2.6,

$$\nabla v_h = \nabla^s u_h \quad \text{and} \quad \nabla v = \nabla^s u \quad \text{a.e. in } \mathbb{R}^n.$$

Thus, we can rephrase (4.7) as

$$\frac{1}{2} \int_\Omega B_\infty(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx \leq \liminf_{h \rightarrow \infty} \frac{1}{2} \int_\Omega A_h(x) \nabla^s u_h(x) \cdot \nabla^s u_h(x) \, dx.$$

On the other hand, since $A_0 \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$, by (4.6) we get

$$\frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx \leq \liminf_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u_h(x) \cdot \nabla^s u_h(x) \, dx.$$

Hence,

$$\begin{aligned}
F_\infty(u) &= \frac{1}{2} \int_\Omega B_\infty(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx + \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx \\
&\leq \liminf_{h \rightarrow \infty} \frac{1}{2} \int_\Omega A_h(x) \nabla^s u_h(x) \cdot \nabla^s u_h(x) \, dx + \liminf_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u_h(x) \cdot \nabla^s u_h(x) \, dx \\
&\leq \liminf_{h \rightarrow \infty} F_h(u_h).
\end{aligned}$$

Γ -limsup inequality. We fix $u \in L^2(\mathbb{R}^n)$ and show the existence of a recovery sequence $(u_h)_h \subset L^2(\mathbb{R}^n)$ such that $(u_h)_h$ strongly converges to u in $L^2(\mathbb{R}^n)$, as $h \rightarrow \infty$, and

$$(4.8) \quad F_\infty(u) \geq \limsup_{h \rightarrow \infty} F_h(u_h).$$

The proof of the Γ -limsup inequality is rather technical and for the readers' convenience, we indicate the main steps below.

- First, we exploit the Riesz potential to move to the local setting and we obtain the existence of a recovery sequence $(v_h)_h$ for the Γ -convergence of the local energies $(G_h)_h$ to G_∞ .
- Then, through a cut-off argument, we adapt the sequence $(v_h)_h$ to the boundary data of our problem and we come back to the nonlocal setting, obtaining the existence of a sequence $(u_h^\varepsilon)_h$ satisfying the Γ -limsup inequality up to a reminder term, which depends on a parameter $\varepsilon > 0$.
- In the last part of the proof, we let the reminder term tend to zero via a diagonal argument, which ensures the existence of a recovery sequence $(u_h)_h$ for our problem.

Without loss of generality, we consider only the case of $u \in H_0^s(\Omega)$, the conclusion being otherwise trivial.

We define

$$v := I_{1-s}u.$$

Then, by Proposition 2.6,

$$(4.9) \quad v \in H^1(\Omega) \quad \text{and} \quad \nabla v = \nabla^s u \quad \text{a.e. in } \mathbb{R}^n.$$

Moreover, by (4.5), there exists a recovery sequence $(v_h)_h \subset H^1(\Omega)$ for v , i.e. such that

$$(4.10) \quad v_h \rightarrow v \text{ strongly in } L^2(\Omega) \text{ as } h \rightarrow \infty \quad \text{and} \quad \lim_{h \rightarrow \infty} G_h(v_h) = G_\infty(v) < \infty$$

(we recall, in fact, that the Γ -liminf and the Γ -limsup inequalities imply that the limit is achieved at least for the recovery sequence). In particular, by the definition of G_h (see (2.19)), $(v_h)_h$ is bounded in $H^1(\Omega)$, which gives that

$$v_h \rightarrow v \quad \text{weakly in } H^1(\Omega) \text{ as } h \rightarrow \infty.$$

Let $\varepsilon > 0$ be fixed and let $K^\varepsilon \Subset \Omega$ be a compact set such that

$$(4.11) \quad \int_{\Omega \setminus K^\varepsilon} |\nabla v(x)|^2 \, dx < \varepsilon.$$

We fix an open set U^ε such that $K^\varepsilon \Subset U^\varepsilon \Subset \Omega$, consider a cut-off function $\varphi^\varepsilon \in C_c^\infty(U^\varepsilon)$ satisfying $0 \leq \varphi^\varepsilon \leq 1$ on U^ε and $\varphi^\varepsilon \equiv 1$ on K^ε and, for any $h \in \mathbb{N}$, we define

$$(4.12) \quad v_h^\varepsilon := \varphi^\varepsilon v_h + (1 - \varphi^\varepsilon)v.$$

By construction,

$$(4.13) \quad v_h^\varepsilon \rightarrow v \quad \text{strongly in } L^2(\Omega) \text{ and weakly in } H^1(\Omega) \text{ as } h \rightarrow \infty.$$

Moreover, by (2.12), (2.21), (4.11), and the convexity of the map $\xi \mapsto A_h(x)\xi \cdot \xi$, it holds that

$$\begin{aligned}
G_h(v_h^\varepsilon) &= \frac{1}{2} \int_{\Omega} A_h(x) [\varphi^\varepsilon(x) \nabla v_h(x) + (1 - \varphi^\varepsilon(x)) \nabla v(x)] \cdot [\varphi^\varepsilon(x) \nabla v_h(x) + (1 - \varphi^\varepsilon(x)) \nabla v(x)] dx \\
&\quad + \int_{\Omega} A_h(x) [\nabla \varphi^\varepsilon(x) (v_h(x) - v(x))] \cdot [\varphi^\varepsilon(x) \nabla v_h(x) + (1 - \varphi^\varepsilon(x)) \nabla v(x)] dx \\
&\quad + \frac{1}{2} \int_{\Omega} A_h(x) [\nabla \varphi^\varepsilon(x) (v_h(x) - v(x))] \cdot [\nabla \varphi^\varepsilon(x) (v_h(x) - v(x))] dx \\
&\leq \frac{1}{2} \int_{\Omega} \varphi^\varepsilon(x) A_h(x) \nabla v_h(x) \cdot \nabla v_h(x) dx + \frac{1}{2} \int_{\Omega} (1 - \varphi^\varepsilon(x)) A_h(x) \nabla v(x) \cdot \nabla v(x) dx \\
&\quad + \Lambda \|\nabla \varphi^\varepsilon\|_{L^\infty(\Omega; \mathbb{R}^n)} \|v_h - v\|_{L^2(\Omega)} (\|\nabla v_h\|_{L^2(\Omega; \mathbb{R}^n)} + \|\nabla v\|_{L^2(\Omega; \mathbb{R}^n)}) \\
&\quad + \frac{\Lambda}{2} \|\nabla \varphi^\varepsilon\|_{L^\infty(\Omega; \mathbb{R}^n)}^2 \|v_h - v\|_{L^2(\Omega)}^2 \\
&\leq G_h(v_h) + \frac{\Lambda}{2} \varepsilon + \Lambda \|\nabla \varphi^\varepsilon\|_{L^\infty(\Omega; \mathbb{R}^n)} \|v_h - v\|_{L^2(\Omega)} (\|\nabla v_h\|_{L^2(\Omega; \mathbb{R}^n)} + \|\nabla v\|_{L^2(\Omega; \mathbb{R}^n)}) \\
&\quad + \frac{\Lambda}{2} \|\nabla \varphi^\varepsilon\|_{L^\infty(\Omega; \mathbb{R}^n)}^2 \|v_h - v\|_{L^2(\Omega)}^2.
\end{aligned}$$

Hence, by (4.10) and the boundedness of $(v_h)_h$ in $H^1(\Omega)$, we conclude that

$$(4.14) \quad \limsup_{h \rightarrow \infty} G_h(v_h^\varepsilon) \leq \lim_{h \rightarrow \infty} G_h(v_h) + \frac{\Lambda}{2} \varepsilon = G_\infty(v) + \frac{\Lambda}{2} \varepsilon.$$

We trivially extend $v_h^\varepsilon - v \in H_0^1(\Omega)$ to a function in $H^1(\mathbb{R}^n)$ and, for any $h \in \mathbb{N}$, we define

$$w_h^\varepsilon := (-\Delta)^{\frac{1-s}{2}} (v_h^\varepsilon - v).$$

By Proposition 2.5, we have that

$$(4.15) \quad w_h^\varepsilon \in H^s(\mathbb{R}^n) \quad \text{and} \quad \nabla^s w_h^\varepsilon = \nabla (v_h^\varepsilon - v) \quad \text{a.e. in } \mathbb{R}^n$$

and, by (2.3), (4.13), and (4.15), there exist two positive constants C and C_ε such that

$$\|w_h^\varepsilon\|_{H^s(\mathbb{R}^n)}^2 = \|w_h^\varepsilon\|_{L^2(\mathbb{R}^n)}^2 + \|\nabla^s w_h^\varepsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)}^2 \leq C \|v_h^\varepsilon - v\|_{H_0^1(\Omega)}^2 \leq C_\varepsilon \quad \text{for all } h \in \mathbb{N}.$$

Therefore, by (2.1), (2.2) and (4.13), for all $\psi \in C_c^\infty(\mathbb{R}^n)$ we get that

$$\int_{\mathbb{R}^n} w_h^\varepsilon(x) \psi(x) dx = \int_{\mathbb{R}^n} (v_h^\varepsilon(x) - v(x)) (-\Delta)^{\frac{1-s}{2}} \psi(x) dx \rightarrow 0 \quad \text{as } h \rightarrow \infty,$$

which yields that

$$(4.16) \quad w_h^\varepsilon \rightarrow 0 \quad \text{weakly in } H^s(\mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

In particular, by (2.3) and (4.13),

$$(4.17) \quad w_h^\varepsilon \rightarrow 0 \quad \text{strongly in } L^2(\mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

Let $\chi^\varepsilon \in C_c^\infty(\Omega)$ satisfy $0 \leq \chi^\varepsilon \leq 1$ on Ω and $\chi^\varepsilon = 1$ on $\overline{U^\varepsilon}$. We define

$$u_h^\varepsilon := u + \chi^\varepsilon w_h^\varepsilon \in H_0^s(\Omega).$$

By (4.16) and (4.17),

$$(4.18) \quad u_h^\varepsilon \rightarrow u \quad \text{strongly in } L^2(\mathbb{R}^n) \text{ and weakly in } H_0^s(\Omega) \text{ as } h \rightarrow \infty.$$

For any $h \in \mathbb{N}$, we also set

$$R_h^\varepsilon := \nabla^s (\chi^\varepsilon w_h^\varepsilon) - \chi^\varepsilon \nabla^s w_h^\varepsilon.$$

By Proposition 2.7, there exists a positive constant C such that

$$\|R_h^\varepsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)} \leq C \|\chi^\varepsilon\|_{W^{1,\infty}(\mathbb{R}^n)} \|w_h^\varepsilon\|_{L^2(\mathbb{R}^n)} \quad \text{for all } h \in \mathbb{N}.$$

Then, by (4.17),

$$(4.19) \quad R_h^\varepsilon \rightarrow 0 \quad \text{strongly in } L^2(\mathbb{R}^n; \mathbb{R}^n) \text{ as } h \rightarrow \infty$$

and, by (4.15),

$$(4.20) \quad \nabla^s u_h^\varepsilon = \nabla^s u + \chi^\varepsilon \nabla^s w_h^\varepsilon + R_h^\varepsilon = \nabla^s u + \chi^\varepsilon \nabla(v_h^\varepsilon - v) + R_h^\varepsilon \quad \text{a.e. in } \mathbb{R}^n.$$

We consider the following decomposition

$$(4.21) \quad F_h(u_h^\varepsilon) = \frac{1}{2} \int_{U^\varepsilon} A_h(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx + \frac{1}{2} \int_{\Omega \setminus U^\varepsilon} A_h(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx \\ + \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx.$$

By (4.19) and (4.20), the last integral in (4.21) satisfies

$$(4.22) \quad \lim_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx \\ = \lim_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) (\nabla^s u(x) + R_h^\varepsilon(x)) \cdot (\nabla^s u(x) + R_h^\varepsilon(x)) dx \\ = \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) \nabla^s u(x) \cdot \nabla^s u(x) dx.$$

Concerning the second integral in (4.21), we note that, by (4.12) and (4.20),

$$\nabla^s u_h^\varepsilon = \nabla^s u + R_h^\varepsilon \quad \text{a.e. in } \Omega \setminus U^\varepsilon.$$

Then, by (2.21), (4.9), and (4.11),

$$(4.23) \quad \limsup_{h \rightarrow \infty} \frac{1}{2} \int_{\Omega \setminus U^\varepsilon} A_h(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx \\ \leq \lim_{h \rightarrow \infty} \frac{\Lambda}{2} \|\nabla^s u + R_h^\varepsilon\|_{L^2(\Omega \setminus U^\varepsilon; \mathbb{R}^n)}^2 = \frac{\Lambda}{2} \|\nabla^s u\|_{L^2(\Omega \setminus U^\varepsilon; \mathbb{R}^n)}^2 \leq \frac{\Lambda}{2} \|\nabla v\|_{L^2(\Omega \setminus K^\varepsilon; \mathbb{R}^n)}^2 \leq \frac{\Lambda}{2} \varepsilon.$$

Finally, for what concerns the first integral in (4.21), we observe that, since $\chi^\varepsilon = 1$ in U^ε , by (4.9) and (4.20) we have

$$\nabla^s u_h^\varepsilon = \nabla v + \nabla(v_h^\varepsilon - v) + R_h^\varepsilon = \nabla v_h^\varepsilon + R_h^\varepsilon \quad \text{a.e. in } U^\varepsilon.$$

Thus, (2.21) and (4.9) imply that

$$(4.24) \quad \frac{1}{2} \int_{U^\varepsilon} A_h(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx \\ = \frac{1}{2} \int_{U^\varepsilon} A_h(x) (\nabla v_h^\varepsilon(x) + R_h^\varepsilon(x)) \cdot (\nabla v_h^\varepsilon(x) + R_h^\varepsilon(x)) dx \\ = \frac{1}{2} \int_{U^\varepsilon} A_h(x) \nabla v_h^\varepsilon(x) \cdot \nabla v_h^\varepsilon(x) dx + \int_{U^\varepsilon} A_h \nabla v_h^\varepsilon(x) \cdot R_h^\varepsilon(x) dx \\ + \frac{1}{2} \int_{U^\varepsilon} A_h R_h^\varepsilon(x) \cdot R_h^\varepsilon(x) dx \\ \leq G_h(v_h^\varepsilon) + \int_{U^\varepsilon} A_h \nabla v_h^\varepsilon(x) \cdot R_h^\varepsilon(x) dx + \frac{1}{2} \int_{U^\varepsilon} A_h R_h^\varepsilon(x) \cdot R_h^\varepsilon(x) dx$$

and, since $(v_h^\varepsilon)_h$ is uniformly bounded in $H^1(\Omega)$, by (4.14), (4.19) and (4.24), we get

$$(4.25) \quad \limsup_{h \rightarrow \infty} \frac{1}{2} \int_{U^\varepsilon} A_h(x) \nabla^s u_h^\varepsilon(x) \cdot \nabla^s u_h^\varepsilon(x) dx \leq G_\infty(v) + \frac{\Lambda}{2} \varepsilon.$$

Therefore, by (4.22), (4.23) and (4.25), we obtain that for all $\varepsilon > 0$

$$(4.26) \quad \limsup_{h \rightarrow \infty} F_h(u_h^\varepsilon) \leq F_\infty(u) + \Lambda \varepsilon.$$

To conclude, we use the following diagonal argument. In view of [16, Definition 4.1 and Remark 4.3], by (4.18) and (4.26), we have that for all $\varepsilon > 0$

$$\Gamma\text{-}\limsup_{h \rightarrow \infty} F_h(u) := \sup_{k \in \mathbb{N}} \limsup_{h \rightarrow \infty} \inf_{z \in B_{\frac{1}{k}}(u)} F_h(z) \leq F_\infty(u) + \Lambda \varepsilon.$$

Hence, by letting $\varepsilon \rightarrow 0$, we conclude that

$$\Gamma\text{-}\limsup_{h \rightarrow \infty} F_h(u) \leq F_\infty(u)$$

and, by the properties of the Γ -lim sup (see e.g. [16, Proposition 8.1]), there exists a sequence $(u_h)_h \subset L^2(\mathbb{R}^n)$ such that $u_h \rightarrow u$ strongly in $L^2(\mathbb{R}^n)$ as $h \rightarrow \infty$ and

$$\limsup_{h \rightarrow \infty} F_h(u_h) = \Gamma\text{-}\limsup_{h \rightarrow \infty} F_h(u) \leq F_\infty(u).$$

This implies the validity of (4.8).

Step 2: Γ -convergence of $(F_h)_h$ implies Γ -convergence of $(G_h)_h$. We assume that

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

and we want to show that

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega).$$

By Proposition 2.19, there exist a not relabeled subsequence and a functional $\widehat{G}_\infty: L^2(\Omega) \rightarrow [0, \infty]$ such that

$$(4.27) \quad (G_h)_h \text{ } \Gamma\text{-converges to } \widehat{G}_\infty \text{ strongly in } L^2(\Omega).$$

Moreover, there exists $\widehat{B}_\infty \in \mathcal{M}^{\text{sym}}(\Omega, \lambda, \Lambda)$ such that

$$\widehat{G}_\infty(v) := \begin{cases} \frac{1}{2} \int_{\Omega} \widehat{B}_\infty(x) \nabla v(x) \cdot \nabla v(x) \, dx & \text{if } v \in H^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H^1(\Omega). \end{cases}$$

By (4.27) and Step 1, we conclude that

$$(F_h)_h \text{ } \Gamma\text{-converges to } \widehat{F}_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

where \widehat{F}_∞ is the nonlocal energies associated with the matrix

$$\widehat{A}_\infty(x) = \begin{cases} \widehat{B}_\infty(x) & \text{if } x \in \Omega, \\ A_0(x) & \text{if } x \in \mathbb{R}^n \setminus \Omega. \end{cases}$$

By the uniqueness of the Γ -limit, we conclude that $\widehat{F}_\infty = F_\infty$, and by Lemma 4.2 we derive that $\widehat{B}_\infty = B_\infty$. Hence, by the Uryshon property of Γ -convergence, we conclude that

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega).$$

□

Remark 4.4. We point out that the nonlocal energies $(F_h)_h$ account for the boundary condition, while the corresponding local energies $(G_h)_h$ do not, as they are finite in $H^1(\Omega)$ instead of $H_0^1(\Omega)$. We have chosen to work with $(G_h)_h$, since it simplifies the proof of the Γ -convergence equivalence between nonlocal energies and local ones. On the other hand, if we define

$$G_h^0(v) := \begin{cases} \frac{1}{2} \int_{\Omega} B_h(x) \nabla v(x) \cdot \nabla v(x) \, dx & \text{if } v \in H_0^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H_0^1(\Omega), \end{cases}$$

and

$$G_\infty^0(v) := \begin{cases} \frac{1}{2} \int_{\Omega} B_\infty(x) \nabla v(x) \cdot \nabla v(x) \, dx & \text{if } v \in H_0^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H_0^1(\Omega), \end{cases}$$

by [16, Theorem 13.12 and Theorem 22.4] we have that

$$(G_h^0)_h \text{ } \Gamma\text{-converges to } G_\infty^0 \text{ strongly in } L^2(\Omega)$$

if and only if

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega).$$

Therefore, Proposition 4.3 is still valid if we replace $(G_h)_h$ and G_∞ , respectively, with $(G_h^0)_h$ and G_∞^0 .

As a consequence of Proposition 2.19 and Proposition 4.3, we can finally prove Theorem 4.1.

Proof of Theorem 4.1. For any $h \in \mathbb{N}$, we define the matrix-valued functions

$$B_h := A_h|_{\Omega} \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega),$$

and we consider the functionals $G_h: L^2(\Omega) \rightarrow [0, \infty]$ associated with B_h , as in (2.19). By Proposition 2.19, there exist a not relabeled subsequence and a matrix $B_{\infty} \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega)$ such that

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_{\infty} \text{ strongly in } L^2(\Omega),$$

where the functional $G_{\infty}: L^2(\Omega) \rightarrow [0, \infty]$ is defined as in (2.20).

We define

$$A_{\infty}(x) := \begin{cases} B_{\infty}(x) & \text{if } x \in \Omega, \\ A_0(x) & \text{if } x \in \mathbb{R}^n \setminus \Omega, \end{cases}$$

and we denote by $F_{\infty}: L^2(\Omega) \rightarrow [0, \infty]$ the nonlocal energy associated with A_{∞} , as in (2.18). Then, $A_{\infty} \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and, by Proposition 4.3, we deduce that

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_{\infty} \text{ strongly in } L^2(\mathbb{R}^n).$$

□

5. EQUIVALENCE BETWEEN NONLOCAL H -CONVERGENCE AND Γ -CONVERGENCE

We conclude this paper with the following equivalence between nonlocal H -convergence of a sequence $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and Γ -convergence of the associated nonlocal energies $(F_h)_h$, introduced in (2.17).

Theorem 5.1. *Let $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. For any $h \in \mathbb{N}$, let $A_h, A_{\infty} \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and $F_h, F_{\infty}: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ be the nonlocal energies, respectively defined in (2.17) and (2.18). Then,*

$$(A_h)_h \text{ } H\text{-converges to } A_{\infty} \text{ in } H_0^s(\Omega)$$

if and only if

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_{\infty} \text{ strongly in } L^2(\mathbb{R}^n).$$

The proof of Theorem 5.1 requires two preliminary results. In Lemma 5.2, we first show the equivalence between the Γ -convergence of the nonlocal energies and the nonlocal G -convergence introduced in Definition 2.21, which corresponds to the sole convergence of the solutions. Later, in Proposition 5.4, we show that the Γ -convergence of the nonlocal energies also implies the convergence of the momenta, as required in Definition 2.13.

The proof of the equivalence between the nonlocal G -convergence and the Γ -convergence of the associated nonlocal energies can be obtained as an application of [16, Theorem 13.12], with $Y = H_0^s(\Omega)$ and $X = L^2(\mathbb{R}^n)$. For the reader's convenience, we provide below a complete proof, following the techniques presented in [16, Theorem 13.5].

Lemma 5.2. *For any $h \in \mathbb{N}$, let $A_h, A_{\infty} \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$ and let $F_h, F_{\infty}: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ be the nonlocal energies, respectively defined in (2.17) and (2.18). Then*

$$(A_h)_h \text{ } G\text{-converges to } A_{\infty} \text{ in } H_0^s(\Omega)$$

if and only if

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_{\infty} \text{ strongly in } L^2(\mathbb{R}^n).$$

Proof. **Γ -convergence implies G -convergence.**

We assume that $(F_h)_h$ Γ -converges to F_{∞} strongly in $L^2(\mathbb{R}^n)$ and we first show that (2.22) holds for every $f \in L^2(\mathbb{R}^n)$. Later, by a density argument, we extend the validity of (2.22) for all $f \in H^{-s}(\Omega)$, which implies the G -convergence of $(A_h)_h$ to A_{∞} in $H_0^s(\Omega)$.

Step 1. We fix $g \in L^2(\mathbb{R}^n)$ and, for any $h \in \mathbb{N}$, we define $F_h^g, F_{\infty}^g: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ as

$$(5.1) \quad F_h^g(u) := F_h(u) + \int_{\mathbb{R}^n} g(x)u(x) dx \quad \text{and} \quad F_{\infty}^g(u) := F_{\infty}(u) + \int_{\mathbb{R}^n} g(x)u(x) dx$$

for all $u \in L^2(\mathbb{R}^n)$. Since we have perturbed with continuity F_h and F_∞ , then

$$(F_h^g)_h \text{ } \Gamma\text{-converges to } F_\infty^g \text{ strongly in } L^2(\mathbb{R}^n),$$

in virtue of [16, Proposition 6.21].

We note that the solutions $w_h \in H_0^s(\Omega)$ and $w_\infty \in H_0^s(\Omega)$ of problems (P_h^g) and (P_∞^g) , whose existence is guaranteed by Lemma 2.12, minimise the energies F_h^g and F_∞^g , respectively, i.e.

$$F_h^g(w_h) = \min_{u \in L^2(\mathbb{R}^n)} F_h^g(u) \quad \text{and} \quad F_\infty^g(w_\infty) = \min_{u \in L^2(\mathbb{R}^n)} F_\infty^g(u).$$

Therefore, by the Fundamental Theorem of Γ -convergence (see e.g. [16, Theorem 7.8]), we get

$$w_h \rightarrow w_\infty \text{ strongly in } L^2(\mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

In particular, since $(w_h)_h \subset H_0^s(\Omega)$ is uniformly bounded, we conclude that

$$(5.2) \quad w_h \rightarrow w_\infty \text{ weakly in } H_0^s(\Omega) \text{ as } h \rightarrow \infty.$$

Step 2. We fix now $f \in H^{-s}(\Omega)$ and denote $u_h, u_\infty \in H_0^s(\Omega)$ the solutions of the problems (P_h^f) and (P_∞^f) , respectively.

Since the embedding $H_0^s(\Omega) \subset L^2(\mathbb{R}^n)$ is continuous and dense, so does the embedding $L^2(\mathbb{R}^n) \subset H^{-s}(\Omega)$. Therefore we can find a sequence $(f_j)_j \subset L^2(\mathbb{R}^n)$ such that

$$f_j \rightarrow f \text{ strongly in } H^{-s}(\Omega) \text{ as } j \rightarrow \infty.$$

For all $j \in \mathbb{N}$, let $u_h^j, u_\infty^j \in H_0^s(\Omega)$ be the solutions of the problems $(P_h^{f_j})$ and $(P_\infty^{f_j})$, respectively. Fixed $g \in H^{-s}(\Omega)$, by Proposition 2.8 and Lemma 2.12, we obtain

$$|\langle g, u_h - u_\infty \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)}| \leq |\langle g, u_h^j - u_\infty^j \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)}| + C \|g\|_{H^{-s}(\Omega)} \|f_j - f\|_{H^{-s}(\Omega)}.$$

Hence, in view of (5.2) in Step 1, by letting first $h \rightarrow \infty$ and then $j \rightarrow \infty$, we obtain (2.22), and so

$$(A_h)_h \text{ } G\text{-converges to } A_\infty \text{ in } H_0^s(\Omega).$$

G-convergence implies Γ -convergence. We now show that

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

in accordance with Definition 2.18, by assuming the G -convergence of the associated operators.

Γ -liminf inequality. We fix $u_h, u \in L^2(\mathbb{R}^n)$, $h \in \mathbb{N}$, such that $u_h \rightarrow u$ as $h \rightarrow \infty$ and, without loss of generality and up to a not relabeled subsequence, we assume that

$$\lim_{h \rightarrow \infty} F_h(u_h) = \liminf_{h \rightarrow \infty} F_h(u_h) < \infty \quad \text{and} \quad \sup_{h \in \mathbb{N}} F_h(u_h) \leq C,$$

for some positive constant C , the conclusion being otherwise trivial.

Then, $(u_h)_h \subset H_0^s(\Omega)$ and there exists another positive constant C such that

$$\|u_h\|_{H_0^s(\Omega)} \leq C \quad \text{for all } h \in \mathbb{N},$$

which gives that also $u \in H_0^s(\Omega)$ and

$$u_h \rightarrow u \quad \text{weakly in } H_0^s(\Omega) \text{ as } h \rightarrow \infty.$$

For any $h \in \mathbb{N}$, define

$$(5.3) \quad f := -\operatorname{div}^s(A_\infty \nabla^s u) \in H^{-s}(\Omega)$$

and consider the (unique) weak solution $w_h \in H_0^s(\Omega)$ of problem (P_h^f) . Since

$$(A_h)_h \text{ } G\text{-converges to } A_\infty \text{ in } H_0^s(\Omega)$$

and, by construction, u is the (unique) weak solution of (P_∞^f) , it holds that

$$w_h \rightarrow u \quad \text{weakly in } H_0^s(\mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

From the one hand, we have that

$$\lim_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n} A_\infty(x) \nabla^s u(x) \cdot (2\nabla^s u_h(x) - \nabla^s w_h(x)) \, dx = F_\infty(u).$$

From the other hand, by (5.3)

$$\begin{aligned}
& \int_{\mathbb{R}^n} A_\infty(x) \nabla^s u(x) \cdot (2\nabla^s u_h(x) - \nabla^s w_h(x)) \, dx \\
&= \langle f, 2u_h - w_h \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} = \int_{\mathbb{R}^n} A_h(x) \nabla^s w_h(x) \cdot (2\nabla^s u_h(x) - \nabla^s w_h(x)) \, dx \\
&\leq \int_{\mathbb{R}^n} A_h(x) \nabla^s u_h(x) \cdot \nabla^s u_h(x) \, dx = 2F_h(u_h),
\end{aligned}$$

being

$$A_h(x) \xi_1 \cdot \xi_1 - A_h(x) \xi_2 \cdot (2\xi_1 - \xi_2) = A_h(x) (\xi_1 - \xi_2) \cdot (\xi_1 - \xi_2) \geq 0$$

for a.e. $x \in \mathbb{R}^n$ and all $\xi_1, \xi_2 \in \mathbb{R}^n$.

Hence,

$$\liminf_{h \rightarrow \infty} F_h(u_h) = \lim_{h \rightarrow \infty} F_h(u_h) \geq \lim_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n} A_\infty(x) \nabla^s u(x) \cdot (2\nabla^s u_h(x) - \nabla^s w_h(x)) \, dx = F_\infty(u),$$

which conclude the proof of the Γ -liminf inequality.

Γ -limsup inequality. We fix $u \in H_0^s(\Omega)$, the conclusion being otherwise trivial by the definition of F_∞ , we set $f := -\operatorname{div}(A_\infty \nabla^s u)$, and we consider $w_h \in H_0^s(\Omega)$, (unique) weak solution of (P_h^f) . By the G -convergence of $(A_h)_h$ towards A_∞ in $H_0^s(\Omega)$ and, by Proposition 2.9,

$$w_h \rightarrow u \quad \text{weakly in } H_0^s(\mathbb{R}^n) \quad \text{and} \quad w_h \rightarrow u \quad \text{strongly in } L^2(\mathbb{R}^n) \quad \text{as } h \rightarrow \infty.$$

Moreover,

$$\begin{aligned}
\lim_{h \rightarrow \infty} F_h(w_h) &= \lim_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n} A_h(x) \nabla^s w_h(x) \cdot \nabla^s w_h(x) \, dx \\
&= \lim_{h \rightarrow \infty} \frac{1}{2} \langle f, w_h \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} = \frac{1}{2} \langle f, u \rangle_{H^{-s}(\Omega) \times H_0^s(\Omega)} \\
&= \frac{1}{2} \int_{\mathbb{R}^n} A_\infty(x) \nabla^s u(x) \cdot \nabla^s u(x) \, dx = F_\infty(u).
\end{aligned}$$

Hence the Γ -limsup inequality also holds (the limit is actually achieved) and this concludes the proof of the Γ -convergence of $(F_h)_h$ towards F_∞ . \square

Remark 5.3. If we consider only the equivalence between the nonlocal G -convergence and the Γ -convergence of the associated nonlocal energies $(F_h)_h$, then the assumption of fixing the matrices $(A_h)_h$ outside the reference domain Ω may be dropped. On the other hand, for the compactness of the Γ -convergence (see Theorem 4.1) and for the following equivalence between the nonlocal H -convergence and the Γ -convergence, we need to consider the subclass $\mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$, for a given $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$.

In view of Lemma 5.2, in order to prove Theorem 5.1 it is enough to show that the Γ -convergence of the nonlocal energies $(F_h)_h$ to F_∞ also implies the convergence of momenta, as required in Definition 2.13. To this aim, we follow the strategies adopted in [17, Lemma 4.11] and [3, Theorem 4.5], and we define the functionals $\mathcal{F}_h, \mathcal{F}_\infty : L^2(\mathbb{R}^n; \mathbb{R}^n) \rightarrow \mathbb{R}$, respectively, as

$$\begin{aligned}
\mathcal{F}_h(\Phi) &:= \frac{1}{2} \int_{\mathbb{R}^n} A_h(x) \Phi(x) \cdot \Phi(x) \, dx \quad \text{for all } \Phi \in L^2(\mathbb{R}^n; \mathbb{R}^n), \\
\mathcal{F}_\infty(\Phi) &:= \frac{1}{2} \int_{\mathbb{R}^n} A_\infty(x) \Phi(x) \cdot \Phi(x) \, dx \quad \text{for all } \Phi \in L^2(\mathbb{R}^n; \mathbb{R}^n).
\end{aligned}$$

Let us consider their Fréchet derivatives \mathcal{F}'_h and \mathcal{F}'_∞ , which are given by

$$\mathcal{F}'_h(\Phi)[\Psi] = \int_{\mathbb{R}^n} A_h(x) \Phi(x) \cdot \Psi(x) \, dx \quad \text{and} \quad \mathcal{F}'_\infty(\Phi)[\Psi] = \int_{\mathbb{R}^n} A_\infty(x) \Phi(x) \cdot \Psi(x) \, dx$$

for all $\Phi, \Psi \in L^2(\mathbb{R}^n; \mathbb{R}^n)$.

Note that \mathcal{F}'_h and \mathcal{F}'_∞ identify the momenta for the functionals F_h and F_∞ , respectively. Indeed, given $(u_h)_h \subset H_0^s(\Omega)$ and $u_\infty \in H_0^s(\Omega)$, then the convergence

$$\mathcal{F}'_h(\nabla^s u_h)[\Psi] \rightarrow \mathcal{F}'_\infty(\nabla^s u_\infty)[\Psi] \quad \text{for all } \Psi \in L^2(\Omega; \mathbb{R}^n)$$

is equivalent to

$$A_h \nabla^s u_h \rightarrow A_\infty \nabla^s u_\infty \quad \text{weakly in } L^2(\mathbb{R}^n; \mathbb{R}^n).$$

We have the following result.

Proposition 5.4. *Let $A_0 \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \mathbb{R}^n)$. For any $h \in \mathbb{N}$, let $A_h, A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and $F_h, F_\infty: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ be the nonlocal energies, respectively defined in (2.17) and (2.18). Assume that*

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

and let $(u_h)_h \in H_0^s(\Omega)$ and $u_\infty \in H_0^s(\Omega)$ satisfy

$$(5.4) \quad u_h \rightarrow u_\infty \text{ strongly in } L^2(\mathbb{R}^n) \text{ as } h \rightarrow \infty \quad \text{and} \quad F_h(u_h) \rightarrow F_\infty(u_\infty) \quad \text{as } h \rightarrow \infty.$$

Then, the convergence the momenta associated with $(F_h)_h$ and F_∞ holds, i.e.

$$(5.5) \quad \mathcal{F}'_h(\nabla^s u_h)[\Psi] \rightarrow \mathcal{F}'_\infty(\nabla^s u_\infty)[\Psi] \quad \text{for all } \Psi \in L^2(\mathbb{R}^n; \mathbb{R}^n) \text{ as } h \rightarrow \infty.$$

Proof. To prove (5.5), it is sufficient to show the following inequality

$$(5.6) \quad \mathcal{F}'(\nabla^s u_\infty)[\Psi] \leq \liminf_{h \rightarrow \infty} \mathcal{F}'_h(\nabla^s u_h)[\Psi] \quad \text{for all } \Psi \in L^2(\mathbb{R}^n; \mathbb{R}^n).$$

Indeed, by replacing Ψ with $-\Psi$, and by the properties of the limit inferior, one can get the desired condition (5.5).

For any $h \in \mathbb{N}$, we define the matrix-valued functions

$$B_h := A_h|_\Omega \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega), \quad B_\infty := A_\infty|_\Omega \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega),$$

and the functionals $G_h, G_\infty: L^2(\Omega) \rightarrow [0, \infty]$, which are the local energies associated with B_h and B_∞ , as in (2.19) and (2.20), respectively. For all $\Phi \in L^2(\Omega; \mathbb{R}^n)$, we define the functionals $\mathcal{G}_h^\Phi, \mathcal{G}_\infty^\Phi: L^2(\Omega; \mathbb{R}^n) \rightarrow [0, \infty]$ as

$$\mathcal{G}_h^\Phi(v) := \begin{cases} \frac{1}{2} \int_\Omega A_h(x) (\nabla v(x) + \Phi(x)) \cdot (\nabla v(x) + \Phi(x)) \, dx & \text{if } v \in H^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H^1(\Omega), \end{cases}$$

$$\mathcal{G}_\infty^\Phi(v) := \begin{cases} \frac{1}{2} \int_\Omega A_\infty(x) (\nabla v(x) + \Phi(x)) \cdot (\nabla v(x) + \Phi(x)) \, dx & \text{if } v \in H^1(\Omega), \\ \infty & \text{if } v \in L^2(\Omega) \setminus H^1(\Omega). \end{cases}$$

By Proposition 4.3, we have

$$(G_h)_h \text{ } \Gamma\text{-converges to } G_\infty \text{ strongly in } L^2(\Omega)$$

and, by [3, Theorem 4.2], we derive that for all $\Phi \in L^2(\Omega; \mathbb{R}^n)$ the Γ -convergence in the same topology is also guaranteed for the sequence $(\mathcal{G}_h^\Phi)_h$, having as Γ -limit the functional \mathcal{G}_∞^Φ . In particular, for all $(v_h)_h$ converging to v strongly in $L^2(\Omega)$, we have

$$\mathcal{G}_\infty^\Phi(v) \leq \liminf_{h \rightarrow \infty} \mathcal{G}_h^\Phi(v_h).$$

Let $\Psi \in L^2(\Omega; \mathbb{R}^n)$ and $(t_i)_i$ be a sequence of positive numbers converging to 0, as $i \rightarrow \infty$. Then, $t_i \Psi \in L^2(\Omega; \mathbb{R}^n)$. Moreover, fixed $u_h, u_\infty \in H_0^s(\Omega)$, $h \in \mathbb{N}$, satisfying (5.4), we define

$$v_h := I_{1-s} u_h \in H^1(\Omega), \quad v_\infty := I_{1-s} u_\infty \in H^1(\Omega),$$

where the $H^1(\Omega)$ Sobolev regularity of v_h and v_∞ is guaranteed by Proposition 2.6.

Then, by the continuity of $I_{1-s}: L^2(\mathbb{R}^n) \rightarrow L^2(\Omega)$, it holds that

$$v_h \rightarrow v_\infty \quad \text{strongly in } L^2(\Omega) \text{ as } h \rightarrow \infty$$

and, by the Γ -convergence of $(\mathcal{G}_h^\Phi)_h$ toward \mathcal{G}^Φ (with $\Phi := t_i \Psi$), we get

$$(5.7) \quad \mathcal{G}_\infty^{t_i \Psi}(v_\infty) \leq \liminf_{h \rightarrow \infty} \mathcal{G}_h^{t_i \Psi}(v_h) \quad \text{for all } i \in \mathbb{N}.$$

Since

$$\nabla v_h = \nabla^s u_h \quad \text{a.e. in } \mathbb{R}^n \quad \text{and} \quad \nabla v_\infty = \nabla^s u_\infty \quad \text{a.e. in } \mathbb{R}^n,$$

by Proposition 2.6, we can rephrase (5.7) as

$$\begin{aligned} & \frac{1}{2} \int_{\Omega} A_\infty(x) (\nabla^s u_\infty(x) + t_i \Psi(x)) \cdot (\nabla^s u_\infty(x) + t_i \Psi(x)) \, dx \\ & \leq \liminf_{h \rightarrow \infty} \frac{1}{2} \int_{\Omega} A_h(x) (\nabla^s u_h(x) + t_i \Psi(x)) \cdot (\nabla^s u_h(x) + t_i \Psi(x)) \, dx. \end{aligned}$$

Moreover, the sequence $(u_h)_h$ is uniformly bounded in $H_0^s(\Omega)$, since

$$\frac{\lambda}{2} \|\nabla^s u_h\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)}^2 \leq F_h(u_h) \rightarrow F_\infty(u_\infty) \quad \text{as } h \rightarrow \infty.$$

Hence, since $u_h \rightarrow u$ strongly in $L^2(\mathbb{R}^n)$, we have

$$\nabla^s u_h + t_i \Psi \rightarrow \nabla^s u_\infty + t_i \Psi \quad \text{weakly in } L^2(\mathbb{R}^n) \quad \text{as } h \rightarrow \infty.$$

Therefore, being $A_0 \in \mathcal{M}(\lambda, \Lambda, \mathbb{R}^n)$, we get

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) (\nabla^s u_\infty(x) + t_i \Psi(x)) \cdot (\nabla^s u_\infty(x) + t_i \Psi(x)) \, dx \\ & \leq \liminf_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) (\nabla^s u_h(x) + t_i \Psi(x)) \cdot (\nabla^s u_h(x) + t_i \Psi(x)) \, dx. \end{aligned}$$

Hence,

$$\begin{aligned} \mathcal{F}_\infty(\nabla^s u_\infty + t_i \Psi) &= \frac{1}{2} \int_{\Omega} A_\infty(x) (\nabla^s u_\infty(x) + t_i \Psi(x)) \cdot (\nabla^s u_\infty(x) + t_i \Psi(x)) \, dx \\ & \quad + \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) (\nabla^s u_\infty(x) + t_i \Psi(x)) \cdot (\nabla^s u_\infty(x) + t_i \Psi(x)) \, dx \\ & \leq \liminf_{h \rightarrow \infty} \frac{1}{2} \int_{\Omega} A_h(x) (\nabla^s u_h(x) + t_i \Psi(x)) \cdot (\nabla^s u_h(x) + t_i \Psi(x)) \, dx \\ & \quad + \liminf_{h \rightarrow \infty} \frac{1}{2} \int_{\mathbb{R}^n \setminus \Omega} A_0(x) (\nabla^s u_h(x) + t_i \Psi(x)) \cdot (\nabla^s u_h(x) + t_i \Psi(x)) \, dx \\ & \leq \liminf_{h \rightarrow \infty} \mathcal{F}_h(\nabla^s u_h + t_i \Psi) \end{aligned}$$

and, since

$$\mathcal{F}_h(\nabla^s u_h) = F_h(u_h) \quad \text{and} \quad \mathcal{F}_\infty(\nabla^s u_\infty) = F_\infty(u_\infty),$$

by (5.4) we have

$$\frac{\mathcal{F}_\infty(\nabla^s u_\infty + t_i \Psi) - \mathcal{F}_\infty(\nabla^s u_\infty)}{t_i} \leq \liminf_{h \rightarrow \infty} \frac{\mathcal{F}_h(\nabla^s u_h + t_i \Psi) - \mathcal{F}_h(\nabla^s u_h)}{t_i} \quad \text{for all } i \in \mathbb{N}.$$

Therefore, there exists an increasing sequence of integers $(h_i)_i \subset \mathbb{N}$ such that

$$(5.8) \quad \frac{\mathcal{F}_\infty(\nabla^s u_\infty + t_i \Psi) - \mathcal{F}_\infty(\nabla^s u_\infty)}{t_i} - \frac{1}{i} \leq \frac{\mathcal{F}_h(\nabla^s u_h + t_i \Psi) - \mathcal{F}_h(\nabla^s u_h)}{t_i}$$

for all $h \geq h_i$. If we set $\varepsilon_h := t_i$ for $h_i \leq h < h_{i+1}$ and $i \in \mathbb{N}$, then, by (5.8)

$$(5.9) \quad \liminf_{h \rightarrow \infty} \frac{\mathcal{F}_\infty(\nabla^s u_\infty + \varepsilon_h \Psi) - \mathcal{F}_\infty(\nabla^s u_\infty)}{\varepsilon_h} \leq \liminf_{h \rightarrow \infty} \frac{\mathcal{F}_{h_i}(\nabla^s u_{h_i} + \varepsilon_h \Psi) - \mathcal{F}_{h_i}(\nabla^s u_{h_i})}{\varepsilon_h}.$$

Note now that, by the nature of the functional \mathcal{F}_∞ , the limit inferior on the left hand side of (5.9), is actually achieved and can be represented by the Fréchet derivative of the functional \mathcal{F}_∞ , i.e.

$$(5.10) \quad \mathcal{F}'_\infty(\nabla^s u_\infty)[\Psi] = \lim_{h \rightarrow \infty} \frac{\mathcal{F}_\infty(\nabla^s u_\infty + \varepsilon_h \Psi) - \mathcal{F}_\infty(\nabla^s u_\infty)}{\varepsilon_h}.$$

For what concerns, instead, the right hand side of (5.9), we note that

$$(5.11) \quad \frac{\mathcal{F}_h(\nabla^s u_h + \varepsilon_h \Psi) - \mathcal{F}_h(\nabla^s u_h)}{\varepsilon_h} = \mathcal{F}'_h(\nabla^s u_h)[\Psi] + \varepsilon_h \mathcal{F}_h(\Psi).$$

Since the last term on the right-hand side of (5.11) converges to 0 as $h \rightarrow \infty$, from (5.9)–(5.11) we finally get (5.6). \square

We can finally prove Theorem 5.1.

Proof of Theorem 5.1. We need to prove that the H -convergence of $(A_h)_h$ to A_∞ in $H_0^s(\Omega)$ implies the Γ -convergence of $(F_h)_h$ in the strong topology of $L^2(\mathbb{R}^n)$, and viceversa.

H -convergence implies Γ -convergence. Since the H -convergence is stronger than the G -convergence, this part follows by Lemma 5.2.

Γ -convergence implies H -convergence. As in Lemma 5.2, we first show that (2.13) and (2.14) hold for every $f \in L^2(\mathbb{R}^n)$. Then, by a density argument, we show that (2.13) and (2.14) are satisfied for all $f \in H^{-s}(\Omega)$, leading to the H -convergence of $(A_h)_h$ to A_∞ .

Step 1. We fix $f \in L^2(\mathbb{R}^n)$. By Lemma 5.2, the Γ -convergence of $(F_h)_h$ towards F_∞ implies the weak convergence in $H_0^s(\Omega)$ of u_h , solutions of the problem (P_h^f) , towards u_∞ , solution of the limit problem (P_∞^f) . Moreover, u_h and u_∞ are also, respectively, the unique minimisers of the nonlocal energies F_h^f and F_∞^f defined in (5.1). Hence, by the Fundamental Theorem of Γ -convergence [16, Theorem 7.8], $F_h^f(u_h) \rightarrow F_\infty^f(u_\infty)$ as $h \rightarrow \infty$, which implies that

$$F_h(u_h) \rightarrow F_\infty(u_\infty) \quad \text{as } h \rightarrow \infty.$$

Therefore, in virtue of Proposition 5.4, it holds that

$$A_h \nabla^s u_h \rightarrow A_\infty \nabla^s u_\infty \quad \text{weakly in } L^2(\mathbb{R}^n; \mathbb{R}^n) \text{ as } h \rightarrow \infty,$$

and (2.13)–(2.14) are satisfied for all $f \in L^2(\mathbb{R}^n)$.

Step 2. We fix now $f \in H^{-s}(\Omega)$. For any $h \in \mathbb{N}$, let $u_h, u_\infty \in H_0^s(\Omega)$ be the solutions of the problems (P_h^f) and (P_∞) , respectively. By Lemma 5.2, we already know the validity of the convergence of the solutions (2.13), and it remains to prove the convergence of the momenta (2.14).

Since the embedding $L^2(\mathbb{R}^n) \subset H^{-s}(\Omega)$ is continuous and dense, there exists a sequence $(f_j)_j \subset L^2(\mathbb{R}^n)$ such that as $j \rightarrow \infty$

$$(f_j)_j \text{ strongly converges to } f \text{ in } H^{-s}(\Omega).$$

For all $j \in \mathbb{N}$, let $u_h^j, u_\infty^j \in H_0^s(\Omega)$ be, respectively, the solutions of the problems $(P_h^{f_j})$ and $(P_\infty^{f_j})$. Fixed $\Phi \in L^2(\mathbb{R}^n; \mathbb{R}^n)$, by Proposition 2.8 and Lemma 2.12, we get

$$\begin{aligned} & \left| \int_{\Omega} (A_h(x) \nabla^s u_h(x) - A_\infty(x) \nabla^s u_\infty(x)) \cdot \Phi(x) \, dx \right| \\ & \leq \left| \int_{\Omega} (A_h(x) \nabla^s u_h^j(x) - A_\infty(x) \nabla^s u_\infty^j(x)) \cdot \Phi(x) \, dx \right| + C \|\Phi\|_{L^2(\mathbb{R}^n; \mathbb{R}^n)} \|f_j - f\|_{H^{-s}(\Omega)}, \end{aligned}$$

and by Step 1, by sending first $h \rightarrow \infty$ and then $j \rightarrow \infty$, we obtain (2.14), leading to the H -convergence of the sequence $(A_h)_h$ to A_∞ in $H_0^s(\Omega)$. \square

As a consequence of Lemma 4.2 and Theorem 5.1, we easily derive the uniqueness of the H -limit in the symmetric case.

Corollary 5.5. *Let $A_0 \in \mathcal{M}^{\text{sym}}(\mathbb{R}^n, \lambda, \Lambda)$. For any $h \in \mathbb{N}$, let $A_h, A_\infty, \widehat{A}_\infty \in \mathcal{M}^{\text{sym}}(\Omega, \lambda, \Lambda, A_0)$ be such that*

$$(5.12) \quad (A_h)_h \text{ } H\text{-converges to } A_\infty \text{ in } H_0^s(\Omega), \quad (A_h)_h \text{ } H\text{-converges to } \widehat{A}_\infty \text{ in } H_0^s(\Omega).$$

Then,

$$(5.13) \quad A_\infty(x) = \widehat{A}_\infty(x) \quad \text{for a.e. } x \in \mathbb{R}^n.$$

Proof. Let us assume (5.12). By Theorem 5.1, we derive that

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n)$$

and

$$(F_h)_h \text{ } \Gamma\text{-converges to } \widehat{F}_\infty \text{ strongly in } L^2(\mathbb{R}^n),$$

where, for any $h \in \mathbb{N}$, $F_h, F_\infty, \widehat{F}_\infty: L^2(\mathbb{R}^n) \rightarrow [0, \infty]$ are the nonlocal energies, respectively associated with $A_h, A_\infty, \widehat{A}_\infty$. Since the Γ -limit is unique, we obtain that $F_\infty = \widehat{F}_\infty$, which implies (5.13), in virtue of Lemma 4.2. \square

We conclude this section providing an alternative proof of Theorem 3.2, purely based on the variational techniques introduced in Theorem 4.1 and Theorem 5.1.

Proof of Theorem 3.2. Let $(A_h)_h \subset \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ and let $(F_h)_h$ be the associated nonlocal energies, as in (2.17). By Theorem 4.1, there exists a not relabeled subsequence of $(A_h)_h$ and $A_\infty \in \mathcal{M}^{\text{sym}}(\lambda, \Lambda, \Omega, A_0)$ such that, if F_∞ denotes the nonlocal energy associated with A_∞ , as in (2.18), then

$$(F_h)_h \text{ } \Gamma\text{-converges to } F_\infty \text{ strongly in } L^2(\mathbb{R}^n).$$

We can therefore conclude that

$$(A_h)_h \text{ } H\text{-converges to } A_\infty \text{ in } H_0^s(\Omega),$$

in virtue of Theorem 5.1. \square

6. CONCLUSIONS AND OPEN PROBLEMS

Through our distributional approach, the H -convergence theory extends to linear operators in fractional divergence form. In what follows, we list some possible future research directions stemming from our results that we believe may be of particular interest to the community.

- (1) A first direction that we are planning to investigate concerns the study of the asymptotic behaviour of *monotone operators* in fractional divergence form with superlinear growth, whose local counterpart is presented e.g in [42, Chapter 11]. A key tool useful to characterise the H -limit still as a monotone operator is the Div-Curl Lemma [26, 27], and the lack of an analogous one in the fractional case precludes to prove the H -compactness for this class of operators by standard techniques.
- (2) In Section 5, we proved that the H -compactness in the symmetric case is equivalent to the Γ -compactness of the associated energies. In [2], the authors show that an analogous result can be obtained also in the case of not necessarily symmetric matrices for which, a priori, there is no natural energy associated with the problem. We conjecture that the techniques used in the aforementioned paper can be adapted in the fractional scenario. As a consequence, we would get the uniqueness of the H -limit also for general matrices.
- (3) Once the H -convergence for elliptic operators has been characterised, it is natural to ask whether this can provide information about the asymptotic behaviour of sequences of parabolic nonlocal operators of the form

$$\partial_t - \text{div}^s(B_h(x)\nabla^s).$$

In [23], the authors show that, whenever the sequence of matrix-valued functions $(A_h)_h$ is independent of time, then the parabolic H -limit $B_\infty(x, t)$ coincides with the elliptic H -limit $A_\infty(x)$, meaning that B_∞ is constant in time. Again, the authors conjecture that a similar discussion can be extended to the nonlocal scenario.

- (4) The most famous application of the H -convergence relies in the periodic homogenisation of operators of the type

$$-\text{div}(a(hx)\nabla u(x)), \quad h \in \mathbb{N},$$

where a is 1-periodic. We conjecture that an extension to the distributional fractional setting through techniques similar to those of this paper may be possible. In particular, we think that in this case the hypothesis of fixing a matrix A_0 outside the reference domain, used in the proof of Theorem 3.1, can be relaxed.

- (5) Recently, the H -convergence has been extended also to the sub-Riemannian framework and, more generally, to operators depending on vector fields, see e.g. [22, 23, 24, 25]. Since the definition of fractional operators is more involved in a general sub-Riemannian setting, we plan to extend Theorem 3.1 at least to the case of Carnot groups.

DECLARATIONS

Acknowledgements. The authors warmly thank G. Dal Maso, C. Kreisbeck, F. Paronetto, F. Solombrino and E. Valdinoci for useful suggestions and discussions on this topic.

Conflict of interest. On behalf of all authors, the corresponding author states that there is no conflict of interest.

Funding information. The authors are member of GNAMPA of the Istituto Nazionale di Alta Matematica (INdAM).

M.C. has been founded by the European Union - NextGenerationEU under the Italian Ministry of University and Research (MUR) National National Centre for HPC, Big Data and Quantum Computing (CN_00000013 – CUP: E13C22001000006). M.C. acknowledges also the support of the MUR - PRIN 2022 project “Variational Analysis of Complex Systems in Materials Science, Physics and Biology” (N. 2022HKBF5C), funded by European Union NextGenerationEU, and of the INdAM - GNAMPA 2025 Project “DISCOVERIES - Difetti e Interfacce in Sistemi Continui: un’Ottica Variazionale in Elasticità con Risultati Innovativi ed Efficaci Sviluppi” (Grant Code: CUP_E5324001950001).

A.C. acknowledges the support of the MUR - PRIN 2022 project “Elliptic and parabolic problems, heat kernel estimates and spectral theory” (N. 20223L2NWK), of the INdAM - GNAMPA 2024 Project “Ottimizzazione e disuguaglianze funzionali per problemi geometrico-spettrali locali e nonlocali” (Grant Code: CUP_E53C23001670001).

A.M. is supported by the Spanish State Research Agency, through the Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M), by MCIN/AEI/10.13039/501100011033 (PID2021-123903NB-I00), by Generalitat de Catalunya (2021-SGR-00087).

M.C. and A.M. acknowledge the support of the INdAM - GNAMPA 2024 Project “Pairing and div-curl lemma: extensions to weakly differentiable vector fields and nonlocal differentiation” (Grant Code: CUP_E53C23001670001).

A.C. and A.M. acknowledge the support of the INdAM - GNAMPA 2025 Project “Metodi variazionali per problemi dipendenti da operatori frazionari isotropi e anisotropi” (Grant Code: CUP_E5324001950001).

This research was supported by the Centre de Recerca Matemàtica of Barcelona (CRM), under the International Programme for Research in Groups (IP4RG).

Data availability statement. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- [1] R. ALICANDRO, A. BRAIDES, M. SOLCI, AND G. STEFANI, *Topological singularities arising from fractional-gradient energies*, submitted paper, arXiv: [2309.10112](https://arxiv.org/abs/2309.10112), (2023).
- [2] N. ANSINI, G. DAL MASO, AND C. I. ZEPPIERI, Γ -convergence and H -convergence of linear elliptic operators, *J. Math. Pures Appl.* (9) **99**, 321–329, (2013).
- [3] N. ANSINI, G. DAL MASO, AND C. I. ZEPPIERI, *New results on Γ -limits of integral functionals*, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **31**, 185–202, (2014).
- [4] J. C. BELLIDO AND A. EVGRAFOV, *A simple characterization of H -convergence for a class of nonlocal problems*, *Rev. Mat. Complut.* **34**, 175–183, (2021).
- [5] A. BRAIDES, Γ -convergence for beginners, *Oxford Lecture Series in Mathematics and its Applications* 22. Oxford University Press, Oxford, 2002.
- [6] A. BRAIDES, G. C. BRUSCA, AND D. DONATI, *Another look at elliptic homogenization*, *Milan J. Math.* **92**, 1–23, (2024).
- [7] A. BRAIDES AND G. DAL MASO, *Validity and failure of the integral representation of Γ -limits of convex non-local functionals*, *J. Funct. Anal.* **286**, Paper No. 110317, (2024).

- [8] E. BRUÈ, M. CALZI, G. E. COMI, AND G. STEFANI, *A distributional approach to fractional Sobolev spaces and fractional variation: asymptotics. II*, C. R., Math., Acad. Sci. Paris **360**, 589–626, (2022).
- [9] L. BĂLILESCU, A. GHOSH, AND T. GHOSH, *H-convergence and homogenization of non-local elliptic operators in both perforated and non-perforated domains*, Z. Angew. Math. Phys. **70**, Paper No. 171, (2019).
- [10] C. BUCUR AND E. VALDINOCI, *Nonlocal diffusion and applications*, Lecture Notes of the Unione Matematica Italiana 20. Springer, Unione Matematica Italiana, Bologna, 2016.
- [11] A. CARBOTTI, S. DIPIERRO, AND E. VALDINOCI, *Local density of solutions to fractional equations*, De Gruyter Studies in Mathematics 74. De Gruyter, Berlin, 2019.
- [12] G. E. COMI AND G. STEFANI, *A distributional approach to fractional Sobolev spaces and fractional variation: existence of blow-up*, J. Funct. Anal. **277**, 3373–3435, (2019).
- [13] G. E. COMI AND G. STEFANI, *Leibniz rules and Gauss-Green formulas in distributional fractional spaces*. J. Math. Anal. Appl. **514**, Paper No. 126312, (2022).
- [14] G. E. COMI AND G. STEFANI, *A distributional approach to fractional Sobolev spaces and fractional variation: asymptotics I*. Rev. Mat. Complut. **36**, 491–569, (2023).
- [15] J. CUETO, C. KREISBECK, AND H. SCHÖNBERGER, *A variational theory for integral functionals involving finite-horizon fractional gradients*, Fract. Calc. Appl. Anal. **26**, 2001–2056, (2023).
- [16] G. DAL MASO, *An introduction to Γ -convergence*, Progress in Nonlinear Differential Equations and their Applications. 8. Birkhäuser, Basel, 1993.
- [17] G. DAL MASO, G. A. FRANCFORT, AND R. TOADER, *Quasistatic crack growth in nonlinear elasticity*, Arch. Ration. Mech. Anal. **176**, 165–225, (2005).
- [18] E. DE GIORGI AND S. SPAGNOLO, *Sulla convergenza degli integrali dell'energia per operatori ellittici del secondo ordine*, Boll. Unione Mat. Ital., IV. Ser. **8**, 391–411, (1973).
- [19] J. FERNÁNDEZ BONDER, A. RITORTO, AND A. M. SALORT, *H-convergence result for nonlocal elliptic-type problems via Tartar's method*, SIAM J. Math. Anal. **49**, 2387–2408, (2017).
- [20] M. KASSMANN, A. PIATNITSKI, AND E. ZHIZHINA, *Homogenization of Lévy-type operators with oscillating coefficients*, SIAM J. Math. Anal. **51**, 3641–3665, (2019).
- [21] C. KREISBECK AND H. SCHÖNBERGER, *Quasiconvexity in the fractional calculus of variations: characterization of lower semicontinuity and relaxation*, Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods **215**, Paper No. 112625, (2022).
- [22] A. MAIONE, *H-convergence for equations depending on monotone operators in Carnot groups*, Electron. J. Differ. Equ. **2021**, Paper No. 13, (2021).
- [23] A. MAIONE, F. PARONETTO, AND E. VECCHI, *G-convergence of elliptic and parabolic operators depending on vector fields*, ESAIM, Control Optim. Calc. Var. **29**, Paper No. 8, (2023).
- [24] A. MAIONE, A. PINAMONTI, AND F. SERRA CASSANO, *Γ -convergence for functionals depending on vector fields. I: Integral representation and compactness*, J. Math. Pures Appl. **139**, 109–142, (2020).
- [25] A. MAIONE, A. PINAMONTI, AND F. SERRA CASSANO, *Γ -convergence for functionals depending on vector fields. II: Convergence of minimizers*, SIAM J. Math. Anal. **54**, 5761–5791, (2022).
- [26] F. MURAT, *Compacite par compensation*, Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser. **5**, 489–507, (1978).
- [27] F. MURAT, *Compacite par compensation II*, Recent methods in non-linear analysis, Proc. int. Meet., Rome 1978, 245–256, (1979).
- [28] S. PATRIZI AND E. VALDINOCI, *Homogenization and Orowan's law for anisotropic fractional operators of any order*, Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods **119**, 3–36, (2015).
- [29] E. SANCHEZ-PALENCIA, *Solutions périodiques par rapport aux variables d'espace et applications*, C. R. Acad. Sci., Paris, Sér. A **271**, 1129–1132, (1970).
- [30] E. SANCHEZ-PALENCIA, *Équations aux dérivées partielles dans un type de milieux hétérogènes*, C. R. Acad. Sci., Paris, Sér. A **272**, 1410–1413, (1971).
- [31] E. SANCHEZ-PALENCIA, *Comportements local et macroscopique d'un type de milieux physiques heterogenes*, Int. J. Engin. Sci. **12**, 331–351, (1974).
- [32] O. SAVIN AND E. VALDINOCI, *Γ -convergence for nonlocal phase transitions*, Ann. Inst. Henri Poincaré, Anal. Non Linéaire **29**, 479–500, (2012).
- [33] C. SBORDONE, *Su alcune applicazioni di un tipo di convergenza variazionale*, Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser. **2**, 617–638, (1975).
- [34] T.-T. SHIEH AND D. E. SPECTOR, *On a new class of fractional partial differential equations*, Adv. Calc. Var. **8**, 321–336, (2015).
- [35] T.-T. SHIEH AND D. E. SPECTOR, *On a new class of fractional partial differential equations II*, Adv. Calc. Var. **11**, 289–307, (2018).
- [36] M. SOLCI, *NonGhinteraction vortices*, SIAM J. Math. Anal. **56**, 3430–3451, (2024).
- [37] S. SPAGNOLO, *Sul limite delle soluzioni di problemi di Cauchy relativi all'equazione del calore*, Ann. Sc. Norm. Super. Pisa, Sci. Fis. Mat., III. Ser. **21**, 657–699, (1967).
- [38] S. SPAGNOLO, *Una caratterizzazione degli operatori differenziali autoaggiunti del 2° ordine a coefficienti misurabili e limitati*, Rend. Sem. Mat. Univ. Padova **39**, 56–64, (1968).
- [39] S. SPAGNOLO, *Sulla convergenza di soluzioni di equazioni paraboliche ed ellittiche*, Ann. Sc. Norm. Super. Pisa, Sci. Fis. Mat., III. Ser. **22**, 571–597, (1968); Errata-Corrige. 763, (1968).

- [40] E. M. STEIN, Singular integrals and differentiability properties of functions, Princeton Mathematical Series 30. Princeton University Press, Princeton, N.J., 1970.
- [41] L. TARTAR, *Homogenisation en hydrodynamique*, Singular Perturb. Bound. Layer Theory, Proc. Conf. Lyon 1976, Lect. Notes Math. **594**, 474–481, (1977).
- [42] L. TARTAR, The general theory of homogenization. A personalized introduction. Lecture Notes of the Unione Matematica Italiana 7. Springer, Berlin, 2009.
- [43] M. WAURICK, *Nonlocal H-convergence*, Calc. Var. Partial. Differ. Equ. **57**, Paper No. 6, (2018).
- [44] M. WAURICK, *Nonlocal H-convergence for topologically nontrivial domains*, J. Funct. Anal. **288**, No. 3, (2025).
- [45] M. ŠILHAVÝ, *Fractional vector analysis based on invariance requirements (critique of coordinate approaches)*, Contin. Mech. Thermodyn. **32**, 207–228, (2020).

(M. Caponi) CONTRIBUTING AUTHOR
 DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E MATEMATICA
 UNIVERSITÀ DEGLI STUDI DELL'AQUILA,
 VIA VETOIO 1, COPPITO, 67100 L'AQUILA, ITALY
Email address: `maicol.caponi@univaq.it`

(A. Carbotti) CORRESPONDING AUTHOR
 DIPARTIMENTO DI MATEMATICA E FISICA
 UNIVERSITÀ DEL SALENTO
 VIA PER ARNESANO, 73100 LECCE, ITALY
Email address: `alessandro.carbotti@unisalento.it`

(A. Maione) CONTRIBUTING AUTHOR
 CENTRE DE RECERCA MATEMÀTICA
 EDIFICI C, CAMPUS BELLATERRA, 08193 BELLATERRA, SPAIN
Email address: `amaione@crm.cat`