

GLOBALLY STABLE QUASISTATIC EVOLUTION IN PLASTICITY WITH SOFTENING

G. DAL MASO, A. DESIMONE, M.G. MORA, AND M. MORINI

ABSTRACT. We study a relaxed formulation of the quasistatic evolution problem in the context of small strain associative elastoplasticity with softening. The relaxation takes place in spaces of generalized Young measures. The notion of solution is characterized by the following properties: global stability at each time and energy balance on each time interval. An example developed in detail compares the solutions obtained by this method with the ones provided by a vanishing viscosity approximation, and shows that only the latter capture a decreasing branch in the stress-strain response.

Keywords: quasistatic evolution, rate independent processes, Prandtl-Reuss plasticity, plasticity with softening, shear bands, incremental problems, relaxation, Young measures

2000 Mathematics Subject Classification: 74C05 (28A33, 74G65, 49J45, 35Q72)

CONTENTS

1. Introduction	1
2. Notation and preliminary results	3
2.1. Mathematical preliminaries	3
2.2. Mechanical preliminaries	4
3. Relaxation of the incremental problems	7
3.1. Convex envelope of the nonelastic part	7
3.2. Relaxation with respect to weak convergence	10
3.3. Relaxation in spaces of Young measures	14
3.4. Some structure theorems	15
4. Globally stable quasistatic evolution for Young measures	22
4.1. Definitions and main result	22
4.2. The incremental minimum problems	23
4.3. Further minimality properties	27
4.4. Energy estimates	29
4.5. Proof of the main theorem	31
4.6. Some properties of the solutions	33
5. An example	40
References	43

1. INTRODUCTION

In the study of quasistatic evolution problems for rate independent systems a classical approach is to approximate the continuous time solution by discrete time solutions obtained by solving incremental minimum problems (see the review paper [14] and the references therein).

In this paper we apply this method to the study of a plasticity problem with softening, where the new feature is given by the presence of some nonconvex energy terms. For a general introduction to the mathematical theory of plasticity we refer to [7], [8], [9], [10], and [12]. To focus on the new difficulty, due to the lack of convexity, we consider the simplest relevant model, namely small strain associative elastoplasticity with no applied forces, where the evolution is driven by a time-dependent boundary condition $\mathbf{w}(t)$, prescribed on a portion Γ_0 of the boundary of the reference configuration $\Omega \subset \mathbb{R}^2$.

The unknowns of the problem are the displacement $u: \Omega \rightarrow \mathbb{R}^2$, the elastic strain $e: \Omega \rightarrow \mathbb{M}_{sym}^{2 \times 2}$ (the set of symmetric 2×2 matrices), the plastic strain $p: \Omega \rightarrow \mathbb{M}_D^{2 \times 2}$ (the set of trace free symmetric 2×2 matrices), and the internal variable $z: \Omega \rightarrow \mathbb{R}$. For every given time $t \in [0, T]$ they are related by the kinematic admissibility conditions: $Eu = e + p$ in Ω (additive decomposition) and $u = \mathbf{w}(t)$ on Γ_0 . The stress depends only on the elastic part e through the usual linear relation $\sigma := \mathbb{C}e$, where \mathbb{C} is the elasticity tensor.

Given a sequence of subdivisions of a time interval $[0, T]$

$$0 = t_k^0 < t_k^1 < \dots < t_k^{k-1} < t_k^k = T,$$

we assume that an approximate solution $(u_k^{i-1}, e_k^{i-1}, p_k^{i-1}, z_k^{i-1})$ is known at time t_k^{i-1} . The approximate solution $(u_k^i, e_k^i, p_k^i, z_k^i)$ at time t_k^i is defined as a solution of the following incremental minimum problem:

$$\inf_{(u, e, p, z) \in \mathcal{A}(\mathbf{w}(t_k^i))} \{ \mathcal{Q}(e) + \mathcal{H}(p - p_k^{i-1}, z - z_k^{i-1}) + \mathcal{V}(z) \}, \quad (1.1)$$

where \mathcal{Q} is the stored elastic energy, \mathcal{H} is the plastic dissipation rate, \mathcal{V} is the softening potential, while $\mathcal{A}(\mathbf{w}(t_k^i))$ is the set of functions (u, e, p, z) such that $Eu = e + p$ in Ω , $u = \mathbf{w}(t_k^i)$ on Γ_0 , and $z \in L^1(\Omega)$.

The details of the definition of \mathcal{Q} , \mathcal{H} , \mathcal{V} , together with the technical assumptions which are needed for our analysis, are given in Section 2. For the present discussion it is sufficient to know that \mathcal{Q} is a quadratic form, \mathcal{H} is positively homogeneous of degree one, and \mathcal{V} is strictly concave with linear growth.

Due to the nonconvexity of the functional the infimum in (1.1) is not attained, in general. To overcome this difficulty, in this paper we consider a relaxed formulation of this approach (see Proposition 4.11). To preserve the continuity of the energy terms it is convenient to cast the relaxed problem in the language of Young measures. An additional difficulty is due to the linear growth of \mathcal{H} and \mathcal{V} , which may cause concentration effects. For this reason we formulate the problem in a suitable space of generalized Young measures (see [4, Section 3]).

The next step in our analysis is the study of the convergence of the relaxed approximate solutions as the time step $t_k^i - t_k^{i-1} \rightarrow 0$ as $k \rightarrow \infty$ (uniformly with respect to i). We prove that, up to a subsequence, these solutions converge to a solution of a quasistatic evolution problem formulated in the framework of generalized Young measures. This is characterized by the usual conditions considered in the variational approach to rate independent evolution problems, namely global stability and energy balance (see Definition 4.6), suitably phrased in the language of Young measures. The notion of dissipation required for this purpose is quite delicate and relies on the theory developed in [4].

We also prove that the barycentres of these Young measure solutions define a function $(\mathbf{u}(t), \mathbf{e}(t), \mathbf{p}(t), \mathbf{z}(t))$, where $(\mathbf{u}(t), \mathbf{e}(t), \mathbf{p}(t))$ is a quasistatic evolution of a perfect plasticity problem (see [3]) corresponding to a relaxed dissipation function, denoted $p \mapsto \mathcal{H}_{\text{eff}}(p, 0)$, which can be computed explicitly in terms of \mathcal{H} and \mathcal{V} . Some other qualitative properties of the solutions are investigated at the end of Section 4.

This result allows to compare the globally stable solutions obtained in this paper with the solutions delivered by the vanishing viscosity approach of [5]. In particular, we study in Section 5 the globally stable evolution corresponding to the same data considered in [5, Section 7]. The main differences are the following. While the globally stable solution involves generalized Young measures, the vanishing viscosity evolution takes place in spaces

of affine functions, since the data in the example are spatially homogeneous. The stress $\sigma(t)$ corresponding to the vanishing viscosity solution exhibits a decreasing branch, which accounts for the softening phenomenon. On the contrary, the stress of the globally stable solution is nondecreasing and, after a critical time, it becomes constantly equal to the asymptotic value of the stress of the viscosity solution.

2. NOTATION AND PRELIMINARY RESULTS

2.1. Mathematical preliminaries. We refer to [5] for the standard notation about measures, matrices and functions with bounded deformation. In particular, for every measure μ the symbols μ^a and μ^s always denote the absolutely continuous and the singular part with respect to Lebesgue measure. The former is always identified with its density. The symbol $\|\cdot\|_2$ denotes norm in L^2 , while $\|\cdot\|_1$ denotes the norm in L^1 , as well as in the space M_b of bounded Radon measures. The symbol $\langle \cdot, \cdot \rangle$ denotes a duality pairing depending on the context.

Generalized Young measures. We refer to [4] for the definition and properties of generalized Young measures and of time dependent systems of generalized Young measures. The underlying measure λ will always be the two-dimensional Lebesgue measure \mathcal{L}^2 . In particular we refer to [4, Section 6] for the definition of barycentre of a generalized Young measure, and to [4, Section 3] for the notion of weak* convergence on the space $GY(\overline{U}; \Xi)$ of generalized Young measures on the closure of an open subset U of \mathbb{R}^2 with values in a finite dimensional Hilbert space Ξ .

Given $\nu \in M_b^+(\overline{U})$, $p \in L^1(\overline{U}; \Xi)$, and $\alpha \in L^\infty(\overline{U})$, let $\alpha\omega_p^\nu$ be the element of $M_*(\overline{U} \times \Xi \times \mathbb{R})$ defined by

$$\langle f, \alpha\omega_p^\nu \rangle = \int_{\overline{U}} \alpha(x) f(x, p(x), 0) d\nu(x) \quad (2.1)$$

for every $f \in C^{hom}(\overline{U} \times \Xi \times \mathbb{R})$. Note that $\alpha\omega_p^\nu$ does not belong to $GY(\overline{U}; \Xi)$ since it does not satisfy the projection property (3.3) of [4].

Given $p \in L^1(U; \Xi)$, let $\mathcal{T}_p: U \times \Xi \times \mathbb{R} \rightarrow U \times \Xi \times \mathbb{R}$ be the map defined by $\mathcal{T}_p(x, \xi, \eta) := (x, \xi + \eta p(x), \eta)$. The translation of $\mu \in GY(\overline{U}; \Xi)$ by p is the image $\mathcal{T}_p(\mu)$ of μ under \mathcal{T}_p , that is,

$$\langle f, \mathcal{T}_p(\mu) \rangle = \langle f(x, \xi + \eta p(x), \eta), \mu(x, \xi, \eta) \rangle \quad (2.2)$$

for every $f \in C^{hom}(\overline{U} \times \Xi \times \mathbb{R})$.

Lemma 2.1. *Let $\mu_k, \mu \in GY(\overline{U}; \Xi)$. Assume that $\mu_k \rightharpoonup \mu$ weakly* in $GY(\overline{U}; \Xi)$. Then*

$$\mathcal{T}_p(\mu_k) \rightharpoonup \mathcal{T}_p(\mu) \quad \text{weakly* in } GY(\overline{U}; \Xi) \quad (2.3)$$

for every $p \in L^1(U; \Xi)$.

Proof. For every $\varepsilon > 0$ there exists $p_\varepsilon \in C_0^0(U; \Xi)$ such that $\|p_\varepsilon - p\|_1 < \varepsilon$. By the definition of weak* convergence

$$\langle f(x, \xi + \eta p_\varepsilon(x), \eta), \mu_k(x, \xi, \eta) \rangle \rightarrow \langle f(x, \xi + \eta p_\varepsilon(x), \eta), \mu(x, \xi, \eta) \rangle.$$

for every $f \in C^{hom}(\overline{U} \times \Xi \times \mathbb{R})$. By the projection property (3.3) of [4] we have

$$\begin{aligned} & |\langle f(x, \xi + \eta p_\varepsilon(x), \eta), \mu_k(x, \xi, \eta) \rangle - \langle f(x, \xi + \eta p(x), \eta), \mu_k(x, \xi, \eta) \rangle| \leq \\ & \leq \langle a\eta |p_\varepsilon(x) - p(x)|, \mu_k(x, \xi, \eta) \rangle = \int_U a |p_\varepsilon(x) - p(x)| dx \leq a\varepsilon, \end{aligned}$$

whenever $|f(x, \xi_1, \eta_1) - f(x, \xi_2, \eta_2)| \leq a(|\xi_1 - \xi_2| + |\eta_1 - \eta_2|)$, and the same inequality holds for μ . Since ε is arbitrary, under the same hypothesis on f we obtain

$$\langle f(x, \xi + \eta p(x), \eta), \mu_k(x, \xi, \eta) \rangle \rightarrow \langle f(x, \xi + \eta p(x), \eta), \mu(x, \xi, \eta) \rangle.$$

The conclusion follows from the density result proved in [4, Lemma 2.4]. \square

2.2. Mechanical preliminaries. We now introduce the mechanical notions used in the paper.

The reference configuration. Throughout the paper Ω is a *bounded connected open set* in \mathbb{R}^2 with C^2 *boundary*. Let Γ_0 be a nonempty relatively open subset of $\partial\Omega$ with a finite number of connected components, and let $\Gamma_1 := \partial\Omega \setminus \Gamma_0$.

On Γ_0 we will prescribe a Dirichlet boundary condition. This will be done by assigning a function $w \in H^{1/2}(\Gamma_0; \mathbb{R}^2)$, or, equivalently, a function $w \in H^1(\Omega; \mathbb{R}^2)$, whose trace on Γ_0 (also denoted by w) is the prescribed boundary value. The set Γ_1 will be the traction free part of the boundary.

Admissible stresses and dissipation. Let K be a closed strictly convex set in $\mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ with C^1 boundary. For every value of the internal variable $\zeta \in \mathbb{R}$, the set

$$K(\zeta) := \{\sigma \in \mathbb{M}_D^{2 \times 2} : (\sigma, \zeta) \in K\} \quad (2.4)$$

is interpreted as the *elastic domain* and its boundary as the *yield surface* corresponding to ζ . We assume that there exist two constants A and B , with $0 < A \leq B < \infty$, such that

$$\{(\sigma, \zeta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : |\sigma|^2 + |\zeta|^2 \leq A^2\} \subset K \subset \{(\sigma, \zeta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : |\sigma|^2 + |\zeta|^2 \leq B^2\}. \quad (2.5)$$

We assume in addition that

$$(\sigma, \zeta) \in K \implies (0, \zeta) \in K, \quad (2.6)$$

$$(\sigma, \zeta) \in K \implies (\sigma, -\zeta) \in K. \quad (2.7)$$

Together with convexity, (2.7) yields

$$(\sigma, \zeta) \in K \implies (\sigma, 0) \in K \iff \sigma \in K(0). \quad (2.8)$$

Let $\pi_{\mathbb{R}} : \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \rightarrow \mathbb{R}$ be the projection onto \mathbb{R} . The hypotheses on K imply that there exists a constant $a_K > 0$ such that

$$\pi_{\mathbb{R}}(K) = [-a_K, a_K]. \quad (2.9)$$

The *support function* $H : \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \rightarrow [0, +\infty)$ of K , defined by

$$H(\xi, \theta) := \sup_{(\sigma, \zeta) \in K} \{\sigma : \xi + \zeta \theta\}, \quad (2.10)$$

will play the role of the *dissipation density*. It turns out that H is convex and positively homogeneous of degree one on $\mathbb{M}_D^{2 \times 2} \times \mathbb{R}$. In particular it satisfies the triangle inequality

$$H(\xi_1 + \xi_2, \theta_1 + \theta_2) \leq H(\xi_1, \theta_1) + H(\xi_2, \theta_2). \quad (2.11)$$

Let Φ be the gauge function of K according to [18, Section 4]. Since Φ^2 is strictly convex and differentiable, and $\frac{1}{2}H^2 = (\frac{1}{2}\Phi^2)^*$, by [18, Theorem 26.3] the function H^2 is strictly convex and differentiable, so that the set $\{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : H(\xi, \theta) \leq 1\}$ is strictly convex with C^1 boundary. The same property holds for the sets

$$\{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : H(\xi, \theta) + c\theta \leq 1\} \quad (2.12)$$

for every $c \in \mathbb{R}$.

From (2.5) it follows that

$$A\sqrt{|\xi|^2 + \theta^2} \leq H(\xi, \theta) \leq B\sqrt{|\xi|^2 + \theta^2}, \quad (2.13)$$

from (2.6) and (2.9) we obtain

$$H(\xi, \theta) \geq H(0, \theta) = a_K|\theta|, \quad (2.14)$$

while (2.7) implies

$$H(\xi, \theta) = H(\xi, -\theta). \quad (2.15)$$

It follows from (2.8) and (2.10) that

$$H(\xi, 0) = \sup_{\sigma \in K(0)} \sigma : \xi, \quad (2.16)$$

so that $H(\cdot, 0)$ is the support function of $K(0)$ in $\mathbb{M}_D^{2 \times 2}$.

Using the theory of convex functions of measures developed in [6], we introduce the functional $\mathcal{H}: M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2}) \times M_b(\bar{\Omega}) \rightarrow \mathbb{R}$ defined by

$$\mathcal{H}(p, z) := \int_{\bar{\Omega}} H\left(\frac{dp}{d\lambda}(x), \frac{dz}{d\lambda}(x)\right) d\lambda(x), \quad (2.17)$$

where $\lambda \in M_b^+(\bar{\Omega})$ is any measure such that $p \ll \lambda$ and $z \ll \lambda$ (the homogeneity of H implies that the integral does not depend on λ). Using [6, Theorem 4] and [19, Chapter II, Lemma 5.2] we can see that $\mathcal{H}(p, z)$ coincides with the integral over $\bar{\Omega}$ of the measure studied in [19, Chapter II, Section 4], hence \mathcal{H} is lower semicontinuous on $M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2}) \times M_b(\bar{\Omega})$ with respect to weak* convergence of measures. It follows from the properties of H that \mathcal{H} satisfies the triangle inequality, i.e.,

$$\mathcal{H}(p_1 + p_2, z_1 + z_2) \leq \mathcal{H}(p_1, z_1) + \mathcal{H}(p_2, z_2) \quad (2.18)$$

for every $p_1, p_2 \in M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2})$ and every $z_1, z_2 \in M_b(\bar{\Omega})$.

The elasticity tensor. Let \mathbb{C} be the *elasticity tensor*, considered as a symmetric positive definite linear operator $\mathbb{C}: \mathbb{M}_{sym}^{2 \times 2} \rightarrow \mathbb{M}_{sym}^{2 \times 2}$. We assume that the orthogonal subspaces $\mathbb{M}_D^{2 \times 2}$ and $\mathbb{R}I$ are invariant under \mathbb{C} . This is equivalent to saying that there exist a symmetric positive definite linear operator $\mathbb{C}_D: \mathbb{M}_D^{2 \times 2} \rightarrow \mathbb{M}_D^{2 \times 2}$ and a constant $\kappa > 0$, called *modulus of compression*, such that

$$\mathbb{C}\xi := \mathbb{C}_D\xi_D + \kappa(\text{tr } \xi)I \quad (2.19)$$

for every $\xi \in \mathbb{M}_{sym}^{2 \times 2}$. Note that when \mathbb{C} is isotropic, we have

$$\mathbb{C}\xi = 2\mu\xi_D + \kappa(\text{tr } \xi)I, \quad (2.20)$$

where $\mu > 0$ is the *shear modulus*, so that our assumptions are satisfied.

Let $Q: \mathbb{M}_{sym}^{2 \times 2} \rightarrow [0, +\infty)$ be the quadratic form associated with \mathbb{C} , defined by

$$Q(\xi) := \frac{1}{2}\mathbb{C}\xi : \xi = \frac{1}{2}\mathbb{C}_D\xi_D : \xi_D + \frac{\kappa}{2}(\text{tr } \xi)^2. \quad (2.21)$$

It turns out that there exist two constants $\alpha_{\mathbb{C}}$ and $\beta_{\mathbb{C}}$, with $0 < \alpha_{\mathbb{C}} \leq \beta_{\mathbb{C}} < +\infty$, such that

$$\alpha_{\mathbb{C}}|\xi|^2 \leq Q(\xi) \leq \beta_{\mathbb{C}}|\xi|^2 \quad (2.22)$$

for every $\xi \in \mathbb{M}_{sym}^{2 \times 2}$. These inequalities imply

$$|\mathbb{C}\xi| \leq 2\beta_{\mathbb{C}}|\xi|. \quad (2.23)$$

The softening potential. Let $V: \mathbb{R} \rightarrow \mathbb{R}$ be a function of class C^2 , which will control the evolution of the internal variable ζ , and consequently of the set $K(\zeta)$ of admissible stresses. We assume that there exist two constants $b_V > 0$ and $M_V > 0$ such that for every $\theta \in \mathbb{R}$ and $\tilde{\theta} \in \mathbb{R} \setminus \{0\}$

$$-M_V \leq V''(\theta) \leq 0, \quad (2.24)$$

$$\lim_{\theta \rightarrow -\infty} V'(\theta) = -\lim_{\theta \rightarrow +\infty} V'(\theta) = b_V, \quad (2.25)$$

$$0 < b_V < a_K, \quad (2.26)$$

$$V^\infty(\tilde{\theta}) < V(\theta + \tilde{\theta}) - V(\theta), \quad (2.27)$$

where a_K is the constant in (2.9), and V^∞ denotes the *recession function* of V , defined by

$$V^\infty(\theta) := \lim_{t \rightarrow +\infty} \frac{V(t\theta)}{t} = -b_V|\theta|. \quad (2.28)$$

Note that (2.25) is satisfied when V is even, while (2.27) is satisfied when V is strictly concave. From (2.27) it follows that for every $R > 0$ there exists a constant $c_R > 0$ such that

$$V'(\theta)\tilde{\theta} - V^\infty(\tilde{\theta}) \geq c_R|\tilde{\theta}| \quad (2.29)$$

for every $\theta, \tilde{\theta} \in \mathbb{R}$ with $|\theta| \leq R$.

From (2.14) and (2.26) it follows that there exists a constant $C_V^K > 0$ such that

$$H(\xi_2 - \xi_1, \theta_2 - \theta_1) + V(\theta_2) - V(\theta_1) \geq C_V^K |\xi_2 - \xi_1| + C_V^K |\theta_2 - \theta_1| \quad (2.30)$$

for every $\xi_1, \xi_2 \in \mathbb{M}_D^{2 \times 2}$ and every $\theta_1, \theta_2 \in \mathbb{R}$ (see [5, Subsection 2.2]).

It is convenient to introduce the function $\mathcal{V}^\infty : M_b(\overline{\Omega}) \rightarrow \mathbb{R}$ defined by

$$\mathcal{V}^\infty(z) := \int_{\overline{\Omega}} V^\infty\left(\frac{dz}{d\lambda}\right) d\lambda,$$

where $\lambda \in M_b^+(\overline{\Omega})$ is any measure such that $z \ll \lambda$, and the function $\mathcal{V} : L^1(\Omega) \rightarrow \mathbb{R}$ defined by

$$\mathcal{V}(z) := \int_{\Omega} V(z(x)) dx.$$

The definition is extended to $M_b(\overline{\Omega})$ by setting

$$\mathcal{V}(z) := \mathcal{V}(z^a) + \mathcal{V}^\infty(z^s)$$

for every $z \in M_b(\overline{\Omega})$.

The prescribed boundary displacements. For every $t \in [0, +\infty)$ we prescribe a *boundary displacement* $\mathbf{w}(t)$ in the space $H^1(\Omega; \mathbb{R}^2)$. This choice is motivated by the fact that we do not want to impose “discontinuous” boundary data, so that, if the displacement develops sharp discontinuities, this is due to energy minimization.

We assume also that $\mathbf{w} \in AC_{loc}([0, +\infty); H^1(\Omega; \mathbb{R}^2))$, which means, by definition, that for every $T > 0$ the function \mathbf{w} belongs to the space $AC([0, T]; H^1(\Omega; \mathbb{R}^2))$ of absolutely continuous functions on $[0, T]$ with values in $H^1(\Omega; \mathbb{R}^2)$, so that the time derivative $\dot{\mathbf{w}}$ belongs to $L^1([0, T]; H^1(\Omega; \mathbb{R}^2))$ and its strain $E\dot{\mathbf{w}}$ belongs to $L^1([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}))$. For the main properties of absolutely continuous functions with values in reflexive Banach spaces we refer to [2, Appendix].

Elastic and plastic strains. Given a displacement $u \in BD(\Omega)$ and a boundary datum $w \in H^1(\Omega; \mathbb{R}^2)$, the *elastic strain* $e \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ and the *plastic strain* $p \in M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$ satisfy the *weak kinematic admissibility conditions*

$$Eu = e + p \quad \text{in } \Omega, \quad (2.31)$$

$$p = (w - u) \odot n \mathcal{H}^1 \quad \text{on } \Gamma_0, \quad (2.32)$$

where n is the outward unit normal, \odot denotes the symmetrized tensor product, and \mathcal{H}^1 is the one-dimensional Hausdorff measure. The condition on Γ_0 shows, in particular, that the prescribed boundary condition w is not attained on Γ_0 whenever a plastic slip occurs at the boundary. It follows from (2.31) and (2.32) that $e = E^a u - p^a$ a.e. in Ω and $p^s = E^s u$ in Ω . Since $\text{tr } p = 0$, it follows from (2.31) that $\text{div } u = \text{tr } e \in L^2(\Omega)$ and from (2.32) that $(w - u) \cdot n = 0$ \mathcal{H}^1 -a.e. on Γ_0 , where the dot denotes the scalar product in \mathbb{R}^2 .

Given $w \in H^1(\Omega; \mathbb{R}^2)$, the set $A(w)$ of *admissible displacements and strains* for the boundary datum w on Γ_0 is defined by

$$A(w) := \{(u, e, p) \in BD(\Omega) \times L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}) \times M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2}) : (2.31), (2.32) \text{ hold}\}. \quad (2.33)$$

The set $A_{reg}(w)$ of *regular admissible displacements and strains* is defined as

$$A_{reg}(w) := A(w) \cap (W_{loc}^{1,1}(\Omega; \mathbb{R}^2) \times L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}) \times L^1(\Omega; \mathbb{M}_D^{2 \times 2})). \quad (2.34)$$

Equivalently, $(u, e, p) \in A_{reg}(w)$ if and only if $u \in W_{loc}^{1,1}(\Omega; \mathbb{R}^2) \cap BD(\Omega)$, $e \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $p \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$, $Eu = e + p$ a.e. on Ω , and $u = w$ \mathcal{H}^1 -a.e. on Γ_0 .

The stress. The *stress* $\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ is given by

$$\sigma := \mathbb{C}e = \mathbb{C}_D e_D + \kappa (\text{tr } e) I, \quad (2.35)$$

and the *stored elastic energy* by

$$\mathcal{Q}(e) = \int_{\Omega} Q(e(x)) dx = \frac{1}{2} \langle \sigma, e \rangle. \quad (2.36)$$

It is well known that \mathcal{Q} is lower semicontinuous on $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ with respect to weak convergence.

If $\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ and $\operatorname{div} \sigma \in L^2(\Omega; \mathbb{R}^2)$, then the trace of the normal component of σ on $\partial\Omega$, denoted by $[\sigma n]$, is defined as the distribution on $\partial\Omega$ such that

$$\langle [\sigma n], \psi \rangle_{\partial\Omega} := \langle \operatorname{div} \sigma, \psi \rangle + \langle \sigma, E\psi \rangle \quad (2.37)$$

for every $\psi \in H^1(\Omega; \mathbb{R}^2)$. It turns out that $[\sigma n] \in H^{-1/2}(\partial\Omega; \mathbb{R}^2)$ (see, e.g., [19, Theorem 1.2, Chapter I]). We say that $[\sigma n] = 0$ on Γ_1 if $\langle [\sigma n], \psi \rangle_{\partial\Omega} = 0$ for every $\psi \in H^1(\Omega; \mathbb{R}^2)$ with $\psi = 0$ \mathcal{H}^1 -a.e. on Γ_0 .

3. RELAXATION OF THE INCREMENTAL PROBLEMS

In this section we study different forms of relaxation of the incremental minimum problems.

3.1. Convex envelope of the nonelastic part. In this subsection, given $(\xi_0, \theta_0) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$, we study the convex envelope of the function

$$F(\xi, \theta) := H(\xi - \xi_0, \theta - \theta_0) + V(\theta). \quad (3.1)$$

Setting $\tilde{\xi} = \xi - \xi_0$ and $\tilde{\theta} = \theta - \theta_0$ and subtracting the constant $V(\theta_0)$, it is enough to study the convex envelope of

$$G(\tilde{\xi}, \tilde{\theta}) := H(\tilde{\xi}, \tilde{\theta}) + V(\tilde{\theta} + \theta_0) - V(\theta_0). \quad (3.2)$$

Let G^∞ be the recession function of G , defined by

$$G^\infty(\tilde{\xi}, \tilde{\theta}) := \lim_{t \rightarrow +\infty} \frac{G(t\tilde{\xi}, t\tilde{\theta})}{t}.$$

By the homogeneity of H it follows that

$$G^\infty(\tilde{\xi}, \tilde{\theta}) = H(\tilde{\xi}, \tilde{\theta}) + V^\infty(\tilde{\theta}). \quad (3.3)$$

Lemma 3.1. *For every $(\tilde{\xi}, \tilde{\theta}) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ we have*

$$\operatorname{co} G(\tilde{\xi}, \tilde{\theta}) = \operatorname{co} G^\infty(\tilde{\xi}, \tilde{\theta}), \quad (3.4)$$

where co denotes the convex envelope in $\mathbb{M}_D^{2 \times 2} \times \mathbb{R}$. In particular, $\operatorname{co} G$ does not depend on θ_0 and is positively homogeneous of degree 1 in $(\tilde{\xi}, \tilde{\theta})$.

Proof. As V is concave, we have $V(\tilde{\theta} + \theta_0) - V(\theta_0) \geq V^\infty(\tilde{\theta})$, which gives

$$G \geq G^\infty \quad \text{and} \quad \operatorname{co} G \geq \operatorname{co} G^\infty. \quad (3.5)$$

Since $G \geq 0$ by (2.30) and $G(0, 0) = 0$, we have $\operatorname{co} G(0, 0) = 0$, so that by convexity

$$\operatorname{co} G \leq (\operatorname{co} G)^\infty, \quad (3.6)$$

where $(\operatorname{co} G)^\infty$ is the recession function of $\operatorname{co} G$, defined by

$$(\operatorname{co} G)^\infty(\tilde{\xi}, \tilde{\theta}) := \lim_{t \rightarrow +\infty} \frac{\operatorname{co} G(t\tilde{\xi}, t\tilde{\theta})}{t}.$$

On the other hand, since $\operatorname{co} G \leq G$, we have $(\operatorname{co} G)^\infty \leq G^\infty$, which implies $(\operatorname{co} G)^\infty \leq \operatorname{co} G^\infty$. Therefore, (3.6) gives $\operatorname{co} G \leq \operatorname{co} G^\infty$, which, together with (3.5), yields (3.4). \square

Let us define $H_{\text{eff}} : \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$H_{\text{eff}} := \text{co } G^\infty. \quad (3.7)$$

By the previous lemma the convex envelope $\text{co } F$ of the function F introduced in (3.1) is given by

$$\text{co } F(\xi, \theta) = H_{\text{eff}}(\xi - \xi_0, \theta - \theta_0) + V(\theta_0). \quad (3.8)$$

As H_{eff} is convex and positively homogeneous of degree 1, it can be written in the form

$$H_{\text{eff}}(\xi, \theta) = \sup_{(\sigma, \zeta) \in K_{\text{eff}}} \{\sigma : \xi + \zeta \theta\}, \quad (3.9)$$

where $K_{\text{eff}} = \{(\sigma, \zeta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : H_{\text{eff}}^*(\sigma, \zeta) \leq 0\}$ (see, e.g., [18, Theorem 13.2]), and $H_{\text{eff}}^* = \chi_{K_{\text{eff}}}$, where for every set $E \subset \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ the indicator function χ_E is defined by $\chi_E(\sigma, \zeta) = 0$ if $(\sigma, \zeta) \in E$, $\chi_E(\sigma, \zeta) = +\infty$ otherwise. Since $H_{\text{eff}} = \text{co } G^\infty$, we have that $H_{\text{eff}}^* = (G^\infty)^*$, so that

$$K_{\text{eff}} = \{(\sigma, \zeta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : (G^\infty)^*(\sigma, \zeta) \leq 0\}.$$

Since $V^\infty(\theta) = \min\{b_V \theta, -b_V \theta\}$ by (2.28), the function G^∞ can be expressed as the minimum of two convex functions, namely

$$G^\infty(\xi, \theta) = \min\{H(\xi, \theta) + b_V \theta, H(\xi, \theta) - b_V \theta\}. \quad (3.10)$$

Therefore

$$(G^\infty)^*(\sigma, \zeta) = \max\{H^*(\sigma, \zeta - b_V), H^*(\sigma, \zeta + b_V)\}.$$

Since $H^* = \chi_K$, we obtain

$$\chi_{K_{\text{eff}}} = (G^\infty)^* = \max\{\chi_{K+(0, b_V)}, \chi_{K-(0, b_V)}\},$$

which implies

$$K_{\text{eff}} = (K + (0, b_V)) \cap (K - (0, b_V)). \quad (3.11)$$

Using (2.26), (3.11), and the strict convexity of K , it is easy to check that K_{eff} is a bounded closed convex set and that

$$(0, 0) \in \overset{\circ}{K}_{\text{eff}} \subset K_{\text{eff}} \subset \overset{\circ}{K}. \quad (3.12)$$

Lemma 3.2. *For every $(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ we have*

$$H_{\text{eff}}(\xi, \theta) = G(\xi, \theta) \iff (\xi, \theta) = (0, 0), \quad (3.13)$$

$$H_{\text{eff}}(\xi, 0) = G^\infty(\xi, 0) \iff \xi = 0. \quad (3.14)$$

Proof. By (2.10), (3.9), and (3.12) we have

$$H_{\text{eff}}(\xi, \theta) < H(\xi, \theta) \quad \text{for every } (\xi, \theta) \neq (0, 0). \quad (3.15)$$

If $H_{\text{eff}}(\xi, \theta) = G(\xi, \theta)$, by (3.5) and (3.7) we have $G(\xi, \theta) = G^\infty(\xi, \theta)$, which gives $V^\infty(\theta) = V(\theta + \theta_0) - V(\theta_0)$ thanks to (3.2) and (3.3). By (2.27) this implies $\theta = 0$, so that $H_{\text{eff}}(\xi, 0) = G(\xi, 0) = H(\xi, 0)$. By (3.15) we deduce that $\xi = 0$. This concludes the proof of (3.13).

On the other hand, if $H_{\text{eff}}(\xi, 0) = G^\infty(\xi, 0)$, from (3.3) we obtain $H_{\text{eff}}(\xi, 0) = H(\xi, 0)$. By (3.15) we deduce $\xi = 0$, which concludes the proof of (3.14). \square

Lemma 3.3. *For every $(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ we have*

$$\text{co } G^\infty(\xi, \theta) = \text{co}_\theta G^\infty(\xi, \theta), \quad (3.16)$$

where co_θ denotes the convex envelope with respect to θ . Moreover there exist $\hat{\theta} \in \mathbb{R}$ and $\alpha \in [\frac{1}{2}, 1]$, such that

$$\theta = \alpha \hat{\theta} + (1 - \alpha)(-\hat{\theta}), \quad (3.17)$$

$$H_{\text{eff}}(\xi, \theta) = G^\infty(\xi, \hat{\theta}). \quad (3.18)$$

Proof. Let

$$\begin{aligned} A^\ominus &:= \{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : \theta \leq 0, H(\xi, \theta) + b_V \theta \leq 1\}, \\ A^\oplus &:= \{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : \theta \geq 0, H(\xi, \theta) - b_V \theta \leq 1\}, \end{aligned}$$

and $A := A^\ominus \cup A^\oplus = \{G^\infty \leq 1\}$ (see (3.10)). By (2.15) we have

$$(\xi, \theta) \in A^\ominus \iff (\xi, -\theta) \in A^\oplus. \quad (3.19)$$

Since G^∞ is positively homogeneous of degree 1, we have $\text{co} A = \{\text{co} G^\infty \leq 1\}$ and $\text{co}_\theta A \subset \{\text{co}_\theta G^\infty \leq 1\}$, where $\text{co}_\theta A$ is the smallest set containing A , which is convex with respect to θ , i.e., its intersections with all lines $\{\xi = \text{const.}\}$ are convex. To prove that $\text{co} A = \text{co}_\theta A$, it is enough to show that $\text{co}_\theta A$ is convex. By (3.19) we have that $(\xi, \theta) \in \text{co}_\theta A$ if and only if there exists $\theta^\oplus \in \mathbb{R}$ such that $|\theta| \leq \theta^\oplus$ and $(\xi, \theta^\oplus) \in A^\oplus$. Since A^\oplus is convex, from this property it is easy to deduce that $\text{co}_\theta A$ is convex, hence $\text{co} A = \text{co}_\theta A$. It follows that

$$\text{co}_\theta A \subset \{\text{co}_\theta G^\infty \leq 1\} \subset \{\text{co} G^\infty \leq 1\} = \text{co} A.$$

This implies that $\{\text{co} G^\infty \leq 1\} = \{\text{co}_\theta G^\infty \leq 1\}$. Since both functions $\text{co} G^\infty$ and $\text{co}_\theta G^\infty$ are positively homogeneous of degree 1, we conclude that $\text{co} G^\infty = \text{co}_\theta G^\infty$.

By homogeneity, to prove (3.17) and (3.18) it is not restrictive to assume that $H_{\text{eff}}(\xi, \theta) = 1$, so that $(\xi, \theta) \in \text{co} A$. From the previous discussion it follows that there exists $\theta^\oplus \in \mathbb{R}$ such that $|\theta| \leq \theta^\oplus$ and $(\xi, \theta^\oplus) \in A^\oplus$. In particular we have $H_{\text{eff}}(\xi, -\theta^\oplus) \leq G^\infty(\xi, -\theta^\oplus) \leq 1$ and $H_{\text{eff}}(\xi, \theta^\oplus) \leq G^\infty(\xi, \theta^\oplus) \leq 1$. By convexity we have $H_{\text{eff}}(\xi, -\theta^\oplus) = H_{\text{eff}}(\xi, \theta^\oplus) = 1$, which implies $G^\infty(\xi, -\theta^\oplus) = G^\infty(\xi, \theta^\oplus) = 1$. To conclude the proof of (3.17) and (3.18) it is enough to take $\hat{\theta} = \theta^\oplus$ if $\theta \geq 0$, and $\hat{\theta} = -\theta^\oplus$ if $\theta < 0$. \square

Lemma 3.4. *Let $(\xi_0, \theta_0) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ with $(\xi_0, \theta_0) \neq (0, 0)$. Assume that $H_{\text{eff}}(\xi_0, \theta_0) = G^\infty(\xi_0, \theta_0)$. Then $\theta_0 \neq 0$ and the common tangent hyperplane to the graphs of H_{eff} and G^∞ at the point $(\xi_0, \theta_0, G^\infty(\xi_0, \theta_0))$ is the graph of the linear function*

$$L(\xi, \theta) := \partial_\xi G^\infty(\xi_0, \theta_0) \xi + \partial_\theta G^\infty(\xi_0, \theta_0) \theta.$$

Let $C := \{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : L(\xi, \theta) = G^\infty(\xi, \theta)\}$. Then either $C = \{(\lambda \xi_0, \lambda \theta_0) : \lambda \geq 0\}$ or $C = \{(\lambda \xi_0, \lambda \theta_0) : \lambda \geq 0\} \cup \{(\lambda \xi_0, -\lambda \theta_0) : \lambda \geq 0\}$.

Proof. The inequality $\theta_0 \neq 0$ follows from (3.14). Therefore G^∞ is differentiable at (ξ_0, θ_0) . Using the convexity of H_{eff} and the inequality $H_{\text{eff}} \leq G^\infty$, we deduce that H_{eff} is differentiable at (ξ_0, θ_0) and its partial derivatives coincide with those of G^∞ . The formula for the tangent hyperplane follows easily from the Euler identity.

By (2.15) and (2.28) we may suppose $\theta_0 > 0$. By the homogeneity of the problem it is not restrictive to assume that $H_{\text{eff}}(\xi_0, \theta_0) = G^\infty(\xi_0, \theta_0) = 1$. Then the set $\{L = 1\}$ is the common tangent hyperplane to the hypersurfaces $\{H_{\text{eff}} = 1\}$ and $\{G^\infty = 1\}$ at the point (ξ_0, θ_0) . As $G^\infty(\xi, \theta) = H(\xi, \theta) - b_V \theta$ for $\theta \geq 0$ and the set $\{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : H(\xi, \theta) - b_V \theta \leq 1\}$ is strictly convex by (2.12), we deduce that $\{L = 1\} \cap \{G^\infty = 1\} \cap \{\theta \geq 0\} = \{(\xi_0, \theta_0)\}$. If the set $\{L = 1\} \cap \{G^\infty = 1\} \cap \{\theta < 0\}$ is empty, then $C = \{(\lambda \xi_0, \lambda \theta_0) : \lambda \geq 0\}$ by homogeneity.

Suppose $\{L = 1\} \cap \{G^\infty = 1\} \cap \{\theta < 0\} \neq \emptyset$. Since $L \leq H_{\text{eff}}$ by convexity, if $(\xi_1, \theta_1) \in \{L = 1\} \cap \{G^\infty = 1\} \cap \{\theta < 0\}$ we have $1 = L(\xi_1, \theta_1) \leq H_{\text{eff}}(\xi_1, \theta_1) \leq G^\infty(\xi_1, \theta_1) = 1$. Therefore, the same argument used for (ξ_0, θ_0) shows that

$$\{L = 1\} \cap \{G^\infty = 1\} \cap \{\theta < 0\} = \{(\xi_1, \theta_1)\}. \quad (3.20)$$

This implies

$$\{L = 1\} \cap \{G^\infty = 1\} = \{(\xi_0, \theta_0), (\xi_1, \theta_1)\}. \quad (3.21)$$

Let us prove that $\xi_1 = \xi_0$ and $\theta_1 = -\theta_0$. Let S be the open segment with endpoints (ξ_0, θ_0) and (ξ_1, θ_1) . As $L = 1$ on the endpoints, it is $L = 1$ on S . As $H_{\text{eff}} = 1$ on the endpoints, by convexity we have $H_{\text{eff}} \leq 1$ on S . On the other hand, since the graph of L is tangent to the graph of H_{eff} , by convexity we also have $L \leq H_{\text{eff}}$. Therefore $H_{\text{eff}} = 1$

on S . By (3.21) we have $G^\infty \neq 1$ on S . As $H_{\text{eff}} \leq G^\infty$, we conclude that $H_{\text{eff}} = 1 < G^\infty$ on S .

Let us fix $(\xi, \theta) \in S$. Then $(\xi, \theta) \in \{H_{\text{eff}} \leq 1\} = \text{co}\{G^\infty \leq 1\}$. As $G^\infty(\xi, \theta) > 1$, by the previous lemma there exist θ^\ominus and θ^\oplus with $\theta^\ominus \leq 0 \leq \theta^\oplus$, such that $\theta^\ominus < \theta < \theta^\oplus$, $(\xi, \theta^\ominus) \in \{G^\infty \leq 1\}$, and $(\xi, \theta^\oplus) \in \{G^\infty \leq 1\}$. As $L \leq H_{\text{eff}}$ by convexity and $H_{\text{eff}} \leq G^\infty$ by definition, we have $L(\xi, \theta^\ominus) \leq 1$ and $L(\xi, \theta^\oplus) \leq 1$. Since $\theta^\ominus < \theta < \theta^\oplus$ and $L(\xi, \theta) = 1$, we deduce from the linearity of L that $L(\xi, \theta^\ominus) = L(\xi, \theta^\oplus) = 1$. Using again the inequality $L \leq G^\infty$, we find $G^\infty(\xi, \theta^\ominus) \geq 1$ and $G^\infty(\xi, \theta^\oplus) \geq 1$. Since the opposite inequality follows from the definition of θ^\ominus and θ^\oplus , we also obtain $G^\infty(\xi, \theta^\ominus) = G^\infty(\xi, \theta^\oplus) = 1$. Therefore, (3.21) yields $\xi = \xi_0 = \xi_1$, $\theta^\ominus = \theta_1$, and $\theta^\oplus = \theta_0$. This implies that the straight line $\{(\xi_0, \theta) : \theta \in \mathbb{R}\}$ belongs to the hyperplane $\{L = 1\}$. Since by (2.15) and (2.28) the point $(\xi_0, -\theta_0)$ belongs to $\{G^\infty = 1\}$, we deduce that $\theta_1 = -\theta_0$ by (3.20). This concludes the proof of the equality $\{L = 1\} \cap \{G^\infty = 1\} = \{(\xi_0, \theta_0), (\xi_0, -\theta_0)\}$, which implies that $C = \{(\lambda\xi_0, \lambda\theta_0) : \lambda \geq 0\} \cup \{(\lambda\xi_0, -\lambda\theta_0) : \lambda \geq 0\}$ by homogeneity. \square

Lemma 3.5. *Let $(\xi_0, \theta_0) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$ with $(\xi_0, \theta_0) \neq (0, 0)$. Assume that there exist $\bar{\theta} \geq |\theta_0|$ such that $H_{\text{eff}}(\xi_0, \theta_0) = G^\infty(\xi_0, \bar{\theta})$. Then $\bar{\theta} > 0$. Let $L : \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \rightarrow \mathbb{R}$ be a linear function such that $L \leq H_{\text{eff}}$ and $L(\xi_0, \theta_0) = H_{\text{eff}}(\xi_0, \theta_0)$, and let $C := \{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : L(\xi, \theta) = G^\infty(\xi, \theta)\}$. Then $C \subset \{(\lambda\xi_0, \lambda\bar{\theta}) : \lambda \geq 0\} \cup \{(\lambda\xi_0, -\lambda\bar{\theta}) : \lambda \geq 0\}$.*

Proof. If $\bar{\theta} = |\theta_0|$, the result follows from the previous lemma. If $\bar{\theta} > |\theta_0|$, the affine function $\theta \mapsto L(\xi_0, \theta)$ is bounded from above by $G^\infty(\xi_0, \bar{\theta})$ at the endpoints of the interval $[-\bar{\theta}, \bar{\theta}]$ (recall that $L \leq H_{\text{eff}} \leq G^\infty$) and coincides with $G^\infty(\xi_0, \bar{\theta})$ at the interior point θ_0 . Therefore, $L(\xi_0, \theta) = H_{\text{eff}}(\xi_0, \theta) = G^\infty(\xi_0, \bar{\theta})$ for every $\theta \in [-\bar{\theta}, \bar{\theta}]$. The result follows by the previous lemma with θ_0 replaced by $\bar{\theta}$. \square

3.2. Relaxation with respect to weak convergence. We begin with a result that can be easily deduced from [1]: every (u, e, p) of the admissible set $A(w)$ introduced in (2.33) can be approximated by triples (u_k, e_k, p_k) in the set $A_{\text{reg}}(w)$ introduced in (2.34), so that u_k satisfies the boundary condition $u_k = w$ \mathcal{H}^1 -a.e. on Γ_0 .

Theorem 3.6. *Let $w \in H^1(\Omega; \mathbb{R}^2)$ and let $(u, e, p) \in A(w)$. Then there exists a sequence $(u_k, e_k, p_k) \in A_{\text{reg}}(w)$ such that $u_k \rightharpoonup u$ weakly* in $BD(\Omega)$, $e_k \rightarrow e$ strongly in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, $p_k \rightharpoonup p$ weakly* in $M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2})$, $\|p_k\|_1 \rightarrow \|p\|_1$, and $\|p_k - p^s\|_1 \rightarrow \|p^s\|_1$.*

Proof. By [1, Theorem 5.2] for every k there exists a function $\psi_k \in W^{1,1}(\Omega; \mathbb{R}^2)$ such that $\|\psi_k\|_1 \leq \frac{1}{k}$, $\psi_k = w - u$ \mathcal{H}^1 -a.e. on Γ_0 , $\|\text{div} \psi_k\|_2 \leq \frac{1}{k}$, and

$$\|E\psi_k\|_1 \leq \frac{1}{\sqrt{2}}\|w - u\|_{1, \Gamma_0} + \frac{1}{k} = \|p^s\|_{1, \Gamma_0} + \frac{1}{k},$$

where $\|\cdot\|_{1, \Gamma_0}$ denotes the norms in $L^1_{\mathcal{H}^1}(\Gamma_0; \mathbb{R}^2)$ and in $M_b(\Gamma_0; \mathbb{M}_D^{2 \times 2})$. We define $v_k := u + \psi_k$ and we note that $v_k = w$ \mathcal{H}^1 -a.e. on Γ_0 . By [1, Theorem 5.1] there exists a sequence v_k^m in $BD(\Omega) \cap W_{\text{loc}}^{1,1}(\Omega; \mathbb{R}^2)$, with $v_k^m = v_k = w$ \mathcal{H}^1 -a.e. on Γ_0 , such that $v_k^m \rightarrow v_k$ strongly in $L^1(\Omega; \mathbb{R}^2)$, $\text{div} v_k^m \rightarrow \text{div} v_k$ strongly in $L^2(\Omega)$, $E v_k^m \rightharpoonup E v_k$ weakly* in $M_b(\bar{\Omega}; \mathbb{M}_{\text{sym}}^{2 \times 2})$, and

$$\lim_{m \rightarrow \infty} \|E v_k^m - E^a u - E \psi_k\|_1 = \lim_{m \rightarrow \infty} \|E v_k^m - E^a v_k\|_1 = \|E^s v_k\|_{1, \Omega} = \|p^s\|_{1, \Omega},$$

where $\|\cdot\|_{1, \Omega}$ denotes the norm in $M_b(\Omega; \mathbb{M}_D^{2 \times 2})$. By approximation it is clear that we can find a sequence $m_k \rightarrow \infty$ such that, setting $u_k := v_k^{m_k}$, we have $u_k \in BD(\Omega) \cap W_{\text{loc}}^{1,1}(\Omega; \mathbb{R}^2)$, $u_k = w$ \mathcal{H}^1 -a.e. on Γ_0 , $u_k \rightharpoonup u$ weakly* in $BD(\Omega)$, $\text{div} u_k \rightarrow \text{div} u$ strongly in $L^2(\Omega)$, and

$$\limsup_{k \rightarrow \infty} \|E u_k - E^a u\|_1 \leq \|p^s\|_1. \quad (3.22)$$

Setting $e_k := e_D + \frac{1}{2} \text{div} u_k I$ and $p_k := E u_k - e_k$, we clearly have that $e_k \rightarrow e$ strongly in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$ and $p_k \rightharpoonup p$ weakly* in $M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2})$. Since $E u_k - E^a u = \frac{1}{2} (\text{div} u_k -$

$\operatorname{div} u)I + p_k - p^a$, from (3.22) it follows that

$$\limsup_{k \rightarrow \infty} \|p_k - p^a\|_1 \leq \|p^s\|_1.$$

By lower semicontinuity this implies that $\|p_k - p^a\|_1 \rightarrow \|p^s\|_1$ and $\|p_k\|_1 \rightarrow \|p\|_1$. \square

To deal with the inner variable z we need a technical lemma concerning the approximation of measures on product spaces.

Lemma 3.7. *Let Ξ_1 and Ξ_2 be finite dimensional Hilbert spaces. Let $p_i \in M_b(\overline{\Omega}; \Xi_i)$ for $i = 1, 2$ and let p_1^k be a sequence in $L^1(\Omega; \Xi_1)$ such that $p_1^k \rightharpoonup p_1$ weakly* in $M_b(\overline{\Omega}; \Xi_1)$ and $\|p_1^k\|_1 \rightarrow \|p_1\|_1$. Then, there exists a sequence p_2^k in $L^1(\Omega; \Xi_2)$ such that $p_2^k \rightharpoonup p_2$ weakly* in $M_b(\overline{\Omega}; \Xi_2)$ and $\|(p_1^k, p_2^k)\|_1 \rightarrow \|(p_1, p_2)\|_1$, where the norms are computed in the product Hilbert structure of $\Xi_1 \times \Xi_2$.*

Proof. First of all we observe that $|p_1^k| \rightharpoonup |p_1|$ weakly* in $M_b(\overline{\Omega})$. We decompose p_2 as

$$p_2 = p_{21} + p_{22},$$

with $p_{21}, p_{22} \in M_b(\overline{\Omega}; \Xi_2)$, $|p_{21}| \ll |p_1|$, and $|p_{22}| \perp |p_1|$.

Let us construct a sequence p_{21}^k in $L^1(\Omega; \Xi_2)$ such that $p_{21}^k \rightharpoonup p_{21}$ weakly* in $M_b(\overline{\Omega}; \Xi_2)$ and $\|(p_1^k, p_{21}^k)\|_1 \rightarrow \|(p_1, p_{21})\|_1$. As $|p_{21}| \ll |p_1|$, we have $p_{21} = \psi|p_1|$ for a suitable density $\psi \in L^1_{|p_1|}(\overline{\Omega}; \Xi_2)$. Let ψ_m be a sequence in $C(\overline{\Omega}; \Xi_2)$ which converges to ψ in $L^1_{|p_1|}(\overline{\Omega}; \Xi_2)$, so that $\langle \sqrt{1 + |\psi_m|^2}, |p_1| \rangle \rightarrow \langle \sqrt{1 + |\psi|^2}, |p_1| \rangle = \|(p_1, p_{21})\|_1$, as $m \rightarrow \infty$. For every m let $p_{21}^{km} := \psi_m |p_1^k|$, so that $p_{21}^{km} \rightharpoonup \psi_m |p_1|$ weakly* in $M_b(\overline{\Omega}; \Xi_2)$ and $\|(p_1^k, p_{21}^{km})\|_1 \rightarrow \langle \sqrt{1 + |\psi_m|^2}, |p_1| \rangle$, as $k \rightarrow \infty$. Let $\mathcal{B}_R := \{p \in M_b(\overline{\Omega}; \Xi_2) : \|p\|_1 \leq R\}$, with $R > \|p_2\|_1$. Since ψ_m converges to ψ in $L^1_{|p_1|}(\overline{\Omega}; \Xi_2)$ we have $\psi_m |p_1| \in \mathcal{B}_R$ for m large enough. As $|p_1^k| \rightharpoonup |p_1|$ weakly* in $M_b(\overline{\Omega})$, for these values of m we also have $p_{21}^{km} \in \mathcal{B}_R$ for k large enough. Since the weak* convergence is metrizable on \mathcal{B}_R , we can construct a sequence $m_k \rightarrow \infty$ such that $p_{21}^k := p_{21}^{km_k}$ satisfies the required properties.

Using convolutions it is easy to construct a sequence p_{22}^k in $L^1(\Omega; \Xi_2)$ such that $p_{22}^k \rightharpoonup p_{22}$ weakly* in $M_b(\overline{\Omega}; \Xi_2)$ and $\|p_{22}^k\|_1 \rightarrow \|p_{22}\|_1$.

Let $p_2^k := p_{21}^k + p_{22}^k$. Then $p_2^k \rightharpoonup p_2$ weakly* in $M_b(\overline{\Omega}; \Xi_2)$. It remains to prove that

$$\limsup_{k \rightarrow \infty} \|(p_1^k, p_2^k)\|_1 \leq \|(p_1, p_2)\|_1. \quad (3.23)$$

By the triangle inequality and by the properties of p_{21}^k and p_{22}^k , we have

$$\begin{aligned} \limsup_{k \rightarrow \infty} \|(p_1^k, p_2^k)\|_1 &\leq \lim_{k \rightarrow \infty} \|(p_1^k, p_{21}^k)\|_1 + \lim_{k \rightarrow \infty} \|(0, p_{22}^k)\|_1 = \\ &= \|(p_1, p_{21})\|_1 + \|(0, p_{22})\|_1 = \|(p_1, p_2)\|_1, \end{aligned}$$

where the last equality follows from the fact that the measures (p_1, p_{21}) and $(0, p_{22})$ are mutually singular. \square

Let $\mathcal{H}_{\text{eff}} : M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2}) \times M_b(\overline{\Omega}) \rightarrow \mathbb{R}$ be the functional defined by (2.17) with H replaced by H_{eff} .

Theorem 3.8. *Let $e_0 \in L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, let $z_0 \in M_b(\overline{\Omega})$, let $w \in H^1(\Omega; \mathbb{R}^2)$, let $(u, e, p) \in A(w)$, and let $z \in M_b(\overline{\Omega})$. Then for every $e_k \rightharpoonup e$ weakly in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, $p_k \rightharpoonup p$ weakly* in $M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, $z_k \rightharpoonup z$ weakly* in $M_b(\overline{\Omega})$, we have*

$$\mathcal{Q}(e_0 + e) + \mathcal{H}_{\text{eff}}(p, z) + \mathcal{V}(z_0) \leq \liminf_{k \rightarrow \infty} \{\mathcal{Q}(e_0 + e_k) + \mathcal{H}(p_k, z_k) + \mathcal{V}(z_0 + z_k)\}. \quad (3.24)$$

Moreover, there exist a sequence $(u_k, e_k, p_k) \in A_{\text{reg}}(w)$ and a sequence $z_k \in L^1(\Omega)$ such that $u_k \rightharpoonup u$ weakly* in $BD(\Omega)$, $e_k \rightarrow e$ strongly in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, $p_k \rightharpoonup p$ weakly* in $M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, $z_k \rightharpoonup z$ weakly* in $M_b(\overline{\Omega})$, and

$$\mathcal{Q}(e_0 + e) + \mathcal{H}_{\text{eff}}(p, z) + \mathcal{V}(z_0) \geq \limsup_{k \rightarrow \infty} \{\mathcal{Q}(e_0 + e_k) + \mathcal{H}(p_k, z_k) + \mathcal{V}(z_0 + z_k)\}. \quad (3.25)$$

Proof. Owing to the lower semicontinuity of \mathcal{Q} and \mathcal{H}_{eff} (see the comments after (2.17) and (2.36)), inequality (3.24) follows from the inequality $\mathcal{H}_{\text{eff}}(p_k, z_k) \leq \mathcal{H}(p_k, z_k) + \mathcal{V}(z_0 + z_k) - \mathcal{V}(z_0)$, which is a consequence of (3.4) and (3.7).

We observe that it is enough to prove (3.25) when z_0 belongs to $L^1(\Omega)$ and is piecewise constant on a suitable triangulation. Indeed, there exists a sequence z_0^n of piecewise constant functions which converge to z_0 strongly in $L^1(\Omega)$. For every n let $(u_k^n, e_k^n, p_k^n, z_k^n)$ be a sequence satisfying the second statement of the theorem as $k \rightarrow \infty$, with z_0 replaced by z_0^n . Then

$$\mathcal{Q}(e_0 + e) + \mathcal{H}_{\text{eff}}(p, z) = \lim_{k \rightarrow \infty} \{ \mathcal{Q}(e_0 + e_k^n) + \mathcal{H}(p_k^n, z_k^n) + \mathcal{V}(z_0^n + z_k^n) - \mathcal{V}(z_0^n) \}. \quad (3.26)$$

By (2.25) and by the definition of \mathcal{V} we have

$$\mathcal{V}(z_k^n + z_0^n) - \mathcal{V}(z_0^n) - (\mathcal{V}(z_k^n + z_0) - \mathcal{V}(z_0)) \leq 2b_V \|z_0^n - z_0^a\|_1.$$

Therefore, for every n

$$\begin{aligned} \limsup_{k \rightarrow \infty} \{ \mathcal{Q}(e_0 + e_k^n) + \mathcal{H}(p_k^n, z_k^n) + \mathcal{V}(z_0 + z_k^n) - \mathcal{V}(z_0) \} &\leq \\ &\leq \mathcal{Q}(e_0 + e) + \mathcal{H}_{\text{eff}}(p, z) + 2b_V \|z_0^n - z_0^a\|_1. \end{aligned} \quad (3.27)$$

By a standard double limit procedure it is then easy to construct a sequence (u_k, e_k, p_k, z_k) satisfying the second statement of the theorem.

Moreover, we may also assume that $(u, e, p) \in A_{\text{reg}}(w)$ and $z \in L^1(\Omega)$. Indeed, in the general case, combining Theorem 3.6 with Lemma 3.7 we can construct a sequence $(u_m, e_m, p_m) \in A_{\text{reg}}(w)$ and a sequence $z_m \in L^1(\Omega)$ such that $u_m \rightharpoonup u$ weakly* in $BD(\Omega)$, $e_m \rightarrow e$ strongly in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, $p_m \rightharpoonup p$ weakly* in $M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, $z_m \rightharpoonup z$ weakly* in $M_b(\overline{\Omega})$, and $\|(p_m, z_m)\|_1 \rightarrow \|(p, z)\|_1$. By [17, Theorem 3] (see also [11, Appendix]) these properties imply that

$$\mathcal{Q}(e_0 + e_m) + \mathcal{H}_{\text{eff}}(p_m, z_m) \longrightarrow \mathcal{Q}(e_0 + e) + \mathcal{H}_{\text{eff}}(p, z)$$

and the conclusion of the theorem can be obtained by a standard double limit procedure.

Let us fix a piecewise constant function $z_0 \in L^1(\Omega)$. Let

$$G_0(x, \xi, \theta) := H(\xi, \theta) + V(\theta + z_0(x)) - V(z_0(x)),$$

let

$$G_1(x, \xi, \theta) := \inf_{(\lambda, \xi_1, \xi_2, \theta_1, \theta_2) \in \Lambda} \{ \lambda G_0(x, \xi + \xi_1, \theta + \theta_1) + (1 - \lambda) G_0(x, \xi + \xi_2, \theta + \theta_2) \},$$

and let

$$G_2(x, \xi, \theta) := \inf_{(\lambda, \xi_1, \xi_2, \theta_1, \theta_2) \in \Lambda} \{ \lambda G_1(x, \xi + \xi_1, \theta + \theta_1) + (1 - \lambda) G_1(x, \xi + \xi_2, \theta + \theta_2) \},$$

where Λ is the set of vectors $(\lambda, \xi_1, \xi_2, \theta_1, \theta_2)$ with $0 \leq \lambda \leq 1$, $\xi_1, \xi_2 \in \mathbb{M}_D^{2 \times 2}$, $\theta_1, \theta_2 \in \mathbb{R}$, $\lambda \xi_1 + (1 - \lambda) \xi_2 = 0$, and $\lambda \theta_1 + (1 - \lambda) \theta_2 = 0$. As G_0 is globally Lipschitz continuous in (ξ, θ) , uniformly with respect to x , it follows that G_1 and G_2 satisfy the same property. Moreover, G_1 and G_2 are piecewise constant in x , uniformly with respect to (ξ, θ) . It is easy to see that

$$\text{co}_{(\xi, \theta)} G_0(x, \xi, \theta) \leq G_2(x, \xi, \theta) \leq \sum_{i=1}^4 \lambda_i G_0(x, \xi_i, \theta_i)$$

whenever $(\xi, \theta) = \sum_{i=1}^4 \lambda_i (\xi_i, \theta_i)$ with $\lambda_i \geq 0$ and $\sum_{i=1}^4 \lambda_i = 1$. By the Carathéodory Theorem we conclude that $G_2 = \text{co}_{(\xi, \theta)} G_0 = H_{\text{eff}}$.

To conclude the proof, using a standard double limit procedure, it is enough to show that for every $i = 1, 2$, $(u, e, p) \in A_{\text{reg}}(w)$, $z \in L^1(\Omega)$, and $\eta > 0$ there exist a sequence

$(u_k, e_k, p_k) \in A_{reg}(w)$ and a sequence $z_k \in L^1(\Omega)$ satisfying the properties of the second statement of the theorem and such that

$$\mathcal{G}_i(p, z) + \eta \geq \limsup_{k \rightarrow \infty} \mathcal{G}_{i-1}(p_k, z_k), \quad (3.28)$$

where

$$\mathcal{G}_i(p, z) := \int_{\Omega} G_i(x, p(x), z(x)) dx$$

for $i = 0, 1, 2$.

Using the approximation argument introduced in [13] we can also assume $z \in C^\infty(\Omega) \cap L^1(\Omega)$, $u \in C^\infty(\Omega; \mathbb{R}^2) \cap BD(\Omega)$, $p \in C^\infty(\Omega; \mathbb{M}_D^{2 \times 2}) \cap L^1(\Omega; \mathbb{M}_D^{2 \times 2})$. Using the Lagrange interpolation on a locally finite grid composed by isosceles right triangles which becomes finer and finer near the boundary, we can replace these functions by new functions u , e , p , and z , with $(u, e, p) \in A_{reg}(w)$, such that u is piecewise affine on this triangulation \mathcal{T} , while e , p , and z are piecewise constant. Since z_0 is piecewise constant, it is not restrictive to assume that $G_i(\cdot, \xi, \theta)$ is piecewise constant on \mathcal{T} , so that $G_i(x, \xi, \theta) = G_{i,T}(\xi, \theta)$ for every $x \in T$ and every $T \in \mathcal{T}$. We may assume that every triangle T of the triangulation \mathcal{T} is relatively compact in Ω .

Let us fix $i = 1, 2$ and $T \in \mathcal{T}$. Then

$$u(x) = \xi_T x + c_T \quad \text{for every } x \in T,$$

where ξ_T is a 2×2 -matrix and $c_T \in \mathbb{R}^2$. Moreover, we have

$$e(x) = e_T, \quad p(x) = p_T, \quad z(x) = z_T \quad \text{for every } x \in T,$$

where $e_T \in \mathbb{M}_{sym}^{2 \times 2}$, $p_T \in \mathbb{M}_D^{2 \times 2}$, and $z_T \in \mathbb{R}$. Then we have $\xi_T = e_T + p_T + \omega_T$, where ω_T is a skew symmetric 2×2 -matrix.

For every $\varepsilon > 0$ there exists $(\lambda_T, p_T^1, p_T^2, z_T^1, z_T^2) \in \Lambda$ such that

$$G_{i,T}(p_T, z_T) + \varepsilon > \lambda_T G_{i-1,T}(p_T + p_T^1, z_T + z_T^1) + (1 - \lambda_T) G_{i-1,T}(p_T + p_T^2, z_T + z_T^2). \quad (3.29)$$

By an algebraic property of $\mathbb{M}_D^{2 \times 2}$ there exist $a_T, b_T \in \mathbb{R}^2$ such that $p_T^2 - p_T^1 = a_T \otimes b_T + q_T$ with q_T a skew symmetric 2×2 -matrix. Note that this is the only point where the dimension two is crucial. By a standard lamination procedure with interfaces orthogonal to b_T we can construct two sequences $v_T^k \in W_{loc}^{1,\infty}(\mathbb{R}^2; \mathbb{R}^2)$ and $z_T^k \in L_{loc}^\infty(\mathbb{R}^2)$ such that $v_T^k(0) = 0$, $v_T^k \rightharpoonup p_T x$ weakly* in $W_{loc}^{1,\infty}(\mathbb{R}^2; \mathbb{R}^2)$, $z_T^k \rightharpoonup z_T$ weakly* in $L_{loc}^\infty(\mathbb{R}^2)$, $Ev_T^k = p_T + p_T^1$ and $z_T^k = z_T + z_T^1$ on A_T^k , $Ev_T^k = p_T + p_T^2$ and $z_T^k = z_T + z_T^2$ on $\mathbb{R}^2 \setminus A_T^k$, and $1_{A_T^k} \rightharpoonup \lambda_T$ weakly* in $L_{loc}^\infty(\mathbb{R}^2)$. Let us define $u_T^k(x) := e_T x + v_T^k(x) + \omega_T x + c_T$. Recalling our definitions we find that $u_T^k \rightharpoonup u$ weakly* in $W^{1,\infty}(T; \mathbb{R}^2)$ and $z_T^k \rightharpoonup z$ weakly* in $L^\infty(T)$.

For every $T \in \mathcal{T}$ and every $\delta > 0$ let T_δ be the triangle similar to T with the same centre and similarity ratio $1 - \delta$, and let $\varphi_T^\delta \in C_c^\infty(T)$ a cut-off function such that $\varphi_T^\delta = 1$ on T_δ and $0 \leq \varphi_T^\delta \leq 1$ on T . Let us fix a finite subset $\mathcal{T}' \subset \mathcal{T}$, let

$$\begin{aligned} \Omega' &:= \bigcup_{T \in \mathcal{T}'} T, & \Omega'_\delta &:= \bigcup_{T \in \mathcal{T}'} T_\delta, \\ u_k &:= \sum_{T \in \mathcal{T}'} \varphi_T^\delta u_T^k + \left(1 - \sum_{T \in \mathcal{T}'} \varphi_T^\delta\right) u, & z_k &:= \sum_{T \in \mathcal{T}'} \varphi_T^\delta z_T^k + \left(1 - \sum_{T \in \mathcal{T}'} \varphi_T^\delta\right) z. \end{aligned}$$

It is clear that $u_k \rightharpoonup u$ weakly* in $BD(\Omega)$ and $u_k = w$ \mathcal{H}^1 -a.e. on Γ_0 . We set

$$\begin{aligned} p_k &:= \sum_{T \in \mathcal{T}'} \varphi_T^\delta Ev_T^k + \left(1 - \sum_{T \in \mathcal{T}'} \varphi_T^\delta\right) p, \\ e_k &:= Eu_k - p_k = e + \sum_{T \in \mathcal{T}'} \nabla \varphi_T^\delta \odot (u_T^k - u). \end{aligned}$$

It follows that $p_k \rightharpoonup p$ weakly* in $L^\infty(\Omega; \mathbb{M}_D^{2 \times 2})$ and $e_k \rightarrow e$ strongly in $L^\infty(\Omega; \mathbb{M}_{sym}^{2 \times 2})$. For every $T \in \mathcal{T}'$ we have $e_k = e_T$ a.e. on T_δ , $p_k = p_T + p_T^1$ a.e. on $T_\delta \cap A_T^k$, $z_k = z_T + z_T^1$ a.e. on $T_\delta \cap A_T^k$, $p_k = p_T + p_T^2$ a.e. on $T_\delta \setminus A_T^k$, and $z_k = z_T + z_T^2$ a.e. on $T_\delta \setminus A_T^k$. Therefore

$$\begin{aligned} \mathcal{G}_{i-1}(p_k, z_k) &\leq \sum_{T \in \mathcal{T}'} G_{i-1,T}(p_T + p_T^1, z_T + z_T^1) \mathcal{L}^2(T_\delta \cap A_T^k) + \\ &\quad + \sum_{T \in \mathcal{T}'} G_{i-1,T}(p_T + p_T^2, z_T + z_T^2) \mathcal{L}^2(T_\delta \setminus A_T^k) + \\ &\quad + \int_{\Omega' \setminus \Omega'_\delta} G_{i-1}(p_k, z_k) dx + \int_{\Omega \setminus \Omega'} G_{i-1}(p, z) dx. \end{aligned}$$

We observe that there exists a constant $C(\mathcal{T}')$ such that $G_{i-1}(p_k, z_k) \leq C(\mathcal{T}')$ a.e. on Ω' for every k . As $1_{A_T^k} \rightharpoonup \lambda_T$ weakly* in $L^\infty(T)$ as $k \rightarrow \infty$, using (3.29) we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} \mathcal{G}_{i-1}(p_k, z_k) &\leq \sum_{T \in \mathcal{T}'} (G_{i,T}(p_T, z_T) + \varepsilon) \mathcal{L}^2(T_\delta) + \\ &\quad + C(\mathcal{T}') \mathcal{L}^2(\Omega' \setminus \Omega'_\delta) + \int_{\Omega \setminus \Omega'} G_{i-1}(p, z) dx \leq \\ &\leq \mathcal{G}_i(p, z) + \varepsilon \mathcal{L}^2(\Omega) + C(\mathcal{T}') \mathcal{L}^2(\Omega' \setminus \Omega'_\delta) + \int_{\Omega \setminus \Omega'} G_{i-1}(p, z) dx, \end{aligned}$$

which gives (3.28) with

$$\eta := \varepsilon \mathcal{L}^2(\Omega) + C(\mathcal{T}') \mathcal{L}^2(\Omega' \setminus \Omega'_\delta) + \int_{\Omega \setminus \Omega'} G_{i-1}(p, z) dx.$$

Passing to the limit first as $\delta \rightarrow 0$, then as $\varepsilon \rightarrow 0$, and finally as $\Omega' \nearrow \Omega$, we can make η arbitrarily small, and this concludes the proof. \square

3.3. Relaxation in spaces of Young measures. The following theorem shows the relationships between the incremental problem in $A_{reg}(\tilde{w})$ with H and V , the same problem in $A(\tilde{w})$ with H_{eff} , and a similar problem in a suitable space of generalized Young measures. The statement of the theorem uses the decomposition $\mu = \bar{\mu}^Y + \hat{\mu}^\infty$ of [4, Theorem 4.3], the notion of translation introduced in (2.2), and the homogeneous function $\{V\}: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$\{V\}(\theta, \eta) := \begin{cases} \eta V(\theta/\eta) & \text{if } \eta > 0, \\ V^\infty(\theta) & \text{if } \eta \leq 0. \end{cases} \quad (3.30)$$

Theorem 3.9. *Let $w_0, \tilde{w} \in H^1(\Omega; \mathbb{R}^2)$, let $(u_0, e_0, p_0) \in A(w_0)$, let $z_0 \in M_b(\bar{\Omega})$, let $\mu_0 \in GY(\bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ such that $\text{bar}(\mu_0) = (p_0, z_0)$. Assume that $\bar{\mu}_0^Y = \delta_{(\bar{p}_0, \bar{z}_0)}$ with $\bar{p}_0 \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$ and $\bar{z}_0 \in L^1(\Omega)$. Then the following equalities hold:*

$$\inf_{(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(\tilde{w}), \tilde{z} \in L^1(\Omega)} [\mathcal{Q}(e_0 + \tilde{e}) + \mathcal{H}(\tilde{p}, \tilde{z}) + \langle \{V\}(\theta_1, \eta), \mathcal{T}_{(\tilde{p}, \tilde{z})}(\mu_0) \rangle] = \quad (3.31)$$

$$= \min_{(\tilde{u}, \tilde{e}, \tilde{p}) \in A(\tilde{w}), \tilde{z} \in M_b(\bar{\Omega})} [\mathcal{Q}(e_0 + \tilde{e}) + \mathcal{H}_{\text{eff}}(\tilde{p}, \tilde{z}) + \langle \{V\}(\theta_0, \eta), \mu_0 \rangle] = \quad (3.32)$$

$$= \inf_{(u, e, \mu) \in B} [\mathcal{Q}(e) + \langle H(\xi_1 - \xi_0, \theta_1 - \theta_0) + \{V\}(\theta_1, \eta), \mu_{t_0 t_1} \rangle], \quad (3.33)$$

where the measure $\mathcal{T}_{(\tilde{p}, \tilde{z})}(\mu_0)$ acts on $(x, \xi_1, \theta_1, \eta)$, the measure μ_0 acts on $(x, \xi_0, \theta_0, \eta)$, while the measure $\mu_{t_0 t_1}$ acts on $(x, \xi_0, \theta_0, \xi_1, \theta_1, \eta)$. Here B denotes the class of all triplets (u, e, μ) , with $u \in BD(\Omega)$, $e \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $\mu \in SGY(\{t_0, t_1\}, \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, such that $\mu_{t_0} = \mu_0$ and $(u, e, p) \in A(w_0 + \tilde{w})$, where $(p, z) := \text{bar}(\mu_{t_1})$.

Proof. We start by showing that the infimum in (3.31) is less than or equal to the minimum in (3.32). Let $(\tilde{u}, \tilde{e}, \tilde{p}) \in A(\tilde{w})$ and $\tilde{z} \in M_b(\bar{\Omega})$ be a minimizer of (3.32). By Theorem 3.8

there exist a sequence $(\tilde{u}_m, \tilde{e}_m, \tilde{p}_m) \in A_{reg}(\tilde{w})$ and a sequence $\tilde{z}_m \in L^1(\Omega)$ such that $\tilde{e}_m \rightarrow \tilde{e}$ strongly in $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $\tilde{p}_m \rightarrow \tilde{p}$ weakly* in $M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, $\tilde{z}_m \rightarrow \tilde{z}$ weakly* in $M_b(\overline{\Omega})$, and

$$\mathcal{H}(\tilde{p}_m, \tilde{z}_m) + \mathcal{V}(\bar{z}_0 + \tilde{z}_m) - \mathcal{V}(\bar{z}_0) \longrightarrow \mathcal{H}_{\text{eff}}(\tilde{p}, \tilde{z}). \quad (3.34)$$

We claim that

$$\mathcal{H}(\tilde{p}_m, \tilde{z}_m) + \langle \{V\}(\theta_1, \eta), \mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}(\mu_0) \rangle \longrightarrow \mathcal{H}_{\text{eff}}(\tilde{p}, \tilde{z}) + \langle \{V\}(\theta_0, \eta), \mu_0 \rangle.$$

Indeed, using the definition of $\mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}$, we have

$$\langle \{V\}(\theta_1, \eta), \mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}(\mu_0) \rangle - \langle \{V\}(\theta_0, \eta), \mu_0 \rangle = \langle \{V\}(\theta_0 + \eta \tilde{z}_m(x), \eta) - \{V\}(\theta_0, \eta), \mu_0 \rangle.$$

As $\{V\}(\theta_0 + \eta \tilde{z}_m(x), \eta) - \{V\}(\theta_0, \eta)$ vanishes for $\eta = 0$, we obtain

$$\langle \{V\}(\theta_1, \eta), \mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}(\mu_0) \rangle - \langle \{V\}(\theta_0, \eta), \mu_0 \rangle = \langle \{V\}(\theta_0 + \eta \tilde{z}_m(x), \eta) - \{V\}(\theta_0, \eta), \bar{\mu}_0^Y \rangle.$$

By the assumption $\bar{\mu}_0^Y = \delta_{(\bar{p}_0, \bar{z}_0)}$ we find

$$\langle \{V\}(\theta_1, \eta), \mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}(\mu_0) \rangle - \langle \{V\}(\theta_0, \eta), \mu_0 \rangle = \mathcal{V}(\bar{z}_0 + \tilde{z}_m) - \mathcal{V}(\bar{z}_0). \quad (3.35)$$

From (3.34) and (3.35) we obtain the claim, which, in turn, together with the strong convergence of \tilde{e}_m to \tilde{e} , shows that the infimum (3.31) is less than or equal to the minimum (3.32).

Let $(u, e, \mu) \in B$. By the Jensen inequality for generalized Young measures (see [4, Theorem 6.5]) we have

$$\begin{aligned} \mathcal{H}_{\text{eff}}(p - p_0, z - z_0) &\leq \langle H_{\text{eff}}(\xi_1 - \xi_0, \theta_1 - \theta_0), \mu_{t_0 t_1} \rangle \leq \\ &\leq \langle H(\xi_1 - \xi_0, \theta_1 - \theta_0) + \{V\}(\theta_1, \eta) - \{V\}(\theta_0, \eta), \mu_{t_0 t_1} \rangle. \end{aligned}$$

Since $(u - u_0, e - e_0, p - p_0) \in A(\tilde{w})$, the minimum (3.32) is less than or equal to the infimum in (3.33).

On the other hand the infimum in (3.31) is greater than or equal to the infimum in (3.33), since for every $(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(\tilde{w})$ and every $\tilde{z} \in L^1(\Omega)$ we can construct a triple $(u, e, \mu) \in B$ by setting $u := u_0 + \tilde{u}$, $e := e_0 + \tilde{e}$, and $\mu_{t_0 t_1} := \mathcal{T}_{(\tilde{p}, \tilde{z})}^1(\mu_0)$, where $\mathcal{T}_{(\tilde{p}, \tilde{z})}^1: \overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R} \rightarrow \overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^2 \times \mathbb{R}$ is defined by

$$\mathcal{T}_{(\tilde{p}, \tilde{z})}^1(x, \xi_0, \theta_0, \eta) := (x, \xi_0, \theta_0, \xi_0 + \eta \tilde{p}(x), \theta_0 + \eta \tilde{z}(x), \eta).$$

This concludes the proof of the theorem. \square

3.4. Some structure theorems. We prove now two structure theorems for generalized Young measures whose action on $H + \{V\}$ equals the relaxed functional \mathcal{H}_{eff} evaluated on their barycentres.

Theorem 3.10. *Let $p_0 \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$, $z_0 \in L^1(\Omega)$, $\mu_1 \in GY(\overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, let $(p_1, z_1) := \text{bar}(\mu_1)$, let λ be the total variation of the measure (p_1^s, z_1^s) , and let $(p_1^\lambda, z_1^\lambda)$ be the Radon-Nikodym derivative of the measure (p_1^s, z_1^s) with respect to λ . Assume that*

$$\begin{aligned} \langle H(\xi_1 - \eta p_0(x), \theta_1 - \eta z_0(x)) + \{V\}(\theta_1, \eta), \mu_1(x, \xi_1, \theta_1, \eta) \rangle &= \\ = \mathcal{H}_{\text{eff}}(p_1 - p_0, z_1 - z_0) + \mathcal{V}(z_0). \end{aligned} \quad (3.36)$$

By Lemma 3.3 there exist $z \in L^1(\Omega)$, with $z(z_1^a - z_0) \geq 0$ a.e. on Ω , and $\alpha \in L^\infty(\Omega)$, with $\frac{1}{2} \leq \alpha \leq 1$ a.e. on Ω , such that

$$z_1^a = \alpha(z_0 + z) + (1 - \alpha)(z_0 - z), \quad (3.37)$$

$$H_{\text{eff}}(p_1^a - p_0, z_1^a - z_0) = H(p_1^a - p_0, z) + V^\infty(z) \quad (3.38)$$

a.e. in Ω , and there exist $z_\lambda \in L_\lambda^1(\overline{\Omega})$, with $z_\lambda z_1^\lambda \geq 0$ λ -a.e. on $\overline{\Omega}$, and $\alpha_\lambda \in L_\lambda^\infty(\overline{\Omega})$, with $\frac{1}{2} \leq \alpha_\lambda \leq 1$ λ -a.e. on $\overline{\Omega}$, such that

$$z_1^\lambda = \alpha_\lambda z_\lambda + (1 - \alpha_\lambda)(-z_\lambda), \quad (3.39)$$

$$H_{\text{eff}}(p_1^\lambda, z_1^\lambda) = H(p_1^\lambda, z_\lambda) + V^\infty(z_\lambda) \quad (3.40)$$

λ -a.e. in $\overline{\Omega}$. Then

$$\mu_1 = \delta_{(p_0, z_0)} + \alpha \omega_{(p_1^a - p_0, z)}^{\mathcal{L}^2} + (1 - \alpha) \omega_{(p_1^a - p_0, -z)}^{\mathcal{L}^2} + \alpha_\lambda \omega_{(p_1^\lambda, z_\lambda)}^\lambda + (1 - \alpha_\lambda) \omega_{(p_1^\lambda, -z_\lambda)}^\lambda,$$

that is, according to (2.1),

$$\begin{aligned} \langle f, \mu_1 \rangle &= \int_{\Omega} f(x, p_0(x), z_0(x), 1) dx + \int_{\Omega} \alpha(x) f(x, p_1^a(x) - p_0(x), z(x), 0) dx + \\ &\quad + \int_{\Omega} (1 - \alpha(x)) f(x, p_1^a(x) - p_0(x), -z(x), 0) dx + \\ &\quad + \int_{\overline{\Omega}} \alpha_\lambda(x) f(x, p_1^\lambda(x), z_\lambda(x), 0) d\lambda(x) + \\ &\quad + \int_{\overline{\Omega}} (1 - \alpha_\lambda(x)) f(x, p_1^\lambda(x), -z_\lambda(x), 0) d\lambda(x) \end{aligned} \quad (3.41)$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$ (see [4, Definition 3.14]).

Proof. According to [4, Remark 4.5] there exist $\lambda_1^\infty \in M_b^+(\overline{\Omega})$, a family $(\mu_1^{x,Y})_{x \in \Omega}$ of probability measures on $\mathbb{M}_D^{2 \times 2} \times \mathbb{R}$, and a family $(\mu_1^{x,\infty})_{x \in \overline{\Omega}}$ of probability measures on $\Sigma := \{(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : |\xi|^2 + |\theta|^2 = 1\}$ such that

$$\begin{aligned} \langle f, \mu_1 \rangle &= \int_{\Omega} \left(\int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} f(x, \xi_1, \theta_1, 1) d\mu_1^{x,Y}(\xi_1, \theta_1) \right) dx + \\ &\quad + \int_{\overline{\Omega}} \left(\int_{\Sigma} f(x, \xi_1, \theta_1, 0) d\mu_1^{x,\infty}(\xi_1, \theta_1) \right) d\lambda_1^\infty(x) \end{aligned} \quad (3.42)$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$. According to [4, Remark 6.3], we also have

$$(p_1^a(x), z_1^a(x)) = \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (\xi_1, \theta_1) d\mu_1^{x,Y}(\xi_1, \theta_1) + \lambda_1^{\infty,a}(x) \int_{\Sigma} (\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1)$$

for \mathcal{L}^2 -a.e. $x \in \Omega$, where $\lambda_1^{\infty,a}$ is the absolutely continuous part of λ_1^∞ . This implies

$$\begin{aligned} (p_1^a(x) - p_0(x), z_1^a(x) - z_0(x)) &= \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (\xi_1 - p_0(x), \theta_1 - z_0(x)) d\mu_1^{x,Y}(\xi_1, \theta_1) + \\ &\quad + \lambda_1^{\infty,a}(x) \int_{\Sigma} (\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1). \end{aligned} \quad (3.43)$$

Let $\lambda_1^{\infty,s}$ be the singular part of λ_1^∞ . By [4, Remark 6.3] we also have that $\lambda \ll \lambda_1^{\infty,s}$ and

$$(p_1^\lambda(x), z_1^\lambda(x)) \frac{d\lambda}{d\lambda_1^{\infty,s}}(x) = \int_{\Sigma} (\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) \quad (3.44)$$

for $\lambda_1^{\infty,s}$ -a.e. $x \in \overline{\Omega}$.

Let G^∞ be defined by (3.3). From (3.36) and (3.42) we obtain

$$\begin{aligned} &\int_{\Omega} H_{\text{eff}}(p_1^a(x) - p_0(x), z_1^a(x) - z_0(x)) dx + \int_{\overline{\Omega}} H_{\text{eff}}(p_1^\lambda(x), z_1^\lambda(x)) d\lambda(x) = \\ &= \int_{\Omega} \left(\int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))) d\mu_1^{x,Y}(\xi_1, \theta_1) \right) dx + \\ &\quad + \int_{\Omega} \left(\int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) \right) \lambda_1^{\infty,a}(x) dx + \\ &\quad + \int_{\overline{\Omega}} \left(\int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) \right) d\lambda_1^{\infty,s}(x). \end{aligned} \quad (3.45)$$

As $H_{\text{eff}} = \text{co } G^\infty$, using the homogeneity of H_{eff} and the Jensen inequality, we deduce from (3.43) that for a.e. $x \in \Omega$ we have

$$\begin{aligned} H_{\text{eff}}(p_1^a(x) - p_0, z_1^a(x) - z_0(x)) &\leq \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} G^\infty(\xi_1 - p_0(x), \theta_1 - z_0(x)) d\mu_1^{x,Y}(\xi_1, \theta_1) + \\ &\quad + \lambda_1^{\infty,a}(x) \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi, \theta) \leq \\ &\leq \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))) d\mu_1^{x,Y}(\xi_1, \theta_1) + \\ &\quad + \lambda_1^{\infty,a}(x) \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1), \end{aligned}$$

where the second inequality follows from (3.5). Analogously, from (3.44) we deduce that

$$H_{\text{eff}}(p_1^\lambda(x), z_1^\lambda(x)) \frac{d\lambda}{d\lambda_1^{\infty,s}}(x) \leq \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1)$$

for $\lambda_1^{\infty,s}$ -a.e. $x \in \overline{\Omega}$. Therefore, we deduce from (3.45) that

$$\begin{aligned} &H_{\text{eff}}(p_1^a(x) - p_0(x), z_1^a(x) - z_0(x)) = \\ &= \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))) d\mu_1^{x,Y}(\xi_1, \theta_1) + \\ &\quad + \lambda_1^{\infty,a}(x) \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) \end{aligned} \quad (3.46)$$

for a.e. $x \in \Omega$,

$$H_{\text{eff}}(p_1^\lambda(x), z_1^\lambda(x)) \frac{d\lambda}{d\lambda_1^{\infty,s}}(x) = \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) \quad (3.47)$$

for $\lambda_1^{\infty,s}$ -a.e. $x \in \overline{\Omega}$.

Let us define $A := \{x \in \Omega : p_1^a(x) = p_0(x), z_1^a(x) = z_0(x)\}$ and $A_\lambda := \{x \in \overline{\Omega} : p_1^\lambda(x) = 0, z_1^\lambda(x) = 0\}$. By (3.46) and (3.47) we have

$$\int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))) d\mu_1^{x,Y}(\xi_1, \theta_1) = 0 \text{ for a.e. } x \in A, \quad (3.48)$$

$$\lambda_1^{\infty,a}(x) \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) = 0 \text{ for a.e. } x \in A, \quad (3.49)$$

$$\int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) = 0 \text{ for } \lambda_1^{\infty,s}\text{-a.e. } x \in A_\lambda. \quad (3.50)$$

By (2.30) and (3.48) $\mu_1^{x,Y}$ is concentrated on $(p_0(x), z_0(x))$, hence

$$\mu_1^{x,Y} = \delta_{(p_0(x), z_0(x))} \text{ for a.e. } x \in A. \quad (3.51)$$

Since G^∞ is strictly positive on Σ and $\mu_1^{x,\infty}$ are probability measures, we deduce from (3.49) and (3.50) that

$$\lambda_1^{\infty,a}(x) = 0 \text{ for a.e. } x \in A, \quad (3.52)$$

$$\lambda_1^{\infty,s}(A_\lambda) = 0. \quad (3.53)$$

Let us consider now $\mu_1^{x,Y}$ for $x \in B := \Omega \setminus A$. For every $x \in B$ let $L(x, \cdot, \cdot) : \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \rightarrow \mathbb{R}$ be a linear function such that $L(x, p_1^a(x) - p_0(x), z_1^a(x) - z_0(x)) = H_{\text{eff}}(p_1^a(x) - p_0(x), z_1^a(x) - z_0(x))$ and $L(x, \xi, \theta) \leq H_{\text{eff}}(\xi, \theta)$ for every $(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$. Using (3.43), (3.46), and the

linearity of L , for a.e. $x \in B$ we obtain

$$\begin{aligned} & \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} L(x, \xi_1 - p_0(x), \theta_1 - z_0(x)) d\mu_1^{x,Y}(\xi_1, \theta_1) + \\ & \quad + \lambda_1^{\infty,a}(x) \int_{\Sigma} L(x, \xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) = \\ = & \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} (H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))) d\mu_1^{x,Y}(\xi_1, \theta_1) + \\ & \quad + \lambda_1^{\infty,a}(x) \int_{\Sigma} G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1). \end{aligned} \quad (3.54)$$

Using (3.5) we find $L(x, \xi_1 - p_0(x), \theta_1 - z_0(x)) \leq H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))$. Therefore, equality (3.54) implies that for a.e. $x \in B$ we have

$$L(x, \xi_1 - p_0(x), \theta_1 - z_0(x)) = H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x)) \quad (3.55)$$

for $\mu_1^{x,Y}$ -a.e. $(\xi_1, \theta_1) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$, and

$$\lambda_1^{\infty,a}(x)L(x, \xi_1, \theta_1) = \lambda_1^{\infty,a}(x)G^\infty(\xi_1, \theta_1) \quad (3.56)$$

for $\mu_1^{x,\infty}$ -a.e. $(\xi_1, \theta_1) \in \Sigma$. As $L(x, \xi_1 - p_0(x), \theta_1 - z_0(x)) \leq H_{\text{eff}}(\xi_1 - p_0(x), \theta_1 - z_0(x)) \leq H(\xi_1 - p_0(x), \theta_1 - z_0(x)) + V(\theta_1) - V(z_0(x))$, we deduce from (3.55) and Lemma 3.2 that $(\xi_1, \theta_1) = (p_0(x), z_0(x))$ for $\mu_1^{x,Y}$ -a.e. $(\xi_1, \theta_1) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$. This implies that $\mu_1^{x,Y}$ is concentrated on $(p_0(x), z_0(x))$. Since $\mu_1^{x,Y}$ is a probability measure, we conclude that

$$\mu_1^{x,Y} = \delta_{(p_0(x), z_0(x))} \quad \text{for a.e. } x \in B. \quad (3.57)$$

We now consider the measures $\mu_1^{x,\infty}$. We first observe that $z(x) \neq 0$ for a.e. $x \in B$ by (3.37), (3.38), and Lemma 3.5. For every $x \in B$ we define

$$\varphi(x) := \sqrt{|p_1^\alpha(x) - p_0(x)|^2 + z(x)^2} \quad \text{and} \quad (\hat{p}(x), \hat{z}(x)) := (p_1^\alpha(x) - p_0(x), z(x))/\varphi(x).$$

By (3.56) and Lemma 3.5 for a.e. $x \in B$ with $\lambda_1^{\infty,a}(x) \neq 0$ we have

$$(\xi_1, \theta_1) \in \{(\hat{p}(x), \hat{z}(x)), (\hat{p}(x), -\hat{z}(x))\} \quad \text{for } \mu_1^{x,\infty}\text{-a.e. } (\xi_1, \theta_1) \in \Sigma,$$

so that

$$\mu_1^{x,\infty} = \beta(x)\delta_{(\hat{p}(x), \hat{z}(x))} + (1 - \beta(x))\delta_{(\hat{p}(x), -\hat{z}(x))} \quad (3.58)$$

for a suitable $\beta(x) \in [0, 1]$. Using (3.43) we find that

$$p_1^\alpha - p_0 = \hat{p} \lambda_1^{\infty,a}, \quad z_1^\alpha - z_0 = (2\beta - 1)\hat{z} \lambda_1^{\infty,a} \quad (3.59)$$

a.e. in B . Since $p_1^\alpha - p_0 = \hat{p}\varphi$, the first equality implies that

$$\lambda_1^{\infty,a} = \varphi \quad \text{a.e. in } \{x \in B : \hat{p}(x) \neq 0\}. \quad (3.60)$$

Since $z_1^\alpha - z_0 = (2\alpha - 1)z = (2\alpha - 1)\varphi\hat{z}$ and $\hat{z} \neq 0$, the second equality in (3.59) implies that $\alpha = \beta$ a.e. in $\{x \in B : \hat{p}(x) \neq 0\}$. Therefore,

$$\mu_1^{x,\infty} = \alpha(x)\delta_{(\hat{p}(x), \hat{z}(x))} + (1 - \alpha(x))\delta_{(\hat{p}(x), -\hat{z}(x))} \quad \text{a.e. in } \{x \in B : \hat{p}(x) \neq 0\}. \quad (3.61)$$

As $H_{\text{eff}}(0, \theta) = G^\infty(0, \theta) = (a_K - b_V)|\theta|$ for every $\theta \in \mathbb{R}$ by (2.14), if $\hat{p}(x) = 0$ we deduce from (3.38) that $z(x) = z_1^\alpha(x) - z_0(x)$ and $\alpha(x) = 1$. Then the second equality in (3.59) implies that

$$(2\beta(x) - 1)\lambda_1^{\infty,a}(x) = |z(x)| \quad \text{a.e. in } \{x \in B : \hat{p}(x) = 0\}. \quad (3.62)$$

Therefore, from (3.46) and (3.58) we deduce that

$$\begin{aligned} G^\infty(0, z(x)) &= H_{\text{eff}}(0, z(x)) = \\ &= \lambda_1^{\infty,a}(x) \left(\beta(x)G^\infty\left(0, \frac{z(x)}{|z(x)|}\right) + (1 - \beta(x))G^\infty\left(0, -\frac{z(x)}{|z(x)|}\right) \right) = \\ &= \frac{1}{2\beta(x) - 1} G^\infty(0, z(x)), \end{aligned}$$

hence $\beta = \alpha = 1$ a.e. in $\{x \in B : \hat{p}(x) = 0\}$. By (3.62) we have $\lambda_1^{\infty,a} = |z| = \varphi$ a.e. in $\{x \in B : \hat{p}(x) = 0\}$. Using also (3.58), (3.60), and (3.61) we conclude that

$$\begin{aligned} \mu_1^{x,\infty} &= \alpha(x)\delta_{(\hat{p}(x), \hat{z}(x))} + (1 - \alpha(x))\delta_{(\hat{p}(x), -\hat{z}(x))} \quad \text{a.e. in } B, \\ \lambda_1^{\infty,a} &= \varphi \quad \text{a.e. in } B. \end{aligned} \quad (3.63)$$

Let us consider now the properties of the measure $\mu_1^{x,\infty}$ for $\lambda_1^{\infty,s}$ -a.e. $x \in \overline{\Omega}$. For every $x \in B_\lambda := \overline{\Omega} \setminus A_\lambda$ we define

$$\varphi_\lambda(x) := \sqrt{|p_1^\lambda(x)|^2 + z_\lambda(x)^2} \quad \text{and} \quad (\hat{p}_\lambda(x), \hat{z}_\lambda(x)) := (p_1^\lambda(x), z_\lambda(x))/\varphi_\lambda(x),$$

and notice that $\varphi_\lambda(x) \geq \sqrt{|p_1^\lambda(x)|^2 + z_1^\lambda(x)^2} = 1$. As in the previous step we consider a linear function $L_\lambda(x, \cdot, \cdot): \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $L_\lambda(x, p_1^\lambda(x), z_1^\lambda(x)) = H_{\text{eff}}(p_1^\lambda(x), z_1^\lambda(x))$ and $L_\lambda(x, \xi, \theta) \leq H_{\text{eff}}(\xi, \theta)$ for every $(\xi, \theta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R}$. Using (3.44), (3.47), and the linearity of L_λ , for $\lambda_1^{\infty,s}$ -a.e. $x \in B_\lambda$ we obtain

$$\int_\Sigma L_\lambda(x, \xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1) = \int_\Sigma G^\infty(\xi_1, \theta_1) d\mu_1^{x,\infty}(\xi_1, \theta_1). \quad (3.64)$$

Since $L_\lambda(x, \xi, \theta) \leq G^\infty(\xi, \theta)$, for $\lambda_1^{\infty,s}$ -a.e. $x \in B_\lambda$ we deduce that

$$L_\lambda(x, \xi_1, \theta_1) = G^\infty(\xi_1, \theta_1)$$

for $\mu_1^{x,\infty}$ -a.e. $(\xi_1, \theta_1) \in \Sigma$. By Lemma 3.5 for $\lambda_1^{\infty,s}$ -a.e. $x \in B_\lambda$ we have $z_\lambda(x) \neq 0$ and

$$(\xi_1, \theta_1) \in \{(\hat{p}_\lambda(x), \hat{z}_\lambda(x)), (\hat{p}_\lambda(x), -\hat{z}_\lambda(x))\} \quad \text{for } \mu_1^{x,\infty}\text{-a.e. } (\xi_1, \theta_1) \in \Sigma,$$

so that

$$\mu_1^{x,\infty} = \beta_\lambda(x)\delta_{(\hat{p}_\lambda(x), \hat{z}_\lambda(x))} + (1 - \beta_\lambda(x))\delta_{(\hat{p}_\lambda(x), -\hat{z}_\lambda(x))} \quad (3.65)$$

for a suitable $\beta_\lambda(x) \in [0, 1]$. Using (3.44) we find that

$$p_1^\lambda \frac{d\lambda}{d\lambda_1^{\infty,s}} = \hat{p}_\lambda, \quad z_1^\lambda \frac{d\lambda}{d\lambda_1^{\infty,s}} = (2\beta_\lambda - 1)\hat{z}_\lambda \quad (3.66)$$

$\lambda_1^{\infty,s}$ -a.e. in B_λ . Since $p_1^\lambda = \hat{p}_\lambda \varphi_\lambda$, the first equality implies that

$$\varphi_\lambda \frac{d\lambda}{d\lambda_1^{\infty,s}} = 1 \quad \lambda_1^{\infty,s}\text{-a.e. in } \{x \in B_\lambda : \hat{p}_\lambda(x) \neq 0\}. \quad (3.67)$$

Since $z_1^\lambda = (2\alpha_\lambda - 1)z_\lambda = (2\alpha_\lambda - 1)\varphi_\lambda \hat{z}_\lambda$, the second equality in (3.66) implies that $\alpha_\lambda = \beta_\lambda$ $\lambda_1^{\infty,s}$ -a.e. in $\{x \in B_\lambda : \hat{p}_\lambda(x) \neq 0\}$. Therefore,

$$\begin{aligned} \mu_1^{x,\infty} &= \alpha_\lambda(x)\delta_{(\hat{p}_\lambda(x), \hat{z}_\lambda(x))} + (1 - \alpha_\lambda(x))\delta_{(\hat{p}_\lambda(x), -\hat{z}_\lambda(x))} \\ &\lambda_1^{\infty,s}\text{-a.e. in } \{x \in B_\lambda : \hat{p}_\lambda(x) \neq 0\}. \end{aligned} \quad (3.68)$$

As $H_{\text{eff}}(0, \theta) = G^\infty(0, \theta)$ for every $\theta \in \mathbb{R}$, if $\hat{p}_\lambda(x) = 0$ we deduce from (3.40) that $z_\lambda(x) = z_1^\lambda(x)$ and $\alpha_\lambda(x) = 1$. Then the second equality in (3.66) implies that

$$2\beta_\lambda(x) - 1 = |z_\lambda(x)| \frac{d\lambda}{d\lambda_1^{\infty,s}}(x) \quad \lambda_1^{\infty,s}\text{-a.e. in } \{x \in B_\lambda : \hat{p}_\lambda(x) = 0\}. \quad (3.69)$$

Therefore, from (3.47) and (3.65) we deduce that

$$\begin{aligned} G^\infty(0, z_\lambda(x)) &= H_{\text{eff}}(0, z_\lambda(x)) = \\ &= \frac{|z_\lambda(x)|}{2\beta_\lambda(x) - 1} (\beta_\lambda(x)G^\infty(0, \frac{z_\lambda(x)}{|z_\lambda(x)|}) + (1 - \beta_\lambda(x))G^\infty(0, -\frac{z_\lambda(x)}{|z_\lambda(x)|})) = \\ &= \frac{1}{2\beta_\lambda(x) - 1} G^\infty(0, z_\lambda(x)), \end{aligned}$$

hence $\beta_\lambda = \alpha_\lambda = 1$ $\lambda_1^{\infty,s}$ -a.e. in $\{x \in B_\lambda : \hat{p}_\lambda(x) = 0\}$. By (3.69) we have $\varphi_\lambda \frac{d\lambda}{d\lambda_1^{\infty,s}} = |z_\lambda| \frac{d\lambda}{d\lambda_1^{\infty,s}} = 1$ $\lambda_1^{\infty,s}$ -a.e. in $\{x \in B_\lambda : \hat{p}_\lambda(x) = 0\}$. Using also (3.53), (3.65), (3.67), and

(3.68) we conclude that

$$\begin{aligned} \mu_1^{x,\infty} &= \alpha_\lambda(x)\delta_{(\hat{p}_\lambda(x), \hat{z}_\lambda(x))} + (1 - \alpha_\lambda(x))\delta_{(\hat{p}_\lambda(x), -\hat{z}_\lambda(x))} \quad \lambda_1^{\infty,s}\text{-a.e. in } \overline{\Omega}, \\ \varphi_\lambda \frac{d\lambda}{d\lambda_1^{\infty,s}} &= 1 \quad \lambda_1^{\infty,s}\text{-a.e. in } \overline{\Omega}. \end{aligned} \quad (3.70)$$

It follows that $\lambda_1^{\infty,s} \ll \lambda$ and that

$$\frac{d\lambda_1^{\infty,s}}{d\lambda} = \varphi_\lambda \quad \lambda\text{-a.e. in } \overline{\Omega}. \quad (3.71)$$

The conclusion follows from (3.42), (3.51), (3.52), (3.57), (3.63), (3.70), and (3.71), using the homogeneity of f . \square

To prove the next theorem we need two technical results.

Lemma 3.11. *Let Ξ_1, Ξ_2 be finite dimensional Hilbert spaces and let $\pi_i: \overline{\Omega} \times \Xi_1 \times \Xi_2 \times \mathbb{R} \rightarrow \overline{\Omega} \times \Xi_i \times \mathbb{R}$, $i = 1, 2$, be the projections defined by $\pi_i(x, \xi_1, \xi_2, \eta) := (x, \xi_i, \eta)$. Let $\mu \in GY(\overline{\Omega}; \Xi_1 \times \Xi_2)$ and let $p \in L^1(\Omega; \Xi_1)$. Assume that $\pi_1(\mu) = \delta_p$ and let $\mu_2 := \pi_2(\mu)$. Then*

$$\langle f(x, \xi_1, \xi_2, \eta), \mu(x, \xi_1, \xi_2, \eta) \rangle = \langle f(x, \eta p(x), \xi_2, \eta), \mu_2(x, \xi_2, \eta) \rangle \quad (3.72)$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \Xi_1 \times \Xi_2 \times \mathbb{R})$.

Proof. Using [4, Definition 3.16] and standard arguments in measure theory, it is enough to prove (3.72) for every $f \in C^{hom}(\overline{\Omega} \times \Xi_1 \times \Xi_2 \times \mathbb{R})$.

By the definition of μ_2 we have

$$\langle f(x, \eta p(x), \xi_2, \eta), \mu_2(x, \xi_2, \eta) \rangle = \langle f(x, \eta p(x), \xi_2, \eta), \mu(x, \xi_1, \xi_2, \eta) \rangle.$$

Therefore, to prove (3.72) it is enough to show that

$$\langle f(x, \xi_1, \xi_2, \eta) - f(x, \eta p(x), \xi_2, \eta), \mu(x, \xi_1, \xi_2, \eta) \rangle = 0.$$

By approximation it suffices to prove this equality when f is Lipschitz continuous with respect to ξ_1, ξ_2, η with a constant L independent of x (see [4, Lemma 2.4]). In this case we have

$$|\langle f(x, \xi_1, \xi_2, \eta) - f(x, \eta p(x), \xi_2, \eta), \mu(x, \xi_1, \xi_2, \eta) \rangle| \leq L \langle |\eta p(x) - \xi_1|, \mu(x, \xi_1, \xi_2, \eta) \rangle.$$

As $\pi_1(\mu) = \delta_p$, we have

$$\langle |\eta p(x) - \xi_1|, \mu(x, \xi_1, \xi_2, \eta) \rangle = \langle |\eta p(x) - \xi_1|, \delta_p(x, \xi_1, \eta) \rangle = 0,$$

which concludes the proof. \square

Corollary 3.12. *Let $\Xi_1, \Xi_2, \pi_1, \pi_2, \mu$, and p be as in Lemma 3.11, and let $\mu_1 := \pi_1(\mu)$ and $\mu_2 := \pi_2(\mu)$. Assume that $\overline{\mu}_1^Y = \delta_p$. Then*

$$\langle f(x, \xi_1, \xi_2, \eta), \overline{\mu}^Y(x, \xi_1, \xi_2, \eta) \rangle = \langle f(x, \eta p(x), \xi_2, \eta), \overline{\mu}_2^Y(x, \xi_2, \eta) \rangle$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \Xi_1 \times \Xi_2 \times \mathbb{R})$.

Proof. It is enough to apply Lemma 3.11 to $\overline{\mu}^Y$, using [4, Lemma 4.8]. \square

Theorem 3.13. *Let $\boldsymbol{\mu} \in SGY(\{t_0, t_1\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, let $(p_0, z_0) := \text{bar}(\boldsymbol{\mu}_{t_0})$, and let $(p_1, z_1) := \text{bar}(\boldsymbol{\mu}_{t_1})$. Assume that*

$$\begin{aligned} \langle H(\xi_1 - \xi_0, \theta_1 - \theta_0) + \{V\}(\theta_1, \eta) - \{V\}(\theta_0, \eta), \boldsymbol{\mu}_{t_0 t_1}(x, \xi_0, \theta_0, \xi_1, \theta_1, \eta) \rangle = \\ = \mathcal{H}_{\text{eff}}(p_1 - p_0, z_1 - z_0) \end{aligned} \quad (3.73)$$

and that

$$\overline{\boldsymbol{\mu}}_{t_0}^Y = \delta_{(\overline{p}_0, \overline{z}_0)} \quad (3.74)$$

with $\overline{p}_0 \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$ and $\overline{z}_0 \in L^1(\Omega)$. Then $\overline{\boldsymbol{\mu}}_{t_1}^Y = \delta_{(\overline{p}_0, \overline{z}_0)}$.

Proof. If $p_0 = \bar{p}_0 \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$, $z_0 = \bar{z}_0 \in L^1(\Omega)$, and $\boldsymbol{\mu}_{t_0} = \delta_{(p_0, z_0)}$, then (3.73) implies (3.36) by Lemma 3.11 with $\mu_1 := \boldsymbol{\mu}_{t_1}$, and the conclusion follows from Theorem 3.10.

We consider now the general case. Let $\phi(x, \xi_0, \theta_0, \xi_1, \theta_1, \eta) := (x, \xi_0, \theta_0, \xi_1 - \xi_0, \theta_1 - \theta_0, \eta)$ and let $\pi(x, \xi_0, \theta_0, \tilde{\xi}, \tilde{\theta}, \eta) := (x, \tilde{\xi}, \tilde{\theta}, \eta)$. We define $\nu := (\pi \circ \phi)(\boldsymbol{\mu}_{t_0 t_1})$ and observe that

$$\text{bar}(\nu) = (p_1 - p_0, z_1 - z_0). \quad (3.75)$$

By (3.73) we have

$$\langle H(\tilde{\xi}, \tilde{\theta}) + \{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta), \phi(\boldsymbol{\mu}_{t_0 t_1}) \rangle = \mathcal{H}_{\text{eff}}(p_1 - p_0, z_1 - z_0), \quad (3.76)$$

where the measure $\phi(\boldsymbol{\mu}_{t_0 t_1})$ acts on the variables $(x, \xi_0, \theta_0, \tilde{\xi}, \tilde{\theta}, \eta)$. Moreover, since $\nu = \pi(\phi(\boldsymbol{\mu}_{t_0 t_1}))$, we have that

$$\langle H(\tilde{\xi}, \tilde{\theta}), \phi(\boldsymbol{\mu}_{t_0 t_1}) \rangle = \langle H(\tilde{\xi}, \tilde{\theta}), \nu \rangle, \quad (3.77)$$

where the measure ν acts on the variables $(x, \tilde{\xi}, \tilde{\theta}, \eta)$. We consider the decomposition

$$\begin{aligned} \langle \{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta), \phi(\boldsymbol{\mu}_{t_0 t_1}) \rangle &= \langle \{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta), \overline{\phi(\boldsymbol{\mu}_{t_0 t_1})}^Y \rangle + \\ &+ \langle \{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta), \widehat{\phi(\boldsymbol{\mu}_{t_0 t_1})}^\infty \rangle \end{aligned} \quad (3.78)$$

given by [4, Theorem 4.3]. As $\boldsymbol{\mu}_{t_0}$ is the image of the measure $\phi(\boldsymbol{\mu}_{t_0 t_1})$ under the map $(x, \xi_0, \theta_0, \tilde{\xi}, \tilde{\theta}, \eta) \mapsto (x, \xi_0, \theta_0, \eta)$, by (3.74) we can apply Corollary 3.12 and we obtain

$$\langle \{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta), \overline{\phi(\boldsymbol{\mu}_{t_0 t_1})}^Y \rangle = \langle \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta), \bar{\nu}^Y \rangle. \quad (3.79)$$

Since by concavity $\{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta) \geq V^\infty(\tilde{\theta})$, we have

$$\begin{aligned} \langle \{V\}(\theta_0 + \tilde{\theta}, \eta) - \{V\}(\theta_0, \eta), \widehat{\phi(\boldsymbol{\mu}_{t_0 t_1})}^\infty \rangle &\geq \langle V^\infty(\tilde{\theta}), \widehat{\phi(\boldsymbol{\mu}_{t_0 t_1})}^\infty \rangle = \\ &= \langle V^\infty(\tilde{\theta}), \pi(\widehat{\phi(\boldsymbol{\mu}_{t_0 t_1})}^\infty) \rangle = \langle V^\infty(\tilde{\theta}), \hat{\nu}^\infty \rangle = \\ &= \langle \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta), \hat{\nu}^\infty \rangle, \end{aligned} \quad (3.80)$$

where the second equality follows from [4, Lemma 4.8], taking into account that $\nu = \pi(\phi(\boldsymbol{\mu}_{t_0 t_1}))$. By (3.76)–(3.80) we obtain

$$\mathcal{H}_{\text{eff}}(p_1 - p_0, z_1 - z_0) \geq \langle H(\tilde{\xi}, \tilde{\theta}) + \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta), \nu \rangle.$$

By the Jensen inequality for generalized Young measures [4, Theorem 6.5] we deduce from (3.75) that

$$\mathcal{H}_{\text{eff}}(p_1 - p_0, z_1 - z_0) = \langle H(\tilde{\xi}, \tilde{\theta}) + \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta), \nu \rangle. \quad (3.81)$$

Let us fix $x \in \Omega$ and let $G: \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by

$$G(\tilde{\xi}, \tilde{\theta}, \eta) := H(\tilde{\xi}, \tilde{\theta}) + \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta).$$

It follows from (3.4) and (3.7) that the function $H_{\text{eff}}(\tilde{\xi}, \tilde{\theta})$ is the convex envelope of $G(\tilde{\xi}, \tilde{\theta}, \eta)$ with respect to $(\tilde{\xi}, \tilde{\theta}, \eta)$. Moreover, by (3.13) we deduce that for every $\eta > 0$ the equality

$$H_{\text{eff}}(\tilde{\xi}, \tilde{\theta}) = H(\tilde{\xi}, \tilde{\theta}) + \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta)$$

holds if and only if $(\tilde{\xi}, \tilde{\theta}) = (0, 0)$. Therefore (3.75), (3.81), and [4, Lemma 6.7] imply that

$$\text{supp } \nu \subset \{(x, 0, 0, \eta) : x \in \bar{\Omega}, \eta \geq 0\} \cup (\bar{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \{0\}),$$

and, in particular,

$$\text{supp } \bar{\nu}^Y \subset \{(x, 0, 0, \eta) : x \in \bar{\Omega}, \eta \geq 0\},$$

hence $\bar{\nu}^Y = \delta_{(0,0)}$.

From the definitions of ν and from [4, Lemma 4.8] it follows that

$$\begin{aligned} \langle f, \delta_{(\bar{p}_0, \bar{z}_0)} \rangle &= \int_{\Omega} f(x, \bar{p}_0(x), \bar{z}_0(x), 1) dx = \langle f(x, \tilde{\xi} + \eta \bar{p}_0(x), \tilde{\theta} + \eta \bar{z}_0(x), \eta), \bar{\nu}^Y \rangle = \\ &= \langle f(x, \xi_1 - \xi_0 + \eta \bar{p}_0(x), \theta_1 - \theta_0 + \eta \bar{z}_0(x), \eta), \bar{\boldsymbol{\mu}}_{t_0 t_1}^Y \rangle \end{aligned} \quad (3.82)$$

for every $f \in C^{hom}(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$. By (3.74) we can apply Corollary 3.12 and we obtain that the last term in the previous formula equals $\langle f(x, \xi_1, \theta_1, \eta), \overline{\boldsymbol{\mu}}_{t_1}^Y \rangle$. Therefore, $\overline{\boldsymbol{\mu}}_{t_1}^Y = \delta_{(\overline{p}_0, \overline{z}_0)}$. \square

4. GLOBALLY STABLE QUASISTATIC EVOLUTION FOR YOUNG MEASURES

4.1. Definitions and main result. We begin with the definition of the set of *admissible triples* in the Young measure formulation, with boundary datum \boldsymbol{w} on Γ_0 .

Definition 4.1. Given a set $\Theta \subset \mathbb{R}$ and a map $\boldsymbol{w}: \Theta \rightarrow H^1(\Omega; \mathbb{R}^2)$, we define $AY(\Theta, \boldsymbol{w})$ as the set of all triples $(\boldsymbol{u}, \boldsymbol{e}, \boldsymbol{\mu})$ with $\boldsymbol{u}: \Theta \rightarrow BD(\Omega)$, $\boldsymbol{e}: \Theta \rightarrow L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $\boldsymbol{\mu} \in SGY(\Theta, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, with the following property: for every finite sequence t_1, \dots, t_m in Θ , with $t_1 < \dots < t_m$, and every $i = 1, \dots, m$ there exist a sequence $(u_k^i, e_k^i, p_k^i) \in A(\boldsymbol{w}(t_i))$ and a sequence $z_k^i \in M_b(\overline{\Omega})$ such that

$$\begin{aligned} u_k^i &\rightharpoonup \boldsymbol{u}(t_i) \quad \text{weakly}^* \text{ in } BD(\Omega), \\ e_k^i &\rightarrow \boldsymbol{e}(t_i) \quad \text{strongly in } L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}), \end{aligned}$$

and

$$\delta_{((p_k^1, z_k^1), \dots, (p_k^m, z_k^m))} \rightharpoonup \boldsymbol{\mu}_{t_1 \dots t_m} \quad \text{weakly}^* \text{ in } GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^m). \quad (4.1)$$

Remark 4.2. Since the weak* convergence in $GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^m)$ implies the convergence of the norms $\|\cdot\|_*$ (see [4, Remark 3.12]), it is not restrictive to assume that

$$\|e_k^i\|_2 \leq \|\boldsymbol{e}(t_i)\|_2 + 1, \quad \|(p_k^i, z_k^i)\|_* \leq \|\boldsymbol{\mu}_{t_i}\|_* + 1$$

for every i and k . As $(u_k^i, e_k^i, p_k^i) \in A(\boldsymbol{w}(t_i))$, there exists a constant C_i , depending only on \boldsymbol{w} , t_i , $\|\boldsymbol{e}(t_i)\|_2$, $\|\boldsymbol{\mu}_{t_i}\|_*$, such that

$$\|u_k^i\|_1 + \|Eu_k^i\|_1 \leq C_i$$

for every k .

Remark 4.3. It follows from [4, Remark 6.4] that

$$\begin{aligned} p_k^i &\rightharpoonup \boldsymbol{p}(t_i) \quad \text{weakly}^* \text{ in } M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2}), \\ z_k^i &\rightharpoonup \boldsymbol{z}(t_i) \quad \text{weakly}^* \text{ in } M_b(\overline{\Omega}), \end{aligned}$$

where $(\boldsymbol{p}(t_i), \boldsymbol{z}(t_i)) := \text{bar}(\boldsymbol{\mu}_{t_i})$. As $(u_k^i, e_k^i, p_k^i) \in A(\boldsymbol{w}(t_i))$, by [3, Lemma 2.1] we conclude that

$$(\boldsymbol{u}(t_i), \boldsymbol{e}(t_i), \boldsymbol{p}(t_i)) \in A(\boldsymbol{w}(t_i)). \quad (4.2)$$

Remark 4.4. The inequalities proved in Remark 4.2 allow to use the metrizability of the weak* topology on bounded subsets of the dual of a separable Banach space and to prove that the set $AY(\Theta, \boldsymbol{w})$ satisfies the following closure property: if $\boldsymbol{u}: \Theta \rightarrow BD(\Omega)$, $\boldsymbol{e}: \Theta \rightarrow L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $\boldsymbol{\mu} \in SGY(\Theta, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, and $(\boldsymbol{u}_k, \boldsymbol{e}_k, \boldsymbol{\mu}_k)$ is a sequence in $AY(\Theta, \boldsymbol{w})$ such that

$$\boldsymbol{u}_k(t) \rightharpoonup \boldsymbol{u}(t) \quad \text{weakly}^* \text{ in } BD(\Omega), \quad (4.3)$$

$$\boldsymbol{e}_k(t) \rightarrow \boldsymbol{e}(t) \quad \text{strongly in } L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}) \quad (4.4)$$

for every $t \in \Theta$, and

$$(\boldsymbol{\mu}_k)_{t_1 \dots t_m} \rightharpoonup \boldsymbol{\mu}_{t_1 \dots t_m} \quad \text{weakly}^* \text{ in } GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^m) \quad (4.5)$$

for every finite sequence t_1, \dots, t_m in Θ , with $t_1 < \dots < t_m$, then $(\boldsymbol{u}, \boldsymbol{e}, \boldsymbol{\mu}) \in AY(\Theta, \boldsymbol{w})$.

More in general, if $\boldsymbol{u}: \Theta \rightarrow BD(\Omega)$, $\boldsymbol{e}: \Theta \rightarrow L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $\boldsymbol{\mu} \in SGY(\Theta, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, and $(\boldsymbol{u}_k, \boldsymbol{e}_k, \boldsymbol{\mu}_k)$ is a sequence in $AY(\Theta, \boldsymbol{w}_k)$ such that (4.3)–(4.5) hold and $\boldsymbol{w}_k(t) \rightarrow \boldsymbol{w}(t)$ strongly in $H^1(\Omega; \mathbb{R}^2)$ for every $t \in \Theta$, then $(\boldsymbol{u}, \boldsymbol{e}, \boldsymbol{\mu}) \in AY(\Theta, \boldsymbol{w})$. This follows from the closure property, observing that $(\boldsymbol{u}_k - \boldsymbol{w}_k + \boldsymbol{w}, \boldsymbol{e}_k - E\boldsymbol{w}_k + E\boldsymbol{w}, \boldsymbol{\mu}_k)$ belongs to $AY(\Theta, \boldsymbol{w})$.

Remark 4.5. Using Theorem 3.6, Lemma 3.7, and [17, Theorem 3] (see also [11, Appendix]), it is easy to see that the definition does not change if we replace $A(\mathbf{w}(t_i))$ by $A_{reg}(\mathbf{w}(t_i))$.

Given $\boldsymbol{\mu} \in SGY([0, +\infty), \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, its dissipation $\mathcal{D}_H(\boldsymbol{\mu}; a, b)$ on the time interval $[a, b] \subset [0, +\infty)$ is defined as

$$\sup \sum_{i=1}^k \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\mu}_{t_0 t_1 \dots t_k}(x, \xi_0, \theta_0, \dots, \xi_k, \theta_k, \eta) \rangle, \quad (4.6)$$

where the supremum is taken over all finite families t_0, t_1, \dots, t_k such that $a = t_0 < t_1 < \dots < t_k = b$. As in the case of the variation $\text{Var}(\boldsymbol{\mu}; a, b)$ considered in [4, Section 8], we have

$$\mathcal{D}_H(\boldsymbol{\mu}; a, b) = \sup \sum_{i=1}^k \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\mu}_{t_{i-1} t_i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle, \quad (4.7)$$

where the supremum is taken over all finite families t_0, t_1, \dots, t_k such that $a = t_0 < t_1 < \dots < t_k = b$.

In the following definition we use the homogeneous function $\{V\}$ defined by (3.30) and the notion of weakly* left-continuous system of generalized Young measures introduced in [4, Definition 7.6].

Definition 4.6. Given $\mathbf{w} \in AC_{loc}([0, +\infty); H^1(\Omega; \mathbb{R}^2))$, a *globally stable quasistatic evolution of Young measures* with boundary datum \mathbf{w} is a triple $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu}) \in AY([0, +\infty), \mathbf{w})$, with $\mathbf{u}, \mathbf{e}, \boldsymbol{\mu}$ weakly* left-continuous, such that the following conditions are satisfied:

(ev1) *global stability*: for every $t \in [0, +\infty)$ we have

$$\begin{aligned} & \mathcal{Q}(\mathbf{e}(t)) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle \leq \\ & \leq \mathcal{Q}(\mathbf{e}(t) + \tilde{\mathbf{e}}) + \mathcal{H}(\tilde{\mathbf{p}}, \tilde{\mathbf{z}}) + \langle \{V\}(\theta + \eta \tilde{\mathbf{z}}(x), \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle \end{aligned}$$

for every $(\tilde{\mathbf{u}}, \tilde{\mathbf{e}}, \tilde{\mathbf{p}}) \in A_{reg}(0)$ and every $\tilde{\mathbf{z}} \in L^1(\Omega)$;

(ev2) *energy balance*: for every $T \in (0, +\infty)$ we have $\text{Var}(\boldsymbol{\mu}; 0, T) < +\infty$ and

$$\begin{aligned} & \mathcal{Q}(\mathbf{e}(T)) + \mathcal{D}_H(\boldsymbol{\mu}; 0, T) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle = \\ & = \mathcal{Q}(\mathbf{e}(0)) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_0(x, \xi, \theta, \eta) \rangle + \int_0^T \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt, \end{aligned}$$

where $\boldsymbol{\sigma}(t) := \mathbb{C}\mathbf{e}(t)$.

We are now in a position to state the main theorem of the paper.

Theorem 4.7. Let $\mathbf{w} \in AC_{loc}([0, +\infty); H^1(\Omega; \mathbb{R}^2))$, $(u_0, e_0, p_0) \in A(\mathbf{w}(0))$, and $z_0 \in M_b(\bar{\Omega})$. Assume that

$$\mathcal{Q}(e_0) + \mathcal{V}(z_0) \leq \mathcal{Q}(e_0 + \tilde{\mathbf{e}}) + \mathcal{H}(\tilde{\mathbf{p}}, \tilde{\mathbf{z}}) + \mathcal{V}(z_0 + \tilde{\mathbf{z}}) \quad (4.8)$$

for every $(\tilde{\mathbf{u}}, \tilde{\mathbf{e}}, \tilde{\mathbf{p}}) \in A_{reg}(0)$ and every $\tilde{\mathbf{z}} \in L^1(\Omega)$. Then there exists a globally stable quasistatic evolution of Young measures $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ with boundary datum \mathbf{w} such that $\mathbf{u}(0) = u_0$, $\mathbf{e}(0) = e_0$, and $\boldsymbol{\mu}_0 = \delta_{(p_0, z_0)}$.

4.2. The incremental minimum problems. The proof of Theorem 4.7 will be obtained by time discretization, using an implicit Euler scheme. Let us fix a sequence of subdivisions $(t_k^i)_{i \geq 0}$ of the half-line $[0, +\infty)$, with

$$0 = t_k^0 < t_k^1 < \dots < t_k^{i-1} < t_k^i \rightarrow +\infty \quad \text{as } i \rightarrow \infty, \quad (4.9)$$

$$\tau_k := \sup_i (t_k^i - t_k^{i-1}) \rightarrow 0 \quad \text{as } k \rightarrow \infty. \quad (4.10)$$

For every k let $w_k^i := \mathbf{w}(t_k^i)$ for $i \geq 0$ and let $\tilde{w}_k^i := \mathbf{w}(t_k^i) - \mathbf{w}(t_k^{i-1})$ for $i \geq 1$.

We define $u_k^i \in BD(\Omega)$, $e_k^i \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, and $\boldsymbol{\mu}_k^i \in SGY(\{t_k^0, \dots, t_k^i\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ by induction on i . We set $u_k^0 := u_0$, $e_k^0 := e_0$, $\boldsymbol{\mu}_k^0 := \delta_{(p_0, z_0)}$, and for $i \geq 1$ we define $(u_k^i, e_k^i, \boldsymbol{\mu}_k^i)$ as a minimizer (see Lemma 4.9 below) of the functional

$$\mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \boldsymbol{\nu}_{t_k^{i-1} t_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle \quad (4.11)$$

over the set A_k^i of all triplets $(u, e, \boldsymbol{\nu})$ with $u \in BD(\Omega)$, $e \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, and $\boldsymbol{\nu} \in SGY(\{t_k^0, \dots, t_k^i\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, with the following property: there exist a sequence $(\tilde{u}_m, \tilde{e}_m, \tilde{p}_m) \in A_{reg}(\tilde{w}_k^i)$ and a sequence $\tilde{z}_m \in L^1(\Omega)$ such that

$$\begin{aligned} u_k^{i-1} + \tilde{u}_m &\rightharpoonup u \quad \text{weakly}^* \text{ in } BD(\Omega), \\ e_k^{i-1} + \tilde{e}_m &\rightarrow e \quad \text{strongly in } L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}), \end{aligned}$$

$$\mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}^i((\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}}) \rightharpoonup \boldsymbol{\nu}_{t_k^0 \dots t_k^i} \quad \text{weakly}^* \text{ in } GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1}),$$

where $\mathcal{T}_{(\tilde{p}, \tilde{z})}^i: \overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^i \times \mathbb{R} \rightarrow \overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1} \times \mathbb{R}$ is defined by

$$\mathcal{T}_{(\tilde{p}, \tilde{z})}^i(x, \xi_0, \theta_0, \dots, \xi_{i-1}, \theta_{i-1}, \eta) := (x, \xi_0, \theta_0, \dots, \xi_{i-1}, \theta_{i-1}, \xi_{i-1} + \eta \tilde{p}(x), \theta_{i-1} + \eta \tilde{z}(x), \eta).$$

We note that if $(u, e, \boldsymbol{\nu}) \in A_k^i$, then

$$\boldsymbol{\nu}_{t_k^0 \dots t_k^{i-1}} = (\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}}, \quad (4.12)$$

$$(u, e, p) \in A(w_k^i), \quad (4.13)$$

where $(p, z) := \text{bar}(\boldsymbol{\nu}_{t_k^i})$. Then we define $(p_k^i, z_k^i) := \text{bar}((\boldsymbol{\mu}_k^i)_{t_k^i})$.

Remark 4.8. The following equalities hold:

$$\begin{aligned} &\inf_{(u, e, \boldsymbol{\nu}) \in A_k^i} [\mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \boldsymbol{\nu}_{t_k^{i-1} t_k^i} \rangle] = \\ &= \inf_{\substack{(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(\tilde{w}_k^i) \\ \tilde{z} \in L^1(\Omega)}} [\mathcal{Q}(e_k^{i-1} + \tilde{e}) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \mathcal{T}_{(\tilde{p}, \tilde{z})}^i((\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}}) \rangle] = \\ &= \inf_{\substack{(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(\tilde{w}_k^i) \\ \tilde{z} \in L^1(\Omega)}} [\mathcal{Q}(e_k^{i-1} + \tilde{e}) + \mathcal{H}(\tilde{p}, \tilde{z}) + \langle \{V\}(\theta_i, \eta), \mathcal{T}_{(\tilde{p}, \tilde{z})}^i((\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}}) \rangle] = \\ &= \inf_{(u, e, \boldsymbol{\nu}) \in B_k^i} [\mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \boldsymbol{\nu}_{t_k^{i-1} t_k^i} \rangle], \end{aligned}$$

where B_k^i is the class of all triplets $(u, e, \boldsymbol{\nu})$, with $u \in BD(\Omega)$, $e \in L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $\boldsymbol{\nu} \in SGY(\{t_k^0, \dots, t_k^i\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, such that $\boldsymbol{\nu}_{t_k^0 \dots t_k^{i-1}} = (\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}}$ and $(u, e, p) \in A(w_k^i)$, where $(p, z) := \text{bar}(\boldsymbol{\nu}_{t_k^i})$. The first two equalities follow from the definition of A_k^i and the continuity properties of the functional (4.11). On the other hand the infimum in the last line is greater than or equal to the infimum in the previous line by Theorem 3.9, and is less than or equal to the infimum in the first line, since $A_k^i \subset B_k^i$ by (4.12) and (4.13).

The existence of a minimizer $(u_k^i, e_k^i, \boldsymbol{\mu}_k^i)$ to (4.11) is guaranteed by the following lemma.

Lemma 4.9. *For every i the functional (4.11) has a minimizer on A_k^i , every minimizer $(u_k^i, e_k^i, \boldsymbol{\mu}_k^i)$ satisfies $\overline{(\boldsymbol{\mu}_k^i)_{t_k^i}}^Y = \delta_{(p_0^i, z_0^i)}$, and*

$$\mathcal{Q}(e_k^i) + \mathcal{H}_{\text{eff}}(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) \leq \mathcal{Q}(e) + \mathcal{H}_{\text{eff}}(p - p_k^{i-1}, z - z_k^{i-1}) \quad (4.14)$$

for every $(u, e, p) \in A(w(t_k^i))$ and every $z \in M_b(\overline{\Omega})$.

Proof. The lemma will be proved by induction on i . Assume that $\boldsymbol{\mu}_k^{i-1}$ is defined and $\overline{(\boldsymbol{\mu}_k^{i-1})_{t_k^{i-1}}}^Y = \delta_{(p_0^{i-1}, z_0^{i-1})}$. We shall prove that the functional (4.11) has a minimizer $(u_k^i, e_k^i, \boldsymbol{\mu}_k^i)$ in A_k^i and

$$\overline{(\boldsymbol{\mu}_k^i)_{t_k^i}}^Y = \delta_{(p_0^i, z_0^i)}. \quad (4.15)$$

Thanks to Remark 4.8 there exists a minimizing sequence $(u_m, e_m, \boldsymbol{\nu}^m)$ in A_k^i with $\boldsymbol{\nu}_{t_k^0 \dots t_k^i}^m = \mathcal{T}_{(\tilde{p}_m, \tilde{z}_m)}^i((\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}})$ and $(u_m, e_m, \tilde{p}_m) \in A_{reg}(\tilde{w}_k^i)$. By (2.30) we have

$$H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta) \geq C_V^K |\xi_i - \xi_{i-1}| + C_V^K |\theta_i - \theta_{i-1}| + \{V\}(\theta_{i-1}, \eta),$$

hence by the compatibility condition (7.2) of [4] the sequence

$$\begin{aligned} \mathcal{Q}(e^m) + C_V^K \langle |\xi_i - \xi_{i-1}| + |\theta_i - \theta_{i-1}|, \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ + \langle \{V\}(\theta_{i-1}, \eta), \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \eta) \rangle \end{aligned}$$

is bounded uniformly with respect to m . By (4.12) we have $\boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m = (\boldsymbol{\mu}_k^{i-1})_{t_k^{i-1}}$, so that

$$\mathcal{Q}(e^m) + C_V^K \langle |\xi_i - \xi_{i-1}| + |\theta_i - \theta_{i-1}|, \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle \quad (4.16)$$

is bounded uniformly with respect to m . Since by [4, Remark 2.9 and (7.2)]

$$\begin{aligned} \|\boldsymbol{\nu}_{t_k^i}^m\|_* &\leq \langle |\xi_i| + |\theta_i|, \boldsymbol{\nu}_{t_k^i}^m(x, \xi_i, \theta_i, \eta) \rangle = \\ &= \langle |\xi_i| + |\theta_i|, \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle \leq \\ &\leq \langle |\xi_i - \xi_{i-1}| + |\theta_i - \theta_{i-1}|, \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ &\quad + \langle |\xi_{i-1}| + |\theta_{i-1}|, \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle = \\ &= \langle |\xi_i - \xi_{i-1}| + |\theta_i - \theta_{i-1}|, \boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ &\quad + \langle |\xi_{i-1}| + |\theta_{i-1}|, (\boldsymbol{\mu}_k^{i-1})_{t_k^{i-1}}(x, \xi_{i-1}, \theta_{i-1}, \eta) \rangle, \end{aligned}$$

it follows from (2.22) and (4.16) and that e^m is bounded in $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ and $\boldsymbol{\nu}_{t_k^i}^m$ is bounded in $GY(\bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$. Using (4.12) and [4, Lemma 7.8] we obtain also that $\boldsymbol{\nu}_{t_k^0 \dots t_k^i}^m$ is bounded in $GY(\bar{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1})$.

Passing to a subsequence, we may assume that $e_m \rightharpoonup e$ weakly in $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ and $\boldsymbol{\nu}_{t_k^0 \dots t_k^i}^m \rightharpoonup \nu_{0 \dots i}$ weakly* in $GY(\bar{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1})$. Let $\boldsymbol{\nu} \in SGY(\{t_k^0, \dots, t_k^i\}, \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ be the system associated with $\nu_{0 \dots i}$ according to [4, Remark 7.9] and let $(p, z) := \text{bar}(\boldsymbol{\nu}_{t_k^i}^m)$. Note that $(p_k^{i-1} + \tilde{p}_m, z_k^{i-1} + \tilde{z}_m) = \text{bar}(\boldsymbol{\nu}_{t_k^i}^m) \rightharpoonup (p, z)$ weakly* in $M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2}) \times M_b(\bar{\Omega})$ by [4, Remark 6.4]. Since $(u_m, e_m, \boldsymbol{\nu}^m) \in A_k^i$ we have $\|Eu^m\|_1 \leq \|e^m\|_1 + \|\text{bar}(\boldsymbol{\nu}_{t_k^i}^m)\|_1$ and $\|\boldsymbol{w}(t_k^i) - u^m\|_{1, \Gamma_0} \leq \|\text{bar}(\boldsymbol{\nu}_{t_k^i}^m)\|_1$. By [19, Proposition 2.4 and Remark 2.5] it follows that u^m is bounded in $BD(\Omega)$. Therefore, passing to a subsequence, we may assume that $u^m \rightharpoonup u$ weakly* in $BD(\Omega)$. By [3, Lemma 2.1] it follows that $(u, e, p) \in A(w_k^i)$, hence $(u, e, \boldsymbol{\nu}) \in B_k^i$.

We claim that

$$e^m \rightarrow e \quad \text{strongly in } L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}). \quad (4.17)$$

Indeed, if not, then we can find a subsequence (not relabelled) such that

$$\mathcal{Q}(e) < \liminf_m \mathcal{Q}(e_m). \quad (4.18)$$

Since the other term in (4.11) is continuous with respect to the weak* convergence of $\boldsymbol{\nu}_{t_k^{i-1} t_k^i}^m$ to $\boldsymbol{\nu}_{t_k^{i-1} t_k^i}$, (4.18) would imply that

$$\begin{aligned} \mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \boldsymbol{\nu}_{t_k^{i-1} t_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle < \\ < \inf_{(\hat{u}, \hat{e}, \hat{\boldsymbol{\nu}}) \in A_k^i} [\mathcal{Q}(\hat{e}) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \hat{\boldsymbol{\nu}}_{t_k^{i-1} t_k^i} \rangle], \end{aligned}$$

which contradicts the equalities in Remark 4.8, since $(u, e, \boldsymbol{\nu}) \in B_k^i$. Therefore, (4.17) is proved.

We deduce from (4.17) that $(u, e, \nu) \in A_k^i$ and that it is a minimizer of (4.11) in A_k^i . From now on we set $(u_k^i, e_k^i, \mu_k^i) := (u, e, \nu)$. By Remark 4.8 and Theorem 3.9 we obtain

$$\begin{aligned} & \min_{(\tilde{u}, \tilde{e}, \tilde{p}) \in A(\tilde{w}_k^i), \tilde{z} \in M_b(\overline{\Omega})} [\mathcal{Q}(e_k^{i-1} + \tilde{e}) + \mathcal{H}_{\text{eff}}(\tilde{p}, \tilde{z}) + \langle \{V\}(\theta_{i-1}, \eta), (\mu_k^{i-1})_{t_k^{i-1}} \rangle] = \\ & = \mathcal{Q}(e_k^i) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), (\mu_k^i)_{t_k^{i-1}t_k^i} \rangle \geq \\ & \geq \mathcal{Q}(e_k^i) + \mathcal{H}_{\text{eff}}(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) + \langle \{V\}(\theta_{i-1}, \eta), (\mu_k^{i-1})_{t_k^{i-1}} \rangle, \end{aligned} \quad (4.19)$$

where the last inequality follows from Jensen inequality. Since $(u_k^i - u_k^{i-1}, e_k^i - e_k^{i-1}, p_k^i - p_k^{i-1}) \in A(\tilde{w}_k^i)$, we deduce that the previous inequalities are in fact equalities. Theorem 3.13 now yields (4.15). Finally, (4.14) easily follows from (4.19). \square

Corollary 4.10. *For every i and k we have*

$$\mathcal{Q}(e_k^i) + \mathcal{H}_{\text{eff}}(p_k^i - p_k^{i-1}, 0) \leq \mathcal{Q}(e) + \mathcal{H}_{\text{eff}}(p - p_k^{i-1}, 0)$$

for every $(u, e, p) \in A(w(t_k^i))$.

Proof. It is enough to take $z = z_k^{i-1}$ in (4.14) and to use the inequality $H_{\text{eff}}(\xi, \theta) \geq H_{\text{eff}}(\xi, 0)$, which follows from the fact that $\theta \mapsto H_{\text{eff}}(\xi, \theta)$ is convex and even. \square

The following theorem shows that the incremental problems can be considered as a relaxed version of incremental problems defined on functions. For different approaches to the relaxation problem in the context of rate-independent processes we refer to [15] and [16].

Proposition 4.11. *Let us fix k . Let (u^m, e^m, p^m) be a sequence in $A_{\text{reg}}(w(0))$ and let z^m be a sequence in $L^1(\Omega)$. For every $i \geq 1$ let us consider two sequences (with respect to the index m) $(\tilde{u}^{i,m}, \tilde{e}^{i,m}, \tilde{p}^{i,m}) \in A_{\text{reg}}(\tilde{w}_k^i)$ and $\tilde{z}^{i,m} \in L^1(\Omega)$. For every multiindex $m_0 \dots m_i$ with $i+1$ components we define*

$$\begin{aligned} u^{m_0 \dots m_i} &:= u^{m_0} + \sum_{j=1}^i \tilde{u}^{j, m_j}, & e^{m_0 \dots m_i} &:= e^{m_0} + \sum_{j=1}^i \tilde{e}^{j, m_j}, \\ p^{m_0 \dots m_i} &:= p^{m_0} + \sum_{j=1}^i \tilde{p}^{j, m_j}, & z^{m_0 \dots m_i} &:= z^{m_0} + \sum_{j=1}^i \tilde{z}^{j, m_j}. \end{aligned}$$

Note that

$$(u^{m_0 \dots m_i}, e^{m_0 \dots m_i}, p^{m_0 \dots m_i}) \in A_{\text{reg}}(w_k^i).$$

Moreover, we define $\mu^{m_0 \dots m_i} \in GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1})$ by

$$\mu^{m_0 \dots m_i} := \delta_{((p^{m_0}, z^{m_0}), (p^{m_0 m_1}, z^{m_0 m_1}), \dots, (p^{m_0 \dots m_i}, z^{m_0 \dots m_i}))}.$$

Suppose that there exist $\hat{e}^i \in L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$ and $\hat{\mu}^i \in SGY(\{t_k^0, \dots, t_k^i\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ such that for every $i \geq 0$

$$\lim_{m_i \rightarrow \infty} \dots \lim_{m_0 \rightarrow \infty} e^{m_0 \dots m_i} = \hat{e}^i, \quad (4.20)$$

$$\lim_{m_i \rightarrow \infty} \dots \lim_{m_0 \rightarrow \infty} \mu^{m_0 \dots m_i} = \hat{\mu}_{t_k^0 \dots t_k^i}^i, \quad (4.21)$$

where in the former formula all limits are with respect to weak convergence in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, while in the latter they are taken in the weak* convergence in $GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1})$. Then for every $i \geq 1$

$$\begin{aligned} & \liminf_{m_i \rightarrow \infty} \dots \liminf_{m_0 \rightarrow \infty} [\mathcal{Q}(e^{m_0 \dots m_i}) + \mathcal{H}(\tilde{p}^{i, m_i}, \tilde{z}^{i, m_i}) + \mathcal{V}(z^{m_0 \dots m_i})] \geq \\ & \geq \mathcal{Q}(\hat{e}^i) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), \hat{\mu}_{t_k^{i-1}t_k^i}^i \rangle, \end{aligned} \quad (4.22)$$

where $\hat{\mu}_{t_k^{i-1}t_k^i}^i$ acts on the variable $(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta)$.

Conversely, if \hat{e}^i and $\hat{\boldsymbol{\mu}}^i$ coincide with the function e_k^i and the measure $\boldsymbol{\mu}_k^i$ obtained in the incremental construction, then there exist two sequences $(u^m, e^m, p^m) \in A_{reg}(\mathbf{w}(0))$, $z^m \in L^1(\Omega)$ and for every $i \geq 1$ two sequences $(\tilde{u}^{i,m}, \tilde{e}^{i,m}, \tilde{p}^{i,m}) \in A_{reg}(\tilde{w}_k^i)$ and $\tilde{z}^{i,m} \in L^1(\Omega)$ such that for every $i \geq 0$ (4.20) holds with respect to strong convergence and (4.21) holds with respect to weak* convergence, while

$$\begin{aligned} & \lim_{m_i \rightarrow \infty} \dots \lim_{m_0 \rightarrow \infty} [\mathcal{Q}(e^{m_0 \dots m_i}) + \mathcal{H}(\tilde{p}^{i,m_i}, \tilde{z}^{i,m_i}) + \mathcal{V}(z^{m_0 \dots m_i})] = \\ & = \mathcal{Q}(e_k^i) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta), (\boldsymbol{\mu}_k^i)_{t_k^{i-1} t_k^i} \rangle \end{aligned} \quad (4.23)$$

for every $i \geq 1$.

Proof. Inequality (4.22) follows from (4.20) and (4.21) by the lower semicontinuity of \mathcal{Q} in the weak topology of $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ and the continuity of the duality product in the weak* topology of $GY(\bar{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^2)$.

By Theorem 3.6 and Lemma 3.7 there exist a sequence $(u^m, e^m, p^m) \in A_{reg}(\mathbf{w}(0))$ and a sequence $z^m \in L^1(\Omega)$ such that $u^m \rightharpoonup u_0$ weakly* in $BD(\Omega)$, $e^m \rightarrow e_0$ strongly in $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$, $p^m \rightharpoonup p_0$ weakly* in $M_b(\bar{\Omega}; \mathbb{M}_D^{2 \times 2})$, $z^m \rightharpoonup z_0$ weakly* in $M_b(\bar{\Omega})$, and $\|(p^m, z^m)\|_1 \rightarrow \|(p_0, z_0)\|_1$. Using [17, Theorem 3] (see also [11, Appendix]) we obtain that $\delta_{(p^m, z^m)} \rightharpoonup \delta_{(p_0, z_0)}$ weakly* in $GY(\bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$.

For every $i \geq 1$, by definition of A_k^i there exist a sequence $(\tilde{u}^{i,m}, \tilde{e}^{i,m}, \tilde{p}^{i,m}) \in A_{reg}(\tilde{w}_k^i)$ and a sequence $\tilde{z}^{i,m} \in L^1(\Omega)$ such that

$$\begin{aligned} & u_k^{i-1} + \tilde{u}^{i,m} \rightharpoonup u_k^i \quad \text{weakly* in } BD(\Omega), \\ & e_k^{i-1} + \tilde{e}^{i,m} \rightarrow e_k^i \quad \text{strongly in } L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}), \end{aligned} \quad (4.24)$$

$$\mathcal{T}_{(\tilde{p}^{i,m}, \tilde{z}^{i,m})}^i((\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}}) \rightharpoonup (\boldsymbol{\mu}_k^i)_{t_k^0 \dots t_k^i} \quad \text{weakly* in } GY(\bar{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1}). \quad (4.25)$$

Condition (4.20) is trivially satisfied thanks to (4.24). To prove (4.21) we observe that for every $i \geq 1$

$$\boldsymbol{\mu}^{m_0 \dots m_i} = \mathcal{T}_{(\tilde{p}^{i,m_i}, \tilde{z}^{i,m_i})}^i(\boldsymbol{\mu}^{m_0 \dots m_{i-1}}).$$

We now proceed by induction on i . Equality (4.21) for $i = 0$ is true by construction. Assume that (4.21) holds for $i - 1$. Then by Lemma 2.1

$$\lim_{m_{i-1} \rightarrow \infty} \dots \lim_{m_0 \rightarrow \infty} \boldsymbol{\mu}^{m_0 \dots m_i} = \mathcal{T}_{(\tilde{p}^{i,m_i}, \tilde{z}^{i,m_i})}^i((\boldsymbol{\mu}_k^{i-1})_{t_k^0 \dots t_k^{i-1}})$$

The conclusion for i follows from (4.25). \square

4.3. Further minimality properties. We now prove that the solutions of the incremental problems satisfy some additional minimality conditions.

Lemma 4.12. *For every i and k and every $t > t_k^i$ we have*

$$\begin{aligned} & \mathcal{Q}(e_k^i) + \langle \{V\}(\theta_i, \eta), (\boldsymbol{\mu}_k^i)_{t_k^i}(x, \xi_i, \theta_i, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e) + \langle H(\xi - \xi_i, \theta - \theta_i), \boldsymbol{\nu}_{t_k^i t}(x, \xi_i, \theta_i, \xi, \theta, \eta) \rangle + \langle \{V\}(\theta, \eta), \boldsymbol{\nu}_t(x, \xi, \theta, \eta) \rangle \end{aligned} \quad (4.26)$$

for every $(u, e, \boldsymbol{\nu}) \in BD(\Omega) \times L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}) \times SGY(\{t_k^0, \dots, t_k^i, t\}, \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ such that

$$\boldsymbol{\nu}_{t_k^0 \dots t_k^i} = (\boldsymbol{\mu}_k^i)_{t_k^0 \dots t_k^i}, \quad (4.27)$$

$$(u, e, p) \in A(\mathbf{w}(t_k^i)), \quad (4.28)$$

where $(p, z) := \text{bar}(\boldsymbol{\nu}_t)$.

Proof. Let us fix $(u, e, \boldsymbol{\nu})$ as in the statement of the lemma, and let $\tilde{\boldsymbol{\nu}}$ be the system in $SGY(\{t_k^0 \dots t_k^i\}, \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ associated with $\pi_{t_k^0 \dots t_k^i t}^{t_k^0 \dots t_k^i t}(\boldsymbol{\nu}_{t_k^0 \dots t_k^i t}) \in GY(\bar{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{i+1})$ according to [4, Remark 7.9]. Since $\boldsymbol{\mu}_k^i$ satisfies (4.12), by (4.27) and (4.28) the triplet

$(u, e, \tilde{\nu})$ satisfies (4.12) and (4.13), hence $(u, e, \tilde{\nu})$ belongs to the set B_k^i defined in Remark 4.8. By minimality we have

$$\begin{aligned} & \mathcal{Q}(e_k^i) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), (\boldsymbol{\mu}_k^i)_{t_k^{i-1}t_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta_i, \eta), (\boldsymbol{\mu}_k^i)_{t_k^i}(x, \xi_i, \theta_i, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \tilde{\boldsymbol{\nu}}_{t_k^{i-1}t_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta_i, \eta), \tilde{\boldsymbol{\nu}}_{t_k^i}(x, \xi_i, \theta_i, \eta) \rangle. \end{aligned}$$

Since $(\boldsymbol{\mu}_k^i)_{t_k^{i-1}t_k^i} = \boldsymbol{\nu}_{t_k^{i-1}t_k^i}$, $\tilde{\boldsymbol{\nu}}_{t_k^{i-1}t_k^i} = \boldsymbol{\nu}_{t_k^{i-1}t}$, and $\tilde{\boldsymbol{\nu}}_{t_k^i} = \boldsymbol{\nu}_t$, we get

$$\begin{aligned} & \mathcal{Q}(e_k^i) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\nu}_{t_k^{i-1}t_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta_i, \eta), (\boldsymbol{\mu}_k^i)_{t_k^i}(x, \xi_i, \theta_i, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\nu}_{t_k^{i-1}t}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta, \eta), \boldsymbol{\nu}_t(x, \xi, \theta, \eta) \rangle. \end{aligned}$$

From the compatibility condition (7.2) of [4] we obtain

$$\begin{aligned} & \mathcal{Q}(e_k^i) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\nu}_{t_k^{i-1}t_k^i t}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \xi, \theta, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta_i, \eta), (\boldsymbol{\mu}_k^i)_{t_k^i}(x, \xi_i, \theta_i, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\nu}_{t_k^{i-1}t_k^i t}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \xi, \theta, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta, \eta), \boldsymbol{\nu}_t(x, \xi, \theta, \eta) \rangle. \end{aligned}$$

By the triangle inequality (2.11) we deduce that

$$\begin{aligned} & \mathcal{Q}(e_k^i) + \langle \{V\}(\theta_i, \eta), (\boldsymbol{\mu}_k^i)_{t_k^i}(x, \xi_i, \theta_i, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e) + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\nu}_{t_k^{i-1}t_k^i t}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \xi, \theta, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta, \eta), \boldsymbol{\nu}_t(x, \xi, \theta, \eta) \rangle, \end{aligned}$$

which gives (4.26) by the compatibility condition (7.2) of [4]. \square

For every i and k we set $\sigma_k^i := \mathbb{C}e_k^i$ and for every $t \in [0, +\infty)$ we consider the piecewise constant interpolations defined by

$$\begin{aligned} \mathbf{u}_k(t) &:= \mathbf{u}_k^i, & \mathbf{e}_k(t) &:= \mathbf{e}_k^i, & \mathbf{p}_k(t) &:= \mathbf{p}_k^i, & \mathbf{z}_k(t) &:= \mathbf{z}_k^i, \\ \boldsymbol{\sigma}_k(t) &:= \boldsymbol{\sigma}_k^i, & \mathbf{w}_k(t) &:= \mathbf{w}_k^i, & [t]_k &:= t_k^i, \end{aligned} \quad (4.29)$$

for $t \in [t_k^i, t_k^{i+1})$. We define also $\boldsymbol{\mu}_k$ as the unique system in $SGY([0, +\infty), \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ whose restrictions to the time intervals $[0, t_k^i]$ coincide with the piecewise constant interpolations of $\boldsymbol{\mu}_k^i \in SGY(\{t_k^0, \dots, t_k^i\}, \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ introduced in [4, Definition 7.10]. As $(\mathbf{p}_k(t), \mathbf{z}_k(t)) = \text{bar}((\boldsymbol{\mu}_k)_t)$, we have also

$$(\mathbf{u}_k(t), \mathbf{e}_k(t), \mathbf{p}_k(t)) \in A(\mathbf{w}_k(t)) \quad (4.30)$$

for every $t \in [0, +\infty)$.

Lemma 4.13. *Let $t, \hat{t} \in [0, +\infty)$ with $t < \hat{t}$. Then*

$$\begin{aligned} & \mathcal{Q}(e_k(t)) + \langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e_k(\hat{t}) - E\mathbf{w}_k(\hat{t}) + E\mathbf{w}_k(t)) + \langle H(\hat{\xi} - \xi, \hat{\theta} - \theta), (\boldsymbol{\mu}_k)_{t\hat{t}}(x, \xi, \theta, \hat{\xi}, \hat{\theta}, \eta) \rangle + \\ & \quad + \langle \{V\}(\hat{\theta}, \eta), (\boldsymbol{\mu}_k)_{\hat{t}}(x, \hat{\xi}, \hat{\theta}, \eta) \rangle \end{aligned} \quad (4.31)$$

for every k .

Proof. Let $u := \mathbf{u}_k(\hat{t}) - \mathbf{w}_k(\hat{t}) + \mathbf{w}_k(t)$ and $e := \mathbf{e}_k(\hat{t}) - E\mathbf{w}_k(\hat{t}) + E\mathbf{w}_k(t)$, and let i be the greatest index such that $t_k^i \leq t$. Since the triplet $(u, e, (\boldsymbol{\mu}_k)_{t_k^0 \dots t_k^i \hat{t}})$ satisfies (4.27) and (4.28), the result follows from Lemma 4.12. \square

Lemma 4.14. *Let $t \in [0, +\infty)$. Then*

$$\begin{aligned} & \mathcal{Q}(\mathbf{e}_k(t)) + \langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle \leq \\ & \leq \mathcal{Q}(\mathbf{e}_k(t) + \tilde{\mathbf{e}}) + \mathcal{H}(\tilde{\mathbf{p}}, \tilde{\mathbf{z}}) + \langle \{V\}(\theta + \eta \tilde{\mathbf{z}}(x), \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle \end{aligned} \quad (4.32)$$

for every $(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(0)$ and every $\tilde{z} \in L^1(\Omega)$.

Proof. Let us fix $(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(0)$ and $\tilde{z} \in L^1(\Omega)$. Let i be the greatest index such that $t_k^i \leq t$ and let $\hat{t} > t_k^i$. We set $\hat{u} := u_k^i + \tilde{u} = \mathbf{u}_k(t) + \tilde{u}$, $\hat{e} := e_k^i + \tilde{e} = \mathbf{e}_k(t) + \tilde{e}$, $\nu_{t_k^0 \dots t_k^{\hat{t}}} := \mathcal{T}_{(\tilde{p}, \tilde{z})}^{i+1}((\boldsymbol{\mu}_k)_{t_k^0 \dots t_k^i}) = \mathcal{T}_{(\tilde{p}, \tilde{z})}^{i+1}((\boldsymbol{\mu}_k)_{t_k^0 \dots t_k^i})$, and we define $\boldsymbol{\nu} \in SGY(\{t_k^0, \dots, t_k^i, \hat{t}\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ as the system associated with $\nu_{t_k^0 \dots t_k^{\hat{t}}}$ according to [4, Remark 7.9]. Since the triplet $(\hat{u}, \hat{e}, \boldsymbol{\nu})$ satisfies (4.27) and (4.28), the conclusion follows from Lemma 4.12. \square

4.4. Energy estimates. We now prove some energy estimates for the solutions of the incremental minimum problems.

Lemma 4.15. *For every $T > 0$ there exists a sequence $\omega_k^T \rightarrow 0^+$ such that*

$$\begin{aligned} & \mathcal{Q}(\mathbf{e}_k(t)) + \mathcal{D}_H(\boldsymbol{\mu}_k; 0, t) + \langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e_0) + \mathcal{V}(z_0) + \int_0^{[t]_k} \langle \boldsymbol{\sigma}_k(t), E\dot{\mathbf{w}}(t) \rangle dt + \omega_k^T \end{aligned} \quad (4.33)$$

for every k and every $t \in [0, T]$.

Proof. Let us fix $T > 0$ and $t \in [0, T]$. Arguing as in [4, Remark 8.5], we can prove that

$$\mathcal{D}_H(\boldsymbol{\mu}_k; 0, t) = \sum_{r=1}^i \langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}), (\boldsymbol{\mu}_k)_{t_k^{r-1} t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle,$$

where i is the largest integer such that $t_k^i \leq t$. Therefore, using the definition of piecewise constant interpolation of a generalized Young measures, we have to show that there exists a sequence $\omega_k^T \rightarrow 0^+$ such that

$$\begin{aligned} & \mathcal{Q}(e_k^i) + \sum_{r=1}^i \langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}), (\boldsymbol{\mu}_k^r)_{t_k^{r-1} t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k^i)_{t_k^i}(x, \xi, \theta, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e_0) + \mathcal{V}(z_0) + \int_0^{t_k^i} \langle \boldsymbol{\sigma}_k(t), E\dot{\mathbf{w}}(t) \rangle dt + \omega_k^T \end{aligned} \quad (4.34)$$

for every k and every i with $t_k^i \leq T$.

Fix an integer r with $1 \leq r \leq i$ and let $\pi^{(r)}: \overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^r \times \mathbb{R} \rightarrow \overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^{r+1} \times \mathbb{R}$ be the map defined by

$$\pi^{(r)}(x, (\xi_0, \theta_0), \dots, (\xi_{r-1}, \theta_{r-1}), \eta) = (x, (\xi_0, \theta_0), \dots, (\xi_{r-1}, \theta_{r-1}), (\xi_{r-1}, \theta_{r-1}), \eta).$$

Let $\hat{u} := u_k^{r-1} - w_k^{r-1} + w_k^r$, $\hat{e} := e_k^{r-1} - Ew_k^{r-1} + Ew_k^r$, and $\hat{\boldsymbol{\mu}}_{t_k^0 \dots t_k^r} := \pi^{(r)}((\boldsymbol{\mu}_k^{r-1})_{t_k^0 \dots t_k^{r-1}})$. Let $\hat{\boldsymbol{\mu}} \in SGY(\{t_k^0, \dots, t_k^r\}, \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ be the system associated with $\hat{\boldsymbol{\mu}}_{t_k^0 \dots t_k^r}$ by [4, Remark 7.9]. It is easy to check that

$$\hat{\boldsymbol{\mu}}_{t_k^0 \dots t_k^{r-1}} = (\boldsymbol{\mu}_k^{r-1})_{t_k^0 \dots t_k^{r-1}}.$$

Let $(\hat{p}, \hat{z}) := \text{bar}(\hat{\boldsymbol{\mu}}_{t_k^r})$. Since $\hat{p} = p_k^{r-1}$, we find that $(\hat{u}, \hat{e}, \hat{\boldsymbol{\mu}}) \in A(w_k^r)$, hence $(\hat{u}, \hat{e}, \hat{\boldsymbol{\mu}})$ belongs to the class B_k^r , introduced in Remark 4.8. By minimality we have

$$\begin{aligned} & \mathcal{Q}(e_k^r) + \langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}) + \{V\}(\theta_r, \eta), (\boldsymbol{\mu}_k^r)_{t_k^{r-1} t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle \leq \\ & \leq \mathcal{Q}(\hat{e}) + \langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}) + \{V\}(\theta_r, \eta), \hat{\boldsymbol{\mu}}_{t_k^{r-1} t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle. \end{aligned} \quad (4.35)$$

As $\hat{\boldsymbol{\mu}}_{t_k^{r-1}t_k^r} = (\pi_{t_k^{r-1}t_k^r}^{t_k^0 \dots t_k^r} \circ \pi^{(r)})((\boldsymbol{\mu}_k^{r-1})_{t_k^0 \dots t_k^{r-1}})$ and

$$(\pi_{t_k^{r-1}t_k^r}^{t_k^0 \dots t_k^r} \circ \pi^{(r)})(x, (\xi_0, \theta_0), \dots, (\xi_{r-1}, \theta_{r-1}), \eta) = (x, (\xi_{r-1}, \theta_{r-1}), (\xi_{r-1}, \theta_{r-1}), \eta),$$

we have

$$\langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}), \hat{\boldsymbol{\mu}}_{t_k^{r-1}t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle = 0$$

and

$$\langle \{V\}(\theta_r, \eta), \hat{\boldsymbol{\mu}}_{t_k^{r-1}t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle = \langle \{V\}(\theta_{r-1}, \eta), (\boldsymbol{\mu}_k^{r-1})_{t_k^{r-1}}(x, \xi_{r-1}, \theta_{r-1}, \eta) \rangle.$$

Therefore (4.35) gives, thanks to the compatibility condition (7.2) of [4],

$$\begin{aligned} & \mathcal{Q}(e_k^r) + \langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}), (\boldsymbol{\mu}_k^r)_{t_k^{r-1}t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta_r, \eta), (\boldsymbol{\mu}_k^r)_{t_k^r}(x, \xi_r, \theta_r, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e_k^{r-1} + Ew_k^r - Ew_k^{r-1}) + \langle \{V\}(\theta_{r-1}, \eta), (\boldsymbol{\mu}_k^{r-1})_{t_k^{r-1}}(x, \xi_{r-1}, \theta_{r-1}, \eta) \rangle, \end{aligned} \quad (4.36)$$

where the quadratic form in the right-hand side can be developed as

$$\mathcal{Q}(e_k^{r-1} + Ew_k^r - Ew_k^{r-1}) = \mathcal{Q}(e_k^{r-1}) + \langle \sigma_k^{r-1}, Ew_k^r - Ew_k^{r-1} \rangle + \mathcal{Q}(Ew_k^r - Ew_k^{r-1}). \quad (4.37)$$

From the absolute continuity of w with respect to t we obtain

$$w_k^r - w_k^{r-1} = \int_{t_k^{r-1}}^{t_k^r} \dot{w}(t) dt,$$

where we use a Bochner integral of a function with values in $H^1(\Omega; \mathbb{R}^2)$. This implies that

$$Ew_k^r - Ew_k^{r-1} = \int_{t_k^{r-1}}^{t_k^r} E\dot{w}(t) dt, \quad (4.38)$$

where we use a Bochner integral of a function with values in $L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$. By (2.22) and (4.38) we get

$$\mathcal{Q}(Ew_k^r - Ew_k^{r-1}) \leq \beta_{\mathbb{C}} \left(\int_{t_k^{r-1}}^{t_k^r} \|E\dot{w}(t)\|_2 dt \right)^2. \quad (4.39)$$

By (4.36)–(4.39) we obtain

$$\begin{aligned} & \mathcal{Q}(e_k^r) + \langle H(\xi_r - \xi_{r-1}, \theta_r - \theta_{r-1}), (\boldsymbol{\mu}_k^r)_{t_k^{r-1}t_k^r}(x, \xi_{r-1}, \theta_{r-1}, \xi_r, \theta_r, \eta) \rangle + \\ & \quad + \langle \{V\}(\theta_r, \eta), (\boldsymbol{\mu}_k^r)_{t_k^r}(x, \xi_r, \theta_r, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e_k^{r-1}) + \langle \{V\}(\theta_{r-1}, \eta), (\boldsymbol{\mu}_k^{r-1})_{t_k^{r-1}}(x, \xi_{r-1}, \theta_{r-1}, \eta) \rangle + \\ & \quad + \int_{t_k^{r-1}}^{t_k^r} \langle \sigma_k^{r-1}, E\dot{w}(t) \rangle dt + \beta_{\mathbb{C}} \left(\int_{t_k^{r-1}}^{t_k^r} \|E\dot{w}(t)\|_2 dt \right)^2 \leq \\ & \leq \mathcal{Q}(e_k^{r-1}) + \langle \{V\}(\theta_{r-1}, \eta), (\boldsymbol{\mu}_k^{r-1})_{t_k^{r-1}}(x, \xi_{r-1}, \theta_{r-1}, \eta) \rangle + \\ & \quad + \int_{t_k^{r-1}}^{t_k^r} \langle \sigma_k^{r-1}, E\dot{w}(t) \rangle dt + \rho_k^T \int_{t_k^{r-1}}^{t_k^r} \|E\dot{w}(t)\|_2 dt, \end{aligned} \quad (4.40)$$

where

$$\rho_k^T := \max_{t_k^r \leq T} \beta_{\mathbb{C}} \int_{t_k^{r-1}}^{t_k^r} \|E\dot{w}(t)\|_2 dt \rightarrow 0$$

by the absolute continuity of the integral. Iterating now inequality (4.40) for $1 \leq r \leq i$, we get (4.34) with $\omega_k^T := \rho_k^T \int_0^T \|E\dot{w}(t)\|_2 dt$. \square

4.5. Proof of the main theorem. Let us fix a sequence of subdivisions $(t_k^i)_{i \geq 0}$ of the half-line $[0, +\infty)$ satisfying (4.9) and (4.10). For every k let $(u_k^i, e_k^i, \boldsymbol{\mu}_k^i)$, $i = 1, \dots, k$, be defined inductively as minimizers of the functional (4.11) on the sets A_k^i , with $(u_k^0, e_k^0, \boldsymbol{\mu}_k^0) = (u_0, e_0, \delta_{(p_0, z_0)})$, and let $\mathbf{u}_k(t)$, $\mathbf{e}_k(t)$, $\boldsymbol{\sigma}_k(t)$, $\mathbf{w}_k(t)$, and $[t]_k$ be defined by (4.29) and let $\boldsymbol{\mu}_k$ be the unique system in $SGY([0, +\infty); \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ whose restrictions to the intervals $[0, t_k^i]$ coincide with the piecewise constant interpolations of $\boldsymbol{\mu}_k^i$ (see [4, Definition 7.10]). Using Lemma 2.1 and the definition of A_k^i we can prove by induction on i that $(\mathbf{u}_k, \mathbf{e}_k, \boldsymbol{\mu}_k) \in AY(\{t_k^0, t_k^1, \dots, t_k^i\}, \mathbf{w}_k)$ for every i and k . This implies that

$$(\mathbf{u}_k, \mathbf{e}_k, \boldsymbol{\mu}_k) \in AY([0, +\infty), \mathbf{w}_k) \quad (4.41)$$

for every k .

Let us prove that for every $T > 0$ there exists a constant C_T , independent of k , such that

$$\sup_{t \in [0, T]} \|\mathbf{e}_k(t)\|_2 \leq C_T, \quad \text{Var}(\boldsymbol{\mu}_k; 0, T) \leq C_T. \quad (4.42)$$

By (2.30) we have

$$\begin{aligned} & \mathcal{D}_H(\boldsymbol{\mu}_k; 0, t) + \langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle \geq \\ & \geq \langle H(\xi - \xi_0, \theta - \theta_0) + \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_{0t}(x, \xi_0, \theta_0, \xi, \theta, \eta) \rangle \geq \\ & \geq \langle C_V^K |\xi - \xi_0| + C_V^K |\theta - \theta_0| + \{V\}(\theta_0, \eta), (\boldsymbol{\mu}_k)_{0t}(x, \xi_0, \theta_0, \xi, \theta, \eta) \rangle = \\ & = C_V^K \langle |\xi - \xi_0| + |\theta - \theta_0|, (\boldsymbol{\mu}_k)_{0t}(x, \xi_0, \theta_0, \xi, \theta, \eta) \rangle + \mathcal{V}(z_0), \end{aligned} \quad (4.43)$$

where the last equality follows from the fact that $(\boldsymbol{\mu}_k)_0 = \delta_{(p_0, z_0)}$. From (2.22), (2.23), (4.33), and (4.43) we deduce that

$$\alpha_C \|\mathbf{e}_k(t)\|_2^2 \leq \beta_C \|e_0\|_2^2 + 2\beta_C \sup_{t \in [0, T]} \|\mathbf{e}_k(t)\|_2 \int_0^T \|E\dot{\mathbf{w}}(t)\|_2 dt + \omega_k^T$$

for every k and every $t \in [0, T]$. The first estimate in (4.42) can be obtained now by using the Cauchy inequality.

By (4.33) and the first inequality in (4.42) we have that

$$\mathcal{D}_H(\boldsymbol{\mu}_k; 0, t) + \langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle \quad (4.44)$$

is bounded uniformly with respect to k and $t \in [0, T]$. By (4.43) this implies the boundedness of

$$\langle |\xi - \xi_0| + |\theta - \theta_0|, (\boldsymbol{\mu}_k)_{0t}(x, \xi_0, \theta_0, \xi, \theta, \eta) \rangle. \quad (4.45)$$

By the compatibility condition (7.2) of [4] and by the equality $(\boldsymbol{\mu}_k)_0 = \delta_{(p_0, z_0)}$ we have

$$\langle |\xi_0| + |\theta_0|, (\boldsymbol{\mu}_k)_{0t}(x, \xi_0, \theta_0, \xi, \theta, \eta) \rangle = \|p_0\|_1 + \|z_0\|_1,$$

which, together with the boundedness of (4.45), gives that $\langle |\xi| + |\theta|, (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle$ is bounded. This implies that $\langle \{V\}(\theta, \eta), (\boldsymbol{\mu}_k)_t(x, \xi, \theta, \eta) \rangle$ is bounded too, so that (2.13) and the boundedness of (4.44) yield the second estimate in (4.42).

By the Helly Theorem for compatible systems of generalized Young measures proved in [4, Theorem 8.10] there exist a subsequence, still denoted $\boldsymbol{\mu}_k$, a set $\Theta \subset [0, +\infty)$, containing 0 and with $[0, +\infty) \setminus \Theta$ at most countable, and a left continuous $\boldsymbol{\mu} \in SGY([0, +\infty), \overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, with

$$\sup_{t \in [0, T]} \|\boldsymbol{\mu}_t\|_* < +\infty, \quad \text{Var}(\boldsymbol{\mu}; 0, T) < +\infty \quad (4.46)$$

for every $T > 0$, such that

$$(\boldsymbol{\mu}_k)_{t_1 \dots t_m} \rightharpoonup \boldsymbol{\mu}_{t_1 \dots t_m} \quad \text{weakly}^* \text{ in } GY(\overline{\Omega}; (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^m) \quad (4.47)$$

for every finite sequence t_1, \dots, t_m in Θ with $t_1 < \dots < t_m$.

Let $\mathbf{p}_k(t) \in M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, $\mathbf{z}_k(t) \in M_b(\overline{\Omega})$, $\mathbf{p}(t) \in M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, and $\mathbf{z}(t) \in M_b(\overline{\Omega})$ be the measures defined by

$$(\mathbf{p}_k(t), \mathbf{z}_k(t)) := \text{bar}((\boldsymbol{\mu}_k)_t) \quad \text{and} \quad (\mathbf{p}(t), \mathbf{z}(t)) := \text{bar}(\boldsymbol{\mu}_t). \quad (4.48)$$

By (4.47) and by [4, Remark 6.4] we have

$$\mathbf{p}_k(t) \rightharpoonup \mathbf{p}(t) \quad \text{weakly}^* \text{ in } M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2}) \quad (4.49)$$

for every $t \in \Theta$.

By Corollary 4.10 the sequence $(\mathbf{u}_k(t), \mathbf{e}_k(t), \mathbf{p}_k(t))$ coincides with the discrete-time approximation of the quasistatic evolution corresponding to the function $\xi \mapsto H_{\text{eff}}(\xi, 0)$ according to [3, Definition 4.2]. Using [3, Theorems 4.5, 4.8, 5.2] we obtain that there exist a subsequence, still denoted $(\mathbf{u}_k, \mathbf{e}_k, \mathbf{p}_k)$, a continuous function $t \mapsto (\mathbf{u}(t), \mathbf{e}(t))$ from $[0, +\infty)$ into $BD(\Omega) \times L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2})$ and an extension of $t \mapsto \mathbf{p}(t)$ to $[0, +\infty)$, still denoted by the same symbol, such that $t \mapsto (\mathbf{u}(t), \mathbf{e}(t), \mathbf{p}(t))$ is a quasistatic evolution of the problem corresponding to the function $\xi \mapsto H_{\text{eff}}(\xi, 0)$, and

$$\mathbf{e}_k(t) \rightarrow \mathbf{e}(t) \quad \text{strongly in } L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}), \quad (4.50)$$

$$\mathbf{u}_k(t) \rightharpoonup \mathbf{u}(t) \quad \text{weakly}^* \text{ in } BD(\Omega), \quad (4.51)$$

for every $t \in [0, +\infty)$.

By Remark 4.4 and by (4.41) the triple $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ belongs to $AY(\Theta, \mathbf{w})$. By the left continuity of \mathbf{u} , \mathbf{e} , $\boldsymbol{\mu}$ we have also $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu}) \in AY([0, +\infty), \mathbf{w})$.

Let us fix $(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{reg}(0)$ and $\tilde{z} \in L^1(\Omega)$. Passing to the limit in (4.32) thanks to Lemma 2.1 we obtain that (ev1) is satisfied for every $t \in \Theta$. By left continuity the same inequality holds for every $t \in [0, +\infty)$.

By (4.42) and by the weak* lower semicontinuity of the dissipation we can pass to the limit in (4.33) and we obtain

$$\begin{aligned} \mathcal{Q}(\mathbf{e}(T)) + \mathcal{D}_H(\boldsymbol{\mu}; 0, T) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle &\leq \\ &\leq \mathcal{Q}(\mathbf{e}_0) + \mathcal{V}(z_0) + \int_0^T \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt, \end{aligned} \quad (4.52)$$

for every $T \in \Theta$. By left continuity the same inequality holds for every $T \in [0, +\infty)$.

Passing to the limit in (4.31), we obtain

$$\begin{aligned} \mathcal{Q}(\mathbf{e}(t)) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle &\leq \\ &\leq \mathcal{Q}(\mathbf{e}(\hat{t})) - \langle \boldsymbol{\sigma}(\hat{t}), E\mathbf{w}(\hat{t}) - E\mathbf{w}(t) \rangle + \mathcal{Q}(E\mathbf{w}(\hat{t}) - E\mathbf{w}(t)) + \\ &+ \langle H(\hat{\xi} - \xi, \hat{\theta} - \theta), \boldsymbol{\mu}_{\hat{t}\hat{t}}(x, \xi, \theta, \hat{\xi}, \hat{\theta}, \eta) \rangle + \langle \{V\}(\hat{\theta}, \eta), \boldsymbol{\mu}_{\hat{t}}(x, \hat{\xi}, \hat{\theta}, \eta) \rangle \end{aligned} \quad (4.53)$$

for every $t, \hat{t} \in \Theta$ with $t < \hat{t}$. By left continuity the same inequality holds for every $t, \hat{t} \in [0, +\infty)$ with $t < \hat{t}$.

Using this inequality, we want to prove that

$$\begin{aligned} \mathcal{Q}(\mathbf{e}(T)) + \mathcal{D}_H(\boldsymbol{\mu}; 0, T) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle &\geq \\ &\geq \mathcal{Q}(\mathbf{e}_0) + \mathcal{V}(z_0) + \int_0^T \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt, \end{aligned} \quad (4.54)$$

for every $T \in (0, +\infty)$.

Let us fix $T \in (0, +\infty)$ and a sequence of subdivisions $(s_k^i)_{0 \leq i \leq k}$ of $[0, T]$ with

$$\begin{aligned} 0 = s_k^0 &< s_k^1 < \dots < s_k^{k-1} < s_k^k = T, \\ \lim_{k \rightarrow \infty} \max_{1 \leq i \leq k} (s_k^i - s_k^{i-1}) &= 0. \end{aligned} \quad (4.55)$$

For every $i = 1, \dots, k$ we apply (4.53) with times s_k^{i-1} and s_k^i , and we obtain

$$\begin{aligned} & \mathcal{Q}(e(s_k^{i-1})) + \langle \{V\}(\theta_{i-1}, \eta), \boldsymbol{\mu}_{s_k^{i-1}}(x, \xi_{i-1}, \theta_{i-1}, \eta) \rangle \leq \\ & \leq \mathcal{Q}(e(s_k^i)) - \langle \boldsymbol{\sigma}(s_k^i), E\mathbf{w}(s_k^i) - E\mathbf{w}(s_k^{i-1}) \rangle + \mathcal{Q}(E\mathbf{w}(s_k^i) - E\mathbf{w}(s_k^{i-1})) + \\ & + \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\mu}_{s_k^{i-1} s_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle + \langle \{V\}(\theta_i, \eta), \boldsymbol{\mu}_{s_k^i}(x, \xi_i, \theta_i, \eta) \rangle. \end{aligned} \quad (4.56)$$

We notice that

$$\begin{aligned} \langle \boldsymbol{\sigma}(s_k^i), E\mathbf{w}(s_k^i) - E\mathbf{w}(s_k^{i-1}) \rangle &= \int_{s_k^{i-1}}^{s_k^i} \langle \boldsymbol{\sigma}(s_k^i), E\dot{\mathbf{w}}(s) \rangle ds, \\ \mathcal{Q}(E\mathbf{w}(s_k^i) - E\mathbf{w}(s_k^{i-1})) &\leq \beta_{\mathbb{C}} \left(\int_{s_k^{i-1}}^{s_k^i} \|E\dot{\mathbf{w}}(s)\|_2 ds \right)^2, \\ \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\mu}_{s_k^{i-1} s_k^i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle &\leq \mathcal{D}_H(\boldsymbol{\mu}; s_k^{i-1}, s_k^i). \end{aligned} \quad (4.57)$$

On $[0, T]$ we define the piecewise constant function $\bar{\boldsymbol{\sigma}}_k(s) := \boldsymbol{\sigma}(s_k^i)$, where i is the smallest index such that $s \leq s_k^i$. Summing the inequalities (4.56) for $1 \leq i \leq k$, we obtain

$$\begin{aligned} \mathcal{Q}(e(T)) + \mathcal{D}_H(\boldsymbol{\mu}; 0, T) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle + \rho_k \int_0^T \|E\dot{\mathbf{w}}(s)\|_2 ds &\geq \\ \geq \mathcal{Q}(e_0) + \mathcal{V}(z_0) + \int_0^T \langle \bar{\boldsymbol{\sigma}}_k(s), E\dot{\mathbf{w}}(s) \rangle ds, \end{aligned} \quad (4.58)$$

where

$$\rho_k := \sup_{1 \leq i \leq k} \beta_{\mathbb{C}} \int_{s_k^{i-1}}^{s_k^i} \|E\dot{\mathbf{w}}(s)\|_2 ds.$$

Now conditions (4.55) and the continuity of $\boldsymbol{\sigma}$ guarantee that $\rho_k \rightarrow 0$ and that $\bar{\boldsymbol{\sigma}}_k \rightarrow \boldsymbol{\sigma}$ strongly in $L^2([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{2 \times 2}))$. Hence, taking the limit as $k \rightarrow \infty$ in (4.58), we obtain inequality (4.54), which, together with (4.52), gives (ev2).

4.6. Some properties of the solutions. We conclude this section by proving some qualitative properties of the Young measure solutions to the evolution problem.

Theorem 4.16. *Let $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ be a globally stable quasistatic evolution of Young measures. For every $t \in [0, +\infty)$ let $(\mathbf{p}(t), \mathbf{z}(t)) := \text{bar}(\boldsymbol{\mu}_t)$. Then $(\mathbf{u}, \mathbf{e}, \mathbf{p})$ is a quasistatic evolution corresponding to the function $\xi \mapsto H_{\text{eff}}(\xi, 0)$ according to [3, Definition 4.2].*

Proof. Let us fix $t \in [0, +\infty)$. We want to prove that

$$\mathcal{Q}(\mathbf{e}(t)) \leq \mathcal{Q}(\mathbf{e}(t) + \tilde{\mathbf{e}}) + \mathcal{H}_{\text{eff}}(\tilde{\mathbf{p}}, 0). \quad (4.59)$$

for every $(\tilde{\mathbf{u}}, \tilde{\mathbf{e}}, \tilde{\mathbf{p}}) \in A(0)$. Let us fix $(\tilde{\mathbf{u}}, \tilde{\mathbf{e}}, \tilde{\mathbf{p}})$, let (μ_t^Y, μ_t^∞) be the pair of measures associated with $\boldsymbol{\mu}_t$ by [4, Theorem 4.3], let $(\mu_t^{x,Y})_{x \in \Omega}$ be the disintegration of μ_t^Y considered in [4, Remark 4.5], let

$$F(x, \tilde{\xi}, \tilde{\theta}) := \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} [H(\tilde{\xi}, \tilde{\theta}) + V(\theta + \tilde{\theta}) - V(\theta)] d\mu_t^{x,Y}(\xi, \theta),$$

let F^∞ be the recession function of F with respect to $(\tilde{\xi}, \tilde{\theta})$, and let $\text{co} F$ be the convex envelope of F with respect to $(\tilde{\xi}, \tilde{\theta})$. It is easy to see that $F^\infty = H + V^\infty$. We claim that

$$\text{co} F = H_{\text{eff}}. \quad (4.60)$$

Indeed, as V is concave, we have $V(\theta + \tilde{\theta}) - V(\theta) \geq V^\infty(\tilde{\theta})$, which gives

$$F(x, \tilde{\xi}, \tilde{\theta}) \geq \int_{\mathbb{M}_D^{2 \times 2} \times \mathbb{R}} [H(\tilde{\xi}, \tilde{\theta}) + V^\infty(\tilde{\theta})] d\mu_t^{x,Y}(\xi, \theta) = H(\tilde{\xi}, \tilde{\theta}) + V^\infty(\tilde{\theta}) \geq H_{\text{eff}}(\tilde{\xi}, \tilde{\theta}),$$

where the intermediate equality follows from the fact that $\mu_t^{x,Y}$ is a probability measure. This implies $\text{co} F \geq H_{\text{eff}}$. The opposite inequality can be obtained arguing as in the proof of (3.4).

By Theorem 3.8 there exist a sequence $(\tilde{u}_k, \tilde{e}_k, \tilde{p}_k) \in A(0)$ and a sequence $\tilde{z}_k \in L^1(\Omega)$ such that $\tilde{p}_k \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$, $\tilde{e}_k \rightarrow \tilde{e}$ strongly in $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, $\tilde{p}_k \rightharpoonup \tilde{p}$ weakly* in $M_b(\overline{\Omega}; \mathbb{M}_D^{2 \times 2})$, $\tilde{z}_k \rightharpoonup 0$ weakly* in $M_b(\overline{\Omega})$, and

$$\mathcal{H}(\tilde{p}_k, \tilde{z}_k) + \mathcal{V}(\tilde{z}_k) \longrightarrow \mathcal{H}_{\text{eff}}(\tilde{p}, 0) + \mathcal{V}(0).$$

Passing to a subsequence, we may assume that $\delta_{(\tilde{p}_k, \tilde{z}_k)}$ converges weakly* to some $\nu \in GY(\overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$. We note that $\text{bar}(\nu) = (\tilde{p}, 0)$ and that

$$\langle H(\tilde{\xi}, \tilde{\theta}) + \{V\}(\tilde{\theta}, \eta), \nu(x, \tilde{\xi}, \tilde{\theta}, \eta) \rangle = \mathcal{H}_{\text{eff}}(\tilde{p}, 0) + \mathcal{V}(0).$$

By Theorem 3.10 we deduce that

$$\nu = \delta_{(0,0)} + \frac{1}{2}\omega_{(\tilde{p}^a, z)}^{\mathcal{C}^2} + \frac{1}{2}\omega_{(\tilde{p}^a, -z)}^{\mathcal{C}^2} + \frac{1}{2}\omega_{(\tilde{p}^\lambda, z_\lambda)}^\lambda + \frac{1}{2}\omega_{(\tilde{p}^\lambda, -z_\lambda)}^\lambda, \quad (4.61)$$

where $\lambda := |\tilde{p}^s|$ and \tilde{p}^λ is the Radon-Nikodym derivative of \tilde{p}^s with respect to $|\tilde{p}^s|$, while z and z_λ are two nonnegative functions such that

$$\begin{aligned} H_{\text{eff}}(\tilde{p}^a, 0) &= H(\tilde{p}^a, z) + V^\infty(z) \quad \text{a.e. in } \Omega \\ H_{\text{eff}}(\tilde{p}^\lambda, 0) &= H(\tilde{p}^\lambda, z_\lambda) + V^\infty(z_\lambda) \quad \lambda\text{-a.e. in } \overline{\Omega}. \end{aligned} \quad (4.62)$$

As $\delta_{(\tilde{p}_k, \tilde{z}_k)} \rightharpoonup \nu$ weakly* in $GY(\overline{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$, we have

$$\int_{\Omega} F(x, \tilde{p}_k, \tilde{z}_k) dx \longrightarrow \langle \{F\}, \nu \rangle,$$

where

$$\{F\}(\tilde{\xi}, \tilde{\theta}, \eta) := \begin{cases} \eta F(\tilde{\xi}/\eta, \tilde{\theta}/\eta) & \text{if } \eta > 0, \\ F^\infty(\tilde{\xi}, \tilde{\theta}) & \text{if } \eta \leq 0. \end{cases}$$

As $F^\infty = H + V^\infty$, using (4.61) and (4.62) we obtain $\langle \{F\}, \nu \rangle = \mathcal{H}_{\text{eff}}(\tilde{p}, 0)$, hence using also the strong convergence of \tilde{e}_k to \tilde{e} , we deduce

$$\mathcal{Q}(e(t) + \tilde{e}_k) + \int_{\Omega} F(x, \tilde{p}_k, \tilde{z}_k) dx \longrightarrow \mathcal{Q}(e(t) + \tilde{e}) + \mathcal{H}_{\text{eff}}(\tilde{p}, 0).$$

From the definition of $F(x, \tilde{\xi}, \tilde{\theta})$ this is equivalent to saying that

$$\mathcal{Q}(e(t) + \tilde{e}_k) + \mathcal{H}(\tilde{p}_k, \tilde{z}_k) + \langle \{V\}(\theta + \eta \tilde{z}_k(x), \eta) - \{V\}(\theta, \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle$$

converges to $\mathcal{Q}(e(t) + \tilde{e}) + \mathcal{H}_{\text{eff}}(\tilde{p}, 0)$. Since

$$\mathcal{Q}(e(t)) \leq \mathcal{Q}(e(t) + \tilde{e}_k) + \mathcal{H}(\tilde{p}_k, \tilde{z}_k) + \langle \{V\}(\theta + \eta \tilde{z}_k(x), \eta) - \{V\}(\theta, \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle$$

by (ev1), we obtain (4.59) by passing to the limit as $k \rightarrow \infty$.

Thanks to [3, Theorem 4.7], to conclude the proof of the theorem it is enough to show that

$$\mathcal{Q}(e(T)) + \mathcal{D}_{H_{\text{eff}}}((\mathbf{p}, 0); 0, T) \leq \mathcal{Q}(e_0) + \int_0^T \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt \quad (4.63)$$

for every $T \in (0, +\infty)$, where $\mathcal{D}_{H_{\text{eff}}}((\mathbf{p}, 0); 0, T)$ is defined as in [3, Section 4]. By (ev2) it suffices to prove that

$$\begin{aligned} \mathcal{D}_{H_{\text{eff}}}((\mathbf{p}, 0); 0, T) &\leq \mathcal{D}_H(\boldsymbol{\mu}; 0, T) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle - \\ &\quad - \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_0(x, \xi, \theta, \eta) \rangle. \end{aligned} \quad (4.64)$$

To show this we fix any subdivision $(t_i)_{0 \leq i \leq k}$ of the interval $[0, T]$. From the definition of \mathcal{D}_H we obtain, using the compatibility condition (7.2) of [4],

$$\begin{aligned} & \mathcal{D}_H(\boldsymbol{\mu}; 0, T) + \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle - \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_0(x, \xi, \theta, \eta) \rangle \geq \\ & \geq \sum_{i=1}^k \langle H(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}) + \{V\}(\theta_i, \eta) - \{V\}(\theta_{i-1}, \eta), \boldsymbol{\mu}_{t_{i-1}t_i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle \geq \\ & \geq \sum_{i=1}^k \langle H_{\text{eff}}(\xi_i - \xi_{i-1}, \theta_i - \theta_{i-1}), \boldsymbol{\mu}_{t_{i-1}t_i}(x, \xi_{i-1}, \theta_{i-1}, \xi_i, \theta_i, \eta) \rangle \geq \\ & \geq \sum_{i=1}^k \mathcal{H}_{\text{eff}}(\mathbf{p}(t_i) - \mathbf{p}(t_{i-1}), \mathbf{z}(t_i) - \mathbf{z}(t_{i-1})), \end{aligned}$$

where the last inequality follows from the Jensen inequality. Recalling that $H_{\text{eff}}(\xi, \theta) \geq H_{\text{eff}}(\xi, 0)$ for every ξ and θ , from the arbitrariness of the subdivision we obtain (4.64). \square

Every globally stable quasistatic evolution of Young measures is absolutely continuous with respect to time, as made precise by the following theorem.

Theorem 4.17. *Let $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ be a globally stable quasistatic evolution of Young measures. Then for every $T \in [0, +\infty)$ the functions \mathbf{u} and \mathbf{e} are absolutely continuous on $[0, T]$ with values in $BD(\Omega)$ and $L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$, respectively, while $\boldsymbol{\mu}$ is absolutely continuous on $[0, T]$ according to [4, Definition 10.1].*

Proof. The assertion on \mathbf{u} and \mathbf{e} follows from Theorem 4.16 and [3, Theorem 5.2].

By (2.30) we have

$$C_V^K \text{Var}(\boldsymbol{\mu}; t_1, t_2) \leq \mathcal{D}_H(\boldsymbol{\mu}; t_1, t_2) + \langle \{V\}(\theta_2, \eta), \boldsymbol{\mu}_{t_2}(x, \xi_2, \theta_2, \eta) \rangle - \langle \{V\}(\theta_1, \eta), \boldsymbol{\mu}_{t_1}(x, \xi_1, \theta_1, \eta) \rangle.$$

It follows from the energy balance (ev2) that the right-hand side of the previous inequality is equal to

$$\mathcal{Q}(\mathbf{e}(t_1)) - \mathcal{Q}(\mathbf{e}(t_2)) + \int_{t_1}^{t_2} \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt,$$

therefore

$$\begin{aligned} & C_V^K \langle |\xi_2 - \xi_1| + |\theta_2 - \theta_1|, \boldsymbol{\mu}_{t_1 t_2}(x, \xi_1, \theta_1, \xi_2, \theta_2, \eta) \rangle \leq \\ & \leq |\mathcal{Q}(\mathbf{e}(t_1)) - \mathcal{Q}(\mathbf{e}(t_2))| + \int_{t_1}^{t_2} |\langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle| dt. \end{aligned}$$

Since the functions $t \mapsto \mathcal{Q}(\mathbf{e}(t))$ and $t \mapsto \int_0^t |\langle \boldsymbol{\sigma}(s), E\dot{\mathbf{w}}(s) \rangle| ds$ are absolutely continuous, we conclude that $\boldsymbol{\mu}$ satisfies [4, Definition 10.1]. \square

Owing to the previous theorem, if $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ is a globally stable quasistatic evolution of Young measures, then $\boldsymbol{\mu}$ has a weak* derivative $\dot{\boldsymbol{\mu}}_t$ at a.e. time $t \in [0, +\infty)$ in the sense of [4, Definition 9.4]. The next theorem deals with the structure of $\dot{\boldsymbol{\mu}}_t$ and shows that the finite part $\overline{\boldsymbol{\mu}}_t^Y$ of $\boldsymbol{\mu}_t$ does not evolve.

Theorem 4.18. *Let $\overline{\mathbf{p}}_0 \in L^1(\Omega; \mathbb{M}_D^{2 \times 2})$, $\overline{\mathbf{z}}_0 \in L^1(\Omega)$, $\mathbf{w} \in AC_{\text{loc}}([0, +\infty); H^1(\Omega; \mathbb{R}^2))$, let $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ be a globally stable quasistatic evolution of Young measures with boundary datum \mathbf{w} such that $\overline{\boldsymbol{\mu}}_0^Y = \delta_{(\overline{\mathbf{p}}_0, \overline{\mathbf{z}}_0)}$, and let $(\mathbf{p}(t), \mathbf{z}(t)) := \text{bar}(\boldsymbol{\mu}_t)$. Denote the total variation of the measure $(\dot{\mathbf{p}}^s(t), \dot{\mathbf{z}}^s(t))$ by $\boldsymbol{\lambda}(t)$, and let $(\dot{\mathbf{p}}^\lambda(t), \dot{\mathbf{z}}^\lambda(t))$ be the Radon-Nikodym derivative of the measure $(\dot{\mathbf{p}}^s(t), \dot{\mathbf{z}}^s(t))$ with respect to $\boldsymbol{\lambda}(t)$. By Lemma 3.3 for a.e. $t \in [0, +\infty)$ there exist $\hat{\mathbf{z}}(t) \in L^1(\Omega)$, with $\hat{\mathbf{z}}(t) \dot{\mathbf{z}}^a(t) \geq 0$ a.e. on Ω , and $\boldsymbol{\alpha}(t) \in L^\infty(\Omega)$, with $0 \leq \boldsymbol{\alpha}(t) \leq 1$ a.e. on Ω , such that*

$$\begin{aligned} \dot{\mathbf{z}}^a(t) &= \boldsymbol{\alpha}(t) \hat{\mathbf{z}}(t) + (1 - \boldsymbol{\alpha}(t))(-\hat{\mathbf{z}}(t)), \\ H_{\text{eff}}(\dot{\mathbf{p}}^a(t), \dot{\mathbf{z}}^a(t)) &= H(\dot{\mathbf{p}}^a(t), \hat{\mathbf{z}}(t)) + V^\infty(\hat{\mathbf{z}}(t)) \end{aligned} \tag{4.65}$$

a.e. in Ω , and there exist $\hat{z}_\lambda(t) \in L^1_{\lambda(t)}(\bar{\Omega})$, with $\hat{z}_\lambda(t) \dot{z}^\lambda(t) \geq 0$ $\lambda(t)$ -a.e. on $\bar{\Omega}$, and $\alpha_\lambda(t) \in L^\infty_{\lambda(t)}(\bar{\Omega})$, with $0 \leq \alpha_\lambda(t) \leq 1$ $\lambda(t)$ -a.e. on $\bar{\Omega}$, such that

$$\begin{aligned} \dot{z}^\lambda(t) &= \alpha_\lambda(t) \hat{z}_\lambda(t) + (1 - \alpha_\lambda(t))(-\hat{z}_\lambda(t)), \\ H_{\text{eff}}(\dot{\mathbf{p}}^\lambda(t), \dot{z}^\lambda(t)) &= H(\dot{\mathbf{p}}^\lambda(t), \hat{z}_\lambda(t)) + V^\infty(\hat{z}_\lambda(t)) \end{aligned} \quad (4.66)$$

$\lambda(t)$ -a.e. in $\bar{\Omega}$. Then

$$\begin{aligned} \dot{\boldsymbol{\mu}}_t &= \delta_{(0,0)} + \alpha(t) \omega_{(\dot{\mathbf{p}}^\alpha(t), \hat{z}(t))}^{\mathcal{L}^2} + (1 - \alpha(t)) \omega_{(\dot{\mathbf{p}}^\alpha(t), -\hat{z}(t))}^{\mathcal{L}^2} + \\ &+ \alpha_\lambda(t) \omega_{(\dot{\mathbf{p}}^\lambda(t), \hat{z}_\lambda(t))}^{\lambda(t)} + (1 - \alpha_\lambda(t)) \omega_{(\dot{\mathbf{p}}^\lambda(t), -\hat{z}_\lambda(t))}^{\lambda(t)} \end{aligned} \quad (4.67)$$

and

$$\mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), \dot{z}(t)) = \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), 0) \quad (4.68)$$

for a.e. $t \in [0, +\infty)$. Moreover,

$$\bar{\boldsymbol{\mu}}_t^Y = \delta_{(\bar{\mathbf{p}}_0, \bar{z}_0)} \quad (4.69)$$

for every $t \in [0, +\infty)$.

Proof. As the system $\boldsymbol{\mu}$ is absolutely continuous with respect to time by Theorem 4.17, we have by [4, Theorem 10.4] that for every $t_1, t_2 \in [0, +\infty)$ with $t_1 < t_2$

$$\mathcal{D}_H(\boldsymbol{\mu}; t_1, t_2) = \int_{t_1}^{t_2} \langle H(\xi, \theta), \dot{\boldsymbol{\mu}}_t(x, \xi, \theta, \eta) \rangle dt, \quad (4.70)$$

where $\dot{\boldsymbol{\mu}}_t$ is the weak* derivative of $\boldsymbol{\mu}$ at time t in the sense of [4, Definition 9.4]. By [4, Remark 9.6] we also have that the maps $t \mapsto \mathbf{p}(t)$ and $t \mapsto \mathbf{z}(t)$ are absolutely continuous and $(\dot{\mathbf{p}}(t), \dot{z}(t)) = \text{bar}(\dot{\boldsymbol{\mu}}_t)$ for a.e. $t \in [0, +\infty)$.

From Theorem 4.16 and [3, Proposition 5.6] it follows that

$$\mathcal{Q}(e(t_2)) + \int_{t_1}^{t_2} \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), 0) dt = \mathcal{Q}(e(t_1)) + \int_{t_1}^{t_2} \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt.$$

By (ev2) and (4.70) we deduce that for every $t_1 < t_2$

$$\int_{t_1}^{t_2} \langle H(\xi, \theta), \dot{\boldsymbol{\mu}}_t \rangle dt + \langle \{V\}(\theta_2, \eta) - \{V\}(\theta_1, \eta), \boldsymbol{\mu}_{t_1 t_2} \rangle = \int_{t_1}^{t_2} \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), 0) dt, \quad (4.71)$$

where the measure $\dot{\boldsymbol{\mu}}_t$ acts on (x, ξ, θ, η) , while $\boldsymbol{\mu}_{t_1 t_2}$ acts on $(x, \xi_1, \theta_1, \xi_2, \theta_2, \eta)$. By concavity we have $\{V\}(\theta_2, \eta) - \{V\}(\theta_1, \eta) \geq V^\infty(\theta_2 - \theta_1)$, so that (4.71) yields

$$\int_{t_1}^{t_2} \langle H(\xi, \theta), \dot{\boldsymbol{\mu}}_t \rangle dt + \langle V^\infty(\theta_2 - \theta_1), \boldsymbol{\mu}_{t_1 t_2} \rangle \leq \int_{t_1}^{t_2} \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), 0) dt. \quad (4.72)$$

Dividing by $t_2 - t_1$ and letting $t_2 \rightarrow t_1 = t$ we obtain

$$\langle H(\xi, \theta) + V^\infty(\theta), \dot{\boldsymbol{\mu}}_t(x, \xi, \theta, \eta) \rangle \leq \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), 0) \leq \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), \dot{z}(t)) \quad (4.73)$$

for a.e. $t \in [0, +\infty)$. Using the Jensen inequality for generalized Young measures [4, Theorem 6.5] we conclude that

$$\langle H(\xi, \theta) + V^\infty(\theta), \dot{\boldsymbol{\mu}}_t(x, \xi, \theta, \eta) \rangle = \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), 0) = \mathcal{H}_{\text{eff}}(\dot{\mathbf{p}}(t), \dot{z}(t)) \quad (4.74)$$

for a.e. $t \in [0, +\infty)$, which shows, in particular, (4.68). By taking the derivative of (4.71) we obtain from (4.74)

$$\lim_{t_2 \rightarrow t_1^+} \langle \frac{1}{t_2 - t_1} (\{V\}(\theta_2, \eta) - \{V\}(\theta_1, \eta)) - V^\infty(\frac{\theta_2 - \theta_1}{t_2 - t_1}), \boldsymbol{\mu}_{t_1 t_2} \rangle = 0. \quad (4.75)$$

As $-V^\infty$ is convex and positively homogeneous, by [4, Theorem 10.4] we have that for every $t \in [0, +\infty)$

$$\langle V^\infty(\theta_1 - \theta_0), \boldsymbol{\mu}_{0t} \rangle \geq \int_0^t \langle V^\infty, \dot{\boldsymbol{\mu}}_s \rangle ds,$$

where $\boldsymbol{\mu}_{0t}$ acts on the variables $(x, \xi_0, \theta_0, \xi_1, \theta_1, \eta)$. Since $s \mapsto \langle \{V\}, \boldsymbol{\mu}_s \rangle$ is absolutely continuous and by (4.75)

$$\langle V^\infty, \dot{\boldsymbol{\mu}}_s \rangle = \frac{d}{ds} \langle \{V\}, \boldsymbol{\mu}_s \rangle$$

for a.e. $s \in [0, +\infty)$, we deduce that

$$\langle V^\infty(\theta_1 - \theta_0), \boldsymbol{\mu}_{0t} \rangle \geq \langle \{V\}(\theta_1, \eta) - \{V\}(\theta_0, \eta), \boldsymbol{\mu}_{0t} \rangle. \quad (4.76)$$

Let $\psi: \overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^2 \times \mathbb{R} \rightarrow \Omega \times \mathbb{R} \times \mathbb{R}$ be defined by

$$\psi(x, \xi_0, \theta_0, \xi_1, \theta_1, \eta) := (x, \theta_1 - \theta_0, \eta),$$

and let $\nu := \psi(\boldsymbol{\mu}_{0t})$. By repeating the arguments used in the proof of (3.78)–(3.80) we obtain

$$\langle \{V\}(\theta_1, \eta) - \{V\}(\theta_0, \eta), \boldsymbol{\mu}_{0t} \rangle \geq \langle \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta), \nu \rangle,$$

where ν acts on $(x, \tilde{\theta}, \eta)$. This inequality, together with (4.76), yields

$$\langle V^\infty, \nu \rangle \geq \langle \{V\}(\eta \bar{z}_0(x) + \tilde{\theta}, \eta) - \{V\}(\eta \bar{z}_0(x), \eta), \nu \rangle,$$

By (2.27) it follows that $\bar{\nu}^Y = \delta_0$. Arguing as in the proof of (3.82), we conclude that

$$\pi_{\mathbb{R}}(\overline{\boldsymbol{\mu}}_t^Y) = \delta_{\bar{z}_0}. \quad (4.77)$$

where $\pi_{\mathbb{R}}: \overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R} \rightarrow \overline{\Omega} \times \mathbb{R} \times \mathbb{R}$ is defined by $\pi_{\mathbb{R}}(x, \xi, \theta, \eta) := (x, \theta, \eta)$.

As the system $\boldsymbol{\mu}$ is absolutely continuous with respect to time, we also have that the system $\overline{\boldsymbol{\mu}}^Y$ given by $\overline{\boldsymbol{\mu}}_{t_1 \dots t_m}^Y := \overline{(\boldsymbol{\mu}_{t_1 \dots t_m})^Y}$ is absolutely continuous, so that its weak* derivative $\dot{\overline{\boldsymbol{\mu}}}_t^Y$ exists for a.e. $t \in [0, +\infty)$. We note that in general the weak* derivative $\dot{\overline{\boldsymbol{\mu}}}_t^Y$ of $\overline{\boldsymbol{\mu}}^Y$ does not coincide with the finite part $(\dot{\boldsymbol{\mu}}_t)^Y$ of $\dot{\boldsymbol{\mu}}_t$. Nevertheless the following identity holds

$$\overline{(\dot{\boldsymbol{\mu}}_t)^Y} = \dot{\overline{\boldsymbol{\mu}}}_t^Y. \quad (4.78)$$

To see this we observe that at every time t where both $\dot{\boldsymbol{\mu}}_t$ and $\dot{\overline{\boldsymbol{\mu}}}_t^Y$ are defined, there also exists the weak* limit of the difference quotients $q_{ts}(\dot{\boldsymbol{\mu}}_{ts}^\infty)$ as $s \rightarrow t^+$ (see [4, Definition 9.1]), which is an element of $M_*^+(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$ supported on $\{\eta = 0\}$ (see [4, Definition 2.8]). The equality (4.78) follows now from the identity

$$\dot{\boldsymbol{\mu}}_t = \dot{\overline{\boldsymbol{\mu}}}_t^Y + \lim_{s \rightarrow t^+} q_{ts}(\dot{\boldsymbol{\mu}}_{ts}^\infty).$$

Using (4.77) and Corollary 3.12 it is easy to see that $\psi(\overline{\boldsymbol{\mu}}_{ts}^Y) = \delta_0$ for every $s > t$, which, in turn, implies

$$\pi_{\mathbb{R}}(\dot{\overline{\boldsymbol{\mu}}}_t^Y) = \delta_0. \quad (4.79)$$

By (4.78), (4.79), and [4, Lemma 4.8] we deduce that $\pi_{\mathbb{R}}(\overline{(\dot{\boldsymbol{\mu}}_t)^Y}) = \delta_0$. It follows that

$$\langle \{V\}(\theta, \eta), \dot{\boldsymbol{\mu}}_t(x, \xi, \theta, \eta) \rangle = \langle V^\infty(\theta), \dot{\boldsymbol{\mu}}_t(x, \xi, \theta, \eta) \rangle + \mathcal{V}(0),$$

hence by (4.74) we have

$$\langle H(\xi, \theta) + \{V\}(\theta, \eta), \dot{\boldsymbol{\mu}}_t(x, \xi, \theta, \eta) \rangle = \mathcal{H}_{\text{eff}}(\dot{\boldsymbol{p}}(t), \dot{\boldsymbol{z}}(t)) + \mathcal{V}(0) \quad (4.80)$$

for a.e. $t \in [0, +\infty)$. Identity (4.67) is now a consequence of Theorem 3.10 (applied with $p_0 = 0$ and $z_0 = 0$).

We now claim that for almost every $t \in [0, +\infty)$

$$\dot{\overline{\boldsymbol{\mu}}}_t^Y = \delta_{(0,0)}. \quad (4.81)$$

Let us fix $t \in [0, +\infty)$ such that $\dot{\boldsymbol{\mu}}_t$ and $\dot{\overline{\boldsymbol{\mu}}}_t^Y$ exist. By (4.67) and (4.78) there exists $\nu_1^\infty \in M_*^+(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$, with support contained in $\{\eta = 0\}$, such that

$$\dot{\overline{\boldsymbol{\mu}}}_t^Y = \delta_{(0,0)} + \nu_1^\infty. \quad (4.82)$$

By (4.82), (4.79), and Lemma 3.11 we infer that

$$\langle f(x, \xi, \theta, \eta), \nu_1^\infty(x, \xi, \theta, \eta) \rangle = \langle f(x, \xi, 0, \eta), \nu_1^\infty(x, \xi, \theta, \eta) \rangle, \quad (4.83)$$

$$\langle f, \dot{\mu}_t^Y \rangle = \int_{\Omega} f(x, 0, 0, 1) dx + \langle f(x, \xi, 0, \eta), \nu_1^\infty(x, \xi, \theta, \eta) \rangle \quad (4.84)$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$. Let $\nu_2^\infty \in M_*^+(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$ be the weak* limit of the difference quotients $q_{ts}(\dot{\mu}_{ts}^\infty)$ as $s \rightarrow t^+$. From (4.67) and (4.84) it follows that

$$\begin{aligned} \langle f, \nu_2^\infty \rangle &= \int_{\Omega} \alpha f(x, \dot{p}^a, \hat{z}, 0) dx + \int_{\Omega} (1 - \alpha) f(x, \dot{p}^a, -\hat{z}, 0) dx + \\ &+ \int_{\overline{\Omega}} \alpha_\lambda f(x, \dot{p}^\lambda, \hat{z}_\lambda, 0) d\lambda + \int_{\overline{\Omega}} (1 - \alpha_\lambda) f(x, \dot{p}^\lambda, -\hat{z}_\lambda, 0) d\lambda - \\ &- \langle f(x, \xi, 0, \eta), \nu_1^\infty(x, \xi, \theta, \eta) \rangle \end{aligned} \quad (4.85)$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$. In the previous formula and in the remaining part of the proof the dependence upon time is omitted, since t is fixed.

We shall prove that $\nu_1^\infty = 0$, so that claim (4.81) will follow from the decomposition (4.82). According to [4, Remark 4.5], there exist $\pi^\infty \in M_b^+(\overline{\Omega})$ and a family $(\nu_1^{x,\infty})_{x \in \overline{\Omega}}$ of probability measures on $\Sigma := \{\xi \in \mathbb{M}_D^{2 \times 2} : |\xi| = 1\}$ such that

$$\langle f(x, \xi, 0, \eta), \nu_1^\infty(x, \xi, \theta, \eta) \rangle = \int_{\overline{\Omega}} \left(\int_{\Sigma} f(x, \xi, 0, 0) d\nu_1^{x,\infty}(\xi) \right) d\pi^\infty(x) \quad (4.86)$$

for every $f \in B_{\infty,1}^{hom}(\overline{\Omega} \times \mathbb{M}_D^{2 \times 2} \times \mathbb{R} \times \mathbb{R})$. We have to prove that $\pi^\infty = 0$. Let us consider the Lebesgue decomposition $\pi^\infty = \pi^{\infty,a} + \pi^{\infty,s}$.

We first prove that $\pi^{\infty,a} = 0$. We argue by contradiction. Assume that there exists a Borel set A , with $\mathcal{L}^2(A) > 0$ and $\lambda(A) = 0$, such that $\pi^{\infty,a}(x) > 0$ for every $x \in A$.

For every Borel set $A' \subset A$ let $f(x, \xi, \theta, \eta) := 1_{A'}(x)|\xi|$. Since ν_2^∞ is positive, by (4.85) and (4.86) we deduce that

$$\int_{A'} |\dot{p}^a| dx \geq \int_{A'} \pi^{\infty,a} dx$$

for every Borel set $A' \subset A$. Therefore, $|\dot{p}^a| > 0$ a.e. on A and there exists $h \in L^\infty(A)$, with $0 < h \leq 1$, such that $\pi^{\infty,a} = h|\dot{p}^a|$ on A .

Since $|\dot{p}^a| > 0$ a.e. on A , by (4.65) and Lemma 3.5 we have that $|\hat{z}| > 0$ a.e. on A . Hence, there exists $M > 0$ such that the set $A_M := \{x \in A : |\dot{p}^a(x)| < M, |\hat{z}(x)| > \frac{1}{M}\}$ has positive Lebesgue measure. Let us consider the function $f(x, \xi, \theta, \eta) := 1_{A_M}(x)(|\xi| - M^2|\theta|)^+$. Using (4.85), (4.86), the fact that $\pi^{\infty,a} = h|\dot{p}^a|$ on A , and the positivity of ν_2^∞ , we obtain

$$\int_{A_M} (|\dot{p}^a| - M^2|\hat{z}|)^+ dx \geq \int_{A_M} h|\dot{p}^a| dx,$$

which gives the contradiction, since the left-hand side vanishes, while the right-hand side is strictly positive. Therefore, $\pi^{\infty,a} = 0$.

It remains to prove that $\pi^{\infty,s} = 0$. We argue by contradiction. Assume that there exists a Borel set A with $\mathcal{L}^2(A) = 0$ and $\pi^{\infty,s}(A) > 0$.

For every Borel set $A' \subset A$ let $f(x, \xi, \theta, \eta) := 1_{A'}(x)|\xi|$. Since ν_2^∞ is positive, by (4.85) and (4.86) we deduce that

$$\int_{A'} |\dot{p}^\lambda| d\lambda \geq \pi^{\infty,s}(A')$$

for every Borel set $A' \subset A$. Therefore, there exists $h_\lambda \in L_\lambda^\infty(A)$, with $0 \leq h_\lambda \leq 1$, such that $\pi^{\infty,s} = h_\lambda |\dot{p}^\lambda| \lambda$ on A . Taking A smaller if needed, we may assume that $\lambda(A) > 0$, $h_\lambda > 0$ and $|\dot{p}^\lambda| > 0$ λ -a.e. on A . We can now argue as before with \mathcal{L}^2 replaced by λ and we obtain a contradiction, which implies $\pi^{\infty,s} = 0$.

This concludes the proof of the fact that $\nu_1^\infty = 0$ and, in turn, of (4.81) by (4.82).

By [4, Theorem 10.4] identity (4.81) yields

$$\text{Var}(\overline{\boldsymbol{\mu}}^Y; 0, t) = \int_0^t \langle \sqrt{|\xi|^2 + \theta^2}, \dot{\boldsymbol{\mu}}_t^Y(x, \xi, \theta, \eta) \rangle dt = 0$$

for every $t \in [0, +\infty)$. By Corollary 3.12 we deduce that

$$\begin{aligned} & \langle |\xi_1 - \eta \overline{p}_0(x)| + |\theta_1 - \eta \overline{z}_0(x)|, \overline{\boldsymbol{\mu}}_t^Y(x, \xi_1, \theta_1, \eta) \rangle = \\ & = \langle |\xi_1 - \xi_0| + |\theta_1 - \theta_0|, \overline{\boldsymbol{\mu}}_{0t}^Y(x, \xi_0, \theta_0, \xi_1, \theta_1, \eta) \rangle \leq \text{Var}(\overline{\boldsymbol{\mu}}^Y; 0, t) = 0, \end{aligned}$$

which easily implies the (4.69). \square

Remark 4.19. We remark that in the previous proof we could not deduce (4.69) from (4.67) simply by “integration” with respect to time. In fact, for a system $\boldsymbol{\mu}$ of generalized Young measures it is not true in general that the knowledge of $\dot{\boldsymbol{\mu}}_t$ and of $\boldsymbol{\mu}_0$ is enough to identify $\boldsymbol{\mu}_t$, as the following example shows.

Let $U := (0, 1)$ and for every $t \in [0, +\infty)$ let $\mathbf{p}(t) \in L^1(U)$ be the characteristic function $1_{(0,t)}$. We now consider the two systems $\boldsymbol{\mu}^1, \boldsymbol{\mu}^2 \in \text{SGY}([0, +\infty), U; \mathbb{R})$ defined by

$$\boldsymbol{\mu}_{t_1 \dots t_m}^1 := \delta_{(\mathbf{p}(t_1), \dots, \mathbf{p}(t_m))} \quad \text{and} \quad \boldsymbol{\mu}_{t_1 \dots t_m}^2 := \delta_{(0, \dots, 0)} + \omega_{(\mathbf{p}(t_1), \dots, \mathbf{p}(t_m))}^{\mathcal{L}^1}$$

for every finite sequence $0 \leq t_1 < t_2 < \dots < t_m$. Note that for every $0 \leq t < t_1$

$$\langle |\theta_1 - \theta|, \boldsymbol{\mu}_{tt_1}^1 \rangle = \langle |\theta_1 - \theta|, \boldsymbol{\mu}_{tt_1}^2 \rangle = \int_0^1 |\mathbf{p}(t_1) - \mathbf{p}(t)| dx = t_1 - t,$$

which shows that both $\boldsymbol{\mu}^1$ and $\boldsymbol{\mu}^2$ are absolutely continuous. Moreover, for every $f \in C^{\text{hom}}(U \times \mathbb{R} \times \mathbb{R})$ and every $0 \leq t < t_1$ we have

$$\begin{aligned} & \langle f(x, \frac{\theta_1 - \theta}{t_1 - t}, \eta), \boldsymbol{\mu}_{tt_1}^1(x, \theta, \theta_1, \eta) \rangle = \int_0^1 f(x, \frac{\mathbf{p}(t_1) - \mathbf{p}(t)}{t_1 - t}, 1) dx = \\ & = \int_0^t f(x, 0, 1) dx + \frac{1}{t_1 - t} \int_t^{t_1} f(x, 1, t_1 - t) dx + \int_{t_1}^1 f(x, 0, 1) dx \end{aligned}$$

and

$$\begin{aligned} & \langle f(x, \frac{\theta_1 - \theta}{t_1 - t}, \eta), \boldsymbol{\mu}_{tt_1}^2(x, \theta, \theta_1, \eta) \rangle = \int_0^1 f(x, 0, 1) dx + \int_0^1 f(x, \frac{\mathbf{p}(t_1) - \mathbf{p}(t)}{t_1 - t}, 0) dx \\ & = \int_0^1 f(x, 0, 1) dx + \frac{1}{t_1 - t} \int_t^{t_1} f(x, 1, 0) dx. \end{aligned}$$

It follows that

$$\begin{aligned} \lim_{t_1 \rightarrow t^+} \langle f(x, \frac{\theta_1 - \theta}{t_1 - t}, \eta), \boldsymbol{\mu}_{tt_1}^1(x, \theta, \theta_1, \eta) \rangle & = \lim_{t_1 \rightarrow t^+} \langle f(x, \frac{\theta_1 - \theta}{t_1 - t}, \eta), \boldsymbol{\mu}_{tt_1}^2(x, \theta, \theta_1, \eta) \rangle = \\ & = \int_0^1 f(x, 0, 1) dx + f(t, 1, 0), \end{aligned}$$

which yields

$$\dot{\boldsymbol{\mu}}_t^1 = \dot{\boldsymbol{\mu}}_t^2 = \delta_0 + \omega_1^{\delta_t}$$

for every $t \in [0, +\infty)$. We point out that, although $\overline{(\dot{\boldsymbol{\mu}}_t^1)}^Y = \delta_0$, the finite part of $\boldsymbol{\mu}_t^1$, which coincides with $\boldsymbol{\mu}_t^1$ itself, evolves in time.

5. AN EXAMPLE

In this section we assume that \mathbb{C} is isotropic, which implies that

$$\mathbb{C}\xi = 2\mu\xi_D + \kappa(\text{tr}\xi)I$$

for some constants $\mu > 0$ and $\kappa > 0$. We also assume that

$$\begin{aligned} K &:= \{(\sigma, \zeta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : |\sigma|^2 + \zeta^2 \leq 1\}, \\ V(\theta) &:= \frac{1}{2} - \frac{1}{2}\sqrt{1 + \theta^2}, \quad \Gamma_0 := \partial\Omega, \quad \Gamma_1 := \emptyset. \end{aligned} \quad (5.1)$$

Let us fix a constant $\theta_0 > 0$ and a 2×2 matrix ξ_0 with $\text{tr}\xi_0 = 0$. We assume that the symmetric part ξ_0^s of ξ_0 is different from 0. We will examine the globally stable quasistatic evolution corresponding to the boundary datum

$$\mathbf{w}(t, x) := t\xi_0 x,$$

and to the initial conditions

$$u_0(x) = 0, \quad e_0(x) = 0, \quad p_0(x) = 0, \quad z_0(x) = \theta_0.$$

Theorem 5.1. *Assume that \mathbb{C} , K , V , Γ_0 , Γ_1 , θ_0 , ξ_0 , ξ_0^s , \mathbf{w} , u_0 , e_0 , p_0 , and z_0 satisfy the conditions considered at the beginning of this section. Let*

$$t_0 := \frac{\sqrt{3}}{4\mu|\xi_0^s|}, \quad (5.2)$$

and let \mathbf{u} , \mathbf{e} , \mathbf{p} , \mathbf{z}_∞ be defined by

$$\begin{aligned} \mathbf{u}(t, x) &:= t\xi_0 x, & \mathbf{e}(t, x) &:= \alpha(t)\xi_0^s, \\ \mathbf{p}(t, x) &:= \beta(t)\xi_0^s, & \mathbf{z}_\infty(t, x) &:= \frac{1}{\sqrt{3}}|\mathbf{p}(t, x)|, \end{aligned}$$

where

$$\alpha(t) := \begin{cases} t & \text{for } t \in [0, t_0], \\ t_0 & \text{for } t \in [t_0, +\infty), \end{cases} \quad \beta(t) := \begin{cases} 0 & \text{for } t \in [0, t_0], \\ t - t_0 & \text{for } t \in [t_0, +\infty), \end{cases}$$

and let $\boldsymbol{\mu} \in \text{SGY}([0, +\infty), \bar{\Omega}; \mathbb{M}_D^{2 \times 2} \times \mathbb{R})$ be the system defined by

$$\begin{aligned} \boldsymbol{\mu}_{t_1 \dots t_m} &:= \delta_{((p_0, z_0), \dots, (p_0, z_0))} + \frac{1}{2}\omega_{((\mathbf{p}(t_1), \mathbf{z}_\infty(t_1)), \dots, (\mathbf{p}(t_m), \mathbf{z}_\infty(t_m)))} + \\ &\quad + \frac{1}{2}\omega_{((\mathbf{p}(t_1), -\mathbf{z}_\infty(t_1)), \dots, (\mathbf{p}(t_m), -\mathbf{z}_\infty(t_m)))} \end{aligned}$$

for every finite sequence t_1, \dots, t_m with $0 \leq t_1 < \dots < t_m$. Then $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu})$ is a globally stable quasistatic evolution of Young measures with boundary datum \mathbf{w} and initial condition $\boldsymbol{\mu}_0 = \delta_{(p_0, z_0)}$.

Proof. By (3.11) and (5.1) we have

$$K_{\text{eff}} = \{(\sigma, \zeta) \in \mathbb{M}_D^{2 \times 2} \times \mathbb{R} : |\sigma| \leq \frac{\sqrt{3}}{2}, |\zeta| \leq \sqrt{1 - |\sigma|^2} - \frac{1}{2}\}.$$

By (3.9) we have

$$H_{\text{eff}}(\xi, 0) = \frac{\sqrt{3}}{2}|\xi| \quad \text{for every } \xi \in \mathbb{M}_D^{2 \times 2}.$$

Thanks to (5.2) the function $(\mathbf{u}, \mathbf{e}, \mathbf{p})$ satisfies condition (b) of [3, Theorem 6.1] for the dissipation function $\xi \mapsto H_{\text{eff}}(\xi, 0)$, therefore it is a quasistatic evolution for the same dissipation function according to [3, Definition 4.2]. It follows, in particular, that

$$\mathcal{Q}(\mathbf{e}(t)) \leq \mathcal{Q}(\mathbf{e}(t) + \tilde{e}) + \mathcal{H}_{\text{eff}}(\tilde{p}, 0) \quad (5.3)$$

for every $t \in [0, +\infty)$ and every $(\tilde{u}, \tilde{e}, \tilde{p}) \in A_{\text{reg}}(0)$. Since H_{eff} is convex and even, we have $\mathcal{H}_{\text{eff}}(\tilde{p}, 0) \leq \mathcal{H}_{\text{eff}}(\tilde{p}, \tilde{z})$ for every $\tilde{z} \in L^1(\Omega)$. From (3.4) and (5.3) it follows that

$$\begin{aligned} \mathcal{Q}(\mathbf{e}(t)) &\leq \mathcal{Q}(\mathbf{e}(t) + \tilde{e}) + \mathcal{H}(\tilde{p}, \tilde{z}) + \mathcal{V}(\tilde{z} + z_0) - \mathcal{V}(z_0) = \\ &= \mathcal{Q}(\mathbf{e}(t) + \tilde{e}) + \mathcal{H}(\tilde{p}, \tilde{z}) + \langle \{V\}(\theta + \eta \tilde{z}(x), \eta) - \{V\}(\theta, \eta), \boldsymbol{\mu}_t(x, \xi, \theta, \eta) \rangle, \end{aligned}$$

which gives the global stability condition (ev1) of Definition 4.6.

Let us prove the energy balance (ev2). Since for $0 \leq t_1 < t_2$ we have

$$\langle \sqrt{|\xi_2 - \xi_1|^2 + |\theta_2 - \theta_1|^2}, \boldsymbol{\mu}_{t_1 t_2}(x, \xi_1, \theta_1, \xi_2, \theta_2, \eta) \rangle = \frac{2}{\sqrt{3}} \|\mathbf{p}(t_2) - \mathbf{p}(t_1)\|_1,$$

we deduce that

$$\mathcal{D}_H(\boldsymbol{\mu}; 0, T) = \frac{2}{\sqrt{3}} \int_0^T \|\dot{\mathbf{p}}(t)\|_1 dt.$$

Therefore, if $T \in (0, t_0]$ condition (ev2) is satisfied, since

$$\mathcal{Q}(e(T)) = \mu T^2 |\xi_0^s|^2 \mathcal{L}^2(\Omega), \quad \mathcal{Q}(e(0)) = 0, \quad \mathcal{D}_H(\boldsymbol{\mu}; 0, T) = 0,$$

$$\langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle = \mathcal{V}(z_0) = \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_0(x, \xi, \theta, \eta) \rangle,$$

$$\int_0^T \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt = \int_0^T 2\mu t |\xi_0^s|^2 \mathcal{L}^2(\Omega) dt = \mu T^2 |\xi_0^s|^2 \mathcal{L}^2(\Omega).$$

If $T \in [t_0, +\infty)$ condition (ev2) is satisfied, since

$$\mathcal{Q}(e(T)) = \mu t_0^2 |\xi_0^s|^2 \mathcal{L}^2(\Omega), \quad \mathcal{D}_H(\boldsymbol{\mu}; 0, T) = \frac{2}{\sqrt{3}} (T - t_0) |\xi_0^s| \mathcal{L}^2(\Omega),$$

$$\langle \{V\}(\theta, \eta), \boldsymbol{\mu}_T(x, \xi, \theta, \eta) \rangle = \mathcal{V}(z_0) + \mathcal{V}^\infty(z_\infty(T)) = \mathcal{V}(z_0) - \frac{1}{2\sqrt{3}} (T - t_0) |\xi_0^s| \mathcal{L}^2(\Omega),$$

$$\mathcal{Q}(e(0)) = 0, \quad \langle \{V\}(\theta, \eta), \boldsymbol{\mu}_0(x, \xi, \theta, \eta) \rangle = \mathcal{V}(z_0),$$

$$\begin{aligned} \int_0^T \langle \boldsymbol{\sigma}(t), E\dot{\mathbf{w}}(t) \rangle dt &= \int_0^{t_0} 2\mu t |\xi_0^s|^2 \mathcal{L}^2(\Omega) dt + \int_{t_0}^T 2\mu t_0 |\xi_0^s|^2 \mathcal{L}^2(\Omega) dt = \\ &= \mu t_0^2 |\xi_0^s|^2 \mathcal{L}^2(\Omega) + 2\mu (T - t_0) t_0 |\xi_0^s|^2 \mathcal{L}^2(\Omega) = \mu t_0^2 |\xi_0^s|^2 \mathcal{L}^2(\Omega) + \frac{\sqrt{3}}{2} (T - t_0) |\xi_0^s| \mathcal{L}^2(\Omega), \end{aligned}$$

where in the last equality we use (5.2).

It remains to prove that $(\mathbf{u}, \mathbf{e}, \boldsymbol{\mu}) \in AY([0, +\infty), \mathbf{w})$. Let us fix a finite sequence t_1, \dots, t_m with $0 \leq t_1 < \dots < t_m$. By an algebraic property of $\mathbb{M}_D^{2 \times 2}$ there exist $a, b \in \mathbb{R}^2$, with $|b| = 1$, and a skew symmetric 2×2 -matrix q such that $\xi_0^s = a \otimes b + q$. Therefore there exists a skew symmetric 2×2 -matrix ω such that $\xi_0 = a \odot b + \omega$.

For every $k \in \mathbb{N}$ and $i \in \mathbb{Z}$ we set

$$\begin{aligned} A_k^i &:= \{x \in \Omega : \frac{i}{k} < b \cdot x \leq \frac{i+1}{k} - \frac{1}{k^2}\}, \\ B_k^i &:= \{x \in \Omega : \frac{i+1}{k} - \frac{1}{k^2} < b \cdot x \leq \frac{i+1}{k} - \frac{1}{2k^2}\}, \\ C_k^i &:= \{x \in \Omega : \frac{i+1}{k} - \frac{1}{2k^2} < b \cdot x \leq \frac{i+1}{k}\}, \\ A_k &:= \bigcup_i A_k^i, \quad B_k := \bigcup_i B_k^i, \quad C_k := \bigcup_i C_k^i, \\ v_k(x) &:= \begin{cases} \frac{i}{k} a + qx & \text{if } x \in A_k^i, \\ (kb \cdot x - i - 1 + \frac{i+1}{k}) a + qx & \text{if } x \in B_k^i \cup C_k^i, \end{cases} \\ \theta_k(x) &:= \begin{cases} 0 & \text{if } x \in A_k^i, \\ \frac{k}{\sqrt{3}} |\xi_0^s| & \text{if } x \in B_k^i, \\ -\frac{k}{\sqrt{3}} |\xi_0^s| & \text{if } x \in C_k^i, \end{cases} \end{aligned}$$

so that $v_k \in W^{1, \infty}(\Omega; \mathbb{R}^2)$, $|v_k(x) - \xi_0^s x| \leq |a|/k$ on Ω , $E v_k(x) = 0$ on A_k , and $E v_k(x) = k \xi_0^s$ on $B_k \cup C_k$. We note that

$$1_{A_k} \rightharpoonup 1, \quad k 1_{B_k} \rightharpoonup \frac{1}{2}, \quad k 1_{C_k} \rightharpoonup \frac{1}{2} \quad \text{weakly}^* \text{ in } M_b(\Omega). \quad (5.4)$$

Let Ω_k be an increasing sequence of open sets, with union equal to Ω , such that $0 < \text{dist}(\Omega_k, \mathbb{R}^2 \setminus \Omega) < 2/\sqrt{k}$, and let $\varphi_k \in C_c^\infty(\Omega)$ be cut-off functions such that $\varphi_k = 1$ on Ω_k , $0 \leq \varphi_k \leq 1$ on $\Omega \setminus \Omega_k$, and $|\nabla \varphi_k| \leq \sqrt{k}$ on Ω .

Let us define

$$\begin{aligned}\mathbf{u}_k(t, x) &:= \varphi_k(x)(\alpha(t)\xi_0^s x + \beta(t)v_k(x) + t\omega x) + (1 - \varphi_k(x))t\xi_0 x, \\ \mathbf{p}_k(t, x) &:= \varphi_k(x)\beta(t)Ev_k(x) + (1 - \varphi_k(x))t\xi_0^s, \\ \mathbf{e}_k(t, x) &:= E\mathbf{u}_k(t, x) - \mathbf{p}_k(t, x) = \varphi_k(x)\alpha(t)\xi_0^s + \beta(t)\nabla\varphi_k(x) \odot (v_k(x) - \xi_0^s x), \\ \mathbf{z}_k(t, x) &:= \theta_0 + \beta(t)\theta_k(x).\end{aligned}$$

Then $\mathbf{u}_k(t, x) \rightarrow t\xi_0 x$ and $\mathbf{e}_k(t, x) \rightarrow \alpha(t)\xi_0^s$ uniformly on Ω for every $t \in [0, +\infty)$. As for p_k and z_k , for every $f \in C^{hom}(\overline{\Omega} \times (\mathbb{M}_D^{2 \times 2} \times \mathbb{R})^m \times \mathbb{R})$ we have

$$\begin{aligned}\int_{\Omega} f(x, p_k(t_1, x), z_k(t_1, x), \dots, p_k(t_m, x), z_k(t_m, x), 1) dx &= \\ &= \int_{A_k \cap \Omega_k} f(x, 0, \theta_0, \dots, 0, \theta_0, 1) dx + \\ &+ \int_{B_k \cap \Omega_k} f(x, \beta(t_1)k\xi_0^s, \theta_0 + \beta(t_1)k\frac{|\xi_0^s|}{\sqrt{3}}, \dots, \beta(t_m)k\xi_0^s, \theta_0 + \beta(t_m)k\frac{|\xi_0^s|}{\sqrt{3}}, 1) dx + \\ &+ \int_{C_k \cap \Omega_k} f(x, \beta(t_1)k\xi_0^s, \theta_0 - \beta(t_1)k\frac{|\xi_0^s|}{\sqrt{3}}, \dots, \beta(t_m)k\xi_0^s, \theta_0 - \beta(t_m)k\frac{|\xi_0^s|}{\sqrt{3}}, 1) dx + \mathcal{R}_k = \\ &= \int_{A_k \cap \Omega_k} f(x, 0, \theta_0, \dots, 0, \theta_0, 1) dx + \\ &+ k \int_{B_k \cap \Omega_k} f(x, \beta(t_1)\xi_0^s, \frac{\theta_0}{k} + \beta(t_1)\frac{|\xi_0^s|}{\sqrt{3}}, \dots, \beta(t_m)\xi_0^s, \frac{\theta_0}{k} + \beta(t_m)\frac{|\xi_0^s|}{\sqrt{3}}, \frac{1}{k}) dx \\ &+ k \int_{C_k \cap \Omega_k} f(x, \beta(t_1)\xi_0^s, \frac{\theta_0}{k} - \beta(t_1)\frac{|\xi_0^s|}{\sqrt{3}}, \dots, \beta(t_m)\xi_0^s, \frac{\theta_0}{k} - \beta(t_m)\frac{|\xi_0^s|}{\sqrt{3}}, \frac{1}{k}) dx + \mathcal{R}_k,\end{aligned}$$

where the remainder \mathcal{R}_k satisfies the estimate

$$|\mathcal{R}_k| \leq c \|f\|_{hom} (\mathcal{L}^2(\Omega \setminus \Omega_k) + k\mathcal{L}^2(B_k \cap (\Omega \setminus \Omega_k)) + k\mathcal{L}^2(C_k \cap (\Omega \setminus \Omega_k))).$$

By (5.4) it follows that

$$\begin{aligned}\lim_{k \rightarrow \infty} \int_{\Omega} f(x, p_k(t_1, x), z_k(t_1, x), \dots, p_k(t_m, x), z_k(t_m, x), 1) dx &= \\ &= \int_{\Omega} f(x, 0, \theta_0, \dots, 0, \theta_0, 1) dx + \\ &+ \frac{1}{2} \int_{\Omega} f(x, \beta(t_1)\xi_0^s, \beta(t_1)\frac{|\xi_0^s|}{\sqrt{3}}, \dots, \beta(t_m)\xi_0^s, \beta(t_m)\frac{|\xi_0^s|}{\sqrt{3}}, 0) dx + \\ &+ \frac{1}{2} \int_{\Omega} f(x, \beta(t_1)\xi_0^s, \beta(t_1)\frac{|\xi_0^s|}{\sqrt{3}}, \dots, \beta(t_m)\xi_0^s, \beta(t_m)\frac{|\xi_0^s|}{\sqrt{3}}, 0) dx.\end{aligned}$$

This proves (4.1) and concludes the proof of the theorem. \square

Remark 5.2. For $t > t_0$ the globally stable quasistatic evolution described in Theorem 5.1 is completely different from the approximable quasistatic evolution presented in [5, Section 7]. The globally stable quasistatic evolution contains the terms

$$\frac{1}{2}\omega_{((\mathbf{p}(t_1), \mathbf{z}_{\infty}(t_1)), \dots, (\mathbf{p}(t_m), \mathbf{z}_{\infty}(t_m)))} + \frac{1}{2}\omega_{((\mathbf{p}(t_1), -\mathbf{z}_{\infty}(t_1)), \dots, (\mathbf{p}(t_m), -\mathbf{z}_{\infty}(t_m)))},$$

which describe oscillations of \mathbf{p} and \mathbf{z} with infinite amplitude and frequency, while the approximable quasistatic evolution is given by ordinary functions, without concentration or oscillation effects. Moreover the stress $\boldsymbol{\sigma}$ of the globally stable quasistatic evolution satisfies

$$\boldsymbol{\sigma}(t) = \frac{\sqrt{3}}{2} \frac{\xi_0^s}{|\xi_0^s|} \quad \text{for every } t > t_0,$$

while the stress σ of the approximable quasistatic evolution satisfies

$$\sigma(t) \rightarrow \frac{\sqrt{3}}{2} \frac{\xi_0^s}{|\xi_0^s|} \quad \text{as } t \rightarrow +\infty,$$

but $|\sigma(t)| > \frac{\sqrt{3}}{2}$ for every $t > t_0$, as shown in [5, Remark 7.7].

Acknowledgments. This work is part of the Project ‘‘Calculus of Variations’’ 2006, supported by the Italian Ministry of University and Research, and of the research project ‘‘Mathematical Challenges in Nanomechanics’’ supported by INdAM.

REFERENCES

[1] Anzellotti G., Giaquinta M.: On the existence of the field of stresses and displacements for an elasto-perfectly plastic body in static equilibrium. *J. Math. Pures Appl.* **61** (1982), 219-244.
 [2] Brezis H.: Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert. North-Holland, Amsterdam-London; American Elsevier, New York, 1973.
 [3] Dal Maso G., DeSimone A., Mora M.G.: Quasistatic evolution problems for linearly elastic - perfectly plastic materials. *Arch. Ration. Mech. Anal.* **180** (2006), 237-291.
 [4] Dal Maso G., DeSimone A., Mora M.G., Morini M.: Time-dependent systems of generalized Young measures. *Netw. Heterog. Media* **2** (2007), 1-36.
 [5] Dal Maso G., DeSimone A., Mora M.G., Morini M.: A vanishing viscosity approach to quasistatic evolution in plasticity with softening. *Arch. Ration. Mech. Anal.*, to appear.
 [6] Goffman C., Serrin J.: Sublinear functions of measures and variational integrals. *Duke Math. J.* **31** (1964), 159-178.
 [7] Han W., Reddy B.D.: Plasticity. Mathematical theory and numerical analysis. Springer Verlag, Berlin, 1999.
 [8] Hill R.: The mathematical theory of plasticity. Clarendon Press, Oxford, 1950.
 [9] Kachanov L.M.: Fundamentals of the theory of plasticity, Dover, 2004.
 [10] Lubliner J.: Plasticity theory. Macmillan Publishing Company, New York, 1990.
 [11] Luckhaus S., Modica L.: The Gibbs-Thompson relation within the gradient theory of phase transitions. *Arch. Ration. Mech. Anal.* **107** (1989), 71-83.
 [12] Martin J.B.: Plasticity. Fundamentals and general results. MIT Press, Cambridge, 1975.
 [13] Meyers N.G., Serrin J.: H=W. *Proc. Nat. Acad. Sci. U.S.A.* **51** (1964), 1055-1056.
 [14] Mielke A.: Evolution of rate-independent systems. In: Evolutionary equations. Vol. II. Edited by C. M. Dafermos and E. Feireisl, 461-559, Handbook of Differential Equations. Elsevier/North-Holland, Amsterdam, 2005.
 [15] Mielke A., Ortiz M.: A class of minimum principles for characterizing the trajectories and the relaxation of dissipative systems. *ESAIM Control Optim. Calc. Var.*, to appear.
 [16] Mielke A., Roubíček T., Stefanelli U.: Γ -limits and relaxations for rate-independent evolutionary problems. *Calc. Var. Partial Differential Equations*, to appear.
 [17] Reshetnyak Yu.G.: Weak convergence of completely additive vector functions on a set. *Siberian Math. J.* **9** (1968), 1039-1045.
 [18] Rockafellar R.T.: Convex Analysis. Princeton University Press, Princeton, 1970.
 [19] Temam R.: Mathematical problems in plasticity. Gauthier-Villars, Paris, 1985. Translation of Problèmes mathématiques en plasticité. Gauthier-Villars, Paris, 1983.

(G. Dal Maso, A. DeSimone, M.G. Mora, and M. Morini) SISSA, VIA BEIRUT 2-4, 34014 TRIESTE, ITALY
E-mail address, Gianni Dal Maso: dalmaso@sissa.it
E-mail address, Antonio DeSimone: desimone@sissa.it
E-mail address, Maria Giovanna Mora: mora@sissa.it
E-mail address, Massimiliano Morini: morini@sissa.it