

COMPACT MANIFOLDS WITH UNBOUNDED NILPOTENT FUNDAMENTAL GROUPS AND POSITIVE RICCI CURVATURE

ELIA BRUÈ, AARON NABER, AND DANIELE SEMOLA

ABSTRACT. It follows from the work of Kapovitch and Wilking that a closed manifold with nonnegative Ricci curvature has an almost nilpotent fundamental group. Leftover questions and conjectures, see [27] and [33], have asked if in this context the fundamental group is actually uniformly almost abelian.

The main goal of this work is to construct examples (M_k^{10}, g_k) with uniformly positive Ricci curvature $\text{Ric}_{g_k} \geq 9$ whose fundamental groups cannot be uniformly virtually abelian.

1. INTRODUCTION

A consequence of the work of Kapovitch and Wilking in [27] for a closed manifold (M^n, g) with $\text{Ric} \geq 0$ is that the fundamental group $\pi_1(M^n)$ has a nilpotent subgroup $N \leq \pi_1(M)$ of uniformly bounded index $[N, \pi_1(M)] \leq C(n)$. An important open question from their work, see [27, page 48], is whether this nilpotent subgroup N can be taken to be abelian. More generally, it is asked whether for a space with $\text{Ric} \geq -(n-1)$ the torsion of the local fundamental group¹ can be taken to be uniformly almost abelian.

The question of Kapovitch and Wilking, and conjectures of Fukaya and Yamaguchi, have been quantitatively refined into conjecture by Pan and Rong, see [33, Conjecture 2.22] and [35, Conjecture 12]; namely, it is conjectured that compact spaces with nonnegative Ricci curvature have fundamental groups which are uniformly almost abelian. The results of [29] and [35] are able to answer these conjectures in the affirmative under the additional hypothesis that the manifolds are noncollapsing; more precisely, the index of the abelian subgroup is bounded in terms of a constant depending on the volume of unit balls in the universal cover.

The main goal of this paper is to answer these questions in the general case with the following construction:

Theorem 1.1. *There exists a sequence of smooth 10-dimensional Riemannian manifolds (M_k^{10}, g_k) such that:*

- (1) $\text{Ric}_{g_k} \geq 9$, and consequently $\text{diam}(M_k) \leq \text{diam}(\widetilde{M}_k) \leq \pi$ with $\widetilde{M}_k$ the universal cover of M_k ;
- (2) The fundamental group $\pi_1(M_k)$ is a degree-two extension of the finite Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$ for each $k \in \mathbb{N}$.

The interest of Theorem 1.1 stems from the fact that the groups $H_3(\mathbb{Z}/k\mathbb{Z})$, see Section 2.2, are not *uniformly virtually abelian* with respect to k . That is, the index of every abelian subgroup $A_k \leq H_3(\mathbb{Z}/k\mathbb{Z})$ diverges as $k \rightarrow \infty$. In particular, this answers the question of Kapovitch and Wilking and conjectures of [33, Conjecture 2.22] and [35, Conjecture 12] in the negative.

¹ $\text{Im}(\pi_1(B_{\epsilon(n)}(p)) \rightarrow \pi_1(B_1(p)))$

As such, Theorem 1.1 also refutes the natural counterpart for Ricci curvature of some long-standing conjectures of Fukaya and Yamaguchi for fundamental groups of manifolds with $\sec \geq 0$. In the context of noncompact manifolds with torsion free fundamental group, see also the excellent example of Wei [37]. One is still left with the original conjecture of Fukaya and Yamaguchi in the context of nonnegative sectional curvature, which is very much open. Additionally, if we compare to [26], we see that the *torsion in the center* conjecture for spaces with nonnegative sectional curvature must fail under the weaker nonnegative Ricci assumption.

Theorem 1.1 is also connected to another series of questions which remain open. In [5] and [6] examples of manifolds with nonnegative Ricci curvature and infinitely generated fundamental group were constructed. The infinite generation was however abelian in nature. For instance, it is unknown if there always exists a normal abelian subgroup $A \leq \pi_1(M)$ such that the quotient $\pi_1(M)/A$ is finitely generated.

The main step in the construction of the spaces M_k^{10} in Theorem 1.1 is based on the construction of simply connected four manifolds M_k^4 with effective isometric actions by $H_3(\mathbb{Z}/k\mathbb{Z})$. These four manifolds even have uniformly bounded Ricci curvature. However, the actions of $H_3(\mathbb{Z}/k\mathbb{Z})$ on the M_k^4 are not free. We will construct the universal covers $\widetilde{M}_k^{10}$ from M_k^4 by looking at the spin bundles of M_k^4 and lifting the actions of $H_3(\mathbb{Z}/k\mathbb{Z})$ to *free* actions. In particular, this also explains the $\mathbb{Z}/2\mathbb{Z}$ extension of Theorem 1.1.

Our precise construction is the following:

Theorem 1.2. *For every $k \in \mathbb{N}$ there exists a closed, simply connected 4-manifold (M_k^4, g_k) with $\text{diam}(M_k) \leq \pi$ and $0 < \text{Ric}_{g_k} \leq 3$, admitting an effective isometric action of the finite Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$.*

Each M_k^4 above is diffeomorphic to the connected sum of $(k-1)$ copies of $S^2 \times S^2$. We address the reader to Section 3 for more details on this and on the geometry of the examples.

Remark 1.3. *Theorem 1.2 is optimal in at least two senses:*

- i) *The family of finite groups Γ that act smoothly and effectively on a sphere S^n for $n = 2, 3$ is uniformly virtually abelian. See for instance [14]. In particular, there are no three dimensional examples as above.*
- ii) *In dimension four the Fukaya-Yamaguchi conjecture has been proved in [7]. More generally, by [30, Theorem 1.2] and [19], there is no family of four dimensional examples as above with positive sectional curvature. See [7] for the argument.*

Broader Mathematical Context. We recall that the fundamental group of every closed manifold with $\text{Ric} > 0$ is finite, as a consequence of Myers' theorem. It is not the case that one has uniform finiteness, the easiest examples being the Lens spaces in dimension three. Similarly, one also knows that the fundamental group of every closed manifold with $\text{Ric} \geq 0$ is virtually abelian, as a corollary of the Cheeger-Gromoll splitting theorem, see [8, Theorem 3].

On the other hand, in the noncompact case Wei constructed a family of complete (M^n, g) with $\text{Ric} \geq 0$ and for which π_1 is a nilpotent lattice [37]. Wilking refined Wei's construction more recently in [38] and proved that any finitely generated virtually nilpotent group is the fundamental group of some complete (noncompact) (M^n, g) with $\text{Ric} \geq 0$, for some $n \in \mathbb{N}$.

For manifolds (M^n, g) with nonnegative sectional curvature, fundamental groups are virtually abelian without any compactness assumption, by the Cheeger-Gromoll soul theorem [9] (combined again with the splitting theorem). Fukaya and Yamaguchi conjectured in [16] that the fundamental group of any (M^n, g) with $\sec \geq 0$ should have an abelian subgroup with index bounded by a dimensional constant $C(n)$. Their conjecture remains open.

Kapovitch, Petrunin, and Tuschmann proved that fundamental groups of almost nonnegatively curved manifolds in any fixed dimension are uniformly virtually nilpotent in [25] after some earlier progress due to Fukaya and Yamaguchi in [16]. They conjectured that one should be able to arrange the finite-index nilpotent subgroup to have torsion contained in the center, see [25, Main Conjecture 6.1.2]. If the “Torsion in the center” conjecture holds, then the Fukaya-Yamaguchi conjecture would be true as well, as observed by Wilking, see [25, Section 6]. We address the reader to [34, 29, 26] for some partial results in the direction of the Fukaya-Yamaguchi conjecture, without the aim of being complete.

Acknowledgments. Part of this work was conducted while D.S. was a Hermann Weyl Instructor. He is grateful to the FIM-Institute for Mathematical Research of ETH Zürich for the support and the excellent working conditions.

2. PRELIMINARIES

2.1. Heisenberg Groups. The 3-dimensional Heisenberg group over a commutative ring $(A, +, \cdot)$ is defined as

$$H_3(A) := \left\{ \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} : a, b, c \in A \right\} < \mathrm{GL}_3(A). \quad (1)$$

In the case that $(A, +)$ is generated by the identity $1 \in A$, we have that $H_3(A)$ is generated as a group by the two elements

$$X := \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad Y := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}. \quad (2)$$

If we set

$$Z := \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3)$$

then we have the relations

$$XYX^{-1}Y^{-1} = Z, \quad XZ = ZX, \quad YZ = ZY. \quad (4)$$

We will mainly be interested in the cases $A = \mathbb{R}$, $A = \mathbb{Z}$, and $A = \mathbb{Z}/k\mathbb{Z}$ with $k \in \mathbb{N}$.

2.2. Heisenberg Nilmanifolds and Group Actions. Fix $k \in \mathbb{Z}$. We consider the free action of $\mathbb{Z}^2$ on $\mathbb{R}^2 \times S^1$ defined by

$$(a, b) \cdot (x, y, z) := (x + a, y + b, e^{-2\pi i k a y} z), \quad (5)$$

for every $(a, b) \in \mathbb{Z}^2$ and $(x, y, z) \in \mathbb{R}^2 \times S^1 \subseteq \mathbb{R}^2 \times \mathbb{C}$. The quotient Nil_k^3 is the so-called *Heisenberg nilmanifold of degree $k \in \mathbb{Z}$* . It admits a free, smooth S^1 -action defined by

$$\theta \cdot [x, y, z] = [x, y, e^{i\theta} z], \quad \theta \in S^1, \quad (6)$$

where $[x, y, z]$ denotes the equivalence class of $(x, y, z) \in \mathbb{R}^2 \times S^1$. This S^1 -action endows every Heisenberg nilmanifold Nil_k^3 with the structure of a principal S^1 -bundle over T^2 , with projection map $\pi_k : \text{Nil}_k^3 \rightarrow T^2$ induced by the natural map $\mathbb{R}^2 \times S^1 \ni (x, y, z) \mapsto (x, y) \in \mathbb{R}^2$.

Assume that $k > 0$ for notational simplicity. A similar construction applies for $k < 0$. The nilmanifold Nil_k^3 admits a free effective action of the finite Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$, whose generators act as

$$\begin{aligned} X \cdot [x, y, z] &:= [x + 1/k, y, e^{-2\pi i y} z], \\ Y \cdot [x, y, z] &:= [x, y + 1/k, z]. \end{aligned} \tag{7}$$

The commutator of the X and Y action gives the action by Z , which generates a cyclic group $\mathbb{Z}/k\mathbb{Z}$ acting by rotation on the S^1 -fibers of Nil_k^3 :

$$Z \cdot [x, y, z] := [x, y, e^{-2\pi i/k} z]. \tag{8}$$

Note that the actions of X and Y are lifted from the torus action

$$\begin{aligned} X \cdot [x, y] &:= [x + 1/k, y], \\ Y \cdot [x, y] &:= [x, y + 1/k], \end{aligned} \tag{9}$$

where $[x, y] \in T^2 = \mathbb{R}^2/\mathbb{Z}^2$ denotes the equivalence class of $(x, y) \in \mathbb{R}^2$.

There are several equivalent ways to build the action of $H_3(\mathbb{Z}/k\mathbb{Z})$ on a nilmanifold Nil_k^3 . The construction we have followed here has been exploited in the past, for instance in [39] (resp. [12]) to construct an example of an irreducible complex surface (resp. smooth, closed 4-manifold) such that the finite subgroups of the group of birational transformations (resp. diffeomorphisms) are not uniformly virtually abelian. See also the more recent [13, Section 2].

3. STRUCTURE OF M_k^4

In this section, we outline the diffeomorphic and metric structure of the manifolds (M_k^4, g_k) built in Theorem 1.2. We will go about the construction carefully and rigorously in the next sections.

3.1. Sketch of Construction. Our construction of M_k^4 is based on a variant of the Gibbons-Hawking ansatz (see for instance [17, 1, 28, 21]) with base space the three-sphere S^3 . We consider two distinct circle fibers F_0 and F_1 of the Hopf fibration $S^3 \rightarrow S^2$, viewed as the fibers corresponding to the north and south poles of S^2 . We remove k points from each fiber, in a way so that discrete Hopf rotation by angle $2\pi/k$ leaves the set invariant. There is a $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z} \subseteq T^2$ action on S^3 , which is a subaction of the Clifford action, which maps this removed set to itself. See Section 4 for more on this.

The next step is to construct the unique (up to orientation) S^1 -bundle whose Chern class is invariant under this $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ action on S^3 , and for which when restricted to each small 2-sphere enclosing one of the removed points is the Hopf bundle. The resulting open four manifold can be smoothly compactified by adding $2k$ points. The Riemannian metric g_k is defined as (a small conformal perturbation of) the Gibbons-Hawking type metric (22), constructed using a Green's function with positive poles corresponding to the points removed from F_0 and negative poles corresponding to the points removed from F_1 , see (17). A key distinction from the classical Gibbons-Hawking construction is the behavior of the Green's function, which changes sign: it is positive near the fiber F_0 and negative near F_1 . This causes a great deal of delicacy and care in the region between the fibers. We will

show this $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ action on S^3 lifts to an (isometric) action by the Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$ on M_k^4 , whose isotropy points are the $2k$ removed points.

Let us now discuss several viewpoints from which to understand the geometry of our M_k^4 :

3.2. M_k^4 as a Singular S^1 Bundle over S^3 . The simply connected manifold (M_k^4, g_k) admits an isometric S^1 -action with $2k$ fixed points corresponding to the points that we removed from the Hopf fibers F_0 and F_1 . The quotient space of the action is naturally homeomorphic to S^3 , essentially by construction, and we shall denote $\pi : M_k^4 \rightarrow S^3$ to be the natural projection. Smooth S^1 actions with finitely many fixed points on closed simply connected 4-manifolds are completely classified up to equivariant diffeomorphism. In particular, by [11], we deduce that M_k^4 is diffeomorphic to the connected sum of $k-1$ copies of $S^2 \times S^2$. Hence, M_k^4 is spin and we can compute the homology as $H_1(M_k^4, \mathbb{Z}) = H_3(M_k^4, \mathbb{Z}) = 0$, and $H_2(M_k^4, \mathbb{Z}) \cong \mathbb{Z}^{2k-2}$.

3.3. M_k^4 as a Singular T^2 Bundle Over S^2 . The map $\pi_{T^2} : M_k^4 \rightarrow S^2$, obtained by composing $\pi : M_k^4 \rightarrow S^3$ and the Hopf map $\pi_{\text{Hopf}} : S^3 \rightarrow S^2$, is a smooth torus fibration away from the preimages of the north and south poles in S^2 . For every $x \in S^2$ different from the north and south pole, the fiber $\pi_{T^2}^{-1}(x)$ is diffeomorphic to a two-torus. On the other hand, the preimages of the south and north poles are homeomorphic to singular Kodaira fibers of type I_k . Topologically, the latter can be viewed as a chain of k spheres $S_1^2, S_2^2, \dots, S_k^2$ with the property that S_i^2 intersects S_{i+1}^2 at exactly one point, forming a loop (where S_{k+1}^2 is identified with S_1^2).

The reader is addressed to [21] and [10] for more on the appearance of singular Kodaira fibers through constructions based on the Gibbons-Hawking ansatz.

3.4. M_k^4 as a Singular Nil_k^3 Bundle Over I . The map $\pi_{\text{Nil}_k^3} : M_k^4 \rightarrow [0, 1]$, obtained by composing $\pi : M_k^4 \rightarrow S^3$ and $S^3 \ni (z_1, z_2) \mapsto |z_1| \in [0, 1]$, is a smooth fibration away from the preimages of 0 and 1. The (generic) fiber is diffeomorphic to the Heisenberg nilmanifold Nil_k^3 of degree k . The preimages of the boundary points are I_k fibers. We refer the reader to [23] for further details on the emergence of fibrations with nilmanifold fibers via constructions based on the Gibbons-Hawking ansatz. Additionally, we point to the earlier works [36] and [22] for the construction of Calabi-Yau metrics on manifolds that locally fiber as nilmanifolds over intervals.

The global action by the finite Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$ preserves the fibers of $\pi_{\text{Nil}_k^3}$. On the regular fibers this action matches the one described in Section 2.2. On the two singular I_k fibers, the two generators of $H_3(\mathbb{Z}/k\mathbb{Z})$ act in the following ways:

- (1) *Cycling of the spheres:* The spheres $S_1^2, \dots, S_k^2$ are permuted cyclically, with the action $S_\ell^2 \rightarrow S_{\ell+1}^2$ for $\ell = 1, \dots, k-1$, and $S_k^2 \rightarrow S_1^2$.
- (2) *Asynchronous rotation:* The sphere S_ℓ^2 is rotated by an angle $2\pi\ell/k$ (with respect to the axis connecting the touching points with the two neighbouring 2-spheres) for each $\ell = 1, \dots, k$.

3.5. Doubling of a Neighborhood of the I_k Fiber. Let $\Sigma \subset S^3$ be the Clifford torus. As we observe in the proof of Lemma 4.4 below, the preimage $\pi^{-1}(\Sigma) \subset M_k^4$ is diffeomorphic to Nil_k^3 . It is not hard to see that the latter disconnects M_k^4 into two open regions, each diffeomorphic to a neighborhood of an I_k fiber.

Thus, M_k^4 can be obtained by gluing together these two neighbourhoods of the I_k fiber along their boundaries, which are diffeomorphic to Nil_k^3 . Importantly, the gluing map is an equivariant diffeomorphism of Nil_k^3 with respect to its S^1 and $H_3(\mathbb{Z}/k\mathbb{Z})$ symmetries. Moreover the gluing map switches the sign of the Chern

class on the base two-torus. This is consistent with the fact that M_k^4 is simply connected, although the two neighbourhoods of the I_k fibers deformation retract onto the singular fibers and, as such, have fundamental groups isomorphic to $\mathbb{Z}$.

4. PROOF OF THEOREM 1.2

The careful construction of M_k^4 begins with the three sphere S^3 . Let us view S^3 as the unit sphere in $\mathbb{R}^4 = \mathbb{C}^2$. We consider the fibers $F_0, F_1 \subset S^3$ of the Hopf fibration defined by

$$F_0 := \{(z_1, 0) : |z_1| = 1\}, \quad F_1 := \{(0, z_2) : |z_2| = 1\}. \quad (10)$$

Let $k \in \mathbb{N}$ be fixed and $\mathcal{B}_k$ be the incomplete manifold obtained by removing the k -th roots of unity from the Hopf fibers F_0 and F_1 . Namely, we set

$$\mathcal{B}_k := S^3 \setminus \{(e^{2\pi i \ell/k}, 0), (0, e^{2\pi i \ell'/k}) : 0 \leq \ell, \ell' \leq k-1\}. \quad (11)$$

Note that there is a natural action of $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ on $\mathcal{B}_k$ given by

$$(\ell, \ell') \cdot (z_1, z_2) := (e^{2\pi i \ell/k} z_1, e^{2\pi i \ell'/k} z_2). \quad (12)$$

The action is effective and isometric with respect to the restriction of the standard metric of S^3 to $\mathcal{B}_k$.

We note that $H_2(\mathcal{B}_k, \mathbb{Z}) \simeq \mathbb{Z}^{2k-1}$ is generated by the homology classes of small 2-spheres enclosing the points $p_\ell^0 := (e^{2\pi i \ell/k}, 0)$ and $p_{\ell'}^1 := (0, e^{2\pi i \ell'/k})$. We enumerate them

$$S_\ell^\alpha = \partial B_r(p_\ell^\alpha), \quad (13)$$

with $\alpha \in \{0, 1\}$, $\ell \in \{0, \dots, k-1\}$ and $r > 0$ sufficiently small chosen sufficiently small so that the collection of balls $B_r(p_\ell^\alpha)$ is disjoint and each ball is diffeomorphic to a three ball.

Let us now define the integral cohomology class $[\omega] \in H^2(\mathcal{B}_k, \mathbb{Z}) \simeq \mathbb{Z}^{2k-1}$ by the condition

$$\int_{S_\ell^\alpha} \omega = (-1)^\alpha 2\pi. \quad (14)$$

Observe that this class is invariant under the $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ action on $\mathcal{B}_k$.

Let us now consider the unique principal S^1 bundle $\tilde{\pi} : \mathcal{N}_k \xrightarrow{S^1} \mathcal{B}_k$ whose Chern class is $[\omega]$. Using (14), we have that $\tilde{\pi}^{-1}(B_r(p_\ell^\alpha) \setminus \{p_\ell^\alpha\})$ is diffeomorphic to a punctured 4-ball for each p_ℓ^α . Hence $\mathcal{N}_k$ can be “completed”

$$M_k^4 \equiv \mathcal{N}_k \cup \{\tilde{p}_\ell^\alpha\}, \quad (15)$$

to a closed 4-manifold M_k^4 by adding $2k$ points. Moreover, this completion gives rise to a smooth extension

$$\pi : M_k^4 \rightarrow S^3, \quad (16)$$

of $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$. The S^1 action on $\mathcal{N}_k$ extends smoothly to M_k with the added points as fixed points. See Section 4.1 for coordinate computations and also [1, 28], [20, Section 2] for analogous arguments.

Note that $[\omega]$ is a primitive class. Since $\mathcal{B}_k$ is simply connected we have that $\mathcal{N}_k$ is simply connected, and hence M_k is simply connected as well.

We are going to define a metric on M_k^4 through a variant of the so-called Gibbons-Hawking ansatz, originally introduced in [17].

Let $u : S^3 \rightarrow [-\infty, +\infty]$ be a solution of

$$-\Delta_{S^3} u = 2\pi \sum_{\ell=0}^{k-1} (\delta_{(e^{2\pi i \ell/k}, 0)} - \delta_{(0, e^{2\pi i \ell/k})}) = 2\pi \sum_{\ell=0}^{k-1} (\delta_{p_\ell^0} - \delta_{p_\ell^1}), \quad (17)$$

where S^3 is endowed with the round metric of radius one. As the right hand side integrates to zero we have that such a function u exists, is smooth away from $\{p_\ell^\alpha\}$, and is unique up to a constant. We normalize it in such a way that it is odd under the involution $\iota : S^3 \rightarrow S^3$, $\iota(z_1, z_2) := (z_2, z_1)$. That is we want the condition $u(z_2, z_1) = -u(z_1, z_2)$. Notice that $u : \mathcal{B}_k \rightarrow \mathbb{R}$ is smooth and harmonic. We can define the two-form

$$\omega = *du, \quad (18)$$

where $*$ denotes the Hodge $*$ operator on S^3 . Note that $d\omega = 0$ on $\mathcal{B}_k$ follows because u is harmonic on $\mathcal{B}_k$, and hence ω defines a 2-cohomology class on $\mathcal{B}_k$.

Lemma 4.1. *The cohomology class $[\omega]$ satisfies (14), and thus represents the Chern class of the principal S^1 bundle $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$ in de Rham cohomology.*

Proof. Clearly, u is invariant under the $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ -action introduced in (12). Hence, $\omega = *du$ is also invariant with respect to this action.

Let $B_r(p_\ell^\alpha)$ be a small three-ball centered at one of the removed points p_ℓ^α with $S_\ell^\alpha = \partial B_r^3(p_\ell^\alpha)$. Observe that

$$d\omega = d * du = *(d * du) = *\Delta u. \quad (19)$$

Hence, by (a distributional formulation of) Stokes' theorem, we get

$$\int_{S_\ell^\alpha} \omega = \int_{B_r(p_\ell^\alpha)} d\omega = \int_{B_r(p_\ell^\alpha)} \Delta u = (-1)^\alpha 2\pi, \quad (20)$$

as claimed. $\square$

Thanks to Lemma 4.1, there exists a connection on the principal S^1 -bundle $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$ with curvature form ω . Let $\theta \in \Omega^1(\mathcal{N}_k)$ be a principal connection 1-form corresponding to this connection. In particular, $d\theta = \tilde{\pi}^*\omega$. While such a connection 1-form is not unique, its uniqueness is ensured up to gauge equivalence as the base space is simply connected.

Let us now consider the function

$$V(x) := \frac{1}{4\pi} \log(e^{4\pi x} + e^{-4\pi x} + 2). \quad (21)$$

We introduce the Riemannian metric h_k on $\mathcal{N}_k$ by defining

$$h_k := V_\Lambda(u_k) \tilde{\pi}^* g_{S^3} + \frac{1}{k^2 \Lambda^2 V_\Lambda(u_k)} \theta \otimes \theta, \quad (22)$$

where

$$\begin{aligned} u_k &:= u/k, \\ V_\Lambda(x) &:= \Lambda^{-1} V(x) + 1, \end{aligned} \quad (23)$$

with $\Lambda > 1$ to be chosen later to be sufficiently large. The introduction of the parameter Λ is somewhat motivated by the expression for the Taub-NUT metrics via the Gibbons-Hawking ansatz.

Equivalently, h_k is the unique principal S^1 bundle metric on the total space of $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$ such that:

- (1) The connection 1-form is θ with $d\theta = \tilde{\pi}^*\omega$; This connection is smooth away from $\{p_\ell^\alpha\}$ but degenerates in the natural S^3 coordinates near $\{p_\ell^\alpha\}$.

- (2) The induced metric on the base space $\mathcal{B}_k$ is $V_\Lambda(u_k)g_{S^3}$; For large Λ the geometry of $\mathcal{B}_k$ is close to S^3 away from $\{p_\ell^\alpha\}$, with singularities near $\{p_\ell^\alpha\}$.
- (3) The S^1 fibers have length $(k\Lambda\sqrt{V_\Lambda(u_k)})^{-1}$; For large Λ the fibers are $\approx \frac{1}{k\Lambda}$ small away from $\{p_\ell^\alpha\}$, with fiber length tending to zero at $\{p_\ell^\alpha\}$.

Remark 4.2 (Choice of V). *The specific choice of V in (21) does not play a central role in the sequel; it is merely convenient for calculations. In principle, any smooth convex function $V : \mathbb{R} \rightarrow (0, +\infty)$, independent of k , with the correct asymptotic behavior $V(x) \sim |x|$ as $x \rightarrow \pm\infty$, would work.*

Despite the obvious singularities of the metric components in the S^3 coordinates, we claim that h_k extends to a smooth Riemannian metric on M_k^4 . This Riemannian metric is invariant under an effective isometric action of the finite Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$. Moreover, the manifolds (M_k^4, h_k) have uniformly bounded diameters, and $0 \leq \text{Ric}_{h_k} \leq 3$ for every $k \in \mathbb{N}$. We will prove these claims, corresponding to Lemma 4.3, Lemma 4.4, and Lemma 4.5 below respectively, in the forthcoming subsections.

Lemma 4.3. *The metric h_k on $\mathcal{N}_k$ extends smoothly to a Riemannian metric, still denoted h_k , on M_k^4 .*

Lemma 4.4. *The finite Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$ acts effectively and isometrically on (M_k^4, h_k) .*

Lemma 4.5. *The manifold (M_k^4, h_k) satisfies*

- (1) $\text{diam}(M_k) \leq \pi + O(k^{-1})$,
- (2) $0 \leq \text{Ric}_{h_k} \leq 2 + O(k^{-1})$.

Moreover, we have that $\text{Ric}_{h_k} > 0$ in $\mathcal{N}_k \subset M_k$. That is, the Ricci tensor is zero only at the $2k$ points $\{\tilde{p}_\ell^\alpha\}$ added to $\mathcal{N}_k$ to form M_k^4 .

Once the proofs of Lemma 4.3, Lemma 4.4, and Lemma 4.5 are completed, to conclude the proof of Theorem 1.2 we will perturb the geometry of (M_k^4, h_k) with a smooth conformal factor $g_k := e^{-2\varphi_k} h_k$. This will allow us to achieve $\text{Ric}_{g_k} > 0$ while maintaining the Ricci and diameter upper bounds, as well as the isometric action of the Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$.

Lemma 4.6. *There exists $\varphi_k \in C^\infty(M_k^4)$ such that $|\varphi_k| \leq O(k^{-1})$ and φ_k is invariant under the action of $H_3(\mathbb{Z}/k\mathbb{Z})$. Moreover, the metric $g_k := e^{-2\varphi_k} h_k$ satisfies $0 < \text{Ric}_{g_k} \leq 2 + O(k^{-1})$.*

Similar conformal perturbation arguments have been used in several other instances, for instance in [2] and [15].

Remark 4.7. *The L^2 -norm of the Riemann tensor of (M_k^4, g_k) is not uniformly bounded with respect to k . Indeed, by Chern-Gauss-Bonnet*

$$\int_{M_k^4} |\text{Riem}_{g_k}|^2 = 32\pi^2 \chi(M_k^4) + \int_{M_k^4} 4|\text{Ric}_{g_k}|^2 - R_{g_k}^2. \quad (24)$$

Although the Ricci and scalar curvature terms are bounded, $\chi(M_k^4) = 2k + 1 \rightarrow \infty$ as $k \rightarrow \infty$. One can also verify that the sectional curvatures are not uniformly bounded, either from below or above.

4.1. Proof of Lemma 4.3. Some variants of the proof below have already appeared in the literature in slightly different contexts, see for instance [1, 28]. It is also clearly enough to check that h_k admits a smooth extension within each $\tilde{\pi}^{-1}(B_r(p_\ell^\alpha))$ for some sufficiently small $0 < r = r(k) < 1$.

Let us first write each $B_r(p_\ell^\alpha) \subset S^3$ in polar coordinates as

$$\begin{aligned} B_r(p_\ell^\alpha) \setminus \{p_\ell^\alpha\} &\approx B_r(0^3) \setminus \{0^3\} \approx (0, r) \times S^2, \\ g_{S^3} &= ds^2 + \sin^2(s)g_{S^2}. \end{aligned} \quad (25)$$

Recall that $\tilde{\pi}^{-1}(B_r(p_\ell^\alpha) \setminus \{p_\ell^\alpha\}) \cup \{\tilde{p}_\ell^\alpha\} \approx \mathbb{R}^4$ is identified with a ball in Euclidean space. We can make this identification a bit more explicitly as follows. Observe that $[\omega]$, the defining cohomology class of the circle bundle, restricts to that of a Hopf bundle over each S^2 by (14). We can write $\tilde{\pi}^{-1}(B_r(p_\ell^\alpha)) \approx \mathbb{R}^4 \setminus \{0^4\}$ in polar coordinates, as in [28], as the projection mapping

$$\pi : (0, r) \times S^3 \ni (s, x) \mapsto \left(\frac{s^2}{2}, \pi_{\text{Hopf}}(x) \right) \in (0, r) \times S^2. \quad (26)$$

We see that in these coordinates π extends smoothly to $\mathbb{R}^4$. Recall that the metric (22) is written

$$\begin{aligned} h_k &:= \frac{1}{k\Lambda} (kV(u/k) + k\Lambda) \left(\pi^* g_{S^3} + \frac{1}{(kV(u/k) + k\Lambda)^2} \theta \otimes \theta \right), \\ g_{S^3} &:= ds^2 + \sin^2(s)g_{S^2}. \end{aligned} \quad (27)$$

We have that u is a solution of $-\Delta_{g_{S^3}} u = 2\pi(-1)^\alpha \delta_{0^3}$, which we can write in these coordinates as

$$u = (-1)^\alpha \frac{1}{2} \frac{\cos(s)}{\sin(s)} + u', \quad (28)$$

where u' is a smooth solution of $\Delta_{S^3} u' = 0$ on $B_r(p_\ell^\alpha)$. The connection 1-form θ is determined by the equation $d\theta = *_S du$. If θ_{Hopf} is the Hopf connection 1-form on S^3 , then $(-1)^\alpha d\theta_{\text{Hopf}} = d\theta - d *_S du'$. Hence, up to gauge transformation, we have that

$$\theta = (-1)^\alpha \theta_{\text{Hopf}} + \theta', \quad (29)$$

where θ' is smooth on $\tilde{\pi}^{-1}(B_r(p_\ell^\alpha)) \approx \mathbb{R}^4$.

Using the explicit polar coordinate expression of the projection $\pi : \mathbb{R}^4 \rightarrow \mathbb{R}^3$, we can express

$$\pi^* g_{S^3} = s^2 ds^2 + \sin^2\left(\frac{s^2}{2}\right) \pi_{\text{Hopf}}^* g_{S^2}. \quad (30)$$

On the other hand, from (28) and the explicit expression of V , we can deduce that

$$kV(u/k) + k\Lambda = \frac{1}{2} \frac{\cos(s)}{\sin(s)} + u'', \quad u'' \in C^\infty(B_{r_0}(0^3)). \quad (31)$$

By combining (29), (30), and (31), we obtain the expansion

$$\begin{aligned} h_k &= \frac{1}{k\Lambda} \left(ds^2 + s^2 \pi_{\text{Hopf}}^* \frac{1}{4} g_{S^2} + s^2 \theta_{\text{Hopf}} \otimes \theta_{\text{Hopf}} \right) + h' \\ &= \frac{1}{k\Lambda} g_{\mathbb{R}^4} + h', \end{aligned} \quad (32)$$

where $h' = O(r^2)$ is smooth. In particular, h_k admits a smooth extension, as claimed. $\square$

4.2. Proof of Lemma 4.4. It suffices to show that $H_3(\mathbb{Z}/k\mathbb{Z})$ acts effectively and isometrically on $(\mathcal{N}_k, h_k)$, as (M_k, h_k) is the metric completion of $(\mathcal{N}_k, h_k)$.

Let $K \cong \mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ be the group of diffeomorphisms of $\mathcal{B}_k$ introduced in (12). Let H be the group of diffeomorphisms $\varphi : \mathcal{N}_k \rightarrow \mathcal{N}_k$ such that:

- (i) φ is equivariant with respect to the S^1 action on $\mathcal{N}_k$, i.e., $R_\theta \circ \varphi = \varphi \circ R_\theta$ for every $\theta \in S^1$. Here we denoted by $R_\theta : \mathcal{N}_k \rightarrow \mathcal{N}_k$ the S^1 action on the principal S^1 bundle $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$;
- (ii) φ is a lift of some element $\tilde{\varphi} \in K$, i.e., $\tilde{\pi} \circ \varphi = \tilde{\varphi} \circ \tilde{\pi}$;
- (iii) $\varphi^* \theta = \theta$, where θ is the connection 1-form on $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$.

Claim 1: H acts isometrically on $(\mathcal{N}_k, h_k)$.

For any $\varphi \in H$ we can compute

$$\varphi^* h_k = \varphi^* \left(V_\Lambda(u_k) \tilde{\pi}^* g_{S^3} + \frac{1}{k^2 \Lambda^2 V_\Lambda(u_k)} \theta \otimes \theta \right) \quad (33)$$

$$= V_\Lambda(u_k \circ \tilde{\varphi}) \tilde{\pi}^* \tilde{\varphi}^* g_{S^3} + \frac{1}{k^2 \Lambda^2 V_\Lambda(u_k \circ \tilde{\varphi})} \varphi^* \theta \otimes \varphi^* \theta \quad (34)$$

$$= V_\Lambda(u_k) \tilde{\pi}^* g_{S^3} + \frac{1}{k^2 \Lambda^2 V_\Lambda(u_k)} \theta \otimes \theta = h_k, \quad (35)$$

where we used (ii), (iii), and the fact that K acts isometrically with respect to the standard metric on S^3 .

Claim 2: Any $\tilde{\varphi} \in K$ admits a lift $\varphi \in H$. Such a lift is unique modulo composition with the action R_θ .

The Claim follows from a fairly standard lifting argument, see for instance [24, Proposition 2.7]. Indeed, as $\tilde{\varphi}^* \omega = \omega$ and $H^2(\mathcal{B}_k, \mathbb{Z})$ is torsion-free we have that $\tilde{\varphi}$ preserves the Chern class of our S^1 bundle and hence lifts to an equivariant diffeomorphism $\hat{\varphi} : \mathcal{N}_k \rightarrow \mathcal{N}_k$. This lift is well defined up to a gauge transformation, so let us observe that $d(\hat{\varphi}^* \theta - \theta) = \omega - \omega = 0$. As $H^1(\mathcal{B}_k, \mathbb{Z}) = 0$ we can write $d(\hat{\varphi}^* \theta - \theta) = -df$ with $f : \mathcal{N}_k \rightarrow \mathbb{R}$. We use f to define our gauge transformation we obtain $\varphi = \hat{\varphi} \cdot e^{if}$, where $\varphi^* \theta - \theta = \hat{\varphi}^* \theta + df - \theta = 0$, as claimed.

Claim 3: H contains a subgroup isomorphic to $H_3(\mathbb{Z}/k\mathbb{Z})$.

The restriction of ω to the Clifford torus $\Sigma := \{|z_1| = |z_2|\} \subset \mathcal{B}_k$ is cohomologous to the $2\pi k$ -multiple of the volume form. Indeed, up to a choice of orientation, from (19) we deduce that

$$\int_\Sigma \omega = \int_{\{|z_1| < |z_2|\}} \Delta_{S^3} u = 2\pi k. \quad (36)$$

Hence $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$ restricts to a principal S^1 bundle over the Clifford torus isomorphic to the degree k Heisenberg nilmanifold Nil_k^3 .

Note that K leaves the Clifford torus Σ invariant. Hence by (ii) we can restrict the action of H on $\mathcal{N}_k$ to an isometric action on a Riemannian Heisenberg nilmanifold Nil_k^3 respecting the principal S^1 bundle structure. As the nilmanifold Nil_k^3 is codimension one and the isometries of H are orientation preserving, we see that the restriction homomorphism $r : H \rightarrow \text{Iso}(\text{Nil}_k^3)$ is injective. Moreover, we see it is an isomorphism onto the subgroup of isometries of Nil_k^3 which are lifts of the $\mathbb{Z}/k\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ action on $\Sigma \approx T^2$, viewed as the base of $S^1 \rightarrow \text{Nil}_k^3 \rightarrow T^2$. It then follows from the construction of Section 2.2 that $r(H)$ (and hence H) has a subgroup isomorphic to $H_3(\mathbb{Z}/k\mathbb{Z})$ acting as in (7). $\square$

4.3. Proof of Lemma 4.5. We will prove first that for each $\varepsilon > 0$ if $\Lambda = \Lambda(\varepsilon) \geq 1$ is sufficiently large, actually independent of k , then $\text{diam}(M_k) \leq \pi + \varepsilon$. In particular, if $\Lambda(k) \rightarrow \infty$ then we have the claimed diameter bound on M_k^4 .

First observe that if $\Lambda \geq \Lambda(\delta)$ then $k^2 \Lambda^2 V_\Lambda \geq \delta^{-2}$, which ensures that the fiber lengths of the principal bundle $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$ are less than δ for every $k \geq 1$. Thus it is sufficient to establish a uniform diameter bound for the metrics $V_\Lambda(u_k)g_{S^3}$ on the base space. We may also focus our attention on $\mathcal{B}_k$, as M_k^4 is the (metric) completion of $(\mathcal{N}_k, h_k)$.

To prove the diameter bound on $V_\Lambda(u_k)g_{S^3}$ we need to estimate u_k . Let us begin with the following:

Claim: $\exists C > 0$, independent of k , so that $|u_k(z)| \leq C + |z_1|^{-1} + |z_2|^{-1}$.

Recall that $u_1 : S^3 \rightarrow [-\infty, +\infty]$ is the unique solution of

$$-\Delta_{S^3} u_1 = 2\pi (\delta_{(1,0)} - \delta_{(0,1)}) , \quad (37)$$

such that $u_1 \circ \iota = -u_1$. Note that we can also identify u_1 as the difference between the Green's functions on S^3 at $(1,0)$ and $(0,1)$. Using the standard estimates on Green's functions in dimension three we have for $(z_1, z_2) \in S^3 \subseteq \mathbb{C} \times \mathbb{C}$ the estimate

$$\left| u_1(z_1, z_2) - \frac{1}{\sqrt{|z_1 - 1|^2 + |z_2|^2}} + \frac{1}{\sqrt{|z_1|^2 + |z_2 - 1|^2}} \right| \leq C . \quad (38)$$

As a consequence we have the (very poor) estimate

$$|u_1(z)| \leq C + |z_2|^{-1} + |z_1|^{-1} , \quad (39)$$

which bounds u_1 uniformly away from the singular fibers F_0 and F_1 . For each $k \geq 2$ we can write

$$u_k(z_1, z_2) = \frac{1}{k} \sum_{\ell=0}^{k-1} u_1(e^{\frac{2\pi i \ell}{k}} z_1, e^{\frac{2\pi i \ell}{k}} z_2) . \quad (40)$$

In particular we arrive at the (less poor, as k gets large) estimate

$$|u_k(z)| \leq C + |z_1|^{-1} + |z_2|^{-1} , \quad (41)$$

which finishes the proof of the claim.

With the claim in hand we can now estimate $V_\Lambda(u_k) := \Lambda^{-1}V(u_k) + 1$. To begin, we can directly plug in the estimate of the claim to get

$$V_\Lambda(u_k(z_1, z_2)) \leq \Lambda^{-1}(C + |z_1|^{-1} + |z_2|^{-1}) + 1 , \quad (42)$$

for every $(z_1, z_2) \in S^3$. Consider for $r > 0$ the set $S^3 \setminus B_r^{S^3}(F_0 \cup F_1)$. Note we have subtracted the r -tube with respect to the S^3 metric around each fiber off. Then using (42) we can choose $\Lambda \geq \Lambda(C, r, \delta) > 0$ sufficiently large that

$$|V_\Lambda(u_k) - 1| \leq \begin{cases} \delta^3 \left(1 + \frac{1}{|z_1|} + \frac{1}{|z_2|}\right) & \text{in } B_r^{S^3}(F_0 \cup F_1) \\ \delta^3 & \text{in } S^3 \setminus B_r^{S^3}(F_0 \cup F_1) . \end{cases} \quad (43)$$

In particular, the metric $V_\Lambda(u_k)g_{S^3}$ is uniformly close to the S^3 metric on the set $S^3 \setminus B_r^{S^3}(F_0 \cup F_1)$, and it is then clear that the diameter of this set is $\leq (1 + \delta)\pi$. To estimate the diameter of $\mathcal{B}_k$ we are left showing that each point of $B_r^{S^3}(F_0)$ (and $B_r^{S^3}(F_1)$) can be connected to the boundary by a curve of small length. To this end we have the following, which focuses on F_0 :

Claim: Let $z_1 \in \mathbb{C}$ be such that $|z_1| = 1$. Consider the curve $\gamma_{z_1} : (0, 2\sqrt{r}) \rightarrow S^3$ defined as

$$\gamma_{z_1}(t) := (\sqrt{1-t^2} z_1, t^2). \quad (44)$$

If $\ell_k(\gamma_{z_1})$ denotes the length of γ_{z_1} with respect to the metric $V_\Lambda(u_k) g_{S^3}$, then

$$\ell_k(\gamma_{z_1}) \leq 6r + 24\delta^{3/2}\sqrt{r}. \quad (45)$$

To prove the claim, let us use (43) and the estimate $|\gamma'_{z_1}(t)| \leq 3t$ to obtain

$$\ell_k(\gamma_{z_1}) = \int_0^{2\sqrt{r}} \sqrt{V_\Lambda(u_k(\gamma_{z_1}(t)))} |\gamma'_{z_1}(t)| dt \quad (46)$$

$$\leq \int_0^{2\sqrt{r}} \left(1 + 4\frac{\delta^{3/2}}{t}\right) \cdot 3t dt \quad (47)$$

$$\leq 6r + 24\delta^{3/2}\sqrt{r}, \quad (48)$$

as claimed.

If we now choose $r = r(\varepsilon) > 0$, $\delta(\varepsilon) > 0$ sufficiently small with $\Lambda > \Lambda(C, r, \varepsilon, \delta) > 0$ sufficiently big, we can combine the estimates of this subsection to arrive at the diameter bound $\text{diam}(M_k^4) \leq \pi + \varepsilon$, as claimed. $\square$

4.4. Proof of Lemma 4.5, Ricci Curvature Bound. Let U and X be a unit vertical vector and a unit horizontal vector for the S^1 bundle metric h_k on $\tilde{\pi} : \mathcal{N}_k \rightarrow \mathcal{B}_k$ respectively. We claim that

$$\text{Ric}_{h_k}(U, U) = \frac{|du_k|^2}{2} \frac{\Lambda}{V(u_k) + \Lambda} \left[\frac{V''(u_k)}{V(u_k) + \Lambda} + \frac{1 - (V'(u_k))^2}{(V(u_k) + \Lambda)^2} \right], \quad (49)$$

$$\text{Ric}_{h_k}(U, X) = 0, \quad (50)$$

$$\text{Ric}_{h_k}(X, X) = \frac{\Lambda}{V(u_k) + \Lambda} \left(2 + \frac{(1 - (V'(u_k))^2)}{2(V(u_k) + \Lambda)} [du_k(X)]^2 \right. \quad (51)$$

$$\left. - \frac{|du_k|^2}{2} \left[\frac{V''(u_k)}{V(u_k) + \Lambda} + \frac{1 - (V'(u_k))^2}{(V(u_k) + \Lambda)^2} \right] \right), \quad (52)$$

where, with a slight abuse of notation, we denoted $du(X) := du(d\tilde{\pi}(X))$.

We postpone the verification of the expressions of the Ricci curvature to the end of the subsection and first exploit them to complete the proof of Lemma 4.5.

Consider the identities

$$1 - (V'(x))^2 = \frac{4}{e^{4\pi x} + e^{-4\pi x} + 2}, \quad V''(x) = \frac{8\pi}{e^{4\pi x} + e^{-4\pi x} + 2}. \quad (53)$$

We can use the asymptotic expansion of u_k near the poles to estimate

$$\frac{|du_k|^2}{e^{4\pi u_k} + e^{-4\pi u_k} + 2} \leq C(k). \quad (54)$$

If we combine with the previous estimate we get

$$\begin{aligned} 0 &< |du_k|^2 V''(u_k) \leq C(k) \\ 0 &< |du_k|^2 (1 - (V'(u_k))^2) \leq C(k). \end{aligned} \quad (55)$$

If we choose $\Lambda \geq \Lambda(k, \delta)$ we then get the estimates

$$\begin{aligned} 0 &< |du_k|^2 \frac{V''(u_k)}{V(u_k) + \Lambda} \leq \delta \\ 0 &< |du_k|^2 \frac{(1 - (V'(u_k))^2)}{V(u_k) + \Lambda} \leq \delta, \end{aligned} \quad (56)$$

as well as

$$0 \leq \frac{\Lambda}{V(u_k) + \Lambda} \leq 1, \quad (57)$$

with strict positivity near the poles. If we plug these into our formulas for the Ricci curvature we arrive at the estimates (away from the poles of u_k)

$$\begin{aligned} 0 < \text{Ric}_{h_k}(U, U) &\leq \delta, \\ \text{Ric}_{h_k}(X, U) &= 0, \\ 0 < \text{Ric}_{h_k}(X, X) &< 2 + 10\delta, \end{aligned} \quad (58)$$

which proves our desired Ricci curvature bounds.

We are left with the verification of the expressions (49), (50), and (51), for the Ricci curvatures of the metric g_k . To this aim, it is convenient to view

$$h_k = \frac{1}{k\Lambda} \left[W_k(u) \tilde{\pi}^* g_{S^3} + \frac{1}{W_k(u)} \theta \otimes \theta \right], \quad (59)$$

for $W_k(x) := k(V(x/k) + \Lambda)$ for each $k \geq 1$, and compute the Ricci curvatures of a general Riemannian circle bundle metric of the form

$$W \tilde{\pi}^* g_{S^3} + \frac{1}{W} \theta \otimes \theta, \quad (60)$$

with $W : S^3 \rightarrow (0, +\infty]$. If ω denotes the curvature 2-form of such circle bundle, U is a unit vertical vector and X any horizontal vector, then there hold

$$\text{Ric}(U, U) = \frac{1}{2} W^{-2} \Delta_{g_{S^3}} W + \frac{1}{2} W^{-3} \left(|\omega|_{g_{S^3}}^2 - |\nabla W|_{g_{S^3}}^2 \right), \quad (61)$$

$$\text{Ric}(U, X) = -\frac{1}{2} W^{-3/2} \left(\delta_{g_{S^3}} \omega(X) - W^{-1} \omega(\nabla^{g_{S^3}} W, X) \right), \quad (62)$$

$$\begin{aligned} \text{Ric}(X, X) &= 2|X|_{g_{S^3}}^2 - \frac{1}{2} \frac{\Delta_{g_{S^3}} W}{W} |X|_{g_{S^3}}^2 \\ &\quad - W^{-2} \frac{1}{2} \left(|\iota_X \omega|_{g_{S^3}}^2 - |dW \wedge X^\flat|_{g_{S^3}}^2 \right). \end{aligned} \quad (63)$$

To prove (61), (62), and (63) we can start from the well-known formulas for the Ricci curvatures of Riemannian circle bundles $\pi : M \rightarrow B$ with fibres length f , curvature 2-form ω , and induced Riemannian metric on the base g_B . Namely

$$\text{Ric}(U, U) = -\frac{\Delta_B f}{f} + \frac{f^2}{2} |\omega|_B^2, \quad (64)$$

$$\text{Ric}(U, X) = \frac{1}{2} \left(-f \delta_B \omega(X) + 3\omega(X, \nabla^B f) \right), \quad (65)$$

$$\text{Ric}(X, X) = \text{Ric}_B(X, X) - \frac{f^2}{2} |\omega(X)|_B^2 - \frac{\nabla_B^2 f(X, X)}{f}, \quad (66)$$

where as above U denotes a unit vertical vector and X denotes any horizontal vector. See for instance [4, Proposition 9.36], or [18].

To get from (64) to (61) it suffices to exploit the formula for the transformation of the Laplacian under a conformal change in dimension 3, i.e.,

$$\Delta_{W g_{S^3}} f = W^{-1} \left(\Delta_{S^3} f + \frac{1}{2} W^{-1} g_{S^3}(\nabla^{g_{S^3}} W, \nabla^{g_{S^3}} f) \right), \quad (67)$$

and the norm of 2-forms

$$|\omega|_{W g_{S^3}}^2 = W^{-2} |\omega|_{g_{S^3}}^2. \quad (68)$$

To get from (65) to (62) we rely on the formula for the transformation of the divergence for 2-forms under conformal changes, i.e.,

$$\delta_{Wg_{S^3}}\omega = W^{-1} \left(\delta_{g_{S^3}}\omega + \frac{1}{2}W^{-1}\iota_{\nabla^{g_{S^3}}W}\omega \right). \quad (69)$$

To obtain (63) from (66) we combine the formulas for the transformation of Ricci curvature and Hessians under conformal changes to establish the identity

$$\begin{aligned} \text{Ric}_{Wg_{S^3}} - \frac{\text{Hess}_{Wg_{S^3}}W^{-1/2}}{W^{-1/2}} &= \text{Ric}_{g_{S^3}} - \frac{1}{2} \left(\frac{\Delta_{g_{S^3}}W}{W} - \frac{|dW|^2}{W^2} \right) g_{S^3} \\ &\quad - 2 \frac{dW}{W} \otimes \frac{dW}{W}. \end{aligned} \quad (70)$$

Under the assumption that $\omega = *du$, where $u : \mathcal{B}_k \rightarrow \mathbb{R}$ is harmonic with respect to the round metric g_{S^3} , (61), (62), and (63) simplify into

$$\text{Ric}(U, U) = \frac{1}{2} \frac{\Delta W}{W^2} + \frac{1}{2} \frac{|du|^2 - |dW|^2}{W^3}, \quad (71)$$

$$\text{Ric}(U, X) = \frac{1}{2} \frac{*(du \wedge dW)(X)}{W^{5/2}}, \quad (72)$$

$$\text{Ric}(X, X) = 2|X|^2 - \frac{1}{2} \frac{\Delta W}{W} |X|^2 - \frac{1}{2} \frac{|du \wedge X^\flat|^2 - |dW \wedge X^\flat|^2}{W^2}. \quad (73)$$

Above, it is understood that norms and Laplacians are computed with respect to the round metric g_{S^3} . If we also assume that, with a slight abuse of notation, $W = W \circ u$, then we can rewrite (71), (72), and (73) as

$$\text{Ric}(U, U) = \frac{1}{2} \frac{|du|^2}{W} \left(\frac{W''}{W} + \frac{1 - (W')^2}{W^2} \right), \quad (74)$$

$$\text{Ric}(U, X) = 0, \quad (75)$$

$$\begin{aligned} \text{Ric}(X, X) &= 2|X|^2 + \frac{1 - (W')^2}{2W^2} (du(X))^2 \\ &\quad - \frac{1}{2} \left(\frac{W''}{W} + \frac{1 - (W')^2}{W^2} \right) |du|^2 |X|^2. \end{aligned} \quad (76)$$

The expressions (49), (50), and (51) can be easily obtained from (74), (75), and (76), recalling that $W_k(x) := k(V(x/k) + \Lambda)$ and scaling to account for the $\frac{1}{k\Lambda}$ factor in (59). $\square$

4.5. Proof of Lemma 4.6. Recall that $M_k \setminus \mathcal{N}_k$ is a set of $2k$ points forming an orbit of the isometric action of the Heisenberg group $H_3(\mathbb{Z}/k\mathbb{Z})$. We define $\rho_k(x) := \frac{1}{2}d_{h_k}(x, M_k \setminus \mathcal{N}_k)^2$, the square of the distance function to this set with respect to the metric h_k . Note that ρ_k is globally $H_3(\mathbb{Z}/k\mathbb{Z})$ -invariant and smooth in a sufficiently small neighborhood of $M_k \setminus \mathcal{N}_k$. Let $\eta_k \in C^\infty(\mathbb{R})$ be a smooth function such that $\eta_k(t) = t$ for $t \leq r_k$ and $\eta_k(t) = 0$ for $t \geq 2r_k$, where $r_k > 0$ is sufficiently small to ensure that $\eta_k(\rho_k) \in C^\infty(M_k)$ and

$$\text{Hess}_{h_k}\rho_k \geq \frac{1}{2}h_k, \quad |d\rho_k|_{h_k} \leq 10^{-2}, \quad \text{when } \rho_k \leq r_k^2. \quad (77)$$

The existence of such r_k follows from the fact that

$$\text{Hess}_{h_k}\rho_k = h_k, \quad d\rho_k = 0, \quad (78)$$

for every $\tilde{p}_\ell^\alpha \in M_k \setminus \mathcal{N}_k$.

Finally, we define $\varphi_k := \varepsilon_k \eta_k(\rho_k)$, where $\varepsilon_k > 0$ is a small parameter to be chosen later. The Ricci curvature of $g_k := e^{-2\varphi_k}h_k$ is given by

$$\text{Ric}_{g_k} = \text{Ric}_{h_k} + 2\text{Hess}_{h_k}\varphi_k + 2d\varphi_k \otimes d\varphi_k + (\Delta_{h_k}\varphi_k - 2|d\varphi_k|_{h_k})h_k. \quad (79)$$

It is clear that $\text{Ric}_{g_k} > 0$ in the set $\{\rho_k \leq r_k/2\}$ as a consequence of (77). On the other hand, $\text{Ric}_{h_k} \geq c > 0$ in the set $\{\rho_k \geq r_k/2\} \subset \mathcal{N}_k$, hence Ric_{g_k} is positive in this region provided $\varepsilon_k > 0$ is small enough. Similarly, we have the upper bound $\text{Ric}_{g_k} \leq \text{Ric}_{h_k} + C\varepsilon_k$. $\square$

5. PROOF OF THEOREM 1.1

The Heisenberg actions in Theorem 1.2 are not free, and thus do not yield the desired fundamental group. One can verify that the quotient of M_k^4 by the Heisenberg action $H_3(\mathbb{Z}/k\mathbb{Z})$ is homeomorphic to the four-sphere S^4 .

To address this issue, consider the oriented frame bundle $\pi_{FM} : FM_k \xrightarrow{\text{SO}(4)} M_k^4$. Given constants $d_k > 0$ there is a unique principle bundle metric g_k^{FM} on FM_k defined by the conditions:

- (1) The right $\text{SO}(4)$ actions on FM_k are isometric. More specifically, let $\{v^a\}$ be an orthonormal basis of $\mathfrak{so}(4)$, the Lie algebra of $\text{SO}(4)$, with respect to the standard Frobenius inner product. If V^a denotes the corresponding pushforward vertical vector fields on FM_k , then the metric satisfies $g_k^{FM}(V^a, V^b) = d_k^2 \delta^{ab}$.
- (2) Consider the natural connection form $\theta \in \Omega^1(FM_k; \mathfrak{so}(4))$ associated with the Levi-Civita connection ∇_k on (M_k, g_k) . If H^j are vector fields on M_k we may lift them to horizontal vector fields on FM_k , sloppily also denoted by H_k . Then $g_k^{FM}(H^j, V^a) = 0$.
- (3) If H^j are horizontal lifts of vector fields on M_k , then $g_k^{FM}(H^j, H^k) = g_k(H^j, H^k)$.

We begin with the usual observation about isometries on M_k and their lifts to FM_k . Namely, note that an isometry on M_k maps an orthonormal frame to an orthonormal frame, and thus lifts to a mapping on FM_k . It is easy to check this lifted mapping is an isometry with respect to a metric as above. Additionally, since an isometry fixes a point and a frame (e.g. its derivative is the identity) if and only if it is globally the identity, we have that the lifted isometric action on FM_k is free. In particular, we can lift our $H_3(\mathbb{Z}/k\mathbb{Z})$ action to a *free* action on FM_k .

To understand the Ricci curvature of FM_k , let us begin with the observation that if d_k is very small then our $\text{SO}(4)$ fibers are very small. In particular, they are very positively curved. We want to see that the positive Ricci curvature of M_k together with the potentially very positive Ricci curvature of the fibers leads to positive Ricci curvature of FM_k .

More precisely, let V be a g^{FM} unit vertical vector with H unit a horizontal vector. Note that the V can be identified as a 2-form in M_k (in local coordinates $\sum_{ij} \theta_{ij}(V) dx_i \wedge dx_j$), and so we can identify $\text{Rm}_k[V]$ as a 2-form. Now recall the formulas (see [32] and also [3, 31, 4]):

$$\begin{aligned} \text{Ric}^{FM}(V, V) &= d_k^{-2} + \frac{1}{4} |\text{Rm}_k[V]|_k^2 \geq d_k^{-2}, \\ \text{Ric}^{FM}(V, H) &= \langle \text{div} \text{Rm}_k[H], V \rangle_{FM}, \implies |\text{Ric}^{FM}(V, H)| \leq C d_k, \\ \text{Ric}^{FM}(H, H) &= \text{Ric}_k(H, H) - \frac{3}{4} d_k^2 \sum_{i=1}^3 |\text{Rm}_k(H, H_i)|_k^2 \geq \epsilon_k - C d_k^2, \end{aligned} \quad (80)$$

where H, H_1, H_2, H_3 form an orthonormal frame of (M_k, g) , $C = C(n, \text{Rm}_k, \nabla \text{Rm}_k)$ and ϵ_k is the positive Ricci lower bound on M_k^4 . In particular, if d_k is sufficiently small (depending on the size of Rm_k and ∇Rm_k and the lower Ricci bound on M_k^4), then positive Ricci on M_k^4 implies positive Ricci on FM_k .

We have thus built a ten dimensional compact manifold with positive Ricci curvature and a free action of $H_3(\mathbb{Z}/k\mathbb{Z})$. However, we are not quite done yet. As M_k is homeomorphic to the connected sum of $(k-1)$ copies of $S^2 \times S^2$ we have that M_k is a spin manifold. In particular, even though M_k is simply connected we have that FM_k is not simply connected. More precisely, $\pi_1(FM_k) = \mathbb{Z}/2\mathbb{Z}$.

In order to produce the examples of Theorem 1.1 we need to define $\widetilde{M}_k^{10}$ to be the double cover of FM_k , which is to say the spin bundle of M_k . As $\widetilde{M}_k^{10}$ is simply connected the action of $H_3(\mathbb{Z}/k\mathbb{Z})$ on FM_k lifts to an action by a $\mathbb{Z}/2\mathbb{Z}$ extension of $H_3(\mathbb{Z}/k\mathbb{Z})$. This now finishes the proof of Theorem 1.1. $\square$

REFERENCES

- [1] ANDERSON, MICHAEL T.; KRONHEIMER, PETER B.; LEBRUN, CLAUDE: *Complete Ricci-flat Kähler manifolds of infinite topological type*. Commun. Math. Phys. **125**, no. 4, 637-642 (1989).
- [2] AUBIN, THIERRY: *Métriques riemanniennes et courbure*. J. Differ. Geom. **4**, 383-424 (1970).
- [3] BÉRARD BERGERY, LIONEL: *Certains fibres à courbure de Ricci positive*. C. R. Acad. Sci., Paris, Sér. A **286**, 929-931 (1978).
- [4] BESSE, ARTHUR L.: *Einstein manifolds*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, Bd. 10. Berlin etc.: Springer-Verlag. XII, 510 p. (1987).
- [5] BRUÈ, ELIA; NABER, AARON; SEMOLA, DANIELE: *Fundamental Groups and the Milnor Conjecture*. Ann. Math. **201**, no. 1, 225-289 (2025).
- [6] BRUÈ, ELIA; NABER, AARON; SEMOLA, DANIELE: *Six Dimensional Counterexample to the Milnor Conjecture*. Preprint, arXiv:2311.12155 (2023).
- [7] BRUÈ, ELIA; NABER, AARON; SEMOLA, DANIELE: *Fukaya-Yamaguchi conjecture in dimension four*. Preprint (2025).
- [8] CHEEGER, JEFF; GROMOLL, DETLEF: *The splitting theorem for manifolds of nonnegative Ricci curvature*. J. Differ. Geom. **6**, 119-128 (1971).
- [9] CHEEGER, JEFF; GROMOLL, DETLEF: *On the structure of complete manifolds of non-negative curvature*. Ann. Math. (2) **96**, 413-443 (1972).
- [10] CHEN, GAO; VIACLOVSKY, JEFF; ZHANG, RUOBING: *Collapsing Ricci-flat metrics on elliptic K3 surfaces*. Commun. Anal. Geom. **28**, no. 8, 2019-2133 (2020).
- [11] CHURCH, PHILIP T. AND LAMOTKE, KLAUS: *Almost free actions on manifolds*. Bull. Austral. Math. Soc. **10**, 177-196 (1974).
- [12] CSIKÓS, BALÁZS; PYBER, LÁSZLÓ; SZABÓ, ENDRE: *Diffeomorphism Groups of Compact 4-manifolds are not always Jordan*. Preprint, arXiv:1411.7524 (2014).
- [13] CSIKÓS, BALÁZS; PYBER, LÁSZLÓ; SZABÓ, ENDRE: *Finite subgroups of the homeomorphism group of a compact topological manifold are almost nilpotent*. Preprint, arXiv:2204.13375 (2022).
- [14] DINKELBACH, JONATHAN; LEEB, BERNHARD: *Equivariant Ricci flow with surgery and applications to finite group actions on geometric 3-manifolds*. Geom. Topol. **13**, no. 2, 1129-1173 (2009).
- [15] EHRLICH, PAUL: *Metric deformations of curvature. I: Local convex deformations*. Geom. Dedicata **5**, 1-23 (1976).
- [16] FUKAYA, KENJI; YAMAGUCHI, TAKAO: *The fundamental groups of almost nonnegatively curved manifolds*. Ann. Math. (2) **136**, no. 2, 253-333 (1992).
- [17] GIBBONS, GARY; HAWKING, STEPHEN: *Gravitational multi-instantons*. Phys. Lett. **78B**, 430-432, (1978).
- [18] GILKEY, PETER B.; PARK, JEONG HYEONG; TUSCHMANN, WILDERICH: *Invariant metrics of positive Ricci curvature on principal bundles*. Math. Z. **227**, no. 3, 455-463 (1998).
- [19] GROMOV, MICHAEL: *Curvature, diameter and Betti numbers*. Comment. Math. Helv. **56**, 179-195 (1981).
- [20] GROSS, MARK: *Topological mirror symmetry*. Invent. Math. **144**, no. 1, 75-137 (2001).
- [21] GROSS, MARK; WILSON, P. M. H.: *Large complex structure limits of K3 surfaces*. J. Differ. Geom. **55**, no. 3, 475-546 (2000).
- [22] HEIN, HANS-JOACHIM: *Gravitational instantons from rational elliptic surfaces*. J. Am. Math. Soc. **25**, no. 2, 355-393 (2012).
- [23] HEIN, HANS-JOACHIM; SUN, SONG; VIACLOVSKY, JEFF; ZHANG, RUOBING: *Nilpotent structures and collapsing Ricci-flat metrics on the K3 surface*. J. Am. Math. Soc. **35**, no. 1, 123-209 (2022).

- [24] HONDA, NOBUHIRO; VIACLOVSKY, JEFF: *Conformal symmetries of self-dual hyperbolic monopole metrics*. Osaka J. Math. **50**, no. 1, 197-249 (2013).
- [25] KAPOVITCH, VITALI; PETRUNIN, ANTON; TUSCHMANN, WILDERICH: *Nilpotency, almost nonnegative curvature, and the gradient flow on Alexandrov spaces*. Ann. Math. (2) **171**, no. 1, 343-373 (2010).
- [26] KAPOVITCH, VITALI; PETRUNIN, ANTON; TUSCHMANN, WILDERICH: *On the torsion in the center conjecture*. Electron. Res. Announc. Math. Sci. **25**, 27-35 (2018).
- [27] KAPOVITCH, VITALI; WILKING, BURKHARD: *Structure of fundamental groups of manifolds with Ricci curvature bounded below*. Preprint arXiv:1105.5955 (2011).
- [28] LEBRUN, CLAUDE: *Explicit self-dual metrics on $\mathbb{CP}_2 \# \dots \# \mathbb{CP}_2$* . J. Differ. Geom. **34**, no. 1, 223-253 (1991).
- [29] MAZUR, MARCIN; RONG, XIAOCHUN; WANG, YUSHENG: *Margulis lemma for compact Lie groups*. Math. Z. **258**, no. 2, 395-406 (2008).
- [30] MUNDET I RIERA, IGNASI: *Finite group actions on homology spheres and manifolds with nonzero Euler characteristic*. J. Topol. **12**, no. 3, 744-758 (2019).
- [31] NASH, JOHN C.: *Positive Ricci curvature on fibre bundles*. J. Differ. Geom. **14**, 241-254 (1979).
- [32] O'NEILL, B.: *The fundamental equations of a submersion*. Mich. Math. J. **13**, 459-469 (1966).
- [33] PAN, JIAYIN; RONG, XIAOCHUN: *Ricci curvature and isometric actions with scaling nonvanishing property*. Preprint arXiv:1808.02329 (2018).
- [34] RONG, XIAOCHUN: *On the fundamental groups of manifolds of positive sectional curvature*. Ann. Math. (2) **143**, no. 2, 397-411 (1996).
- [35] SANTOS-RODRÍGUEZ, JAIME; ZAMORA-BARRERA, SERGIO: *On fundamental groups of RCD spaces*. J. Reine Angew. Math. **799**, 249-286 (2023).
- [36] TIAN, GANG; YAU, SHING TUNG: *Complete Kähler manifolds with zero Ricci curvature. I*. J. Am. Math. Soc. **3**, no. 3, 579-609 (1990).
- [37] WEI, GUOFANG: *Examples of complete manifolds of positive Ricci curvature with nilpotent isometry groups*. Bull. Am. Math. Soc., New Ser. **19**, no. 1, 311-313 (1988).
- [38] WILKING, BURKHARD: *On fundamental groups of manifolds of nonnegative curvature*. Differ. Geom. Appl. **13**, no. 2, 129-165 (2000).
- [39] ZARHIN, YURI G.: *Theta groups and products of abelian and rational varieties*. Proc. Edinb. Math. Soc., II. Ser. **57**, no. 1, 299-304 (2014).