

# EQUILIBRIA OF AGGREGATION-DIFFUSION MODELS WITH NONLINEAR POTENTIALS

FRANCESCO BOZZOLA AND EDOARDO MAININI

ABSTRACT. We consider an evolution model with nonlinear diffusion of porous medium type in competition with a nonlocal drift term favoring mass aggregation. The distinguishing trait of the model is the choice of a nonlinear  $(s, p)$  Riesz potential for describing the overall aggregation effect. We investigate radial stationary states of the dynamics, showing their relation with extremals of suitable Hardy-Littlewood-Sobolev inequalities. In the case that aggregation does not dominate over diffusion, radial stationary states also relate to global minimizers of a homogeneous free energy functional featuring the  $(s, p)$  energy associated to the nonlinear potential. In the limit as the fractional parameter  $s$  tends to zero, the nonlocal interaction term becomes a backward diffusion and we describe the asymptotic behavior of the stationary states.

## CONTENTS

1. Introduction	1
2. Main results	3
2.1. $(s, p)$ potential, stationary states and HLS inequalities	3
2.2. Asymptotic behavior of stationary states as $s \rightarrow 0$	5
3. Preliminaries	6
3.1. Notation and functional framework	6
3.2. Basics on Riesz potentials	7
4. Extremals of the HLS type inequality	10
4.1. Existence: the Lieb-Oxford method	10
4.2. Euler-Lagrange equation	14
4.3. Regularity properties	16
5. Minimizers of the energy functional	21
6. The limit $s \rightarrow 0$	26
6.1. The limit functional	26
6.2. The limiting diffusion dominated regime	28
6.3. The limiting fair competition regime	32
References	36

## 1. INTRODUCTION

We are interested in stationary solutions of aggregation-diffusion models of the form

$$(1.1) \quad \partial_t \rho = \Delta \rho^m - \chi \operatorname{div}(\rho \nabla \mathcal{S}(\rho)) \quad \text{in } (0, +\infty) \times \mathbb{R}^N,$$

where  $\rho = \rho(x, t)$  represents a mass density whose evolution is driven by a porous medium diffusion ( $m > 1$ ) and a nonlocal interaction modeled by a potential  $\mathcal{S}$  that accounts for long range effects. Here,  $\chi > 0$  is the sensitivity constant measuring the interaction strength.

---

*Date:* April 8, 2026.

*2010 Mathematics Subject Classification.* 35K44, 35R11, 49K20.

*Key words and phrases.* Aggregation-diffusion equations, nonlinear Riesz potential, stationary states.

Equations of the form (1.1) typically appear in mathematical biology as macroscopic models of interacting particles/agents [9, 17, 40, 43], such as the Keller-Segel model of chemotaxis [25, 27, 29, 30, 31, 41, 42]. These models usually feature linear potentials in convolution form, i.e.,  $\mathcal{S}(\rho)$  is the convolution of  $\rho$  with some suitable radial convolution kernel accounting for mutual interaction forces.

Among the most relevant modeling examples is the Newtonian or the Riesz (attractive) potential, appearing in the Keller-Segel model and its many variants, which is given by

$$(1.2) \quad \mathcal{S}(\rho) = K_s * \rho.$$

Here, the kernel  $K_s$  is defined for  $0 < s < N/2$  as

$$(1.3) \quad K_s(x) := c_{N,s} |x|^{2s-N}, \quad c_{N,s} := \pi^{-\frac{N}{2}} 2^{-2s} \frac{\Gamma(N/2 - s)}{\Gamma(s)},$$

and in terms of Fourier transform we have  $\hat{K}_s(\xi) = |\xi|^{-2s}$  (with  $\hat{u}(\xi) := \int_{\mathbb{R}^N} e^{-ix \cdot \xi} u(x) dx$ ). The particular case of the Newtonian potential corresponds to  $s = 1$  if  $N \geq 3$ . With the choice (1.2), the free energy of the system is

$$\mathcal{F}_s(\rho) = \frac{1}{m-1} \int_{\mathbb{R}^N} \rho^m - \frac{\chi}{2} \int_{\mathbb{R}^N} \rho K_s * \rho,$$

featuring the competition among the diffusion term and the total interaction energy associated to the mean field potential. Functional  $\mathcal{F}_s$  has to be analyzed among mass densities in the following class (defined for any given  $M > 0$ )

$$(1.4) \quad \mathcal{Y}_M = \mathcal{Y}_{M,m} := \left\{ \rho \in L^1_+(\mathbb{R}^N) \cap L^m(\mathbb{R}^N) : \int_{\mathbb{R}^N} x \rho(x) dx = 0, \int_{\mathbb{R}^N} \rho(x) dx = M \right\},$$

which naturally arises by taking into account that the evolution problem is formally preserving mass, center of mass and positivity.  $\mathcal{F}_s$  is a Lyapunov functional for the dynamics. In fact, (1.1)-(1.2) can be seen as the gradient flow of  $\mathcal{F}_s$  with respect to the square Wasserstein distance, see [10, 28]. In the search for stationary solutions to the evolution problem (1.1)-(1.2), it is therefore natural to look for minimizers (if existing) of  $\mathcal{F}_s$  over  $\mathcal{Y}_M$  and, more generally, for critical points satisfying suitable Euler-Lagrange equations. We also stress a crucial property of functional  $\mathcal{F}_s$ , which is the homogeneity with respect to the mass invariant dilations

$$(1.5) \quad \rho^\lambda(x) := \lambda^N \rho(\lambda x), \quad x \in \mathbb{R}^N, \lambda > 0.$$

Indeed, we have

$$\mathcal{F}_s(\rho^\lambda) = \frac{\lambda^{N(m-1)}}{m-1} \int_{\mathbb{R}^N} \rho^m - \frac{\chi \lambda^{N-2s}}{2} \int_{\mathbb{R}^N} \rho K_s * \rho.$$

As a consequence, aggregation and diffusion are in balance if  $m = 2 - 2s/N$ , which is called the *fair competition regime* [10]. If  $m$  is below this threshold, aggregation dominates and concentrating all the mass at a single point (that is, letting  $\lambda \rightarrow +\infty$ ) is energetically favorable.

The classical Keller-Segel model [30] of chemotaxis, in its simplest mathematical formulation [4, 8, 24, 29, 48] is a fair competition model, formally obtained by letting  $N = 2$ ,  $s = 1$  so that the convolution kernel is the Newtonian kernel (in dimension 2 it is understood that  $K_1(x) = -\frac{1}{2\pi} \log|x|$ ), and by letting the diffusion be linear  $m = m_c = 1$  (the diffusion term in

the free energy becomes  $\int_{\mathbb{R}^N} \rho \log \rho$ ). It is well known that a critical mass  $M_c$  exists in such a model (whose explicit value is  $8\pi/\chi$ ), and that global-in-time solutions for the associated Cauchy problem exist if the mass is not above the critical mass, while blow up in finite time occurs if  $M > M_c$ . Moreover, stationary states exist only if  $M = 8\pi/\chi$ , see [5, 9, 14]. The above properties of the classical Keller-Segel model generalize to fair competition models in higher dimension: it has been proven in the Newtonian potential case  $s = 1, N \geq 3$  in [6] that a critical mass  $M_c$  still appears for  $m = 2 - 2/N$ , that its value can be written in terms of the best constant of suitable Hardy-Littlewood-Sobolev (HLS) inequalities, and that stationary states exist only if  $M = M_c$ . The validity of analogous properties for more singular Riesz potentials  $0 < s < 1$  has been shown in [10, 11], still in the fair competition regime  $m = 2 - 2s/N$ . On the other hand, the diffusion dominated regime has been considered in [19], and in such case stationary states exist for every choice of the mass  $M > 0$  and can be obtained as minimizers of  $\mathcal{F}_s$  over  $\mathcal{Y}_M$ . In the aggregation dominated regime  $m < 2 - 2s/N$  the free energy  $\mathcal{F}_s$  is not bounded from below over  $\mathcal{Y}_M$  (whatever the choice of  $M > 0$ ), but stationary states of the dynamics can still be obtained, as seen in [20], as solutions to the Euler-Lagrange equation associated with the free energy (see also [3] for the Newtonian case  $s = 1$ ).

## 2. MAIN RESULTS

**2.1.  $(s, p)$  potential, stationary states and HLS inequalities.** In this work we shall investigate the nonlinear potential counterpart of the previous results about stationary states, by considering an interaction described by the nonlinear Riesz potential, which has been introduced in [38], see also [1], [37, Section 4.2], [39, Section 5.4] and the references therein. We let

$$(2.1) \quad \mathcal{S} = \mathcal{K}_{s,p},$$

where  $1 < p < \infty$ ,  $0 < sp < N$ , and  $\mathcal{K}_{s,p}$  stands for the *nonlinear  $(s, p)$  Riesz potential* given by

$$\mathcal{K}_{s,p}(\rho) := K_{s/2} * (K_{s/2} * \rho)^{p'-1}.$$

Here,  $p'$  is the conjugate exponent of  $p$ , i.e.,  $1/p + 1/p' = 1$ . The total interaction energy of the mass density  $\rho$  associated to the nonlinear potential  $\mathcal{K}_{s,p}$  (the  $(s, p)$  energy) is given by

$$\mathcal{I}_{s,p}(\rho) = \frac{1}{p'} \int_{\mathbb{R}^N} \rho \mathcal{K}_{s,p}(\rho) = \frac{1}{p'} \int_{\mathbb{R}^N} (K_{s/2} * \rho)^{p'},$$

where the second equality is due to Plancherel theorem, which also implies that for  $\varepsilon \rightarrow 0$

$$\mathcal{I}_{s,p}(\rho + \varepsilon\varphi) = \mathcal{I}_{s,p}(\rho) + \varepsilon \int_{\mathbb{R}^N} (K_{s/2} * \rho)^{p'-1} K_{s/2} * \varphi + o(\varepsilon) = \mathcal{I}_{s,p}(\rho) + \varepsilon \int_{\mathbb{R}^N} \mathcal{K}_{s,p}(\rho) \varphi + o(\varepsilon)$$

for every test function  $\varphi$ , showing that indeed  $\mathcal{K}_{s,p}$  is the functional derivative of  $\mathcal{I}_{s,p}$ . The free energy is therefore

$$\mathcal{F}_{s,p}(\rho) = \frac{1}{m-1} \int_{\mathbb{R}^N} \rho^m dx - \frac{\chi}{p'} \mathcal{I}_{s,p}(\rho)$$

and the evolution equation (1.1)-(2.1) is formally its Wasserstein gradient flow. The composition property of  $K_s$  shows that for  $p = 2$  we are reduced to the linear potential case:

$\mathcal{K}_{s,2}(\rho) = K_s * \rho$  and  $\mathcal{F}_{s,2}(\rho) = \mathcal{F}_s(\rho)$ . For  $p \neq 2$  the free energy is still a homogeneous functional, satisfying

$$\mathcal{F}_{s,p}(\rho^\lambda) = \frac{\lambda^{N(m-1)}}{m-1} \int_{\mathbb{R}^N} \rho^m - \lambda^{N(m_c-1)} \frac{\chi}{p'} \int_{\mathbb{R}^N} (K_{s/2} * \rho)^{p'},$$

where

$$(2.2) \quad m_c := p' - \frac{sp'}{N}$$

is the critical exponent. Therefore, we still recognize three regimes according to the value of the diffusion exponent  $m$ : we are in the *diffusion dominated regime* if  $m > m_c$ , in the *fair competition regime* if  $m = m_c$ , and in the *aggregation dominated regime* if  $m < m_c$ .

We perform the analysis of stationary states of (1.1)-(2.1). As in the linear potential case, we show that a critical mass appears only if  $m = m_c$ . Moreover, we show that stationary states are strictly related to optimizers of the following Hardy-Littlewood-Sobolev (HLS) type inequality, stating that if

$$m > (p_s^*)', \quad \text{where} \quad p_s^* := \frac{Np}{N-sp},$$

there exists a constant  $H > 0$  such that

$$(2.3) \quad \|K_{s/2} * h\|_{p'}^{p'} \leq H \|h\|_1^{p' \vartheta_0} \|h\|_m^{p'(1-\vartheta_0)} \quad \text{for every } h \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N),$$

where  $0 < \vartheta_0 < 1$  is given by

$$\vartheta_0 := 1 - \frac{m'}{p_s^*} = \frac{1}{(p_s^*)'} \frac{m - (p_s^*)'}{m-1}.$$

We shall prove existence and regularity properties of optimizers of (2.3), which will be shown to be solutions of the nonlocal equation

$$(2.4) \quad \rho^{m-1} = a (\mathcal{K}_{s,p}(\rho) - \mathcal{C})_+ \quad \text{in } \mathbb{R}^N$$

for suitable values of the positive constants  $a, \mathcal{C}$ . We notice that for  $p = 2$ , in terms of  $u := K_s * \rho$  the above equation becomes the fractional semilinear PDE

$$(-\Delta)^s u = a^{\frac{1}{m-1}} (u - \mathcal{C})_+^{\frac{1}{m-1}},$$

which is the fractional plasma equation investigated in [20]. The terminology for such a semilinear equation is due to the fact that the nonlinearity in the right hand side, where  $(x)_+ := \max\{x, 0\}$ , appears in some classical models of plasma physics [50, 51]. The following is our first main result, which provides the main properties of the HLS optimizers. In the case that  $m \geq m_c$ , these results can be translated in a statement about minimizers of the free energy  $\mathcal{F}_{s,p}$ . In this regard, a critical mass appears for  $m = m_c$ , given by

$$(2.5) \quad M_c := \left( \frac{p_s^*}{\chi H_{m_c, s, p}^*} \right)^{\frac{N}{sp'}},$$

where  $H_{m, s, p}^*$  is the best constant in (2.3).

**Theorem 2.1.** *Let  $1 < p < \infty$ ,  $0 < sp < N$ ,  $m > (p_s^*)'$ . The best constant in the HLS inequality (2.3) is attained. Each optimizer is radially nonincreasing (up to translation), compactly supported, Hölder regular in  $\mathbb{R}^N$  and smooth in the interior of its support. It satisfies (2.4) for suitable values of the constants  $a > 0, \mathcal{C} > 0$ .*

*If  $m > m_c$ , then for each optimizer  $h$  of the HLS inequality (2.3) there exists a unique scaling factor  $\lambda > 0$  such that  $h^\lambda$  is (up to translation) a minimizer of functional  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$ , where  $M = \|h\|_1$ ; conversely for every  $M > 0$  minimizers of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  exist and are optimizers of (2.3).*

*If  $m = m_c$ , then each optimizer  $h$  of the HLS inequality (2.3) having mass  $M_c$  is (up to translation) a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_{M_c}$  and  $\mathcal{F}_{s,p}(h) = 0$ ; conversely minimizers of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  exist if and only if  $M = M_c$  and are optimizers of (2.3).*

Existence of optimizers of (2.3) is a standard application of Riesz rearrangement inequalities along with compactness theorems for radially decreasing functions. For an optimizer, the constants  $a, \mathcal{C}$  can be explicitly expressed, as well as the optimal dilation factor  $\lambda$  in the case  $m > m_c$ , as seen through the proof. It is not difficult to check that for every  $M > 0$  the infimum of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  equals  $-\infty$  if  $(p_s^*)' < m < m_c$ . However, an optimizer of the HLS inequality is still satisfying (2.4), hence after a suitable mass invariant dilation it satisfies the Euler-Lagrange equation

$$(2.6) \quad \rho^{m-1} = \frac{m-1}{m} (\chi \mathcal{K}_{s,p}(\rho) - \mathcal{Q})_+ \quad \text{in } \mathbb{R}^N$$

associated with functional  $\mathcal{F}_{s,p}$ , where  $\mathcal{Q} > 0$  is a constant playing the role of Lagrange multiplier for the mass constraint. As such, it is (up to translation) a radially nonincreasing stationary state for (1.1)-(2.1) as we discuss in Section 5. About the regularity properties in Theorem 2.1, we mention that boundedness of optimizers has been proved in [19] by a purely variational argument in the case  $p = 2$ ,  $m > m_c$ , which consists in the construction of a suitable bounded competitor for every unbounded candidate. Such an argument seems not straightforward in the nonlinear potential setting, therefore we prove boundedness by classical bootstrap methods, based on HLS inequalities and on (2.4), that are working for every  $m > (p_s^*)'$ . We stress that Theorem 2.1 generalizes the previous results in the literature about inequality (2.3): in the case  $p = 2$  it is also called the Lane-Emden inequality and has been studied in [10, 15]. Interestingly, other generalizations have been recently investigated in [26], in relation with the Choquard equation, which still leads to radially decreasing compactly supported optimizers for suitable choices of the parameters therein.

**2.2. Asymptotic behavior of stationary states as  $s \rightarrow 0$ .** As observed in [28] by considering that  $K_s$  is an approximate identity for small  $s$ , the aggregation term can be considered as an approximation of a backward diffusion process, so that the evolution model (1.1)-(2.1) formally becomes the forward-backward diffusion equation

$$\partial_t \rho = \Delta \rho^m - \frac{\chi}{p'} \Delta \rho^{p'}$$

Similarly, the associated free energy  $\mathcal{F}_{s,p}$  formally becomes, in the limit  $s \rightarrow 0$ , the following functional featuring the competition of  $L^m$  and  $L^{p'}$  norms

$$(2.7) \quad \mathcal{F}_0(\rho) = \frac{1}{m-1} \int_{\mathbb{R}^N} \rho^m - \frac{\chi}{p'} \int_{\mathbb{R}^N} \rho^{p'}$$

Clearly, the minimization problem for functional  $\mathcal{F}_0$  in the class  $\mathcal{Y}_M$  is strongly influenced by the sign of  $m - p'$ , which is reflected in the fact that the critical exponent  $m_c$  from (2.2) is equal to  $p'$  if  $s = 0$ . If  $m < p'$ , then functional  $\mathcal{F}_{s,p}$  does not have minimizers over  $\mathcal{Y}_M$  for every small enough  $s$ , and moreover  $m < (p_s^*)'$  for every small enough  $s$ , so that we are not in the range of parameters of Theorem 2.1. Therefore, in our second main result, which is the following, we restrict to  $m \geq p'$ . The result for  $p = 2, m > 2$  is given in [28].

**Theorem 2.2.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m \geq p'$ ,  $M > 0$ . Let  $\rho_s$  be a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ .*

*If  $m > p'$ , then  $\rho_s \rightarrow \rho_0$  strongly in  $L^q(\mathbb{R}^N)$  for every  $1 < q < +\infty$  as  $s \rightarrow 0$ , where  $\rho_0$  is the unique radially decreasing minimizer of  $\mathcal{F}_0$  over  $\mathcal{Y}_M$ , which is the characteristic function of a ball.*

*Else if  $m = p'$ , we have  $\inf_{\mathcal{Y}_M} \mathcal{F}_s \rightarrow \inf_{\mathcal{Y}_M} \mathcal{F}_0$  as  $s \rightarrow 0$ . Moreover,  $\rho_s \rightarrow 0$  uniformly on  $\mathbb{R}^N$  if  $0 < \chi < p$ , and  $\rho_s \rightarrow M\delta_0$  in the sense of measures if  $\chi > p$ .*

Let us conclude this section with a discussion on possible further extensions and open problems. First of all, uniqueness (up to translations) of stationary states of given mass (or of optimizers of the HLS inequality (2.3) up to the natural scaling) would require a further, deep analysis. It has been proved in the case  $p = 2$  by different methods in [12, 15, 20, 21], and each of them could be suitable for treating the nonlinear potential case as well. The stability result of the HLS inequality in [15] could also be potentially generalized to  $p \neq 2$ . Second, radiality of every stationary solution to (1.1)-(2.1) is not guaranteed. Such a property has been proven in [19] in the linear potential case  $p = 2$  under some restrictions on  $m, s$  (building on the result from [18] for  $s = 1$ ). It remains an open problem to extend such result for the case  $p \neq 2$ . It would prove that all the steady states of the dynamics are actually radially decreasing. Moreover, it would also be interesting to investigate stationary states of the dynamics, meant as solutions to (2.6), in the regime  $1 < m \leq (p_s^*)'$ : in this range radially decreasing solutions are expected to exist only for  $\mathcal{Q} = 0$  and to be smooth, positive and vanishing at infinity, since this behavior has been proven for  $p = 2$  in [20].

### 3. PRELIMINARIES

**3.1. Notation and functional framework.** The dimension of the ambient space  $\mathbb{R}^N$  will be  $N \geq 1$ . For  $x_0 \in \mathbb{R}^N$  and  $r > 0$ , the symbol  $B_r(x_0)$  stands for the euclidean  $N$ -dimensional open ball

$$B_r(x_0) = \{x \in \mathbb{R}^N : |x - x_0| < r\}.$$

As usual, we will denote with  $|\cdot|$  the  $N$ -dimensional Lebesgue measure. For  $1 \leq p \leq \infty$ , the standard Lebesgue spaces are denoted by  $L^p_{\text{loc}}$  and  $L^p$ , and we will use the shortcut notation  $\|\cdot\|_p$  for the  $L^p(\mathbb{R}^N)$  norms. For an open set  $\Omega \subseteq \mathbb{R}^N$ , the notation  $W^{1,p}(\Omega)$  and  $BV(\Omega)$  stand respectively for the usual Sobolev space and the usual space of bounded variation functions on  $\Omega$ . We use the following notation for the Hölder spaces

$$C^{0,\alpha}(\Omega) := \left\{ u \in C^0(\Omega) \cap L^\infty(\Omega) : \sup_{\substack{x,y \in \Omega, \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} < \infty \right\}.$$

We say that  $u \in C^{0,\alpha}_{\text{loc}}(\Omega)$  if  $u \in C^{0,\alpha}(\Omega')$  for every open set  $\Omega'$  that is compactly contained in  $\Omega$ . In particular  $C^{0,1}(\Omega)$  is the space of bounded Lipschitz functions on  $\Omega$ .

For every  $\rho \in L^1(\mathbb{R}^N)$  and  $\lambda > 0$ , the *mass invariant dilation of  $\rho$  by factor  $\lambda$*  is given by (1.5). Since

$$\|\rho\|_1 = \|\rho^\lambda\|_1,$$

if  $\rho \in \mathcal{Y}_M$  then also  $\rho^\lambda \in \mathcal{Y}_M$ , for every  $\lambda > 0$ , where  $\mathcal{Y}_M$  is defined by (1.4).

With an abuse of notation, we will say that a radially symmetric function  $\rho \in L^1(\mathbb{R}^N)$  is nonincreasing if its radial profile is nonincreasing. The *radially symmetric nonincreasing rearrangement* of a function  $\rho \in L^1(\mathbb{R}^N)$  will be denoted by  $\rho^*$ . For the precise definition and its properties, we refer the reader to [35, Chapter 3]. We recall that the convolution among two nonnegative radially nonincreasing functions on  $\mathbb{R}^N$  is still radially nonincreasing on  $\mathbb{R}^N$ , see [13]. In particular, if  $f$  is radially nonincreasing nonnegative, so is  $K_s * f$ .

The Fourier transform of the Riesz kernel  $K_s$  defined in (1.3) is given by (see for example [47, Lemma 1, Chapter V] or also [39, Theorem 2.8] and [49, Proposition 12.10])

$$\widehat{K}_s(\xi) = |\xi|^{-2s}.$$

Moreover, for the normalization constant  $c_{N,s}$  in (1.3) we have the following limiting behavior

$$(3.1) \quad \lim_{s \rightarrow 0} \frac{c_{N,s}}{s} = \frac{\Gamma\left(\frac{N}{2}\right)}{\pi^{\frac{N}{2}}} = \frac{2}{N \omega_N}.$$

**3.2. Basics on Riesz potentials.** We now recall some facts we will need throughout the whole paper.

**Lemma 3.1.** *Let  $1 \leq q < r \leq \infty$ ,  $0 < sq < N$  and  $sr > N$ . For every  $h \in L^q(\mathbb{R}^N) \cap L^r(\mathbb{R}^N)$  we have*

$$\|K_{s/2} * h\|_\infty \leq \alpha_s \|h\|_q + \beta_s \|h\|_r,$$

for some positive constant  $\alpha_s = \alpha(N, q, s)$  and  $\beta_s = \beta(N, r, s) > 0$ . Moreover, we have

$$(3.2) \quad \lim_{s \rightarrow 0} \frac{\alpha_s}{s} = \begin{cases} \pi^{-\frac{N}{2}} \Gamma\left(\frac{N}{2}\right) (\omega_N (q-1))^{\frac{q-1}{q}} & \text{if } q > 1, \\ \pi^{-\frac{N}{2}} \Gamma\left(\frac{N}{2}\right) & \text{if } q = 1, \end{cases}$$

and

$$(3.3) \quad \lim_{s \rightarrow 0} \beta_s = 1 \quad \text{if } r = \infty.$$

*Proof.* Case  $q > 1$ . Our assumptions imply that

$$q' > \frac{N}{N-s} \quad \text{and} \quad r' < \frac{N}{N-s}.$$

Then, for every  $x \in \mathbb{R}^N$ , Hölder's inequality yields

$$(3.4) \quad \begin{aligned} \frac{(K_{s/2} * h)(x)}{c_{N,s/2}} &= \int_{\mathbb{R}^N \setminus B_1(x)} \frac{h(y)}{|x-y|^{N-s}} dy + \int_{B_1(x)} \frac{h(y)}{|x-y|^{N-s}} dy \\ &\leq \left( \int_{\mathbb{R}^N \setminus B_1} \frac{dy}{|y|^{(N-s)q'}} \right)^{\frac{1}{q'}} \|h\|_q + \left( \int_{B_1} \frac{dy}{|y|^{(N-s)r'}} \right)^{\frac{1}{r'}} \|h\|_r \\ &= \left( \frac{N \omega_N}{(N-s)q' - N} \right)^{\frac{1}{q'}} \|h\|_q + \left( \frac{N \omega_N}{N - (N-s)r'} \right)^{\frac{1}{r'}} \|h\|_r, \end{aligned}$$

which gives the desired conclusion with

$$\alpha_s = \alpha(N, s, q) := c_{N,s/2} \left( \frac{N\omega_N}{(N-s)q' - N} \right)^{\frac{1}{q'}},$$

$$\beta_s = \beta(N, s, r) := c_{N,s/2} \left( \frac{N\omega_N}{N - (N-s)r'} \right)^{\frac{1}{r'}}.$$

By recalling (3.1), we get the claimed asymptotic behaviors (3.2)-(3.3) for  $\alpha_s$  and  $\beta_s$ .

*Case  $q = 1$ .* We pass to the limit for  $q \searrow 1$  in (3.4) obtaining

$$\frac{(K_{s/2} * h)(x)}{c_{N,s/2}} \leq \|h\|_1 + \left( \frac{N\omega_N}{N - (N-s)r'} \right)^{\frac{1}{r'}} \|h\|_r.$$

This yields our claimed estimate with  $\alpha_s = c_{N,s/2}$  and  $\beta_s$  as before. By recalling (3.1), we eventually get the desired asymptotic behaviors.  $\square$

Next we introduce the *Hardy-Littlewood-Sobolev type inequalities* that are crucial in this work.

**Lemma 3.2** (Hardy-Littlewood-Sobolev type inequality). *Let  $1 < q < \infty$  and  $0 < sq < N$ . For every  $h \in L^q(\mathbb{R}^N)$ , we have*

$$K_{s/2} * h \in L^{q_s^*}(\mathbb{R}^N), \quad \text{where } q_s^* = \frac{Nq}{N-sq}.$$

*More precisely, there exists a sharp constant  $\bar{H}_s = \bar{H}(N, s, q) > 0$  such that*

$$(3.5) \quad \|K_{s/2} * h\|_{q_s^*} \leq \bar{H}_s \|h\|_q, \quad \text{with } \limsup_{s \rightarrow 0} \bar{H}_s \leq 1.$$

*In particular, for every  $h \in L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$ ,  $m > q$ , we have*

$$(3.6) \quad \|K_{s/2} * h\|_{q_s^*} \leq \bar{H}_s \|h\|_1^\vartheta \|h\|_m^{1-\vartheta}, \quad \text{where } \vartheta = \frac{1}{q} \frac{m-q}{m-1}.$$

*Proof.* Inequality (3.5) follows by using in duality the well-known *Hardy-Littlewood-Sobolev Inequality* [35, Theorem 4.3]. Indeed, with the notation therein used, if we plug the following

$$r := q, \quad \lambda := N - s \quad \text{and so} \quad p := (q_s^*)',$$

we get that

$$\int_{\mathbb{R}^N} \varphi (K_{s/2} * h) dx = c_{N,s/2} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\varphi(x) h(y)}{|x-y|^{N-s}} dx dy \leq C(N, q, s) c_{N,s/2} \|h\|_q,$$

for every  $\varphi \in L^{(q_s^*)'}(\mathbb{R}^N)$  with  $\|\varphi\|_{(q_s^*)'} = 1$ , which allows to conclude. The constant  $C(N, q, s)$  denotes the sharp constant of [35, Theorem 4.3(1)] and as shown therein we have

$$\begin{aligned} \bar{H}_s &= \bar{H}(N, q, s) := C(N, q, s) c_{N,s/2} \\ &\leq c_{N,s/2} \frac{N}{s} \omega_N^{(N-s)/N} \frac{1}{q(q_s^*)'} \left( \left( \frac{(N-s)/N}{1-1/(q_s^*)'} \right)^{(N-s)/N} + \left( \frac{(N-s)/N}{1-1/q} \right)^{(N-s)/N} \right). \end{aligned}$$

By (3.1), we infer that

$$\limsup_{s \rightarrow 0} \bar{H}_s \leq 1.$$

Eventually, by the interpolation inequality in  $L^p$ -spaces, we also get (3.6).  $\square$

For our purposes, it will be convenient to rewrite (3.5) and (3.6) with  $q$  replaced by  $(p_s^*)'$ , given that  $p$  is the nonlinear Riesz potential exponent appearing in functional  $\mathcal{F}_{s,p}$ . It reads as follows

**Corollary 3.3.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . We have*

$$(3.7) \quad \|K_{s/2} * h\|_{p'} \leq H_s \|h\|_{(p_s^*)'} \quad \text{for every } h \in L^{(p_s^*)'}(\mathbb{R}^N),$$

where the sharp constant  $H_s = H(N, s, p) > 0$  satisfies

$$(3.8) \quad H_s \leq c_{N,s/2} \frac{N}{s} \omega_N^{(N-s)/N} \frac{1}{p(p_s^*)'} \left( \left( \frac{(N-s)/N}{1-1/p} \right)^{(N-s)/N} + \left( \frac{(N-s)/N}{1-1/(p_s^*)'} \right)^{(N-s)/N} \right),$$

in particular  $\limsup_{s \rightarrow 0} H_s \leq 1$ . Moreover, if  $m > (p_s^*)'$  we have

$$(3.9) \quad \|K_{s/2} * h\|_{p'} \leq H_s \|h\|_1^{\vartheta_0} \|h\|_m^{1-\vartheta_0} \quad \text{for every } h \in L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N),$$

where  $0 < \vartheta_0 < 1$  is given by

$$(3.10) \quad \vartheta_0 = \vartheta_0(m, N, p, s) := \frac{1}{(p_s^*)'} \frac{m - (p_s^*)'}{m - 1} = 1 - \frac{m'}{p_s^*}.$$

*Proof.* We have

$$(p_s^*)' = \frac{Np}{N(p-1) + sp} \in (1, N/s),$$

thus the exponent  $q := (p_s^*)'$  satisfies the assumptions of Lemma 3.2. Since  $q_s^* = p'$ , inequality (3.6) can be rewritten as (3.7) where the sharp constant  $H_s = H(N, s, p) := \bar{H}(N, s, (p_s^*)') > 0$  satisfies (3.8). By the interpolation inequality in  $L^p$ -spaces we also get (3.9).  $\square$

The following theorem is due to Kurokawa. It will be used in Section 6 to establish convergence results for minimizers of  $\mathcal{F}_{s,p}$  as  $s \rightarrow 0$ , see Theorem 6.7 and Proposition 6.8. We recall its elegant proof below.

**Theorem 3.4** ([32]). *Let  $1 < q < p$ . For every  $h \in L^q(\mathbb{R}^N) \cap L^p(\mathbb{R}^N)$ , we have*

$$\lim_{s \rightarrow 0} \|K_{s/2} * h - h\|_p = 0.$$

*Proof.* Since  $h \in L^p(\mathbb{R}^N)$ , for every  $\varepsilon > 0$  we can find  $0 < \delta < 1$  such that

$$(3.11) \quad \int_{\mathbb{R}^N} |h(x-y) - h(x)|^p dx < \varepsilon, \quad \text{for } |y| < \delta,$$

see for instance [23, Proposition 17.1]. We set

$$K_{s/2}(x) = c_{N,s/2} |x|^{s-N} \mathbf{1}_{B_\delta}(x) + c_{N,s/2} |x|^{s-N} \mathbf{1}_{B_\delta^c}(x) =: K_{s/2}^0(x) + K_{s/2}^\infty(x),$$

so we have

$$(3.12) \quad \|K_{s/2} * h - h\|_p \leq \|K_{s/2}^0 * h - h\|_p + \|K_{s/2}^\infty * h\|_p.$$

For the first addendum, since

$$(3.13) \quad \int_{B_\delta} |y|^{s-N} dy = N \omega_N \int_0^\delta \varrho^{s-1} d\varrho = N \omega_N \frac{\delta^s}{s},$$

by adding and subtracting  $c_{N,s/2} \int_{B_\delta} |y|^{s-N} h(x) dy$ , we get

$$\begin{aligned} \|K_{s/2}^0 * h - h\|_p &\leq c_{N,s/2} \left( \int_{\mathbb{R}^N} \left| \int_{B_\delta} |y|^{s-N} (h(x-y) - h(x)) dy \right|^p dx \right)^{\frac{1}{p}} \\ &\quad + \left( c_{N,s/2} - \frac{s}{N\omega_N\delta^s} \right) \left( \int_{\mathbb{R}^N} \left| \int_{B_\delta} |y|^{s-N} h(x) dy \right|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

By Minkowski's integral inequality, (3.11) and again (3.13) we get

$$\|K_{s/2}^0 * h - h\|_p \leq N\omega_N \frac{\delta^s}{s} \left[ c_{N,s/2} \varepsilon + \left( c_{N,s/2} - \frac{s}{N\omega_N\delta^s} \right) \|h\|_p \right].$$

By (3.1), we then obtain

$$(3.14) \quad \limsup_{s \rightarrow 0} \|K_{s/2}^0 * h - h\|_p \leq \varepsilon.$$

To estimate the second term in the right hand side of (3.12) for small  $s$ , we take  $\beta, r$  so that

$$(3.15) \quad 0 < \beta < N \left( \frac{1}{q} - \frac{1}{p} \right) \quad \text{and} \quad r_\beta^* = p,$$

in particular  $q < r < p$ . Without loss of generality, we can assume that  $0 < s < \beta$  and observe that

$$|y|^{s-N} = |y|^{(s-\beta)+(\beta-N)} \leq \delta^{s-\beta} |y|^{\beta-N} \leq \delta^{-\beta} |y|^{\beta-N}, \quad \text{for } |y| \geq \delta,$$

being  $0 < \delta < 1$ . This entails that

$$\|K_{s/2}^\infty * h\|_p = c_{N,s/2} \left( \int_{\mathbb{R}^N} \left| \int_{B_\delta^c} |y|^{s-N} h(x-y) dy \right|^p dx \right)^{\frac{1}{p}} \leq c_{N,s/2} \delta^{-\beta} \|K_{\beta/2} * |h|\|_p.$$

Since  $h \in L^r(\mathbb{R}^N)$ , by (3.15) and by the Hardy-Littlewood-Sobolev inequality (3.5) we then obtain that  $\|K_{\beta/2} * |h|\|_p < \infty$  and so

$$\lim_{s \rightarrow 0} \|K_{s/2}^\infty * h\|_p = 0,$$

by (3.1). By spending this information and (3.14) in (3.12), we get the desired conclusion.  $\square$

#### 4. EXTREMALS OF THE HLS TYPE INEQUALITY

**4.1. Existence: the Lieb-Oxford method.** In this section, we discuss the existence of extremals of the Hardy-Littlewood-Sobolev type inequality (3.9).

Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > (p_s^*)'$  and  $\vartheta_0 = 1 - m'/p_s^*$ . The following quantity

$$(4.1) \quad H_{m,s}^* := \sup \left\{ \frac{\|K_{s/2} * h\|_{p'}^{p'}}{\|h\|_1^{p'\vartheta_0} \|h\|_m^{p'(1-\vartheta_0)}} : h \in L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N) \setminus \{0\} \right\},$$

is the sharp constant in the Hardy-Littlewood-Sobolev type inequality (3.9) raised to the power  $p'$ . By (3.8) this quantity is finite, since we have  $H_{m,s}^* \leq H_s^{p'}$ .

**Remark 4.1.** The quotient defining  $H_{m,s}^*$  given by

$$h \mapsto \frac{\|K_{s/2} * h\|_{p'}^{p'}}{\|h\|_1^{p'\vartheta_0} \|h\|_m^{p'(1-\vartheta_0)}}, \quad h \in L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N) \setminus \{0\},$$

is *invariant* under the actions of the two families of transformations

$$\mu \mapsto \mu h, \quad \text{for } \mu > 0, \quad \text{and} \quad \lambda \mapsto \lambda^N h(\lambda x), \quad \text{for } \lambda > 0.$$

This property will be crucially exploited in Lemma 4.2 below.

We now infer the existence of extremals for (4.1) by using a classical argument by Lieb and Oxford (see for instance [36, Appendix A] and [34, Theorem 2.5]).

**Lemma 4.2** (Existence of extremals). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > (p_s^*)'$ . There exists a radially symmetric and nonincreasing function  $h_s \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  realizing the supremum in (4.1) and satisfying  $\|h_s\|_1 = \|h_s\|_m = 1$ .*

*Moreover, every function  $h_s \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  attaining the supremum in (4.1) is such that  $h_s = h_s^*(\cdot - y)$ , for some  $y \in \mathbb{R}^N$ .*

**Proof. Step 1: reduction to normalized radially symmetric and nonincreasing functions.** Let  $(h_j)_{j \in \mathbb{N}}$  be a maximizing sequence of feasible competitors for  $H_{m,s}^*$ , i.e.

$$\lim_{j \rightarrow \infty} \frac{\|K_{s/2} * h_j\|_{p'}^{p'}}{\|h_j\|_1^{p'\vartheta_0} \|h_j\|_m^{p'(1-\vartheta_0)}} = H_{m,s}^*, \quad \text{with } h_j \in L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N) \setminus \{0\}, \quad \text{for } j \in \mathbb{N}.$$

We can assume that  $h_j \geq 0$ , since  $K_{s/2} * h_j \leq K_{s/2} * |h_j|$  pointwisely. We can further assume that

$$(4.2) \quad \|h_j\|_1 = \|h_j\|_m = 1, \quad \text{for } j \in \mathbb{N}.$$

This is not restrictive, since we could replace each approximant  $h_j$  with a rescaled version given by

$$\tilde{h}_j(x) = \lambda_j h_j(\mu_j x), \quad \text{with } \mu_j = \left( \frac{\|h_j\|_1}{\|h_j\|_m} \right)^{\frac{1}{N} \frac{m}{m-1}}, \quad \lambda_j = \frac{\mu_j^N}{\|h_j\|_1}, \quad \text{for } j \in \mathbb{N}.$$

Indeed, we have

$$\|K_{s/2} * \tilde{h}_j\|_{p'}^{p'} = \int_{\mathbb{R}^N} \left| \int_{\mathbb{R}^N} |x-y|^{s-N} \lambda_j h(\mu_j y) dy \right|^{p'} dx = \frac{\lambda^{p'}}{\mu^{s p' + N}} \|K_{s/2} * h_j\|_{p'}^{p'},$$

and

$$\|\tilde{h}_j\|_1 = \frac{\lambda}{\mu^N} \|h_j\|_1, \quad \|\tilde{h}_j\|_m = \frac{\lambda}{\mu^{\frac{N}{m}}} \|h_j\|_m,$$

for every  $j \in \mathbb{N}$ . This yields

$$\begin{aligned} \frac{\|K_{s/2} * \tilde{h}_j\|_{p'}^{p'}}{\|\tilde{h}_j\|_1^{p'\vartheta_0} \|\tilde{h}_j\|_m^{p'(1-\vartheta_0)}} &= \frac{\lambda_j^{p'}}{\lambda_j^{p'\vartheta_0} \lambda_j^{p'(1-\vartheta_0)}} \frac{\mu_j^{N p'\vartheta_0 + \frac{N}{m} p'(1-\vartheta_0)}}{\mu_j^{s p' + N}} \frac{\|K_{s/2} * h_j\|_{p'}^{p'}}{\|h_j\|_1^{p'\vartheta_0} \|h_j\|_m^{p'(1-\vartheta_0)}} \\ &= \frac{\|K_{s/2} * h_j\|_{p'}^{p'}}{\|h_j\|_1^{p'\vartheta_0} \|h_j\|_m^{p'(1-\vartheta_0)}}, \end{aligned}$$

for every  $j \in \mathbb{N}$ .

Eventually, we claim that it is not restrictive to assume that  $h_j$  is radially nonincreasing, for every  $j \in \mathbb{N}$ . Indeed, take  $\varphi_j \in L^p(\mathbb{R}^N)$  with  $\|\varphi_j\|_p = 1$  such that

$$\|K_{s/2} * h_j\|_{p'} = \int_{\mathbb{R}^N} \varphi_j (K_{s/2} * h_j) dx.$$

We denote with  $\varphi_j^*$  and  $h_j^*$  the radially symmetric nonincreasing rearrangement of  $\varphi_j$  and  $h_j$ , respectively. By using the Riesz's rearrangement inequality [35, Theorem 3.7], we get

$$\|K_{s/2} * h_j\|_{p'} = c_{N,s/2} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\varphi_j(x) h_j(y)}{|x-y|^{N-s}} dx dy \leq c_{N,s/2} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\varphi_j^*(x) h_j^*(y)}{|x-y|^{N-s}} dx dy.$$

Since  $\|\varphi_j^*\|_p = 1$ , passing to the supremum on the right hand side we obtain

$$\|K_{s/2} * h_j\|_{p'} \leq \|K_{s/2} * h_j^*\|_{p'},$$

which proves our claim.

**Step 2: the supremum is achieved.** Thanks to *Step 1* we can assume that  $h_j$  is a nonnegative radially symmetric nonincreasing function with  $\|h_j\|_1 = \|h_j\|_m = 1$ , for every  $j \in \mathbb{N}$ . By using spherical coordinates, by the monotonicity of  $h_j$ , we can infer that for every  $R > 0$  we have

$$1 = \|h_j\|_1 \geq N \omega_N \int_0^R \xi_j(r) r^{N-1} dr \geq N \omega_N \xi_j(R) \int_0^R r^{N-1} dr = \omega_N \xi_j(R) R^N,$$

where  $\xi_j(|x|) := h_j(x)$  is the one-dimensional radial profile of  $h_j$ , for every  $j \in \mathbb{N}$ . Similarly, we have

$$1 = \|h_j\|_m^m \geq \omega_N \xi_j(R)^m R^N,$$

that is

$$(4.3) \quad \sup_{(R,\infty)} |\xi_j| \leq \frac{1}{\omega_N} \min \left\{ \frac{1}{R^N}, \frac{1}{R^{N/m}} \right\} =: \omega(R), \quad \text{for every } R > 0, j \in \mathbb{N}.$$

Lebesgue's differentiation theorem for monotone functions (see [2, Corollary 3.29] for instance) entails that

$$\int_R^\infty |\xi_j'(r)| dr \leq \omega(R), \quad \text{for every } R > 0, j \in \mathbb{N}.$$

By means of a diagonal argument and *Helly's Selection Theorem* (see [23, Proposition 19.1c]), we can extract a subsequence (not relabeled) of nonincreasing functions  $(\xi_j)_j$  converging everywhere to a nonincreasing function  $\xi$  in  $(R, \infty)$ , for every rational number  $R > 0$ . This implies that

$$(4.4) \quad \lim_{j \rightarrow \infty} h_j(x) = \lim_{j \rightarrow \infty} \xi_j(|x|) = \xi(|x|) =: h_s(x), \quad \text{for every } x \in \mathbb{R}^N \setminus \{0\}.$$

By collecting the previous information, we infer that  $h_s$  is a radially nonincreasing function satisfying

$$(4.5) \quad 0 \leq h_s(x) \leq \omega(|x|) =: v(x), \quad \text{for every } x \in \mathbb{R}^N \setminus \{0\}.$$

Observe that

$$(4.6) \quad v \in L^q(\mathbb{R}^N), \quad \text{for every } 1 < q < m.$$

By using (4.2) and (4.4), Fatou's Lemma entails that

$$(4.7) \quad \|h_s\|_1 \leq 1 \quad \text{and} \quad \|h_s\|_m \leq 1,$$

thus in particular  $h_s \in L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$ . Moreover, we have

$$(4.8) \quad (K_{s/2} * h_s)(x) = \lim_{j \rightarrow \infty} (K_{s/2} * h_j)(x) \leq (K_{s/2} * v)(x), \quad \text{a.e. } x \in \mathbb{R}^N.$$

Indeed, from (4.3) we get the inequality in (4.8) and

$$(K_{s/2} * h_j)(x) = c_{N,s} \int_{\mathbb{R}^N} \frac{h_j(y)}{|x-y|^{N-s}} dy \leq c_{N,s} \int_{\mathbb{R}^N} \frac{v(y)}{|x-y|^{N-s}} dy = (K_{s/2} * v)(x).$$

From (4.6) and the properties of the Riesz potential (see [47, Theorem 1, Chapter V])

$$(K_{s/2} * v)(x) < \infty, \quad \text{a.e. } x \in \mathbb{R}^N.$$

Then, by using (4.4) and Lebesgue's Dominated Convergence Theorem, we obtain the equality in (4.8). Observe that, by using (4.6) and Corollary 3.3, we have  $K_{s/2} * v \in L^{p'}(\mathbb{R}^N)$ . By using Lebesgue's Dominated Convergence Theorem, by (4.8) and by recalling (4.2), we infer that

$$H_{m,s}^* = \lim_{j \rightarrow \infty} \|K_{s/2} * h_j\|_{p'}^{p'} = \|K_{s/2} * h_s\|_{p'}^{p'} \leq \frac{\|K_{s/2} * h_s\|_{p'}^{p'}}{\|h_s\|_1^{p' \vartheta_0} \|h_s\|_m^{p'(1-\vartheta_0)}},$$

where in the last inequality we also used (4.7). Then, the maximality of  $H_{m,s}^*$  among functions in  $L^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  entails that

$$\|h_s\|_1^{p' \vartheta_0} \|h_s\|_m^{p'(1-\vartheta_0)} = 1.$$

This combined with (4.7) gives that  $\|h_s\|_1 = \|h_s\|_m = 1$ .

To complete the proof, we are only left out to prove that every other nonnegative extremal of (4.1) must be radially nonincreasing up to translations. Assume that  $h_s \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N) \setminus \{0\}$  satisfies equality in (4.1). In light of Corollary 3.3 we know that  $K_{s/2} * h_s \in L^{p'}(\mathbb{R}^N)$ , thus we take  $\varphi \in L^p(\mathbb{R}^N)$  such that

$$\|K_{s/2} * h_s\|_{p'} = \int_{\mathbb{R}^N} \varphi(K_{s/2} * h_s) dx.$$

By using the Riesz's rearrangement inequality in strict form [35, Theorem 3.9], we get

$$\|K_{s/2} * h_s\|_{p'} = c_{N,s/2} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\varphi(x) h_s(y)}{|x-y|^{N-s}} dx dy \leq \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\varphi^*(x) h_s^*(y)}{|x-y|^{N-s}} dx dy$$

with equality holding only if  $\varphi = \varphi^*(\cdot - y)$  and  $h_s = h_s^*(\cdot - y)$ , for some  $y \in \mathbb{R}^N$ . This proves our claim and ends the proof.  $\square$

**Remark 4.3** (Extremals in  $\mathcal{Y}_M$ ). Under the assumptions of Lemma 4.2, we can infer that for every prescribed mass  $M > 0$  there exists a nonnegative radially symmetric nonincreasing function  $h_{s,M} \in \mathcal{Y}_M$  realizing the supremum in (4.1). More precisely, it is obtained as

$$h_{s,M} := M h_s,$$

where  $h_s$  is an extremal of (3.9) provided by Lemma 4.2 (thus satisfying  $\|h_s\|_1 = 1$ ) with barycenter at the origin.

**4.2. Euler-Lagrange equation.** The Euler-Lagrange equation satisfied by nonnegative extremals of (3.9) is derived below. We share arguments from [16, Theorem 3.1].

**Lemma 4.4.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > (p_s^*)'$ . For every extremal  $h_0 \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  of the HLS (3.9), we have*

$$(4.9) \quad \mathcal{A}_s h_0^{m-1} = \left( \mathcal{K}_{s,p}(h_0) - \mathcal{C}_s \right)_+ \quad \text{in } \mathbb{R}^N,$$

where

$$(4.10) \quad \mathcal{A}_s = \frac{m'}{p_s^*} \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_m^m}, \quad \mathcal{C}_s = \left( 1 - \frac{m'}{p_s^*} \right) \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_1} > 0.$$

*Proof.* We set

$$\mathcal{G}_s(h) := H_{m,s}^* \|h\|_1^{p' \vartheta_0} \|h\|_m^{p'(1-\vartheta_0)} - \|K_{s/2} * h\|_{p'}^{p'}, \quad \text{for } h \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N) \setminus \{0\},$$

where  $H_{m,s}^*$  and  $\vartheta_0$  are respectively given by (4.1) and (3.10). Let  $h_0$  be as in the statement. We shall make perturbations of  $h_0$  that preserve positivity. We take  $\varphi \in C_0^\infty(\mathbb{R}^N)$  and set

$$(4.11) \quad \psi := \varphi h_0, \quad \varepsilon_0 := \frac{1}{2\|\varphi\|_\infty}, \quad h_\varepsilon := h_0 + \varepsilon \psi \geq 0, \quad \text{for } 0 \leq |\varepsilon| < \varepsilon_0.$$

By the optimality of  $h_0$ , we have

$$\mathcal{G}_s(h_\varepsilon) \geq \mathcal{G}(h_0) = 0, \quad \text{for every } 0 \leq |\varepsilon| < \varepsilon_0,$$

this entails that

$$(4.12) \quad \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \mathcal{G}_s(h_\varepsilon) = 0.$$

We have

$$(4.13) \quad \mathcal{G}_s(h_\varepsilon) = H_{m,s}^* \|h_0 + \varepsilon \psi\|_1^{p' \vartheta_0} \|h_0 + \varepsilon \psi\|_m^{p'(1-\vartheta_0)} - \|K_{s/2} * (h_0 + \varepsilon \psi)\|_{p'}^{p'}.$$

We expand to the first order, with respect to the variable  $\varepsilon$ , the three integral terms appearing in the rightmost term. For the first one, it is clear that

$$\int_{\mathbb{R}^N} (h_0 + \varepsilon \psi) dx = \int_{\mathbb{R}^N} h_0 dx + \varepsilon \int_{\mathbb{R}^N} \psi dx, \quad \text{for } 0 < |\varepsilon| < \varepsilon_0.$$

For the second integral term, we have

$$(h_0 + \varepsilon \psi)^m - h_0^m = \varepsilon \int_0^1 m (h_0 + \varepsilon t \psi)^{m-1} \psi dt, \quad \text{a.e. in } \mathbb{R}^N, \quad \text{for } 0 < |\varepsilon| < \varepsilon_0.$$

By integrating over  $\mathbb{R}^N$ , dividing by  $\varepsilon$  and using Fubini theorem we obtain

$$(4.14) \quad \int_{\mathbb{R}^N} \frac{(h_0 + \varepsilon \psi)^m - h_0^m}{\varepsilon} dx = \int_0^1 \mathcal{H}_\varepsilon(t) dt,$$

where we set

$$\mathcal{H}_\varepsilon(t) = m \int_{\mathbb{R}^N} (h_0 + \varepsilon t \psi)^{m-1} \psi dx, \quad \text{for } t \in [0, 1].$$

By Hölder's inequality and (4.11), we infer that

$$|\mathcal{H}_\varepsilon(t)| \leq m \|h_0 + \varepsilon t \psi\|_{\frac{m}{m'}}^{\frac{m}{m'}} \|\psi\|_m \leq m (\|\psi\|_m + \varepsilon_0 \|\psi\|_m)^{\frac{m}{m'}} \|\psi\|_m, \quad \text{for } t \in [0, 1],$$

for every  $0 < |\varepsilon| < \varepsilon_0$ . By using Lebesgue's Dominated Convergence Theorem in (4.14)

$$\int_{\mathbb{R}^N} (h_0 + \varepsilon \psi)^m dx = \int_{\mathbb{R}^N} h_0^m dx + \varepsilon \int_{\mathbb{R}^N} m h_0^{m-1} \psi dx + o(\varepsilon), \quad \text{as } \varepsilon \rightarrow 0.$$

For the third integral term, we have a.e. in  $\mathbb{R}^N$  the following identity

$$(K_{s/2} * (h_0 + \varepsilon \psi))^{p'} - (K_{s/2} * h_0)^{p'} = \varepsilon p' \int_0^1 (K_{s/2} * (h_0 + \varepsilon t \psi))^{p'-1} (K_{s/2} * \psi) dt,$$

for  $0 < |\varepsilon| < \varepsilon_0$ . By integrating over  $\mathbb{R}^N$ , dividing by  $\varepsilon$  and by Fubini's theorem

$$(4.15) \quad \int_{\mathbb{R}^N} \frac{1}{\varepsilon} \left[ (K_{s/2} * (h_0 + \varepsilon \psi))^{p'} - (K_{s/2} * h_0)^{p'} \right] dx = \int_0^1 \mathcal{K}_\varepsilon(t) dt,$$

where we set

$$\mathcal{K}_\varepsilon(t) = p' \int_{\mathbb{R}^N} (K_{s/2} * (h_0 + \varepsilon t \psi))^{p'-1} (K_{s/2} * \psi) dx, \quad \text{for } t \in [0, 1].$$

By Hölder's inequality and the HLS-type inequality (3.9), we get

$$\begin{aligned} |\mathcal{K}_\varepsilon(t)| &\leq p' \|K_{s/2} * (h_0 + \varepsilon t \psi)\|_{p'}^{\frac{p'}{p}} \|K_{s/2} * \psi\|_{p'} \\ &\leq p' H_{m,s}^* \|h_0 + \varepsilon t \psi\|_1^{\frac{p'}{p} \vartheta_0} \|h_0 + \varepsilon t \psi\|_m^{\frac{p'}{p} (1-\vartheta_0)} \|\psi\|_1^{\vartheta_0} \|\psi\|_m^{1-\vartheta_0}, \end{aligned}$$

where  $H_{m,s}^*$  and  $\vartheta_0$  are respectively given by (4.1) and (3.10). By Minkowski's inequality, we further have

$$|\mathcal{K}_\varepsilon(t)| \leq p' H_{m,s}^* (\|h_0\|_1 + \varepsilon_0 \|\psi\|_1)^{\frac{p'}{p} \vartheta_0} (\|h_0\|_m + \varepsilon_0 \|\psi\|_m)^{\frac{p'}{p} (1-\vartheta_0)} \|\psi\|_1^{\vartheta_0} \|\psi\|_m^{1-\vartheta_0},$$

for  $t \in [0, 1]$ , for  $0 < |\varepsilon| < \varepsilon_0$ . Thus we can use Lebesgue's Dominated Convergence Theorem in (4.15), obtaining

$$\begin{aligned} \int_{\mathbb{R}^N} (K_{s/2} * (h_0 + \varepsilon \psi))^{p'} dx &= \int_{\mathbb{R}^N} (K_{s/2} * h_0)^{p'} dx + \varepsilon p' \int_{\mathbb{R}^N} (K_{s/2} * h_0)^{p'-1} (K_{s/2} * \psi) dx + o(\varepsilon) \\ &= \int_{\mathbb{R}^N} (K_{s/2} * h_0)^{p'} dx + \varepsilon p' \int_{\mathbb{R}^N} K_{s/2} * (K_{s/2} * h_0)^{p'-1} \psi dx + o(\varepsilon), \end{aligned}$$

as  $\varepsilon \rightarrow 0$ , where the last identity follows from Plancherel's theorem. By collecting the previous asymptotic expansions and by using that

$$H_{m,s}^* = \frac{\|K_{s/2} * h_0\|_{p'}^{\frac{p'}{p}}}{\|h_0\|_1^{\frac{p'}{p} \vartheta_0} \|h_0\|_m^{\frac{p'}{p} (1-\vartheta_0)}},$$

from (4.13) we get

$$\begin{aligned} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \mathcal{G}_s(h_\varepsilon) &= p' \vartheta_0 \frac{\|K_{s/2} * h_0\|_{p'}^{\frac{p'}{p}}}{\|h_0\|_1} \int_{\mathbb{R}^N} \psi dx + p' (1 - \vartheta_0) \frac{\|K_{s/2} * h_0\|_{p'}^{\frac{p'}{p}}}{\|h_0\|_m^m} \int_{\mathbb{R}^N} h_0^{m-1} \psi dx \\ &\quad - p' \int_{\mathbb{R}^N} K_{s/2} * (K_{s/2} * h_0)^{p'-1} \psi dx. \end{aligned}$$

By recalling (4.11) and (4.12), this entails that

$$\int_{\mathbb{R}^N} \left( p' \vartheta_0 \frac{\|K_{s/2} * h_0\|_{p'}^{\frac{p'}{p}}}{\|h_0\|_1} + p' (1 - \vartheta_0) \frac{\|K_{s/2} * h_0\|_{p'}^{\frac{p'}{p}}}{\|h_0\|_m^m} h_0^{m-1} - p' K_{s/2} * (K_{s/2} * h_0)^{p'-1} \right) \varphi h_0 dx = 0,$$

for every  $\varphi \in C_0^\infty(\mathbb{R}^N)$ . By the positivity of  $h_0$  on its support (which is either a ball or  $\mathbb{R}^N$ , as a consequence of Lemma 4.2) and by recalling the expression of  $\vartheta_0$  given by (3.10), we obtain

$$(4.16) \quad \frac{m'}{p_s^*} \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_m^m} h_0^{m-1} = \mathcal{K}_{s,p}(h_0) - \left(1 - \frac{m'}{p_s^*}\right) \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_1} \quad \text{in } \text{supp}(h_0).$$

In order to deduce a condition outside the support of  $h_0$ , we take any nonnegative function  $\varphi \in C_0^\infty(\mathbb{R}^N) \setminus \{0\}$  and set

$$h_\varepsilon := h_0 + \varepsilon \varphi, \quad \text{for } \varepsilon \geq 0.$$

Since  $\varphi \geq 0$  and  $\varepsilon \geq 0$ , we have  $h_\varepsilon \geq 0$  thus by minimality of  $h_0$  for  $\mathcal{G}$  we infer

$$\lim_{\varepsilon \rightarrow 0^+} \frac{\mathcal{G}(h_\varepsilon) - \mathcal{G}(h_0)}{\varepsilon} \geq 0.$$

By arguing as before, we get

$$\int_{\mathbb{R}^N} \left( p' \vartheta_0 \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_1} + p' (1 - \vartheta_0) \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_m^m} h_0^{m-1} - p' K_{s/2} * (K_{s/2} * h_0)^{p'-1} \right) \varphi \, dx \geq 0,$$

for every nonnegative function  $\varphi \in C_0^\infty(\mathbb{R}^N)$ . This entails that

$$(4.17) \quad p' \vartheta_0 \frac{\|K_{s/2} * h_0\|_{p'}^{p'}}{\|h_0\|_1} - p' K_{s/2} * (K_{s/2} * h_0)^{p'-1}(x) \geq 0 \quad \text{in } \text{supp}(h_0)^c.$$

The proof is thereby complete, in light of (4.16) and (4.17).  $\square$

**Remark 4.5.** We can express the constants  $\mathcal{A}_s$  and  $\mathcal{C}_s$  in (4.10) in terms of  $H_{m,s}^*$ : we have

$$\mathcal{A}_s = \frac{m'}{p_s^*} H_{m,s}^* \|h\|_m^{p'-m} \left( \frac{\|h\|_m}{\|h\|_1} \right)^{p' \left( \frac{m'}{p_s^*} - 1 \right)}, \quad \mathcal{C}_s = \left( 1 - \frac{m'}{p_s^*} \right) H_{m,s}^* \|h\|_1^{p'-1} \left( \frac{\|h\|_m}{\|h\|_1} \right)^{\frac{p' m'}{p_s^*}}.$$

**4.3. Regularity properties.** Next we show that any extremal of the HLS inequality (3.9) has compact support and it is bounded. We rely on a *bootstrap argument* based on the combination of the HLS inequality (3.5) and Lemma 3.1.

**Lemma 4.6** ( $L^\infty$ -bound and compactness of the support). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > (p_s^*)'$ . For every extremal  $h_s \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  of the HLS inequality (3.9), we have that  $h_s \in L^\infty(\mathbb{R}^N)$ . Moreover, the support of  $h_s$  is compact.*

*Proof.* We start by proving that  $\text{supp}(h_s)$  is compact. Recall that by Lemma 4.2, the function  $h_s$  is radially symmetric and nonincreasing, thus the support of  $h_s$  and its superlevel sets are either balls or  $\mathbb{R}^N$ . By contradiction, assume that  $\text{supp}(h_s) = \mathbb{R}^N$ . Our assumptions entail that  $h_s \in L^{(p_s^*)'}(\mathbb{R}^N)$  and so, by using twice the HLS inequality (3.5), we infer that  $\mathcal{K}_{s,p}(h_s) \in L^{p_s^*}(\mathbb{R}^N)$ . We claim that  $\lim_{|x| \rightarrow \infty} \mathcal{K}_{s,p}(h_s)(x) = 0$ . Indeed, since  $h_s$  is a nonnegative radially symmetric and nonincreasing function, so is the function  $\mathcal{K}_{s,p}(h_s)$ , see [13]. By spending the previous information, we get

$$(4.18) \quad \mathcal{K}_{s,p}(h_s)(x)^{p_s^*} \omega_N |x|^N \leq \int_{B_{|x|}} \mathcal{K}_{s,p}(h_s)(y)^{p_s^*} \, dy \leq \|K_{s/2} * h_s\|_{p_s^*}^{p_s^*},$$

yielding our claim. In particular, we infer that  $\{\mathcal{K}_{s,p}(h_s) > \mathcal{C}_s\} = B_{R_0}$  for some  $0 \leq R_0 < \infty$ , where  $\mathcal{C}_s > 0$  is given by (4.10). Then, by using (4.9), we get that  $h_s = 0$  on  $\mathbb{R}^N \setminus B_{R_0}$ , which is a contradiction.

We now prove that  $h_s \in L^\infty(\mathbb{R}^N)$ . We set  $m_1 := m$  and distinguish two cases according to whether  $s m_1 \geq N$  or  $s m_1 < N$ .

*Case 1:  $s m_1 \geq N$ .* Since  $h_s \in L^1(\mathbb{R}^N) \cap L^{m_1}(\mathbb{R}^N)$ , in particular  $h_s \in L^r(\mathbb{R}^N)$  for every  $1 < r < N/s$ . By Corollary 3.3, we infer that

$$K_{s/2} * h_s \in L^\theta(\mathbb{R}^N), \quad \text{for every } N/(N-s) < \theta < \infty,$$

and so

$$(K_{s/2} * h_s)^{p'-1} \in L^{\theta(p-1)}(\mathbb{R}^N), \quad \text{for every } N/(N-s) < \theta < \infty.$$

Since  $s p < N$ , we have

$$\lim_{\theta \rightarrow \frac{N}{N-s}} \theta(p-1) < N/s.$$

These considerations entail that

$$(K_{s/2} * h_s)^{p'-1} \in L^{\theta_1}(\mathbb{R}^N) \cap L^{\theta_2}(\mathbb{R}^N), \quad \text{for some } \theta_1 < N/s \text{ and } \theta_2 > N/s.$$

By Lemma 3.1, we then obtain  $K_{s/2} * (K_{s/2} * h_s)^{p'-1} \in L^\infty(\mathbb{R}^N)$ . In turn, from the Euler-Lagrange equation Lemma 4.4, we conclude that  $h_s \in L^\infty(\mathbb{R}^N)$ .

*Case 2:  $s m_1 < N$ .* Since  $h_s \in L^1(\mathbb{R}^N) \cap L^{m_1}(\mathbb{R}^N)$ , from the HLS inequality (3.5), we infer that  $K_{s/2} * h_s \in L^r(\mathbb{R}^N) \cap L^{(m_1)_s^*}(\mathbb{R}^N)$ , for every  $N/(N-s) < r < (m_1)_s^*$ . This entails that

$$(K_{s/2} * h_s)^{p'-1} \in L^{r(p-1)}(\mathbb{R}^N) \cap L^{(m_1)_s^*(p-1)}(\mathbb{R}^N), \quad \text{for every } N/(N-s) < r < (m_1)_s^*.$$

If  $(m_1)_s^*(p-1) \geq N/s$ , by arguing as in *Case 1*, we conclude that  $h_s \in L^\infty(\mathbb{R}^N)$  and we stop. If otherwise  $(m_1)_s^*(p-1) < N/s$ , by the HLS inequality (3.5) we infer

$$K_{s/2} * (K_{s/2} * h_s)^{p'-1} \in L^{((m_1)_s^*(p-1))_s^*}(\mathbb{R}^N).$$

In turn, from the Euler-Lagrange equation, this implies that

$$h_s \in L^{m_2}(\mathbb{R}^N),$$

where we have set

$$m_2 := (m-1) ((m_1)_s^*(p-1))_s^* = \frac{N(m-1)(p-1)m_1}{N-s p m_1},$$

and observe that  $m_2 > m_1$ , being this condition equivalent to  $m > (p_s^*)'$ . In general, let  $k \in \mathbb{N} \setminus \{0\}$  and assume that  $m_i < N/s$  for every  $1 \leq i \leq k$ . We define

$$(4.19) \quad m_{k+1} := (m-1) ((m_k)_s^*(p-1))_s^* = \frac{N(m-1)(p-1)m_k}{N-s p m_k}.$$

We want to prove by induction that  $m_{k+1} > m_k$ . Our inductive assumption reads as

$$(4.20) \quad m_{i+1} > m_i, \quad \text{for every } 1 \leq i \leq k-1.$$

Since  $m > (p_s^*)'$ , we have

$$m > \frac{(N-s m) p}{N(p-1)} = \frac{(N-s m_1) p}{N(p-1)} > \frac{(N-s m_k) p}{N(p-1)},$$

where in the last inequality we used (4.20). In particular

$$m > \frac{(N - s m_k) p}{N(p-1)} \iff m_{k+1} > m_k,$$

as we can infer by recalling (4.19). We now claim that

$$(4.21) \quad \lim_{k \rightarrow \infty} m_k = +\infty.$$

This would entail that for some  $\bar{k}$  we must have  $m_{\bar{k}} \geq N/s$ , thus, by arguing as in *Case 1*, this would also end the proof. Since  $m_k \geq m$ , we have

$$\frac{m_{k+1}}{m_k} = \frac{N(m-1)(p-1)}{N - s p m_k} \geq \frac{N(m-1)(p-1)}{N - s p m} = \frac{m_2}{m_1}, \quad \text{for every } k \geq 1.$$

Moreover, as we have already observed

$$\frac{m_2}{m_1} > 1 \iff m > (p_s^*)'.$$

The last two facts entail that

$$\liminf_{k \rightarrow \infty} \frac{m_{k+1}}{m_k} > 1,$$

and so our claim (4.21).  $\square$

In the next lemma, we can readily adapt the argument of [19, Theorem 8] to infer Hölder regularity for extremals of (3.9).

**Lemma 4.7** (Hölder regularity). *Let  $1 < p < \infty$  and  $0 < s p < N$ . Let  $m > (p_s^*)'$ . For every extremal  $h_s \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  of the HLS inequality (3.9), we have*

- for  $0 < s < 1/2$

$$\begin{cases} h_s \in C^{0,1}(\mathbb{R}^N), & \text{if } m \leq 2, \\ h_s \in C^{0, \frac{1}{m-1}}(\mathbb{R}^N), & \text{if } 2 < m < m^*, \\ h_s \in C^{0,\gamma}(\mathbb{R}^N), & \text{if } m^* \leq m, \end{cases}$$

$$\text{where } m^* = \frac{2-2s}{1-2s} \text{ and } \gamma \in \left(0, \frac{2s}{m-2}\right).$$

- for  $s \geq 1/2$

$$h_s \in C^{0,\gamma}(\mathbb{R}^N), \quad \text{where } \gamma = \min \left\{ 1, \frac{1}{m-1} \right\}.$$

Moreover,  $h_s$  has  $C^\infty$ -regularity in the interior of its support.

*Proof.* First, we assume  $0 < s < 1/2$ . We will take advantage of the embeddings between Bessel potential spaces, fractional Sobolev spaces and Hölder spaces. We briefly recall that for  $1 \leq q < \infty$  the fractional Sobolev space  $W^{s,q}(\mathbb{R}^N)$  is given by

$$W^{s,q}(\mathbb{R}^N) = \left\{ u \in L^q(\mathbb{R}^N) : [u]_{W^{s,q}(\mathbb{R}^N)} < \infty \right\},$$

where  $[\cdot]_{W^{s,q}(\mathbb{R}^N)}$  denotes the Gagliardo-Slobodeckii seminorm

$$[u]_{W^{s,q}(\mathbb{R}^N)} := \left( \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^q}{|x - y|^{N+sq}} dx dy \right)^{\frac{1}{q}}.$$

The Bessel potential spaces  $\mathcal{L}^{s,q}(\mathbb{R}^N)$ , where  $1 < q < \infty$ , are defined through the Fourier transform, see for instance [47, Section V.3], [52, Section 2.2.2], [45, Section 27.3]. They can be characterized as

$$\mathcal{L}^{s,q}(\mathbb{R}^N) = \{u \in L^q(\mathbb{R}^N) : u = K_{s/2} * h, \text{ for some } h \in L^q(\mathbb{R}^N)\},$$

see for example [46, Theorem 2] (or also [45, Theorem 26.8, Theorem 27.3]). By recalling [47, Theorem 5, pag. 155] and [22, Theorem 4.47], the following continuous embeddings holds true:

$$(4.22) \quad \mathcal{L}^{s,q}(\mathbb{R}^N) \hookrightarrow W^{s,q}(\mathbb{R}^N), \quad \text{for } q \geq 2,$$

and

$$(4.23) \quad W^{s,q}(\mathbb{R}^N) \hookrightarrow C^{0,\gamma}(\mathbb{R}^N), \quad \text{where } \gamma = s - N/q, \text{ for } q > N/s.$$

Let  $h_s$  be as in the statement. By Lemma 4.2, we can assume it is radially nonincreasing. By Lemma 3.1, Lemma 4.4 and Lemma 4.6, we have that

$$(4.24) \quad h_s \in L^1(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N) \quad \text{and} \quad K_{s/2} * h_s \in L^q(\mathbb{R}^N), \quad \text{for every } N/(N-s) < q \leq \infty.$$

In particular, since  $N/(N-s) < N/s$  for every  $0 < s < 1/2$ , we have that

$$K_{s/2} * h_s \in \mathcal{L}^{s,q}(\mathbb{R}^N), \quad \text{for every } q > N/s.$$

In view of the embeddings (4.22) and (4.23), this entails that

$$K_{s/2} * h_s \in C^{0,\gamma}(\mathbb{R}^N), \quad \text{where } \gamma = s - N/q, \text{ for } q > N/s.$$

Since  $h_s$  is a nonnegative radially nonincreasing function, so is of  $K_{s/2} * h_s$  (see [13]). By reasoning as in (4.18), we then infer that  $\lim_{|x| \rightarrow \infty} (K_{s/2} * h_s)(x) = 0$ . Moreover, the function  $K_{s/2} * h_s$  attains only strictly positive values, and so it is bounded away from zero on compact sets. For these reasons we get  $(K_{s/2} * h_s)^{p'-1} \in C_{\text{loc}}^{0,\gamma}(\mathbb{R}^N)$ . Then, by taking into account (4.24) and the identity

$$(-\Delta)^{s/2} \mathcal{K}_{s,p}(h_s) = (K_{s/2} * h_s)^{p'-1} \quad \text{in } \mathbb{R}^N,$$

we are in position to apply [44, Corollary 3.5]. This yields the Hölder estimate

$$(4.25) \quad \|\mathcal{K}_{s,p}(h_s)\|_{C^{0,\gamma+s}(B_{1/2})} \leq c \left( \|\mathcal{K}_{s,p}(h_s)\|_{L^\infty(\mathbb{R}^N)} + \|(K_{s/2} * h_s)^{p'-1}\|_{C^{0,\gamma}(B_1)} \right),$$

for a constant  $c = c(N, s) > 0$  (notice that since  $s < 1/2$ , then  $\gamma + s$  is not an integer as  $\gamma = s - N/q$ ). By suitably rescaling the last inequality, we get

$$\begin{aligned} & \|\mathcal{K}_{s,p}(h_s)\|_{L^\infty(B_{R/2})} + R^{\gamma+s} [\mathcal{K}_{s,p}(h_s)]_{C^{0,\gamma+s}(B_{R/2})} \\ & \leq c \left( \|\mathcal{K}_{s,p}(h_s)\|_{L^\infty(\mathbb{R}^N)} + R^s \|(K_{s/2} * h_s)^{p'-1}\|_{L^\infty(B_R)} + R^{\gamma+s} [(K_{s/2} * h_s)^{p'-1}]_{C^{0,\gamma}(B_R)} \right), \end{aligned}$$

for every  $R > 0$ . In particular, this entails

$$(4.26) \quad \|\mathcal{K}_{s,p}(h_s)\|_{C^{0,\gamma+s}(B_{R/2})} \leq C \left( \|\mathcal{K}_{s,p}(h_s)\|_{L^\infty(\mathbb{R}^N)} + \|(K_{s/2} * h_s)^{p'-1}\|_{C^{0,\gamma}(B_R)} \right),$$

for a suitable constant  $C = C(N, s, R, \gamma) > 0$ , and so

$$\mathcal{K}_{s,p}(h_s) \in C^{0,\gamma+s}(B_{R/2}), \quad \text{where } \gamma = s - N/q, \text{ for } q > N/s.$$

From the Euler-Lagrange equation provided by Lemma 4.4, this entails that

$$h_s^{m-1} \in C^{0,\gamma+s}(B_{R/2}), \quad \text{for every } R > 0.$$

By using the fact that  $h_s$  has compact support, we infer

$$(4.27) \quad h_s \in C^{0,(\gamma+s)\alpha}(\mathbb{R}^N), \quad \text{where } \alpha = \min \left\{ 1, \frac{1}{m-1} \right\} \text{ and } \gamma = s - N/q, \text{ for } q > N/s.$$

Now we distinguish three cases.

*Case 1:*  $(p_s^*)' < m \leq 2$ . By using (4.24) and (4.27), from [44, Corollary 3.5] we have that

$$K_{s/2} * h_s \in C_{\text{loc}}^{0,\gamma+2s}(\mathbb{R}^N),$$

if  $\gamma+2s$  is not an integer. Since  $K_{s/2} * h_s$  is bounded and bounded away from zero on compact sets, as before we deduce

$$(K_{s/2} * h_s)^{p'-1} \in C_{\text{loc}}^{0,\gamma+2s}(\mathbb{R}^N).$$

If  $\gamma+2s > 1$ , we get  $(K_{s/2} * h_s)^{p'-1} \in C_{\text{loc}}^{0,1}(\mathbb{R}^N)$  and so also  $\mathcal{K}_{s,p}(h_s) \in C_{\text{loc}}^{0,1}(\mathbb{R}^N)$ . Thus, in light of Lemma 4.4, using that  $m \leq 2$  and the compactness of the support of  $h_s$ , we get  $h_s \in C^{0,1}(\mathbb{R}^N)$  as desired. On the other hand, if  $\gamma+2s < 1$  we newly apply [44, Corollary 3.5] and (4.26) obtaining that

$$\mathcal{K}_{s,p}(h_s) \in C_{\text{loc}}^{0,\gamma+3s}(\mathbb{R}^N),$$

if  $\gamma+3s$  is not an integer. Observe that we gained  $2s$  derivatives starting from (4.27), and the gain in regularity depends therefore on  $s$  but not on  $p$ . In other words, the regularity gain provided by the nonlinear potential  $\mathcal{K}_{s,p}$  does not depend on  $p$  and it is the same of the linear potential  $\mathcal{K}_{s,2}$ . Thus, the proof gets reduced to the case  $p = 2$  which is given in [19, Theorem 8]. For this reason, we just sketch the conclusion of the argument, omitting some details. We take an integer  $j \geq 1$  such that

$$(4.28) \quad \frac{1}{2(j+1)} < s < \frac{1}{2j},$$

and set  $\gamma_j := \gamma + (j-1)2s = 2sj - N/q$ , where  $q > N/s$  is chosen large enough so that  $1-2s < \gamma_j < 1$ . This is a feasible choice thanks to (4.28). By iterating the previous argument  $j$  times starting from (4.27), we get  $\mathcal{K}_{s,p}(h_s) \in C_{\text{loc}}^{0,\gamma_j+2s}(\mathbb{R}^N) \subseteq C_{\text{loc}}^{0,1}(\mathbb{R}^N)$ , being  $\gamma_j+2s > 1$  by construction. By using the Euler-Lagrange equation provided by Lemma 4.4 and the compactness of  $\text{supp}(h_s)$  from Lemma 4.6 and since  $m \leq 2$ , we conclude that  $h_s \in C^{0,1}(\mathbb{R}^N)$ .

*Case 2:*  $2 < m < m^*$ . Starting from (4.27), we can improve the Hölder regularity of  $h_s$  by a bootstrap argument, as in the previous case. We give the details of one iteration, in order to clarify that the same argument used in the proof of [19, Theorem 8] still holds. In light of (4.24) and (4.27), we can apply [44, Corollary 3.5] to infer that

$$(K_{s/2} * h_s)^{p'-1} \in C_{\text{loc}}^{0,\frac{\gamma+s}{m-1}+s}(\mathbb{R}^N), \quad \text{where } \gamma = s - N/q, \text{ for } q > N/s,$$

if  $(\gamma+s)/(m-1)+s$  is not an integer. By newly applying [44, Corollary 3.5], we obtain

$$\mathcal{K}_{s,p}(h_s) \in C_{\text{loc}}^{0,\frac{\gamma+s}{m-1}+2s}(\mathbb{R}^N),$$

if  $(\gamma+s)/(m-1)+2s$  is not an integer. By reasoning as in the previous case, if  $\gamma+2s/(m-1)+2s > 1$ , we have  $h_s^{m-1} \in C^{0,1}(\mathbb{R}^N)$  and so  $h_s \in C^{0,\frac{1}{m-1}}(\mathbb{R}^N)$ . On the other hand, if  $\gamma+2s/(m-1)+2s < 1$ , by always using the Euler-Lagrange equation provided by Lemma

4.4 and the fact that  $h_s$  has compact support, we obtain  $h_s^{m-1} \in C^{0, \frac{\gamma+s}{m-1} + 2s}(\mathbb{R}^N)$ , which entails that

$$h_s \in C^{0, \gamma_1}(\mathbb{R}^N), \quad \text{where } \gamma_1 = \frac{\gamma + s}{(m-1)^2} + \frac{2s}{m-1},$$

if  $(\gamma + s)/(m-1) + 2s$  is not an integer, where  $\gamma = s - N/q$ , for  $q > N/s$ . In general, by iterating this argument following [19, Theorem 8], we can improve the Hölder regularity of  $h_s$  to infer that  $h_s^{m-1} \in C^{0,1}(\mathbb{R}^N)$ , which yields  $h_s \in C^{0, \frac{1}{m-1}}(\mathbb{R}^N)$ , as desired.

*Case 3:  $m \geq m^*$ .* We can proceed with the same bootstrap argument, however without reaching Lipschitz regularity of  $h_s^{m-1}$ . We observe that [19, Remark 2], to which we refer, still holds and gives the desired result.

In order to conclude, we observe that the case  $1/2 \leq s$  is simpler than the case  $0 < s < 1/2$  and can be treated in the same way, up to some minor modifications, as done in [19, Theorem 8]. Eventually, by using the same argument of [19, Theorem 10], from Lemma 4.4 and Lemma 4.6, we obtain the  $C^\infty$ -regularity of  $h_s$  in the interior of its support.  $\square$

We end this section by remarking that the extremals of (4.1) are always in  $W^{1,1}(\mathbb{R}^N)$ :

**Corollary 4.8.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > (p_s^*)'$ . For every function  $h_s \in L_+^1(\mathbb{R}^N) \cap L^m(\mathbb{R}^N)$  attaining the supremum in (4.1), we have  $h_s \in W^{1,1}(\mathbb{R}^N)$ .*

*Proof.* The desired conclusion follows from Lemma 4.2, Lemma 4.4 and Lemma 4.6 by arguing as in [28, Proposition 2.10].  $\square$

## 5. MINIMIZERS OF THE ENERGY FUNCTIONAL

In this section, we analyze minimizers of functional  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  thus concluding the proof of Theorem 2.1. We start by considering the *diffusion dominated regime*, that is the case  $m > m_c$ .

**Proposition 5.1** (Diffusion dominated regime). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > m_c$  and let  $\chi, M > 0$ . Define the functional*

$$\mathcal{Y}_M \ni \rho \mapsto \Lambda(\rho) := \left( \frac{\|K_{s/2} * \rho\|_{p'}^{p'(m-1)}}{\|\rho\|_m^{m(m_c-1)}} \right)^{\frac{1}{m-m_c}},$$

which is invariant by mass-invariant dilations. Let moreover

$$\kappa := \left( \frac{\chi}{p'} \right)^{\frac{m-1}{m-m_c}} \left( \frac{p'}{p_s^*} \right)^{\frac{m_c-1}{m-m_c}} \left( \frac{m_c - m}{m-1} \right).$$

For every  $\rho \in \mathcal{Y}_M$ , there exists a unique positive number  $\lambda_*(\rho)$ , called the *optimal dilation factor* of  $\rho$ , such that

$$\mathcal{F}_{s,p}(\rho^\lambda) \geq \mathcal{F}_{s,p}(\rho^{\lambda_*(\rho)}) = \kappa \Lambda(\rho) \quad \text{for every } \lambda > 0, \quad \text{with equality only if } \lambda = \lambda_*(\rho).$$

It is expressed as

$$(5.1) \quad \lambda_*(\rho) = \left( \frac{\chi}{p_s^*} \frac{\|K_{s/2} * \rho\|_{p'}^{p'}}{\|\rho\|_m^m} \right)^{\frac{1}{N(m-m_c)}}.$$

*Proof.* Let  $\rho \in \mathcal{Y}_M$ . For  $\lambda > 0$ , we consider the function given by

$$(5.2) \quad \lambda \mapsto f_\rho(\lambda) := \mathcal{F}_{s,p}(\rho^\lambda) = \frac{\lambda^{N(m-1)}}{m-1} \|\rho\|_m^m - \lambda^{N(m_c-1)} \frac{\chi}{p'} \|K_{s/2} * \rho\|_{p'}^{p'}.$$

Recall that, since  $1 < p < N/s$ , we have  $N/(N-s) < p' < \infty$ , thus  $m_c = p'(1 - s/N) > 1$ . By optimizing with respect to  $\lambda$ , we get

$$(5.3) \quad \frac{d}{d\lambda} \mathcal{F}(\rho^\lambda) = N \lambda^{N(m-1)-1} \|\rho\|_m^m - \frac{N(m_c-1)}{p'} \chi \lambda^{N(m_c-1)-1} \|K_{s/2} * \rho\|_{p'}^{p'} = 0.$$

The unique extremal is given by (5.1). Clearly, at  $\lambda_*(\rho)$ , the function given by (5.2) attains a global minimum, and notice also that we have

$$\lim_{\lambda \rightarrow +\infty} \mathcal{F}_{s,p}(\rho^\lambda) = +\infty \quad \text{and} \quad \lim_{\lambda \rightarrow 0} \mathcal{F}_{s,p}(\rho^\lambda) = 0.$$

Furthermore, by using (5.1) we can write

$$\begin{aligned} \mathcal{F}_{s,p}(\rho^{\lambda_*(\rho)}) &= \frac{1}{m-1} \left( \frac{\chi}{p_s^*} \frac{\|K_{s/2} * \rho\|_{p'}^{p'}}{\|\rho\|_m^m} \right)^{\frac{m-1}{m-m_c}} \|\rho\|_m^m - \frac{\chi}{p'} \left( \frac{\chi}{p_s^*} \frac{\|K_{s/2} * \rho\|_{p'}^{p'}}{\|\rho\|_m^m} \right)^{\frac{m_c-1}{m-m_c}} \|K_{s/2} * \rho\|_{p'}^{p'} \\ &= \kappa \Lambda(\rho) < 0, \end{aligned}$$

where  $\Lambda$  and  $\kappa$  are defined as in the statement.  $\square$

For the *fair competition regime*, that is  $m = m_c$ , we have the following two sided-estimate for the energy, which extends [6, Proposition 3.4].

**Proposition 5.2** (Fair competition regime). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $\chi, M > 0$ . For every  $\rho \in \mathcal{Y}_M$ , we have*

$$\frac{\chi}{p'} H_{m_c,s}^* \left( M_c^{p' \frac{s}{N}} - M^{p' \frac{s}{N}} \right) \|\rho\|_{m_c}^{m_c} \leq \mathcal{F}_{s,p}(\rho) \leq \frac{\chi}{p'} H_{m_c,s}^* \left( M_c^{p' \frac{s}{N}} + M^{p' \frac{s}{N}} \right) \|\rho\|_{m_c}^{m_c},$$

where  $H_{m_c,s}^*$  is given by (4.1) and  $M_c$  is given by (2.5).

*Proof.* For every  $\rho \in \mathcal{Y}_M$ , by recalling (4.1), we get

$$\begin{aligned} \mathcal{F}_{s,p}(\rho) &= \frac{\|\rho\|_{m_c}^{m_c}}{m_c-1} - \frac{\chi}{p'} \|K_{s/2} * \rho\|_{p'}^{p'} \geq \left( \frac{1}{m_c-1} - \frac{\chi}{p'} H_{m_c,s}^* M^{p' \frac{s}{N}} \right) \|\rho\|_{m_c}^{m_c} \\ &= \frac{\chi}{p'} H_{m_c,s}^* \left( M_c^{p' \frac{s}{N}} - M^{p' \frac{s}{N}} \right) \|\rho\|_{m_c}^{m_c}, \end{aligned}$$

where  $M_c$  is given by (2.5). On the other hand, by recalling (2.2), we also have

$$\mathcal{F}_{s,p}(\rho) \leq \frac{\chi}{p'} H_{m_c,s}^* \left( M_c^{p' \frac{s}{N}} + M^{p' \frac{s}{N}} \right) \|\rho\|_{m_c}^{m_c},$$

which yields the claimed estimate.  $\square$

We can now recognize the infimum of the energy functional  $\mathcal{F}_{s,p}$  on  $\mathcal{Y}_M$ , corresponding to different values of the diffusion exponent  $m$ . This is the content of the following proposition.

**Proposition 5.3** (Infimum of  $\mathcal{F}_{s,p}$ ). *Let  $1 < p < \infty$ ,  $0 < sp < N$  and let  $\chi > 0$ . For every  $M > 0$ , we have*

$$\inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_{s,p}(\rho) = \begin{cases} -\infty, & \text{if } 1 < m < m_c, \\ \nu_s, & \text{if } m = m_c, \\ \mu_s, & \text{if } m > m_c, \end{cases}$$

for some  $\mu_s = \mu(N, p, s, \chi, m, M) < 0$ , where  $\nu_s = \nu_s(N, p, s, \chi, M)$  is given by

$$\nu_s = \begin{cases} 0, & \text{if } 0 < M \leq M_c, \\ -\infty, & \text{if } M > M_c, \end{cases}$$

being  $M_c$  the critical mass introduced in (2.5).

*Proof.* As in the beginning of the proof of Proposition 5.1, we take  $\rho \in \mathcal{Y}_M$  and consider the function  $f_\rho(\lambda) := \mathcal{F}_{s,p}(\rho^\lambda)$ ,  $\lambda > 0$ , whose expression is given by (5.2) for every  $m > 1$ .

If  $1 < m < m_c$ , by sending  $\lambda \nearrow \infty$  we infer that, for every  $M > 0$ ,

$$\inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_{s,p}(\rho) = \lim_{\lambda \rightarrow \infty} f_\rho(\lambda) = -\infty.$$

If  $m > m_c$ , from Proposition 5.1 and by recalling (2.2) and (3.10), we get

$$(5.4) \quad \Lambda(\rho)^{\frac{m-m_c}{m-1}} = \frac{\|K_{s/2} * \rho\|_{p'}^{p'}}{\|\rho\|_m^{\frac{m-p'}{m-1} \frac{p'}{s}}} = \frac{\|K_{s/2} * \rho\|_{p'}^{p'}}{\|\rho\|_m^{p'(1-\vartheta_0)}} = M^{p'\vartheta_0} \left( \frac{\|K_{s/2} * \rho\|_{p'}^{p'}}{M^{p'\vartheta_0} \|\rho\|_m^{p'(1-\vartheta_0)}} \right).$$

Since  $m > m_c > (p_s^*)'$ , the Hardy-Littlewood-Sobolev type inequality (3.9) entails therefore that

$$\sup_{\rho \in \mathcal{Y}_M} \Lambda(\rho) \in (0, +\infty)$$

for every  $M > 0$ . In turn, by Proposition 5.1

$$\inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_{s,p}(\rho) = \inf_{\rho \in \mathcal{Y}_M} (\kappa \Lambda(\rho)) = \kappa \sup_{\rho \in \mathcal{Y}_M} \Lambda(\rho)$$

since  $\kappa < 0$ , thus

$$\mu_s = \mu(N, p, s, \chi, m, M) := \inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_{s,p}(\rho) \in (-\infty, 0).$$

If  $m = m_c$ , we need to distinguish two cases.

*Case  $0 < M \leq M_c$ .* We take  $\rho \in \mathcal{Y}_M$  and test the energy  $\mathcal{F}_{s,p}$  with its mass invariant dilations  $\rho^\lambda$  given by (1.5). By the change of variable formula, we have

$$\|\rho^\lambda\|_{m_c}^{m_c} = \lambda^{N(m_c-1)} \|\rho\|_{m_c}^{m_c}, \quad \text{for } \lambda > 0.$$

Since  $m_c > 1$ , using Proposition 5.2 and sending  $\lambda \searrow 0$  we get

$$(5.5) \quad \inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_{s,p}(\rho) = \lim_{\lambda \rightarrow 0} \mathcal{F}_{s,p}(\rho^\lambda) = 0.$$

*Case  $M > M_c$ .* Let  $h_s$  be the unit mass extremal (with barycenter at the origin) of the HLS-type inequality (3.9), provided by Lemma 4.2. We set for every  $\lambda > 0$

$$\rho_\lambda(x) := M \lambda^N h_s(\lambda x) \in \mathcal{Y}_M.$$

By recalling (2.2) and (2.5), we have

$$\begin{aligned} \mathcal{F}_{s,p}(\rho_\lambda) &= \frac{\|\rho_\lambda\|_{m_c}^{m_c}}{m_c - 1} - \frac{\chi}{p'} \|K_{s/2} * \rho_\lambda\|_{p'}^{p'} = \lambda^{2N m_c - N} \frac{M^{m_c}}{m_c - 1} - \frac{\chi}{p'} \lambda^{(N-s)p' - N} M^{p'} H_{m_c, s}^* \\ &= \lambda^{p'(N-s)} \frac{p_s^*}{p'} M^{p'} \left[ \frac{1}{M^{\frac{s p'}{N}}} - \frac{1}{M_c^{\frac{s p'}{N}}} \right] < 0, \end{aligned}$$

from which, by sending  $\lambda \rightarrow \infty$ , we get the desired conclusion.  $\square$

**Remark 5.4.** By the previous proposition, we infer that there are no minimizers in  $\mathcal{Y}_M$  of the energy functional  $\mathcal{F}_{s,p}$  in the aggregation dominated regime  $m \in (1, m_c)$ , whatever the value of the mass  $M > 0$ . In the fair competition regime,  $m = m_c$ , still there are no minimizers in  $\mathcal{Y}_M$  for prescribed mass  $M \in (M_c, \infty)$ . Also for values of the mass  $M \in (0, M_c)$ , there are no minimizers of  $\mathcal{F}_{s,p}$  in  $\mathcal{Y}_M$ , in light of the leftmost inequality in Proposition 5.2.

**Remark 5.5.** By inspecting the proof of Proposition 5.1 we can infer that, for every  $m > 1$ , any critical point  $\rho$  of  $\mathcal{F}_{s,p}$  necessarily satisfies the relevant identity

$$(5.6) \quad \frac{\chi}{p_s^*} \|K_{s/2} * \rho\|_{p'}^{p'} = \|\rho\|_m^m.$$

Indeed, (5.2) holds for every  $m > 1$ . Therefore imposing criticality of  $\rho$  only with respect to mass invariant dilations, i.e. imposing (5.3), yields (5.6). Notice that if  $1 < m < m_c$  then  $\rho$  is a maximum, and not a minimum, in the family  $\{\rho^\lambda\}_{\lambda > 0}$ . Notice also that, in the case  $m = m_c$ , (5.6) is equivalent to  $\mathcal{F}_{s,p}(\rho) = 0$ .

Next, we discuss the relation between extremals of (3.9) and minimizers of  $\mathcal{F}_{s,p}$ .

**Corollary 5.6** (Extremals HLS vs minimizers of  $\mathcal{F}_{s,p}$ ). *Let  $1 < p < \infty$  and  $0 < s p < N$ .*

*Let  $m \geq m_c$  and let  $M > 0$ . If  $\rho_s \in \mathcal{Y}_M$  attains the infimum of the energy functional  $\mathcal{F}_{s,p}$ , among functions in  $\mathcal{Y}_M$ , then  $\rho_s$  is an extremal of the Hardy-Littlewood-Sobolev type inequality (3.9), that is  $\rho_s$  attains the supremum in (4.1).*

*Viceversa, for  $m > m_c$ , if  $M > 0$  and  $\rho_s \in \mathcal{Y}_M$  is an extremal of the Hardy-Littlewood-Sobolev type inequality (3.9), then its optimal dilation, in the sense of Proposition 5.1, attains the infimum of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$ . If  $m = m_c$  and  $\rho_s \in \mathcal{Y}_{M_c}$  is an extremal of the Hardy-Littlewood-Sobolev type inequality (3.9), then  $\rho_s$  is a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_{M_c}$ .*

*Proof.* Assume first that  $m > m_c$ ,  $M > 0$ . Let  $\rho_s \in \mathcal{Y}_M$  be a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$ . By Proposition 5.1, we infer that  $\rho_s$  maximizes  $\Lambda$  among functions in  $\mathcal{Y}_M$ , where  $\Lambda$  is the functional defined therein. In particular, by Remark 4.1 and by (5.4),  $\rho_s$  is an extremal of the Hardy-Littlewood-Sobolev type inequality (3.9). On the other hand, let  $\rho_s \in \mathcal{Y}_M$  be an extremal of the Hardy-Littlewood-Sobolev inequality (3.9) (existence of an extremal in  $\mathcal{Y}_M$  is guaranteed by Remark 4.3). Therefore it also satisfies

$$\Lambda(\rho_s) \geq \Lambda(\rho) \quad \text{for every } \rho \in \mathcal{Y}_M,$$

in view of (5.4) and Remark 4.1. Its optimal dilation in the sense of Proposition 5.1, which is still an extremal of (3.9) in light of Remark 4.1, is given by

$$\tilde{\rho}_s := \rho_s^{\lambda_*(\rho_s)},$$

for  $\lambda_*(\rho_s)$  as in (5.1). We claim that  $\tilde{\rho}_s$  minimizes the energy functional  $\mathcal{F}_{s,p}$  among all functions in  $\mathcal{Y}_M$ . Indeed, for every  $\rho \in \mathcal{Y}_M$ , by using the maximality of  $\rho_s$ , the invariance by dilations property of  $\Lambda$  and Proposition 5.1, we have

$$\mathcal{F}_{s,p}(\tilde{\rho}_s) = \mathcal{F}_{s,p}(\rho_s^{\lambda_*(\rho_s)}) = \kappa \Lambda(\rho_s) \leq \kappa \Lambda(\rho) \leq \mathcal{F}_{s,p}(\rho)$$

where the first inequality comes by recalling that  $\kappa$  is negative. This proves the claim.

If  $m = m_c$ , take any minimizer  $\rho_s \in \mathcal{Y}_{M_c}$  of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_{M_c}$ . By Proposition 5.3 we have  $\mathcal{F}_{s,p}(\rho_s) = 0$ , that is

$$(5.7) \quad \frac{1}{m_c - 1} \|\rho_s\|_{m_c}^{m_c} = \frac{\chi}{p'} \|K_{s/2} * \rho_s\|_{p'}^{p'}.$$

By recalling (2.5) and (4.1), this entails that

$$(5.8) \quad \frac{\|K_{s/2} * \rho_s\|_{p'}^{p'}}{M_c^{\frac{p'_s}{N}} \|\rho_s\|_{m_c}^{m_c}} = H_{m_c, s}^*$$

as desired. On the other hand, let  $\rho_s \in \mathcal{Y}_{M_c}$  be an extremal of (3.9). Then, always by recalling (2.5), (5.8) implies (5.7) that is  $\mathcal{F}_{s,p}(\rho_s) = 0$ , proving that  $\rho_s$  minimizes  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_{M_c}$  in view of Proposition 5.3.  $\square$

The equation satisfied by the global minimizers of  $\mathcal{F}_{s,p}$ , provided by Corollary 5.6, reads as follows.

**Lemma 5.7.** *Let  $1 < p < \infty$  and  $0 < sp < N$ , and let  $\chi, M > 0$ . For  $m > m_c$ , if  $\rho_s \in \mathcal{Y}_M$  is a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  then it solves*

$$(5.9) \quad \frac{m}{m-1} \rho_s^{m-1} = \left( \chi \mathcal{K}_{s,p}(\rho_s) - \mathcal{D}_s \right)_+ \quad \text{in } \mathbb{R}^N,$$

with

$$(5.10) \quad \mathcal{D}_s := \left( \frac{p_s^* - m'}{M} \right) \|\rho_s\|_m^m = \left( \frac{p_s^* - m'}{M} \right) \frac{\chi}{p_s^*} \|K_{s/2} * \rho_s\|_{p'}^{p'}.$$

*In particular, it is bounded with compact support, radially nonincreasing,  $W^{1,1}(\mathbb{R}^N)$  and satisfies the Hölder regularity properties of Lemma 4.7. The same holds for  $m = m_c$ , by taking  $M = M_c$ .*

*Proof.* Let  $\rho_s \in \mathcal{Y}_M$  be as in the statement. By Corollary 5.6, we have that  $\rho_s$  is an extremal of the Hardy-Littlewood-Sobolev type inequality (3.9). In light of Lemma 4.4, it satisfies (4.9). If  $m > m_c$ , Proposition 5.1 implies that  $\rho_s$  coincides with its optimal dilation, i.e.,  $\lambda_*(\rho_s) = 1$ , where  $\lambda_*$  is given by (5.1), so that (5.6) holds. If  $m = m_c$ , then Remark 5.4 and Lemma 5.3 imply  $M = M_c$  and  $\mathcal{F}_{s,p}(\rho_s) = 0$  which in turn directly yields (5.6). By inserting (5.6) in (4.10), we see that (4.9) becomes (5.9)-(5.10). By Corollary 4.8, Lemma 4.6 and Lemma 4.7 we then infer the desired conclusions.  $\square$

**Remark 5.8.** For future purposes we record the following identities, valid for  $m > m_c$ , involving the constant  $\mathcal{D}_s$  appearing in the Euler-Lagrange equation (5.9) satisfied by a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$ :

$$\mathcal{D}_s = \left( \frac{p_s^* - m'}{M} \right) \|\rho_s\|_m^m = \left( \frac{p_s^* - m'}{M} \right) \frac{\chi}{p_s^*} \|K_{s/2} \rho_s\|_{p'}^{p'} = \left( \frac{p_s^* - m'}{M} \right) \frac{(m_c - 1)(m - 1)}{m_c - m} \mathcal{F}_{s,p}(\rho_s),$$

which follows from (5.10) and the definition of  $\mathcal{F}_{s,p}$ .

**Remark 5.9.** The conclusion of Lemma 5.7 holds also for continuous radially decreasing critical points of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  in the case  $(p_s^*)' < m < m_c$ . These are defined as continuous radially decreasing solutions to (5.9), with  $\mathcal{D}_s$  still expressed by (5.10). Indeed, a first variation argument along the line of Lemma 4.4 proves that the Euler-Lagrange equation that is necessarily satisfied by a continuous radially decreasing critical point  $\rho$  of  $\mathcal{F}_{s,p}$ , constrained to  $\mathcal{Y}_M$ , is of the form (2.6), for some suitable constant  $\mathcal{Q}$  having the role of Lagrange multiplier for the mass constraint. Moreover  $\mathcal{Q}$  necessarily coincides with  $\mathcal{D}_s$  from (5.10) in view of the criticality condition (5.6). Existence of such critical points, for every mass  $M > 0$ , is deduced from the existence of radially decreasing extremals of the HLS inequality (3.9) having mass  $M$  and satisfying (5.6), which is guaranteed by Lemma 4.2 and Remark 4.1: notice indeed that an extremal of mass  $M$  satisfies (5.6) after taking a mass invariant dilation (thus preserving extremality), see Remark 5.5. A HLS extremal having mass  $M$  and satisfying (5.6) does satisfy (5.9)-(5.10), thanks to Lemma 4.4, as seen by plugging (5.6) in (4.9)-(4.10). The further regularity of such critical points is then deduced in the same way starting from the Euler-Lagrange equation, see Lemma 4.6, Lemma 4.7 and Corollary 4.8.

**Proof of Theorem 2.1.** The first part of Theorem 2.1 follows by Lemma 4.2, Lemma 4.4, Lemma 4.6 and Lemma 4.7. The second part follows by Corollary 5.6 and Lemma 5.7.  $\square$

## 6. THE LIMIT $s \rightarrow 0$

This section is devoted to study the asymptotic behavior of the minimizers  $\rho_s$ , provided by Corollary 5.6, as  $s$  tends to 0. Since the critical exponent  $m_c$  given by (2.2) tends to  $p'$ , as  $s$  goes to zero, we need discuss separately three cases according to whether  $m < p'$ ,  $m = p'$  or  $m > p'$ .

If  $m < p'$ , we have  $m < (p_s^*)'$  for  $s$  small enough. By Proposition 5.3, we get  $\inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_{s,p} = -\infty$  thus there are no minimizers of  $\mathcal{F}_{s,p}$ , and not even stationary states obtained from extremals of the HLS inequality according to Remark 5.9. The case  $m = p'$  (that we call the limiting fair competition regime) will be discussed in Section 6.3. The limiting diffusion dominated regime  $m > p'$  is the most interesting one and it will be treated in the first part of this section. The main property is contained in Theorem 6.7 below: we will prove that if  $m > p'$ , any family of minimizers  $(\rho_s)_{s \in (0, N/p)}$  of the free energy functional  $\mathcal{F}_{s,p}$  provided by Corollary 5.6 strongly converges as  $s \searrow 0$  to the unique minimizer in  $\mathcal{Y}_M$  of a limit functional  $\mathcal{F}_0$ . In addition,  $\mathcal{F}_{s,p}$   $\Gamma$ -converges to  $\mathcal{F}_0$  on  $\mathcal{Y}_M$ , with respect to the strong convergence in  $L^{p'}(\mathbb{R}^N)$ , see Proposition 6.8.

**6.1. The limit functional.** Concerning the limit functional  $\mathcal{F}_0$  defined by (2.7) we have the following

**Proposition 6.1.** *Let  $1 < p < \infty$  and  $m > p'$ . There exists a unique radially symmetric and nonincreasing minimizer of  $\mathcal{F}_0$  in  $\mathcal{Y}_M$ , given by*

$$(6.1) \quad \rho_0(x) = \left(\frac{\chi}{p}\right)^{\frac{1}{m-p'}} 1_{B_{R_0}}(x), \quad \text{where } R_0 = \left(\frac{M}{\omega_N} \left(\frac{p}{\chi}\right)^{\frac{1}{m-p'}}\right)^{\frac{1}{N}}.$$

*Proof.* Let  $\rho \in \mathcal{Y}_M$  and let  $\rho_\lambda$  be its mass invariant dilation given by (1.5). We have

$$\mathcal{F}_0(\rho^\lambda) = \frac{\lambda^{N(m-1)}}{m-1} \|\rho\|_m^m - \frac{\chi}{p'} \lambda^{N(p'-1)} \|\rho\|_{p'}^{p'},$$

and

$$\frac{d}{d\lambda} \mathcal{F}_0(\rho^\lambda) = N \lambda^{N(m-1)-1} \|\rho\|_m^m - \frac{\chi}{p} N \lambda^{N(p'-1)-1} \|\rho\|_{p'}^{p'} = N \lambda^{N(p'-1)-1} \left[ \lambda^{N(m-p')} \|\rho\|_m^m - \frac{\chi}{p} \|\rho\|_{p'}^{p'} \right],$$

for  $\lambda > 0$ . By optimizing in  $\lambda$ , we find that

$$(6.2) \quad \lambda_*(\rho) := \left( \frac{\chi}{p} \frac{\|\rho\|_{p'}^{p'}}{\|\rho\|_m^m} \right)^{\frac{1}{N(m-p')}},$$

is the unique global minimum of  $\lambda \mapsto \mathcal{F}_0(\rho^\lambda)$ , for  $\lambda > 0$ . We then have

$$\mathcal{F}_0(\rho^{\lambda_*(\rho)}) = \kappa \Lambda(\rho),$$

where

$$\kappa = - \left(\frac{\chi}{p}\right)^{\frac{m-1}{m-p'}} \quad \text{and} \quad \Lambda(\rho) = \frac{\|\rho\|_{p'}^{p' \frac{m-1}{m-p'}}}{\|\rho\|_m^m \frac{p'-1}{m-p'}}.$$

In order to minimize the functional  $\mathcal{F}_0$  on  $\mathcal{Y}_M$  we can equivalently maximize  $\Lambda$ . Moreover, by symmetrization, we can look for maximizer of  $\Lambda$  in the restricted class of

$$\tilde{\mathcal{Y}}_M := \{\rho \in \mathcal{Y}_M : \rho \text{ is radially symmetric and nonincreasing}\}.$$

By using Hölder's inequality, we get that for  $\rho \in \tilde{\mathcal{Y}}_M$

$$\|\rho\|_{p'}^{p'} \leq M^{\frac{m-p'}{m-1}} \|\rho\|_m^m \frac{p'-1}{m-1} \iff \Lambda(\rho) \leq M,$$

and the equality is satisfied by a function  $\rho_0$  if and only if  $\rho_0(x) = c 1_F(x)$ , for some measurable set  $F \subseteq \mathbb{R}^N$ , see [35, Theorem 2.3 (ii.b)] for example. Since  $\rho_0 \in \tilde{\mathcal{Y}}_M$  and by using that  $\|\rho_0\|_1 = M$ , we infer that  $F = B_R$ , for some  $R > 0$ , and  $c = M/|B_R|$ . Moreover, since  $\rho_0$  minimizes  $\mathcal{F}_0$  it must coincide with its optimal dilation given by (6.2), i.e.  $\lambda_*(\rho_0) = 1$ . This entails that

$$M = \Lambda(\rho_0) = \left(\frac{p}{\chi}\right)^{\frac{m-1}{m-p'}} \frac{M^m}{|B_R|^{m-1}},$$

from which we infer (6.1). □

In the next result we infer information on the limiting behavior of the minimum value of  $\mathcal{F}_{s,p}$ , by testing the energy  $\mathcal{F}_{s,p}$  with  $\rho_0$ .

**Corollary 6.2.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > p'$  and let  $\chi, M > 0$ . If  $\rho_s \in \mathcal{Y}_M$  is a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ , then we have*

$$\limsup_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) < 0 \quad \text{and} \quad \limsup_{s \rightarrow 0} \mathcal{D}_s > 0$$

where  $\mathcal{D}_s > 0$  is the constant related to  $\rho_s$  appearing in Lemma 5.7.

*Proof.* Let  $\rho_0$  be as in Proposition 6.1. By using Theorem 3.4, we have

$$\lim_{s \rightarrow 0} \|K_{s/2} * \rho_0 - \rho_0\|_{p'} = 0.$$

By the minimality of  $\rho_s$  and by using the explicit expression of  $\rho_0$ , this entails that

$$\limsup_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) \leq \lim_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_0) = \frac{M}{m-1} \left(\frac{\chi}{p}\right)^{\frac{m-1}{m-p'}} - M \frac{\chi}{p'} \left(\frac{\chi}{p}\right)^{\frac{p'-1}{m-p'}} < 0,$$

where the last inequality follows since  $m > p'$ . By recalling Remark 5.8, we also get the announced asymptotic behavior of  $\mathcal{D}_s$ .  $\square$

**Remark 6.3.** Concerning the limit functional  $\mathcal{F}_0$  in the case  $m = p'$ , for any  $M > 0$  it is clear that  $\inf_{\mathcal{Y}_M} \mathcal{F}_0 = 0$  if  $0 < \chi \leq p$  and that  $\inf_{\mathcal{Y}_M} \mathcal{F}_0 = -\infty$  if  $\chi > p$ . These properties are obtained by taking dilations  $\rho^\lambda$  for any given  $\rho \in \mathcal{Y}_M$  and by sending  $\lambda$  to  $+\infty$  and to 0, respectively. The infimum is not realized, except for the trivial case  $p = \chi$ .

**6.2. The limiting diffusion dominated regime.** Next, we discuss the limiting behavior of the minimizers for  $s \searrow 0$ , in the case  $m > p'$ .

**Proposition 6.4** (Equiboundedness of  $\rho_s$ ). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > p'$  and let  $\chi, M > 0$ . Let  $\rho_s \in \mathcal{Y}_M$  be a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ . Then there exists  $s_0 \in (0, N/p)$  such that*

$$\sup_{s \in (0, s_0)} \|\rho_s\|_\infty < \infty \quad \text{and} \quad \sup_{s \in (0, s_0)} \|\rho_s\|_m < \infty.$$

*Proof.* From Corollary 5.6 and Lemma 5.7 we know that  $\rho_s$  is a radially symmetric non-increasing Hölder continuous function. This yields that

$$(6.3) \quad \|\rho_s\|_\infty = \rho_s(0) \quad \text{and} \quad \rho_s(x) \leq \frac{M}{\omega_N |x|^N} \quad \text{for every } x \in \mathbb{R}^N \setminus \{0\},$$

for every  $s$ . The first fact is clear, and the second one follows since we have

$$M \geq \int_{B_{|x|}} \rho_s dy \geq \int_{B_{|x|}} \rho(x) dy = \rho_s(x) \omega_N |x|^N, \quad \text{for every } x \in \mathbb{R}^N \setminus \{0\}.$$

By using the Euler-Lagrange equation (4.9) and Hölder's inequality we get

$$\begin{aligned}
\frac{m'}{\chi} \rho_s(0)^{m-1} &\leq K_{s/2} * (K_{s/2} * \rho_s)^{p'-1}(0) \\
&= c_{N,s/2} \int_{B_1} |y|^{s-N} (K_{s/2} * \rho_s)^{p'-1}(y) dy + c_{N,s/2} \int_{B_1^c} |y|^{s-N} (K_{s/2} * \rho_s)^{p'-1}(y) dy \\
&\leq c_{N,s/2} \left( \frac{N\omega_N}{s} \right) \|K_{s/2} * \rho_s\|_\infty^{p'-1} + c_{N,s/2} \frac{N\omega_N}{(N-s)p' + N} \|K_{s/2} * \rho_s\|_{p'}^{\frac{p'}{p}} \\
&\leq c_{N,s/2} \left( \frac{N\omega_N}{s} \right) (\alpha_s M + \beta_s \rho_s(0))^{p'-1} + c_{N,s/2} \frac{N\omega_N}{(N-s)p' + N} H_s^{p'-1} \|\rho_s\|_{(p_s^*)}^{p'-1},
\end{aligned}$$

where in the last line we used Lemma 3.1 with data  $q := 1$  and  $r := \infty$  and the HLS inequality (3.7). By spending again (6.3), we infer

$$\int_{\mathbb{R}^N} \rho_s^{(p_s^*)'} dx = \int_{B_1} \rho_s^{(p_s^*)'} dx + \int_{B_1^c} \rho_s^{(p_s^*)'} dx \leq \omega_N \rho_s(0)^{(p_s^*)'} + \left( \frac{M}{\omega_N} \right)^{(p_s^*)'} \frac{1}{N((p_s^*)' - 1)}$$

By collecting the last two inequalities, we get

$$\begin{aligned}
\frac{m'}{\chi} \rho_s(0)^{m-1} &\leq c_{N,s/2} \left( \frac{N\omega_N}{s} \right) (\alpha_s M + \beta_s \rho_s(0))^{p'-1} \\
&\quad + c_{N,s/2} \frac{N\omega_N H_s^{p'-1}}{(N-s)p' + N} \left( \rho_s(0)^{(p_s^*)'} \omega_N + \left( \frac{M}{\omega_N} \right)^{(p_s^*)'} \frac{1}{N((p_s^*)' - 1)} \right)^{\frac{p'-1}{(p_s^*)'}}.
\end{aligned}$$

By contradiction, we assume now that

$$\limsup_{s \rightarrow 0} \rho_s(0) = +\infty,$$

and we divide both sides of the previous inequality by  $\rho_s(0)^{p'-1}$ . We get

$$\begin{aligned}
\frac{m'}{\chi} \rho_s(0)^{m-p'} &\leq c_{N,s/2} \left( \frac{N\omega_N}{s} \right) \left( \frac{\alpha_s M}{\rho_s(0)} + \beta_s \right)^{p'-1} \\
&\quad + c_{N,s/2} \frac{N\omega_N H_s^{p'-1}}{(N-s)p' + N} \left( \omega_N + \left( \frac{M}{\omega_N} \right)^{(p_s^*)'} \frac{\rho_s(0)^{-(p_s^*)'}}{N((p_s^*)' - 1)} \right)^{\frac{p'-1}{(p_s^*)'}}.
\end{aligned}$$

By taking the limsup as  $s \searrow 0$  on both sides of the last estimate, we get that the leftmost term diverges to  $+\infty$ , being  $m > p'$ . On the other hand, by recalling the asymptotic behaviors of  $\alpha_s, \beta_s, H_s$  and  $c_{N,s/2}$  given respectively in Lemma 3.1, Corollary 3.3 and formula (3.1), we infer that the rightmost term stays finite. This yields a contradiction and so there must exist  $s_0 > 0$  such that  $S := \sup_{s \in (0, s_0)} \rho_s(0) < \infty$ . We conclude the proof by observing that we can also infer the equiboundedness of  $\|\rho_s\|_m$  for  $s \in (0, s_0)$ , since we have  $\|\rho_s\|_m \leq S^{\frac{1}{m'}} M^{\frac{1}{m}}$ , for  $s \in (0, s_0)$ , in light of the interpolation inequality in  $L^p$  spaces.  $\square$

**Proposition 6.5** (Equiboundedness of  $\text{supp}(\rho_s)$ ). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > p'$  and let  $\chi, M > 0$ . Let  $\rho_s \in \mathcal{Y}_M$  be a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ . Then there exist  $s_0 \in (0, N/p)$  and  $R_0 > 0$  such that  $\text{supp}(\rho_s) \subseteq B_{R_0}$  for every  $0 < s < s_0$ .*

*Proof.* Let  $s_0$  given by Proposition 6.4. We set  $B_{R_s} = \text{supp}(\rho_s)$  and by contradiction, we assume that

$$\limsup_{s \rightarrow 0} R_s = +\infty.$$

This entails that  $R_s > 1$ , for small value of  $s$ . We preliminary observe that, from Corollary 3.3 and Proposition 6.4, we have

$$(6.4) \quad \|K_{s/2} * \rho_s\|_{p'} \leq H_s \|\rho_s\|_{(p_s^*)'} \leq H_s M^\vartheta \|\rho_s\|_\infty^{1-\vartheta} \leq L, \quad \text{for } s \in (0, s_0),$$

where  $\vartheta = 1/(p_s^*)'$  and  $L > 0$  is a constant depending only on  $s_0$ . We take  $x \in \partial B_{R_s}$ , and by using Lemma 5.7 we get

$$(6.5) \quad \begin{aligned} \frac{1}{\chi} \mathcal{D}_s &= K_{s/2} * (K_{s/2} * \rho_s)^{p'-1}(x) = c_{N,s/2} \int_{B_1} |y|^{s-N} ((K_{s/2} * \rho_s)(x-y))^{p'-1} dy \\ &+ c_{N,s/2} \int_{B_1^c} |y|^{s-N} ((K_{s/2} * \rho_s)(x-y))^{p'-1} dy \\ &=: \mathcal{A}_1 + \mathcal{A}_2. \end{aligned}$$

We estimate the last two integrals separately, starting from the second one. By using Hölder's inequality (we observe that, by assumption  $p' > N/(N-s)$ , so we have  $(s-N)p' + N < 0$ ) we have

$$\begin{aligned} \mathcal{A}_2 &\leq c_{N,s/2} \left( \int_{B_1^c} |y|^{(s-N)p'} dy \right)^{\frac{1}{p'}} \left( \int_{B_1^c} ((K_{s/2} * \rho_s)(x-y))^{p'} dy \right)^{\frac{1}{p'}} \\ &\leq c_{N,s/2} \frac{N\omega_N}{(N-s)p' + N} \|K_{s/2} * \rho_s\|_{p'}^{\frac{p'}{p}}. \end{aligned}$$

So by using (3.1) and (6.4) we get

$$(6.6) \quad \lim_{s \rightarrow 0} \mathcal{A}_2 = 0.$$

We now consider the first integral

$$\mathcal{A}_1 = c_{N,s/2} \int_{B_1} |y|^{s-N} ((K_{s/2} * \rho_s)(x-y))^{p'-1} dy$$

Since  $\rho_s$  is nonincreasing, so is  $K_{s/2} * \rho_s$  (see [13]). Then, by using also (6.4), we get

$$L \geq \left( \int_{\mathbb{R}^N} |K_{s/2} * \rho_s|^{p'} dy \right)^{\frac{1}{p'}} \geq \left( \int_{B_{|x|}} |K_{s/2} * \rho_s|^{p'} dy \right)^{\frac{1}{p'}} \geq (\omega_N |x|^N)^{\frac{1}{p'}} (K_{s/2} * \rho_s)(x),$$

for every  $x \in \mathbb{R}^N \setminus \{0\}$ . Since  $|x| = R_s > 1$ , by the triangle inequality we have

$$|x-y| \geq |x| - |y| \geq R_s - 1, \quad \text{for } y \in B_1,$$

thus

$$\begin{aligned} \mathcal{A}_1 &\leq L \omega_N^{-\frac{1}{p}} c_{N,s/2} \int_{B_1} |y|^{s-N} |x-y|^{-\frac{N}{p}} dy \leq \frac{L \omega_N^{-\frac{1}{p}}}{(R_s - 1)^{\frac{N}{p}}} c_{N,s/2} \int_{B_1} |y|^{s-N} dy \\ &= \frac{L \omega_N^{-\frac{1}{p}}}{(R_s - 1)^{\frac{N}{p}}} c_{N,s/2} \left( \frac{N\omega_N}{s} \right). \end{aligned}$$

By recalling (3.1) and (6.2), we eventually get that  $\lim_{s \rightarrow 0} \mathcal{A}_1 = 0$ , and so also  $\limsup_{s \rightarrow 0} \mathcal{D}_s = 0$ , by (6.5) and (6.6). This contradicts Corollary 6.2 and gives the desired result.  $\square$

**Proposition 6.6.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > p'$ , let  $\chi, M > 0$  and let  $\rho_s \in \mathcal{Y}_M$  be a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ . There exists  $s_0 \in (0, N/p)$  such that the family of minimizers  $(\rho_s)_{s \in (0, s_0)}$  is equibounded in  $W^{1,1}(\mathbb{R}^N)$ . Moreover, if  $(s_k)_{k \in \mathbb{N}} \subseteq (0, s_0)$  converges to 0, then the family  $(\rho_{s_k})_{k \in \mathbb{N}}$  admits limit points in the strong  $L^1(\mathbb{R}^N)$  topology, and if  $\rho$  is a limit point along a not relabeled subsequence we have  $\rho \in \mathcal{Y}_M \cap L^\infty(\mathbb{R}^N)$  and*

$$\lim_{k \rightarrow \infty} \|\rho_{s_k} - \rho\|_q = 0, \quad \text{for every } q \in [1, \infty).$$

*Proof.* The proof is the same of [28, Lemma 3.7] by using Proposition 6.4 and Proposition 6.5.  $\square$

The main result of this section regarding the case  $m > p'$  reads as follows

**Theorem 6.7.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $m > p'$ , let  $\chi, M > 0$  and let  $\rho_s \in \mathcal{Y}_M$  be a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ . Then*

$$\lim_{s \rightarrow 0} \|\rho_s - \rho_0\|_q = 0, \quad \text{for every } q \in [1, \infty),$$

where  $\rho_0$  is given by (6.1).

*Proof.* By using Proposition 6.6, there exists a sequence  $(s_k)_{k \in \mathbb{N}}$  converging to 0 and a function  $\rho \in \mathcal{Y}_M \cap L^\infty(\mathbb{R}^N)$  such that

$$(6.7) \quad \lim_{k \rightarrow \infty} \|\rho_{s_k} - \rho\|_m = 0.$$

By the triangle inequality and the HLS-type inequality (3.9), we get

$$\begin{aligned} \|K_{s_k/2} * \rho_{s_k} - \rho\|_{p'} &\leq \|K_{s_k/2} * (\rho_{s_k} - \rho)\|_{p'} + \|K_{s_k/2} * \rho - \rho\|_{p'} \\ &\leq H_s (2M)^{\vartheta_0} \|\rho_{s_k} - \rho\|_m^{1-\vartheta_0} + \|K_{s_k/2} * \rho - \rho\|_{p'}. \end{aligned}$$

where  $\vartheta_0$  is given by (3.10), it depends on  $s_k$  and converges to  $1 - m'/p > 0$  as  $k \rightarrow +\infty$ . By Theorem 3.4, we have

$$\lim_{k \rightarrow \infty} \|K_{s_k/2} * \rho - \rho\|_{p'} = 0,$$

thus from the previous inequality, from (6.7) and from the bound on  $H_s$  by Corollary 3.3, we get

$$\lim_{k \rightarrow \infty} \|K_{s_k/2} * \rho_{s_k} - \rho\|_{p'} = 0.$$

This entails that

$$\lim_{k \rightarrow \infty} \mathcal{F}_{s_k,p}(\rho_{s_k}) = \frac{\|\rho\|_m^m}{m-1} - \frac{\chi}{p'} \|\rho\|_{p'} = \mathcal{F}_0(\rho).$$

For every  $\tilde{\rho} \in \mathcal{Y}_M$ , by using the previous equality, the minimality of  $\rho_{s_k}$  and Theorem 3.4, we infer

$$\mathcal{F}_0(\rho) = \lim_{k \rightarrow \infty} \mathcal{F}_{s_k,p}(\rho_{s_k}) \leq \lim_{k \rightarrow \infty} \mathcal{F}_{s_k,p}(\tilde{\rho}) = \mathcal{F}_0(\tilde{\rho}).$$

This proves that  $\rho$  must be a minimizer of  $\mathcal{F}_0$  in  $\mathcal{Y}_M$  and so by Proposition 6.1 it must coincide with  $\rho_0$ . By the arbitrariness of  $(s_k)_{k \in \mathbb{N}}$ , we eventually get that the whole family  $(\rho_s)$  strongly converges to  $\rho_0$  in  $L^q(\mathbb{R}^N)$ , for every  $q \in [1, \infty)$ .  $\square$

We further have the following  $\Gamma$ -convergence result.

**Proposition 6.8.** *Let  $1 < p < \infty$  and  $\chi, M > 0$ . Let  $m > p'$ . For  $s \rightarrow 0$ , the functional  $\mathcal{F}_{s,p}$   $\Gamma$ -converges to  $\mathcal{F}_0$  on  $\mathcal{Y}_M$  with respect to the strong convergence in  $L^{p'}(\mathbb{R}^N)$ .*

*Proof.* Let  $(\rho_s)_{s \in (0, N/p)} \subset \mathcal{Y}_M$  and  $\rho \in \mathcal{Y}_M$  be such that

$$(6.8) \quad \lim_{s \rightarrow 0} \|\rho_s - \rho\|_{p'} = 0.$$

By the triangle inequality and the Hardy-Littlewood-Sobolev inequality (3.7) we have

$$\begin{aligned} \|K_{s/2} * \rho_s - \rho\|_{p'} &\leq \|K_{s/2} * (\rho_s - \rho)\|_{p'} + \|K_{s/2} * \rho - \rho\|_{p'} \\ &\leq H_s \|\rho_s - \rho\|_{(p_s^*)'} + \|K_{s/2} * \rho - \rho\|_{p'} \\ &\leq H_s \|\rho_s - \rho\|_1^\vartheta \|\rho_s - \rho\|_{p'}^{1-\vartheta} + \|K_{s/2} * \rho - \rho\|_{p'}, \end{aligned}$$

where  $\vartheta = 1 - p/p_s^*$ . By Theorem 3.4, we have

$$\lim_{s \rightarrow 0} \|K_{s/2} * \rho - \rho\|_{p'} = 0,$$

which entails that

$$\lim_{s \rightarrow 0} \|K_{s/2} * \rho_s - \rho\|_{p'} = 0,$$

where we also used our assumption (6.8). By Fatou's lemma and by spending the last information, we infer

$$\mathcal{F}_0(\rho) \leq \liminf_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s).$$

On the other hand, let  $\rho \in \mathcal{Y}_M$  we set  $\rho_s := \rho$ , for every  $s$ . By using again Theorem 3.4 we have

$$\limsup_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) = \mathcal{F}_0(\rho),$$

From the last two facts we obtain the claimed result.  $\square$

**6.3. The limiting fair competition regime.** We now analyze the limiting behavior of the minimizers in the remaining case  $m = p'$  thus concluding the proof of Theorem 2.2. We treat separately the cases  $\chi \neq p$  and  $\chi = p$ , starting from the first one.

**Theorem 6.9.** *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $M > 0$  and  $m = p'$ . If  $\rho_s \in \mathcal{Y}_M$  is a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ , then we have*

$$\lim_{s \rightarrow 0} \|\rho_s\|_\infty = - \lim_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) = \begin{cases} 0, & \text{if } 0 < \chi < p, \\ +\infty, & \text{if } \chi > p. \end{cases}$$

Moreover, if  $\chi > p$  we have  $\rho_s \rightarrow M\delta_0$  in the sense of measures as  $s \rightarrow 0$ .

*Proof.* By Corollary 5.6, we have that  $\rho_s$  is radially symmetric and nonincreasing. We discuss separately two cases.

If  $\boxed{0 < \chi < p}$ , we argue by contradiction and assume that

$$\limsup_{s \rightarrow 0} \rho_s(0) > 0.$$

By arguing as in the proof of Proposition 6.4, we infer that

$$\begin{aligned} \frac{p}{\chi} \rho_s(0)^{p'-1} &\leq c_{N,s/2} \left( \frac{N\omega_N}{s} \right) (\alpha_s M + \beta_s \rho_s(0))^{p'-1} \\ &\quad + c_{N,s/2} \frac{N\omega_N H_s^{p'-1}}{(N-s)p' + N} \left( \rho_s(0)^{(p_s^*)'} \omega_N + \left( \frac{M}{\omega_N} \right)^{(p_s^*)'} \frac{1}{N((p_s^*)' - 1)} \right)^{\frac{p'-1}{(p_s^*)'}}. \end{aligned}$$

By dividing both sides of the previous inequality by  $\rho_s(0)^{p'-1}$ , we can rewrite it as

$$\begin{aligned} \frac{p}{\chi} - c_{N,s/2} \left( \frac{N\omega_N}{s} \right) \left( \frac{\alpha_s M}{\rho_s(0)} + \beta_s \right)^{p'-1} \\ \leq c_{N,s/2} \frac{N\omega_N H_s^{p'-1}}{(N-s)p' + N} \left( \omega_N + \left( \frac{M}{\omega_N \rho_s(0)} \right)^{(p_s^*)'} \frac{1}{N((p_s^*)' - 1)} \right)^{\frac{p'-1}{(p_s^*)'}} = o(s). \end{aligned}$$

By sending  $s \rightarrow 0$  and by using (3.1), (3.2) and (3.3), we get a contradiction. Therefore  $\rho_s$  converges uniformly to zero as  $s \rightarrow 0$ , which also implies  $\mathcal{F}_{s,p}(\rho_s) \rightarrow 0$ , by Corollary 3.3.

If  $\boxed{\chi > p}$ , we consider the limit functional  $\mathcal{F}_0$ , given by (2.7) with  $m = p'$ . For  $\rho \in \mathcal{Y}_M$ , we have

$$\lim_{\lambda \rightarrow \infty} \mathcal{F}_0(\rho^\lambda) = \lim_{\lambda \rightarrow \infty} \lambda^{N(p'-1)} \left( \frac{1}{p'-1} - \frac{\chi}{p'} \right) \|\rho\|_{p'}^{p'} = -\infty,$$

where  $\rho^\lambda \in \mathcal{Y}_M$  is the mass invariant dilation of  $\rho$  by factor  $\lambda$ , given by (1.5). This entails that

$$\inf_{\rho \in \mathcal{Y}_M} \mathcal{F}_0 = -\infty.$$

Then if  $\beta < 0$ , we can take  $\bar{\rho} \in \mathcal{Y}_M$  such that  $\mathcal{F}_0(\bar{\rho}) < \beta$ . By using that

$$\lim_{s \rightarrow 0} \|K_{s/2} * \bar{\rho} - \bar{\rho}\|_{p'} = 0,$$

thanks to Theorem 3.4, and by the minimality of  $\rho_s$  we then obtain

$$\limsup_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) \leq \limsup_{s \rightarrow 0} \mathcal{F}_0(\bar{\rho}) = \mathcal{F}_0(\bar{\rho}) < \beta.$$

By the arbitrariness of  $\beta$  and by using Remark 5.8, this yields

$$(6.9) \quad \lim_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) = -\infty \quad \text{and} \quad \lim_{s \rightarrow 0} \mathcal{D}_s = +\infty.$$

By contradiction, we assume that

$$R_0 := \limsup_{s \rightarrow 0} R_s > 0,$$

where we set  $R_s = \text{supp}(\rho_s)$ , as in Proposition 6.5. We take a sequence  $(s_k)_{k \in \mathbb{N}} \subseteq (0, 1)$  converging to zero and such that

$$(6.10) \quad \lim_{k \rightarrow \infty} R_{s_k} = \limsup_{s \rightarrow 0} R_s = R_0.$$

We set  $\bar{R} = R_0/2$  and  $B_{R_{s_k}} := \text{supp}(\rho_{s_k})$ . For  $x \in \partial B_{R_{s_k}}$  by using Lemma 5.7 we have

$$\begin{aligned}
(6.11) \quad \frac{1}{\chi} \mathcal{D}_{s_k} &= K_{s_k/2} * (K_{s_k/2} * \rho_{s_k})^{p'-1}(x) \\
&= c_{N,s_k/2} \int_{B_{\bar{R}}} |y|^{s_k-N} ((K_{s_k/2} * \rho_{s_k})(x-y))^{p'-1} dy \\
&\quad + c_{N,s_k/2} \int_{B_{\bar{R}}^c} |y|^{s_k-N} ((K_{s_k/2} * \rho_{s_k})(x-y))^{p'-1} dy =: \mathcal{A}_1 + \mathcal{A}_2.
\end{aligned}$$

For  $\mathcal{A}_1$ , by arguing as in Proposition 6.5, we have that

$$(\omega_N |x|^N)^{\frac{1}{p'}} (K_{s_k/2} * \rho_{s_k})(x) \leq L, \quad \text{for every } x \in \mathbb{R}^N \setminus \{0\},$$

and, since  $|x| = R_{s_k} > \bar{R}$  for  $k$  large enough, by the triangle inequality we have

$$|x-y| \geq |x| - |y| \geq R_{s_k} - \bar{R}, \quad \text{for } y \in B_{\bar{R}},$$

This entails that

$$(6.12) \quad \mathcal{A}_1 \leq c_{N,s_k/2} \left( \frac{N\omega_N}{s_k} \right) \frac{L\omega_N^{-\frac{1}{p}}}{(R_{s_k} - \bar{R})^{\frac{N}{p}}}.$$

For  $\mathcal{A}_2$ , by using Hölder's inequality we have

$$\begin{aligned}
(6.13) \quad \mathcal{A}_2 &\leq c_{N,s_k/2} \left( \int_{B_1^c} |y|^{(s_k-N)p'} dy \right)^{\frac{1}{p'}} \left( \int_{B_1^c} ((K_{s_k/2} * \rho_{s_k})(x-y))^{p'} dy \right)^{\frac{1}{p'}} \\
&\leq c_{N,s_k/2} \frac{N\omega_N}{(N-s_k)p' + N} \|K_{s_k/2} * \rho_{s_k}\|_{p'}^{\frac{p'}{p}} \\
&\leq E c_{N,s_k/2} \left( \frac{\mathcal{D}_{s_k}}{s_k} \right)^{\frac{1}{p}},
\end{aligned}$$

where in the last equality we used Remark 5.8 and we set

$$E = E(N, p, \chi, M) := \frac{N\omega_N}{(N-1)p' + N} \left( \frac{M}{\chi} \frac{N}{p} \right)^{\frac{1}{p}}.$$

By spending (6.12) and (6.13) in (6.11) and by dividing for  $\mathcal{D}_{s_k}^{\frac{1}{p}}$ , we get

$$\frac{1}{\chi} \mathcal{D}_{s_k}^{1-\frac{1}{p}} \leq c_{N,s_k/2} \left( \frac{N\omega_N}{s_k} \right) \frac{L\omega_N^{-\frac{1}{p}}}{(R_{s_k} - \bar{R})^{\frac{N}{p}}} + E c_{N,s_k/2} \left( \frac{1}{s_k} \right)^{\frac{1}{p}}.$$

In light of (3.1), (6.9) and (6.10), by sending  $k \rightarrow \infty$ , the last inequality yields a contradiction. This implies that

$$\lim_{s \rightarrow 0} R_s = 0,$$

thus, since  $\rho_s \in \mathcal{Y}_M$ , we must have that  $\rho_s$  converges to a point mass at the origin as  $s \rightarrow 0$ , as desired. Along with (6.9), this concludes the proof.  $\square$

**Proposition 6.10** (Case  $\chi = p$ ). *Let  $1 < p < \infty$  and  $0 < sp < N$ . Let  $M > 0$ ,  $m = p'$  and  $\chi = p$ . If  $\rho_s \in \mathcal{Y}_M$  is a minimizer of  $\mathcal{F}_{s,p}$  over  $\mathcal{Y}_M$  for every  $s \in (0, N/p)$ , then we have*

$$\lim_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) = 0 \quad \text{and} \quad \lim_{s \rightarrow 0} \mathcal{D}_s = 0.$$

*Proof.* We recall that by Proposition 5.3, we have

$$\limsup_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) \leq \limsup_{s \rightarrow 0} \mu_s \leq 0.$$

We claim that

$$(6.14) \quad \limsup_{s \rightarrow 0} \|\rho_s\|_{p'} < \infty.$$

Since we have

$$\mathcal{F}_{s,p}(\rho_s) = -\frac{sp}{N-sp} \|\rho_s\|_{p'}^{p'},$$

by Remark 5.8, this would entail that

$$\liminf_{s \rightarrow 0} \mathcal{F}_{s,p}(\rho_s) = 0,$$

and so the desired result. By recalling (4.1) and Corollary 5.6, we have that

$$H_s^{p'} \geq H_{p',s}^* = \frac{\|K_{s/2} * \rho_s\|_{p'}^{p'}}{M^{p' \frac{s}{N}} \|\rho_s\|_{p'}^{p'(1-\frac{s}{N})}} = \frac{p_s^*}{p} \left( \frac{\|\rho_s\|_{p'}^{p'}}{M^{p'}} \right)^{\frac{s}{N}},$$

where  $H_s$  is given as in Corollary 3.3 and the last equality follows from (5.6). Our claim (6.14), will follow by proving that

$$(6.15) \quad \limsup_{s \rightarrow 0} H_s^{\frac{1}{s}} < \infty.$$

By (3.8), we have

$$(6.16) \quad \begin{aligned} H_s^{\frac{1}{s}} &\leq \mathcal{B}_s^{\frac{1}{s}} \left( \frac{N-s}{N} \right)^{\frac{N-s}{Ns}} \left( \frac{(p')^{\frac{N-s}{N}} + (p_s^*)^{\frac{N-s}{N}}}{(p_s^*)' p} \right)^{\frac{1}{s}} \\ &\leq \mathcal{B}_s^{\frac{1}{s}} \left( \frac{N}{N-sp} \right)^{\frac{1}{s}} \leq \mathcal{B}_s^{\frac{1}{s}} \left( 1 + \frac{sp}{N-p} \right)^{\frac{1}{s}}, \end{aligned}$$

where we set

$$\mathcal{B}_s := \frac{c_{N,s/2}}{s} N \omega_N^{\frac{N-s}{N}}.$$

By (3.1), we have  $\lim_{s \rightarrow 0} \mathcal{B}_s = 1$ , and we can write

$$\mathcal{B}_s = \frac{1}{2} \frac{\pi^{-\frac{N}{2}} 2^{-s} \Gamma(\frac{N-s}{2})}{\Gamma(\frac{s}{2} + 1)} N \left( \frac{\pi^{\frac{N}{2}}}{\Gamma(\frac{N}{2} + 1)} \right)^{\frac{N-s}{N}},$$

for every  $s \in (0, N/p)$ . Since  $\mathcal{B}_s$  is smooth in a neighborhood of  $s = 0$ , we have

$$\lim_{s \rightarrow 0} \frac{\mathcal{B}_s - 1}{s} < \infty.$$

By spending this information in (6.16), we get (6.15) which in turn implies (6.14). Eventually, by recalling Remark 5.8 we infer the desired asymptotic behavior also for  $\mathcal{D}_s$ .  $\square$

**Proof of Theorem 2.2.** If  $m > p'$ , the claimed result is contained in Theorem 6.7. If  $m = p'$ , the conclusion follows by Theorem 6.9, Proposition 6.10 and Remark 6.3.  $\square$

**Acknowledgments.** The authors wish to warmly thank Bruno Volzone for fruitful discussions and Pedro Miguel Campos for having provided us the reference [32]. The authors are members of the *Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). F.B. is partially supported by the “INdAM - GNAMPA Project Ottimizzazione Spettrale, Geometrica e Funzionale”, CUP E5324001950001. E.M. is supported by the MIUR- PRIN project 202244A7YL.

#### REFERENCES

- [1] D. R. Adams and L. I. Hedberg, *Function spaces and potential theory*, Grundlehren der mathematischen Wissenschaften, 314, Springer, Berlin, 1996;
- [2] L. Ambrosio, N. Fusco, D. Pallara, *Functions of bounded variation and free discontinuity problems*. Oxford Math. Monogr. The Clarendon Press, Oxford University Press, New York, 2000.
- [3] S. Bian and J.-G. Liu. Dynamic and steady states for multi-dimensional Keller-Segel model with diffusion exponent  $m > 0$ . *Commun. Math. Phys.*, 323(3):1017–1070, 2013.
- [4] A. Blanchet, V. Calvez, J. A. Carrillo, *Convergence of the mass-transport steepest descent scheme for the sub-critical Patlak-Keller-Segel model*. SIAM J. Numer. Anal. **46** (2008) 691–721.
- [5] A. Blanchet, E. Carlen, J. A. Carrillo, *Functional inequalities, thick tails and asymptotics for the critical mass Patlak-Keller-Segel model*. J. Func. Anal. **262**, (2012) 2142-2230.
- [6] A. Blanchet, J.A. Carrillo, P. Laurençot, *Critical mass for a Patlak-Keller-Segel model with degenerate diffusion in higher dimensions*. Calc. Var. Partial Differential Equations **35** (2009), 133–168.
- [7] A. Blanchet, J.A. Carrillo, N. Masmoudi, *Infinite time aggregation for the critical Patlak-Keller-Segel model in  $\mathbb{R}^2$* , Comm. Pure Appl. Math. **61** (10) (2008), 1449–1481.
- [8] A. Blanchet, J. Dolbeault, B. Perthame, *Two-dimensional Keller-Segel model: optimal critical mass and qualitative properties of the solutions*. Electron. J. Differential Equations **44** (2006), 32 pp. (electronic).
- [9] M. Burger, V. Capasso, D. Morale, *On an aggregation model with long and short range interactions*, Nonlinear Anal. Real World Appl. **8** (2007), no. 3, 939–958.
- [10] V. Calvez, J. A. Carrillo, and F. Hoffmann. Equilibria of homogeneous functionals in the fair-competition regime. *Nonlinear Anal.*, 159:85–128, 2017.
- [11] V. Calvez, J. A. Carrillo, and F. Hoffmann. The geometry of diffusing and self-attracting particles in a one-dimensional fair-competition regime. In *Nonlocal and nonlinear diffusions and interactions: new methods and directions*, volume 2186 of *Lecture Notes in Math.*, pages 1–71. Springer, Cham, 2017.
- [12] V. Calvez, J. A. Carrillo, and F. Hoffmann. Uniqueness of stationary states for singular Keller-Segel type models. *arXiv preprint arXiv:1905.07788*, 2019.
- [13] J. A. Canizo, <https://canizo.org/convolution/>.
- [14] E.A. Carlen, A. Figalli, *Stability for a GNS inequality and the log-HLS inequality, with application to the critical mass Keller–Segel equation*, Duke Math. J. **162** (3) (2013) 579–625.
- [15] E. Carlen, M. Lewin, E.H. Lieb, R. Seiringer, *Stability estimate for the Lane-Emden inequality*, Calc. Var. Partial Differential Equations **64** (2025), no. 7, Paper No. 226.
- [16] J. A. Carrillo, D. Castorina and B. Volzone, *Ground states for diffusion dominated free energies with logarithmic interaction*, SIAM J. Math. Anal. **47** (2015), no. 1, 1–25;
- [17] J. A. Carrillo, M. Fornasier, G. Toscani, and F. Vecil Particle, kinetic, and hydrodynamic models of swarming. In *G. Naldi, L. Pareschi, G. Toscani (eds) Mathematical Modeling of Collective Behavior in Socio-Economic and Life Sciences. Modeling and Simulation in Science, Engineering and Technology*, Birkhäuser Boston, pages 297–336, 2010.
- [18] J. A. Carrillo, S. Hittmeir, B. Volzone, and Y. Yao. Nonlinear aggregation-diffusion equations: radial symmetry and long time asymptotics. *Invent. Math.*, 218 (3): 889–977, 2019.
- [19] J. A. Carrillo, F. Hoffmann, E. Mainini, B. Volzone, *Ground states in the diffusion-dominated regime*, Calc. Var. Partial Differential Equations **57** (2018), no. 5, Paper No. 127, 28 pp.;
- [20] H. Chan, M.d.M. González, Y. Huang, E. Mainini, B. Volzone. Uniqueness of entire ground states for the fractional plasma problem. *Calc. Var. Partial Differential Equations*, 59: 195 (2020).

- [21] M. G. Delgadino, X. Yan, and Y. Yao. Uniqueness and non-uniqueness of steady states of aggregation-diffusion equations. *arXiv preprint arXiv:1908.09782*, 2019.
- [22] F. Demengel and G. Demengel, *Functional spaces for the theory of elliptic partial differential equations*, translated from the 2007 French original by Reinie Ern e, Universitext, Springer, London, 2012 EDP Sci., Les Ulis, 2012;
- [23] E. DiBenedetto, *Real analysis*, second edition, Birkh user Advanced Texts: Basler Lehrb ucher, Birkh user/Springer, New York, 2016;
- [24] J. Dolbeault, B. Perthame, *Optimal critical mass in the two dimensional Keller-Segel model in  $\mathbb{R}^2$* , C. R. Math. Acad. Sci. Paris **339** (2004) 611–616.
- [25] H. Gajewski, K. Zacharias, *Global behaviour of a reaction–diffusion system modelling chemotaxis* Math. Nachr. **195** (1998), 77–114.
- [26] D. Greco, Y. Huang, Z. Liu, V. Moroz, *Ground states of a non-local variational problem and Thomas-Fermi limit for the Choquard equation*, J. London Math. Soc. 111: e70115.
- [27] D. Horstmann, *From 1970 until present: the Keller-Segel model in chemotaxis and its consequences*, I. Jahresber. Deutsch. Math.-Verein. **105** (2003), 103–165
- [28] Y. Huang, E. Mainini, J. L. V azquez, B. Volzone, Nonlinear aggregation-diffusion equations with Riesz potentials, J. Funct. Anal. **287** (2024), no. 2, Paper No. 110465, 59 pp.;
- [29] W. J ager, S. Luckhaus, *On explosions of solutions to a system of partial differential equations modelling chemotaxis*, Trans. Amer. Math. Soc. **329** (1992) 819–824.
- [30] E. Keller, L. Segel, *Initiation of slime mold aggregation viewed as an instability*, J. Theoret. Biol. **26** (1970) 399–415.
- [31] E. Keller, L. Segel, *Model for chemotaxis*, J. Theor. Biol. **30** (1971), 225–234.
- [32] T. Kurokawa, *On the Riesz and Bessel kernels as approximations of the identity*, Sci. Rep. Kagoshima Univ. No. 30 (1981), 31–45;
- [33] G. Leoni, *A first course in fractional Sobolev spaces*, Graduate Studies in Mathematics, 229, Amer. Math. Soc., Providence, RI, (2023);
- [34] E. H. Lieb, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math. (2) **118** (1983), no. 2, 349–374;
- [35] E. H. Lieb, M. Loss, *Analysis*. Second edition. Graduate Studies in Mathematics, **14**. American Mathematical Society, Providence, RI, 2001.
- [36] E. Lieb, S. Oxford, *Improved lower bound on the indirect Coulomb energy*, Int. J. Quantum Chem., **19** (1980), pp. 427–439.
- [37] V. Maz’ya, *Sobolev spaces with applications to elliptic partial differential equations*. Second, revised and augmented edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **342**. Springer, Heidelberg, 2011.
- [38] V. G. Maz’ya, V. P. Havin, *Nonlinear potential theory*, Usp. Mat. Nauk **27** (1972), 67–138 (in Russian). English translation: Russ. Math. Surv. **27** (1972), 71–148.
- [39] Y. Mizuta, *Potential theory in Euclidean spaces*, GAKUTO International Series. Mathematical Sciences and Applications, 6, Gakkotosho, Tokyo, 1996;
- [40] A. Mogilner, L. Edelstein-Keshet, *A nonlocal model for a swarm*, J. Math. Biol. **38** (1999), no. 6, 534–570.
- [41] V. Nanjundiah, *Chemotaxis, signal relaying and aggregation morphology*, J. Theor. Biol. **42** (1973), 63–105.
- [42] C.S. Patlak, *Random walk with persistence and external bias*, Bull. Math. Biophys. **15** (1953), 311–338.
- [43] B. Perthame, *Transport Equations in Biology*, Frontiers in Mathematics. Birkh user, Basel (2006).
- [44] X. Ros-Oton and J. Serra, *Regularity theory for general stable operators*, J. Differential Equations **260** (2016), no. 12, 8675–8715;
- [45] S. G. Samko, A. A. Kilbas and O. I. Marichev, *Fractional integrals and derivatives*, translated from the 1987 Russian original, Gordon and Breach, Yverdon, 1993;
- [46] E. M. Stein, *The characterization of functions arising as potentials*, Bull. Amer. Math. Soc. **67** (1961), 102–104;
- [47] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30, Princeton Univ. Press, Princeton, NJ, 1970;
- [48] Y. Sugiyama, *Global existence in sub-critical cases and finite time blow-up in super-critical cases to degenerate Keller-Segel systems*, Differential Integral Equations **19** (8) (2006) 841–876.

- [49] A. Taheri, *Function spaces and partial differential equations. Vol. 1. Classical analysis*, Oxford Lecture Series in Mathematics and its Applications, 40, Oxford Univ. Press, Oxford, 2015;
- [50] R. Temam. A non-linear eigenvalue problem: the shape at equilibrium of a confined plasma. *Arch. Rational Mech. Anal.*, 60(1):51–73, 1975/76.
- [51] R. Temam. Remarks on a free boundary value problem arising in plasma physics. *Comm. Partial Differential Equations*, 2(6):563–585, 1977.
- [52] H. Triebel, *Theory of function spaces*, reprint of 1983 edition, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2010;

(F. Bozzola) DIME DIPARTIMENTO DI INGEGNERIA MECCANICA, ENERGETICA, GESTIONALE E DEI TRASPORTI  
UNIVERSITÀ DI GENOVA  
VIA ALLA OPERA PIA 15, 16145 GENOVA, ITALY  
*Email address:* [francesco.bozzola@edu.unige.it](mailto:francesco.bozzola@edu.unige.it)

(E. Mainini) DIME DIPARTIMENTO DI INGEGNERIA MECCANICA, ENERGETICA, GESTIONALE E DEI TRASPORTI  
UNIVERSITÀ DI GENOVA  
VIA ALLA OPERA PIA 15, 16145 GENOVA, ITALY  
*Email address:* [edoardo.mainini@unige.it](mailto:edoardo.mainini@unige.it)