

Γ -convergence of free discontinuity problems for circle-valued maps in the linear regime

Giovanni Bellettini*

Roberta Marziani[†]

Riccardo Scala[‡]

October 19, 2026

Abstract

We investigate the Γ -convergence of Ambrosio–Tortorelli-type functionals for circle-valued functions, in the case of volume terms with linear growth. We show the emergence of a nonlocal Γ -limit, which is due to the topological structure of the target space, and discuss compactness of minimal liftings. Our results extend the analysis of a previous work on the quadratic case. Applications to the area functional are given.

Key words: Minimizing liftings, Γ -convergence, $\mathbb{S}^1$ -valued maps, free boundary problems.

AMS (MOS) subject classification: 49Q15, 49Q20, 49J45, 58E12.

1 Introduction

Let $n \geq 1$ and $\Omega \subset \mathbb{R}^n$ be a connected and simply connected bounded open set with Lipschitz boundary, and let $\mathbb{S}^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$ denote the unit circle. For $u : \Omega \rightarrow \mathbb{S}^1$ and $v : \Omega \rightarrow [0, 1]$, consider the family of functionals defined, for $\varepsilon > 0$, by

$$\mathrm{AT}_\varepsilon^{\mathbb{S}^1}(u, v) := \int_{\Omega} \left(v^2 |\nabla u|^2 + \varepsilon |\nabla v|^2 + \frac{(1-v)^2}{4\varepsilon} \right) dx.$$

As in the classical Ambrosio–Tortorelli approximation [6, 7], one observes a decoupling, as $\varepsilon \rightarrow 0^+$, between the bulk term $\int_{\Omega} v^2 |\nabla u|^2 dx$ and the phase-field term $\int_{\Omega} (\varepsilon |\nabla v|^2 + (1-v)^2/\varepsilon) dx$. However, in sharp contrast with the classical setting, [13] shows that for $\mathbb{S}^1$ -valued maps the Γ -limit of $\mathrm{AT}_\varepsilon^{\mathbb{S}^1}$ depends on the choice of the functional domain. More precisely, if one considers $\{(u, v) \in W^{1,1}(\Omega; \mathbb{S}^1) \times W^{1,2}(\Omega) \text{ with } v|\nabla u| \in L^2(\Omega)\}$, as domain, then $\mathrm{AT}_\varepsilon^{\mathbb{S}^1}$ Γ -converges in L^1 to the Mumford–Shah functional for circle-valued maps, namely to

$$\mathrm{MS}_{\mathbb{S}^1}(u) := \int_{\Omega} |\nabla u|^2 dx + \mathcal{H}^{n-1}(S_u), \quad u \in \mathrm{SBV}^2(\Omega; \mathbb{S}^1).$$

*Dipartimento di Scienze Matematiche, Informatiche e Fisiche, via delle Scienze 206, 33100 Udine UD, Italy, and ICTP International Centre for Theoretical Physics, Mathematics Section, Strada Costiera 11, I-34151 Trieste Italy. E-mail: giovanni.bellettini@uniud.it

[†]Dipartimento di Matematica, Università degli studi di Trento, 38123 Povo (TN), Italy. E-mail: roberta.marziani@unitn.it

[‡]Dipartimento di Ingegneria dell’Informazione e Scienze Matematiche, Università di Siena, 53100 Siena, Italy. E-mail: riccardo.scala@unisi.it

If, instead, one restricts to the smaller domain $\{(u, v) \in W^{1,2}(\Omega; \mathbb{S}^1) \times W^{1,2}(\Omega)\}$, then $\text{AT}_\varepsilon^{\mathbb{S}^1}$ Γ -converges in L^1 to the nonlocal functional

$$\text{MS}_{\text{lift}}(u) := \int_{\Omega} |\nabla u|^2 dx + m_2[u], \quad u \in SBV^2(\Omega; \mathbb{S}^1),$$

where, denoting by S_φ the jump set of a function φ ,

$$m_2[u] := \inf \{ \mathcal{H}^{n-1}(S_\varphi) : \varphi \in GSBV^2(\Omega), e^{i\varphi} = u \text{ a.e. in } \Omega \}. \quad (1.1)$$

Since $S_u \subseteq S_\varphi$ for any lifting φ of u , it follows $\text{MS}_{\text{lift}}(u) \geq \text{MS}_{\mathbb{S}^1}(u)$, possibly with the strict inequality for appropriate maps u .

This dichotomy, sometimes referred to as the Lavrentiev phenomenon, reflects the influence of the topological structure of $\mathbb{S}^1$, in particular the possibility of a nonzero degree for circle-valued maps. In the first case, one has more freedom in the construction of the recovery sequence, and it is possible to preserve the degree throughout the approximation procedure. In contrast, in the second case, since a map $u \in W^{1,2}(\Omega; \mathbb{S}^1)$ has zero topological degree whenever Ω is simply connected (see, e.g., [18]), one is forced to approximate maps of nonzero degree by maps of zero degree, which entails an additional energetic cost. This phenomenon is clearly illustrated by the example of the vortex map described in [13] and also occurs in other variational problems concerning the relaxation of the area functional in codimension two [9] and the study of ferromagnetic spin systems [22].

The minimisation problem in (1.1) can be formulated for any $u \in SBV^p(\Omega; \mathbb{S}^1)$ with $p > 1$ by minimizing over all liftings $\varphi \in GSBV^p(\Omega)$; we denote the corresponding value by $m_p[u]$. In [13] it is proven that $m_p[u]$ admits a minimizer which in general does not belong to $SBV^p(\Omega)$, and that such minimisers are related either to suitably defined optimal transport problems (when $n = 2$) or to certain solutions of the Plateau problem (when $n > 2$).

Linear growth setting. In the present paper we consider a variant where the quadratic term $|\nabla u|^2$ is replaced by a term of the form $f(|\nabla u|)$, with f having linear growth at infinity (see Section 3 for the precise assumptions). Similarly to the unconstrained case [2, 3], the linear growth introduces a genuine interaction between the bulk term and the phase-field term which is absent in the quadratic case. In addition, the resulting Γ -limit is defined on the larger space $BV(\Omega; \mathbb{S}^1)$ and thus has a Cantor part.

More precisely, by allowing also for more general phase-field terms (cf. Section 3), our main result Theorem 3.1 states that if one restricts to the domain $(u, v) \in W^{1,2}(\Omega; \mathbb{S}^1) \times W^{1,2}(\Omega)$, then the Γ -limit is the nonlocal functional

$$\int_{\Omega} f(|\nabla u|) dx + |D^c u|(\Omega) + m_g[u], \quad u \in BV(\Omega; \mathbb{S}^1),$$

where $m_g[u]$ is expressed as a minimization problem associated to the cost function g defined in (2.3), that is

$$m_g[u] := \inf \left\{ \int_{S_\varphi} g(|\varphi^+ - \varphi^-|) d\mathcal{H}^{n-1} : \varphi \in GBV(\Omega), e^{i\varphi} = u \text{ a.e. in } \Omega \right\}. \quad (1.2)$$

We also prove that $m_g[u]$ is actually a minimum, i.e., we prove the existence of a minimal lifting for $m_g[u]$. As a byproduct of this, we establish compactness results for sequences of liftings in the linear-growth setting (Theorems 4.2 and 4.5), extending the results of [13]

to the full BV -framework. In particular, we show that sequences of liftings of a given map in $BV(\Omega; \mathbb{S}^1)$, for which we control the portion of jump set with jump amplitude greater than $\sigma > 0$ are precompact with respect to the local convergence modulo 2π (see Definition 4.1), and moreover, that the same holds for sequences of liftings corresponding to $u_k \xrightarrow{*} u$ in $BV(\Omega; \mathbb{S}^1)$.

The linear growth case entails substantial differences in the proof strategy. First, compactness results for liftings in the class GBV are required, which rely on slicing arguments and measure-theoretic localization techniques. Moreover, the genuine interaction between the bulk and the phase-field terms prevents a direct adaptation of the quadratic-growth analysis and necessitates a localized approach to Γ -convergence. These tools allow us to simplify several arguments with respect to the quadratic case, while at the same time extending the theory to a broader and more flexible variational framework. In addition, our analysis reveals natural connections with optimal transport problems and with the relaxation of the area functional in $\mathbb{S}^1$. We briefly discuss these relations below and defer a complete investigation to future work.

Connections with the relaxation of the area functional in $\mathbb{S}^1$. The relaxed area in $\mathbb{S}^1$ of the graphs of L^1 -functions $u: \Omega \rightarrow \mathbb{S}^1$ is defined by

$$\mathcal{A}^{\mathbb{S}^1}(u, \Omega) := \inf \left\{ \liminf_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla u_k|^2} \, dx : u_k \in C^1(\Omega; \mathbb{S}^1), \, u_k \rightarrow u \text{ in } L^1 \right\}.$$

This functional was characterised in terms of Cartesian currents (cf. [28]). Alternatively one could propose a characterisation using lifting theory. For instance, if we further assume that Ω is simply connected, then we show that (see Theorem 7.1)

$$\mathcal{A}^{\mathbb{S}^1}(u, \Omega) = \int_{\Omega} \sqrt{1 + |\nabla u|^2} \, dx + |D^c u|(\Omega) + m[u],$$

where

$$m[u] := \min \left\{ \int_{S_{\varphi}} |\varphi^+ - \varphi^-| \, d\mathcal{H}^{n-1} : \varphi \in BV(\Omega), \, e^{i\varphi} = u \text{ a.e. in } \Omega \right\}.$$

The similarity with (1.2) (when $f(t) = \sqrt{1 + t^2}$) suggests the possibility of approximating $\mathcal{A}^{\mathbb{S}^1}$ through suitably chosen linear-growth Ambrosio–Tortorelli type functionals. Indeed, using a diagonal argument, we provide such an approximation in Theorem 7.2, opening interesting perspectives for future research related to the relaxation of the area functional in codimension two (see [1, 8–12, 14, 15, 20, 31, 32] and references therein for related results).

Connections with optimal transport. The quantity $m_g[u]$ admits a natural interpretation as an optimal transport problem with cost g . Assume for simplicity $\Omega \subset \mathbb{R}^2$ and let $u \in W^{1,1}(\Omega; \mathbb{S}^1)$ have distributional Jacobian of the form

$$\text{Det}(\nabla u) = \pi \sum_{i=1}^N (\delta_{x_i} - \delta_{y_i}), \quad x_i, y_i \in \Omega.$$

Setting $\mu := \sum_{i=1}^N \delta_{x_i}$ and $\nu := \sum_{i=1}^N \delta_{y_i}$ one can show that

$$m_g[u] = \inf \int_R g(\theta) \, d\mathcal{H}^1,$$

where the infimum is taken over all integer multiplicity 1-currents $T = (R, \theta, \tau)$ with $\partial T = \mu - \nu$. We refer to [13] for the discussion on the equivalence between the minimization problem $m_g[u]$ and the optimal transport problem with cost function $g \equiv 1$. For optimal transport problems in this setting we refer to [19, 27, 29, 30] and references therein.

Let us also mention that, due to its property of approximation of the Mumford-Shah type energy of circle-valued maps, as observed in [13], our Ambrosio-Tortorelli functional has a strict connection with analysis of Ginzburg-Landau energies and dislocation mechanics (we refer to [16] for the former topic). Specifically, the Mumford-Shah functional for $\mathbb{S}^1$ -valued maps has been successfully used to model the appearance of dislocations in 2-dimensional domains (see [23, 25] for more details; we also refer to the introduction of [13] for a more exhaustive discussion on the link with vortex singularities).

Content of the paper. In Section 2 we collect some notation and recall some useful results needed to prove our main results. In Section 3 we state our main result Theorem 3.1. In Section 4 we state two compactness results for sequences of liftings (Theorem 4.2 and Theorem 4.5). We also state and prove existence of a solution to (1.2). Section 5.2 is devoted to the proof of Theorems 4.2 and 4.5. In Section 6 we provide the proof of Theorem 3.1. Eventually, in Section 7 we prove Theorems 7.1 and 7.2.

2 Notation and preliminaries

We start to recall some notions concerning BV and GBV functions [5] and lifting theory [18]. In what follows: $\partial^* A$ denotes the reduced boundary of a finite perimeter set $A \subset \mathbb{R}^n$, $|\cdot|$ and $\mathcal{H}^{n-1}$ denote the Lebesgue measure and the $(n-1)$ -dimensional Hausdorff measure in $\mathbb{R}^n$, respectively, and χ_A denotes the characteristic function of the set $A \subset \mathbb{R}^n$. If $\Omega \subseteq \mathbb{R}^n$ is an open set we denote by $\mathcal{A}(\Omega)$ the family of all open subsets of Ω .

We recall the following lemma [17, Proposition 1.16]:

Lemma 2.1 (Localisation). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with Lipschitz boundary. Let $\mu: \mathcal{A}(\Omega) \rightarrow [0, +\infty)$ be a superadditive set function on disjoint open sets, λ be a positive measure on Ω and, for $k \in \mathbb{N}$, $h_k: \Omega \rightarrow [0, +\infty]$ be a Borel function such that*

$$\mu(A) \geq \int_A h_k d\lambda \quad \text{for every } A \in \mathcal{A}(\Omega) \text{ and } k \in \mathbb{N}.$$

Then

$$\mu(A) \geq \int_A \sup_{k \geq 1} h_k d\lambda \quad \text{for every } A \in \mathcal{A}(\Omega).$$

2.1 BV and GBV functions

Let $U \subset \mathbb{R}^n$ be open and bounded, and $m \geq 1$ be an integer. We denote by $BV(U; \mathbb{R}^m)$ the space of vector-valued $L^1(U; \mathbb{R}^m)$ -functions with bounded variation in U , and with $|\cdot|_{BV}$ and $\|\cdot\|_{BV}$ the BV seminorm and norm in U , respectively, i.e.

$$|u|_{BV} := |Du|(U), \quad \|u\|_{BV} := \|u\|_{L^1} + |u|_{BV},$$

see [5]. We recall that if $u \in BV(U; \mathbb{R}^m)$ then its distributional gradient is a $\mathbb{R}^{m \times n}$ -valued Radon measure with finite total variation, and it can be decomposed as

$$Du = \nabla u \mathcal{L}^n + D^c u + \llbracket u \rrbracket \otimes \nu_u \mathcal{H}^{n-1} \llcorner S_u,$$

where ∇u is the approximate gradient of u , $D^c u$ the Cantor part, S_u is the approximate jump set of u , $\llbracket u \rrbracket = u^+ - u^-$ is the jump opening, ν_u is a unit normal to the jump set S_u of u and $\otimes$ stands for the tensor product, see [5, Def. 3.67]. We say that $u \in BV(U; \mathbb{R}^m)$ is a special function of bounded variation in U if $D^c u = 0$. We denote by $SBV(U; \mathbb{R}^m)$ the space of vector-valued *special functions with bounded variation in U* . For $p > 1$ we define also the space

$$SBV^p(\Omega; \mathbb{R}^m) = \{u \in SBV(\Omega; \mathbb{R}^m) : |\nabla u| \in L^p(\Omega), \mathcal{H}^{n-1}(S_u) < +\infty\}.$$

A measurable function $u : U \rightarrow \mathbb{R}^m$ belongs to the space of *generalised functions with bounded variation in U* , that is, $u \in GBV(U; \mathbb{R}^m)$, if $\phi \circ u \in BV_{\text{loc}}(U)$ for any $\phi \in C^1(\mathbb{R}^m)$ with $\nabla \phi$ compactly supported.¹ Note that also in GBV we can define the approximate gradient, the Cantor part and the jump set. Moreover $GBV(\Omega; \mathbb{R}^m) \cap L^\infty(\Omega; \mathbb{R}^m) = BV(\Omega; \mathbb{R}^m) \cap L^\infty(\Omega; \mathbb{R}^m)$. Analogously one can define $GSBV(U; \mathbb{R}^m)$ and $GSBV^p(U; \mathbb{R}^m)$. If $m = 1$ we write $BV(U) = BV(U; \mathbb{R})$, and $GBV(U) = GBV(U; \mathbb{R})$.

We set

$$BV(U; \mathbb{S}^1) = \{u \in BV(U; \mathbb{R}^2) : |u| = 1 \text{ a.e. in } U\}.$$

Eventually, a (finite or countable) family $\{E_i\}$ of finite perimeter subsets of a finite perimeter set $F \subseteq \mathbb{R}^n$ is called a Caccioppoli partition of F if the sets E_i are pairwise disjoint, their union is F , and the sum of their perimeters is finite.

2.2 Approximation and compactness for the vector Mumford-Shah functional

We recall a result due to Alicandro-Braides-Shah [2], subsequently extended to the vector case by Alicandro-Focardi [3]. Let $m \geq 1$, and

$$\mathcal{D} := \{(u, v) \in W^{1,2}(\Omega; \mathbb{R}^m) \times W^{1,2}(\Omega) : 0 \leq v \leq 1\}.$$

Consider the family of functionals $F_\varepsilon : L^1(\Omega; \mathbb{R}^m) \times L^1(\Omega) \rightarrow [0, +\infty]$ given by

$$F_\varepsilon(u, v) := \begin{cases} \int_{\Omega} \left(\psi(v) f(|\nabla u|) + \varepsilon |\nabla v|^2 + \frac{W(v)}{\varepsilon} \right) dx & \text{if } (u, v) \in \mathcal{D}, \\ +\infty & \text{otherwise in } L^1(\Omega; \mathbb{R}^m) \times L^1(\Omega), \end{cases} \quad (2.1)$$

where

- (a) $\psi : [0, 1] \rightarrow [0, 1]$ is an increasing lower semicontinuous function with $\psi(0) = 0$, $\psi(1) = 1$ and $\psi(t) > 0$ if $t \in (0, 1)$;
- (b) $f : [0, +\infty) \rightarrow [0, +\infty)$ is a convex increasing function such that

$$\lim_{t \rightarrow +\infty} \frac{f(t)}{t} = 1;$$

- (c) $W : [0, 1] \rightarrow [0, +\infty)$ is a continuous function such that $W(s) = 0$ if and only if $s = 1$.

¹Recall that $f \in BV_{\text{loc}}(U)$ if $f \in BV(U')$ for every $U' \subset U$ open with $\overline{U'}$ compact in U .

Set also

$$c_W(t) := 2 \int_t^1 \sqrt{W(s)} \, ds, \quad (2.2)$$

and

$$g(z) := \min\{\psi(t)z + 2c_W(t) : 0 \leq t \leq 1\}. \quad (2.3)$$

Then the following theorem holds.

Theorem 2.2. *Let $\Omega \subseteq \mathbb{R}^n$ be an open set with Lipschitz boundary. Then $F_\varepsilon \Gamma - L^1$ -converge to the functional F defined as*

$$F(u, v) := \begin{cases} \int_\Omega f(|\nabla u|) \, dx + |D^c u|(\Omega) + \int_{S_u} g(|u^+ - u^-|) \, d\mathcal{H}^{n-1} & \text{if } u \in GBV(\Omega; \mathbb{R}^m), v = 1 \text{ a.e.} \\ +\infty & \text{otherwise in } L^1(\Omega; \mathbb{R}^m) \times L^1(\Omega) \end{cases}$$

as $\varepsilon \rightarrow 0^+$.

Remark 2.3. By inspecting the proof of Theorem 2.2 one actually deduces the following properties:

- (i) **Lower bound on open sets.** The lower bound inequality holds on every $A \in \mathcal{A}(\Omega)$. Precisely, let $F_\varepsilon(\cdot, \cdot, A)$ and $F(\cdot, \cdot, A)$ denote the localised functionals on the open set $A \subseteq \Omega$ and suppose $(u_\varepsilon, v_\varepsilon) \rightarrow (u, v)$ in $L^1(\Omega; \mathbb{R}^m) \times L^1(\Omega)$. Then

$$\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, v_\varepsilon, A) \geq F(u, v, A);$$

- (ii) **g is a truncating function.** The function g in (2.3) satisfies the following properties:

1. g is increasing, $g(0) = 0$ and $\lim_{z \rightarrow +\infty} g(z) = 2c_W(0)$;
2. g is subadditive, i.e., $g(z_1 + z_2) \leq g(z_1) + g(z_2)$ for all $z_1, z_2 \in [0, +\infty)$;
3. g is Lipschitz-continuous with Lipschitz constant 1;
4. $g(z) \leq z$ for all $z \in [0, +\infty)$ and $\lim_{z \rightarrow 0^+} g(z)/z = 1$;
5. For any $T > 0$ there exists $c_T > 0$ such that $z \leq c_T g(z)$ for all $z \in [0, T]$;
6. For $\sigma > 0$ there exists $C_\sigma > 0$ such that $g(z) \geq C_\sigma$ for all $z \geq \sigma > 0$.

Properties 1 – 5 follow by [2, Remark 4.2]. Property 6 follows taking $C_\sigma = g(\sigma)$.

2.3 Liftings of $\mathbb{S}^1$ -valued maps

Let $\Omega \subset \mathbb{R}^n$ be a bounded connected and simply connected open set with Lipschitz boundary, and $u : \Omega \rightarrow \mathbb{S}^1$ be a measurable function. A *lifting* of u is a measurable function $\varphi : \Omega \rightarrow \mathbb{R}$ such that

$$u(x) = e^{i\varphi(x)} \quad \text{for a.e. } x \in \Omega.$$

Given a Borel set $B \subseteq \Omega$, we say that $\varphi : B \rightarrow \mathbb{R}$ is a lifting of u on B if the previous equality holds a.e. on B . Clearly, if φ is a lifting of u , then so is $\varphi + 2\pi m$ for all $m \in \mathbb{Z}$.

If u has some regularity, a natural question is whether φ can be chosen with the same regularity. The answer is partially positive, see [18, 24] for more details: we recall in particular that if $n \geq 2$ and $u \in W^{1,p}(\Omega; \mathbb{S}^1)$ for some $p \in [2, +\infty)$, then, being Ω simply connected, u has a lifting $\varphi \in W^{1,p}(\Omega)$, and φ is unique (mod 2π). Moreover, for $n = 2$

and $p \in [1, 2)$ there exists $u \in W^{1,p}(\Omega; \mathbb{S}^1)$ for which there are no liftings in $W^{1,p}(\Omega)$, see [18, Theorem 1.2, Remark 1.9]. Indeed it can be shown [18, pages 17-19] that there are no liftings of the vortex map $u_V(x) = x/|x|$ in $W^{1,1}(B_1)$ (and thus there are no liftings in $W^{1,p}(B_1)$ for all $p \in [1, 2)$).

Next we recall the

Theorem 2.4 (Davila-Ignat). *Let $u \in BV(\Omega; \mathbb{S}^1)$. Then there exists a lifting $\varphi \in BV(\Omega)$ such that $\|\varphi\|_{L^\infty} \leq 2\pi$ and $|\varphi|_{BV} \leq 2|u|_{BV}$.*

Proof. See [24, Theorem 1.1], and also [18, Theorem 1.4]. $\square$

As usual, for any lifting $\varphi \in SBV(\Omega)$ of u we write² $S_\varphi = S_\varphi^I \cup S_\varphi^f$ where

$$S_\varphi^I := \{x \in S_\varphi : \llbracket \varphi \rrbracket(x) \in 2\pi\mathbb{Z}\}, \quad S_\varphi^f := S_\varphi \setminus S_\varphi^I.$$

In particular $S_\varphi^f = S_u$; moreover $|\nabla u| = |\nabla \varphi|$ a.e. in Ω . If $u \in BV(\Omega; \mathbb{S}^1)$, also $|D^c u|(\Omega) = |D^c \varphi|(\Omega)$.

Let $u \in BV(\Omega; \mathbb{S}^1)$ and consider the minimum problem

$$\inf\{|\varphi|_{BV} : \varphi \in BV(\Omega), e^{i\varphi} = u \text{ a.e. in } \Omega\}.$$

Then there exists a minimizer $\varphi \in BV(\Omega)$ such that $|\varphi|_{BV} \leq 2|u|_{BV}$ and $0 \leq \int_\Omega \varphi \, dx \leq 2\pi|\Omega|$ [18, page 25]. Moreover, as already mentioned in the introduction, in [13] it was proven that for any $p > 1$ and any $u \in SBV^p(\Omega; \mathbb{S}^1)$ the minimisation problem

$$m_p[u] := \inf\{\mathcal{H}^{n-1}(S_\varphi) : \varphi \in GSBV^p(\Omega), e^{i\varphi} = u \text{ a.e. in } \Omega\}$$

admits a solution. Here, we are instead concerned with the existence of minimal lifting for a minimisation problem which depends on the cost function g in (2.3) (cf. (1.2) for its precise definition).

3 Statement of the main result

Let $n \geq 2$ and $\Omega \subset \mathbb{R}^n$ be a connected bounded open set with Lipschitz boundary. To the pair $u : \Omega \rightarrow \mathbb{S}^1$, $v : \Omega \rightarrow [0, 1]$ we associate the functional

$$F_\varepsilon(u, v) := \int_\Omega \left(\psi(v)f(|\nabla u|) + \varepsilon|\nabla v|^2 + \frac{W(v)}{\varepsilon} \right) dx,$$

where $\varepsilon \in (0, 1]$, ψ , f and W satisfy properties (a), (b) and (c) in Section 2.2.

We consider the functional domain given by

$$\mathcal{D}_{\mathbb{S}^1} := \{(u, v) \in W^{1,2}(\Omega; \mathbb{S}^1) \times W^{1,2}(\Omega) : 0 \leq v \leq 1\} \subset L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega). \quad (3.1)$$

For $\varepsilon \in (0, 1]$ let us consider the corresponding family of functionals

$$F_\varepsilon^{\mathbb{S}^1} : L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega) \rightarrow [0, +\infty]$$

given by

$$F_\varepsilon^{\mathbb{S}^1}(u, v) := \begin{cases} F_\varepsilon(u, v) & \text{if } (u, v) \in \mathcal{D}_{\mathbb{S}^1}, \\ +\infty & \text{otherwise in } L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega). \end{cases} \quad (3.2)$$

The main result of the present paper reads as follows.

² S_φ^I stands for the “integer” part of the jump, and S_φ^f for the “fractional” part.

Theorem 3.1 (Γ -convergence of $F_\varepsilon^{\mathbb{S}^1}$). *Provided Ω is simply-connected, we have*

$$\Gamma - L^1 \lim_{\varepsilon \rightarrow 0^+} F_\varepsilon^{\mathbb{S}^1} = F_{\text{lift}},$$

where

$$F_{\text{lift}}(u, v) := \begin{cases} \int_{\Omega} f(|\nabla u|) \, dx + |D^c u|(\Omega) + m_g[u] & \text{if } u \in BV(\Omega; \mathbb{S}^1), v = 1 \text{ a.e.}, \\ +\infty & \text{otherwise in } L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega), \end{cases} \quad (3.3)$$

and $m_g[u]$ is defined in (1.2). In addition, $m_g[u]$ is a minimum.

We note that the admissible set in (1.2) is nonempty, according to Theorem 2.4. As an outcome of our analysis, we derive compactness results for sequences of liftings (see Theorems 4.2 and 4.5). In particular, Theorem 4.5 will play a key role in establishing the lower bound in Theorem 3.1.

4 On g -minimizing liftings

In this section we establish the existence of solutions to the minimization problem (1.2), for g as in (2.3) and $u \in BV(\Omega; \mathbb{S}^1)$, as stated in Corollary 4.3. For later use, we introduce the following notion of convergence.

Definition 4.1 (Local convergence modulo 2π). *Let $(\varphi_k)_{k \geq 1} \subset GBV(\Omega)$ and $\varphi_\infty \in GBV(\Omega)$. We say that (φ_k) converges locally modulo 2π to φ_∞ (in Ω) if the following holds: there exist a Caccioppoli partition $\{E_i\}_{i \geq 1}$ of Ω and sequences $(d_k^{(i)})_{k \geq 1} \subset \mathbb{Z}$ for $i \in \mathbb{N}$ such that*

$$\begin{aligned} \lim_{k \rightarrow +\infty} (\varphi_k(x) - 2\pi d_k^{(i)}) &= \varphi_\infty(x) \quad \forall i \in \mathbb{N}, \text{ for a.e. } x \in E_i, \\ \lim_{k \rightarrow +\infty} |\varphi_k(x) - 2\pi d_k^{(i)}| &= +\infty \quad \forall i \in \mathbb{N}, \text{ for a.e. } x \in \Omega \setminus E_i. \end{aligned} \quad (4.1)$$

Theorem 4.2 (Compactness). *Let $0 \leq \sigma \leq \pi$. Let $u \in BV(\Omega; \mathbb{S}^1)$ and $(\varphi_k)_{k \geq 1} \subset GBV(\Omega)$ be a sequence of liftings of u in Ω with*

$$H := \sup_{k \in \mathbb{N}} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) < +\infty, \quad (4.2)$$

where

$$S_{\varphi_k}^\sigma := \{x \in S_{\varphi_k} : |\llbracket \varphi_k(x) \rrbracket| \geq \sigma\}. \quad (4.3)$$

Then there exist a (not-relabelled) subsequence of indices k and a lifting $\varphi_\infty \in GBV(\Omega)$ of u in Ω such that (φ_k) converges locally modulo 2π to φ_∞ .

A first consequence of Theorem 4.2 is the following result.

Corollary 4.3 (Existence of a minimizer to $m_g[u]$). *Let g be as in (2.3). Let $u \in BV(\Omega; \mathbb{S}^1)$ and let $(\varphi_k)_{k \geq 1} \subset GBV(\Omega)$ be a sequence of liftings of u in Ω with*

$$C := \sup_{k \in \mathbb{N}} \int_{S_{\varphi_k}} g(|\varphi_k^+ - \varphi_k^-|) \, d\mathcal{H}^{n-1} < +\infty. \quad (4.4)$$

Then the following holds:

(i) *Compactness:* There exist a (not-relabelled) subsequence of indices k and a lifting $\varphi_\infty \in GBV(\Omega)$ of u in Ω such that (φ_k) converges locally modulo 2π to φ_∞ ;

(ii) *Lower semicontinuity:*

$$\liminf_{k \rightarrow +\infty} \int_{S_{\varphi_k}} g(|\varphi_k^+ - \varphi_k^-|) d\mathcal{H}^{n-1} \geq \int_{S_{\varphi_\infty}} g(|\varphi_\infty^+ - \varphi_\infty^-|) d\mathcal{H}^{n-1}.$$

As a consequence, the infimum defining $m_g[u]$ is attained in $GBV(\Omega)$.

Proof. (i). Let $0 < \sigma \leq \pi$ be fixed. Remark 2.3-(ii) and (4.4) yield

$$C \geq \int_{S_{\varphi_k}} g(|\varphi_k^+ - \varphi_k^-|) d\mathcal{H}^{n-1} \geq C_\sigma \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma).$$

and the assertion follows from Theorem 4.2.

(ii). Let φ_∞ , $\{E_i\}_{i \geq 1}$ and $(d_k^{(i)})_{k \geq 1}$ be as in (4.1). We exploit the fact that g arises from the Γ -limit of the functionals in (2.1). Fix $N, i \in \mathbb{N}$ and set

$$\varphi_k^{i,N} := ((\varphi_k - 2\pi d_k^{(i)}) \wedge N) \vee (-N).$$

Then $\varphi_k^{i,N} \rightarrow \varphi_\infty^{i,N}$ in $L^1(\Omega)$ with $\varphi_\infty^{i,N} := (\varphi_\infty \wedge N) \vee (-N)$ in E_i and $\varphi_\infty^{i,N} = \pm N$ in two suitable finite perimeter subsets of Ω the complement (in Ω) of the union of which is E_i .

Define

$$G(\phi, A) := F(\phi, 1, A) \quad \text{for } \phi \in GBV(\Omega), A \in \mathcal{A}(\Omega)$$

where $F(\cdot, \cdot, A)$ is as in Remark 2.3-(i) with $f(t) = t$. From Theorem 2.2 and properties of Γ -convergence it follows that $G(\cdot, A)$ is lower semicontinuous in the L^1 -topology, so that

$$\liminf_{k \rightarrow +\infty} G(\varphi_k, A) \geq \liminf_{k \rightarrow +\infty} G(\varphi_k^{i,N}, A) \geq G(\varphi_\infty^{i,N}, A), \quad A \in \mathcal{A}(\Omega). \quad (4.5)$$

Next we define the superadditive set function

$$\mu(A) := \liminf_{k \rightarrow +\infty} G(\varphi_k, A) \quad \forall A \in \mathcal{A}(\Omega), \quad (4.6)$$

and the positive measure

$$\begin{aligned} \lambda(B) := G(\varphi_\infty, B) &= \int_B |\nabla \varphi_\infty| dx + |D^c \varphi_\infty|(B) + \int_{(S_{\varphi_\infty} \setminus \Sigma) \cap B} g(|\varphi_\infty^+ - \varphi_\infty^-|) d\mathcal{H}^{n-1} \\ &\quad + 2c_W(0) \mathcal{H}^{n-1}(\Sigma \cap B), \end{aligned}$$

for every Borel set $B \subseteq \Omega$, where we have denoted

$$\Sigma := \left(\bigcup_{i=1}^{+\infty} \partial^* E_i \right) \setminus \partial \Omega. \quad (4.7)$$

We identify E_i with the set of points of density 1 for χ_{E_i} . Define also, for $i \geq 1$,

$$h_1^i := \begin{cases} 1 & \text{in } E_i \setminus (\Sigma \cup S_{\varphi_\infty}) \\ 0 & \text{otherwise in } \Omega \end{cases},$$

and

$$h_2^i := \begin{cases} 1 & \text{in } S_{\varphi_\infty} \cap E_i \\ 0 & \text{otherwise in } \Omega \end{cases}, \quad h_3^i := \begin{cases} 1 & \text{in } (\partial^* E_i) \setminus \partial\Omega \\ 0 & \text{otherwise in } \Omega. \end{cases}$$

By construction we have

$$\lim_{N \rightarrow +\infty} |(\varphi_\infty^{i,N})^+ - (\varphi_\infty^{i,N})^-| = |\varphi_\infty^+ - \varphi_\infty^-| \quad \mathcal{H}^{n-1} \text{ a.e. in } (S_{\varphi_\infty} \cap E_i),$$

and

$$\lim_{N \rightarrow +\infty} |(\varphi_\infty^{i,N})^+ - (\varphi_\infty^{i,N})^-| = +\infty \quad \mathcal{H}^{n-1} \text{ a.e. in } \partial^* E_i,$$

hence by monotonicity of g and the monotone convergence theorem we arrive at

$$\lim_{N \rightarrow +\infty} \int_{(S_{\varphi_\infty} \cap E_i) \cap A} g(|(\varphi_\infty^{i,N})^+ - (\varphi_\infty^{i,N})^-|) d\mathcal{H}^{n-1} \geq \int_A h_2^i d\lambda$$

and

$$2c_W(0)\mathcal{H}^{n-1}(A \cap \partial^* E_i) = \lim_{N \rightarrow \infty} \int_{A \cap \partial^* E_i} g(|(\varphi_\infty^{i,N})^+ - (\varphi_\infty^{i,N})^-|) d\mathcal{H}^{n-1} \geq \int_A h_3^i d\lambda.$$

Whence, by (4.5) we conclude

$$\mu(A) \geq \int_A h_2^i d\lambda, \quad \mu(A) \geq \int_A h_3^i d\lambda, \quad A \in \mathcal{A}(\Omega).$$

On the other hand, still by (4.5) it is easily seen that

$$\mu(A) \geq \int_A h_1^i d\lambda \quad \forall i \geq 1, \quad A \in \mathcal{A}(\Omega).$$

Therefore, since

$$\sup_{j=1,2,3, i \in \mathbb{N}} h_j^i \equiv 1,$$

using Lemma 2.1 we infer

$$\mu(A) \geq \lambda(A) \quad \text{for all } A \in \mathcal{A}(\Omega).$$

As a consequence we get

$$\liminf_{k \rightarrow +\infty} G(\varphi_k, \Omega) \geq G(\varphi_\infty, \Omega). \quad (4.8)$$

Since $\varphi_k, \varphi_\infty$ are liftings of u in Ω it holds $|\nabla \varphi_k| = |\nabla \varphi_\infty| = |\nabla u|$ and $|D^c \varphi_k|(\Omega) = |D^c \varphi_\infty|(\Omega) = |D^c u|(\Omega)$, hence (4.8) in turn implies

$$\liminf_{k \rightarrow +\infty} \int_{S_{\varphi_k}} g(|\varphi_k^+ - \varphi_k^-|) d\mathcal{H}^{n-1} \geq \int_{S_{\varphi_\infty}} g(|\varphi_\infty^+ - \varphi_\infty^-|) d\mathcal{H}^{n-1},$$

and the proof of (ii) is concluded. $\square$

From the proof above we also obtain the following:

Corollary 4.4. *In the same hypotheses and notations of Corollary 4.3, we have*

$$\liminf_{k \rightarrow +\infty} \int_{S_{\varphi_k}} g(|\varphi_k^+ - \varphi_k^-|) d\mathcal{H}^{n-1} \geq \int_{S_{\varphi_\infty} \setminus \Sigma} g(|\varphi_\infty^+ - \varphi_\infty^-|) d\mathcal{H}^{n-1} + 2c_W(0)\mathcal{H}^{n-1}(\Sigma),$$

where we recall $2c_W(0) = \sup g$ and Σ is defined in (4.7).

Now we state a generalization of Theorem 4.2 to a sequence (u_k) , needed to show Theorem 3.1; the proof will be given in the next section.

Theorem 4.5 (Generalised compactness). *Let $u, u_k \in BV(\Omega; \mathbb{S}^1)$ be such that*

$$u_k \xrightarrow{*} u \quad \text{in } BV(\Omega; \mathbb{S}^1), \quad (4.9)$$

and let $\varphi_k \in GBV(\Omega)$ be a lifting of u_k in Ω , for all $k \geq 1$. Suppose that for some $0 \leq \sigma \leq \pi$

$$H := \sup_{k \in \mathbb{N}} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) < +\infty. \quad (4.10)$$

Then there exist a (not relabelled) subsequence of indices k and a lifting $\varphi_\infty \in GBV(\Omega)$ of u in Ω such that φ_k converges locally modulo 2π to φ_∞ .

5 Proofs of the compactness Theorems 4.2 and 4.5

We split the proof of Theorems 4.2 and 4.5 into a number of intermediate lemmas.

Lemma 5.1 (Localized compactness). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with Lipschitz boundary, and $u \in BV(\Omega; \mathbb{S}^1)$. Let $0 \leq \sigma \leq \pi$ and $(\varphi_k)_{k \geq 1} \subset GBV(\Omega)$ be a sequence of liftings of u satisfying (4.2). Let $F \subset \Omega$ be a nonempty finite perimeter set in Ω . Then, for a not-relabelled subsequence, there exist a sequence $(d_k)_{k \geq 1} \subset \mathbb{Z}$, a finite perimeter set*

$$E \subseteq F$$

in Ω , and a function $\varphi_\infty \in GBV(\Omega)$, such that:

$$\varphi_\infty \text{ is a lifting of } u \text{ in } E, \quad \varphi_\infty = 0 \quad \text{in } \Omega \setminus E, \quad (5.1)$$

and

$$\begin{aligned} \lim_{k \rightarrow +\infty} (\varphi_k(x) - 2\pi d_k) &= \varphi_\infty(x) && \text{for a.e. } x \in E, \\ \lim_{k \rightarrow +\infty} |\varphi_k(x) - 2\pi d_k| &= +\infty && \text{for a.e. } x \in F \setminus E, \end{aligned} \quad (5.2)$$

$$|E| \geq \frac{n^n \omega_n |F|^n}{2^n (2H + \mathcal{H}^{n-1}(\partial^* F))^n} > 0, \quad (5.3)$$

If furthermore

$$|\varphi_k(x) - 2\pi d_k| \rightarrow +\infty \quad \text{for a.e. } x \in \Omega \setminus E, \quad (5.4)$$

and $\{x \in \Omega : \varphi_k(x) - 2\pi d_k \rightarrow \pm\infty\}$ are finite perimeter sets in Ω , then

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma \cap A) \geq \mathcal{H}^{n-1}(A \cap \partial^* E) \quad \text{for any } A \in \mathcal{A}(\Omega), \quad (5.5)$$

and

$$\mathcal{H}^{n-1}(F \cap \partial^* E) = \mathcal{H}^{n-1}(F \cap \partial^*(F \setminus E)) \leq C. \quad (5.6)$$

Proof. Following the lines of the proof of [13, Lemma 4.1] and noticing that [13, equation 4.8] still holds if we replace S_{φ_k} and S_{φ_1} with $S_{\varphi_k}^\sigma$ and $S_{\varphi_1}^\sigma$ we can prove that there exist a sequence $(d_k)_{k \geq 1} \subset \mathbb{Z}$, a measurable set $E \subset F$, and a measurable function $\varphi_\infty: \Omega \rightarrow \mathbb{R}$ such that (5.1)–(5.3) hold true for a not-relabelled subsequence.

We next show that E has finite perimeter and $\varphi_\infty \in GBV(\Omega)$. For each $k \geq 1$ we define the auxiliary function

$$\widehat{\varphi}_k := \begin{cases} \varphi_k - 2\pi d_k & \text{in } F \\ k & \text{in } \Omega \setminus F. \end{cases}.$$

Since F has finite perimeter we have $(\widehat{\varphi}_k) \subset GBV(\Omega)$. Moreover

$$\sup_{k \in \mathbb{N}} \mathcal{H}^{n-1}(S_{\widehat{\varphi}_k}^\sigma) \leq H + \mathcal{H}^{n-1}(\partial^* F),$$

and

$$\begin{aligned} \lim_{k \rightarrow +\infty} \widehat{\varphi}_k(x) &= \varphi_\infty(x) & \text{for a.e. } x \in E, \\ \lim_{k \rightarrow +\infty} |\widehat{\varphi}_k(x)| &= +\infty & \text{for a.e. } x \in \Omega \setminus E. \end{aligned}$$

Now we proceed by adapting a slicing argument in [4] (see also [21]). For $\mathcal{H}^{n-1}$ -a.e. $\xi \in \mathbb{S}^{n-1} \subset \mathbb{R}^n$ and $y \in \Pi^\xi := \{y \in \mathbb{R}^n: y \cdot \xi = 0\}$ we introduce the following notation:

$$A_y^\xi := \{t \in \mathbb{R}: y + t\xi \in A\}, \quad a_y^\xi(t) := a(y + t\xi),$$

for any $A \subseteq \mathbb{R}^n$ and $a: A \rightarrow \mathbb{R}$. In particular $(\widehat{\varphi}_k)_y^\xi$ is a map of one variable in $GBV(\Omega_y^\xi)$ for a.e. $y \in \Pi^\xi$.

Recalling that $S_{(\widehat{\varphi}_k)_y^\xi}^\sigma := \{t \in S_{(\widehat{\varphi}_k)_y^\xi}: |[(\widehat{\varphi}_k)_y^\xi(t)]| \geq \sigma\}$ and $S_{\widehat{\varphi}_k}^\sigma := \{x \in S_{\widehat{\varphi}_k}: |[(\widehat{\varphi}_k)(x)]| \geq \sigma\}$, we have

$$S_{(\widehat{\varphi}_k)_y^\xi}^\sigma \subset (S_{\widehat{\varphi}_k}^\sigma)_y^\xi.$$

This together with Fatou's lemma implies

$$\begin{aligned} & \int_{\Pi^\xi} \liminf_{k \rightarrow +\infty} \left(|D(\widehat{\varphi}_k)_y^\xi|(\Omega_y^\xi \setminus S_{(\widehat{\varphi}_k)_y^\xi}^\sigma) + \mathcal{H}^0(\Omega_y^\xi \cap S_{(\widehat{\varphi}_k)_y^\xi}^\sigma) \right) d\mathcal{H}^{n-1}(y) \\ & \leq \int_{\Pi^\xi} \liminf_{k \rightarrow +\infty} \left(|D(\widehat{\varphi}_k)_y^\xi|(\Omega_y^\xi \setminus S_{(\widehat{\varphi}_k)_y^\xi}^\sigma) + \mathcal{H}^0\left(\Omega_y^\xi \cap (S_{\widehat{\varphi}_k}^\sigma)_y^\xi\right) \right) d\mathcal{H}^{n-1}(y) \\ & \leq \liminf_{k \rightarrow +\infty} \int_{\Pi^\xi} |D(\widehat{\varphi}_k)_y^\xi|(\Omega_y^\xi \setminus S_{(\widehat{\varphi}_k)_y^\xi}^\sigma) + \mathcal{H}^0\left(\Omega_y^\xi \cap (S_{\widehat{\varphi}_k}^\sigma)_y^\xi\right) d\mathcal{H}^{n-1}(y) \\ & \leq \liminf_{k \rightarrow +\infty} \left(|D\widehat{\varphi}_k|(\Omega \setminus S_{\widehat{\varphi}_k}^\sigma) + \int_{\Omega \cap S_{\widehat{\varphi}_k}^\sigma} |\nu_{\widehat{\varphi}_k} \cdot \xi| d\mathcal{H}^{n-1} \right) \\ & \leq c|Du|(\Omega) + \sigma \mathcal{H}^{n-1}(\partial^* F) + c \liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\widehat{\varphi}_k}^\sigma) < +\infty, \end{aligned}$$

where the last inequality follows from the fact that $(S_{\widehat{\varphi}_k} \setminus S_{\widehat{\varphi}_k}^\sigma) \subset S_u \cup \partial^* F$, and since $\sigma \leq \pi$, one has $|[(\widehat{\varphi}_k)(x)]| \leq c|[(u)(x)]|$ for a.e. $x \in S_{\widehat{\varphi}_k} \setminus S_{\widehat{\varphi}_k}^\sigma$ and some $c > 0$. Thus for $\mathcal{H}^{n-1}$ -a.e. $\xi \in \mathbb{S}^{n-1}$ and $\mathcal{H}^{n-1}$ -a.e. $y \in \Pi^\xi$ we can consider a not relabelled subsequence (depending on ξ and y) such that

$$\sup_k \left[|D(\widehat{\varphi}_k)_y^\xi|(\Omega_y^\xi \setminus S_{(\widehat{\varphi}_k)_y^\xi}^\sigma) + \mathcal{H}^0(\Omega_y^\xi \cap S_{(\widehat{\varphi}_k)_y^\xi}^\sigma) \right] < +\infty.$$

Hence for not relabelled subsequence (depending on ξ and y) there exists $m \in \mathbb{N}$ depending on ξ and y such that $\mathcal{H}^0 \left(\Omega_y^\xi \cap S_{(\widehat{\varphi}_k)_y^\xi}^\sigma \right) = m$ for all $k \geq 1$, and the jump points converge as well in $\overline{\Omega_y^\xi}$ as $k \rightarrow +\infty$ for $\mathcal{H}^{n-1}$ -a.e. $\xi \in \mathbb{S}^{n-1}$ and $\mathcal{H}^{n-1}$ -a.e. $y \in \Pi^\xi$. Thus, using the fact that $\widehat{\varphi}_k \rightarrow +\infty$ in $\Omega \setminus E$, and therefore on $\Omega_y^\xi \setminus E_y^\xi$, it is possible to show that

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^0 \left(A_y^\xi \cap S_{(\widehat{\varphi}_k)_y^\xi}^\sigma \right) \geq \mathcal{H}^0 \left(A_y^\xi \cap \partial^* E_y^\xi \right)$$

(see [4] for details). By integrating over Π^ξ and using once more Fatou's lemma we conclude

$$\begin{aligned} \liminf_{k \rightarrow +\infty} \int_{A \cap S_{\widehat{\varphi}_k}^\sigma} |\nu_{\widehat{\varphi}_k} \cdot \xi| d\mathcal{H}^{n-1} &\geq \int_{\Pi^\xi} \liminf_{k \rightarrow +\infty} \mathcal{H}^0 \left(A_y^\xi \cap S_{(\widehat{\varphi}_k)_y^\xi}^\sigma \right) d\mathcal{H}^{n-1}(y) \\ &\geq \int_{\Pi^\xi} \mathcal{H}^0 \left(\partial^* E_y^\xi \right) d\mathcal{H}^{n-1}(y), \end{aligned}$$

for $\mathcal{H}^{n-1}$ -a.e. $\xi \in \mathbb{S}^{n-1}$ and every open set $A \subseteq \Omega$. As a consequence we deduce that E has finite perimeter in Ω , since by integrating over $\mathbb{S}^{n-1}$,

$$\alpha_n \liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(A \cap S_{\widehat{\varphi}_k}^\sigma) \geq \alpha_n \mathcal{H}^{n-1}(A \cap \partial^* E),$$

where $\alpha_n := \int_{\mathbb{S}^{n-1}} |\nu \cdot \xi| d\mathcal{H}^{n-1}(\nu)$, which in turn implies

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(A \cap S_{\widehat{\varphi}_k}^\sigma) \geq \mathcal{H}^{n-1}(A \cap \partial^* E). \quad (5.7)$$

To prove that $\varphi_\infty \in GBV(\Omega)$ we argue as follows. For $N \in \mathbb{N}$ we set $\widehat{\varphi}_k^N := (\widehat{\varphi}_k \wedge N) \vee (-N)$, so that

$$|D\widehat{\varphi}_k^N|(\Omega) \leq |Du|(\Omega) + 2N\mathcal{H}^{n-1}(S_{\widehat{\varphi}_k^N}) \leq c.$$

Hence $(\widehat{\varphi}_k^N)$ converges weakly* in $BV(\Omega)$ to $\widehat{\varphi}_\infty^N$ as $k \rightarrow +\infty$, where

$$\widehat{\varphi}_\infty^N := \begin{cases} (\varphi_\infty \wedge N) \vee (-N) & \text{in } E, \\ \pm N & \text{in } \Omega \setminus E, \end{cases} \quad \text{so that } \widehat{\varphi}_\infty^N \in BV(\Omega).$$

Hence observing that in the whole of Ω we have $(\varphi_\infty \wedge N) \vee (-N) = \widehat{\varphi}_\infty^N \chi_E \in BV(\Omega)$ we conclude $\varphi_\infty \in GBV(\Omega)$. Eventually if in addition (5.4) holds then (5.5) can be deduced in the same way as (5.7) with $\varphi_k - 2\pi d_k$ in place of $\widehat{\varphi}_k$. This additionally implies

$$P(E, F) \leq H,$$

which shows (5.6). □

Lemma 5.2. *Let $0 \leq \sigma \leq \pi$. Let $(\varphi_k)_{k \geq 1} \subset GBV(\Omega)$ be a sequence of functions satisfying*

$$H := \sup_{k \in \mathbb{N}} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) < +\infty. \quad (5.8)$$

Let $N \geq 1$ be an integer, $E_1, \dots, E_N \subset \Omega$ be pairwise disjoint nonempty finite perimeter sets with the following property: for any $i = 1, \dots, N$,

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma \cap A) \geq \mathcal{H}^{n-1}(A \cap \partial^* E_i) \text{ for any } A \in \mathcal{A}(\Omega).$$

Then

$$H \geq \liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) \geq \mathcal{H}^{n-1}(\Omega \cap (\cup_{i=1}^N \partial^* E_i)).$$

Proof. We define the superadditive set function $\mu: \mathcal{A}(\Omega) \rightarrow [0, +\infty)$ as

$$\mu(A) := \liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(A \cap S_{\varphi_k}^\sigma),$$

and the positive measure

$$\lambda(B) := \mathcal{H}^{n-1}(B \cap (\cup_{i=1}^N \partial^* E_i)) \quad \forall \text{ Borel set } B \subseteq \Omega.$$

Moreover for $i = 1, \dots, N$ we set

$$h_i := \begin{cases} 1 & \text{in } \partial^* E_i \\ 0 & \text{otherwise in } \Omega \end{cases}.$$

Clearly we have

$$\mu(A) \geq \int_A h_i \, d\lambda \quad \forall i = 1, \dots, N.$$

Therefore, since

$$h := \sup_{i=1, \dots, N} h_i = \begin{cases} 1 & \text{in } \cup_{i=1}^N \partial^* E_i \\ 0 & \text{otherwise in } \Omega \end{cases},$$

by Lemma 2.1 it follows

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma \cap A) = \mu(A) \geq \int_A h \, d\lambda = \mathcal{H}^{n-1}(A \cap (\cup_{i=1}^N \partial^* E_i)),$$

which implies the thesis by taking $A = \Omega$. □

5.1 Proof of Theorem 4.2

We adapt the arguments of [13, Theorem 3.1].

Base step $M = 1$. We set

$$F_1 := \Omega.$$

By Lemma 5.1 we find, after extracting a not-relabelled subsequence, a finite perimeter set $E_1 \subseteq F_1$, a sequence $(d_k^{(1)})_{k \geq 1} \subset \mathbb{Z}$ and a function $\varphi_\infty^{(1)} \in GBV(\Omega)$, such that

$$\varphi_\infty^{(1)} \text{ is a lifting of } u \text{ in } E_1, \quad \varphi_\infty^{(1)} = 0 \quad \text{in } F_1 \setminus E_1,$$

and

$$|E_1| \geq \frac{n^n \omega_n |F_1|^n}{2^n (2H + \mathcal{H}^{n-1}(\partial\Omega))^n}.$$

Moreover

$$(\varphi_k(x) - 2\pi d_k^{(1)}) \rightarrow \varphi_\infty^{(1)}(x) \quad \text{for a.e. } x \in E_1,$$

$$|\varphi_k(x) - 2\pi d_k^{(1)}| \rightarrow +\infty \quad \text{for a.e. } x \in F_1 \setminus E_1.$$

Since $F_1 = \Omega$, estimate (5.5) yields

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma \cap A) \geq \mathcal{H}^{n-1}(A \cap \partial^* E_1) \text{ for any open set } A \subseteq F_1.$$

Inductive step $M \rightsquigarrow M+1$. Let $M \geq 2$. Suppose we have pairwise disjoint nonempty finite perimeter sets, $E_1, \dots, E_M \subset \Omega$, and define inductively

$$F_1 := \Omega, \quad F_i := \Omega \setminus \bigcup_{j=1}^{i-1} E_j \quad \text{for } i = 2, \dots, M,$$

so that $E_i \subset F_i$ for all $i = 1, \dots, M$. Assume that, still for all $i = 1, \dots, M$, the following holds:

- (i) There exists a function $\varphi_\infty^{(i)} \in GBV(\Omega)$ which is a lifting of u in E_i , and $\varphi_\infty^{(i)} = 0$ in $\Omega \setminus E_i$;
- (ii) The set E_i satisfies

$$|E_i| \geq \frac{n^n \omega_n |F_i|^n}{2^n (2H + \mathcal{H}^{n-1}(\partial^* F_i))^n} \geq \frac{n^n \omega_n |F_i|^n}{2^n (3H + \mathcal{H}^{n-1}(\partial \Omega))^n};$$

- (iii) There exists a sequence $(d_k^{(i)})_k \subset \mathbb{Z}$, such that

$$\begin{aligned} (\varphi_k(x) - 2\pi d_k^{(i)}) &\rightarrow \varphi_\infty^{(i)}(x) \quad \text{for a.e. } x \in E_i, \\ |\varphi_k(x) - 2\pi d_k^{(i)}| &\rightarrow +\infty \quad \text{for a.e. } x \in \Omega \setminus E_i, \\ \liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma \cap A) &\geq \mathcal{H}^{n-1}(A \cap \partial^* E_i) \quad \text{for any } A \in \mathcal{A}(\Omega). \end{aligned} \tag{5.9}$$

Set, for $M \geq 1$,

$$F_{M+1} := \Omega \setminus \left(\bigcup_{i=1}^M E_i \right),$$

If $F_{M+1} = \emptyset$ nothing remains to prove. Therefore, we may suppose $F_{M+1} \neq \emptyset$. From Lemma 5.2

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) \geq \mathcal{H}^{n-1}(\Omega \cap (\cup_{i=1}^M \partial^* E_i)), \tag{5.10}$$

and thus, since $\partial^*(\cup_{i=1}^M E_i) \subseteq \cup_{i=1}^M \partial^* E_i$,

$$\mathcal{H}^{n-1}\left(\Omega \cap \partial^*(\cup_{i=1}^M E_i)\right) = \mathcal{H}^{n-1}(\Omega \cap \partial^* F_{M+1}) \leq H. \tag{5.11}$$

Applying Lemma 5.1 to the set $F = F_{M+1}$, we obtain a finite perimeter set

$$E_{M+1} \subseteq F_{M+1},$$

a function $\varphi_\infty^{M+1} \in GBV(\Omega)$ and a sequence $(d_k^{M+1})_{k \geq 1} \subset \mathbb{Z}$, such that

$$\varphi_\infty^{M+1} = 0 \quad \text{in } \Omega \setminus E_{M+1}, \quad \varphi_\infty^{M+1} \text{ is a lifting of } u \text{ in } E_{M+1},$$

and

$$|E_{M+1}| \geq \frac{n^n \omega_n |F_{M+1}|^n}{2^n (2H + \mathcal{H}^{n-1}(\partial^* F_{M+1}))^n}, \tag{5.12}$$

together with

$$(\varphi_k(x) - 2\pi d_k^{M+1}) \rightarrow \varphi_\infty^{M+1}(x) \quad \text{for a.e. } x \in E_{M+1},$$

$$|\varphi_k(x) - 2\pi d_k^{M+1}| \rightarrow +\infty \quad \text{for a.e. } x \in F_{M+1} \setminus E_{M+1},$$

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma \cap B) \geq \mathcal{H}^{n-1}(B \cap \partial^* E_{M+1}) \text{ for any open set } B \subseteq F_{M+1}.$$

Gathering (5.11) and (5.12) we have

$$|E_{M+1}| \geq \frac{n^n \omega_n |F_{M+1}|^n}{2^n (2H + \mathcal{H}^{n-1}(\partial^* F_{M+1}))^n} \geq \frac{n^n \omega_n |F_{M+1}|^n}{2^n (3H + \mathcal{H}^{n-1}(\partial\Omega))^n} > 0.$$

Moreover by (5.9), also

$$|\varphi_k(x) - 2\pi d_k^{M+1}| \rightarrow +\infty \quad \text{for a.e. } x \in \Omega \setminus E_{M+1}.$$

Thus properties (i)-(iii) are preserved at level $M+1$.

Conclusion. Iterating the above construction and extracting a diagonal subsequence, we obtain a sequence $(E_i)_{i \geq 1}$ of mutually disjoint finite perimeter sets in Ω such that for every $M \geq 1$, $E_1, \dots, E_M$ satisfy properties (i)-(iii). From (5.10),

$$\liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) \geq \mathcal{H}^{n-1}(\Omega \cap (\cup_{i=1}^m \partial^* E_i)) \quad \forall m \geq 1. \quad (5.13)$$

To conclude, we show that

$$|\Omega \setminus (\cup_{i=1}^{+\infty} E_i)| = 0.$$

Since $\sum_{m=1}^{+\infty} |E_m| < +\infty$, the sequence $(|E_m|)$ tends to zero as $m \rightarrow +\infty$, and using

$$|E_m| \geq \frac{n^n \omega_n |F_m|^2}{2^n (3H + \mathcal{H}^{n-1}(\partial\Omega))^2}$$

we deduce that $|F_m| \rightarrow 0$. In particular,

$$|\Omega \setminus (\cup_{i=1}^{+\infty} E_i)| = \lim_{m \rightarrow +\infty} |\Omega \setminus (\cup_{i=1}^{m-1} E_i)| = \lim_{m \rightarrow +\infty} |F_m| = 0.$$

Define

$$\varphi_\infty(x) := \varphi_\infty^{(i)}(x) \quad \text{if } x \in E_i \text{ for some } i \in \mathbb{N}.$$

Then φ_∞ is a lifting of u in Ω . We now show that

$$\varphi_\infty \in GBV(\Omega). \quad (5.14)$$

First observe that by letting $m \rightarrow \infty$ in (5.13) we deduce that

$$H \geq \liminf_{k \rightarrow +\infty} \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) \geq \mathcal{H}^{n-1}(\Omega \cap (\cup_{i=1}^\infty \partial^* E_i)). \quad (5.15)$$

Now, for each $N \in \mathbb{N}$ we define the auxiliary sequence

$$\Phi_k^N := ((\varphi_k - 2\pi d_k^{(i)}) \wedge N) \vee (-N) \quad \text{in } E_i \text{ for all } i \geq 1.$$

Thus, noticing that $S_{\Phi_k^N} \subset S_u \cup S_{\varphi_k}^\sigma \cup (\cup_{i=1}^\infty \partial^* E_i)$ we have from (5.15)

$$|D\Phi_k^N|(\Omega) \leq \int_\Omega |\nabla u| dx + |D^c u|(\Omega) + \int_{S_u} |u^+ - u^-| d\mathcal{H}^{n-1} + 2N \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) \leq C,$$

with $C > 0$ depending on N . Therefore, up to a subsequence, (Φ_k^N) converges to $(\varphi_\infty \wedge N) \vee (-N)$ weakly* in $BV(\Omega)$ as $k \rightarrow +\infty$. By the arbitrariness of N we deduce (5.14). $\square$

5.2 Proof of Theorem 4.5

Let $\bar{\varphi}, \bar{\varphi}_k \in BV(\Omega) \cap L^\infty(\Omega)$ be liftings of u and of u_k in Ω , respectively with

$$|\bar{\varphi}|_{BV} \leq 2|u|_{BV} \leq C, \quad |\bar{\varphi}_k|_{BV} \leq 2|u_k|_{BV} \leq C$$

(see Theorem 2.4). The constant $C > 0$ can be found using the assumption (4.9) and so u_k are uniformly bounded in $BV(\Omega)$. In particular, for $\sigma \in [0, \pi]$ as in the statement, there is a positive constant $\bar{C}$ such that

$$\sup_{k \geq 1} \{ \mathcal{H}^{n-1}(S_{\bar{\varphi}_k}^\sigma) + \mathcal{H}^{n-1}(S_{\bar{\varphi}}^\sigma) \} \leq \bar{C}.$$

Furthermore, $\bar{\varphi}_k$ have equibounded BV -norm, and so there is a (not relabelled) subsequence such that

$$\bar{\varphi}_k \rightarrow \bar{\varphi}_\infty \text{ weakly}^* \text{ in } BV(\Omega); \quad (5.16)$$

since up to a subsequence the same convergence holds pointwise a.e., and also $u_k \rightarrow u$ a.e. on Ω , we deduce that $\bar{\varphi}_\infty$ must be a lifting of u .

Thanks to the fact that $\sigma \leq \pi$, the functions

$$v_k := \varphi_k - \bar{\varphi}_k, \quad k \geq 1$$

take values in $2\pi\mathbb{Z}$, and belong to $GSBV(\Omega; 2\pi\mathbb{Z})$, since the jump sets satisfy $S_{v_k} \subseteq S_{\bar{\varphi}_k}^\sigma \cup S_{\varphi_k}^\sigma$, and so

$$\mathcal{H}^{n-1}(S_{v_k}) \leq \mathcal{H}^{n-1}(S_{\bar{\varphi}_k}^\sigma) + \mathcal{H}^{n-1}(S_{\varphi_k}^\sigma) < \bar{C} < +\infty,$$

for all $k \geq 1$. Trivially v_k are liftings of the same function $f \equiv (1, 0) \in \mathbb{S}^1$, and so we can apply Theorem 4.2 to obtain that, for a non-relabelled subsequence, there exist a Caccioppoli partition $\{E_i\}_{i \in \mathbb{N}}$ of Ω , sequences of integers $(d_k^{(i)})_{k \geq 1}$, and a function $v_\infty \in GSBV(\Omega; 2\pi\mathbb{Z})$ such that, for all $i \in \mathbb{N}$,

$$\begin{aligned} (\varphi_k - \bar{\varphi}_k - 2\pi d_k^{(i)}) &\rightarrow v_\infty \text{ pointwise a.e. in } E_i, \\ |\varphi_k - \bar{\varphi}_k - 2\pi d_k^{(i)}| &\rightarrow +\infty \text{ pointwise a.e. in } \Omega \setminus E_i. \end{aligned}$$

From this and (5.16) we conclude

$$\begin{aligned} (\varphi_k - 2\pi d_k^{(i)}) &\rightarrow v_\infty + \bar{\varphi}_\infty \text{ pointwise a.e. in } E_i, \\ |\varphi_k - 2\pi d_k^{(i)}| &\rightarrow +\infty \text{ pointwise a.e. in } \Omega \setminus E_i. \end{aligned}$$

This is the thesis, just by setting $\varphi_\infty := \bar{\varphi}_\infty + v_\infty$, which is a lifting of u since v_∞ takes values in $2\pi\mathbb{Z}$. $\square$

6 Γ -convergence of functionals on $\mathbb{S}^1$ -valued maps

The proof of Theorem 3.1 is delicate. In particular the lower bound inequality requires a local argument which relies on the compactness result for liftings (Theorem 4.5). For convenience we introduce the localised Modica-Mortola-type (or Allen-Cahn type) functionals

$$\text{MM}_\varepsilon(v, A) := \int_A \left(\varepsilon |\nabla v|^2 + \frac{W(v)}{\varepsilon} \right) dx \quad \forall v \in W^{1,2}(\Omega),$$

for every open set $A \subseteq \Omega$, where W is defined in (c) of Section 2.2.

6.1 Proof of Theorem 3.1

Step 1: Lower bound. Let $\varepsilon_k \searrow 0$ as $k \rightarrow +\infty$. We have to show that

$$\liminf_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) \geq F_{\text{lift}}(u, v) \quad (6.1)$$

for every sequence $((u_k, v_k))_{k \geq 1} \subset L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$ converging to (u, v) in $L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$. We may assume

$$\sup_{k \in \mathbb{N}} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) \leq C < +\infty$$

so that $(u_k, v_k) \in \mathcal{D}_{\mathbb{S}^1}$, $v = 1$ a.e. in Ω and, up to a not relabelled subsequence,

$$\liminf_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) = \lim_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k).$$

From Theorem 2.2 it follows that

$$\liminf_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) \geq \int_{\Omega} f(|\nabla u|) dx + |D^c u|(\Omega) + \int_{S_u} g(|u^+ - u^-|) d\mathcal{H}^{n-1}.$$

In particular, from Remark 2.3-(ii)-5 and the fact that $|u^+ - u^-| \leq 2$ a.e. (since u is $\mathbb{S}^1$ -valued) we deduce $u \in BV(\Omega; \mathbb{S}^1)$.

For every $k \geq 1$, since Ω simply connected, we can choose a lifting $\varphi_k \in W^{1,2}(\Omega)$ of u_k in Ω . Using that $|\nabla u_k| = |\nabla \varphi_k|$ we have

$$+\infty > C \geq F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) = \int_{\Omega} \left(\psi(v_k) f(|\nabla \varphi_k|) + \varepsilon_k |\nabla v_k|^2 + \frac{W(v_k)}{\varepsilon_k} \right) dx.$$

Thus, from the coarea formula, for any $k \geq 1$,

$$C \geq \text{MM}_{\varepsilon_k}(v_k, \Omega) \geq \int_{\Omega} \sqrt{W(v_k)} |\nabla v_k| dx = \int_0^1 \sqrt{W(t)} \mathcal{H}^{n-1}(\partial^* F_k^t) dt \quad (6.2)$$

where $F_k^t := \{x \in \Omega : v_k(x) \leq t\}$. Now, fix $\eta', \eta'' \in (0, 1)$ with $\eta' < \eta''$. By (6.2) and the mean value theorem there exist $t(k) \in (\eta', \eta'')$ and $c(\eta', \eta'') > 0$ independent of k such that

$$C \geq c(\eta', \eta'')(\eta'' - \eta') \mathcal{H}^{n-1}(\partial^* F_k^{t(k)}). \quad (6.3)$$

Moreover, using also that ψ is increasing,

$$C \geq \int_{\Omega} \psi(v_k) f(|\nabla \varphi_k|) dx \geq C \psi(\eta') \int_{\Omega} \chi_{\Omega \setminus F_k^{t(k)}} |\nabla \varphi_k| dx. \quad (6.4)$$

Setting

$$\phi_k := \varphi_k \chi_{\Omega \setminus F_k^{t(k)}} \in SBV^2(\Omega),$$

we have

$$S_{\phi_k} \subseteq \partial^* F_k^{t(k)} \quad \text{and} \quad \nabla \phi_k = \nabla \varphi_k \chi_{\Omega \setminus F_k^{t(k)}}.$$

Therefore, from (6.4) and (6.3),

$$\int_{\Omega} |\nabla \phi_k| dx + \mathcal{H}^{n-1}(S_{\phi_k}) \leq C(\eta', \eta'') \quad (6.5)$$

for some $C(\eta', \eta'') > 0$ depending on η', η'' and independent of k . Define

$$\bar{u}_k := e^{i\phi_k} = u_k \chi_{\Omega \setminus F_k^{t(k)}} + (1, 0) \chi_{F_k^{t(k)}} \quad \forall k \in \mathbb{N},$$

which, by (6.3) and (6.4), are uniformly bounded in $BV(\Omega; \mathbb{S}^1)$. This and

$$|F_k^{t(k)}| \rightarrow 0, \quad (6.6)$$

which is a consequence of $t(k) < \eta'' < 1$ and $v_k \rightarrow 1$ a.e., imply that the sequence $(\bar{u}_k)$ weakly* converges to u in $BV(\Omega; \mathbb{S}^1)$. Hence, using (6.5), we can apply Theorem 4.5 to the sequence $(\phi_k)_{k \geq 1}$ and get, for a not-relabelled subsequence, a lifting $\varphi_\infty \in GBV(\Omega)$ of u in Ω , such that (ϕ_k) converges locally modulo 2π to φ_∞ . Namely, there exists a Caccioppoli partition $\{E_i\}_{i \in \mathbb{N}}$ of Ω , sequences $(d_k^{(i)})_{k \geq 1} \subset \mathbb{Z}$ for any integer $i \geq 1$ with the following properties:

$$\begin{aligned} \lim_{k \rightarrow +\infty} (\phi_k(x) - 2\pi d_k^{(i)}) &= \varphi_\infty(x) \quad \forall i \in \mathbb{N}, \text{ for a.e. } x \in E_i, \\ \lim_{k \rightarrow +\infty} |\phi_k(x) - 2\pi d_k^{(i)}| &= +\infty \quad \forall i \in \mathbb{N}, \text{ for a.e. } x \in \Omega \setminus E_i. \end{aligned}$$

Again, using (6.6), the same holds for φ_k , i.e.,

$$\begin{aligned} \lim_{k \rightarrow +\infty} (\varphi_k(x) - 2\pi d_k^{(i)}) &= \varphi_\infty(x) \quad \forall i \in \mathbb{N}, \text{ for a.e. } x \in E_i, \\ \lim_{k \rightarrow +\infty} |\varphi_k(x) - 2\pi d_k^{(i)}| &= +\infty \quad \forall i \in \mathbb{N}, \text{ for a.e. } x \in \Omega \setminus E_i. \end{aligned} \quad (6.7)$$

In particular, to prove the validity of (6.1) it is sufficient to show

$$\liminf_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) \geq \int_{\Omega} f(|\nabla \varphi_\infty|) dx + |D^c \varphi_\infty|(\Omega) + \int_{S_u} g(|\varphi_\infty^+ - \varphi_\infty^-|) d\mathcal{H}^{n-1}.$$

since, being φ_∞ a lifting of u , we have

$$\int_{\Omega} f(|\nabla \varphi_\infty|) dx = \int_{\Omega} f(|\nabla u|) dx, \quad |D^c \varphi_\infty|(\Omega) = |D^c u|(\Omega),$$

and

$$\int_{S_u} g(|\varphi_\infty^+ - \varphi_\infty^-|) d\mathcal{H}^{n-1} \geq m_g[u].$$

For fixed i and k and any integer $K \geq 1$ we consider the truncated function

$$\varphi_k^{i,K} := ((\varphi_k - 2\pi d_k^{(i)}) \wedge K) \vee (-K) \in W^{1,2}(\Omega),$$

and

$$\phi_k^{i,K} := \varphi_k^{i,K} \chi_{\Omega \setminus F_k^{t(k)}}.$$

According to the inclusion $S_{\phi_k^{i,K}} \subseteq \partial^* F_k^{t(k)}$ and the inequality $|\llbracket \phi_k^{i,K} \rrbracket| \leq 2K$, we have

$$\begin{aligned} |D\phi_k^{i,K}|(\Omega) &= \int_{\Omega} |\nabla \phi_k| \chi_{\{|\varphi_k - 2\pi d_k^{(i)}| < K\}} dx + \int_{S_{\phi_k^{i,K}}} |\llbracket \phi_k^{i,K} \rrbracket| d\mathcal{H}^{n-1} \\ &\leq \int_{\Omega} |\nabla \phi_k| dx + K \mathcal{H}^{n-1}(\partial^* F_k^{t(k)}) \leq C. \end{aligned}$$

Hence, up to a subsequence (depending on K and on i), $(\phi_k^{i,K})$ converges to some $\phi_\infty^{i,K}$ in $L^1(\Omega)$ as $k \rightarrow +\infty$. Moreover from (6.7) it holds

$$\phi_\infty^{i,K} := (\varphi_\infty \wedge K) \vee (-K) \quad \text{in } E_i.$$

Let $F_i^\pm$ be such that

$$\Omega \setminus E_i = F_i^+ \cup F_i^- \quad \text{and} \quad \phi_\infty^{i,K} = \pm K \text{ in } F_i^\pm.$$

From (6.6), it follows that $(\varphi_k^{i,K})$ converges to $\phi_\infty^{i,K}$ in $L^1(\Omega)$ as $k \rightarrow +\infty$. Hence the sequence $((\varphi_k^{i,K}, v_k))$ converges to $(\phi_\infty^{i,K}, 1)$ in $L^1(\Omega) \times L^1(\Omega)$ and

$$\int_A \left(\psi(v_k) f(|\nabla \varphi_k^{i,K}|) + \varepsilon_k |\nabla v_k|^2 + \frac{W(v_k)}{\varepsilon_k} \right) dx \leq C,$$

for any open set $A \subseteq \Omega$, for some $C = C(A) > 0$ independent of k . From Theorem 2.2 and Remark 2.3-(i) for every $i \in \mathbb{N}$

$$\begin{aligned} & \liminf_{k \rightarrow +\infty} \int_A \left(\psi(v_k) f(|\nabla \varphi_k|) + \varepsilon_k |\nabla v_k|^2 + \frac{W(v_k)}{\varepsilon_k} \right) dx \\ & \geq \liminf_{k \rightarrow +\infty} \int_A \left(\psi(v_k) f(|\nabla \varphi_k^{i,K}|) + \varepsilon_k |\nabla v_k|^2 + \frac{W(v_k)}{\varepsilon_k} \right) dx \\ & \geq \int_A f(|\nabla \phi_\infty^{i,K}|) dx + |D^c \phi_\infty^{i,K}|(A) + \int_{S_{\phi_\infty^{i,K}} \cap A} g(|(\phi_\infty^{i,K})^+ - (\phi_\infty^{i,K})^-|) d\mathcal{H}^{n-1}. \end{aligned} \tag{6.8}$$

In particular Remark 2.3 together with $|(\phi_\infty^{i,K})^+ - (\phi_\infty^{i,K})^-| \leq 2K$ imply

$$\phi_\infty^{i,K} \in BV(\Omega).$$

Consider the bounded positive measure given by

$$\lambda(B) := \int_B f(|\nabla \varphi_\infty|) dx + |D^c \varphi_\infty|(B) + \mathcal{H}^{n-1}(S_{\varphi_\infty} \cap B) \quad \forall \text{ Borel set } B \subseteq \Omega.$$

For each $i, K \in \mathbb{N}$ we define

$$h^{i,K} := \begin{cases} g(|(\phi_\infty^{i,K})^+ - (\phi_\infty^{i,K})^-|) & \text{in } S_{\phi_\infty^{i,K}} \\ 1 & \text{otherwise in } \Omega. \end{cases}$$

Thus from (6.8) we have

$$\liminf_{k \rightarrow +\infty} \int_A \left(\psi(v_k) f(|\nabla \varphi_k|) + \varepsilon_k |\nabla v_k|^2 + \frac{W(v_k)}{\varepsilon_k} \right) dx \geq \int_A h^{i,K} d\lambda \quad \forall i, K \in \mathbb{N},$$

for all $A \in \mathcal{A}(\Omega)$. Next, as $|(\phi_\infty^{i,K})^+ - (\phi_\infty^{i,K})^-| \rightarrow |(\phi_\infty^i)^+ - (\phi_\infty^i)^-|$ as $K \rightarrow \infty$ pointwise $\mathcal{H}^1$ -a.e. on S_{φ_∞} , and then observing that

$$\sup_{i, K \in \mathbb{N}} h^{i,K} = \begin{cases} g(|\varphi_\infty^+ - \varphi_\infty^-|) & \text{in } S_{\varphi_\infty}, \\ 1 & \text{in } \Omega \setminus S_{\varphi_\infty}, \end{cases}$$

we conclude by Lemma 2.1.

Step 2: Upper bound. Let $\varepsilon_k \searrow 0$ and $u \in BV(\Omega; \mathbb{S}^1)$. We have to find a sequence $((u_k, v_k)) \subset \mathcal{D}_{\mathbb{S}^1}$ converging to $(u, 1)$ in $L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$ such that

$$\limsup_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) \leq F_{\text{lift}}(u, 1).$$

By Corollary 4.3 we can select a jump minimizing lifting $\varphi \in GBV(\Omega)$ of u in Ω , and

$$\int_{S_\varphi} g(|\varphi^+ - \varphi^-|) d\mathcal{H}^{n-1} = m_g[u]. \quad (6.9)$$

By Theorem 2.2 there exist $(\varphi_k, v_k) \in W^{1,2}(\Omega) \times W^{1,2}(\Omega)$ such that $(\varphi_k, v_k) \rightarrow (\varphi, 1)$ in $L^1(\Omega) \times L^1(\Omega)$ and

$$\limsup_{k \rightarrow +\infty} F_{\varepsilon_k}(\varphi_k, v_k) \leq F(\varphi, 1). \quad (6.10)$$

Next we let $u_k := e^{i\varphi_k} \in W^{1,2}(\Omega; \mathbb{S}^1)$, so that $(u_k, v_k) \rightarrow (u, 1)$ in $L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$. Moreover, from (6.9), (6.10), $|\nabla u_k| = |\nabla \varphi_k|$ and $|\nabla u| = |\nabla \varphi|$ we get

$$F_{\varepsilon_k}(\varphi_k, v_k) = F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k), \quad F(\varphi, 1) = F_{\text{lift}}(u, 1)$$

and so

$$\limsup_{k \rightarrow +\infty} F_{\varepsilon_k}^{\mathbb{S}^1}(u_k, v_k) \leq F_{\text{lift}}(u, 1).$$

□

6.2 An extension of Theorem 3.1

We consider the domain $\widehat{\mathcal{D}}_{\mathbb{S}^1} \supsetneq \mathcal{D}_{\mathbb{S}^1}$, defined as

$$\widehat{\mathcal{D}}_{\mathbb{S}^1} := \left\{ (u, v) \in W_{\text{loc}}^{1,2}(\Omega; \mathbb{S}^1) \times W_{\text{loc}}^{1,2}(\Omega) : 0 \leq v \leq 1 \right\} \subset L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega). \quad (6.11)$$

and the extended energy

$$\widehat{F}_{\varepsilon}^{\mathbb{S}^1}(u, v) := \begin{cases} F_{\varepsilon}(u, v) & \text{if } (u, v) \in \widehat{\mathcal{D}}_{\mathbb{S}^1}, \\ +\infty & \text{otherwise in } L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega). \end{cases} \quad (6.12)$$

Then, the following result holds:

Theorem 6.1 (Γ -convergence of $\widehat{F}_{\varepsilon}^{\mathbb{S}^1}$). *Provided Ω is simply-connected, we have*

$$\Gamma - L^1 \lim_{\varepsilon \rightarrow 0^+} \widehat{F}_{\varepsilon}^{\mathbb{S}^1} = F_{\text{lift}}.$$

The proof of the upper bound in Theorem 6.1 is straightforward since $\widehat{\mathcal{D}}_{\mathbb{S}^1} \supsetneq \mathcal{D}_{\mathbb{S}^1}$. To obtain the lower bound, it suffices to follow the proof of Theorem 3.1 for a sequence $A_k \subset\subset \Omega$ with $A_k \nearrow \Omega$, and apply the locality property in Remark 2.3(ii). The thesis follows by a diagonal argument.

7 Connections with the relaxed area in $\mathbb{S}^1$

The relaxation of the area functional in $\mathbb{S}^1$ for the graph of an L^1 -function $u: \Omega \rightarrow \mathbb{S}^1$ is defined as

$$\mathcal{A}^{\mathbb{S}^1}(u, \Omega) := \inf \left\{ \liminf_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla u_k|^2} dx : u_k \in C^1(\Omega; \mathbb{S}^1), u_k \rightarrow u \text{ in } L^1 \right\}.$$

In [28, page 613] it is observed that $\mathcal{A}^{\mathbb{S}^1}(u, \cdot)$ is not subadditive as a set function, therefore it can not admit an integral representation and a characterization using Cartesian currents is provided in [28, page 612, Theorem 1]. We recall that, as a consequence of [28, page 612, Theorem 1], $\mathcal{A}^{\mathbb{S}^1}(u, \Omega)$ is finite if and only if $u \in BV(\Omega; \mathbb{S}^1)$.

Here we propose a characterization using lifting theory when Ω is simply connected and leave the general case for future developments. The expression we obtain clearly reveals the nonlocal nature of $\mathcal{A}^{\mathbb{S}^1}(u, \cdot)$.

Theorem 7.1 (Characterization of $\mathcal{A}^{\mathbb{S}^1}$). *Let $\Omega \subset \mathbb{R}^n$ be a bounded connected simply connected open set with Lipschitz boundary. Then*

$$\mathcal{A}^{\mathbb{S}^1}(u, \Omega) = \int_{\Omega} \sqrt{1 + |\nabla u|^2} dx + |D^c u|(\Omega) + m[u], \quad u \in BV(\Omega; \mathbb{S}^1), \quad (7.1)$$

where

$$m[u] := \min \left\{ \int_{S_{\varphi}} |\varphi^+ - \varphi^-| d\mathcal{H}^{n-1} : \varphi \in BV(\Omega), e^{i\varphi} = u \text{ a.e. in } \Omega \right\}. \quad (7.2)$$

We preliminarily observe that if $u \in W^{1,1}(\Omega; \mathbb{S}^1) \cap C^1(\Omega; \mathbb{S}^1)$, then there exists a lifting $\varphi \in W^{1,1}(\Omega) \cap C^1(\Omega)$ of u such that $-\pi < \oint_{\Omega} \varphi dx \leq \pi$. Indeed, the existence of a lifting $\hat{\varphi} \in W^{1,1}(\Omega) \cap C^1(\Omega)$ follows by well-known results [18]. To achieve the latter property it suffices to add to $\hat{\varphi}$ a suitable integer multiple of 2π .

Proof of Theorem 7.1. Step 1: Lower bound. Let $u_k \in C^1(\Omega; \mathbb{S}^1)$ be such that $u_k \rightarrow u$ in L^1 . We need to prove that

$$\liminf_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla u_k|^2} dx \geq \int_{\Omega} \sqrt{1 + |\nabla u|^2} dx + |D^c u|(\Omega) + m[u]. \quad (7.3)$$

Assume without loss of generality that

$$\sup_k \int_{\Omega} \sqrt{1 + |\nabla u_k|^2} dx < +\infty.$$

Since $\int_{\Omega} \sqrt{1 + |\nabla u_k|^2} dx \geq \int_{\Omega} |\nabla u_k| dx$ and $|u_k| = 1$ we have $u_k \in W^{1,1}(\Omega; \mathbb{S}^1)$. For each $k \in \mathbb{N}$ select a lifting $\varphi_k \in W^{1,1}(\Omega) \cap C^1(\Omega)$ of u_k in Ω with $-\pi \leq \oint_{\Omega} \varphi_k dx \leq \pi$. We have

$$\int_{\Omega} |\nabla \varphi_k| dx = \int_{\Omega} |\nabla u_k| dx \leq C.$$

Furthermore, by Poincaré inequality

$$\int_{\Omega} |\varphi_k| dx \leq \int_{\Omega} \left| \varphi_k - \oint_{\Omega} \varphi_k dy \right| dx + |\Omega| \left| \oint_{\Omega} \varphi_k dx \right| \leq C \int_{\Omega} |\nabla \varphi_k| dx + |\Omega| \pi \leq C.$$

Thus, up to a not relabelled subsequence, $\varphi_k \xrightarrow{*} \varphi$ in $BV(\Omega)$, where φ is a lifting of u in Ω . By [1] we conclude

$$\begin{aligned} \liminf_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla u_k|^2} dx &= \liminf_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla \varphi_k|^2} dx \\ &\geq \int_{\Omega} \sqrt{1 + |\nabla \varphi|^2} dx + |D^c \varphi|(\Omega) + \int_{S_{\varphi}} |\varphi^+ - \varphi^-| d\mathcal{H}^{n-1} \\ &\geq \int_{\Omega} \sqrt{1 + |\nabla u|^2} dx + |D^c u|(\Omega) + m[u]. \end{aligned}$$

Step 2: Upper bound. Let $u \in BV(\Omega; \mathbb{S}^1)$. We need to find $u_k \in C^1(\Omega; \mathbb{S}^1)$ such that $u_k \rightarrow u$ in L^1 and

$$\limsup_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla u_k|^2} dx \leq \int_{\Omega} \sqrt{1 + |\nabla u|^2} dx + |D^c u|(\Omega) + m[u]. \quad (7.4)$$

By [19, page 25] there exists a lifting $\tilde{\varphi} \in BV(\Omega)$ of u in Ω such that

$$|D\tilde{\varphi}|(\Omega) = \inf \{ |D\varphi|(\Omega) : \varphi \in BV(\Omega), e^{i\varphi} = u \text{ a.e. in } \Omega \}.$$

Moreover $|D\tilde{\varphi}|(\Omega) \leq 2|Du|(\Omega)$ and $0 \leq \int_{\Omega} \tilde{\varphi} dx \leq 2\pi$. From the identities

$$\int_{\Omega} |\nabla \varphi| dx = \int_{\Omega} |\nabla u| dx \quad \text{and} \quad |D^c \varphi|(\Omega) = |D^c u|(\Omega)$$

it follows that

$$\int_{S_{\tilde{\varphi}}} |\tilde{\varphi}^+ - \tilde{\varphi}^-| d\mathcal{H}^{n-1} \leq \int_{S_{\varphi}} |\varphi^+ - \varphi^-| d\mathcal{H}^{n-1}$$

for any lifting $\varphi \in BV(\Omega)$ of u in Ω . Hence

$$\int_{S_{\tilde{\varphi}}} |\tilde{\varphi}^+ - \tilde{\varphi}^-| d\mathcal{H}^{n-1} = m[u].$$

Now by [26] there exists a sequence $(\varphi_k) \subset C^1(\Omega)$ converging to φ in L^1 and such that

$$\limsup_{k \rightarrow +\infty} \int_{\Omega} \sqrt{1 + |\nabla \varphi_k|^2} dx \leq \int_{\Omega} \sqrt{1 + |\nabla \tilde{\varphi}|^2} dx + |D^c \tilde{\varphi}|(\Omega) + \int_{S_{\tilde{\varphi}}} |\tilde{\varphi}^+ - \tilde{\varphi}^-| d\mathcal{H}^{n-1}.$$

Therefore the maps $u_k := e^{i\varphi_k} \in C^1(\Omega; \mathbb{S}^1)$ converge to u in L^1 and satisfy (7.4). $\square$

We conclude the paper by providing a phase-field approximation of Ambrosio-Tortorelli type of $\mathcal{A}^{\mathbb{S}^1}(u, \Omega)$. Let $\mathcal{D}_{\mathbb{S}^1}$ be as in (3.1). For $\varepsilon, h > 0$ consider the family of functionals $\mathcal{F}_{\varepsilon, h}^{\mathbb{S}^1} : L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega) \rightarrow [0, +\infty]$ given by

$$\mathcal{F}_{\varepsilon, h}^{\mathbb{S}^1}(u, v) := \int_{\Omega} v^2 \sqrt{1 + |\nabla u|^2} dx + \int_{\Omega} \left(\varepsilon |\nabla v|^2 + \frac{(1-v)^2}{4h\varepsilon} \right) dx, \quad (7.5)$$

when $(u, v) \in \widehat{\mathcal{D}_{\mathbb{S}^1}}$ and $+\infty$ otherwise. With a little abuse of notation, we denote by

$$\mathcal{A}^{\mathbb{S}^1} : L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega) \rightarrow [0, +\infty]$$

the following functional: $\mathcal{A}^{\mathbb{S}^1}(u, v) = \mathcal{A}^{\mathbb{S}^1}(u)$ if $u \in BV(\Omega)$ and $v = 1$ almost everywhere in Ω , and $\mathcal{A}^{\mathbb{S}^1}(u, v) = +\infty$ if either $u \notin BV(\Omega; \mathbb{S}^1)$ or v is not almost everywhere equal to 1 in Ω .

Theorem 7.2. *Let $\Omega \subset \mathbb{R}^n$ be a bounded connected and simply connected open set with Lipschitz boundary. Let $h_\varepsilon := h(\varepsilon)$ be positive, with*

$$\lim_{\varepsilon \rightarrow 0} \frac{\varepsilon}{h_\varepsilon} = 0 \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0^+} h_\varepsilon = 0.$$

Then $\mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}$ Γ -converges to $\mathcal{A}^{\mathbb{S}^1}$ in the L^1 -topology.

Proof. Step 1: Lower bound. Let $(u_\varepsilon, v_\varepsilon) \rightarrow (u, v)$ in $L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$. We need to prove that

$$\liminf_{\varepsilon \rightarrow 0^+} \mathcal{F}_\varepsilon^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) \geq \mathcal{A}^{\mathbb{S}^1}(u, v). \quad (7.6)$$

Assume, without loss of generality, that

$$\sup_\varepsilon \mathcal{F}_\varepsilon^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) < +\infty.$$

Let $h > 0$ be fixed. Then for ε small enough we have $h \geq h_\varepsilon$, so that

$$\liminf_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon).$$

Moreover from Theorem 3.1 and (2.3) we have that

$$\liminf_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) \geq \int_\Omega \sqrt{1 + |\nabla u|^2} \, dx + |D^c u|(\Omega) + m_{g_h}[u],$$

where

$$m_{g_h}[u] = \inf \left\{ \int_{S_\varphi} g_h(|\varphi^+ - \varphi^-|) \, d\mathcal{H}^{n-1} : \varphi \in GBV(\Omega), \, e^{i\varphi} = u \text{ a.e. in } \Omega \right\},$$

with

$$g_h(t) := \frac{t}{1 + ht}, \quad t \geq 0.$$

Hence

$$\liminf_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) \geq \int_\Omega \sqrt{1 + |\nabla u|^2} \, dx + |D^c u|(\Omega) + m_{g_h}[u].$$

By the arbitrariness of $h > 0$, we obtain

$$\liminf_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) \geq \int_\Omega \sqrt{1 + |\nabla u|^2} \, dx + |D^c u|(\Omega) + \sup_{h > 0} m_{g_h}[u].$$

Now, we claim that

$$\sup_{h > 0} m_{g_h}[u] = \lim_{h \rightarrow 0^+} m_{g_h}[u] = m[u]. \quad (7.7)$$

Indeed, since $g_h(t) \leq t$ for all $t \geq 0$, we immediately have $m_{g_h}[u] \leq m[u]$ for all $h > 0$, hence $\limsup_{h \rightarrow 0^+} m_{g_h}[u] \leq m[u]$.

To prove the converse inequality, for every $h > 0$ let $\varphi_h \in GBV(\Omega)$ be a jump-minimizing lifting for $m_{g_h}[u]$, which exists by Corollary 4.3, so that

$$m_{g_h}[u] = \int_{S_{\varphi_h}} \frac{|\varphi_h^+ - \varphi_h^-|}{1 + h|\varphi_h^+ - \varphi_h^-|} \, d\mathcal{H}^{n-1}.$$

Fix $h_0 > 0$. For any $h \in (0, h_0)$ we have $g_h(t) \geq g_{h_0}(t)$ for all $t \geq 0$, which yields

$$\sup_{h \in (0, h_0)} \mathcal{H}^{n-1}(S_{\varphi_h}^\pi) \leq (1 + h_0\pi) \sup_{h \in (0, h_0)} m_{g_h}[u] \leq (1 + h_0\pi)m[u] < +\infty,$$

where $S_{\varphi_h}^\pi$ is defined in (4.3). Thus, we can apply Theorem 4.2 to the family $(\varphi_h)_{h \in (0, h_0)}$. For a sequence $h_k \rightarrow 0^+$, (φ_{h_k}) converges locally modulo 2π to a lifting $\varphi_0 \in GBV(\Omega)$ of u in Ω . By the lower semicontinuity result in Corollary 4.3-(ii) and the Fatou's lemma, we infer

$$\liminf_{k \rightarrow +\infty} m_{h_k}[u] = \liminf_{k \rightarrow +\infty} \int_{S_{\varphi_{h_k}}} \frac{|\varphi_{h_k}^+ - \varphi_{h_k}^-|}{1 + h_k|\varphi_{h_k}^+ - \varphi_{h_k}^-|} d\mathcal{H}^{n-1} \geq \int_{S_{\varphi_0}} |\varphi_0^+ - \varphi_0^-| d\mathcal{H}^{n-1} \geq m[u].$$

This shows claim (7.7) and concludes the proof of the lower bound.

Step 2: Upper bound. Let $u \in BV(\Omega; \mathbb{S}^1)$. We need to construct a sequence $((u_\varepsilon, v_\varepsilon))_{\varepsilon > 0} \subset \mathcal{D}_{\mathbb{S}^1}$ converging to $(u, 1)$ in $L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$ such that

$$\limsup_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) \leq \mathcal{A}^{\mathbb{S}^1}(u, 1).$$

By definition of the relaxed area $\mathcal{A}^{\mathbb{S}^1}(u, \Omega)$, there exists a sequence $(u_\varepsilon) \subset C^1(\Omega; \mathbb{S}^1)$ such that $u_\varepsilon \rightarrow u$ in $L^1(\Omega; \mathbb{S}^1)$ and

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} \sqrt{1 + |\nabla u_\varepsilon|^2} dx = \mathcal{A}^{\mathbb{S}^1}(u, \Omega). \quad (7.8)$$

Since $u_\varepsilon \in C^1(\Omega; \mathbb{S}^1)$ we have $u_\varepsilon \in W_{\text{loc}}^{1,2}(\Omega; \mathbb{S}^1)$. Setting $v_\varepsilon \equiv 1$ on Ω , we have $(u_\varepsilon, v_\varepsilon) \in \widehat{\mathcal{D}_{\mathbb{S}^1}}$ for all $\varepsilon > 0$, and $(u_\varepsilon, v_\varepsilon) \rightarrow (u, 1)$ in $L^1(\Omega; \mathbb{S}^1) \times L^1(\Omega)$. Moreover, since $v_\varepsilon \equiv 1$, the phase-field terms vanish identically, yielding

$$\mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) = \int_{\Omega} \sqrt{1 + |\nabla u_\varepsilon|^2} dx.$$

Invoking (7.8), we obtain

$$\limsup_{\varepsilon \rightarrow 0^+} \mathcal{F}_{\varepsilon, h_\varepsilon}^{\mathbb{S}^1}(u_\varepsilon, v_\varepsilon) = \lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} \sqrt{1 + |\nabla u_\varepsilon|^2} dx = \mathcal{A}^{\mathbb{S}^1}(u, \Omega),$$

which completes the proof. $\square$

Data availability statement

No new data was created or analysed in this work.

Conflict of interest statement

The authors have no conflict of interest to declare that are relevant to the content of this article.

Acknowledgements

R. Marziani has received funding from the European Union's Horizon research and innovation program under the Marie Skłodowska-Curie agreement No 101150549. The research that led to the present paper was partially supported by the INdAM-GNAMPA project 2025 CUP E5324001950001. The first author benefits from the support of the GNAMPA of INdAM. The first and third authors also acknowledge the partial financial support of the PRIN project 2022PJ9EFL "Geometric Measure Theory: Structure of Singular Measures, Regularity Theory and Applications in the Calculus of Variations", PNRR Italia Domani, funded by the European Union via the program NextGenerationEU (Missione 4 Componente 2) CUP:E53D23005860006. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

References

- [1] E. Acerbi, G. Dal Maso, *New lower semicontinuity results for polyconvex integrals*, Calc. Var. Partial Differential Equations **2** (1994), 329–371.
- [2] R. Alicandro, A. Braides, J. Shah, *Free-discontinuity problems via functionals involving the L^1 -norm of the gradient and their approximation*, Interfaces Free Bound. **1** (1999), 17–37.
- [3] R. Alicandro, M. Focardi, *Variational approximation of free-discontinuity energies with linear growth*, Commun. Contemp. Math. **4** (2002), 685–723.
- [4] S. Almi, E. Tasso, *A new proof of compactness in $G(S)BD$* , Adv. Calc. Var. **16** (2023), 637–650.
- [5] L. Ambrosio, N. Fusco, D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 2000.
- [6] L. Ambrosio, V. M. Tortorelli, *Approximation of functionals depending on jumps by elliptic functionals via Γ -convergence*, Comm. Pure Appl. Math. **43** (1990), 999–1036.
- [7] L. Ambrosio, V. M. Tortorelli, *On the approximation of free-discontinuity problems*, Boll. Un. Mat. Ital. (7) **6-B** (1992), 105–123.
- [8] G. Bellettini, A. Elshorbagy, M. Paolini, R. Scala, *On the relaxed area of the graph of discontinuous maps from the plane to the plane taking three values with no symmetry assumptions*, Ann. Mat. Pura Appl. **199** (2020), 445–477.
- [9] G. Bellettini, A. Elshorbagy, R. Scala, *The L^1 -relaxed area of the graph of the vortex map: optimal lower bound*, Nonlinear Anal. **256** (2025), Paper No. 113803.
- [10] G. Bellettini, A. Elshorbagy, R. Scala, *Relaxation of the area of the vortex map: a non-parametric Plateau problem for a catenoid containing a segment*, J. Funct. Anal. **289** (2025), 110247.

- [11] G. Bellettini, A. Elshorbagy, R. Scala, *The L^1 -relaxed area of the graph of the vortex map: optimal upper bound*, Adv. Calc. Var. (2025) <https://doi.org/10.1515/acv-2024-0101>.
- [12] G. Bellettini, R. Marziani, R. Scala, *A non-parametric Plateau problem with partial free boundary*, J. Echolle Polytechnique – Mathématiques **11** (2024), 1035-1098.
- [13] G. Bellettini, R. Marziani, R. Scala, *On jump minimizing liftings for $\mathbb{S}^1$ -valued maps and connections with Ambrosio-Tortorelli-type Γ -limits*, J. Math. Pures Appl. **217** (2027), 103992.
- [14] G. Bellettini, M. Paolini, *On the area of the graph of a singular map from the plane to the plane taking three values*, Adv. Calc. Var. **3** (2010), 371-386.
- [15] G. Bellettini, R. Scala, G. Scianna, *L^1 -relaxed area of graphs of $\mathbb{S}^1$ -valued Sobolev maps and its countably subadditive envelope*, Rev. Mat. Iberoam. **40** (2024), 2135-2178.
- [16] F. Bethuel, H. Brezis, F. Hélein, Ginzburg-Landau Vortices, Progress in Nonlinear Differential Equations and Their Applications, Birkhäuser Boston, Springer (1994).
- [17] A. Braides, Approximation of Free-Discontinuity Problems, Lecture Notes in Math. **1694**, Springer Verlag, Berlin, 1998.
- [18] H. Brezis, P. Mironescu, Sobolev Maps to the Circle, Progress in Nonlinear Differential Equations and their Applications **96**, Birkhäuser/Springer, New York, 2021.
- [19] H. Brezis, P. Mironescu, *The Plateau problem from the perspective of optimal transport*, Comptes Rendus. Mathématique, **357**(7) (2019), 597–612.
- [20] S. Carano, *Relaxed area of 0-homogeneous maps in the strict BV-convergence*, Ann. Mat. Pura Appl. **203** (2024), 2057–2074.
- [21] A. Chambolle, V. Crismale, *Compactness and lower semicontinuity in GSBD*, J. Eur. Math. Soc. (JEMS) **23** (2021), 701–719.
- [22] M. Cicalese, G. Orlando, M. Ruf, *Emergence of concentration effects in the variational analysis of the N-clock model*, Comm. Pure Appl. Math. **75** (2022), 2279-2342.
- [23] V. Crismale, L. De Luca, R. Scala, *Approximation of topological singularities through free discontinuity functionals: the critical and super-critical regimes*, Calc. Var. Partial Differential Equations **65** (2026), Art. 5.
- [24] J. Dávila, R. Ignat, *Lifting of BV functions with values in $\mathbb{S}^1$* , C. R. Math. Acad. Sci. Paris **337** (2003), 159–164.
- [25] L. De Luca, R. Scala, N. Van Goethem, *A new approach to topological singularities via a weak notion of Jacobian for functions of bounded variation*, Indiana Univ. Math. J. **73** (2024), 723–779.
- [26] G. Dal Maso, *Integral representation of Γ -limits of variational integrals*, Manuscripta Mat. **30** (1980), 387-416.
- [27] L. Ferrari, C. Dirks, B. Wirth, *Phase field approximations of branched transportation problems*, Calc. Var. Partial Differential Equations **59** (2020), Art. 37, 34 pp.

- [28] M. Giaquinta, G. Modica, J. Souček, Cartesian Currents in the Calculus of Variations II. Variational Integrals, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Vol. 38, Springer-Verlag, Berlin-Heidelberg, 1998.
- [29] A. Marchese, A. Massaccesi, S. Stuvard, R. Tione, *A multi-material transport problem with arbitrary marginals*, *Calc. Var. Partial Differential Equations* **60** (2021), 1-49.
- [30] A. Marchese, A. Massaccesi, R. Tione, *A multimaterial transport problem and its convex relaxation via rectifiable G -currents*, *SIAM J. Math. Anal.* **51** (2019), 1965-1998.
- [31] R. Scala, *Optimal estimates for the triple junction function and other surprising aspects of the area functional*, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **20**, (2020), 491-564.
- [32] R. Scala, G. Scianna, *On the L^1 -relaxed area of graphs of BV piecewise constant maps taking three values*, *Adv. Calc. Var.* **18** (2015), 339–365.