

On the evolution of subcritical regions for the Perona-Malik equation

Marina Ghisi

Università degli Studi di Pisa
Dipartimento di Matematica “Leonida Tonelli”
PISA (Italy)
e-mail: ghisi@dm.unipi.it

Massimo Gobbino

Università degli Studi di Pisa
Dipartimento di Matematica Applicata “Ulisse Dini”
PISA (Italy)
e-mail: m.gobbino@dma.unipi.it

Abstract

The Perona-Malik equation is a celebrated example of forward-backward parabolic equation. The forward behavior takes place in the so-called subcritical region, in which the gradient of the solution is smaller than a fixed threshold. In this paper we show that this subcritical region evolves in a different way in the following three cases: dimension one, radial solutions in dimension greater than one, general solutions in dimension greater than one.

In the first case subcritical regions increase, but there is no estimate on the expansion rate. In the second case they expand with a positive rate and *always spread over the whole domain* after a finite time, depending only on the (outer) radius of the domain. As a by-product, we obtain a non-existence result for global-in-time classical radial solutions with large enough gradient. In the third case we show an example where subcritical regions do *not* expand.

Our proofs exploit comparison principles for suitable degenerate and non-smooth free boundary problems.

Mathematics Subject Classification 2000 (MSC2000): 35K55, 35K65, 35R35.

Key words: Perona-Malik equation, forward-backward parabolic equation, degenerate parabolic equation, moving domains, subsolutions and supersolutions.

1 Introduction

In this paper we consider the Perona-Malik equation

$$u_t(x, t) - \operatorname{div} \left(\frac{\nabla u(x, t)}{1 + |\nabla u(x, t)|^2} \right) = 0 \quad \forall (x, t) \in \Omega \times [0, T), \quad (1.1)$$

where $\Omega \subseteq \mathbb{R}^n$ is an open set and $T > 0$. This equation is the formal gradient flow of the functional

$$PM(u) := \frac{1}{2} \int_{\Omega} \log(1 + |\nabla u(x)|^2) dx.$$

The convex-concave behavior of the integrand makes (1.1) a forward-backward partial differential equation of parabolic type. The forward (or *subcritical*) region is the set of points (x, t) where $|\nabla u(x, t)| < 1$, the backward (or *supercritical*) region is the set of points where $|\nabla u(x, t)| > 1$.

This equation was introduced by P. PERONA and J. MALIK [17] in the context of image denoising. They considered equation (1.1) in a rectangle $\Omega \subseteq \mathbb{R}^2$, with an initial condition $u(x, 0) = u_0(x)$ representing the grey level of a (noisy) picture, and Neumann boundary conditions. For increasing values of t the functions $x \rightarrow u(x, t)$ are thought as successively restored versions of $u_0(x)$.

The heuristic idea is that small disturbances, corresponding to small values of the gradient, are smoothed out because of the diffusion which takes place in the forward regions. On the other hand, sharp edges correspond to large values of the gradient and therefore they are expected to be enhanced by the backward character of the equation in supercritical regions. This phenomenology has been actually observed in numerical experiments, which also reveal an unexpected stability (see [7, 8, 14]). This discrepancy between the practical efficacy of (1.1) and its analytical ill-posedness has been called “Perona-Malik paradox” after S. KICHENASSAMY’s paper [16].

In the last fifteen years the paradox has been investigated in numerous papers. Several authors proved well posedness results for approximations of (1.1) obtained via space discretization [4, 9] or convolution [5], time delay [1], fractional derivatives [13], fourth order regularization [2], simplified nonlinearities [3]. The behavior of such approximations as the suitable parameter goes to 0 is a much more challenging problem. As far as we know, results in this direction have been obtained only for the semidiscrete scheme in dimension one. In this case the authors [9] (see also [4]) proved that approximated solutions converge to a limit, and under reasonable assumptions this limit is a classical solution of (1.1) inside its subcritical region.

Despite of these partial progresses, a solution of the paradox is still far away. We remind that a solution of the paradox is a notion of weak solution for (1.1) which exists for large classes of initial data (for example in $BV(\Omega)$ or $SBV(\Omega)$), is reasonably stable, and to which reasonable approximations converge.

In this direction, K. ZHANG [19] (see also [6, 18]) showed that the class of Lipschitz solutions is far from being a solution to the paradox. Indeed he proved that for any nonconstant smooth initial condition, even if subcritical, the Neumann boundary value problem admits infinitely many (pathological) Lipschitz solutions.

Classical solutions (namely solutions which are at least of class C^1) have also been investigated in the last decade. B. KAWOHL and N. KUTEV [15] observed that global-in-time classical solutions exist if the initial condition is subcritical, while in [16] it is remarked that local-in-time classical solutions cannot exist unless the initial condition is very regular in its supercritical region. Moreover the authors proved in [10] that in dimension one there exists a dense set of initial data for which the Cauchy problem with Neumann boundary conditions admits a local-in-time classical solution of class $C^{2,1}$ (namely with two continuous derivatives with respect to space variables, and one continuous derivative with respect to time). On the other hand, such solutions *cannot* be global if the initial condition has a nonempty supercritical region (see [15] and [12]).

Quite surprisingly, things are not so drastic in dimension greater than one. Indeed the authors proved in [11] that *global-in-time* radial solutions of class $C^{2,1}$ *do* exist for some classes of initial data with nonempty supercritical region.

For classical solutions one can define the family of open sets

$$I^-(t) := \{x \in \Omega : |\nabla u(x, t)| < 1\} \quad \forall t \in [0, T).$$

This is the family of subcritical regions. Its behavior as t varies is the object of this paper. We point out that this definition is purely local, in the sense that it does not depend on the boundary conditions. We show three situations in which subcritical regions evolve in a different way. As far as we know, these different behaviors had not been explicitly reported in numerical experiments. The reason is maybe that in an unstable framework it is always difficult to distinguish what depends on the model itself, and what depends on the implementation. In any case, we leave to numerical analysts and applied mathematicians any comment about the significance of these results for the model and its practical applications.

The one dimensional case In dimension one subcritical regions don't shrink, namely

$$I^-(s) \subseteq I^-(t) \quad \text{whenever } 0 \leq s \leq t < T. \quad (1.2)$$

This fact had already been proved in [15] under some structural assumptions on the initial condition, afterwards removed in [12].

In this paper we present an alternative proof (see Theorem 2.1) based on a comparison principle, which only requires C^1 regularity, and works substantially for all equations which are the gradient flow of a nonconvex functional. This proof gives us the opportunity to show, in a simpler setting, the method which is fundamental in the next case, when the result was not known before.

This result is optimal in the sense that it may happen that the subcritical region is the same for every $t \in [0, T)$. An example is provided by the local-in-time solutions constructed in [10].

The radial case Let us consider a radial solution of (1.1) defined in a ball or an annulus. Then (1.2) holds true also in this case. If moreover $I^-(0) \neq \emptyset$, then the inclusion is strict whenever $s < t$, and there exists $T_0 > 0$ such that $I^-(t) = \Omega$ for every $t > T_0$. The value of T_0 depends on the outer radius of Ω , but is independent on u . In other words, supercritical and critical regions disappear after a finite time depending only on Ω .

If the solution survives and remains regular up to T_0 , then it becomes subcritical and there are no more obstructions to global existence. This is actually what happens in the classes of global-in-time radial solutions constructed in [11].

In Theorem 2.3 we show that this is not always the case. If the initial condition has a supercritical region where the gradient is large enough, and this supercritical region is surrounded by subcritical regions, then there is no C^1 classical solution with $T > T_0$, independently on the boundary conditions. The reason is that the maximum of $|\nabla u(x, t)|$ in this supercritical region is a function of time which cannot decrease too fast. On the other hand, the supercritical region must disappear after T_0 , and thus this maximum has not enough time to decrease from its large initial value up to 1.

The nonradial case Roughly speaking, in the radial case each interface between the subcritical and the supercritical region is a circle which evolves with velocity proportional to the inverse of its radius. This reminded us of the mean curvature motion, and in a first moment led us to suspect that this interface could evolve in a similar way also for nonradial solutions. In particular it seemed reasonable that this interface could evolve in such a way that supercritical regions tend to shrink, at least where they are convex.

In Theorem 2.4 we show that this is not the case. We prove indeed that there exists an initial condition u_0 in two variables with the following properties. The supercritical region of u_0 is convex in a neighborhood of the origin, and any local-in-time solution starting from u_0 , independently on the boundary conditions, has a supercritical region which expands in a neighborhood of the origin. In particular any such solution does *not* satisfy (1.2).

Evolution of supercritical regions If subcritical regions expand, then supercritical regions shrink. This is true, but not obvious. Indeed it could happen that both the subcritical and the supercritical region expand at the expense of the critical region where $|\nabla u(x, t)| = 1$. Nevertheless one can prove that this is not the case. It is enough to apply the techniques of this paper to the equation obtained by reversing the time. In this way the role of subcritical and supercritical regions is just exchanged.

Connection with free boundary problems The evolution of subcritical regions is itself a free boundary problem. Unfortunately it involves a forward-backward equation. After some variable changes we reduce ourselves to more standard situations. Roughly speaking, in the new variable v we end up with equations such as

$$v_t = \sqrt{v} v_{xx}, \quad v_t = \sqrt{v} \{v_{xx} + A + \text{lower order terms}\},$$

where $A > 0$. These equations are satisfied where $v > 0$, and we are interested in the evolution of the region where v is positive.

The good news are that these equations are forward parabolic. The bad news are that they are *degenerate* and they involve a nonlinear term which is *not Lipschitz* continuous. This complicates things when using comparison principles.

For the first equation we show (see Theorem 2.5) that the region where v is positive does not shrink with time. This is enough to threat the Perona-Malik equation in dimension one. For the second equation we show (see Theorem 2.6) that the region where v is positive expands with a positive rate depending on A . This implies our conclusions for the radial Perona-Malik equation.

This paper is organized as follows. In section 2 we state our results for the Perona-Malik equation and the related free boundary problems. In section 3 we prove these results.

2 Statements

Throughout this paper we assume that $\varphi \in C^\infty(\mathbb{R})$ is an even function, hence in particular $\varphi'(0) = 0$. We also assume that

$$\varphi''(\sigma) > 0 \quad \forall \sigma \in [0, 1), \quad (2.1)$$

$$\varphi''(1) = 0, \quad (2.2)$$

$$\varphi''(\sigma) < 0 \quad \forall \sigma > 1. \quad (2.3)$$

These assumptions imply that $\varphi'(1) > 0$ and $\varphi'''(1) \leq 0$. In some statement we also need the stronger assumption

$$\varphi'''(1) < 0. \quad (2.4)$$

These assumptions are consistent with the concrete case $\varphi(\sigma) = 2^{-1} \log(1 + \sigma^2)$ of the Perona-Malik equation. We refer to Figure 1 for the typical behavior of $\varphi'(\sigma)$.

We consider the following equation

$$u_t = \operatorname{div} \left(\varphi'(|\nabla u|) \frac{\nabla u}{|\nabla u|} \right), \quad (2.5)$$

which is the natural generalization of (1.1). We believe and we hope that this generality simplifies the presentation, and shows more clearly which properties of the nonlinearity are essential in each step. For the sake of generality one could also weaken the regularity assumptions on φ (we never consider more than 3 derivatives), replace the threshold $\sigma = 1$ in (2.2) with any σ_1 , and weaken (2.1) and (2.3) by asking only that φ'' is positive in a left-hand neighborhood of σ_1 and negative in a right-hand neighborhood of σ_1 .

2.1 Main results

Let us state our results on the evolution of subcritical regions. The first result concerns the one dimensional case, where (2.5) reduces to

$$u_t = (\varphi'(u_x))_x \quad (2.6)$$

This form of the equation is suitable for C^1 solutions, because it involves only first order derivatives. When the solution is of class $C^{2,1}$, equation (2.6) can of course be rewritten as $u_t = \varphi''(u_x)u_{xx}$.

Theorem 2.1 *Let $\varphi \in C^\infty(\mathbb{R})$ be a function satisfying (2.1) through (2.3). Let $x_1 \leq x_3 < x_4 \leq x_2$ and $T > 0$ be real numbers.*

Let $u \in C^1((x_1, x_2) \times [0, T])$ be a function satisfying (2.6) in $(x_1, x_2) \times [0, T]$, and

$$|u_x(x, 0)| < 1 \quad \forall x \in (x_3, x_4). \quad (2.7)$$

Then $|u_x(x, t)| < 1$ for every $(x, t) \in (x_3, x_4) \times [0, T]$.

We point out that in Theorem 2.1 we don't need assumption (2.4), which was used in all previous results of the same type.

Our second result concerns the radial case. Let $r := |x|$ be the radial variable, and let $u(r, t)$ be a radial solution. In this case (1.1) becomes

$$u_t = (\varphi'(u_r))_r + (n-1)\frac{\varphi'(u_r)}{r}, \quad (2.8)$$

where n is the space dimension. From now on we assume for simplicity that $n = 2$. The general case is completely analogous.

Theorem 2.2 *Let $\varphi \in C^\infty(\mathbb{R})$ be a function satisfying (2.1) through (2.4). Let $0 < r_1 \leq r_3 < r_4 \leq r_2$ and $T > 0$ be real numbers. Let $u \in C^1((r_1, r_2) \times [0, T])$ be a function satisfying (2.8) (with $n = 2$) in $(r_1, r_2) \times [0, T]$, and*

$$|u_r(r, 0)| < 1 \quad \forall r \in (r_3, r_4).$$

Let $k_0 := r_2^{-1}\sqrt{2\varphi'(1)|\varphi'''(1)|}$, and let us set

$$\mathcal{D} := \{(r, t) \in (r_1, r_2) \times [0, T] : r_3 - k_0 t < r < r_4 + k_0 t\}. \quad (2.9)$$

Then $|u_r(r, t)| < 1$ for every $(r, t) \in \mathcal{D}$.

In other words, this result says that in the radial case the subcritical region expands with a rate which is bounded from below by a positive constant k_0 . Figure 2 shows the shape of the set \mathcal{D} . The slope of the slanted lines depends on k_0 . It is clear that when $t > (r_2 - r_1)/k_0$ every nonempty initial subcritical region (r_3, r_4) has invaded the whole interval (r_1, r_2) .

The third result concerns the nonexistence of global-in-time C^1 radial solutions if the gradient of the initial condition is too large. We point out that this result is independent on the boundary conditions.

Theorem 2.3 *Let $\varphi \in C^\infty(\mathbb{R})$ be a function satisfying (2.1) through (2.4). Let $0 \leq r_1 < r_3 < r_4 < r_5 < r_2$ and $T > 0$ be real numbers, and let $u \in C^1((r_1, r_2) \times [0, T])$ be a solution of (2.8) (with $n = 2$) such that*

$$|u_r(r_3, 0)| < 1, \quad |u_r(r_5, 0)| < 1, \quad u_r(r_4, 0) > 1 + \frac{r_2(r_2 - r_1)}{r_1^2} \sqrt{\frac{\varphi'(1)}{2|\varphi'''(1)|}}. \quad (2.10)$$

Then we have that

$$T \leq \frac{r_2(r_2 - r_1)}{\sqrt{2\varphi'(1)|\varphi'''(1)|}}.$$

Our last result is a counterexample to the expansion of subcritical regions for non-radial solutions. Note that the condition we impose on the initial condition u_0 depends only on the Taylor expansion of u_0 of order 3 in a neighborhood of the origin.

Theorem 2.4 *Let $\varphi \in C^\infty(\mathbb{R})$ be a function such that $\varphi'(1) > 0$ and $\varphi''(1) = 0$. Let $\Omega \subseteq \mathbb{R}^2$ be any open set such that $(0, 0) \in \Omega$. Let $u_0 : \Omega \rightarrow \mathbb{R}$ be any function of class C^3 such that*

$$u_0(x, y) = \frac{\sqrt{2}}{2}x + \frac{\sqrt{2}}{2}y + k_1x^2 + k_2y^2 + h_1x^3 + h_2y^3 + o((x^2 + y^2)^{3/2}) \quad (2.11)$$

as $(x, y) \rightarrow (0, 0)$. This clearly implies that $|\nabla u_0(0, 0)| = 1$. Moreover one can choose the parameters k_1, k_2, h_1, h_2 in such a way that the following properties hold true.

(1) *There exist $\delta > 0, a > 0$, and a convex function $g : (-\delta, \delta) \rightarrow (-a, a)$ such that*

$$\begin{aligned} & \{(x, y) \in (-\delta, \delta) \times (-a, a) : |\nabla u_0(x, y)| > 1\} = \\ & = \{(x, y) \in (-\delta, \delta) \times (-a, a) : y > g(x)\}. \end{aligned}$$

(2) *Let $T > 0$, and let $u \in C^2(\Omega \times [0, T])$ be a function satisfying (2.5), and the initial condition $u(x, y, 0) = u_0(x, y)$ for every $(x, y) \in \Omega$. Then there exists $t_0 \in (0, T)$ such that*

$$|\nabla u(0, 0, t)| > 1 \quad \forall t \in (0, t_0).$$

In other words, at time $t = 0$ the origin lies in the interface which separates the subcritical and the (locally convex) supercritical region, while for $t \in (0, t_0)$ the origin has been absorbed by the supercritical region.

2.2 Heuristics

In this section we present simple “proofs” of Theorem 2.1 and Theorem 2.2.

Let us start with Theorem 2.1. Let us assume that u is smooth enough, and that one component of the interface between the subcritical and the supercritical region is represented by a smooth curve $(\alpha(t), t)$. Just to fix ideas, let us assume that the subcritical region lies on the left of the interface, namely where $r < \alpha(t)$. Taking the time derivative of the relation $u_x(\alpha(t), t) = 1$ we obtain that

$$\begin{aligned} 0 &= u_{xx}(\alpha(t), t)\alpha'(t) + u_{xt}(\alpha(t), t) \\ &= u_{xx}\alpha'(t) + \varphi''(u_x)u_{xxx} + \varphi'''(u_x)u_{xx}^2, \end{aligned}$$

where all the partial derivatives of u in the last line are computed in the point $(\alpha(t), t)$. Recalling that $\varphi''(u_x(\alpha(t), t)) = 0$, we have therefore that

$$\alpha'(t) = -\varphi'''(1)u_{xx}(\alpha(t), t).$$

Now we have that $\varphi'''(1) \leq 0$, and $u_{xx}(\alpha(t), t) \geq 0$ because the subcritical region lies on the left of $r = \alpha(t)$. We have thus proved that $\alpha'(t) \geq 0$, hence the subcritical region tends to expand.

This “proof” is *not* rigorous for several reasons: we assumed that u is of class C^3 , we assumed that the interface is a smooth curve, we divided by u_{xx} which could be 0. Nevertheless we believe that this simple argument is quite explicative.

Let us consider now Theorem 2.2. As before, we assume that the interface is given by a smooth curve $(\alpha(t), t)$, the subcritical region being on the left. Taking the time derivative of the relation $u_r(\alpha(t), t) = 1$ we obtain that

$$\begin{aligned} 0 &= u_{rr}(\alpha(t), t)\alpha'(t) + u_{rt}(\alpha(t), t) \\ &= u_{rr}\alpha'(t) + \varphi''(u_r)u_{rrr} + \varphi'''(u_r)u_{rr}^2 + \frac{\varphi''(u_r)}{\alpha(t)}u_{rr} - \frac{\varphi'(u_r)}{\alpha^2(t)}, \end{aligned}$$

where all the partial derivatives of u in the last line are computed in the point $(\alpha(t), t)$. Recalling that $\varphi''(u_r(\alpha(t), t)) = 0$, we have therefore that

$$\alpha'(t) = \frac{\varphi'(1)}{\alpha^2(t)} \cdot \frac{1}{u_{rr}} - \varphi'''(1)u_{rr} = \frac{\varphi'(1)}{\alpha^2(t)} \cdot \frac{1}{u_{rr}} + |\varphi'''(1)|u_{rr}.$$

Applying the inequality between the arithmetic and the geometric mean, we thus obtain that

$$\alpha'(t) \geq \frac{2\sqrt{\varphi'(1)|\varphi'''(1)|}}{\alpha(t)} \geq \frac{2\sqrt{\varphi'(1)|\varphi'''(1)|}}{r_2}.$$

This “proves” that the subcritical region expands with a rate which is bounded from below by a positive constant. The value of this constant is quite similar to the constant k_0 of Theorem 2.2.

2.3 Free boundary problems

Our proofs of Theorem 2.1 and Theorem 2.2 rely on the following two results for free boundary problems involving degenerate and nonlipschitz parabolic equations. We state them independently because they could be interesting in themselves.

Theorem 2.5 *Let $x_1 \leq x_3 < x_4 \leq x_2$, and let $c_0 > 0$ and $T > 0$ be real numbers. Let $g : (0, c_0) \rightarrow (0, +\infty)$ be a continuous function. Let $v : (x_1, x_2) \times [0, T) \rightarrow \mathbb{R}$ be a function such that*

(v1) *v is continuous in $(x_1, x_2) \times [0, T)$;*

(v2) *$v(x, t) \geq 0$ for every $(x, t) \in (x_1, x_2) \times [0, T)$;*

(v3) *$v(x, 0) > 0$ for every $x \in (x_3, x_4)$;*

(v4) *the partial derivative $v_x(x, t)$ exists for every $(x, t) \in (x_1, x_2) \times (0, T)$;*

(v5) *setting*

$$\mathcal{P} := \{(x, t) \in (x_1, x_2) \times (0, T) : 0 < v(x, t) < c_0\},$$

we have that $v \in C^{2,1}(\mathcal{P})$, and

$$v_t(x, t) = g(v(x, t))v_{xx}(x, t) \quad \forall (x, t) \in \mathcal{P}.$$

Then $v(x, t) > 0$ for every $(x, t) \in (x_3, x_4) \times [0, T)$.

Theorem 2.6 *Let $r_1 \leq r_3 < r_4 \leq r_2$, and let c_0, c_1, T, G, A be positive real numbers. Let $g : (0, c_0) \rightarrow (0, +\infty)$ be a continuous function such that*

$$\lim_{\sigma \rightarrow 0^+} \frac{g(\sigma)}{\sqrt{\sigma}} = G. \quad (2.12)$$

Let $f : (r_1, r_2) \times (0, T) \times [-c_1, c_1]^2 \rightarrow \mathbb{R}$ be a function such that $f(r, t, 0, 0) = 0$ uniformly in (r, t) , namely

$$\lim_{\sigma \rightarrow 0^+} \sup \{ |f(r, t, p, q)| : (r, t, p, q) \in (r_1, r_2) \times (0, T) \times [-\sigma, \sigma]^2 \} = 0. \quad (2.13)$$

Let $v : (r_1, r_2) \times [0, T) \rightarrow \mathbb{R}$ be a function such that

(v1) *v is continuous in $(r_1, r_2) \times [0, T)$;*

(v2) *$v(r, t) \geq 0$ for every $(r, t) \in (r_1, r_2) \times [0, T)$;*

(v3) *$v(r, 0) > 0$ for every $r \in (r_3, r_4)$;*

(v4) the partial derivative $v_r(r, t)$ exists for every $(r, t) \in (r_1, r_2) \times (0, T)$;

(v5) setting

$$\mathcal{P} := \{(r, t) \in (r_1, r_2) \times (0, T) : 0 < v(r, t) < c_0\},$$

we have that $v \in C^{2,1}(\mathcal{P})$, and

$$v_t \geq g(v) \{v_{rr} + f(r, t, v, v_r) + A\} \quad \forall (r, t) \in \mathcal{P}. \quad (2.14)$$

Finally, let \mathcal{D} be the set defined as in (2.9) with $k_0 := G\sqrt{A}$.

Then $v(r, t) > 0$ for every $(r, t) \in \mathcal{D}$.

3 Proofs

3.1 Proof of Theorem 2.1

We limit ourselves to prove that $u_x(x, t) < 1$ for every $(x, t) \in (x_3, x_4) \times [0, T)$. The proof of the symmetric inequality $u_x(x, t) > -1$ is completely analogous.

Let us introduce some notation. Let us consider any function $h \in C^1(\mathbb{R})$ which is nondecreasing and such that $h(\sigma) = \varphi'(\sigma)$ for every $\sigma \in [0, 1]$, $h(\sigma) = \varphi'(1)$ for every $\sigma \geq 1$, and $h(\sigma)$ is constant for $\sigma \leq -1/2$. Figure 1 shows the typical graph of such a function h . Note that condition (2.2) is essential for the C^1 regularity of h .




Figure 1: Typical graph of functions φ' , h , and g

The function h , as well as the function φ' , is invertible as a function from $(0, 1)$ to $(0, \varphi'(1))$. We can therefore define $g : (0, \varphi'(1)) \rightarrow \mathbb{R}$ by setting

$$g(\sigma) := \varphi''(h^{-1}(\varphi'(1) - \sigma)) \quad \forall \sigma \in (0, \varphi'(1)).$$

It is not difficult to see that the function g is well defined, positive, and continuous (but not Lipschitz continuous). In the case of the Perona-Malik equation with some computations one finds that $g(\sigma) = (\sigma - \sigma^2)^{1/2} + 2(\sigma - \sigma^2)$. Its graph is shown in Figure 1.

Let us finally set

$$v(x, t) := \varphi'(1) - h(u_x(x, t)) \quad \forall (x, t) \in (x_1, x_2) \times [0, T].$$

We claim that g and v satisfy the assumptions of Theorem 2.5. If we prove this claim, then we can conclude that $v(x, t) > 0$ in $(x_3, x_4) \times [0, T]$. This is equivalent to say that $h(u_x(x, t)) < \varphi'(1)$, which in turn is equivalent to say that $u_x(x, t) < 1$ in the same region.

So we only need to show that v fulfils assumptions (v1) through (v5) of Theorem 2.5.

Properties (v1) through (v3) The continuity of v easily follows from the continuity of h and u_x . Moreover $v(x, t) \geq 0$ in $(x_1, x_2) \times [0, T]$ because $h(\sigma) \leq \varphi'(1)$ for every $\sigma \in \mathbb{R}$. Due to (2.7) and the fact that $\varphi'(\sigma) < \varphi'(1)$ when $\sigma < 1$, we have that v satisfies (v3).

Property (v4) It is well known that u is of class C^∞ where $|u_x(x, t)| \neq 1$ (because of the standard interior regularity theory for parabolic equations). Therefore the existence of the partial derivative $v_x(x, t)$ is trivial when $|u_x(x, t)| \neq 1$. The existence of $v_x(x, t)$ is trivial also when $u_x(x, t) = -1$ because $h(\sigma)$ is constant for $\sigma \leq 1/2$.

Let us consider now a point (x_0, t_0) with $u_x(x_0, t_0) = 1$. We claim that in this point $v_x(x_0, t_0)$ exists and is equal to 0. Let us assume that this is not the case. Then there exists a sequence $\delta_k \rightarrow 0$ such that

$$\left| \frac{v(x_0 + \delta_k, t_0) - v(x_0, t_0)}{\delta_k} \right| \geq \nu > 0 \quad \forall k \in \mathbb{N}. \quad (3.1)$$

Up to subsequences, we can always assume that either $u_x(x_0 + \delta_k, t_0) > 1$ for every $k \in \mathbb{N}$, or $u_x(x_0 + \delta_k, t_0) \leq 1$ for every $k \in \mathbb{N}$. In the first case the fraction in (3.1) is always 0, which is incompatible with the condition stated therein. In the second case the fraction in (3.1) can be rewritten as

$$-\frac{h(u_x(x_0 + \delta_k, t_0)) - h(u_x(x_0, t_0))}{\delta_k} = -\frac{\varphi'(u_x(x_0 + \delta_k, t_0)) - \varphi'(u_x(x_0, t_0))}{\delta_k}.$$

When $\delta_k \rightarrow 0$, this quotient tends to $(\varphi'(u_x))_x(x_0, t_0)$, and we know that this derivative exists because u is a solution of (2.6) of class C^1 . In order to find a contradiction, it is enough to show that this derivative is equal to 0, and this is true because it is the derivative in $x = x_0$ of the function $x \rightarrow \varphi'(u_x(x, t_0))$, which attains its maximum for $x = x_0$.

Property (v5) Let us set

$$\mathcal{P} := \{(x, t) \in (x_1, x_2) \times (0, T) : 0 < v(x, t) < \varphi'(1)\}. \quad (3.2)$$

From the properties of h it follows that $0 < u_x(x, t) < 1$ in \mathcal{P} , hence u is of class C^∞ in \mathcal{P} . Since h and φ' coincide in $(0, 1)$, we have therefore that

$$\begin{aligned} v_t &= -h'(u_x)u_{xt} = -h'(u_x)(\varphi'(u_x))_{xx} = -\varphi''(u_x)(h(u_x))_{xx} = \varphi''(u_x)(-h(u_x))_{xx} = \\ &= \varphi''(u_x)(\varphi'(1) - h(u_x))_{xx} = \varphi''(u_x)v_{xx} \end{aligned}$$

in \mathcal{P} . Moreover in \mathcal{P} we can express u_x in function of v as $u_x = h^{-1}(\varphi'(1) - v)$. In conclusion

$$v_t = \varphi''(u_x)v_{xx} = \varphi''(h^{-1}(\varphi'(1) - v))v_{xx} = g(v)v_{xx},$$

which proves that v satisfies (v5).

3.2 Proof of Theorem 2.2

The argument is similar to the proof of Theorem 2.1. We define h , g , and v as in that proof, and we claim that the assumptions of Theorem 2.6 are satisfied.

Properties of g As in the proof of Theorem 2.1 it is quite easy to show that $g : (0, \varphi'(1)) \rightarrow (0, +\infty)$ is a continuous function. It remains to compute the limit in (2.12). Since we deal with positive functions, we can square the numerator and the denominator. Applying the variable change $\tau := h^{-1}(\varphi'(1) - \sigma)$ and De L'Hôpital's rule, we have therefore that

$$\begin{aligned} \lim_{\sigma \rightarrow 0^+} \frac{[g(\sigma)]^2}{\sigma} &= \lim_{\sigma \rightarrow 0^+} \frac{[\varphi''(h^{-1}(\varphi'(1) - \sigma))]^2}{\sigma} = \lim_{\tau \rightarrow 1^-} \frac{[\varphi''(\tau)]^2}{\varphi'(1) - \varphi'(\tau)} = \\ &= \lim_{\tau \rightarrow 1^-} \frac{2\varphi''(\tau)\varphi'''(\tau)}{-\varphi''(\tau)} = -2\varphi'''(1) = 2|\varphi'''(1)|, \end{aligned}$$

which proves (2.12) with $G := \sqrt{2|\varphi'''(1)|}$.

Properties of v The proof of (v1) through (v4) is analogous to the proof of the corresponding properties in Theorem 2.1. In order to prove (v5), let us consider the set \mathcal{P} defined in analogy with (3.2). As in the previous case we have that $0 < u_r(r, t) < 1$ in this set, hence v is regular and satisfies

$$v_t = -h'(u_r)u_{rt} = -\varphi''(u_r) \left[(\varphi'(u_r))_r + \frac{\varphi'(u_r)}{r} \right]_r. \quad (3.3)$$

Once again $\varphi''(u_r) = \varphi''(h^{-1}(\varphi'(1) - v)) = g(v)$. Moreover

$$\left[(\varphi'(u_r))_r + \frac{\varphi'(u_r)}{r} \right]_r = (\varphi'(u_r))_{rr} + \frac{(\varphi'(u_r))_r}{r} - \frac{\varphi'(u_r)}{r^2} = -v_{rr} - \frac{v_r}{r} - \frac{\varphi'(1)}{r^2} + \frac{v}{r^2}.$$

Plugging these identities into (3.3) we obtain that

$$v_t = g(v) \left\{ v_{rr} + \frac{v_r}{r} - \frac{v}{r^2} + \frac{\varphi'(1)}{r^2} \right\} \geq g(v) \{v_{rr} + f(r, t, v, v_r) + A\},$$

where

$$f(r, t, p, q) := \frac{q}{r} - \frac{p}{r^2}, \quad A := \frac{\varphi'(1)}{r_2^2}.$$

We have thus proved that v satisfies the differential inequality (2.14) in \mathcal{P} with a constant $A > 0$, and a function $f(r, t, p, q)$ satisfying (2.13).

Conclusion From Theorem 2.6 we deduce that v is positive in the region \mathcal{D} defined according to (2.9) with $k_0 = G\sqrt{A} = r_2^{-1}\sqrt{2\varphi'(1)|\varphi'''(1)|}$, which in turn implies that $u_r < 1$ in the same region. The proof of the symmetric inequality $u_r > -1$ is completely analogous.

3.3 Proof of Theorem 2.3

Let us assume that a solution exists with

$$T > T_0 := \frac{r_2(r_2 - r_1)}{\sqrt{2\varphi'(1)|\varphi'''(1)|}}.$$

Let us set

$$M(t) := \max \{u_r(r, t) : r \in [r_3, r_5]\} \quad \forall t \in [0, T].$$

Due to the first two inequalities in (2.10) we know that the subcritical region is nonempty at time $t = 0$. Applying Theorem 2.2 we have therefore that the subcritical region expands, and coincides with the whole interval (r_1, r_2) as soon as

$$t > \frac{r_2 - r_1}{k_0} = T_0.$$

In particular this means that

$$M(T_0) \leq 1. \tag{3.4}$$

On the other hand we claim that

$$M(t) \geq M(0) - \frac{\varphi'(1)}{r_1^2}t \quad \forall t \in [0, T_0]. \tag{3.5}$$

If we prove this claim, then setting $t = T_0$ and exploiting the last inequality in (2.10), we find that

$$\begin{aligned} M(T_0) &\geq M(0) - \frac{\varphi'(1)}{r_1^2}T_0 \\ &> 1 + \frac{r_2(r_2 - r_1)}{r_1^2} \sqrt{\frac{\varphi'(1)}{2|\varphi'''(1)|}} - \frac{\varphi'(1)}{r_1^2} \cdot \frac{r_2(r_2 - r_1)}{\sqrt{2\varphi'(1)|\varphi'''(1)|}} \\ &= 1, \end{aligned}$$

which contradicts (3.4).

Proof of (3.5) The argument is similar to the usual comparison principles. Setting for simplicity $v(r, t) := u_r(r, t)$, we have that v is a solution of

$$v_t = \varphi''(v)v_{rr} + \varphi'''(v)v_r^2 + \frac{\varphi''(v)}{r}v_r - \frac{\varphi'(1)}{r^2} \quad (3.6)$$

in the subset of $(r_1, r_2) \times [0, T]$ where $|v| \neq 1$. Let us set

$$w(t) := M(0) - \varepsilon - \frac{\varphi'(1)}{r_1^2}t \quad \forall t \in [0, T_0], \quad (3.7)$$

where $\varepsilon > 0$ is small enough so that $w(T_0) > 1$, hence $w(t) > 1$ for every $t \in [0, T_0]$. We claim that

$$M(t) \geq w(t) \quad \forall t \in [0, T_0], \quad (3.8)$$

from which (3.5) follows by letting $\varepsilon \rightarrow 0^+$. Let us prove (3.8) by contradiction. Let us assume that $M(t) < w(t)$ for some $t \in [0, T_0]$, and let us set

$$t_0 := \inf \{t \in [0, T_0] : v(r, t) < w(t) \quad \forall r \in [r_3, r_5]\}.$$

Since $M(0) > w(0)$, we have that $t_0 > 0$. Moreover, due to the continuity of v and w , there exists $r_0 \in [r_3, r_5]$ such that $v(r_0, t_0) = w(t_0)$, and

$$v(r, t_0) - w(t_0) \leq 0 \quad \forall r \in [r_3, r_5].$$

Since subcritical regions don't shrink, we have that $v(r_3, t_0) < 1$ and $v(r_5, t_0) < 1$, while $w(t_0) > 1$. This shows in particular that $r_0 \neq r_3$ and $r_0 \neq r_5$. Now we know that r_0 is a maximum point for the function $r \rightarrow v(r, t_0) - w(t_0)$, and r_0 is contained in the open interval (r_3, r_5) , hence

$$v_r(r_0, t_0) = 0 \quad \text{and} \quad v_{rr}(r_0, t_0) \leq 0. \quad (3.9)$$

Let us consider now time derivatives. Since $v(r_0, t_0) = w(t_0) > 1$, we can use (3.6). Exploiting also (3.7) and (3.9) we obtain that

$$\begin{aligned} v_t(r_0, t_0) - w_t(t_0) &= \varphi''(v(r_0, t_0))v_{rr}(r_0, t_0) - \frac{\varphi'(1)}{r_0^2} + \frac{\varphi'(1)}{r_1^2} \\ &> \varphi''(v(r_0, t_0))v_{rr}(r_0, t_0). \end{aligned}$$

Since $\varphi''(v(r_0, t_0)) \leq 0$, we can conclude that

$$v_t(r_0, t_0) - w_t(t_0) > \varphi''(v(r_0, t_0))v_{rr}(r_0, t_0) \geq 0.$$

This implies that $v(r_0, t) - w(t) > 0$ for every t in a suitable right-hand neighborhood of t_0 , which contradicts the definition of t_0 .

3.4 Proof of Theorem 2.4

Let us set

$$k_1 := n, \quad k_2 := 1, \quad h_1 := n^3, \quad h_2 := -n^2. \quad (3.10)$$

We claim that the conclusions of statement (1) and statement (2) hold true provided that n is large enough.

Statement (1) Let us set for simplicity $v_0(x, y) := |\nabla u_0(x, y)|^2$. Let us assume that

$$v_{0y}(0, 0) > 0. \quad (3.11)$$

Then the implicit function theorem implies that the set $v_0(x, y) > 1$ can be represented, in a neighborhood of $(0, 0)$, as $y > g(x)$, where g is a suitable function defined in a neighborhood of $x = 0$. Such a function satisfies

$$\begin{aligned} g(0) &= 0, & g'(0) &= -\frac{v_{0x}(0, 0)}{v_{0y}(0, 0)}, \\ g''(0) &= -\frac{1}{v_{0y}^3} \{v_{0x}^2 v_{0yy} + v_{0y}^2 v_{0xx} - 2v_{0x} v_{0y} v_{0xy}\}, \end{aligned} \quad (3.12)$$

where in (3.12) all partial derivatives of v_0 are computed in $(0, 0)$. In particular g is convex in a neighborhood of 0 if the right-hand side of (3.12) is positive.

From (2.11) we have that, up to higher order terms,

$$u_{0x}(x, y) = \frac{\sqrt{2}}{2} + 2k_1x + 3h_1x^2, \quad u_{0y}(x, y) = \frac{\sqrt{2}}{2} + 2k_2y + 3h_2y^2,$$

hence

$$v_0(x, y) = 1 + 2\sqrt{2}(k_1x + k_2y) + \left(4k_1^2 + 3\sqrt{2}h_1\right)x^2 + \left(4k_2^2 + 3\sqrt{2}h_2\right)y^2.$$

All the derivatives appearing in (3.11) and (3.12) can be easily computed. It follows that condition (3.11) is equivalent to $k_2 > 0$, while $g''(0) > 0$ if and only if

$$8k_1^2k_2^2 + 3\sqrt{2}(k_1^2h_2 + k_2^2h_1) < 0.$$

Both conditions are satisfied if the values of the parameters are given by (3.10) and n is large enough.

Statement (2) Let us set for simplicity $v(x, y, t) := |\nabla u(x, y, t)|^2$. Thesis is proved if we show that

$$v_t = 2u_x u_{tx} + 2u_y u_{ty} > 0$$

in the point $(x, y, t) = (0, 0, 0)$. We can therefore deduce the value of $v_t(0, 0, 0)$ from the Taylor expansion of $u_0(x, y)$.

In order to compute u_t , we recall that $u_t = \Psi_{1x} + \Psi_{2y}$, where

$$\Psi_1 := \varphi' \left((u_x^2 + u_y^2)^{1/2} \right) \frac{u_x}{(u_x^2 + u_y^2)^{1/2}}, \quad \Psi_2 := \varphi' \left((u_x^2 + u_y^2)^{1/2} \right) \frac{u_y}{(u_x^2 + u_y^2)^{1/2}}.$$

With some computations we obtain that, up to higher order terms,

$$(u_x^2 + u_y^2)^{1/2} = 1 + \sqrt{2}(k_1 x + k_2 y) + \left[k_1^2 + \frac{3\sqrt{2}}{2} h_1 \right] x^2 + \left[k_2^2 + \frac{3\sqrt{2}}{2} h_2 \right] y^2 - 2k_1 k_2 x y,$$

hence

$$\varphi' \left((u_x^2 + u_y^2)^{1/2} \right) = \varphi'(1) + (k_1 x + k_2 y)^2 \varphi'''(1),$$

and therefore

$$\begin{aligned} \Psi_1 &= \frac{\sqrt{2}}{2} \varphi'(1) + \varphi'(1) (k_1 x - k_2 y) + \frac{1}{2} \left[3\varphi'(1) (h_1 - \sqrt{2} k_1^2) + \sqrt{2} \varphi'''(1) k_1^2 \right] x^2 \\ &\quad + \frac{1}{2} \left[\varphi'(1) (\sqrt{2} k_2^2 - 3h_2) + \sqrt{2} \varphi'''(1) k_2^2 \right] y^2 + \sqrt{2} (\varphi'(1) + \varphi'''(1)) k_1 k_2 x y. \end{aligned}$$

The expression for Ψ_2 is just the symmetric one. It follows that, up to higher order terms,

$$\begin{aligned} u_t(x, y, 0) &= (k_1 + k_2) \varphi'(1) \\ &\quad + \left\{ \left(3h_1 + \sqrt{2} k_1 (k_2 - 3k_1) \right) \varphi'(1) + \sqrt{2} \varphi'''(1) k_1 (k_1 + k_2) \right\} x \\ &\quad + \left\{ \left(3h_2 + \sqrt{2} k_2 (k_1 - 3k_2) \right) \varphi'(1) + \sqrt{2} \varphi'''(1) k_2 (k_1 + k_2) \right\} y, \end{aligned}$$

hence

$$v_t(0, 0, 0) = \varphi'(1) \left\{ 3\sqrt{2}(h_1 + h_2) + 4k_1 k_2 - 6k_1^2 - 6k_2^2 \right\} + 2\varphi'''(1)(k_1 + k_2)^2.$$

From this expression it is easy to see that $v_t(0, 0, 0) > 0$ if the values of the parameters are given by (3.10) and n is large enough.

3.5 Proof of Theorem 2.5

Let $x_\star \in (x_3, x_4)$ be any point. We have to prove that

$$v(x_\star, t) > 0 \quad \forall t \in [0, T]. \quad (3.13)$$

To this end we fix some notation. First of all we choose real numbers x_5 and x_6 such that $x_3 < x_5 < x_\star < x_6 < x_4$. Then we consider the functions

$$\psi(x) := (x - x_5)(x_6 - x), \quad w(x, t) := e^{-\lambda t} (\delta^2 \psi(x) + \delta \psi^2(x)),$$

where λ and δ are positive parameters. We claim that when λ is large enough and δ is small enough we have that

$$v(x, t) \geq w(x, t) \quad \forall (x, t) \in [x_5, x_6] \times [0, T]. \quad (3.14)$$

Since w is positive in $(x_5, x_6) \times [0, T)$, and $x_\star \in (x_5, x_6)$, this is enough to prove (3.13). In order to prove (3.14) we first establish some properties of w .

Properties of w Let us show that w fulfils the following properties.

- (w1) $w \in C^\infty([x_5, x_6] \times [0, +\infty))$;
- (w2) $w(x, t) > 0$ for every $(x, t) \in (x_5, x_6) \times [0, +\infty)$;
- (w3) $w_x(x_5, t) = \delta^2 e^{-\lambda t} \psi'(x_5) > 0$ for every $t \geq 0$;
- (w4) $w_x(x_6, t) = \delta^2 e^{-\lambda t} \psi'(x_6) < 0$ for every $t \geq 0$;
- (w5) if δ is small enough we have that $w(x, 0) < v(x, 0)$ for every $x \in [x_5, x_6]$;
- (w6) if δ is small enough we have that $w(x, t) < c_0$ for every $(x, t) \in (x_5, x_6) \times [0, +\infty)$;
- (w7) if δ is small enough and λ is large enough, then w satisfies

$$w_t(x, t) < g(w(x, t))w_{xx}(x, t) \quad \forall (x, t) \in (x_5, x_6) \times [0, +\infty). \quad (3.15)$$

Properties (w1) through (w4) easily follow from the definition of w and ψ . Property (w5) follows from the fact that the infimum of $v(x, 0)$ for $x \in [x_5, x_6]$ is strictly positive due to (v3). Property (w6) is almost trivial. In order to prove (w7) we recall that $\psi''(x) = -2$, hence

$$w_t(x, t) = -\lambda e^{-\lambda t} (\delta^2 \psi(x) + \delta \psi^2(x)), \quad w_{xx}(x, t) = e^{-\lambda t} (2\delta[\psi'(x)]^2 - 4\delta\psi(x) - 2\delta^2).$$

Plugging these identities in (3.15), we are left to prove that

$$-\lambda (\delta\psi + \psi^2) < g(w) (2[\psi']^2 - 4\psi - 2\delta) \quad \forall (x, t) \in (x_5, x_6) \times [0, +\infty). \quad (3.16)$$

To this end we fix once for all two real numbers x_7 and x_8 such that $x_5 < x_7 < x_8 < x_6$, and

$$\inf \{2[\psi'(x)]^2 - 4\psi(x) : x \in [x_5, x_7] \cup [x_8, x_6]\} > 0.$$

This is possible because in the endpoints of the interval $[x_5, x_6]$ one has that $\psi = 0$ and $\psi' \neq 0$. Now we distinguish two cases. When $x \in [x_5, x_7] \cup [x_8, x_6]$ the left-hand side of (3.16) is negative, while the left-hand side is positive provided that δ is small enough, independently on x . When $x \in [x_7, x_8]$ the right-hand side may be negative, but also the left-hand side is strictly negative because in this interval ψ is bounded from below by a positive constant. In other words, in $[x_7, x_8] \times [0, +\infty)$ inequality (3.16) holds true if we choose

$$\lambda > \sup \left\{ \frac{g(w(x, t)) (4\psi(x) + 2\delta)}{\delta\psi(x) + \psi^2(x)} : (x, t) \in [x_7, x_8] \times [0, +\infty) \right\}.$$

We point out that the supremum is finite. This completes the proof of (3.16).

Proof of (3.14) Let us choose positive values of δ and λ in such a way that w satisfies (w1) through (w7). Now we argue more or less as in the proof of the classical comparison results. Let us assume that (3.14) is not true, and let us set

$$t_0 := \inf \{t \in [0, T) : \exists x \in [x_5, x_6] \text{ such that } v(x, t) - w(x, t) < 0\}.$$

From (w5) we have that $t_0 > 0$. Moreover, from the definition of t_0 it follows that

$$v(x, t) - w(x, t) \geq 0 \quad \forall (x, t) \in [x_5, x_6] \times [0, t_0]. \quad (3.17)$$

Finally, from the continuity of v and w we deduce that there exists $x_0 \in [x_5, x_6]$ such that $v(x_0, t_0) - w(x_0, t_0) = 0$.

We claim that $x_0 \neq x_5$. Indeed let us assume by contradiction that $x_0 = x_5$. Then $w(x_0, t_0) = 0$, hence also $v(x_0, t_0) = 0$. By (v2) it follows that x_0 is a minimum point for the function $x \rightarrow v(x, t_0)$. By (v4) we have therefore that $v_x(x_0, t_0) = 0$. Keeping (w3) into account, we deduce that

$$(v - w)_x(x_0, t_0) = v_x(x_0, t_0) - w_x(x_0, t_0) < 0. \quad (3.18)$$

On the other hand, from (3.17) we know also that $x_0 = x_5$ is a minimum point for the function $x \rightarrow v(x, t_0) - w(x, t_0)$ restricted to the interval $[x_5, x_6]$. Since the minimum point is the left-hand endpoint of the interval, we deduce that $(v - w)_x(x_0, t_0) \geq 0$, which contradicts (3.18).

In a symmetric way we prove that $x_0 \neq x_6$. So we are left with the case in which $t_0 > 0$ and $x_0 \in (x_5, x_6)$. In this case by (w2) and (w6) we have that $0 < v(x_0, t_0) = w(x_0, t_0) < c_0$, hence both v and w are smooth in a neighborhood of this point and fulfil (v5) and (w7), respectively. In particular, since x_0 is always a minimum point of the

function $x \rightarrow v(x, t_0) - w(x, t_0)$, and now x_0 is in the interior of the interval (x_5, x_6) , we have that

$$v_x(x_0, t_0) = w_x(x_0, t_0) \quad \text{and} \quad v_{xx}(x_0, t_0) \geq w_{xx}(x_0, t_0). \quad (3.19)$$

Let us consider now the time derivatives. On the one hand, from (3.17) we deduce that $(v - w)_t(x_0, t_0) \leq 0$. On the other hand, from (v5), (w7), and (3.19) we have that

$$\begin{aligned} (v - w)_t(x_0, t_0) &> g(v(x_0, t_0))v_{xx}(x_0, t_0) - g(w(x_0, t_0))w_{xx}(x_0, t_0) \\ &= g(w(x_0, t_0)) (v_{xx}(x_0, t_0) - w_{xx}(x_0, t_0)) \\ &\geq 0. \end{aligned}$$

This rules out the last case and completes the proof of (3.14).

3.6 Proof of Theorem 2.6

The strategy is similar to the proof of Theorem 2.5. The main difference is that in this case we have to cope with moving domains.

Let (r_*, t_*) be any point of \mathcal{D} . We have to prove that

$$v(r_*, t_*) > 0. \quad (3.20)$$

To this end we fix some notation. First of all it is not difficult to see that there exists real numbers r_5, r_6, k such that

$$|k| < G\sqrt{A}, \quad r_1 < r_5 + kt_* < r_* < r_6 + kt_* < r_2.$$

Then we consider the set

$$\mathcal{D}_* := \{(r, t) \in (r_1, r_2) \times [0, T) : t \leq t_*, r_5 + kt \leq r \leq r_6 + kt\} \subseteq \mathcal{D}.$$

We refer to Figure 2 for a representation of the set \mathcal{D}_* (corresponding in that case to some $k < 0$) and its relation with (r_*, t_*) and \mathcal{D} . Note that the slope of the slanted lines bounding \mathcal{D}_* is larger than the slope of the slanted lines limiting \mathcal{D} . This is just because $|k| < G\sqrt{A}$.

Due to this inequality, there exists $\varepsilon_0 \in (0, \min\{1, G, A/2\})$ such that

$$|k| < (1 - \varepsilon_0)(G - \varepsilon_0)\sqrt{A - 2\varepsilon_0}. \quad (3.21)$$

From now on, ε_0 is a fixed positive constant. Due to (2.12) there exists also $c_2 \in (0, c_0)$ such that

$$g(\sigma) \geq (G - \varepsilon_0)\sqrt{\sigma} \quad \forall \sigma \in (0, c_2). \quad (3.22)$$

Finally we consider the functions

$$\psi(r) := (r - r_5)(r_6 - r), \quad w(r, t) := \delta^3 \psi(r - kt) + \delta[\psi(r - kt)]^{3/2},$$




Figure 2: the sets \mathcal{D} and \mathcal{D}_*

where δ is a positive parameter. We claim that when δ is small enough we have that

$$v(r, t) \geq w(r, t) \quad \forall (r, t) \in \mathcal{D}_*. \quad (3.23)$$

This inequality, applied with $(r, t) = (r_*, t_*)$, implies (3.20). In order to prove (3.23) we first establish some properties of w .

Properties of w Let $\text{Int}(\mathcal{D}_*)$ denote the set of points $(r, t) \in \mathcal{D}_*$ with $r_5 + kt < r < r_6 + kt$. Let us show that w fulfils the following properties.

- (w1) $w \in C^1(\mathcal{D}_*) \cap C^\infty(\text{Int}(\mathcal{D}_*))$;
- (w2) $w(r, t) > 0$ for every $(r, t) \in \text{Int}(\mathcal{D}_*)$;
- (w3) $w_r(r_5 + kt, t) = \delta^3 \psi'(r_5) > 0$ for every $t \in [0, t_*]$;
- (w4) $w_r(r_6 + kt, t) = \delta^3 \psi'(r_6) < 0$ for every $t \in [0, t_*]$;
- (w5) if δ is small enough we have that $w(r, 0) < v(r, 0)$ for every $r \in [r_5, r_6]$;
- (w6) if δ is small enough we have that $w(r, t) < c_2$ for every $(r, t) \in \mathcal{D}_*$;
- (w7) if δ is small enough, then w satisfies

$$w_t < g(w) \{w_{rr} + f(r, t, w, w_r) + A\} \quad \forall (r, t) \in \text{Int}(\mathcal{D}_*). \quad (3.24)$$

Properties (w1) through (w4) easily follow from the definition of w and ψ . Property (w5) follows from the fact that the infimum of $v(r, 0)$ for $r \in [r_5, r_6]$ is strictly positive due to (v3). Property (w6) is almost trivial. In order to prove (w7) we recall that $\psi''(r) = -2$, hence (for simplicity we set $y := r - kt$, and we observe that $y \in [r_5, r_6]$)

$$w_t(r, t) = -k\delta^3 \psi'(y) - \frac{3}{2}k\delta [\psi(y)]^{1/2} \psi'(y),$$

$$w_r(r, t) = \delta^3 \psi'(y) + \frac{3}{2} \delta [\psi(y)]^{1/2} \psi'(y),$$

$$w_{rr}(r, t) = -2\delta^3 - 3\delta [\psi(y)]^{1/2} + \frac{3}{4} \delta [\psi(y)]^{-1/2} [\psi'(y)]^2.$$

When $\delta \rightarrow 0^+$ we have that w and w_r tend to zero uniformly in \mathcal{D}_* . Thanks to (2.13) we have therefore that

$$|f(r, t, w(r, t), w_r(r, t))| \leq \varepsilon_0 \quad \forall (r, t) \in \mathcal{D}_*,$$

provided that δ is small enough. In an analogous way we have also that

$$w_{rr}(r, t) > \frac{3}{4} \delta [\psi(y)]^{-1/2} [\psi'(y)]^2 - \varepsilon_0$$

provided that δ is small enough. It follows that

$$w_{rr} + f(r, t, w, w_r) + A > A - 2\varepsilon_0 + \frac{3}{4} \delta [\psi(y)]^{-1/2} [\psi'(y)]^2 > 0$$

in $\text{Int}(\mathcal{D}_*)$. Moreover, from (3.22), (w2) and (w6) we have that

$$g(w(r, t)) \geq (G - \varepsilon_0) \sqrt{w(r, t)} \quad \forall (r, t) \in \text{Int}(\mathcal{D}_*),$$

and in conclusion

$$g(w) \{w_{rr} + f(r, t, w, w_r) + A\} > (G - \varepsilon_0) \sqrt{w} \left\{ A - 2\varepsilon_0 + \frac{3}{4} \delta \frac{[\psi'(y)]^2}{[\psi(y)]^{1/2}} \right\}.$$

Therefore inequality (3.24) is proved if we show that

$$\delta^3 |k\psi'| + \frac{3}{2} \delta |k\psi'| \sqrt{\psi} \leq (G - \varepsilon_0) \sqrt{\delta^3 \psi + \delta [\psi]^{3/2}} \left\{ A - 2\varepsilon_0 + \frac{3}{4} \delta \frac{[\psi']^2}{\sqrt{\psi}} \right\}, \quad (3.25)$$

where the argument of ψ and ψ' is any $y \in (r_5, r_6)$. Let us consider the right-hand side of (3.25) multiplied by $(1 - \varepsilon_0)$. Applying the inequality between the arithmetic mean and the geometric mean, and recalling (3.21), we obtain that

$$\begin{aligned} (1 - \varepsilon_0) \cdot (\text{right-hand side}) &\geq (1 - \varepsilon_0)(G - \varepsilon_0) \sqrt{\delta [\psi]^{3/2}} \cdot \left\{ A - 2\varepsilon_0 + \frac{3}{4} \delta \frac{[\psi']^2}{\sqrt{\psi}} \right\} \\ &\geq (1 - \varepsilon_0)(G - \varepsilon_0) \sqrt{\delta [\psi]^{3/2}} \cdot 2 \left[(A - 2\varepsilon_0) \cdot \frac{3}{4} \delta \frac{[\psi']^2}{\sqrt{\psi}} \right]^{1/2} \\ &= (1 - \varepsilon_0)(G - \varepsilon_0) \sqrt{A - 2\varepsilon_0} \cdot \delta |\psi'| \sqrt{\psi} \cdot \sqrt{3} \\ &\geq \frac{3}{2} \delta |k\psi'| \sqrt{\psi}. \end{aligned}$$

In order to prove (3.25) it is therefore enough to show that

$$\delta^3 |k\psi'| \leq \varepsilon_0(G - \varepsilon_0) \sqrt{\delta^3 \psi + \delta[\psi]^{3/2}} \left\{ A - 2\varepsilon_0 + \frac{3}{4} \delta \frac{[\psi']^2}{\sqrt{\psi}} \right\},$$

which in turn is true if we show that

$$\delta^{3/2} |k\psi'| \leq \varepsilon_0(G - \varepsilon_0) \sqrt{\psi} \left\{ A - 2\varepsilon_0 + \frac{3}{4} \delta \frac{[\psi']^2}{\sqrt{\psi}} \right\}.$$

To this end, we fix once for all two real numbers r_7 and r_8 such that

$$r_5 < r_7 < \frac{r_5 + r_6}{2} < r_8 < r_6.$$

When $y \in (r_5, r_7] \cup [r_8, r_6)$ we have that $|\psi'(y)|$ is bounded from below by a positive constant. Therefore

$$\delta^{3/2} |k\psi'(y)| \leq \varepsilon_0(G - \varepsilon_0) \frac{3}{4} \delta [\psi'(y)]^2$$

provided that δ is small enough. When $y \in [r_7, r_8]$ we have that $\psi(y)$ is bounded from below by a positive constant, hence in this case

$$\delta^{3/2} |k\psi'(y)| \leq \varepsilon_0(G - \varepsilon_0)(A - 2\varepsilon_0) \sqrt{\psi(y)}$$

provided that δ is small enough. This completes the proof of (3.25) and shows that w satisfies (w7) whenever δ is small enough.

Proof of (3.23) The argument is analogous to the proof of the corresponding inequality in Theorem 2.5. Let us choose a positive value of δ in such a way that w satisfies (w1) through (w7). Let us assume that (3.23) is not true, and let us set

$$t_0 := \inf \{t \in [0, t_*] : \exists r \in [r_5 + kt, r_6 + kt] \text{ such that } v(r, t) - w(r, t) < 0\}.$$

From (w5) we have that $t_0 > 0$. Moreover, from the definition of t_0 we have that

$$v(r, t) - w(r, t) \geq 0 \quad \forall (r, t) \in \mathcal{D}_* \text{ with } t \leq t_0. \quad (3.26)$$

Finally, due to the continuity of v and w , we deduce also that there exists $r_0 \in [r_5 + kt_0, r_6 + kt_0]$ such that $v(r_0, t_0) - w(r_0, t_0) = 0$.

We claim that $r_0 \neq r_5 + kt_0$. Indeed let us assume by contradiction that $r_0 = r_5 + kt_0$. Then $w(r_0, t_0) = 0$, hence also $v(r_0, t_0) = 0$. By (v2) it follows that r_0 is a minimum point for the function $r \rightarrow v(r, t_0)$. By (v4) we have therefore that $v_r(r_0, t_0) = 0$. Keeping (w3) into account, we deduce that

$$(v - w)_r(r_0, t_0) = v_r(r_0, t_0) - w_r(r_0, t_0) < 0. \quad (3.27)$$

On the other hand, from (3.26) we know also that $r_0 = r_5 + kt_0$ is a minimum point for the function $r \rightarrow v(r, t_0) - w(r, t_0)$ restricted to the interval $[r_5 + kt_0, r_6 + kt_0]$. Since the minimum point is the left-hand endpoint of the interval, we deduce that $(v - w)_r(r_0, t_0) \geq 0$, which contradicts (3.27).

In a symmetric way we prove that $r_0 \neq r_6 + kt_0$. So we are left with the case in which $t_0 > 0$ and $r_0 \in (r_5 + kt_0, r_6 + kt_0)$. In this case $0 < v(r_0, t_0) = w(r_0, t_0) < c_2$, hence both v and w are smooth in a neighborhood of this point and fulfil (v5) and (w7), respectively. In particular, since r_0 is always a minimum point of the function $r \rightarrow v(r, t_0) - w(r, t_0)$, and now r_0 is in the interior of the interval $(r_5 + kt_0, r_6 + kt_0)$, we have that

$$v_r(r_0, t_0) = w_r(r_0, t_0) \quad \text{and} \quad v_{rr}(r_0, t_0) \geq w_{rr}(r_0, t_0). \quad (3.28)$$

Let us consider now the time derivatives. On the one hand, from (3.26) we deduce that $(v - w)_t(r_0, t_0) \leq 0$. On the other hand, from (v5), (w7), and (3.28), in the point (r_0, t_0) we have that

$$\begin{aligned} (v - w)_t &> g(v) \{v_{rr} + f(r, t, v, v_r) + A\} - g(w) \{w_{rr} + f(r, t, w, w_r) + A\} \\ &= g(w) \{v_{rr} + f(r, t, w, w_r) + A\} - g(w) \{w_{rr} + f(r, t, w, w_r) + A\} \\ &= g(w) (v_{rr} - w_{rr}) \geq 0. \end{aligned}$$

This rules out the last case and completes the proof of (3.23).

References

- [1] H. AMANN; Time-delayed Perona-Malik type problems. *Acta Math. Univ. Comenian. (N.S.)* **76** (2007), no. 1, 15–38.
- [2] G. BELLETTINI, G. FUSCO; The Γ -limit and the related gradient flow for singular perturbation functionals of Perona-Malik type. *Trans. Amer. Math. Soc.* **360** (2008), no. 9, 4929–4987.
- [3] G. BELLETTINI, M. NOVAGA, E. PAOLINI; Global solutions to the gradient flow equation of a nonconvex functional. *SIAM J. Math. Anal.* **37** (2006), no. 5, 1657–1687.
- [4] G. BELLETTINI, M. NOVAGA, M. PAOLINI, C. TORNESE; Convergence of discrete schemes for the Perona-Malik equation. *J. Differential Equations* **245** (2008), no. 4, 892–924.
- [5] F. CATTÉ, P. L. LIONS, J. M. MOREL, T. COLL; Image selective smoothing and edge detection by nonlinear diffusion. *SIAM J. Numer. Anal.* **29** (1992), no. 1, 182–193.

- [6] Y. CHEN, K. ZHANG; Young measure solutions of the two-dimensional Perona-Malik equation in image processing. *Commun. Pure Appl. Anal.* **5** (2006), no. 3, 615–635.
- [7] S. ESEDOGLU; Stability properties of the Perona-Malik scheme. *SIAM J. Numer. Anal.* **44** (2006), no. 3, 1297–1313.
- [8] S. ESEDOGLU, J. B. GREER; Upper bounds on the coarsening rate of discrete, ill-posed nonlinear diffusion equations. *Comm. Pure Appl. Math.* **62** (2009), no. 1, 57–81.
- [9] M. GHISI, M. GOBBINO; Gradient estimates for the Perona-Malik equation. *Math. Ann.* **337** (2007), no. 3, 557–590.
- [10] M. GHISI, M. GOBBINO; A class of local classical solutions for the one-dimensional Perona-Malik equation. *Trans. Amer. Math. Soc.* **361** (2009), no. 12, 6429–6446.
- [11] M. GHISI, M. GOBBINO; An example of global transcritical solution for the Perona-Malik equation. Preprint. [arXiv:0907.0772](https://arxiv.org/abs/0907.0772) [math.AP]
- [12] M. GOBBINO; Entire solutions of the one-dimensional Perona-Malik equation. *Comm. Partial Differential Equations* **32** (2007), no. 4-6, 719–743.
- [13] P. GUIDOTTI; A new nonlocal nonlinear diffusion of image processing, *J. Differential Equations* **246** (2009), no. 12, 4731-4742.
- [14] P. GUIDOTTI, J. V. LAMBERS; Two new nonlinear nonlocal diffusions for noise reduction. *J. Math. Imaging Vision* **33** (2009), no. 1, 25–37.
- [15] B. KAWOHL, N. KUTEV; Maximum and comparison principle for one-dimensional anisotropic diffusion. *Math. Ann.* **311** (1998), no. 1, 107–123.
- [16] S. KICHENASSAMY; The Perona-Malik paradox. *SIAM J. Appl. Math.* **57** (1997), no. 5, 1328–1342.
- [17] P. PERONA, J. MALIK; Scale space and edge detection using anisotropic diffusion, *IEEE Trans. Pattern Anal. Mach. Intell.* **12** (1990), 629–639.
- [18] S. TAHERI, Q. TANG, K. ZHANG; Young measure solutions and instability of the one-dimensional Perona-Malik equation. *J. Math. Anal. Appl.* **308** (2005), no. 2, 467–490.
- [19] K. ZHANG; Existence of infinitely many solutions for the one-dimensional Perona-Malik model. *Calc. Var. Partial Differential Equations* **26** (2006), no. 2, 171–199.